

# A coherent categorification of the based ring of the lowest two-sided cell

Stefan Dawydiak \*

December 1, 2021

## Abstract

We give a coherent categorification of  $J_0$ , the based ring of the lowest two sided cell of an affine Weyl group, equipped with a monoidal functor from the category of coherent sheaves on the derived Steinberg variety. We show that our categorification acts on natural coherent categorifications of the Iwahori invariants of the Schwartz space of the basic affine space. In low rank cases, we construct complexes that lift the basis elements  $t_w$  of  $J_0$  and their structure constants.

**Keywords**— Asymptotic Hecke algebra, Iwahori-Hecke algebra, flag variety, Steinberg variety

## 1 Introduction

Let  $\tilde{W}$  be an affine Weyl group. Its group algebra  $\mathbb{C}[\tilde{W}]$  is deformed by the affine Hecke algebra  $H_{\text{aff}} = H(\tilde{W})$  of  $\tilde{W}$ . In turn, Lusztig defined the asymptotic Hecke algebra  $J$ , a based ring with basis  $t_w$ ,  $w \in \tilde{W}$  and structure constants determined from certain “leading terms” of the structure constants of  $H_{\text{aff}}$ . Further, he provided a morphism of algebras  $\phi: H_{\text{aff}} \hookrightarrow J \otimes_{\mathbb{Z}} \mathbb{Z}[\mathbf{q}^{\pm 1/2}]$  and showed it was an algebra after a mild completion. Thus  $H_{\text{aff}}$  can be viewed as a subalgebra of  $J$ , and  $J$  can be viewed as a subalgebra of a completion  $\mathcal{H}_{\text{aff}}$  of  $H_{\text{aff}}$ . While the morphism  $\phi$  is an essential part of Lusztig’s exploration of  $J$ , until recently there have been few compelling reasons to adopt the perspective of  $J \otimes_{\mathbb{Z}} \mathbb{Z}[\mathbf{q}^{\pm 1/2}]$  as a subalgebra of  $\mathcal{H}_{\text{aff}}$ .

The algebra  $H_{\text{aff}}$  appears in many areas of mathematics in many guises, but one of the most prominent relates to the representation theory of  $p$ -adic groups. Let  $F$  be a local non-archimedean field and  $q$  be the cardinality of the residue field of  $F$ . Let  $\mathbf{G}^{\vee}$  be a connected reductive algebraic group defined and split over  $F$ , with Langlands dual group  $\mathbf{G}$  taken over  $\mathbb{C}$  with  $G = \mathbf{G}(\mathbb{C})$ . For the purposes of harmonic analysis on  $\mathbf{G}^{\vee}(F)$ , it is natural to consider  $H_{\text{aff}}$ , very much an algebraic object, as a subalgebra of the larger, analytically-characterized Harish-Chandra Schwartz algebra  $\mathcal{C}(G^{\vee})^I$ .

In [BK18], Braverman and Kazhdan gave an interpretation of  $J$  in terms of harmonic analysis, casting  $J$  as an algebraic version of  $\mathcal{C}(G^{\vee})^I$  (they also defined a ring  $\mathcal{J}$  doing the same for the full algebra  $\mathcal{C}(G^{\vee})$ ) by defining an map  $J \rightarrow \mathcal{C}(G^{\vee})^I$ . In [Daw21], the author showed that this morphism was essentially the specialization of  $\phi^{-1}$  for  $\mathbf{q} = q$ , and in particular was an injection.

Lusztig gave a categorification of  $J$  in [Lus97] in terms of perverse sheaves on the affine flag variety. In the spirit of the definition of  $J$  as a ring, the underlying category is again  $\text{Perv}(\mathcal{F}l)$ , but with monoidal structure given by *truncated convolution* as opposed to convolution. In this paper we provide a categorification of a large direct summand  $J_0$  of  $J$  that is compatible with the perspective of [BK18]. Namely, we obtain a natural categorification of the action of  $J_0$  on the unitary principal series, and produce a completely new category whose  $K$ -theory is  $J_0$ , as opposed a new monoidal structure. Our main result is

---

\*Department of Mathematics, University of Toronto, Toronto, ON M5S 2E4 Canada; email [stefand@math.utoronto.ca](mailto:stefand@math.utoronto.ca)

**Theorem 1.** *Let  $\mathcal{B}$  be the derived zero section of the Springer resolution  $\tilde{\mathcal{N}} \rightarrow \mathcal{N}$ . Then*

1. *the category*

$$\mathcal{J}_0 := D^b \text{Coh}_G(\mathcal{B} \times \mathcal{B})_!$$

*has a monoidal structure given by convolution, and admits a natural monoidal functor*

$$D^b \text{Coh}_{G \times \mathbb{C}^\times}(\text{St}) \rightarrow D^b \text{Coh}_{G \times \mathbb{C}^\times}(\mathcal{B} \times \mathcal{B})_!$$

*such that*

2. *the induced morphism*

$$H_{\text{aff}} \rightarrow K_0(D^b \text{Coh}_{G \times \mathbb{C}^\times}(\mathcal{B} \times \mathcal{B})_!) \rightarrow K_{G \times \mathbb{C}^\times}(\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit)$$

*is conjugate to  $\phi_0$ ;*

3. *In the special case when  $G$  has universal cover equal to  $\text{SL}_2$  or  $\text{SL}_3$ , there exists a family of objects  $\{t_w\}_{w \in \mathfrak{c}_0}$  in  $\mathcal{J}_0$ , such that if  $t_w t_x = \sum_z \gamma_{w,x,z^{-1}} t_z$  in  $J_0$ , then*

$$t_w \star t_x = \bigoplus_z t_z^{\oplus \gamma_{w,x,z^{-1}}}$$

*in  $\mathcal{J}_0$  and such that the image in  $K_G(\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit)$  of the class  $[t_w]$  under the above morphism is  $[t_w]$ .*

4. *The category  $\mathcal{J}_0$  acts on  $\text{Coh}_T(\mathcal{B})$  and on  $\text{Coh}_G(\mathcal{B} \times \mathcal{B})$ ;*

*Proof.* The four propositions below each prove one statement of the theorem. □

The algebra  $J$  is very close to being a direct sum of matrix algebras. In type  $A$ , this is the main result of Xi's monograph [Xi02], and Bezrukavnikov-Ostrik in [BO04] showed this up to central extensions in general. (It is however now known [BDD21] that this partial result is sharp; the central extensions do in fact appear in general.)

Therefore the last two items are particularly relevant:  $J$  is most interesting as a based algebra admitting a morphism from  $H_{\text{aff}}$ , and can be quite simple in isolation. Our categorification captures the failure of  $\phi_0$  to be surjective, as explained in Remark 2 and the discussion preceeding it. We hope to remove the very restrictive current hypothesis on item 3 in a future version of this paper; see remark 1 for an explanation of why it is currently necessary.

## 1.1 Acknowledgements

The author thanks Alexander Braverman for introducing him to the notion of singular support and its consequences for convolution, Dylan Butson and Kostya Tolmachov for many patient and helpful conversations about the basics of derived algebraic geometry and the coherent Hecke category, and Andy Ramirez-Coté for a helpful discussion. This research was supported by NSERC.

## 2 Functions and algebras

In this section we will recall the various algebras whose categorifications we will discuss in Section 3. There is no new material in this section, although we could not find a recollection of all the relationships below in one place in the existing literature.

## 2.1 The affine Hecke algebra

Let  $\mathbf{G}$  be a connected simply-connected reductive group defined over  $\mathbb{C}$ . Let  $X^*$  be the character lattice of  $\mathbf{G}$ , and  $\tilde{W} = W \ltimes X^*$ , where  $W$  is the finite Weyl group of  $\mathbf{G}$ . Let  $H = H_{\text{aff}}$  be the corresponding affine Hecke algebra over  $\mathcal{A} = \mathbb{Z}[\mathbf{q}^{1/2}, \mathbf{q}^{-1/2}]$  with standard basis  $\{T_w\}_{w \in \tilde{W}}$ . The multiplication in this, the Coxeter presentation, of  $H$  is determined by  $T_w T_{w'} = T_{ww'}$  when  $\ell(ww') = \ell(w) + \ell(w')$  and the quadratic relation  $(T_s + 1)(T_s - \mathbf{q}) = 0$  for all  $s \in S$ , where  $S \subset \tilde{W}$  is the set of simple reflections. The Coxeter presentation is well-suited to studying the action of  $H$  on admissible representations, and to the constructible categorification of  $H$ .

There is a second presentation of  $H$ , due to Bernstein (and Bernstein-Zelevinskii in type  $A$ ), which appears naturally in coherent descriptions of  $H$ , both on the level of  $K$ -theory and the level of categories.

**Definition 1.** The *Bernstein presentation* of  $H$  is the presentation with basis  $\{T_w \theta_\lambda\}_{w \in W, \lambda \in X^*}$ , where

- For  $w \in W$ ,  $T_w$  is the same basis element as in the Coxeter presentation.
- If  $\lambda \in \tilde{W}$  is an antidominant character, and hence corresponds to a *geometrically* dominant cocharacter in the sense of [CG97], then

$$\theta_\lambda = \mathbf{q}^{-\frac{\ell(\lambda)}{2}} T_\lambda.$$

- If  $\lambda \in \tilde{W}$  is a dominant character, and hence corresponds to an *geometrically* antidominant cocharacter, then

$$\theta_\lambda = \mathbf{q}^{\frac{\ell(\lambda)}{2}} T_\lambda^{-1}.$$

The sets  $\{\theta_\lambda, \theta_\lambda T_{s_0}\}_{\lambda \in X^*}$  and  $\{\theta_\lambda, T_s \theta_\lambda\}_{\lambda \in X^*}$  are each  $\mathcal{A}$ -bases. The relations are as follows:

- The same quadratic relation for  $T_{s_0}$ ;
- For any cocharacters  $\lambda, \lambda'$ , we have  $\theta_\lambda \theta_{\lambda'} = \theta_{\lambda + \lambda'}$ .
- The Bernstein relation

$$\theta_{\frac{\alpha}{2}} T_{s_\alpha^\vee} - T_{s_\alpha^\vee} \theta_{-\frac{\alpha}{2}} = (q - 1) \frac{\theta_{\frac{\alpha}{2}} - \theta_{-\frac{\alpha}{2}}}{1 - \theta_{-\alpha}} = (q - 1) \theta_{\frac{\alpha}{2}}.$$

*Example 1.* Let  $\mathbf{G}^\vee = \text{PGL}_2$ , so that  $\tilde{W} = \mathfrak{S}_2 \ltimes X$ , where  $X$  is the character lattice of  $\mathbf{G} = \text{SL}_2$ . Write  $s_0$  for the finite simple reflection, and  $s_1$  for the affine simple reflection in  $\tilde{W}$ . Then the generators of the Bernstein subalgebra are as follows:

1. If  $\lambda = -n\alpha^\vee = (s_1 s_0)^n \in \tilde{W}$ , and hence corresponds to a *geometrically* dominant cocharacter in the sense of [CG97], then

$$\theta_{(s_1 s_0)^n} = \theta_{-n\alpha^\vee} = \mathbf{q}^{-n} T_{(s_1 s_0)^n}.$$

2. If  $\lambda = n\alpha^\vee = (s_0 s_1)^n \in \tilde{W}$ , and hence corresponds to an *geometrically* antidominant cocharacter, then

$$\theta_{(s_0 s_1)^n} = \theta_{n\alpha^\vee} = \mathbf{q}^n T_{(s_0 s_1)^n}^{-1}.$$

In particular, under the geometric choice of dominance, we have  $\rho = -1$ .

## 2.2 The asymptotic Hecke algebra

**Definition 2.** *Lusztig's  $a$ -function*  $a: \tilde{W} \rightarrow \mathbb{N}$  is defined such that  $a(w)$  is minimal such that  $q^{\frac{a(w)}{2}} h_{x,y,w} \in \mathcal{A}^+$  for all  $x, y \in \tilde{W}$ .

It is known that  $a$  is constant on two-sided cells of  $\tilde{W}$  and that

$$a(\mathbf{c}) = \dim \mathcal{B}_u$$

where  $u$  is the unipotent conjugacy class in  $G$  corresponding to  $\mathbf{c}$  under Lusztig's bijection. It is also known that  $a(w) \leq \ell(w)$  for all  $w \in \tilde{W}$ .

In [Lus87] Lusztig defined an associative algebra  $J$  over  $\mathbb{Z}$  equipped with an injection  $\phi: H \hookrightarrow J \otimes_{\mathbb{Z}} \mathcal{A}$  which becomes an isomorphism after taking a certain completion of both sides. As an abelian group,  $J$  has a basis  $\{t_w\}_{w \in W}$ . Recalling the Kazhdan-Lusztig basis elements

$$C_w = \sum_{y \leq w} (-1)^{\ell(w) - \ell(y)} \mathbf{q}^{\frac{\ell(w)}{2} - \ell(y)} P_{y,w}(\mathbf{q}^{-1}) T_y,$$

the structure constants of  $J$  are obtained from those in  $H$  written in the  $\{C_w\}_{w \in W}$ -basis under the following procedure. Taking the structure constants

$$C_x C_y = \sum_{z \in W} h_{x,y,z} C_z$$

for  $h_{x,y,z} \in \mathcal{A}$ , Lusztig then defines the integer  $\gamma_{x,y,z}$  by the condition

$$q^{\frac{a(z)}{2}} h_{x,y,z^{-1}} - \gamma_{x,y,z} \in q\mathcal{A}^+.$$

The product in  $J$  is then defined as

$$t_x t_y = \sum_z \gamma_{x,y,z^{-1}} t_z.$$

One then defines

$$\phi(C_w) = \sum_{\substack{z \in W, d \in \mathcal{D} \\ a(z) = a(d)}} h_{x,d,z} t_z,$$

where  $\mathcal{D} \subset \tilde{W}$  is the set of distinguished involutions. The elements  $t_d$  for distinguished involutions  $d$  are orthogonal idempotents. Moreover,  $J = \bigoplus_{\mathbf{c}} J_{\mathbf{c}}$  is a direct sum of two-sided ideals indexed by two-sided cells  $\mathbf{c} \subset \tilde{W}$ . The unit element in each summand is  $\sum_{\mathcal{D} \cap \mathbf{c}} t_d$ , and the unit element of  $J$  is  $\sum_{d \in \mathcal{D}} t_d$ .

### 2.2.1 The lowest two-sided cell

Let  $\mathbf{c}_0$  be the lowest two-sided cell, also called the “big cell.” It can be characterized by containing the longest element of  $W$ , and we have  $a(\mathbf{c}_0) = \ell(w_0)$ . The summand  $J_0 := J_{\mathbf{c}_0}$  is particularly well-understood, and has historically been the first summand for which any structure-theoretic result is achieved (consider, for example, the progression [Xi90], [Xi02], [BO04]).

By [Xi90] and [Nie11], we have the following description of  $\mathbf{c}_0 \subset \tilde{W}$ . Let  $\tilde{\mathbf{c}}_0$  be the lowest cell of the affine Weyl group of the universal covering group  $\tilde{G}$  of  $G$  with maximal torus  $\tilde{T}$ . Then  $\mathbf{c}_0 = \tilde{\mathbf{c}}_0 \cap \tilde{W}$ , and

$$\tilde{\mathbf{c}}_0 = \left\{ f^{-1} w_0 \chi g \mid f, g \in \Sigma, \chi \in X^*(\tilde{T})^+ \right\},$$

where  $\Sigma = \{w x_w \mid x \in W\} \subset \tilde{W}(\tilde{G})$ , where

$$x_w = w^{-1} \left( \prod_{\substack{\alpha \in \Delta \\ w^{-1}(\alpha) < 0}} \varpi_{\alpha} \right) \in X^*(\tilde{T}), \quad (1)$$

where  $\varpi_{\alpha}$  is the fundamental dominant weight corresponding to  $\alpha$ . We have

$$t_{f^{-1} w_0 \lambda g} t_{(f')^{-1} w_0 \nu g'} = 0$$

if  $g \neq f'$ , and

$$t_{f^{-1}w_0\lambda g} t_{g^{-1}w_0\nu g'} = \sum_{\mu} m_{\lambda,\nu}^{\mu} t_{f^{-1}w_0\mu g'},$$

where  $m_{\lambda,\nu}^{\mu}$  is the multiplicity of  $V(\mu)$  in  $V(\lambda) \otimes V(\nu)$ .

### 2.2.2 On a theorem of Steinberg

Steinberg showed in [Ste75] that  $K_{\tilde{T}}(\text{pt})$  is a free  $K_{\tilde{G}}(\text{pt})$  module with basis  $\{x_w\}_{w \in W}$ . Under the isomorphism  $K_T(\text{pt}) \simeq K_{\tilde{G}}(\mathcal{B}^{\vee})$ , the  $x_w$  define an  $K_{\tilde{G}}(\text{pt})$ -basis  $\{\mathcal{F}_w\}_w$  of the latter ring, where  $\mathcal{F}_w = \mathcal{O}_{\mathcal{B}^{\vee}}(x_w)$ , and [KL87] show that the natural pairing

$$\langle -, - \rangle : K_{\tilde{G}}(\mathcal{B}^{\vee}) \otimes_{K_{\tilde{G}}(\text{pt})} K_{\tilde{G}}(\mathcal{B}^{\vee}) \rightarrow K_{\tilde{G}}(\text{pt})$$

is nondegenerate. While the dual basis is employed often in the literature starting from *loc. cit.*, we are not aware of an explicit description of it. We provide one here in very low rank cases in type A. The lack of a description in other cases of the dual basis elements as classes in  $K$ -theory of some natural objects of  $\text{Coh}_{\tilde{G}}(\mathcal{B}^{\vee})$  is the only obstruction to proving Proposition 3.

**Lemma 1.** *Let  $\tilde{G} = \text{SL}_2$  or  $\text{SL}_3$ . The collection  $\mathcal{G}_w = \mathcal{O}(y_w)[\ell(w)]$ , where*

$$y_w = \left( w^{-1} \prod_{\substack{\alpha \in \Delta \\ w^{-1}(\alpha) > 0}} \varpi_{\alpha} \right) \rho^{-1}$$

*defines the basis dual to Steinberg's basis of  $K_{\tilde{G}}(\mathcal{B}^{\vee})$  under the above pairing. Then*

*Example 2.* In type  $A_1$  and additive notation, we have  $x_1 = 0$  and  $x_{s_{\alpha}} = s_{\alpha}(\varpi_{\alpha}) = 1 - 2 = -1$ . In this case the Steinberg basis is self-dual, with  $y_1 = \varpi_{\alpha} - \rho = 1 - 1 = 0$  and  $y_{s_{\alpha}} = s_{\alpha}(0) - \rho = -1$ .

The lemma can be proved by direct computation, for example by computer.

*Remark 1.* The classes  $[\mathcal{G}_w]$  case to pair correctly with the Steinberg basis classes starting for  $G = \text{SL}_4$ , in a way apparently governed by singularities of Schubert cells. For example, for  $\text{SL}_4$ , one has

$$\langle [\mathcal{F}_w], [\mathcal{G}_1] \rangle = \text{triv}_{\text{SL}_4}$$

where  $w = 1$ , or when  $w = \sigma$  is the product of the two permutations in  $\mathfrak{S}_4$  that index singular Schubert varieties. In this case the element dual to  $[\mathcal{F}_1]$  is  $[\mathcal{G}_1] + [\mathcal{G}_{\sigma}]$ . We hope to produce natural complexes in a future version of this paper that will lift these sums and pair correctly.

## 3 Sheaves and categories

All categories, functors, and schemes are derived unless indicated otherwise. We emphasize especially that all fibre products are derived (although frequently this consideration will have no effect). Sections 3.1 and 3.3 recall the necessary material to define the category  $\mathcal{J}_0$ ; they contain no new material.

### 3.1 Derived schemes

Unless otherwise indicated, by to “apply base-change” we mean to apply Proposition 2.2.2 (b) of [GR17].

If  $X$  is classical, then  $\text{Coh}(X)$  and  $\text{Coh}_G(X)$  are the usual bounded derived categories. We write  $\text{pt} := \text{Spec } \mathbb{C}$ , and  $\mathbf{Rep}(G) := \text{Coh}_G(\text{pt})$ . We will often use silently the fact if  $f: X \rightarrow Y$  is a morphism of smooth locally-Noetherian schemes, then the pullback functor  $f^*$  preserves coherence. The classical schemes we work with will of course be exclusively locally-Noetherian, and the flag variety and bundles over it are smooth.

### 3.2 The scheme of singularities and singular support

Given a coherent sheaf  $\mathcal{F}$  on a scheme  $X$  Arinkin and Gaiitsgory in [AG15] define a classical scheme  $\text{SingSupp}(\mathcal{F})$ , the singular support of  $\mathcal{F}$ . The singular support serves in particular to measure the extent to which an object of  $\text{Coh}(X)$  fails to lie in  $\text{Perf}(X)$ . We will require only very special cases of the theory of singular support.

Let  $X$  be a quasi-smooth derived scheme. The classical scheme  $\text{Sing}(X)$  measure how far from being smooth  $X$  is. Let  $T^*(X)$  be the cotangent complex of  $X$  and  $T(X)$  its dual. Then one defines

$$\text{Sing}(X) := \mathbf{Spec} \left( \text{Sym}_{\mathcal{O}_X^\heartsuit} H^1(T(X)) \right) \rightarrow X^\heartsuit.$$

The scheme of singularities is affine over  $X^\heartsuit$ , but is not in general a vector bundle. The singular support will be a conical subset of  $\text{Sing}(X)$ . In general if a morphism

$$f: X \rightarrow Y$$

exhibits  $f^{-1}(\text{pt})$  as quasi-smooth, and  $x \in X$ , then

$$\text{Sing}(X)_x = \text{coker}(df_x)^*.$$

Note that if  $f: V \rightarrow W$  is linear, then the dg-algebra of functions on the derived scheme  $f^{-1}(0)$  is

$$\mathcal{O}_{\ker f} \otimes \text{Sym}(\text{coker}(f)^*[1]).$$

As derived schemes will appear below with approximately the same frequency as their truncations, there is no notational savings to be had by adopting either the convention that all schemes are derived unless otherwise indicated, or the opposite convention. To match our convention about functors, we declare that in any case where a derived scheme and its classical truncation appears, the derived scheme will be without decoration, as will all classical schemes that appear without any derived enhancement.

### 3.3 Categorification of $H_{\text{aff}}$ , Bezrukavnikov's equivalence

Let  $\tilde{\mathcal{N}} = T^*(\mathcal{B}^\heartsuit)$ , and denote the Steinberg variety by  $\text{St} = \tilde{\mathcal{N}} \times_{\mathfrak{g}}^L \tilde{\mathcal{N}}$ . It is naturally a derived scheme. The category  $\text{Coh}_{G \times \mathbb{G}_m}(\text{St})$  is monoidal under convolution of sheaves. Moreover,

Defining composite morphism

$$\tilde{\mathcal{N}} \times \tilde{\mathcal{N}} \xrightarrow{i} \mathfrak{g} \oplus \mathfrak{g} \xrightarrow{f} \mathfrak{g}$$

$$(x, y) \longmapsto x - y$$

we see that  $\text{St}$  fits into the pullback diagram

$$\begin{array}{ccc} \text{St} & \longrightarrow & \tilde{\mathcal{N}} \times \tilde{\mathcal{N}} \\ \downarrow & & \downarrow f \circ i \\ \text{pt} & \longrightarrow & \mathfrak{g}. \end{array}$$

Therefore  $\text{St}$  is quasi-smooth, and  $\text{Sing}(\text{St})$  is defined. We will compute its fibres over  $\text{St}^\heartsuit$ . Recall that for any variety  $X$ , if  $\xi$  is a cotangent vector at  $x \in X$ , then

$$T_{(x, \xi)}(T^*X) \simeq T_x X \oplus T_x^* X.$$

Thus

$$T_{(\mathfrak{b}, x)}(\tilde{\mathcal{N}}) = \mathfrak{g}/\mathfrak{b} \oplus \mathfrak{n}.$$

The morphism  $f \circ i$  induces the differential

$$\begin{aligned} (\mathfrak{g}/\mathfrak{b}_1 \oplus \mathfrak{n}_1) \oplus (\mathfrak{g}/\mathfrak{b}_2 \oplus \mathfrak{n}_2) &\rightarrow \mathfrak{g} \\ ((x_1, y_1), (x_2, y_2)) &\mapsto y_1 - y_2. \end{aligned}$$

Therefore

$$\mathrm{coker} \left( d_{((x_1, y_1), (x_2, y_2))} \right) (f \circ i) = \mathfrak{g}/(\mathfrak{n}_1 \oplus \mathfrak{n}_2),$$

Generically this quotient is the Cartan subalgebra  $\mathfrak{h}$ . Over the diagonal component of  $\mathrm{St}$ , it is the opposite Borel subalgebra. Thus

$$\mathrm{Sing}(\mathrm{St})_{(X, \mathfrak{b}_1, \mathfrak{b}_2)} = (\mathfrak{g}/(\mathfrak{n}_1 \oplus \mathfrak{n}_2))^*$$

### 3.4 Categorification of $J_0$

#### 3.4.1 Derived enhancement of the flag variety

Let  $\mathcal{B}^\heartsuit$  be the classical flag variety of  $\mathbf{G}$ . Let  $|\mathcal{E}|$  be the total space of the quotient

$$0 \rightarrow \tilde{\mathcal{N}} \rightarrow \mathcal{B}^\heartsuit \times \mathfrak{g} \rightarrow |\mathcal{E}| \rightarrow 0$$

and define

$$\mathcal{B} = \mathbf{Spec}(\mathrm{Sym}_{\mathcal{O}_{\mathcal{B}^\heartsuit}} \mathcal{E}[1]) = \mathbf{Spec} \left( \mathrm{Sym}_{\mathcal{O}_{\mathcal{B}^\heartsuit}} \left( \mathcal{B}^\heartsuit \times \mathfrak{g}/\tilde{\mathcal{N}} \right)^* [1] \right).$$

It is naturally a derived scheme with classical truncation  $\mathcal{B}^\heartsuit$ , with a morphism  $i: \mathcal{B} \rightarrow \tilde{\mathcal{N}}$ , and hence a morphism

$$i_{\mathrm{der}}: \mathcal{B} \times \mathcal{B} \rightarrow \mathrm{St}.$$

By construction, we have a pullback diagram

$$\begin{array}{ccc} \mathcal{B} & \longrightarrow & \tilde{\mathcal{N}} \\ \downarrow & & \downarrow \\ \{0\} & \longrightarrow & \mathcal{B}^\heartsuit \times \mathfrak{g}, \end{array} \tag{2}$$

where  $\mathcal{B}^\heartsuit \times \mathfrak{g} \rightarrow \mathcal{B}^\heartsuit$  is the trivial bundle with fibre  $\mathfrak{g}$  and  $\{0\}$  is its zero-section. Therefore  $\mathcal{B}$  is a quasi-smooth DG-scheme in the sense of [AG15]. The description of  $\mathcal{B}$  as a fibre product yields a similar description of  $\mathcal{B} \times \mathcal{B}$ . Indeed, the diagram

$$\begin{array}{ccccc} \tilde{\mathcal{N}} & \longrightarrow & \mathfrak{g} & \longleftarrow & \tilde{\mathcal{N}} \\ \downarrow & & \downarrow \mathrm{id} & & \downarrow \\ \mathcal{B}^\heartsuit \times \mathfrak{g} & \longrightarrow & \mathfrak{g} & \longleftarrow & \mathcal{B}^\heartsuit \times \mathfrak{g} \\ \uparrow & & \uparrow & & \uparrow \\ \{0\} & \longrightarrow & \mathrm{pt} & \longleftarrow & \{0\} \end{array} \tag{3}$$

gives immediately the description

$$\begin{array}{ccc} \mathcal{B} \times \mathcal{B} & \xrightarrow{i_{\mathrm{der}}} & \mathrm{St} \\ \downarrow & & \downarrow p_{\mathrm{St}} \\ \{0\} & \xrightarrow{i_{\{0\}}} & \mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit \times \mathfrak{g}. \end{array} \tag{4}$$

where now  $\{0\}$  means the zero-section of the trivial bundle  $\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit \times \mathfrak{g}$ .

The category  $\mathrm{DGCoh}_G(\mathcal{B} \times \mathcal{B})$  is a module category over the monoidal category  $\mathbf{Rep}(G)$ , via

$$V \cdot \mathcal{F} = \pi^* V \otimes_{\mathcal{O}_{\mathcal{B} \times \mathcal{B}}} \mathcal{F},$$

for  $V \in \mathbf{Rep}(G)$ , where  $\pi: \mathcal{B} \times \mathcal{B} \rightarrow \mathrm{Spec} \mathbb{C}$ . The same procedure makes  $\mathcal{J}_0$  into a module category over  $\mathbf{Rep}(G)$ .

**Lemma 2.** *If  $\mathcal{F} \in \mathrm{Coh}_{G \times \mathbb{G}_m}(\mathrm{St})$ , then*

$$\mathrm{SingSupp}(i_{\mathrm{der}}^* \mathcal{F}) \subseteq \Delta \tilde{\mathfrak{g}} \subset \tilde{\mathfrak{g}} \times \tilde{g}.$$

*Proof.* We see to apply [AG15] Proposition 7.1.3. We have

$$\begin{array}{ccc} \mathrm{Spec} \mathrm{Sym}_{\mathcal{O}_{\mathcal{B}^\heartsuit} \times \mathcal{B}^\heartsuit}}(i_{\mathrm{der}}^* T(\mathrm{St})[1]) & \xrightarrow{\mathrm{Sing}(i_{\mathrm{der}})} & \mathrm{Sing}(\mathcal{B} \times \mathcal{B}) \\ & \searrow & \swarrow \\ & \mathrm{St}^\heartsuit & \end{array}$$

Fibrewise, the singular codifferential is the linear map

$$\mathrm{Sing}(\mathrm{St})_{(0, \mathfrak{b}_1, \mathfrak{b}_2)} = (\mathfrak{g}/(\mathfrak{n}_1 \oplus \mathfrak{n}_2))^* \rightarrow (\mathfrak{g}/\tilde{\mathcal{N}}_1)^* \oplus (\mathfrak{g}/\mathfrak{n}_2)^* = \mathfrak{b}_1 \oplus \mathfrak{b}_2 = \mathrm{Sing}(\mathcal{B} \times \mathcal{B})_{(\mathfrak{b}_1, \mathfrak{b}_2)} \quad (5)$$

induced by the map, *i.e.* the direct sum of projections

$$\mathfrak{g}/\mathfrak{n}_1 \oplus \mathfrak{g}/\mathfrak{n}_2 \twoheadrightarrow \mathfrak{g}/(\mathfrak{n}_1 \oplus \mathfrak{n}_2).$$

This implies that (5) is simply the diagonal embedding, and the lemma follows.  $\square$

### 3.4.2 Definition of the category $\mathcal{J}_0$

We now define the category  $\mathcal{J}_0$ , the main point being the condition we impose on the singular supports of its objects. In our case

$$\mathrm{Sing}(\mathcal{B}) = \mathbf{Spec} \left( \mathrm{Sym}_{\mathcal{O}_{\mathcal{B}^\heartsuit}} \mathcal{E}[2] \right) \rightarrow \mathcal{B}^\heartsuit.$$

Koszul duality gives an equivalence

$$\mathrm{KD}: \mathrm{Coh}(\mathcal{B}) \rightarrow \mathrm{Sym}_{\mathcal{O}_{\mathcal{B}^\heartsuit}} \mathcal{E}[2] - \mathrm{mod}^{\mathrm{f.g.}},$$

and following [AG15], we set

$$\mathrm{SingSupp}(\mathcal{F}) = \mathrm{supp}(\mathrm{KD}(\mathcal{F})) \subset V.$$

Thus for  $\mathcal{F} = \mathcal{F}_1 \boxtimes \mathcal{F}_2 \in \mathrm{Coh}(\mathcal{B} \times \mathcal{B})$ ,  $\mathrm{SingSupp}(\mathcal{F})$  is just the usual support of some other sheaf on the total space of the bundle  $\mathrm{Sing}(\mathcal{B}) \times \mathrm{Sing}(\mathcal{B})$ . We note that  $\mathrm{Sing}(\mathcal{B})$  is none other than the bundle  $\tilde{\mathfrak{g}}$ . Indeed, the fibres of  $\mathcal{B}$  are

$$\mathrm{Spec} \mathrm{Sym}_{\mathbb{C}}((\mathfrak{g}/\mathfrak{n})^*[1]) = \mathrm{Spec} \mathrm{Sym}_{\mathbb{C}}(\mathfrak{b}[1])$$

and Koszul duality identifies

$$\mathrm{Sym}_{\mathbb{C}}(\mathfrak{b}[1]) - \mathbf{Mod} \simeq \mathrm{Sym}_{\mathbb{C}}(\mathfrak{b}^*[2]) - \mathbf{Mod},$$

and it makes sense to take the support of a module on the right-hand side on the scheme  $\mathfrak{b}$ , by defining the support to be the support of the cohomology over the classical ring  $\mathrm{Sym}_{\mathbb{C}} \mathfrak{b}^*$ .

We define  $\mathcal{J}_0$  to be the full subcategory of  $\mathrm{Coh}_G(\mathcal{B} \times \mathcal{B})$  with objects  $\mathcal{F}$  such that the projection

$$\mathrm{SingSupp}(\mathcal{F}) \rightarrow \mathrm{Sing}(\mathcal{B})$$



onto the first factor is a proper morphism. We write

$$\mathcal{J}_0 := \mathrm{Coh}_G(\mathcal{B} \times \mathcal{B})_!$$

and

$$\mathcal{J}_{0\mathcal{A}} := \mathrm{Coh}_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B})_!,$$

where  $\mathbb{G}_m$  acts trivially on  $\mathcal{B}$ .

There are two obvious ways that the projection onto the first factor can be proper: either  $\mathrm{KD}(\mathcal{F})$  is of form  $\Delta_* \mathcal{F}'$  where  $\Delta$  is the diagonal, or  $\mathrm{KD}(\mathcal{F}) = \mathcal{F}'_1 \boxtimes \mathcal{F}'_2$  with  $\mathrm{supp}(\mathcal{F}_2)$  contained in the zero section. This latter case arises precisely from sheaves  $\mathcal{F}_1 \boxtimes \mathcal{F}_2 \in \mathcal{J}_0$  such that  $\mathcal{F}_2$  is perfect. These are essentially the only examples that we will encounter: the image of  $i_{\mathrm{der}}^*$  consists of sheaves of the first type (this is especially easy to see for those sheaves whose images in  $K$ -theory are contained in  $Z(J_0) = \phi_0(Z(H_{\mathrm{aff}}))$ ), and the sheaves  $\mathcal{F}_w$  for that we define in Section 3.4.3 are all examples of the second kind.

*Remark 2.* This fact, together with the second statement of the main theorem, can be viewed as a categorification of the fact that  $\phi_0$  is injective but not surjective.

Consider the pairing operation defined by

$$(\mathcal{F}, \mathcal{G}) \mapsto \pi_*(\mathcal{F} \otimes \mathcal{G})$$

where  $\pi: \mathcal{B} \rightarrow \mathrm{pt}$ . In general, this operation does not define a functor

$$\langle -, - \rangle : \mathrm{Coh}_G(\mathcal{B}) \times \mathrm{Coh}_G(\mathcal{B}) \rightarrow \mathbf{Rep}(G)$$

but it will do so when it comes to convolution of objects of  $\mathcal{J}_0$ .

**Proposition 1.** *The category  $\mathcal{J}_0$  is a monoidal category under convolution of sheaves, and the pullback  $i_{\mathrm{der}}^*$  defines a monoidal functor*

$$i_{\mathrm{der}}^* : \mathrm{DGCoh}_{G \times \mathbb{G}_m}(\mathrm{St}) \rightarrow \mathcal{J}_{0\mathcal{A}}$$

such that

$$\mathrm{SingSupp}(i_{\mathrm{der}}^* \mathcal{F}) \subset \Delta \tilde{\mathfrak{g}}$$

for all  $\mathcal{F}$ .

Additionally,

1. If  $\mathcal{F}_1 \boxtimes \mathcal{G}$  and  $\mathcal{G}' \boxtimes \mathcal{F}_2$  are in  $\mathcal{J}_0$ , then  $\langle \mathcal{G}, \mathcal{G}' \rangle \in \mathbf{Rep}(G)$  and

$$\mathcal{F}_1 \boxtimes \mathcal{G} \star \mathcal{G}' \boxtimes \mathcal{F}_2 = \langle \mathcal{G}, \mathcal{G}' \rangle \mathcal{F}_1 \boxtimes \mathcal{F}_2;$$

2. If  $V_1, V_2 \in \mathbf{Rep}G$ , then

$$(V_1 \cdot \mathcal{F}) \star (V_2 \cdot \mathcal{G}) = (V_1 \otimes_{\mathbb{C}} V_2) \cdot \mathcal{F} \star \mathcal{G}$$

for  $\mathcal{F}, \mathcal{G} \in \mathcal{J}_0$ .

*Remark 3.* The category  $\mathrm{Coh}_G(\mathcal{B} \times \mathcal{B})$  is not monoidal.

*Proof.* Let  $\mathcal{F} \in \mathcal{J}_0$  and let  $\mathcal{G} \in \mathrm{Coh}(\mathcal{B} \times \mathcal{B})$ . Then their convolution will be coherent if

$$\mathcal{F}_{12} \boxtimes \mathcal{O}_{\mathcal{B}} \otimes \mathcal{O}_{\mathcal{B}} \boxtimes \mathcal{G}_{23},$$

is coherent, where the subscripts  $ij$  indicate which factors inside  $\mathcal{B} \times \mathcal{B} \times \mathcal{B}$  a given sheaf sits on. Noting that  $\mathrm{SingSupp}(\mathcal{O}_{\mathcal{B}}) = \{0\}$ , we have

$$\mathrm{SingSupp}(\mathcal{F}_{12} \boxtimes \mathcal{O}_{\mathcal{B}}) \cap \mathrm{SingSupp}(\mathcal{O}_{\mathcal{B}} \boxtimes \mathcal{G}_{23}) \subset \{0\} \times V \times \{0\}. \quad (6)$$

We claim that this intersection is in fact contained in the zero section of  $V \times V \times V$ . First, projection from to the first coordinate  $\mathrm{SingSupp}(\mathcal{F}_{12}) \times \{0\}$  is a proper morphism, and so the same is true for projection

from the intersection;  $\{0\} \times \text{SingSupp}(\mathcal{G}_{23})$  is closed. As  $\text{SingSupp}(\mathcal{F}_{12})$  is a conical subset, it now follows that the intersection is contained in  $\{0\} \times \{0\} \times \{0\}$ . The claim now follows from [AG15] Proposition 7.2.2. (b).

We now use that projection to the first factor from  $\text{SingSupp}(\mathcal{G}_{23})$  is also proper. We have

$$\text{Sing}(p_{13}): \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}} \times \mathcal{B} \rightarrow \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}}$$

is the inclusion of the zero section into the second coordinate, and is the identity on the other coordinates.

Define

$$Y_2 = \{(x_1, x_3) \mid (x_1, z) \in \text{SingSupp}(\mathcal{F}_{12}), (z, x_3) \in \text{SingSupp}(\mathcal{G}_{23}) \text{ for some } z \in \{0\}\}.$$

Then

$$\begin{aligned} & \text{Sing}(p_{13})^{-1}(\{(x_1, x_2, x_3) \mid (x_1, x_2) \in \text{SingSupp}(\mathcal{F}_{12}) (x_2, x_3) \in \text{SingSupp}(\mathcal{G}_{23})\}) \\ &= \{(x_1, z, x_3) \mid (x_1, z) \in \text{SingSupp}(\mathcal{F}_{12}) (x_2, z) \in \text{SingSupp}(\mathcal{G}_{23}), z \in \{0\}\} \\ &= Y_2 \times_{\mathcal{B} \times \mathcal{B}} \mathcal{B} \times \mathcal{B} \times \mathcal{B}. \end{aligned}$$

It follows from [AG15], Proposition 7.1.3 (b) that  $\text{SingSupp}(\mathcal{F} \star \mathcal{G}) \subset Y_2$ . It therefore suffices to show that projection  $Y_2 \rightarrow \tilde{\mathfrak{g}}$  is proper. Indeed, though, we have

$$p_1^{-1}(K) \subset K \times p_{2,\mathcal{G}} p_{1,\mathcal{F}}^{-1}(\{0\}),$$

for any  $K$ , where  $p_{i,\mathcal{G}}$  is the projection  $\text{SingSupp}(\mathcal{G}) \rightarrow \tilde{\mathfrak{g}}$  onto the  $i$ -th factor.

Therefore  $\mathcal{J}_0$  is a monoidal category. The same of course goes for  $\mathcal{J}_{0,\mathcal{A}}$ .

Now we show the first formula. It is easy to see that if  $\mathcal{F} \boxtimes \mathcal{G} \in \mathcal{J}_0$ , then  $\text{SingSupp}(\mathcal{G})$  must be contained in the zero section, *i.e.* that  $\mathcal{G}$  must be perfect. Then,  $\langle \mathcal{G}, \mathcal{G}' \rangle$  is coherent because  $\mathcal{G} \otimes \mathcal{G}'$  is and the map to  $\text{pt}$  is proper. Its pullback to  $\mathcal{B} \times \mathcal{B}$  is then perfect. The remainder of the formula is obtained by carrying out the calculations in Lemma 5.2.28 of [CG97]. The required projection formula and base-change are provided by Lemma 3.2.4 and Proposition 2.2.2 (b) of [GR17], respectively.

We next claim that if  $\mathcal{F} \in \text{DGCoh}_{G \times \mathbb{G}_m}(\text{St})$ , then  $i_{\text{der}}^* \mathcal{F}$  is coherent, and the projection

$$p: \text{SingSupp}(i_{\text{der}}^* \mathcal{F}) \subset \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$$

onto the first factor is proper. Coherence follows again from (3). Indeed, we need only show that the pushforward of  $i_{\text{der}}^* \mathcal{F}$  to  $\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit$  is coherent, and base-change says that this equals  $i_{\{0\}}^* p_{\text{St}*} \mathcal{F}$ . By hypothesis  $p_{\text{St}*} \mathcal{F}$  is coherent, and hence by smoothness of  $\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit \times \mathfrak{g}$  the pullback is also coherent. We must check that  $i_{\text{der}}^* \mathcal{F} \in \mathcal{J}_0$ . Indeed, this follows immediately from Lemma 2, which says that  $\text{SingSupp}(i_{\text{der}}^* \mathcal{F}) \subset \Delta V$ .

We now check that  $i_{\text{der}}^*$  is monoidal. Diagrams 2 and

$$\begin{array}{ccccc} \tilde{\mathcal{N}} & \longrightarrow & \mathfrak{g} & \longleftarrow & \mathfrak{g} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{B}^\heartsuit & \longrightarrow & \text{pt} & \longleftarrow & \mathfrak{g} \\ \uparrow & & \uparrow & & \uparrow \\ \mathcal{B}^\heartsuit & \longrightarrow & \text{pt} & \longleftarrow & \text{pt}. \end{array}$$

imply that

$$\mathcal{B} = \tilde{\mathcal{N}} \times_{\mathcal{B}^\heartsuit \times \mathfrak{g}} \mathcal{B}^\heartsuit \simeq \tilde{\mathcal{N}} \times_{\mathfrak{g}} \text{pt}.$$

Thus

$$\begin{aligned} \mathcal{B} \times \mathcal{B} \times_{\text{St}} \tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}} &\simeq \mathcal{B} \times \mathcal{B} \times_{\mathfrak{g}} \tilde{\mathcal{N}} \\ &\simeq \mathcal{B} \times \mathcal{B} \times \text{pt} \times_{\mathfrak{g}} \tilde{\mathcal{N}} \\ &\simeq \mathcal{B} \times \mathcal{B} \times \mathcal{B}, \end{aligned}$$

and we can apply base-change to the pullback diagram

$$\begin{array}{ccc} \mathcal{B} \times \mathcal{B} \times \mathcal{B} & \xrightarrow{i \times i \times i} & \tilde{\mathcal{N}} \times_{\mathfrak{g}}^L \tilde{\mathcal{N}} \times_{\mathfrak{g}}^L \tilde{\mathcal{N}} \\ \downarrow \pi_{ij} & & \downarrow p_{ij} \\ \mathcal{B} \times \mathcal{B} & \xrightarrow{i_{\text{der}}} & \text{St}. \end{array} \quad (7)$$

With diagram (7) in hand, the remainder is entirely formal. Indeed, according to the definition of convolution on  $\text{St}$ , we compute as follows: Let  $\mathcal{F}, \mathcal{G} \in \text{DGCoh}_{G \times \mathbb{G}_m}(\text{St})$ , and let  $p_{ij}, \pi_{ij}$  be the projections

$$p_{ij}: \mathcal{B} \times \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B} \times \mathcal{B}$$

and

$$\pi_{ij}: \tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}} \rightarrow \text{St}.$$

Then

$$i_{\text{der}}^*(\mathcal{F} \star \mathcal{G}) = i_{\text{der}}^* p_{13*} (p_{12}^* \mathcal{F} \otimes p_{23}^* \mathcal{G}) \quad (8)$$

$$\simeq \pi_{13*} (i \times i \times i)^* (p_{12}^* \mathcal{F} \otimes p_{23}^* \mathcal{G}) \quad (9)$$

$$\simeq \pi_{13*} (\pi_{12}^* i_{\text{der}}^* \mathcal{F} \otimes \pi_{23}^* i_{\text{der}}^* \mathcal{G}) \quad (10)$$

$$= i_{\text{der}}^* \mathcal{F} \star i_{\text{der}}^* \mathcal{G}.$$

We used base-change for the diagram (7) with  $ij = 13$  between lines (8) and (9), and just commutativity of (7) for  $ij = 12$  and  $ij = 23$  on line (10).  $\square$

Recalling that  $a(\mathbf{c}_0) = \dim \mathcal{B}$ , any quasicoherent sheaf on  $\mathcal{B}^\heartsuit$  has cohomology only in degrees at most  $a(\mathbf{c}_0)$ . This reflects that, on the level of  $K$ -theory of the influence of the  $a$ -function on the multiplication in  $J$ .

### 3.4.3 The sheaves $t_w$

Xi, in [Xi90] for  $G$  simply-connected, and Nie in [Nie11] in general gave a description in  $K$ -theory of the elements  $t_w$  for  $w \in \mathbf{c}_0$ . We recall this construction below in Section 2.2.1; here we follow it on the level of categories in the special case  $G = \text{SL}_2$  or  $\text{SL}_3$ , where it can be carried out almost verbatim. As remarked above, we hope to move beyond these two special cases in a future version of this paper.

Recalling the equivalence

$$\text{Ind}_B^G: \text{Coh}_B(\text{pt}) \rightarrow \text{Coh}_G(\mathcal{B}^\heartsuit),$$

we define

$$\mathcal{F}_w = \text{Ind}_B^G \text{Infl}_T^B(x_w),$$

where  $x_w \in \text{Coh}_T(\text{pt})$  is as in (1) and

$$\text{Infl}_T^B: \text{Coh}_T(\text{pt}) \rightarrow \text{Coh}_B(\text{pt})$$

is inflation. Likewise, define

$$\mathcal{G}_w = \text{Ind}_B^G \text{Infl}_T^B(y_w)$$

where  $y_w$  is the dual basis from Lemma 1.

Now if  $w = fw_0g^{-1}$ , define

$$t_w = \mathcal{F}_f \boxtimes p^* \mathcal{G}_g,$$

and if  $w = fw_0\chi g^{-1}$ , define

$$t_w = V(\chi) t_{fw_0g^{-1}},$$

where we view  $\mathcal{F}_f$  as pushed forward under the inclusion of the zero section of  $\mathcal{B}$ . Clearly  $t_w \in \text{Coh}_G(\mathcal{B} \times \mathcal{B})$ . Moreover, as  $\mathcal{B}^\heartsuit$  is smooth,  $\mathcal{G}_g$  is perfect, and hence  $p^* \mathcal{G}_g$  is perfect. Therefore  $\text{SingSupp}(\mathcal{G})$  is contained in the zero section of  $\text{Sing}(\mathcal{B})$ , and  $t_w \in \mathcal{I}_0$ .

*Remark 4.* At this point we see the failure of  $\phi_0$  to be surjective reflected categorically: the sheaves  $\mathcal{E}_w$  all perfect in the second coordinate, while this need not be true of sheaves restricted from  $\text{St}$ . Moreover, it is now obvious that, under the natural modification of point 1 in Proposition 1,  $\mathcal{F}_{0,\mathcal{A}}$  will act on  $\text{Coh}_{T \times \mathbb{G}_m}(\mathcal{B})$  (note the absence of any condition on singular support), a categorification of the Schwartz space of the basic affine space. We defer investigation of this action to a later work.

By Proposition 1 (or using Proposition 7.2.2. (b) of [AG15] directly),  $\langle \mathcal{E}_g, \mathcal{F}_f \rangle$  is defined for all  $g, f$  and takes values in  $\mathbf{Rep}(G)$ . In fact, it agrees with the pairing on the classical truncation given by the same procedure:

$$\langle p^* \mathcal{E}_g, \mathcal{F}_{f'} \rangle = \pi_* (p^* \mathcal{E}_g \otimes \mathcal{F}_{f'}) = \pi_*^\heartsuit p_* (p^* \mathcal{E}_g \otimes \mathcal{F}_{f'}) = \pi_*^\heartsuit (\mathcal{E}_g \otimes \mathcal{F}_{f'}) = \langle \mathcal{E}_g, \mathcal{F}_{f'} \rangle_\heartsuit$$

where

$$\begin{array}{ccc} \mathcal{B} & \xrightarrow{p} & \mathcal{B}^\heartsuit \\ & \searrow \pi & \swarrow \pi^\heartsuit \\ & \text{pt.} & \end{array}$$

Therefore by the Borel-Weil-Bott theorem, we have

$$\mathcal{F}_f \boxtimes p^* \mathcal{E}_g \star \mathcal{F}_{f'} \boxtimes p^* \mathcal{E}_{g'} = \begin{cases} \mathcal{F}_f \boxtimes \mathcal{E}_{g'} & \text{if } g = f' \\ 0 & \text{otherwise} \end{cases}.$$

Moreover, as  $G$  is reductive, we have again by Proposition 1 that

$$(V(\lambda) \mathcal{F}_f \boxtimes p^* \mathcal{E}_g) \star (V(\nu) \mathcal{F}_{g'} \boxtimes p^* \mathcal{E}_{g'}) = (V(\lambda) \otimes V(\nu)) \mathcal{F}_f \boxtimes p^* \mathcal{E}_{g'} = \bigoplus_{\mu} V(\mu) (\mathcal{F}_f \boxtimes p^* \mathcal{E}_g)^{\oplus m_{\lambda, \nu}^{\mu}}.$$

## 4 $K$ -theory

In this section we show that the functor  $\iota^*$  categorifies Lusztig's homomorphism  $\phi_0$ . By  $K$ -theory we shall always mean simply the Grothendieck group.

### 4.1 $K$ -theory of classical schemes and Lusztig's homomorphism

We first relate Lusztig's homomorphism to a construction in  $K$ -theory of classical schemes in [CG97]. There is no new material in this section; when  $G$  is simply-connected the relationship is given by [Xi16], and the analogous result in general follows from [BO04]. In order to perform calculations, though, we must devote significant space to fixing conventions.

Recall that the  $K_{G \times \mathbb{G}_m}(\text{St}) \simeq H$  as  $\mathcal{A}$ -algebras. We will use the explicit isomorphism given by Chriss and Ginzburg in [CG97], Theorem 7.2.5.

Recall that by [BO04],  $J_0 \simeq K_G(\mathbf{Y} \times \mathbf{Y})$  for a centrally-extended set  $\mathbf{Y}$  of cardinality  $\#W$ . Moreover, by 5.5 (a) of *loc. cit.*, the stabilizer of every  $y \in \mathbf{Y}$  is  $G$ . When  $G$  is simply-connected, it has no nontrivial central extensions, and hence in this case  $J_0 \simeq \text{Mat}_{\#W}(R(G))$  is a matrix ring, as first shown in [Xi90]. Combining these results, we obtain an injection  $\varphi_1: J_0 \hookrightarrow \text{Mat}_{\#W}(R(\tilde{G}))$ , where  $\tilde{G} \twoheadrightarrow G$  is a simply-connected.

In parallel, when  $G$  is simply-connected, the external tensor product gives an isomorphism  $K_G(\mathcal{B} \times \mathcal{B}) \simeq K_G(\mathcal{B}) \otimes_{R(G)} K_G(\mathcal{B}) \simeq \text{Mat}_{\#W}(R(G))$ , by [CG97], Theorem 6.2.4. This theorem does not hold when  $G$  is not simply-connected. Indeed, the trivial and Steinberg representations of  $H$  give two one-dimensional representations of  $J_0$  for  $\mathbf{G} = \text{SL}_2$ . On the other hand, the external tensor product still gives an inclusion

$$\psi_1: K^G(\mathcal{B} \times \mathcal{B}) \hookrightarrow K^{\tilde{G}}(\mathcal{B} \times \mathcal{B}) \simeq \text{Mat}_{\#W}(R(\tilde{G})).$$

(Note that  $G$  and  $\tilde{G}$  have canonically isomorphic flag varieties and Weyl groups.)

By [Nie11], we have an isomorphism  $\sigma: J_0 \rightarrow K_G(\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit)$  regardless of whether  $G$  is simply-connected or not.

**Lemma 3** ([Xi16]). *The following diagram of  $\mathcal{A}$ -algebras*

$$\begin{array}{ccccccc}
K_{G \times \mathbb{G}_m}(\mathrm{St}) & \longrightarrow & K_{G \times \mathbb{G}_m}(\tilde{\mathcal{N}} \times \tilde{\mathcal{N}}) & \xrightarrow{\tilde{\iota}^* \circ \tilde{p}^*} & K_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B}) & \xhookrightarrow{\psi_1} & \mathrm{Mat}_{\#W \times \#W}(R(\tilde{G} \times \mathbb{G}_m)) \\
\uparrow \sim & & & & & & \uparrow \mathrm{Ad}(A) \\
\mathbf{H} & \xrightarrow{\phi_0} & J_0 \otimes_{\mathbb{Z}} \mathcal{A} & \xhookrightarrow{\varphi_1} & \mathrm{Mat}_{\#W \times \#W}(R(\tilde{G} \times \mathbb{G}_m)) & & 
\end{array}$$

commutes, where  $A$  is the change-of-basis matrix from the  $Z(H(\tilde{W}(\tilde{G})))$ -basis  $\{\theta_{e_w} C \mid w \in W\}$  of  $H(\tilde{W}(\tilde{G}))$  to the  $Z(H(\tilde{W}(\tilde{G})))$ -basis  $\{C_{d_w w_0} \mid w \in W\}$ , where  $e_w$  and  $d_w$  are as in [Xi16].

#### 4.1.1 $K$ -theory of derived schemes and Lusztig's homomorphism

Koszul duality identifies with  $\mathrm{DGCoh}_{G \times \mathbb{G}_m}(\mathcal{B}_{\mathrm{der}}) \mathrm{Coh}_{G \times \mathbb{G}_m}(\tilde{\mathfrak{g}}[2])$ .

We now establish the relationship in  $K$ -theory between the monoidal functor  $i_{\mathrm{der}}^*$  from Section 3.4.1 and Lusztig's morphism  $\phi$ . Write

$$K_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B}) := K_0(\mathrm{DGCoh}_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B})_!)$$

and

$$K_{G \times \mathbb{G}_m}(\mathrm{St}) := K_0(\mathrm{DGCoh}_{G \times \mathbb{G}_m}(\mathrm{St})).$$

and write  $K(X^\heartsuit) := K_0(\mathrm{Coh}(X^\heartsuit))$  whenever  $X^\heartsuit$  is a classical scheme, and similarly for equivariant  $K$ -theory.

If  $X$  is a derived scheme with classical truncation  $X^\heartsuit$ , we may define a morphism

$$K(X) \rightarrow K(X^\heartsuit)$$

by

$$[\mathcal{F}] \mapsto \sum_i (-1)^i [\pi_i(\mathcal{F})], \quad (11)$$

where  $\pi_i(F)$  is viewed as a  $\pi_0(\mathcal{O}_X)$ -module.

Recalling that the derived structure on  $X$  is to be thought of as “higher nilpotents,” this morphism is identical in spirit to identifying  $K_0(\mathrm{Coh}(\mathrm{Spec} A))$  and  $K_0(\mathrm{Coh}(\mathrm{Spec} A_{\mathrm{red}}))$ , where  $A$  is Noetherian and  $A_{\mathrm{red}}$  is reduced. Indeed, the map (11) is also an isomorphism of abelian groups, and both isomorphisms are consequences of dévissage; see e.g. [Toë14].

**Lemma 4.** *Pushforward by bundle projection  $p: \mathrm{St} \rightarrow \mathrm{St}^\heartsuit$  induces the above map on  $G \times \mathbb{G}_m$ -equivariant  $K$ -theory. This map is an isomorphism of rings.*

*Proof.* By the remarks preceding the lemma, it suffices to show that  $p_*$  respects convolution in  $K$ -theory. By definition, we have

$$p_*([\mathcal{F}]) \star p_*([\mathcal{G}]) = \sum_{i,j} (-1)^{i+j} \pi_i(\mathcal{F}) \star \pi_j(\mathcal{G}), \quad (12)$$

whereas

$$p_*([\mathcal{F}] \star [\mathcal{G}]) = p_* p_{13*} (p_{12}^* \mathcal{F} \otimes_{q_2^* \mathcal{O}_{\mathrm{St}}} p_{23}^* \mathcal{G}) = p_{13*}^\heartsuit p_{3*} (p_{12}^* \mathcal{F} \otimes_{q_2^* \mathcal{O}_{\mathrm{St}}} p_{23}^* \mathcal{G})$$

by commutativity of the diagram

$$\begin{array}{ccc}
(\tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}}) \times_{\tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}}} (\tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}} \times_{\mathfrak{g}} \tilde{\mathcal{N}}) & \xrightarrow{p_{13}} & \mathrm{St} \\
\downarrow p_3 & & \downarrow p_{\mathrm{St}} \\
\tilde{\mathcal{N}} \times_{\mathfrak{g}} \times_{\mathfrak{g}} \tilde{\mathcal{N}} & \xrightarrow{p_{13}^\heartsuit} & \mathrm{St}^\heartsuit.
\end{array}$$

We have

$$p_{13*}^\heartsuit p_{3*}(p_{12}^* \mathcal{F} \otimes_{q_2^* \mathcal{O}_{\text{St}}} p_{23}^* \mathcal{G}) = p_{13*}^\heartsuit \sum_n (-1)^n \pi_n (p_{12}^* \mathcal{F} \otimes_{q_2^* \mathcal{O}_{\text{St}}} p_{23}^* \mathcal{G}), \quad (13)$$

and so for (12) to equal (13), we need

$$\pi_n (p_{12}^* \mathcal{F} \otimes_{q_2^* \mathcal{O}_{\text{St}}} p_{23}^* \mathcal{G}) = \sum_{i+j=n} p_{12}^{\heartsuit*} \pi_i(\mathcal{F}) \otimes_{\tilde{\mathcal{N}} \times \tilde{\mathcal{N}} \times \tilde{\mathcal{N}}} p_{23}^{\heartsuit*} \pi_j(\mathcal{G}),$$

which follows from the Künneth formula.  $\square$

In the case of  $K_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B})$ , we can define another map to the  $K$ -theory of the truncation. Let  $p_{\text{der}}: \mathcal{B} \rightarrow \mathcal{B}^\heartsuit$  be the bundle projection morphism, and let  $i: \mathcal{B}^\heartsuit \rightarrow \mathcal{B}$  be the inclusion of the zero section. Then define  $\Phi$  to be the composite

$$\Phi: K(\mathcal{J}_0) \xrightarrow{\text{id} \times i^*} K_G(\mathcal{B} \times \mathcal{B}^\heartsuit) \xrightarrow{p_* \times \text{id}} K_G(\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit).$$

That is, if  $\mathcal{F} \boxtimes \mathcal{G} \in \mathcal{J}_0$ , then

$$\Phi([\mathcal{F} \boxtimes \mathcal{G}]) = [p_* \mathcal{F}] \boxtimes [i^* \mathcal{G}].$$

*Remark 5.* It is necessary that the source of  $(\text{id} \times i)^*$  (which we will show makes sense a functor) is  $\mathcal{J}_0$  and not all of  $\text{Coh}_G(\mathcal{B} \times \mathcal{B})$ ; the functor

$$i^*: \text{QCoh}(\mathcal{B}) \rightarrow \text{QCoh}(\mathcal{B}^\heartsuit)$$

does not preserve coherence in general.

**Lemma 5.** *The morphism  $\Phi$  is well-defined and is a surjective morphism of  $R(G \times \mathbb{G}_m)$ -algebras.*

*Proof.* By the Künneth formula, we have  $\text{Sing}(\mathcal{B} \times \mathcal{B}^\heartsuit) \simeq \tilde{\mathfrak{g}} \times \tilde{\mathcal{N}}^*$ . Then

$$\text{Sing}(\text{id} \times i): \tilde{\mathfrak{g}} \times \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}} \times \tilde{\mathcal{N}}^*$$

is given by the identity in the first coordinate, and then zero map

$$(X, \mathfrak{b}) \mapsto ((Y, \mathfrak{b}) \mapsto B(X, Y) = 0),$$

in the second coordinate, where  $B$  is the Killing form (although one may of course also take the trace form).

Therefore  $\ker \text{Sing}(\text{id} \times i) = \{0\} \times \tilde{\mathfrak{g}}$ . Now let  $\mathcal{F} \in \mathcal{J}_0$ . The argument is essentially the same as in the proof of Proposition 1. The set-theoretic intersection

$$\text{SingSupp}(\mathcal{F}) \times_{\mathcal{B} \times \mathcal{B}} \mathcal{B} \times \mathcal{B}^\heartsuit \cap \ker \text{Sing}(\text{id} \times i) = \text{SingSupp}(\mathcal{F}) \cap (\{0\} \times \tilde{\mathfrak{g}})$$

is contained in the zero-section of  $\text{Sing}(\mathcal{B} \times \mathcal{B})$ . Indeed,  $\mathcal{F} \in \mathcal{J}_0$  and  $\mathcal{B}^\heartsuit$  is compact, so the above intersection must be compact. As  $\text{SingSupp}(\mathcal{F})$  is conical, we see the intersection must be contained in the zero-section. We conclude by [AG15], Proposition 7.2.2 (d) that  $(\text{id} \times i)^*$  is well-defined. Obviously  $p_* \times \text{id}$  is well-defined, and hence  $\Phi$  is well-defined. As each of  $(\text{id} \times i)^*$  and  $(p \times \text{id})_*$  are  $R(G)$ -linear (the latter by the projection formula), so is  $\Phi$ .

Finally, we show that  $\Phi$  is a morphism of rings. Using linearity, we compute

$$\Phi([\mathcal{F}_1] \boxtimes [\mathcal{G}]) \star \Phi([\mathcal{G}'] \boxtimes [\mathcal{F}_2]) = p_*[\mathcal{F}_1] \boxtimes i_{\text{der}}^*[\mathcal{G}] \star p_*[\mathcal{G}'] \boxtimes i_{\text{der}}^*[\mathcal{F}_2] = \langle i_{\text{der}}^*[\mathcal{G}], p_*[\mathcal{G}'] \rangle_{\mathcal{B}^\heartsuit} \cdot p_*[\mathcal{F}_1] \boxtimes i_{\text{der}}^*[\mathcal{F}_2],$$

whereas

$$\Phi([\mathcal{F}_1] \boxtimes [\mathcal{G}] \star [\mathcal{G}'] \boxtimes [\mathcal{F}_2]) = \langle [\mathcal{G}], [\mathcal{G}'] \rangle_{\mathcal{B}} \cdot p_*[\mathcal{F}_1] \boxtimes i_{\text{der}}^*[\mathcal{F}_2].$$

Write  $\pi: \mathcal{B}^\heartsuit \rightarrow \text{Spec } \mathbb{C}$ . Then

$$\langle i_{\text{der}}^*[\mathcal{G}], p_*[\mathcal{G}'] \rangle_{\mathcal{B}^\heartsuit} = \pi_*(i_{\text{der}}^*[\mathcal{G}] \otimes_{\mathcal{O}_{\mathcal{B}^\heartsuit}} p_*[\mathcal{G}']) = \pi_* p_*(p^* i_{\text{der}}^*[\mathcal{G}] \otimes_{\mathcal{O}_{\mathcal{B}}} [\mathcal{G}']) = \langle [\mathcal{G}], [\mathcal{G}'] \rangle_{\mathcal{B}}$$

by the formulation of the projection formula in [GR17], Lemma 3.2.4. Surjectivity follows as  $\Phi$  has a section  $\Psi$  defined  $[\mathcal{F}] \boxtimes [\mathcal{G}] \mapsto i_*[\mathcal{F}] \boxtimes p^*[\mathcal{G}]$ . This completes the proof.  $\square$

**Lemma 6.** We have  $\Phi(\mathcal{O}_{\Delta\mathcal{B}}(\lambda)) = \mathcal{O}_{\Delta\mathcal{B}^\heartsuit}(\lambda)$

*Proof.* This is a local computation that amounts to the map

$$\mathbb{C}[x, \epsilon] \otimes \mathbb{C}[y, \delta] / (x - y, \epsilon - \delta) \rightarrow \mathbb{C}[x, y]$$

quotienting by  $\delta$  and leaving the first factor untouched, where  $|x| = |y| = 0$  and  $|\epsilon| = |\delta| = -1$ . One sees that quotienting by  $\delta$  also kills  $\epsilon$ .  $\square$

**Proposition 2.** The following diagram of  $R(\mathbb{G}_m)$ -algebras

$$\begin{array}{ccccccc} K_{G \times \mathbb{G}_m}(\mathrm{St}_{\mathrm{der}}) & \xrightarrow{i_{\mathrm{der}}^*} & K_{G \times \mathbb{G}_m}(\mathcal{B}_{\mathrm{der}} \times \mathcal{B}_{\mathrm{der}}) & & & & \\ \downarrow p_{\mathrm{St}*} & & \downarrow \Phi & & & & \\ K_{G \times \mathbb{G}_m}(\mathrm{St}) & \longrightarrow & K_{G \times \mathbb{G}_m}(\tilde{\mathcal{N}} \times \tilde{\mathcal{N}}) & \xrightarrow{\bar{\iota}^* \circ \bar{p}_*} & K_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B}) & \xrightarrow{\psi_1 \otimes \mathrm{id}_{\mathcal{A}}} & \mathrm{Mat}_{\#W \times \#W}(R(\tilde{G} \times \mathbb{G}_m)) \\ \sim \uparrow & & & & & & \uparrow \mathrm{Ad}(A) \\ \mathbf{H} & \xrightarrow{\phi_0} & J_0 \otimes_{\mathbb{Z}} \mathcal{A} & \xrightarrow{\phi_1 \otimes \mathrm{id}_{\mathcal{A}}} & \mathrm{Mat}_{\#W \times \#W}(R(\tilde{G} \times \mathbb{G}_m)) \end{array}$$

commutes.

We will first describe the middle morphism on the  $K$ -theory of the classical schemes. Consider the diagrams

$$\begin{array}{ccc} \tilde{\mathcal{N}} & \xrightarrow{\pi_{\Delta}} & \mathcal{B}_{\Delta}^{\heartsuit} \\ \downarrow \iota_{\Delta} & & \downarrow \Delta \\ \mathrm{St}^{\heartsuit} & & \mathcal{B} \\ \downarrow \bar{p} & & \downarrow \\ \mathcal{B}^{\heartsuit} \times \tilde{\mathcal{N}} & \xrightarrow{\mathrm{id} \times \pi} & \mathcal{B} \times \mathcal{B}, \end{array} \quad (14)$$

which is Cartesian, and the diagram

$$\begin{array}{ccc} \mathbb{P}^1 \times \mathbb{P}^1 & & \\ \downarrow \bar{\iota} & & \\ T^*\mathbb{P}^1 \times \mathbb{P}^1 & \xleftarrow{\bar{p}} & \mathrm{St} = T^*\mathbb{P}^1 \times_{\tilde{N}^\vee} T^*\mathbb{P}^1 \\ & \nwarrow \bar{\iota} & \uparrow j \\ & & \mathbb{P}^1 \times \mathbb{P}^1. \end{array} \quad (15)$$

Let

$$\mathcal{O}_{\lambda} := [\iota_{\Delta*} \pi_{\Delta}^* \mathcal{O}_{\mathcal{B}^{\heartsuit}}(\lambda)].$$

**Lemma 7.** We have

$$\bar{\iota}^* \circ \bar{p}_*(\mathcal{O}_{\lambda}) = \Delta_* \mathcal{O}_{\mathcal{B}^{\heartsuit}}(\lambda),$$

and in the case when  $G = \mathrm{SL}_2$ , we have

$$\bar{\iota}^* \circ \bar{p}_*(-q^{1/2} j_* \mathcal{O}(0, -2)) = -q^{\frac{1}{2}} \mathcal{O}(0, -2) + q^{-\frac{1}{2}} \mathcal{O}(0, 0).$$

*Proof.* By diagram (15)

$$\bar{\iota}^* \circ \bar{p}_*(-q^{1/2} j_* \mathcal{O}(0, -2)) = -q^{\frac{1}{2}} \bar{\iota}^* \mathcal{O}(0, -2) = -q^{\frac{1}{2}} \lambda \otimes \mathcal{O}(0, -2)$$

by [CG97], Lemma 5.4.9, where  $\lambda = [\mathrm{Sym}(\mathbb{P}^1 \times T\mathbb{P}^1[1])] = [\mathcal{O}(0, 0)] - q^{-1}[\mathcal{O}(0, 2)]$ . Here we have multiplied  $\mathcal{O}(0, 2)$  by the character  $q^{-1}$ , giving its fibres trivial  $\mathbb{C}^\times$ -action, which restores equivariance of the complex defining the class  $\lambda$ . (When confronted with a linear map  $V \rightarrow W$  where  $W$  has trivial  $\mathbb{C}^\times$ -action and  $V$  is scaled by a character, one restores equivariance by tensoring  $V$  with the inverse character.) Thus we have

$$\bar{\iota}^* \circ \bar{p}_*(-q^{1/2}j_*\mathcal{O}(0, -2)) = -q^{\frac{1}{2}}(\mathcal{O}(0, 0) - q^{-1}\mathcal{O}(0, 2)) \otimes \mathcal{O}(0, -2) = -q^{\frac{1}{2}}\mathcal{O}(0, -2) + q^{-\frac{1}{2}}\mathcal{O}(0, 0).$$

To prove the second formula, apply base-change diagram (14) and use that, according to the Thom isomorphism theorem,  $((\mathrm{id} \times \pi)^*)^{-1} = \bar{\iota}^*$ . Then we have

$$\bar{\iota}^* \circ \bar{p}_*(\mathcal{O}_\lambda) = \bar{\iota}^*(\bar{p} \circ \iota_\Delta)_*\pi_\Delta^*\mathcal{O}(\lambda)$$

by base-change we have  $(\mathrm{id} \times \pi)^*\Delta_* = (\bar{p} \circ \iota_\Delta)_*\pi_\Delta^*$ , hence  $\Delta_* = ((\mathrm{id} \times \pi)^*)^{-1}(\bar{p} \circ \iota_\Delta)_*\pi_\Delta^*$ .  $\square$

*Proof of Proposition 2.* That the bottom square commutes is the combination of the main results of [Xi16] and [Nie11].

By Proposition 1, the morphism  $i_{\mathrm{der}}^*$  induces a morphism on  $K$ -theory as above. Hence by Lemma 5 and the above discussion, all the morphisms in the diagram are well-defined morphisms of algebras.

We first show the diagram commutes on the Bernstein subalgebra. Recalling from Section 3.3 that the diagonal component of the Steinberg variety is a classical rather than a derived scheme, and so  $\mathcal{O}_\lambda$  is naturally an element of  $K_{G \times \mathbb{G}_m}(\mathrm{St})$  for which  $p_{\mathrm{St}*}\mathcal{O}_\lambda = \mathcal{O}_\lambda$  with the right-hand side regarded as an object of  $K_{G \times \mathbb{G}_m}(\mathrm{St}^\heartsuit)$ . Then by Lemma 7, it suffices to show that  $\Phi([i^*\mathcal{O}_\lambda]) = [\Delta_*\mathcal{O}_{\mathcal{B}^\heartsuit}(\lambda)]$ . Indeed, though, we have

$$i^*(\mathcal{O}_\lambda) = [\mathcal{O}_{\Delta\mathcal{B}}(\lambda)],$$

as the structure sheaf of the diagonal pulls back to the structure sheaf of the diagonal. By Lemma 6 we have  $\Phi([\mathcal{O}_{\Delta\mathcal{B}}]) = [\mathcal{O}_{\Delta\mathcal{B}^\heartsuit}]$ , and likewise for the twists. Thus the diagram commutes for the Bernstein subalgebra.

Assume temporarily that  $G = \mathrm{SL}_2$ . Let  $s$  be the finite simple reflection. Consider the class

$$i_{\{0\}*}[\mathcal{O}_{\mathbb{P}^1}] \boxtimes i_{\{0\}*}[\mathcal{O}_{\mathbb{P}^1}(-2)]$$

in  $K_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B})$ . Base-change and dévissage applied to Diagram (4) shows that

$$i^*(i_{\{0\}*}[\mathcal{O}_{\mathbb{P}^1}] \boxtimes i_{\{0\}*}[\mathcal{O}_{\mathbb{P}^1}(-2)]) = (i_{\mathcal{B}} \times i_{\mathcal{B}})_*i_{\{0\}, \mathcal{B}^\heartsuit}^*i_{\mathrm{St}*}j_*\mathcal{O}(0, -2), \quad (16)$$

where  $j$  is as in diagram (15),

$$\begin{aligned} i_{\mathrm{St}}: \mathrm{St}^\heartsuit &\rightarrow \mathrm{St}, \\ i_{\{0\}, \mathcal{B}^\heartsuit}: \mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit &\rightarrow \mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit \times \mathfrak{g}, \end{aligned}$$

and

$$i_{\mathcal{B}}: \mathcal{B}^\heartsuit \rightarrow \mathcal{B}.$$

Now pasting the diagram

$$\begin{array}{ccccc} \mathrm{St} & \longleftarrow & \mathrm{St}^\heartsuit & \xleftarrow{j} & \mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit \\ \downarrow & & & \swarrow i_{\{0\}, \mathcal{B}^\heartsuit} & \\ \mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit \times \mathfrak{g} & & & & \end{array}$$

next to (3), we get that (16) equals

$$(i_{\mathcal{B}} \times i_{\mathcal{B}})_*i_{\{0\}, \mathcal{B}^\heartsuit}^*i_{\{0\}, \mathcal{B}^\heartsuit}* \mathcal{O}(0, -2).$$

Applying  $\Phi$  then gives

$$\mathcal{O} \boxtimes (i_{\mathcal{B}}^*i_{\mathcal{B}*}([\mathrm{Sym}(\mathcal{O}_{\mathcal{B}^\heartsuit}^{\oplus 3}[1]) \otimes \mathcal{O}(-2)]).$$

For the same reason, we also have

$$i_{\mathcal{B}}^*i_{\mathcal{B}*} = - \otimes \mathrm{Sym}(\mathcal{E}[2]).$$



Further, by definition there is a short exact sequence of vector bundles

$$0 \longrightarrow \tilde{\mathcal{N}} \longrightarrow \mathcal{B}^\heartsuit \times \mathfrak{g} \longrightarrow \left( \mathcal{B}^\heartsuit \times \mathfrak{g} / \tilde{\mathcal{N}} \right) = \operatorname{Spec} \operatorname{Sym}((\mathcal{B}^\heartsuit \times \mathfrak{g} / \tilde{\mathcal{N}})^*) = \operatorname{Spec} \operatorname{Sym}(\mathcal{E}) \longrightarrow 0$$

corresponding to the distinguished triangle of modules

$$\mathcal{E} \rightarrow \mathcal{O}_{\mathcal{B}^\heartsuit}^{\oplus 3} \rightarrow \mathcal{O}(2) \rightarrow \mathcal{E}[1] \rightarrow .$$

Therefore in  $K_{G \times \mathbb{G}_m}(\mathcal{B}^\heartsuit)$  we have

$$[\operatorname{Sym}(\mathcal{E}[2])] \otimes [\operatorname{Sym}(\mathcal{O}_{\mathcal{B}^\heartsuit}^{\oplus 3}[1])] = [\operatorname{Sym}(\mathcal{O}(2)[1])].$$

This says precisely that the diagram commutes when  $G = \operatorname{SL}_2$ .

Returning to general  $G$ , let  $s$  be a simple reflection. Let  $\bar{Y}_s \subset \mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit$  be the closure of the  $G$ -orbit labelled by  $s$ , and let

$$\pi_s : T_{\bar{Y}_s}^*(\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit) \rightarrow \bar{Y}_s$$

be the conormal bundle. Put  $\mathcal{Q}_s = \pi_s^* \Omega_{\bar{Y}_s / \mathcal{B}^\heartsuit}^1$ . Then by equation 7.6.34 in [CG97], it suffices to show that

$$\Phi(i^* i_{\operatorname{St}*} \mathcal{Q}_s) = \mathcal{O}_{\mathcal{B}^\heartsuit} \boxtimes i_{X_s*}(q[\mathcal{O}(2)] - [\mathcal{O}]),$$

where  $i_{X_s}$  is the inclusion of the Schubert variety  $X_s \simeq \mathbb{P}^1$  into  $\mathcal{B}^\heartsuit$ . That is, the image of  $\mathcal{Q}_s$  is just the pushforward of the answer in the  $\operatorname{SL}_2$  case. But it is clear that this is indeed the case.  $\square$

We have now nearly proved

**Proposition 3.** *For  $G = \operatorname{SL}_2$  or  $\operatorname{SL}_3$ , there exists a family of objects  $\{t_w\}_{w \in \mathbf{c}_0}$  in  $\mathcal{J}_0$ , such that that if  $t_w t_x = \sum_z \gamma_{w,x,z^{-1}} t_z$  in  $J_0$ , then*

$$t_w \star t_x = \bigoplus_z t_z^{\oplus \gamma_{w,x,z^{-1}}}$$

in  $\mathcal{J}_0$  and such that  $\Phi([t_w]) = [t_w]$ .

*Proof.* The discussion in Section 3.4.3 proves all but the last statement of the proposition. Finally, by Proposition 2, if  $w = f w_0 g^{-1}$  we have

$$\Phi([t_w]) = \Phi([\mathcal{F}_f] \boxtimes [p^* \mathcal{G}_g]) = [\mathcal{F}_f] \boxtimes i^* p^* [\mathcal{G}_g] = [\mathcal{F}_f] \boxtimes [\mathcal{G}_g].$$

The general claim follows by linearity over  $K^G(\operatorname{pt})$  and the parameterization in Section 3.4.3.  $\square$

## 4.2 The Schwartz space of the basic affine space

The Schwartz space of the basic affine space  $\mathcal{S}$  was defined by Braverman-Kazhdan in [BK99] to organize the principal series representations of  $\mathbf{G}^\vee(F)$  in a way insensitive to the poles of intertwining operators. In [BK18], Braverman and Kazhdan gave the following description of  $J_0$  in terms of the Iwahori-invariants  $\mathcal{S}^I$  of  $\mathcal{S}$ . In *loc. cit.* it was shown that  $\mathcal{S}^I$  is isomorphic to  $K_{T \times \mathbb{G}_m}(\mathcal{B}^\heartsuit)$  as an  $H \otimes \mathbb{C}[\tilde{W}]$ -module, and in [BK18], it was proven that  $J_0 \simeq \operatorname{End}_{\tilde{W}}(\mathcal{S}^I)$ , where the action of  $\tilde{W}$  is as defined in *loc. cit.*

*Example 3.* Let  $\mathbf{G} = \operatorname{SL}_2$ , with  $\tilde{W} = \langle s_0, s_1 \mid s_0^2 = s_1^2 = 1 \rangle$  and  $s_0$  the finite simple reflection. Then we have that  $K_{T \times \mathbb{G}_m}(\mathbb{P}^1)$  has basis  $\{[\mathcal{O}_{\mathbb{P}^1}], [\mathcal{O}_{\mathbb{P}^1}(-1)]\}$ , and we have  $t_{s_0} = [\mathcal{O}_{\mathbb{P}^1}] \boxtimes [\mathcal{O}_{\mathbb{P}^1}]$  and  $t_{s_1} = [\mathcal{O}(-1)] \boxtimes [\mathcal{O}(-1)][1]$  under the identification in Lemma 3. The basis elements corresponding to the two distinguished involutions in  $\mathbf{c}_0$  act by projectors, with  $t_{s_0}$  preserving  $\mathcal{O}_{\mathbb{P}^1}$  and killing  $\mathcal{O}_{\mathbb{P}^1}(-1)$ , and vice-versa for  $t_{s_1}$ .

Recalling that  $K_{G \times \mathbb{G}_m}(\mathcal{B}^\heartsuit \times \mathcal{B}^\heartsuit) \simeq K_{T \times \mathbb{G}_m}(\mathcal{B}^\heartsuit)$ , we see that we have two natural coherent categorifications of  $\mathcal{S}^I$ , and that  $\mathcal{J}_0$  acts on both of them:

**Proposition 4.** *The category  $\mathcal{J}_0$  acts on  $\operatorname{Coh}_{G \times \mathbb{G}_m}(\mathcal{B} \times \mathcal{B})$  and on  $\operatorname{Coh}_{T \times \mathbb{G}_m}(\mathcal{B})$ .*

*Proof.* This is a porism of Proposition 1. Indeed, the proof that if  $\mathcal{F}, \mathcal{G} \in \mathcal{J}_0$  then  $\mathcal{F} \star \mathcal{G} \in \operatorname{Coh}_G(\mathcal{B} \times \mathcal{B})$  used nothing about  $\operatorname{SingSupp}(\mathcal{G})$ . The proof for  $\operatorname{Coh}_T(\mathcal{B})$  is entirely similar.  $\square$

# References

- [AG15] D. Arinkin and D. Gaitsgory, *Singular support of coherent sheaves and the geometric Langlands conjecture*, Sel. Math. New Ser. **21** (2015), 1–199.
- [BDD21] R. Bezrukavnikov, S. Dawydiak, and G. Dobrovolska, *On the structure of the affine asymptotic Hecke algebras* (2021), available at [arXiv:2110.15903](#).
- [BK18] A. Braverman and D. Kazhdan, *Remarks on the asymptotic Hecke algebra* (2018), available at [arXiv:1704.03019](#).
- [BK99] ———, *On the schwartz space of the basic affine space*, Sel. Math., New Ser. **5** (1999Apr), no. 1, 1.
- [BO04] R. Bezrukavnikov and V. Ostrik, *On tensor categories attached to cells in affine Weyl groups II*, Representation Theory of Algebraic Groups and Quantum Groups, 2004, pp. 101–119.
- [CG97] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser, Boston, 1997.
- [Daw21] S. Dawydiak, *Denominators in Lusztig’s asymptotic Hecke algebra via the Plancherel formula* (2021), available at [arXiv:2110.07148](#).
- [GR17] D. Gaitsgory and N. Rozenblyum, *A Study in Derived Algebraic Geometry vol. I*, Math. Surveys Monogr., vol. 221, Amer. Math. Soc., Providence, RI, 2017.
- [KL87] D. Kazhdan and G. Lusztig, *Proof of the Deligne-Langlands conjectures for Hecke algebras*, Invent. Math. **87** (1987), 153–215.
- [Lus87] G. Lusztig, *Cells in affine Weyl groups, II*, J. Algebra **109** (1987), 536–548.
- [Lus97] ———, *Cells in affine Weyl groups and tensor categories*, Adv. Math. **129** (1997), 85–98.
- [Nie11] S. Nie, *The convolution algebra structure on  $K^{G^\vee}(\mathcal{B} \times \mathcal{B})$*  (2011), available at [arXiv:1111.1868](#).
- [Ste75] R. Steinberg, *On a theorem of Pittie*, Topology **14** (1975), 173–177.
- [Toë14] B. Toën, *Derived algebraic geometry*, EMS Surv. Math. Sci. **1** (2014), no. 2, 153–240.
- [Xi02] N. Xi, *The based ring of two-sided cells of affine Weyl groups of type  $\tilde{A}_{n-1}$* , Mem. Amer. Math. Soc., vol. 157, Amer. Math. Soc., Providence, RI., 2002.
- [Xi16] ———, *The based ring of the lowest two-sided cell of an affine weyl group, III*, Algebr Represent Theor **19** (2016), 1467–1475.
- [Xi90] ———, *The based ring of the lowest two-sided cell of an affine weyl group*, J. Algebra **134** (1990), 356–368.