

Gluon string breaking and meson spectrum in the holographic Soft Wall model

Sergey Afonin¹ and Timofey Solomko

Saint Petersburg State University, 7/9 Universitetskaya nab., St.Petersburg, 199034, Russia

Abstract

We propose a general method for finding a string-like meson spectrum which is based on a certain condition for the breaking of closed gluon string. A model is constructed that successfully realizes the proposed method. The model is based on the holographic Soft Wall model for QCD and the use of the Wilson confinement criterion. We applied our approach to the vector and scalar cases and obtained numerical predictions for the intercepts of corresponding Regge like radial meson spectra. A good agreement is obtained both with the existing experimental data and with some other known phenomenological approaches. Remarkably, our closed string breaking condition has two branches. We argue that they should correspond to states of opposite parity. The Wilson confinement criterion leads then to a natural mass splitting between parity partners. The constructed model represents thus the first example of bottom-up holographic model in which the effects of chiral symmetry breaking emerge automatically, i.e., without any ad hoc constructions based on the standard preconceived ideas from the chiral perturbation theory or the quark model.

1 Introduction and general idea

Some time ago, Andreev and Zakharov made an interesting observation [1] that the Cornell heavy-quark potential,

$$V(r) = -\frac{\kappa}{r} + \sigma r + C, \quad (1.1)$$

can be reproduced within the so-called Soft-Wall (SW) holographic model [2, 3]. Later the SW model became a popular AdS/QCD approach to modeling the non-perturbative strong interactions — the hadron Regge spectroscopy, hadron form-factors, distribution amplitudes, QCD thermodynamics, etc. (many modern references are given in the recent paper [4]). In the present Letter, we extend the approach of Ref. [1] and use it to construct a model for calculation of the intercept of linear radial spectrum of light mesons.

The linear Regge trajectories arise in the string approach to QCD. The string picture of QCD, in its turn, looks most naturally in the limit of large number of colors (called also planar), $N_c \rightarrow \infty$, of QCD [5, 6]. QCD simplifies considerably in the planar limit, in particular, the spectrum consists of infinite number of narrow (the decay width is suppressed by a $\mathcal{O}(N_c^{-1})$ factor) colorless glueballs and mesons with $\mathcal{O}(1)$ masses. The appearance of holographic QCD approach, which is inherently a large- N_c approach, has awoken interest in finding a string description of strong interactions.

If the string picture is valid, the spectrum should be string-like, at least in the first approximation. Let the radial spectrum of glueballs be

$$M_{\text{gl}}^2(n) = a(n+1), \quad n = 0, 1, 2, \dots \quad (1.2)$$

¹E-mail: s.afonin@spbu.ru.

A concrete value of intercept is not relevant and we consider the simplest one. The spectrum of mesons is expected to be different,

$$M_{\text{mes}}^2(n) = a'(n + 1 + b), \quad (1.3)$$

i.e., with a different slope and intercept. The latter depends on the quantum numbers of quark-antiquark pair. Note in passing that the spectrum (1.3) qualitatively arises in the planar two-dimensional QCD [7] and also it seems to hold semiquantitatively in real light mesons with universal slope [8–16], with the intercept dependent on spin-parity in a certain way [13–16].

When the quarks are in the fundamental representation of $SU(N_c)$ gauge group, the gluons absolutely dominate in the strong dynamics, thus the quark effects are $\mathcal{O}(1/N_c)$ suppressed and can be completely neglected in the first approximation. Imagine that we know the glueball spectrum (1.2) in the leading order in $N_c \rightarrow \infty$. We will consider the following problem: How can we get the spectrum (1.3) from (1.2) in this leading order in N_c where the quark dynamics is suppressed? This problem is partly relevant to the AdS/QCD models because the holographic approach is formulated exclusively in the leading order in $N_c \rightarrow \infty$.

Within the string picture of pure gluodynamics, the glueballs represent closed strings. If we stretch such a string, its energy grows linearly,

$$E \simeq \sigma r + \text{const}, \quad (1.4)$$

where σ is the string tension. The introduction of quarks into consideration leads to the possibility of breaking of closed strings to open strings. The stretching of a closed string costs then same energy as the stretching of two open strings with quark-antiquark pair at the ends, see Fig. 1. Thus, the tension of open string is expected to be twice smaller than that of the closed string,

$$\sigma_{\text{mes}} = \frac{1}{2}\sigma_{\text{gl}}. \quad (1.5)$$

The radial spectra (1.2) and (1.3) should emerge after quantization of the corresponding strings. The slopes of those spectra are proportional to the string tension (this is evident from the mass dimension), for instance, $a' = 2\pi\sigma$ in the case of Nambu-Goto string.

Suppose that the quarks are introduced into gluodynamics leading thus to the possibility of open string creation. How can one deduce the spectrum of quark-antiquark pairs from the spectrum (1.2)? The first guess follows from (1.5) and consists in the replacement $a \rightarrow a' = a/2$. But this is not enough because the intercept is expected to change as well, $n + 1 \rightarrow n + 1 + b$. Suppose further that we have a dynamical model for planar gluodynamics which is able to reproduce the confinement behavior (1.4) at large distances and can incorporate the dependence $\sigma(b)$ on the external parameter b that we relate with quark effects. The closed gluon string tension $\sigma(b)$ should be a decreasing function of b because the fermions contribute to the strong dynamics with opposite sign. And if at some point $b = b_0$ the tension $\sigma(b)$ is halved we get a




Figure 1: A schematic view of a closed gluon string vs. two open strings.

tension that, according to (1.5), is characteristic to the open string. Our basic idea is that the realization of the corresponding condition,

$$\sigma(b_0) = \frac{1}{2}\sigma(0), \quad (1.6)$$

triggers the closed string breaking and formation of open string(s).

We thus propose the following answer to the question above: In order to find the string-like radial meson spectrum (1.3) from the glueball spectrum (1.2) we should find the value of $b = b_0$ from the closed string breaking condition (1.6). Since $a \sim \sigma$, the meson spectrum will be given by

$$M_{\text{mes}}^2(n) = \frac{1}{2}a(n+1+b_0). \quad (1.7)$$

Below we construct an explicit model realizing the proposed idea for prediction of string-like meson spectrum in the large- N_c limit and briefly compare our results with the phenomenology and some other approaches.

The paper is organized as follows. The SW holographic model and its generalization to the arbitrary intercept parameter is recalled in Section 2. In Section 3, we first give a brief review on calculation of linearly rising potential within the SW model, then extend the corresponding technique to the generalized SW model, and after that construct our model for finding the intercept in the case of vector mesons. The results obtained are compared with the existing experimental data on light mesons and argued to be consistent both with these data and with some known phenomenological approaches. The scalar case is analyzed in Section 4. We conclude in Section 5. Some technical details are given in Appendix.

2 The SW holographic model

Our starting point will be the well-known phenomenological SW holographic model. The original SW model put forward by Son et al. [2] is defined by the action

$$S = \int d^4x dz \sqrt{g} e^{-cz^2} \mathcal{L}, \quad (2.1)$$

where g_{MN} is the metric of the Poincare patch ($z > 0$) of the AdS₅ space

$$ds^2 = g^{MN} dx_M dx_N = \frac{R^2}{z^2} (\eta^{\mu\nu} dx_\mu dx_\nu - dz^2), \quad \eta_{\mu\nu} = \text{diag}\{1, -1, -1, -1\}, \quad (2.2)$$

R is the radius of the AdS₅ space and z is the fifth (called holographic) coordinate. The function e^{-cz^2} represents the dilaton background (c is a constant) and \mathcal{L} is the Lagrangian density of some free field in AdS₅ space which, by assumption, is dual on the AdS₅ boundary to some QCD operator. The model is defined in the probe approximation, i.e., the metric is not backreacted by matter fields and dilaton (there exist also various dynamical SW approaches, we will follow the standard probe approximation — the backreaction is assumed to be suppressed in the large- N_c limit within which the holographic approach is formulated [4]). In the case of free vector fields, the mass spectrum of the model is given by [2]

$$M^2(n) = 4|c|(n+1). \quad (2.3)$$

There exists another formulation of SW model proposed independently by Andreev [3], in which the dilaton background is replaced by a modified metric,

$$g_{MN} = \text{diag} \left\{ \frac{R^2}{z^2} h, \dots, \frac{R^2}{z^2} h \right\}, \quad h = e^{-2cz^2}. \quad (2.4)$$

The vector spectrum of this model coincides with (2.3). Other formulations of SW model are possible. Various interrelations between different formulations were thoroughly studied in the recent work [4]. In particular, if a SW model is given by the action

$$S = \int d^4x dz \sqrt{g} B(z) \mathcal{L}, \quad (2.5)$$

it can be reformulated as a SW model with rescaled fields

$$S = \int d^4x dz \sqrt{\tilde{g}} \tilde{\mathcal{L}}, \quad (2.6)$$

and modified metric

$$\tilde{g}_{MN} = B^{(3/2-J)^{-1}} g_{MN}. \quad (2.7)$$

Here J denotes the value of spin of 5D field in the Lagrangian density \mathcal{L} .

As was shown in [17] and expanded upon in [4], the vector SW model (2.1) can be generalized to the case of arbitrary intercept in the spectrum (2.3),

$$M^2(n) = 4|c|(n+1+b), \quad (2.8)$$

where the parameter b regulates the value of intercept. The action of generalized vector SW model takes the form

$$S = \int d^4x dz \sqrt{g} e^{-cz^2} U^2(b, 0, cz^2) \mathcal{L}, \quad (2.9)$$

where U is the Tricomi hypergeometric function.

3 Holographic Wilson loop, gluon confinement and meson string

A practical way to calculate the potential between two sources, such as (1.4), is based on a holographic interpretation of the expectation value of the Wilson loop suggested by Maldacena [18]. Assume that we have a rectangular Wilson loop located in the 4D boundary of the 5D AdS space, schematically pictured in the Fig. 2. Such Wilson loop can be related to the propagation of a massive quark [18]. Then, in the limit of $T \rightarrow \infty$, its expectation value is proportional to [18]

$$\langle W(\mathcal{C}) \rangle \sim e^{-TE(r)}, \quad (3.1)$$

where $E(r)$ corresponds to the energy of the quark-antiquark pair. On the other hand, as was proposed in [18], it can be computed as

$$\langle W(\mathcal{C}) \rangle \sim e^{-S}, \quad (3.2)$$




Figure 2: A Wilson loop.

where S represents the area of a string world-sheet which produces the loop \mathcal{C} . From comparing (3.1) and (3.2) one can calculate the energy as

$$E = \frac{S}{T}. \quad (3.3)$$

A particular implementation of this idea within the SW holographic model was developed by Andreev and Zakharov in [1]. They considered the modified Euclidean AdS₅ space with the metric (2.4), in which the sign of exponent in h was argued to be opposite in the Euclidean case²,

$$g_{MN} = \text{diag} \left\{ \frac{R^2}{z^2} h, \dots, \frac{R^2}{z^2} h \right\}, \quad h = e^{2cz^2}. \quad (3.4)$$

The given setup was shown to result in the linearly rising energy at large distances,

$$E \underset{r \rightarrow \infty}{\simeq} \sigma r + \text{Const} + \mathcal{O}(1/r), \quad \sigma \equiv \frac{R^2}{\pi\alpha'} ec, \quad (3.5)$$

where $1/\alpha'$ is the tension of fundamental string (see (3.8) below). The Coulomb behavior,

$$E \underset{r \rightarrow 0}{\simeq} \frac{\kappa}{r} + \text{Const} + \mathcal{O}(r), \quad (3.6)$$

was also obtained at small distances. Thus the structure of the Cornell potential (1.1) was reproduced.

Our first goal is to extend this result to the case of generalized SW model with arbitrary intercept parameter. As is seen from (2.9) and (2.7), we must apply the analysis above to the following extension of modified Euclidean AdS metric (3.4)

$$g_{MN} = \text{diag} \left\{ \frac{R^2}{z^2} h, \dots, \frac{R^2}{z^2} h \right\}, \quad h = e^{2cz^2} U^4(b, 0, cz^2). \quad (3.7)$$

In what follows we apply the analysis of [1] to the case of metric (3.7).

The starting point is the Nambu-Goto action

$$S = \frac{1}{2\pi\alpha'} \int d^2\xi \sqrt{\det g_{MN} \partial_\alpha X^M \partial_\beta X^N}, \quad (3.8)$$

where g_{MN} is the warped metric defined above. This action describes the area of string world-sheet for which the expectation value of Wilson loop should be calculated. If we choose $\xi_1 = t$ and $\xi_2 = x$ as the parametrization of the world-sheet then, after integration over t from 0 to T , the action can be rewritten as

$$S = \frac{TR^2}{2\pi\alpha'} \int_{-r/2}^{r/2} dx \frac{h}{z^2} \sqrt{1 + z'^2}, \quad (3.9)$$

where $z' = \frac{dz}{dx}$. Since the action is translation invariant due to the absence of the explicit dependence of the Lagrangian on x , there exists a conserving quantity, which in this case is equal to

$$\frac{h}{z^2} \frac{1}{\sqrt{1 + z'^2}} = \text{Const}. \quad (3.10)$$

²Our definition of c differs from the one in [1] by the factor of 4, namely $h = e^{cz^2/2}$ in [1]. The latter leads to the absence of factor 4 in the spectrum (2.3).

Let us denote

$$z_0 \equiv z|_{x=0}, \quad h_0 \equiv h|_{z=z_0}. \quad (3.11)$$

Since we know that $z = 0$ at the ends of the Wilson loop, $x = \pm r/2$, and the system is symmetric, z reaches the maximum value of $z = z_0$ at $x = 0$. Using $z'|_{x=0} = 0$ we obtain from (3.10)

$$\frac{h_0}{z_0^2} = \frac{h}{z^2} \frac{1}{\sqrt{1+z'^2}} \Rightarrow z' = \sqrt{\frac{h^2 z_0^4}{h_0^2 z^4} - 1} \Rightarrow dx = dz \frac{h_0 z^2}{h z_0^2} \frac{1}{\sqrt{1 - \frac{h_0^2 z^4}{h^2 z_0^4}}}, \quad (3.12)$$

and one can then write

$$\int_{-r/2}^{r/2} dx = 2 \int_0^{z_0} dz \frac{h_0 z^2}{h z_0^2} \frac{1}{\sqrt{1 - \frac{h_0^2 z^4}{h^2 z_0^4}}}. \quad (3.13)$$

The factor of 2 appears due to two equal contributions from integration from $x = -r/2$ to 0 and from 0 to $x = r/2$. After defining the new variables

$$v \equiv \frac{z}{z_0}, \quad \lambda \equiv cz_0^2, \quad (3.14)$$

and substituting $h(z)$ we get the final expression for the distance r ,

$$r = 2\sqrt{\frac{\lambda}{c}} \int_0^1 dv \frac{U^4(b, 0, \lambda)}{U^4(b, 0, \lambda v^2)} \frac{v^2 e^{2\lambda(1-v^2)}}{\sqrt{1 - v^4 e^{4\lambda(1-v^2)} \frac{U^8(b, 0, \lambda)}{U^8(b, 0, \lambda v^2)}}}. \quad (3.15)$$

The expression for the energy can be obtained from the equation (3.3) and the action (3.9) by replacing integration over x with the integration over z using (3.13). The resulting expression, however, is divergent at $v = z = 0$ and thus requires regularization. After imposing the cutoff $\varepsilon \rightarrow 0$ we get the regularized energy

$$E_R = \frac{R^2}{\pi\alpha'\varepsilon} + E, \quad (3.16)$$

with the energy integral equal to

$$E = \frac{R^2}{\pi\alpha'} \sqrt{\frac{c}{\lambda}} \left[\int_0^1 \frac{dv}{v^2} \left(\frac{e^{2\lambda v^2} U^4(b, 0, \lambda v^2)}{\sqrt{1 - v^4 e^{4\lambda(1-v^2)} \frac{U^8(b, 0, \lambda)}{U^8(b, 0, \lambda v^2)}}} - D \right) - D \right], \quad (3.17)$$

where we introduced the regularization constant $D \equiv U^4(b, 0, 0)$. At this point we have expressed the potential in terms of the parametric functions $r(\lambda)$ and $E(\lambda)$. In order to find the asymptotics $r \rightarrow \infty$ we need to express it in terms of λ .

Let us start by noting that these integrals must be real-valued which requires the expression under the square root to be non-negative in the entire integration range $v \in [0, 1]$,

$$1 - v^4 e^{4\lambda(1-v^2)} \frac{U^8(b, 0, \lambda)}{U^8(b, 0, \lambda v^2)} \geq 0. \quad (3.18)$$

In its turn, this condition is satisfied if the minimum value of this expression is non-negative within the same range. The minima are given by the roots of the derivative of (3.18). For $v \neq 0$, this leads to the equation

$$1 - 2\lambda v^2 - 4\lambda v^2 \frac{U'(b, 0, \lambda v^2)}{U(b, 0, \lambda v^2)} = 0, \quad U(b, 0, \lambda v^2) \neq 0, \quad (3.19)$$

where $U'(b, 0, x) = \partial_x U(b, 0, x)$. Denoting

$$x \equiv \lambda v^2, \quad (3.20)$$

the numerical calculations show that the integrals are real-valued if

$$\lambda < x. \quad (3.21)$$

In Fig. 3, we display an example of behavior of the expression (3.18) as the function of two arguments, λ and v . The expression is non-negative in the region to the left of the thick vertical line, hence, the integrals are real-valued for all $v \in [0, 1]$ in that region.

It is interesting to remark that similar conditions can be obtained from the Sonnenschein confinement criterion [19] which imposes the following requirements on the g_{00} element of the warped metric

$$\partial_z g_{00}|_{z=z_0} = 0, \quad g_{00}|_{z=z_0} \neq 0, \quad (3.22)$$

in order for the background to be dual to a confining theory in the sense of area law behavior of Wilson loop. In our case, using the previous notation $\lambda = cz_0^2$ and after certain simplifications, the second of Sonnenschein conditions takes the form

$$U(b, 0, \lambda) \neq 0. \quad (3.23)$$

The first condition reads as follows

$$\frac{U^4(b, 0, \lambda)}{(\lambda/c)^{3/2}} \left(1 - 2\lambda - 4\lambda \frac{U'(b, 0, \lambda)}{U(b, 0, \lambda)} \right) = 0. \quad (3.24)$$

Due to the first condition the only possible case when this equality can be true is

$$1 - 2\lambda - 4\lambda \frac{U'(b, 0, \lambda)}{U(b, 0, \lambda)} = 0. \quad (3.25)$$

The conditions (3.23) and (3.25) coincide with the equation (3.19).

Going back to the integrals we note that since r is a growing function of λ , for obtaining the large distance asymptotics we can consider the “large” λ asymptotics. Since λ is bounded from above, this means finding the asymptotics $\lambda \rightarrow x$.

Further we follow the procedure outlined in [1]. Since the integrals diverge in the upper bound $v = 1$, we expand the expression under the square root near $v = 1$. The integrals transform into

$$r \underset{v \rightarrow 1}{=} 2\sqrt{\frac{\lambda}{c}} \int_0^1 \frac{dv}{\sqrt{A(b, \lambda)(1-v) + B(b, \lambda)(v-1)^2}}, \quad (3.26)$$




Figure 3: The contour plots of the expression under the square root inside the integrals, see (3.18), in the case of $b \approx -0.48$. The black vertical line is $\lambda \approx 0.16$. The horizontal lines display the region where (3.18) is non-negative. The region of negative (3.18) is marked by vertical lines.

$$E \underset{v \rightarrow 1}{=} \frac{R^2}{\pi\alpha'} \sqrt{\frac{c}{\lambda}} e^{2\lambda U^4(b, 0, \lambda)} \int_0^1 \frac{dv}{\sqrt{A(b, \lambda)(1-v) + B(b, \lambda)(v-1)^2}}, \quad (3.27)$$

where A and B are some functions of b and λ which do not depend on v (see Appendix). We do not need the analytical expressions for r and E (see Appendix) as we are interested in the behavior $E(r)$. Substituting (3.26) into (3.27) we get

$$E \underset{r \rightarrow \infty}{\sim} \frac{R^2}{2\pi\alpha'} \frac{e^{2x U^4(b, 0, x)}}{x} cr, \quad (3.28)$$

where $\lambda \rightarrow x$ while x is the solution of the equation

$$1 - 2x - 4x \frac{U'(b, 0, x)}{U(b, 0, x)} = 0, \quad (3.29)$$

which can be found numerically.

As follows from (1.4), the coefficient in front of r in (3.28) represents the string tension,

$$\sigma(b) = \frac{R^2}{2\pi\alpha'} \frac{e^{2x U^4(b, 0, x)}}{x} c, \quad (3.30)$$

where the dependence on the parameter b is hidden inside x by virtue of the equation (3.29). Since $U(0, 0, x) = 1$, the equation (3.29) yields $x = 1/2$ at $b = 0$ and thus (3.30) reproduces the string tension (3.5) of Andreev-Zakharov analysis in [1],

$$\sigma(0) = \frac{R^2}{\pi\alpha'} ec. \quad (3.31)$$

We approached to the key point of our analysis. Following the discussion in the Section 1, we interpret the case of $b = 0$ as corresponding to pure gluodynamics that absolutely dominates in the large- N_c limit — the limit where the holographic models are supposedly defined. Thus the case of $b = 0$ (zero “quark” external parameter) is interpreted as the case of closed gluon string with pure gluon sources. Now we impose the closed string breaking condition (1.6). From (3.30) and (3.31) we get the equation

$$\frac{e^{2x-1} U^4(b, 0, x)}{2x} = \frac{1}{2}. \quad (3.32)$$

The equations (3.29) (the reality condition) and (3.32) (the closed string breaking condition) form a system of two equations for two parameters b and x . As we consider the vector case, the solutions b_0 for the intercept parameter are interpreted as our prediction for the vector spectrum of quark-antiquark pairs.

The closed string breaking condition (3.32) has two branches and each one is intersected by the reality condition (3.29), see Fig. 4 (the seeming intersection at $x = 0$ is not a well defined solution as $x = 0$ is a singularity line³).




Figure 4: The string breaking (blue solid) and reality (orange dashed) conditions in the vector case.

³Actually the emerging point $(x, b) = (0, -1)$ seems to require a separate study. The arising spectrum would correspond to the deep ultraviolet limit $z \rightarrow 0$ (as $x = cz^2$) and, according to (1.3), would contain a massless vector particle. A physical interpretation (the photon?) is unclear.

The first numerical solution (intersection with the lower branch) yields

$$b_0 \approx -0.483, \quad x \approx 0.162. \quad (3.33)$$

The obtained value of b_0 in (3.33) is remarkably close to $b = -0.5$ which arises for vector mesons in various approaches. Namely, according to (1.3) the predicted radial spectrum of ρ -mesons (and of their isoscalar counterparts ω) is close to

$$M_\rho^2(n) = 2M_\rho^2 \left(n + \frac{1}{2} \right), \quad (3.34)$$

where M_ρ is the mass of the ground ρ -meson state $\rho(770)$ [20]. The spectrum (3.34) emerged for the first time in the Veneziano dual amplitudes after imposing the Adler self-consistency condition [21–23] (the pion scattering amplitude is zero at zero momentum). The relation (3.34) also appeared in the finite energy QCD sum rules [24] and in the QCD sum rules in the large- N_c limit [25–27]. This relation arises naturally in the light-front holographic approach [28]. The experimental data on the form-factor of charged pion $F_\pi(Q^2)$ have the most satisfactory resonance description if the radial vector spectrum is close to the behavior $m_\rho^2(n) \sim n + 1/2$ as in (3.34) (see, e.g., [4, 28]). Within the framework of essentially the same holographic model, a very close value of b_0 was recently obtained from fitting the mean pion charge radius squared [4]. We see thus that the proposed new approach passes an important phenomenological test.

Taking the standard mean value for the slope of meson radial trajectories $a' = 4|c| \approx 1.14 \text{ GeV}^2$ [9] in (1.3), the predicted large- N_c masses are $M(n) \approx \{0.77, 1.32, 1.69, 2.00, 2.27, \dots\} \text{ GeV}$. It looks natural to match them to the resonances $\rho(770)$, $\rho(1450)$ (it is a broad resonance region including possible $\rho(1270)$ [20]), $\rho(1700)$, $\rho(2000)$, and $\rho(2270)$, correspondingly [20].

The second numerical solution (intersection with the upper branch) gives

$$b_0 \approx 0.341, \quad x \approx 0.851. \quad (3.35)$$

The same input for the slope leads to the radial spectrum $M(n) \approx \{1.24, 1.63, 1.95, 2.22, \dots\} \text{ GeV}$. This spectrum is surprisingly close to the experimental spectrum of axial-vector mesons a_1 — the chiral partners of ρ -mesons: $a_1(1260)$ (the mass is $1230 \pm 40 \text{ GeV}$), $a_1(1640)$, $a_1(1930)$, and $a_1(2270)$, correspondingly [20].

We propose that the observed coincidence is not accidental but points at a new way of incorporation of the Chiral Symmetry Breaking (CSB) into the bottom-up holographic approach.

We remind the reader that the AdS_5 space locally looks like the 5D Minkowski space, i.e., a space with one time and four flat space directions. The space rotations in four dimensions contain two independent three-dimensional rotations, $SO(4) = SO(3) \times SO(3)$. Correspondingly, the spin is marked by two numbers. In the vector case, this leads to the existence of two spin-1 mesons — the “Left” one A_L^M with spin $(1, 0)$ and the “Right” one A_R^M with spin⁴ $(0, 1)$. From the viewpoint of representations of Lorentz group, $SO(3)$ is the rotational subgroup of the little group for 5D massless particles (the extension of $SO(2)$ for 4D massless particles) which coincides with the little group for 4D massive particles. The ensuing equality of physical degrees of freedom makes possible a direct Kaluza-Klein like projection of 5D massless particles to 4D massive ones. But the 5D states have no definite space parity (this notion appears only for an odd number of space dimensions), hence, the dimensional reduction must be performed together with constructing

⁴There is also the representation $(1/2, 1/2)$ but such a state does not carry helicity λ (according to a Weinberg theorem, the Lorentz representation (a, b) carries the helicity $\lambda = |b - a|$) describing just a derivative of a scalar field with only one physical degree of freedom.

states of definite space parity. In the spin-1 case under consideration, the corresponding states are the usual vector, $V^\mu = (A_R^\mu + A_L^\mu)/\sqrt{2}$, and the axial-vector, $A^\mu = (A_R^\mu - A_L^\mu)/\sqrt{2}$, as now the P -reflection interchanges (1, 0) and (0, 1). The Lagrangian density for free vector fields in (2.1) becomes

$$\mathcal{L} = -\frac{1}{4}F^2 = -\frac{1}{4}(F_L^2 + F_R^2) = -\frac{1}{4}(F_V^2 + F_A^2). \quad (3.36)$$

The field V_μ for quark-antiquark states is supposedly dual to the QCD operator $\bar{q}\gamma_\mu q$ or its isovector partner $\bar{q}\gamma_\mu\vec{\tau}q$ and A_μ is dual to $\bar{q}\gamma_\mu\gamma_5 q$ or $\bar{q}\gamma_\mu\gamma_5\vec{\tau}q$. Since these operators have equal mass dimension $\Delta = 3$, for description of mass splitting between parity partners due to the CSB effects one needs some *ad hoc* modification that usually has no motivation inside the bottom-up holographic approach itself. The most known examples are incorporation of pions in the spirit of chiral perturbation theory in the pioneering Hard-Wall models [29, 30] and incorporation of quark angular momentum in the light-front holographic QCD [28].

We conjecture that the existence of two branches in the closed string breaking condition (3.32) corresponds to the possibility of having two kinds of spin-1 states, the vector and axial mesons, after the dimensional reduction of 5D fields A_L^M and A_R^M . We obtain then a description of mass splitting between the parity partners without any *ad hoc* modifications based on a “pre-understanding” of CSB from other approaches. Remarkably, the arising description turns out to be quantitative.

4 Scalar case

The original Andreev-Zakharov analysis [1] was based on the vector SW model. It is interesting to apply this analysis and our extension to the scalar case (the SW model describing the scalar fields). According to the general design of generalized SW models with arbitrary intercept, the Tricomi function in (2.9) is replaced by $U(b, -1, cz^2)$ in the scalar case [4] and the SW scalar spectrum is (see, e.g., [4])

$$M^2(n) = 4|c|(n + 3/2 + b). \quad (4.1)$$

The transformation (2.7) then leads to the following background function in the metric (3.7),

$$h = e^{2cz^2/3}U^{4/3}(b, -1, cz^2). \quad (4.2)$$

Repeating the same steps as in the vector case, we obtain the following expressions for the distance $r(\lambda)$ and energy $E(\lambda)$

$$r = 2\sqrt{\frac{\lambda}{c}} \int_0^1 dv \frac{U^{4/3}(b, -1, \lambda)}{U^{4/3}(b, -1, \lambda v^2)} \frac{v^2 e^{2\lambda(1-v^2)/3}}{\sqrt{1 - v^4 e^{4\lambda(1-v^2)/3} \frac{U^{8/3}(b, -1, \lambda)}{U^{8/3}(b, -1, \lambda v^2)}}}, \quad (4.3)$$

$$E = \frac{R^2}{\pi\alpha'} \sqrt{\frac{c}{\lambda}} \left[\int_0^1 \frac{dv}{v^2} \left(\frac{e^{2\lambda v^2/3} U^{4/3}(b, -1, \lambda v^2)}{\sqrt{1 - v^4 e^{4\lambda(1-v^2)/3} \frac{U^{8/3}(b, -1, \lambda)}{U^{8/3}(b, -1, \lambda v^2)}}} - D \right) - D \right], \quad (4.4)$$

with the regularization constant $D \equiv U^{4/3}(b, -1, 0)$. They lead to the large distance asymptotics for the energy

$$E \underset{r \rightarrow \infty}{\sim} \frac{R^2}{2\pi\alpha'} \frac{e^{2x/3} U^{4/3}(b, -1, x)}{x} cr. \quad (4.5)$$

The parameter x has the same meaning as in the vector case — it is one of the roots of derivative of the expression under the square root in the integrals above. The scalar counterpart of the reality condition (3.29) becomes

$$1 - \frac{2}{3}x - \frac{4}{3}x \frac{U'(b, -1, x)}{U(b, -1, x)} = 0, \quad (4.6)$$

and the one for the string tension (3.30) is

$$\sigma(b) \equiv \frac{R^2}{2\pi\alpha'} \frac{e^{2x/3} U^{4/3}(b, -1, x)}{x} c. \quad (4.7)$$

The particular case $b = 0$ corresponds to

$$\sigma(0) = \frac{R^2}{3\pi\alpha'} ec. \quad (4.8)$$

It differs from (3.31) by the factor of $1/3$. This difference arises from the formulation of SW model in the form of (3.4): If, for the purpose of describing the Regge form of spectrum, one introduces an IR modification in the metric, the slope becomes spin-dependent (actually this was the reason to introduce the IR modification in the form of dilaton background e^{-cz^2} in the original paper by Son et al. [2]). According to the transformation (2.7), the function e^{cz^2} in the Euclidean dilaton background becomes e^{2cz^2} in the spin-1 case and $e^{2cz^2/3}$ in the spin-0 one resulting in the factor of $1/3$ that emerged in (4.8).

As follows from (4.7) and (4.8), our condition for the closed gluon string breaking (1.6) (which does not depend on a general constant factor in σ) now reads

$$\frac{3e^{2x/3-1}U^{4/3}(b, -1, x)}{2x} = \frac{1}{2}. \quad (4.9)$$

As in the vector case, this condition has two branches, see Fig. 5, and we may try to interpret them as arising for scalar states of opposite space parity, namely the lower branch should correspond to mesons of negative parity and the upper one to heavier mesons of positive parity.

The reality condition (4.6) intersects the upper branch in the point

$$b_0 \approx 0.394, \quad x \approx 1.906. \quad (4.10)$$

The scalar sector is notoriously difficult for theoretical descriptions, especially the scalar isoscalar resonances f_0 having the vacuum quantum numbers. Many of them are believed to be not dominated by the quark-antiquark component [20]. If true, this means that a pattern of spectrum emerging in the large- N_c limit can be very different from the reality. Since in our holographic approach this limit is supposed to be built-in, the comparison with the isovector scalar mesons a_0 looks more reasonable but the corresponding experimental data are scarce. Nevertheless, the spectrum (4.1) with the standard mean slope $4|c| = 1.14 \text{ GeV}^2$ [9] and the intercept b_0 given by the solution (4.10) predicts for the ground state




Figure 5: The string breaking (blue solid) and reality (orange dashed) conditions in the scalar case.

the mass $M(0) \approx 1.47$ GeV which matches perfectly the mass of $a_0(1450)$ -meson (its experimental mass is 1474 ± 19 GeV [20]). This agrees with a widespread opinion about tetraquark nature of $a_0(980)$ (together with its isoscalar counterpart $f_0(980)$) and that the lowest quark-antiquark a_0 -meson is $a_0(1450)$ [20].

The lower branch in Fig. 5 is intersected by the reality condition (4.6) in the point

$$b_0 \approx -1.859, \quad x \approx 0.227. \quad (4.11)$$

The ground state of spectrum (4.1) is then tachyonic. This looks like a failure of our approach. On the other hand, the scalar SW spectrum (4.1) was derived for the fields supposedly dual to the scalar QCD operators of canonical dimension $\Delta = 3$, which are $\bar{q}q$ and $\bar{q}\gamma_5 q$. If we do not specify Δ , the SW scalar spectrum is [31]

$$M^2(n) = 4|c|(n + \Delta/2 + b). \quad (4.12)$$

As is known from the old Gell-Mann–Oakes–Renner relation, the pion mass is related with the renormalization-group invariant QCD operator $m_q \bar{q}q$ that has the mass dimension $\Delta = 4$. The spectrum (4.12) yields then $M(0) \approx 0.4$ GeV which already somewhat resembles the physical pion mass⁵.

5 Conclusion

A vital problem for any successful phenomenological model is fixing its parameters from some theoretical principles. We considered a generalized version of phenomenologically successful holographic Soft Wall model [2] that contains an extra “intercept” parameter b contributing both to the Regge like mass spectrum and to other observables like hadron form factors [4], and proposed a theoretical way for calculation of this parameter within the framework of holographic approach itself. Our method is based on a holographic realization of the Wilson confinement criterion and the introduced closed string breaking condition. We applied this method to the vector and scalar cases and found a very good agreement with many other phenomenological approaches and with the experimental data on light unflavored mesons. We argued that within the framework of the proposed holographic approach, the effects of chiral symmetry breaking seems to be built-in, at least the mass splitting between the parity partners may be described without incorporation of additional modifications.

The proposed model can be extended to the case of arbitrary integer spin. It would be also interesting to apply our approach, with certain, perhaps serious, modifications, to the actively developing holographic Higgs sector — it is not excluded that the condition for the creation of quark-antiquark states that we used might have a counterpart for the creation of fermion-antifermion pairs from the Higgs field and this could be exploited for a holographic prediction, e.g., of the Higgs mass. Also our approach could find applications in studies of the Z -string stability in the two Higgs doublet extension of the Standard Model [32].

Acknowledgements

This research was funded by the Russian Science Foundation grant number 21-12-00020.

⁵Similarly to the vector case, a singular solution seems to appear in the point $(x, b) = (0, -2)$, see Fig. 5, which for $\Delta = 4$ in (4.12) would lead to a massless pseudoscalar particle in the deep ultraviolet limit. A physical interpretation (the pion?) is again unclear.

Appendix

Introducing the notation

$$s(b, \lambda; v) \equiv 1 - v^4 e^{4\lambda(1-v^2)} \frac{U^8(b, 0, \lambda)}{U^8(b, 0, \lambda v^2)}, \quad (5.1)$$

the functions A and B in the expressions (3.26) and (3.27) can be written as

$$A(b, \lambda) = -s'_v(b, \lambda; 1) = 4 \left[1 - 2\lambda - 4\lambda \frac{U'(b, 0, \lambda)}{U(b, 0, \lambda)} \right], \quad (5.2)$$

$$B(b, \lambda) = \frac{s''_{vv}(b, \lambda; 1)}{2} = -2 \left[16\lambda^2 - 18\lambda + 3 + 72\lambda^2 \frac{U'(b, 0, \lambda)^2}{U(b, 0, \lambda)^2} + 4\lambda \frac{(16\lambda - 9)U'(b, 0, \lambda) - 2\lambda U''(b, 0, \lambda)}{U(b, 0, \lambda)} \right]. \quad (5.3)$$

The integral in the expressions (3.26) and (3.27) can be calculated analytically using [33]

$$\int \frac{du}{\sqrt{Au + Bu^2}} = \frac{1}{\sqrt{B}} \ln \left(2\sqrt{B(Au + Bu^2)} + 2Bu + A \right). \quad (5.4)$$

This integral produces logarithmic in λ asymptotics.

References

- [1] O. Andreev and V. I. Zakharov, *Heavy-quark potentials and AdS/QCD*, Phys. Rev. D **74** (2006), 025023, [hep-ph/0604204]
- [2] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, *Linear confinement and AdS/QCD*, Phys. Rev. D **74** (2006), 015005, [hep-ph/0602229]
- [3] O. Andreev, *1/q**2 corrections and gauge/string duality*, Phys. Rev. D **73** (2006), 107901, [hep-th/0603170]
- [4] S. S. Afonin and T. D. Solomko, *Theory of bottom-up holographic models for linear Regge trajectories of light mesons*, [2106.01846]
- [5] G. 't Hooft, *A Planar Diagram Theory for Strong Interactions*, Nucl. Phys. B **72** (1974), 461
- [6] E. Witten, *Baryons in the 1/n Expansion*, Nucl. Phys. B **160** (1979), 57-115
- [7] G. 't Hooft, *A Two-Dimensional Model for Mesons*, Nucl. Phys. B **75** (1974), 461-470
- [8] A. V. Anisovich, V. V. Anisovich and A. V. Sarantsev, *Systematics of q anti-q states in the (n, M**2) and (J, M**2) planes*, Phys. Rev. D **62** (2000), 051502, [hep-ph/0003113]
- [9] D. V. Bugg, *Four sorts of meson*, Phys. Rept. **397** (2004), 257-358, [hep-ex/0412045]
- [10] E. Klempt and A. Zaitsev, *Glueballs, Hybrids, Multiquarks. Experimental facts versus QCD inspired concepts*, Phys. Rept. **454** (2007), 1-202, [0708.4016]

- [11] D. M. Li, B. Ma, Y. X. Li, Q. K. Yao and H. Yu, *Meson spectrum in Regge phenomenology*, Eur. Phys. J. C **37** (2004), 323-333, [hep-ph/0408214]
- [12] M. Shifman and A. Vainshtein, *Highly Excited Mesons, Linear Regge Trajectories and the Pattern of the Chiral Symmetry Realization*, Phys. Rev. D **77** (2008), 034002, [0710.0863]
- [13] S. S. Afonin, *Experimental indication on chiral symmetry restoration in meson spectrum*, Phys. Lett. B **639** (2006), 258-262, [hep-ph/0603166]
- [14] S. S. Afonin, *Light meson spectrum and classical symmetries of QCD*, Eur. Phys. J. A **29** (2006), 327-335, [hep-ph/0606310]
- [15] S. S. Afonin, *Towards understanding spectral degeneracies in nonstrange hadrons. Part I. Mesons as hadron strings versus phenomenology*, Mod. Phys. Lett. A **22** (2007), 1359-1372, [hep-ph/0701089]
- [16] S. S. Afonin, *Properties of new unflavored mesons below 2.4-GeV*, Phys. Rev. C **76** (2007), 015202, [0707.0824]
- [17] S. S. Afonin, *Generalized Soft Wall Model*, Phys. Lett. B **719** (2013), 399-403, [1210.5210]
- [18] J. M. Maldacena, *Wilson loops in large N field theories*, Phys. Rev. Lett. **80** (1998), 4859-4862, [hep-th/9803002]
- [19] J. Sonnenschein, *Stringy confining Wilson loops*, PoS **tmr2000** (2000), 008, [hep-th/0009146]
- [20] P. A. Zyla *et al.* [Particle Data Group], *Review of Particle Physics*, PTEP **2020** (2020) no.8, 083C01
- [21] M. Ademollo, G. Veneziano and S. Weinberg, *Quantization conditions for regge intercepts and hadron masses*, Phys. Rev. Lett. **22** (1969), 83-85
- [22] P. D. B. Collins, *Regge theory and particle physics*, Phys. Rept. **1** (1971), 103-234
- [23] P. D. B. Collins, *An Introduction to Regge Theory and High-Energy Physics*, (Cambridge University Press, Cambridge, 1977)
- [24] N. V. Krasnikov and A. A. Pivovarov, *The Use of Finite Energy Sum Rules for the Description of Resonances in QCD*, Phys. Lett. B **112** (1982), 397
- [25] S. S. Afonin, *Quark condensate and deviations from string - like behavior of meson spectra*, Phys. Lett. B **576** (2003), 122-126, [hep-ph/0309337]
- [26] S. S. Afonin, A. A. Andrianov, V. A. Andrianov and D. Espriu, *Matching Regge theory to the OPE*, JHEP **04** (2004), 039, [hep-ph/0403268]
- [27] S. S. Afonin and D. Espriu, *Qualitative solution of QCD sum rules*, JHEP **09** (2006), 047, [hep-ph/0602219]
- [28] S. J. Brodsky, G. F. de Teramond, H. G. Dosch and J. Erlich, *Light-Front Holographic QCD and Emerging Confinement*, Phys. Rept. **584** (2015), 1, [1407.8131]

- [29] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, *QCD and a holographic model of hadrons*, Phys. Rev. Lett. **95** (2005), 261602, [hep-ph/0501128]
- [30] L. Da Rold and A. Pomarol, *Chiral symmetry breaking from five dimensional spaces*, Nucl. Phys. B **721** (2005), 79-97, [hep-ph/0501218]
- [31] S. S. Afonin, *Towards reconciling the holographic and lattice descriptions of radially excited hadrons*, Eur. Phys. J. C **80** (2020), 723, [2008.05610]
- [32] M. Eto, Y. Hamada and M. Nitta, *Stable Z-strings with topological polarization in two Higgs doublet model*, [2111.13345]
- [33] D. Zwillinger, *Table of Integrals, Series, and Products*, Academic Press (2014)