Hyperbolic energy and Maskit gluings*

Piotr T. Chruściel^{†‡} Erwann Delay[§]¶ Raphaela Wutte*^{††}

December 2, 2021

Abstract

We derive a formula for the energy of asymptotically locally hyperbolic (ALH) manifolds obtained by a gluing at infinity of two ALH manifolds. As an application we show that there exist three dimensional conformally compact ALH manifolds without boundary, with connected conformal infinity of higher genus, with constant negative scalar curvature and with negative mass.

Contents

1	Introduction	2
2	Localised boundary-gluing of ALH metrics	4
3	Boundary-gluing of Birmingham-Kottler solutions	6
4	Hyperbolic mass4.1 The definition	
5	Higher genus solutions with negative mass5.1 Genus two5.2 Higher genus	14 14 25
6	A lower bound on mass?	26
\mathbf{A}	A punctured torus	27

^{*}Preprint UWThPh-2021-25

[†]University of Vienna, Faculty of Physics

[‡]EMAIL piotr.chrusciel@univie.ac.at, URL homepage.univie.ac.at/piotr.chrusciel

[§] Avignon Université, Laboratoire de Mathématiques d'Avignon, F-84916 Avignon

 $[\]P Aix Marseille Université – F.R.U.M.A.M.- CNRS
– F-13331 Marseille$

EMAIL Erwann.Delay@univ-avignon.fr, URL https://erwanndelay.wordpress.com

^{**}TU Wien, Faculty of Physics, Institute of Theoretical Physics

^{††}EMAIL rwutte@hep.itp.tuwien.ac.at

В	Static KIDs	28
	B.1 Birmingham-Kottler	28
	B.1.1 $m_c = 0, n \neq 2 \dots \dots$	31
	B.1.2 $m_c \neq 0, n \neq 2$	32
	B.1.3 $n = 2$	32
	B.2 Horowitz-Myers	33
C	Transformation behaviour of the mass aspect tensor under conformal trans-	
	formations	3 5
	C.1 Relation to coordinates in the main text	37
	C.2 $n=3$	38
	C.3 $n=4$	39

1 Introduction

In [33] Isenberg, Lee and Stavrov, inspired by [35], have shown how to glue together two asymptotically locally hyperbolic (ALH) general relativistic initial data sets by performing a boundary connected sum, which they referred to as a "Maskit gluing". The resulting initial data set has a conformal boundary at infinity which is a connected sum of the original ones. A variation of this construction has been presented in [13]. It is of interest to analyse the properties of the initial data sets so obtained. The aim of this work is to address this question in the time symmetric case, with vanishing extinsic curvature tensor K_{ij} .

We start by a short presentation of the boundary-gluing construction of [13] in Section 2. The construction involves a certain amount of freedom which we make precise, showing that the gluing results in whole families of new ALH metrics. As a particular case, in Section 3 we apply the construction to Birmingham-Kottler metrics. This provides new families of ALH metrics with minimal surface boundaries with more than one component, and with locally explicit metric when the mass parameter is zero. One thus obtains time-symmetric vacuum initial data for spacetimes containing several apparent horizons; such initial data sets evolve to spacetimes with multiple black holes.

Next, an important global invariant of asymptotically hyperbolic general relativistic initial data sets is the total energy-momentum vector $\mathbf{m} \equiv (m_{\mu})$ [18, 19, 42] (compare [2, 20, 27]) when the conformal metric at infinity is that of the round sphere (we talk of AH metrics then), and the total mass m for the remaining topologies at infinity; this is reviewed in Section 4. It turns out that the formulae for the mass after the gluings of both [33] and [13] are relatively simple. This is analysed in Section 4.4, where we derive formulae (4.37)-(4.40). This is the first main result of this paper.

A quick glance at (4.38) suggests very strongly that the boundary-gluing of two Horowitz-Myers metrics, which both have negative mass aspect functions, will lead to a manifold with higher-genus boundary at infinity and with negative mass. This turns out, however, to be subtle because of correction terms that are inherent to the constructions. In fact, negativity of the total mass is far from clear for the Isenberg-Lee-Stavrov gluings, because these authors use the conformal method, which changes the mass integrand in a way which appears difficult to control in the neck region. Things are clearer when the localised boundary-gluing of [13] is used, and in Section 5 we show that negativity indeed holds. We thus construct three dimensional conformally compactifiable ALH manifolds with constant scalar curvature, without boundaries at finite distance, with connected boundary at infinity of higher genus topology and with negative mass. This is the second main result of this paper. As already hinted to, such manifolds correspond to time-symmetric Cauchy surfaces in vacuum spacetimes with negative cosmological




Table 1.1: Mass inequalities for asymptotically Birmingham-Kottler metrics. A double question mark indicates that no results are available; a single one indicates existence of partial results. The shorthand "bdry" refers to a black-hole boundary. "Good spin" denotes a topology where the manifold is spin and the spin structure admits asymptotic Killing spinors. The case "other conformal infinity" includes higher genus topologies when the boundary is two-dimensional, but also e.g. quotients of spheres in higher dimensions. Reference [0] is this work. Finally, μ is the mass aspect function. The critical value of the mass $m_{\rm crit}$, assuming it exists, is expected to be determined by the conformal structure of the boundary at infinity.

constant. Here one should keep in mind that the Horowitz-Myers metrics have toroidal topology at infinity, and that the higher-genus Birmingham-Kottler metrics with negative mass are either nakedly singular, or have a totally geodesic boundary, or acquire a conformal boundary at infinity with two components, and contain an apparent horizon, after a doubling across the boundary.

It is likely that a Maskit gluing of a Horowitz-Myers soliton with a spherical Kottler ("Schwarz-schild-anti de Sitter") black hole with small mass will provide an example of a conformally compact vacuum black hole with toroidal infinity and negative mass. This can almost certainly be established by our methods but does not directly follow from the arguments below, where symmetry between the gluing-components is used. We plan to return to this question in a near future.

To put our work in perspective, the currently known or conjectured lower bounds for the mass of asymptotically Birmingham-Kottler manifolds are summarised in Table 1.1. In particular we note the four-dimensional negative-mass ALH metrics of [21, 38] (compare [9, 24]), where conformal infinity is a quotient of a sphere. We further note that while there exists a notion of mass for general locally asymptotically hyperbolic metrics with constant scalar curvature which admit asymptotic static potentials [27], nothing is known about the sign of the mass for those which are not asymptotically Birmingham-Kottler. See also [29] for a discussion of the issues occurring in this context.

2 Localised boundary-gluing of ALH metrics

Before entering the subject, some comments on our terminology are in order. We say that a metric g on a manifold without boundary M has a $conformal \ completion \ (\bar{M}, \bar{g})$ if \bar{M} is a manifold with boundary such that $\bar{M} = M \cup \partial \bar{M}$, and if there exists a function $\Omega \geq 0$ on \bar{M} which vanishes precisely on $\partial \bar{M}$, with $d\Omega$ without zeros on $\partial \bar{M}$, and with $g = \Omega^{-2}\bar{g}$ on M. This definition generalises to manifolds M with boundary, in which case $\partial \bar{M}$ will be the union of the original boundaries of ∂M , where Ω is strictly positive, and the new ones where Ω vanishes; the new ones are referred to as boundaries at conformal infinity. We say that (M,g) is conformally compactifiable when \bar{M} is compact. We say that (M,g) is asymptotically locally hyperbolic (ALH) if the scalar curvature of g approaches a constant when the boundary at infinity is approached. We say that an ALH manifold is asymptotically hyperbolic (AH) if the conformal class of \bar{g} on the conformal boundary at infinity is that of a round sphere. Asymptotically Birmingham-Kottler metrics are defined as metrics which asymptote to the Birmingham-Kottler metrics of Section 3 below. The Birmingham-Kottler metrics themselves are ALH, which can be seen by setting $\Omega = r$ in (3.1), and noting that they have constant scalar curvature since they solve the time-symmetric general relativistic scalar constraint equation with (negative) cosmological constant.

The differentiability requirements of \bar{g} at the conformal boundary at infinity often need to be added in the definitions above, and depend upon the problem at hand. Here we will be interested in a class of manifolds with well defined mass, as will be made precise in Section 4.1.

The reader is warned that there is no consistency in the literature concerning this terminology. While our definition of ALH coincides with that of several authors, some other authors use AH for what we call ALH here. However, we find it natural to reserve the name AH for the special case where the metric is asymptotic to that of hyperbolic space.

Our analysis here is motivated by the localised boundary-gluings of ALH manifolds, or initial data sets, as in [13, Section 3.5]. In this section we present a somewhat more precise version of these gluings.

We start with points p_1 , p_2 , lying on the conformal boundary of two ALH vacuum initial data sets (M_1, g_1, K_1) and (M_2, g_2, K_2) . (An identical construction applies when p_1 and p_2 belong to the same manifold; then the construction provides instead a handle connecting a neighborhood of p_1 with a neighborhood of p_2 . Instead of vacuum initial data one can also take e.g. data satisfying the dominant energy condition; the construction will preserve this. Further, if $K_1 \equiv 0 \equiv K_2$, then one can have $K \equiv 0$ throughout the construction.) We assume that both (M_1, g_1, K_1) and (M_2, g_2, K_2) have extrinsic curvature tensors asymptoting to zero, and have well-defined total energy-momentum, cf. Section 4. As shown in [13] for deformations of data sets preserving the vacuum condition or for scalar curvature deformations preserving an inequality, or in [14, Appendix A] for deformations of data sets preserving the dominant energy condition, for all $\varepsilon > 0$ sufficiently small we can construct new initial data sets $(M_i, g_{i,\varepsilon}, K_{i,\varepsilon})$, i = 1, 2, such that the metrics coincide with the hyperbolic metric in coordinate half-balls $\mathcal{U}_{1,\varepsilon}$ of radius ε around p_1 and $\mathcal{U}_{2,\varepsilon}$ around p_2 , and the $K_{i,\varepsilon}$'s are zero there. Here the coordinate half-balls refer to coordinates on the upper half-space model of hyperbolic space \mathcal{H}^n , in which we have

$$\mathcal{H}^n = \mathbb{R}^n_+ = \left\{ x = (\omega, z) \in \mathbb{R}^{n-1} \times]0, \infty[\right\}, \tag{2.1}$$

with the metric

$$b = \frac{|d\omega|^2 + dz^2}{z^2} \,. \tag{2.2}$$

In order to avoid a proliferation of indices we now choose ε so that the deformation described above has been carried out, with corresponding ALH metrics $(M_i, g_{i,\varepsilon}, K_{i,\varepsilon})$, i = 1, 2, and coordinate half-balls $\mathcal{U}_{i,\varepsilon}$, which we write from now on as (M_i, g_i, K_i) , and \mathcal{U}_i . It should be kept in




Figure 2.1: The "thin component" $\mathcal{U}_{\mathfrak{h}}$ and its boundary \mathfrak{h} in the half-space model, where hyperbolic space is represented as a half-space $\{z>0\}$, and the conformal boundary at infinity is the hyperplane $\{z=0\}$,. After an "exotic gluing" has been performed, the metric becomes exactly hyperbolic inside $\mathcal{U}_{\mathfrak{h}}$.

mind that different values of ε lead to different initial data sets near the gluing region, but the data sets remain the original ones, hence identical, away from the gluing region, which can be chosen as small as desired.

The above shows in which sense the exterior curvature tensor K is irrelevant for the problem at hand. Therefore, from now on, we will only consider initial data sets with $K \equiv 0$ in the gluing region.

The gluing construction uses hyperbolic hyperplanes \mathfrak{h} in the hyperbolic-space region of (M_i, g_i) . These are defined, in the half-space model, as half-spheres with centres on the hyperplane $\{z=0\}$ in the coordinates of (2.2) which, for our purposes, are entirely contained in the half-balls \mathscr{U}_i . The conditionally compact, in \mathbb{R}^n , component of \mathcal{H}^n separated by \mathfrak{h} will be referred to as the *thin component*, denoted by $\mathscr{U}_{\mathfrak{h}}$, and the remaining one will be called the *fat component*; see Figure 2.1.

In what follows we will also invoke the *Poincaré ball* model, which represents the *n*-dimensional hyperbolic space \mathcal{H}^n as the open unit ball B^n endowed with the metric

$$b = \frac{4}{(1 - |\vec{x}|^2)^2} \delta, \qquad (2.3)$$

where δ is the Euclidean metric.

A basic fact is that for every hyperbolic hyperplane \mathfrak{h} as above there exist two isometries of the hyperbolic space, which we denote by $\Lambda_{\mathfrak{h},\pm}$, such that $\Lambda_{\mathfrak{h},+}$ maps \mathfrak{h} to the equatorial hyperplane of the Poincaré ball, with the fat region being mapped to the upper hemisphere, while $\Lambda_{\mathfrak{h},-}$ again maps the hyperbolic hyperplane to the equatorial hyperplane but it maps the fat region into the lower hemisphere. Using physics terminology, an example of $\Lambda_{\mathfrak{h},+}$ is provided by a boost along the axis passing through the origin of the Poincare ball and the barycenter of the hyperbolic hyperplane. Given $\Lambda_{\mathfrak{h},+}$, a map $\Lambda_{\mathfrak{h},-}$ can be obtained by applying to $\Lambda_{\mathfrak{h},+}$ a rotation by π around any axis lying on the equatorial plane.

It should be clear that there are many such pairs $\Lambda_{\mathfrak{h},\pm}$: consider, e.g., the isometries $R_{\pm}\Lambda_{\mathfrak{h},\pm}$, where the R_{\pm} 's are some rotations along the axis joining the north pole and the south pole.

Let \mathscr{P} denote the collection of pairs of isometries (Λ_1, Λ_2) of hyperbolic space with the following property: There exist hyperbolic hyperplanes $\mathfrak{h}_i \subset \mathring{\mathscr{U}}_i$, i = 1, 2, such that Λ_1 is a $\Lambda_{\mathfrak{h}_1,i,+}$ and Λ_2 is a $\Lambda_{\mathfrak{h}_2,-}$. Here $\mathring{\mathscr{U}}_i$ denotes the interior of \mathscr{U}_i .

For each such pair of hyperbolic hyperplanes \mathfrak{h}_i the manifolds $M_i \setminus \mathscr{U}_{\mathfrak{h}_i}$ are manifolds with a non-compact boundary component $\partial \mathscr{U}_{\mathfrak{h}_i} = \mathfrak{h}_i$ extending to the conformal boundary at infinity

of M_i , with the hyperbolic metric near the boundary \mathfrak{h}_i .

Given $(\Lambda_1, \Lambda_2) \in \mathscr{P}$ we construct a boundary-glued manifold M_{Λ_1, Λ_2} by gluing the boundaries \mathfrak{h}_i as follows: We map the thin complement $\mathscr{U}_{\mathfrak{h}_{1,i}}$ of $\mathfrak{h}_{1,i}$ to the lower half of the Poincaré ball using Λ_1 . We map the thin complement $\mathscr{U}_{\mathfrak{h}_2}$ of \mathfrak{h}_2 to the upper half of the Poincaré ball using Λ_2 . We then identify the two manifolds with boundary $M_i \setminus \mathscr{U}_{\mathfrak{h}_i}$ along the equatorial plane of the Poincaré ball using the identity map.

The metrics on $M_i \setminus \mathcal{U}_{\mathfrak{h}_i}$ coincide with the original ones (one can think of the maps $\Lambda_{\mathfrak{h}_i}$ as changes of coordinates), hence are ALH there by hypothesis. Both metrics are exactly hyperbolic at both sides of the gluing boundary, namely the equatorial plane of the Poincaré ball, and extend smoothly there. Hence for every pair $(\Lambda_1, \Lambda_2) \in \mathscr{P}$ the manifold M_{Λ_1, Λ_2} is a smooth ALH manifold.

3 Boundary-gluing of Birmingham-Kottler solutions

An obvious candidate to which our construction can be applied is the space-part of the Birmingham-Kottler (BK) metrics,

$$g = f^{-1}dr^2 + r^2h_k$$
, $f = r^2 + k - \frac{2m_c}{r^{n-2}}$, (3.1)

where m_c is a constant which will be referred to as the *coordinate mass parameter*, and where h_k is an Einstein metric on an (n-1) dimensional manifold N^{n-1} with scalar curvature equal to

$$R(h_k) = k(n-1)(n-2), \qquad k \in \{0, \pm 1\}.$$

The associated spacetime metric

$$-fdt^2 + g$$

is a solution of the (Lorentzian) vacuum Einstein equations with a negative cosmological constant. The collection of (N^{n-1}, h_k) 's is quite rich: one should, e.g., keep in mind the existence of many Einstein metrics on higher dimensional spheres, including exotic ones [7].

We will assume that m_c is in a range so that f has positive zeros, with the largest one, denoted by r_0 , of first order. The metric is then smooth on the product manifold $[r_0, \infty) \times N^{n-1}$, with a totally geodesic boundary at $r = r_0$.

The construction of Section 2 applied to two such manifolds,

$$(M_i = [r_i, \infty) \times N_i^{n-1}, g_i),$$

leads to manifolds with a conformal boundary at infinity with connected-sum topology $N_1^{n-1}\#N_2^{n-1}$ and a totally geodesic boundary which has two connected components, one diffeomorphic to N_1^{n-1} and the second to N_2^{n-1} .

One can double each of the original manifolds across their totally geodesic boundaries, in which case the doubled manifolds have no boundaries but each has a conformal infinity with two components. Performing our construction on a chosen pair of boundaries at infinity one obtains an ALH manifold with three boundary components, one with topology N_1^{n-1} , one with topology N_2^{n-1} , and a third one with topology $N_1^{n-1}\#N_2^{n-1}$.

One can iterate the construction, obtaining ALH manifolds with an arbitrary number of components of the boundary at infinity, and an arbitrary number of totally geodesic compact boundary components. The maximal globally hyperbolic development of the resulting time-symmetric general relativistic vacuum Cauchy data will have a Killing vector field defined in a neighborhood of each such boundary (but not globally in general), which becomes the bifurcation

surface of a bifurcate Killing horizon for this vector field. The case $m_c = 0$ is of special interest here, as then the metric is exactly hyperbolic everywhere, so that the construction of [12], which glues-in an exactly hyperbolic half-ball, is trivial. All the boundary gluing constructions described here apply without further due to this case. It is likely that some of the metrics constructed above evolve to the spacetimes considered in [1, 31], it might be of some interest to explore this.

4 Hyperbolic mass

4.1 The definition

We recall the definition of hyperbolic mass from [18]. We consider a family of Riemannian metrics g which approach a "background metric" b with constant scalar curvature -n(n-1) as the conformal boundary at infinity is approached. We assume that b is equipped with a non-empty set of solutions of the *static Killing Initial Data (KID)* equations:

$$\mathring{\Delta}V - nV = 0, \tag{4.1}$$

$$\mathring{D}_i \mathring{D}_j V = V(\mathring{R}_{ij} + nb_{ij}), \qquad (4.2)$$

where \mathring{R}_{ij} denotes the Ricci tensor of the metric b, \mathring{D} the Levi-Civita connection of b, the operator $\mathring{\Delta} := b^{k\ell}\mathring{D}_k\mathring{D}_\ell$ is the Laplacian of b, and we use D for the Levi-Civita connection of g. Nontrivial triples (M,b,V), where V solves (4.1)-(4.2), are called static Killing Initial Data. Strictly speaking, for the purpose of defining the mass only the geometry of a neighborhood of the conformal boundary at infinity of M is relevant. A couple (b,V) will also be called a static KID when M or its conformal boundary are implicitly understood. The functions V will be interchangeably referred to as static potentials or static KIDs. The (finite dimensional) vector space of static potentials of (M,b) will be denoted by \mathscr{K}_b . All static potentials for the BK and Horowitz-Myers metrics are derived in Appendix B.

Under conditions on the convergence of g to b spelled-out in (4.6)-(4.10) below, well defined global geometric invariants can be extracted from the integrals [18]

$$H(V,b) := \lim_{R \to \infty} \int_{r=R} \mathbb{U}^i(V) dS_i$$
 (4.3)

where $V \in \mathcal{K}_b$, with

$$\mathbb{U}^{i}(V) := 2\sqrt{\det g} \left(V g^{i[k} g^{j]l} \mathring{D}_{j} g_{kl} + D^{[i} V g^{j]k} (g_{jk} - b_{jk}) \right), \tag{4.4}$$

and where dS_i are the hypersurface forms $\partial_i | dx^1 \wedge \cdots \wedge dx^n$.

From now on, until explicitly indicated otherwise the background metric b will be a BK metric with $m_c = 0$. In the coordinates of (3.1) we set

$$M_{\text{ext}} := [R, \infty) \times N^{n-1}, \tag{4.5}$$

for some large $R \in \mathbb{R}^+$, and we consider the following orthonormal frame $\{f_i\}_{i=1,n}$ on M_{ext} :

$$f_i = r^{-1} \epsilon_i, \quad i = 1, \dots, n-1, \quad f_n = \sqrt{r^2 + k} \, \partial_r,$$
 (4.6)

where the ϵ_i 's form an orthonormal frame for the metric h_k . We set

$$g_{ij} := g(f_i, f_j), \quad b_{ij} := b(f_i, f_j) = \begin{cases} 1, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$
 (4.7)

Assuming that

$$\int_{M_{\text{ext}}} \left(\sum_{i,j} |g_{ij} - b_{ij}|^2 + \sum_{i,j,k} |f_k(g_{ij})|^2 \right) r \, d\mu_g < \infty \,, \tag{4.8a}$$

$$\int_{M_{\text{ext}}} |R_g - R_b| \ r \ d\mu_g < \infty \,, \tag{4.8b}$$

$$\exists C > 0 \text{ such that } C^{-1}b(X, X) \le g(X, X) \le Cb(X, X), \tag{4.9}$$

where $d\mu_g$ denotes the measure associated with the metric g, one finds that the limit in (4.3) exists and is finite. If moreover one requires that

$$\sum_{i,j} |g_{ij} - b_{ij}| + \sum_{i,j,k} |f_k(g_{ij})| = \begin{cases} o(r^{-n/2}), & \text{if } n > 2, \\ O(r^{-1-\epsilon}), & \text{if } n = 2, \text{ for some } \epsilon > 0, \end{cases}$$
(4.10)

then the "mass integrals" (4.3) are well defined, in the following sense: Consider any two background metrics b_i , i = 1, 2, of the form (3.1), with the same boundary manifold (N^{n-1}, h_k) with $m_c = 0$, in particular with the same value of k. Assume that g approaches both b_1 and b_2 at the rates presented above. Then there exists an isometry Φ of b_1 such that

$$H(V, b_1) = H(V \circ \Phi, b_2).$$
 (4.11)

Consider a background b of the form (3.1). The function

$$V_{(0)}(r) = \sqrt{r^2 + k} \tag{4.12}$$

satisfies the static KID equations (4.1). Assuming (4.8)-(4.10), a somewhat lengthy calculation shows that the mass integral

$$H(V_{(0)}) := H(\sqrt{r^2 + k}, b)$$

equals [20]

$$H(V_{(0)}) = \lim_{R \to \infty} R^{n-1}(R^2 + k) \times \int_{\{r=R\}} \left(-\sum_{i=1}^{n-1} \left\{ \frac{\partial e_{ii}}{\partial r} + \frac{k e_{ii}}{r(r^2 + k)} \right\} + \frac{(n-1)e_{nn}}{r} \right) d^{n-1} \mu_{h_k} . \tag{4.13}$$

Here

$$e_{ij} := g_{ij} - b_{ij}$$

denotes the frame components in a b-ON frame, with the n'th component corresponding to the direction orthogonal to the level sets of r.

In spacetime dimension 3+1, which will be of main interest below, under the decay conditions spelled above this simplifies to

$$H(V_{(0)}) = \lim_{R \to \infty} R^4 \int_{\{r=R\}} \left(\frac{2e_{33}}{r} - \sum_{i=1}^2 \frac{\partial e_{ii}}{\partial r} \right) d^2 \mu_{h_k} . \tag{4.14}$$

A class of ALH metrics of interest are those for which

$$e_{ij} = r^{-n}\mu_{ij} + o(r^{-n}),$$
 (4.15)

where the μ_{ij} 's depend only upon the coordinates x^A on ∂M . One can further specialise the coordinates in which $\mu_{nn} \equiv 0$, but this choice might be unnecessarily restrictive for some calculations. Metrics with this asymptotics are dense in the space of all AH metrics in suitable circumstances [22, Theorem 5.3]. The tensor μ_{ij} will be referred to as the mass aspect tensor.

For metrics in which (4.15) holds, Equation (4.13) reads

$$H(V_{(0)}) = \int_{\partial M} \left((n-1)\mu_{nn} + n \sum_{i=1}^{n-1} \mu_{ii} \right) d^{n-1}\mu_{h_k}.$$
 (4.16)

An elegant formula for mass can be derived by integration by parts in (4.3)-(4.4), leading to the following: To every KID (b, V) and conformal-boundary component ∂M one associates a mass

$$m = m(V) = m(V, \partial M) \equiv H(V, b)$$

by the formula [30] (compare [5, Equation (IV.40)])

$$m(V, \partial M) = -\lim_{x \to 0} \int_{\{x\} \times \partial M} D^j V(R^i{}_j - \frac{R}{n} \delta^i_j) d\sigma_i, \qquad (4.17)$$

where R_{ij} is the Ricci tensor of the metric g, R its trace, and we have ignored an overall dimension-dependent positive multiplicative factor which is typically included in the physics literature. Here ∂M is a component of conformal infinity, x is a coordinate near ∂M so that ∂M is given by the equation $\{x = 0\}$, and $d\sigma_i := \sqrt{\det g} dS_i$.

As a special case, consider a triple (M, g, \hat{V}) satisfying (4.1)-(4.2); thus both (M, g, \hat{V}) and (M, b, V) are static KIDs. Assume that g asymptotes to b as above, and that \hat{V} asymptotes to V. Using the methods of [19] one checks that V can be replaced by \hat{V} in the integrand of (4.17). One has

$$D^{j}\hat{V}(R^{i}{}_{j} - \frac{R}{n}\delta^{i}{}_{j}) = \frac{1}{\hat{V}}D^{j}\hat{V}(D_{i}D_{j}\hat{V} - \frac{1}{n}\Delta\hat{V}\delta^{i}{}_{j})$$

$$= \frac{1}{\hat{V}}D^{j}\hat{V}(D_{i}D_{j}\hat{V} - \hat{V}\delta^{i}{}_{j})$$

$$= \frac{1}{2\hat{V}}D^{i}(|d\hat{V}|^{2}) - D^{i}\hat{V}. \tag{4.18}$$

Letting r = 1/x, (4.17) becomes

$$m(\hat{V}) = \lim_{R \to \infty} \int_{r=R} \left(D^i \hat{V} - \frac{1}{2\hat{V}} D^i (|d\hat{V}|^2) \right) d\sigma_i.$$
 (4.19)

As an application, and for further use, we consider (4.19) for a Horowitz-Myers metric,

$$g = -r^2 dt^2 + \frac{dr^2}{r^2 - \frac{2m_c}{r^2}} + (r^2 - \frac{2m_c}{r^{n-2}})d\psi^2 + r^2 \mathsf{h}_{ab} d\theta^a d\theta^b , \qquad (4.20)$$

where $h_{ab}d\theta^ad\theta^b$, $a,b \in \{1,\ldots n-2\}$ is a flat (n-2)-dimensional metric, $m_c>0$ is a constant and ψ has a suitable period to guarantee smoothness at $r^n=2m_c$. The background is taken to be g with $m_c=0$, thus $\hat{V}=r$, $|d\hat{V}|^2=(r^2-\frac{2m_c}{r^{n-2}})$, which gives an integrand in (4.19) equal to

$$(D^r \hat{V} - \frac{1}{2\hat{V}} D^r (|d\hat{V}|^2)) r^{n-1} \sqrt{\det h} = -2(n-2) m_c (1 - \frac{2m_c}{r^n}),$$
 (4.21)

and thus total mass

$$m = -2(n-2)m_{\rm c}\,, (4.22)$$

where we have assumed that the area of the conformal boundary at infinity in the metric $d\psi^2 + h_{ab}dx^adx^b$ has been normalised to 1.

4.2 The spherical case

The question then arises, what happens with the mass under the boundary gluings of Section 2. The case when both manifolds have conformal boundary with spherical topology and with a metric conformal to the standard round metric is simplest, because then the maps $(\Lambda_1, \Lambda_2) \in \mathscr{P}$ act globally on collar neighbourhoods of N_1^{n-1} and N_2^{n-1} in M_1 and M_2 . The initial energy-momenta \mathbf{m}_1 of N_1^{n-1} and \mathbf{m}_2 of N_2^{n-1} are transformed to $\Lambda_1 \mathbf{m}_1$ of $\Lambda_2 \mathbf{m}_2$. Since the mass integrands are zero in the neck region, where the metric is exactly hyperbolic, one finds that the energy-momentum \mathbf{m} of the boundary-glued manifold is additive:

$$\mathbf{m} = \Lambda_1 \mathbf{m}_1 + \Lambda_2 \mathbf{m}_2 \,. \tag{4.23}$$

This is true for all dimensions $n \geq 3$.

A more detailed presentation of the spherical case can be found in [14], and note that most of the work there arose from the necessity to control the direction of the momenta \mathbf{m}_{a} .

4.3 Conformal rescaling of h_k

The remaining cases, as well as the Isenberg-Lee-Stavrov gluings, require more effort. Some preliminary remarks are in order.

After the gluing has been done, the initial "boundary metric" h_k of (3.1) will have to be conformally rescaled in general. Thus, there will be a function $\psi > 0$, defined on a subset of the boundary at infinity, such that

$$h_k = \psi^2 h_{\bar{k}} \,, \tag{4.24}$$

for some constant $\bar{k} \in \{0, \pm 1\}$.

The following can be justified by considerations revolving around the Yamabe problem and the Obata equation.

First, if (N^{n-1}, h_k) is a round sphere, there exist globally defined non-trivial such conformal factors with $k = \bar{k} = 1$, and they all arise from conformal isometries of the sphere.

Next, when (N^{n-1}, h_k) is a closed manifold with k = 0 and (4.24) holds globally, then \bar{k} must also be zero and ψ must be constant. Of course any constant will do, and there does not seem to be a geometrically preferred value for this constant. When calculating the mass of the Horowitz-Myers instanton we will normalise the volume of (N^{n-1}, h_k) to one, which removes the ambiguity.

In all other cases $\psi \equiv 1$ and $k = \bar{k}$ is the only possibility for globally defined functions ψ on closed manifolds.

However, we will also need (4.24) on subsets of N^{n-1} . Then non-trivial functions ψ are possible with $k \neq \bar{k}$ in dimension n = 3. By way of example, let

$$N^2 = \mathbb{T}^2 # \mathbb{T}^2 \,, \tag{4.25}$$

with the gluing occurring across a closed geodesic, cf. Figure 4.1. If we cut the connecting neck along the closed geodesic, each of the factors in (4.25) can be viewed as a torus $\mathbb{T}^2 \setminus D^2$ with an open disc removed, and there we can write $(k=0, \bar{k}=-1)$

$$h_{-1} = e^{\omega} h_0 \,, \tag{4.26}$$

where ω is a solution of the two-dimensional Yamabe equation,

$$\Delta_{h_0}\omega = 2e^{\omega}\,, (4.27)$$




Figure 4.1: Gluing two solid tori along a disc, the boundary of which becomes a closed geodesic on the boundary of the glued manifold.

with vanishing Neumann data at ∂D^2 .

Now, our definition of mass requires that the glued manifold admits non-trivial static potentials. This is guaranteed when both the initial manifolds (N_i^{n-1}, h_{k_i}) , i = 1, 2, and the new metric on $N_1^{n-1} \# N_2^{n-1}$ are Einstein manifolds. The gluing construction itself does not care about this, but the existence of a static potential on a collar neighborhood of the glued manifold is not clear, except when the boundary is two dimensional, or when at least one boundary metric is a round sphere in all dimensions. The existence of further higher dimensional such examples is unlikely, see [8].

Here one should keep in mind that there exist large families of smoothly compactifiable stationary vacuum solutions (M, g) of (Riemannian) Einstein equations such that (N^{n-1}, h_k) is not Einstein [3, 4, 11, 15, 16]. Each such metric comes equipped with a definition of mass [27] for metrics with the same conformal infinity. Whether or not our weighted addition formula (4.37) below applies in these more general circumstances is not clear; we plan to address this in the future.

Returning to our main line of thought, suppose thus that (4.24) holds on a subset of N^{n-1} . We extend ψ to a neighbourhood of the conformal boundary by requiring $\partial_r \psi = 0$. Let us denote by (\bar{r}, \bar{x}^A) coordinates such that the new background metric $h_{\bar{k}}$ takes the form

$$\bar{b} := \frac{d\bar{r}^2}{\bar{r}^2 + \bar{k}} + \bar{r}^2 h_{\bar{k}} \,. \tag{4.28}$$

We can write

$$b = \frac{dr^2}{r^2 + k} + r^2 h_k = \frac{dr^2}{r^2 + k} + r^2 \psi^2 h_{\bar{k}} = \frac{d\bar{r}^2}{\bar{r}^2 + \bar{k}} + \bar{r}^2 (h_{\bar{k}} + \delta h), \qquad (4.29)$$

with

$$\delta h(\partial_r, \cdot) = 0$$
.

Let us denote by x^A the local coordinates at ∂M . As shown in Appendix C.1 below (cf. (C.21) and (C.24)) we find for large r

$$\bar{x}^A = x^A + \frac{\mathcal{D}^A \psi}{2\psi r^2} + O(r^{-4}),$$
 (4.30)

$$\bar{r} = \psi r \left(1 + \frac{\left(-\mathscr{D}^A \mathscr{D}_A \log \psi + (n-2) \mathscr{D}^A \log \psi \mathscr{D}_A \log \psi \right)}{2(n-1)r^2} + O(r^{-4}) \right), \tag{4.31}$$

where \mathscr{D} is the Levi-Civita connection of h_k . The pattern is, that the expansion jumps by two powers of r up to the threshold associated with the mass aspect tensor.

It follows from (4.30)-(4.31) (see (C.14), Appendix C) that an asymptotic expansion of the initial metric of the form

$$g(f_i, f_i) - b_{ij} = O(r^{-n}) (4.32)$$

(which is observed in BK metrics) will not be preserved by the above transformations, except possibly 1) in dimension n=3 or 2) when $k=\bar{k}$ and the conformal factor arises from a conformal isometry of h_k . This is addressed in Appendix C. In particular it is shown there, in space-dimension three and in the gauge $\mu_{ri}=0$, that the mass aspect tensor μ_{ij} of (4.15) transforms as

$$\mu_{AB} \mapsto \bar{\mu}_{AB} = \psi \mu_{AB} \,. \tag{4.33}$$

More generally, one can define the mass-aspect tensor as the coefficient of r^{-n} in the asymptotic expansion of the frame components of the metric. Then (4.33) remains valid in odd space dimensions because of the structure (4.30)-(4.31) of the expansion of the coordinate transformation, with powers jumping by two. But (4.33) does not hold in even space dimensions: in Appendix C.3 one can find the transformation formula in dimension n = 4.

4.4 Gluing in three space dimensions

We consider the mass of the manifold obtained by boundary-gluing of two three-dimensional manifolds (M_a, g_a) , a = 1, 2. This includes the Horowitz-Myers metrics which are conformally compactifiable, have no boundary, and have negative mass, as well as Kottler metrics which are conformally compactifiable and have totally geodesic boundaries.

Let us denote by V_a the static potential of the asymptotic backgrounds on M_a , a = 1, 2. For the spherical components of the boundary at infinity, if any, one only needs to consider the static potential $\sqrt{1+r^2}$, as follows from (4.36) below.

We start with the boundary-gluing of Section 2. Letting x = 1/R in (4.17), and using the fact that the metric is exactly hyperbolic in \mathcal{U}_a we have

$$m_{a} := m(V_{a}, \partial M_{a}) = -\lim_{x \to 0} \int_{\{x\} \times \partial M_{a}} D^{j} V_{a} (R^{i}{}_{j} - \frac{R}{n} \delta^{i}_{j}) d\sigma_{i}$$

$$= -\lim_{x \to 0} \int_{\{x\} \times (\partial M_{a} \setminus \partial \mathscr{U}_{a})} D^{j} V_{a} (R^{i}{}_{j} - \frac{R}{n} \delta^{i}_{j}) d\sigma_{i}.$$

$$(4.34)$$

Let us write r for the radial coordinate as in (3.1) on M_1 , so that the static potential entering into the definition of the mass of (M_1, g_1) reads

$$V_1 = \sqrt{r^2 + k} \,. \tag{4.35}$$

Let us write (\bar{M}, \bar{g}) for the boundary-glued manifold, \bar{V} for the associated static potential, and \bar{r} for the radial coordinate on \bar{M} . Viewing $M_1 \setminus \mathcal{U}_1$ as a subset of \bar{M} , using (4.31) with ψ there denoted by ψ_1 here, we can write on $M_1 \setminus \mathcal{U}_1$

$$\bar{V} = \sqrt{\bar{r}^2 + \bar{k}} = \bar{r} + O(\bar{r}^{-1}) = \psi_1 r + O(r^{-1}) = V_1 \psi_1 + O(r^{-1}). \tag{4.36}$$

Here ψ_1 has been extended to the interior of \bar{M}_1 by requiring $\partial_r \psi_1 = 0$.

An identical formula holds on $M_2 \setminus \mathcal{U}_2$, with a conformal factor which we denote by ψ_2 .

Assuming in addition to (4.6)-(4.10) that the *b*-norm of $R^i{}_j - \frac{R}{n} \delta^i_j$ decays as $o(r^{2-n})$ for large r, the above leads to

$$\bar{m} := m(\bar{V}, \partial \bar{M}) = -\lim_{x \to 0} \int_{\{x\} \times \partial \bar{M}} D^j \bar{V}(R^i{}_j - \frac{R}{n} \delta^i_j) d\sigma_i$$

$$= -\sum_{a=1}^2 \lim_{x \to 0} \int_{\{x\} \times \partial M_a} D^j (\psi_a V_a) (R^i{}_j - \frac{R}{n} \delta^i_j) d\sigma_i. \tag{4.37}$$

Comparing with (4.34), we see that the mass of the glued manifold is the sum of mass-type integrals, where the original integrands over ∂M_a are adjusted by the conformal factor relating the metric on the glued boundary to the original one.

The above applies also to the case where the mass aspect tensor is well-defined, giving a simpler formula

$$\bar{m} = -\sum_{a=1}^{2} \int_{\partial M_a} \psi_a \Big((n-1)\mu_{nn} + n \sum_{i=1}^{n-1} \mu_{ii} \Big) d^{n-1} \mu_{h_k} \,. \tag{4.38}$$

It should, however, be pointed-out that the gluing-in of an exactly hyperbolic region as in [13] is *not known* to lead to a metric with well defined mass-aspect tensor in the region where the deformation of the metric is carried-out, so that in our analysis below we have to use (4.37).

The difference in the mass formulae between the boundary-gluing of Section 2 and that of [33] is essentially notational, concerning the integration domains in (4.37) and (4.38). Indeed, after the gluing construction presented in Section 4.2 of [33] has been done, the boundary manifold there is obtained by removing discs $D(\epsilon_a)$ of coordinate radii ϵ_a from ∂M_a , and connecting the boundaries of the discs with a neck $S^1 \times [0,1]$. (The radii ϵ_a are taken to be one in [33, Section 4.2], which is due to a previous rescaling of the coordinates.) For the purpose of the formulae below let us cut the boundary neck $S^1 \times [0,1]$ into two pieces $N_1 := S^1 \times [0,1/2]$ and $N_2 := S^1 \times [1/2,1]$, and let $\Omega_a = (\partial M_a \setminus D(\epsilon_a)) \cup N_a$. In the construction of [33] each Ω_a comes naturally equipped with a constant scalar curvature metric which coincides with the original one on $\partial M_a \setminus D(\epsilon_a)$. The relative conformal factors ψ_a are defined on the Ω_a 's with respect to these metrics. We then have

$$\bar{m} := m(\bar{V}, \partial \bar{M})$$

$$= -\sum_{a=1}^{2} \lim_{x \to 0} \int_{\{x\} \times \Omega_a} D^j(\psi_a V_a) (R^i{}_j - \frac{R}{n} \delta^i_j) d\sigma_i.$$

$$(4.39)$$

Whenever a mass aspect tensor is globally defined on the glued manifold, this last formula coincides with

$$\bar{m} = -\sum_{a=1}^{2} \int_{\Omega_a} \psi_a \Big((n-1)\mu_{nn} + n \sum_{i=1}^{n-1} \mu_{ii} \Big) d^{n-1} \mu_{h_k} \,. \tag{4.40}$$

It should be clear how this generalises to the boundary-gluing of several three-dimensional manifolds.

4.5 Noncompact boundaries

When defining hyperbolic mass, it is usual to assume, and we did, that the conformal boundary at infinity is compact. This is done for convenience: when the boundary manifold is compact, to obtain convergence of the integrals defining mass it suffices to impose decay conditions of the metric g to the asymptotic background b. If the boundary manifold were not compact, one would need to impose further decay conditions in the non-compact directions at the boundary. However, it is not clear what conditions are relevant for physically interesting fields.

Now, our "exotic gluings" construction in [13] creates an open neighbourhood, say \mathcal{O} , of a subset of the conformal boundary at infinity, such that the metric is conformal to the hyperbolic one in \mathcal{O} . One can then create non-trivial non-compact boundary manifolds by removing closed sets, say Σ , from the boundary at infinity inside $\partial M \cap \mathcal{O}$ and changing the asymptotic background

b to a new background \bar{b} on $\partial M \setminus \Sigma$. A simple non-trivial example is provided by taking ∂M to be a two-dimensional torus, Σ to be a finite collection of points $p_i \in \partial M$ (e.g., one point), and replacing the original background with a flat conformal metric h_0 on ∂M by a complete hyperbolic metric with cusps at the (removed) points p_i . Then, instead of measuring the mass of M with respect to the original background with a toroidal boundary we can define a mass with respect to the new background \bar{b} , as in (4.28)-(4.29).

Since the removed set Σ is contained in the set \mathscr{O} , where the metric is conformal to the hyperbolic metric, the convergence of the new mass integrals readily follows from the convergence of the original mass: the support of the boundary integrals near infinity is included in a neighborhood of the compact set $\partial M \setminus \overline{\mathscr{O}}$. The arguments presented in Section 4.4 readily lead to the following formula for the new mass:

$$m(\bar{V}, \partial \bar{M} \setminus \Sigma) = -\lim_{x \to 0} \int_{\{x\} \times (\partial \bar{M} \setminus \Sigma)} D^{j}(\psi V) (R^{i}{}_{j} - \frac{R}{n} \delta^{i}_{j}) d\sigma_{i};$$

$$(4.41)$$

recall that ψ is the conformal factor which relates the original metric h_k to the new one as $h_{\bar{k}} = \psi^{-2} h_k$.

5 Higher genus solutions with negative mass

5.1 Genus two

In this section we wish to show that:

PROPOSITION 5.1 There exist conformally compactifiable three-dimensional ALH manifolds (M, g) without boundary, with genus-two topology at infinity, and with negative mass.

REMARK 5.2 In order to obtain the above conclusion we glue together two Horowitz-Myers instantons. However, all the action happens near the conformal infinity, with the geometry away from the conformal boundary being irrelevant. We show in fact that the gluing-together, as carried-out in the proof below, of two copies of an ALH manifold with toroidal infinity and mass aspect function which does not change sign results in an ALH manifold with higher-genus infinity and with total mass which has the same sign as the original one.

PROOF: The manifold (M, g) will be obtained by a boundary-gluing of two copies of a Horowitz-Myers instanton. More precisely, in the construction described in Section 2 we take (M_a, g_a) , a = 1, 2, to be two copies of a Horowitz-Myers instanton in which the hyperbolic metric has been glued in an ϵ -neighborhood of a boundary point p. We use the coordinates of (4.20) with polar coordinates for the boundary metric

$$h_0 = d\rho^2 + \rho^2 d\varphi^2$$

on ∂M_1 , with p located at the origin of these coordinates. Such coordinates can always be defined, covering a disc $D(\rho_0)$ for some $\rho_0 > 0$. Without loss of generality we can rescale the metric h_0 on ∂M if necessary so that the coordinates are defined on the unit disc D(1). (This might rescale the total mass by a positive factor; one can always scale back at the end of the construction.)

The metric g_1 is thus a Horowitz-Myers metric outside the half-ball $\mathcal{U}_{1,2\epsilon}$ of coordinate radius 2ϵ , and is exactly hyperbolic inside the half-ball $\mathcal{U}_{1,\epsilon}$ of coordinate radius ϵ ; see Figure 5.1. For reasons that will become clear in the proof we need to consider a sequence of boundary gluings indexed by a parameter $i \to \infty$. For $i \geq 8/\epsilon$ we choose the hyperbolic hyperplane $\mathfrak{h}_{1,i} \equiv \mathfrak{h}_{1,i}$




Figure 5.1: The sets $\mathscr{U}_{1,\epsilon} \subset \mathscr{U}_{1,2\epsilon}$ and their boundaries.

of Section 2 to be a half-sphere of radius 1/i. The manifold (M_2, g_2) of Section 2 is taken to be an identical copy of (M_1, g_1) . We choose any pair (Λ_1, Λ_2) as in Section 2 to obtain the boundary-glued manifold $M := M_{\Lambda_1, \Lambda_2}$.

By construction the boundary ∂M of the new manifold is a doubling of

$$\hat{M}_i := \partial M_1 \setminus \mathscr{U}_{1,1/i}$$

across its boundary $\partial \hat{M}_i$. We will often view both \hat{M}_i and its double as subsets of ∂M .

Consider the conformal class of metrics on ∂M induced by g. This conformal class depends upon i but is independent of ϵ , keeping in mind that $\epsilon \geq 8/i$. In this class there exists a unique metric h_{-1} with constant scalar curvature equal to minus two which can be constructed as follows: Let \hat{h} be any representative of the conformal class of metrics on ∂M_1 induced by g which is

1. invariant under reflection across $\partial \hat{M}_i$, so that the coordinate circle

$$S_{1/i} = \partial \hat{M}_i$$

of coordinate radius 1/i is a totally geodesic boundary for (\hat{M}_i, \hat{h}) , and which is

2. invariant under rotations near $S_{1/i}$.

While any metric \hat{h} as in 1. and 2. is adequate, a useful example is provided by the metric which on ∂M_1 equals $\chi^2 h_0$, where $\chi \geq 1/2$ is any smooth function on $\partial M_1 \setminus \{p\}$ which equals to $1/\rho$ on $D(1) \setminus \{0\}$, where ρ is the coordinate radius in the local coordinates on ∂M_1 . This can be accompanied by the introduction of a new coordinate

$$\hat{\rho} = \frac{\log(\rho)}{\log(i)} + 1, \qquad (5.1)$$

so that the flat metric

$$h_0 = d\rho^2 + \rho^2 d\varphi^2$$
, $\rho \in [1/i, 1]$

becomes

$$h_0 = \rho^2 (\frac{d\rho^2}{\rho^2} + d\varphi^2) = \rho^2 (\log^2(i)d\hat{\rho}^2 + d\varphi^2), \qquad \hat{\rho} \in [0, 1],$$
 (5.2)

thus $\hat{h} = \log^2(i)d\hat{\rho}^2 + d\varphi^2$ on $[1, i] \times S^1$.

In this coordinate system the reflection across $\partial \hat{M}_i$ is the map $(\hat{\rho}, \varphi) \mapsto (-\hat{\rho}, \varphi)$, after extending the range of $\hat{\rho}$ from [0, 1] to [-1, 1].

Given a function f defined on $\overline{D(1) \setminus D(1/i)}$ in the original coordinates (ρ, φ) , the mirror-symmetric doubling of f is defined as the function

$$[-1,1] \times S^1 \ni (\hat{\rho},\varphi) \mapsto \hat{f}(\hat{\rho},\varphi) = \begin{cases} f(e^{\log(i)(\hat{\rho}-1)},\varphi) & \hat{\rho} \ge 0; \\ f(e^{\log(i)(-\hat{\rho}-1)},\varphi) & \hat{\rho} < 0. \end{cases}$$
(5.3)

The mirror-symmetric doubling of a tensor field is obtained by transforming the tensor field to the coordinates $(\hat{\rho}, \varphi)$ and mirror-doubling all its coordinate-components. For example, after doubling of \hat{M}_i , the tensor field $\hat{h} = \log^2(i)d\hat{\rho}^2 + d\varphi^2$ just defined above on $[0,1] \times S^1$ maintains the same form $\log^2(i)d\hat{\rho}^2 + d\varphi^2$ on $[-1,1] \times S^1$ after doubling.

Another example, which plays an important role in our proof below, is provided by the daunting-looking metric (cf., e.g., [36])

$$e^{\omega_{*,i}} h_0 := \left(\frac{\pi}{\log(i^2)\sin(\pi \frac{\log \rho}{\log(i^2)})\rho}\right)^2 (d\rho^2 + \rho^2 d\varphi^2), \qquad \rho \in (1/i^2, 1), \tag{5.4}$$

which has constant negative scalar curvature equal to -2. The conformal factor $e^{\omega_{*,i}}$ tends to infinity at $\rho = 1/i^2$ and at $\rho = 1$. In the coordinates $(\hat{\rho}, \varphi)$ the metric (5.4) reads

$$e^{\omega_{*,i}}h_0 = \left(\frac{\pi}{2\cos(\pi\hat{\rho}/2)}\right)^2 (d\hat{\rho}^2 + \frac{1}{\log^2(i)}d\varphi^2), \qquad \hat{\rho} \in (-1,1),$$
 (5.5)

which is manifestly mirror-invariant. Note that the circle $\hat{\rho} = 0$ is a closed geodesic minimising length, of length $\pi^2/\log(i)$.

Let $u_i: \partial M \to \mathbb{R}^+$ be the unique solution of the two-dimensional Yamabe equation

$$\Delta_{\hat{h}} u_i = -Re^{u_i} + \hat{R} \,, \tag{5.6}$$

with R = -2, and where \hat{R} is the scalar curvature of the metric \hat{h} , so that the metric $e^{u_i}\hat{h}$ has scalar curvature R. It is important in what follows that the function u_i is independent of ϵ .

(As uniqueness is likewise important in our analysis, let us provide an argument: let \tilde{u}_i and \hat{u}_i be two solutions, rename the metric $e^{\hat{u}_i}\hat{h}$ to \hat{h} , then from \tilde{u}_i one obtains a solution of (5.6) with $R = \hat{R} = -2$. Multiplying the resulting equation by u_i and integrating over ∂M one obtains

$$\int |du_i|^2 = 2 \int (1 - e^{u_i}) u_i.$$
 (5.7)

Since the integrand of the right-hand side is negative except at $u_i = 0$, we find $u_i \equiv 0$, hence $\tilde{u}_i \equiv \hat{u}_i$.)

Uniqueness implies that u_i is invariant under reflection across $\partial \hat{M}_i$. Hence u_i has vanishing normal derivative on $S_{1/i}$.

As \hat{h} is conformal to h_0 on \hat{M}_i , there exists a function \hat{u} so that we have

$$\hat{h} = e^{\hat{u}} h_0$$

on \hat{M}_i . Then the metric

$$e^{u_i+\hat{u}}h_0$$

defined on \hat{M}_i , has scalar curvature equal to minus two. The mirror-symmetric doubling of \hat{h} across $S_{1/i}$ coincides with the metric h_{-1} on ∂M_1 .

Now, by construction, \hat{u} is rotation-invariant near $S_{1/i}$. The fact that $S_{1/i}$ is totally geodesic is equivalent to the vanishing of the radial derivative of $e^{\hat{u}/2}\rho$ on $S_{1/i}$:

$$0 = \partial_{\rho}(e^{\hat{u}/2}\rho)|_{S_{1/i}} \qquad \Longrightarrow \qquad \partial_{\rho}\hat{u}|_{S_{1/i}} = -2i. \tag{5.8}$$

From what has been said we have

$$\partial_{\rho} u_i|_{S_{1/i}} = 0. \tag{5.9}$$

Since h_0 is flat, we conclude that the function

$$\omega_i := u_i + \hat{u}$$

satisfies on \hat{M}_i the equation

$$\Delta_{h_0}\omega_i = 2e^{\omega_i}\,, (5.10)$$

with Neumann boundary data at the coordinate circle $S_{1/i}$ of radius 1/i centered at the origin:

$$\partial_{\rho}\omega_i|_{S_{1/i}} = -2i. \tag{5.11}$$

Now, while the function ω_i depends only upon i, the metric on M depends upon both ϵ and i. This has the unfortunate consequence that the sign of the mass of M is not clear. In what follows we will determine this sign for i large and ϵ small. For this we need to understand what happens with the mass integrand (4.37) when i tends to infinity and ϵ tends to zero. The idea of the argument is to show that the sequence of functions ω_i , or at least a subsequence thereof, converges to a limit, with sufficient control of the limit to guarantee control of the mass. The needed convergence result is perhaps contained in [43], but it is not completely apparent to us that this is the case, so we provide a direct argument, different from the one in [43].

The maximum principle shows that ω_i has no interior maximum on the compact manifold with boundary $\partial M_1 \setminus D(a)$ for $a \in [1/i, 1]$, where D(a) denotes an open coordinate disc of radius a lying on the boundary ∂M_1 and centered at p.

Integrating (5.10) over \hat{M}_i and using (5.11) one finds

$$\int_{\hat{M}_i} e^{\omega_i} d\mu_{h_0} = 2\pi \tag{5.12}$$

which is essentially the same as the Gauss-Bonnet theorem.

We continue the proof of Proposition 5.1 with the (well known, cf. e.g. [36]) observation, that solutions of the equation

$$\Delta\omega = 2e^{\omega} \tag{5.13}$$

satisfy a comparison principle, which we formulate in the simplest form sufficient for our purposes: If both ω and $\hat{\omega}$ satisfy (5.13) in a bounded domain Ω with boundary, and if $\hat{\omega} > \omega$ near the boundary, then $\hat{\omega} > \omega$ on Ω . This comparison is particularly useful for functions $\hat{\omega}$ which tend to infinity when $\partial\Omega$ is approached. Indeed, if ω is continuous on the closure $\bar{\Omega}$ of Ω , the function $\omega - \hat{\omega}$ is then negative near $\partial\Omega$, and satisfies the equation

$$\Delta_{h_0}(\omega - \hat{\omega}) = 2e^{\omega} - 2e^{\hat{\omega}}. \tag{5.14}$$

If $\omega > \hat{\omega}$ somewhere, then $\omega - \hat{\omega}$ would have a positive maximum at some point $q \in \Omega$ away from the boundary. Since the function $x \mapsto 2e^x$ is increasing, the right-hand side of (5.14) would be positive at q. But the left-hand side is nonpositive at a maximum, a contradiction. We rephrase this loosely as

$$\hat{\omega} > \omega \text{ on } \partial\Omega \implies \hat{\omega} > \omega \text{ on } \Omega.$$
 (5.15)

For 0 < a < b let

$$\Gamma(a,b) := D(b) \setminus D(a)$$
.

The comparison principle allows us to prove:

Lemma 5.3 On $\Gamma(1/i, 1)$ it holds that

$$e^{w_i} \le e^{\omega_{*,i}} := \left(\frac{\pi}{\log(i^2)\sin(\pi \frac{\log \rho}{\log(i^2)})\rho}\right)^2.$$
 (5.16)

Remark 5.4 On the circle $\rho = 1/i$ we have

$$e^{\omega_{*,i}} = \frac{\pi^2 i^2}{4 \log^2(i)},$$

which is unbounded in i, but the hyperbolic length ℓ_i of $S_{1/i}$ equals

$$\ell_i = \frac{1}{i} \int_{\varphi \in [0,2\pi]} (e^{\omega_i/2})_{|\rho=1/i} d\varphi \le \frac{1}{i} \int_{\varphi \in [0,2\pi]} (e^{\omega_{*,i}/2})_{|\rho=1/i} d\varphi = \frac{\pi^2}{\log i},$$

so that ℓ_i approaches zero as i tends to infinity.

PROOF OF LEMMA 5.3: The metric (compare (5.4))

$$e^{\omega_{*,\sqrt{t}}}h_0 = \left(\frac{\pi}{|\log t|}\csc(\pi \frac{\log|z|)}{\log|t|}\right)^2 \frac{|dz|^2}{|z|^2}$$
 (5.17)

has constant negative scalar curvature equal to -2. Furthermore, the circle $S_{1/i}$ is minimal when $t-i^2$.

$$\partial_{\rho}(\rho e^{\frac{\omega_{*,i}}{2}})|_{\rho=1/i} = 0.$$
 (5.18)

The conformal factor $e^{\omega_{*,i}}$ tends to infinity at $\rho = 1/i^2$ and at $\rho = 1$.

The change of the complex variable $z\mapsto w=-1/(i^2\bar{z})$, where \bar{z} is the complex conjugate of z, reproduces the metric on $w\in\Gamma(1/i^2,1)$ and exchanges the regions on both sides of the geodesic $|z|=|w|=1/i.^1$ Our doubled metric on ∂M is likewise symmetric relatively to this geodesic. The conformal factor e^{ω_i} , extended to $\rho\in(1/i^2,1/i)$ using the map just described, provides a smooth solution of the Yamabe equation (5.13) on $\Gamma(1/i^2,1)$ (compare the discussion around (5.3)). It takes finite values both at $\rho=1/i^2$ and at $\rho=1$. The result follows from the comparison principle.

COROLLARY 5.5 For any $\rho_1 \in (0,1)$, there exists a constant $\hat{c} = \hat{c}(\rho_1)$ such that

$$\omega_i < \hat{c}$$

on $\hat{M}_i \setminus D(\rho_1)$, independently of i.

¹The alternative map $z \mapsto w = 1/(i^2 z)$, preferable from the point of view of complex geometry, could also be used, but $z \mapsto w = -1/(i^2 \bar{z})$ fits better with (5.3).

Proof: In view of

$$e^{w_i} \le e^{\omega_{*,i}} = \left(\frac{\pi}{\log(i^2)\sin(\pi \frac{\log \rho_1}{\log(i^2)})\rho_1}\right)^2 \longrightarrow_{i \to +\infty} \frac{1}{\rho_1^2 \log^2(\rho_1)},\tag{5.19}$$

the ω_i 's are bounded by a constant $\hat{c}(\rho_1)$ independently of i on $S(\rho_1)$ for $\rho_1 \in (0,1]$. The result follows now from the maximum principle.

The corollary gives an estimation of the conformal factors from above. In order to prove convergence of the sequence ω_i , away from the puncture, we also need to bound the sequence of conformal factors away from zero. The next lemma provides a first step towards this. At its heart lies the inequality (5.22), which shows that the area does not concentrate near the minimal geodesic $S_{1/i}$.

Lemma 5.6 For all ε sufficiently small and all i sufficiently large we have the bounds

$$\inf_{\partial \hat{M}_i \setminus D(\varepsilon/2)} (\omega_i - \log 2) \le \log \frac{\pi}{|\partial M_1 \setminus D(\varepsilon/2)|_{h_0}} \le \sup_{\partial \hat{M}_i \setminus D(\varepsilon/2)} \omega_i, \tag{5.20}$$

and note that the middle term is independent of i.

Proof: We have

$$\int_{\Gamma(a,b)} e^{\omega_{*,i}} d\mu_{h_0} = -\frac{2\pi^2 \cot\left(\frac{\pi \log(\rho)}{\log(i^2)}\right)}{\log(i^2)} \bigg|_a^b, \tag{5.21}$$

and note that

$$\int_{\Gamma(1/i,\varepsilon/2)} e^{\omega_{*,i}} d\mu_{h_0} = \frac{2\pi^2 \cot\left(\frac{\pi \log(2/\varepsilon)}{\log(i^2)}\right)}{\log(i^2)} \to_{i\to\infty} \frac{2\pi}{\log(2/\varepsilon)}, \tag{5.22}$$

Equation (5.12) gives

$$2\pi = \int_{\hat{M}_i} e^{\omega_i} d\mu_{h_0} = \int_{\partial \hat{M}_i \setminus D(\varepsilon/2)} e^{\omega_i} d\mu_{h_0} + \int_{\Gamma(1/i,\varepsilon/2)} e^{\omega_i} d\mu_{h_0}. \tag{5.23}$$

The estimate (5.16) shows that

$$\int_{\Gamma(1/i,\varepsilon/2)} e^{\omega_i} d\mu_{h_0} \leq \int_{\Gamma(1/i,\varepsilon/2)} e^{\omega_{*,i}} d\mu_{h_0} = \frac{2\pi^2 \cot\left(\frac{\pi \log(2/\varepsilon)}{\log(i^2)}\right)}{\log(i^2)}.$$
 (5.24)

Equation (5.22) makes it clear that there exists ε_0 such that for all $2/i < \varepsilon \le \varepsilon_0$ the last term in (5.23) is in $(0, \pi)$, which implies

$$\pi \le \int_{\partial \hat{M}_i \setminus D(\varepsilon/2)} e^{\omega_i} d\mu_{h_0} < 2\pi.$$
 (5.25)

The conclusion readily follows using

$$\int_{\partial \hat{M}_i \setminus D(\varepsilon/2)} e^{\inf \omega_i} d\mu_{h_0} \le \int_{\partial \hat{M}_i \setminus D(\varepsilon/2)} e^{\omega_i} d\mu_{h_0} \le \int_{\partial \hat{M}_i \setminus D(\varepsilon/2)} e^{\sup \omega_i} d\mu_{h_0}.$$
 (5.26)

We continue with:

Lemma 5.7 There exists a smooth function

$$\omega_{\infty}: \partial M_1 \setminus \{p\} \to \mathbb{R}$$

such that a subsequence of $\{\omega_{i_j}\}_{j\in\mathbb{N}}$ converges uniformly, together with any number of derivatives, to ω_{∞} on every compact subset of $\partial M_1 \setminus \{p\}$.

PROOF: Let K be any compact subset of $\partial M_1 \setminus \{p\}$, there exists $\rho_K > 0$ such that $K \subset \partial M_1 \setminus D(\rho_K)$. It thus suffices to prove the result with $K = \partial M_1 \setminus D(\rho_K)$, which will be assumed from now on.

Let $K_1 = \partial M_1 \setminus D(\rho_K/2)$. Taking $\varepsilon = \rho_K$ in Lemma 5.6 ensures that (5.20) holds on K_1 for all $i \geq i_1$ for some $i_1 < \infty$.

Let $i \geq i_1$, by Corollary 5.5 there exists a constant c_1 , independent of i, such that

$$v_i := -\omega_i \ge c_1 \tag{5.27}$$

on K_1 . Define

$$\hat{v}_i := v_i - c_1 + 1$$
.

It holds that $\hat{v}_i \geq 1$ on K_1 . Moreover, \hat{v}_i satisfies the equation²

$$\Delta_{h_0} \hat{v}_i = \psi_i \hat{v}_i \,, \tag{5.28}$$

where

$$0 \ge \psi_i := -2\frac{e^{-v_i}}{\hat{v}_i} \ge -2e^{-c_1}. \tag{5.29}$$

By Harnack's inequality there exists a constant $C_1 = C_1(K, K_1) > 0$ such that on K we have

$$\sup_{K} \hat{v}_i \le C_1 \inf_{K_1} \hat{v}_i. \tag{5.30}$$

This, together with the definition of \hat{v}_i , shows that

$$\sup_{K} v_i \le C_1 \inf_{K_1} v_i + d_1 = C_1(-\sup_{K_1} \omega_i) + d_1,$$
(5.31)

for some constant d_1 . Equation (5.20) gives

$$-\sup_{K_1} \omega_i \le -\ln \frac{\pi}{|\partial M_1 \setminus D(\varepsilon/2)|_{h_0}} =: c_2,$$
(5.32)

From (5.31) we obtain

$$\sup_{K} v_{i} \le C_{1}c_{2} + d_{1} \implies \inf_{K} \omega_{i} = -\sup_{K} v_{i} \ge -(C_{1}c_{2} + d_{1}). \tag{5.33}$$

This, together with (5.27), shows that that for every compact subset K of $\partial M_1 \setminus \{p\}$ there exists a constant \hat{C}_K such that

$$-\hat{C}_K \le \omega_i \le \hat{C}_K \,. \tag{5.34}$$

Elliptic estimates, together with a standard diagonalisation argument, show that there exists a subsequence ω_{i_j} which converges uniformly on every compact subset of $\partial M_1 \setminus \{p\}$ to a solution ω_{∞} of (5.10) on $\partial M_1 \setminus \{p\}$. Convergence of derivatives follows again from elliptic estimates. \square

²We are grateful to Yanyan Li for providing the argument leading to (5.31).

We continue the proof of Proposition 5.1. By [10] there exists $\alpha > -2$ such that for small ρ we have

$$\omega_{\infty} = \alpha \log \rho + O(1) \implies e^{\omega_{\infty}} \sim \rho^{\alpha},$$
 (5.35)

or

$$\omega_{\infty} = -2\log(-\rho\log\rho) + O(1) \qquad \Longrightarrow \qquad e^{\omega_{\infty}} \sim \rho^{-2}\log^{-2}\rho. \tag{5.36}$$

(In fact, one can show that only (5.36) is relevant for our construction, but this is not needed for what follows.)

Consider the mass $m_{i,\epsilon}$ of M. By construction, M is the union of two copies of $M_1 \setminus \mathscr{U}_{1,1/i}$, identified along the boundary $\mathfrak{h}_{1,i}$. Since the metric is exactly hyperbolic in $\mathscr{U}_{1,\epsilon}$, the mass integrand vanishes in $\mathcal{U}_{1,\epsilon} \setminus \mathcal{U}_{1,1/i}$. From (4.37) we have

$$m_{i,\epsilon} = -2 \lim_{x \to 0} \int_{\{x\} \times \partial M_1} D^j(\psi_i V_1) (R^k{}_j - \frac{R}{n} \delta^k_j) d\sigma_k$$
$$= -2 \lim_{x \to 0} \int_{\{x\} \times (\partial M_1 \setminus D(\epsilon))} D^j(\psi_i V_1) (R^k{}_j - \frac{R}{n} \delta^k_j) d\sigma_k, \qquad (5.37)$$

where

$$\psi_i = e^{-\omega_i/2} \,. \tag{5.38}$$

We will need to estimate the derivatives of the functions ψ_i . As a step towards this, for $y \in \overline{D(4) \setminus D(1/2)}$ we set

$$f_{i,\epsilon}(y) = \begin{cases} \omega_i(\epsilon y) - \alpha \log(\epsilon |y|), & \text{under } (5.35); \\ \omega_i(\epsilon y) + 2\log(-\epsilon |y|\log(\epsilon |y|)), & \text{under } (5.36). \end{cases}$$
(5.39)

Each of the functions $f_{i,\epsilon}$ satisfies on $D(4) \setminus \overline{D(1/2)}$ the equation

$$\Delta_{h_0} f_{i,\epsilon} = \begin{cases} 2\epsilon^{2+\alpha} |y|^{\alpha} e^{f_{i,\epsilon}}, & \text{under (5.35);} \\ \frac{2e^{f_{i,\epsilon}}}{|y|^2 (\log(\epsilon|y|))^2} - \frac{2}{|y|^2 (\log(\epsilon|y|))^2}, & \text{under (5.36).} \end{cases}$$
(5.40)

For every ϵ there exists $i_0(\epsilon)$ such that the functions $f_{i,\epsilon}$ are bounded uniformly in ϵ and i for $i \geq i_0$. This leads to an estimate on the derivatives of $f_{i,\epsilon}$ on $D(2) \setminus D(1)$ (see [25, Section 3.4]):

$$|\partial f_{i,\epsilon}| + |\partial \partial f_{i,\epsilon}| \le C, \tag{5.41}$$

for some constant C independent of ϵ and of i provided that $i \geq i_0(\epsilon)$.

In view of (5.37)-(5.38), we will need estimates for $e^{-\omega_i/2}$. From (5.41) one obtains on $\overline{D(2\epsilon) \setminus D(\epsilon)}$, for $i \geq i_0(\epsilon)$,

$$|\partial_A e^{-\omega_i/2}| \le C\epsilon^{-1} e^{-\omega_i/2}, \qquad |\partial_A \partial_B e^{-\omega_i/2}| \le C\epsilon^{-2} e^{-\omega_i/2}, \tag{5.42}$$

for some possibly different constant C which is independent of ϵ and i when $i \geq i(\epsilon)$.

Directly from [10], or replacing v_i by v_{∞} in the argument just given, one finds

$$|\partial_A e^{-\omega_{\infty}/2}| \le C\rho^{-1} e^{-\omega_{\infty}/2}, \qquad |\partial_A \partial_B e^{-\omega_{\infty}/2}| \le C\rho^{-2} e^{-\omega_{\infty}/2}, \tag{5.43}$$

for possibly yet another constant C.

Near $\partial M_1 \setminus \mathcal{U}_{1,2\epsilon}$ in M the metric g is the Horowitz-Myers metric (4.20) with n=3. Using the fact that the Ricci tensor of g is block-diagonal (cf. (B.60), Appendix B.2 below), so that

the $D^i\psi_i$ terms in the integrand give no contribution on the level sets of x, together with (4.21) with n=3 we obtain

$$m_{i,\epsilon} = -2m_{c} \int_{\partial M_{1} \setminus D(2\epsilon)} \psi_{i} d\mu_{h_{0}}$$

$$-2 \lim_{x \to 0} \int_{\{x\} \times (\overline{D(2\epsilon) \setminus D(\epsilon/2)})} D^{j}(\psi_{i} V_{1}) (R^{k}{}_{j} - \frac{R}{n} \delta^{k}_{j}) d\sigma_{k} . \qquad (5.44)$$

$$=:(*)$$

Since $m_c > 0$ the first line is strictly negative, and we claim that the second line can be made arbitrarily small by choosing i sufficiently large and ϵ sufficiently small. For this we apply the divergence theorem on the set

$$\Omega_{\epsilon} := \{0 \le x \le \epsilon/100\} \times (\overline{D(2\epsilon) \setminus D(\epsilon/2)}).$$

Letting

$$X^{k} := (R^{k}{}_{j} - \frac{R}{n} \delta^{k}_{j}) D^{j}(\psi_{i} V_{1}), \qquad (5.45)$$

where ψ_i has been extended away from ∂M_1 by requiring $\partial_r \psi_i = 0$, we have

$$\int_{\Omega_{\epsilon}} D_k \mathbb{X}^k = (*) + \underbrace{\int_{\Omega_{\epsilon} \cap \{x = \epsilon/100\}} \mathbb{X}^k d\sigma_k}_{=:a} + \underbrace{\int_{[0,\epsilon/100] \times S_{2\epsilon}} \mathbb{X}^k d\sigma_k}_{=:b} + \underbrace{\int_{[0,\epsilon/100] \times S_{\epsilon/2}} \mathbb{X}^k d\sigma_k}_{=:c} . \quad (5.46)$$

On $[0, \epsilon/100] \times S_{\epsilon/2}$ the metric is exactly hyperbolic, thus c = 0.

On $[0, \epsilon/100] \times S_{2\epsilon}$ the metric is exactly Horowitz-Myers, and (4.18) together with (4.21) show that b = 0.

From the construction in [13] (see also [14, Remark 3.1]) for any $s \in [3/2, 2)$ we have the following estimates in Ω_{ϵ} (recall that D is the covariant derivative of g and r = 1/x)

$$|g - b|_b + |Db|_b + |D^2b|_b = O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}).$$
 (5.47)

Let $\bar{V} = \psi_i r \equiv \psi_i V_1$. Then $d\bar{V} = \psi_i dr + r \partial_A \psi_i dx^A$ so that $|d\bar{V}|_b = \sqrt{r^2 \psi_i^2 + |d\psi_i|_{h_0}^2}$. We thus have

$$|d\bar{V}|_b \le \psi_i r + |d\psi_i|_{h_0}, \quad |R^i{}_j - \frac{R}{n} \delta^i_j|_b = O(r^{-3}) + o(\varepsilon^{3-s} r^{-s}),$$
 (5.48)

so that at $r = 100/\epsilon$ we find

$$a = \int_{\overline{D(2\epsilon)\setminus D(\epsilon/2)}} \left(O(1)\psi_i + O(\epsilon)|d\psi_i|_{h_0} \right) d\mu_{h_0} . \tag{5.49}$$

Let us turn our attention to the integral over Ω_{ϵ} . A calculation shows that

$$\mathring{D}\mathring{D}\bar{V} = \bar{V}b + r\partial_A\partial_B\psi_i\,dx^Adx^B$$

(recall that \mathring{D} is the covariant derivative of the metric $b = dr^2/r^2 + r^2h_0$), so that

$$|\mathring{D}\mathring{D}\bar{V} - \bar{V}b|_b = \frac{\sqrt{h_0^{AB}h_0^{CD}\partial_A\partial_C\psi_i\partial_B\partial_D\psi_i}}{r} \equiv \frac{|\partial^2\psi_i|_{h_0}}{r} \,.$$

Thus, with the error terms for tensors in b-norm unless explicitly indicated otherwise,

$$D_{k}D_{j}\bar{V} = \mathring{D}_{k}\mathring{D}_{j}\bar{V} + (O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}))D\bar{V}$$

$$= \bar{V}b_{kj} + O(|\partial^{2}\psi_{i}|_{h_{0}}r^{-1}) + (O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}))D\bar{V}$$

$$= \bar{V}(g_{kj} + O(r^{-3}) + o(\varepsilon^{3-s}r^{-s})) + O(|\partial^{2}\psi_{i}|_{h_{0}}r^{-1})$$

$$+ (O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}))(\psi_{i}r + |d\psi_{i}|_{h_{0}})$$

$$= \bar{V}g_{kj} + O(|\partial^{2}\psi_{i}|_{h_{0}})r^{-1} + \psi_{i}(O(r^{-2}) + o(\varepsilon^{3-s}r^{1-s}))$$

$$+ |d\psi_{i}|_{h_{0}}(O(r^{-3}) + o(\varepsilon^{2-s}r^{-s})).$$
(5.50)

Next, the curvature scalar R is constant, which implies that $D_i R^i{}_j = 0$ by the (twice contracted) second Bianchi identity. We thus find

$$D_{k}X^{k} = \underbrace{(R^{k}_{j} - \frac{R}{n}\delta^{k}_{j})}_{O(r^{-3}) + o(\varepsilon^{3-s}r^{-s})} D_{k}D^{j}\bar{V}$$

$$= (O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}))(O(r^{-2}) + o(\varepsilon^{3-s}r^{-s+1}))\psi_{i}$$

$$+ (O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}))(O(r^{-3}) + o(\varepsilon^{3-s}r^{-s}))|d\psi_{i}|_{h_{0}}$$

$$+ (O(r^{-4}) + o(\epsilon^{3-s}r^{-s-1}))|\partial^{2}\psi_{i}|_{h_{0}}$$

$$= (O(r^{-5}) + o(\varepsilon^{3-s}r^{-s-2}) + o(\varepsilon^{6-2s}r^{-2s+1}))\psi_{i}$$

$$+ (O(r^{-6}) + o(\varepsilon^{3-s}r^{-s-3}) + o(\varepsilon^{6-2s}r^{-2s}))|d\psi_{i}|_{h_{0}}$$

$$+ (O(r^{-4}) + o(\epsilon^{3-s}r^{-s-1}))|\partial^{2}\psi_{i}|_{h_{0}}, \qquad (5.51)$$

leading to

$$\int_{\Omega_{\epsilon}} D_k \mathbb{X}^k = \int_{\overline{D(2\epsilon) \setminus D(\epsilon/2)}} \left(O(\epsilon^3) \psi_i + O(\epsilon^4) |d\psi_i|_{h_0} + O(\epsilon^2) |\partial^2 \psi_i|_{h_0} \right) d\mu_{h_0} . \tag{5.52}$$

Collecting terms in (5.46) leads to the following form of (5.44)

$$m_{i,\epsilon} = -2m_{\rm c} \int_{\partial M_1 \setminus D(2\epsilon)} \psi_i d\mu_{h_0}$$

$$+ \int_{\overline{D(2\epsilon) \setminus D(\epsilon/2)}} \left(O(1)\psi_i + O(\epsilon) |d\psi_i|_{h_0} + O(\epsilon^2) |\partial^2 \psi_i|_{h_0} \right) d\mu_{h_0} .$$
(5.53)

Let $\eta > 0$ and $\psi_{\infty} = e^{-\omega_{\infty}/2}$. By (5.35)-(5.36) and (5.43) we can choose ϵ small enough so that

$$\int_{\overline{D(2\epsilon)\backslash D(\epsilon/2)}} \left(O(1)\psi_{\infty} + O(\epsilon) |d\psi_{\infty}|_{h_0} + O(\epsilon^2) |\partial^2 \psi_{\infty}|_{h_0} \right) d\mu_{h_0} < \eta/2.$$
 (5.54)

The function ψ_i tends uniformly on $\overline{D(2\epsilon) \setminus D(\epsilon/2)}$ to ψ_{∞} when i tends to infinity. This, together with (5.42), shows that we can choose i large enough so that

$$\int_{\overline{D(2\epsilon)\backslash D(\epsilon/2)}} \left(O(1)|\psi_{\infty} - \psi_{i}| \right. \\
+ O(\epsilon)|d(\psi_{\infty} - \psi_{i})|_{h_{0}} + O(\epsilon^{2})|\partial^{2}(\psi_{\infty} - \psi_{i})|_{h_{0}} \right) d\mu_{h_{0}} < \eta/2.$$
(5.55)




Figure 5.2: A punctured torus with a hyperbolic metric; the figure is misleading in that the cusp region is actually infinitely long.

This shows that the integral over the annulus $\overline{D(2\epsilon) \setminus D(\epsilon/2)}$ tends to zero as ϵ tends to zero and i tends to infinity. Since the first integral in (5.44) tends to

$$-2m_{\rm c} \int_{\partial M_1 \setminus D(2\epsilon)} \psi_{\infty} d\mu_{h_0} < 0 \tag{5.56}$$

when i tends to infinity, we conclude that $m_{i,\epsilon}$ is indeed negative when ϵ is chosen small enough and i is chosen large enough. This finishes the proof or Proposition 5.1.

It is of interest to enquire about the shape of the glued boundary-manifold. With some work one can show that, as i tends to infinity, the metric on the conformal boundary converges (uniformly on compact sets away from the puncture) to the cusp metric on each of the two copies of the punctured torus (cf. Figure 5.2), with asymptotic behaviour near the puncture approximated by

$$\frac{1}{\rho^2 \log^2(\rho)} (dx^2 + dy^2) = \frac{1}{\rho^2 \log^2(\rho)} (d\rho^2 + \rho^2 d\varphi^2).$$
 (5.57)

This seems to follow from [43], and we note that a proof using our methods can be obtained in any case using the results in [26, 28]. Since

$$\int \frac{1}{\rho \log \rho} d\rho = \log(-\log \rho) \to_{\rho \to 0} \infty,$$

the connecting necks become longer and longer, with a circumference $\approx 2\pi/|\log \rho| = 2\pi/\log i$ tending to zero when $i \to \infty$, see Figure 5.3. In fact one can show that under the double limit when i tends to infinity first, and then ϵ tends to zero, the mass of M approaches (compare (4.41))

$$-2m_{c}\lim_{x\to 0} \int_{\{x\}\times(\partial M_{1})} D^{j}(e^{-\omega_{\infty}/2}V)(R^{i}{}_{j} - \frac{R}{n}\delta^{i}_{j}) d\sigma_{i}, \qquad (5.58)$$

when two general AH metrics with a toroidal conformal infinity are glued; this reduces to

$$-2m_{\rm c} \int_{\partial M_1} e^{-\omega_{\infty}/2} d\mu_{h_0} , \qquad (5.59)$$




Figure 5.3: Two punctured tori, connected by a thin neck. In the limit $i \to \infty$ the necks become longer and longer and their circumference shrinks to zero.




Figure 5.4: The gluing is done at the discs, whose boundaries become closed geodesics on the boundary, indicated in red in the figure. The construction is symmetric under the reflection across the horizontal plane passing through the center of the figure.

in the HM case.

In the case of the square torus with area equal one, discussed in Appendix A, we have the bound (cf. (A.2))

$$e^{-\omega_{\infty}/2} \le \frac{\Gamma(1/4)^2}{4\pi^{3/2}} \approx 0.59\,,$$
 (5.60)

which leads to a rough lower estimate of the mass m of the HM glued manifold, for small ϵ and large i,

$$-1.18 m_{\rm c} < m < 0. ag{5.61}$$

An upper bound can be obtained by numerically integrating the inverse of the right-hand side of (A.1), leading to

$$-1.18 \, m_{\rm c} < m < -0.67 \, m_{\rm c} \,. \tag{5.62}$$

(The integral of the inverse of the left-hand side of (A.1) gives a worse lower bound $\approx -4.02 \, m_{\rm c.}$)

5.2 Higher genus

It should be clear that one can iterate the construction of the last section to obtain three dimensional conformally compact ALH manifolds without boundary, with constant scalar curvature, negative mass, and a conformal infinity of arbitrary genus. One possible way of doing this proceeds as follows: Let (M_1, g_1) be a three dimensional ALH metric with constant scalar curvature and toroidal boundary at infinity; e.g., but not necessarily, a Horowitz-Myers instanton. Let $N \in \mathbb{N}$ and let $\{p_i\}_{i=1}^N$ be a collection of points lying on the conformal boundary ∂M_1 of M_1 . Let \mathcal{U}_i be any pairwise-disjoint family of coordinate half-balls of coordinate radius ϵ_i centered at the points p_i . Let (M_2, g_2) be an identical copy of (M_1, g_1) . The manifold (M, g) is taken to be a boundary-gluing of (M_1, g_1) with (M_2, g_2) , where each $\mathcal{U}_i \subset M_1$ is glued with its partner in M_2 in a symmetric way; Figure 5.4 illustrates what happens at the conformal boundary at infinity. Solving the Yamabe equation on ∂M , for $j \in \mathbb{N}$ one obtains a function ω_j on

$$\partial \hat{M}_i := \partial M_1 \setminus \left(\cup_{i=1}^N \partial \mathscr{U}_i \right)$$

such that the constant-Gauss-curvature representative of the conformal metric on ∂M equals

$$e^{\omega_j}h_0$$

on $\partial \hat{M}_i$, with ω_j satisfying the Neumann boundary condition (5.11) on each of the coordinate circles $S(p_i)_{1/j}$ centred at p_i :

$$\partial_{\rho}\omega_j|_{S_{1/j}} = -2j. \tag{5.63}$$

This leads to the integral identity

$$\int_{\hat{M}_i} e^{\omega_j} d\mu_{h_0} = 4N\pi \,, \tag{5.64}$$

The analysis of Section 5.1 applies near each of the punctures p_i . Letting $m(\epsilon_1, \dots, \epsilon_N, j)$ denote the mass of (M, g) we conclude that if we started with a Horowitz-Myers metric we will have

$$m(\epsilon_1, \cdots, \epsilon_N, j) < 0$$
 (5.65)

for all j larger than j_0 , for some j_0 that depends upon $\max \epsilon_i$.

From (5.64) we find

$$|\partial M|_{h_{-1}} = 2 \int_{\hat{M}_i} e^{\omega} d\mu_{h_0} = 4N\pi.$$
 (5.66)

Equivalently, since the scalar curvature $R_{h_{-1}}$ of the metric h_{-1} equals -2 we have

$$4\pi\chi(\partial M) = \int_{\partial M} R_{h_{-1}} d\mu_{h_{-1}} = -4 \int_{\hat{M}_i} e^{\omega} d\mu_{h_0} = -8N\pi, \qquad (5.67)$$

which shows that $\chi(\partial M) = -2N$, and thus ∂M has genus N+1.

A variation of the above is the following: Let $\psi:\partial M_1\to\partial M_1$ be an isometry of (M_1,h_0) such that $\psi\circ\psi$ is the identity map. Let N be even, say $N=2\hat{N}$, and suppose that the collection $\{p_i\}_{i=1}^N$ of N distinct points on ∂M_1 takes the form $\{q_i,\psi(q_i)\}_{i=1}^{\hat{N}}$. We can then pairwise glue, as above, neighborhoods of the points q_i with neighborhoods of their ψ -symmetric partners $\psi(q_i)$. Thus ∂M is obtained by adding to ∂M_1 a family of \hat{N} necks connecting the points q_i and $\psi(q_i)$; see Figure 5.5. If we choose ψ and the necks in a way compatible with the constructions described in Section 5.1, one obtains a family of ALH manifolds with constant scalar curvature, with conformal infinity of genus $\hat{N}+1$, with masses $m(\epsilon_1,\cdots,\epsilon_{\hat{N}},j)$ which are negative for j large enough.

6 A lower bound on mass?

It has been conjectured [32, 44] that the Horowitz-Myers instantons have the lowest mass within the class of ALH manifolds without boundary and with the same conformal metric at their toroidal infinity. If the Horowitz-Myers conjecture is wrong, then our construction would provide higher-genus constant-scalar-curvature metrics which have arbitrarily negative mass at fixed conformal class at infinity.

As such, one can envision a construction which cuts a metric with conformal infinity of higher genus into pieces, producing a finite-number of ALH metrics with toroidal infinity which would assemble together to the original one by a boundary-gluing in the spirit of this work. The validity of the Horowitz-Myers conjecture would then provide a lower bound for the mass of each summand. It is thus tempting to formulate the following conjecture




Figure 5.5: The symmetry ψ is a reflection across the vertical plane passing through the center of the figure. The gluing is done at the discs, whose boundaries become closed geodesics on the boundary, indicated in red in the figure. The construction is symmetric under the reflection across the horizontal plane passing through the center of the figure.

Conjecture 6.1 Let (M, g) be an n-dimensional conformally compact ALH manifold, without boundary with well defined mass m and with scalar curvature satisfying $R \ge -n(n-1)$. There exists a constant m_0 , which depends only upon the conformal class of the metric at infinity and which is negative if this conformal class is not that of the round sphere, such that

$$m \ge m_0. \tag{6.1}$$

It is also tempting to conjecture existence of unique static metrics as above which saturate the inequality if m_0 is optimal.

A A punctured torus

Recall that the model for the cusp (at $t = -\infty$ or $\rho = 0$) is :

$$dt^{2} + e^{2t}d\varphi^{2} = \frac{1}{\rho^{2}\log^{2}(\rho)}(d\rho^{2} + \rho^{2}d\varphi^{2}),$$

with $e^{-t} = -\log(\rho)$. Here the Gauss curvature is -1, thus the Ricci scalar equals -2.

The simplest hyperbolic one-punctured torus, also called "the hyperbolic torus with one cusp", is $(\mathbb{R}^2\backslash\mathbb{Z}^2)/\mathbb{Z}^2$ with a hyperbolic metric. Let us use the notation of [39] (keeping in mind a different scaling of the hyperbolic metric there) in order to compare the hyperbolic metric and the flat metric on this torus. For a domain D in \mathbb{C} , the complete hyperbolic metric on D with Gauss curvature equal to -1, as obtained using the Riemann mapping theorem applied to the universal covering space, is denoted by $4\rho_D^2(z)|dz|^2$. If $D \subset D'$ then by the comparison principle (5.15), also known as the Nevanlinna principle, we have $\rho_D \geq \rho_{D'}$. Let us set

$$D_0 = \hat{\mathbb{C}} \setminus \{0, 1, \infty\}, \quad \mathbb{D}^* = B(0, 1) \setminus \{0\}, \quad \mathbb{Z}[i] = \{m + in, m, n \in \mathbb{Z}\} = \mathbb{Z}^2, \quad \tilde{D} = \mathbb{C} \setminus \mathbb{Z}[i].$$

Because $\mathbb{D}^* \subset \tilde{D} \subset D_0$, by the Hempel inequality [28] (see, e.g., [39], inequality (1.2)) we have

$$\frac{1}{|z|(|\log(|z|)| + C_1)} \le 2\rho_{D_0}(z) \le 2\rho_{\tilde{D}}(z) \le 2\rho_{\mathbb{D}^*}(z) = \frac{1}{|z||\log(|z|)|}, \tag{A.1}$$

where $C_1 = \Gamma(1/4)^4/4\pi^2 \sim 4.37688$.

Let $p: \mathbb{C} \longrightarrow T_0 = \mathbb{C}/\mathbb{Z}[i]$ be the canonical projection and denote by [z] = p(z) the equivalence class of z. Let $D = T_0 \setminus \{[0]\}$, then D is proper subdomain of T_0 and $p^{-1}(D) = \tilde{D}$. On D, $\rho_D([z]) = \rho_{\tilde{D}}(z)$ is well defined. This defines the "hyperbolic density" ρ_{X_0} on $X_0 = \tilde{D}/\mathbb{Z}[i]$, which is the simplest one-punctured hyperbolic torus.

Because of the symmetries of D, the inequalities (A.1) on $\{x + iy, x, y \in [-1/2, 1/2]\}\setminus\{0\}$ provide an estimate of ρ_{X_0} throughout X_0 .

It follows from (A.1) that near the origin we have

$$2\rho_{\tilde{D}} = \frac{1}{|z||\log(|z|)|} \left(1 + O(\frac{1}{|\log(|z|)|} \right).$$

Note also that by [39] the minimum of ρ_{X_0} is attained at $\frac{1+i}{2}$ and is given by his formula (1.3):

$$\inf 2\rho_{X_0} = \frac{4\pi^{3/2}}{\Gamma(1/4)^2} \approx 1.695.$$
 (A.2)

B Static KIDs

In this appendix we find all static potentials (KIDs) for the Birmingham-Kottler metric and the Horowitz-Myers metric in (n+1)-dimensions. We start by computing the KIDs for the Birmingham-Kottler metric (B.1). We find that for $m_c \neq 0$ and $n \neq 2$ the solution to the KID equations is given by (B.45). Otherwise, the solution to the KID equations takes the form (B.27). For $m_c = 0$ and $n \neq 2$ β is of the form (B.36) or (B.37) and Ω is a solution to the differential equations (B.38). In the case n = 2 β is of the form (B.47) or (B.48) and Ω takes the form (B.50). Next, we consider the Horowitz-Myers metric (B.51), which for $m_c = 0$ is of Birmingham-Kottler type. For $m_c \neq 0$ we have that the solution to the KID equation is given by (B.76).

B.1 Birmingham-Kottler

The Birmingham-Kottler metric in (n+1)-dimensions, $n \geq 2$, which we denote by \tilde{g} , reads

$$\tilde{g} = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2\mathring{h}_{AB}dx^A dx^B,$$
(B.1)

where $\mathring{h}_{AB} = \mathring{h}_{AB}(x^C)$ where A, B = 1, ..., n-1, and with

$$f(r) = r^2 + k - \frac{2m_c}{r^{n-2}} \tag{B.2}$$

and

$$R(\mathring{h}) = k(n-1)(n-2)$$
. (B.3)

For n=2, R(h) vanishes and thus we set k=0 without loss of generality. This then yields that

$$f(r) = r^2 - 2m_c \tag{B.4}$$

for n=2. The metric (B.1) fulfills the Einstein equations with negative cosmological constant

$$\Lambda = -\frac{n(n-1)}{2} \,. \tag{B.5}$$

Using this normalisation of Λ , the Einstein equations read

$$R_{ab}(\tilde{g}) = -n\tilde{g}_{ab}, \qquad R(\tilde{g}) = -n(n+1), \tag{B.6}$$

where a, b run from 1, ..., n + 1. To compute the KIDs we need the Ricci scalar and the Ricci tensor of the spatial part of \tilde{g} , which we denote as g in what follows. These can be obtained from the Gauss-Codazzi equation [41]

$$R_{abcd}(g) = R_{abcd}^{\parallel}(\tilde{g}) - K_{ac}K_{bd} + K_{bc}K_{ad}. \tag{B.7}$$

where K_{ab} is the extrinsic curvature of the level sets of t (zero in our case)

$$K_{ab} = g_a{}^c \tilde{D}_c \tilde{n}_b \,, \tag{B.8}$$

with \tilde{n}_a being the normal one-form with the normalization chosen such that $\tilde{n}^a \tilde{n}_a = -1$. The superscript " \parallel " means that the respective tensor is projected to the submanifold, e.g. $R^{\parallel}_{abcd}(\tilde{g}) = g_a{}^e g_b{}^f g_c{}^g g_d{}^h R_{efgh}(\tilde{g})$. Contracting with g^{ac} we obtain

$$R_{ab}(g) = R_{ab}^{\parallel}(\tilde{g}) + \left(R_{cadb}(\tilde{g})\tilde{n}^c\tilde{n}^d\right)^{\parallel} - KK_{ab} + K_{ac}K^c_b \tag{B.9}$$

with [37, p. 518]

$$\left(R_{cadb}(\tilde{g})\tilde{n}^c\tilde{n}^d\right)^{\parallel} = -\mathcal{L}_{\tilde{n}}K_{ab} + K_{ac}K^c_{\ b} + D_{(a}a_{b)} + a_aa_b \tag{B.10}$$

where $a_a = \tilde{n}^b \tilde{D}_b \tilde{n}_a$, D_a denotes the covariant derivative of the metric g and symmetrization is defined with a factor 1/2 such that

$$R_{ab}(g) = R_{ab}^{\parallel}(\tilde{g}) - \mathcal{L}_{\tilde{n}}K_{ab} + 2K_{ac}K^{c}_{b} - KK_{ab} + D_{(a}a_{b)} + a_{a}a_{b}.$$
(B.11)

The normal one-form to t = const. surfaces reads

$$\tilde{n} = \sqrt{f}dt \tag{B.12}$$

such that $\tilde{n}_a \tilde{g}^{ab} \tilde{n}_b = -1$. Finally,

$$\tilde{g}_{ab} = g_{ab} - \tilde{n}_a \tilde{n}_b \tag{B.13}$$

yielding that in coordinates t, r, x^A , the tensor field g reads

$$g_{ab} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1/f(r) & 0 \\ 0 & 0 & r^2 \mathring{h}_{AB} \end{pmatrix}, \qquad g^a{}_b = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1_{AB} \end{pmatrix}$$
(B.14)

The only-nonvanishing Christoffel symbols (up to symmetries)

$$\tilde{\Gamma}^{c}{}_{ab} = \frac{1}{2}\tilde{g}^{cd} \left(\tilde{\partial}_{a}\tilde{g}_{bd} + \tilde{\partial}_{b}\tilde{g}_{ad} - \tilde{\partial}_{d}\tilde{g}_{ab} \right)$$
(B.15)

are

$$\tilde{\Gamma}^{t}_{tr} = \frac{\partial_{r} f}{2f}, \quad \tilde{\Gamma}^{r}_{rr} = -\frac{\partial_{r} f}{2f}, \quad \tilde{\Gamma}^{r}_{tt} = -\frac{f \partial_{r} f}{2}, \quad \tilde{\Gamma}^{r}_{AB} = -r \mathring{h}_{AB} f(r), \quad (B.16)$$

$$\tilde{\Gamma}^{C}{}_{rA} = \frac{\delta^{C}{}_{A}}{r} , \quad \tilde{\Gamma}^{C}{}_{AB} = \frac{1}{2} \mathring{h}^{CD} \left(\partial_{A} \mathring{h}_{BD} + \partial_{B} \mathring{h}_{AD} - \partial_{D} \mathring{h}_{AB} \right) . \tag{B.17}$$

As already mentioned, in our case all components of K_{ab} vanish, so that the expression of the Ricci tensor of the constant t submanifold (B.11) simplifies to

$$R_{ab}(g) = R_{ab}^{\parallel}(\tilde{g}) + D_{(a}a_{b)} + a_{a}a_{b}. \tag{B.18}$$

We have that $a = a_r dr$ with

$$a_r = \tilde{n}^t \tilde{D}_t \tilde{n}_r = \frac{1}{2f} \partial_r f \tag{B.19}$$

and

$$D_r a_r = -\frac{1}{4f^2} (\partial_r f)^2 + \frac{1}{2f} \partial_r \partial_r f, \qquad D_A a_B = \frac{\partial_r f}{2} r \mathring{h}_{AB}.$$
 (B.20)

The only non-vanishing components of the Ricci tensor of the metric induced on the submanifolds of constant t are

$$R_{rr} = -\frac{n}{f} + \frac{1}{2f} \partial_r \partial_r f, \qquad R_{AB} = \left(-nr^2 + \frac{\partial_r f}{2}r\right) \mathring{h}_{AB}$$
 (B.21)

and thus

$$R = R_{rr}g^{rr} + R_{AB}g^{AB} = -n(n-1), (B.22)$$

where we have used the explicit form of f(r). Now we consider the KID equation

$$A_{ij} := D_i D_j V - V \left(R_{ij} - \frac{R}{n-1} g_{ij} \right) = D_i D_j V - V (R_{ij} + n g_{ij}) = 0$$
 (B.23)

where $i, j \in 1, ...n - 1$. We have that

$$A_{rr} = \partial_r \partial_r V + \frac{1}{2f} \partial_r f \partial_r V - \frac{V}{2f} \partial_r \partial_r f = 0,$$
(B.24)

$$A_{rA} = \partial_r \partial_A V - \frac{\delta_A{}^C}{r} \partial_C V = \partial_r \partial_A V - \frac{1}{r} \partial_A V = 0, \qquad (B.25)$$

$$A_{AB} = \partial_A \partial_B V - \Gamma^C{}_{AB} \partial_C V + r \mathring{h}_{AB} f \partial_r V - \frac{V \partial_r f}{2} r \mathring{h}_{AB} = 0.$$
 (B.26)

Integrating the second equation in r we have that

$$V(r, x^{1}, ..., x^{n-1}) = \beta(r) + r\Omega(x^{1}, ... x^{n-1}),$$
(B.27)

for some functions β and Ω . Plugging (B.27) into A_{rr} we find

$$\Omega(x^1, \dots x^{n-1}) \frac{(\partial_r f - r \partial_r \partial_r f)}{f} + \frac{\partial_r f \partial_r \beta - \beta \partial_r \partial_r f + 2f \partial_r \partial_r \beta}{f} = 0.$$
 (B.28)

Thus, the differential equation can only be fulfilled if either $\Omega(x^1,...x^{n-1})$ is constant, which can then without loss of generality be taken to be zero as any constant can be reabsorbed into a redefinition of $\beta(r)$, or if

$$\partial_r f - r \partial_r \partial_r f \equiv 2m_c (n-2) n r^{1-n} = 0$$
(B.29)

which can only be fulfilled if $m_c = 0$ or n = 2.

The remaining part of (B.28) reads

$$\partial_r f \partial_r \beta - \beta \partial_r \partial_r f + 2f \partial_r \partial_r \beta = 0.$$
 (B.30)

Changing the dependent variable β to

$$\beta(r) = \sqrt{f(r)}\beta_2(r) \tag{B.31}$$

we obtain the differential equation

$$3\partial_r f \partial_r \beta_2 + 2f \partial_r \partial_r \beta_2 = 0. (B.32)$$

From this we obtain that

$$\beta_2(r) = \tilde{c} - \hat{c} \int \frac{dr}{f(r)^{\frac{3}{2}}},$$
(B.33)

which yields that

$$\beta(r) = \tilde{c}\sqrt{f(r)} - \hat{c}\sqrt{f(r)} \int \frac{dr}{f(r)^{\frac{3}{2}}}.$$
 (B.34)

B.1.1 $m_c = 0, n \neq 2$

In the case $m_{\rm c} = 0$ the function f reduces to

$$f(r) = r^2 + k$$
, (B.35)

and (B.34) readily integrates to

$$\beta(r) = -\frac{\hat{c}r}{k} + \tilde{c}\sqrt{r^2 + k}$$
(B.36)

if $k \neq 0$, and to

$$\beta(r) = \frac{\hat{c}}{2r} + \tilde{c}r \tag{B.37}$$

if k = 0. The differential equations (B.26) reduce to

$$D_A D_B \Omega \equiv \partial_A \partial_B \Omega - \Gamma_{AB}^C \partial_C \Omega = \mathring{h}_{AB} \left(\hat{c} - k \Omega \right) .$$
(B.38)

This is an overdetermined system of equations, and all triples $(M, \mathring{h}, \Omega)$ satisfying (B.38), where $\Omega \not\equiv 0$ and (M, \mathring{h}) a complete Riemannian manifold, were classified in [40].

In the special case

$$\mathring{h}_{AB}dx^{A}dx^{B} = d\psi^{2} + d\theta^{1}^{2} + \dots + d\theta^{n-2}^{2}$$
(B.39)

the constant k is zero and all Christoffel symbols of the boundary metric \mathring{h} vanish, so that (B.38) becomes

$$\partial_A \partial_B \Omega = \hat{c} \, \delta_{AB} \,. \tag{B.40}$$

We find

$$V(r) = \tilde{c}r + \frac{\hat{c}}{2r} + r\left(c\psi + \hat{c}\frac{\psi^2}{2}\right) + r\sum_{I=1}^{n-2} c_I \theta^I + \hat{c}\frac{|\theta|^2}{2},$$
(B.41)

where c and c_I are constants.

B.1.2 $m_c \neq 0, n \neq 2$

In the generic case $m_c \neq 0$ the function $\Omega(x^1, ... x^{n-1}) = 0$ and the function V reduces to

$$V(r, x^{1}, ...x^{n-1}) = \beta(r) = \tilde{c}\sqrt{f(r)} - \hat{c}\sqrt{f(r)} \int \frac{dr}{f(r)^{\frac{3}{2}}}.$$
 (B.42)

The final set of equations that needs to be solved is (B.26), which in this case reduces to

$$A_{AB} = r\mathring{h}_{AB}f\partial_r V - \frac{V\partial_r f}{2}r\mathring{h}_{AB} = 0.$$
 (B.43)

Contracting with \mathring{h}^{AB} yields

$$rf\partial_r V - \frac{V\partial_r f}{2}r = 0 (B.44)$$

which implies that $\hat{c} = 0$ and thus

$$V(r, x^1, ..., x^{n-1}) = \tilde{c}\sqrt{f(r)}$$
 (B.45)

B.1.3 n = 2

As already discussed, in the case n=2 the function f(r) reduces to

$$f(r) = r^2 - 2m_c \,, \tag{B.46}$$

which has the same functional form as (B.35) but with k in (B.35) now being replaced by $-2m_c$. Again, (B.34) thus readily integrates to

$$\beta(r) = \frac{\hat{c}r}{2m_o} + \tilde{c}\sqrt{r^2 - 2m_c} \tag{B.47}$$

if $m_c \neq 0$, and to

$$\beta(r) = \frac{\hat{c}}{2r} + \tilde{c}r \tag{B.48}$$

if $m_c = 0$. The differential equations (B.26) reduce to just one differential equation as the boundary metric is one-dimensional

$$D_A D_A \Omega = \mathring{h}_{AA} \left(\hat{c} + 2m_c \Omega \right) \tag{B.49}$$

with A=1. In a coordinate system $x^1=\psi$ in which \mathring{h}_{11} equals 1 the solutions are

$$\Omega = \begin{cases}
\frac{\hat{c}}{2}\psi^2 + a\psi + b, & m_c = 0; \\
ae^{\sqrt{2m_c}\psi} + be^{-\sqrt{2m_c}\psi} - \frac{\hat{c}}{2m_c}, & m_c > 0; \\
a\sin(\sqrt{-2m_c}\psi) + b\cos(\sqrt{-2m_c}\psi) - \frac{\hat{c}}{2m_c}, & m_c < 0,
\end{cases}$$
(B.50)

with constants $a, b \in \mathbb{R}$.

B.2 Horowitz-Myers

We consider the Horowitz-Myers metrics, which take the form

$$ds^{2} = \frac{1}{f(r)}dr^{2} + f(r)d\psi^{2} + r^{2}\left(-dt^{2} + (d\theta^{1})^{2} + (d\theta^{2})^{2} + \dots + (d\theta^{n-2})^{2}\right)$$
(B.51)

with

$$f(r) = r^2 - \frac{2m_c}{r^{n-2}},$$
 (B.52)

with $n \geq 2$. The spatial part of the metric reads

$$g_{ij}dx^{i}dx^{j} = \frac{1}{f(r)}dr^{2} + f(r)d\psi^{2} + r^{2}\left((d\theta^{1})^{2} + (d\theta^{2})^{2} + \dots + (d\theta^{n-2})^{2}\right).$$
 (B.53)

The indices i, j run from 1 to n.

The only non-vanishing Christoffel symbols (up to symmetries) for (B.53) are

$$\Gamma^{r}{}_{rr} = -\frac{\partial_{r}f}{2f}, \quad \Gamma^{r}{}_{\psi\psi} = -\frac{f\partial_{r}f}{2}, \quad \Gamma^{r}{}_{\theta^{I}\theta^{I}} = -rf, \quad \Gamma^{\psi}{}_{\psi r} = \frac{\partial_{r}f}{2f}, \quad \Gamma^{\theta^{I}}{}_{\theta^{I}r} = \frac{1}{r}.$$
 (B.54a)

Next we compute

$$D_i D_j V(r, \psi, \theta^1, \dots \theta^{n-2}) = \partial_i \partial_j V(r, \psi, \theta^1, \dots \theta^{n-2}) - \Gamma^k_{ij} \partial_k V(r, \psi, \theta^1, \dots \theta^{n-2}). \tag{B.55}$$

We have that

$$D_r D_r V = \partial_r \partial_r V + \frac{1}{2f} \partial_r f \partial_r V, \qquad D_r D_\psi V = \partial_r \partial_\psi V - \frac{1}{2} \frac{\partial_r f}{f} \partial_\psi V, \qquad (B.56a)$$

$$D_r D_{\theta^I} V = \partial_r \partial_{\theta^I} V - \frac{1}{r} \partial_{\theta^I} V , \qquad D_{\psi} D_{\psi} V = \partial_{\psi} \partial_{\psi} V + \frac{1}{2} f \partial_r f \partial_r V , \qquad (B.56b)$$

$$D_{\theta}D_{\theta}V = \partial_{\theta}\partial_{\theta}V, \qquad D_{\theta}D_{\theta}V = \partial_{\theta}\partial_{\theta}V + rf\partial_{r}V\delta_{IJ}, \qquad (B.56c)$$

where I, J run from from 1 to n-2. The only non-vanishing components of the Riemann tensor

$$R_{ijk}^{\ l} = \partial_j \Gamma^l_{ik} - \partial_i \Gamma^l_{jk} + \left(\Gamma^m_{ik} \Gamma^l_{mj} - \Gamma^m_{jk} \Gamma^l_{mi}\right) \tag{B.57}$$

read (up to the ones obtained from antisymmetry in the first index pair)

$$R_{r\psi r}{}^{\psi} = -\frac{\partial_r \partial_r f}{2f}, \qquad R_{r\psi \psi}{}^r = -\frac{f \partial_r \partial_r f}{2}, \qquad R_{r\theta^I r}{}^{\theta^I} = -\frac{\partial_r f}{2fr}, \quad R_{r\theta^I \theta^I}{}^r = \frac{r \partial_r f}{2}, \quad (B.58a)$$

$$R_{\psi\theta^I\psi}^{\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ } = -\frac{f\partial_r f}{2r} \,, \qquad R_{\psi\theta^I\theta^I}^{\ \ \ \ \ \ \ \ } = \frac{r\partial_r f}{2} \,, \qquad R_{\theta^I\theta^J\theta^I}^{\ \ \ \ \ \ \ \ } = -f \qquad \text{with } I \neq J \,.$$
 (B.58b)

From this we obtain that the only non-vanishing components of the Ricci-tensor

$$R_{ij} = R_{ikj}^{\ k} \tag{B.59}$$

read

$$R_{rr} = -\frac{\partial_r \partial_r f}{2rf} - \frac{(n-2)}{2} \frac{\partial_r f}{rf}, \qquad R_{\psi\psi} = -\frac{1}{2} f \partial_r \partial_r f - \frac{(n-2)}{2} \frac{f \partial_r f}{r}, \qquad (B.60a)$$

$$R_{\theta^I \theta^I} = -(n-3)f - r\partial_r f. \tag{B.60b}$$

This leads to the following formula for the Ricci scalar

$$R = -n(n-1). (B.60c)$$

The KID equations,

$$A_{ij} := D_i D_j V - V \left(R_{ij} - \frac{R}{n-1} g_{ij} \right) = D_i D_j V - V (R_{ij} + n g_{ij}) = 0,$$
 (B.60d)

thus become

$$A_{rr} = \partial_r \partial_r V + \frac{1}{2f} \partial_r f \partial_r V - V \left(-\frac{\partial_r \partial_r f}{2rf} - \frac{(n-2)}{2} \frac{\partial_r f}{rf} + \frac{n}{f} \right) = 0,$$
 (B.61a)

$$A_{r\psi} = A_{\psi r} = \partial_r \partial_{\psi} V - \frac{1}{2} \frac{\partial_r f}{f} \partial_{\psi} V = 0,$$
(B.61b)

$$A_{r\theta^I} = A_{\theta^I r} = \partial_r \partial_{\theta^I} V - \frac{1}{r} \partial_{\theta^I} V = 0,$$
(B.61c)

$$A_{\psi\psi} = \partial_{\psi}\partial_{\psi}V + \frac{1}{2}f\partial_{r}f\partial_{r}V - V\left(-\frac{1}{2}f\partial_{r}\partial_{r}f - \frac{(n-2)}{2}\frac{f\partial_{r}f}{r} + nf\right) = 0,$$
 (B.61d)

$$A_{\psi\theta^I} = \partial_{\psi}\partial_{\theta^I}V = 0, \tag{B.61e}$$

$$A_{\theta^I \theta^J} = \partial_{\theta^I} \partial_{\theta^J} V + \delta_{IJ} \left(r f \partial_r V - V \left(-(n-3)f - r \partial_r f + n r^2 \right) \right) = 0.$$
 (B.61f)

Solving (B.61e) and (B.61f) with $I \neq J$ we obtain that

$$V(r, \psi, \theta^1, \dots \theta^{n-2}) = \alpha(r, \psi) + \alpha_1(r, \theta^1) + \alpha_2(r, \theta^2) + \dots + \alpha_{n-2}(r, \theta^{n-2}).$$
 (B.62)

Plugging this into the equation (B.61c) we get that

$$\alpha_I(r, \theta^I) = \beta_I(r) + r\hat{\gamma}_I(\theta^I). \tag{B.63}$$

Solving (B.61b) yields that

$$\alpha(r,\psi) = \tilde{\beta}(r) + \sqrt{f(r)}\hat{\gamma}(\psi). \tag{B.64}$$

Setting $\beta(r) = \tilde{\beta}(r) + \sum_{I=1}^{n-2} \beta_I(r)$ we have

$$V(r, \psi, \theta^{1}, ...\theta^{n-2}) = \beta(r) + \sqrt{f(r)}\hat{\gamma}(\psi) + r \sum_{I=1}^{n-2} \hat{\gamma}_{I}(\theta^{I}),$$
 (B.65)

with the last term omitted when n=2. From (B.61f) we obtain (by pairwise substraction)

$$\partial_{\theta^I}\partial_{\theta^I}V = \partial_{\theta^J}\partial_{\theta^J}V \tag{B.66}$$

for any I, J which is equivalent to

$$\partial_I \partial_I \hat{\gamma}_I(\theta^I) = \partial_J \partial_J \gamma_J(\theta^J). \tag{B.67}$$

From this we obtain that (the θ^I -independent integration constant can be absorbed into $\beta(r)$)

$$V(r, \psi, \theta^{1}, ... \theta^{n-2}) = \beta(r) + \sqrt{f(r)} \hat{\gamma}(\psi) + rc_{I} \theta^{I} + \frac{\hat{c}}{2} |\theta|^{2}.$$
 (B.68)

We turn now our attention to the equation

$$A_{\psi\psi} - f(r)^{2} A_{rr} = 0 = \partial_{\psi} \partial_{\psi} V - f^{2} \partial_{r} \partial_{r} V$$

$$= \frac{1}{4} \sqrt{f(r)} \left(\hat{\gamma}(\psi) \left((\partial_{r} f)^{2} - 2f \partial_{r} \partial_{r} f \right) - 4f^{\frac{3}{2}} \partial_{r} \partial_{r} \beta + 4 \partial_{\psi} \partial_{\psi} \hat{\gamma} \right) , \qquad (B.69)$$

from which we find

$$\hat{\gamma}(\psi)\left((\partial_r f)^2 - 2f\partial_r \partial_r f\right) + 4\partial_\psi \partial_\psi \hat{\gamma} = 4f^{\frac{3}{2}} \partial_r \partial_r \beta. \tag{B.70}$$

For the function $f(r) = r^2 - \frac{2m_c}{r^{n-2}}$ we have that

$$((\partial_r f)^2 - 2f\partial_r \partial_r f) = 4nm_c r^{2-2n} \left((-1+n)r^n - (-2+n)m_c \right). \tag{B.71}$$

If $m_c = 0$ the spatial parts of the TK and the HM metrics coincide, and this case has already been solved in Section B.1.2.

In the case $m_c \neq 0$ we see from (B.71) that $\hat{\gamma}(\psi)$ has to be a constant. This constant is irrelevant as it can be absorbed by a redefinition of $\beta(r)$, which is why we set it to zero in the following. Equation (B.70) then reduces to

$$\partial_r^2 \beta = 0. (B.72)$$

Hence, we have that $\beta(r) = r\tilde{c}_1 + \tilde{c}_2$ which yields

$$V(r, \psi, \theta^{1}, \dots, \theta^{n-2}) = r\tilde{c}_{1} + \tilde{c}_{2} + rc_{I}\theta^{I} + \frac{\hat{c}}{2}|\theta|^{2}.$$
(B.73)

Plugging this into the equation (B.61a), $A_{rr} = 0$, together with the explicit form of f(r) we obtain

$$A_{rr} = -\frac{(r^n + (-2+n)m_c)}{r^2(r^n - 2m_c)}\tilde{c}_2 = 0,$$
(B.74)

from which follows that $\tilde{c}_2 = 0$. The differential equation (B.61d) $A_{\psi\psi} = 0$ is then automatically fulfilled. Plugging all of this into (B.61f) $A_{\theta^I\theta^I} = 0$ we obtain

$$A_{\theta^I\theta^I} = r\hat{c} = 0, \tag{B.75}$$

from which follows that $\hat{c} = 0$. Hence, we have that

$$V(r, \psi, \theta^1, ... \theta^{n-2}) = r\tilde{c}_1 + r \sum_{I=1}^{n-2} (c_I \theta^I) , \qquad (B.76)$$

with the last term omitted when n=2.

C Transformation behaviour of the mass aspect tensor under conformal transformations

Consider a metric of the following Fefferman-Graham form

$$g = x^{-2}(dx^2 + h_{AB}dx^A dx^B),$$
 (C.1)

where the coordinate functions h_{AB} depend upon both x and the local coordinates x^A on the boundary $\{x=0\}$. Here, the index A runs from 1, ... n-1. In the following we will consider $n \geq 3$. Let us further write a Taylor expansion

$$h_{AB} = (1 - kx^2/4)^2 \mathring{h}_{AB}(x^C) + x^n \mu_{AB}(x^C) + o(x^n), \qquad (C.2)$$

where k is a constant. Here, the scalar curvature of \mathring{h}_{AB} is given by

$$R(\mathring{h}) = k(n-1)(n-2)$$
. (C.3)

Let $\phi = \phi(x^C)$ be a function on the boundary, set

$$\bar{h}_{AB}|_{x=0} = \phi^2 h_{AB} \,,$$
 (C.4)

thus (C.1) takes the form

$$g = x^{-2} (dx^2 + (\phi^{-2}\bar{h}_{AB} + O(x))dx^A dx^B).$$
 (C.5)

We wish to rewrite this as

$$g = y^{-2} (dy^2 + \bar{h}_{AB} d\bar{x}^A d\bar{x}^B).$$
 (C.6)

with

$$\bar{h}_{AB} = (1 - \bar{k}y^2/4)^2 \phi^2(\bar{x}^C) \mathring{h}_{AB}(\bar{x}^C) + y^n \bar{\mu}_{AB}(\bar{x}^C) + o(y^n), \qquad (C.7)$$

where $\bar{k} \in \{0, \pm 1\}$.

$$\bar{R}(\phi^2 \mathring{h}) = \bar{k}(n-1)(n-2)$$
. (C.8)

By matching powers in Taylor expansions, this is equivalent to finding a function y with a Taylor expansion

$$y = \phi x \left(1 + \phi_1 x + \phi_2 x^2 + \phi_3 x^3 + \dots \right)$$
 (C.9a)

and new boundary coordinates \bar{x}^A with Taylor expansions

$$\bar{x}^A = x^A + \phi_1^A x + \phi_2^A x^2 + \phi_3^A x^3 + \phi_4^A x^4 \dots$$
 (C.9b)

In order to determine y as a function of the original coordinates, we note that

$$g^{yy} \equiv g(dy, dy) = y^2 \qquad \Longleftrightarrow \qquad |d(\log y)|_g^2 = 1. \tag{C.10}$$

This equation says that the integral curves of $d(\log y)$ are affinely parameterised geodesics. The solutions can be found by shooting geodesics orthogonally, in the metric x^2g , to the conformal boundary, with suitable boundary conditions determined by the function ϕ . Hence smooth solutions always exist, which justifies the existence of the expansion (C.9a). Further, we see that the equation for y can be solved independently of the equation for the \bar{x}^A 's. Next, the equation

$$0 = g(dy, d\bar{x}^A) \tag{C.11}$$

says that the coordinates \bar{x}^A are constant along the integral curves of dy. So, when a smooth function y is known, one obtains smooth functions \bar{x}^A by solving (C.11) along the integral curves of dy, which again justifies the expansion (C.9b), and provides a prescription how to find the expansion functions ψ_n^A .

We find that

$$\phi_1 = 0, \qquad \phi_2 = -\frac{\mathcal{D}_A \phi \mathcal{D}^A \phi}{4\phi^2}, \qquad \phi_3 = 0,$$
 (C.12a)

and

$$\phi_1^A = 0, \qquad \phi_2^A = -\frac{1}{2} \frac{\mathscr{D}^A \phi}{\phi}, \qquad \phi_3^A = 0,$$
 (C.12b)

where \mathscr{D} is the covariant derivative of the metric \mathring{h} and the indices A are raised with \mathring{h}^{AB} . With this it holds that

$$g = x^{-2} (dx^2 + (h_{AB} + O(x^2)) dx^A dx^B) = y^{-2} (dy^2 + (\bar{h}_{AB} + O(y)^2) d\bar{x}^A d\bar{x}^B)$$
(C.13)

However, for general \mathring{h}_{AB} it is not possible to bring the $y^2 d\bar{x}^A d\bar{x}^B$ term into the form (C.7) by changing coordinates as (C.9) after the conformal transformation. Indeed, one finds that the terms of order y^0 in the coordinate-transformed metric read

$$-k\frac{\mathring{h}_{AB}}{2} + \mathscr{D}_{A}\mathscr{D}_{B}\log\phi - (\mathscr{D}_{A}\log\phi)(\mathscr{D}_{B}\log\phi) + \frac{1}{2}(\mathscr{D}^{C}\log\phi)(\mathscr{D}_{C}\log\phi)\mathring{h}_{AB}. \tag{C.14}$$

It follows that for (C.7) to hold true the following expression must vanish

$$\frac{\phi^2}{2}(-\bar{k}\phi^2 + k)\mathring{h}_{AB} - \phi \mathcal{D}_A \mathcal{D}_B \phi + 2(\mathcal{D}_A \phi)(\mathcal{D}_B \phi) - \frac{1}{2}(\mathcal{D}^C \phi)(\mathcal{D}_C \phi)\mathring{h}_{AB} = 0.$$
 (C.15)

The trace of (C.15) is equivalent to the transformation of the Ricci scalar under conformal transformations (compare e.g. [41, Appendix D])

$$\bar{R} = \phi^{-2} \left(R - 2(n-2) \mathcal{D}^A \mathcal{D}_A \log \phi - (n-3)(n-2) (\mathcal{D}^A \log \phi) (\mathcal{D}_A \log \phi) \right). \tag{C.16}$$

where we have used the explicit expressions for the Ricci scalars (C.3) and (C.8).

C.1 Relation to coordinates in the main text

The coordinates x, x^A in (C.1) are related to the coordinates in the main text r, x^A , Section 4

$$r = \frac{1}{x} - \frac{kx}{4} \,, \tag{C.17a}$$

$$x = \frac{2}{r + \sqrt{r^2 + k}} = \frac{1}{r} - \frac{k}{4r^3} + \frac{k^2}{8r^5} + O(r^{-7})$$
 (C.17b)

Under this change of coordinates the line element

$$g = x^{-2}(dx^2 + \left((1 - kx^2/4)^2\mathring{h}_{AB}(x^C) + x^n\mu_{AB}(x^C) + o(x^n)\right)dx^Adx^B)$$
 (C.18)

changes as

$$g = \frac{dr^2}{r^2 + k} + r^2 \left(\mathring{h}_{AB}(x^c) + r^{-n}\mu_{AB}(x^C) + o(r^{-n})\right) dx^A dx^B$$
 (C.19a)
(C.19b)

Using the relations (C.17) we have that

$$\bar{x}^A = x^A - \frac{1}{2} \frac{\mathscr{D}^A \phi}{\phi} x^2 + O(x^4)$$
 (C.20)

which, using

$$\psi := \phi^{-1} \,,$$

translates to

$$\bar{x}^A = x^A + \frac{\mathcal{D}^A \psi}{2\psi r^2} + O(r^{-4}),$$
 (C.21)

while the expansion in y

$$y = \phi x \left(1 - \frac{\mathcal{D}^A \phi \mathcal{D}_A \phi}{4\phi^2} x^2 + \mathcal{O}(x^4) \right)$$
 (C.22)

leads to an expansion in \bar{r}

$$\bar{r} = \psi r \left(1 + \frac{\left(\mathcal{D}^A \psi \mathcal{D}_A \psi + k \psi^2 - \bar{k} \right)}{4 \psi^2 r^2} + O(r^{-4}) \right). \tag{C.23}$$

Using (C.16) we can also write

$$\bar{r} = \psi r \left(1 + \frac{\left(-\mathcal{D}^A \mathcal{D}_A \log \psi + (n-2) \mathcal{D}^A \log \psi \mathcal{D}_A \log \psi \right)}{2(n-1)r^2} + O(r^{-4}) \right). \tag{C.24}$$

C.2 n = 3

For n=3, the boundary metric is two-dimensional. Thus we have the relation that

$$R_{AB} = \frac{R}{2}\mathring{h}_{AB} = k\mathring{h}_{AB}, \qquad \bar{R}_{AB} = \frac{\bar{R}}{2}\phi^2\mathring{h}_{AB} = \bar{k}\phi^2\mathring{h}_{AB}.$$
 (C.25)

Reconsidering (C.15) and using the expression for the Ricci tensor (C.25) we get

$$\bar{R}_{AB} = R_{AB} - 2\mathcal{D}_A \mathcal{D}_B \log \phi + 2\mathcal{D}_A \log \phi \mathcal{D}_B \log \phi - \mathring{h}_{AB} \mathcal{D}^C \log \phi \mathcal{D}_C \log \phi. \tag{C.26}$$

This coincides with the transformation behaviour of the Ricci tensor in two dimensions if and only if

$$\mathring{h}_{AB} \left(\mathscr{D}^C \mathscr{D}_C \log \phi - \mathscr{D}^C \log \phi \mathscr{D}_C \log \phi \right) = 2 \left(\mathscr{D}_A \mathscr{D}_B \log \phi - \mathscr{D}_A \log \phi \mathscr{D}_B \log \phi \right). \tag{C.27}$$

Using (C.15) the $y^2 d\bar{x}^A d\bar{x}^B$ term can be brought into the correct form. For n=3 we find that the new mass aspect tensor $\bar{\mu}_{AB}$ takes the form

$$\bar{\mu}_{AB} = \frac{\mu_{AB}}{\phi} = \psi \mu_{AB} \,, \tag{C.28}$$

where $\psi = \phi^{-1}$, which is the variable sometimes used in the main text. From

$$\dot{h}^{AB}\mu_{AB} = \phi^3 \bar{h}_{AB}|_{x=0} \bar{\mu}_{AB}, \qquad \sqrt{\det \dot{h}} = \phi^{-2} \sqrt{\det \bar{h}}|_{x=0}, \qquad \bar{x}^A = x^A|_{x=0}$$
 (C.29)

we conclude that

$$\int \mathring{h}^{AB} \mu_{AB} \sqrt{\det \mathring{h}} \, dx^1 dx^2 = \int \left(\phi \bar{h}^{AB} \bar{\mu}_{AB} \sqrt{\det \bar{h}} \right) \big|_{\bar{x}=0} \, d\bar{x}^1 d\bar{x}^2 \,, \tag{C.30}$$

as well as

$$\int \left(\bar{h}^{AB} \bar{\mu}_{AB} \sqrt{\det \bar{h}} \right) \big|_{\bar{x}=0} d\bar{x}^1 d\bar{x}^2 = \int \phi^{-1} \mathring{h}^{AB} \mu_{AB} \sqrt{\det \mathring{h}} dx^1 dx^2 . \tag{C.31}$$

C.3 n = 4

We consider now the transformation formulae for the higher-order terms in the expansion of the metric. These are irrelevant for the mass aspect tensor when n=3 but become relevant in higher dimensions. Because the powers of x in the expansion of y and \bar{x}^A jump by two, and so do the powers in the expansion of the metric up to the mass-aspect level x^{n-2} , when $n \geq 4$ the terms of order x in the physical metric remain zero after the change of coordinates. The terms of order x^2 change in a non-trivial way, which is relevant for the mass aspect function in space-dimension n=4, and which we determine now.

The explicit form of the coefficients ϕ_1, ϕ_2, ϕ_3 and $\phi_1^A, \phi_2^A, \phi_3^A$ in (C.12) holds true for arbitrary dimensions $n \geq 3$. However, to determine the transformation behaviour of the mass aspect in n = 4 we need to determine further coefficients. We have that

$$\phi_4^A = -\frac{\bar{k}}{8}\phi \mathcal{D}^A \phi + \frac{1}{8\phi^2} \mathcal{D}^B \phi \mathcal{D}^C \phi \mathcal{D}_C \mathcal{D}_B x^A, \quad \phi_4 = -\frac{\bar{k}}{16} \mathcal{D}_C \phi \mathcal{D}^C \phi + \frac{1}{16\phi^4} (\mathcal{D}_C \phi \mathcal{D}^C \phi)^2, \quad (C.32)$$

where x^A in the first equation (C.32) is treated as a scalar for every A. In dimension n=4 the coefficients ϕ_5^A vanish

$$\phi_5^A = 0, \qquad \phi_5 = 0.$$
 (C.33)

This leads to

$$\phi^{2}\bar{\mu}_{AB} = \mu_{AB} + \frac{1}{16}\mathring{h}_{AB}\left(k^{2} - \bar{k}^{2}\phi^{4}\right) - \frac{\bar{k}}{4}\phi\mathcal{D}_{A}\mathring{D}_{B}\phi - \frac{3}{8}\bar{k}\mathring{h}_{AB}\mathcal{D}^{C}\phi\mathcal{D}_{C}\phi + \frac{3}{4}\bar{k}\mathcal{D}_{A}\phi\mathcal{D}_{B}\phi$$

$$- \frac{1}{4\phi^{2}}\mathcal{D}^{C}\mathcal{D}_{A}\phi\mathcal{D}_{C}\mathcal{D}_{B}\phi + \frac{3}{2\phi^{3}}\mathcal{D}^{C}\mathcal{D}_{(A}\phi\mathcal{D}_{B)}\phi\mathcal{D}_{C}\phi - \frac{1}{4\phi^{3}}\mathring{h}_{AB}\mathcal{D}^{C}\phi\mathcal{D}_{D}\mathcal{D}_{C}\phi\mathcal{D}^{D}\phi$$

$$- \frac{1}{2\phi^{3}}\mathcal{D}_{A}\mathcal{D}_{B}\phi\mathcal{D}^{C}\phi\mathcal{D}_{C}\phi - \frac{3}{4\phi^{4}}\mathcal{D}_{A}\phi\mathcal{D}_{B}\phi\mathcal{D}^{C}\phi\mathcal{D}_{C}\phi + \frac{3}{16\phi^{4}}(\mathcal{D}^{C}\phi\mathcal{D}_{C}\phi)^{2}\mathring{h}_{AB}$$

$$- \frac{1}{4\phi^{2}}R_{ACDB}\mathcal{D}^{C}\phi\mathcal{D}^{D}\phi. \tag{C.34}$$

This is best computed using coordinates in which the metric is diagonal, which can always be done locally in dimension three [23]. Here and elsewhere, the symmetrization is defined as $M_{(AB)} = \frac{1}{2} (M_{AB} + M_{BA})$. Contracting (C.34) with \mathring{h}^{AB} we obtain

$$\phi^{2}\bar{\mu}_{AB}\mathring{h}^{AB} - \mu_{A}^{A} = \frac{3}{16}(k^{2} - \bar{k}^{2}\phi^{4}) - \frac{\bar{k}}{4}\phi\mathscr{D}^{A}\mathscr{D}_{A}\phi - \frac{3}{8}\bar{k}(\mathscr{D}^{A}\phi\mathscr{D}_{A}\phi) - \frac{1}{4\phi^{2}}(\mathscr{D}^{A}\mathscr{D}^{B}\phi)(\mathscr{D}_{A}\mathscr{D}_{B}\phi) + \frac{3}{4\phi^{3}}\mathscr{D}^{A}\phi\mathscr{D}^{B}\phi\mathscr{D}_{A}\mathscr{D}_{B}\phi - \frac{1}{2\phi^{3}}(\mathscr{D}^{A}\mathscr{D}_{A}\phi)(\mathscr{D}^{B}\phi\mathscr{D}_{B}\phi) - \frac{3}{16\phi^{4}}(\mathscr{D}^{A}\phi\mathscr{D}_{A}\phi)^{2} + \frac{1}{4\phi^{2}}R_{AB}\mathscr{D}^{A}\phi\mathscr{D}^{B}\phi.$$
(C.35)

Using the transformation behaviour of the Ricci scalar (C.16),

$$\bar{R} = \phi^{-2} \left(R - \frac{4}{\phi} \mathcal{D}^A \mathcal{D}_A \phi + \frac{2}{\phi^2} \mathcal{D}^A \phi \mathcal{D}_A \phi \right), \tag{C.36}$$

and the explicit expression of the Ricci scalars (C.3) and (C.8),

$$R = 6k, \qquad \bar{R} = 6\bar{k}, \tag{C.37}$$

for n = 4 we find that

$$\bar{k} = \frac{3k\phi^2 + (\mathcal{D}^A\phi\mathcal{D}_A\phi) - 2\phi\mathcal{D}^A\mathcal{D}_A\phi}{3\phi^4},$$
(C.38)

which yields

$$\phi^{2}\bar{\mu}_{AB}\mathring{h}^{AB} - \mu_{A}^{A} = -\frac{1}{3\phi^{4}}(\mathscr{D}^{A}\phi\mathscr{D}_{A}\phi)^{2} - \frac{1}{4\phi^{3}}(\mathscr{D}^{A}\phi\mathscr{D}_{A}\phi)(\mathscr{D}^{B}\mathscr{D}_{B}\phi) + \frac{(\mathscr{D}^{A}\mathscr{D}_{A}\phi)^{2}}{12\phi^{2}}$$
$$-\frac{1}{4\phi^{2}}(\mathscr{D}^{A}\mathscr{D}^{B}\phi)(\mathscr{D}_{A}\mathscr{D}_{B}\phi) + \frac{3}{4\phi^{3}}\mathscr{D}^{A}\phi\mathscr{D}^{B}\phi\mathscr{D}_{A}\mathscr{D}_{B}\phi$$
$$+\frac{1}{4\phi^{2}}(R_{AB} - 2k\mathring{h}_{AB})\mathscr{D}^{A}\phi\mathscr{D}^{B}\phi. \tag{C.39}$$

Making the change of variable

$$\phi = \exp(u) \tag{C.40}$$

(so that u here corresponds to $+\omega/2$ in (4.26)), Equation (C.39) becomes

$$\exp(2u)\bar{\mu}_{AB}\mathring{h}^{AB} - \mu_{A}^{A} = -\frac{1}{12}(\mathscr{D}^{A}u\mathscr{D}_{A}u)(\mathscr{D}^{B}\mathscr{D}_{B}u) + \frac{1}{12}(\mathscr{D}^{A}\mathscr{D}_{A}u)(\mathscr{D}^{B}\mathscr{D}_{B}u) -\frac{1}{4}(\mathscr{D}^{A}\mathscr{D}^{B}u)(\mathscr{D}_{A}\mathscr{D}_{B}u) + \frac{1}{4}(\mathscr{D}^{A}\mathscr{D}^{B}u)(\mathscr{D}_{A}u\mathscr{D}_{B}u) +\frac{1}{4}(R_{AB} - 2k\mathring{h}_{AB})\mathscr{D}^{A}u\mathscr{D}^{B}u.$$
 (C.41)

Using

$$\sqrt{\det \mathring{h}} = \phi^{-3} \sqrt{\det \overline{h}}|_{x=0}, \quad \overline{x}^A = x^A|_{x=0},$$
 (C.42)

and (C.41) we find that

$$\int \left(\bar{\mu}_{AB}\bar{h}^{AB}\sqrt{\det\bar{h}}\right)|_{x=0} d\bar{x}^1 d\bar{x}^2 d\bar{x}^3 = \int e^{-u} \left(\mu^A_A + C(x^A)\right) \sqrt{\det\mathring{h}} dx^1 dx^2 dx^3 \qquad (C.43)$$

with

$$C(x^{A}) = \frac{1}{6} (\mathscr{D}_{A} \mathscr{D}_{B} \mathscr{D}^{A} u) (\mathscr{D}^{B} u) + \frac{1}{12} \left(R_{AB} - \frac{R}{2} \mathring{h}_{AB} \right) \mathscr{D}^{A} u \mathscr{D}^{B} u. \tag{C.44}$$

ACKNOWLEDGEMENTS: We are grateful to Michael Anderson, Greg Galloway, Colin Guillarmou, Jack Lee, Yanyan Li, Rafe Mazzeo, Bruno de Oliveira and Laurent Véron for useful discussions. PTC was supported in part by the Austrian Science Fund (FWF) under project P29517-N27 and by the Polish National Center of Science (NCN) 2016/21/B/ST1/00940. Part of this research was performed while PTC was visiting the Institute for Pure and Applied Mathematics (IPAM) at UCLA, which is supported by the National Science Foundation (Grant No. DMS-DMS-1925919). ED was supported by the grant ANR-17-CE40-0034 of the French National Research Agency ANR (project CCEM). RW was supported by the Austrian Science Fund FWF under the Doctoral Program W1252-N27 Particles and Interactions. RW thanks the University of Amsterdam for hospitality during the completion of this project.

References

- S. Åminneborg, I. Bengtsson, S. Holst, and P. Peldán, Making anti-de Sitter black holes, Class. Quantum Grav. 13 (1996), 2707–2714, gr-qc/9604005 [gr-qc]. MR 1413147
- [2] L.F. Abbott and S. Deser, Stability of gravity with a cosmological constant, Nucl. Phys. **B195** (1982), 76–96.

- [3] M.T. Anderson, P.T. Chruściel, and E. Delay, Non-trivial, static, geodesically complete vacuum spacetimes with a negative cosmological constant, Jour. High Energy Phys. 10 (2002), 063, arXiv:gr-qc/0211006. MR 1951922
- [4] ______, Non-trivial, static, geodesically complete spacetimes with a negative cosmological constant. II. $n \ge 5$, AdS/CFT correspondence: Einstein metrics and their conformal boundaries, IRMA Lect. Math. Theor. Phys., vol. 8, Eur. Math. Soc., Zürich, 2005, arXiv:gr-qc/0401081, pp. 165–204. MR MR2160871
- [5] H. Barzegar, P.T. Chruściel, and M. Hörzinger, Energy in higher-dimensional spacetimes, Phys. Rev. D 96 (2017), 124002, 25 pp., arXiv:1708.03122 [gr-qc].
- [6] H. Barzegar, P.T. Chruściel, M. Hörzinger, M. Maliborski, and L. Nguyen, On the energy of asymptotically Horowitz-Myers metrics, Phys. Rev. D 101 (2019), 024007, 16, arXiv:1907.04019 [gr-qc].
- [7] C.P. Boyer, K. Galicki, and J. Kollár, Einstein metrics on spheres, Ann. of Math. (2) 162 (2005), 557–580. MR 2178969
- [8] R. Bryant, https://mathoverflow.net/questions/396848/einstein-metrics-on-connected-sums/397631#397631, 2021.
- [9] J. Chen and X. Zhang, Metrics of Eguchi-Hanson types with the negative constant scalar curvature, Jour. Geom. Phys. 161 (2021), 104010, 10, arXiv:2007.15964 [math.DG]. MR 4180104
- [10] K.S. Chou and T.Y.-H. Wan, Correction to: "Asymptotic radial symmetry for solutions of $\Delta u + e^u = 0$ in a punctured disc" [Pacific J. Math. **163** (1994), no. 2, 269–276; MR1262297 (95a:35038)], Pacific J. Math. **171** (1995), 589–590. MR 1372247
- [11] P.T. Chruściel and E. Delay, Non-singular, vacuum, stationary spacetimes with a negative cosmological constant, Ann. Henri Poincaré 8 (2007), 219–239. MR MR2314449
- [12] _____, Gluing constructions for asymptotically hyperbolic manifolds with constant scalar curvature, Commun. Anal. Geom. 17 (2009), 343–381, arXiv:0711.1557[gr-qc]. MR 2520913 (2011a:53052)
- [13] _____, Exotic hyperbolic gluings, Jour. Diff. Geom. 108 (2018), 243–293, arXiv:1511.07858 [gr-qc].
- [14] _____, The hyperbolic positive energy theorem, (2019), arXiv:1901.05263v1 [math.DG].
- [15] P.T. Chruściel, E. Delay, and P. Klinger, Non-singular spacetimes with a negative cosmological constant: III. Stationary solutions with matter fields, Phys. Rev. D95 (2017), 104039, arXiv:1701.03718 [gr-qc].
- [16] _____, Non-singular spacetimes with a negative cosmological constant: IV. Stationary black hole solutions with matter fields, Class. Quantum Grav. 35 (2017), no. 3, 035007, 15, arXiv:1708.04947 [gr-qc]. MR 3755963
- [17] P.T. Chruściel and G.J. Galloway, Positive mass theorems for asymptotically hyperbolic Riemannian manifolds with boundary, (2021), arXiv:2107.05603 [gr-qc].

- [18] P.T. Chruściel and M. Herzlich, The mass of asymptotically hyperbolic Riemannian manifolds, Pacific Jour. Math. 212 (2003), 231–264, arXiv:math/0110035 [math.DG]. MR MR2038048 (2005d:53052)
- [19] P.T. Chruściel and G. Nagy, The mass of spacelike hypersurfaces in asymptotically anti de Sitter spacetimes, Adv. Theor. Math. Phys. 5 (2001), 697–754, arXiv:gr-qc/0110014.
- [20] P.T. Chruściel and W. Simon, Towards the classification of static vacuum spacetimes with negative cosmological constant, Jour. Math. Phys. 42 (2001), 1779–1817, arXiv:gr-qc/0004032.
- [21] R. Clarkson and Robert B. Mann, Soliton solutions to the Einstein equations in five dimensions, Phys. Rev. Lett. 96 (2006), 051104, arXiv:hep-th/0508109.
- [22] M. Dahl and A. Sakovich, A density theorem for asymptotically hyperbolic initial data satisfying the dominant energy condition, (2015), arXiv:1502.07487 [math.DG].
- [23] D.M. DeTurck and D. Yang, Existence of elastic deformations with prescribed principal strains and triply orthogonal systems, Duke Math. Jour. 51 (1984), 243–260. MR 747867
- [24] D. Dold, Global dynamics of asymptotically locally AdS spacetimes with negative mass, Class. Quantum Grav. **35** (2018), 095012, 26 pp., arXiv:1711.06700 [gr-qc]. MR 3788372
- [25] D. Gilbarg and N.S. Trudinger, *Elliptic partial differential equations of second order*, Springer, Berlin, 1983.
- [26] C. Guillarmou, S. Moroianu, and F. Rochon, Renormalized volume on the Teichmüller space of punctured surfaces, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 17 (2017), no. 1, 323–384. MR 3676051
- [27] S. de Haro, S.N. Solodukhin, and K. Skenderis, *Holographic reconstruction of spacetime* and renormalization in the AdS/CFT correspondence, Commun. Math. Phys. **217** (2001), 595–622, arXiv:hep-th/0002230.
- [28] J. Hempel, 3-manifolds, Princeton University Press, Princeton, 1976, Annals of Mathematics Studies No 86.
- [29] M. Herzlich, Mass formulae for asymptotically hyperbolic manifolds, AdS/CFT correspondence: Einstein metrics and their conformal boundaries, IRMA Lect. Math. Theor. Phys., vol. 8, Eur. Math. Soc., Zürich, 2005, pp. 103–121. MR 2160869
- [30] ______, Computing asymptotic invariants with the Ricci tensor on asymptotically flat and asymptotically hyperbolic manifolds, Ann. Henri Poincaré 17 (2016), 3605–3617, arXiv:1503.00508 [math.DG]. MR 3568027
- [31] S. Holst and P. Peldán, Black holes and causal structure in anti-de Sitter isometric spacetimes, Class. Quantum Grav. 14 (1997), 3433–3452, gr-qc/9705067 [gr-qc]. MR 1492284
- [32] G.T. Horowitz and R.C. Myers, The AdS/CFT correspondence and a new positive energy conjecture for general relativity, Phys. Rev. D 59 (1998), 026005, arXiv:hep-th/9808079.
- [33] J. Isenberg, J.M. Lee, and I. Stavrov Allen, Asymptotic gluing of asymptotically hyperbolic solutions to the Einstein constraint equations, Ann. Henri Poincaré 11 (2010), 881–927, arXiv:0910.1875 [math-dg]. MR 2736526 (2012h:58037)

- [34] D.A. Lee and A. Neves, The Penrose inequality for asymptotically locally hyperbolic spaces with nonpositive mass, Commun. Math. Phys. **339** (2015), 327–352. MR 3370607
- [35] R. Mazzeo and F. Pacard, Maskit combinations of Poincaré-Einstein metrics, Adv. Math. **204** (2006), arXiv:math/0211099 [math.DG].
- [36] R. Mazzeo and M. Taylor, Curvature and uniformization, Israel Jour. Math. 130 (2002), 323–346. MR MR1919383 (2003j:30063)
- [37] C.W. Misner, K. Thorne, and J.A. Wheeler, Gravitation, Freeman, San Fransisco, 1973.
- [38] H. Pedersen, Eguchi-Hanson metrics with cosmological constant, Classical Quantum Gravity 2 (1985), no. 4, 579–587. MR 795103
- [39] T. Sugawa, Estimates of hyperbolic metric with applications to Teichmüller spaces, Kyungpook Math. J. **42** (2002), 51–60. MR 1915906
- [40] Y. Tashiro, Complete Riemannian manifolds and some vector fields, Trans. Amer. Math. Soc. 117 (1965), 251–275. MR 174022
- [41] R.M. Wald, General relativity, University of Chicago Press, Chicago, 1984.
- [42] X. Wang, Mass for asymptotically hyperbolic manifolds, Jour. Diff. Geom. 57 (2001), 273–299. MR MR1879228 (2003c:53044)
- [43] S.A. Wolpert, The hyperbolic metric and the geometry of the universal curve, Jour. Diff. Geom. 31 (1990), 417–472. MR 1037410
- [44] E. Woolgar, The rigid Horowitz-Myers conjecture, Jour. High Energy Phys. 3 (2016), 104, arXiv:1602.06197 [math.DG]. MR 3650713