Anisotropic Caffarelli-Kohn-Nirenberg type inequalities

YanYan Li* and Xukai Yan[†]

Abstract

Caffarelli, Kohn and Nirenberg considered in 1984 the interpolation inequalities

$$|||x|^{\gamma_1}u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2}\nabla u||_{L^p(\mathbb{R}^n)}^a |||x|^{\gamma_3}u||_{L^q(\mathbb{R}^n)}^{1-a}$$

in dimension $n \geq 1$, and established necessary and sufficient conditions for which to hold under natural assumptions on the parameters. Motivated by our study of the asymptotic stability of solutions to the Navier-Stokes equations, we consider a more general and improved anisotropic version of the interpolation inequalities

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\mathbb{R}^n)}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(\mathbb{R}^n)}^{1-a}$$

in dimensions $n \geq 2$, where $x = (x', x_n)$ and $x' = (x_1, ..., x_{n-1})$, and give necessary and sufficient conditions for which to hold under natural assumptions on the parameters. Moreover we extend the Caffarelli-Kohn-Nirenberg inequalities from $q \geq 1$ to q > 0. This extension, together with a nonlinear Poincaré inequality which we obtain in this paper, has played an important role in our proof of the above mentioned anisotropic interpolation inequalities.

1 Introduction

Let $n \geq 1$, $s, p, q, \alpha, \mu, \beta, a$ be real numbers satisfying

$$s > 0, \quad p, q \ge 1, \quad 0 \le a \le 1,$$
 (1.1)

$$\frac{1}{s} + \frac{\gamma_1}{n} > 0, \quad \frac{1}{p} + \frac{\gamma_2}{n} > 0, \quad \frac{1}{q} + \frac{\gamma_3}{n} > 0,$$
 (1.2)

$$\frac{1}{s} + \frac{\gamma_1}{n} = a\left(\frac{1}{p} + \frac{\gamma_2 - 1}{n}\right) + (1 - a)\left(\frac{1}{q} + \frac{\gamma_3}{n}\right),\tag{1.3}$$

^{*}Department of Mathematics, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854, USA. Email: yyli@math.rutgers.edu, is partially supported by NSF Grants DMS-1501004, DMS-2000261, and Simons Fellows Award 677077.

[†]Department of Mathematics, Oklahoma State University, 401 Mathematical Sciences Building, Stillwater, OK 74078, USA. Email: xuyan@okstate.edu, is partially supported by AMS-Simons Travel Grant and AWM-NSF Travel Grant 1642548.

$$\gamma_1 \le a\gamma_2 + (1-a)\gamma_3,\tag{1.4}$$

$$\frac{1}{s} \le \frac{a}{p} + \frac{1-a}{q}$$
 if $a = 0$ or $a = 1$ or $\frac{1}{s} + \frac{\gamma_1}{n} = \frac{1}{p} + \frac{\gamma_2 - 1}{n} = \frac{1}{q} + \frac{\gamma_3}{n}$. (1.5)

Caffarelli, Kohn and Nirenberg established the following classical interpolation inequalities.

Theorem A. ([10], see also [9]) For $n \ge 1$, let $s, p, q, \gamma_1, \gamma_2, \gamma_3$ and a satisfy (1.1) and (1.2). Then there exists some positive constant C such that

$$|||x|^{\gamma_1}u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2}\nabla u||_{L^p(\mathbb{R}^n)}^a |||x|^{\gamma_3}u||_{L^q(\mathbb{R}^n)}^{1-a}$$
(1.6)

holds for all $u \in C_c^1(\mathbb{R}^n)$ if and only if (1.3)-(1.5) hold. Furthermore, on any compact set in the parameter space in which (1.1) and (1.2) hold, the constant C is bounded.

Given (1.1), condition (1.2) holds if and only if $|||x|^{\gamma_1}u||_{L^s(\mathbb{R}^n)}$, $|||x|^{\gamma_2}\nabla u||_{L^p(\mathbb{R}^n)}$ and $|||x|^{\gamma_3}u||_{L^q(\mathbb{R}^n)}$ are finite for all $u \in C_c^{\infty}(\mathbb{R}^n)$.

The above theorem is the same as the theorem in [10], though the formulation of the conditions is somewhat different.

Lin [21] generalized (1.6) to include derivatives of any order. Badiale and Tarantello [1] derived a cylindrical Sobolev-Hardy type inequality. Bahouri, Chemin and Gallagher [4, 5] obtained refined Hardy inequalities. Nguyen and Squassina [24, 25] generalized the Caffarelli-Kohn-Nirenberg inequalities to fractional Sobolev spaces. Best constants and the existence (and nonexistence) of extremal functions of (1.6) have been studied extensively, see Catrina and Wang [11], Dolbeault, Esteban and Loss [12], and the references therein. Partly motivated by works of Bourgain, Brezis and Mironescu [2, 3] and Maz'ya and Shaposhnikova [22], Frank and Seiringer [13] identified best constants for fractional Hardy type inequalities. Sharp Sobolev and isoperimetric inequalities with monomial weights, and related problems, are studied by Cabré, Ros-Oton and Serra, see [7, 8].

In this paper, we prove the following theorem on anisotropic Caffarelli-Kohn-Nirenberg type inequalities.

For $n \geq 2$, let $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β be real numbers satisfying

$$s, q > 0, \quad p \ge 1, \quad 0 \le a \le 1,$$
 (1.7)

$$\frac{1}{s} + \frac{\alpha}{n-1} > 0, \quad \frac{1}{p} + \frac{\mu}{n-1} > 0, \quad \frac{1}{q} + \frac{\beta}{n-1} > 0,$$
 (1.8)

$$\frac{1}{s} + \frac{\alpha + \gamma_1}{n} > 0, \quad \frac{1}{p} + \frac{\mu + \gamma_2}{n} > 0, \quad \frac{1}{q} + \frac{\beta + \gamma_3}{n} > 0,$$
 (1.9)

$$\frac{1}{s} + \frac{\gamma_1 + \alpha}{n} = a\left(\frac{1}{p} + \frac{\gamma_2 + \mu - 1}{n}\right) + (1 - a)\left(\frac{1}{q} + \frac{\gamma_3 + \beta}{n}\right),\tag{1.10}$$

$$\gamma_1 \le a\gamma_2 + (1-a)\gamma_3,\tag{1.11}$$

$$\gamma_1 + \alpha \le a(\gamma_2 + \mu) + (1 - a)(\gamma_3 + \beta),$$
 (1.12)

$$\frac{1}{s} + \frac{\alpha}{n-1} \ge a\left(\frac{1}{p} + \frac{\mu - 1}{n-1}\right) + (1-a)\left(\frac{1}{q} + \frac{\beta}{n-1}\right),\tag{1.13}$$

$$\frac{1}{s} \le \frac{a}{p} + \frac{1-a}{q} \text{ if } a = 0 \text{ or } a = 1 \text{ or } \frac{1}{p} + \frac{\gamma_2 + \mu - 1}{n} = \frac{1}{q} + \frac{\gamma_3 + \beta}{n} = \frac{1}{s} + \frac{\gamma_1 + \alpha}{n},$$

$$\text{or } \frac{1}{s} + \frac{\alpha}{n-1} = a\left(\frac{1}{p} + \frac{\mu - 1}{n-1}\right) + (1-a)\left(\frac{1}{q} + \frac{\beta}{n-1}\right).$$
(1.14)

Throughout the paper, we denote $x = (x', x_n)$, where $x' = (x_1, ..., x_{n-1})$. We have the following theorem.

Theorem 1.1. For $n \geq 2$, let $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β be real numbers satisfying (1.7)-(1.9). Then there exists some positive constant C such that

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\mathbb{R}^n)}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(\mathbb{R}^n)}^{1-a}$$
(1.15)

holds for all $u \in C_c^1(\mathbb{R}^n)$ if and only if (1.10)-(1.14) hold. Furthermore, on any compact set in the parameter space in which (1.7)-(1.9) hold, the constant C is bounded.

Given (1.7), conditions (1.8) and (1.9) hold if and only if $||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(\mathbb{R}^n)}$, $||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\mathbb{R}^n)}$ and $||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(\mathbb{R}^n)}$ are finite for all $u \in C_c^{\infty}(\mathbb{R}^n)$.

Inequality (1.15) was proved in [1] in the special cases when $n \ge 3$, a = 1, $\gamma_1 = \gamma_2 = \mu = 0$, $1/s + \alpha/n = 1/p - 1/n$, $1 , <math>-1 \le \alpha \le 0$ and $(1/p - 1/n)(n - 1) + \alpha/n > 0$; and proved in [19] in the special cases when $n \ge 2$, a = 1, $1 \le s = p < n$, $\alpha p > 1 - n$, $\mu p > 1 - n$, $(\alpha + \gamma_1)p > -n$, $\gamma_1 + \alpha = \gamma_2 + \mu - 1$ and $\gamma_1 \le \gamma_2$.

Taking $\alpha = \mu = \beta = 0$, inequality (1.15) is an improvement of the Caffarelli-Kohn-Nirenberg inequalities (Theorem A) from $q \ge 1$ to q > 0.

When $\alpha < 0$ and $\mu, \beta > 0$, inequality (1.15) strengthens

$$|||x|^{\gamma_1 + \alpha} u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2 + \mu} \nabla u||_{L^p(\mathbb{R}^n)}^a |||x|^{\gamma_3 + \beta} u||_{L^q(\mathbb{R}^n)}^{1 - a}$$

which is given by (1.6). In particular, when $n \ge 3$, a = 1, s = p = 2, $\gamma_1 = \gamma_2 = \alpha = -1/2$, and $\mu = 1/2$, inequality (1.15) takes the form

$$\int_{\mathbb{R}^n} \frac{|u|^2}{|x||x'|} dx \le C \int_{\mathbb{R}^n} |\nabla u|^2 \frac{|x'|}{|x|} dx, \tag{1.16}$$

which strengthens the Hardy inequality

$$\int_{\mathbb{R}^n} \frac{|u|^2}{|x|^2} dx \le C \int_{\mathbb{R}^n} |\nabla u|^2 dx.$$

Inequality (1.16) was among the special cases proved in [19] as mentioned above.

The necessity of (1.10) is proved by dimensional analysis. The necessity of (1.11) and (1.12) are deduced from the fact that if (1.15) holds for u, then it also holds for $u(x) \mapsto u(x_1 + S, x_2, ..., x_{n-1}, x_n + T)$ for all S, T > 0. The necessity of (1.13) and (1.14) are more delicate.

A main ingredient of the proof of Theorem 1.1, even in the case when $q \ge 1$, is the above mentioned improvement of Theorem A from $q \ge 1$ to q > 0, which is stated as the following theorem.

Theorem 1.2. For $n \ge 1$, let $s, p, q, \gamma_1, \gamma_2, \gamma_3$, and a satisfy (1.7) and (1.2). Then there exists some positive constant C, such that

$$|||x|^{\gamma_1}u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2}\nabla u||_{L^p(\mathbb{R}^n)}^a|||x|^{\gamma_3}u||_{L^q(\mathbb{R}^n)}^{1-a}$$
(1.17)

holds for all $u \in C_c^1(\mathbb{R}^n)$ if and only if (1.3)-(1.5) hold. Furthermore, on any compact set in the parameter space in which (1.7) and (1.2) hold, the constant C is bounded.

Our proof of the sufficiency part of Theorem 1.2, which yields the extension from $q \ge 1$ to q > 0, is quite different from the proof of Theorem A in [10].

Another main ingredient of the proof of Theorem 1.1 is the following nonlinear Poincaré inequality.

Theorem 1.3 (A nonlinear Poincaré inequality). For $n \geq 1$ and $0 < \lambda < \infty$, assume $1 \leq p \leq \infty$ if $1 \leq \lambda < \infty$, and $\max\{1, n/(1+n\lambda)\} \leq p \leq \infty$ if $0 < \lambda < 1$. Let (M, g) be an n-dimensional smooth compact Riemannian manifold without boundary, $\Omega = M$ or $\Omega \subset M$ be an open connected set with Lipschitz boundary, and $S \subset \Omega$ has positive measure |S|. Then there exists some positive constant C, depending only on p, λ , Ω and S, such that for every nonnegative $w \in W^{1,p}(\Omega)$,

$$\|w - \left(f_S w^{1/\lambda}\right)^{\lambda}\|_{L^p(\Omega)} \le C\|\nabla w\|_{L^p(\Omega)},\tag{1.18}$$

On the other hand, if $0 < \lambda < 1$ and $0 or <math>0 < \lambda < \infty$ and 0 , there does not exist C for which (1.18) holds.

Theorem 1.3 gives necessary and sufficient conditions on $(\lambda, p) \in (0, \infty) \times (0, \infty]$ for (1.18) to hold.

Corollary 1.1. For $n \geq 1$, $1 \leq p \leq \infty$, and $0 < q < \infty$, let (M,g) be an n-dimensional smooth compact Riemannian manifold without boundary, $\Omega \subset M$ be an open connected set with Lipschitz boundary, and $S \subset \Omega$ has positive measure |S|. Then there exists some positive constant C, depending only on p, q, Ω and S, such that for every nonnegative $w \in W^{1,p}(\Omega)$,

$$||w||_{L^p(\Omega)} \le \left(\oint_S w^q \right)^{1/q} |\Omega|^{1/q} + C ||\nabla w||_{L^p(\Omega)}.$$
 (1.19)

Theorem 1.1 grows out of our study in [19] on the stability of solutions to Navier-Stokes equations. In joint work with L. Li [17, 18], we classified (-1)-homogeneous axisymmetric no-swirl solutions of the 3D incompressible stationary Navier-Stokes equations which are smooth in \mathbb{R}^3 away from the symmetry axis. All such solutions u are of the following three mutually exclusive types:

- Type 1. Landau solutions, which satisfy $\sup_{|x|=1} |\nabla u(x)| < \infty$;
- Type 2. Solutions satisfying $0 < \limsup_{|x|=1, x' \to 0} |x'| |\nabla u(x)| < \infty;$
- Type 3. Solutions satisfying $\limsup_{|x|=1,x'\to 0} |x'|^2 |\nabla u(x)| > 0$.

Karch and Pilarczyk [15] proved the asymptotic stability of Landau solutions under L^2 -perturbations. In [19], we proved the asymptotic stability of Type 2 solutions under L^2 -perturbations. An important ingredient in our proof is the following improved version of Hardy's inequality

$$\int \frac{|u|^2}{|x||x'|} \le C \int |\nabla u|^2, \tag{1.20}$$

a weaker form of (1.16). We expect that Theorem 1.1 will be useful in the study of the asymptotic stability of Type 3 solutions and the stability of Type 2 solutions in other function spaces. For related results on the stability of singular solutions to the Navier-Stokes equations, see [6], [16], [20], [26] and the reference therein.

In the special case when a=1 and $1 \le s=p < n$, Theorem 1.1 says that under the conditions

$$p \ge 1, \quad \frac{1}{p} + \frac{\mu}{n-1} > 0, \quad \frac{1}{p} + \frac{\gamma_1 + \alpha}{n} > 0,$$
 (1.21)

$$\gamma_1 + \alpha = \gamma_2 + \mu - 1, \quad \gamma_1 \le \gamma_2, \tag{1.22}$$

inequality

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^p(\mathbb{R}^n)} \le C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\mathbb{R}^n)}$$
(1.23)

holds for $u \in C_c^1(\mathbb{R}^n)$. This was proved in [19] when $1 \leq p < n$ as mentioned earlier. The proof there applies to $p \geq n$ as well, and we present the proof concisely below.

The necessity of $\gamma_1 + \alpha = \gamma_2 + \mu - 1$ follows from a dimensional analysis argument. The necessity of $\gamma_1 \leq \gamma_2$ can also be seen easily: take a unit ball B_i centered at $x_i = (x_i', i)$ with $2 < |x_i'| < 3$ and let $u_i(\cdot) = v(\cdot + x_i)$.

Let $x = (r, \theta)$ in spherical coordinates. Since $(\gamma_1 + \alpha)p + n - 1 > -1$, we have, for each fixed θ , that

$$\int_0^\infty r^{(\gamma_1+\alpha)p+n-1}|u|^p dr \le C \int_0^\infty r^{(\gamma_1+\alpha+1)p+n-1}|\partial_r u|^p dr. \tag{1.24}$$

For any $0 < \epsilon < \pi/4$, let $K_{\epsilon} := \{x \in \mathbb{R}^n \mid |x'| \le |x| \sin \epsilon\}$. Integrate the above in θ on $(\mathbb{R}^n \setminus K_{\epsilon}) \cap \mathbb{S}^{n-1}$, we have, using $|x| \sin \epsilon \le |x'| \le |x|$ in $\mathbb{R}^n \setminus K_{\epsilon}$ and $\gamma_1 + \alpha + 1 = \gamma_2 + \mu$, that

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^p(\mathbb{R}^n\setminus K_{\epsilon})} \le C|||x|^{\gamma_2}|x'|^{\mu}\partial_r u||_{L^p(\mathbb{R}^n\setminus K_{\epsilon})}.$$
(1.25)

Since $\int_{K_{2\epsilon}\backslash K_{\epsilon}} |x|^{\gamma_1 p} |x'|^{\alpha p} |u|^p dx = \int_{\epsilon}^{2\epsilon} \int_{\partial K_{\delta}} |x|^{\gamma_1 p} |x'|^{\alpha p} |\nabla u|^p \frac{|x|}{\cos \delta} d\sigma(x) d\delta$, there is some $\epsilon < \delta < 2\epsilon$, such that

$$|||x|^{\gamma_1}|x'|^{\alpha}|x|^{1/p}u||_{L^p(\partial K_s)} \le C|||x|^{\gamma_2}|x'|^{\mu}\partial_r u||_{L^p(K_{2s}\setminus K_s)}. \tag{1.26}$$

Next, let $x = (r, \theta) = (r, \theta_1, \omega)$, where r = |x|, $\omega \in \mathbb{S}^{n-2}$ and θ_1 is the polar angle, i.e. the angle between the x_n -axis and the ray from the origin to x. Then $|x'| = |x| \sin \theta_1$, and

$$|u(r,\theta_1,\omega)|^p - |u(r,\delta,\omega)|^p = -\int_{\theta_1}^{\delta} \partial_t |u(r,t,\omega)|^p dt \le \int_{\theta_1}^{\delta} p|u|^{p-1} |\partial_t u| dt.$$

We multiply the above by $\theta_1^{\alpha p+n-2}$ and integrate in θ_1 over $[0, \delta]$. We know from (1.21) and (1.22) that $\alpha p + n - 1 > 0$. So we have

$$\int_0^\delta \theta_1^{\alpha p+n-2} \int_{\theta_1}^\delta p|u|^{p-1} |\partial_t u| dt d\theta_1 \le C \int_0^\delta \theta_1^{\alpha p+n-1} |u|^{p-1} |\partial_{\theta_1} u| d\theta_1,$$

and

$$\int_0^\delta \theta_1^{\alpha p+n-2} |u(r,\theta_1,\omega)|^p d\theta_1 \le C|u(r,\delta,\omega)|^p + C \int_0^\delta \theta_1^{\alpha p+n-1} |u|^{p-1} |\partial_{\theta_1} u| d\theta_1.$$

Using Cauchy-Schwarz inequality, we have

$$\int_0^\delta \theta_1^{\alpha p+n-2} |u(r,\theta_1,\omega)|^p d\theta_1 \leq C|u(r,\delta,\omega)|^p + C \int_0^\delta \theta_1^{(\alpha+1)p+n-2} |\partial_{\theta_1} u(r,\theta_1,\omega)|^p d\theta_1.$$

Multiplying the above by $r^{(\gamma_1+\alpha)p+n-1}$ and integrating in r and ω , we have, using the fact that $\gamma_1+\alpha+1=\gamma_2+\mu,\ \theta_1^{\alpha+1}\leq\theta_1^{\mu}$ on $[0,\delta]$ in view of $\mu\leq\alpha+1$, and $|\partial_{\theta_1}u|/r\leq|\nabla u|$, that

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^p(K_{\delta})} \le C|||x|^{\gamma_1}|x'|^{\alpha}|x|^{1/p}u||_{L^p(K_{\delta})} + C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(K_{\delta})}.$$

Inequality (1.23) follows from (1.25), (1.26) and the above.

For the sufficiency part of Theorem 1.1 when $s \neq p$ or 0 < a < 1, the proof is more involved. Let us look at a simple case where n = 2, a = 1, $\gamma_1 = \gamma_2 = \gamma_3 = 0$, s = 2, p = 1, $\mu = \alpha > -1/2$. The following lemma is a slightly stronger version of Theorem 1.1 in this case.

Proposition 1.1. For $\alpha > -1/2$, there exists some constant C depending only on α such that

$$|||x_1|^{\alpha}u||_{L^2([0,1]^2)}^2 \le C|||x_1|^{\alpha}\partial_{x_1}u||_{L^1([0,1]^2)}|||x_1|^{\alpha}\partial_{x_2}u||_{L^1([0,1]^2)}$$
(1.27)

holds for all $u \in C^1([0,1]^2)$ satisfying $u(1,x_2) = u(x_1,1) = 0, \ 0 \le x_1, x_2 \le 1$.

Proof. Make a change of variables $y_1 = x_1^{2\alpha+1}$, $y_2 = x_2$, and $\tilde{u}(y_1, y_2) = u(x_1, x_2)$. Then (1.27) is equivalent to

$$\|\tilde{u}\|_{L^{2}([0,1]^{2})}^{2} \leq C\||y_{1}|^{\beta}\partial_{y_{1}}\tilde{u}\|_{L^{1}([0,1]^{2})}\||y_{1}|^{-\beta}\partial_{y_{2}}\tilde{u}\|_{L^{1}([0,1]^{2})},\tag{1.28}$$

for $\beta < 1/2$ (where $\beta := \alpha/(2\alpha + 1)$) and $\tilde{u} \in C^1([0,1]^2)$ satisfying $\tilde{u}(y_1, 1) = \tilde{u}(1, y_2) = 0, \ 0 \le y_1, y_2 \le 1$.

For $k \in \mathbb{N}$, let $R_k = [2^{-k-1}, 2^{-k}]$,

$$A_k := \|\tilde{u}\|_{L^2(R_k \times [0,1])}, \quad P_k := \|y_1^{\beta} \partial_{y_1} \tilde{u}\|_{L^1(R_k \times [0,1])}, \quad Q_k := \|y_1^{-\beta} \partial_{y_2} \tilde{u}\|_{L^1(R_k \times [0,1])},$$

and

$$P:=\||y_1|^\beta\partial_{y_1}\tilde{u}\|_{L^1([0,1]^2)},\quad Q:=\||y_1|^{-\beta}\partial_{y_2}\tilde{u}\|_{L^1([0,1]^2)}.$$

For any $y_1 \in R_k$, $\xi \in R_{k-1}$ and $y_2 \in [0,1]$, we have

$$\begin{split} |\tilde{u}(y_1, y_2)|^2 &= |\tilde{u}(y_1, y_2)| |\tilde{u}(\xi, y_2) - \int_{y_1}^{\xi} \partial_{\eta} \tilde{u}(\eta, y_2) d\eta | \\ &\leq |\tilde{u}(y_1, y_2)| |\tilde{u}(\xi, y_2)| + C y_1^{-\beta} |\tilde{u}(y_1, y_2)| \int_{y_1}^{\xi} \eta^{\beta} |\partial_{\eta} \tilde{u}(\eta, y_2)| d\eta \\ &\leq |\tilde{u}(y_1, y_2)| |\tilde{u}(\xi, y_2)| + C \int_{0}^{1} y_1^{-\beta} |\partial_{y_2} \tilde{u}(y_1, y_2)| dy_2 \int_{R_k \cup R_{k-1}} \eta^{\beta} |\partial_{\eta} \tilde{u}(\eta, y_2)| d\eta. \end{split}$$

Taking $\int_0^1 f_{R_{k-1}} \int_{R_k} dy_1 d\xi dy_2$ of the above and using Hölder's inequality, we have

$$A_{k} \leq \frac{1}{|R_{k-1}|} \int_{0}^{1} \left(\int_{R_{k}} |\tilde{u}(y_{1}, y_{2})| dy_{1} \right) \left(\int_{R_{k-1}} |\tilde{u}(\xi, y_{2})| d\xi \right) dy_{2} + CQ_{k}P$$

$$\leq \sqrt{\frac{|R_{k}|}{|R_{k-1}|}} \sqrt{A_{k}A_{k-1}} + CQ_{k}P \leq \frac{1}{2\sqrt{2}} (A_{k} + A_{k-1}) + CQ_{k}P.$$

Thus

$$A_k \leq \theta A_{k-1} + CQ_k P$$
,

where $\theta = 1/(2\sqrt{2} - 1) < 1$. Suming over $k \ge 1$ gives

$$\sum_{k=1}^{\infty} A_k \le \frac{1}{1-\theta} A_0 + CPQ \le CPQ. \tag{1.29}$$

where we have used $|\tilde{u}(y_1,y_2)| \leq \int_0^1 |\partial_{y_1}\tilde{u}(y_1,\cdot)|$ and $|\tilde{u}(y_1,y_2)| \leq \int_0^1 |\partial_{y_2}\tilde{u}(\cdot,y_2)|$.

For the sufficiency part of Theorem 1.1 in general, our first consideration was for $q \geq 1$. We were able to prove the sufficiency part of Theorem 1.1 for $q \geq 1$ in dimension n provided Theorem A for q > 0 in dimension n - 1, with the help of the nonlinear Poincaré's inequality (Theorem 1.3). We also proved Theorem A for q > 0 in dimension n = 1 and therefore proved Theorem 1.1 for $q \geq 1$ in dimension n = 2 as well as Theorem 1.1 for axisymmetric u and $q \geq 1$ in dimensions $n \geq 3$.

Next we established the sufficiency part of Theorem 1.1 for $q \ge 1$ in dimensions $n \ge 3$. A key step is to prove (1.15) on a cylinder $D := \{x \in \mathbb{R}^n \mid |x'| \le 1, 0 \le x_n \le 1\}$ when $\gamma_1 = \gamma_2 = \gamma_3 = 0$. For simplicity, one may consider $u \in C_c^1(D)$ and the estimate is

$$|||x'|^{\alpha}u||_{L^{s}(D)} \le C|||x'|^{\mu}\nabla u||_{L^{p}(D)}^{a}|||x'|^{\beta}u||_{L^{q}(D)}^{1-a}.$$
(1.30)

The left hand side of the above can be written as

$$|||x'|^{\alpha}u||_{L^{s}(D)} = |||x'|^{\frac{\alpha s}{\bar{s}}}|u|^{\frac{s}{\bar{s}}}||_{L^{\bar{s}}(D)}^{\frac{\bar{s}}{\bar{s}}}| \leq C|||x'|^{\frac{\alpha s}{\bar{s}}}(|u|^{\frac{s}{\bar{s}}} - |u^{*}|^{\frac{s}{\bar{s}}})||_{L^{\bar{s}}(D)}^{\frac{\bar{s}}{\bar{s}}} + C|||x'|^{\frac{\alpha s}{\bar{s}}}|u^{*}|^{\frac{\bar{s}}{\bar{s}}}||_{L^{\bar{s}}(D)}^{\frac{\bar{s}}{\bar{s}}} = C(I_{1} + I_{2}),$$

where $1/\bar{s} = 1/s + 1 - 1/p$ and $u^*(x', x_n) = \int_{|y'| = |x'|} u(y', x_n) d\sigma(y')$. Since Theorem 1.1 for $q \ge 1$ holds for axisymmetric u, so does (1.30). Thus I_2 is bounded by the right

hand side of (1.30). The estimate that I_1 is bounded by the right hand side of (1.30) follows from a variant of the Caffarelli-Kohn-Nirenberg inequalities (see Theorem 6.2), using the fact that $0 < \bar{s} < s$.

Later we proved Theorem A for q > 0 in all dimensions and in turn proved Theorem 1.1. This is the proof presented in this paper.

In Section 2, we prove the necessity part of Theorem 1.1 and 1.2. In Section 3, we prove Theorem 1.3 and Corollary 1.1. In Section 4, we prove the sufficiency part of Theorem 1.2 by establishing Theorem 4.1, a more general result including inequalities on cones. In Section 5, we prove the sufficiency part of Theorem 1.1. In Section 6, we give two variants of Theorem A and Theorem 1.1. Some properties of the parameters used in the proofs are given in the appendix.

2 Proof of the necessity parts of Theorem 1.1 and 1.2

In this section, we prove the necessity parts of Theorem 1.1 and 1.2. We first prove the necessity part of Theorem 1.1 by the following lemma.

Lemma 2.1. For $n \geq 2$, let $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β satisfy (1.7)-(1.9). If there exists a constant C such that (1.15) holds for all u in $C_c^{\infty}(\mathbb{R}^n)$, then (1.10)-(1.14) hold.

Proof. Let C denote a positive constant depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β which may vary from line to line. We prove (1.10)-(1.14) one by one.

Proof of (1.10): Fixing a $v \in C_c^{\infty}(\mathbb{R}^n) \setminus \{0\}$ and plugging $u(x) := v(\lambda x), \ \lambda > 0$, into (1.15), we have

$$\lambda^{-nA_1} \||x|^{\gamma_1} |x'|^{\alpha} v\|_{L^s(\mathbb{R}^n)} \le C \lambda^{-nA_2} \||x|^{\gamma_2} |x'|^{\mu} \nabla v\|_{L^p(\mathbb{R}^n)}^a \||x|^{\gamma_3} |x'|^{\beta} v\|_{L^q(\mathbb{R}^n)}^{1-a},$$

where A_1 and A_2 are the left and right hand side of (1.10) respectively. Sending λ to 0 and ∞ in the above, we obtain (1.10).

Proof of (1.11) and (1.12): Fixing a $v \in C_c^{\infty}(B_1(0)) \setminus \{0\}$, we consider $u(x) := v(x-x_0)$ where $x_0 = (S, 0, ..., 0, R)$ and S, R > 0. Then $u \in C_c^{\infty}(B_1(x_0))$, and u satisfies (1.15). Choose S = 2 and R large. For $x \in B_1(x_0)$, we have $2 \le |x'| \le 3$ and $R/2 \le |x| \le 2R$. Plugging u into (1.15), we have,

$$R^{\gamma_1} \le C R^{a\gamma_2 + (1-a)\gamma_3}$$

for some constant C independent of R. Inequality (1.11) follows, since R can be arbitrarily large.

Now we choose large S and R = 0. For $x \in B_1(x_0)$, both |x'| and |x| are in [S/2, S]. Plugging u into (1.15), we have

$$S^{\gamma_1 + \alpha} \le C S^{a(\gamma_2 + \mu) + (1 - a)(\gamma_3 + \beta)}.$$

Inequality (1.12) follows from the above, since S can be arbitrarily large.

Next, to prove (1.13) and (1.14), we fix a $g \in C_c^{\infty}(1,4)$ satisfying

$$g(t) = \begin{cases} 0, & t \le 1 \text{ or } t \ge 4, \\ 1, & 2 \le t \le 3. \end{cases}$$

Proof of (1.13): For $0 < \epsilon < 1$, let

$$f_1(\rho) = \begin{cases} 0, & \rho \ge 2\epsilon, \\ 2\epsilon - \rho, & \epsilon \le \rho \le 2\epsilon, \\ \epsilon, & \rho \le \epsilon. \end{cases}$$

Then $u(x) := f_1(|x'|)g(x_n)$ satisfies (1.15). We have supp $u \subset \{|x'| \le 2\epsilon, 1 \le x_n \le 4\}$. For any x in supp u, $1 \le |x| \le 5$. Then (1.15) for this u is equivalent to

$$|||x'|^{\alpha}u||_{L^{s}(\mathbb{R}^{n})} \leq C|||x'|^{\mu}\nabla u||_{L^{p}(\mathbb{R}^{n})}^{a}|||x'|^{\beta}u||_{L^{q}(\mathbb{R}^{n})}^{1-a}$$

for some constant C independent of ϵ . By calculation,

$$\int_{\mathbb{R}^n} ||x'|^{\alpha} u|^s dx \ge \frac{1}{C} \int_{5\epsilon/4 < |x'| < 7\epsilon/4} ||x'|^{\alpha} f_1(|x'|)|^s dx' \ge \frac{1}{C} \epsilon^{(\alpha+1)s+n-1},$$

$$\int_{\mathbb{R}^n} ||x'|^{\mu} \nabla u|^p dx \leq C \int_{|x'| < 2\epsilon} |x'|^{p\mu} (|f_1'(|x'|)|^p + |f_1(|x'|)|^p) dx' \leq C \epsilon^{\mu p + n - 1},$$

$$\int_{\mathbb{R}^n} ||x'|^{\beta} u|^q dx \le C \int_{|x'| \le 2\epsilon} ||x'|^{\beta} f_1(|x'|)|^q dx' \le C \epsilon^{(\beta+1)q+n-1}.$$

Thus we have

$$\epsilon^{\alpha+1+(n-1)/s} \le C\epsilon^{a(\mu+(n-1)/p)+(1-a)(\beta+1+(n-1)/q)}$$
.

Inequality (1.13) follows, since ϵ can be arbitrarily small.

Proof of (1.14): We divide the proof into two cases.

Case 1.
$$a = 0$$
 or $a = 1$ or $1/p + (\gamma_2 + \mu - 1)/n = 1/q + (\gamma_3 + \beta)/n = 1/s + (\gamma_1 + \alpha)/n$.

We first prove the inequality in (1.14) when $1/p + (\gamma_2 + \mu - 1)/n = 1/q + (\gamma_3 + \beta)/n = 1/s + (\gamma_1 + \alpha)/n$. For $0 < \epsilon < 1$, let

$$f_2(r) = \begin{cases} r^{-\alpha - \gamma_1 - n/s + \epsilon}, & 0 < r \le 1, \\ 1, & 1 \le r \le 2, \\ \frac{4 - r}{2}, & 2 \le r \le 4, \\ 0, & r \ge 4. \end{cases}$$
 (2.1)

Then $u(x) := f_2(|x|)g(|x_n|/|x'|)$ satisfies (1.15) by the approximation of $u_\delta(x) := f_2(\sqrt{|x|^2 + \delta^2})g(|x_n|/|x'|)$. By computation,

$$\int_{\mathbb{R}^n} ||x|^{\gamma_1} |x'|^{\alpha} u|^s dx \ge \frac{1}{C} \int_{0 < |x| \le 1, 2 \le |x_n|/|x'| \le 3} ||x|^{\gamma_1 + \alpha} f_2|^s dx \ge \frac{1}{C} \int_0^1 r^{\epsilon s - 1} dr \ge \frac{1}{C} \epsilon^{-1}.$$
(2.2)

Notice that for $1 \le |x_n|/|x'| \le 4$, $|\nabla g(|x_n|/|x'|)| \le C|g'(|x_n|/|x'|)|/|x'| \le C|x|^{-1}$, we have, using the fact that $1/s + (\gamma_1 + \alpha)/n = 1/p + (\gamma_2 + \mu - 1)/n > 0$, that

$$\int_{\mathbb{R}^{n}} ||x|^{\gamma_{2}}|x'|^{\mu} \nabla u|^{p} dx \leq C \int_{|x| \leq 1, 1 \leq |x_{n}|/|x'| \leq 4} |x|^{p(\gamma_{2} + \mu)} (|\nabla f_{2}|^{p} + |x|^{-p} |f_{2}|^{p}) dx
\leq C \int_{0}^{4} r^{(\gamma_{2} + \mu)p + n - 1 + (-\alpha - \gamma_{1} - n/s - 1 + \epsilon)p} dr
= C \int_{0}^{4} r^{\epsilon p - 1} dr \leq C \epsilon^{-1}.$$
(2.3)

Similarly, using the fact $1/s + (\gamma_1 + \alpha)/n = 1/q + (\gamma_3 + \beta)/n > 0$, we have

$$\int_{\mathbb{R}^n} ||x|^{\gamma_3} |x'|^{\beta} u|^q dx \le C \int_{|x| \le 1, 1 \le |x_n|/|x'| \le 4} ||x|^{\gamma_3 + \beta} f_2(x)|^q dx \le C \epsilon^{-1}. \tag{2.4}$$

By (1.15), (2.2), (2.3) and (2.4), we have

$$\epsilon^{-1/s} < C\epsilon^{-a/p - (1-a)/q}$$

for arbitrarily small ϵ . So the inequality in (1.14) holds.

Now we turn to a=0 or a=1. In view of (1.10), when a=0, we have $1/s+(\gamma_1+\alpha)/n=1/q+(\gamma_3+\beta)/n$, and when a=1, we have $1/s+(\gamma_1+\alpha)/n=1/p+(\gamma_2+\mu-1)/n$. The inequality in (1.14) follows from the same proof as above.

Case 2. $0 < a < 1, 1/p + (\gamma_2 + \mu - 1)/n \neq 1/q + (\gamma_3 + \beta)/n$, and

$$\frac{1}{s} + \frac{\alpha}{n-1} = a\left(\frac{1}{p} + \frac{\mu - 1}{n-1}\right) + (1-a)\left(\frac{1}{q} + \frac{\beta}{n-1}\right). \tag{2.5}$$

If (2.5) holds, then either Case 1 or Case 2 holds.

We divide the proof of Case 2 into two subcases.

Subcase 2.1.
$$1/p + (\mu - 1)/(n - 1) = 1/q + \beta/(n - 1)$$
.

In this subcase, we have, in view of (2.5), that $1/s + \alpha/(n-1) = 1/p + (\mu-1)/(n-1) = 1/q + \beta/(n-1)$. For $0 < \epsilon < 1$, let

$$f_3(\rho) = \begin{cases} \rho^{-\alpha - (n-1)/s + \epsilon}, & 0 < \rho \le 1, \\ 1, & 1 \le \rho \le 2, \\ \frac{4 - \rho}{2}, & 2 \le \rho \le 4, \\ 0, & \rho \ge 4. \end{cases}$$

Let $u(x) := f_3(|x'|)g(x_n)$. Then it satisfies (1.15) by the approximation of $u_{\delta}(x) := f_3(\sqrt{|x'|^2 + \delta^2})g(x_n)$. By computation, we have

$$\int_{\mathbb{R}^n} ||x|^{\gamma_1} |x'|^{\alpha} u|^s dx \ge \frac{1}{C} \int_{0 \le |x'| \le 1} ||x'|^{\alpha} f_3(|x'|)|^s dx' \ge \frac{1}{C} \int_0^1 \rho^{-1+\epsilon s} d\rho \ge \frac{1}{C} \epsilon^{-1}. \quad (2.6)$$

Since $1/s + \alpha/(n-1) = 1/p + (\mu - 1)/(n-1) > 0$, we have

$$\int_{\mathbb{R}^{n}} ||x|^{\gamma_{2}}|x'|^{\mu} \nabla u|^{p} dx \leq C \int_{|x'| \leq 1} |x'|^{\mu p} (|\nabla f|^{p} + |f|^{p}) dx'
\leq C \int_{0}^{4} \rho^{\mu p + (\alpha - (n-1)/s - 1 + \epsilon)p + n - 2} d\rho
= C \int_{0}^{4} \rho^{\epsilon p - 1} d\rho \leq C \epsilon^{-1}.$$
(2.7)

Similarly, since $1/s + \alpha/(n-1) = 1/q + \beta/(n-1) > 0$, we have

$$\int_{\mathbb{R}^n} ||x|^{\gamma_3} |x'|^{\beta} u|^q dx \le C \int_{|x'| \le 1} ||x'|^{\beta} f_3(x')|^q dx' \le C\epsilon^{-1}. \tag{2.8}$$

So by (1.15), (2.6), (2.7) and (2.8), we have

$$\epsilon^{-1/s} < C \epsilon^{-a/p - (1-a)/q}$$

for arbitrarily small ϵ . So the inequality in (1.14) follows in this subcase.

Subcase 2.2.
$$1/p + (\mu - 1)/(n - 1) \neq 1/q + \beta/(n - 1)$$
.

Introduce the spherical coordinates in \mathbb{R}^n_+ : r = |x|, $\theta = x/|x|$. Let $\theta' = x'/|x|$, we have $\theta = (\theta', \sqrt{1 - |\theta'|^2})$. For simplicity, we denote $x = (r, \theta')$. Fix $\delta > 0$ small, and let $R_0 := \{x \in \mathbb{R}^n_+ \mid 1 < r < 2, \delta < |\theta'| < 2\delta\}$. Fix a function $u \in C_c^{\infty}(R_0) \setminus \{0\}$, and let

$$u_j(r, \theta') = 2^{(b_1\kappa + d_1)j} u(2^{\kappa j}r, 2^j \theta'), \quad j \ge 1,$$

where $b_1 = n/s + \gamma_1 + \alpha$, $d_1 = (n-1)/s + \alpha$, and κ is some j-independent constant to be determined later. Then $u_j \in C_c^{\infty}(R_j)$, where

$$R_j := \{ x \in \mathbb{R}^n \mid 2^{-\kappa j} < r < 2^{-\kappa j+1}, \ 2^{-j}\delta < |\theta'| < 2^{-j+1}\delta \}.$$

Denote

$$I_0 := \||x|^{\gamma_1}|x'|^{\alpha}u\|_{L^s(R_0)}, \quad A_0 := \||x|^{\gamma_2}|x'|^{\mu}\nabla u\|_{L^p(R_0)}, \quad B_0 := \||x|^{\gamma_3}|x'|^{\beta}u\|_{L^q(R_0)}.$$

In the following, the notation $A \simeq B$ means $B/C \le A \le CB$, and $A \lesssim B$ means $A \le CB$, for some C > 1 depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β .

For any $\kappa \in \mathbb{R}$, we have

$$|||x|^{\gamma_1}|x'|^{\alpha}u_j||_{L^s(R_j)} \simeq I_0.$$
 (2.9)

Another computation gives

$$|||x|^{\gamma_2}|x'|^{\mu}\nabla u_j||_{L^p(R_i)} \lesssim 2^{(b_1\kappa + d_1 - b_2\kappa - d_2)j}A_0,$$
 (2.10)

where $b_2 = n/p + \gamma_2 + \mu - 1$ and $d_2 = (n-1)/p + \mu - 1$. Since we are in the case when $1/p + (\mu - 1)/(n - 1) \neq 1/q + \beta/(n - 1)$ and $1/p + (\gamma_2 + \mu - 1)/n \neq 1/q + (\gamma_3 + \beta)/n$, we have, using (1.10) and (2.5), that $b_1 \neq b_2$ and $d_1 \neq d_2$. Now we fix

$$\kappa := \frac{d_2 - d_1}{b_1 - b_2} \in \mathbb{R} \setminus \{0\},\,$$

so that

$$|||x|^{\gamma_2}|x'|^{\mu}\nabla u_j||_{L^p(R_j)} \lesssim A_0.$$
 (2.11)

Using (1.10), (2.5), and the definition of κ , we have

$$|||x|^{\gamma_3}|x'|^{\beta}u_j||_{L^q(R_i)} \simeq B_0.$$
 (2.12)

For any positive integer $m, w := \sum_{j=1}^m u_j \in C_c^{\infty}(\mathbb{R}^n)$. Since $(\sup u_j) \cap (\sup u_i) = \phi$ for $i \neq j$, we have, by (2.9), (2.11) and (2.12), that

$$|||x|^{\gamma_1}|x'|^{\alpha}w||_{L^s(\mathbb{R}^n)}^s \simeq mI_0^s, \quad |||x|^{\gamma_2}|x'|^{\mu}\nabla w||_{L^p(\mathbb{R}^n)}^p \lesssim mA_0^p, \quad |||x|^{\gamma_3}|x'|^{\beta}w||_{L^q(\mathbb{R}^n)}^q \simeq mB_0^q.$$

If w satisfies (1.15), then we have, by the above, that

$$m^{1/s} \le C \frac{A_0^a B_0^{1-a}}{I_0} m^{a/p + (1-a)/q}.$$

Since $I_0, A_0, B_0 > 0$ and m can be arbitrarily large, we have $1/s \le a/p + (1-a)/q$. (1.14) is proved.

Proof of the necessity part of Theorem 1.2: Let $n \geq 1$, $s, p, q, a, \gamma_1, \gamma_2$ and γ_3 satisfy (1.7) and (1.2). We show that if (1.17) holds for all $u \in C_c^{\infty}(\mathbb{R})$, then (1.3)-(1.5) hold. This is the same as in [10] when $q \geq 1$, while the proof there applies to q > 0 as well.

Since the formulation of our conditions is somewhat different from that in [10], we present a proof of the necessity of (1.3)-(1.5) using similar arguments as in the proof of Lemma 2.1. Condition (1.3) follows from a dimensional analysis argument as in the proof of (1.10) with $\alpha = \mu = \beta = 0$. Set $\alpha = \mu = \beta = 0$ and $x_0 = (0, ..., R)$ in the proof of (1.11), the same arguments give (1.4).

To prove (1.5), let $u = f_2(|x|)$ where f_2 is given by (2.1) with $\alpha = 0$, and insert u into (1.17). When 0 < a < 1, we have $1/s + \gamma_1/n = 1/p + (\gamma_2 - 1)/n = 1/q + \gamma_3/n$. Similar to (2.2)-(2.4), we have

$$|||x|^{\gamma_1}u||_{L^s(\mathbb{R})} \ge C\epsilon^{-1/s}, \quad |||x|^{\gamma_2}u'||_{L^p(\mathbb{R})} \le C\epsilon^{-1/p}, \quad |||x|^{\gamma_3}u||_{L^q(\mathbb{R})} \le C\epsilon^{-1/q}.$$

Using (1.17) and the above, we have $e^{-1/s} \leq Ce^{-a/p-(1-a)/q}$ for arbitrarily small e, thus the inequality in (1.5) follows. In view of (1.3), we have $1/s + \gamma_1/n = 1/q + \gamma_3/n$ when a = 0, and $1/s + \gamma_1/n = 1/p + (\gamma_2 - 1)/n$ when a = 1. The inequality in (1.5) when a = 0 or a = 1 follows from the same proof for 0 < a < 1.

3 A nonlinear Poincaré inequality

In this section, we give the proof of Theorem 1.3.

Proof of Theorem 1.3: We divide the proof into three steps.

Step 1. We prove (1.18) under the hypotheses of the theorem.

For $\lambda = 1$, Theorem 1.3 is a generalized Poincare inequality (see e.g. Lemma 1.1.11 of [23]). In the rest of Step 1 we assume $\lambda \neq 1$.

Since $C^1(\bar{\Omega})$ is dense in $W^{1,p}(\Omega)$, we may assume without loss of generality that $w \in C^1(\bar{\Omega})$ and w > 0 in $\bar{\Omega}$. Let $u := w^{1/\lambda}$, then inequality (1.18) takes an equivalent formulation: for all $u \in C^1(\bar{\Omega})$ and u > 0 in $\bar{\Omega}$,

$$||v||_{L^p(\Omega)} \le C||\nabla v||_{L^p(\Omega)}$$
, where $v := u^{\lambda} - (\bar{u})^{\lambda}$ and $\bar{u} := \oint_S u$.

We prove (1.18) by contradiction argument. Suppose the contrary, there exists a sequence of positive functions $\{u_i\} \in C^1(\overline{\Omega})$ such that

$$v_j := (u_j)^{\lambda} - (\bar{u}_j)^{\lambda} \tag{3.1}$$

satisfies

$$1 = ||v_j||_{L^p(\Omega)} > j||\nabla v_j||_{L^p(\Omega)}, \tag{3.2}$$

where $\bar{u}_i := \int_S u_i > 0$.

By (3.2) and the compact embedding of $W^{1,p}(\Omega)$ to $L^p(\Omega)$, there exists some $v \in W^{1,p}(\Omega)$ such that, after passing to a subsequence (still denoted by $\{v_j\}$), $v_j \to v$ in $W^{1,p}(\Omega)$, $v_j \to v$ in $L^p(\Omega)$ and q.e. in Ω , $\|\nabla v\|_{L^p(\Omega)} = 0$, and $\|v\|_{L^p(\Omega)} = 1$. Now we have, using (3.2), that

$$||v_j - v||_{W^{1,p}(\Omega)} \to 0.$$
 (3.3)

We divide into two cases, $\lambda > 1$ and $0 < \lambda < 1$.

Case 1. $\lambda > 1$.

In this case the function $s \to s^{\lambda}$ is convex, and therefore

$$\bar{v}_j \ge \left(\oint_S u_j \right)^{\lambda} - (\bar{u}_j)^{\lambda} = 0.$$

Thus, by (3.3), we have $\bar{v}_j = \int_S v_j \to v > 0$.

Passing to another subsequence if necessary, we either have $\bar{u}_j \to \alpha \in [0, \infty)$ or $\bar{u}_j \to \infty$.

If $\bar{u}_i \to \alpha \in [0, \infty)$, we have

$$u_i \to (v + \alpha^{\lambda})^{1/\lambda}$$
 a.e. in Ω .

By Fatou's lemma,

$$|S|(v+\alpha^{\lambda})^{1/\lambda} = \int_{S} \liminf_{j \to \infty} u_j \le \liminf_{j \to \infty} \int_{S} u_j = \alpha |S|.$$

A contradiction, since v > 0 and $\alpha \ge 0$. So inequality (1.18) holds.

In the rest of Case 1, we assume $\bar{u}_i \to \infty$.

Denote $a_j := \bar{u}_j \to \infty$, and write

$$0 \le u_j = a_j + \eta_j. \tag{3.4}$$

Then

$$\int_{S} \eta_{j} = 0, \quad \forall j, \tag{3.5}$$

and, by (3.1),

$$v_j = (a_j + \eta_j)^{\lambda} - (a_j)^{\lambda}.$$

We will show that this leads to a contradiction.

Write $v_j^+(\theta) = \max\{v_j(\theta), 0\}, \ v_j^-(\theta) = \max\{-v_j(\theta), 0\}, \ \theta \in \overline{\Omega}$. Then $v_j = v_j^+ - v_j^-$. By (3.3) and the positivity of v, we have

$$\|v_i^+ - v\|_{L^1(\Omega)} \to 0, \quad \|v_i^-\|_{L^1(\Omega)} \to 0.$$
 (3.6)

Lemma 3.1.

$$(a_j)^{\lambda-1} \int_{\Omega} \eta_j^- \to 0$$
, and $(a_j)^{\lambda-1} \int_{S} |\eta_j| \to 0$.

Proof. Write

$$v_j^- = (a_j)^{\lambda} - (a_j - \eta_j^-)^{\lambda} = (a_j)^{\lambda} \left(1 - \left(1 - \frac{\eta_j}{a_j}\right)^{\lambda}\right),$$

and recall from (3.4) that $0 \le w^- \le a_j$. Since $\lambda \ge 1$, we have the following elementary inequality:

$$g(t) := 1 - (1 - t)^{\lambda} - t \ge 0, \quad \forall \ 0 \le t \le 1.$$

Indeed, the above inequality holds due to the concavity of g in [0,1] ($g''(t) = -(\lambda - 1)(1-t)^{\lambda-2} < 0$ for all 0 < t < 1) and the fact that g(0) = g(1) = 0.

Now we have, using (3.6) and the above, that

$$\circ(1) = \int_{\Omega} v_j^- = (a_j)^{\lambda} \left(1 - \left(1 - \frac{\eta_j^-}{a_j} \right)^{\lambda} \right) \ge (a_j)^{\lambda} \int_{\Omega} \frac{\eta_j^-}{a_j} = (a_j)^{\lambda - 1} \int_{\Omega} \eta_j^-.$$

Lemma 3.1 follows from the above and (3.5).

Lemma 3.2. There exists some positive constant C independent of j such that

$$\int_{S} (\eta_{j}^{+})^{\lambda} \geq \frac{1}{C}, \quad \forall \ j.$$

Proof. We will use the following elementary inequality: for $\lambda \geq 1$, there exists some positive constant C, depending only on λ , such that

$$(1+t)^{\lambda} - 1 \le C(t^{\lambda} + t), \quad \forall \ t \ge 0.$$

With this constant C, we have, using (3.6), that

$$v|S| + o(1) = \int_{S} v_{j}^{+} = (a_{j})^{\lambda} \int_{S} \left(\left(1 + \frac{\eta_{j}^{+}}{a_{j}} \right)^{\lambda} - 1 \right)$$

$$\leq C(a_{j})^{\lambda} \int_{S} \left(\left(\frac{\eta_{j}^{+}}{a_{j}} \right)^{\lambda} + \frac{\eta_{j}^{+}}{a_{j}} \right) = C \int_{S} \left((\eta_{j}^{+})^{\lambda} + (a_{j})^{\lambda - 1} \eta_{j}^{+} \right).$$

Lemma 3.2 follows from the above in view of Lemma 3.1.

Lemma 3.3. For every $\epsilon > 0$, $(a_j)^{\lambda-1} |\{\eta_j > \epsilon\}| \to 0$.

Proof. Since

$$v_j \ge (a_j + \epsilon)^{\lambda} - (a_j)^{\lambda} > 0 \quad \text{on } \{\eta_j > \epsilon\},$$

we have, using (3.6), that

$$v|\Omega| + o(1) = \int_{\Omega} v_j^+ \ge \int_{\{\eta_j > \epsilon\}} \left((a_j + \epsilon)^{\lambda} - (a_j)^{\lambda} \right) = \left((a_j + \epsilon)^{\lambda} - (a_j)^{\lambda} \right) |\{\eta_j > \epsilon\}|.$$

Lemma 3.3 follows from the above since $\lambda > 1$ and $a_i \to \infty$.

Lemma 3.4. For every $\epsilon > 0$, $\int_{\Omega} [(\eta_j - \epsilon)^+]^{\lambda} \to 0$ as $j \to \infty$.

Proof. For $\epsilon > 0$, denote $\xi_j := [(\eta_j - \epsilon)^+]^{\lambda}$. By Lemma 3.3, $|\{\xi_j = 0\}| \to |\Omega| > 0$ as $j \to \infty$. Apply a generalized Poincaré inequality (see e.g. Lemma 7.16 and Lemma 7.12 in [14]—writing Ω as the union of finitely many convex open sets and apply these lemmas on each of the convex open sets) and use (3.2) and the fact that $\lambda \geq 1$, we have

$$\int_{\Omega} \xi_j \le C \int_{\Omega} |\nabla \xi_j| \le C \int_{\{\eta_j > \epsilon\}} \left[(\eta_j - \epsilon)^+ \right]^{\lambda - 1} |\nabla \eta_j^+|$$

$$\le C \int_{\{\eta_j > \epsilon\}} (\eta_j^+)^{\lambda - 1} |\nabla \eta_j^+| \le C \int_{\{\eta_j > \epsilon\}} |\nabla v_j| \to 0.$$

Lemma 3.4 is established.

For every $\epsilon > 0$, write $\eta_j = (\eta_j - \epsilon) + \epsilon \le (\eta_j - \epsilon)^+ + \epsilon$. Thus

$$(\eta_j^+)^{\lambda} \le 2^{\lambda} \left[(\eta_j - \epsilon)^+ \right]^{\lambda} + 2^{\lambda} \epsilon^{\lambda}.$$

It follows, using Lemma 3.2 and Lemma 3.4, that

$$0 < \frac{1}{C} \le \int_{S} (\eta_{j}^{+})^{\lambda} \le C \int_{S} \left[(\eta_{j} - \epsilon)^{+} \right]^{\lambda} + 2^{\lambda} \epsilon^{\lambda} |S| \le o(1) + C \epsilon^{\lambda}.$$

Sending j to ∞ , we have from the above that $0 < 1/C \le C\epsilon^{\lambda}$. Sending ϵ to 0, we have $0 < 1/C \le 0$, a contradiction. Estimate (1.18) is established in Case 1.

Case 2. $0 < \lambda < 1$.

Recall that $p \ge n/(1+n\lambda)$. Since $0 < \lambda < 1$, the function $s \to s^{\lambda}$ is concave, and we have

$$\bar{v}_j \le \left(\oint_S u_j \right)^{\lambda} - (\bar{u}_j)^{\lambda} = 0.$$

Thus, by (3.3), we have

$$\bar{v}_j = \oint_S v_j \to v < 0. \tag{3.7}$$

Fix a $\delta > 0$ satisfying $1 + \delta \leq \min\{2, 1/\lambda\}$. We will make use of the following elementary fact: For $0 < \lambda < 1$, there exists some positive constant C, depending only on λ and δ , such that

$$\left|(1+t)^{1/\lambda} - 1 - \frac{1}{\lambda}t\right| \le C(|t|^{1+\delta} + |t|^{1/\lambda}), \quad \forall -1 \le t < \infty.$$

By (3.1),

$$u_j = \left(v_j + (\bar{u}_j)^{\lambda}\right)^{1/\lambda}.\tag{3.8}$$

Integrating the above over S gives, with C given by the one in (3), that

$$0 = \frac{1}{|S|} \int_{S} \left(\left[v_j + (\bar{u}_j)^{\lambda} \right]^{1/\lambda} - \bar{u}_j \right) = (\bar{u}_j) \frac{1}{|S|} \int_{S} \left(\left[1 + (\bar{u}_j)^{-\lambda} v_j \right]^{1/\lambda} - 1 \right)$$

$$\leq \frac{1}{\lambda} (\bar{u}_j)^{1-\lambda} \bar{v}_j + \frac{C\bar{u}_j}{\lambda |S|} \int_{S} \left(\left| (\bar{u}_j)^{-\lambda} v_j \right|^{1+\delta} + \left| (\bar{u}_j)^{-\lambda} v_j \right|^{1/\lambda} \right).$$

$$(3.9)$$

Since $1 + \delta \leq 1/\lambda$, $W^{1,p}(\Omega)$ embeds into $L^{1/\lambda}(\Omega)$ and $L^{1+\delta}(\Omega)$ by the assumption on p. By this and (3.3), we have

$$||v_j - v||_{L^{1+\delta}(\Omega)} \le C||v_j - v||_{L^{1/\lambda}(\Omega)} \le C||v_j - v||_{W^{1,p}(\Omega)} \to 0.$$
 (3.10)

We deduce from (3.9), using (3.7) and (3.10), that

$$|v| + \circ(1) = -\bar{v}_j \le C\left((\bar{u}_j)^{-\delta\lambda} \int_{\Omega} |v_j|^{1+\delta} + (\bar{u}_j)^{\lambda-1} \int_{\Omega} |v_j|^{1/\lambda}\right) \le C\left((\bar{u}_j)^{-\delta\lambda} + (\bar{u}_j)^{\lambda-1}\right).$$

Since $v \neq 0$, we have the boundedness of $\{\bar{u}_j\}$. Passing to a subsequence, $\bar{u}_j \to \alpha$ for some $\alpha \in [0, \infty)$. Integrating (3.8) over S and using (3.10) and $\bar{u}_j \to \alpha$, we have

$$\alpha + \circ(1) = \oint_S \left(v_j + (\bar{u}_j)^{\lambda} \right)^{1/\lambda} = \left(v + \alpha^{\lambda} \right)^{1/\lambda} + \circ(1).$$

It follows that $\alpha = (v + \alpha^{\lambda})^{1/\lambda}$ which implies that v = 0. A contradiction. Estimate (1.18) is established in Case 2. Step 1 is completed.

Step 2. Inequality (1.18) does not hold if $0 < \lambda < 1$ and 0 .

For simplicity, we let $\Omega \subset \mathbb{R}^n$ be a bounded open set, and $S \subset \Omega$ has positive Lebesgue measure. Take a Lebesgue point \bar{x} of S, i.e. $\lim_{r\to 0^+} |B_r(\bar{x})\cap S|/|B_r(\bar{x})|=1$. For convenience, $\bar{x}=0$ is the Lebesgue point. For small $\epsilon>0$ and large $\alpha>1$, let

$$v(x) = \begin{cases} -1, & \text{if } |x| \ge \epsilon, \\ -1 + \alpha \left(1 - \frac{|x|}{\epsilon}\right), & \text{if } |x| \le \epsilon. \end{cases}$$

In the following, C denotes some positive constant independent of α and ϵ . A calculation gives

$$\int_{\Omega} |v|^p = |\Omega \setminus B_{\epsilon}| + \int_{B_{\epsilon}} |v|^p \ge |\Omega| + \frac{1}{C} \alpha^p \epsilon^n,$$

$$\int_{\Omega} |\nabla v|^p = \alpha^p \epsilon^{n-p} |B_1|,$$

$$\int_{S} (v+1)^{1/\lambda} = \alpha^{1/\lambda} \epsilon^n \int_{\{|y| \le 1, \epsilon y \in S\}} (1-|y|)^{1/\lambda} dy,$$

where $|O(\alpha^p \epsilon^n)| \leq C \alpha^p \epsilon^n$.

Since 0 is a Lebesgue point of S, we have

$$\lim_{\epsilon \to 0^+} \frac{|\{|y| \le 1, \ \epsilon y \in S\}|}{\{|y| \le 1\}} = \lim_{\epsilon \to 0^+} \frac{|B_{\epsilon}(0) \cap S|}{|B_{\epsilon}(0)|} = 1.$$

It follows that

$$\lim_{\epsilon \to 0^+} \int_{\{|y| < 1, \ \epsilon y \in S\}} (1 - |y|)^{1/\lambda} dy = \int_{|y| < 1} (1 - |y|)^{1/\lambda} dy > 0.$$

Now we fix the value of α so that $\int_S (v+1)^{1/\lambda} = |S|$. So $\alpha \leq C\epsilon^{-n\lambda}$. Consider

$$u := (v+1)^{1/\lambda}.$$

Then $\bar{u} = \int_S u = 1$, $u \ge 0$, $v = u^{\lambda} - \bar{u}^{\lambda}$. Using $p < n/(1 + n\lambda)$,

$$\int_{\Omega} |\nabla v|^p \le C\alpha^p \epsilon^{n-p} \le C\epsilon^{n-(1+n\lambda)p} \to 0.$$

This and (3.9) violate (1.18) for any choice of C. Step 2 is completed.

Step 3. Inequality (1.18) does not hold if $0 < \lambda < \infty$ and 0 .

For simplicity, we take $\Omega = S = [-1, 1]^n$. For $\alpha > 0$ small, let

$$f(x_1) := \begin{cases} |x_1|^{\alpha}, & x_1 < 0, \\ -|x_1|^{\alpha}, & x_1 \ge 0, \end{cases}$$

and

$$w(x) := (2 + f(x_1))^{\lambda}.$$

Then $w \in W^{1,p}([-1,1]^n)$ and $w \ge 1$.

By the definition of w, we have $f_{[-1,1]^n} w^{1/\lambda} = 2$. Let $v = w - (f_{[-1,1]^n} w^{1/\lambda})^{\lambda}$. We have, for some constant C > 0 depending only on λ and p, that

$$\int_{[-1,1]^n} |v(x)|^p dx \ge \int_{[1/2,1]^n} |v(x)|^p dx = \int_{[1/2,1]^n} |(2+|x_1|^\alpha)^\lambda - 2^\lambda|^p dx \ge \frac{1}{C}.$$
 (3.11)

On the other hand, by the assumption that 0 , we have

$$\int_{[-1,1]^n} |\nabla v|^p dx = \int_{[-1,1]^n} |\lambda (2 + f(x_1))^{\lambda - 1} f'(x_1)|^p dx$$

$$\leq C \int_{-1}^1 \alpha^p |x_1|^{(\alpha - 1)p} dx_1 \leq C \alpha^p \to 0$$

as $\alpha \to 0$. This and (3.11) violate (1.18). Step 3 is completed. Theorem 1.3 is proved.

Proof of Corollary 1.1: If $q \leq 1$, then, by Theorem 1.3 with $\lambda = 1/q$, we have

$$||w||_{L^{p}(\Omega)} \leq \left(\int_{S} w^{q}\right)^{1/q} \cdot |\Omega|^{1/p} + ||w - \left(\int_{S} w^{q}\right)^{1/q}||_{L^{p}(\Omega)} \leq \left(\int_{S} w^{q}\right)^{1/q} |\Omega|^{1/p} + C||\nabla w||_{L^{p}(\Omega)}.$$

If q > 1, then (1.19) follows from the result for q = 1 and Hölder's inequality. The corollary is proved.

4 Extension of the Caffarelli-Kohn-Nirenberg inequalities from $q \ge 1$ to q > 0

In this section, we prove Theorem 1.2. The necessity part has been established in Section 2. The sufficiency part follows from the following theorem which includes the inequalities on cones.

Let $\mathbb{S}^{n-1} := \{x \in \mathbb{R}^n \mid |x| = 1\}$ be the unit sphere in \mathbb{R}^n . For any $\Omega \subset \mathbb{S}^{n-1}$ with nonempty Lipschitz boundary, denote the cone

$$K := \{ rx \mid r \ge 0, \ x \in \Omega \}. \tag{4.1}$$

Theorem 4.1. Let $n \geq 1$, $K = \mathbb{R}^n$ or K be as above, and $s, p, q, \gamma_1, \gamma_2, \gamma_3, a$ satisfy (1.7) and (1.2)-(1.5). Then there exists some positive constant C such that for all $u \in C_c^{0,1}(\overline{K})$

$$|||x|^{\gamma_1}u||_{L^s(K)} \le C|||x|^{\gamma_2}\nabla u||_{L^p(K)}^a |||x|^{\gamma_3}u||_{L^q(K)}^{1-a}. \tag{4.2}$$

Furthermore, on any compact set in the parameter space in which (1.7) and (1.2) hold, the constant C is bounded.

Lemma 4.1. Let $n \geq 1$, $0 < r_1 < r_2 < \infty$, $K = \mathbb{R}^n$ or K be given by (4.1), $s, p, q, a, \gamma_1, \gamma_2$ and γ_3 satisfy (1.7), (1.3), (1.4), $1/s + \gamma_1/n > 0$, and $1/s \leq a/p + (1-a)/q$. Then there exists some positive constant C, depending only on $s, q, a, \gamma_1, \gamma_2, \gamma_3, r_1, r_2$ and Ω , such that for all $u \in C^{0,1}(K \cap B_{r_2})$,

$$|||x|^{\gamma_1}u||_{L^s(K\cap B_{r_1})} \le C|||x|^{\gamma_1}u||_{L^s(K\cap B_{r_2}\setminus B_{r_1})} + C|||x|^{\gamma_2}\nabla u||_{L^p(K\cap B_{r_2})}^a|||x|^{\gamma_3}u||_{L^q(K\cap B_{r_1})}^{1-a}.$$

$$(4.3)$$

Proof. For simplicity, we only prove (4.3) for $r_1 = 1$ and $r_2 = 2$. The general case can be proved similarly.

For a = 0, we deduce from (1.3), (1.4) and $1/s \le a/p + (1-a)/q$ that $\gamma_1 = \gamma_3$ and s = q, thus (4.3) is obvious. In the rest of the proof we assume $0 < a \le 1$. Without loss of generality, assume $u \ge 0$.

Step 1. We prove (4.3) for p = 1 and $\gamma_1 = 0$.

Let

$$R_k := \{ x \in K \mid 2^{k-1} \le |x| \le 2^k \}, \quad k \in \mathbb{Z}.$$

Denote

$$A_k := \int_{R_k} |u|^s dx, \quad M_k := \int_{R_k} ||x|^{\gamma_2} \nabla u(x)| dx, \quad N_k := \int_{R_k} ||x|^{\gamma_3} u|^q dx.$$

We first establish for any $0 < \epsilon < 2^{an} - 1$ that

$$A_k \le \theta A_{k+1} + C(M_k + M_{k+1})^{as} N_k^{(1-a)s/q}, \quad k \in \mathbb{Z},$$
 (4.4)

where

$$\theta := \frac{a(1+\epsilon)}{2^{an} - (1+\epsilon)(1-a)},$$

and C depends only on $s, q, a, \gamma_1, \gamma_2, \gamma_3, r_1, r_2, K$ and ϵ . Since K is a cone, by (1.3) and scaling, we only need to prove (4.4) for k = 0.

Let $\bar{u} = \int_{R_1} u(y) dy$. For any $0 < \epsilon < 2^{an} - 1$, $x \in R_0$ and $\xi \in R_1$, we have

$$|u(x)|^{s} = |u(x)|^{(1-a)s}|u(x)|^{as}$$

$$\leq |u(x)|^{(1-a)s}(|u(x) - \bar{u}| + |\bar{u} - u(\xi)| + |u(\xi)|)^{as}$$

$$\leq (1+\epsilon)|u(x)|^{(1-a)s}|u(\xi)|^{as} + C|u(x)|^{(1-a)s}(|u(x) - \bar{u}| + |\bar{u} - u(\xi)|)^{as}.$$
(4.5)

Taking $\int_{R_1} \int_{R_0} dx d\xi$ of the above and using Hölder's inequality, we have

$$A_{0} \leq (1+\epsilon) \oint_{R_{1}} |u(\xi)|^{as} d\xi \int_{R_{0}} |u(x)|^{(1-a)s} dx + C \int_{R_{0}} |u(x)|^{(1-a)s} |u(x) - \bar{u}|^{as} dx$$

$$+ C \int_{R_{0}} |u(x)|^{(1-a)s} dx \oint_{R_{1}} |\bar{u} - u(\xi)|^{as} d\xi$$

$$\leq (1+\epsilon) \frac{|R_{0}|^{a}}{|R_{1}|^{a}} \left(\int_{R_{1}} |u(\xi)|^{s} d\xi \right)^{a} \left(\int_{R_{0}} |u(x)|^{s} dx \right)^{1-a} + C \int_{R_{0}} |u(x)|^{(1-a)s} |u(x) - \bar{u}|^{as} dx$$

$$+ C \int_{R_{0}} |u(x)|^{(1-a)s} dx \oint_{R_{1}} |\bar{u} - u(\xi)|^{as} d\xi$$

$$=: (1+\epsilon) \frac{|R_{0}|^{a}}{|R_{1}|^{a}} A_{0}^{1-a} A_{1}^{a} + C(I_{1} + I_{2}).$$

$$(4.6)$$

Since p=1, by (1.3) and (1.4), we have $1/s \ge a(1-1/n)+(1-a)/q$. Since we are in the case $1/s \le a/p+(1-a)/q$, we have $a(1-1/n)+(1-a)/q \le 1/s \le a+(1-a)/q$. Thus there exist some $1 \le t \le n/(n-1)$ ($1 \le t \le \infty$ when n=1) such that 1/s = a/t + (1-a)/q. Then by Hölder's inequality, Sobolev inequality and Poincaré's inequality, we have

$$I_{1} \leq C \|u - \bar{u}\|_{L^{t}(R_{0} \cup R_{1})}^{as} \|u\|_{L^{q}(R_{0})}^{(1-a)s}$$

$$\leq C (\|u - \bar{u}\|_{L^{1}(R_{0} \cup R_{1})} + \|\nabla(u - \bar{u})\|_{L^{1}(R_{0} \cup R_{1})})^{as} N_{0}^{(1-a)s/q}$$

$$\leq C (M_{0} + M_{1})^{as} N_{0}^{(1-a)s/q}. \tag{4.7}$$

Similarly, we have

$$I_{2} \leq C \|u - \bar{u}\|_{L^{t}(R_{1})}^{as} \|u\|_{L^{q}(R_{0})}^{(1-a)s}$$

$$\leq C (\|u - \bar{u}\|_{L^{1}(R_{1})} + \|\nabla(u - \bar{u})\|_{L^{1}(R_{1})})^{as} N_{0}^{(1-a)s/q}$$

$$\leq C M_{1}^{as} N_{0}^{(1-a)s/q}.$$

$$(4.8)$$

By (4.6), (4.7), (4.8) and the fact that $|R_0|/|R_1| = 2^{-n}$, we have, for any $0 < \epsilon < 2^{an} - 1$, that

$$A_0 \le (1+\epsilon) \frac{|R_0|^a}{|R_1|^a} A_0^{1-a} A_1^a + C(M_0 + M_1)^{as} N_0^{(1-a)s/q}$$

$$\le (1+\epsilon) 2^{-an} ((1-a)A_0 + aA_1) + C(M_0 + M_1)^{as} N_0^{(1-a)s/q}$$

Thus

$$A_0 \le \frac{a(1+\epsilon)}{2^{an} - (1+\epsilon)(1-a)} A_1 + C(M_0 + M_1)^{as} N_0^{(1-a)s/q}.$$

So (4.4) holds for k=0, and therefore holds for all $k\in\mathbb{Z}$.

Since a > 0, $2^{an} > 1$, and $0 < \epsilon < 2^{an} - 1$, we have $0 < \theta < 1$.

For $c, d \ge 0$, $c + d \ge 1$, and sequences $x_n, y_n \ge 0$, $n \ge 1$, we have

$$\sum_{n=1}^{\infty} x_n^c y_n^d \le \left(\sum_{n=1}^{\infty} x_n\right)^c \left(\sum_{n=1}^{\infty} y_n\right)^d. \tag{4.9}$$

Take the sum of (4.4) over $k \le 0$, by the fact $1/s \le a + (1-a)/q$ and (4.9) with c = (1-a)s/q and d = as, we have, that

$$\sum_{k \le 0} A_k \le \theta A_1 + \theta \sum_{k \le 0} A_k + C \sum_{k \le 0} (M_k + M_{k+1})^{as} N_k^{(1-a)s/q}$$

$$\le \theta A_1 + \theta \sum_{k \le 0} A_k + C \left(\sum_{k \le 0} (M_k + M_{k+1}) \right)^{as} \left(\sum_{k \le 0} N_k \right)^{(1-a)s/q}.$$

So

$$\int_{K \cap B_{1}} |u|^{s} dx = \sum_{k \leq 0} A_{k} \leq \frac{\theta}{1 - \theta} A_{1} + C \left(\sum_{k \leq 0} (M_{k} + M_{k+1}) \right)^{as} \left(\sum_{k \leq 0} N_{k} \right)^{(1 - a)s/q} \\
\leq \frac{\theta}{1 - \theta} \int_{R_{1}} |u|^{s} dx + C \left(\int_{K \cap B_{2}} ||x|^{\gamma_{2}} \nabla u| dx \right)^{as} \left(\int_{K \cap B_{1}} ||x|^{\gamma_{3}} u|^{q} dx \right)^{(1 - a)s/q}.$$
(4.10)

Thus when p = 1 and $\gamma_1 = 0$, (4.3) follows from (4.10).

Step 2. We prove (4.3) for p = 1 and $\gamma_1 \neq 0$.

We will reduce it to Step 1. Make a change of variables $y = |x|^{\gamma_1 s/n} x$, and define $\tilde{u}(y) := u(x)$, $\tilde{\gamma}_1 = 0$, $\tilde{\gamma}_2 = (\gamma_2 n + \gamma_1 s(1-n))/(\gamma_1 s + n)$ and $\tilde{\gamma}_3 = (\gamma_3 q - \gamma_1 s)n/(\gamma_1 s + n)q$.

We have s, q > 0 from (1.7) and $1/s + \tilde{\gamma}_1/n = 1/s > 0$. By computation and using (1.3),

$$a\left(1 + \frac{\tilde{\gamma}_2 - 1}{n}\right) + (1 - a)\left(\frac{1}{q} + \frac{\tilde{\gamma}_3}{n}\right) = \frac{n}{\gamma_1 s + n}\left(a\left(1 + \frac{\gamma_2 - 1}{n}\right) + (1 - a)\left(\frac{1}{q} + \frac{\gamma_3}{n}\right)\right) = \frac{1}{s}.$$
(4.11)

Next, by (1.3) and (1.4) with p = 1, we have $1/s \ge a(1 - 1/n) - (1 - a)/q$. Use this and (4.11), we have

$$a\tilde{\gamma}_2 + (1-a)\tilde{\gamma}_3 = n\left(\frac{1}{s} - a\left(1 - \frac{1}{n}\right) - \frac{1-a}{a}\right) \ge 0.$$

So we have verified (1.7), (1.3), (1.4), and $1/s + \tilde{\gamma}_1/n > 0$ with $\tilde{\gamma}_1 = 0$. By this and the fact that $1/s \leq a/p + (1-a)/q$, apply Step 1 to $\tilde{u}(y)$ and $\tilde{\gamma}_1, \tilde{\gamma}_3, \tilde{\gamma}_3$, we have

$$\|\tilde{u}\|_{L^{s}(K\cap B_{1})} \leq C\|\tilde{u}\|_{L^{s}(K\cap B_{R}\setminus B_{1})} + C\||y|^{\tilde{\gamma}_{2}}\nabla \tilde{u}\|_{L^{1}(K\cap B_{R})}^{a}\||y|^{\tilde{\gamma}_{3}}\tilde{u}\|_{L^{q}(K\cap B_{1})}^{1-a}.$$
(4.12)

Since $\gamma_1 s/n + 1 > 0$, we have, with $R := 2^{\gamma_1 s/n + 1} > 1$, that

$$\int_{K \cap B_1} ||x|^{\gamma_1} u(x)|^s dx = \frac{n}{\gamma_1 s + n} \int_{K \cap B_1} |\tilde{u}(y)|^s dy,
\int_{K \cap B_2} ||x|^{\gamma_2} \nabla u(x)| dx = \int_{K \cap B_R} ||y|^{\tilde{\gamma}_2} \nabla \tilde{u}(y)| dy,
\int_{K \cap B_1} ||x|^{\gamma_3} u(x)|^q dx = \frac{n}{\gamma_1 s + n} \int_{K \cap B_1} ||y|^{\tilde{\gamma}_3} \tilde{u}(y)|^q dy.$$

By (4.12) and the above, we have (4.3).

Step 3. We prove (4.3) for p > 1.

Let $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\gamma}_1, \bar{\gamma}_2$ and $\bar{\gamma}_3$ be defined by

$$\frac{1}{\bar{s}} = \frac{1}{s} + \frac{1}{p'}, \quad \bar{p} = 1, \quad \frac{1}{\bar{q}} = \frac{s}{\bar{s}q}, \quad \bar{a} = \frac{as}{(1-a)\bar{s} + as},
\bar{\gamma}_1 = \frac{\gamma_1 s}{\bar{s}}, \quad \bar{\gamma}_2 = \frac{\gamma_1 s}{p'} + \gamma_2, \quad \bar{\gamma}_3 = \frac{\gamma_3 s}{\bar{s}},$$

where 1/p + 1/p' = 1.

It can be verified that $0 < \bar{s} < s$, and $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3$ satisfy (1.7), (1.3), (1.4), $1/\bar{s} + \bar{\gamma}_1/n > 0$, and $1/\bar{s} \le \bar{a}\bar{p} + (1-\bar{a})/\bar{q}$ (for detail of the verification, see Lemma 7.1). So we can apply Step 2 to $|u|^{s/\bar{s}}$ to obtain, using Hölder's inequality and Young's

inequality, that

$$\begin{aligned} & \||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{1})}^{s/\bar{s}} = \||x|^{\bar{\gamma}_{1}}|u|^{s/\bar{s}}\|_{L^{\bar{s}}(K\cap B_{1})} \\ & \leq C\||x|^{\bar{\gamma}_{1}}|u|^{s/\bar{s}}\|_{L^{\bar{s}}(K\cap B_{2}\setminus B_{1})} + C\||x|^{\bar{\gamma}_{2}}\nabla|u|^{s/\bar{s}}\|_{L^{1}(K\cap B_{2})}^{\bar{a}}\||x|^{\bar{\gamma}_{3}}|u|^{s/\bar{s}}\|_{L^{\bar{q}}(K\cap B_{1})}^{1-\bar{a}} \\ & \leq C\||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{2}\setminus B_{1})}^{s/\bar{s}} + C\||x|^{\bar{\gamma}_{2}}|u|^{s/\bar{s}-1}|\nabla u|\|_{L^{1}(K\cap B_{2})}^{\bar{a}}\||x|^{\gamma_{3}}u\|_{L^{q}(K\cap B_{1})}^{(1-\bar{a})q/\bar{q}} \\ & \leq C\||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{2}\setminus B_{1})}^{s/\bar{s}} + C\||x|^{\bar{\gamma}_{2}-\gamma_{2}}|u|^{s/\bar{s}-1}\|_{L^{p'}(K\cap B_{2})}^{\bar{a}}\||x|^{\gamma_{2}}\nabla u\|_{L^{p}(K\cap B_{2})}^{\bar{a}}\||x|^{\gamma_{3}}u\|_{L^{q}(K\cap B_{1})}^{(1-\bar{a})s/\bar{s}} \\ & \leq C\||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{2}\setminus B_{1})}^{s/\bar{s}} + C\||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{2})}^{\bar{a}s/p'}\||x|^{\gamma_{2}}\nabla u\|_{L^{p}(K\cap B_{2})}^{\bar{a}}\||x|^{\gamma_{3}}u\|_{L^{q}(K\cap B_{1})}^{(1-\bar{a})s/\bar{s}} \\ & \leq C\||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{2}\setminus B_{1})}^{s/\bar{s}} + \frac{1}{2}\||x|^{\gamma_{1}}u\|_{L^{s}(K\cap B_{1})}^{s/\bar{s}} + C\left(\||x|^{\gamma_{2}}\nabla u\|_{L^{p}(K\cap B_{2})}^{\bar{a}} \\ & \cdot \||x|^{\gamma_{3}}u\|_{L^{q}(K\cap B_{1})}^{(1-\bar{a})s/\bar{s}}\right)^{1/(1-\bar{a}\bar{s}/p')}. \end{aligned}$$

Inequality (4.2) follows from the above and the definitions of \bar{a} and \bar{s} . Lemma 4.1 is proved.

Proof of Theorem 4.1. Without loss of generality, we assume $u \geq 0$. By (1.3) and scaling, we may assume $\sup u \subset B_1$.

For a = 0, we deduce from (1.3), (1.4) and (1.5) that $\gamma_1 = \gamma_3$ and s = q, thus (4.2) is obvious. In the rest of the proof we assume $0 < a \le 1$.

Case 1.
$$1/s \le a/p + (1-a)/q$$
.

In this case, inequality (4.2) follows from Lemma 4.1 with $r_1 = 1$ and $r_2 = 2$.

Case 2.
$$1/s > a/p + (1-a)/q$$
.

Case 2 can be reduced to Case 1 by section (V) in [10] - this reduction is the same for q > 0 even though $q \ge 1$ was assumed in the paper. For reader's convenience, we include such an argument here.

By (1.3) and (1.5), $1/p + (\gamma_2 - 1)/n \neq 1/q + \gamma_3/n$. Thus there exist some positive constants λ_1 and λ_2 , such that $\hat{u}(x) = \lambda_1 u(\lambda_2 x)$ satisfies $|||x|^{\gamma_2} \nabla \hat{u}||_{L^p(K)} = 1$ and $|||x|^{\gamma_3} \hat{u}||_{L^q(K)} = 1$. We claim that there exist some $0 \leq a', a'' \leq 1$, such that

$$\begin{aligned} |||x|^{\gamma_{1}}\hat{u}||_{L^{s}(K)} &\leq C\left(|||x|^{\gamma_{2}}\nabla u||_{L^{p}(K)}^{a'}|||x|^{\gamma_{3}}u||_{L^{q}(K)}^{1-a'} + |||x|^{\gamma_{2}}\nabla u||_{L^{p}(K)}^{a''}|||x|^{\gamma_{3}}u||_{L^{q}(K)}^{1-a''}\right) \\ &= 2C|||x|^{\gamma_{2}}\nabla \hat{u}||_{L^{p}(K)}^{a}|||x|^{\gamma_{3}}\hat{u}||_{L^{q}(K)}^{1-a}. \end{aligned}$$

$$(4.13)$$

Then by scaling, we have that (4.2) holds for u.

To see (4.13) when $n \geq 2$, notice that by (1.7), (1.8)-(1.5), it can be directly verified that $s, p, q, a, \gamma_1, \gamma_2$ and γ_3 satisfy (1.7)-(1.14) with $\alpha = \mu = \beta = 0$. Then (4.13) follows from Lemma 5.3 with $\alpha = \mu = \beta = 0$. If n = 1, we can obtain (4.13) by the same proof as that of Lemma 5.3, where we set $\alpha = \mu = \beta = 0$ and choose $\alpha' = \alpha'' = 0$ there. Theorem 4.1 is proved.

Proof of Theorem 1.2. The necessity part has been proved in Section 2. The sufficiency part follows from Theorem 4.1 with $K = \mathbb{R}^n$.

5 Proof of the sufficiency part of Theorem 1.1

In this section, we prove the sufficiency part of Theorem 1.1.

We first prove the sufficiency part of Theorem 1.1 when $1/s \le a/p + (1-a)/q$. We make use of Theorem 1.2 (rather, its variants Theorem 4.1 and Lemma 4.1) and Theorem 1.3.

For $\delta, h > 0$, denote $B'_{\delta} = \{x' \in \mathbb{R}^{n-1} \mid |x'| \le \delta\}, \ D^h_{\delta} = B'_{\delta} \times [0, h] \text{ and } D_{\delta} = D^1_{\delta}.$

Lemma 5.1. Let $n \geq 2$, $0 < \delta_1 < \delta_2 < \infty$, h > 0, s, p, q, a, α, μ and β satisfy (1.7)-(1.14) with $\gamma_1 = \gamma_2 = \gamma_3 = 0$. Then there exists some positive constant C, depending only on $s, p, q, a, \alpha, \mu, \beta, \sigma$, δ_1, δ_2 and h, such that for all $u \in C^{0,1}(D^h_{\delta_2})$

$$|||x'|^{\alpha}u||_{L^{s}(D_{\delta_{1}}^{h})} \leq C|||x'|^{\alpha}u||_{L^{s}(D_{\delta_{2}}^{h}\setminus D_{\delta_{1}}^{h})} + C|||x'|^{\mu}\nabla u||_{L^{p}(D_{\delta_{2}}^{h})}^{a}|||x'|^{\beta}u||_{L^{q}(D_{\delta_{2}}^{h})}^{1-a}.$$
(5.1)

Proof. Since $\gamma_1 = \gamma_2 = \gamma_3 = 0$, we deduce from (1.10) and (1.13) that $1/s - a/p - (1-a)/q \ge (a(\mu-1) + (1-a)\beta - \alpha)/(n-1) = \frac{n}{n-1}(1/s - a/p - (1-a)/q)$, i.e.

$$\frac{1}{s} \le \frac{a}{p} + \frac{1-a}{q}.\tag{5.2}$$

Let $x = (r', \theta', x_n)$ be the cylindrical coordinates where r' = |x'| and $\theta' = x'/|x'|$. For simplicity, we only prove the lemma when h = 1, $\delta_1 = 1$ and $\delta_2 = 2$. The general case can be proved similarly.

If a = 0, by (1.10), (1.12) and (1.14), we have s = q and $\alpha = \beta$, and therefore (5.1) is obvious. In the rest of proof we assume $0 < a \le 1$.

Step 1. We prove inequality (5.1) when p=1.

By (5.2), we have, in view of p = 1 and $a \le 1$, that

$$\frac{1}{s} - \frac{1-a}{q} \le a \le 1. \tag{5.3}$$

Case 1. 1/s - (1-a)/q = 1.

By (5.3), we have a=1 and s=1. Because of this, (1.10), and the fact that s=p=1 and $\gamma_1=\gamma_2=\gamma_3=0$, we have $\alpha=\mu-1$. Let

$$\hat{s} = 1$$
, $\hat{p} = 1$, $\hat{q} = q$, $\hat{a} = 1$, $\hat{\gamma}_1 = \alpha$, $\hat{\gamma}_2 = \alpha + 1$, $\hat{\gamma}_3 = 0$

It is easy to verify that $\hat{s}, \hat{p}, \hat{q}, \hat{a}, \hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_3$ satisfy (1.7), (1.2)-(1.5) and $1/\hat{s} \leq \hat{a}/\hat{p} + (1-\hat{a})/\hat{q}$. Apply Lemma 4.1 to $u(\cdot, x_n)$ for each fixed $0 \leq x_n \leq 1$, with $K = \mathbb{R}^{n-1}$ and $s, p, q, a, \gamma_1, \gamma_2, \gamma_3$ replaced by $\hat{s}, \hat{p}, \hat{q}, \hat{a}, \hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_3$, we have, with notation $\nabla' = \nabla_{x'}$, that

$$\int_{B_1'} |x'|^{\alpha} |u(x',x_n)| dx' \le C \int_{B_2' \setminus B_1'} |x'|^{\alpha} |u(x',x_n)| dx' + C \int_{B_2'} |x'|^{\alpha+1} |\nabla' u(x',x_n)| dx'.$$

Integrate the above in x_n on [0,1], we have (5.1) in this case, i.e.

$$|||x'|^{\alpha}u||_{L^{1}(D_{1})} \leq C|||x'|^{\alpha}u||_{L^{1}(D_{2}\setminus D_{1})} + C|||x'|^{\alpha+1}\nabla' u||_{L^{1}(D_{2})}.$$

Case 2. 1/s - (1-a)/q < 1.

Let

$$b = \frac{1}{a} \left(\frac{1}{s} - \frac{1-a}{q} \right), \quad \lambda = \frac{a(1-b)}{1-ab}.$$
 (5.4)

Since a>0, b is well defined. In the definition of λ above, we have used the assumption that ab=1/s-(1-a)/q<1. By (5.2) with p=1, we have $b\leq 1$. By (1.10) and (1.12), we have $1/s-(a(1/p-1/n)+(1-a)/q)=(a(\gamma_2+\mu)+(1-a)(\gamma_3+\beta)-(\gamma_1+\alpha))/n\geq 0$. Thus when p=1, we have $b=(1/s-(1-a)/q)/a\geq (n-1)/n$. So $(n-1)/n\leq b\leq 1$. Consequently, we have $0\leq \lambda\leq 1$ in view of $0< a\leq 1$.

Let

$$\hat{a} = ab, \quad \hat{s} = s, \quad \hat{p} = 1, \quad \frac{1}{\hat{q}} = \lambda + \frac{1 - \lambda}{q},$$

$$\hat{\gamma}_1 = \alpha, \quad \hat{\gamma}_2 = \mu, \quad \hat{\gamma}_3 = \lambda \mu + (1 - \lambda)\beta.$$
(5.5)

We have shown that $0 < \hat{a} < 1$. Using (1.7)-(1.14) and the assumption that 1/s - (1-a)/q < 1, it can be verified that $\hat{s}, \hat{p}, \hat{q}, \hat{a}, \hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_3$ satisfy (1.7), (1.2)-(1.5) with $\hat{p} = 1$, $1/\hat{s} \le \hat{a}/\hat{p} + (1-\hat{a})/\hat{q}$ and with n replaced by n-1. For the details of the verification, see Lemma 7.2 and its proof.

Let $m = \min\{1, q, s\}$ and $1 < \delta < 2$ be some fixed number, set

$$v := u - \left(\int_{D_2 \setminus D_1} u^m \right)^{1/m}.$$

Apply Lemma 4.1 with p=1 to $v(\cdot,x_n)$ for each fixed $0 \le x_n \le 1$, with $r_1=1, r_2=2$, and $s, p, q, a, \gamma_1, \gamma_2, \gamma_3$ replaced by $\hat{s}, \hat{p}, \hat{q}, \hat{a}, \hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_3$, we have

$$|||x'|^{\alpha}v(\cdot,x_n)||_{L^s(B'_1)}$$

$$\leq C|||x'|^{\alpha}v(\cdot,x_n)||_{L^s(B'_2\setminus B'_1)} + C|||x'|^{\mu}\nabla'v(\cdot,x_n)||_{L^1(B'_2)}^{ab}|||x'|^{\hat{\gamma}_3}v(\cdot,x_n)||_{L^{\hat{q}}(B'_1)}^{1-ab}.$$

$$(5.6)$$

Using the definition of $\hat{\gamma}_3$, \hat{q} and λ in (5.4) and (5.5), and the fact that $0 \le \lambda \le 1$, we apply Hölder's inequality to estimate the last term in (5.6) as follows.

$$|||x'|^{\hat{\gamma}_{3}}v(\cdot,x_{n})||_{L^{\hat{q}}(B'_{1})}^{1-ab} = ||||x'|^{\mu}v(\cdot,x_{n})|^{\lambda} \cdot ||x'|^{\beta}v(\cdot,x_{n})|^{1-\lambda}||_{L^{\hat{q}}(B'_{1})}^{1-ab}$$

$$\leq |||x'|^{\mu}v(\cdot,x_{n})||_{L^{1}(B'_{1})}^{\lambda(1-ab)}|||x'|^{\beta}v(\cdot,x_{n})||_{L^{q}(B'_{1})}^{(1-\lambda)(1-ab)}$$

$$\leq |||x'|^{\mu}v(\cdot,x_{n})||_{L^{1}(B'_{1})}^{a(1-b)}|||x'|^{\beta}v(\cdot,x_{n})||_{L^{q}(B'_{1})}^{(1-a)}.$$

$$(5.7)$$

Next, we estimate the term $\int_{B_1'} |x'|^{\mu} |v| dx'$ in the above. Notice that

$$|v(x',x_n)| \le C \int_0^1 |v_{x_n}(x',t)| dt + C \int_0^1 |v(x',t)| dt, \quad \forall \ (x',x_n) \in D_2.$$

So, for each $x_n \in [0,1]$, we have

$$\int_{B_1'} |x'|^{\mu} |v(x', x_n)| dx' \le C \int_{B_1'} \int_0^1 |x'|^{\mu} |v_{x_n}(x', t)| dt dx' + C \int_{B_1'} \int_0^1 |x'|^{\mu} |v(x', t)| dt dx'.$$
(5.8)

Applying Lemma 4.1 in dimension n-1, we have, for every x_n in [0,1], that

$$\int_{B_1'} |x'|^{\mu} |v(x', x_n)| dx' \le C \int_{B_2' \setminus B_1'} |x'|^{\mu} |v(x', x_n)| dx' + C \int_{B_2'} |x'|^{\mu+1} |\nabla' v(x', x_n)| dx'.$$

Integrating the above in x_n over [0,1], and then inserting it into (5.8), we have

$$\int_{B_1'} |x'|^{\mu} |v(x', x_n)| dx' \le C(\||x'|^{\mu} v\|_{L^1(D_2 \setminus D_1)} + \||x'|^{\mu} \nabla v\|_{L^1(D_2)}). \tag{5.9}$$

Putting (5.6), (5.7) and (5.9) together, we have

$$\begin{aligned} |||x'|^{\alpha}v(\cdot,x_n)||_{L^s(B'_1)}^s &\leq C|||x'|^{\alpha}v(\cdot,x_n)||_{L^s(B'_2\setminus B'_1)}^s + C|||x'|^{\mu}\nabla'v(\cdot,x_n)||_{L^1(B'_2)}^{abs} \\ & \cdot |||x'|^{\beta}v(\cdot,x_n)||_{L^q(B'_1)}^{(1-a)s} \left(|||x'|^{\mu}\nabla v||_{L^1(D_2)} + |||x'|^{\mu}v||_{L^1(D_2\setminus D_1)}\right)^{a(1-b)s}. \end{aligned}$$

Integrating the above in x_n over [0,1], applying Hölder's inequality, and followed by Young's inequality, we have, using abs + (1-a)s/q = 1, that

$$\begin{split} \||x'|^{\alpha}v\|_{L^{s}(D_{1})}^{s} &\leq C\||x'|^{\alpha}v\|_{L^{s}(D_{2}\backslash D_{1})}^{s} + C\||x'|^{\mu}\nabla'v\|_{L^{1}(D_{2})}^{abs}\||x'|^{\beta}v\|_{L^{q}(D_{1})}^{(1-a)s} \big(\||x'|^{\mu}\nabla v\|_{L^{1}(D_{2})}^{1} + \||x'|^{\mu}v\|_{L^{1}(D_{2}\backslash D_{1})}\big)^{a(1-b)s} \\ &\leq C\||x'|^{\alpha}v\|_{L^{s}(D_{2}\backslash D_{1})}^{s} + C\big(\||x'|^{\mu}\nabla v\|_{L^{1}(D_{2})} + \||x'|^{\mu}\nabla'v\|_{L^{1}(D_{2})}^{b} \\ & \cdot \||x'|^{\mu}v\|_{L^{1}(D_{2}\backslash D_{1})}^{1-b}\big)^{as}\||x'|^{\beta}v\|_{L^{q}(D_{1})}^{(1-a)s} \\ &\leq C\||x'|^{\alpha}v\|_{L^{s}(D_{2}\backslash D_{1})}^{s} + C\big(\||x'|^{\mu}\nabla v\|_{L^{1}(D_{2})} + \||x'|^{\mu}v\|_{L^{1}(D_{2}\backslash D_{1})}\big)^{as} \\ & \cdot \||x'|^{\beta}v\|_{L^{q}(D_{1})}^{(1-a)s}. \end{split}$$

By the definition of v and the above, using $m \leq s, q$, we have

$$|||x'|^{\alpha}u||_{L^{s}(D_{1})} \leq C|||x'|^{\alpha}u||_{L^{s}(D_{2}\setminus D_{1})} + C(|||x'|^{\mu}\nabla u||_{L^{1}(D_{2})}^{a} + |||x'|^{\mu}v||_{L^{1}(D_{2}\setminus D_{1})}^{a})|||x'|^{\beta}u||_{L^{q}(D_{2})}^{1-a}.$$

$$(5.10)$$

Since $m \leq 1$ and $1 \leq |x'| \leq 2$ in $D_2 \setminus D_1$, we apply Theorem 1.3 to obtain

$$|||x'|^{\mu}v||_{L^{1}(D_{2}\backslash D_{1})} \leq C||v||_{L^{1}(D_{2}\backslash D_{1})} \leq C||\nabla u||_{L^{1}(D_{2}\backslash D_{1})} \leq C|||x'|^{\mu}\nabla u||_{L^{1}(D_{2}\backslash D_{1})}.$$
(5.11)

By (5.10) and (5.11), we have

$$|||x'|^{\alpha}u||_{L^{s}(D_{1})} \leq C|||x'|^{\alpha}u||_{L^{s}(D_{2}\setminus D_{1})} + C|||x'|^{\mu}\nabla u||_{L^{1}(D_{2})}^{a}|||x'|^{\beta}u||_{L^{q}(D_{2})}^{1-a}.$$

The lemma is proved for p = 1.

Step 2. We prove inequality (5.1) when p > 1.

Let $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\alpha}, \bar{\mu}$ and $\bar{\beta}$ be defined by

$$\begin{split} &\frac{1}{\bar{s}} = \frac{1}{s} + \frac{1}{p'}, \quad \bar{p} = 1, \quad \frac{1}{\bar{q}} = \frac{s}{\bar{s}q}, \quad \bar{a} = \frac{as}{(1-a)\bar{s} + as}, \\ &\bar{\alpha} = \frac{\alpha s}{\bar{s}}, \quad \bar{\mu} = \frac{\alpha s}{p'} + \mu, \quad \bar{\beta} = \frac{\beta s}{\bar{s}}, \end{split}$$

where 1/p + 1/p' = 1. It can be verified that $0 < \bar{s} < s$, and $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\alpha}, \bar{\mu}, \bar{\beta}$ satisfy (1.7)-(1.14) with $\gamma_1 = \gamma_2 = \gamma_3 = 0$ and $s, p, q, a, \alpha, \mu, \beta$ replaced by $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\alpha}, \bar{\mu}, \bar{\beta}$ respectively. For the details of the verification, see Lemma 7.1 and its proof. For $u \in C^{0,1}(D_2)$, we have $|u|^{s/\bar{s}} \in C^{0,1}(D_2)$. Apply (5.1) with p = 1 to $|u|^{s/\bar{s}}$, we have, using Hölder's inequality and Young's inequality, that

$$\begin{split} & \||x'|^{\alpha}u\|_{L^{s}(D_{1})}^{s/\bar{s}} = \||x'|^{\bar{\alpha}}|u|^{s/\bar{s}}\|_{L^{\bar{s}}(D_{1})} \\ & \leq C\||x'|^{\bar{\alpha}}|u|^{s/\bar{s}}\|_{L^{\bar{s}}(D_{2}\backslash D_{1})} + C\||x'|^{\bar{\mu}}\nabla|u|^{s/\bar{s}}\|_{L^{1}(D_{2})}^{\bar{a}}\||x'|^{\bar{\beta}}|u|^{s/\bar{s}}\|_{L^{\bar{q}}(D_{2})}^{1-\bar{a}} \\ & = C\||x'|^{\alpha}u\|_{L^{s}(D_{2}\backslash D_{1})}^{s/\bar{s}} + C\||x'|^{\bar{\mu}-\mu}|u|^{s/\bar{s}-1} \cdot |x'|^{\mu}|\nabla u|\|_{L^{1}(D_{2})}^{\bar{a}}\||x'|^{\beta}u\|_{L^{q}(D_{2})}^{(1-\bar{a})q/\bar{q}} \\ & \leq C\||x'|^{\alpha}u\|_{L^{s}(D_{2}\backslash D_{1})}^{s/\bar{s}} + C\||x'|^{\alpha}u\|_{L^{s}(D_{2})}^{\bar{a}s/p'}\||x'|^{\mu}\nabla u\|_{L^{p}(D_{2})}^{\bar{a}}\||x'|^{\beta}u\|_{L^{q}(D_{2})}^{(1-\bar{a})s/\bar{s}} \\ & \leq C\||x'|^{\alpha}u\|_{L^{s}(D_{2}\backslash D_{1})}^{s/\bar{s}} + \frac{1}{2}\||x'|^{\alpha}u\|_{L^{s}(D_{2})}^{s/\bar{s}} + C\Big(\||x'|^{\mu}\nabla u\|_{L^{p}(D_{2})}^{\bar{a}}\||x'|^{\beta}u\|_{L^{q}(D_{2})}^{(1-\bar{a})s/\bar{s}}\Big)^{1/(1-\bar{a}\bar{s}/p')}. \end{split}$$

Inequality (5.1) follows from the above and the definitions of \bar{a} and \bar{s} . Lemma 5.1 is proved.

Remark 5.1. In the proof of Lemma 5.1, when a=1 or when 0 < a < 1 and $1/s + 1 - 1/p \le q/s$, we can use Theorem A and the classical Poincaré's inequality instead of Theorem 1.2 and 1.3.

For $0 \le r_1 < r_2 \le \infty$ and $\epsilon > 0$, let

$$K_{r_1, r_2, \epsilon} := \{ x \in \mathbb{R}^n \mid r_1 \le |x| < r_2, \quad |x'| \le \epsilon |x| \}.$$
 (5.12)

Lemma 5.2. Let $n \geq 2$, $0 \leq r_1 < r_2 \leq \infty$, $0 < \epsilon_1 < \epsilon_2 \leq 1$, $K_{\epsilon_i} := K_{r_1,r_2,\epsilon_i}$, i = 1, 2, and let $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β be real numbers satisfying (1.7)-(1.14) with $1/s \leq a/p + (1-a)/q$. Then there exists some positive constant C, depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu, \beta, \epsilon_1, \epsilon_2, r_1$ and r_2 , such that for all $u \in C^1(\bar{K}_{\epsilon_2})$,

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(K_{\epsilon_1})} \le C|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(K_{\epsilon_2}\setminus K_{\epsilon_1})} + C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(K_{\epsilon_2})}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(K_{\epsilon_2})}^{1-a}.$$
(5.13)

Furthermore, on any compact set in the parameter space in which (1.7)-(1.9) hold, the constant C is bounded.

Remark 5.2. Consider more general cones $K_{r_1,r_2,\Omega} = \{rx \mid r_1 \leq r \leq r_2, x \in \Omega\}$ for some open set $\Omega \subset \mathbb{S}^{n-1}$ with Lipschitz boundary. For open sets $\Omega_1 \subset \bar{\Omega}_1 \subset \Omega_2 \subset \mathbb{S}^{n-1}$ with Lipschitz boundaries, Lemma 5.2 still holds with K_{r_1,r_2,ϵ_i} replaced by K_{r_1,r_2,Ω_i} , i = 1, 2.

Proof of Lemma 5.2.

For $\epsilon > 0$, denote $K_{\epsilon} := K_{r_1, r_2, \epsilon}$, and let $K_{\epsilon}^+ := K_{\epsilon} \cap \{x_n \geq 0\}$, $K_{\epsilon}^- := K_{\epsilon} \cap \{x_n < 0\}$. We will only prove (5.13) with K_{ϵ_i} , i = 1, 2, replaced by $K_{\epsilon_i}^+$. The estimate on $K_{\epsilon_i}^-$ is similar.

Case 1. $0 < r_1 < r_2 < \infty$.

In this case, there exists a diffeomorphism $y = \Phi(x)$ from $K_{\epsilon_i}^+$ to $D_{\epsilon_i} = B'_{\epsilon_i} \times [0,1]$, satisfying $|y'|/C \leq |x'| \leq C|y'|$. Let $\tilde{\mu} = \mu + \gamma_2 - \gamma_1/a + (1-a)\gamma_3/a$. By (1.11) we have $\tilde{\mu} \geq \mu$. Notice we are in the case $1/s \leq a/p + (1-a)/q$, it can be verified that $s, p, q, a, \alpha, \tilde{\mu}, \beta$ satisfy (1.7)-(1.14) with μ replaced by $\tilde{\mu}$ and $\gamma_1 = \gamma_2 = \gamma_3 = 0$. Applying Lemma 5.1 to $\hat{u} = u \circ \Phi^{-1}$, we have

$$|||y'|^{\alpha} \hat{u}||_{L^{s}(D_{\epsilon_{1}})} \leq C|||y'|^{\alpha} \hat{u}||_{L^{s}(D_{\epsilon_{2}} \setminus D_{\epsilon_{1}})} + C|||y'|^{\tilde{\mu}} \nabla \hat{u}||_{L^{p}(D_{\epsilon_{2}})}^{a}|||y'|^{\beta} \hat{u}||_{L^{q}(D_{\epsilon_{2}})}^{1-a}$$

$$\leq C|||y'|^{\alpha} \hat{u}||_{L^{s}(D_{\epsilon_{2}} \setminus D_{\epsilon_{1}})} + C|||y'|^{\mu} \nabla \hat{u}||_{L^{p}(D_{\epsilon_{2}})}^{a}|||y'|^{\beta} \hat{u}||_{L^{q}(D_{\epsilon_{2}})}^{1-a}.$$

Inequality (5.13) follows immediately.

Case 2. $r_1 = 0$ or $r_2 = \infty$.

Working with $u(\lambda x)$ instead of u(x), we only need to treat the cases when $r_1 = 0$ and $r_2 = 1$, or $r_1 = 1$ and $r_2 = \infty$, or $r_1 = 0$ and $r_2 = \infty$. Let $R_k := \{x \in \mathbb{R}^n \mid 2^{k-1} \le |x| < 2^k\}$, $k \in \mathbb{Z}$. By Case 1, (1.10) and scaling, we have, for every $k \in \mathbb{Z}$, that

$$|||x|^{\gamma_{1}}|x'|^{\alpha}u||_{L^{s}(R_{k}\cap K_{\epsilon_{1}})}^{s} \leq C|||x|^{\gamma_{1}}|x'|^{\alpha}u||_{L^{s}(R_{k}\cap K_{\epsilon_{2}}\setminus K_{\epsilon_{1}})}^{s} + C|||x|^{\gamma_{2}}|x'|^{\mu}\nabla u||_{L^{p}(R_{k}\cap K_{\epsilon_{2}})}^{as} \cdot |||x|^{\gamma_{3}}|x'|^{\beta}u||_{L^{q}(R_{k}\cap K_{\epsilon_{2}})}^{(1-a)s}.$$

$$(5.14)$$

When $r_1 = 0$ and $r_2 = \infty$, take the sum of (5.14) over all $k \in \mathbb{Z}$, we have, using $as/p + \frac{(1-a)s}{a} \ge 1$ and (4.9), that

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(K_{\epsilon_1})}^s$$

$$\leq C \||x|^{\gamma_1}|x'|^{\alpha}u\|_{L^s(K_{\epsilon_2}\backslash K_{\epsilon_1})}^s + C\sum_{k=-\infty}^{\infty} \||x|^{\gamma_2}|x'|^{\mu}\nabla u\|_{L^p(R_k\cap K_{\epsilon_2})}^{as}\||x|^{\gamma_3}|x'|^{\beta}u\|_{L^q(R_k\cap K_{\epsilon_2})}^{(1-a)s}$$

$$\leq C \||x|^{\gamma_1}|x'|^{\alpha}u\|_{L^s(K_{\epsilon_2}\backslash K_{\epsilon_1})}^s + C \||x|^{\gamma_2}|x'|^{\mu}\nabla u\|_{L^p(K_{\epsilon_2})}^{as} \||x|^{\gamma_3}|x'|^{\beta}u\|_{L^q(K_{\epsilon_2})}^{(1-a)s}.$$

So (5.13) is proved when $r_1 = 0$ and $r_2 = \infty$. Inequality (5.13) for $r_1 = 0$ and $r_2 = 1$ follows by summing (5.14) over $k \leq 0$. For $r_1 = 1$ and $r_2 = \infty$, we sum (5.14) over $k \geq 0$. Lemma 5.2 is proved.

Proof of the sufficiency part of Theorem 1.1 when $1/s \le a/p + (1-a)/q$.

Fix $\epsilon > 0$ small, let K_{ϵ} be the cone defined by (5.12) with $r_1 = 0$ and $r_2 = \infty$.

By (1.7), (1.9), (1.10), (1.12) and (1.14), we have that $s, p, q, a, \gamma_1 + \alpha, \gamma_2 + \mu, \gamma_3 + \beta$ satisfy (1.7) and (1.2)-(1.5) with $\gamma_1, \gamma_2, \gamma_3$ replaced by $\gamma_1 + \alpha, \gamma_2 + \mu, \gamma_3 + \beta$ respectively. Then by Theorem 4.1, we have

$$|||x|^{\gamma_1+\alpha}u||_{L^s(\mathbb{R}^n\setminus K_{\epsilon})} \le C|||x|^{\gamma_2+\mu}\nabla u||_{L^p(\mathbb{R}^n\setminus K_{\epsilon})}^a|||x|^{\gamma_3+\beta}u||_{L^q(\mathbb{R}^n\setminus K_{\epsilon})}^{1-a}.$$

Since $\epsilon |x| \leq |x'| \leq |x|$ for x in $\mathbb{R}^n \setminus K_{\epsilon}$, we have

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(\mathbb{R}^n\setminus K_{\epsilon})} \le C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\mathbb{R}^n\setminus K_{\epsilon})}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(\mathbb{R}^n\setminus K_{\epsilon})}^{1-a}.$$
(5.15)

By Lemma 5.2,

It follows from (5.15) and (5.16) that

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(\mathbb{R}^n)} \le C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\mathbb{R}^n)}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(\mathbb{R}^n)}^{1-a}.$$
(5.17)

The sufficiency part of Theorem 1.1 is proved when $1/s \le a/p + (1-a)/q$.

Next, we prove the sufficiency part of Theorem 1.1 when 1/s > a/p + (1-a)/q. We reduce it to the case 1/s = a/p + (1-a)/q by the following lemma. This reduction procedure is analogous to the arguments in Section (V) in [10].

Lemma 5.3. Let $n \geq 2$, Ω be a bounded open set in \mathbb{R}^n and $u \in C^{0,1}(\Omega)$. Assume that for any $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu, \beta$ satisfying (1.7)-(1.14) and 1/s = a/p + (1-a)/q, there exists some constant C, depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β , such that

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(\Omega)} \le C'|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(\Omega)}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(\Omega)}^{1-a}.$$
 (5.18)

Then for any $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β satisfying (1.7)-(1.14) with 1/s > a/p + (1-a)/q, there exists some constant C and $0 \le a', a'' \le 1$, depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu, \beta, \Omega$ and C', such that

$$|||x|^{\gamma_{1}}|x'|^{\alpha}u||_{L^{s}(\Omega)} \leq C\Big(|||x|^{\gamma_{2}}|x'|^{\mu}\nabla u||_{L^{p}(\Omega)}^{a'}|||x|^{\gamma_{3}}|x'|^{\beta}u||_{L^{q}(\Omega)}^{1-a'} + |||x|^{\gamma_{2}}|x'|^{\mu}\nabla u||_{L^{p}(\Omega)}^{a''}|||x|^{\gamma_{3}}|x'|^{\beta}u||_{L^{q}(\Omega)}^{1-a''}\Big).$$

$$(5.19)$$

Proof. For $u \in C^{0,1}(\Omega)$, we assume (5.18) holds, and we will prove (5.19). Let C denote a positive constant depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu, \beta, \Omega$ and C' which may vary from line to line. Condition (1.14) and 1/s > a/p + (1-a)/q imply 0 < a < 1. Denote $A := \||x|^{\gamma_2}|x'|^{\mu}\nabla u\|_{L^p(\mathbb{R}^n)}$ and $B := \||x|^{\gamma_3}|x'|^{\beta}u\|_{L^q(\mathbb{R}^n)}$.

For constants $0 \le a', a'' \le 1, \alpha', \alpha''$, we define $s', s'', \gamma'_1, \gamma''_1$ by

$$\frac{1}{s'} = \frac{a'}{p} + \frac{1 - a'}{q}, \quad \gamma_1' + \alpha' = a'(\gamma_2 + \mu - 1) + (1 - a')(\gamma_3 + \beta),
\frac{1}{s''} = \frac{a''}{p} + \frac{1 - a''}{q}, \quad \gamma_1'' + \alpha'' = a''(\gamma_2 + \mu - 1) + (1 - a'')(\gamma_3 + \beta).$$
(5.20)

Let $\zeta(x)$ be a smooth function satisfying $\zeta(x) = 1$ for $|x| \le 1$, $\zeta(x) = 0$ for $|x| \ge 2$ and $|\nabla \xi(x)| \le 3$. We have

$$|||x|^{\gamma_1}|x_1|^{\alpha}u||_{L^s(\mathbb{R}^n)} \le |||x|^{\gamma_1}|x_1|^{\alpha}\zeta u||_{L^s(\mathbb{R}^n)} + |||x|^{\gamma_1}|x_1|^{\alpha}(1-\zeta)u||_{L^s(\mathbb{R}^n)} =: I_1 + I_2.$$
(5.21)

We estimate

$$I_{1} \leq \||x|^{\gamma'_{1}}|x_{1}|^{\alpha'}u\|_{L^{s'}(\mathbb{R}^{n})} \left(\int_{|x| \leq 2} \left| |x|^{\gamma_{1} - \gamma'_{1}}|x'|^{\alpha - \alpha'} \right|^{ss'/(s' - s)} \right)^{1/s - 1/s'}, \tag{5.22}$$

and

$$I_{2} \leq \||x|^{\gamma_{1}''}|x_{1}|^{\alpha''}u\|_{L^{s''}(\mathbb{R}^{n})} \left(\int_{|x|\geq 1} \left| |x|^{\gamma_{1}-\gamma_{1}''}|x'|^{\alpha-\alpha''} \right|^{ss''/(s''-s)} \right)^{1/s-1/s''}.$$
 (5.23)

by Hölder's inequality, provided

$$\frac{1}{s'} < \frac{1}{s} \text{ and } \frac{1}{s''} < \frac{1}{s}.$$
 (5.24)

The second integrals on the right hand sides in (5.22) and (5.23) are finite if

$$\frac{1}{s'} + \frac{\gamma_1' + \alpha'}{n} < \frac{1}{s} + \frac{\gamma_1 + \alpha}{n} < \frac{1}{s''} + \frac{\gamma_1'' + \alpha''}{n},\tag{5.25}$$

$$\frac{1}{s'} + \frac{\alpha'}{n-1} < \frac{1}{s} + \frac{\alpha}{n-1}$$
 and $\frac{1}{s''} + \frac{\alpha''}{n-1} < \frac{1}{s} + \frac{\alpha}{n-1}$. (5.26)

By the assumption of the lemma, we will have

$$|||x|^{\gamma_1'}|x_1|^{\alpha'}u||_{L^{s'}(\mathbb{R}^n)} \le CA^{a'}B^{1-a'}, \quad |||x|^{\gamma_1''}|x_1|^{\alpha''}u||_{L^{s''}(\mathbb{R}^n)} \le CA^{a''}B^{1-a''}, \quad (5.27)$$

provided (1.7)-(1.14) with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ or $s'', a'', \gamma''_1, \alpha''$ respectively.

So by (5.21)-(5.23) and (5.27), to prove (5.19), we only need to choose appropriate a', a'', α' and α'' such that (5.24)-(5.26) are satisfied, and (1.7)-(1.14) hold with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ or $s'', \alpha'', \gamma''_1, \alpha''$ respectively.

The choice of a' and α' and the choice of a'' and α'' can be made independently and analogously. We always require a' and a'' to be close to a and in particular 0 < a', a'' < 1. By (5.20), conditions (1.7), (1.10) and (1.14) always hold with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ or $s'', a'', \gamma''_1, \alpha''$ respectively. By (5.20), we have

$$a'(\gamma_2 + \mu) + (1 - a')(\gamma_3 + \beta) - (\gamma_1' + \alpha') = a'.$$

By the above requirement on a' and a'', we have (1.12) with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ respectively. Similarly, we have (1.12) with s, a, γ_1, α there replaced by $s'', a'', \gamma''_1, \alpha''$ respectively.

By (5.20), we have

$$\frac{1}{s'} + \frac{\gamma_1' + \alpha'}{n} = a'(\frac{1}{n} + \frac{\gamma_2 + \mu - 1}{n}) + (1 - a')(\frac{1}{a} + \frac{\gamma_3 + \beta}{n}).$$

By (1.9) and (1.10), the right hand side of the above is strictly positive when a' = a. Thus as long as we choose a' close enough to a, we have $1/s' + (\gamma'_1 + \alpha')/n > 0$, and therefore (1.9) holds with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ respectively. Similarly,

we have (1.9) with s, a, γ_1, α there replaced by $s'', a'', \gamma_1'', \alpha''$ respectively, as long as we choose a'' close enough to a.

Moreover, by the assumption 1/s > a/p + (1-a)/q and the definition of s' and s'' in (5.20), we have that (5.24) hold as long as a' and a'' are close enough to a. By (1.14), (1.10) and the assumption that 1/s > a/p + (1-a)/q, we have $1/p + (\gamma_2 + \mu - 1)/n \neq 1/q + (\gamma_3 + \beta)/n$. For (5.25) to hold, we only need to require

$$0 < a' < a < a'' < 1, \quad \text{if } \frac{1}{p} + \frac{\gamma_2 + \mu - 1}{n} > \frac{1}{q} + \frac{\gamma_3 + \beta}{n},$$

$$1 > a' > a > a'' > 0, \quad \text{if } \frac{1}{p} + \frac{\gamma_2 + \mu - 1}{n} < \frac{1}{q} + \frac{\gamma_3 + \beta}{n}.$$

It remains to show that we can further require $a', a'', \alpha', \alpha''$ to satisfy additional properties, such that (5.26) is satisfied, and (1.8), (1.11) and (1.13) hold with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ or $s'', a'', \gamma''_1, \alpha''$ respectively. By the definition of 1/s' and 1/s'' in (5.20), equation (5.26) holds provided

$$\alpha' < G(a'), \quad \alpha'' < G(a''), \tag{5.28}$$

where $G(\theta) = (n-1)(1/s - \theta/p - (1-\theta)/q) + \alpha$. By the definition of 1/s' and 1/s'' in (5.20), equation (1.8) holds with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ or $s'', a'', \gamma''_1, \alpha''$ respectively, provided

$$\alpha' > F_1(a'), \quad \alpha'' > F_1(a'),$$
 (5.29)

where $F_1(\theta) = -(n-1)(\theta/p + (1-\theta)/q)$. By the definition of $\gamma_1' + \alpha'$ and $\gamma_1'' + \alpha''$ in (5.20), equation (1.11) holds with s, a, γ_1, α there replaced by $s', a', \gamma_1', \alpha'$ or $s'', a'', \gamma_1'', \alpha''$ respectively, provided

$$\alpha' > F_2(a'), \quad \alpha'' > F_2(a''),$$
 (5.30)

where $F_2(\theta) = \theta(\mu - 1) + (1 - \theta)\beta$. By the definition of 1/s' and 1/s'' in (5.20), equation (1.13) holds with s, a, γ_1, α there replaced by $s', a', \gamma'_1, \alpha'$ or $s'', a'', \gamma''_1, \alpha''$ respectively, provided (5.30). So we only need to further require $a', a'', \alpha', \alpha''$ to satisfy (5.28)-(5.30).

By (1.8), $1/s + \alpha/(n-1) > 0$, so we have $F_1(a) < G(a)$. By (1.13), (1.14) and the assumption that 1/s > a/p + (1-a)/q, the inequality in (1.13) is strict, and therefore $F_2(a) < G(a)$. So as long as a' and a'' are close enough to a, we can find α' and α'' to satisfy (5.28)-(5.30). Lemma 5.3 is proved.

Proof of the sufficiency part of Theorem 1.1 when 1/s > a/p + (1-a)/q.

In this case, by (1.10) and (1.14), we must have $1/p + (\gamma_2 + \mu - 1)/n \neq 1/q + (\gamma_3 + \beta)/n$. So there exist some constants C and λ , such that $\hat{u} = Cu(\lambda x)$ satisfies $\||x|^{\gamma_2}|x'|^{\mu}\nabla \hat{u}\|_{L^p(\mathbb{R}^n)} = 1$ and $\||x|^{\gamma_3}|x'|^{\beta}\hat{u}\|_{L^q(\mathbb{R}^n)} = 1$. By Theorem 1.1 for $1/s \leq a/p + (1-a)/q$, (1.15) holds for \hat{u} and all $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu, \beta$ satisfying (1.7)-(1.14) and $1/s \leq a/p + (1-a)/q$. Then by Lemma 5.3, when 1/s > a/p + (1-a)/q, we have

$$|||x|^{\gamma_1}|x'|^{\alpha}\hat{u}||_{L^p(\mathbb{R}^n)} \leq C|||x|^{\gamma_2}|x'|^{\mu}\nabla\hat{u}||_{L^p(\mathbb{R}^n)}^a|||x|^{\gamma_3}|x'|^{\beta}\hat{u}||_{L^q(\mathbb{R}^n)}^{1-a}.$$

Then (1.15) holds for u by scaling.

The sufficiency part of Theorem 1.1 is proved.

6 Two variants of Theorem 1.1 and Theorem A

We have the following variant of Theorem 1.1.

Theorem 6.1. Let $n \geq 2$, $0 \leq r_1 < r_2 \leq \infty$, $\epsilon > 0$, $K := K_{r_1,r_2,\epsilon}$ be defined as (5.12), and $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β be real numbers satisfying (1.7)-(1.14). Then there exists some positive constant C, depending only on $\epsilon, s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β , such that for all $u \in C^1(\bar{K})$ with u = 0 on ∂K ,

$$|||x|^{\gamma_1}|x'|^{\alpha}u||_{L^s(K)} \le C|||x|^{\gamma_2}|x'|^{\mu}\nabla u||_{L^p(K)}^a|||x|^{\gamma_3}|x'|^{\beta}u||_{L^q(K)}^{1-a}.$$

$$(6.1)$$

Furthermore, on any compact set in the parameter space in which (1.7)-(1.9) hold, the constant C is bounded.

Proof. Extend u to be zero outside K. When $1/s \le a/p + (1-a)/q$, apply Lemma 5.2 to u with $K_{\epsilon_1} = K$ and K_{ϵ_2} be a larger cone containing K, we obtain (6.1).

Now we consider the case when 1/s > a/p + (1-a)/q. By (1.10) and (1.14), we have $1/p + (\gamma_2 + \mu - 1)/n \neq 1/q + (\gamma_3 + \beta)/n$. So there exist some constants C and λ , such that $\hat{u} = Cu(\lambda x)$ satisfies $||x|^{\gamma_2}|x'|^{\mu}\nabla \hat{u}||_{L^p(K)} = 1$ and $||x|^{\gamma_3}|x'|^{\beta}\hat{u}||_{L^q(K)} = 1$. Since we have proved (6.1) when 1/s = a/p + (1-a)/q, we can apply Lemma 5.3 to \hat{u} to obtain, for some $0 \leq a', a'' \leq 1$, that

$$\begin{aligned} & |||x|^{\gamma_{1}}|x'|^{\alpha}\hat{u}||_{L^{s}(K)} \\ & \leq C\Big(|||x|^{\gamma_{2}}|x'|^{\mu}\nabla u||_{L^{p}(K)}^{a'}|||x|^{\gamma_{3}}|x'|^{\beta}u||_{L^{q}(K)}^{1-a'} + |||x|^{\gamma_{2}}|x'|^{\mu}\nabla u||_{L^{p}(K)}^{a''}|||x|^{\gamma_{3}}|x'|^{\beta}u||_{L^{q}(K)}^{1-a''}\Big) \\ & = 2C|||x|^{\gamma_{2}}|x'|^{\mu}\nabla\hat{u}||_{L^{p}(K)}^{a}|||x|^{\gamma_{3}}|x'|^{\beta}\hat{u}||_{L^{q}(K)}^{1-a}. \end{aligned}$$

The following is a variant of Theorem A.

Theorem 6.2. Let $n \geq 1$, R > 0, $B_R = \{x \in \mathbb{R}^n \mid |x| \leq R\}$, $0 < \lambda < \infty$, Assume $s, p, q, a, \gamma_1, \gamma_2, \gamma_3$ satisfy (1.7), (1.2)-(1.5). Moreover, assume $1 \leq p \leq \infty$ if $1 \leq \lambda < \infty$, and $\max\{1, (n-1)/(1+(n-1)\lambda)\} \leq p \leq \infty$ if $0 < \lambda < 1$. Then there exists some positive constant C, depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3$ and λ , such that for every nonnegative $w \in W^{1,1}(B_R)$, $v := w - (\int_{\partial B_{|x|}} w^{1/\lambda})^{\lambda}$ satisfies

$$|||x|^{\gamma_1}v||_{L^s(B_R)} \le C|||x|^{\gamma_2}\nabla v||_{L^p(B_R)}^a |||x|^{\gamma_3}v||_{L^q(B_R)}^{1-a}.$$
(6.2)

Furthermore, on any compact set in the parameter space in which (1.7) and (1.2) hold, the constant C is bounded.

Proof. Let C denote a positive constant depending only on $s, p, q, a, \gamma_1, \gamma_2, \gamma_3$ and λ , which may vary from line to line.

For a = 0, we deduce from (1.3), (1.4) and (1.5) that $\gamma_1 = \gamma_3$ and s = q, thus (6.2) is obvious. In the rest of the proof we assume $0 < a \le 1$.

Case 1.
$$1/s \le a/p + (1-a)/q$$
.

Let

$$R_k := \{ x' \in B_1 \mid \frac{1}{2^k} \le |x'| \le \frac{1}{2^{k-1}} \}, \quad k \in \mathbb{Z}.$$

We first prove that

$$|||x|^{\gamma_1}v||_{L^s(R_k)} \le C|||x|^{\gamma_2}\nabla v||_{L^p(R_k)}^a |||x|^{\gamma_3}v||_{L^q(R_k)}^{1-a}, \quad k \in \mathbb{Z}.$$

$$(6.3)$$

By scaling, using (1.3), we only need to prove (6.3) for k = 1.

Let \bar{s}, \bar{q} and \bar{a} be defined as in (7.1). Since a > 0, we have $\bar{a} > 0$. Define $t \in (0, \infty]$ by

$$\frac{1}{t} = \frac{1}{\bar{a}}(\frac{1}{\bar{s}} - \frac{1-\bar{a}}{\bar{q}}), \quad \text{if } \frac{1}{\bar{s}} - \frac{1-\bar{a}}{\bar{q}} > 0,$$

and $t = \infty \text{ if } 1/\bar{s} - (1 - \bar{a})/\bar{q} = 0.$

In the current case we have $1/s \le a/p + (1-a)/q$. By (1.3) and (1.4), we have $1/s \ge a(1/p-1/n) + (1-a)/q$. By the same arguments as in part (g) in the proof of Lemma 7.1, we have $1/\bar{s} \le \bar{a} + (1-\bar{a})/\bar{q}$ and $1/\bar{s} - (1-\bar{a})/\bar{q} \ge \bar{a}(n-1)/n \ge 0$, and therefore $(n-1)/n \le 1/t \le 1$. We have proved that

$$1 \le t \le \frac{n}{n-1}$$
 for $n \ge 2$ and $1 \le t \le \infty$ for $n = 1$.

By Hölder's inequality, provided $1/\bar{s} = \bar{a}/t + (1-\bar{a})/\bar{q}$, $0 < \bar{a} \le 1$, $1 \le t \le \infty$, and $\bar{q} > 0$, we have,

$$||v||_{L^{s}(R_{1})}^{s/\bar{s}} = ||v|^{s/\bar{s}}||_{L^{\bar{s}}(R_{1})} \le ||v|^{s/\bar{s}}||_{L^{t}(R_{1})}^{\bar{a}}||v|^{s/\bar{s}}||_{L^{\bar{q}}(R_{1})}^{1-\bar{a}} = ||v|^{s/\bar{s}}||_{L^{t}(R_{1})}^{\bar{a}}||v||_{L^{q}(R_{1})}^{(1-\bar{a})s/\bar{s}}.$$

$$(6.4)$$

where we have used the definition of \bar{q} in the last step.

Since $1 \le t \le n/(n-1)$, we apply Hölder's inequality and Sobolev's inequality to obtain

$$|||v|^{s/\bar{s}}||_{L^{t}(R_{1})} \leq C|||v|^{s/\bar{s}}||_{L^{\frac{n}{n-1}}(R_{1})} \leq C(|||v|^{s/\bar{s}}||_{L^{1}(R_{1})} + ||\nabla|v|^{s/\bar{s}}||_{L^{1}(R_{1})})$$

$$\leq C\left(|||v|^{s/\bar{s}-1}||_{L^{p'}(R_{1})}||v||_{L^{p}(R_{1})} + |||v|^{s/\bar{s}-1}||_{L^{p'}(R_{1})}||\nabla v||_{L^{p}(R_{1})}\right)$$

$$\leq C||v||_{L^{s}}^{s/p'}(||v||_{L^{p}(R_{1})} + ||\nabla v||_{L^{p}(R_{1})}),$$

$$(6.5)$$

where in the last step we have used the fact that $(s/\bar{s}-1)p'=s$ from the definition of \bar{s} .

Since $p \ge 1$ when $1 \le \lambda < \infty$ and $\max\{1, n/(1+n\lambda)\} \le p \le \infty$ when $0 < \lambda < 1$, we have, by Theorem 1.3, that

$$||v||_{L^p(R_1)}^p \le \int_{1/2}^1 ||v||_{L^p(\partial B_\rho)}^p d\rho \le C \int_{1/2}^1 ||\nabla_{tan}v||_{L^p(\partial B_\rho)}^p d\rho \le ||\nabla v||_{L^p(R_1)}^p.$$

By (6.4), (6.5) and the above, we have

$$\|v\|_{L^{s}(R_{1})}^{s/\bar{s}} \leq \||v|^{s/\bar{s}}\|_{L^{t}(R_{1})}^{\bar{a}}\|v\|_{L^{q}(R_{1})}^{(1-\bar{a})s/\bar{s}} \leq C\|v\|_{L^{s}}^{\bar{a}s/p'}\|\nabla v\|_{L^{p}(R_{1})}^{\bar{a}}\|v\|_{L^{q}(R_{1})}^{(1-\bar{a})s/\bar{s}}.$$

Using the definition of \bar{a} and \bar{s} in (7.1), we deduce from the above that

$$||v||_{L^s(R_1)} \le C||\nabla v||_{L^p(R_1)}^a ||v||_{L^q(R_1)}^{1-a}.$$

We have proved (6.3) for k = 1.

Since we are in the case $as/p + (1-a)s/q \ge 1$, we can use (4.9) to deduce from (6.3) that

$$\begin{split} \sum_{k=-\infty}^{\infty} \int_{R_k} ||x|^{\gamma_1} v|^s dx &\leq C \sum_{k=-\infty}^{\infty} \left(\int_{R_k} ||x|^{\gamma_2} \nabla v|^p dx \right)^{as/p} \left(\int_{R_k} ||x|^{\gamma_3} v|^q dx' \right)^{(1-a)s/q} \\ &\leq C \left(\sum_{k=-\infty}^{\infty} \int_{R_k} ||x|^{\gamma_2} \nabla v|^p dx \right)^{as/p} \left(\sum_{k=-\infty}^{\infty} \int_{R_k} ||x|^{\gamma_3} v|^q dx \right)^{(1-a)s/q} \\ &\leq C |||x^{\gamma_2} \nabla v||_{L^p(B_R)}^{as} |||x|^{\gamma_3} v||_{L^q(B_R)}^{(1-a)s}. \end{split}$$

We have proved (6.2) in Case 1.

Case 2.
$$1/s > a/p + (1-a)/q$$
.

By (1.3) and (1.5), we have $1/p + (\gamma_2 - 1)/n \neq 1/q + \gamma_3/n$. Thus there exist some positive constants λ_1 and λ_2 , such that $\hat{v}(x) = \lambda_1 v(\lambda_2 x)$ satisfies $||x|^{\gamma_2} \nabla \hat{v}||_{L^p(K)} = 1$ and $||x|^{\gamma_3} \hat{v}||_{L^q(K)} = 1$. By Case 1, arguing as in the paragraph below (4.13), we can find $0 \leq a', a'' \leq 1$ such that

$$\begin{aligned} ||x|^{\gamma_1} \hat{v}||_{L^s(K)} &\leq C \left(||x|^{\gamma_2} \nabla v||_{L^p(K)}^{a'} ||x|^{\gamma_3} v||_{L^q(K)}^{1-a'} + ||x|^{\gamma_2} \nabla v||_{L^p(K)}^{a''} ||x|^{\gamma_3} v||_{L^q(K)}^{1-a''} \right) \\ &= 2C ||x|^{\gamma_2} \nabla \hat{v}||_{L^p(K)}^a ||x|^{\gamma_3} \hat{v}||_{L^q(K)}^{1-a}. \end{aligned}$$

Inequality (6.2) follows.

7 Appendix: some facts about the parameters

In this section, we prove some properties of the parameters $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β which we use in earlier sections.

Let $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β be real numbers satisfying (1.7), define $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3, \bar{\alpha}, \bar{\mu}$ and $\bar{\beta}$ by

$$\frac{1}{\bar{s}} = \frac{1}{s} + \frac{1}{p'}, \quad \bar{p} = 1, \quad \frac{1}{\bar{q}} = \frac{s}{q\bar{s}}, \quad \bar{a} = \frac{as}{(1-a)\bar{s} + as},$$

$$\bar{\gamma}_1 = \frac{\gamma_1 s}{\bar{s}}, \quad \bar{\gamma}_2 = \frac{\gamma_1 s}{p'} + \gamma_2, \quad \bar{\gamma}_3 = \frac{\gamma_3 s}{\bar{s}},$$

$$\bar{\alpha} = \frac{\alpha s}{\bar{s}}, \quad \bar{\mu} = \frac{\alpha s}{p'} + \mu, \quad \bar{\beta} = \frac{\beta s}{\bar{s}},$$
(7.1)

where 1/p + 1/p' = 1. Clearly, $0 < \bar{s} < s$.

Lemma 7.1. (i) If $n \geq 1$, $s, p, q, a, \gamma_1, \gamma_2$ and γ_3 satisfy (1.7), (1.2)-(1.5), then $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\gamma}_1, \bar{\gamma}_2$ and $\bar{\gamma}_3$ also satisfy (1.7) and (1.2)-(1.5).

(ii) If $n \geq 2$, $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu$ and β satisfy (1.7)-(1.14), then $\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3, \bar{\alpha}, \bar{\mu}$ and $\bar{\beta}$ also satisfy (1.7)-(1.14).

(iii) Assume (1.7) holds, then $1/s \le a/p + (1-a)/q$ if and only if $1/\bar{s} \le \bar{a}/\bar{p} + (1-\bar{a})/\bar{q}$, and $1/s \ge a(1/p - 1/n) + (1-a)/q$ if and only if $1/\bar{s} \ge \bar{a}(1-1/n) + (1-\bar{a})/\bar{q}$.

Proof. For convenience, denote $\Lambda = (s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \mu, \beta)$ and $\bar{\Lambda} = (\bar{s}, \bar{p}, \bar{q}, \bar{a}, \bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3, \bar{\alpha}, \bar{\mu}, \bar{\beta})$. By (7.1), it is clear that $\bar{\Lambda}$ satisfies (1.7).

Now we prove the following statements (a)-(h), which imply (i)-(iii).

(a) If $n \geq 2$ and (1.8) holds for Λ , then (1.8) also holds for $\bar{\Lambda}$.

This follows from

$$\begin{split} &\frac{1}{\bar{s}} + \frac{\bar{\alpha}}{n-1} = \frac{s}{\bar{s}} \left(\frac{1}{s} + \frac{\alpha}{n-1} \right), \\ &\frac{1}{\bar{p}} + \frac{\bar{\mu}}{n-1} = \frac{1}{p} + \frac{\mu}{n-1} + \frac{s}{p'} \left(\frac{1}{s} + \frac{\alpha}{n-1} \right), \\ &\frac{1}{\bar{q}} + \frac{\bar{\beta}}{n-1} = \frac{s}{\bar{s}} \left(\frac{1}{q} + \frac{\beta}{n-1} \right), \end{split}$$

(b) If $n \ge 1$ and (1.9) holds for Λ , then (1.9) also holds for $\bar{\Lambda}$. This follows from

$$\frac{1}{\bar{s}} + \frac{\bar{\gamma}_1 + \bar{\alpha}}{n} = \frac{s}{\bar{s}} \left(\frac{1}{s} + \frac{\gamma_1 + \alpha}{n} \right),$$

$$\frac{1}{\bar{p}} + \frac{\bar{\gamma}_2 + \bar{\mu}}{n} = \frac{1}{p} + \frac{\gamma_2 + \mu}{n} + \frac{s}{p'} \left(\frac{1}{s} + \frac{\gamma_1 + \alpha}{n} \right),$$

$$\frac{1}{\bar{q}} + \frac{\bar{\gamma}_3 + \bar{\beta}}{n} = \frac{s}{\bar{s}} \left(\frac{1}{q} + \frac{\gamma_3 + \beta}{n} \right).$$
(7.2)

(c) Let $n \geq 1$, then Λ satisfies (1.10) if and only if $\bar{\Lambda}$ satisfies (1.10). Using the definition of \bar{s} , $\bar{\gamma}_1$ and $\bar{\alpha}$, we have

$$\frac{1}{\bar{s}} + \frac{\bar{\gamma}_1 + \bar{\alpha}}{n} = \frac{s}{\bar{s}} \left(1 + \frac{as}{p'} \right)^{-1} \left(1 + \frac{as}{p'} \right) \left(\frac{1}{s} + \frac{\gamma_1 + \alpha}{n} \right)
= \frac{s}{\bar{s}} \left(1 - a + \frac{s}{\bar{s}} a \right)^{-1} \left(\frac{1}{s} + \frac{\gamma_1 + \alpha}{n} + a \frac{s}{p'} \left(\frac{1}{s} + \frac{\gamma_1 + \alpha}{n} \right) \right).$$
(7.3)

By the definition of $\bar{\gamma}_2$, $\bar{\gamma}_3$, $\bar{\mu}$ and $\bar{\beta}$, we have

$$\bar{a}\left(1 + \frac{\bar{\gamma}_2 + \bar{\mu} - 1}{n}\right) + (1 - \bar{a})\left(\frac{1}{\bar{q}} + \frac{\bar{\gamma}_3 + \bar{\beta}}{n}\right) \\
= \frac{s}{\bar{s}}\left(1 - a + \frac{s}{\bar{s}}a\right)^{-1}\left(a\left(\frac{1}{p} + \frac{\gamma_2 + \mu - 1}{n}\right) + (1 - a)\left(\frac{1}{q} + \frac{\gamma_3 + \beta}{n}\right) + a\frac{s}{p'}\left(\frac{1}{s} + \frac{\gamma_1 + \alpha}{n}\right)\right). \tag{7.4}$$

By (7.3) and (7.4), we have (c).

(d) Let $n \geq 1$, then Λ satisfies (1.11) if and only if $\bar{\Lambda}$ satisfies (1.11).

This follows from the fact

$$\bar{a}\bar{\gamma}_{2} + (1-\bar{a})\bar{\gamma}_{3} - \bar{\gamma}_{1} = \frac{s}{(1-a)\bar{s} + as} \left(a\gamma_{2} + (1-a)\gamma_{3} + a\gamma_{1} \left(\frac{s}{\bar{s}} - 1 \right) \right) - \frac{s}{\bar{s}}\gamma_{1}$$
$$= \frac{s}{(1-a)\bar{s} + as} (a\gamma_{2} + (1-a)\gamma_{3} - \gamma_{1}).$$

(e) Λ satisfies (1.12) if and only if $\bar{\Lambda}$ satisfies (1.12). This is because of

$$\bar{a}(\bar{\gamma}_{2} + \bar{\mu}) + (1 - \bar{a})(\bar{\gamma}_{3} + \bar{\beta}) - (\bar{\gamma}_{1} + \bar{\alpha})$$

$$= \frac{s}{(1 - a)\bar{s} + as} \left(a(\gamma_{2} + \mu) + (1 - a)(\gamma_{3} + \beta) + a(\gamma_{1} + \alpha) \left(\frac{s}{\bar{s}} - 1 \right) \right) - \frac{s}{\bar{s}} (\gamma_{1} + \alpha)$$

$$= \frac{s}{(1 - a)\bar{s} + as} (a(\gamma_{2} + \mu) + (1 - a)(\gamma_{3} + \beta) - (\gamma_{1} + \alpha)).$$

(f) Let $n \geq 2$, then Λ satisfies (1.13) if and only if $\bar{\Lambda}$ satisfies (1.13). Using the definition of \bar{s} and $\bar{\alpha}$, we have

$$\frac{1}{\bar{s}} + \frac{\bar{\alpha}}{n-1} = \frac{s}{\bar{s}} \left(1 + \frac{as}{p'} \right)^{-1} \left(1 + \frac{as}{p'} \right) \left(\frac{1}{s} + \frac{\alpha}{n-1} \right)
= \frac{s}{\bar{s}} \left(1 - a + \frac{s}{\bar{s}} a \right)^{-1} \left(\frac{1}{s} + \frac{\alpha}{n-1} + a \frac{s}{p'} \left(\frac{1}{s} + \frac{\alpha}{n-1} \right) \right).$$
(7.5)

By the definition of $\bar{\mu}$ and $\bar{\beta}$, and the second and third equations in (??), we have

$$\bar{a}\left(\frac{1}{\bar{p}} + \frac{\bar{\mu} - 1}{n - 1}\right) + (1 - \bar{a})\left(\frac{1}{\bar{q}} + \frac{\bar{\beta}}{n - 1}\right) \\
= \frac{s}{\bar{s}}\left(1 - a + \frac{s}{\bar{s}}a\right)^{-1}\left(a\left(\frac{1}{p} + \frac{\mu - 1}{n - 1}\right) + (1 - a)\left(\frac{1}{q} + \frac{\beta}{n - 1}\right) + a\frac{s}{p'}\left(\frac{1}{s} + \frac{\alpha}{n - 1}\right)\right). \tag{7.6}$$

So (f) follows from (7.5) and (7.6).

(g) $1/s \le a/p + (1-a)/q$ if and only if $1/\bar{s} \le \bar{a} + (1-\bar{a})/\bar{q}$, and $1/s \ge a(1/p-1/n) + (1-a)/q$ if and only if $1/\bar{s} \ge \bar{a}(1-1/n) + (1-\bar{a})/\bar{q}$.

The first part follows from

$$\bar{a} + \frac{1 - \bar{a}}{\bar{q}} - \frac{1}{\bar{s}} = \frac{s}{\bar{s}(1 - a + as/\bar{s})} \left(a + \frac{1 - a}{q} - \frac{1 - a + as/\bar{s}}{s} \right)
= \frac{s}{\bar{s}(1 - a + as/\bar{s})} \left(a + \frac{1 - a}{q} - \frac{1 + as/p'}{s} \right)
= \frac{s}{\bar{s}(1 - a + as/\bar{s})} \left(\frac{a}{p} + \frac{1 - a}{q} - \frac{1}{s} \right).$$
(7.7)

The second part follows from (7.7) and the definition of \bar{a} , through the following computation

$$\bar{a}\left(1 - \frac{1}{n}\right) + \frac{1 - \bar{a}}{\bar{q}} - \frac{1}{\bar{s}} = \bar{a} + \frac{1 - \bar{a}}{\bar{q}} - \frac{1}{\bar{s}} - \frac{\bar{a}}{n} = \frac{s}{\bar{s}(1 - a + as/\bar{s})} \left(a\left(\frac{1}{p} - \frac{1}{n}\right) + \frac{1 - a}{q} - \frac{1}{s}\right).$$

(h) Λ satisfies (1.14) if and only if $\bar{\Lambda}$ satisfies (1.14), and Λ satisfies (1.5) if and only if $\bar{\Lambda}$ satisfies (1.5).

By the definition of $\bar{\Lambda}$, we have a=0 if and only if $\bar{a}=0$, and a=1 if and only if $\bar{a}=1$. By (7.2) and using $s/\bar{s}=1+s/p'$, we have

$$\frac{1}{p} + \frac{\gamma_2 + \mu - 1}{n} = \frac{1}{q} + \frac{\gamma_3 + \beta}{n} = \frac{1}{s} + \frac{\gamma_1 + \alpha}{n}$$

if and only if

$$\frac{1}{\bar{p}} + \frac{\bar{\gamma}_2 + \bar{\mu} - 1}{n} = \frac{1}{\bar{q}} + \frac{\bar{\gamma}_3 + \bar{\beta}}{n} = \frac{1}{\bar{s}} + \frac{\bar{\gamma}_1 + \bar{\alpha}}{n}.$$

By (7.5) and (7.6),

$$\frac{1}{s} + \frac{\alpha}{n-1} = a\left(\frac{1}{p} + \frac{\mu - 1}{n-1}\right) + (1 - a)\left(\frac{1}{q} + \frac{\beta}{n-1}\right)$$

if and only if

$$\frac{1}{\bar{s}} + \frac{\bar{\alpha}}{n-1} = \bar{a}\left(1 + \frac{\bar{\mu}-1}{n-1}\right) + (1-\bar{a})\left(\frac{1}{\bar{q}} + \frac{\bar{\beta}}{n-1}\right).$$

(h) then follows from the above in view of the first part of (g).

Now (i) follows from the fact that Λ satisfies (1.7), (b)-(d) and (h). (ii) follows from the fact that $\bar{\Lambda}$ satisfies (1.7), (a)-(e) and (h). (iii) follows from (g).

Lemma 7.2. Let $n \ge 2$, $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \beta$ and μ satisfy (1.7)-(1.14) with $\gamma_1, \gamma_2, \gamma_3 = 0$, a > 0, p = 1, and 1/s - (1-a)/q < 1. Then the parameters $\hat{s}, \hat{p}, \hat{q}, \hat{a}, \hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_3$, defined by (5.5), satisfy (1.7), (1.2)-(1.4) with n replaced by n - 1, and $1/\hat{s} \le \hat{a}/\hat{p} + (1-\hat{a})/\hat{q}$.

Proof. Assume $s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \beta, \mu$ satisfy (1.7)-(1.14) with $\gamma_1, \gamma_2, \gamma_3 = 0$. For convenience, denote $\Lambda = (s, p, q, a, \gamma_1, \gamma_2, \gamma_3, \alpha, \beta, \mu)$ and $\widehat{\Lambda} = (\hat{s}, \hat{p}, \hat{q}, \hat{a}, \hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_3)$.

Let b and λ be defined by (5.4). By the arguments below (5.4), we have $0 < \hat{a} < 1$ and $0 \le \lambda \le 1$. By this and the definition of $\hat{s}, \hat{p}, \hat{q}, \hat{a}$, (1.7) holds for $\hat{\Lambda}$. Also, by the definition (5.5) of $\hat{\Lambda}$ and (1.8) for Λ , (1.2) holds for $\hat{\Lambda}$ with n replaced by n-1.

Next, by the definition of Λ , λ and b, we have

$$\hat{a}\left(1 + \frac{\hat{\gamma}_2 - 1}{n - 1}\right) + (1 - \hat{a})\left(\frac{1}{\hat{q}} + \frac{\hat{\gamma}_3}{n - 1}\right)$$

$$= \hat{a}\left(1 + \frac{\mu - 1}{n - 1}\right) + (1 - \hat{a})\left(\lambda\left(1 + \frac{\mu}{n - 1}\right) + (1 - \lambda)\left(\frac{1}{q} + \frac{\beta}{n - 1}\right)\right)$$

$$= ab(1 + \frac{\mu - 1}{n - 1}) + (1 - ab)\left(\frac{a(1 - b)}{1 - ab}(1 + \frac{\mu}{n - 1}) + \frac{1 - a}{1 - ab}(\frac{1}{q} + \frac{\beta}{n - 1})\right)$$

$$= a\left(b(1 + \frac{\mu - 1}{n - 1}) + (1 - b)(1 + \frac{\mu}{n - 1})\right) + (1 - a)(\frac{1}{q} + \frac{\beta}{n - 1})$$

$$= a\left(1 + \frac{\mu}{n - 1} - \frac{b}{n - 1}\right) + (1 - a)\left(\frac{1}{q} + \frac{\beta}{n - 1}\right)$$

$$= \frac{n}{n - 1}\left(a\left(1 + \frac{\mu - 1}{n}\right) + (1 - a)\left(\frac{1}{q} + \frac{\beta}{n}\right) - \frac{1}{ns}\right),$$
(7.8)

where the definition of b is used in the last step. On the other hand, we have, by using the definition of $\hat{\Lambda}$ and λ , that

$$\frac{1}{\hat{s}} + \frac{\hat{\gamma}_1}{n-1} = \frac{1}{s} + \frac{\alpha}{n-1}.$$

Since Λ satisfies (1.10), by the above and (7.8), we have (1.3) holds for $\widehat{\Lambda}$.

Using the definition of $\widehat{\Lambda}$ and the fact that $\gamma_1 = \gamma_2 = \gamma_3 = 0$, we have, by using the definition of $\widehat{\Lambda}$ and λ , that

$$\hat{a}\hat{\gamma}_2 + (1 - \hat{a})\hat{\gamma}_3 - \hat{\gamma}_1 = a\mu + (1 - a)\beta - \alpha = a(\mu + \gamma_2) + (1 - a)(\beta + \gamma_3) - (\alpha + \gamma_1).$$

In view of (1.12) for Λ , (1.4) holds for $\widehat{\Lambda}$.

Finally, by the definition of λ and \hat{a} , we have

$$\hat{a} + \frac{1 - \hat{a}}{\hat{q}} = a + \frac{1 - a}{q}.$$

In view of (i) and the assumption p=1, we have $1/\hat{s} \leq \hat{a}/\hat{p} + (1-\hat{a})/\hat{q}$.

References

- [1] M. Badiale and G. Tarantello, A Sobolev-Hardy inequality with applications to a nonlinear elliptic equation arising in astrophysics, *Arch. Ration. Mech. Anal.* 163 (2002), 259–293.
- [2] J. Bourgain, H. Brezis and P. Mironescu, Another look at Sobolev spaces, Optimal control and partial differential equations. IOS, Amsterdam, (2001), 439–455.
- [3] J. Bourgain, H. Brezis and P. Mironescu, Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications, J. Anal. Math. 87 (2002), 77–101.
- [4] H. Bahouri, J.-Y. Chemin and I. Gallagher, Inégalités de Hardy précisées, (French) [Sharper Hardy inequalities], C. R. Math. Acad. Sci. Paris. 341 (2005), 89-92.
- [5] H. Bahouri, J.-Y. Chemin and I. Gallagher, Refined Hardy inequalities, *Ann. Sc Norm. Super. Pisa Cl. Sci.* 5 (2006), 375-391.
- [6] M. Cannone, G. Karch, D. Pilarczyk and G. Wu, Stability of singular solutions to the Navier-Stokes system, arXiv:2012.12714 [math.AP], 2020.
- [7] X. Cabré and X. Ros-Oton, Sobolev and isoperimetric inequalities with monomial weights, *Journal of Differential Equations* 255 (2013), 4312-4336.
- [8] X. Cabré, X. Ros-Oton and J. Serra, Sharp isoperimetric inequalities via the ABP method, Journal of the European Mathematical Society 18 (2016), 2971-2998.
- [9] L. Caffarelli, R. Kohn and L. Nirenberg, Partial regularity of suitable weak solutions of the Navier-Stokes equations, Comm. Pure Appl. Math. 35 (1982), 771-831.

- [10] L. Caffarelli, R. Kohn and L. Nirenberg, First order interpolation inequalities with weights, *Composition Math.* 53 (1984), 259–275.
- [11] F. Catrina and Z.Q. Wang, On the Caffarelli-Kohn-Nirenberg inequalities: Sharp constants, existence (and nonexistence), and symmetry of extremal functions, *Comm. Pure Appl. Math.* 54 (2001), 229-258.
- [12] J. Dolbeault, M.J. Esteban and M. Loss, Rigidity versus symmetry breaking via nonlinear flows on cylinders and Euclidean spaces, *Invent. math.* 206 (2016), 397–440.
- [13] R. Frank and R. Seiringer, Non-linear ground state representations and sharp Hardy inequalities, *J. Funct. Anal.* 255 (2008), 3407–3430.
- [14] D. Gilbarg and N. Trudinger, Elliptic partial differential equations of second order, Reprint of the 1998 edition, Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [15] G. Karch and D. Pilarczyk, Asymptotic stability of Landau solutions to Navier-Stokes system, *Arch. Ration. Mech. Anal.* 202 (2011), 115-131.
- [16] G. Karch, D. Pilarczyk and M.E. Schonbek, L^2 -asymptotic stability of singular solutions to the Navier-Stokes system of equations in \mathbb{R}^3 , *J. Math. Pures Appl.* 108 (2017), 14-40.
- [17] L. Li, Y.Y. Li and X. Yan, Homogeneous solutions of stationary Navier-Stokes equations with isolated singularities on the unit sphere. I. One singularity, *Arch. Ration. Mech. Anal.* 227 (2018), 1091–1163.
- [18] L. Li, Y.Y. Li and X.Yan, Homogeneous solutions of stationary Navier-Stokes equations with isolated singularities on the unit sphere. II. Classification of axisymmetric no-swirl solutions, *Journal of Differential Equations* 264 (2018), 6082-6108.
- [19] Y.Y. Li and X. Yan, Asymptotic stability of homogeneous solutions to the Navier-Stokes equations in R³, Journal of Differential Equations 297 (2021), 226-245.
- [20] Y. Y. Li, J. Zhang and T. Zhang, Asymptotic stability of Landau solutions to Navier-Stokes system under L^p -perturbations, arXiv:2012.14211 [math.AP], 2020.
- [21] C.S. Lin, Interpolation inequalities with weights, Comm. Partial Differential Equations 11 (1986), 1515–1538.
- [22] V.G. Maz'ya and T. Shaposhnikova, On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces, J. Funct. Anal. 195 (2002), 230–238; Erratum, J. Funct. Anal. 201 (2003), 298–300.
- [23] V.G. Maz'ja, Sobolev spaces, Translated from the Russian by T. O. Shaposhnikova. Springer Series in Soviet Mathematics. *Springer-Verlag*, Berlin, 1985.
- [24] H. Nguyen and M. Squassina, Fractional Caffarelli–Kohn–Nirenberg inequalities, *J. Funct. Anal.* 274 (2018), 2661–2672.

- [25] H. Nguyen and M. Squassina, On Hardy and Caffarelli-Kohn-Nirenberg inequalities. *J. Anal. Math.* 139 (2019), 773–797.
- [26] J. Zhang and T. Zhang, Global well-posedness of perturbed Navier-Stokes system around Landau-solutions, preprint, May 2021.