# CP violation phase and decomposition relations in Higgs BSM amplitudes

Ke-Yao Feng<sup>1a</sup>, Xia Wan<sup>1b</sup>, You-Kai Wang<sup>1c</sup>

<sup>1</sup>School of physics and Information Technology, Shaanxi Normal University, Xi'an 710119, China

## Abstract

We define a CP violation phase angle  $\xi$  to quantify the mixture of CP-even and CP-odd states for Higgs boson in new physics beyond Standard Model (BSM) firstly, and then show it explicitly in  $H \to \gamma\gamma$ ,  $H \to \gamma\ell\ell$  and  $H \to 4\ell$  amplitudes. The analytical form gives a good explanation why the CP violation phase could be observed and the interference between CP-even and CP-odd parts exist in  $H \to 4\ell$  process, but not in  $H \to \gamma\gamma$  and  $H \to \gamma\ell\ell$  processes. To understand the analytical structure of these BSM amplitudes, we introduce a new method of decomposing  $H \to \gamma\ell\ell$  and  $H \to 4\ell$  amplitudes into  $H \to \gamma\gamma$  amplitudes. For a comparison, by using the on-shell scattering amplitude approach we study the recursion relations of amplitudes and get a consistent result independently.

<sup>&</sup>lt;sup>a</sup> fengkeyao@snnu.edu.cn

<sup>&</sup>lt;sup>b</sup> wanxia@snnu.edu.cn

<sup>&</sup>lt;sup>c</sup> wangyk@snnu.edu.cn

## I. INTRODUCTION

There are two kinds of CP violation sources in Standrad Model (SM), one is weak CPviolation in Cabibbo-Kobayashi-Maskawa matrix [1, 2], the other is strong CP violation related to topological charge in QCD vacuum [3–5]. Both of them have relations with Higgs Yukawa couplings. The CKM matrix originates from general Yukawa coupling matrices for three generations [6, 7], the  $\theta$  angle in QCD vacuum could rotate to the complex phase of mass matrix by chiral transformation [8]. Even though, the SM Higgs boson is a CP-even scalar with CP-conserving interactions. By contrast, in new physics beyond SM (BSM), CP violation usually relates to Higgs bosons. One reason is there exist scalars and pseudoscalars instead of one single scalar in SM. A mixture of scalar and pseudoscalar is natural. Such as in Two Higgs Doublet Model (THDM) [9], Minimal Supersymmetric Standard model (MSSM) [10], and Composite Higgs Model [11], pseudoscalar always appears and there are no simple rule to forbid a mixture between scalar and pseudoscalar. Except for the theoretical naturalness and generality, one practical motivation for new CP violation source comes from the matter-antimatter asymmetry observed in our Universe [12–14]. In electroweak Baryogenesis mechanism [15], CP violation plus sphaleron transition [16] could produce baryon and lepton number violation during electroweak phase transition, but the CP violation ratio in SM is too small to fulfil the quantity of the matter-antimatter asymmetry observed [17-19]. Therefore new *CP* violation source must be added to get electroweak Baryogenesis.

We choose two model-independent framework to study CP violation: one is a traditional way through standard model effective field theory (SMEFT), the other is using on-shell scattering amplitude method to analyze amplitudes.

SMEFT provides a systematic way to describe BSM [20–22]. Based on Lorentz symmetry and gauge symmetries of SM, non-renormalizable operators of SM fields are written in Lagrangian to represent BSM new physics [21]. Then according to Feynman diagram, BSM amplitudes could be calculated and combined into SM software package to make simulation for experimental comparison. In this way, scattering amplitude is an intermediate object and its analytical expression is not illuminating. That is, although sometimes it looks having simple compact mathematical structures, the reason is not very clear.

On-shell method is a novel tool to deal with amplitudes directly, with no Lagrangian and

Feynman diagram needed [23]. It starts from on-shell particle states instead of field, sets up constraints, exploits analytical properties such as poles and branch cuts, then gets an available amplitude. Specifically, a 3-point massless (or 1 massive 2 massless) amplitude could be fixed by locality and little group scaling [23–25], then a n + 1-point tree amplitude could be constructed from n-point amplitudes through recursion relations. In this way all tree amplitudes could be obtained and they have clear mathematical structures.

We focus on  $H \to \gamma\gamma$ ,  $H \to \gamma\ell\ell$  and  $H \to 4\ell$  processes to analyze their BSM amplitudes. At Large Hadron collider (LHC), the  $H \to \gamma\gamma$  and  $H \to ZZ \to 4\ell$  processes are Higgs discovery channel [26, 27], which have the advantage of clean background and relative large signal. They are also golden channels for precise measurement of Higgs properties [28–31]. In our previous research, we notice that CP violation phase could not be probed solely in  $H \to \gamma\gamma$  or  $H \to \gamma\ell\ell$  processes without interference from background [32, 33]. By contrast, in  $H \to ZZ \to 4\ell$  processes CP violation could be probed solely through its kinematic angles [31, 34–36]. These could be explained clearly at amplitude level after we get a compact formula. Furthermore, we explore the relations between these BSM amplitudes through two independent ways in this paper.

The outline of this paper is as follows. In section II we calculate BSM amplitudes in SMEFT framework. The  $H \to \gamma\gamma$ ,  $H \to \gamma\ell\ell$  and  $H \to 4\ell$  processes correspond seperately to 3, 4, 5-point amplitudes. In section III we deduce decomposition relations for these amplitudes. In section IV we reproduce these BSM amplitudes in on-shell scattering amplitude approach. Section V is a summary and discussion.

#### II. BSM AMPLITUDES IN SMEFT

#### A. SMEFT

In SMEFT [20–22], the complete form of higher-dimensional operators can be written as

$$\mathcal{L} = \mathcal{L}_{\rm SM} + \frac{1}{\Lambda} \sum_{k} C_k^5 \mathcal{O}_k^5 + \frac{1}{\Lambda^2} \sum_{k} C_k^6 \mathcal{O}_k^6 + \mathcal{O}(\frac{1}{\Lambda^3}) , \qquad (1)$$

where  $\Lambda$  is energy scale of new physics, and  $C_k^i$  with i = 5, 6 are Wilson loop coefficients.

BSM HVV (V represents  $\gamma$ , Z/W boson) vertices start from dimension-six operators  $\mathcal{O}_k^6$ .

In Warsaw basis [21], they are

$$\mathcal{O}_{\Phi D}^{6} = (\Phi^{\dagger} D^{\mu} \Phi)^{*} (\Phi^{\dagger} D^{\mu} \Phi),$$
  

$$\mathcal{O}_{\Phi W}^{6} = \Phi^{\dagger} \Phi W_{\mu\nu}^{I} W^{I\mu\nu}, \quad \mathcal{O}_{\Phi B}^{6} = \Phi^{\dagger} \Phi B_{\mu\nu} B^{\mu\nu}, \quad \mathcal{O}_{\Phi WB}^{6} = \Phi^{\dagger} \tau^{I} \Phi W_{\mu\nu}^{I} B^{\mu\nu},$$
  

$$\mathcal{O}_{\Phi \tilde{W}}^{6} = \Phi^{\dagger} \Phi \tilde{W}_{\mu\nu}^{I} W^{I\mu\nu}, \quad \mathcal{O}_{\Phi \tilde{B}}^{6} = \Phi^{\dagger} \Phi \tilde{B}_{\mu\nu} B^{\mu\nu}, \quad \mathcal{O}_{\Phi \tilde{W}B}^{6} = \Phi^{\dagger} \tau^{I} \Phi \tilde{W}_{\mu\nu}^{I} B^{\mu\nu}, \quad (2)$$

where  $\Phi$  is a doublet representation under the  $SU(2)_L$  group and the aforementioned Higgs field H is one of its four components;  $D_{\mu} = \partial_{\mu} - igW^I_{\mu}T^I - ig'YB_{\mu}$ , where g and g' are coupling constants,  $T^I = \tau^I/2$ , where  $\tau^I$  are Pauli matrices, Y is the  $U(1)_Y$  generator;  $W^I_{\mu\nu} = \partial_{\mu}W^I_{\nu} - \partial_{\nu}W^I_{\mu} - g\epsilon^{IJK}W^J_{\mu}W^K_{\nu}$ ,  $B_{\mu\nu} = \partial_{\mu}B_{\nu} - \partial_{\nu}B_{\mu}$ ,  $\tilde{X}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu\rho\sigma}X^{\rho\sigma}$ ,

After spontaneous symmetry breaking, we get HVV effective interactions,

$$\mathcal{L}^{int} = -\frac{c_{VV}}{v} H V^{\mu\nu} V_{\mu\nu} - \frac{\dot{c}_{VV}}{v} H V^{\mu\nu} \tilde{V}_{\mu\nu} , \qquad (3)$$

where v = 246 GeV is the electroweak vacuum expectation value,  $c_{VV}$ ,  $\tilde{c}_{VV}$  are real numbers that originate from Wilson loop coefficients, V represents vector boson. A detailed formula between  $c_{VV}$ ,  $\tilde{c}_{VV}$  and Wilson loop coefficients  $C_k^6$  could be found in Ref. [34].

The CP violation phase could be defined as

$$\xi \equiv \tan^{-1}(\tilde{c}_{VV}/c_{VV}) , \text{ when } \operatorname{Arg}(\tilde{c}_{VV}/c_{VV}) = 0 \text{ or } \pi , \qquad (4)$$

where  $\xi = 0$   $(\frac{\pi}{2})$  represents a pure *CP*-even (-odd) *HVV* vertex.  $\xi \neq 0$  means *CP* violation and  $\xi = \frac{\pi}{2}$  corresponds to maximal *CP* violation if other Higgs vertices are supposed to be *CP*-even. In amplitudes we will see that  $\xi$  appears as a phase, which changes sign under *CP* transformation. That is why we name it as *CP* violation phase. Meanwhile,

$$c_{VV}^S \equiv \sqrt{c_{VV}^2 + \tilde{c}_{VV}^2} \tag{5}$$

could be defined as the amplitude moduli, which is proportional to signal strength in collider experiment.

## B. Helicity amplitudes

Feynman diagrams with effective HVV couplings are shown in Fig. 1. In calculation we adopt the conventions in [37, 38]:

$$\langle ij \rangle \equiv \langle i^{-}|j^{+}\rangle = \overline{u_{-}(p_{i})}u_{+}(p_{j}), \quad [ij] \equiv \langle i^{+}|j^{-}\rangle = \overline{u_{+}(p_{i})}u_{-}(p_{j}),$$
  
$$\langle ij \rangle [ji] = 2p_{i} \cdot p_{j}, \quad s_{ij} = (p_{i} + p_{j})^{2}, \quad \epsilon_{\mu}^{\pm}(p_{i},q) = \pm \frac{\langle q^{\mp}|\gamma_{\mu}|p_{i}^{\mp}\rangle}{\sqrt{2}\langle q^{\mp}|p_{i}^{\pm}\rangle}, \tag{6}$$




FIG. 1: Feynman diagrams of  $H \to \gamma\gamma$ ,  $H \to V\gamma \to \ell\ell\gamma$  and  $H \to VV \to 2\ell 2\ell'$  from left to right. Each HVV vertex is dotted as an effective coupling.

where  $p_i$  are momentum of external legs, q is the reference momentum that reflect the freedom of gauge transformation,  $\epsilon^{\pm}(p_i, q)$  is for outgoing photons with  $\pm$  helicities. After some calculation, the helicity amplitudes are as follows.

• For process  $H \to \gamma \gamma$ ,

$$\mathcal{M}(2^{+}_{\gamma}, 3^{+}_{\gamma}) = \frac{2c^{S}_{\gamma\gamma}}{v}e^{i\xi}[23]^{2} ,$$
  

$$\mathcal{M}(2^{-}_{\gamma}, 3^{-}_{\gamma}) = \frac{2c^{S}_{\gamma\gamma}}{v}e^{-i\xi}\langle 23\rangle^{2} ,$$
  

$$\mathcal{M}(2^{+}_{\gamma}, 3^{-}_{\gamma}) = 0 ,$$
  

$$\mathcal{M}(2^{-}_{\gamma}, 3^{+}_{\gamma}) = 0 ,$$
  
(7)

where we use  $\mathcal{M}(2^{h_2}_{\gamma}, 3^{h_3}_{\gamma})$  to represent  $\mathcal{M}(1^{h_1}_H, 2^{h_2}_{\gamma}, 3^{h_3}_{\gamma})$  since  $h_1$  is trivially zero for all cases,  $h_i$ s are helicities of external legs with momentum outgoing. The results show that the helicities of two photons should keep same sign because the spin of Higgs is zero and total angular momenta conserves. Under *CP* transformation  $\mathcal{M}(2^+_{\gamma}, 3^+_{\gamma})$ changes to  $\mathcal{M}(2^-_{\gamma}, 3^-_{\gamma})$ . Analytically it corresponds that  $\langle ij \rangle$  changes to [ij]. Thus in Eq. (7), a general nonzero  $\xi$  represents *CP* violation.

• For process  $H \to V\gamma \to \ell \ell \gamma$ ,

$$\mathcal{M}(2^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, 4^{-}_{\gamma}) = f^{-}_{V}(s_{23}) \times \frac{2c^{S}_{\gamma V}}{v} e^{-i\xi} [23] \langle 24 \rangle^{2},$$
  

$$\mathcal{M}(2^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, 4^{+}_{\gamma}) = f^{-}_{V}(s_{23}) \times \frac{2c^{S}_{\gamma V}}{v} e^{i\xi} \langle 23 \rangle [34]^{2},$$
  

$$\mathcal{M}(2^{+}_{\ell^{-}}, 3^{-}_{\ell^{+}}, 4^{+}_{\gamma}) = f^{+}_{V}(s_{23}) \times \frac{2c^{S}_{\gamma V}}{v} e^{i\xi} \langle 23 \rangle [24]^{2},$$
  

$$\mathcal{M}(2^{+}_{\ell^{-}}, 3^{-}_{\ell^{+}}, 4^{-}_{\gamma}) = f^{+}_{V}(s_{23}) \times \frac{2c^{S}_{\gamma V}}{v} e^{-i\xi} [23] \langle 34 \rangle^{2},$$
  
(8)

where  $s_{23} = (p_2 + p_3)^2$ ,  $f_V^-(s) = \sqrt{2} e l_V P_V(s)$  and  $f_V^+(s) = -\sqrt{2} e r_V P_V(s)$ ,  $P_V(s) = \frac{1}{s - M_V^2}$  is the propagator of the gauge boson,  $l_V$  and  $r_V$  are the left-handed and right-handed couplings between vector boson and leptons, and leptons are supposed to be massless. The remaining helicity amplitudes are equal to zero and thus not listed.

• For process  $H \to VV \to 2\ell 2\ell'$ ,

$$\mathcal{M}(2^{-}_{\ell^{-}},3^{+}_{\ell^{+}},4^{-}_{\ell^{\prime-}},5^{+}_{\ell^{\prime+}}) = f^{-}_{V}(s_{23})f^{-}_{V}(s_{45})\frac{2c^{S}_{VV}}{v}\left(e^{i\xi}\langle 23\rangle\langle 45\rangle[35]^{2} + e^{-i\xi}[23][45]\langle 24\rangle^{2}\right),$$

$$\mathcal{M}(2^{-}_{\ell^{-}},3^{+}_{\ell^{+}},4^{+}_{\ell^{\prime-}},5^{-}_{\ell^{\prime+}}) = f^{-}_{V}(s_{23})f^{+}_{V}(s_{45})\frac{2c^{S}_{VV}}{v}\left(e^{i\xi}\langle 23\rangle\langle 45\rangle[34]^{2} + e^{-i\xi}[23][45]\langle 25\rangle^{2}\right),$$

$$\mathcal{M}(2^{+}_{\ell^{-}},3^{-}_{\ell^{+}},4^{-}_{\ell^{\prime-}},5^{+}_{\ell^{\prime+}}) = f^{+}_{V}(s_{23})f^{-}_{V}(s_{45})\frac{2c^{S}_{VV}}{v}\left(e^{i\xi}\langle 23\rangle\langle 45\rangle[25]^{2} + e^{-i\xi}[23][45]\langle 34\rangle^{2}\right),$$

$$\mathcal{M}(2^{+}_{\ell^{-}},3^{-}_{\ell^{+}},4^{+}_{\ell^{\prime-}},5^{-}_{\ell^{\prime+}}) = f^{+}_{V}(s_{23})f^{+}_{V}(s_{45})\frac{2c^{S}_{VV}}{v}\left(e^{i\xi}\langle 23\rangle\langle 45\rangle[24]^{2} + e^{-i\xi}[23][45]\langle 35\rangle^{2}\right),$$

$$(9)$$

where VV could be  $\gamma\gamma$ , or ZZ, or  $\gamma Z$ , or  $W^+W^-$ , but when it represents  $\gamma Z$  or  $W^+W^-$ , the original Lagrangian in Eq. (3) should be scaled by a factor of 2 on the whole to make the formula consistent. The remaining helicity amplitudes are equal to zero.

#### III. DECOMPOSITION OF AMPLITUDES

#### A. Proof

Amplitudes in Eq. (7)(8) and (9) have similar structures. In  $H \to \gamma \gamma$  and  $H \to \ell \ell \gamma$ processes, there is only one term for each helicity amplitude. *CP* violation phase shows as a global phase. However, in  $H \to 4\ell$  process, two terms appear and *CP* violation phases have reverse signs. To explore how amplitudes change when external legs increase, we get this decomposition relation.

It is based on two key points. One point is the BSM HVV vertex from Eq. (3) is bilinear to the momenta of vector bosons, which is

$$\Gamma^{\mu\nu}(k,k') = -i\frac{4}{v} [c_{VV} \ (k^{\nu}k'^{\mu} - k \cdot k'g^{\mu\nu}) + \tilde{c}_{VV} \ \epsilon^{\mu\nu\rho\sigma}k_{\rho}k'_{\sigma}] , \qquad (10)$$

where k, k' are the momenta of the two vector bosons. So when  $k = p_2 + p_3$  or  $k' = p_4 + p_5$ ,

or both, where  $p_i$ s are momentum of external legs, we have

$$\Gamma^{\mu\nu}(k,k') = \Gamma^{\mu\nu}(p_2 + p_3,k') = \Gamma^{\mu\nu}(p_2,k') + \Gamma^{\mu\nu}(p_3,k')$$

$$= \Gamma^{\mu\nu}(p_2 + p_3,k') = \Gamma^{\mu\nu}(p_2,k') + \Gamma^{\mu\nu}(p_3,k')$$
(11)
(12)

$$=1^{\mu\nu}(p_2+p_3,p_4+p_5)=1^{\mu\nu}(p_2,p_4)+1^{\mu\nu}(p_2,p_5)+1^{\mu\nu}(p_3,p_4)+1^{\mu\nu}(p_3,p_5).$$
 (12)

The other point is that the current  $J_{\mu}$  of  $V \to \ell^+ \ell^-$  in Fig. 2 has same form as a polarization vector of photon (see Eq. (6)), except for an extra number,

$$J^{\pm}_{\mu}(s_{23}) = \frac{f^{\mp}_{V}(s_{23})}{\sqrt{2}} \langle 2^{\mp} | \gamma_{\mu} | 3^{\mp} \rangle \tag{13}$$

$$= \pm f_V^{\mp}(s_{23}) \langle 2^{\mp} | 3^{\pm} \rangle \epsilon_{\mu}^{\pm}(3,2)$$
 (14)

$$= \pm f_V^{\mp}(s_{23}) \langle 2^{\pm} | 3^{\mp} \rangle \epsilon_{\mu}^{\mp}(2,3), \tag{15}$$

where  $\epsilon_{\mu}^{\pm}(3,2) \equiv \epsilon_{\mu}^{\pm}(p_3,p_2)$  could be considered as a polarization vector of photon with external momentum of  $p_3$  and  $p_2$  is the chosen reference momentum. Similarly,  $\epsilon_{\mu}^{\pm}(2,3)$ represents a photon with external momentum of  $p_2$  and reference momentum of  $p_3$ . In principle,  $J_{\mu}$  is a gauge-dependent quantity. As we ignore the mass of leptons, it could be considered as a gauge-independent quantity in our proof.




FIG. 2: The current  $J_{\mu}$  of  $V \to \ell^- \ell^+$ .

Based on these equations, we could decompose amplitudes of  $H \to V \gamma \to \ell \ell \gamma$  as

$$\mathcal{M}(2^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, 4^{-}_{\gamma}) = -i\Gamma^{\mu\nu}(p_{2} + p_{3}, p_{4})J^{+}_{\mu}(s_{23})\epsilon^{-}_{\nu}(4, q)$$

$$= -i\Gamma^{\mu\nu}(p_{2}, p_{4})f^{l}_{V}(s_{23})[23]\epsilon^{-}_{\mu}(2, 3)\epsilon^{-}_{\nu}(4, q)$$

$$-i\Gamma^{\mu\nu}(p_{3}, p_{4})f^{l}_{V}(s_{23})\langle 23\rangle\epsilon^{+}_{\mu}(3, 2)\epsilon^{-}_{\nu}(4, q)$$

$$= f^{l}_{V}(s_{23}) \times ([23]\mathcal{M}(2^{-}_{\gamma}, 4^{-}_{\gamma}) + \langle 23\rangle\mathcal{M}(3^{+}_{\gamma}, 4^{-}_{\gamma})), \qquad (16)$$

where Eq. (15) is inserted for the first term and Eq. (14) for the second. In the last step, the reference momenta of photons are different, which do not affect the form of  $\mathcal{M}(\gamma, \gamma)$  because the vertex  $\Gamma^{\mu\nu}$  satisfy Ward identity. The other helicity amplitudes of  $H \to V\gamma \to \ell\ell\gamma$  have similar decomposition. An illustrating diagram for Eq. (16) is shown in Fig. 3. Each amplitude of  $H \to V\gamma \to \ell\ell\gamma$  is composed of two amplitudes of  $H \to \gamma\gamma$ . It degenerate to one term because the amplitude of  $H \to \gamma\gamma$  with reverse helicities is equal to zero. So the CP violation phase keeps as a global phase in  $H \to V\gamma \to \ell\ell\gamma$  process.




FIG. 3: Decomposition of amplitudes of  $H \to V\gamma \to \ell \ell \gamma$ .

Next we prove that decomposition relation is also suitable for process  $H \to VV \to 4\ell$ . That is

$$\mathcal{M}(2_{\ell^{-}}^{-}, 3_{\ell^{+}}^{+}, 4_{\ell^{\prime-}}^{-}, 5_{\ell^{\prime+}}^{+})$$

$$= -i\Gamma^{\mu\nu}(p_{2} + p_{3}, p_{4} + p_{5})J_{\mu}^{+}(s_{23})J_{\mu}^{+}(s_{45})$$

$$= f_{V}^{l}(s_{23})f_{V}^{l}(s_{45}) \times ($$

$$-i\Gamma^{\mu\nu}(p_{2}, p_{4})[23][45]\epsilon^{-}(2, 3)\epsilon^{-}(4, 5)$$

$$-i\Gamma^{\mu\nu}(p_{2}, p_{5})[23]\langle 45\rangle\epsilon^{-}(2, 3)\epsilon^{+}(5, 4)$$

$$-i\Gamma^{\mu\nu}(p_{3}, p_{4})\langle 23\rangle[45]\epsilon^{+}(3, 2)\epsilon^{-}(4, 5)$$

$$-i\Gamma^{\mu\nu}(p_{3}, p_{5})\langle 23\rangle\langle 45\rangle\epsilon^{+}(3, 2)\epsilon^{+}(5, 4) ) \qquad (17)$$

$$= f_{V}^{l}(s_{23})f_{V}^{l}(s_{45}) \times ($$

$$[23][45]\mathcal{M}(2_{\gamma}^{-}, 4_{\gamma}^{-}) + [23]\langle 45\rangle\mathcal{M}(2_{\gamma}^{-}, 5_{\gamma}^{+})$$

$$+\langle 23\rangle[45]\mathcal{M}(3_{\gamma}^{+}, 4_{\gamma}^{-}) + \langle 23\rangle\langle 45\rangle\mathcal{M}(3_{\gamma}^{+}, 5_{\gamma}^{+}) ), \qquad (18)$$

where in Eq. (17) one term of  $H \to 4\ell$  decompose to four terms, in Eq. (18) four terms degenerate to two terms since reverse-sign  $H \to \gamma\gamma$  amplitudes are zero. The illustrating diagrams are

## **B.** *CP* violation phase in amplitudes

From the decomposition relations, we see that the amplitudes of  $H \to \gamma \gamma$  are basis for other amplitudes. Since  $\mathcal{M}(+, -) = \mathcal{M}(-, +) = 0$ , the left basis are  $\mathcal{M}(+, +)$  and  $\mathcal{M}(-, -)$ .




FIG. 4: Decomposition of amplitudes of  $H \to VV \to 4\ell$ .

CP violation phases are reverse in the two bases. In  $H \to \gamma\gamma$  and  $H \to V\gamma \to \ell\ell\gamma$  processes, the CP violation phase is a common phase in each amplitude. So generally speaking it is an unobservable phase if one doesn't consider interference between this amplitude and the background [32, 33]. In  $H \to 4\ell$  process, two bases coexist in each amplitude, thus the CP violation phase appear as a physical observable. Meanwhile, it means that the interference between CP-even term and CP-odd term exists at differential cross section level after squaring the amplitude. So the interference could be probed through kinematic angles [31, 34–36]. An obvious effect is a shift of azimuthal angle caused by the interference between CP-even and CP-odd term [39].

#### IV. BSM AMPLITUDES FROM ON-SHELL APPROACH

In on-shell approach, the amplitude is not derived from Lagrangian and Feynman rules, instead it is constructed directly from on-shell particle states. In this section firstly we introduce spinor variables for particles, secondly we show how amplitudes of  $H\gamma\gamma$  are represented and fixed, thirdly we get amplitudes of  $H\gamma\ell\ell$  and  $H4\ell$  through recursion relations.

#### A. Spinor variables

The right-handed and left-handed spinors in Eq. (6) have their two-component versions [37, 38]:

$$|i_{\alpha}\rangle \equiv \lambda_{i\alpha} \equiv u_{+}(p_{i}) \equiv |i^{+}\rangle, \quad |i^{\dot{\alpha}}] \equiv \tilde{\lambda}_{i}^{\dot{\alpha}} \equiv u_{-}(p_{i}) \equiv |i^{-}\rangle,$$
$$\langle i^{\alpha}| \equiv \lambda_{i}^{\alpha} \equiv \overline{u_{-}(p_{i})} \equiv \langle i^{-}|, \quad [i_{\dot{\alpha}}| \equiv \tilde{\lambda}_{i\dot{\alpha}} \equiv \overline{u_{+}(p_{i})} \equiv \langle i^{+}|, \qquad (19)$$

where the spinor indices can be raised or lowered by antisymmetric tensors  $\epsilon^{\alpha\beta}$  and  $\epsilon_{\alpha\beta}$ ,

$$\lambda^{\alpha} = \epsilon^{\alpha\beta} \lambda_{\beta}, \quad \lambda_{\alpha} = \epsilon_{\alpha\beta} \lambda^{\beta} .$$
 (20)

In this notation,

$$\langle ij \rangle \equiv \lambda_i^{\alpha} \lambda_{j\alpha}, \quad [ij] \equiv \tilde{\lambda}_{i\dot{\alpha}} \tilde{\lambda}_j^{\dot{\alpha}} .$$
 (21)

An on-shell momentum of a massless particle is represented as

$$p_{\alpha\dot{\alpha}} \equiv p_{\mu}\sigma^{\mu}_{\alpha\dot{\alpha}} = \lambda_{\alpha}\tilde{\lambda}_{\dot{\alpha}},\tag{22}$$

where  $\sigma^{\mu} = (1, \vec{\sigma})$  with  $\vec{\sigma}$  being the Pauli matrices.

## **B.** Amplitude of $H\gamma\gamma$

A general three point amplitude with one massive and two massless particle interaction is shown in Fig. 5 [25], where  $\alpha_i$ , i = 1, 2, ..., 2S are indicies of spinors, S represents spin of the massive particle,  $h_2$ ,  $h_3$  are helicities of two massless particles.




FIG. 5: A general one massive and two massless particle interaction. The subscript S represents spin of the massive particle,  $h_2$ ,  $h_3$  are helicities of two massless particles.

For amplitude of  $H\gamma\gamma$ , as the massive particle H is a scalar with zero spin, we don't need to take care about its spinors, which makes the formula much simpler. A general ansatz is [25, 40, 41]

$$\mathcal{M}_3(1_H, 2^{h_2}_{\gamma}, 3^{h_3}_{\gamma}) = e^{i\xi^{h_2, h_3}} \frac{g}{m^{h_2 + h_3 - 1}} [23]^{h_2 + h_3} , \qquad (23)$$

where  $\xi^{h_2,h_3}$  represents a heicity-related phase, g represents an overall coupling constant, m is the mass of the Higgs boson. As  $\langle 23 \rangle [32] = (p_2 + p_3)^2 = p_1^2 = m^2$ ,  $\langle 23 \rangle = \frac{m^2}{[32]}$ . The little group scaling [23, 24] requires  $h_2 + h_3 = 2h_2 = 2h_3$ , so  $\mathcal{M}(2^+_{\gamma}, 3^-_{\gamma}) = \mathcal{M}(2^-_{\gamma}, 3^+_{\gamma}) = 0$ . The non-zero amplitudes are only  $\mathcal{M}(2^+_{\gamma}, 3^+_{\gamma})$  and  $\mathcal{M}(2^-_{\gamma}, 3^-_{\gamma})$ . It doen't lose generality to require  $\xi^{+,+} = -\xi^{-,-} = \xi'$  since their equal part could be absorbed into the redefinition of g. Another CP violation assumption could be  $|\mathcal{M}(2^+_{\gamma}, 3^+_{\gamma})| \neq |\mathcal{M}(2^-_{\gamma}, 3^-_{\gamma})|$ , however it seems more weird than the different phase assumption, we neglect this possibility in our study. Then

$$\mathcal{M}_3(1_H, 2^+_{\gamma}, 3^+_{\gamma}) = e^{i\xi'} \frac{g}{m} [23]^2 , \qquad (24)$$

$$\mathcal{M}_3(1_H, 2^-_{\gamma}, 3^-_{\gamma}) = e^{-i\xi'} \frac{g}{m} \langle 23 \rangle^2 , \qquad (25)$$

which is equal to Eq. (7) as long as we require  $\frac{g}{m} = \frac{2c_{\gamma\gamma}^S}{v}$  and  $\xi' = \xi$ .

## C. Amplitudes of $H \to \gamma \ell \ell$




FIG. 6: Factorization of  $H \to \gamma \ell \ell$ . We take the mediate particle as  $\gamma$  for simplicity.

The amplitudes of  $H \to \gamma \ell \ell$  could be built from three point amplitudes by recursion relations. For the amplitude of  $H \to \gamma \ell \ell$ , a factorization way is  $H \to \gamma V, V \to \ell \ell$ . Figure 6 shows this factorization. The mediate particle is taken as  $\gamma$  to avoid amplitude of massive particles, its momentum is marked as "I". We shift momenta of the 2, 4 external particles according to BCFW recursion relation approach [42–44]. That is

$$|\hat{2}] = |2], \quad |\hat{4}] = |4] + z|2], \quad |\hat{4}\rangle = |4\rangle, \quad |\hat{2}\rangle = |2\rangle - z|4\rangle,$$
(26)

where z is a complex number, and shifted momenta are hatted.

The corresponding analytical formula are

$$\mathcal{M}(1_{H}, 2^{h_{2}}_{\ell^{-}}, 3^{h_{3}}_{\ell^{+}}, 4^{h_{4}}_{\gamma}) = P_{\gamma}(s_{23})\mathcal{M}(1_{H}, \hat{4}^{h_{4}}_{\gamma}, \hat{I}^{-}_{\gamma})\mathcal{M}(\hat{I}^{+}_{\gamma}, \hat{2}^{h_{2}}_{\ell^{-}}, 3^{h_{3}}_{\ell^{+}}) + P_{\gamma}(s_{23})\mathcal{M}(1_{H}, \hat{4}^{h_{4}}_{\gamma}, \hat{I}^{+}_{\gamma})\mathcal{M}(\hat{I}^{-}_{\gamma}, \hat{2}^{h_{2}}_{\ell^{-}}, 3^{h_{3}}_{\ell^{+}}) , \qquad (27)$$

where  $\hat{p}_I = p_1 + \hat{p}_4 = \hat{p}_2 + p_3$  is momentum of the mediate photon,  $P_{\gamma}(s_{23}) = 1/s_{23} = 1/(p_2 + p_3)^2$  is the propagator with unshifted momenta.

The helicity amplitudes of  $\gamma \ell^- \ell^+$  are three point amplitudes with massless particles, which are fully fixed by little group scaling and dimension analysis,

$$\mathcal{M}(1_{\gamma}^{-}, 2_{\ell^{-}}^{-}, 3_{\ell^{+}}^{+}) = \tilde{e} \frac{\langle 12 \rangle^{2}}{\langle 23 \rangle} , \qquad (28)$$

$$\mathcal{M}(1_{\gamma}^{-}, 2_{\ell^{-}}^{+}, 3_{\ell^{+}}^{-}) = \tilde{e} \frac{\langle 13 \rangle^{2}}{\langle 23 \rangle} , \qquad (29)$$

$$\mathcal{M}(1^+_{\gamma}, 2^-_{\ell^-}, 3^+_{\ell^+}) = \tilde{e} \frac{[13]^2}{[23]} , \qquad (30)$$

$$\mathcal{M}(1^+_{\gamma}, 2^+_{\ell^-}, 3^-_{\ell^+}) = \tilde{e} \frac{[12]^2}{[23]} , \qquad (31)$$

where  $\tilde{e} = -\sqrt{2}e$ , Eqs. (28)(29) correspond to [23] = 0 solution and Eqs. (30)(31) correspond to  $\langle 23 \rangle = 0$  solution.

After inserting Eq. (25) and Eq. (28) into Eq. (27), we get

$$\mathcal{M}(1_{H}, 2^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, 4^{-}_{\gamma}) = \tilde{e}P_{\gamma}(s_{23}) \times \frac{2c^{S}_{\gamma V}}{v} e^{-i\xi} \frac{\langle \hat{I}\hat{4} \rangle^{2} [\hat{I}3]^{2}}{[\hat{2}3]},$$
  
$$= \tilde{e}P_{\gamma}(s_{23}) \times \frac{2c^{S}_{\gamma V}}{v} e^{-i\xi} [23] \langle 24 \rangle^{2}, \qquad (32)$$

where the last equation is because

$$\langle \hat{4}\hat{I} \rangle [\hat{I}3] = \langle \hat{4}|\hat{p}_{I}|3] = \langle \hat{4}|\hat{p}_{2} + p_{3}|3] = \langle \hat{4}|\hat{p}_{2}|3] = \langle \hat{4}\hat{2} \rangle [\hat{2}3] = \langle 42 \rangle [23] .$$
(33)

So Eq. (32) is the same formula as the one derived in SMEFT (see Eq. (8)). It is worthy to notice that because  $P_{\gamma}(s_{23}) = \frac{1}{\langle 23 \rangle [32]}$ , Eq. (32) is proportional to  $\frac{\langle 24 \rangle^2}{\langle 23 \rangle}$  and thus has a singularity when  $\langle 23 \rangle = 0$ .

If we take the propagator V as a Z boson, we should consider a  $H\gamma Z$  amplitude together with a  $Z\gamma\gamma$  amplitude. The  $H\gamma Z$  amplitude is an amplitude with two massive one massless particles, and the  $Z\gamma\gamma$  amplitude is an amplitude with one massive two massless particles. They are complex than the  $H\gamma\gamma$  amplitude since the spin of Z is 1. These two amplitudes should use bolded spinor variables [25, 40], we leave this bolded formula in future work.

## **D.** Amplitudes of $H \to 4\ell$

The amplitude of  $H \to 4\ell$  is a five point amplitude, we could factorize it into two parts: a four point amplitude plus a three point amplitude. Each amplitude split into four parts as shown in Fig. 7.




FIG. 7: Factorization of  $H \rightarrow 4\ell$ . The external legs are arranged in clockwise order.

In formula, it is

$$\mathcal{M}_{5}(1_{H}, 2^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, 4^{-}_{\ell^{\prime-}}, 5^{+}_{\ell^{\prime+}}) = P_{\gamma}(s_{23})\mathcal{M}(1_{H}, \hat{4}^{-}_{\ell^{\prime-}}, 5^{+}_{\ell^{\prime+}}, \hat{I}^{-}_{\gamma})\mathcal{M}(\hat{I}^{+}_{\gamma}, \hat{2}^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}) + P_{\gamma}(s_{23})\mathcal{M}(1_{H}, \hat{4}^{-}_{\ell^{\prime-}}, 5^{+}_{\ell^{\prime+}}, \hat{I}^{+}_{\gamma})\mathcal{M}(\hat{I}^{-}_{\gamma}, \hat{2}^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}) + P_{\gamma}(s_{45})\mathcal{M}(1_{H}, \hat{2}^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, \hat{I}^{+}_{\gamma})\mathcal{M}(\hat{I}^{-}_{\gamma}, \hat{4}^{-}_{\ell^{\prime-}}, 5^{+}_{\ell^{\prime+}}) + P_{\gamma}(s_{45})\mathcal{M}(1_{H}, \hat{2}^{-}_{\ell^{-}}, 3^{+}_{\ell^{+}}, \hat{I}^{-}_{\gamma})\mathcal{M}(\hat{I}^{+}_{\gamma}, \hat{4}^{-}_{\ell^{\prime-}}, 5^{+}_{\ell^{\prime+}}),$$
(34)

which corresponds to diagram A, B, C, D respectively. Diagram A and B correspond to (1,4,5) + (2,3) factorization, Diagram C and D correspond to (1,2,3) + (4,5) factorization. We assumed  $\ell \neq \ell'$  for generality, so the factorizations of (1,2,5) + (3,4) and (1,3,4) + (2,5) are absent to keep flavor symmetry. Next we calculate these four diagrams separately.

The formula for diagram A is

$$P_{\gamma}(s_{23})\mathcal{M}(1_{H}, \hat{4}_{\ell'^{-}}, 5_{\ell'^{+}}^{+}, \hat{I}_{\gamma}^{-})\mathcal{M}(\hat{I}_{\gamma}^{+}, \hat{2}_{\ell^{-}}^{-}, 3_{\ell^{+}}^{+})$$

$$= \frac{2c_{\gamma\gamma}^{S}}{v}e^{-i\xi}P_{\gamma}(s_{23})P_{\gamma}(s_{\hat{4}5})[\hat{4}5]\langle\hat{4}\hat{I}\rangle^{2}\frac{[\hat{I}3]^{2}}{[\hat{2}3]}$$

$$= \frac{2c_{\gamma\gamma}^{S}}{v}e^{-i\xi}P_{\gamma}(s_{23})P_{\gamma}(s_{45})[\hat{4}5][23]\langle 24\rangle^{2}, \qquad (35)$$

where in the last step we have used

$$\langle \hat{4}\hat{I} \rangle [\hat{I}3] = \langle \hat{4}|\hat{p}_2 + p_3|3] = \langle \hat{4}|\hat{p}_2|3] = \langle \hat{4}\hat{2} \rangle [\hat{2}3] = \langle 42 \rangle [23]$$
 (36)

as in Eq. (33) and

$$P_{\gamma}(s_{\hat{4}5})[\hat{4}5] = \frac{-1}{\langle \hat{4}5 \rangle [\hat{4}5]} [\hat{4}5] = \frac{-1}{\langle 45 \rangle [45]} [45] = P_{\gamma}(s_{45})[45] , \qquad (37)$$

 $\langle \hat{2}3 \rangle = 0$  is chosen for three-point amplitude, which is also required in diagram B.

The formula for diagram B is

$$P_{\gamma}(s_{23})\mathcal{M}(1_{H}, \hat{4}_{\ell'^{-}}, 5_{\ell'^{+}}^{+}, \hat{I}_{\gamma}^{+})\mathcal{M}(\hat{I}_{\gamma}^{-}, \hat{2}_{\ell^{-}}^{-}, 3_{\ell^{+}}^{+})$$

$$= \frac{2c_{\gamma\gamma}^{S}}{v}e^{i\xi}P_{\gamma}(s_{23})P_{\gamma}(s_{45})\langle\hat{4}5\rangle[5\hat{I}]^{2} \times \frac{\langle\hat{I}\hat{2}\rangle^{2}}{\langle\hat{2}3\rangle}$$

$$= \frac{2c_{\gamma\gamma}^{S}}{v}e^{i\xi}P_{\gamma}(s_{23})P_{\gamma}(s_{45})\langle\hat{4}5\rangle[5\hat{I}]^{2} \times 0$$

$$= 0, \qquad (38)$$

where the 3-point amplitude is equal to zero because  $\langle \hat{2}3 \rangle = \langle \hat{I}\hat{2} \rangle = 0$  .

The formula for diagram C is

$$P_{\gamma}(s_{45})\mathcal{M}(1_{H}, \hat{2}_{\ell^{-}}^{-}, 3_{\ell^{+}}^{+}, \hat{I}_{\gamma}^{+})\mathcal{M}(\hat{I}_{\gamma}^{-}, \hat{4}_{\ell^{\prime-}}^{-}, 5_{\ell^{\prime+}}^{+}) \\ = \frac{2c_{\gamma\gamma}^{S}}{v}e^{i\xi}P_{\gamma}(s_{45})P_{\gamma}(s_{23})\langle\hat{2}3\rangle[3\hat{I}]^{2} \times \frac{\langle\hat{I}\hat{4}\rangle^{2}}{\langle\hat{4}5\rangle} \\ = \frac{2c_{\gamma\gamma}^{S}}{v}e^{i\xi}P_{\gamma}(s_{45})P_{\gamma}(s_{23})\langle\hat{2}3\rangle\langle 45\rangle[35]^{2} , \qquad (39)$$

where

$$[3\hat{I}]\langle\hat{I}\hat{4}\rangle = [3\hat{5}]\langle\hat{5}\hat{4}\rangle \tag{40}$$

and

$$P_{\gamma}(s_{\hat{2}3})[\hat{2}3] = \frac{-1}{\langle \hat{2}3 \rangle [\hat{2}3]} [\hat{2}3] = \frac{-1}{\langle 23 \rangle [23]} [23] = P_{\gamma}(s_{23})[23]$$
(41)

are used.  $[\hat{4}5] = 0$  is required.

The formula for diagram D is

$$P_{\gamma}(s_{45})\mathcal{M}(1_{H}, \hat{2}_{\ell^{-}}, 3_{\ell^{+}}^{+}, \hat{I}_{\gamma}^{-})\mathcal{M}(\hat{I}_{\gamma}^{+}, \hat{4}_{\ell^{\prime-}}^{-}, 5_{\ell^{\prime+}}^{+})$$

$$= \frac{2c_{\gamma\gamma}^{S}}{v}e^{-i\xi}P_{\gamma}(s_{45})P_{\gamma}(s_{\hat{2}3})[\hat{2}3]\langle\hat{2}\hat{I}\rangle^{2}\frac{[\hat{I}5]^{2}}{[\hat{4}5]}$$

$$= 0 , \qquad (42)$$

where  $[\hat{4}5] = [\hat{I}5] = 0$  makes the three point amplitude zero.

After summing up the results of four parts, that is adding Eq.s (35)(38)(39)(42) together, we get

$$\mathcal{M}(1_H, 2^-_{\ell^-}, 3^+_{\ell^+}, 4^-_{\ell^{\prime-}}, 5^+_{\ell^{\prime+}}) = \frac{2c^S_{\gamma\gamma}}{v} e^{-i\xi} P_{\gamma}(s_{23}) P_{\gamma}(s_{45}) [45] [23] \langle 24 \rangle^2 , \qquad (43)$$

$$+\frac{2c_{\gamma\gamma}^{S}}{v}e^{i\xi}P_{\gamma}(s_{45})P_{\gamma}(s_{23})\langle 23\rangle\langle 45\rangle[35]^{2},\qquad(44)$$

which has the same form as the one derived in SMEFT (see Eq. (9)). So we get a consistent result from the on shell approach. Boundary contributions here are supposed to be zero.

## V. SUMMARY AND DISCUSSION

The helicity amplitudes of  $H \to \gamma\gamma$ ,  $H \to \ell\ell\gamma$  and  $H \to 4\ell$  in beyond Standard Model are analyzed in two ways. One way is under field theory framework, we decompose the amplitudes of  $H \to \ell\ell\gamma$  and  $H \to 4\ell$  into the amplitudes of  $H \to \gamma\gamma$ . In the decomposition relations, we explain why CP violation phase is a global phase in amplitudes of  $H \to \gamma\gamma$ and  $H \to \ell\ell\gamma$  while it is not in amplitudes of  $H \to 4\ell$ . The other way is through the on-shell scattering amplitude approach. The 3-point amplitude of  $H\gamma\gamma$  is the start point, then the 4-point amplitude of  $H\ell\ell\gamma$  and the 5-point amplitude  $H4\ell$  are obtained through recursion relations. We get a consistent result in two ways.

Nevertheless, there are some differences. The decomposition relations are suitable for both massless and massive mediate particles, but it needs a precondition that the HVVvertex is bilinear to the momenta of vector bosons. In our proof this is inherited from the dimension-six operators in SMEFT. So for other Lorentz structures of HVV vertex (e.g. from dimension-eight operators in SMEFT) the decomposition relations are not valid. By contrast, in on-shell scattering amplitude approach the recursion relations for massless mediate particle are obtained without such precondition. That is because a bilinear  $H\gamma\gamma$ vertex is automatically fulfilled to all orders due to their particle properties. How about recursion relations for massive mediate particle? The decomposition relation tell us that if we could pick out its bilinear HVV component, we could get a similar recursion relation as massless mediate particle. Even though in on-shell approach massive amplitudes are complex and its recursion relations are difficult, we get an illuminating possibility.

#### ACKNOWLEDGMENTS

We are grateful to Chih-Hao Fu, Kang Zhou for useful discussion about BCFW recursion relations. X.W. thank Bo Feng for scattering amplitude school 2021 in Hangzhou, China. The work is supported by the National Natural Science Foundation of China under Grant No. 11847168, the Fundamental Research Funds for the Central Universities of China under Grant No. GK202003018, and the Natural Science Foundation of Shannxi Province, China (2019JM-431, 2019JQ-739).

- [1] N. Cabibbo, "Unitary Symmetry and Leptonic Decays," Phys. Rev. Lett. 10 (1963) 531–533.
- M. Kobayashi and T. Maskawa, "CP-Violation in the Renormalizable Theory of Weak Interaction," *Progress of Theoretical Physics* 49 (02, 1973) 652-657, https://academic.oup.com/ptp/article-pdf/49/2/652/5257692/49-2-652.pdf. https://doi.org/10.1143/PTP.49.652.
- [3] G. 't Hooft, "Symmetry Breaking Through Bell-Jackiw Anomalies," Phys. Rev. Lett. 37 (1976) 8–11.
- [4] G. 't Hooft, "Computation of the Quantum Effects Due to a Four-Dimensional Pseudoparticle," *Phys. Rev. D* 14 (1976) 3432–3450. [Erratum: Phys.Rev.D 18, 2199 (1978)].
- [5] R. D. Peccei and H. R. Quinn, "CP Conservation in the Presence of Instantons," *Phys. Rev. Lett.* 38 (1977) 1440–1443.
- [6] R. S. Chivukula and H. Georgi, "Composite Technicolor Standard Model," *Phys. Lett. B* 188 (1987) 99–104.
- M. E. Peskin and D. V. Schroeder, An Introduction to quantum field theory, P.721-724.
   Addison-Wesley, Reading, USA, 1995.
- [8] R. D. Peccei, "The Strong CP problem and axions," Lect. Notes Phys. 741 (2008) 3-17, arXiv:hep-ph/0607268.
- [9] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, "Theory and phenomenology of two-Higgs-doublet models," *Phys. Rept.* 516 (2012) 1–102, arXiv:1106.0034 [hep-ph].
- [10] A. Djouadi, "The Anatomy of electro-weak symmetry breaking. II. The Higgs bosons in the

minimal supersymmetric model," Phys. Rept. 459 (2008) 1-241, arXiv:hep-ph/0503173.

- [11] G. Panico and A. Wulzer, *The Composite Nambu-Goldstone Higgs*, vol. 913. Springer, 2016.
   arXiv:1506.01961 [hep-ph].
- [12] G. Steigman, "Observational tests of antimatter cosmologies," Ann. Rev. Astron. Astrophys. 14 (1976) 339–372.
- [13] G. Steigman, "When Clusters Collide: Constraints On Antimatter On The Largest Scales," JCAP 10 (2008) 001, arXiv:0808.1122 [astro-ph].
- [14] A. G. Cohen and D. B. Kaplan, "Thermodynamic Generation of the Baryon Asymmetry," *Phys. Lett. B* 199 (1987) 251–258.
- [15] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, "On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe," *Phys. Lett. B* 155 (1985) 36.
- [16] F. R. Klinkhamer and N. S. Manton, "A Saddle Point Solution in the Weinberg-Salam Theory," Phys. Rev. D 30 (1984) 2212.
- [17] M. B. Gavela, P. Hernandez, J. Orloff, and O. Pene, "Standard model CP violation and baryon asymmetry," *Mod. Phys. Lett. A* 9 (1994) 795–810, arXiv:hep-ph/9312215.
- M. B. Gavela, P. Hernandez, J. Orloff, O. Pene, and C. Quimbay, "Standard model CP violation and baryon asymmetry. Part 2: Finite temperature," Nucl. Phys. B 430 (1994) 382-426, arXiv:hep-ph/9406289.
- [19] P. Huet and E. Sather, "Electroweak baryogenesis and standard model CP violation," *Phys. Rev. D* 51 (1995) 379-394, arXiv:hep-ph/9404302.
- [20] W. Buchmuller and D. Wyler, "Effective Lagrangian Analysis of New Interactions and Flavor Conservation," Nucl. Phys. B268 (1986) 621–653.
- [21] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, "Dimension-Six Terms in the Standard Model Lagrangian," JHEP 10 (2010) 085, arXiv:1008.4884 [hep-ph].
- [22] I. Brivio and M. Trott, "The Standard Model as an Effective Field Theory," *Phys. Rept.* 793 (2019) 1–98, arXiv:1706.08945 [hep-ph].
- [23] H. Elvang and Y.-t. Huang, Scattering Amplitudes in Gauge Theory and Gravity. Cambridge University Press, 4, 2015.
- [24] C. Cheung, TASI Lectures on Scattering Amplitudes, pp. 571-623. 2018. arXiv:1708.03872 [hep-ph].
- [25] N. Arkani-Hamed, T.-C. Huang, and Y.-t. Huang, "Scattering Amplitudes For All Masses

and Spins," arXiv:1709.04891 [hep-th].

- [26] CMS Collaboration, S. Chatrchyan et al., "Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC," Phys. Lett. B 716 (2012) 30-61, arXiv:1207.7235 [hep-ex].
- [27] ATLAS Collaboration, G. Aad *et al.*, "Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC," *Phys. Lett. B* 716 (2012) 1–29, arXiv:1207.7214 [hep-ex].
- [28] J. M. Campbell, R. K. Ellis, and C. Williams, "Bounding the Higgs width at the LHC using full analytic results for gg → e<sup>-</sup>e<sup>+</sup>μ<sup>-</sup>μ<sup>+</sup>," JHEP 04 (2014) 060, arXiv:1311.3589 [hep-ph].
- [29] F. Caola and K. Melnikov, "Constraining the Higgs boson width with ZZ production at the LHC," Phys. Rev. D88 (2013) 054024, arXiv:1307.4935 [hep-ph].
- [30] L. J. Dixon and Y. Li, "Bounding the Higgs Boson Width Through Interferometry," Phys. Rev. Lett. 111 (2013) 111802, arXiv:1305.3854 [hep-ph].
- [31] CMS Collaboration, A. M. Sirunyan *et al.*, "Measurements of the Higgs boson width and anomalous *HVV* couplings from on-shell and off-shell production in the four-lepton final state," *Phys. Rev. D* 99 (2019) no. 11, 112003, arXiv:1901.00174 [hep-ex].
- [32] X. Wan and Y.-K. Wang, "Probing the CP violating Hγγ coupling using interferometry," Chin. Phys. C 43 (2019) no. 7, 073101, arXiv:1712.00267 [hep-ph].
- [33] X. Chen, G. Li, and X. Wan, "Probe CP violation in  $H \to \gamma Z$  through forward-backward asymmetry," *Phys. Rev.* D96 (2017) no. 5, 055023, arXiv:1705.01254 [hep-ph].
- [34] H.-R. He, X. Wan, and Y.-K. Wang, "Anomalous H → ZZ → 4ℓ decay and its interference effects on gluon-gluon contribution at the LHC," Chin. Phys. C 44 (2020) no. 12, 123101, arXiv:1902.04756 [hep-ph].
- [35] CMS Collaboration, S. Chatrchyan *et al.*, "Measurement of the properties of a Higgs boson in the four-lepton final state," *Phys. Rev.* D89 (2014) no. 9, 092007, arXiv:1312.5353
   [hep-ex].
- [36] I. Anderson et al., "Constraining Anomalous HVV Interactions at Proton and Lepton Colliders," Phys. Rev. D 89 (2014) no. 3, 035007, arXiv:1309.4819 [hep-ph].
- [37] L. J. Dixon, "Calculating scattering amplitudes efficiently," in QCD and beyond. Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics, TASI-95, Boulder,

USA, June 4-30, 1995, pp. 539-584. 1996. arXiv:hep-ph/9601359 [hep-ph]. http://www-public.slac.stanford.edu/sciDoc/docMeta.aspx?slacPubNumber=SLAC-PUB-7106.

- [38] L. J. Dixon, "A brief introduction to modern amplitude methods," in Theoretical Advanced Study Institute in Elementary Particle Physics: Particle Physics: The Higgs Boson and Beyond, pp. 31-67. 2014. arXiv:1310.5353 [hep-ph].
- [39] Y.-S. Lu, X. Wan, Y.-K. Wang, and X.-Y. You, "Study of HZZ anomalous coupling by differential distributions," in preparation.
- [40] Y. Shadmi and Y. Weiss, "Effective Field Theory Amplitudes the On-Shell Way: Scalar and Vector Couplings to Gluons," JHEP 02 (2019) 165, arXiv:1809.09644 [hep-ph].
- [41] L. J. Dixon, E. W. N. Glover, and V. V. Khoze, "MHV rules for Higgs plus multi-gluon amplitudes," JHEP 12 (2004) 015, arXiv:hep-th/0411092.
- [42] R. Britto, F. Cachazo, and B. Feng, "New recursion relations for tree amplitudes of gluons," Nucl. Phys. B 715 (2005) 499-522, arXiv:hep-th/0412308.
- [43] R. Britto, F. Cachazo, B. Feng, and E. Witten, "Direct proof of tree-level recursion relation in Yang-Mills theory," *Phys. Rev. Lett.* 94 (2005) 181602, arXiv:hep-th/0501052.
- [44] B. Feng and M. Luo, "An Introduction to On-shell Recursion Relations," Front. Phys. (Beijing) 7 (2012) 533-575, arXiv:1111.5759 [hep-th].