

Approximating Length-Restricted Means under Dynamic Time Warping

Maike Buchin ✉ 

Faculty of Informatics, Ruhr University Bochum, Germany

Anne Driemel ✉

Hausdorff Center for Mathematics, Germany

University of Bonn, Germany

Koen van Greevenbroek ✉ 

Department of Computer Science, UiT The Arctic University of Norway, Tromsø, Norway

Ioannis Psarros ✉ 

Athena Research Center, Greece.

Dennis Rohde ✉ 

Faculty of Informatics, Ruhr University Bochum, Germany

Abstract

We study variants of the mean problem under the p -Dynamic Time Warping (p -DTW) distance, a popular and robust distance measure for sequential data. In our setting we are given a set of finite point sequences over an arbitrary metric space and we want to compute a mean point sequence of given length that minimizes the sum of p -DTW distances, each raised to the q^{th} power, between the input sequences and the mean sequence. In general, the problem is NP-hard and known not to be fixed-parameter tractable in the number of sequences. On the positive side, we show that restricting the length of the mean sequence significantly reduces the hardness of the problem. We give an exact algorithm running in polynomial time for constant-length means. We explore various approximation algorithms that provide a trade-off between the approximation factor and the running time. Our approximation algorithms have a running time with only linear dependency on the number of input sequences. In addition, we use our mean algorithms to obtain clustering algorithms with theoretical guarantees.

2012 ACM Subject Classification Theory of computation → Design and analysis of algorithms; Mathematics of computing → Probabilistic algorithms; Theory of computation → Computational geometry

Keywords and phrases Dynamic Time Warping, Clustering, Time Series Averaging, Sample Mean

Funding *Ioannis Psarros*: The author was partially supported by the European Union’s Horizon 2020 Research and Innovation programme, under the grant agreement No. 957345: “MORE”. Part of this work was done while the author was a postdoctoral researcher at the Hausdorff Center for Mathematics and the University of Bonn, Germany.

1 Introduction

The p -Dynamic Time Warping distance – in short p -DTW – is a popular distance measure for temporal data sequences, which has been studied and applied extensively over the past decades. While it was first applied to speech recognition [30], it also showed effective for other kinds of sequential data and now there is a broad variety of applications in numerous domains, cf. [4, 13, 26, 1, 37, 25, 21]. Its particular strength is the ability to handle differences in the length and in the temporal properties (e.g. phase or sampling rates) of the data. Furthermore, it is not sensitive to outliers in the sequences, e.g. from measurement errors or noise. It is based on monotonic alignments of the sequences, i.e., every element of the first sequence is paired to an element of the second sequence in a monotonic fashion (along the temporal axis). To compensate differences in length, samplings rates or in phase, several elements of the first sequence can be paired to a single element of the second sequence and vice versa. Such an alignment is called a warping and the p -DTW is the p^{th} root of the sum of distances, each raised to the p^{th} power, between all pairs of elements determined by an optimal warping, i.e., a warping that minimizes said quantity. The distances between elements are determined by an underlying space, which itself is determined by the application at hand. The p -DTW can be computed by a dynamic program with running time quadratic in the lengths of the given sequences, and it cannot be computed in strongly subquadratic time unless the Exponential Time Hypothesis is false [8, 2]. Apart from its apparent benefits, p -DTW has the drawback that it is not a metric since it does not fulfill the identity of indiscernibles nor the triangle inequality. This rules out a wealth of techniques that were developed for proper metrics.

In this work, we consider the problem of computing a mean under p -DTW. Here, we are given a finite set of n point sequences over an arbitrary metric space, each of complexity, i.e., the number of elements of the sequence, bounded by a number m and we want to compute a point sequence (the mean) that minimizes the sum of p -DTW distances, each raised to the q^{th} power, between the given sequences and the mean.¹ We call this problem the *unrestricted* (p, q) -mean problem. It is known to be NP-hard [12, 10] and all known algorithms that solve it either suffer from exponential running time, only work for binary alphabets, or are of heuristic nature [28, 6, 32].

We show that when we restrict the complexity of the mean to be bounded by some constant ℓ – we call this the *restricted* (p, q) -mean problem – the problem becomes tractable, i.e., there exist polynomial time approximation algorithms. This restriction also comes with a practical motivation, i.e., to suppress overfitting, see also the discussion in [5].

1.1 Related Work

Among many practical approaches for the problem of computing a mean, one very influential heuristic is the DTW Barycentric Average (DBA) method, as formalized by Petitjean, Ketterlin and Gançarski [28]. The core idea behind DBA is a Lloyd’s style (k -means) iterative strategy, which has been rediscovered many times for this problem in the past (see e.g. [29, 3, 18]). DBA iteratively improves the solution as follows: given a candidate average sequence $c = (c_1, \dots, c_\ell)$, it first computes the warpings between c and all input sequences, and then given each set of input vertices S_i matched with the same vertex c_i , it substitutes c_i with the mean of S_i . DBA has inspired many recent solutions that are successful in practice [33, 24, 20, 15, 27]. However, it does not give any guarantees. Just like the k -means

¹ For $q = 1$ this is an adaption of the Euclidean median and for $q = 2$ an adaption of the Euclidean mean.

algorithm, it may even converge to a local optimum that is arbitrarily far from the global optimum in terms of the (p, q) -mean target function.

There are few results in the literature with formal guarantees on the running time or the quality of the solution. Brill et al. [6, 7] presented an algorithm for solving the unrestricted $(2, 2)$ -mean problem defined over \mathbb{Q} with the Euclidean distance, with an asymptotic bound on the time complexity. Their algorithm is based on dynamic programming, and computes the (unrestricted) $(2, 2)$ -mean, in time $O(m^{2n+1}2^n n)$. The algorithm can be slightly modified, to compute a restricted $(2, 2)$ -mean. Brill et al. [6, 7] also show that the unrestricted $(2, 2)$ -mean problem defined over $\{0, 1\}$ with the Euclidean distance can be solved in $O(nm^3)$ time. This was later improved by Schaar, Froese and Niedermeier [32] to $O(nm^{1.87})$ time.

All previous hardness results concern the exact computation of the (p, q) -mean. Bulteau, Froese and Niedermeier [12] proved that the $(2, 2)$ -mean problem defined over \mathbb{Q} with the Euclidean distance is NP-hard and W[1]-hard with the number of input sequences n as the parameter. Moreover, they show that the problem cannot be solved in time $O(f(n)) \cdot m^{o(n)}$ for any computable function f unless the Exponential Time Hypothesis (ETH) fails. Buchin, Driemel and Struijs [10] presented an alternative proof of the above statements, which more generally applies to the unrestricted (p, q) -mean problem for any $p, q \in \mathbb{N}$.

1.2 Overview of Results

In this section we give an overview of our results.² In Section 2.1, we present an exact algorithm with polynomial running time for the problem of computing a restricted 2-mean in Euclidean space. Our approach is based on a decomposition of the solution space by an arrangement of polynomial surfaces such that each cell corresponds to a set of means with uniquely defined optimal warpings to all input sequences. The algorithm has running time $(nm)^{2^{O(d\ell)}}$, where the input are n sequences of m points in \mathbb{R}^d . Note that the running time is doubly-exponential in ℓ and d . In the remainder of the paper, our goal is to improve upon this with the help of approximation and randomization techniques. We will show that linear dependency n and singly-exponential dependency on ℓ and d are possible.

In Section 2.2, we present a randomized constant-factor approximation algorithm for the restricted p -mean problem that works for sequences from any fixed metric space. As such, this result is applicable to the classical median problem under the 1-DTW distance and the classical mean problem under the 2-DTW distance. The main idea is to uniformly sample from the union of points of all given sequences, then enumerate all sequences of complexity ℓ from the sampled vertices and return the sequence with the lowest cost. We also show how to derandomize the algorithm.

In Section 2.4 we present a $(1 + \varepsilon)$ -approximation algorithm for the restricted $(p, 1)$ -mean problem. This algorithm is based on an exhaustive search over a carefully constructed set of candidate means. The crucial ingredients for this are presented in Section 2.3: an efficient approximation algorithm for simplification under p -DTW and a weak triangle inequality for p -DTW. The result holds for the important case of sequences that stem from a Euclidean space. A nice property of this result is that it provides a complete trade-off between the approximation factor and the running time.

Finally, in Section 3, we briefly discuss an application of the newly developed techniques to the problem of clustering for p -DTW distances. In particular, we can use the random sampling

² An earlier version of this manuscript claimed hardness of approximation for the problem of computing the mean under dynamic time warping. However, the proof turned out to be flawed. We leave it as an open problem to show hardness of approximation for this problem.

techniques developed in Sections 2.2 and 2.3, in combination with a known algorithmic scheme that reduces the computation of k -medians to a problem of computing 1-median candidates for an arbitrary subset of the input set.

1.3 Preliminaries

In the following $d \in \mathbb{N}$ is an arbitrary constant. For $n \in \mathbb{N}$ we define $[n] = \{1, \dots, n\}$. Let $\mathcal{X} = (X, \rho)$ be a metric space. For $x \in X$ and $r \in \mathbb{R}_{\geq 0}$ we denote by $B(x, r) = \{y \in X \mid \rho(x, y) \leq r\}$ the ball of radius r centered at x . We define sequences of points over \mathcal{X} .

► **Definition 1.** A point sequence over \mathcal{X} is a tuple $(\sigma_1, \dots, \sigma_m) \in X^m$, where $m \in \mathbb{N}_{>1}$ is called its complexity, denoted by $|\sigma|$, and $\sigma_1, \dots, \sigma_m$ are called its vertices.

By $X^* = \bigcup_{i=1}^{\infty} X^i$ we define the set of all point sequences over \mathcal{X} and by $X^{\leq m} = \bigcup_{i=1}^m X^i$ we define the subset of point sequences of complexity at most m . The concatenation of a point sequence $\pi = (\pi_1, \dots, \pi_m)$ with a sequence $\tau = (\tau_1, \dots, \tau_m)$ is denoted by $\pi \oplus \tau$ and is defined as the point sequence $(\pi_1, \dots, \pi_m, \tau_1, \dots, \tau_m)$. We define the p -Dynamic Time Warping distance.

► **Definition 2.** For $m_1, m_2 \in \mathbb{N}_{>1}$, let \mathcal{W}_{m_1, m_2} denote the set of all (m_1, m_2) -warpings, that is, the set of all sequences $(i_1, j_1), \dots, (i_n, j_n)$ with

- $i_1 = j_1 = 1, i_n = m_1, j_n = m_2$ and
- $(i_k - i_{k-1}, j_k - j_{k-1}) \in \{(0, 1), (1, 0), (1, 1)\}$ for each $k \in \{2, \dots, n\}$.

For $p \in [1, \infty)$ and two point sequences $\sigma = (\sigma_1, \dots, \sigma_{m_1}) \in X^{m_1}, \tau = (\tau_1, \dots, \tau_{m_2}) \in X^{m_2}$ the p -Dynamic Time Warping distance is defined as

$$d_{\text{DTW}_p}(\sigma, \tau) = \min_{W \in \mathcal{W}_{m_1, m_2}} \left(\sum_{(i,j) \in W} \rho(\sigma_i, \tau_j)^p \right)^{1/p}.$$

Here we assume that $\rho(\cdot, \cdot)$ can be evaluated in constant time.³ Let $\sigma = (\sigma_1, \dots, \sigma_{m_1}), \tau = (\tau_1, \dots, \tau_{m_2})$ be point sequences over \mathcal{X} of complexity m_1 and m_2 . We call a warping $W \in \arg \min_{W \in \mathcal{W}_{m_1, m_2}} \left(\sum_{(i,j) \in W} \rho(\sigma_i, \tau_j)^p \right)^{1/p}$ an optimal p -warping between σ and τ .

► **Definition 3.** The restricted (p, q) -mean problem is defined as follows, where $\ell \in \mathbb{N}_{>1}$ and $p, q \in [1, \infty)$ are fixed (constant) parameters of the problem: given a set $T = \{\tau_1, \dots, \tau_n\} \subseteq X^{\leq m}$ of point sequences, compute a point sequence $c \in X^{\leq \ell}$, such that $\text{cost}_p^q(T, c) = \sum_{i=1}^n d_{\text{DTW}_p}(c, \tau_i)^q$ is minimal.

If p is clear from the context, we drop it from our notation. By the *unrestricted* (p, q) -mean problem we define the problem that is similar to the restricted (p, q) -mean problem with the only difference that we compute a mean sequence $c \in X^*$. We mainly study the case that $p = q$. Therefore, we shorthand call the restricted, respectively unrestricted, (p, p) -mean problem the restricted, respectively unrestricted, p -mean problem. We emphasize that these problems are prevalent in the literature.

³ This restriction is only for the sake of simplicity of presentation. Our results can be easily extended to metric spaces that do not have a constant-time distance oracle.

2 Tractability of the Restricted Mean Problem

We study exact and approximation algorithms for the restricted p -mean and $(p, 1)$ -mean problems.

2.1 Exact Computation of a Restricted 2-Mean in Euclidean Space

In the following, we make use of a simplified structure of the solution space, which holds in case $p = q$. This is captured in the notion of sections, which we define as follows.

► **Definition 4** (sections). *Let $T = \{\tau_1, \dots, \tau_n\} \subseteq X^{\leq m}$ be a set of point sequences and $c = (c_1, \dots, c_\ell) \in X^\ell$ be a point sequence. For $i \in [n]$ and $p \in [1, \infty)$, let W_i be an optimal p -warping between c and τ_i . For $j \in [\ell]$ we define the j^{th} section of c with respect to T (and W_1, \dots, W_n) as follows: $S_j(c, T, W_1, \dots, W_n) = \{\tau_{i,k} \mid i \in [n], (j, k) \in W_i\}$, where $\tau_{i,k}$ is the k^{th} vertex of τ_i .*

If T is clear from the context, we omit it from the notation. Also, we will always omit W_1, \dots, W_n from the notation, because the specific choice of optimal p -warplings is not of interest. We will then write $S_j^p(T, c)$ to clarify that the sections are defined with respect to optimal p -warplings. An immediate consequence of this definition is the following identity:

$$\text{cost}_p^p(T, c) = \sum_{j=1}^{\ell} \sum_{v \in S_j^p(c, T)} \rho(c_j, v)^p,$$

where ℓ denotes the complexity of c .

A central observation is that the vertices of an optimal restricted p -mean $c = (c_1, \dots, c_{\ell'})$ must minimize the sum of distances, each raised to the p^{th} power, to the vertices in their section, i.e., for all $j \in [\ell']$: $c_j \in \arg \min_{w \in X} \sum_{v \in S_j^p(c, T)} \rho(w, v)^p$. Using this, we obtain the following result.

► **Theorem 5.** *There exists an algorithm that, given a set $T \subset (\mathbb{Q}^d)^{\leq m}$ of n point sequences, computes an optimal restricted 2-mean (defined over the Euclidean distance) in time $(nm)^{2^{O(d\ell)}}$.*

The proof is deferred to Appendix A.1. For the sake of simplicity, suppose that we want to compute the best mean of complexity exactly $\ell' \in [\ell]$. To compute the optimal restricted 2-mean, it suffices to find the best mean for every $\ell' \in [\ell]$. For a fixed $\ell' \in [\ell]$, the main idea is to compute for any two warplings between a point sequence of complexity ℓ' and an input point sequence, a polynomial function whose sign indicates which of the warplings yields a smaller distance between the sequences. These functions are then used to define an arrangement that partitions the space $(\mathbb{R}^d)^{\ell'}$. The trick is that while there is an infinite number of point sequences in $(\mathbb{R}^d)^{\ell'}$, to each input point sequence there are only $O(m^{2\ell'})$ warplings and in each face of the arrangement the point sequences have the same optimal warplings to the input point sequences. Therefore, for an arbitrary point sequence from each face of the arrangement, we can compute the optimal warplings to the input sequences and then use the resulting sections to compute an optimal point sequence for these warplings, obtaining the optimal mean of complexity exactly ℓ' when we eventually hit the face containing it. We use the cylindrical algebraic decomposition algorithm to compute the arrangement and obtain an element of each face.

We start by describing a simple approximation algorithm that reveals the basic idea underlying the following algorithms. The algorithm relies on the following observation. If p -DTW is defined over a metric, then the triangle inequality holds for the point-to-point-distances in the sum that defines the p -DTW distance (albeit not for p -DTW distance itself). Assume for simplicity that $p = 1$. In this case, there always exists a 2-approximate mean that is formed by points from the input sequences. Enumerating all possible such sequences, then, if the input consists of n point sequences of length m , leads to an algorithm with running time in $O((nm)^{\ell+1})$, where ℓ denotes the largest allowed complexity of the mean. This approach also extends to other variants of the mean problem for different choices of p and q (with varying approximation factors). One obvious disadvantage of this simple algorithm is the high running time. In the following, we use similar observations as above and show that the dependency on the number of input sequences n can be improved to linear while still achieving approximation factors close to 2.

We present a randomized constant-factor approximation algorithm for the restricted p -mean problem. The approximation factor of the algorithm depends on p , and the best it can achieve is $2 + \varepsilon$ for $p = 1$ and $4 + \varepsilon$ for $p = 2$, which resemble the famous Euclidean median and mean problems. The idea of the algorithm is to obtain for each $j \in [\ell']$ from the corresponding section $S_j^p(c, T)$ of an optimal restricted p -mean $c = (c_1, \dots, c_{\ell'})$ one of the closest input vertices to c_j . The obtained vertices in the corresponding order form an approximate restricted p -mean. We formalize the idea in the following lemma. The proof is deferred to Appendix A.2.

Now we present the first algorithm. The idea is to uniformly sample from the set of all vertices of all point sequences, to obtain at least one vertex from each ball guaranteed by the previous lemma, with high probability. After the sampling, the algorithm enumerates all point sequences of at most ℓ elements from the sample and returns a point sequence with lowest cost.

```

1: procedure MEAN-C( $T = \{\tau_1 = (\tau_{1,1}, \dots, \tau_{1,|\tau_1|}), \dots, \tau_n = (\tau_{n,1}, \dots, \tau_{n,|\tau_n|})\}, \delta, \varepsilon, p$ )
2:    $P \leftarrow \bigcup_{i=1}^n \bigcup_{j=1}^{|\tau_i|} \{\tau_{i,j}\}$ 
3:    $S \leftarrow \text{sample} \left[ \frac{m(\ln(\ell) + \ln(1/\delta))}{\varepsilon/(2^{p-1} + \varepsilon)} \right]$  points from  $P$  uniformly and independently at random
      with replacement
4:    $C \leftarrow S^{\leq \ell}$ 
5:   return an arbitrary element from  $\arg \min_{c \in C} \text{cost}_p^p(T, c)$ 

```

The correctness of Algorithm 1 follows by an application of Lemma 6. The proof of the following theorem can be found in Appendix A.2.

► **Theorem 7.** *Given a set $T = \{\tau_1, \dots, \tau_n\} \subseteq X^{\leq m}$ of point sequences, three parameters $\delta \in (0, 1)$, $\varepsilon \in (0, \infty)$ and $p \in [1, \infty)$, Algorithm 1 returns with probability at least $1 - \delta$ a $(2^p + \varepsilon)$ -approximate restricted p -mean for T , in time $O\left(nm^{\ell+1} \ln(1/\delta)^\ell \left(1 + \frac{2^{p-1}}{\varepsilon}\right)^\ell\right)$.*

2.2.2 Derandomization

In this section, we consider finite metric spaces (X, ρ) for which the set of all metric balls $\mathcal{B} = \{B(x, r) \mid x \in X, r \in \mathbb{R}_{\geq 0}\}$ forms a range space (X, \mathcal{B}) with bounded VC dimension \mathcal{D} . We present a deterministic algorithm for the restricted p -mean problem which is applicable under the additional assumption that there is a subsystem oracle for (X, \mathcal{B}) . We show that this is the case for the Euclidean metric. Note that our algorithm depends on the existence of a subsystem oracle, which is not always obvious for a given metric.

We formally define range spaces and the associated concepts. A range space is defined as a pair of sets (X, \mathcal{R}) , where X is the *ground set* and $\mathcal{R} \subseteq 2^X$ is the *range set*. For $Y \subseteq X$, we denote $\mathcal{R}|_Y = \{R \cap Y \mid R \in \mathcal{R}\}$ and if $\mathcal{R}|_Y$ contains all subsets of Y , then Y is *shattered* by \mathcal{R} . A measure of the combinatorial complexity of such a range space is the VC dimension.

► **Definition 8** (VC dimension). *The Vapnik-Chervonenkis dimension [31, 34, 35] of (X, \mathcal{R}) is the maximum cardinality of a shattered subset of X .*

Range spaces need not to be finite and can be discretized by means of ε -nets.

► **Definition 9** (ε -net). *A set $N \subset X$ is an ε -net for (X, \mathcal{R}) if for any range $R \in \mathcal{R}$, $R \cap N \neq \emptyset$ if $|R \cap X| \geq \varepsilon|X|$.*

To compute ε -nets deterministically, we need a subsystem oracle, which we now define.

► **Definition 10** (subsystem oracle). *Let (X, \mathcal{R}) be a finite range space. A subsystem oracle is an algorithm which for any $Y \subseteq X$, lists all sets in $\mathcal{R}|_Y$ in time $O(|Y|^{\mathcal{D}+1})$, where \mathcal{D} is the VC dimension of (X, \mathcal{R}) .*

We use the following theorem to obtain ε -nets when provided with a subsystem oracle.

► **Theorem 11** ([9, Theorem 2.1]). *Let (X, \mathcal{R}) be a range space with finite ground set and VC dimension \mathcal{D} , and $\varepsilon > 0$ be a given parameter. Assume that there is a subsystem oracle for (X, \mathcal{R}) . Then an ε -net of size $O\left(\frac{\mathcal{D}}{\varepsilon} \log \frac{\mathcal{D}}{\varepsilon}\right)$ can be computed deterministically in time $O(\mathcal{D}^{3\mathcal{D}}) \cdot \left(\frac{1}{\varepsilon} \log \frac{1}{\varepsilon}\right)^{\mathcal{D}} \cdot |X|$.*

The following algorithm is a modification of Algorithm 1 where the sampling step is substituted for a computation of an (ε/m) -net of the set of all vertices of all given point sequences. Since the balls guaranteed by Lemma 6 are of appropriate size, the (ε/m) -net stabs all of them and by enumeration of all point sequences of at most ℓ points from the (ε/m) -net, we again find a good approximate restricted p -mean.

■ Algorithm 2 Restricted p -Mean Constant-Factor Approximation

-
- 1: **procedure** MEAN-C-D($T = \{\tau_1 = (\tau_{1,1}, \dots, \tau_{1,|\tau_1|}), \dots, \tau_n = (\tau_{n,1}, \dots, \tau_{n,|\tau_n|})\}, \varepsilon, p$)
 - 2: $\varepsilon' \leftarrow \frac{\varepsilon}{2^{p-1} + \varepsilon}$, $P \leftarrow \bigcup_{i=1}^n \bigcup_{j=1}^{|\tau_i|} \{\tau_{i,j}\}$
 - 3: $S \leftarrow$ compute an (ε'/m) -net of (P, \mathcal{B})
 - 4: $C \leftarrow S^{\leq \ell}$
 - 5: **return** an arbitrary element from $\arg \min_{c \in C} \text{cost}_p^T(T, c)$
-

The correctness of Algorithm 2 follows from Definition 9. The proofs of the following statements can be found in Appendix A.2.

► **Theorem 12.** *Given a set $T \subseteq X^{\leq m}$ of n point sequences, parameters $\varepsilon \in (0, \infty)$, and $p \in [1, \infty)$, Algorithm 2 returns a $(2^p + \varepsilon)$ -approximate restricted p -mean for T .*

We now turn to the Euclidean setting. First, we prove that there exists a subsystem oracle for (X, \mathcal{B}) , when $X \subset \mathbb{R}^d$ is a finite subset of the d -dimensional Euclidean space.

► **Lemma 13.** *There is a subsystem oracle for the range space (X, \mathcal{B}) , where X is a finite subset of \mathbb{R}^d .*

Then we can analyze the running time of Algorithm 2 in the Euclidean setting.

► **Theorem 14.** *Given a set $T \subset (\mathbb{R}^d)^{\leq m}$ of n point sequences, parameters $\varepsilon \in (0, \infty)$, and $p \in [1, \infty)$, Algorithm 2 can be implemented to run in $O\left(nm \left(\left(\frac{m}{\varepsilon'} \log \frac{m}{\varepsilon'}\right)^{d+1} + \left(\frac{m}{\varepsilon'} \log \frac{m}{\varepsilon'}\right)^\ell\right)\right)$ deterministic time, where $\varepsilon' = \frac{\varepsilon}{2^{p-1} + \varepsilon}$.*

2.3 Simplifications and the Triangle Inequality

In this section, we give an efficient approximation algorithm for simplification under p -DTW and show that a weak triangle inequality for p -DTW holds. These results are then used in Section 2.4, where we provide an algorithm for the restricted $(p, 1)$ -mean problem, which achieves an approximation factor of $(1 + \varepsilon)$, for any $\varepsilon \in (0, \infty)$. The result holds for the important case of sequences lying in the Euclidean space.

In this section, we focus on approximating a minimum-error simplification of a point sequence with respect to the DTW distance, and we show that a weak triangle inequality holds. This improves upon previous similar statements [19] when one bounds the DTW distance of two short point sequences. We will later apply these two results in the analysis of one of our mean approximation algorithms. In particular, we will bound the expected cost of a mean obtained by randomly sampling an input point sequence and then computing its approximate minimum-error simplification.

2.3.1 Minimum-Error Simplification

We first define the notion of simplification of a point sequence under the p -DTW distance.

► **Definition 15.** *Let $\pi \in X^*$. An (α, ℓ) -simplification of π , under the d_{DTW_p} distance, is a point sequence $\tilde{\pi} \in X^{\leq \ell}$ such that*

$$\forall \pi' \in X^{\leq \ell} : d_{\text{DTW}_p}(\pi, \tilde{\pi}) \leq \alpha \cdot d_{\text{DTW}_p}(\pi, \pi').$$

We design a dynamic programming solution for the problem of computing a simplification. Each subproblem is parameterized by the length of a prefix of the input point sequence and the maximum length of a simplification of that prefix. A detailed description of our algorithm and the analysis can be found in Section A.3.

► **Theorem 16.** *There is an algorithm that given as input $\pi \in X^m$, computes a $(2, \ell)$ -simplification of π under the d_{DTW_p} , in time $O(m^4 \ell)$.*

2.3.2 Weak Triangle Inequality

While DTW is not a metric and it is known that the triangle inequality fails for certain instances, there is a weak version of the triangle inequality that is satisfied. In particular, Lemire [19] shows that given $x, y, z \in X^m$, and $p \in [1, \infty)$, we have $d_{\text{DTW}_p}(x, z) \leq m^{1/p} \cdot (d_{\text{DTW}_p}(x, y) + d_{\text{DTW}_p}(y, z))$. We slightly generalize the above inequality in a way that implies a better bound for the distance between two short point sequences using the distances to a potentially longer point sequence. All missing proofs can be found in Appendix A.3.

► **Lemma 17.** *For any $m_1, m_2 \in \mathbb{N}$, let $x, z \in X^{\leq m_1}$, $y \in X^{m_2}$, and $p \in [1, \infty)$. Then,*

$$d_{\text{DTW}_p}(x, z) \leq m_1^{1/p} \cdot (d_{\text{DTW}_p}(x, y) + d_{\text{DTW}_p}(y, z)).$$

The following theorem uses the weak triangle inequality and provides an upper bound on the expected cost of the restricted $(p, 1)$ -mean obtained by first sampling an input point sequence uniformly at random and then computing an (α, ℓ) -simplification of this point sequence. This theorem will be useful in the next section, where we design an approximation scheme for the mean problem that relies on a first rough estimation of the cost.

► **Theorem 18.** *Let $T = \{\tau_1, \dots, \tau_n\} \subseteq X^{\leq m}$ be a set of point sequences and let $p \in [1, \infty)$. Let π be a point sequence picked uniformly at random from T , and let $\tilde{\pi}$ be an (α, ℓ) -simplification of π under d_{DTW_p} , where $\ell \leq m$. Then,*

$$\mathbb{E}_\pi [\text{cost}_p^1(T, \tilde{\pi})] \leq (2 + \alpha) m^{1/p} \ell^{1/p} \cdot \text{OPT}_\ell,$$

where OPT_ℓ denotes the cost of the optimal restricted $(p, 1)$ -mean of T .

Proof. Let c be an optimal $(p, 1)$ -mean of T with cost OPT_ℓ . Then,

$$\begin{aligned} \mathbb{E}_\pi [\text{cost}(T, \tilde{\pi})] &= \mathbb{E}_\pi \left[\sum_{i=1}^n d_{\text{DTW}_p}(\tau_i, \tilde{\pi}) \right] \leq \mathbb{E}_\pi \left[m^{\frac{1}{p}} \sum_{i=1}^n (d_{\text{DTW}_p}(\tau_i, c) + d_{\text{DTW}_p}(c, \tilde{\pi})) \right] \quad (1) \\ &= m^{1/p} \cdot (\text{OPT}_\ell + n \cdot \mathbb{E}_\pi [d_{\text{DTW}_p}(c, \tilde{\pi})]) \\ &\leq m^{1/p} \cdot (\text{OPT}_\ell + n \cdot \ell^{1/p} \cdot \mathbb{E}_\pi [d_{\text{DTW}_p}(c, \pi) + d_{\text{DTW}_p}(\pi, \tilde{\pi})]) \quad (2) \\ &\leq m^{1/p} \cdot (\text{OPT}_\ell + (1 + \alpha) \cdot n \cdot \ell^{1/p} \cdot \mathbb{E}_\pi [d_{\text{DTW}_p}(c, \pi)]) \\ &= m^{1/p} \cdot \left(\text{OPT}_\ell + (1 + \alpha) \cdot n \cdot \ell^{1/p} \cdot \sum_{\pi \in T} d_{\text{DTW}_p}(c, \pi) \cdot \frac{1}{n} \right) \\ &= m^{1/p} \cdot (\text{OPT}_\ell + (1 + \alpha) \cdot \ell^{1/p} \cdot \text{OPT}_\ell) \leq (2 + \alpha) m^{1/p} \ell^{1/p} \cdot \text{OPT}_\ell, \end{aligned}$$

where in Step (1) and Step (2), we applied Lemma 17. ◀

2.4 Approximation Scheme for Point Sequences in Euclidean Spaces

Now we study the restricted $(p, 1)$ -mean problem for point sequences in the Euclidean space. This is exactly the problem of computing one median point sequence of complexity at most ℓ , under the p -DTW distance. Formally, input point sequences belong to $(\mathbb{R}^d)^{\leq m}$ and we are interested in computing a median point sequence in $(\mathbb{R}^d)^\ell$. The distance between any two points $x, y \in \mathbb{R}^d$ is measured by the Euclidean distance $\|x - y\|$. For any $x \in \mathbb{R}^d$, $r > 0$, $B(x, r)$ denotes the Euclidean ball $\{y \in \mathbb{R}^d \mid \|x - y\| \leq r\}$. We also use Euclidean grids:

► **Definition 19 (grid).** *Given a number $r \in \mathbb{R}_+$, for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ we define by $G(r, x) = (\lfloor x_1/r \rfloor \cdot r, \dots, \lfloor x_d/r \rfloor \cdot r)$ the r -grid-point of x . Let $P \subseteq \mathbb{R}^d$ be a subset of \mathbb{R}^d . The grid of cell width r that covers P is the set $\mathbb{G}(P, r) = \{G(r, x) \mid x \in P\}$.*

A grid partitions \mathbb{R}^d into cubic regions. For any $r \in \mathbb{R}_+$, $x \in P$, we have $\|x - G(r, x)\| \leq r\sqrt{d}$.

2.4.1 Algorithm

We build upon ideas developed in Section 2.3 and we design a $(1 + \varepsilon)$ -approximation algorithm for the restricted $(p, 1)$ -mean problem. The algorithm is randomized and succeeds with probability $1 - \delta$, where δ is a user-defined parameter.

The high-level idea is the following. Given a set T of n point sequences, we first compute a rough estimate of the optimal cost. To do so, we sub-sample a sufficiently large number of input sequences that we store in a set S , and we compute a $(2, \ell)$ -simplification for each one of them. We detect a sequence in S whose simplification minimizes the restricted $(p, 1)$ -mean cost; this cost is denoted by R . The results of Section 2.3 imply that with good probability, R is a $O((m\ell)^{1/p})$ approximation of the optimal cost. We can now use R to “guess” a refined estimate for the restricted $(p, 1)$ -mean cost which is within a constant factor from the optimal, by enumerating multiples of 2 in the interval $[\Omega(R(m\ell)^{-1/p}), R]$. Assuming that we have such an estimate, we can use it to fine-tune a grid, which is then intersected with balls centered at the points of sequences in S . We use the resulting grid points to compute a set of candidate solutions. The idea here is that with good probability one of the point sequences in S is very close to the optimal solution, so one of the candidate solutions will be a good approximation. The complete pseudocode can be found in Algorithm 3.

Algorithm 3 Restricted $(p, 1)$ -mean $(1 + \varepsilon)$ -Approximation

```

1: procedure MED-APPR( $T = \{\tau_1 = (\tau_{1,1}, \dots, \tau_{1,|\tau_1|}), \dots, \tau_n = (\tau_{n,1}, \dots, \tau_{n,|\tau_n|})\}, \varepsilon, p, \delta$ )
2:    $S \leftarrow$  sample  $\lceil \log(2/\delta) \rceil$  point sequences from  $T$  uniformly and independently
      at random with replacement
3:    $\mathcal{R} \leftarrow \emptyset, C \leftarrow \emptyset$ 
4:   for each  $\tau_i \in S$  do
5:      $\tau'_i \leftarrow (2, \ell)$ -simplification of  $\tau_i$  under  $d_{\text{DTWP}}$ 
6:      $\mathcal{R} \leftarrow \mathcal{R} \cup \{\text{cost}_p^1(T, \tau'_i)\}$ 
7:    $R \leftarrow \min \mathcal{R}$ 
8:    $\beta \leftarrow 2 \cdot \left( \frac{68m^{1/p}}{\varepsilon} + 5 \right)^d$ 
9:    $I_R \leftarrow \left\{ \frac{R \cdot 2^{-i}}{n} \mid i = 0, \dots, \lceil 3 + \log(m\ell)/p \rceil \right\}$ 
10:  for each  $r \in I_R$  do
11:     $\gamma \leftarrow \frac{\varepsilon \cdot r}{(2m)^{1/p} \sqrt{d}}$ 
12:    for each  $\tau_i \in S$  do
13:       $\mathcal{B}(\tau_i, 4r) \leftarrow \bigcup_{j=1}^{|\tau_i|} B(\tau_{i,j}, 4r)$ 
14:       $N \leftarrow \mathbb{G}(\mathcal{B}(\tau_i, 4r), \gamma)$ 
15:      if  $|N| \leq \ell \cdot \beta$  then
16:         $C \leftarrow C \cup N^{\leq \ell}$ 
17:  return an arbitrary element of  $\arg \min_{c \in C} \text{cost}_p^1(T, c)$ 

```

2.4.2 Analysis

Now we analyze the running time and correctness of Algorithm 3. Missing proofs can be found in Appendix A.4. We begin with a bound on the probability that R is a rough approximation of the optimal median cost.

► **Lemma 20.** *Let c be an optimal restricted $(p, 1)$ -mean of T . With probability at least*

$$1 - \delta/2,$$

$$R \leq 8m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c).$$

Next, we bound the probability that a point sequence in the sample S is conveniently close to the optimal median.

► **Lemma 21.** *Let c be an optimal restricted $(p, 1)$ -mean of T . With probability at least $1 - \delta/2$, there exists a $\tau_i \in S$ such that $d_{\text{DTW}_p}(\tau_i, c) < (2/n) \cdot \text{cost}(T, c)$.*

The set I_R contains a value r such that nr is within a factor of 2 from the optimal cost.

► **Lemma 22.** *Let c be an optimal restricted $(p, 1)$ -mean of T . If $R \leq 8m^{1/p}\ell^{1/p} \text{cost}(T, c)$, then there exists $r \in I_R$ such that $\text{cost}(T, c) \in [nr, 2nr]$.*

The following is an upper bound on the number of grid cells needed to cover a Euclidean ball. Similar bounds appear often in the literature, but they are typically asymptotic and not sufficient for our needs. Therefore, we prove an exact (non-asymptotic) upper bound.

► **Lemma 23.** *Let $x \in \mathbb{R}^d$, $r > 0$, $\gamma > 0$.*

$$|\mathbb{G}(B(x, 8r), \gamma)| \leq 2 \cdot \left(\frac{34r}{\gamma\sqrt{d}} + 5 \right)^d.$$

We now focus on the iteration of the algorithm with $r \in I_R$, $\tau_i \in S$, such that r satisfies the property guaranteed by Lemma 22 and τ_i satisfies the property guaranteed by Lemma 21. We claim that in that iteration, an $(1 + \varepsilon)$ -approximate median is inserted to C .

► **Lemma 24.** *Let c be an optimal restricted $(p, 1)$ -mean of T . Let r^* be such that $\text{cost}(T, c) \in [nr^*, 2nr^*]$ and let $\gamma^* = \frac{\varepsilon r^*}{(2m)^{1/p}\sqrt{d}}$. If $\tau_i \in S$ is such that $d_{\text{DTW}_p}(\tau_i, c) \leq (2/n) \cdot \text{cost}(T, c)$ then*

- i) $|\mathbb{G}(\mathcal{B}(\tau_i, 4r^*), \gamma^*)| \leq \ell \cdot 2 \cdot \left(\frac{34r^*}{\gamma^*\sqrt{d}} + 5 \right)^d$ and
- ii) *there exists $c' \in \mathbb{G}(\mathcal{B}(\tau_i, 4r^*), \gamma^*)^{\leq \ell}$ such that $\text{cost}(T, c') \leq (1 + \varepsilon) \cdot \text{cost}(T, c)$.*

The correctness of our algorithm follows by combining the above.

► **Lemma 25.** *Given a set $T \subset (\mathbb{R}^d)^{\leq m}$, $\varepsilon > 0$, $p \in [1, \infty)$, $\delta \in (0, 1)$, Algorithm 3 returns a $(1 + \varepsilon)$ -approximate restricted $(p, 1)$ -mean with probability of success $1 - \delta$.*

Proof. Let c be an optimal restricted $(p, 1)$ -mean of T . Applying a union bound over the events of Lemma 20 and Lemma 21, we conclude that with probability at least $1 - \delta$, we have $R \leq 8m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c)$, and there exists a $\tau_i \in S$ such that $d_{\text{DTW}_p}(\tau_i, c) < (2/n) \cdot \text{cost}(T, c)$. We show correctness assuming that the above two events hold. By Lemma 22 we know that there exists an $r^* \in I_R$ such that $\text{cost}(T, c) \in [nr^*, 2nr^*]$.

We focus on the iteration where r^* is considered. Let γ^* be the value of γ in that iteration and let N^* be the set N in that iteration. By Lemma 24 i), $|N^*| \leq \ell\beta$ and all point sequences of complexity at most ℓ defined by points in N^* will be considered as possible solutions. By Lemma 24 ii), there is a point sequence in $(N^*)^{\leq \ell}$ which is a $(1 + \varepsilon)$ -approximate solution. ◀

Finally, we bound the running time of Algorithm 3.

► **Theorem 26.** *Given a set $T \subset (\mathbb{R}^d)^{\leq m}$ of n point sequences, $\varepsilon \in (0, m^{1/p}]$, $p \in [1, \infty)$, $\delta \in (0, 1)$, Algorithm 3 returns a $(1 + \varepsilon)$ -approximate restricted $(p, 1)$ -mean with probability of success $1 - \delta$ and has running time in $O\left(\left(m^4 + nm \cdot \left(\frac{m^{1/p}}{\varepsilon}\right)^{d\ell} \cdot \frac{\log(m)}{p}\right) \cdot \log\left(\frac{1}{\delta}\right)\right)$.*

Proof. Correctness follows from Lemma 25. It remains to bound the running time. For each one of the point sequences in S , we compute its $(2, \ell)$ -simplification in $O(dm^4\ell)$ time using Theorem 16 and its median cost in $O(dnm\ell)$ time. Hence, the total time needed to compute \mathcal{R} and then R is $O((m^4 + nm) \cdot d\ell \log(1/\delta))$. The set I_R has cardinality $|I_R| = O(\log(m\ell)/p)$. For each value $r \in I_R$, we add at most $\sum_{i=1}^{\ell} |N|^i \cdot |S| \leq \ell \left(\ell \cdot 2 \cdot (68m^{1/p}\varepsilon^{-1} + 5)^d \right)^{\ell} \cdot |S|$ candidates. For each candidate point sequence in C , we compute the cost in time $O(dnm\ell)$. Since d and ℓ are considered constants, the total running time is $O\left(\left(m^4 + nm \cdot (m^{1/p}\varepsilon^{-1})^{d\ell} \cdot \log(m)/p\right) \cdot \log(1/\delta)\right)$. \blacktriangleleft

3 Application to Clustering

We can apply the results of Sections 2.2 and 2.3 to the problem of clustering of point sequences. The details can be found in Appendix A.5.

► **Definition 27** ((k, ℓ, p, q) -clustering). *The (k, ℓ, p, q) -clustering problem is defined as follows, where $k \in \mathbb{N}$, $\ell \in \mathbb{N}_{>1}$ and $p, q \in [1, \infty)$ are fixed (constant) parameters of the problem: given a set $T = \{\tau_1, \dots, \tau_n\} \subseteq X^{\leq m}$ of point sequences, compute a set $C \subseteq X^{\leq \ell}$ of k point sequences, such that $\text{cost}_p^q(T, C) = \sum_{i=1}^n \min_{c \in C} d_{\text{DTW}_p}(c, \tau_i)^q$ is minimal.*

Solving an instance of the (k, ℓ, p, q) -clustering problem is equivalent to solving an instance of the k -median problem, where the distance between any center c and any other element x is measured by $d_{\text{DTW}_p}(x, c)^q$. To solve the k -median problem, we apply a generic framework, as described in [10], which reduces the clustering problem to a problem of computing a set that contains medians for sufficiently large subsets of the input, with good probability. To solve this latter problem, we adapt our algorithms from Sections 2.2 and 2.3. Algorithm 1 can be easily modified to return the set of candidates, instead of returning the best among them. We show that setting parameters appropriately yields an algorithm that satisfies the properties required by the above-mentioned framework and leads to a randomized algorithm for the (k, ℓ, p, p) -clustering problem with approximation factor in $O(2^p)$, probability of success $1 - \delta$ and running time in $O((2^p k m \ln(\ell/\delta))^{\ell(k+2)} \cdot nm)$, assuming that the time needed to compute the distance between two points is constant (see Theorem 33 in Appendix A.5). Similarly, the random sampling method implied by Theorem 18 can be used to produce a sufficiently large sample of candidates, which leads to a randomized algorithm for the $(k, \ell, p, 1)$ -clustering problem with approximation factor in $O(m^{1/p}\ell^{1/p})$, probability of success $1 - \delta$ and running time in $O((k \log(1/\delta))^{k+2} \cdot nm + m^4(k \log(1/\delta))^{k+2})$, assuming again constant time for distance computations of points (see Theorem 35 in Appendix A.5).

4 Conclusions

We have studied mean problems for point sequences under the p -DTW distance and devised exact and approximation algorithms for several relevant problem variants where the complexity of the mean is restricted by a parameter ℓ . Our exact algorithm runs in polynomial time for constant ℓ and d . The running times of our approximation algorithms depend only linearly on the number of input sequences. The dependency on the length of the input sequences, however, is high; the dependency on the parameter ℓ is even exponential. We hope that the algorithmic techniques developed in this paper will inspire further work on the topic. In particular, we think the weak triangle inequality and the simplification algorithm could be of great use. In contrast, a proof of hardness of approximation for the central problem studied in this paper is not in sight. We leave this as an open problem.

References

- 1 John Aach and George M. Church. Aligning gene expression time series with time warping algorithms. *Bioinformatics*, 17(6):495–508, 2001.
- 2 Amir Abboud, Arturs Backurs, and Virginia Vassilevska Williams. Tight hardness results for LCS and other sequence similarity measures. In Venkatesan Guruswami, editor, *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015*, pages 59–78. IEEE Computer Society, 2015.
- 3 Waleed H. Abdulla, David Chow, and Gary Sin. Cross-words reference template for dtw-based speech recognition systems. In *TENCON 2003. Conference on Convergent Technologies for Asia-Pacific Region*, volume 4, pages 1576–1579 Vol.4, 2003.
- 4 Donald J. Berndt and James Clifford. Using dynamic time warping to find patterns in time series. In Usama M. Fayyad and Ramasamy Uthrusamy, editors, *Knowledge Discovery in Databases: Papers from the 1994 AAAI Workshop, Seattle, Washington, USA, July 1994. Technical Report WS-94-03*, pages 359–370. AAAI Press, 1994.
- 5 Milutin Brankovic, Kevin Buchin, Koen Klaren, André Nusser, Aleksandr Popov, and Sampson Wong. (k, ℓ) -Medians Clustering of Trajectories Using Continuous Dynamic Time Warping. In Chang-Tien Lu, Fusheng Wang, Goce Trajcevski, Yan Huang, Shawn D. Newsam, and Li Xiong, editors, *SIGSPATIAL '20: 28th International Conference on Advances in Geographic Information Systems, Seattle, WA, USA, November 3-6, 2020*, pages 99–110. ACM, 2020.
- 6 Markus Brill, Till Fluschnik, Vincent Froese, Brijnesh J. Jain, Rolf Niedermeier, and David Schultz. Exact mean computation in dynamic time warping spaces. In *Proceedings of the 2018 SIAM International Conference on Data Mining, SDM 2018, May 3-5, 2018, San Diego Marriott Mission Valley, San Diego, CA, USA*, pages 540–548, 2018.
- 7 Markus Brill, Till Fluschnik, Vincent Froese, Brijnesh J. Jain, Rolf Niedermeier, and David Schultz. Exact mean computation in dynamic time warping spaces. *Data Min. Knowl. Discov.*, 33(1):252–291, 2019.
- 8 Karl Bringmann and Marvin Künnemann. Quadratic conditional lower bounds for string problems and dynamic time warping. In Venkatesan Guruswami, editor, *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015*, pages 79–97. IEEE Computer Society, 2015.
- 9 Hervé Brönnimann, Bernard Chazelle, and Jiri Matousek. Product range spaces, sensitive sampling, and derandomization. *SIAM Journal on Computing*, 28(5):1552–1575, 1999.
- 10 Kevin Buchin, Anne Driemel, and Martijn Struijs. On the hardness of computing an average curve. In *17th Scandinavian Symposium and Workshops on Algorithm Theory, SWAT 2020, June 22-24, 2020, Tórshavn, Faroe Islands*, pages 19:1–19:19, 2020.
- 11 Maike Buchin, Anne Driemel, and Dennis Rohde. Approximating (k, ℓ) -median clustering for polygonal curves. In Dániel Marx, editor, *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021*, pages 2697–2717. SIAM, 2021.
- 12 Laurent Bulteau, Vincent Froese, and Rolf Niedermeier. Tight hardness results for consensus problems on circular strings and time series. *SIAM J. Discret. Math.*, 34(3):1854–1883, 2020.
- 13 E. G. Caiani, A. Porta, G. Baselli, M. Turiel, S. Muzzupappa, F. Pieruzzi, C. Crema, A. Malliani, and S. Cerutti. Warped-average template technique to track on a cycle-by-cycle basis the cardiac filling phases on left ventricular volume. In *Computers in Cardiology 1998. Vol. 25 (Cat. No.98CH36292)*, pages 73–76, 1998.
- 14 George E. Collins. Quantifier elimination for real closed fields by cylindrical algebraic decomposition. In H. Brakhage, editor, *Automata Theory and Formal Languages*, pages 134–183, Berlin, Heidelberg, 1975. Springer Berlin Heidelberg.
- 15 Shreyasi Datta, Chandan K. Karmakar, and Marimuthu Palaniswami. Averaging methods using dynamic time warping for time series classification. In *2020 IEEE Symposium Series on Computational Intelligence (SSCI)*, pages 2794–2798, 2020.

- 16 Herbert Edelsbrunner, Joseph O'Rourke, and Raimund Seidel. Constructing arrangements of lines and hyperplanes with applications. *SIAM Journal on Computing*, 15(2):341–363, May 1986.
- 17 Sariel Har-peled. *Geometric Approximation Algorithms*. American Mathematical Society, USA, 2011.
- 18 Ville Hautamäki, Pekka Nykanen, and Pasi Franti. Time-series clustering by approximate prototypes. In *2008 19th International Conference on Pattern Recognition*, pages 1–4, 2008.
- 19 Daniel Lemire. Faster retrieval with a two-pass dynamic-time-warping lower bound. *Pattern Recognition*, 42(9):2169 – 2180, 2009.
- 20 Yu-Tao Liu, Yong'an Zhang, and Ming Zeng. Adaptive global time sequence averaging method using dynamic time warping. *IEEE Transactions on Signal Processing*, 67:2129–2142, 2019.
- 21 Alexander De Luca, Alina Hang, Frederik Brudy, Christian Lindner, and Heinrich Hussmann. Touch me once and i know it's you!: implicit authentication based on touch screen patterns. In Joseph A. Konstan, Ed H. Chi, and Kristina Höök, editors, *CHI Conference on Human Factors in Computing Systems, CHI '12, Austin, TX, USA - May 05 - 10, 2012*, pages 987–996. ACM, 2012.
- 22 Jirí Matousek. *Lectures on discrete geometry*, volume 212 of *Graduate texts in mathematics*. Springer, 2002.
- 23 Michael Mitzenmacher and Eli Upfal. *Probability and Computing: Randomization and Probabilistic Techniques in Algorithms and Data Analysis*. Cambridge University Press, USA, 2nd edition, 2017.
- 24 Marion Morel, Catherine Achard, Richard Kulpa, and Séverine Dubuisson. Time-series averaging using constrained dynamic time warping with tolerance. *Pattern Recognition*, 74:77–89, 2018.
- 25 Lindasalwa Muda, Mumtaj Begam, and I. Elamvazuthi. Voice recognition algorithms using mel frequency cepstral coefficient (mfcc) and dynamic time warping (dtw) techniques, 2010. [arXiv:1003.4083](https://arxiv.org/abs/1003.4083).
- 26 Mario E. Munich and Pietro Perona. Continuous dynamic time warping for translation-invariant curve alignment with applications to signature verification. In *Proceedings of the International Conference on Computer Vision, Kerkyra, Corfu, Greece, September 20-25, 1999*, pages 108–115. IEEE Computer Society, 1999.
- 27 Manabu Okawa. Time-series averaging and local stability-weighted dynamic time warping for online signature verification. *Pattern Recognition*, 112:107699, 2021.
- 28 François Petitjean, Alain Ketterlin, and Pierre Gançarski. A global averaging method for dynamic time warping, with applications to clustering. *Pattern Recognit.*, 44(3):678–693, 2011.
- 29 Lawrence Rabiner and Jay Wilpon. Considerations in applying clustering techniques to speaker independent word recognition. In *ICASSP '79. IEEE International Conference on Acoustics, Speech, and Signal Processing*, volume 4, pages 578–581, 1979.
- 30 Hiroaki Sakoe and Seibi Chiba. Dynamic programming algorithm optimization for spoken word recognition. *IEEE Transactions on Acoustics, Speech, and Signal Processing*, 26(1):43–49, 1978.
- 31 Norbert Sauer. On the density of families of sets. *Journal of Combinatorial Theory Series A*, 13:145–147, 1972.
- 32 Nathan Schar, Vincent Froese, and Rolf Niedermeier. Faster binary mean computation under dynamic time warping. In *31st Annual Symposium on Combinatorial Pattern Matching, CPM 2020, June 17-19, 2020, Copenhagen, Denmark*, pages 28:1–28:13, 2020.
- 33 David Schultz and Brijnesh Jain. Nonsmooth analysis and subgradient methods for averaging in dynamic time warping spaces. *Pattern Recognition*, 74:340–358, 2018.
- 34 Saharon Shelah. A combinatorial problem; stability and order for models and theories in infinitary languages. *Pacific Journal of Mathematics*, 41(1), 1972.
- 35 Vladimir Vapnik and Alexey Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability and its Applications*, 16:264–280, 1971.

- 36 Edmund T. Whittaker and George N. Watson. *A Course of Modern Analysis*. Cambridge Mathematical Library. Cambridge University Press, 4 edition, 1996.
- 37 Yunyue Zhu and Dennis E. Shasha. Warping indexes with envelope transforms for query by humming. In Alon Y. Halevy, Zachary G. Ives, and AnHai Doan, editors, *Proceedings of the 2003 ACM SIGMOD International Conference on Management of Data, San Diego, California, USA, June 9-12, 2003*, pages 181–192. ACM, 2003.

A

 Appendix

A.1 Missing Proof of Section 2.1

► **Theorem 5.** *There exists an algorithm that, given a set $T \subset (\mathbb{Q}^d)^{\leq m}$ of n point sequences, computes an optimal restricted 2-mean (defined over the Euclidean distance) in time $(nm)^{2^{O(d\ell)}}$.*

Before we prove the theorem we note that the employed cylindrical algebraic decomposition algorithm only works for polynomials with real algebraic coefficients. Therefore, here we assume that the input point sequences have rational coordinates. This is indeed a realistic assumption since physical computers are not capable of storing arbitrary real numbers.

Proof of Theorem 5. To simplify our exposition, we restrict ourselves to means of complexity exactly $\ell' \in [\ell]$, i.e. in the rest of the proof we describe an algorithm for computing an optimal mean of complexity exactly ℓ' . The complete algorithm consists of iteratively computing the optimal mean of complexity ℓ' , for each $\ell' \in [\ell]$.

For each $\tau = (\tau_1, \dots, \tau_{|\tau|}) \in T$ and all $W_1, W_2 \in \mathcal{W}_{\ell', |\tau|}$ we define for $c = (c_1, \dots, c_{\ell'}) \in (\mathbb{R}^d)^{\ell'}$ the polynomial function

$$P_{\tau, W_1, W_2}(c) = \left(\sum_{(i,j) \in W_1} \|c_i - \tau_j\|^2 \right) - \left(\sum_{(i,j) \in W_2} \|c_i - \tau_j\|^2 \right).$$

Clearly, iff W_1 yields a smaller distance between c and τ than W_2 , then $P_{\tau, W_1, W_2}(c) < 0$ and iff W_2 yields a smaller distance between c and τ than W_1 , then $P_{\tau, W_1, W_2}(c) > 0$. Iff $P_{\tau, W_1, W_2}(c) = 0$, both yield the same distance.

Let $F = \{P_{\tau, W_1, W_2} \mid \tau \in T, W_1, W_2 \in \mathcal{W}_{\ell', |\tau|}\}$ be the set of these polynomials. The central observation is that if all functions in F have the same sign for any $c_1, c_2 \in (\mathbb{R}^d)^{\ell'}$, then c_1 and c_2 have the same optimal 2-warplings to the point sequences in T . To see this, for each $\tau \in T$ let $W_\tau \in \mathcal{W}_{\ell', |\tau|}$ be an optimal 2-warpling between c_1 and τ . Clearly, $P_{\tau, W_\tau, W}(c_1) \leq 0$ for all $\tau \in T$ and $W \in \mathcal{W}_{\ell', |\tau|}$. Now, if all functions in F have the same sign for c_1 and c_2 it must be that $P_{\tau, W_\tau, W}(c_2) \leq 0$ for all $\tau \in T$ and $W \in \mathcal{W}_{\ell', |\tau|}$. Thus, W_τ is an optimal 2-warpling between c_2 and τ for each $\tau \in T$.

Now, we compute an arrangement of the zero sets of the polynomials in F (cf. [22]), i.e., a partition of $(\mathbb{R}^d)^{\ell'}$ into regions where all functions in F have the same sign. For this purpose we use the cylindrical algebraic decomposition algorithm [14], which also yields a sample from each face of the arrangement and has running time $O(|F|^{f(\ell' \cdot d)})$ for some function $f \in 2^{O(d\ell')}$. For each sample c from some face of the arrangement we first compute the optimal 2-warplings between c and the input point sequences $\tau \in T$ in time $O(nm)$. Second we compute all sections $S_j(c)$ of c and stores the point sequence $c' = (c'_1, \dots, c'_{\ell'})$ consisting of the means $c'_j = \frac{1}{|S_j(c)|} \sum_{v \in S_j(c)} v$ for $j = 1, \dots, \ell'$. This takes time $O(nm)$.

At some point, we obtain a sample from the face containing the optimal $(\ell', 2, 2)$ -mean $c^* = (c_1^*, \dots, c_{\ell'}^*)$ (where c_j^* is the mean of $S_j(c^*)$ for each $j \in [\ell']$), which we return when we

finally return the point sequence c' that minimizes the objective function. This takes time $O(nmA)$, where A is the number of cells in the arrangement.

To conclude the proof, note that for each $\tau \in T$ we have that $|\mathcal{W}_{\ell', |\tau|}| \leq m^{2\ell'}$, thus $|F| \leq nm^{4\ell'}$. Hence, $A \leq \left(\frac{100nm^{4\ell'}}{\ell'd}\right)^{\ell'd}$ by [22, Theorem 6.2.1].

As we have already mentioned, we iteratively run the above algorithm to compute means of complexity ℓ' , for each $\ell' \in [\ell]$, in order to find an optimal restricted 2-mean. Each iteration runs in $(nm)^{2^{O(d\ell')}} \leq (nm)^{2^{O(d\ell)}}$. Since ℓ is constant, the running time is in $(nm)^{2^{O(d\ell)}}$. ◀

A.2 Missing Proofs of Section 2.2

► **Lemma 6.** *Let $T = \{\tau_1 = (\tau_{1,1}, \dots, \tau_{1,|\tau_1|}), \dots, \tau_n = (\tau_{n,1}, \dots, \tau_{n,|\tau_n|})\} \subseteq X^{\leq m}$ be a set of point sequences and let $P = \bigcup_{i=1}^n \bigcup_{j=1}^{|\tau_i|} \{\tau_{i,j}\}$. For any $p \in [1, \infty)$, $\ell \in \mathbb{N}_{>1}$ and $\varepsilon \in (0, \infty)$ there exists an $\ell' \leq \ell$ and balls $B_1, \dots, B_{\ell'} \subseteq P$, of cardinality at least $\frac{\varepsilon n}{2^{p-1} + \varepsilon}$ each, such that any point sequence $c' = (c'_1, \dots, c'_{\ell'})$, with $c'_i \in B_i$ for each $i \in [\ell']$, is a $(2^p + \varepsilon)$ -approximate restricted p -mean for T .*

Proof. Let $c = (c_1, \dots, c_{\ell'}) \in X^{\ell'}$ be an optimal restricted p -mean for T and for $j \in [\ell']$ let $S_j = \{s_{j,1}, \dots, s_{j,n_j}\} = S_j^p(c)$ for brevity. Define $\Delta(S_j) = \sum_{v \in S_j} \rho(c_j, v)^p$. We immediately

have $\text{cost}(T, c) = \sum_{j=1}^{\ell'} \Delta(S_j)$. Now, for $j \in [\ell']$, let π_j be a permutation of the index set $[n_j]$, such that

$$\rho(c_j, s_{j,\pi_j^{-1}(1)})^p \leq \dots \leq \rho(c_j, s_{j,\pi_j^{-1}(n_j)})^p.$$

Let $\varepsilon' = \frac{\varepsilon}{2^{p-1} + \varepsilon}$. For the sake of simplicity, we assume that $\varepsilon'n$ is integral. Further, for $j \in [\ell']$, we define $C_j = \{s_{j,\pi_j^{-1}(1)}, \dots, s_{j,\pi_j^{-1}(\varepsilon'n)}\}$. We have that $\rho(c_j, s_{j,\pi_j^{-1}(\varepsilon'n)})^p \leq \frac{\Delta(S_j)}{|S_j| - (\varepsilon'n - 1)}$ by the fact that $\rho(c_j, s_{j,\pi_j^{-1}(\varepsilon'n)})^p$ is maximal, iff $\rho(c_j, s')^p = 0$ for each $s' \in C_j \setminus \{s_{j,\pi_j^{-1}(\varepsilon'n)}\}$ and $\rho(c_j, s')^p = \rho(c_j, s_{j,\pi_j^{-1}(\varepsilon'n)})^p$ for each $s' \in S_j \setminus C_j$. For $j \in [\ell']$, we now define

$$B_j = \{x \in P \mid \rho(c_j, x)^p \leq \rho(c_j, s_{j,\pi_j^{-1}(\varepsilon'n)})^p\}$$

and by definition we have $\rho(c_j, x)^p \leq \frac{\Delta(S_j)}{|S_j| - \varepsilon'n + 1} \leq \frac{\Delta(S_j)}{|S_j| - \varepsilon'n}$ for each $x \in B_j$ and $j \in [\ell']$. Then let $c' = (c'_1, \dots, c'_{\ell'})$ be a point sequence with $c'_j \in B_j$ for each $j \in [\ell']$. We bound its cost:

$$\begin{aligned} \text{cost}(T, c') &= \sum_{j=1}^{\ell'} \sum_{v \in S_j} \rho(c'_j, v)^p \leq \sum_{j=1}^{\ell'} \sum_{v \in S_j} (\rho(c_j, v) + \rho(c_j, c'_j))^p \\ &\leq \sum_{j=1}^{\ell'} \sum_{v \in S_j} 2^{p-1} (\rho(c_j, v)^p + \rho(c_j, c'_j)^p) \leq 2^{p-1} \sum_{j=1}^{\ell'} \sum_{v \in S_j} \left(\rho(c_j, v)^p + \frac{\Delta(S_j)}{|S_j| - \varepsilon'n} \right) \\ &\leq 2^{p-1} \text{cost}(T, c) + 2^{p-1} \sum_{j=1}^{\ell'} \sum_{v \in S_j} \frac{\Delta(S_j)}{(1 - \varepsilon')|S_j|} = \left(2^{p-1} + \frac{2^{p-1}}{1 - \varepsilon'} \right) \text{cost}(T, c) \\ &= (2^p + \varepsilon) \text{cost}(T, c). \end{aligned}$$

The first inequality follows from the triangle-inequality and the last inequality holds, because a vertex from each $\tau_i \in T$ must be warped to each $c_j \in c$, thus $|S_j| \geq n$ for each $j \in [\ell']$. ◀

► **Theorem 7.** *Given a set $T = \{\tau_1, \dots, \tau_n\} \subseteq X^{\leq m}$ of point sequences, three parameters $\delta \in (0, 1)$, $\varepsilon \in (0, \infty)$ and $p \in [1, \infty)$, Algorithm 1 returns with probability at least $1 - \delta$ a $(2^p + \varepsilon)$ -approximate restricted p -mean for T , in time $O\left(nm^{\ell+1} \ln(1/\delta)^\ell \left(1 + \frac{2^{p-1}}{\varepsilon}\right)^\ell\right)$.*

Proof. For the given ε , let $\varepsilon' = \frac{\varepsilon}{2^{p-1} + \varepsilon}$ and let $B_1, \dots, B_{\ell'}, \ell' \leq \ell$, be the balls guaranteed by Lemma 6. Recall that each ball has size at least $\varepsilon'n$. For each $i \in [\ell']$ and $s \in S$ we have $\Pr[s \notin B_i] \leq (1 - \frac{\varepsilon'n}{|P|}) \leq (1 - \frac{\varepsilon'n}{nm}) = (1 - \frac{\varepsilon'}{m}) \leq \exp(-\varepsilon'/m)$.

By independence, for each $i \in [\ell']$ we have $\Pr[B_i \cap S = \emptyset] \leq \exp(-\varepsilon'/m)^{\lceil \frac{m(\ln(\ell) - \ln(\delta))}{\varepsilon'} \rceil} \leq \delta/\ell$. Using a union bound we conclude that with probability at least $1 - \delta$, S contains at least one element of B_i , for each $i \in [\ell']$, and thus Algorithm 1 returns a $(2^p + \varepsilon)$ -approximate restricted p -mean for T with probability at least $1 - \delta$ by Lemma 6.

The running time of the algorithm is dominated by computing the cost of all point sequences of complexity at most ℓ over S . Since $|S^{\leq \ell}|$ is in $O\left(\frac{\ln(1/\delta)^\ell m^\ell}{(\varepsilon')^\ell}\right) = O\left(\ln(1/\delta)^\ell m^\ell \frac{(2^{p-1} + \varepsilon)^\ell}{\varepsilon^\ell}\right)$ and every distance can be computed in time $O(m)$, this takes time $O\left(\ln(1/\delta)^\ell m^{\ell+1} n \frac{(2^{p-1} + \varepsilon)^\ell}{\varepsilon^\ell}\right)$. \blacktriangleleft

► **Theorem 12.** *Given a set $T \subseteq X^{\leq m}$ of n point sequences, parameters $\varepsilon \in (0, \infty)$, and $p \in [1, \infty)$, Algorithm 2 returns a $(2^p + \varepsilon)$ -approximate restricted p -mean for T .*

Proof. By Lemma 6, for any $\varepsilon \in (0, \infty)$ there exist balls $B_1, \dots, B_{\ell'} \subseteq P$, $\ell' \leq \ell$ of cardinality at least $\varepsilon'n$ each, such that any point sequence $c' = (c'_1, \dots, c'_{\ell'})$, with $c'_i \in B_i$ for each $i \in [\ell']$, is a $(2^p + \varepsilon)$ -approximate restricted p -mean for T , where $\varepsilon' = \frac{\varepsilon}{2^{p-1} + \varepsilon}$. Since we compute an (ε'/m) -net of P and $|P| \leq nm$, S contains at least one point from each of $B_1, \dots, B_{\ell'}$ by Definition 9. Hence, $S^{\leq \ell}$ contains a $(2^p + \varepsilon)$ -approximate restricted p -mean for T . \blacktriangleleft

► **Lemma 13.** *There is a subsystem oracle for the range space (X, \mathcal{B}) , where X is a finite subset of \mathbb{R}^d .*

Proof. The VC dimension of (P, \mathcal{B}) is bounded by $d + 1$, see [17]. For any $Y \subseteq P$, we need to compute the set \mathcal{B}_Y explicitly in time $O(|Y|^{d+2})$. We first apply the standard lifting $\phi: (x_1, \dots, x_d) \mapsto (x_1, \dots, x_d, \sum_{i=1}^d x_i^2)$. A point $p \in Y$ belongs to some ball $B \in \mathcal{B}$, with center $c = (c_1, \dots, c_d) \in \mathbb{R}^d$ and radius $r > 0$, if and only if $\phi(p)$ lies below the hyperplane h_B , where h_B is the hyperplane defined by the equation $\langle a_B, x \rangle = b_B$, where $a_B = (2c_1, 2c_2, \dots, 2c_d, 1)$ and $b_B = r^2 - \sum_{i=1}^d c_i^2$. Notice that h_B is nonvertical by definition. Then we dualize: for any point $\phi(p) = (y_1, \dots, y_{d+1})$, $D(\phi(p)) = \{(x_1, \dots, x_{d+1}) \in \mathbb{R}^{d+1} \mid x_{d+1} = \sum_{i=1}^d x_i y_i - y_{d+1}\}$ is a nonvertical hyperplane in \mathbb{R}^{d+1} and for any nonvertical hyperplane h_B , $D^{-1}(h_B)$ is a point in \mathbb{R}^{d+1} . A standard fact about duality is that a point $\phi(p)$ lies below a hyperplane h_B if and only if the hyperplane $D(\phi(p))$ lies above point $D^{-1}(h_B)$. Finally we construct the arrangement of hyperplanes in the dual space in time $O(|Y|^{d+1})$, using the algorithm in [16]. For each of the at most $O(|Y|^{d+1})$ cells, we return a subset $X \subseteq Y$ corresponding to the hyperplanes lying above. The overall running time is $O(|Y|^{d+2})$. \blacktriangleleft

► **Theorem 14.** *Given a set $T \subset (\mathbb{R}^d)^{\leq m}$ of n point sequences, parameters $\varepsilon \in (0, \infty)$, and $p \in [1, \infty)$, Algorithm 2 can be implemented to run in $O\left(nm \left(\left(\frac{m}{\varepsilon'} \log \frac{m}{\varepsilon'}\right)^{d+1} + \left(\frac{m}{\varepsilon'} \log \frac{m}{\varepsilon'}\right)^\ell\right)$ deterministic time, where $\varepsilon' = \frac{\varepsilon}{2^{p-1} + \varepsilon}$.*

Proof. The VC dimension of the range space (P, \mathcal{B}) is bounded by $d + 1$, see [17]. By Lemma 13, we can use Theorem 11 to compute an (ε'/m) -net S of (P, \mathcal{B}) , with size $|S| = O\left(\frac{m}{\varepsilon'} \log \left(\frac{m}{\varepsilon'}\right)\right)$, in time $O\left(nm \left(\frac{m}{\varepsilon'} \log \left(\frac{m}{\varepsilon'}\right)\right)^{d+1}\right)$. We then compute the d_{DTW_p} distance of any of the $|S^{\leq \ell}|$ candidates with the n input point sequences in time $O(|S|^\ell \cdot nm)$. \blacktriangleleft

A.3 Details of Section 2.3

A.3.1 Minimum-error Simplification

We present a dynamic programming solution for the problem of computing an approximate simplification. Our algorithm can be seen as a special case of the result of Brill et al. [6, 7] for computing a mean of restricted complexity, but since our statement is different, we include a proof for completeness.

■ **Algorithm 4** 2-Approximate Simplification

```

1: procedure 2-APPROXIMATE-SIMPLIFICATION( $\pi = (x_1, \dots, x_m), \ell, p$ )
2:   Initialize  $m \times \ell$  table  $D$  with elements in  $\mathbb{R}$ 
3:   Initialize  $m \times \ell$  table  $C$  with elements in  $\mathbb{R}^{\leq \ell}$ 
4:    $P \leftarrow \{x_1, \dots, x_m\}$ 
5:   for each  $i = 1, \dots, m$  do
6:     for each  $j = 1, \dots, \ell$  do
7:       if  $j = 1$  then
8:          $x^* \leftarrow \arg \min_{x \in P} \sum_{k=1}^i \rho(x_k, x)^p$ 
9:          $D(i, j) \leftarrow \sum_{k=1}^i \rho(x_k, x^*)^p$ 
10:         $C(i, j) \leftarrow (x^*)$ 
11:       else
12:         $i' \leftarrow \arg \min_{i' \leq i} \left( D(i', j-1) + \min_{x \in P} \sum_{k=i'}^i \rho(x_k, x)^p \right)$ 
13:         $x^* \leftarrow \arg \min_{x \in P} \sum_{k=i'}^i \rho(x_k, x)^p$ 
14:         $D(i, j) \leftarrow D(i', j-1) + \sum_{k=i'}^i \rho(x_k, x^*)^p$ 
15:         $C(i, j) \leftarrow C(i', j-1) \oplus (x^*)$ 
16:    $j^* \leftarrow \arg \min_{j \in [\ell]} D(m, j)$ 
17:   return  $C(m, j^*)$ 

```

► **Lemma 28.** *Let $\mathcal{X} = (X, \rho)$ be a metric space. Given as input a point sequence $\pi = (x_1, \dots, x_m) \in X^m$, Algorithm 4 returns a point sequence from $\{x_1, \dots, x_m\}^{\leq \ell}$, which minimizes the d_{DTW_p} distance to π , among all point sequences in $\{x_1, \dots, x_m\}^{\leq \ell}$.*

Proof. We show that $C(m, j^*)$ satisfies

$$d_{\text{DTW}_p}(\pi, C(m, j^*)) = \min_{\pi' \in P^{\leq \ell}} d_{\text{DTW}_p}(\pi, \pi').$$

We claim that there is a point sequence $\tilde{\pi} \in P^{\leq \ell}$ such that

$$d_{\text{DTW}_p}(\pi, \tilde{\pi}) = \min_{\pi' \in P^{\leq \ell}} d_{\text{DTW}_p}(\pi, \pi'),$$

and such that the optimal warping between π and $\tilde{\pi}$ does not match two vertices of $\tilde{\pi}$ with the same vertex of π . To see this, consider an optimal warping W between π and some point sequence $\pi' = (p'_1, \dots, p'_j) \in P^{\leq \ell}$. Let $(t_1, t_2) \in W$ and $(t_1, t_2 + 1) \in W$. If $(t_1 - 1, t_2) \in W$ then removing (t_1, t_2) yields a new warping with a cost at most equal to the cost of W . Similarly, if $(t_1 + 1, t_2 + 1) \in W$ then removing $(t_1, t_2 + 1)$ from W yields a new warping with a cost at most equal to the cost of W . If $(t_1 - 1, t_2) \notin W$, then we can remove p'_{t_2} from π' . If $(t_1 + 1, t_2 + 1) \notin W$, then we can remove p'_{t_2+1} from π' . We conclude that there exists a point sequence $\pi'' \in P^{\leq \ell}$ such that $d_{\text{DTW}_p}(\pi, \pi'') \leq d_{\text{DTW}_p}(\pi, \pi')$, and an optimal warping W between π and π'' for which there are no $t_1 \in [m], t_2 \in [\ell]$ such that both $(t_1, t_2) \in W$ and $(t_1, t_2 + 1) \in W$.

For each $i \in [m]$, let $\pi|_i = (x_1, \dots, x_i)$. By construction, each $D(i, j)$ stores the minimum distance between $\pi|_i$ and any point sequence x from P^j , where the distance is attained by a warping that does not match two vertices of x to the same vertex of π . Hence, $D(m, j^*)$ stores the minimum distance between π and any point sequence in $P^{\leq \ell}$, and $C(m, j^*)$ stores a point sequence from $P^{\leq \ell}$ with distance $D(m, j^*)$ from π . \blacktriangleleft

► **Lemma 29.** *Let $\mathcal{X} = (X, \rho)$ be a metric space. Given as input a point sequence $\pi \in X^m$, Algorithm 4 returns a $(2, \ell)$ -simplification under the d_{DTW_p} distance.*

Proof. Let $P = \{x_1, \dots, x_m\}$. By Lemma 28, $C(m, j^*)$ is a point sequence in $P^{\leq \ell}$ that minimizes the distance to π , among all point sequences in $P^{\leq \ell}$. We show that $C(m, j^*)$ is a $(2, \ell)$ -simplification. Let $\pi^* = (x_1^*, \dots, x_{\ell'}^*)$ be a $(1, \ell)$ -simplification of π , and let $\tilde{\pi}^* = (\tilde{x}_1^*, \dots, \tilde{x}_{\ell'}^*)$, where for each $i \in [\ell']$, $\tilde{x}_i^* := \arg \min_{x \in P} \rho(x, x_i^*)$. Let $W^* \in \mathcal{W}_{m, \ell'}$ be an optimal warping of π and π^* . Then,

$$\begin{aligned}
d_{\text{DTW}_p}^1(\pi, C(m, j^*)) &\leq d_{\text{DTW}_p}^1(\pi, \tilde{\pi}^*) \\
&= \min_{W \in \mathcal{W}_{m, \ell}} \left(\sum_{(i, j) \in W} \rho(x_i, \tilde{x}_j^*)^p \right)^{1/p} \\
&\leq \left(\sum_{(i, j) \in W^*} \rho(x_i, \tilde{x}_j^*)^p \right)^{1/p} \\
&\leq \left(\sum_{(i, j) \in W^*} (\rho(x_i, x_j^*) + \rho(x_j^*, \tilde{x}_j^*))^p \right)^{1/p} \\
&\leq \left(\sum_{(i, j) \in W^*} 2^p (\rho(x_i, x_j^*))^p \right)^{1/p} \\
&= 2 \cdot d_{\text{DTW}_p}(\pi, \pi^*),
\end{aligned} \tag{3}$$

where in Step (3) we applied the triangle inequality. \blacktriangleleft

► **Theorem 16.** *There is an algorithm that given as input $\pi \in X^m$, computes a $(2, \ell)$ -simplification of π under the d_{DTW_p} , in time $O(m^4 \ell)$.*

Proof. Correctness of Algorithm 4 follows from Lemma 29. It remains to bound the running time of the algorithm. To do so, we consider the operations taking place in the body of the nested loop. For each i, j , we iterate over $O(m)$ values for i' and for each value of i' we compute $\min_{x \in P} \sum_{k=i'}^i \rho(x_k, x)^p$ in time $O((i - i') \cdot m) = O(m^2)$. Hence, the total running time is $O(m^4 \ell)$. \blacktriangleleft

A.3.2 Weak Triangle Inequality

► **Lemma 17.** *For any $m_1, m_2 \in \mathbb{N}$, let $x, z \in X^{\leq m_1}$, $y \in X^{m_2}$, and $p \in [1, \infty)$. Then,*

$$d_{\text{DTW}_p}(x, z) \leq m_1^{1/p} \cdot (d_{\text{DTW}_p}(x, y) + d_{\text{DTW}_p}(y, z)).$$

Proof. Let $W_{xz} \in \mathcal{W}_{|x|, |z|}$ be the optimal warping between x and z . Let $W_{xy} \in \mathcal{W}_{|x|, |y|}$ be the optimal warping between x and y and $W_{yz} \in \mathcal{W}_{|y|, |z|}$ be the optimal warping between

y and z . Let $S_{xz} = \{(i, k, j) \in [|x|] \times [|y|] \times [|z|] \mid (i, k) \in W_{xy} \text{ and } (k, j) \in W_{yz}\}$ and $W'_{xz} = \{(i, j) \in [|x|] \times [|z|] \mid \exists k (i, k, j) \in S_{xz}\}$. Then,

$$\begin{aligned} d_{\text{DTW}_p}(x, z) &= \left(\sum_{(i,j) \in W_{xz}} \rho(x_i, z_j)^p \right)^{1/p} \\ &\leq \left(\sum_{(i,j) \in W'_{xz}} \rho(x_i, z_j)^p \right)^{1/p} \\ &\leq \left(\sum_{(i,k,j) \in S_{xz}} (\rho(x_i, y_k) + \rho(y_k, z_j))^p \right)^{1/p} \\ &\leq \left(\sum_{(i,k,j) \in S_{xz}} \rho(x_i, y_k)^p \right)^{1/p} + \left(\sum_{(i,k,j) \in S_{xz}} \rho(y_k, z_j)^p \right)^{1/p} \\ &\leq m_1^{1/p} \cdot d_{\text{DTW}_p}(x, y) + m_1^{1/p} \cdot d_{\text{DTW}_p}(y, z), \end{aligned}$$

where the second inequality holds by the triangle inequality and the third inequality holds by Minkowski's inequality. \blacktriangleleft

A.4 Missing Proofs of Section 2.4

► **Lemma 20.** *Let c be an optimal restricted $(p, 1)$ -mean of T . With probability at least $1 - \delta/2$,*

$$R \leq 8m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c).$$

Proof. For any point sequence $\tau_i \in T$, let τ'_i be a $(2, \ell)$ -simplification. Let τ_j be a randomly sampled point sequence from T . By Theorem 18,

$$\mathbb{E} [\text{cost}(T, \tau'_j)] \leq 4m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c).$$

By Markov's inequality, $\Pr [\text{cost}(T, \tau'_j) \geq 8m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c)] \leq \frac{1}{2}$. Hence, the probability that $R \geq 8m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c)$ is equal to

$$\Pr [\forall \tau_i \in S : \text{cost}(T, \tau'_i) \geq 8m^{1/p}\ell^{1/p} \cdot \text{cost}(T, c)] \leq \frac{1}{2^{|S|}} \leq \frac{\delta}{2}.$$

\blacktriangleleft

► **Lemma 21.** *Let c be an optimal restricted $(p, 1)$ -mean of T . With probability at least $1 - \delta/2$, there exists a $\tau_i \in S$ such that $d_{\text{DTW}_p}(\tau_i, c) < (2/n) \cdot \text{cost}(T, c)$.*

Proof. Let τ_i be a randomly sampled point sequence from T :

$$\mathbb{E}_{\tau_i} [d_{\text{DTW}_p}(\tau_i, c)] = \sum_{i=1}^n d_{\text{DTW}_p}(\tau_i, c) \cdot \frac{1}{n} = \frac{\text{cost}(T, c)}{n}.$$

By Markov's inequality, $\Pr [d_{\text{DTW}_p}(\tau_i, c) > 2 \cdot \frac{\text{cost}(T, c)}{n}] \leq \frac{1}{2}$. Hence,

$$\Pr \left[\forall \tau_i \in S : d_{\text{DTW}_p}(\tau_i, c) > 2 \cdot \frac{\text{cost}(T, c)}{n} \right] \leq \frac{1}{2^{|S|}} \leq \frac{\delta}{2}.$$

\blacktriangleleft

► **Lemma 22.** *Let c be an optimal restricted $(p, 1)$ -mean of T . If $R \leq 8m^{1/p}\ell^{1/p} \text{cost}(T, c)$, then there exists $r \in I_R$ such that $\text{cost}(T, c) \in [nr, 2nr]$.*

Proof. Since R is the cost of a curve of complexity at most ℓ , we have that $\text{cost}(T, c) \leq R$. By assumption, $\text{cost}(T, c) \geq R/(8m^{1/p}\ell^{1/p})$. By the definition of I_R , there exists $j \geq 0$ such that

$$2^{-(j+1)} \cdot \frac{R}{n} \leq \frac{\text{cost}(T, c)}{n} \leq 2^{-j} \cdot \frac{R}{n}.$$

Hence, the lemma is true for $r = 2^{-(j+1)} \cdot \frac{R}{n}$. ◀

► **Lemma 23.** *Let $x \in \mathbb{R}^d$, $r > 0$, $\gamma > 0$.*

$$|\mathbb{G}(B(x, 8r), \gamma)| \leq 2 \cdot \left(\frac{34r}{\gamma\sqrt{d}} + 5 \right)^d.$$

Proof. We use Binet's second expression [36] for the Gamma function $\ln \Gamma(z)$:

$$\ln \Gamma(z) = z \ln(z) - z + \frac{1}{2} \ln \left(\frac{2\pi}{z} \right) + \int_0^\infty \frac{2 \arctan \left(\frac{t}{z} \right)}{e^{2\pi t} - 1} dt.$$

Since $\arctan(x) \geq 0$ for $x \geq 0$ and $e^{2\pi x} - 1 \geq 0$ for $x \geq 0$, we have the following inequality:

$$\begin{aligned} \ln \Gamma(z) &\geq z \ln(z) - z + \frac{1}{2} \ln \left(\frac{2\pi}{z} \right) \\ \iff \ln \Gamma(z) &\geq \ln(z^z) - \ln(e^z) + \ln \left(\sqrt{\frac{2\pi}{z}} \right) \\ \iff \Gamma(z) &\geq z^z e^{-z} \sqrt{\frac{2\pi}{z}} \\ \iff \Gamma(z) &\geq \sqrt{2\pi} z^{z-\frac{1}{2}} e^{-z}. \end{aligned} \tag{4}$$

We apply a standard volumetric argument to upper bound $|\mathbb{G}(B(x, 8r), \gamma)|$.

$$\begin{aligned} |\mathbb{G}(B(x, 8r), \gamma)| &\leq \frac{\text{vol}(B(x, 8r + \gamma\sqrt{d}))}{\gamma^d} = \frac{\pi^{d/2}}{\Gamma(\frac{d}{2} + 1)} \cdot \frac{(8r + \gamma\sqrt{d})^d}{\gamma^d} \\ &\leq \frac{\pi^{d/2} e^{d/2+1}}{\sqrt{2\pi} \left(\frac{d}{2} + 1 \right)^{d/2+1/2}} \cdot \frac{(8r + \gamma\sqrt{d})^d}{\gamma^d} \\ &\leq \frac{2^{d/2+1/2} \pi^{d/2} e^{d/2+1}}{\sqrt{2\pi} \cdot d^{d/2+1/2}} \cdot \frac{(8r + \gamma\sqrt{d})^d}{\gamma^d} \\ &\leq \frac{e \cdot (4.2)^d}{\sqrt{\pi} \cdot d^{d/2}} \cdot \frac{(8r + \gamma\sqrt{d})^d}{\gamma^d} \\ &\leq 2 \cdot \left(\frac{34r}{\gamma\sqrt{d}} + 5 \right)^d, \end{aligned} \tag{5}$$

where in (5) we used (4). ◀

► **Lemma 24.** *Let c be an optimal restricted $(p, 1)$ -mean of T . Let r^* be such that $\text{cost}(T, c) \in [nr^*, 2nr^*]$ and let $\gamma^* = \frac{\varepsilon r^*}{(2m)^{1/p}\sqrt{d}}$. If $\tau_i \in S$ is such that $d_{\text{DTW}_p}(\tau_i, c) \leq (2/n) \cdot \text{cost}(T, c)$ then*

- i) $|\mathbb{G}(\mathcal{B}(\tau_i, 4r^*), \gamma^*)| \leq \ell \cdot 2 \cdot \left(\frac{34r^*}{\gamma^* \sqrt{d}} + 5\right)^d$ and
 ii) there exists $c' \in \mathbb{G}(\mathcal{B}(\tau_i, 4r^*), \gamma^*)^{\leq \ell}$ such that $\text{cost}(T, c') \leq (1 + \varepsilon) \cdot \text{cost}(T, c)$.

Proof. Let $c = (c_1, \dots, c_{\ell'})$, where $\ell' \leq \ell$. To prove i), notice that $d_{\text{DTW}_p}(\tau_i, c) \leq 4r^*$, which implies that for any vertex $\tau_{i,j}$ of τ_i , there exists a vertex c_z of c such that $\tau_{i,j} \in B(c_z, 4r^*)$. By the triangle inequality $B(\tau_{i,j}, 4r^*) \subseteq B(c_z, 8r^*)$. Hence,

$$\begin{aligned} \mathcal{B}(\tau_i, 4r^*) \subseteq \bigcup_{z=1}^{\ell} B(c_z, 8r^*) &\implies |\mathbb{G}(\mathcal{B}(\tau_i, 4r^*), \gamma^*)| \leq \left| \mathbb{G}\left(\bigcup_{z=1}^{\ell'} B(c_z, 8r^*), \gamma^*\right) \right| \\ &\leq \sum_{z=1}^{\ell} |\mathbb{G}(B(c_z, 8r^*), \gamma^*)|. \end{aligned}$$

By Lemma 23, we obtain

$$|\mathbb{G}(\mathcal{B}(\tau_i, 4r^*), \gamma^*)| \leq \ell \cdot 2 \cdot \left(\frac{34r^*}{\gamma^* \sqrt{d}} + 5\right)^d.$$

To prove ii), notice that all vertices of c are contained in $\mathcal{B}(\tau_i, 4r^*)$. Hence, for each point c_z there exists a grid point $\tilde{c}_z \in \mathbb{G}(\mathcal{B}(\tau_i, r^*), \gamma^*)$ such that $\|c_z - \tilde{c}_z\| \leq \gamma^* \sqrt{d}$. We will show that the point sequence $\tilde{c} = (\tilde{c}_1, \dots, \tilde{c}_{\ell'})$ is a $(1 + \varepsilon)$ -approximation. For each $i \in [n]$, W_i^* denotes the optimal warping of τ_i with c .

$$\begin{aligned} \text{cost}(T, \tilde{c}) &= \sum_{i=1}^n d_{\text{DTW}_p}(\tau_i, \tilde{c}) \\ &= \sum_{i=1}^n \min_{W \in \mathcal{W}_{|\tau_i|, \ell}} \left(\sum_{(k,j) \in W} \|\tau_{i,k} - \tilde{c}_j\|^p \right)^{1/p} \\ &\leq \sum_{i=1}^n \left(\sum_{(k,j) \in W_i^*} \|\tau_{i,k} - \tilde{c}_j\|^p \right)^{1/p} \\ &\leq \sum_{i=1}^n \left(\sum_{(k,j) \in W_i^*} (\|\tau_{i,k} - c_j\| + \|c_j - \tilde{c}_j\|)^p \right)^{1/p} \\ &\leq \sum_{i=1}^n \left(\left(\sum_{(k,j) \in W_i^*} \|\tau_{i,k} - c_j\|^p \right)^{1/p} + \left(\sum_{(k,j) \in W_i^*} \|c_j - \tilde{c}_j\|^p \right)^{1/p} \right) \\ &\leq \sum_{i=1}^n \left(d_{\text{DTW}_p}(\tau_i, c) + |W_i^*|^{1/p} \cdot \gamma^* \sqrt{d} \right) \\ &\leq \sum_{i=1}^n \left(d_{\text{DTW}_p}(\tau_i, c) + \frac{\text{cost}(T, c) \cdot \varepsilon}{n} \right) \\ &= (1 + \varepsilon) \cdot \text{cost}(T, c), \end{aligned}$$

where the second inequality follows from the triangle inequality, and the third inequality follows from Minkowski's inequality. We also make use of the fact that $|W_i^*| \leq 2m$. \blacktriangleleft

A.5 Applications to Clustering

In this section we apply the results of Sections 2.2 and 2.3 to the problem of clustering of point sequences.

Solving a (k, ℓ, p, q) -clustering problem is equivalent to solving a k -median problem, where the distance between any center c and any other element x is measured by $d_{\text{DTW}_p}(x, c)^q$. To solve the k -medians problem, one can apply the following two theorems, which are proven in [10], and appear slightly rephrased here. The two theorems provide sufficient conditions for a solution to the k -medians problem. We specialize the statements to our case of interest, the d_{DTW_p} distances, raised to the power of q .

► **Theorem 30** (Theorem 7.2 [11]). *Let $T = \{\tau_1, \dots, \tau_n\} \subset X^{\leq m}$, $\alpha \in [1, \infty)$, $\beta \in [1, \infty)$, $\delta \in (0, 1)$, and let $T' \subseteq T$ be an arbitrary subset such that $|T'| \geq |T| \cdot \beta^{-1}$. Suppose that there is an algorithm **Candidates** that given as input T, α, β, δ outputs $C \subset X^{\leq \ell}$ such that with probability at least $1 - \delta$, C contains a point sequence c such that $\text{cost}_p^q(T', c) \leq \alpha \cdot \text{cost}_p^q(T', c^*)$, where c^* is a restricted p -mean of T' .*

*Then, there is an algorithm **k-clustering** that given as input $(T, \emptyset, k, \beta, \delta)$, where $\beta \in (2k, \infty)$, $\delta \in (0, 1)$, $p, q \in [1, \infty)$, returns with probability at least $1 - \delta$ a set $C = \{c_1, \dots, c_k\} \subset X^{\leq \ell}$ with $\text{cost}_p^q(T, C) \leq \left(1 + \frac{4k}{\beta - 2k}\right) \cdot \alpha \cdot \text{cost}_p^q(T, C^*)$, where C^* is an optimal solution to the (k, ℓ, p, q) -clustering problem with input T .*

► **Theorem 31** (Theorem 7.3 [11]). *Let $T_1(n, \alpha, \beta, \delta)$ denote the worst-case running time of **Candidates** for an arbitrary input-set $T \subset X^{\leq m}$ with $|T| = n$ and let $C(n, \alpha, \beta, \delta)$ denote the maximum number of candidates it returns. If T_1 and C are non-decreasing in n , then **k-clustering** has running time in*

$$O\left(C(n, \alpha, \beta, \delta)^{k+2} \cdot nm \cdot T_p + C(n, \alpha, \beta, \delta)^{k+1} \cdot T_1(n, \alpha, \beta, \delta)\right),$$

where T_p denotes the worst-case running time needed to compute the distance between two points in X .

A.5.1 (k, ℓ, p, p) -Clustering

In this section, we apply the result of Section 2.2 to design a randomized algorithm for the (k, ℓ, p, p) -clustering problem. The following algorithm is an adaptation of Algorithm 1.

■ **Algorithm 5** $(1, \ell, p, p)$ -clustering approximate candidates

-
- 1: **procedure** CAND1($T = \{\tau_1 = (\tau_{1,1}, \dots, \tau_{1,|\tau_1|}), \dots, \tau_n = (\tau_{n,1}, \dots, \tau_{n,|\tau_n|})\}, \beta, \delta, \varepsilon, p$)
 - 2: $P \leftarrow \bigcup_{i=1}^n \bigcup_{j=1}^{|\tau_i|} \{\tau_{i,j}\}$
 - 3: $S \leftarrow \text{sample } \lceil (2^p \varepsilon^{-1} + 1) \beta m \ln(\ell/\delta) \rceil$ points from P uniformly and independently at random with replacement
 - 4: **return** $S^{\leq \ell}$
-

► **Lemma 32.** *Let $T \subset X^{\leq m}$, $\beta > 1$, $\delta \in (0, 1)$, $\varepsilon > 0$, $p \geq 1$. Let $T' \subseteq T$ such that $|T'| \geq |T| \cdot \beta^{-1}$ and let C be a set obtained by running Algorithm 5 with input (T, β, p, δ) . Let c be an optimal p -mean of T' . With probability at least $1 - \delta$, there exists $\tau' \in C$ such that*

$$\text{cost}_p^1(T', \tau') \leq (2^p + \varepsilon) \cdot \text{cost}_p^1(T', c).$$

Proof. Let $c \in X^{\ell'}$, $\ell' \leq \ell$. By Lemma 6 applied on T' , we have that there exist sets $B_1, \dots, B_{\ell'} \subseteq P$, each of cardinality at least $\left(\frac{\varepsilon}{2^p + \varepsilon}\right) \cdot |T'|$ such that any point sequence

$c' = (c'_1, \dots, c'_{\ell'})$ with $\forall i \in [\ell'] : c'_i \in B_i$, is a $(2^p + \varepsilon)$ -approximate restricted p -mean of T' . We upper bound the probability that S does not contain any point from a fixed B_i :

$$\Pr[\forall x \in B_i : x \notin S] \leq \left(\frac{|P| - \left(\frac{\varepsilon}{2^p + \varepsilon}\right) \cdot |T'|}{|P|} \right)^{|S|} \leq \left(1 - \left(\frac{\varepsilon}{(2^p + \varepsilon)\beta m} \right) \right)^{|S|} \leq \frac{\delta}{\ell}$$

Then, by a union bound we have that the probability that there exists $i \in [\ell']$ such that $\forall x \in B_i : x \notin S$, is at most δ . Hence, with probability at least $1 - \delta$, there is a point sequence $c' \in S^{\leq \ell}$ which is a $(2^p + \varepsilon)$ -approximate restricted p -mean of T' . ◀

► **Theorem 33.** *There is an algorithm that given a set $T \subset X^{\leq m}$ of n point sequences, $p \in [1, \infty)$, $\beta \in (2k, \infty)$ and $\delta \in (0, 1)$, returns with probability at least $1 - \delta$ a set $C = \{c_1, \dots, c_k\}$ with $\text{cost}_p^1(T, C) \leq \left(1 + \frac{4k}{\beta - 2k}\right) \cdot (2^p + \varepsilon) \cdot \text{cost}_p^1(T, C^*)$, where C^* is an optimal set of k -medians of T , under d_{DTW_p} . The algorithm has running time in*

$$O\left((2^p \varepsilon^{-1} + 1)\beta m \ln(\ell/\delta))^{\ell(k+2)} \cdot nm\right),$$

assuming that the time needed to compute the distance between two points of X is constant.

Proof. We plug Algorithm 5 into Theorems 30 and 31. By Lemma 32, for any $T' \subseteq T$ with $|T'| \geq |T|\beta^{-1}$ Algorithm 5 returns a set of point sequences which contains a $(2^p + \varepsilon)$ -approximate restricted p -mean of T' . Therefore, by Theorem 30 the clustering algorithm is correct. The running time of Algorithm 5 is upper bounded by $O(|S|^\ell + nm)$. The running time then follows by Theorem 31. ◀

A.5.2 k -medians under p -DTW

In this section, we apply the random sampling bound developed in Section 2.3 to design a randomized algorithm for the $(k, \ell, p, 1)$ -clustering problem, that is the problem of computing k -medians of complexity at most ℓ , under d_{DTW_p} . We achieve an approximation factor in $O(m^{1/p} \ell^{1/p})$.

The main idea is that one can use random sampling and approximate simplifications, to obtain a simple algorithm for computing a set of 1-median candidates. Those candidates are guaranteed, up to some user-defined probability, to contain a point sequence which is an approximate 1-median for a fixed but unknown subset of the input.

■ **Algorithm 6** $(1, \ell, p, 1)$ -clustering approximate candidates

-
- 1: **procedure** CAND2($T = \{\tau_1, \dots, \tau_n\}, \beta, p, \delta$)
 - 2: $S \leftarrow$ sample $\lceil 2\beta \cdot \log(2/\delta) \rceil$ point sequences from T uniformly and independently at random with replacement
 - 3: $C \leftarrow \emptyset$
 - 4: **for each** $\tau \in S$ **do**
 - 5: $\tau' \leftarrow (2, \ell)$ -simplification of τ , under d_{DTW_p}
 - 6: $C \leftarrow C \cup \{\tau'\}$
 - 7: **return** C
-

► **Lemma 34.** *Let $T \subset X^{\leq m}$, $\beta > 1$, $p \geq 1$, $\delta \in (0, 1)$. Let $T' \subseteq T$ such that $|T'| \geq |T| \cdot \beta^{-1}$ and let C be a set obtained by running Algorithm 6 with input (T, β, p, δ) . Let c be an optimal restricted $(p, 1)$ -mean of T' . With probability at least $1 - \delta$, there exists $\tau' \in C$ such that*

$$\text{cost}_p^1(T', \tau') \leq 8 \cdot m^{1/p} \ell^{1/p} \cdot \text{cost}_p^1(T', c).$$

Proof. We use a standard Chernoff bound (see [23, Theorem 4.5]) to upper bound the probability that $|S \cap T'| \leq |S|/(2\beta)$. Notice that $\mathbb{E}[|S \cap T'|] \geq |S| \cdot \beta^{-1}$. Hence,

$$\Pr \left[|S \cap T'| \leq \frac{|S|}{2\beta} \right] \leq \exp \left(-\frac{|S|}{8\beta} \right) \leq \frac{\delta}{2}. \quad (6)$$

Let $\mathcal{E}_{T'}$ be the event that $|S \cap T'| > \frac{|S|}{2\beta}$. We condition the rest of the proof on the event $\mathcal{E}_{T'}$. Let τ' be a $(2, \ell)$ -simplification of any point sequence $\tau \in S \cap T'$. Then, by Theorem 18,

$$\mathbb{E}_{\tau} [\text{cost}_p^1(T', \tau') \mid \mathcal{E}_{T'}] \leq 4m^{1/p}\ell^{1/p} \cdot \text{cost}_p^1(T', c).$$

By Markov's inequality, $\Pr [\text{cost}_p^1(T', \tau') \geq 8m^{1/p}\ell^{1/p} \cdot \text{cost}_p^1(T', c) \mid \mathcal{E}_{T'}] \leq \frac{1}{2}$. Hence, by independence of the random sampling,

$$\Pr \left[\forall \tau \in S \cap T' : \text{cost}_p^1(T', \tau') \geq 8m^{1/p}\ell^{1/p} \cdot \text{cost}_p^1(T', c) \mid \mathcal{E}_{T'} \right] \leq \frac{1}{2^{|S|/(2\beta)}} \leq \frac{\delta}{2}. \quad (7)$$

A union bound using inequalities (6), (7) completes the proof. \blacktriangleleft

► **Theorem 35.** *There is an algorithm that given a set $T \subset X^{\leq m}$ of n point sequences, $p \in [1, \infty)$, $\beta \in (2k, \infty)$ and $\delta \in (0, 1)$, returns with probability at least $1 - \delta$ a set $C = \{c_1, \dots, c_k\}$ with $\text{cost}_p^1(T, C) \leq \left(1 + \frac{4k}{\beta - 2k}\right) \cdot (8m^{1/p}\ell^{1/p}) \cdot \text{cost}_p^1(T, C^*)$, where C^* is an optimal set of k -medians of T , under d_{DTW_p} . The algorithm has running time in*

$$O((\beta \log(1/\delta))^{k+2} \cdot nm + (\beta \log(1/\delta))^{k+1} \cdot \beta m^4 \log(1/\delta)),$$

assuming that the distance between two points of X can be computed in constant time.

Proof. We plug Algorithm 6 into Theorems 30 and 31. Lemma 34 guarantees that with probability at least $1 - \delta$, there exists a point sequence in the set C , returned by Algorithm 6, which is an $(8m^{1/p}\ell^{1/p})$ -approximate 1-median to an arbitrary subset $T' \subset T$, as required by Theorem 30. Let T_ρ be time needed to compute the distance between two elements of X . Using Theorem 16 to compute simplifications, Algorithm 6 needs $O(T_\rho \cdot \beta m^4 \ell \log(1/\delta))$ time to compute C . Taking into account the time needed to read the input, and assuming that d, ℓ are constants, the total running time of Algorithm 6 is in $O(nm + T_\rho \cdot \beta m^4 \log(1/\delta))$. Therefore, by Theorem 30, there is an algorithm that returns an $\left(1 + \frac{4k}{\beta - 2k}\right) \cdot (8m^{1/p}\ell^{1/p})$ -approximate solution to the k -medians problem, and by Theorem 31 the algorithm has running time in $O((\beta \log(1/\delta))^{k+2} \cdot nm \cdot T_\rho + (\beta \log(1/\delta))^{k+1} \cdot (nm + T_\rho \cdot \beta m^4 \log(1/\delta)))$. \blacktriangleleft