

Symmetries and Conservation Laws in Hořava Gravity

Deniz O. Devecioğlu ^{*1} and Mu-In Park ^{†2}

¹*School of Physics, Huazhong University of Science
and Technology, Wuhan, Hubei, 430074, China*

²*Center for Quantum Spacetime, Sogang University, Seoul, 121-742, Korea*

(Dated: December 2, 2021)

Abstract

Hořava gravity has been proposed as a renormalizable quantum gravity without the ghost problem through anisotropic scaling dimensions which break Lorentz symmetry in UV. In the Hamiltonian formalism, due to the Lorentz-violating terms, the constraint structure looks quite different from that of general relativity (GR) but we have recently found that “*there exists the case where we can recover the same number of degrees of freedom as in GR*”, in a rather general set-up. In this paper, we study its Lagrangian perspectives and examine the full diffeomorphism (*Diff*) symmetry and its associated conservation laws in Hořava gravity. Surprisingly, we find that the *full Diff* symmetry in the action can also be recovered when a certain condition, called “super-condition”, which *super-selects* the Lorentz-symmetric sector in Hořava gravity, is satisfied. This indicates that the broken Lorentz symmetry, known as “foliation-preserving” *Diff*, is just an *apparent* symmetry of the Hořava gravity action and rather its “*full action symmetry can be as large as the Diff in GR*”. The super-condition exactly corresponds to the *tertiary* constraint in Hamiltonian formalism which is the second-class constraint and provides a non-trivial realization of the Lorentz symmetry otherwise being absent apparently. From the recovered Lorentz symmetry in the action, we obtain the conservation laws with the Noether currents as in covariant theories. The general formula for the conserved Noether charges reproduces the mass of four-dimensional static black holes with an *arbitrary* cosmological constant in Hořava gravity, and is independent of ambiguities associated with the choice of asymptotic boundaries. We also discuss several challenging problems, including its implications to Hamiltonian formalism, black hole thermodynamics, radiations from colliding black holes.

Keywords: Horava Gravity, Lorentz Symmetry, Conservation Laws, Constraint Algebra, Black hole thermodynamics

* E-mail address: dodeve@gmail.com

† E-mail address: muinpark@gmail.com, Corresponding author

The renormalizable gravity without the ghost problem has been studied by considering a gravity action *à la* Hořava, Lifshitz, and DeWitt (HLD) [1–3] in $(D + 1)$ -dimensions (up to some boundary terms),

$$S = \int_{\mathcal{M}} dt d^D x \sqrt{g} N \left\{ \frac{2}{\kappa^2} (K^{ij} K_{ij} - \lambda K^2) - \mathcal{V}[g^{ij}, R^i{}_{jkl}, \nabla_i] \right\} \quad (1)$$

with the higher-spatial-derivative potential \mathcal{V} , satisfying $[\mathcal{V}] \geq D + z$ for the (power-counting) renormalizability, while keeping only the second-time-derivatives with the terms of $K^{ij} K_{ij}$, K^2 in the kinetic part, through the *anisotropic* scaling dimensions, $[t] = -1, [\mathbf{x}] = -z$ for the dynamical critical exponent $z > 1$. Here,

$$K_{ij} \equiv \frac{1}{2N} (\dot{g}_{ij} - \nabla_i N_j - \nabla_j N_i) \quad (2)$$

is the extrinsic curvature [the overdot $(\dot{})$ denotes the time derivative $\partial_t \equiv \partial_0 = ()_{,0}$] and $R^i{}_{jkl}, \nabla_i$ are the Riemann tensor, the spatial covariant derivative for D -dimensional spatial metric g_{ij} on the hypersurface with the ADM decomposition

$$ds^2 = -N^2 dt^2 + g_{ij} (dx^i + N^i dt) (dx^j + N^j dt). \quad (3)$$

The peculiar property of the HLD action (1) is that the Lorentz, *i.e.*, *diffeomorphism* (*Diff*) symmetry in general relativity (GR) is broken into the “foliation-preserving” diffeomorphism (*Diff_F*) from either the DeWitt’s λ parameter in IR ($\lambda \neq 1$) [2], or the higher-derivative terms in UV ($z \geq D$) for power-counting renormalizability [1, 3, 4]. However, it has been unclear how to *canonically* describe the sudden change (reduction) of symmetry beyond the GR limit, $\lambda \rightarrow 1, -\mathcal{V} \rightarrow \Lambda + (2/\kappa^2)R$, by tracing the missing (Lorentz) symmetry all the way down to UV.

On the other hand, we have recently found that the constraint structure in the Hamiltonian formalism looks quite different due to Lorentz-violating terms but “*there exists the case where we can recover the same number of degrees of freedom as in GR [5], at the fully non-linear level*”. This appears to mismatch with the *apparently* broken symmetry in the action and so it suggests some *unbroken/enhanced* symmetry in order to be consistent with the Hamiltonian formalism.

In this paper, in order to clarify this problem, we examine the full action symmetry and its associated conservation laws in Hořava gravity. Surprisingly, we find that the *full Diff* symmetry can be recovered in the Hořava gravity action when the “super-condition” of $\mathcal{L}_0 \equiv \nabla_i \Omega^i = 0$ is satisfied, which exactly corresponds to the *tertiary* constraint in Hamiltonian formalism which is the second-class constraint. This provides a non-trivial realization of the *Diff* symmetry otherwise being absent apparently. From the recovered Lorentz symmetry in the action on the *super-selected* sector, which is still (considered as) an off-shell condition, we obtain the conservation laws with the Noether currents as in covariant theories. The general formula for the conserved Noether charge reproduces the mass of four-dimensional static black hole with an *arbitrary* cosmological constant in Hořava gravity, and is independent of ambiguities associated with the choice of asymptotic boundaries.

To this end, we start by considering the potential $\mathcal{V}[g^{ij}, R^i_{jkl}]$, which is an arbitrary function of metric g_{ij} and curvature invariants only, *eg.*,

$$-\mathcal{V} = \Lambda + \xi R + \alpha R^n + \beta (R_{ij}R^{ij})^s + \gamma (R^i_{jkl}R_i{}^{jkl})^r + \dots, \quad (4)$$

but without (covariant) derivatives, for simplicity. In order that the construction of a renormalizable action (1) is *not* spoiled by the mixing of space and time (derivatives) in the general coordinate transformations, which could produce ghosts of higher-time derivatives from the higher-spatial-derivative terms in the potential, we need to further constrain the allowed coordinate transformations into the foliation-preserving diffeomorphism ($Diff_{\mathcal{F}}$) [3],

$$\delta_{\xi} t = \tilde{\xi}^0(t), \quad \delta_{\xi} x^i = \xi^i(t, \mathbf{x}), \quad (5)$$

$$\delta_{\xi} N = (N\tilde{\xi}^0)_{,0} + \xi^k \nabla_k N, \quad (6)$$

$$\delta_{\xi} N_i = \tilde{\xi}^0_{,0} N_i + \xi^j_{,0} g_{ij} + \nabla_i \xi^j N_j + N_{i,0} \tilde{\xi}^0 + \nabla_j N_i \xi^j, \quad (7)$$

$$\delta_{\xi} g_{ij} = \nabla_i \xi^k g_{kj} + \nabla_j \xi^k g_{ki} + g_{ij,0} \tilde{\xi}^0. \quad (8)$$

Then, from some straightforward computations [3], one can show that the standard action (1) is invariant under $Diff_{\mathcal{F}}$ (6) - (8)

$$\delta_{\xi} S = \int dt d^D x \left\{ \partial_t [\tilde{\xi}^0(t) \mathcal{L}] + \partial_i [\xi^i(\mathbf{x}, t) \mathcal{L}] \right\}, \quad (9)$$

which reflects the scalar density nature of the Lagrangian density \mathcal{L} , defined by $S \equiv \int dt d^D x \mathcal{L}$. Here, each term in the potential as well as in the kinetic part of (1) is “separately” invariant for an arbitrary λ ¹ and all the other parameters in the potential \mathcal{V} so that the higher-time-derivative terms, as well as the mixing terms between the time and spatial-derivative terms, from the (Lorentz) transformation of the potential with higher-spatial derivative terms would not occur.

If we consider Einstein-Hilbert (EH) action in GR with $\lambda = 1$ and $-\mathcal{V} = \Lambda + (2/\kappa^2)R$, then there is an “*accidental*” symmetry enhancement which mixes each term in the action [6] so that we can recover the full $Diff$ [2] [$\xi^{\mu} \equiv (\xi^0, \xi^i)$],

$$\delta_{\xi} S_{EH} = \int dt d^D x \partial_{\mu} [\xi^{\mu}(\mathbf{x}, t) \mathcal{L}_{EH}] \quad (10)$$

with

$$\delta_{\xi} t = \xi^0(t, \mathbf{x}), \quad \delta_{\xi} x^i = \xi^i(t, \mathbf{x}), \quad (11)$$

$$\delta_{\xi} N = (N\xi^0)_{,0} - N\nabla_i \xi^0 g^{ij} N_j + \xi^k \nabla_k N, \quad (12)$$

$$\delta_{\xi} N_i = \xi^0_{,0} N_i + \xi^j_{,0} g_{ij} + \nabla_i \xi^0 (g^{kl} N_k N_l - N^2) + \nabla_i \xi^j N_j + N_{i,0} \xi^0 + \nabla_j N_i \xi^j, \quad (13)$$

$$\delta_{\xi} g_{ij} = \nabla_i \xi^0 N_j + \nabla_j \xi^0 N_i + \nabla_i \xi^k g_{kj} + \nabla_j \xi^k g_{ki} + g_{ij,0} \xi^0. \quad (14)$$

¹ For the case $\lambda = 1/D$, a separate consideration is needed [3]. We will briefly discuss about this in the discussion No. 4.

This shows a *sudden reduction* of the *Diff* symmetry beyond the GR limit in Hořava gravity but it is not quite satisfactory due to lack of canonical understanding of missing (Lorentz) symmetry. On the other hand, in the Hamiltonian formalism, the symmetries of an action are revealed in the existence of constraints between the field variables and their conjugate momenta, which being the *canonical* generators of the symmetry transformations. In EH case, there are $2(D + 1)$ *first-class* constraints which generate the full *Diff* transformations (12) - (14) so that we have $(D + 1)(D - 2)$ physical graviton (transverse traceless) modes. Whereas, in HLD case with the action (1), the constraint structure is quite different, having the *second-class* constraints also due to Lorentz-violating terms, but we have recently found [5] that

“there exists the case (called Case A) where the same number of degrees of freedom can be recovered as in GR, at the “fully non-linear” level.”

This may suggest that, even though its “apparent” symmetry is just the Lorentz-violating *Diff_F*, the “full” symmetry of HLD action can be as large as *Diff* in GR, in order to be consistent with the Hamiltonian analysis!

In order to examine this, which may fill the gap in those two sharply different symmetries in (5) - (8) and (11) - (14), we study the full *Diff* with an arbitrary $\xi^\mu(\mathbf{x}, t)$ to see if the Hořava gravity action (1) can be invariant in a non-trivial way, just beyond the *apparent* symmetry of *Diff_F*. To this end in a canonical way, we start by considering the variation of the action (1) with the potential $\mathcal{V}[g^{ij}, R^i{}_{jkl}]$, under the arbitrary variations of the ADM variables,

$$\delta S = \int dt d^D x \left[-\mathcal{H} \delta N - \mathcal{H}^i \delta N_i + \mathbf{E}^{ij} \delta g_{ij} + \partial_\mu \Theta^\mu (\delta N_i, \delta g_{ij}) \right], \quad (15)$$

where we have the bulk terms with

$$\mathcal{H} \equiv -\frac{\delta S}{\delta N} = \sqrt{g} \left[\left(\frac{2}{\kappa^2} \right) (K_{ij} K^{ij} - \lambda K^2) + \mathcal{V} \right], \quad (16)$$

$$\mathcal{H}^i \equiv -\frac{\delta S}{\delta N_i} = -2\sqrt{g} \left(\frac{2}{\kappa^2} \right) \nabla_j (K^{ij} - \lambda g^{ij} K), \quad (17)$$

$$\mathbf{E}^{ij} \equiv \frac{\delta S}{\delta g_{ij}} = E_{(0)}^{ij} - \sqrt{g} \left[NP^{iklm} R^j{}_{klm} + \frac{1}{2} N g^{ij} \mathcal{V}[g^{ij}, R^i{}_{jkl}] - 2 \nabla_k \nabla_l (NP^{iklj}) \right], \quad (18)$$

and the boundary terms $\Theta^\mu(\delta N_i, \delta g_{ij})$ with

$$\Theta^0 \equiv \sqrt{g} \left(\frac{2}{\kappa^2} \right) (K^{ij} - \lambda g^{ij} K) \delta g_{ij}, \quad (19)$$

$$\begin{aligned} \Theta^i \equiv & \sqrt{g} \left(\frac{2}{\kappa^2} \right) (2N^l G^{ijkl} K_{km} \delta g_{jl} - N^i G^{lkmn} K_{mn} \delta g_{jl} - 2G^{kjil} K_{kj} \delta N_l) \\ & + 2\sqrt{g} P^{jkil} N \nabla_k \delta g_{lj} - 2\sqrt{g} \delta g_{lj} \nabla_k (P^{jkl} N), \end{aligned} \quad (20)$$

where $G^{ijkl} \equiv \delta^{ijkl} - \lambda g^{ij} g^{kl}$ is the DeWitt metric [2]. Here, the tensor

$$P_i{}^{jkl} \equiv \left(\frac{\partial \mathcal{L}}{\partial R^i{}_{jkl}} \right)_{g^{mn}} = - \left(\frac{\partial \mathcal{V}}{\partial R^i{}_{jkl}} \right)_{g^{mn}}, \quad (21)$$

by treating g^{ij} and $R^i{}_{jkl}$ are independent fields [7], has the same symmetries in the indices as those of Riemann tensor $R^i{}_{jkl}$. (For the explicit forms for \mathbf{E}^{ij} and Θ^i , see Appendix A).

Plugging *Diff* transformations (12) - (14) into the arbitrary variation (15) and doing some straightforward computations, we obtain the action transformation [8],

$$\delta_\xi S = \int dt d^D x \left[-\mathcal{H} \delta_\xi N - \mathcal{H}^i \delta_\xi N_i + \mathbf{E}^{ij} \delta_\xi g_{ij} + \partial_\mu \Theta^\mu (\delta_\xi N_i, \delta_\xi g_{ij}) \right] \quad (22)$$

$$= \int dt d^D x \left[\xi^0 \mathcal{I}_0 + \xi^i \mathcal{I}_i + \partial_\mu \Psi^\mu (\delta_\xi N_i, \delta_\xi g_{ij}) \right], \quad (23)$$

where

$$\mathcal{I}_0 \equiv N \dot{\mathcal{H}} - \nabla_m (N N^m \mathcal{H}) + N_i \dot{\mathcal{H}}^i + \nabla_m [\mathcal{H}^m (g^{jl} N_j N_l - N^2)] + \dot{g}_{ij} \mathbf{E}^{ij} - 2 \nabla_m (N_i \mathbf{E}^{mi}), \quad (24)$$

$$\mathcal{I}_i \equiv (g_{ij} \mathcal{H}^j)_{,0} + \nabla_m (\mathcal{H}^m N_i) - \mathcal{H} \nabla_i N - \mathcal{H}^j \nabla_i N_j - 2 g_{ij} \nabla_m \mathbf{E}^{jm}, \quad (25)$$

$$\Psi^0 \equiv -\xi^0 (N \mathcal{H} + N_i \mathcal{H}^i) - \xi^j g_{ij} \mathcal{H}^i + \Theta^0, \quad (26)$$

$$\Psi^i \equiv \xi^0 [N N^i \mathcal{H} - \mathcal{H}^i (g^{lj} N_l N_j - N^2) + 2 N_j \mathbf{E}^{ij}] + \xi^j (-N_j \mathcal{H}^i + 2 g_{jl} \mathbf{E}^{il}) + \Theta^i. \quad (27)$$

Here, it is important to note that the formal expressions of (24) - (27) are generally valid for arbitrary potential $\mathcal{V}[g^{ij}, R^i{}_{jkl}]$ though the explicit expressions of \mathcal{H} , \mathcal{H}^i , \mathbf{E}^{ij} , Θ^μ , Ψ^μ may depend on the potential form.

Now, by expressing the time-derivatives in terms of extrinsic curvature K_{ij} in (2) and using the definitions (16) - (18), *but without using the dynamical equations of motion* $\mathbf{E}^{ij} = 0$, *nor the constraints* $\mathcal{H} \approx 0$, $\mathcal{H}^i \approx 0$, we obtain

$$\begin{aligned} \mathcal{I}_0 &= \nabla_i \left\{ 2N^2 \left[\nabla_j \pi^{ij} + \left(\frac{\kappa^2}{2} \right) \left(\frac{2\lambda}{\lambda D - 1} (\pi \nabla_l P^{kl}{}_{i} - P^{kl}{}_{i} \nabla_l \pi) + 2P_{jkl}{}^i \nabla^k \pi^{jl} - 2\pi^{jl} \nabla^k P_{jkl}{}^i \right) \right] \right\} \\ &\equiv \nabla_i \Omega^i, \end{aligned} \quad (28)$$

$$\mathcal{I}_i = 0, \quad (29)$$

$$\Psi^0 = \xi^0 \mathcal{L} + \partial_i \mathcal{U}^{0i}, \quad (30)$$

$$\Psi^i = \xi^i \mathcal{L} + \Sigma^i + \partial_0 \mathcal{U}^{i0} + \partial_j \mathcal{U}^{ij}, \quad (31)$$

where Σ^i and $\mathcal{U}^{\mu\nu} = -\mathcal{U}^{\nu\mu}$, which is called the ‘‘super-potential’’ [9, 10], are given by

$$\begin{aligned} \Sigma^i &= 2N^2 \left[\left(\frac{\kappa^2}{2} \right) \left(\frac{2\lambda}{\lambda D - 1} (P^{li}{}_{lk} \nabla^k (\xi^0 \pi) - \xi^0 \pi \nabla^k P^{li}{}_{lk}) + 2\xi^0 \pi^{jl} \nabla^k P_{jkl}{}^i - 2P_{jkl}{}^i \nabla^k (\xi^0 \pi^{jl}) \right) \right. \\ &\quad \left. + \pi^{ij} \nabla_j \xi^0 - \xi^0 \nabla_j \pi^{ij} \right], \end{aligned} \quad (32)$$

$$\mathcal{U}^{0i} = -\mathcal{U}^{i0} = 2\sqrt{g} (\xi^0 N_j + \xi_j) \left(\frac{2}{\kappa^2} \right) (K^{ij} - \lambda g^{ij} K), \quad (33)$$

$$\mathcal{U}^{ij} = -\mathcal{U}^{ji}. \quad (34)$$

(See Appendix A for the computational details of (28) - (31), and the explicit forms for \mathcal{U}^{ij} of (34)). Here, we note that $\mathcal{I}_i = 0$ identically, which is the analogue of the spatial component

of the (contracted) Bianchi identity $\hat{\nabla}_\mu G^{\mu i} = 0$ in GR, for the Einstein tensor $G^{\mu\nu} = \hat{R}^{\mu\nu} - (1/2)\hat{g}^{\mu\nu}\hat{R}$ with the $(D+1)$ -dimensional Ricci tensor $\hat{R}^{\mu\nu}$, Ricci scalar \hat{R} [11] and covariant derivative $\hat{\nabla}_\mu$: This is due to the first-class nature of momentum constraint $\mathcal{H}_i \approx 0$, the generator of the spatial *Diff*, even in HLD gravity [5]. In \mathcal{I}_0 , we have replaced the extrinsic curvature with the canonical momenta $\pi_{ij} = (2/\kappa^2)\sqrt{g}(K_{ij} - \lambda K g_{ij})$ in order to compare with the Hamiltonian analysis.

In GR case, we have $\Sigma^i = 0$ and $\mathcal{I}_0 = 0$, which corresponds to the time-component of Bianchi identity, $\hat{\nabla}_\mu G^{\mu 0} = 0$ and hence obtain the usual Lorentz invariance of (10) from (23). For the *Diff* _{\mathcal{F} in Hořava gravity case, the parameter $\xi^0(t) \equiv \tilde{\xi}^0(t)$ for the $\xi^0 \mathcal{I}_0$ in (23) can be factorized out from the space integration and the $\xi^0 \mathcal{I}_0$ term turns into a (spatial) boundary term, but it is “exactly canceled” by another boundary term $\nabla_i \Sigma^i$ so that we can obtain the invariance of (9) under *Diff* _{\mathcal{F} .}}

For the full *Diff* with an arbitrary $\xi^0(\mathbf{x}, t)$, on the other hand, the $\mathcal{I}_0 (\neq 0)$ term can not be removed from the bulk terms anymore and may result the non-invariance of the HLD action generally, confirming the usual belief of its generic Lorentz violation. Therefore, the only way for obtaining the full *Diff* invariance of the HLD action, *if it exists*, would be to consider a “super-condition”,

$$\mathcal{I}_0 \equiv \nabla_i \Omega^i = 0, \quad (35)$$

which *super-selects* the Lorentz-invariant sector in HLD gravity. Then, it is remarkable that the super-condition can be written as

$$\mathcal{I}_0 = \Omega - \nabla_i (N^2 \mathcal{H}^i) = 0, \quad (36)$$

where

$$\Omega \equiv \nabla_i (N^2 C^i) \quad (37)$$

reduces to the *tertiary* constraint $\Omega \approx 0$ in Hamiltonian formalism using the dynamical equations of motion², *i.e.*, in *on-shell*, from $\dot{\mathcal{H}} \approx \Omega/N \approx 0$, with the Hamiltonian constraint $\mathcal{H} \approx 0$ and the momentum constraint $\mathcal{H}^i \equiv -2\nabla_j \pi^{ij} \approx 0$. In other words, the super-condition (35) is *on-shell* equivalent to the tertiary constraint in HLD gravity, which is trivial in GR, up to the Hamiltonian and momentum constraints, $\mathcal{H} \approx 0, \mathcal{H}^i \approx 0$, which are the secondary constraints.

³ This may justify our introduction of the super-condition (35) but it is important to note that

² $\dot{g}_{ij} = \{g_{ij}, H_C\}$, $\dot{\pi}^{ij} = \{\pi^{ij}, H_C\}$ with the canonical Hamiltonian $H_C = \int dx^D \{N\mathcal{H} + N_i \mathcal{H}^i\}$ [5].

³ From this result, one can use (24) as an off-shell, Lagrangian definition of the *tertiary* constraints via the terms of $N\dot{\mathcal{H}}$ and $N_i \dot{\mathcal{H}}^i$, which are usually quite cumbersome in Hamiltonian formalism. Actually, using the new definition in this paper we have generalized the Hamiltonian analysis on tertiary constraint for $\dot{\mathcal{H}}$ in [5], where we have considered only an arbitrary function of R in the potential (4). Moreover, one can find easily the exactly same constraint algebra for the Hamiltonian and momentum constraints as in the previous case [5], $\{\langle \eta \mathcal{H} \rangle, \langle \zeta \mathcal{H} \rangle\} = \langle \langle \eta \nabla_i \zeta - \zeta \nabla_i \eta \rangle C^i \rangle$, $\{\langle \eta \mathcal{H} \rangle, \langle \zeta^i \mathcal{H}_i \rangle\} = -\langle \zeta^i \nabla_i \eta \mathcal{H} \rangle$, $\{\langle \eta^i \mathcal{H}_i \rangle, \langle \zeta^j \mathcal{H}_j \rangle\} = \langle \langle \eta^i \nabla_i \zeta^j - \zeta^i \nabla_i \eta^j \rangle \mathcal{H}_j \rangle$, for the smeared constraints, $\langle \eta \mathcal{H} \rangle \equiv \int d^D x \eta \mathcal{H}$, *etc.*

the super-condition is *assumed* to be valid even *off-shell*⁴, *i.e.* prior to considering the “classical geometry dynamics” which extremises the action. In this way, the *full Diff* can be maintained all the way down to UV, even beyond the GR limit. Here, the super-condition $\mathcal{I}_0 = 0$ in HLD gravity corresponds to the temporal component of Bianchi identity, $\hat{\nabla}_\mu G^{\mu 0} = 0$ in GR.

Finally, from (15), (23), and with the help of the super condition $\mathcal{I}_0 = 0$ and the Bianchi identity $\mathcal{I}_i = 0$, we can obtain the Noether currents ($\Sigma^0 = 0$),

$$\begin{aligned}\mathcal{J}^\mu(\delta_\xi g) &\equiv \Theta^\mu(\delta_\xi N_i, \delta_\xi g_{ij}) - \Psi^\mu(\delta_\xi N_i, \delta_\xi g_{ij}) \\ &= \Theta^\mu - \xi^\mu \mathcal{L} - \Sigma^\mu - \partial_\nu \mathcal{M}^{\mu\nu},\end{aligned}\tag{38}$$

which satisfies

$$\partial_\mu \mathcal{J}^\mu(\delta_\xi g) = \mathcal{H} \delta_\xi N + \mathcal{H}^i \delta_\xi N_i - \mathbf{E}^{ij} \delta_\xi g_{ij}.\tag{39}$$

Note that the Noether currents satisfies the usual conservation laws *on-shell*, *i.e.* $\mathcal{H} = \mathcal{H}^i = \mathbf{E}^{ij} = 0$, for an arbitrary *Diff* transformation, but *off-shell* for Killing vectors ξ^μ , $\delta_\xi N = \delta_\xi N_i = \delta_\xi g_{ij} = 0$. The second term Σ^μ is due to the *apparent* non-covariance of the Horava action.⁵ Moreover, the last part in the Noether current (38), corresponds to the identically conserved or *off-shell current* $\mathcal{J}_{\text{off}} \equiv \partial_\nu \mathcal{M}^{\mu\nu}$ [9, 10, 13]. Then, the conserved charge passing through a hypersurface Σ is given by

$$Q(\xi) = \int_\Sigma d^D x \sqrt{g} n_\mu \mathcal{J}^\mu(\delta_\xi g)\tag{40}$$

for the unit normal vector n^μ of Σ and the physically *measurable* charge can be obtained by subtracting the background charge $\bar{Q}(\xi) \equiv \int_\Sigma d^D x \sqrt{\bar{g}} \bar{n}_\mu \bar{\mathcal{J}}^\mu(\delta_\xi \bar{g})$, *i.e.*, $Q(\xi)_{\text{phys}} \equiv Q(\xi) - \bar{Q}(\xi)$ generally, where the bars denote the background quantities.

In order to check the general Noether charge formula (40), let us consider the static metric ansatz

$$ds^2 = -N^2(r) dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_k^2\tag{41}$$

by which the original Horava gravity action in 3 + 1 dimensions [3] reduces to the case with the potential form of (4), due to the vanishing Cotton tensor, $C^{ij} \equiv \epsilon^{ijkl} \nabla_k (R^j_l - \delta^j_l R/4) = 0$. Here, $d\Omega_k^2$ denotes the line element for two-dimensional surface with a constant scalar curvature, $R^{(2)} = 2k$ for spherical, plane, and hyperbolic topologies with $k = +1, 0, -1$, respectively.

Then, for the time-like Killing vector $\xi^\mu = (1, 0, 0, 0)$ and the normal vector $n_\mu = (-N, 0, 0, 0)$, the only non-vanishing contributions in the Noether charge (40) come from the

⁴ From the form of the Bianchi identity in GR, $\bar{\nabla}_\mu G^{\mu\nu} = \bar{\nabla}_0 G^0_0 + \bar{\nabla}_i G^i_0 = (\mathcal{H}_{GR}/2\sqrt{\bar{g}})_{,0} + \dots + \nabla_i (\mathcal{H}^i/2N\sqrt{\bar{g}}) = 0$ with $G^0_0 = \mathcal{H}_{GR}/2\sqrt{\bar{g}}$, $G^i_0 = \mathcal{H}^i/2\sqrt{\bar{g}}N$ for the Hamiltonian constraint in GR, $\mathcal{H}_{GR} \approx 0$, the super-condition suggests *formally* the same Bianchi identity but now with $G^0_0 \equiv \mathcal{H}/2\sqrt{\bar{g}}$ in HLD gravity also. This provides a more fundamental *off-shell* reason for the appearance of the tertiary constraint via $\dot{\mathcal{H}}$.

⁵ The non-covariance term appears also in Chern-Simons theories [12].

second term in the current (38) and, after the angular integrations, is given by

$$\begin{aligned}
Q(\xi^0) &= \Omega_k \int_0^r dr r^2 \left(\frac{N}{\sqrt{f}} \right) \xi^0 \mathcal{L} \\
&= -\hat{\alpha} \xi^0 \left(\frac{N}{\sqrt{f}} \right) \left[-\frac{\lambda}{r} (f-k)^2 + 2(\omega - \Lambda_W) r (f-k) - \Lambda_W^2 r^3 \right] \Big|_0^r \\
&\quad + \hat{\alpha} \int_0^r dr \partial_r \left(\xi^0 \frac{N}{\sqrt{f}} \right) \left[-\frac{\lambda}{r} (f-k)^2 + 2(\omega - \Lambda_W) r (f-k) - \Lambda_W^2 r^3 \right] \\
&\quad - \hat{\alpha} (\lambda - 1) \int_0^r dr \xi^0 \left(\frac{N}{\sqrt{f}} \right) \left[\frac{(f-k)^2}{r^2} + \frac{(\partial_r f)^2}{2} \right]
\end{aligned} \tag{42}$$

in the usual conventions, with $\hat{\alpha} \equiv \Omega_k \kappa^2 \mu^2 / 8(3\lambda - 1)$, the IR-modification parameter ω , and D -dimensional cosmological constant parameter Λ_W [3, 14–18]. Here, it is important that we need to (1) first, change the Lagrangian into the total derivatives (the second line in the above formula) plus the bulk terms (the third and fourth lines), and then (2) second, compute the charge by plugging the known solutions: If we first plug the solutions into the Noether charge formula (42), we obtain the trivially vanishing charge because the Lagrangian in the charge formula is proportional to the Hamiltonian constraint $\mathcal{H} \approx 0$, which is solved by the solutions.⁶

Now, by plugging the general static vacuum solution for arbitrary cosmological constant parameter Λ_W and IR parameter ω with $\lambda = 1$ [16, 18], whose uniqueness is guaranteed by the corresponding Birkhoff's theorem [20] (for more general cases, see the discussion No. 8 below),

$$N^2 = f = k + (\omega - \Lambda_W) r^2 + \epsilon \sqrt{r[\omega(\omega - 2\Lambda_W) r^3 + \beta]}, \tag{43}$$

where $\epsilon = \pm 1$ and β is an integration constant⁷, we can obtain

$$Q(\xi^0) = \hat{\alpha} \beta, \tag{44}$$

which exactly agrees with the mass in the conventional Hamiltonian approach [17, 21, 22]. Note that the mass $\mathcal{M} \equiv Q(\xi^0)$ satisfies the first law of black hole thermodynamics

$$\delta \mathcal{M} = T_H \delta \mathcal{S} \tag{45}$$

with the black hole temperature T_H and the entropy \mathcal{S} with a logarithmic term, up to an arbitrary constant \mathcal{S}_0 ,

$$T_H \equiv \frac{\hbar \kappa|_H}{2\pi} = \frac{\hbar(3\Lambda_W^2 r_H^4 + 2k(\omega - \Lambda_W) r_H^2 - k^2)}{8\pi r_H (k + (\omega - \Lambda_W) r_H^2)}, \tag{46}$$

$$\mathcal{S} = \frac{4\pi \hat{\alpha}}{\hbar} ((\omega - \Lambda_W) r_H^2 + 2k \ln r_H) + \mathcal{S}_0 \tag{47}$$

⁶ This looks tricky but this kind of prescription seems to be essential to get the right answer. (For some related discussions, see also [19]) In particular, for $\lambda = 1$ and $(N/\sqrt{f}) = \text{constant}$, our charge (42) agrees with the (generalized) Misner-Sharp mass [17].

⁷ $\epsilon = -1$ (+1) represent an asymptotically flat or anti-de Sitter (de Sitter) with $\omega, \mu^2 > 0$ ($\omega, \mu^2 < 0$) [18].

for the surface gravity $\kappa_H = (1/2)\partial_r f|_{r_H}$ at the black hole horizon r_H .

Two remarkable properties of this result are as follows. First, the result (42) does not depend on the boundary $(D - 1)$ -hypersurface only if there is a time-like Killing vector inside the boundary. This means that the boundary needs not to be an asymptotic infinity even in asymptotically de-Sitter space as well as in flat or anti-de Sitter space (for similar results in covariant theories, see [25, 26]). Second, related to the first property, there are no divergences in anti-de Sitter space, and it is independent on the ambiguities associated with the choice of asymptotic boundary at $r \rightarrow \infty$ in de Sitter space [27]. So, for the asymptotically de Sitter black hole, the boundary can be any region between the outer black hole horizon r_+ and the cosmological horizon r_{++} , *i.e.* $r_+ < r < r_{++}$.

In conclusion, we have shown that the Lorentz symmetry, which is represented by *Diff* symmetry, is preserved on the super-selected sector of $\mathcal{I}_0 \equiv \nabla_i \Omega^i = 0$ even in HLD gravity action where the explicit Lorentz violating terms are introduced for (power-counting) renormalizability. This indicates that the full *Diff* symmetry of HLD action can be as large as the *Diff* in GR and, from the obtained full *Diff* symmetry, we find the conservation laws with the Noether currents as in covariant theories [10]. Several further remarks about challenging problems are in order.

1. The super-condition $\mathcal{I}_0 \equiv \nabla_i \Omega^i = 0$ is similar to the Maxwell's equation $\nabla_i B^i = 0$ for the magnetic field B^i without (magnetic) monopoles. If we define the $(D - 2)$ -form "currents" for D -dimensional space with the component $J^{i\dots n} = \epsilon^{i\dots njk} \nabla_j \Omega_k$, it satisfies the (spatial) conservation laws $\nabla_i J^{i\dots n} = 0$ as in the equations $j^i = \epsilon^{imn} \partial_m B_n$ for the magneto statistics with the electric currents j^i . Then, we can solve Ω^i in terms of the currents $J_{i\dots n}$, which are the additional data for a complete specification of Ω^i . For example, if the super-condition $\nabla_i \Omega^i = 0$ and $\epsilon^{i\dots njk} \nabla_j \Omega_k = J^{i\dots n} = 0$ are satisfied for the *whole-space* region, *i.e.* without singularities, then $\Omega^i = 0$ would be the only solution and this would correspond to "Case A" in the Hamiltonian analysis of [5] where the degrees of freedom in HLD gravity are the same as in GR at the fully non-linear level. Otherwise, Ω^i would be non-vanishing generally due to either (a) non-trivial topology/cohomology, or (b) singularities, or from (c) non-vanishing current $J^{i\dots n} \neq 0$. This latter case would correspond to "Case B" in [5] where an extra scalar graviton mode exists in Hamiltonian analysis of HLD gravity. It would be interesting to find the generic (formal) solution of Ω^i in curved space, corresponding to Biot-Savart's law in electromagnetism in Minkowski space-time.

2. From the invariance of the action (23), we have obtained the tertiary constraints $\nabla_i (N^2 C^i) = 0$ in Hamiltonian formalism via \mathcal{H} and $\dot{\mathcal{H}}^i$ in the super-condition $\mathcal{I}_0 = 0$. As have been noted above, the action invariance does not necessarily mean the same degrees of freedom as in GR, which is manifestly Lorentz invariant. In Hamiltonian formalism, we need to find a complete set of constraints to completely specify the degrees of freedom. This implies that we need more consistency analysis in Lagrangian formalism, corresponding to the preservation of constraints in Hamiltonian formalism. We suspect that the higher-order invariance of $\delta_\xi \delta_\eta \dots \delta_\zeta S = 0$ would be important in HLD gravity and needs to be considered in order to obtain the complete set of constraints, consistently with the Hamiltonian formalism.

3. Our formulation about the gravity sector is self-contained and independent on the matter sector. If we now consider matter action S_m as well, which may have non-relativistic higher-derivative terms also in accordance with the HLD gravity, the additional contributions to the action transformation are $\int dt d^D x [\xi^0 \sqrt{\hat{g}} \hat{\nabla}^\mu T_{\mu 0} + \xi^i \sqrt{\hat{g}} \hat{\nabla}^\mu T_{\mu i}]$ with the energy-momentum tensors for matters $T_{\mu\nu} = -\frac{2}{\sqrt{\hat{g}}} \frac{\delta S_m}{\delta \hat{g}^{\mu\nu}}$, for the $(D+1)$ -dimensional metric $\hat{g}_{\mu\nu}$ and its associated covariant derivatives $\hat{\nabla}^\mu$, together with the matter contributions to the boundary terms Θ^μ and Ψ^μ . But, from the super-condition $\mathcal{I}_0 = 0$ and the Bianchi identity $\mathcal{I}_i = 0$ in the gravity sector, which indicating their geometrical origin, the consistent theory with the full *Diff* is possible only for the covariantly-conserved matters, *i.e.* $\hat{\nabla}^\mu T_{\mu\nu} = 0$, regardless higher-derivatives in HLD gravity. In other words, only the energy-conserving matters $\hat{\nabla}^\mu T_{\mu 0} = 0$, as well as the momentum-conserving matters $\hat{\nabla}^\mu T_{\mu i} = 0$, can be consistent with the HLD gravity. Actually, there seems to exist some evidence for the covariant conservation laws of matter's energy momentum tensors and of the effective energy-momentum tensors from the higher-derivative terms, separately, for spherically symmetric case [28].⁸ It would be interesting to see whether the covariant form of the conservation laws holds generally, as another super-condition in matter sectors.

4. For the special value of IR Lorentz-violation parameter $\lambda = 1/D$, the theory has *anisotropic* Weyl symmetry additionally [3] but a separate consideration is needed. Based on the Hamiltonian analysis [5], which gives the same degrees of freedom as in GR, its full symmetry would be also as large as that of GR, though its details of the symmetry are different. It would be interesting to clarify the full action symmetry also and its connection to the Case A.

5. From the obtained Lorentz symmetry of HLD action for the super-selected sector of $\nabla_i \Omega^i = 0$, one can consider the corresponding *Ward-like* identity $\langle \delta_\xi \mathcal{F} \rangle - (i/\hbar) \int \xi \langle \mathcal{F} \partial_\mu \mathcal{J}^\mu \rangle dt d^D x = 0$ for a *Diff* invariant observable \mathcal{F} , from the *Diff* invariance of the path-integral measure with the first and second-class constraints [29]. The proof of renormalizability for HLD gravity from the gravitational Ward-like identity would be a challenging problem.

6. In our formulation, we have considered the arbitrary potential $\mathcal{V}[g^{ij}, R^i_{jkl}]$ without derivatives ∇_i . The inclusion of derivatives in the potential, *i.e.* $\mathcal{V}[g^{ij}, R^i_{jkl}, \nabla_i]$ would be more desirable to describe the most general systems as in the action (1). The computations are straightforward but, due to the complications, we have not yet succeeded in obtaining the canonical form of (23). However, we believe that the formulation itself should not depend on the existence of derivatives in the potential so that there should be no fundamental problem to get the corresponding canonical forms.

⁸ One can *formally* write the Horava gravity's equations of motion into a *covariant* Einstein's equation $G^{\mu\nu} = 8\pi G T_{\text{eff}}^{\mu\nu}$ by considering the higher-derivative contributions as the effective energy-momentum tensor $T_{\text{eff}}^{\mu\nu}$. Then, from the usual (covariant) Bianchi identity on the Einstein tensor $\hat{\nabla}_\mu G^{\mu\nu} = 0$, one can find the covariant conservation laws $\hat{\nabla}_\mu T_{\text{eff}}^{\mu\nu} = 0$ [28]. However, the geometric origin of this identity/property is still unknown.

7. The *Diff* invariance of HLD gravity sheds a new light on the very meaning of black hole entropy and its thermodynamical laws, due to the revival of Lorentz invariant concept of the event horizon, which has been essential to give an absolute meaning to black hole entropy as a *measure of observable ignorance* inside the event horizon as well as the role of the *universal horizons* in the presence of the (Lorentz-invariant) event horizons [30]. In particular, for the $k = 1$ black holes in an asymptotically flat/AdS space, the logarithmic correction to the usual Bekenstein-Hawking entropy implies the “positive” *minimum* of horizon radius r_H for the *positive* black hole entropy \mathcal{S} , which is consistent with the existence of a positive minimum for the mass \mathcal{M} where the Hawking temperature T_H vanishes [18].⁹ Moreover, in that case, the black hole entropy \mathcal{S} increases (the second law of black hole thermodynamics) by $\Delta\mathcal{S} = (8\pi\hat{\alpha}/\hbar)[(\omega - \Lambda_W)r_H + 2k/r_H]\Delta r_H = T_H\Delta\mathcal{M}$ for the increased mass $\Delta\mathcal{M} > 0$. From the associated increase of area¹⁰ $A_H = 4\pi r_H^2$, one can compute the upper bound to the energy of the gravitational radiations when one black hole captures another. For the asymptotically flat black holes ($\Lambda_W = 0$) [15] with the area ($\beta \equiv 4\omega M$)

$$A_H = 8\pi M^2[1 + (1 - (2\omega M^2)^{-1})^{1/2} - (4\omega M^2)^{-1}], \quad (48)$$

the increased area gives the inequality ($m \equiv \omega^{1/2}M$)

$$m_3^2[1 + (1 - (2m_3^2)^{-1})^{1/2}] > m_1^2[1 + (1 - (2m_1^2)^{-1})^{1/2}] + m_2^2[1 + (1 - (2m_2^2)^{-1})^{1/2}] - 1/4. \quad (49)$$

Then, the energy emitted in gravitational or other form of radiations is $m_1 + m_2 - m_3$ and its efficiency $\epsilon \equiv (m_1 + m_2 - m_3)/(m_1 + m_2)$ is limited by (49). The highest limit on ϵ is $1 - 3/4\sqrt{2} \approx 0.47$ which occurs when $m_1 = m_2 = 2^{-1/2}$, which are the minimum values for positive Hawking temperature $T_H > 0$, and $m_3 = 3/4$. On the other hand, when particles or fields which do not have horizons, impinge on a single black hole, one finds

$$m_2^2[1 + (1 - (2m_2^2)^{-1})^{1/2}] > m_1^2[1 + (1 - (2m_1^2)^{-1})^{1/2}]. \quad (50)$$

Note that m_2 can *not* be less than m_1 . This means that *one can not extract energy from a black hole* and there is *no analogue of the Penrose process* for Kerr or charged black hole in GR [31, 32]. This is basically due to fact that one can not turn-off the parameter ω arbitrarily, in contrast to the rotation parameter a in a Kerr black hole or the charge parameter e in a Reissner-Norström black hole even though they look similar in the black hole area formula (48) [33].¹¹

⁹ One can choose S_0 so that the two minimum horizon radii agree. This choice achieves the third law of black hole thermodynamics, *i.e.*, $\mathcal{S} = 0$ at $T_H = 0$.

¹⁰ This indicates the energy conditions, especially the weak energy conditions (WEC) and the null energy conditions (NEC) are not violated by the higher-derivative contributions to the (effective) energy-momentum tensors. For example, for the asymptotically flat case, *i.e.*, $\Lambda_W = 0$, see [28].

¹¹ Actually, (50) is exactly the same as that of the Kerr black hole case with the identification $a = 2^{-1}$, whereas (49) as that of Reissner-Norström black hole with $e = 2^{-1}$ in [33]. (For some earlier discussions on the similarity, see [23, 24])

These results are *quantitatively* (the lower highest limit $\epsilon = 1 - 2^{-1/2} \approx 0.29$ for two non-rotating initial black holes with the same masses [33]) and *qualitatively* (no energy extractions via an analogue of the Penrose process [31, 32]) different from GR black holes which could be tested experimentally in the near future.

8. The ambiguities associated with the choice of asymptotic boundary at $r \rightarrow \infty$ is absent in the charge formula (42) when we consider $\lambda = 1$ solution (43), where the second and the third (bulk) terms in (42) vanish. Note that this case covers a wide range of static (vacuum) solutions with higher curvatures, including those in GR [26]. However, for $\lambda \neq 1$ generally [34] which is beyond the GR limit, we still seems to need asymptotic boundary at $r \rightarrow \infty$, even for the asymptotically de Sitter black hole, in order to obtain its “finite” physical mass. The intimate physical connection between the IR-Lorentz violation and the need of an infinite boundary (of our universe) is still unclear. However, for the asymptotically de Sitter black hole, the choice of the asymptotic boundary at $r \rightarrow \infty$ might not be quite nonsensical because the possible communications between inside and outside of the cosmological horizon from the Lorentz violating effect.

Acknowledgments

We would like to thank Gokhan Alkac and Sang-Heon Yi for useful correspondences. DOD was supported by the National Natural Science Foundation of China under Grant No. 11875136 and the Major Program of the National Natural Science Foundation of China under Grant No. 11690021. MIP was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2020R1A2C1010372, 2020R1A6A1A03047877).

Appendix A: Computational details of (28) - (31), and explicit form of \mathcal{U}^{ij}

Here, we describe the computational details of (28) - (31), and the explicit form of \mathcal{U}^{ij} .

First, in order to compute \mathcal{I}_0 in (28) from (24), without using the dynamical equations of motion $\mathbf{E}^{ij} = 0$ nor the constraints $\mathcal{H} \approx 0, \mathcal{H}^i \approx 0$, we first consider the time derivative $\partial_t (\equiv (\dot{\quad}))$ of the potential \mathcal{V} , which appears in the term $N\dot{\mathcal{H}}$ in (24),

$$\begin{aligned} \frac{d\mathcal{V}}{dt} &= \frac{\partial\mathcal{V}}{\partial g_{ij}} \partial_t g_{ij} + \frac{\partial\mathcal{V}}{\partial R^i{}_{jkl}} \partial_t R^i{}_{jkl} \\ &= \dot{g}_{ij} P^{ilnp} R^j{}_{lnp} - P_i{}^{jkl} \dot{R}_{jkl}^i, \end{aligned} \quad (\text{A1})$$

where we have used [7]

$$\left(\frac{\partial\mathcal{V}}{\partial g^{ij}} \right)_{R^m{}_{nkl}} = -P_i{}^{lnp} R_{jlnp}. \quad (\text{A2})$$

Expressing the time derivatives in terms of extrinsic curvature via its definition (2), we have the relation

$$\begin{aligned}\frac{d\mathcal{V}}{dt} &= (2NK_{ij} + \nabla_i N_j + \nabla_j N_i) P^{ilnp} R^j_{lnp} - P_i^{jkl} \dot{R}^i_{jkl} \\ &= (2NK_{ij} + \nabla_i N_j + \nabla_j N_i) P^{ilnp} R^j_{lnp} - P_i^{jkl} (\nabla_k H^i_{jl} - \nabla_l H^i_{jk}),\end{aligned}\quad (\text{A3})$$

where we have used a useful relation,

$$\dot{R}^i_{jkl} = \nabla_k H^i_{jl} - \nabla_l H^i_{jk} \quad (\text{A4})$$

with

$$H^l_{ij} \equiv \nabla_i (NK_j{}^l) + \nabla_j (NK_i{}^l) - \nabla_l (NK_{ij}) + \nabla_{(i} \nabla_{j)} N^l - R^l_{(ij)}{}^m N_m. \quad (\text{A5})$$

With all these identities and (16) - (18), we can compute (24) as

$$\mathcal{I}_0 = \nabla_i \left\{ 2N^2 \left[\nabla_j \pi^{ij} + \frac{\kappa^2}{2} \left(\frac{2\lambda}{\lambda D - 1} (\pi \nabla_l P^{kl}{}^i - P^{kl}{}^i \nabla_l \pi) + 2P_{jkl}{}^i \nabla^k \pi^{jl} - 2\pi^{jl} \nabla^k P_{jkl}{}^i \right) \right] \right\} \quad (\text{A6})$$

$$\equiv \nabla_i \Omega^i, \quad (\text{A7})$$

in terms of the canonical momenta $\pi_{ij} = (2/\kappa^2)\sqrt{g}(K_{ij} - \lambda K g_{ij})$.

As an explicit example, if we consider the potential (4)

$$\begin{aligned}-\mathcal{V}[g^{ij}, R^i_{jkl}] &= \Lambda + \xi R + \alpha R^n + \beta (R_{ij} R^{ij})^s + \gamma (R^i_{jkl} R_i{}^{jkl})^r \\ &= \Lambda + \xi \delta_p^k g^{ql} R^p_{qkl} + \alpha (\delta_i^k g^{jl} R^i_{jkl})^n \\ &\quad + \beta (\delta_i^k R^i_{jkl} \delta_p^q R^p_{mqn} g^{mj} g^{ln})^s + \gamma (R^i_{jkl} R_i{}^{jkl})^r,\end{aligned}\quad (\text{A8})$$

$P_i{}^{jkl}$ is given by

$$P_i{}^{jkl} = \xi \delta_i^{[k} g^{l]j} + \alpha n R^{n-1} \delta_i^{[k} g^{l]j} + \beta s \mu^{s-1} (\delta_i^{[k} R^{l]j} + g^{j[l} R^{k]}_i) + 2\gamma r \rho^{r-1} R_i{}^{jkl}, \quad (\text{A9})$$

where we denote $\mu \equiv R_{ij} R^{ij}$, $\rho \equiv R_{ijkl} R^{ijkl}$.

By plugging (A9) into (A6), we obtain

$$\begin{aligned}
\mathcal{I}_0 &\equiv \nabla_i \Omega^i = \nabla_i \left[\Omega_{(0)}^i + \tilde{\xi} \Omega_{(1)}^i + \tilde{\alpha} \Omega_{(2)}^i + \tilde{\beta} \Omega_{(3)}^i + \tilde{\gamma} \Omega_{(4)}^i \right], \\
\Omega_{(0)}^i &= 2N^2 \nabla_j \pi^{ij}, \\
\Omega_{(1)}^i &= 2N^2 \left[\frac{(\lambda - 1)}{(\lambda D - 1)} \nabla^i \pi - \nabla_j \pi^{ij} \right], \\
\Omega_{(2)}^i &= 2nN^2 \left[\frac{(\lambda - 1)}{(\lambda D - 1)} (R^{n-1} \nabla^i \pi - \pi \nabla^i R^{n-1}) - (R^{n-1} \nabla_j \pi^{ij} - \pi^{ij} \nabla_j R^{n-1}) \right], \\
\Omega_{(3)}^i &= 2sN^2 \left[\frac{(2\lambda - 1)}{(\lambda D - 1)} (\mu^{s-1} R^{ij} \nabla_j \pi - \pi \nabla_j (\mu^{s-1} R^{ij})) - \frac{\lambda}{(\lambda D - 1)} (\mu^{s-1} R \nabla^i \pi - \pi \nabla^i (\mu^{s-1} R)) \right. \\
&\quad + (\mu^{s-1} R^{jk} \nabla^i \pi_{jk} - \pi_{jk} \nabla^i (\mu^{s-1} R^{jk})) - (\mu^{s-1} R^{ij} \nabla_k \pi_j^k - \pi_j^k \nabla_k (\mu^{s-1} R^{ij})) \\
&\quad \left. - (\mu^{s-1} R^{jk} \nabla_k \pi_j^i - \pi_j^i \nabla_k (\mu^{s-1} R^{jk})) \right], \\
\Omega_{(4)}^i &= 2rN^2 \left[\frac{4\lambda}{\lambda D - 1} (\pi \nabla_k (\rho^{r-1} R^{ik}) - \rho^{r-1} R^{ik} \nabla_k \pi) + 4\rho^{r-1} R_{jkl}{}^i \nabla^k \pi^{jl} - 4\pi^{jl} \nabla^k (\rho^{r-1} R_{jkl}{}^i) \right],
\end{aligned} \tag{A10}$$

where $\mu \equiv R_{ij} R^{ij}$, $\rho \equiv R_{ijkl} R^{ijkl}$, $(\tilde{\xi}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}) \equiv (\kappa^2/2)(\xi, \alpha, \beta, \gamma)$. On the other hand, if we consider \mathcal{I}_i in (25) similarly, one can find that it vanishes identically $\mathcal{I}_i \equiv 0$, which proves (29), as in GR or general covariant theories.

Similarly, if we consider Ψ^0 and Ψ^i in (26), (27), respectively, one can find that, from (19) and (20) as well as (16) - (18),

$$\begin{aligned}
\Psi^0 &\equiv -\xi^0 (N\mathcal{H} + N_i \mathcal{H}^i) - \xi^j g_{ij} \mathcal{H}^i + \Theta^0 \\
&= \xi^0 \mathcal{L} + \partial_i \left[2\sqrt{g} (\xi^0 N_j + \xi_j) \left(\frac{2}{\kappa^2} \right) G^{ijkl} K_{kl} \right]
\end{aligned} \tag{A11}$$

and

$$\begin{aligned}
\Psi^i &\equiv \xi^0 [N N^i \mathcal{H} - \mathcal{H}^i (g^{jk} N_j N_k - N^2) + 2N_j \mathcal{H}^{ij}] + \xi^j (-N_j \mathcal{H}^i + 2g_{jl} \mathbf{E}^{il}) + \Theta^i \\
&= \xi^i \mathcal{L} + \Sigma^i(\xi^0) - \partial_0 \left[2\sqrt{g} (\xi^0 N_j + \xi_j) \left(\frac{2}{\kappa^2} \right) G^{ijkl} K_{kl} \right] + \partial_j [\mathcal{A}^{ij}(\xi^0) + \mathcal{B}^{ij}(\xi^m)],
\end{aligned} \tag{A12}$$

where

$$\begin{aligned}
\Sigma^i(\xi^0) &= 2N^2 \left[\left(\frac{\kappa^2}{2} \right) \left(\frac{2\lambda}{\lambda D - 1} (P^{li}{}_{lk} \nabla^k (\xi^0 \pi) - \xi^0 \pi \nabla^k P^{li}{}_{lk}) + 2\xi^0 \pi^{jl} \nabla^k P_{jkl}{}^i - 2P_{jkl}{}^i \nabla^k (\xi^0 \pi^{jl}) \right) \right. \\
&\quad \left. + \pi^{ij} \nabla_j \xi^0 - \xi^0 \nabla_j \pi^{ij} \right]
\end{aligned} \tag{A13}$$

and \mathcal{A}^{ij} , \mathcal{B}^{ij} are antisymmetric tensors as

$$\begin{aligned}\mathcal{A}^{ij}(\xi^0) &\equiv 2\sqrt{g} \left[\left(\frac{2}{\kappa^2} \right) 2\xi^0 N_m N^{[j} G^{i]mkl} K_{kl} + P^{ijkl} (2\xi^0 N_l \nabla_k N + N \nabla_l (\xi^0 N_k)) + 4\xi^0 N N^l \nabla^k P_{[kl}^{j]i} \right], \\ \mathcal{B}^{ij}(\xi^m) &\equiv 2\sqrt{g} \left[\left(\frac{2}{\kappa^2} \right) 2\xi_m N^{[j} G^{i]mkl} K_{kl} + 4\xi^l \nabla^k (N P_{kl}^{[j} i]) + 2N P_{kl}^{[j} i] \nabla^k \xi^l \right].\end{aligned}\quad (\text{A14})$$

Then, one can write $\Psi^\mu \equiv \xi^\mu \mathcal{L} + \Sigma^\mu(\xi^0) + \partial_\nu \mathcal{U}^{\mu\nu}$ with $\Sigma^0 \equiv 0$ and the ‘‘super-potential’’ $\mathcal{U}^{\mu\nu}$, which is anti-symmetric $\mathcal{U}^{\mu\nu} = -\mathcal{U}^{\nu\mu}$ [9, 10] and given by

$$\mathcal{U}^{0i} = -\mathcal{U}^{i0} \equiv 2\sqrt{g}(\xi^0 N_j + \xi_j) \left(\frac{2}{\kappa^2} \right) (G^{ijkl} K_{kl}), \quad (\text{A15})$$

$$\mathcal{U}^{ij} = -\mathcal{U}^{ji} \equiv \mathcal{A}^{ij}(\xi^0) + \mathcal{B}^{ij}(\xi^m), \quad (\text{A16})$$

proving (30) and (31).

If we consider the potential (A8) with P_i^{jkl} tensor (A9), as an explicit example, one can find \mathbf{E}^{ij} in (18) as

$$\begin{aligned}\mathbf{E}^{ij} &\equiv E_{(0)}^{ij} + \xi E_{(1)}^{ij} + \alpha E_{(2)}^{ij} + \beta E_{(3)}^{ij} + \gamma E_{(4)}^{ij}, \quad (\text{A17}) \\ E_{(0)}^{ij} &= \sqrt{g} \left(\frac{2}{\kappa^2} \right) \left[-N^i \nabla_k K^{jk} - N^j \nabla_k K^{ik} + K^{ik} \nabla^j N_k + K^{jk} \nabla^i N_k + N^k \nabla_k K^{ij} \right. \\ &\quad \left. + 2N K^{ik} K^j_k - N K K^{ij} + \frac{1}{2} g^{ij} N K^{kl} K_{kl} - g^{ik} g^{jl} \dot{K}_{kl} \right] \\ &\quad + \lambda \sqrt{g} \left[\frac{1}{2} N g^{ij} K^2 + N^j \nabla^i K + N^i \nabla^j K - g^{ij} N^k \nabla_k K - g^{ij} K^{kl} \dot{g}_{kl} + g^{ij} g^{kl} \dot{K}_{kl} \right], \\ E_{(1)}^{ij} &= \sqrt{g} \left[N \left(-R^{ij} + \frac{1}{2} R g^{ij} + \frac{\Lambda}{\xi} g^{ij} \right) + (g^{il} g^{jk} - g^{ij} g^{kl}) \nabla_l \nabla_k N \right], \\ E_{(2)}^{ij} &= \sqrt{g} \left[N \left(-n R^{n-1} R^{ij} + \frac{1}{2} R^n g^{ij} \right) + n (g^{il} g^{jk} - g^{ij} g^{kl}) \nabla_l \nabla_k (N R^{n-1}) \right], \\ E_{(3)}^{ij} &= \sqrt{g} \left[N \left(-2s \mu^{s-1} R^{ik} R^j_k + \frac{1}{2} \mu^s g^{ij} \right) \right. \\ &\quad \left. + s (g^{ik} g^{jm} g^{ln} + g^{jk} g^{mi} g^{nl} - g^{kl} g^{mi} g^{nj} - g^{ij} g^{km} g^{ln}) \nabla_l \nabla_k (N \mu^{s-1} R_{mn}) \right], \\ E_{(4)}^{ij} &= \sqrt{g} \left[N \left(-2r \rho^{r-1} R^{iklm} R^j_{klm} + \frac{1}{2} \rho^r g^{ij} \right) + 4r \nabla_k \nabla_l (\rho^{r-1} N R^{iklj}) \right],\end{aligned}$$

and Θ^i in (20) as

$$\begin{aligned}
\Theta^i &\equiv \Theta_{(0)}^i + \xi \Theta_{(1)}^i + \alpha \Theta_{(2)}^i + \beta \Theta_{(3)}^i + \gamma \Theta_{(4)}^i, \\
\Theta_{(0)}^i &= \sqrt{g} \left(\frac{2}{\kappa^2} \right) [2N^l G^{ijklm} K_{km} \delta g_{jl} - N^i G^{lkmn} K_{mn} \delta g_{jl} - 2G^{kjil} K_{kj} \delta N_l], \\
\Theta_{(1)}^i &= 2\sqrt{g} g^{jk} g^{li} [N \nabla_k \delta g_{lj} - (\nabla_k N) \delta g_{lj}], \\
\Theta_{(2)}^i &= 4n\sqrt{g} [N g^{il} g^{kj} R^{n-1} \nabla_k \delta g_{lj} - \nabla_k (N g^{il} g^{kj} R^{n-1}) \delta g_{lj}], \\
\Theta_{(3)}^i &= 4s\sqrt{g} [N \mu^{s-1} g_{[l}^{[i} R_{k]}^{j]} \nabla^k \delta g_{lj}^l - \nabla^k (N \mu^{s-1} g_{[l}^{[i} R_{k]}^{j]} \delta g_{lj}^l)], \\
\Theta_{(4)}^i &= 4r\sqrt{g} \rho^{r-1} R^{jkil} N \nabla_k \delta g_{lj} - 4r\sqrt{g} \delta g_{lj} \nabla_k (\rho^{r-1} N R^{ijkl}).
\end{aligned} \tag{A18}$$

For Σ^i in (32), we find

$$\begin{aligned}
\Sigma^i(\xi^0) &\equiv \Sigma_{(0)}^i + \tilde{\xi} \Sigma_{(1)}^i + \tilde{\alpha} \Sigma_{(2)}^i + \tilde{\beta} \Sigma_{(3)}^i + \tilde{\gamma} \Sigma_{(4)}^i, \\
\Sigma_{(0)}^i &= 2N^2 \left(\frac{2}{\kappa^2} \right) [-\xi^0 \nabla_i \pi^{ij} + \pi^i_j \nabla^j \xi^0], \\
\Sigma_{(1)}^i &= 2N^2 [-\hat{\lambda} \nabla^i (\xi^0 \pi) + \xi^0 \nabla_i \pi^{ij} + \pi^i_j \nabla^j \xi^0], \\
\Sigma_{(2)}^i &= 2nN^2 [-\hat{\lambda} (R^{n-1} \nabla^i (\xi^0 \pi) - \xi^0 \pi \nabla^i R^{n-1})], \\
\Sigma_{(3)}^i &= 2sN^2 \left[\tilde{\lambda} (\mu^{s-1} R \nabla^i (\xi^0 \pi) - \xi^0 \pi \nabla^i (R \mu^{s-1})) \right. \\
&\quad \left. - \bar{\lambda} (\mu^{s-1} R^{ij} \nabla_j (\xi^0 \pi) - \xi^0 \pi \nabla_j (\mu^{s-1} R^{ij})) + (\xi^0 \pi^{jk} \nabla^i (\mu^{s-1} R_{jk}) - \mu^{s-1} R_{jk} \nabla^i (\mu^{s-1} R^{ij})) \right. \\
&\quad \left. + (\mu^{s-1} R_j^k \nabla^j (\xi^0 \pi_k) - \xi^0 \pi_k \nabla^j (\mu^{s-1} R_j^k)) + (\mu^{s-1} R^{ij} \nabla_k (\xi^0 \pi_j^k) - \xi^0 \pi_j^k \nabla_k (\mu^{s-1} R^{ij})) \right], \\
\Sigma_{(4)}^i &= 2rN^2 \left[4\tilde{\lambda} (\rho^{r-1} R^{ij} \nabla_j (\xi^0 \pi) - \xi^0 \pi \nabla_j (\rho^{r-1} R^{ij})) \right. \\
&\quad \left. + 4 (\rho^{r-1} R_{jkl}^i \nabla^l (\xi^0 \pi^{jk}) - \xi^0 \pi^{jk} \nabla^l (\rho^{r-1} R_{jkl}^i)) \right],
\end{aligned} \tag{A19}$$

where $\hat{\lambda} \equiv (\lambda - 1)/(\lambda D - 1)$, $\tilde{\lambda} \equiv \lambda/(\lambda D - 1)$, $\bar{\lambda} \equiv \hat{\lambda} + \tilde{\lambda} = (2\lambda - 1)/(\lambda D - 1)$, and \mathcal{A}^{ij} , \mathcal{B}^{ij} as

$$\begin{aligned}
\mathcal{A}^{ij}(\xi^0) &\equiv \mathcal{A}_{(0)}^{ij} + \xi \mathcal{A}_{(1)}^{ij} + \alpha \mathcal{A}_{(2)}^{ij} + \beta \mathcal{A}_{(3)}^{ij} + \gamma \mathcal{A}_{(4)}^{ij}, \\
\mathcal{A}_{(0)}^{ij} &= 2\sqrt{g} \left(\frac{2}{\kappa^2} \right) [2\xi^0 N_m N^{[j} G^{i]mkl} K_{kl}], \\
\mathcal{A}_{(1)}^{ij} &= 2\sqrt{g} [g^{ik} g^{lj} (2\xi^0 N_l \nabla_k N + N \nabla_l (\xi^0 N_k))], \\
\mathcal{A}_{(2)}^{ij} &= 2n\sqrt{g} [R^{n-1} g^{ik} g^{lj} (2\xi^0 N_l \nabla_k N + N \nabla_l (\xi^0 N_k)) + 4\xi^0 N N_l g^{il} g^{kj} \nabla_k R^{n-1}], \\
\mathcal{A}_{(3)}^{ij} &= 2s\sqrt{g} [2s\mu^{s-1} g_{[l}^{[j} R_{k]}^{i]} (2\xi^0 N^l \nabla^k N + N \nabla^l (\xi^0 N^k)) + 8\xi^0 N N^l g_{[l}^{[i} \nabla^{k]} (\mu^{s-1} R_{k]}^j)], \\
\mathcal{A}_{(4)}^{ij} &= 2r\sqrt{g} [8\xi^0 N N^l \nabla^k (\rho^{r-1} R_{kl}^{[j} \nabla^{i]}) + 2\rho^{r-1} R^{ijkl} (2\xi^0 N_l \nabla_k N + N \nabla_l (\xi^0 N_k))],
\end{aligned} \tag{A20}$$

and

$$\begin{aligned}
\mathcal{B}^{ij}(\xi^m) &\equiv \mathcal{B}_{(0)}^{ij} + \xi \mathcal{B}_{(1)}^{ij} + \alpha \mathcal{B}_{(2)}^{ij} + \beta \mathcal{B}_{(3)}^{ij} + \gamma \mathcal{B}_{(4)}^{ij}, \\
\mathcal{B}_{(0)}^{ij} &= 2\sqrt{g} \left(\frac{2}{\kappa^2} \right) [2\xi_m N^{[j} G^{i]mkl} K_{kl}], \\
\mathcal{B}_{(1)}^{ij} &= 2\sqrt{g} [2g^{l[i} g^{j]k} (N \nabla^k \xi_l - 2\xi_l \nabla^k N)], \\
\mathcal{B}_{(2)}^{ij} &= 2n\sqrt{g} [2g^{l[i} g^{j]k} R^{n-1} N \nabla^k \xi_l + 4\xi_l \nabla^k (g^{l[j} g^{i]k} R^{n-1} N)], \\
\mathcal{B}_{(3)}^{ij} &= 2s\sqrt{g} [4\mu^{s-1} g^{[j[k} R^{i]l]} N \nabla_k \xi_l + 8\xi_l \nabla_k (\mu^{s-1} g^{[j[k} R^{i]l]} N)], \\
\mathcal{B}_{(4)}^{ij} &= 2r\sqrt{g} [4\rho^{r-1} R^{k[ij]l} N \nabla_k \xi_l + 8\xi_l \nabla_k (\rho^{r-1} R^{k[ji]l} N)].
\end{aligned} \tag{A21}$$

-
- [1] E. M. Lifshitz, Zh. Eksp. Teor. Fiz., **11**, 255 & 269 (1941).
[2] B. S. DeWitt, Phys. Rev. **160**, 1113 (1967).
[3] P. Horava, Phys. Rev. D **79**, 084008 (2009).
[4] M. Visser, arXiv:0912.4757 [hep-th].
[5] D. O. Devecioglu and M. I. Park, Eur. Phys. J. C **80**, 597 (2020) [Erratum: Eur. Phys. J. C **80**, 764 (2020)].
[6] M. I. Park, Class. Quant. Grav. **28**, 015004 (2011) [arXiv:0910.1917 [hep-th]].
[7] V. Iyer and R. M. Wald, Phys. Rev. D **50**, 846 (1994). [gr-qc/9403028];
T. Padmanabhan, Phys. Rev. D **84**, 124041 (2011) [arXiv:1109.3846 [gr-qc]].
[8] N. Kiriushcheva, P. G. Komorowski and S. V. Kuzmin, arXiv:1108.6105 [gr-qc].
[9] P. von Freud, Ann. Math. **40**, 417 (1939).
[10] P. G. Bergman, Phys. Rev. **75**, 680 (1949); *ibid.*, **112**, 287 (1958); P. G. Bergmann and R. Schiller, *ibid.*, **89**, 4 (1953).
[11] A. Anderson and J. W. York, Jr., Phys. Rev. Lett. **81**, 1154 (1998). [arXiv:gr-qc/9807041 [gr-qc]].
[12] P. Oh and M. I. Park, Mod. Phys. Lett. A **14**, 231 (1999) [arXiv:hep-th/9805178 [hep-th]];
M. I. Park, Nucl. Phys. B **544**, 377 (1999) [arXiv:hep-th/9811033 [hep-th]];
S. N. Solodukhin, Phys. Rev. D **74**, 024015 (2006) [arXiv:hep-th/0509148 [hep-th]];
Y. Tachikawa, Class. Quant. Grav. **24**, 737 (2007) [arXiv:hep-th/0611141 [hep-th]];
W. Kim, S. Kulkarni and S. H. Yi, Phys. Rev. D **88**, 124004 (2013) [arXiv:1310.1739 [hep-th]].
[13] W. Kim, S. Kulkarni, and S.-H. Yi, Phys. Rev. Lett. **111**, 081101 (2013) [Erratum: *ibid.* **112**, 079902 (2014)].
[14] H. Lu, J. Mei and C. N. Pope, Phys. Rev. Lett. **103**, 091301 (2009) [arXiv:0904.1595 [hep-th]].
[15] A. Kehagias and K. Sfetsos, Phys. Lett. B **678**, 123 (2009) [arXiv:0905.0477 [hep-th]].
[16] M. I. Park, JHEP **09**, 123 (2009) [arXiv:0905.4480 [hep-th]].
[17] R. G. Cai, L. M. Cao and N. Ohta, Phys. Lett. B **679**, 504 (2009) [arXiv:0905.0751 [hep-th]];
R. G. Cai and N. Ohta, Phys. Rev. D **81**, 084061 (2010) [arXiv:0910.2307 [hep-th]].
[18] C. Argüelles, N. Grandi and M. I. Park, JHEP **10**, 100 (2015) [arXiv:1508.04380 [hep-th]].
[19] S. M. Carroll, *Spacetime and Geometry* (Cambridge University Press, 2019).
[20] D. O. Devecioglu and M. I. Park, Phys. Rev. D **99**, 104068 (2019) [arXiv:1804.05698 [hep-th]].

- [21] W. Donnelly and T. Jacobson, *Phys. Rev. D* **84**, 104019 (2011) [arXiv:1106.2131 [hep-th]].
- [22] J. Bellorin and A. Restuccia, *Phys. Rev. D* **84**, 104037 (2011) [arXiv:1106.5766 [hep-th]].
- [23] Y. S. Myung, *Phys. Lett. B* **678**, 127 (2009) [arXiv:0905.0957 [hep-th]].
- [24] A. Castillo and A. Larranaga, *Electron. J. Theor. Phys.* **8**, 83 (2011) [arXiv:0906.4380 [gr-qc]].
- [25] N. Deruelle and J. Katz, *Class. Quant. Grav.* **22**, 421 (2005) [arXiv:gr-qc/0410135 [gr-qc]].
- [26] B. P. Dolan, *Class. Quant. Grav.* **36**, 077001 (2019) [arXiv:1808.09081 [gr-qc]].
- [27] M. I. Park, *Phys. Lett. B* **440**, 275 (1998) [arXiv:hep-th/9806119 [hep-th]].
- [28] E. J. Son and W. Kim, *Phys. Rev. D* **83**, 124012 (2011) [arXiv:1011.3596 [gr-qc]].
- [29] P. Senjanovic, *Annals Phys.* **100**, 227 (1976) [Erratum: *Annals Phys.* **209**, 248 (1991)].
- [30] D. Blas and S. Sibiryakov, *Phys. Rev. D* **84**, 124043 (2011) [arXiv:1110.2195 [hep-th]].
- [31] R. Penrose, *Riv. Nuovo Cim.* **1**, 252 (1969).
- [32] D. Christodoulou and R. Ruffini, *Phys. Rev. D* **4**, 3552 (1971).
- [33] S. W. Hawking, *Phys. Rev. Lett.* **26**, 1344 (1971); *Commun. Math. Phys.* **25**, 152-166 (1972).
- [34] E. B. Kiritsis and G. Kofinas, *JHEP* **01**, 122 (2010) [arXiv:0910.5487 [hep-th]].