# ALMOST MINIMIZERS FOR A SINGULAR SYSTEM WITH FREE BOUNDARY

DANIELA DE SILVA, SEONGMIN JEON, AND HENRIK SHAHGHOLIAN

ABSTRACT. In this paper we study vector-valued almost minimizers of the energy functional

$$\int_D \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \, dx.$$

We establish the regularity for both minimizers and the "regular" part of the free boundary. The analysis of the free boundary is based on Weiss-type mono-tonicity formula and the epiperimetric inequality for the energy minimizers.

## Contents

1. Introduction	1
1.1. Singular cooperative system	1
1.2. Almost minimizers	2
1.3. Main results	2
1.4. Notation	3
2. Regularity of almost minimizers	4
3. Weiss-type monotonicity formula	7
4. Growth estimates	8
5. Non-degeneracy	13
6. 2-Homogeneous blowups	15
7. Regularity of the regular set	21
Appendix A. Example of almost minimizers	23
References	24

## 1. INTRODUCTION

1.1. Singular cooperative system. Let  $D \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an open set and consider the problem of minimizing the energy

(1.1) 
$$\int_D \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \, dx$$

among all functions  $\mathbf{u} = (u_1, \dots, u_m) \in W^{1,2}(D; \mathbb{R}^m), m \ge 1$ , satisfying  $\mathbf{u} = \mathbf{g}$ on  $\partial D$ , where  $\mathbf{g} : \partial B \to \mathbb{R}^m$  is a prescribed boundary data. This problem can be viewed as a vector-valued version of the classical *obstacle problem*. The energy minimizers are solutions of the singular system

(1.2) 
$$\Delta \mathbf{u} = \frac{\mathbf{u}}{|\mathbf{u}|} \chi_{\{|\mathbf{u}|>0\}},$$

Date: December 2, 2021.

<sup>2010</sup> Mathematics Subject Classification. Primary 35R35; Secondary 35J60.

Key words and phrases. Almost minimizers, Singular system, Regular set, Weiss-type monotonocity formula, Epiperimetric inequality.

DD is supported by a NSF grant (RTG 1937254). HS is supported by Swedish Research Council.

where  $\chi_{\{|\mathbf{u}|>0\}}$  is the characteristic function of  $\{|\mathbf{u}|>0\}$ . The regularity of both solutions  $\mathbf{u}$  of (1.2) and their free boundaries  $\partial\{|\mathbf{u}|>0\}$  was established in [1], with the help of monotonicity formulas and epiperimetric inequalities.

1.2. Almost minimizers. Given  $r_0 > 0$ , we say that a function  $\omega : [0, r_0) \rightarrow [0, \infty)$  is a gauge function or a modulus of continuity if  $\omega$  is monotone nondecreasing and  $\omega(0+) = 0$ .

**Definition 1.** Let  $r_0 > 0$  be a constant and  $\omega(r)$  be a gauge function. We say that  $\mathbf{u} \in W^{1,2}(B_1; \mathbb{R}^m)$  is an almost minimizer for the functional  $\int_D (|\nabla \mathbf{u}|^2 + 2|\mathbf{u}|)$ , with gauge function  $\omega(r)$ , if for any ball  $B_r(x_0) \in D$  with  $0 < r < r_0$  and for any  $\mathbf{v} \in \mathbf{u} + W_0^{1,2}(B_r(x_0); \mathbb{R}^m)$  we have

$$\int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) dx \le (1 + \omega(r)) \int_{B_r(x_0)} \left( |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| \right) dx.$$

These almost minimizers can be seen as perturbations of minimizers, or solutions of (1.2). Some examples of almost minimizers can be found in Appendix A.

The notion of almost minimizer was first introduced for the standard Dirichlet energy functional in [2]. Recently, almost minimizers for Alt-Caffarelli-type functionals were studied in [4], [3], [5], [6] etc., and for the thin obstacle problems were considered in [8], [9], [10].

In this paper, we are interested in the regularity of almost minimizers of (1.1) as well as the analysis of their free boundary.

1.3. Main results. Due to the technical nature of the problem, we restrict ourselves to the case when the gauge function  $\omega(r) = r^{\alpha}$ ,  $0 < \alpha < 2$ . Because we are concerned with local properties of almost minimizers and free boundaries, we simply assume that D is the unit ball  $B_1$  and the constant  $r_0 = 1$  in Definition 1.

We now state our first main result. Here and henceforth we use notation from Subsection 1.4.

**Theorem 1** (Regularity of almost minimizers). Let  $\mathbf{u} \in W^{1,2}(B_1; \mathbb{R}^m)$  be an almost minimizer of (1.1) in  $B_1$ . Then  $\mathbf{u} \in C^{1,\alpha/2}(B_1)$ . Moreover, for any  $K \subseteq B_1$ ,

$$\|\mathbf{u}\|_{C^{1,\alpha/2}(K)} \le C(n,\alpha,K)(E(\mathbf{u},1)^{1/2}+1).$$

Next, we consider a class of *half-space* solutions

 $\mathbb{H} := \{ \frac{\max(x \cdot \nu, 0)^2}{2} \mathbf{e} : \nu \text{ is a unit vector in } \mathbb{R}^n \text{ and } \mathbf{e} \text{ is a unit vector in } \mathbb{R}^m \}.$ 

Members of  $\mathbb{H}$  are 2-homogeneous global solutions of (1.2). For

$$M(\mathbf{v}) := \int_{B_1} \left( |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| \right) - 2 \int_{\partial B_1} |\mathbf{v}|^2,$$

we define

$$\frac{\beta_n}{2} := M\left(\frac{\max(x \cdot \nu, 0)^2}{2}\mathbf{e}\right).$$

Now, we state a Weiss-type monotonicity formula for almost minimizers, which plays a significant role in the analysis of the free boundary.

**Theorem 2** (Weiss-type monotonicity formula). Let **u** be an almost minimizer of (1.1) in  $B_1$ . For  $x_0 \in B_{1/2}$ , set

$$W(\mathbf{u}, x_0, t) := \frac{e^{at^{\alpha}}}{t^{n+2}} \left[ \int_{B_t(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) - \frac{2(1 - bt^{\alpha})}{t} \int_{\partial B_t(x_0)} |\mathbf{u}|^2 \right],$$

with

$$= \frac{n+2}{\alpha}, \quad b = \frac{n+4}{\alpha}.$$

Then  $W(\mathbf{u}, x_0, t)$  is nondecreasing in t for  $0 < t < t_0$ . Moreover, if  $x_0 \in \Gamma(\mathbf{u}) \cap B_{1/2}$  is a free boundary point, then we have either

(1.3) 
$$W(\mathbf{u}, x_0, 0+) = \beta_n/2 \quad or \quad W(\mathbf{u}, x_0, 0+) \ge \bar{\beta}_n$$

for some  $\bar{\beta}_n > \beta_n/2$ .

We are now able to define regular free boundary points.

a

**Definition 2.** We say that a free boundary point  $x_0 \in \Gamma(\mathbf{u})$  is regular if

$$W(\mathbf{u}, x_0, 0+) = \beta_n/2.$$

We denote by  $\mathfrak{R}_{\mathbf{u}}$  the set of all regular free boundary points of  $\mathbf{u}.$ 

Our main result concerning the regularity of the free boundary is as follows.

**Theorem 3** (Regularity of the regular set).  $\mathcal{R}_{\mathbf{u}}$  is a relatively open subset of the free boundary  $\Gamma(\mathbf{u})$  and locally a  $C^{1,\gamma}$ -manifold for some  $\gamma = \gamma(n, \alpha, \kappa) > 0$ , where  $\kappa$  is a constant as in Theorem 7.

Our proof is based on the Weiss-type monotonicity formula above and the epiperimetric inequality for solutions of (1.2). The proof uses a procedure similar to the one for solutions developed in [1].

1.4. Notation. Throughout this paper  $\mathbb{R}^n$  will be equipped with the Euclidean inner product  $x \cdot y$  and the induced norm |x|. For  $x_0 \in \mathbb{R}^n$  and r > 0,  $B_r(x_0)$  means the open *n*-dimensional ball of radius r, centered at  $x_0$  with boundary  $\partial B_r(x_0)$ . We typically drop the center from the notation if it is the origin.

When we consider a given set, we indicate by  $\nu$  the unit outward normal vector to the boundary and by  $\partial_{\theta} \mathbf{u} := \nabla \mathbf{u} - (\nabla \mathbf{u} \cdot \nu)\nu$  the surface derivative of a given function  $\mathbf{u}$ .

In integrals, we usually drop the variable and the measure of the integration if it is with respect to the Lebesgue measure or the surface measure. For example,

$$\int_{B_r} \mathbf{u} = \int_{B_r} \mathbf{u}(x) \, dx, \quad \int_{\partial B_r} \mathbf{u} = \int_{\partial B_r} \mathbf{u}(x) \, dS_x,$$

where  $S_x$  stands for the surface measure.

For a domain  $D \subset \mathbb{R}^n$  and an integrable function  $\mathbf{u},$  we denote its integral mean value by

$$\langle \mathbf{u} \rangle_D := \oint_D \mathbf{u} = \frac{1}{|D|} \int_D \mathbf{u}.$$

In particular, when  $D = B_r(x_0)$ , we simply write

$$\langle \mathbf{u} \rangle_{x_0,r} := \langle \mathbf{u} \rangle_{B_r(x_0)}.$$

For  $\mathbf{u} \in W^{1,2}(B_r)$ , we set

$$E(\mathbf{u},r) := \int_{B_r} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right).$$

If  $\mathbf{u} \in W^{1,2}(B_1)$ ,  $x_0 \in B_{1/2}$  and 0 < r < 1/2, we denote the 2-homogeneous recaling of  $\mathbf{u}$  by

$$\mathbf{u}_{x_0,r}(x) := \frac{\mathbf{u}(x_0 + rx)}{r^2}, \quad x \in B_{1/(2r)},$$

and the 2-homogeneous replacement of **u** in  $B_r$  (or equivalently  $\mathbf{u}_{x_0,r}$  in  $B_1$ ) by

$$\mathbf{c}_{x_0,r} := |x|^2 \mathbf{u}_{x_0,r}\left(\frac{x}{|x|}\right) = \frac{|x|^2}{r^2} \mathbf{u}\left(x_0 + \frac{r}{|x|}x\right), \quad x \in \mathbb{R}^n.$$

When  $x_0 = 0$ , we simply write  $\mathbf{u}_r$  and  $\mathbf{c}_r$  for  $\mathbf{u}_{0,r}$  and  $\mathbf{c}_{0,r}$ , respectively. We also indicate by  $\Gamma(\mathbf{u}) := \partial\{|\mathbf{u}| > 0\} \cap \{\nabla \mathbf{u} = 0\}$  the free boundary. Lastly, it may be worth noting that in the body of a proof, a constant may change from line to line and the dependence of the parameter of the problem is explicitly noted.

## 2. Regularity of almost minimizers

In this section, we follow the argument in [2] to obtain the regularity of almost minimizers.

**Proposition 1.** Let **u** be an almost minimizer in  $B_1$ . Then, there is  $C_0 = C_0(n) > 1$  such that

(2.1) 
$$\int_{B_{\rho}(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) \le C_0 \left[ \left( \frac{\rho}{r} \right)^n + r^{\alpha} \right] \int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) + C_0 r^{n+2}$$

whenever  $B_{\rho}(x_0) \subset B_r(x_0) \subset B_1$ .

*Proof.* Let  $\mathbf{h} \in W^{1,2}(B_r(x_0); \mathbb{R}^m)$  be such that  $\Delta \mathbf{h} = \mathbf{0}$  in  $B_r(x_0)$  and v = u on  $\partial B_r(x_0)$ . Since  $|\mathbf{h}|$  and  $|\nabla \mathbf{h}|^2$  are subharmonic, they satisfy the sub-mean value properties

(2.2) 
$$\int_{B_{\rho}(x_0)} |\mathbf{h}| \le \left(\frac{\rho}{r}\right)^n \int_{B_r(x_0)} |\mathbf{h}|, \quad \int_{B_{\rho}(x_0)} |\nabla \mathbf{h}|^2 \le \left(\frac{\rho}{r}\right)^n \int_{B_r(x_0)} |\nabla \mathbf{h}|^2.$$

Moreover, it follows from the weak equation  $\int_{B_r(x_0)} \nabla \mathbf{h} \nabla (\mathbf{u} - \mathbf{h}) = 0$  and the almost minimizing property of  $\mathbf{u}$  that

(2.3)  
$$\int_{B_r(x_0)} |\nabla(\mathbf{u} - \mathbf{h})|^2 = \int_{B_r(x_0)} |\nabla \mathbf{u}|^2 - |\nabla \mathbf{h}|^2$$
$$\leq \int_{B_r(x_0)} r^{\alpha} |\nabla \mathbf{h}|^2 + 2(1 + r^{\alpha}) |\mathbf{h}| - 2|\mathbf{u}|$$
$$= \int_{B_r(x_0)} r^{\alpha} |\nabla \mathbf{h}|^2 + 2(1 + r^{\alpha}) (|\mathbf{h}| - |\mathbf{u}|) + 2r^{\alpha} |\mathbf{u}|$$
$$\leq \int_{B_r(x_0)} r^{\alpha} |\nabla \mathbf{u}|^2 + 2(1 + r^{\alpha}) |\mathbf{h} - \mathbf{u}| + 2r^{\alpha} |\mathbf{u}|,$$

where in the last inequality we used the fact that **h** is the energy minimizer of the Dirichlet integral  $\int_{B_r(x_0)} |\nabla \mathbf{h}|^2$ . We also apply Poincarè inequality and Young's inequality to obtain

$$5\int_{B_r(x_0)} |\mathbf{u} - \mathbf{h}| \le C(n)r \int_{B_r(x_0)} |\nabla(\mathbf{u} - \mathbf{h})| \le \frac{1}{2} \int_{B_r(x_0)} |\nabla(\mathbf{u} - \mathbf{h})|^2 + C(n)r^{n+2}.$$

From this and (2.3), we have

$$\int_{B_r(x_0)} \left( |\nabla(\mathbf{u} - \mathbf{h})|^2 + 5|\mathbf{u} - \mathbf{h}| \right)$$
  
$$\leq \int_{B_r(x_0)} \left( \frac{1}{2} |\nabla(\mathbf{u} - \mathbf{h})|^2 + 2(1 + r^{\alpha})|\mathbf{u} - \mathbf{h}| + r^{\alpha} |\nabla\mathbf{u}|^2 + 2r^{\alpha} |\mathbf{u}| \right) + C(n)r^{n+2},$$

From  $B_r(x_0) \subset B_1$ , we see that 0 < r < 1, thus  $2(1+r^{\alpha}) \leq 4$ . Then, it follows that (2.4)

$$\int_{B_r(x_0)} \left( |\nabla(\mathbf{u} - \mathbf{h})|^2 + |\mathbf{u} - \mathbf{h}| \right) \le C(n) r^\alpha \int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) + C(n) r^{n+2}.$$

Now, by combining (2.2) and (2.4), we obtain that for  $0 < \rho < r < r_0$ ,

$$\begin{split} &\int_{B_{\rho}(x_{0})} \left( |\nabla \mathbf{u}|^{2} + |\mathbf{u}| \right) \\ &\leq 2 \int_{B_{\rho}(x_{0})} \left( |\nabla \mathbf{h}|^{2} + |\nabla(\mathbf{u} - \mathbf{h})|^{2} + |\mathbf{h}| + |\mathbf{u} - \mathbf{h}| \right) \\ &\leq 2 \left( \frac{\rho}{r} \right)^{n} \int_{B_{r}(x_{0})} \left( |\nabla \mathbf{h}|^{2} + |\mathbf{h}| \right) + 2 \int_{B_{\rho}(x_{0})} \left( |\nabla(\mathbf{u} - \mathbf{h})|^{2} + |\mathbf{u} - \mathbf{h}| \right) \\ &\leq 4 \left( \frac{\rho}{r} \right)^{n} \int_{B_{r}(x_{0})} \left( |\nabla \mathbf{u}|^{2} + 2|\mathbf{u}| + |\nabla(\mathbf{u} - \mathbf{h})|^{2} + 2|\mathbf{u} - \mathbf{h}| \right) \\ &\quad + 2 \int_{B_{\rho}(x_{0})} \left( |\nabla(\mathbf{u} - \mathbf{h})|^{2} + |\mathbf{u} - \mathbf{h}| \right) \\ &\leq 4 \left( \frac{\rho}{r} \right)^{n} \int_{B_{r}(x_{0})} \left( |\nabla \mathbf{u}|^{2} + 2|\mathbf{u}| \right) + 10 \int_{B_{r}(x_{0})} \left( |\nabla(\mathbf{u} - \mathbf{h})|^{2} + |\mathbf{u} - \mathbf{h}| \right) \\ &\leq C(n) \left[ \left( \frac{\rho}{r} \right)^{n} + r^{\alpha} \right] \int_{B_{r}(x_{0})} \left( |\nabla \mathbf{u}|^{2} + |\mathbf{u}| \right) + C(n)r^{n+2}. \end{split}$$

From here, we deduce the almost Lipschitz regularity of  $\mathbf{u}$  with the help of the following lemma, whose proof can be found in [7].

**Lemma 1.** Let  $r_0 > 0$  be a positive number and let  $\varphi : (0, r_0) \to (0, \infty)$  be a nondecreasing function. Let  $a, \beta$ , and  $\gamma$  be such that  $a > 0, \gamma > \beta > 0$ . There exist two positive numbers  $\varepsilon = \varepsilon(a, \gamma, \beta), c = c(a, \gamma, \beta)$  such that, if

$$\varphi(\rho) \le a \Big[ \Big( \frac{\rho}{r} \Big)^{\gamma} + \varepsilon \Big] \varphi(r) + b r^{\beta}$$

for all  $\rho$ , r with  $0 < \rho \le r < r_0$ , where  $b \ge 0$ , then one also has, still for  $0 < \rho < r < r_0$ ,

$$\varphi(\rho) \le c \Big[ \Big(\frac{\rho}{r}\Big)^{\beta} \varphi(r) + b \rho^{\beta} \Big].$$

**Theorem 4.** Let  $\mathbf{u}$  be an almost minimizer in  $B_1$ . Then  $\mathbf{u} \in C^{0,\sigma}(B_1)$  for all  $0 < \sigma < 1$ . Moreover, for any  $K \subseteq B_1$ ,

$$\|\mathbf{u}\|_{C^{0,\sigma}(K)} \le C\left(E(\mathbf{u},1)^{1/2}+1\right)$$

with  $C = C(n, \alpha, \sigma, K)$ .

c

*Proof.* For given  $K \in B_1$  and  $x_0 \in K$ , take  $\delta = \delta(n, \alpha, \sigma, K) > 0$  such that  $\delta < \text{dist}(K, \partial B_1)$  and  $\delta^{\alpha} \leq \varepsilon(C_0, n, n+2\sigma-2)$ , where  $C_0 = C_0(n)$  is as in Proposition 1 and  $\varepsilon = \varepsilon(C_0, n, n+2\sigma-2)$  is as in Lemma 1. Then, by (2.1), for  $0 < \rho < r < \delta$ ,

$$\int_{B_{\rho}(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) \le C_0 \left[ \left( \frac{\rho}{r} \right)^n + \varepsilon \right] \int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) + C_0 r^{n+2\sigma-2}$$

By applying Lemma 1, we obtain

$$\int_{B_{\rho}(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) \le C(n,\sigma) \left[ \left( \frac{\rho}{r} \right)^{n+2\sigma-2} \int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) + \rho^{n+2\sigma-2} \right].$$

Taking  $r \nearrow \delta(n, \alpha, \sigma, K)$ , we get

(2.5) 
$$\int_{B_{\rho}(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) \leq C(n, \alpha, \sigma, K) \left( E(\mathbf{u}, 1) + 1 \right) \rho^{n+2\sigma-2},$$

for  $0 < \rho < \delta$ . In particular, we have

$$\int_{B_{\rho}(x_0)} |\nabla \mathbf{u}|^2 \le C(n, \alpha, \sigma, K) \left( E(\mathbf{u}, 1) + 1 \right) \rho^{n+2\sigma-2},$$

and by Morrey space embedding we conclude  $\mathbf{u} \in C^{0,\sigma}(K)$  with

$$\|\mathbf{u}\|_{C^{0,\sigma}(K)} \le C(n,\alpha,\sigma,K) \left( E(\mathbf{u},1)^{1/2} + 1 \right).$$

Now we prove Theorem 1 using the above almost Lipschitz estimate of almost minimizers.

Proof of Theorem 1. For  $K \in B_1$ , fix a small  $r_0 = r_0(n, \alpha, K) > 0$  to be chosen later. Particularly, we ask  $r_0 < \operatorname{dist}(K, \partial B_1)$ . For  $x_0 \in K$  and  $0 < r < r_0$ , let  $\mathbf{h} \in W^{1,2}(B_r(x_0); \mathbb{R}^m)$  be a harmonic function such that  $\mathbf{h} = \mathbf{u}$  on  $\partial B_r(x_0)$ . Then, by (2.4) and (2.5) with  $\sigma = 1 - \alpha/4 \in (0, 1)$ ,

(2.6) 
$$\int_{B_{r}(x_{0})} |\nabla(\mathbf{u} - \mathbf{h})|^{2} \leq C(n) r^{\alpha} \int_{B_{r}(x_{0})} \left( |\nabla \mathbf{u}|^{2} + |\mathbf{u}| \right) + C(n) r^{n+2} \\ \leq C(n, \alpha, K) \left( E(\mathbf{u}, 1) + 1 \right) r^{n+\alpha/2} + C(n) r^{n+2} \\ \leq C(n, \alpha, K) \left( E(\mathbf{u}, 1) + 1 \right) r^{n+\alpha/2},$$

for  $0 < r < r_0(n, \alpha, K)$ . Note that since **h** is harmonic in  $B_r(x_0)$ , for  $0 < \rho < r$ 

$$\int_{B_{\rho}(x_0)} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_0,\rho}|^2 \le \left(\frac{\rho}{r}\right)^{n+2} \int_{B_r(x_0)} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_0,r}|^2.$$

Moreover, by Jensen's inequality,

$$\begin{split} &\int_{B_{\rho}(x_{0})} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_{0},\rho}|^{2} \\ &\leq 3 \int_{B_{\rho}(x_{0})} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_{0},\rho}|^{2} + |\nabla (\mathbf{u} - \mathbf{h})|^{2} + |\langle \nabla (\mathbf{u} - \mathbf{h}) \rangle_{x_{0},\rho}|^{2} \\ &\leq 3 \int_{B_{\rho}(x_{0})} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_{0},\rho}|^{2} + 6 \int_{B_{\rho}(x_{0})} |\nabla (\mathbf{u} - \mathbf{h})|^{2}, \end{split}$$

and similarly,

$$\int_{B_r(x_0)} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_0, r}|^2 \le 3 \int_{B_r(x_0)} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_0, r}|^2 + 6 \int_{B_r(x_0)} |\nabla (\mathbf{u} - \mathbf{h})|^2.$$

Now, we use the above inequalities to obtain

$$\begin{split} &\int_{B_{\rho}(x_{0})} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_{0},\rho}|^{2} \\ &\leq 3 \int_{B_{\rho}(x_{0})} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_{0},\rho}|^{2} + 6 \int_{B_{\rho}(x_{0})} |\nabla (\mathbf{u} - \mathbf{h})|^{2} \\ &\leq 3 \left(\frac{\rho}{r}\right)^{n+2} \int_{B_{r}(x_{0})} |\nabla \mathbf{h} - \langle \nabla \mathbf{h} \rangle_{x_{0},r}|^{2} + 6 \int_{B_{\rho}(x_{0})} |\nabla (\mathbf{u} - \mathbf{h})|^{2} \\ &\leq 9 \left(\frac{\rho}{r}\right)^{n+2} \int_{B_{r}(x_{0})} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_{0},r}|^{2} + 24 \int_{B_{r}(x_{0})} |\nabla (\mathbf{u} - \mathbf{h})|^{2} \\ &\leq 9 \left(\frac{\rho}{r}\right)^{n+2} \int_{B_{r}(x_{0})} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_{0},r}|^{2} + C(n,\alpha,K) \left(E(\mathbf{u},1)+1\right) r^{n+\alpha/2}. \end{split}$$

Thus, applying Lemma 1 we get

$$\int_{B_{\rho}(x_{0})} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_{0},\rho}|^{2} \leq C(n,\alpha) \left(\frac{\rho}{r}\right)^{n+\alpha/2} \int_{B_{r}(x_{0})} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_{0},r}|^{2} + C(n,\alpha,K) \left(E(\mathbf{u},1)+1\right) \rho^{n+\alpha/2},$$

for  $0 < \rho < r < r_0$ . Taking  $r \nearrow r_0(n, \alpha, K)$ , we have

$$\int_{B_{\rho}(x_0)} |\nabla \mathbf{u} - \langle \nabla \mathbf{u} \rangle_{x_0,\rho}|^2 \le C(n,\alpha,K) \left( E(\mathbf{u},1) + 1 \right) \rho^{n+\alpha/2}$$

By Campanato space embedding, we obtain  $\nabla \mathbf{u} \in C^{0,\alpha/4}(K)$  with

$$\|\nabla \mathbf{u}\|_{C^{0,\alpha/4}(K)} \le C(n,\alpha,K)(E(\mathbf{u},1)^{1/2}+1).$$

In particular, we have

$$\|\nabla \mathbf{u}\|_{L^{\infty}(K)} \le C(n, \alpha, K)(E(\mathbf{u}, 1)^{1/2} + 1),$$

for any  $K \subseteq B_1$ . With this estimate, we can improve (2.6)

$$\begin{split} \int_{B_r(x_0)} |\nabla(\mathbf{u} - \mathbf{h})|^2 &\leq C(n) r^{\alpha} \int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + |\mathbf{u}| \right) + C(n) r^{n+2} \\ &\leq C(n, \alpha, K) \left( E(\mathbf{u}, 1) + 1 \right) r^{n+\alpha} + C(n) r^{n+2} \\ &\leq C(n, \alpha, K) \left( E(\mathbf{u}, 1) + 1 \right) r^{n+\alpha}, \end{split}$$

and by repeating the process above we conclude that  $\nabla \mathbf{u} \in C^{1,\alpha/2}(K)$  with

$$\|\nabla \mathbf{u}\|_{C^{0,\alpha/2}(K)} \le C(n,\alpha,K)(E(\mathbf{u},1)^{1/2}+1).$$

## 3. Weiss-type monotonicity formula

In the rest of the paper, we study the free boundary of almost minimizers. This section is devoted to proving the monotonicity of the Weiss-type functional introduced in Theorem 2. It is obtained by comparing almost minimizers with homogeneous functions of degree 2 with some algebraic manipulation. This argument goes back to [11] in the case of the classical obstacle problem, and recently to [9] for almost minimizers of the thin obstacle problem.

**Theorem 5.** Let **u** be an almost minimizer in  $B_1$ . Then, for  $x_0 \in B_{1/2}$  and  $0 < t < t_0(n, \alpha)$ ,

$$\frac{d}{dt}W(\mathbf{u}, x_0, t) \ge \frac{e^{at^{\alpha}}}{t^{n+2}} \int_{\partial B_t(x_0)} \left| \partial_{\nu} \mathbf{u} - \frac{2(1-bt^{\alpha})}{t} \mathbf{u} \right|^2$$

In particular,  $W(\mathbf{u}, x_0, t)$  is nondecreasing in t for  $0 < t < t_0$ .

*Proof.* The proof can be obtained with similar arguments as in [9], and we sketch it below. Assume without loss of generality that  $x_0 = 0$ . Then, for the 2-homogeneous replacement  $\mathbf{c}_r$  of  $\mathbf{u}$  in  $B_r$ , a standard computation (see Theorem 5.1 in [9]) gives that

$$\int_{B_r} |\nabla \mathbf{c}_r|^2 = \frac{r}{n+2} \int_{\partial B_r} \left( |\nabla \mathbf{u}|^2 - |\partial_\nu \mathbf{u}|^2 + \frac{4}{r^2} |\mathbf{u}|^2 \right)$$

Using that **u** is almost minimizing and  $\mathbf{u} = \mathbf{c}_r$  on  $\partial B_r$ , we deduce that,

$$\begin{split} \int_{\partial B_r} |\mathbf{u}|^2 &= \frac{r^2}{4} \left( \frac{n+2}{r} E(\mathbf{c}_r, r) - 2\frac{n+2}{r} \int_{B_r} |\mathbf{c}_r| - \int_{\partial B_r} \left( |\nabla \mathbf{u}|^2 - |\partial_{\nu} \mathbf{u}|^2 \right) \right) \\ &\geq \frac{r^2}{4} \left( \frac{n+2}{r} (1-r^{\alpha}) E(\mathbf{u}, r) - 2\frac{n+2}{r} \int_{B_r} |\mathbf{c}_r| - \int_{\partial B_r} \left( |\nabla \mathbf{u}|^2 - |\partial_{\nu} \mathbf{u}|^2 \right) \right) \\ &\geq \frac{r^2}{4} \left( \frac{n+2}{r} (1-r^{\alpha}) E(\mathbf{u}, r) - 2 \int_{\partial B_r} |\mathbf{u}| - \int_{\partial B_r} \left( |\nabla \mathbf{u}|^2 - |\partial_{\nu} \mathbf{u}|^2 \right) \right), \end{split}$$

where in the last line we used the definition of  $\mathbf{c}_r$ . From this we deduce that, (3.1)

$$E(\mathbf{u},r) \le \frac{r}{(n+2)(1-r^{\alpha})} \left(\frac{4}{r^2} \int_{\partial B_r} |\mathbf{u}|^2 + 2 \int_{\partial B_r} |\mathbf{u}| + \int_{\partial B_r} \left( |\nabla \mathbf{u}|^2 - |\partial_{\nu} \mathbf{u}|^2 \right) \right).$$

 $\overline{7}$ 

We now set

$$\psi(r) := \frac{2e^{ar^{\alpha}}(1-br^{\alpha})}{r^{n+3}},$$

so that

$$W(r) := W(\mathbf{u}, 0, r) = e^{ar^{\alpha}} r^{-n-2} E(\mathbf{u}, r) - \psi(r) \int_{\partial B_r} |\mathbf{u}|^2$$

and

(3.2) 
$$\frac{d}{dr}W(r) = \frac{d}{dr}(e^{ar^{\alpha}}r^{-n-2})E(\mathbf{u},r) + e^{ar^{\alpha}}r^{-n-2}\int_{\partial B_{r}}(|\nabla \mathbf{u}|^{2} + 2|\mathbf{u}|) + \Phi(r),$$
with

$$\Phi(r) := -\frac{d}{dr} \left( \psi(r) \int_{\partial B_r} |\mathbf{u}|^2 \right).$$

On the other hand,

$$\frac{d}{dr}(e^{ar^{\alpha}}r^{-n-2})E(\mathbf{u},r) = -(n+2)e^{ar^{\alpha}}r^{-n-3}(1-r^{\alpha})E(\mathbf{u},r),$$

which in view of (3.1) gives

$$\begin{split} \frac{d}{dr}(e^{ar^{\alpha}}r^{-n-2})E(\mathbf{u},r) &\geq e^{ar^{\alpha}}r^{-n-2}\left(\int_{\partial B_{r}}(|\partial_{\nu}\mathbf{u}|^{2}-|\nabla\mathbf{u}|^{2})-2\int_{\partial B_{r}}|\mathbf{u}|\right)\\ &-4e^{ar^{\alpha}}r^{-n-4}\int_{\partial B_{r}}|\mathbf{u}|^{2}. \end{split}$$

This, combined with (3.2) implies

$$\frac{d}{dr}W(r) \ge e^{ar^{\alpha}}r^{-n-2}\int_{\partial B_r}|\partial_{\nu}\mathbf{u}|^2 - 4e^{ar^{\alpha}}r^{-n-4}\int_{\partial B_r}|\mathbf{u}|^2$$
$$-2\psi(r)\int_{\partial B_r}\mathbf{u}\cdot\partial_{\nu}\mathbf{u} - (4e^{ar^{\alpha}}r^{-n-4} + \psi'(r) + \frac{n-1}{r}\psi(r))\int_{\partial B_r}|\mathbf{u}|^2$$

It is easy to see that, for  $r < r_0 = r_0(n, \alpha)$ 

$$-\frac{e^{ar^{\alpha}}}{r^{n+2}}(4e^{ar^{\alpha}}r^{-n-4} + \psi'(r) + \frac{n-1}{r}\psi(r)) \ge \psi(r)^2$$

which combined with the inequality above and the definition of  $\psi$  implies the desired claim.  $\hfill \Box$ 

## 4. Growth estimates

In this section we show that almost minimizers have quadratic growth away from the free boundary. We begin with a weak quadratic growth estimate.

**Lemma 2.** Let  $\mathbf{u} \in W^{1,2}(B_1)$  be an almost minimizer in  $B_1$ . Then for any  $0 < \varepsilon < 1$ , there exist C > 0 and  $r_0 > 0$ , depending on n,  $\alpha$ ,  $\varepsilon$ ,  $E(\mathbf{u}, 1)$ , such that

$$\sup_{B_r(x)} \left( \frac{|\mathbf{u}|}{r^{2-\varepsilon}} + \frac{|\nabla \mathbf{u}|}{r^{1-\varepsilon}} \right) \le C,$$

for any  $x \in \Gamma(\mathbf{u}) \cap B_{1/2}$  and  $0 < r < r_0$ .

*Proof.* Let  $M \geq E(\mathbf{u}, 1)$ , and note that

$$\|\mathbf{u}\|_{C^{1,\alpha/2}(B_{3/4})} \le C(n,\alpha)(E(\mathbf{u},1)^{1/2}+1) \le C(n,\alpha,M).$$

Assume by contradiction that the lemma is not true for almost minimizers with energy bounded by M. Then we can find a sequence  $\{\mathbf{u}_j\}_{j=1}^{\infty}$  of almost minimizers

with  $E(\mathbf{u}_j, 1) \leq M$ , a sequence  $\{x_j\}_{j=1}^{\infty} \subset \Gamma(\mathbf{u}_j) \cap B_{1/2}$  of free boundary points and a sequence of positive radii  $\{r_j\}_{j=1}^{\infty} \subset (0, 1), r_j \searrow 0$ , such that for large j

$$\sup_{B_{r_j}(x_j)} \left( \frac{|\mathbf{u}_j|}{r_j^{2-\varepsilon}} + \frac{|\nabla \mathbf{u}_j|}{r_j^{1-\varepsilon}} \right) = j, \quad \sup_{B_r(x_j)} \left( \frac{|\mathbf{u}_j|}{r^{2-\varepsilon}} + \frac{|\nabla \mathbf{u}_j|}{r^{1-\varepsilon}} \right) \le j \quad \text{for any } r_j \le r \le 1/4.$$

Define the function

$$\widetilde{\mathbf{u}_j}(x) := \frac{\mathbf{u}_j(r_j x + x_j)}{j r_j^{2-\varepsilon}}, \quad x \in B_{\frac{1}{4r_j}}.$$

Then

$$\sup_{B_1} \left( |\widetilde{\mathbf{u}_j}| + |\nabla \widetilde{\mathbf{u}_j}| \right) = 1 \quad \text{and} \ \sup_{B_R} \left( \frac{|\widetilde{\mathbf{u}_j}|}{R^{2-\varepsilon}} + \frac{|\nabla \widetilde{\mathbf{u}_j}|}{R^{1-\varepsilon}} \right) \le 1 \quad \text{for any } 1 \le R \le \frac{1}{4r_j}.$$

Now we claim that there exists a harmonic function  $\widetilde{\mathbf{u}_0} \in C^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^m)$  such that over a subsequence

$$\widetilde{\mathbf{u}_j} \to \widetilde{\mathbf{u}_0} \quad \text{in } C^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^m).$$

Indeed, for a fixed R > 1 and a ball  $B_{\rho}(z) \subset B_R$ , we have

$$\int_{B_{\rho}(z)} \left( |\nabla \widetilde{\mathbf{u}_{j}}|^{2} + \frac{2r_{j}^{\varepsilon}}{j} |\widetilde{\mathbf{u}_{j}}| \right) = \frac{1}{j^{2}r_{j}^{n+2-2\varepsilon}} \int_{B_{r_{j}\rho}(r_{j}z+x_{j})} \left( |\nabla \mathbf{u}_{j}|^{2} + 2|\mathbf{u}_{j}| \right).$$

This implies that each  $\widetilde{\mathbf{u}_{j}}$  is an almost minimizer of a functional

$$J_{B_{\rho}(z)}(\widetilde{\mathbf{u}_{j}}) = \int_{B_{\rho}(z)} \left( |\nabla \widetilde{\mathbf{u}_{j}}|^{2} + \frac{2r_{j}^{\varepsilon}}{j} |\widetilde{\mathbf{u}_{j}}| \right),$$

with a gauge function  $\omega(\rho) = (r_j \rho)^{\alpha} \leq \rho^{\alpha}$ . Let **h** be the harmonic replacement of  $\widetilde{\mathbf{u}}_j$  in  $B_{\rho}(z)$ . Then, since  $\frac{r_j^{\epsilon}}{j} \leq 1$ , we can obtain the equivalent of (2.3):

$$\begin{split} \int_{B_{\rho}(z)} |\nabla(\widetilde{\mathbf{u}_{j}} - \mathbf{h})| &= \int_{B_{\rho}(z)} \left( |\nabla\widetilde{\mathbf{u}_{j}}|^{2} - |\nabla\mathbf{h}|^{2} \right) \\ &\leq \int_{B_{\rho}(z)} \rho^{\alpha} |\nabla\mathbf{h}|^{2} + \frac{2r_{j}^{\varepsilon}}{j} \left( (1 + \rho^{\alpha}) |\mathbf{h}| - |\widetilde{\mathbf{u}_{j}}| \right) \\ &= \int_{B_{\rho}(z)} \rho^{\alpha} |\nabla\mathbf{h}|^{2} + \frac{2r_{j}^{\varepsilon}}{j} (1 + \rho^{\alpha}) (|\mathbf{h}| - |\widetilde{\mathbf{u}_{j}}|) + \frac{2r_{j}^{\varepsilon}}{j} \rho^{\alpha} |\widetilde{\mathbf{u}_{j}}| \\ &\leq \int_{B_{\rho}(z)} \rho^{\alpha} |\nabla\widetilde{\mathbf{u}_{j}}|^{2} + 2(1 + \rho^{\alpha}) |\mathbf{h} - \widetilde{\mathbf{u}_{j}}| + 2\rho^{\alpha} |\widetilde{\mathbf{u}_{j}}|. \end{split}$$

With this estimate at hand, we can proceed as in the proofs of Proposition 1, Theorem 4 and Theorem 1 to obtain

$$\|\widetilde{\mathbf{u}_j}\|_{C^{1,\alpha/2}(\overline{B_{R/2}};\mathbb{R}^m)} \le C(n,\alpha,R) \left( E(\widetilde{\mathbf{u}_j},R)^{1/2} + 1 \right) \le C(n,\alpha,M,R).$$

Thus, over a subsequence,

$$\widetilde{\mathbf{u}_j} \to \widetilde{\mathbf{u}_0} \quad \text{in } C^1(B_{R/2}; \mathbb{R}^m).$$

Letting  $R \to \infty$  and using Cantor's diagonal argument, we have that

$$\widetilde{\mathbf{u}_j} \to \widetilde{\mathbf{u}_0} \quad \text{in } C^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^m).$$

Now, to verify that  $\widetilde{\mathbf{u}_0}$  is harmonic, we fix R > 1 and observe that for large j and for the harmonic replacement  $\mathbf{h}_j$  of  $\widetilde{\mathbf{u}_j}$  in  $B_R$ ,

(4.1) 
$$\int_{B_R} \left( |\nabla \widetilde{\mathbf{u}_j}|^2 + \frac{2r_j^{\varepsilon}}{j} |\widetilde{\mathbf{u}_j}| \right) \le (1 + (r_j R)^{\alpha}) \int_{B_R} \left( |\nabla \mathbf{h}_j|^2 + \frac{2r_j^{\varepsilon}}{j} |\mathbf{h}_j| \right).$$

From

$$\|\mathbf{h}_j\|_{C^{1,\alpha/2}(\overline{B_R};\mathbb{R}^m)} \le C(n,R) \|\widetilde{\mathbf{u}_j}\|_{C^{1,\alpha/2}(\overline{B_R};\mathbb{R}^m)} \le C(n,\alpha,M,R),$$

we have that up to a subsequence

$$\mathbf{h}_j \to \mathbf{h}_0 \quad \text{in } C^1(\overline{B_R}; \mathbb{R}^m),$$

for some harmonic function  $\mathbf{h}_0 \in C^1(\overline{B_R}; \mathbb{R}^m)$ . Thus, by taking  $j \to \infty$  in (4.1), we obtain

$$\int_{B_R} |\nabla \widetilde{\mathbf{u}_0}|^2 \le \int_{B_R} |\nabla \mathbf{h}_0|^2.$$

Since  $\mathbf{h}_0$  is a harmonic function in  $B_R$  with  $\widetilde{\mathbf{u}_0}$ -boundary value,  $\widetilde{\mathbf{u}_0}$  should be equal to  $\mathbf{h}_0$  in  $B_R$  and hence is harmonic there. Since R > 1 is arbitrary,  $\widetilde{\mathbf{u}_0}$  is harmonic in  $\mathbb{R}^n$ . This finishes the proof of the claim.

Now, we observe that

$$\sup_{B_1}(|\widetilde{\mathbf{u}_0}| + |\nabla\widetilde{\mathbf{u}_0}|) = 1 \quad \text{and} \ \sup_{B_R}\left(\frac{|\widetilde{\mathbf{u}_0}|}{R^{2-\varepsilon}} + \frac{|\nabla\widetilde{\mathbf{u}_0}|}{R^{1-\varepsilon}}\right) \le 1 \quad \text{for any } R \ge 1.$$

Moreover, notice that from  $x_j \in \Gamma(\mathbf{u}_j)$ , we have  $|\widetilde{\mathbf{u}_j}(0)| = |\nabla \widetilde{\mathbf{u}_j}(0)| = 0$ , and thus  $|\widetilde{\mathbf{u}_0}(0)| = |\nabla \widetilde{\mathbf{u}_0}(0)| = 0$ . To arrive at a contradiction, we note that  $|\nabla \widetilde{\mathbf{u}_0}(x)| \leq |x|^{1-\varepsilon}$  for  $|x| \geq 1$  and  $|\nabla \widetilde{\mathbf{u}_0}(0)| = 0$  and apply Liouville's theorem to obtain  $\nabla \widetilde{\mathbf{u}_0} \equiv 0$  in  $\mathbb{R}^n$ . This means that  $\widetilde{\mathbf{u}_0}$  is a constant vector, which contradicts that  $\sup_{B_1} |\widetilde{\mathbf{u}_0}| = \sup_{B_1}(|\widetilde{\mathbf{u}_0}| + |\nabla \widetilde{\mathbf{u}_0}|) = 1$  and  $|\widetilde{\mathbf{u}_0}(0)| = 0$ .

Now we are going to establish the quadratic growth of almost minimizers at free boundary points (Lemma 3 and Lemma 4) with the help of the weak quadratic growth estimate (Lemma 2) and the monotonicity of the Weiss-type energy (Theorem 5).

**Lemma 3.** Let **u** be an almost minimizer in  $B_1$  with  $x_0 \in \Gamma(\mathbf{u}) \cap B_{1/2}$ . Then, for C > 0 and  $r_0 > 0$ , depending only on n,  $\alpha$ ,  $E(\mathbf{u}, 1)$ , we have

$$\int_{B_r(x_0)} |\mathbf{u}| \le C r^{n+2}, \quad 0 < r < r_0.$$

*Proof.* From the monotonicity formula,

$$W(\mathbf{u}, x_0, r) \le W(\mathbf{u}, x_0, 1/2) \le C(n, \alpha) E(u, 1).$$

Thus

$$\frac{2e^{ar^{\alpha}}}{r^{n+2}} \int_{B_{r}(x_{0})} |\mathbf{u}| \\
= W(\mathbf{u}, x_{0}, r) + \frac{2e^{ar^{\alpha}}(1 - br^{\alpha})}{r^{n+3}} \int_{\partial B_{r}(x_{0})} |\mathbf{u}|^{2} - \frac{e^{ar^{\alpha}}}{r^{n+2}} \int_{B_{r}(x_{0})} |\nabla \mathbf{u}|^{2} \\
(4.2) \leq C(n, \alpha)E(\mathbf{u}, 1) + e^{ar^{\alpha}} \left(\frac{2}{r^{n+3}} \int_{\partial B_{r}(x_{0})} |\mathbf{u}|^{2} - \frac{1}{r^{n+2}} \int_{B_{r}(x_{0})} |\nabla \mathbf{u}|^{2}\right) \\
= C(n, \alpha)E(\mathbf{u}, 1) \\
+ e^{ar^{\alpha}} \left(\frac{2}{r^{n+3}} \int_{\partial B_{r}(x_{0})} |\mathbf{u} - S_{x_{0}}\mathbf{p}|^{2} - \frac{1}{r^{n+2}} \int_{B_{r}(x_{0})} |\nabla(\mathbf{u} - S_{x_{0}}\mathbf{p})|^{2}\right) \\
\leq C(n, \alpha)E(\mathbf{u}, 1) + \frac{2e^{ar^{\alpha}}}{r^{n+3}} \int_{\partial B_{r}(x_{0})} |\mathbf{u} - S_{x_{0}}\mathbf{p}|^{2}$$

for each  $\mathbf{p} = (p_1, \ldots, p_m) \in \mathcal{H}$ , where  $\mathcal{H}$  is the set of all  $\mathbf{p} = (p_1, \ldots, p_m)$  such that each component  $p_j$  is 2-homogeneous harmonic polynomial, and  $S_{x_0}f(x) := f(x - x_0)$ .

For each  $x_0 \in \Gamma(\mathbf{u})$ , let  $\mathbf{p}_{x_0,r} = \mathbf{p}_{x_0,r,\mathbf{u}}$  be the minimizer of  $\int_{\partial B_r(x_0)} |\mathbf{u} - S_{x_0}\mathbf{p}|^2$  in  $\mathcal{H}$ . Then it is easy to see that

$$\int_{\partial B_r(x_0)} (\mathbf{u} - S_{x_0} \mathbf{p}_{x_0,r}) S_{x_0} \mathbf{q} = 0 \quad \text{for every } \mathbf{q} \in \mathcal{H}.$$

Now, the lemma follows once we prove that there are  $C = C(n, \alpha, E(\mathbf{u}, 1))$  and  $r_0 = r_0(n, \alpha, E(\mathbf{u}, 1)) > 0$  such that

(4.3) 
$$\int_{\partial B_r(x_0)} |\mathbf{u} - S_{x_0} \mathbf{p}_{x_0, r}|^2 \le C r^{n+3}, \quad 0 < r < r_0.$$

Towards this, we assume to the contrary that there are a sequence of almost minimizers  $\mathbf{u}_k$ , a sequence of points  $x_k \in \Gamma(\mathbf{u}_k) \cap B_{1/2}$ , and  $r_k \to 0$  such that  $E(\mathbf{u}_k, 1)$ is uniformly bounded and

$$M_k := \frac{1}{r_k^{n+3}} \int_{\partial B_{r_k}(x_k)} |\mathbf{u}_k - S_{x_k} \mathbf{p}_{x_k, r_k}|^2 \to \infty.$$

Denote

$$\mathbf{v}_k(x) := (\mathbf{u}_k)_{x_k, r_k}(x) = \frac{\mathbf{u}_k(x_k + r_k x)}{r_k^2} \quad \text{and } \mathbf{w}_k(x) := \frac{\mathbf{v}_k(x) - \mathbf{p}_{x_k, r_k}(x)}{M_k^{1/2}}, \quad x \in B_1$$

For each k, let  $\mathbf{h}_k$  be the harmonic function in  $B_1$  such that  $\mathbf{h}_k = \mathbf{v}_k$  on  $\partial B_1$ , and write

$$\mathbf{w}_k = \frac{\mathbf{h}_k - \mathbf{p}_{x_k, r_k}}{M_k^{1/2}} + \frac{\mathbf{v}_k - \mathbf{h}_k}{M_k^{1/2}} := \mathbf{w}_k^1 + \mathbf{w}_k^2.$$

Then

$$\int_{\partial B_1} |\mathbf{w}_k^1|^2 = \frac{1}{M_k} \int_{\partial B_1} |\mathbf{v}_k - \mathbf{p}_{x_k, r_k}|^2 = \frac{1}{M_k r_k^{n+3}} \int_{\partial B_{r_k}(x_k)} |\mathbf{u}_k - S_{x_k} \mathbf{p}_{x_k, r_k}|^2 = 1$$

and

$$\begin{split} \int_{B_1} |\nabla \mathbf{w}_k^1|^2 &- 2 \int_{\partial B_1} |\mathbf{w}_k^1|^2 = \frac{1}{M_k} \left( \int_{B_1} |\nabla (\mathbf{h}_k - \mathbf{p}_{x_k, r_k})|^2 - 2 \int_{\partial B_1} |\mathbf{h}_k - \mathbf{p}_{x_k, r_k}|^2 \right) \\ &= \frac{1}{M_k} \left( \int_{B_1} |\nabla \mathbf{h}_k|^2 - 2 \int_{\partial B_1} |\mathbf{h}_k|^2 \right) \\ &\leq \frac{1}{M_k} \left( \int_{B_1} |\nabla \mathbf{v}_k|^2 - 2 \int_{\partial B_1} |\mathbf{v}_k|^2 \right) \\ &= \frac{1}{M_k} \left( \frac{1}{r_k^{n+2}} \int_{B_{r_k}(x_k)} |\nabla \mathbf{u}_k|^2 - \frac{2}{r_k^{n+3}} \int_{\partial B_{r_k}(x_k)} |\mathbf{u}_k|^2 \right) \\ &\leq \frac{1}{M_k} W(\mathbf{u}_k, x_k, r_k) \leq \frac{1}{M_k} W(\mathbf{u}_k, x_k, 1/2) \\ &\leq \frac{C(n, \alpha)}{M_k} E(\mathbf{u}_k, 1) \to 0 \quad \text{as } k \to \infty. \end{split}$$

Thus  $\{\mathbf{w}_k^1\}_{k\in\mathbb{N}}$  is bounded in  $W^{1,2}(B_1;\mathbb{R}^m)$ , hence there is  $\mathbf{w}_0^1 \in W^{1,2}(B_1;\mathbb{R}^m)$  such that up to a subsequence

$$\begin{aligned} \mathbf{w}_k^1 &\to \mathbf{w}_0^1 \quad \text{weakly in } W^{1,2}(B_1; \mathbb{R}^m), \\ \mathbf{w}_k^1 &\to \mathbf{w}_0^1 \quad \text{strongly in } L^2(\partial B_1; \mathbb{R}^m). \end{aligned}$$

It follows that

(4.4) 
$$\int_{B_1} |\nabla \mathbf{w}_0^1|^2 \le 2 \int_{\partial B_1} |\mathbf{w}_0^1|^2 = 2,$$

(4.5) 
$$\int_{\partial B_1} \mathbf{w}_0^1 \cdot \mathbf{q} = 0 \quad \text{for every } \mathbf{q} \in \mathcal{H}$$

Moreover, since each  $\mathbf{w}_k^1$  is harmonic in  $B_1$ ,  $\mathbf{w}_0^1$  is also harmonic in  $B_1$ . It also follows from  $C^{1,\alpha}$ -estimate of harmonic functions  $\mathbf{w}_k^1$  that

 $\mathbf{w}_k^1 \to \mathbf{w}_0^1$  in  $C_{\mathrm{loc}}^1(B_1; \mathbb{R}^m)$ .

Next, to deal with  $\mathbf{w}_k^2 = \frac{\mathbf{v}_k - \mathbf{h}_k}{M_k^{1/2}}$ , we observe that  $\mathbf{v}_k(x) = \frac{\mathbf{u}_k(x_0 + r_k x)}{r_k^2}$  is an almost minimizer with a gauge function  $\omega(\rho) = (\rho r_k)^{\alpha}$ . Thus, by (2.4) and Lemma 2 with  $\varepsilon = \alpha/4$ ,

$$\begin{split} \int_{B_1} \left( |\nabla \mathbf{w}_k^2|^2 + |\mathbf{w}_k^2| \right) &\leq \frac{1}{M_k^{1/2}} \int_{B_1} \left( |\nabla (\mathbf{v}_k - \mathbf{h}_k)|^2 + |\mathbf{v}_k - \mathbf{h}_k| \right) \\ &\leq \frac{C(n)}{M_k^{1/2}} \left( r_k^\alpha \int_{B_1} \left( |\nabla \mathbf{v}_k|^2 + |\mathbf{v}_k| \right) + 1 \right) \\ &\leq \frac{C(n)}{M_k^{1/2}} \left( C(n, \alpha, E(\mathbf{u}_k, 1)) r_k^{\alpha/2} + 1 \right) \to 0 \quad \text{as } k \to \infty. \end{split}$$

Thus, up to another subsequence,

$$\mathbf{w}_k^2 \to 0$$
 strongly in  $W^{1,2}(B_1; \mathbb{R}^m)$ .

From  $x_k \in \Gamma(\mathbf{u}_k)$ , we have  $|\mathbf{w}_k(0)| = |\nabla \mathbf{w}_k(0)| = 0$ , thus

$$|\mathbf{w}_{0}^{1}(0)| = \lim_{k \to \infty} |\mathbf{w}_{k}^{1}(0)| = \lim_{k \to \infty} |\mathbf{w}_{k}(0)| = 0$$

and

$$|\nabla \mathbf{w}_0^1(0)| = \lim_{k \to \infty} |\nabla \mathbf{w}_k^1(0)| = \lim_{k \to \infty} |\nabla \mathbf{w}_k(0)| = 0.$$

From here, we can repeat the argument in Theorem 2 in [1] to get the contradiction. Indeed, by Lemma 4.1 in [12], each component  $z_j$  of  $\mathbf{w}_0^1$  satisfies

$$2\int_{\partial B_1} z_j^2 \le \int_{B_1} |\nabla z_j|^2.$$

This, together with (4.4), gives

$$2\int_{\partial B_1} z_j^2 = \int_{B_1} |\nabla z_j|^2.$$

By Lemma 4.1 in [12] again, each  $z_j$  is a 2-homogeneous harmonic polynomial. Therefore, we have  $\mathbf{w}_0^1 \in \mathcal{H}$ , and this contradicts (4.5).

**Lemma 4.** Let **u** be an almost minimizer in  $B_1$  with  $x_0 \in \Gamma(\mathbf{u}) \cap B_{1/2}$ . Then, for C and  $r_0$  as in Lemma 3, we have for  $0 < r < r_0$ 

$$\int_{B_r(x_0)} |\nabla \mathbf{u}|^2 \le Cr^{n+2} \qquad and \qquad \int_{\partial B_r(x_0)} |\mathbf{u}|^2 \le Cr^{n+3}.$$

*Proof.* For simplicity we assume  $x_0 = 0$ . For  $0 < r < r_0$ , let **h** be a harmonic replacement of **u** in  $B_r$ . For 2-homogeneous recalings  $\mathbf{u}_r$  and  $\mathbf{h}_r$  of **u** and **h**, respectively, by (2.4) and Lemma 2 with  $\varepsilon = \alpha/4$ , we get

$$\begin{split} \int_{B_1} \left( |\nabla (\mathbf{u}_r - \mathbf{h}_r)|^2 + |\mathbf{u}_r - \mathbf{h}_r| \right) &\leq C(n) r^{\alpha} \int_{B_1} \left( |\nabla \mathbf{u}_r|^2 + |\mathbf{u}_r| \right) + C(n) \\ &\leq C(n, \alpha, E(\mathbf{u}, 1)). \end{split}$$

This, combined with Lemma 3, gives

$$\int_{B_1} |\mathbf{h}_r| \le \int_{B_1} |\mathbf{u}_r - \mathbf{h}_r| + \int_{B_1} |\mathbf{u}_r| \le C(n, \alpha, E(\mathbf{u}, 1)).$$

Since  $\mathbf{h}_r$  is harmonic, we also have

$$\int_{B_{1/2}} |\nabla \mathbf{h}_r|^2 \le C(n) \left( \int_{B_1} |\mathbf{h}_r| \right)^2 \le C(n, \alpha, E(\mathbf{u}, 1)).$$

Therefore,

$$\int_{B_{1/2}} |\nabla \mathbf{u}_r|^2 \le 2 \int_{B_{1/2}} |\nabla \mathbf{h}_r|^2 + 2 \int_{B_{1/2}} |\nabla (\mathbf{u}_r - \mathbf{h}_r)|^2 \le C(n, \alpha, E(\mathbf{u}, 1)).$$

This gives the first estimate. For the second one, we observe that

$$\begin{aligned} \frac{2}{r^{n+3}} \int_{\partial B_r} |\mathbf{u}|^2 &- \frac{1}{r^{n+2}} \int_{B_r} |\nabla \mathbf{u}|^2 \\ &= \frac{2}{r^{n+3}} \int_{\partial B_r} |\mathbf{u} - \mathbf{p}_{0,r}|^2 - \frac{1}{r^{n+2}} \int_{B_r} |\nabla (\mathbf{u} - \mathbf{p}_{0,r})|^2 \\ &\leq \frac{2}{r^{n+3}} \int_{\partial B_r} |\mathbf{u} - \mathbf{p}_{0,r}|^2 \leq C(n, \alpha, E(\mathbf{u}, 1)), \end{aligned}$$

where the last inequality follows from (4.3). This, combined with the first estimate, implies the second one.  $\hfill \Box$ 

Finally, combining Lemma 3, Lemma 4 and Theorem 1 we obtain the following.

**Corollary 1.** Let **u** be an almost minimizer in  $B_1$ . Then, for  $x_0 \in \Gamma(\mathbf{u}) \cap B_{1/2}$ ,  $0 < r < r_0, r_0 = r_0(n, \alpha, E(\mathbf{u}, 1)), \mathbf{u}_{x_0, r} \in C^{1, \alpha/2}(B_1)$  and

(4.6) 
$$\|\mathbf{u}_{x_0,r}\|_{C^{1,\alpha/2}(B_1)} \le C(n,\alpha, E(\mathbf{u},1))$$

**Remark 1.** We remark that in Lemma 3, Lemma 4, and in Corollary 1, it is possible to obtain a constant C independent of  $E(\mathbf{u}, 1)$ . This can be achieved by a standard compactness argument.

#### 5. Non-degeneracy

In this section we prove that almost minimizers satisfy a non-degeneracy property. Precisely, we have the following theorem.

**Theorem 6** (Non-Degeneracy). Let **u** be an almost minimizer in  $B_1$ . There exist constants  $c_0 = c_0(n, m, \alpha, E(\mathbf{u}, 1)) > 0$  and  $r_0 = r_0(n, m, \alpha) > 0$  such that if  $x_0 \in \Gamma(\mathbf{u}) \cap B_{1/2}$  and  $0 < r < r_0$ , then

$$\sup_{\mathsf{B}_r(x_0)} |\mathbf{u}| \ge c_0 r^2.$$

The proof of Theorem 6 relies on the following lemma.

**Lemma 5.** Let **u** be an almost minimizer in  $B_1$ . Then, there exist small constants  $\varepsilon_0 = \varepsilon_0(n,m) > 0$  and  $r_0 = r_0(n,m,\alpha) > 0$  such that for  $0 < r < r_0$ , if  $E(\mathbf{u}_{x_0,r}, 1) \leq \varepsilon_0$  then  $E(\mathbf{u}_{x_0,r/2}, 1) \leq \varepsilon_0$ .

*Proof.* For simplicity we may assume  $x_0 = 0$ . For  $0 < r < r_0$  to be specified later, let  $\mathbf{v}_r$  be a solution of  $\Delta \mathbf{v}_r = \frac{\mathbf{v}_r}{|\mathbf{v}_r|} \chi_{\{|\mathbf{v}_r|>0\}}$  in  $B_1$  with  $\mathbf{v}_r = \mathbf{u}_r$  on  $\partial B_1$ . We claim that if  $\varepsilon_0 = \varepsilon_0(n,m) > 0$  is small, then  $\mathbf{v}_r \equiv \mathbf{0}$  in  $B_{1/2}$ . Indeed, if not, then  $\sup_{B_{3/4}} |\mathbf{v}_r| \ge c_0(n)$  by the non-degeneracy of the solution  $\mathbf{v}_r$ . Thus

 $|\mathbf{v}_r(z_0)| \ge c_0(n)$  for some  $z_0 \in \overline{B_{3/4}}$ . From  $E(\mathbf{v}_r, 1) \le E(\mathbf{u}_r, 1) \le \varepsilon_0$  together with the estimate for the solution  $\mathbf{v}_r$ ,

$$\sup_{B_{7/8}} |\nabla \mathbf{v}_r| \le C_1(n,m) (E(\mathbf{v}_r,1)+1)$$

we have that

$$\sup_{B_{7/8}} |\nabla \mathbf{v}_r| \le C(n,m),$$

hence

$$|\mathbf{v}_r| \ge \frac{c_0(n)}{2}$$
 in  $B_{\rho_0}(z_0)$ 

for some small  $\rho_0 = \rho_0(n, m) > 0$ . This gives that

$$c(n,m) \leq \int_{B_{\rho_0}(z_0)} |\mathbf{v}_r| \leq E(\mathbf{v}_r,1) \leq \varepsilon_0,$$

which is a contradiction if  $\varepsilon_0 = \varepsilon_0(n, m)$  is small.

Now, we use again that  $E(\mathbf{v}_r, 1) \leq \varepsilon_0$  together with the fact that  $\mathbf{u}_r$  is an almost minimizer in  $B_1$  with gauge function  $\omega(\rho) = (r\rho)^{\alpha}$  to get

$$\int_{B_1} \left( |\nabla \mathbf{u}_r|^2 + 2|\mathbf{u}_r| \right) \le (1 + r^{\alpha}) \int_{B_1} \left( |\nabla \mathbf{v}_r|^2 + 2|\mathbf{v}_r| \right)$$
$$\le \varepsilon_0 r^{\alpha} + \int_{B_1} \left( |\nabla \mathbf{v}_r|^2 + 2|\mathbf{v}_r| \right)$$

for  $0 < r < r_0$ . Then

$$\begin{split} \int_{B_1} |\nabla(\mathbf{u}_r - \mathbf{v}_r)|^2 &= \int_{B_1} \left( |\nabla \mathbf{u}_r|^2 - |\nabla \mathbf{v}_r|^2 + 2\nabla(\mathbf{v}_r - \mathbf{u}_r)\nabla \mathbf{v}_r \right) \\ &= \int_{B_1} \left( |\nabla \mathbf{u}_r|^2 - |\nabla \mathbf{v}_r|^2 \right) - 2 \int_{B_1} \left( (\mathbf{v}_r - \mathbf{u}_r) \frac{\mathbf{v}_r}{|\mathbf{v}_r|} \chi_{\{|\mathbf{v}_r| > 0\}} \right) \\ &\leq \varepsilon_0 r^\alpha + 2 \int_{B_1} (|\mathbf{v}_r| - |\mathbf{u}_r|) - 2 \int_{B_1} \left( |\mathbf{v}_r| - \mathbf{u}_r \frac{\mathbf{v}_r}{|\mathbf{v}_r|} \chi_{\{|\mathbf{v}_r| > 0\}} \right) \\ &\leq \varepsilon_0 r^\alpha. \end{split}$$

Combining this with Poincaré's inequality and Hölder's inequality, we obtain

$$\int_{B_1} \left( |\nabla (\mathbf{u}_r - \mathbf{v}_r)|^2 + 2|\mathbf{u}_r - \mathbf{v}_r| \right) \le C(n) r^{\alpha/2}$$

Since  $\mathbf{v}_r \equiv \mathbf{0}$  in  $B_{1/2}$ , we see that for  $0 < r < r_0(n, m, \alpha)$ ,

$$E(\mathbf{u}_{r}, 1/2) = \int_{B_{1/2}} \left( |\nabla \mathbf{u}_{r}|^{2} + 2|\mathbf{u}_{r}| \right) \le C(n) r^{\alpha/2} \le \frac{\varepsilon_{0}}{2^{n+2}}$$

Therefore, we conclude that

$$E(\mathbf{u}_{r/2}, 1) = 2^{n+2} E(\mathbf{u}_r, 1/2) \le \varepsilon_0.$$

Lemma 5 immediately implies the following integral form of non-degeneracy.

**Lemma 6.** Let  $\mathbf{u}$ ,  $\varepsilon_0$  and  $r_0$  be as in the preceeding lemma. If  $x_0 \in \overline{\{|\mathbf{u}| > 0\}} \cap B_{1/2}$ and  $0 < r < r_0$ , then

(5.1) 
$$\int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \ge \varepsilon_0 r^{n+2}.$$

*Proof.* By the continuity of  $\mathbf{u}$ , it is enough to prove (5.1) for  $x_0 \in \{|\mathbf{u}| > 0\} \cap B_{1/2}$ . Towards a contradiction, we suppose that  $\int_{B_r(x_0)} (|\nabla \mathbf{u}|^2 + 2|\mathbf{u}|) \leq \varepsilon_0 r^{n+2}$ , or equivalently  $E(\mathbf{u}_{x_0,r}, 1) \leq \varepsilon_0$ . Then, by the previous lemma we have  $E(\mathbf{u}_{x_0,r/2^k}, 1) \leq \varepsilon_0$  for all  $k \in \mathbb{N}$ . From  $|\mathbf{u}(x_0)| > 0$ , we see that  $|\mathbf{u}| > c_0 > 0$  in  $B_{r/2^k}(x_0)$  for large k. Therefore,

$$\varepsilon_0 \ge E(\mathbf{u}_{x_0, r/2^k}, 1) = \frac{1}{(r/2^k)^{n+2}} \int_{B_{r/2^k}(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right)$$
$$\ge \frac{1}{(r/2^k)^{n+2}} \int_{B_{r/2^k}(x_0)} 2c_0 = \frac{C(n)c_0}{(r/2^k)^2} \to \infty \quad \text{as } k \to \infty.$$

This is a contradiction, as desired.

We are now ready to prove Theorem 6.

Proof of Theorem 6. Assume by contradiction that

$$u_{x_0,r}(x) < c_0, \text{ in } B_1,$$

with  $c_0$  small, to be made precise later. Let  $\epsilon_0, r_0$  be the constants in Lemma 5, and let  $r < r_0$ . Then, by interpolation together with estimate (4.6),

$$\begin{aligned} \|\nabla \mathbf{u}_{x_0,r}\|_{L^{\infty}(B_{1/2})} &\leq \epsilon \|\mathbf{u}_{x_0,r}\|_{C^{1,\alpha/2}(B_{3/4})} + K(\epsilon)\|\mathbf{u}_{x_0,r}\|_{L^{\infty}(B_{3/4})} \\ &\leq \epsilon C(n,\alpha, E(\mathbf{u},1)) + K(\epsilon)c_0 \leq \frac{\epsilon_0}{2^{n+3}}, \end{aligned}$$

by choosing  $\epsilon = \frac{\epsilon_0}{2^{n+4}C}$  and  $c_0 \leq \epsilon_0/(2^{n+4}K(\epsilon))$ . Thus, if  $c_0 < \epsilon_0/2^{n+3}$ , then  $E(\mathbf{u}_{x_0,r}, \frac{1}{2}) < \frac{\epsilon_0}{2^{n+2}}$ , which contradicts Lemma 6.

## 6. 2-Homogeneous blowups

In this section we consider 2-homogeneous rescalings and blowups of almost minimizers and complete the proof of Theorem 2. We then obtain the polynomial decay estimate of Weiss-type energies, with the help of the epiperimetric inequality applied to solutions of (1.2).

**Lemma 7.** Suppose  $\mathbf{u}$  is an almost minimizer in  $B_1$  and  $x_0 \in \Gamma(\mathbf{u}) \cap B_{1/2}$ . Then, for 2-homogeneous rescalings  $\mathbf{u}_{x_0,t}$ , there exists  $\mathbf{u}_{x_0,0} \in C^1_{\text{loc}}(\mathbb{R}^n;\mathbb{R}^m)$  such that over a subsequence  $t = t_j \to 0+$ ,

$$\mathbf{u}_{x_0,t_i} \to \mathbf{u}_{x_0,0} \quad in \ C^1_{\mathrm{loc}}(\mathbb{R}^n;\mathbb{R}^m).$$

Moreover,  $\mathbf{u}_{x_0,0}$  is a nonzero 2-homogeneous global solution.

*Proof.* For simplicity we assume  $x_0 = 0$ .

Step 1. We first prove the  $C^1$ -convergence. From Corollary 1, there is  $\mathbf{u}_0 \in C^1(B_{R/2}; \mathbb{R}^m)$  such that over a subsequence  $t = t_j \to 0+$ ,

$$\mathbf{u}_{t_i} \to \mathbf{u}_0$$
 in  $C^1(B_{R/2}; \mathbb{R}^m)$ .

By letting  $R \to \infty$  and using a Cantor's diagonal argument, we obtain that for another subsequence  $t = t_j \to 0+$ ,

$$\mathbf{u}_{t_i} \to \mathbf{u}_0$$
 in  $C^1_{\mathrm{loc}}(\mathbb{R}^n;\mathbb{R}^m)$ 

Step 2. It follows from the non-degeneracy,  $\sup_{B_1} |\mathbf{u}_{t_j}| \geq c_0 > 0$ , and the  $C^1$ convergence of  $\mathbf{u}_{t_j}$  to  $\mathbf{u}_0$  that  $\mathbf{u}_0$  is nonzero. To show that  $\mathbf{u}_0$  is a global solution,
for fixed R > 1 and small  $t_j$ , let  $\mathbf{v}_{t_j}$  be the solution in  $B_R$  with  $\mathbf{v}_{t_j} = \mathbf{u}_{t_j}$  on  $\partial B_R$ .
Then, by elliptic theory,

$$\|\mathbf{v}_{t_j}\|_{C^{1,\alpha/2}(\overline{B_R};\mathbb{R}^m)} \le C(n,m,R)(\|\mathbf{u}_{t_j}\|_{C^{1,\alpha/2}(\overline{B_R};\mathbb{R}^m)} + 1) \le C(n,m,\alpha,R,E(\mathbf{u},1)).$$

Thus, there exists a solution  $\mathbf{v}_0 \in C^1(\overline{B_R}; \mathbb{R}^m)$  such that

$$\mathbf{v}_{t_j} \to \mathbf{v}_0$$
 in  $C^1(\overline{B_R}; \mathbb{R}^m)$ .

Moreover, we use again that  $\mathbf{u}_{t_j}$  is an almost minimizer in  $B_{1/2t_j}$  with a gauge function  $\omega(\rho) = (t_j \rho)^{\alpha}$  to have

$$\int_{B_R} \left( |\nabla \mathbf{u}_{t_j}|^2 + 2|\mathbf{u}_{t_j}| \right) \le (1 + (t_j R)^{\alpha}) \int_{B_R} \left( |\nabla \mathbf{v}_{t_j}|^2 + 2|\mathbf{v}_{t_j}| \right).$$

By taking  $t_j \to 0$  and using the C<sup>1</sup>-convergence of  $\mathbf{u}_{t_i}$  and  $\mathbf{v}_{t_i}$ , we obtain

$$\int_{B_R} \left( |\nabla \mathbf{u}_0|^2 + 2|\mathbf{u}_0| \right) \le \int_{B_R} \left( |\nabla \mathbf{v}_0|^2 + 2|\mathbf{v}_0| \right)$$

Since  $\mathbf{v}_{t_j} = \mathbf{u}_{t_j}$  on  $\partial B_R$ , we also have  $\mathbf{v}_0 = \mathbf{u}_0$  on  $\partial B_R$ . This means that  $\mathbf{u}_0 = \mathbf{v}_0$  is an energy minimizer, or a solution in  $B_R$ . Since R > 1 is arbitrary, we conclude that  $\mathbf{u}_0$  is a global solution.

Step 3. Now, we prove that  $\mathbf{u}_0$  is 2-homogeneous. Fix  $0 < r < R < \infty$ . By Theorem 5, we have that for small  $t_i$ ,

$$W(\mathbf{u}, Rt_{j}) - W(\mathbf{u}, rt_{j})$$

$$= \int_{rt_{j}}^{Rt_{j}} \frac{d}{d\rho} W(\mathbf{u}, \rho) d\rho$$

$$\geq \int_{rt_{j}}^{Rt_{j}} \frac{1}{\rho^{n+4}} \int_{\partial B_{\rho}} |x \cdot \nabla \mathbf{u} - 2(1 - b\rho^{\alpha})\mathbf{u}|^{2} dS_{x} d\rho$$

$$(6.1)$$

$$= \int_{r}^{R} \frac{t_{j}}{(t_{j}\sigma)^{n+4}} \int_{\partial B_{t_{j}\sigma}} |x \cdot \nabla \mathbf{u} - 2(1 - b(t_{j}\sigma)^{\alpha})\mathbf{u}|^{2} dS_{x} d\sigma$$

$$= \int_{r}^{R} \frac{1}{t_{j}^{4}\sigma^{n+4}} \int_{\partial B_{\sigma}} |t_{j}x \cdot \nabla \mathbf{u}(t_{j}x) - 2(1 - b(t_{j}\sigma)^{\alpha})\mathbf{u}(t_{j}x)|^{2} dS_{x} d\sigma$$

$$= \int_{r}^{R} \frac{1}{\sigma^{n+4}} \int_{\partial B_{\sigma}} |x \cdot \nabla \mathbf{u}_{t_{j}} - 2(1 - b(t_{j}\sigma)^{\alpha})\mathbf{u}_{t_{j}}|^{2} dS_{x} d\sigma.$$

Note that in (4.2) we showed

$$\frac{2e^{ar^{\alpha}}}{r^{n+2}}\int_{B_r}|\mathbf{u}| \le W(\mathbf{u},r) + \frac{2e^{ar^{\alpha}}}{r^{n+3}}\int_{\partial B_r}|\mathbf{u}-\mathbf{p}_{0,r}|^2.$$

Since we also proved in the proof of Lemma 3 that

$$\frac{2e^{ar^{\alpha}}}{r^{n+3}}\int_{\partial B_r}|\mathbf{u}-\mathbf{p}_{0,r}|^2 \le C_0,$$

we see that  $W(\mathbf{u}, 0+) \geq -C_0$  and thus  $W(\mathbf{u}, 0+) \in (-\infty, \infty)$ . Using this and taking  $t_j \to 0+$  in (6.1), we get

$$0 = W(\mathbf{u}, 0+) - W(\mathbf{u}, 0+) \ge \int_{r}^{R} \frac{1}{\sigma^{n+3}} \int_{\partial B_{\sigma}} |x \cdot \nabla \mathbf{u}_{0} - 2\mathbf{u}_{0}|^{2} dS_{x} d\sigma.$$

Taking  $r \to 0+$  and  $R \to \infty$ , we conclude that  $x \cdot \nabla \mathbf{u}_0 - 2\mathbf{u}_0 = 0$  in  $\mathbb{R}^n$ , which implies that  $\mathbf{u}_0$  is 2-homogeneous in  $\mathbb{R}^n$ .

Proof of Theorem 2. The monotonicity of the Weiss energy  $W(\mathbf{u}, x_0, \cdot)$  follows from Theorem 5. To prove (1.3), we observe that for  $\mathbf{u}_{x_0,t} = \frac{\mathbf{u}(tx+x_0)}{t^2}$ ,

$$W(\mathbf{u}, x_0, t) = e^{at^{\alpha}} \left( \frac{1}{t^{n+2}} \int_{B_t(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) - \frac{2(1 - bt^{\alpha})}{t^{n+3}} \int_{\partial B_t(x_0)} |\mathbf{u}|^2 \right)$$
$$= e^{at^{\alpha}} \left( \int_{B_1} \left( |\nabla \mathbf{u}_{x_0, t}|^2 + 2|\mathbf{u}_{x_0, t}| \right) - 2(1 - bt^{\alpha}) \int_{\partial B_1} |\mathbf{u}_{x_0, t}|^2 \right).$$

By Lemma 7, we can take a subsequence  $t = t_j \to 0+$  such that  $\mathbf{u}_{x_0,t_j} \to \mathbf{u}_{x_0,0}$  in  $C^1_{\text{loc}}(\mathbb{R}^n;\mathbb{R}^m)$  for some nonzero 2-homogeneous solution  $\mathbf{u}_{x_0,0}$ , which gives

$$W(\mathbf{u}, x_0, 0+) = M(\mathbf{u}_{x_0, 0}).$$

Therefore, by Corollary 1 in [1],

$$W(\mathbf{u}, x_0, 0+) = \beta_n/2 \text{ or } W(\mathbf{u}, x_0, 0+) \ge \bar{\beta}_n.$$

The following is the epiperimetric inequality from [1].

**Theorem 7.** There exists  $\kappa \in (0,1)$  and  $\delta > 0$  such that if **c** is a homogeneous function of degree 2 satisfying  $\|\mathbf{c} - \mathbf{h}\|_{W^{1,2}(B_1;\mathbb{R}^m)} + \|\mathbf{c} - \mathbf{h}\|_{L^{\infty}(B_1;\mathbb{R}^m)} \leq \delta$  for some  $\mathbf{h} \in \mathbb{H}$ , then there is  $\mathbf{v} \in W^{1,2}(B_1;\mathbb{R}^m)$  such that  $\mathbf{v} = \mathbf{c}$  on  $\partial B_1$  and

$$M(\mathbf{v}) \le (1-\kappa)M(\mathbf{c}) + \kappa \frac{\beta_n}{2}.$$

**Lemma 8.** Let **u** be an almost minimizer in  $B_1$  and  $x_0 \in \mathcal{R}_{\mathbf{u}} \cap B_{1/2}$ . Suppose that the epiperimetric inequality holds with  $\kappa \in (0,1)$  for every homogeous replacement  $\mathbf{c}_{x_0,r}$ . Then

$$W(\mathbf{u}, x_0, r) - W(\mathbf{u}, x_0, 0+) \le C(n, \alpha, E(\mathbf{u}, 1))r^{\delta}, \quad 0 < r < r_0 = r_0(n, \alpha)$$

for some  $\delta = \delta(n, \alpha, \kappa) > 0$ .

*Proof.* For simplicity, we assume  $x_0 = 0$ , and write  $W(r) = W(\mathbf{u}, 0, r)$  and  $W(0+) = W(\mathbf{u}, 0, 0+)$ . We define

$$e(r) := W(r) - W(0+)$$
  
=  $\frac{e^{ar^{\alpha}}}{r^{n+2}} \int_{B_r} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) - \frac{2(1 - br^{\alpha})e^{ar^{\alpha}}}{r^{n+3}} \int_{\partial B_r} |\mathbf{u}|^2 - W(0+).$ 

Then, from the identities

$$\frac{d}{dr}\left(\frac{e^{ar^{\alpha}}}{r^{n+2}}\right) = -\frac{(n+2)(1-r^{\alpha})}{r} \cdot \frac{e^{ar^{\alpha}}}{r^{n+2}}$$

and

$$\frac{d}{dr}\left(\frac{2(1-br^{\alpha})e^{ar^{\alpha}}}{r^{n+3}}\right) = \frac{-2(1-br^{\alpha})e^{ar^{\alpha}}(n+3+O(r^{\alpha}))}{r^{n+4}},$$

we have

$$\begin{split} e'(r) &= -\frac{(n+2)(1-r^{\alpha})}{r} \frac{e^{ar^{\alpha}}}{r^{n+2}} \int_{B_{r}} \left( |\nabla \mathbf{u}|^{2} + 2|\mathbf{u}| \right) + \frac{e^{ar^{\alpha}}}{r^{n+2}} \int_{\partial B_{r}} \left( |\nabla \mathbf{u}|^{2} + 2|\mathbf{u}| \right) \\ &+ \frac{2(1-br^{\alpha})e^{ar^{\alpha}}(n+3+O(r^{\alpha}))}{r^{n+4}} \int_{\partial B_{r}} |\mathbf{u}|^{2} \\ &- \frac{2(1-br^{\alpha})e^{ar^{\alpha}}}{r^{n+3}} \left( \int_{\partial B_{r}} 2\mathbf{u}\partial_{\nu}\mathbf{u} + \frac{n-1}{r} \int_{\partial B_{r}} |\mathbf{u}|^{2} \right) \\ &\geq -\frac{n+2}{r} \left( e(r) + \frac{2(1-br^{\alpha})e^{ar^{\alpha}}}{r^{n+3}} \int_{\partial B_{r}} |\mathbf{u}|^{2} + W(0+) \right) \\ &+ \frac{e^{ar^{\alpha}}}{r} \left[ \frac{1}{r^{n+1}} \int_{\partial B_{r}} \left( |\nabla \mathbf{u}|^{2} + 2|\mathbf{u}| \right) + \frac{8(1-br^{\alpha})+O(r^{\alpha})}{r^{n+3}} \int_{\partial B_{r}} |\mathbf{u}|^{2} \\ &- \frac{2(1-br^{\alpha})}{r^{n+2}} \int_{\partial B_{r}} 2\mathbf{u}\partial_{\nu}\mathbf{u} \right] \\ &\geq -\frac{n+2}{r} \left( e(r) + W(0+) \right) \\ &+ \frac{(1-br^{\alpha})e^{ar^{\alpha}}}{r} \left[ \frac{1}{r^{n+1}} \int_{\partial B_{r}} \left( |\nabla \mathbf{u}|^{2} + 2|\mathbf{u}| \right) \\ &+ \frac{-2n+4+O(r^{\alpha})}{r^{n+3}} \int_{\partial B_{r}} |\mathbf{u}|^{2} - \frac{4}{r^{n+2}} \int_{\partial B_{r}} \mathbf{u}\partial_{\nu}\mathbf{u} \right]. \end{split}$$

Combining this with the computation

$$\begin{split} &\int_{\partial B_r} \left( \frac{1}{r^{n+1}} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) + \frac{-2n + 4 + O(r^{\alpha})}{r^{n+3}} |\mathbf{u}|^2 - \frac{4}{r^{n+2}} \mathbf{u} \partial_{\nu} \mathbf{u} \right) \\ &= \int_{\partial B_1} \left( |\nabla \mathbf{u}_r|^2 + 2|\mathbf{u}_r| + (-2n + 4 + O(r^{\alpha}))|\mathbf{u}_r|^2 - 4\mathbf{u}_r \partial_{\nu} \mathbf{u}_r \right) \\ &= \int_{\partial B_1} \left( |\partial_{\nu} \mathbf{u}_r - 2\mathbf{u}_r|^2 + |\partial_{\theta} \mathbf{u}_r|^2 + 2|\mathbf{u}_r| - (2n + O(r^{\alpha}))|\mathbf{u}_r|^2 \right) \\ &\geq \int_{\partial B_1} \left( |\partial_{\theta} \mathbf{c}_r|^2 + 2|\mathbf{c}_r| - (2n + O(r^{\alpha}))|\mathbf{c}_r|^2 \right) \\ &= \int_{\partial B_1} \left( |\nabla \mathbf{c}_r|^2 + 2|\mathbf{c}_r| - (2n + 4 + O(r^{\alpha}))|\mathbf{c}_r|^2 \right) \\ &= (n+2) \left[ \int_{B_1} \left( |\nabla \mathbf{c}_r|^2 + 2|\mathbf{c}_r| \right) - (2 + O(r^{\alpha})) \int_{\partial B_1} |\mathbf{c}_r|^2 \right] \\ &= (n+2)M(\mathbf{c}_r) + O(r^{\alpha}) \int_{\partial B_1} |\mathbf{u}_r|^2, \end{split}$$

we get

$$e'(r) \ge -\frac{n+2}{r}(e(r)+W(0+)) + \frac{(1-br^{\alpha})e^{ar^{\alpha}}(n+2)}{r}M(\mathbf{c}_r) + \frac{O(r^{\alpha})}{r}\int_{\partial B_1}|\mathbf{u}_r|^2.$$

To estimate  $M(\mathbf{c}_r)$ , we use the almost minimizing property of  $\mathbf{u}$  and that the epiperimetric inequality  $M(\mathbf{v}) \leq (1-\kappa)M(\mathbf{c}_r) + \kappa \frac{\beta_n}{2}$  holds for some  $\mathbf{v} \in W^{1,2}(B_1; \mathbb{R}^m)$ 

19

with  $\mathbf{v} = \mathbf{c}_r = \mathbf{u}_r$  on  $\partial B_1$  to obtain

$$(1-\kappa)M(\mathbf{c}_r) + \kappa W(0+) \ge M(\mathbf{v})$$
  
= 
$$\int_{B_1} \left( |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| \right) - 2 \int_{\partial B_1} |\mathbf{v}|^2$$
  
$$\ge \frac{1}{1+r^{\alpha}} \int_{B_1} \left( |\nabla \mathbf{u}_r|^2 + 2|\mathbf{u}_r| \right) - 2 \int_{\partial B_1} |\mathbf{u}_r|^2$$
  
= 
$$\frac{e^{-ar^{\alpha}}}{1+r^{\alpha}} W(r) + O(r^{\alpha}) \int_{\partial B_1} |\mathbf{u}_r|^2.$$

This gives

$$M(\mathbf{c}_r) \ge \frac{\frac{e^{-ar^{\alpha}}}{1+r^{\alpha}}W(r) - \kappa W(0+)}{1-\kappa} + O(r^{\alpha}) \int_{\partial B_1} |\mathbf{u}_r|^2.$$

Thus

$$\begin{split} e'(r) &\geq -\frac{n+2}{r}(e(r)+W(0+)) + \frac{(1-br^{\alpha})e^{ar^{\alpha}}(n+2)}{r} \left(\frac{\frac{e^{-ar^{\alpha}}}{1+r^{\alpha}}W(r) - \kappa W(0+)}{1-\kappa}\right) \\ &+ \frac{O(r^{\alpha})}{r} \int_{\partial B_{1}} |\mathbf{u}_{r}|^{2}. \end{split}$$

Note that from Lemmas 3 and 4, we have

$$W(0+) \leq W(r) \leq C(n,\alpha,E(\mathbf{u},1))$$

and

$$\int_{\partial B_1} |\mathbf{u}_r|^2 = \frac{1}{r^{n+3}} \int_{\partial B_r} |\mathbf{u}|^2 \le C(n, \alpha, E(\mathbf{u}, 1))$$

for  $0 < r < r_0(n, \alpha)$ . Using these estimates, we obtain

$$\begin{aligned} e'(r) &\geq -\frac{n+2}{r}(e(r) + W(0+)) + \frac{n+2}{r} \left( \frac{W(r) - \kappa W(0+)}{1 - \kappa} \right) \\ &+ \frac{O(r^{\alpha})}{r} \left( W(r) + W(0+) + \int_{\partial B_1} |\mathbf{u}_r|^2 \right) \\ &\geq -\frac{n+2}{r} e(r) + \frac{n+2}{r} \left( \frac{W(r) - W(0+)}{1 - \kappa} \right) - C_0 r^{\alpha - 1} \\ &= \left( \frac{(n+2)\kappa}{1 - \kappa} \right) \frac{e(r)}{r} - C_0 r^{\alpha - 1}, \quad 0 < r < r_0(n, \alpha), \end{aligned}$$

for some  $C_0 = C_0(n, \alpha, E(\mathbf{u}, 1))$ . Now, we take  $\delta = \delta(n, \alpha, \kappa)$  such that  $0 < \delta < \min\{\frac{(n+2)\kappa}{1-\kappa}, \alpha\}$ . Then, using the above differential inequality for e(r) and that  $e(r) = W(r) - W(0+) \ge 0$ , we obtain

$$\frac{d}{dr} \left[ e(r)r^{-\delta} + \frac{C_0}{\alpha - \delta} r^{\alpha - \delta} \right] = r^{-\delta} \left[ e'(r) - \frac{\delta}{r} e(r) \right] + C_0 r^{\alpha - \delta - 1}$$
$$\geq r^{-\delta} \left[ \left( \frac{(n+2)\kappa}{1 - \kappa} - \delta \right) \frac{e(r)}{r} - C_0 r^{\alpha - 1} \right] + C_0 r^{\alpha - \delta - 1}$$
$$\geq 0.$$

Thus

$$e(r)r^{-\delta} \le e(r)r^{-\delta} + \frac{C_0}{\alpha - \delta}r^{\alpha - \delta} \le e(r_0)r_0^{-\delta} + \frac{C_0}{\alpha - \delta}r_0^{\alpha - \delta},$$

and hence we conclude that

$$W(r) - W(0+) = e(r) \le C(n, \alpha, E(\mathbf{u}, 1))r^{\delta}.$$

Now, we consider an auxiliary function

$$\phi(r) := e^{-(2b/\alpha)r^{\alpha}}r^2, \quad r > 0,$$

which is a solution of the differential equation

$$\phi'(r) = 2 \phi(r) \frac{1 - br^{\alpha}}{r}, \quad r > 0.$$

For  $x_0 \in B_{1/2}$ , we define the 2-almost homogeneous rescalings by

$$\mathbf{u}_{x_0,r}^{\phi}(x) := \frac{\mathbf{u}(rx+x_0)}{\phi(r)}, \quad x \in B_{1/(2r)}.$$

Lemma 9 (Rotation estimate). Under the same assumptions as in Lemma 8,

$$\int_{\partial B_1} |\mathbf{u}_{x_0,t}^{\phi} - \mathbf{u}_{x_0,s}^{\phi}| \le C(n,\alpha,\kappa,E(\mathbf{u},1))t^{\delta/2}, \quad s < t < t_0(n,\alpha).$$

*Proof.* Without loss of generality, assume  $x_0 = 0$ . Then, for  $\mathbf{u}_r^{\phi} = \mathbf{u}_{0,r}^{\phi}$ ,

$$\begin{aligned} \frac{d}{dr}\mathbf{u}_r^{\phi}(x) &= \frac{\nabla \mathbf{u}(rx) \cdot x}{\phi(r)} - \frac{\mathbf{u}(rx)[\phi'(r)/\phi(r)]}{\phi(r)} \\ &= \frac{1}{\phi(r)} \left( \nabla \mathbf{u}(rx) \cdot x - \frac{2(1-br^{\alpha})}{r} \mathbf{u}(rx) \right). \end{aligned}$$

Thus, for  $0 < r < t_0 = t_0(n, \alpha)$ , using Theorem 5,

$$\begin{split} \left( \int_{\partial B_1} \left[ \frac{d}{dr} \mathbf{u}_r^{\phi}(\xi) \right]^2 dS_{\xi} \right)^{1/2} \\ &= \frac{1}{\phi(r)} \left( \int_{\partial B_1} \left| \nabla \mathbf{u}(r\xi) \cdot \xi - \frac{2(1 - br^{\alpha})}{r} \mathbf{u}(r\xi) \right|^2 dS_{\xi} \right)^{1/2} \\ &= \frac{1}{\phi(r)} \left( \frac{1}{r^{n-1}} \int_{\partial B_r} \left| \nabla \mathbf{u}(x) \cdot \nu - \frac{2(1 - br^{\alpha})}{r} \mathbf{u}(x) \right|^2 dS_x \right)^{1/2} \\ &\leq \frac{1}{\phi(r)} \left( \frac{1}{r^{n-1}} \frac{r^{n+2}}{e^{ar^{\alpha}}} \frac{d}{dr} W(\mathbf{u}, r) \right)^{1/2} = \frac{e^{cr^{\alpha}}}{r^{1/2}} \left( \frac{d}{dr} W(\mathbf{u}, r) \right)^{1/2}, \quad c = \frac{2b}{\alpha} - \frac{a}{2} \end{split}$$

Using this and Lemma 8, we can compute

$$\begin{split} \int_{\partial B_1} |\mathbf{u}_t^{\phi} - \mathbf{u}_s^{\phi}| &\leq \int_{\partial B_1} \int_s^t \left| \frac{d}{dr} \mathbf{u}_r^{\phi} \right| \, dr = \int_s^t \int_{\partial B_1} \left| \frac{d}{dr} \mathbf{u}_r^{\phi} \right| \, dr \\ &\leq C_n \int_s^t \left( \int_{\partial B_1} \left| \frac{d}{dr} \mathbf{u}_r^{\phi} \right|^2 \right)^{1/2} \, dr \\ &\leq C_n \left( \int_s^t \frac{1}{r} \, dr \right)^{1/2} \left( \int_s^t r \int_{\partial B_1} \left| \frac{d}{dr} \mathbf{u}_r^{\phi} \right|^2 \, dr \right)^{1/2} \\ &\leq C_n e^{ct^{\alpha}} \left( \log \frac{t}{s} \right)^{1/2} \left( \int_s^t \frac{d}{dr} W(\mathbf{u}, r) \, dr \right)^{1/2} \\ &\leq C \left( \log \frac{t}{s} \right)^{1/2} (W(\mathbf{u}, t) - W(\mathbf{u}, s))^{1/2} \\ &\leq C \left( \log \frac{t}{s} \right)^{1/2} t^{\delta/2}, \quad 0 < t < t_0(n, \alpha). \end{split}$$

Now, by a standard dyadic argument, we conclude that

$$\int_{\partial B_1} |\mathbf{u}_t^{\phi} - \mathbf{u}_s^{\phi}| \le C t^{\delta/2}.$$

## 7. Regularity of the regular set

In this last section we prove one of the most important results in this paper, the  $C^{1,\gamma}$  regularity of the regular set.

**Lemma 10.** Let  $C_h$  be a compact subset of  $\mathbb{R}_u$ . Then for small r, the 2-homogeneous replacements  $\mathbf{c}_{x_0,r}$  are uniformly (in  $x_0 \in C_h$ ) close to  $\mathbb{H}$  in the  $C^1(\overline{B_1}; \mathbb{R}^m)$ -topology.

*Proof.* We first claim that  $\mathbf{u}_{x_0,r}$  is uniformly close to  $\mathbb{H}$  in  $C^1(\overline{B_1}; \mathbb{R}^m)$ . Indeed, assume towards a contradiction that there exist  $x_j \in C_h, r_j \to 0$  and  $\varepsilon_0 > 0$  such that

(7.1) 
$$\operatorname{dist}(\mathbf{u}_{x_i,r_i},\mathbb{H}) \ge \varepsilon_0,$$

where the distance is measured in the  $C^1(\overline{B_1}; \mathbb{R}^m)$ -norm. Following the argument in the proof of Lemma 7 with  $\mathbf{u}_{x_j,r_j}$  in the place of  $\mathbf{u}_{t_j}$ , we obtain that up to a subsequence,

$$\mathbf{u}_{x_i,r_i} \to \mathbf{w} \quad \text{in } C^1_{\text{loc}}(\mathbb{R}^n;\mathbb{R}^m)$$

for some nonzero 2-homogeneous global solution  $\mathbf{w} \in C^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^m)$ . Then, by Corollary 1 in [1], we have either  $M(\mathbf{w}) = \beta_n/2$  or  $M(\mathbf{w}) = \overline{\beta}_n$ . Moreover, by Dini's theorem, there is  $r_0 > 0$  such that

$$W(\mathbf{u}, x_j, r) < 1/2(\beta_n/2 + \overline{\beta}_n)$$

for all  $0 < r < r_0$  and all  $x_j \in C_h$ . Thus

$$W(\mathbf{w},0,1) = \lim_{j \to \infty} W(\mathbf{u}_{x_j,r_j},0,1) = \lim_{j \to \infty} W(\mathbf{u},x_j,r_j) = \beta_n/2,$$

and hence  $\mathbf{w} \in \mathbb{H}$  by Corollary 1 in [1]. This contradicts (7.1), and we conclude that  $\mathbf{u}_{x_0,r}$  is uniformly close to  $\mathbb{H}$  in  $C^1(\overline{B_1}; \mathbb{R}^m)$ .

Now, to show that  $\mathbf{c}_{x_0,r}$  is close to  $\mathbb{H}$ , let  $\varepsilon > 0$  be given. Then there is  $r_{\varepsilon} > 0$  such that if  $x_0 \in C_h$  and  $0 < r < r_{\varepsilon}$ , then  $\|\mathbf{u}_{x_0,r} - \mathbf{h}_{x_0,r}\|_{C^1(\overline{B_1};\mathbb{R}^m)} < \varepsilon$  for some  $\mathbf{h}_{x_0,r} \in \mathbb{H}$ . Since  $\mathbf{c}_{x_0,r} = |x|^2 \mathbf{u}_{x_0,r} \left(\frac{x}{|x|}\right)$  is 2-homogeneous  $C^{1,\alpha/2}$ -regular, we have for some universal constant C > 0

$$\begin{aligned} \|\mathbf{c}_{x_0,r} - \mathbf{h}_{x_0,r}\|_{C^1(\overline{B_1};\mathbb{R}^m)} \\ &\leq C\left(\|\mathbf{c}_{x_0,r} - \mathbf{h}_{x_0,r}\|_{L^{\infty}(\partial B_1;\mathbb{R}^m)} + \|\nabla_{\theta}(\mathbf{c}_{x_0,r} - \mathbf{h}_{x_0,r})\|_{L^{\infty}(\partial B_1;\mathbb{R}^m)}\right). \end{aligned}$$

Moreover, from the fact that  $\mathbf{c}_{x_0,r}$  coincides with  $\mathbf{u}_{x_0,r}$  on  $\partial B_1$ , we conclude that

$$\|\mathbf{c}_{x_0,r} - \mathbf{h}_{x_0,r}\|_{C^1(\overline{B_1};\mathbb{R}^m)} = C\|\mathbf{u}_{x_0,r} - \mathbf{h}_{x_0,r}\|_{C^1(\overline{B_1};\mathbb{R}^m)} < C\varepsilon,$$

as desired.

**Lemma 11.** Let  $\mathbf{u}$  be an almost minimizer in  $B_1$ ,  $C_h$  a compact subset of  $\mathcal{R}_{\mathbf{u}}$ , and  $\delta$  as in Lemma 8. Then for every  $x_0 \in C_h$  there is a unique blowup  $\mathbf{u}_{x_0,0} \in \mathbb{H}$ . Moreover, there exists  $r_0 > 0$  and C > 0 such that

$$\int_{\partial B_1} |\mathbf{u}_{x_0,r}^{\phi} - \mathbf{u}_{x_0,0}| \le Cr^{\delta/2}$$

for all  $0 < r < r_0$  and  $x_0 \in C_h$ .

*Proof.* By the definition of  $\mathcal{R}_{\mathbf{u}}$  and Corollary 1 in [1], every blowup  $\mathbf{u}_{x_0,0}$  at  $x_0$  is in  $\mathbb{H}$ . Moreover, by Lemma 9 and Lemma 10,

$$\int_{\partial B_1} |\mathbf{u}_{x_0,r}^{\phi} - \mathbf{u}_{x_0,s}^{\phi}| \le Cr^{\delta/2}, \quad s < r < r_0.$$

Note that  $\mathbf{u}_{x_0,t}^{\phi}(x) = \frac{\mathbf{u}(tx+x_0)}{\phi(t)}$  and  $\lim_{t\to 0} \frac{\phi(t)}{t^2} = 1$ . Thus, if  $\mathbf{u}_{x_0,0}$  is the limit of  $\mathbf{u}_{x_0,t_j}$  for  $t_j \to 0$ , then it is also the limit of  $\mathbf{u}_{x_0,t_j}^{\phi}$ . Taking  $s = t_j \to 0$  in the above estimate and passing to the limit, we get

$$\int_{\partial B_1} |\mathbf{u}_{x_0,r}^{\phi} - \mathbf{u}_{x_0,0}| \le Cr^{\delta/2}$$

Finally, to prove the uniqueness of blowup, let  $\tilde{\mathbf{u}}_{x_0,0}$  be another blowup. Then,

$$\int_{\partial B_1} |\tilde{\mathbf{u}}_{x_0,0} - \mathbf{u}_{x_0,0}| = 0$$

By the homogeneity, we conclude that  $\mathbf{u}_{x_0,0} = \tilde{\mathbf{u}}_{x_0,0}$  in  $\mathbb{R}^n$ .

Now we are ready to prove the main result on the regularity of the regular set.

Proof of Theorem 3. The relative openness of the regular set immediately follows from the fact that  $x \mapsto W(\mathbf{u}, x, 0+)$  is upper-semicontinous and (1.3). For the regularity of  $\mathcal{R}_{\mathbf{u}}$ , we follow the argument in Theorem 5 in [1].

Step 1. Let  $x_0 \in \mathcal{R}_{\mathbf{u}}$ . By Lemma 11, there exists  $\rho_0 > 0$  such that  $B_{2\rho_0}(x_0) \subset B_1$ ,  $B_{2\rho_0}(x_0) \cap \Gamma(\mathbf{u}) = B_{2\rho_0}(x_0) \cap \mathcal{R}_{\mathbf{u}}$  and

$$\int_{\partial B_1} |\mathbf{u}_{x_1,r}^{\phi} - \frac{1}{2} \mathbf{e}(x_1) \max(x \cdot \nu(x_1), 0)^2| \le Cr^{\delta/2},$$

for any  $x_1 \in \Gamma(\mathbf{u}) \cap \overline{B_{\rho_0}(x_0)}$  and for any  $0 < r < \rho_0$ . We then claim that  $x_1 \mapsto \nu(x_1)$  and  $x_1 \mapsto \mathbf{e}(x_1)$  are Hölder continuous of order  $\gamma$  on  $\Gamma(\mathbf{u}) \cap \overline{B_{\rho_1}(x_0)}$  for some  $\gamma = \gamma(n, \alpha, \kappa) > 0$  and  $\rho_1 \in (0, \rho_0)$ . Indeed, we observe that for  $x_1$  and  $x_2$  near  $x_0$  and for small r > 0,

$$\frac{1}{2} \int_{\partial B_{1}} |\mathbf{e}(x_{1}) \max(x \cdot \nu(x_{1}), 0)^{2} - \mathbf{e}(x_{2}) \max(x \cdot \nu(x_{2}), 0)^{2} | dS_{x} \\
\leq 2Cr^{\delta/2} + \int_{\partial B_{1}} |\mathbf{u}_{x_{1},r}^{\phi} - \mathbf{u}_{x_{2},r}^{\phi}| \\
\leq 2Cr^{\delta/2} + \frac{1}{\phi(r)} \int_{\partial B_{1}} \int_{0}^{1} |\nabla \mathbf{u}(rx + (1 - t)x_{1} + tx_{2})| |x_{1} - x_{2}| dt dS_{x} \\
\leq 2Cr^{\delta/2} + C \frac{|x_{1} - x_{2}|}{r^{2}}.$$

Now, if we take  $\rho_1$  small and  $x_1, x_2 \in \overline{B_{\rho_1}(x_0)} \cap \Gamma(\mathbf{u})$  and choose  $r = |x_1 - x_2|^{\frac{2}{4+\delta}}$ , then

$$\frac{1}{2} \int_{\partial B_1} |\mathbf{e}(x_1) \max(x \cdot \nu(x_1), 0)^2 - \mathbf{e}(x_2) \max(x \cdot \nu(x_2), 0)^2| \le C |x_1 - x_2|^{\gamma}, \quad \gamma = \frac{\delta}{4 + \delta}$$

Moreover, we also have (see the proof of Theorem 5 in [1])

$$\frac{1}{2} \int_{\partial B_1} |\mathbf{e}(x_1) \max(x \cdot \nu(x_1), 0)^2 - \mathbf{e}(x_2) \max(x \cdot \nu(x_2), 0)^2 \\ \ge c(n)(|\nu(x_1) - \nu(x_2)| + |\mathbf{e}(x_1) - \mathbf{e}(x_2)|),$$

which readily implies the Hölder continuity of  $x_1 \mapsto \nu(x_1)$  and  $x_1 \mapsto \mathbf{e}(x_1)$ .

23

Step 2. We claim that for every  $\varepsilon \in (0, 1)$ , there exists  $\rho_{\varepsilon} \in (0, \rho_1)$  such that for  $x_1 \in \Gamma(\mathbf{u}) \cap \overline{B_{\rho_1}(x_0)}$  and  $y \in \overline{B_{\rho_{\varepsilon}}(x_1)}$ ,

(7.2) 
$$\mathbf{u}(y) = 0 \quad \text{if } (y - x_1) \cdot \nu(x_1) < -\varepsilon |y - x_1|,$$

(7.3) 
$$|\mathbf{u}(y)| > 0 \quad \text{if } (y - x_1) \cdot \nu(x_1) > \varepsilon |y - x_1|.$$

Indeed, if (7.2) does not hold, then we can take a sequence  $\Gamma(\mathbf{u}) \cap \overline{B_{\rho_1}(x_0)} \ni x_j \to \bar{x}$ and a sequence  $y_j - x_j \to 0$  as  $j \to \infty$  such that

$$|\mathbf{u}(y_j)| > 0$$
 and  $(y_j - x_j) \cdot \nu(x_j) < -\varepsilon |y_j - x_j|$ 

Then we consider  $\mathbf{u}_j(x) := \frac{\mathbf{u}(x_j+|y_j-x_j|x)}{|y_j-x_j|^2}$  and observe that for  $z_j := \frac{y_j-x_j}{|y_j-x_j|} \in \partial B_1$ ,  $|\mathbf{u}_j(z_j)| > 0$  and  $z_j \cdot \nu(x_j) < -\varepsilon |z_j|$ . Moreover, over a subsequence,  $\mathbf{u}_j \to \mathbf{u}_0 = \frac{1}{2}\mathbf{e}(\bar{x})\max(x\cdot\nu(\bar{x}),0)^2$  in  $C^1_{\mathrm{loc}}(\mathbb{R}^n;\mathbb{R}^m)$  (see the proof of Lemma 10). Then, for  $K := \{z \in \partial B_1 : z \cdot \nu(\bar{x}) \leq -\varepsilon/2|z|\}$ , we have that  $z_j \in K$  for large j by Step 1. We also consider a bigger compact set  $\tilde{K} := \{z \in \mathbb{R}^n : 1/2 \leq |z| \leq 2, z \cdot \nu(\bar{x}) \leq -\varepsilon/4|z|\}$ , and let  $t := \min\{\mathrm{dist}(K, \partial \tilde{K}), r_0\}$ , where  $r_0 = r_0(n, m, \alpha)$  is as in Lemma 5, so that  $B_t(z_j) \subset \tilde{K}$ . By applying Lemma 6, we obtain

$$\sup_{\tilde{K}} \left( |\nabla \mathbf{u}_j|^2 + 2|\mathbf{u}_j| \right) \ge C(n,t) \int_{B_t(z_j)} \left( |\nabla \mathbf{u}_j|^2 + 2|\mathbf{u}_j| \right) \ge c(n,m,\alpha,\varepsilon),$$

which gives

$$\sup_{\tilde{K}} \left( |\nabla \mathbf{u}_0| + |\mathbf{u}_0| \right) > 0.$$

However, this is a contradiction since  $\mathbf{u}_0(x) = \frac{1}{2}\mathbf{e}(\bar{x}) \max(x \cdot \nu(\bar{x}), 0)^2 = 0$  in  $\bar{K}$ . On the other hand, if (7.3) is not true, then we take a sequence  $\Gamma(\mathbf{u}) \cap \overline{B_{\rho_1}(x_0)} \ni x_j \to \bar{x}$  and a sequence  $y_j - x_j \to 0$  such that  $|\mathbf{u}(y_j)| = 0$  and  $(y_j - x_j) \cdot \nu(x_j) > \varepsilon |y_j - x_j|$ . For  $\mathbf{u}_j$ ,  $\mathbf{u}_0(x) = \frac{1}{2}\mathbf{e}(\bar{x}) \max(x \cdot \nu(\bar{x}), 0)^2$  and  $z_j$  as above, we will have that  $\mathbf{u}_j(z_j) = 0$  and  $z_j \in K' := \{z \in \partial B_1 : z \cdot \nu(\bar{x}) \ge \varepsilon/2|z|\}$ . We may assume  $z_j \to z_0 \in K'$ , and obtain  $\mathbf{u}_0(z_0) = 0$ , which is a contradiction.

Step 3. By rotations we may assume that  $\nu(x_0) = \mathbf{e}_n$  and  $\mathbf{e}(x_0) = \mathbf{e}_1$ . Fixing  $\varepsilon = \varepsilon_0$ , by Step 2 and the standard arguments, we conclude that there exists a Lipschitz function  $g : \mathbb{R}^{n-1} \to \mathbb{R}$  such that for some  $\rho_{\varepsilon_0} > 0$ ,

$$B_{\rho_{\varepsilon_0}}(x_0) \cap \{\mathbf{u} = 0\} = B_{\rho_{\varepsilon_0}}(x_0) \cap \{x_n \le g(x')\},\$$
  
$$B_{\rho_{\varepsilon_0}}(x_0) \cap \{|\mathbf{u}| > 0\} = B_{\rho_{\varepsilon_0}}(x_0) \cap \{x_n > g(x')\}.$$

Now, taking  $\varepsilon \to 0$ , we can see that  $\Gamma(\mathbf{u})$  is differentiable at  $x_0$  with normal  $\nu(x_0)$ . Recentering at any  $\bar{x} \in B_{\rho_{\varepsilon_0}}(x_0) \cap \Gamma(\mathbf{u})$  and using the Hölder continuity of  $\bar{x} \mapsto \nu(\bar{x})$ , we conclude that g is  $C^{1,\gamma}$ . This completes the proof.

APPENDIX A. EXAMPLE OF ALMOST MINIMIZERS

**Example 1.** Let **u** be a solution of the system

$$\Delta \mathbf{u} + \mathbf{b}(x)\nabla \mathbf{u} - \frac{\mathbf{u}}{|\mathbf{u}|} \chi_{\{|\mathbf{u}|>0\}} = \mathbf{0} \quad in \ B_1,$$

where  $\mathbf{b} \in L^p(B_1; \mathbb{R}^m)$ , p > n, is the velocity field. Then  $\mathbf{u}$  is an almost minimizer with a gauge function  $\omega(r) = C(n, p, \|\mathbf{b}\|_{L^p(B_1; \mathbb{R}^m)})r^{1-n/p}$ .

*Proof.* Let  $B_r(x_0) \in B_1$  and  $\mathbf{v} \in \mathbf{u} + W_0^{1,2}(B_r(x_0))$ . Then  $\mathbf{u}$  satisfies the weak equation

$$\int_{B_r(x_0)} \left( \nabla \mathbf{u} \cdot \nabla (\mathbf{u} - \mathbf{v}) - (\mathbf{u} - \mathbf{v}) \mathbf{b} \nabla \mathbf{u} + (\mathbf{u} - \mathbf{v}) \frac{\mathbf{u}}{|\mathbf{u}|} \chi_{\{|\mathbf{u}| > 0\}} \right) = 0.$$

Thus

$$\int_{B_r(x_0)} |\nabla \mathbf{u}|^2 = \int_{B_r(x_0)} \left( \nabla \mathbf{u} \cdot \nabla \mathbf{v} + (\mathbf{u} - \mathbf{v}) \mathbf{b} \nabla \mathbf{u} + (\mathbf{v} - \mathbf{u}) \frac{\mathbf{u}}{|\mathbf{u}|} \chi_{\{|\mathbf{u}|>0\}} \right)$$
$$\leq \int_{B_r(x_0)} \left( 1/2 |\nabla \mathbf{u}|^2 + 1/2 |\nabla \mathbf{v}|^2 + (\mathbf{u} - \mathbf{v}) \mathbf{b} \nabla \mathbf{u} + |\mathbf{v}| - |\mathbf{u}| \right).$$

It follows that

(A.1) 
$$\int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \le \int_{B_r(x_0)} \left( |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| + 2(\mathbf{u} - \mathbf{v})\mathbf{b}\nabla\mathbf{u} \right).$$

To estimate the last term, we apply Hölder's inequality to have

 $\int_{B_r(x_0)} 2(\mathbf{u} - \mathbf{v}) \mathbf{b} \nabla \mathbf{u} \leq 2 \|\mathbf{u} - \mathbf{v}\|_{L^{p^*}(B_r(x_0);\mathbb{R}^m)} \|\mathbf{b}\|_{L^p(B_r(x_0);\mathbb{R}^m)} \|\nabla \mathbf{u}\|_{L^2(B_r(x_0);\mathbb{R}^m)},$ with  $p^* = 2p/(p-1)$ . Since  $\mathbf{u} - \mathbf{v} \in W_0^{1,2}(B_r(x_0);\mathbb{R}^m)$ , we have by the Sobolev's inequality

$$\|\mathbf{u} - \mathbf{v}\|_{L^{p^*}(B_r(x_0);\mathbb{R}^m)} \le Cr^{\gamma} \|\nabla(\mathbf{u} - \mathbf{v})\|_{L^2(B_r(x_0);\mathbb{R}^m)}, \quad \gamma = 1 - n/p,$$

and thus

$$\begin{split} \int_{B_r(x_0)} 2(\mathbf{u} - \mathbf{v}) \mathbf{b} \nabla \mathbf{u} &\leq Cr^{\gamma} \| \nabla \mathbf{u} \|_{L^2(B_r(x_0);\mathbb{R}^m)} \| \nabla (\mathbf{u} - \mathbf{v}) \|_{L^2(B_r(x_0);\mathbb{R}^m)} \\ &\leq Cr^{\gamma} \left( \| \nabla \mathbf{u} \|_{L^2(B_r(x_0);\mathbb{R}^m)}^2 + \| \nabla (\mathbf{u} - \mathbf{v}) \|_{L^2(B_r(x_0);\mathbb{R}^m)}^2 \right) \\ &\leq Cr^{\gamma} \int_{B_r(x_0)} \left( | \nabla \mathbf{u} |^2 + | \nabla \mathbf{v} |^2 \right). \end{split}$$

Combining this and (A.1) yields

$$\int_{B_r(x_0)} \left( (1 - Cr^{\gamma}) |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \le \int_{B_r(x_0)} \left( (1 + Cr^{\gamma}) |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| \right).$$

This implies

$$(1 - Cr^{\gamma}) \int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \le (1 + Cr^{\gamma}) \int_{B_r(x_0)} \left( |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| \right)$$

and hence we conclude that for  $0 < r < r_0(n, p, \|\mathbf{b}\|_{L^p(B_1; \mathbb{R}^m)})$ 

$$\int_{B_r(x_0)} \left( |\nabla \mathbf{u}|^2 + 2|\mathbf{u}| \right) \le (1 + Cr^{\gamma}) \int_{B_r(x_0)} \left( |\nabla \mathbf{v}|^2 + 2|\mathbf{v}| \right),$$

with  $C = C(n, p, \|\mathbf{b}\|_{L^{p}(B_{1};\mathbb{R}^{m})}).$ 

#### References

- J. Andersson, H. Shahgholian, N. Uraltseva and G. Weiss, Equilibrium points of a singular cooperative system with free boundary, Adv. Math. 280 (2015), 743–771.
- [2] G. Anzellotti, On the C<sup>1,α</sup>-regularity of ω-minima of quadratic functionals, Boll. Un. Mat. Ital. C (6) 2 (1983), no. 1, 195–212.
- [3] G. David, M. Engelstein, T. Toro, Free boundary regularity for almost-minimizers, Adv. Math. 350 (2019), 1109–1192.
- [4] G. David, T. Toro, Regularity of almost minimizers with free boundary, Calc. Var. Partial Differential Equations 54 (2015), no. 1, 455–524.
- [5] D. De Silva, O. Savin, Thin one-phase almost minimizers, Nonlinear Anal. 193 (2020), 111507, 23 pp.
- [6] D. De Silva, O. Savin, Almost minimizers of the one-phase free boundary problem, Comm. Partial Differential Equations 45 (2020), no. 8, 913–930.
- [7] Q. Han and F. Lin, *Elliptic partial differential equations*, Second edition. Courant Lecture Notes in Mathematics, 1. Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2011. x+147 pp.

- [8] S. Jeon, A. Petrosyan, Almost minimizers for certain fractional variational problems, Algebra i Analiz 32 (2020), no. 4, 166–199; reprinted in St. Petersburg Math. J. 32 (2021), no. 4, 729–751.
- [9] S. Jeon and A. Petrosyan, Almost minimizers for the thin obstacle problem, Calc. Var. Partial Differential Equations 60 (2021) Paper No. 124, 59.
- [10] S. Jeon, A. Petrosyan, M. Smit Vega Garcia, Almost minimizers for the thin obstacle problem with variable coefficients, Preprint
- G.S. Weiss, A homogeneity improvement approach to the obstacle problem, Invent. Math. 138 (1999), no. 1, 23–50.
- [12] G.S. Weiss, An obstacle-problem-like equation with two phases: pointwise regularity of the solution and an estimate of the Hausdorff dimension of the free boundary, Interfaces Free Bound. 3 (2001), no. 2, 121–128.

(Daniela De Silva) DEPARTMENT OF MATHEMATICS BARNARD COLLEGE, COLUMBIA UNIVERSITY NEW YORK, NY, 10027 Email address, Daniela De Silva: desilva@math.columbia.edu

(Seongmin Jeon) DEPARTMENT OF MATHEMATICS KTH ROYAL INSTITUTE OF TECHNOLOGY 100 44 STOCKHOLM, SWEDEN *Email address*, Seongmin Jeon: seongmin@kth.se

(Henrik Shahgholian) DEPARTMENT OF MATHEMATICS KTH ROYAL INSTITUTE OF TECHNOLOGY 100 44 STOCKHOLM, SWEDEN *Email address*, Henrik Shahgholian: henriksh@kth.se