

Observational constraints on Starobinsky $f(R)$ cosmology from cosmic expansion and structure growth data

P. Bessa¹, M. Campista^{2,3}, A. Bernui³

¹ International PhD Program in Astrophysics, Cosmology and Gravitation (PPGCosmo), Espírito Santo, Brazil

² Instituto Politécnico, Universidade Federal do Rio de Janeiro (UFRJ-Campus Macaé), Macaé, Rio de Janeiro, Brazil

³ Coordenação de Astronomia e Astrofísica, Observatório Nacional (ON), Rio de Janeiro, Brazil

Received: date / Revised version: date

Abstract. The unknown physical nature of the Dark Energy motivates in cosmology the study of modifications of the gravity theory at large distances. One of these types of modifications are the theories known as $f(R)$ gravity. In this paper we use observational data to both constraint and test the Starobinsky $f(R)$ model [57], using updated measurements from the dynamics of the expansion of the universe, $H(z)$, and the growth rate of cosmic structures, $[f\sigma_8](z)$, where the distinction between the concordance Λ CDM model and modified gravity models, $f(R)$, become clearer. We use MCMC likelihood analyses to explore the parameters space of the $f(R)$ model using $H(z)$ and $[f\sigma_8](z)$ data, both individually and jointly, and further check which of the models better fit the joint data. To further test the Starobinsky model, we use a method proposed by Linder [38], where the data from the observables is jointly binned in redshift space. This allows to further explore the model's parameter that better fits the data in comparison to the Λ CDM model. Our analyses show that the result from the MCMC run using $[f\sigma_8](z)$ data alone fits the analyzed model better than using the joint data. At the end, we reaffirm that this joint analysis is able to break the degenerescence between modified gravity models through the $f\sigma_8$ observable, as the original author proposed; our results show that the $f(R)$ Starobinsky model although not favored by the available data, cannot be discarded and is a possible alternative to the Λ CDM model.

PACS. PACS-key describing text of that key – PACS-key describing text of that key

1 Introduction

The search for an explanation to the current phase of accelerated expansion of the universe is one of the most important paradigms in modern cosmology [3]. With the available observational information, the concordance model of cosmology that best fits the data is the flat Λ CDM [4] model, where the universe is filled in with cold dark matter (CDM) and a dark energy (DE) component –in the form of a cosmological constant Λ – in addition to the standard baryonic and electromagnetic ingredients. The main concern with the DE is that its physical interpretation is still unknown. Clearly, other plausible causes of the observed accelerated expansion should be explored [36, 13, 39, 62, 20, 25]. Because Λ CDM is based on Einstein's general relativity (GR), one possible way is to explore cosmological models not based on GR [40, 15, 53, 51, 52, 14, 47, 17, 7, 16]. However, there is a degeneracy problem between models supported by GR and models based on other metric theories, generally termed Modified Gravity (MG) theories. This degenerescence cannot be broken using data from the dynamical evolution of the Universe alone, as the same dynamics can be explained by different MG theories as well as by evolving DE fluids, $\omega = \omega(a)$, where a is the

scale factor [32, 24, 43]. The best observable to discriminate between MG theories and Λ CDM is the growth rate of cosmic structures, $f(a)$, defined as [58]

$$f(a) \equiv \frac{d \ln \delta}{d \ln a}, \quad (1)$$

where δ is the matter density contrast. The growth rate $f = f(a)$ has indeed the potential to constrain models based on MG theories and Λ CDM DE (based on GR) from the measure of the growth index, γ , when one parametrizes f as [40]

$$f(a) = \Omega_m(a)^\gamma. \quad (2)$$

In the Λ CDM concordance model, $\gamma \simeq 0.55$, and for some MG models γ evolves with time: $\gamma \simeq 0.41 - 0.21z$ [12]. In fact, changing the gravitational theory will affect the way how matter clumps at all scales, beyond the expansion of the homogeneous universe. As such, alternative cosmological models based on MG theories make very different predictions for the matter clustering and evolution.

Direct measurements of $f(a)$, or equivalently of $f(z)$, where the scale factor and the redshift are related by $a = 1/(1+z)$, are difficult to obtain. Instead the data available comes from the measurements of the galaxy redshift

space distortions in the form of a product of two quantities evolving with time: $[f\sigma_8](z) \equiv f(z)\sigma_8(z)$, where σ_8 is the root-mean-square linear fluctuation of the matter distribution at the scale of 8 Mpc/ h , h the dimensionless Hubble parameter, where $H_0 = 100 h$ km/s/Mpc. Notice that the letter f refers to the growth rate of cosmic structures, that is, it represents the function $f(a)$ or $f(z)$, but it is commonly used in the form $f(R)$ to refer to MG theories. In what follows the symbol $f(R)$, as a function of R , only refers to MG theories, while the letter f alone or the function $f(a)$, or $f(z)$, refers to the cosmic growth rate. In this paper we use this observational parameter, $[f\sigma_8]$ and the background evolution of the universe, H to explore the parameter space and test a model of $f(R)$ gravity, namely, the Starobinsky $f(R)$ model. We use a Markov Chain Monte Carlo (MCMC) method to explore the parameter space of the theory with the selected datasets for $H(z)$ and $[f\sigma_8](z)$, both individually and jointly. At the end, we further extend our analysis with a joint study of the observables proposed by Linder [38] and, in the context of Modified Gravity, more recently by Matsumoto [44].

This work is organized as follows. A brief description of the Starobinsky cosmological model [57] is presented in section 2. In section 3 we provide details of the two data sets used in the analyses, and in section 4 we describe the methodology of the analyses performed, show our results, and provide their statistical interpretation. The conclusions obtained from these analyses, and the final remarks of our work are addressed in section 5.

2 Starobinsky $f(R)$ model cosmology

2.1 Background space-time

A general $f(R)$ theory in the metric formalism that is minimally coupled to matter has an action of the form [28]

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa} f(R) + \mathcal{L}_m(g_{\mu\nu}) \right], \quad (3)$$

where we include in the action the matter Lagrangian density, \mathcal{L}_m , and define $\kappa \equiv 8\pi G$ (we adopt units with $c = 1$). This allows us to use geometrized units more easily when needed. In this work, we shall define the function $f(R)$ as in the model introduced by Starobinsky [57], of the form

$$f(R) \equiv R + \lambda R_0 \left(\left(1 + \frac{R^2}{R_0^2} \right)^{-n} - 1 \right), \quad (4)$$

where we shall consider the case $n = 1$. R_0 and λ are model parameters related to the observed DE density and a characteristic curvature respectively [22], which can be constrained by the measured cosmological parameters using equation (13).

The variation with respect to the metric gives rise to the field equations of this model, given by the extended

Einstein equations

$$F(R)R_{\mu\nu} - \frac{1}{2}f(R)g_{\mu\nu} - [\nabla_\mu \nabla_\nu - g_{\mu\nu} \square]F(R) = \kappa T_{\mu\nu}, \quad (5)$$

$$F(R)R - 2f(R) + 3\square f(R) = \kappa T, \quad (6)$$

where we define $F(R) \equiv df(R)/dR$, $T \equiv g^{\mu\nu}T_{\mu\nu}$, and the second equation is the trace of the first one, useful to derive the dynamics of the $f(R)$ function, which is more complicated than in GR where one has an algebraic relation between R and T [56].

To model our cosmological space-time we adopt the Friedmann-Lemaître-Robertson-Walker (FLRW) for the background space-time filled with a perfect fluid, with metric and stress-energy tensor given by, respectively,

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 - Kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right], \quad (7)$$

$$T^{\mu\nu} = (\rho + P)u^\mu u^\nu + Pg^{\mu\nu}, \quad (8)$$

where we take, as usual, K as the scalar curvature of the 3-space, and ρ and P , are the density and pressure of the perfect fluid, respectively. u^μ is the 4-velocity vector of an observer comoving with the fluid; $a = a(t)$ is the dimensionless scale factor, with $a(t_0) = 1$, t_0 is today's cosmic time, and we always use geometrized units, unless otherwise noted. Because we shall compare our analyses with those of the flat Λ CDM model, we consider the flat FLRW space-time, i.e., $K = 0$.

One can analyse the dynamical evolution of our cosmological model using equations (7) and (8) in equation (6), obtaining the modified Friedmann equations

$$H^2 = \frac{\kappa}{3F(R)} \left[\rho + \frac{RF(R) - f(R)}{2} \right], \quad (9)$$

$$2\dot{H} + 3H^2 = \frac{\kappa}{F(R)} [P + \dot{R}F''(R) + 2H\dot{R}F'(R)] \quad (10)$$

$$+ \ddot{R}F'(R) + \frac{1}{2}(f(R) - Rf(R)), \quad (11)$$

where $\dot{}$ means derivative with respect to the cosmic time t . These equations must obey certain constraints in order that our model be stable and correctly reproduces the late-time acceleration of the universe, without deviating from Λ CDM at early times (associated with $R \gg R_0$). An analysis of such constraints on general $f(R)$ theories can be found in [56]. Following [57], it suffices that (4) should satisfy

$$F(R) > 0, \quad F'(R) > 0, \quad R \geq R_1, \quad \lambda > \frac{8\sqrt{3}}{3}, \quad n = 1, \quad (12)$$

where R_1 is the Ricci curvature of a de Sitter point in the space of solutions. More stringent constraints on n and R_1 can be obtained by likelihood analyses on various data sets for the Starobinsky and other $f(R)$ models, like the ones performed in ref.[21].

It will be convenient to write eq. (4) in terms of the measured cosmological parameters $\Omega_{m0} = \frac{8\pi G\rho_{m0}}{3H_0^2}$ and $\Omega_{\Lambda0} = \frac{8\pi G\rho_{\Lambda}}{3H_0^2}$. In this case, one can find the correct way to write the parameters by making the model return to Λ CDM at high z limit [57, 22]

$$f(R) = R + 6\lambda H_0^2(1 - \Omega_m) \quad (13)$$

$$\times \left(\left(1 + \frac{R^2}{[6H_0^2(1 - \Omega_m)]^2} \right)^{-n} - 1 \right) \quad (14)$$

This relation allows us to constrain R_0 and λ using data linked to the observables Ω_{m0} and H_0 .

2.2 Metric perturbations

In order to go beyond the background cosmological observables and test our theory against structure growth we need to introduce metric perturbations. Following the general perturbative procedure for scalar-tensor theories found in [60], in the usual Newtonian (or comoving) gauge, the perturbed metric is given by

$$ds^2 = -(1 + 2\Psi)dt^2 + (1 - 2\Phi)a^2(t)\delta_{ij}dx^i dx^j, \quad (15)$$

where $i, j = 1, 2, 3$, and Ψ, Φ are the Bardeen potentials which satisfy $\Psi = \Phi$ in the absence of anisotropic stress. The physical processes we are interested in, and the cosmological observables associated with them (i.e., accelerated cosmic expansion and growth of cosmic structures), are all well within the scale of the sub-horizon approximation $k \gg a^2 H^2$. In this case, we can write the evolution equation for the matter density contrast, δ , as

$$\ddot{\delta} + 2H\dot{\delta} + \frac{k^2}{a^2}\Phi = 4\pi G\mu(k, a)\rho_m\delta \quad (16)$$

where G is the Newtonian gravitational constant. The $\mu(k, a)$ factor is written as

$$\mu(k, a) \equiv \frac{1}{8\pi F(R)} \left(\frac{1 + 4\frac{k^2}{a^2 R} m}{1 + 3\frac{k^2}{a^2 R} m} \right), \quad (17)$$

where m is a parameter that quantifies the deviation from the Λ CDM model

$$m \equiv \frac{RF'(R)}{F(R)} \implies m|_{\Lambda CDM} = 0.$$

Equation (16) allow us to analyse how our cosmological model should behave at the limits of very small and very large scales; since R is of order $H(z)$, we obtain

$$\lim_{k \gg a^2 H^2} \mu(k) = \frac{4}{3} \frac{1}{8\pi F(R)}, \quad (18)$$

$$\lim_{k \ll a^2 H^2} \mu(k) = \frac{1}{8\pi F(R)}, \quad (19)$$

and we observe that, at very small scales, the modification of GR gives an extra factor of $\frac{4}{3}$ to the force term of

the equation, therefore gravity becomes stronger and cosmic structures form faster. In the opposite limit, at large scales, the equation (16) is the same as in GR [60], now with the term $1/F(R)$, which arises naturally in a theory that couples to gravity. In the language of Scalar-Tensor theory, this would be the term that couples gravity to the scalar field.

The regime of cosmic structure formation lies between these scales, so that we need the full equation (16) to account for the physics in this regime. However, when the Hubble scale is not large enough in comparison to the perturbations (high redshift), or when we have scales of order H_0 (low redshift), the behavior of structure formation should have the Λ CDM model as a limit.

It is useful to rewrite equation (16) as an ODE on the variable $N \equiv \ln a$, also called the e-fold number, which can be obtained after a simple chain rule derivation

$$\delta'' + \left(2 + \frac{H'}{H} \right) \delta' - \frac{3}{2}\Omega_m(a)\mu(k, a)\delta = 0, \quad (20)$$

where $'$ means derivative with respect to N , and $\Omega_m(a)$ is the density parameter of (dark + baryonic) matter as a function of the scale factor a , defined above. This is the equation to be solved in order to find the linear growth of our model, using the definition of the growth function (1) in the same way as in GR.

3 Observational Data

There are in the literature several compilations of $H(z)$ and $[f\sigma_8](z)$ observational data. In this section, we are going to present the datasets we are considering to impose observational constraints in the Starobinsky model.

3.1 $H(z)$ data

The history of expansion of the universe is probed by the observational Hubble parameter $H(z)$. It can be measured by several independent methodologies, most of them, based on distance measurements of galactic objects as supernovae and quasars. However, it is known that in some approaches the cosmological distance measurements depend on a fiducial model, which makes the use of these data problematic when the objective is to constraint the free parameters of cosmological models in alternative scenarios.

In this work, since we are considering a gravity model of type $f(R)$, we are going to use in our analyses, the $H(z)$ measurements obtained by the differential age technique, that is independent of fiducial model. This technique, also known as cosmic chronometers (CC), was proposed by Jimenez and Loeb in [34].

The basic idea of the CC data consists in the spectroscopic determination of age difference between two passively evolving early-type galaxies. The assumption of old galaxies to realize the age difference measurements is important to assure that the galaxies were formed at the

same time, although are localized in slightly different redshift.

The Hubble parameter is directly related to the measured quantity dz/dt by

$$H(z) = -\frac{dz}{dt} \frac{1}{(1+z)}. \quad (21)$$

In this work we use the compilation of 31 measurements of $H(z)$, covering the range $0.07 < z < 1.965$ [65]. The data is shown in Table 1 and in Fig. 1.

Table 1: The 31 Hubble parameter data points, $H(z)$, and their respective 1σ errors $\sigma_H(z)$ from the CC data [65]. The units for both $H(z)$, $\sigma_H(z)$ are km/s/Mpc.

z	$H(z)$	$\sigma_H(z)$
0.07	69.0	19.6
0.09	69.0	12.0
0.12	68.6	26.2
0.17	83.0	8.0
0.179	75.0	4.0
0.199	75.0	5.0
0.2	72.9	29.6
0.27	77.0	14.0
0.28	88.8	36.6
0.352	83.0	14.0
0.3802	83.0	13.5
0.4	95.0	17.0
0.4004	77.0	10.2
0.4247	87.1	11.2
0.4497	92.8	12.9
0.47	89.0	49.6
0.4783	80.9	9.0
0.48	97.0	62.0
0.593	104.0	13.0
0.68	92.0	8.0
0.781	105.0	12.0
0.875	125.0	17.0
0.88	90.0	40.0
0.9	117.0	23.0
1.037	154.0	20.0
1.3	168.0	17.0
1.363	160.0	33.6
1.43	177.0	18.0
1.53	140.0	14.0
1.75	202.0	40.0
1.965	186.5	50.4

3.2 $[f\sigma_8](z)$ data

Precise measurements of $[f\sigma_8](z)$ can be done using the Redshift Space Distortions (RSD) approach, by studying the peculiar velocities caused by local gravitational potentials that introduce distortions in the two-point correlation function of cosmic objects [35]. In fact, the calculation of the anisotropic two-point correlation function,




Fig. 1: The CC Hubble parameter data, $H(z)$, displayed in Table 1.

$\xi(s, \mu)$ [31], allows us to measure $f\sigma_8$, where σ_8 is the root-mean-square linear fluctuation of the matter distribution at the scale of 8 Mpc/h (for other approaches to study matter clustering see, e.g., [11, 41, 26, 27]). The literature reports diverse compilations of measurements of $[f\sigma_8](z)$ (see, e.g., [48, 5]) which we update here. Our compilation, shown in Table 2 and in Fig. 2, takes into account:

1. We consider $[f\sigma_8](z)$ data obtained from disjoint or uncorrelated redshift bins when the measurements concern the same cosmological tracer, and data from possibly correlated redshift bins when different cosmological tracers were analysed;
2. We consider the latest measurement of $[f\sigma_8](z)$ when the same survey collaboration performed two or more measurements corresponding to several data releases.




Fig. 2: The $[f\sigma_8](z)$ data displayed in Table 2.

3.3 Joint data

The binned data given in Table 3 was obtained binning simultaneously the $[f\sigma_8](z)$ and $H(z)$ (from the Table 2

Table 2: Updated compilation of 26 $[f\sigma_8](z)$ data with their respective 1σ errors, $\sigma_{f\sigma_8}$.

Survey	z	$[f\sigma_8] \pm \sigma_{f\sigma_8}$	Reference	Cosmological tracer
ALFALFA	0.013	0.46 ± 0.06	[9]	HI line sources
SNIa+IRAS	0.02	0.398 ± 0.065	[61]	galaxies+SNIa
SNIa+6dFGS	0.02	0.428 ± 0.0465	[33]	galaxies+SNIa
6dFGS	0.025	0.39 ± 0.11	[1]	voids
6dFGS	0.067	0.423 ± 0.055	[18]	galaxies
SDSS-veloc	0.10	0.370 ± 0.130	[29]	galaxies
SDSS-IV	0.15	0.53 ± 0.16	[5]	MGS
BOSS-LOWZ	0.32	0.384 ± 0.095	[59]	LOWZ and CMASS samples
SDSS-IV	0.38	0.500 ± 0.047	[5]	BOSS galaxy sample
WiggleZ	0.44	0.413 ± 0.080	[19]	ELG
BOSS-CMASS	0.57	0.453 ± 0.022	[46]	voids+galaxies
SDSS-CMASS	0.59	0.488 ± 0.060	[23]	CMASS galaxy sample
VIPERS PDR-2	0.60	0.550 ± 0.120	[54]	galaxies
SDSS-IV	0.70	0.448 ± 0.043	[5]	eBOSS DR16 LRG
WiggleZ	0.73	0.437 ± 0.072	[19]	ELG
SDSS-IV	0.74	0.50 ± 0.11	[8]	eBOSS DR16 Voids
Vipers v7	0.76	0.440 ± 0.040	[64]	galaxies
SDSS-IV	0.85	0.52 ± 0.10	[8]	eBOSS DR16 Voids
SDSS-IV	0.85	0.315 ± 0.095	[5]	eBOSS DR16 ELG
VIPERS PDR-2	0.86	0.400 ± 0.110	[54]	galaxies
SDSS-IV	0.978	0.379 ± 0.176	[66]	eBOSS DR14 Quasar
Vipers v7	1.05	0.280 ± 0.080	[64]	galaxies
FastSound	1.40	0.482 ± 0.116	[50]	ELG
SDSS-IV	1.48	0.30 ± 0.13	[8]	eBOSS DR16 Voids
SDSS-IV	1.48	0.462 ± 0.045	[5]	eBOSS DR16 Quasar
SDSS-IV	1.944	0.364 ± 0.106	[66]	eBOSS DR14 Quasar

and the Table 1, respectively) in 5 redshift bins: $(0.0, 0.30]$, $(0.30, 0.60]$, $(0.60, 0.85]$, $(0.85, 1.4]$, $(1.4, 2.0]$, with mean redshifts: $\bar{z} = 0.15, 0.45, 0.725, 1.125, 1.7$. The number of data pairs $(f\sigma_8, H)$ in each bin were: $(7, 9)$, $(6, 10)$, $(6, 2)$, $(4, 6)$, $(3, 4)$, respectively. The values in these 5 bins for $[f\sigma_8](z)$ and $H(z)$, and their errors, correspond to their variance weighted means [45]. This binned data will be used in the χ^2 joint analysis.

Table 3: Binned data of $[f\sigma_8](z)$ and $H(z)$, obtained by calculating the variance weighted mean in each redshift bin (see the text for details).

\bar{z}	$[f\sigma_8](z)$	$H(z)$
0.15	0.4284 ± 0.0338	75.367 ± 5.741
0.45	0.4647 ± 0.0288	88.880 ± 6.717
0.725	0.4433 ± 0.0313	98.500 ± 7.071
1.125	0.3852 ± 0.0602	135.667 ± 10.247
1.70	0.3753 ± 0.0541	176.375 ± 15.300

4 Analyses and Results

In this section we describe the parameter space analysis of the $f(R)$ model 4, using the selected observational data described in the previous section 3. To obtain the theoretical predictions for $H(z)$ and $f(z)$ from the chosen model (3), we integrate equations (9) and (20) numerically and use their results in the assembling of the likelihood function. This gives us our statistical analyses.

The standard Bayesian inference will be considered for the parameter estimation. To investigate how appropriate the data is to constrain the parameter space of the model, they will be considered separately and later combined.

4.1 $H(z)$ and $[f\sigma_8](z)$ observational constraints

Recently, the tension between different estimates of the Hubble constant has drawn the attention of cosmologists. The local determination of the Hubble constant from SHOES collaboration [55] is $H_0^{\text{SHOES}} = 74.03 \pm 1.42$ km/s/Mpc while the value inferred by the Planck collaboration [4], from the Cosmic Microwave Background (CMB) analysis in a flat Λ CDM framework is $H_0^{\text{Planck}} = 67.36 \pm 0.54$ km/s/Mpc from the last Planck collaboration results [4].

The divergence in H_0 measurements is the main reason to look for alternative approaches to perform analyses that are independent of the H_0 value [49]. In our analyses we shall consider $H(z)/H_0$ instead of $H(z)$ data, and for this we use $H_0 = H_0^{\text{Planck}}$ [4].

For the model (4) and $H(z)$ data, the Likelihood function is given by:

$$\mathcal{L}(z|\theta) \propto -\frac{1}{2} \sum_i^{N_H} \frac{(H_{th}^i(z|\theta) - H_{obs}^i(z))^2}{\sigma_i^2}, \quad (22)$$

where N_H is the number of points in the dataset, $\theta = (\Omega_{m0}, \lambda)$ are the free parameters of the model, $H_{th}^i(z|\theta)$ is the theoretical Hubble parameter at redshift z_i , $H_{obs}^i(z)$ and σ are the observation error values of the Hubble parameter given in 1.

The theoretical Hubble parameter depends on the current Hubble parameter H_0 . To eliminate this dependency, we will follow the approach introduced by [6], that consists of marginalization of the likelihood over H_0 parameters

$$\mathcal{L} = \Gamma - \frac{B^2}{A} + \ln A - 2 \ln \left[1 + \text{erf} \left(\frac{B}{\sqrt{2A}} \right) \right], \quad (23)$$

where, by the definition of $E(z) \equiv H_{th}/H_0$,

$$A = \sum_{i=1}^N \frac{E^2(z_i)}{\sigma_i^2}, \quad (24)$$

$$B = \sum_{i=1}^N \frac{E(z_i)H_{obs}(z_i)}{\sigma_i^2}, \quad (25)$$

and

$$\Gamma = \sum_{i=1}^N \frac{H_{obs}^2(z_i)}{\sigma_i^2}. \quad (26)$$

Similarly, the Likelihood function for $[f\sigma_8](z)$ data is given by:

$$\mathcal{L}(z|\theta) \propto -\frac{1}{2} \sum_i^{N_f} \frac{(f\sigma_8^{ith}(z|\theta) - f\sigma_8^{iobs}(z))^2}{\sigma_i^2}, \quad (27)$$

where we have the same parametric space $\theta = (\Omega_{m0}, \lambda)$, $f\sigma_8^{ith}(z|\theta)$ is the theoretical growth function given by the model with parameters θ at redshift z_i ; $f\sigma_8^{iobs}(z)$ and σ are the observed values, given by the data in Table 2; N_f is the size of the $f\sigma_8$ dataset. The $f\sigma_8$ function does not have H_0 as a free parameter and therefore does not need to be marginalized. The parametric space of both observables $H(z)$ and $[f\sigma_8](z)$ are the same and the joint likelihood function is given by the product of the individual likelihoods according to

$$\mathcal{L} = \mathcal{L}_H \times \mathcal{L}_{f\sigma_8}. \quad (28)$$

The Bayesian inference of the parameters is obtained through the expected values of the posterior density $p(\theta|z)$ and since the posteriori distribution is unknown for the

model, i.e., it can not be approximated by a normal or gaussian distribution, we are considering the Bayes Theorem that establishes

$$p(\theta|z) = \mathcal{L}(z|\theta) \times \Pi(\theta) \quad (29)$$

where $\Pi(\theta)$ is the prior.

The parametric space of the variables of the model will be explored following the methodology of Markov chain Monte Carlo (MCMC) and the Metropolis–Hastings algorithm to generate a set of samples from a posteriori distribution. To implement the MCMC, we are using a Python open source code, `emcee` [30]

The posteriori distribution was generated for the following flat prior intervals: $\lambda = [0.001, 1.8]$ and $\Omega_{m0} = [0.2, 0.6]$ for $H(z)$ data set; $\lambda = [0.001, 2.0]$ and $\Omega_{m0} = [0.15, 0.3]$ for $f\sigma_8$ data set; and finally, $\lambda = [0.001, 2.0]$ and $\Omega_{m0} = [0.2, 0.4]$ for the combined data set.




Fig. 3: MCMC simulations for $f(R)$ model, considering $H(z)$ data and the flat priors: $\lambda^{-1} = [0.001, 1.8]$ and $\Omega_{m0} = [0.2, 0.6]$.

We summarize our results in Table 4, and the parameter space contours obtained can be seen in figures 3, 4 and 5.

4.2 $[f\sigma_8 - H]$ diagram analyses

Recently, Linder [38] and Matsumoto et al. [44] have proposed a joint analysis of the $H(z)$ and $[f\sigma_8](z)$ cosmological observables as a way to break the degeneracy of DE and MG models, also allowing to recognize the redshift




Fig. 4: MCMC simulations for $f(R)$ model, considering $[f\sigma_8](z)$ data and the flat priors: $\lambda^{-1} = [0.001, 2.0]$ and $\Omega_{m0} = [0.15, 0.3]$.




Fig. 5: MCMC simulations for $f(R)$ model, considering joint analyses of the $H(z)$ and $[f\sigma_8](z)$ data and the flat priors: $\lambda^{-1} = [0.001, 2.0]$ and $\Omega_{m0} = [0.2, 0.4]$.

regime where the tested model most affects the growth of cosmic structures.

In Moresco et al. [45], the authors used a joint statistical analysis of both these observables and applied the method of Linder [38] to analyze individually the parameters of Λ CDM + Σm_ν and w CDM (for cosmological pa-

data	Ω_{m0}	λ^{-1}
H	$0.346^{+0.086}_{-0.073}$	1.1936 (unconstrained)
$f\sigma_8$	0.204 ± 0.023	$1.460^{+0.47}_{-0.52}$
$H + f\sigma_8$	$0.335^{+0.097}_{-0.063}$	1.761 (unconstrained)

Table 4: Results of our likelihood analyses for the cosmological parameters and their uncertainties.

rameter analyses with massive neutrinos see, e.g., [42]). The authors obtained 1σ constraints on the parameters from growth structure data at low redshifts ($z < 2$) and the last Planck (2018) data release. Of particular interest was the degeneracy between models with massive neutrinos and a modified growth parameter γ , according to this analysis the models investigated provide a better fit in relation to the flat Λ CDM model, as constrained by the Planck mission [2]. The authors then simulated data points in the $f\sigma_8 - H$ plane to distinguish between the models that better fit the data, and found that they could be distinguished with high statistical significance using the $f\sigma_8 - H$ plane. This shows that the joint analysis proposed by Linder [38] and Matsumoto et al. [44] can be contrasted with data to distinguish models that are degenerate in the cosmological parameters fits. In particular, we are interested in study how the Starobinsky $f(R)$ model (see the section 2.1) can be constrained by using $[f\sigma_8](z)$ and $H(z)$ data in the diagram proposed by the aforementioned authors.




Fig. 6: Comparison of the evolution of the ratio of observables in the Starobinsky $f(R)$ models and their Λ CDM values. Each point is the theoretical prediction for a given redshift. Different colored curves correspond to different values of the model parameter, $\lambda^{-1} = 0.3, 0.7, 1$, with $\Omega_{m0} = 0.3$

One can build an $f(R)$ modified gravity model that mimics the background evolution of Λ CDM, like the one presented in section 2.1 and the well-known Hu-Sawicki

model [32]. Both models have a similar cosmological evolution, whereas the growth of structures is quite different to the Λ CDM model, owing to the scale dependence of the solution $\delta(k, a)$ of equation (20). In the refs. [38, 44] it is shown how the $f\sigma_8 - H$ plot can help distinguish between MG and Λ CDM-type models (i.e., Λ CDM or w CDM or w_0w_a CDM models with different parameters). To perform a similar analysis for our model, we fix the scale of our perturbations at $0.1h/\text{Mpc}$, around the scale where measurements and homogeneity assumptions in the Λ CDM model are made [9, 10] and a scale where linear perturbation theory is valid in $f(R)$ theories and nonlinear effects can be disregarded [37].

In figure 6, we see the separate plot of the ratio of $H(z)$ and $[f\sigma_8]/[f\sigma_{8\Lambda\text{CDM}}]$ for our Starobinsky $f(R)$ model and their Λ CDM values, respectively. For $H/H_{\Lambda\text{CDM}}$ the curves show similarity, also requiring a high amount of precision to distinguish the $< 2\%$ absolute difference of the values, whereas for $[f\sigma_8](z)$ there is a high degeneracy in some redshift intervals between the same model with different parameters. This degeneracy makes it difficult to use data from this redshift interval to constrain the true λ parameter of the model, while also, again, requiring a higher degree of precision to distinguish between the curves in this interval. For higher values of redshift, the degeneracy between the $f\sigma_8$ curves worsens, as they need to converge to the Λ CDM value by construction. Thus the redshift interval that allows us to constrain the true parameter of the model with good precision and low degeneracy between the curves is highly dependent on our ability to measure with high precision the local $H(z)$ history and the value of the model parameter. Using the $f\sigma_8/f\sigma_{8\Lambda\text{CDM}} \times H/H_{\Lambda\text{CDM}}$ joint analysis, devised by [44] and the similar $H(z)/H_0 \times [f\sigma_8](z)$ from [38] to plot the curves for our model, we see that the degeneracy between the different values of the parameters is manifestly broken considering the whole parameters space, as shown in figure 7.




Fig. 7: Curves on the $f\sigma_8/f\sigma_{8\Lambda\text{CDM}} \times H/H_{\Lambda\text{CDM}}$ plane for different values of the λ parameter. The curves, while having a similar profile, have different curvatures while also having different start and end points. From left to right we have $\lambda^{-1} = 0.3, 0.5, 0.7, 1$.




Fig. 8: $[f\sigma_8](z) \times H(z)/H_0$ curves for the models Λ CDM and Starobinsky $f(R)$ with different values for the parameter λ .

With the degeneracy broken, the only limiting factor on distinguishing the curves in the parameter space is the precision and the confidence interval of these measurements. Figure 8 shows how the parameter space can be further distinguished, including the Λ CDM model. While the expansion history of our (and general) $f(R)$ models follows the Λ CDM model closely enough that it can't be distinguished from the standard model at the background level, when one conjoins just the $f\sigma_8$ evolution with the $H(z)$ expansion of the models, the degeneracy is clarified. Although there is still a degree of degeneracy between the curves, with the data spanning a large enough interval of $H(z)/H_0$, the difficulty in distinguishing the models lies once again in the precision of the data.

In [38] the author gives an in depth analysis of the relevant redshift range to better distinguish dark energy models using this joint analysis of observables, also giving a preview of what future surveys will be able to give precise enough measurements to further break the degeneracy between models. An extensive analysis using similar methods with other MG models can be found in [44]. In [45] a similar analysis is done considering parameter extensions of the concordance Λ CDM model, with a likelihood

approach. Here we'll follow [45] in using the $f\sigma_8 - H$ diagram to further distinguish between the parameters of our model.

4.2.1 Goodness of fit between models

We have produced a set of 5 binned data pairs, $(f\sigma_8(z), H(z))$, for equal number of redshift bins (see table 3); they were obtained by calculating the variance weighted mean in each redshift bin. We use this data to test a goodness of fit procedure, *via* a χ^2 methodology, to check which set of parameters best fits the joint data, obtained from the MCMC runs in 4.1. This analysis complements our likelihood analysis where we obtained best fit parameters from the three different sets of data described in 3. Here, we use the binned data to check which of the pair of parameters obtained from the exploration of parameter space best fits this joint data. Hereafter we'll call each of the three different best fits obtained from the MCMC run a "model"; the model with the parameters given by the run on just the $H(z)$ data, the one from just $f\sigma_8(z)$ data and the one from the run on both datasets, referring to table 4.

We devised a simple 2D minimum squared weighted deviation (also a type of χ^2), using the binned data from table 3. If, for each data point, we have the individual error on each variable, we have

- $\sigma_{f\sigma_8} \equiv [f\sigma_8](z)$ error,
- $\sigma_H \equiv H(z)$ error,

and since the data for both observables are independent measurements, i.e., they are not correlated, we may write

$$\sigma^2 \equiv \sigma_{\text{joint}}^2 = \sigma_H^2 + \sigma_{f\sigma_8}^2, \quad (30)$$

as the joint error for both variables (termed variance in quadrature).

For each model we have a predicted value $(f\sigma_8(z_i), H(z_i))$ at redshift z_i , and a data point (x_i, y_i) in the $f\sigma_8 \times H(z)$ plane. The squared residual between the model prediction $P_i \equiv (f\sigma_8(z_i), H(z_i))$ and the observed data $O_i \equiv (x_i, y_i)$ is given by

$$(P_i - O_i)^2 = \|(f\sigma_8(z_i) - x_i, H(z_i) - y_i)\|^2, \quad (31)$$

where $\|\cdot\|$ is the distance between the two data points $(f\sigma_8(z_i), H(z_i))$ and (x_i, y_i) in the $f\sigma_8 \times H$ plane.

The χ^2 is then defined as the sum of the square of the residuals averaged by the joint error for each data

$$\chi^2 \equiv \sum_i^n \frac{(P_i - O_i)^2}{\sigma_i^2}. \quad (32)$$

The reduced mean squared weighted deviation is then, for each model analyzed, the ratio between χ^2 and the number of degrees of freedom ν , $\nu = n - m = 5 - 2 = 3$ for the Starobinsky models, and $\nu = 5 - 1 = 4$ for the flat Λ CDM model

$$\chi_\nu^2 = \frac{\chi^2}{\nu}. \quad (33)$$




Fig. 9: Goodness of fit test for binned data, from table 3. The blue curve corresponds to Λ CDM and the orange one to the Starobinsky model with the best-fit parameter λ , found from the minimization of eq. (33) in the MCMC analyses.

In our case the flat Λ CDM model has one free parameter $\Omega_{\Lambda 0}$ (where $\Omega_{m 0}$ is constrained by the algebraic relation $\Omega_{\Lambda 0} + \Omega_{m 0} = 1$), one less than the Starobinsky models analysed (3 cases, each with the best adjusted parameter from a different dataset), i.e., $\Omega_{m 0}$ and λ^{-1} .

Each MCMC parameter space exploration results in a best fit of the model via the likelihood minimization performed by the MCMC method. We performed three sets of MCMC runs, each on a different data set, described in section 3; here we treat each result as a different model, and test the model against the binned data (see table 3).

We then calculate (33) for each of our models and the minimization of this parameter gives a naive estimate of the model that best fits the data in the $f\sigma_8 \times H$ plane. The results of this naive analysis allows us to clearly differentiate between models with different values of λ obtained from the MCMC run, which can be seen in table 5. The model that best fits the data according to this criteria is plotted in figure 9

From the χ^2 statistics, we see that the Λ CDM model is, from this joint data set, the best fit to the data, while the model resulting from the MCMC run on the $f\sigma_8$ data

Model	Ω_{m0}	λ^{-1}	$\chi^2_{\nu} (\times 10^{-4})$
Starobinsky + λ best-fit with $H(z)$	0.335	1.76	13.6
Starobinsky + λ best-fit with $[f\sigma_8](z)$	0.204	1.46	2.1
Starobinsky + λ best-fit with joint data	0.337	1.19	10.9
Λ CDM	0.28	–	0.8

Table 5: Goodness of fit test, measured by χ^2_{ν} through the analysis of the binned data in table 3, to compare the behavior of the Λ CDM and 3 Starobinsky models (each with a best-fitted λ parameter provided by the MCMC runs considering 3 data sets).

only gives the least χ^2 from the models explored using the MCMC method. While one could say that Λ CDM fits the joint data better than any version of the Starobinsky models, i.e., models with the parameter λ^{-1} adjusted by MCMC analyses, some observation on the methods and data are in order. First, it's interesting to note that the $f\sigma_8$ data alone gives a better fit to the joint data than the result from the MCMC run on data from both $[f\sigma_8](z)$ and $H(z)$ data. This could be the case for diverse reasons; the most reasonable one being that the $H(z)$ is actually worsening the results from the $[f\sigma_8](z)$ data when analysed together. As expected theoretically, background quantities don't constrain modified gravity models suitably; the MCMC run on the $H(z)$ data alone, found in 3, was not able to constraint the free parameters analysing the $[f\sigma_8](z)$ data.

Secondly, the binned data presented in table 3, built from the $H(z)$ and $[f\sigma_8](z)$ datasets, gives us 5 data points for goodness of fit analysis. As a test on whether the binning chose affects the χ^2 statistics, we considered other binned sets and found similar results. The advantage of this approach is that it allows to break the degeneracy between models with dissimilar λ parameter, but due to the large errors at high- z the analysis is not as accurate as desirable to clearly discriminate between the Λ CDM and the Starobinsky models. In fact, more data points and tighter error constraints are needed to improve the significance of the goodness of fit. This means not only precise measurements on these observables, in particular at high- z , but also data in more redshift intervals. From our MCMC analysis, we observe that the data on $[f\sigma_8](z)$ was able to better constraint the parameters of the $f(R)$ model, thus results from upcoming surveys such as the DESI, SKA, and EUCLID telescopes [63] will be able to probe the cosmic structures growth with precision to greatly improve the significance of this kind of analysis.

It is important to note that the conjoined analysis found in table 5, even if with low statistical significance, was able to differentiate between different models inside the same $f(R)$ theory, which we saw had high degeneracy in the $H(z)$ observables, and some degeneracy in $[f\sigma_8](z)$. Thus the proposal of [38, 44] of breaking this degeneracy between models using the conjoined data is effectual.

5 Conclusions and Final Remarks

In this paper we have used the cosmic expansion and the structure growth data to constrain the free parameters of a relevant $f(R)$ gravity model, namely the Starobinsky model. We've used the bayesian based Monte Carlo Markov Chain method to explore the parameter space of this model from both sets of data individually and using a joint likelihood analysis. Furthermore, we also used the recent method proposed by Linder [38] and Matsumoto et al. [44] to complement the result of the MCMC statistical analysis. This novel approach, based on a joint data analyses from H and $f\sigma_8$, has proven able to give further distinction between different models which were shown to have degeneracy in the theoretical predictions of the $H(z)$ and $[f\sigma_8](z)$ observables in the range of the free parameters of the $f(R)$ model.

At the end, our results show that the $f(R)$ Starobinsky model, although it has a worse fit to the structure growth data than the Λ CDM model, cannot be discarded given the current data available in the literature. More –and more precise– data are needed in order to determine the preferred model with good statistical significance. Therefore, the Starobinsky model cannot be discarded as a possible alternative to the Λ CDM model.

A possible continuation of this work would be to use this joint analysis to compare diverse cosmologically motivated $f(R)$ models, such a the one in [32] and the one in this paper, and break the degeneracy between different $f(R)$ theories. The results of this paper show that this method is promising for further distinction between modified gravity models that are degenerate in certain cosmological probes. The extension of this analysis is left for future work.

Acknowledgements

PB thanks CAPES for a PhD fellowship; AB thanks a CNPq fellowship. MC acknowledges the financial support from Programa de Capacitação Institucional (PCI/MCTI), that subsidized part of this work. The authors are grateful for the computational facilities of the Observatório Nacional.

References

1. I. Achitouv, C. Blake, P. Carter, J. Koda, and F. Beutler. Consistency of the growth rate in different environments with the 6-degree field galaxy survey: Measurement of the void-galaxy and galaxy-galaxy correlation functions. *Physical Review D*, 95(8), Apr 2017.
2. P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al. Planck2015 results. *Astronomy & Astrophysics*, 594:A13, Sep 2016.
3. P. A. R. Ade et al. Planck 2015 results. XIV. Dark energy and modified gravity. *Astron. Astrophys.*, 594:A14, 2016.
4. N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. [Erratum: *Astron. Astrophys.* 652, C4 (2021)].
5. Shadab Alam, Marie Aubert, Santiago Avila, Christophe Balland, Julian E. Bautista, Matthew A. Bershady, Dmitry Bizyaev, Michael R. Blanton, Adam S. Bolton, Jo Bovy, and et al. Completed sdss-iv extended baryon oscillation spectroscopic survey: Cosmological implications from two decades of spectroscopic surveys at the apache point observatory. *Physical Review D*, 103(8), Apr 2021.
6. Fotios K. Anagnostopoulos and Spyros Basilakos. Constraining the dark energy models with $h(z)$ data: An approach independent of h_0 . *Physical Review D*, 97(6), Mar 2018.
7. Fotios K. Anagnostopoulos, David Benisty, Spyros Basilakos, and Eduardo I. Guendelman. Dark energy and dark matter unification from dynamical space time: observational constraints and cosmological implications. *JCAP*, 06:003, 2019.
8. Marie Aubert, Marie-Claude Cousinou, Stéphanie Escoffier, Adam J. Hawken, Seshadri Nadathur, Shadab Alam, Julian Bautista, Etienne Burtin, Arnaud de Mattia, Héctor Gil-Marín, Jiamin Hou, Eric Jullo, Richard Neveux, Graziano Rossi, Alex Smith, Amélie Tamone, and Mariana Vargas Magaña. The completed sdss-iv extended baryon oscillation spectroscopic survey: Growth rate of structure measurement from cosmic voids, 2020.
9. F. Avila, A. Bernui, E. de Carvalho, and C. P. Novaes. The growth rate of cosmic structures in the local Universe with the ALFALFA survey. *Mon. Not. Roy. Astron. Soc.*, 505(3):3404–3413, 2021.
10. F. Avila, A. Bernui, R. C. Nunes, E. de Carvalho, and C. P. Novaes. The homogeneity scale and the growth rate of cosmic structures. *Monthly Notices of the Royal Astronomical Society*, 2021.
11. F. Avila, C. P. Novaes, A. Bernui, E. de Carvalho, and J. P. Nogueira-Cavalcante. The angular scale of homogeneity in the Local Universe with the SDSS blue galaxies. *Mon. Not. Roy. Astron. Soc.*, 488(1):1481–1487, 2019.
12. Spyros Basilakos. The λ cdm growth rate of structure revisited. *International Journal of Modern Physics D*, 21(07):1250064, 2012.
13. Spyros Basilakos and Fotios K. Anagnostopoulos. Growth index of matter perturbations in the light of dark energy survey. *The European Physical Journal C*, 80(3), Mar 2020.
14. Ronaldo C. Batista. Impact of dark energy perturbations on the growth index. *Phys. Rev. D*, 89(12):123508, 2014.
15. H.Ā. Benaoum, Weiqiang Yang, Supriya Pan, and Eleonora Di Valentino. Modified Emergent Dark Energy and its Astronomical Constraints. *arXiv e-prints*, page arXiv:2008.09098, August 2020.
16. David Benisty, Eduardo Guendelman, and Zbigniew Haba. Unification of dark energy and dark matter from diffusive cosmology. *Phys. Rev. D*, 99(12):123521, 2019. [Erratum: *Phys. Rev. D* 101, 049901 (2020)].
17. David Benisty and Eduardo I. Guendelman. Unified dark energy and dark matter from dynamical spacetime. *Phys. Rev. D*, 98(2):023506, 2018.
18. Florian Beutler, Chris Blake, Matthew Colless, D. Heath Jones, Lister Staveley-Smith, Gregory B. Poole, Lachlan Campbell, Quentin Parker, Will Saunders, and Fred Watson. The 6df galaxy survey: $z \approx 0$ measurements of the growth rate and σ_8 . *Monthly Notices of the Royal Astronomical Society*, 423(4):3430–3444, Jun 2012.
19. Chris Blake, Sarah Brough, Matthew Colless, Carlos Contreras, Warrick Couch, Scott Croom, Darren Croton, Tamara M. Davis, Michael J. Drinkwater, Karl Forster, and et al. The wigglez dark energy survey: joint measurements of the expansion and growth history at $z < 1$. *Monthly Notices of the Royal Astronomical Society*, 425(1):405–414, Jul 2012.
20. Alexander Bonilla, Suresh Kumar, and Rafael C. Nunes. Measurements of h_0 and reconstruction of the dark energy properties from a model-independent joint analysis. *The European Physical Journal C*, 81(2), Feb 2021.
21. Vincenzo F Cardone, Stefano Camera, and Antonaldo Diaferio. An updated analysis of two classes of $f(r)$ theories of gravity. *Journal of Cosmology and Astroparticle Physics*, 2012(02):030–030, Feb 2012.
22. Yow-Chun Chen, Chao-Qiang Geng, Chung-Chi Lee, and Hongwei Yu. Matter Power Spectra in Viable $f(R)$ Gravity Models with Dynamical Background. *The European Physical Journal C*, 79(2):93, jan 2019.
23. Chia-Hsun Chuang, Francisco Prada, Marcos Pellejero-Ibanez, Florian Beutler, Antonio J. Cuesta, Daniel J. Eisenstein, Stéphanie Escoffier, Shirley Ho, Francisco-Shu Kitaura, Jean-Paul Kneib, and et al. The clustering of

- galaxies in the sdss-iii baryon oscillation spectroscopic survey: single-probe measurements from cmass anisotropic galaxy clustering. *Monthly Notices of the Royal Astronomical Society*, 461(4):3781–3793, Jun 2016.
24. Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis. Modified Gravity and Cosmology. jun 2011.
 25. Zahra Davari and Sohrab Rahvar. Mog cosmology without dark matter and the cosmological constant. *Monthly Notices of the Royal Astronomical Society*, 507(3):3387–3399, Aug 2021.
 26. E. de Carvalho, A. Bernui, G. C. Carvalho, C. P. Novaes, and H. S. Xavier. Angular Baryon Acoustic Oscillation measure at $z = 2.225$ from the SDSS quasar survey. *JCAP*, 04:064, 2018.
 27. E. de Carvalho, A. Bernui, H. S. Xavier, and C. P. Novaes. Baryon acoustic oscillations signature in the three-point angular correlation function from the SDSS-DR12 quasar survey. *Mon. Not. Roy. Astron. Soc.*, 492(3):4469–4476, 2020.
 28. Antonio de Felice and Shinji Tsujikawa. $f(R)$ theories. Technical report, 2010.
 29. Martin Feix, Adi Nusser, and Enzo Branchini. Growth rate of cosmological perturbations at $z \approx 0.1$ from a new observational test. *Physical Review Letters*, 115(1), Jun 2015.
 30. Daniel Foreman-Mackey, David W. Hogg, Dustin Lang, and Jonathan Goodman. emcee: The mcmc hammer. *Publications of the Astronomical Society of the Pacific*, 125(925):306–312, Mar 2013.
 31. A. J. S. Hamilton. Measuring Omega and the Real Correlation Function from the Redshift Correlation Function. *apjl*, 385:L5, January 1992.
 32. Wayne Hu and Ignacy Sawicki. Models of $f(R)$ Cosmic Acceleration that Evade Solar-System Tests. may 2007.
 33. Dragan Huterer, Daniel L. Shafer, Daniel M. Scolnic, and Fabian Schmidt. Testing λ cdm at the lowest redshifts with sn ia and galaxy velocities. *Journal of Cosmology and Astroparticle Physics*, 2017(05):015–015, May 2017.
 34. Raul Jimenez and Abraham Loeb. Constraining cosmological parameters based on relative galaxy ages. *The Astrophysical Journal*, 573(1):37–42, jul 2002.
 35. Nick Kaiser. Clustering in real space and in redshift space. *mnras*, 227:1–21, July 1987.
 36. Lavrentios Kazantzidis and Leandros Perivolaropoulos. Evolution of the $f\sigma_8$ tension with the Planck15/ Λ CDM determination and implications for modified gravity theories. *Phys. Rev. D*, 97(10):103503, 2018.
 37. Tomi Koivisto. Matter power spectrum in $f(R)$ gravity. *Physical Review D - Particles, Fields, Gravitation and Cosmology*, 73(8):1–5, 2006.
 38. Eric V. Linder. Cosmic growth and expansion conjoined. *Astroparticle Physics*, 86:41–45, 2017.
 39. Eric V. Linder. Limited modified gravity. *Journal of Cosmology and Astroparticle Physics*, 2020(10):042–042, Oct 2020.
 40. Eric V. Linder and Robert N. Cahn. Parameterized beyond-einstein growth. *Astroparticle Physics*, 28(4):481 – 488, 2007.
 41. Gabriela A. Marques and Armando Bernui. Tomographic analyses of the CMB lensing and galaxy clustering to probe the linear structure growth. *JCAP*, 05:052, 2020.
 42. Gabriela A. Marques, Jia Liu, José Manuel Zorrilla Matilla, Zoltán Haiman, Armando Bernui, and Camila P. Novaes. Constraining neutrino mass with weak lensing Minkowski Functionals. *JCAP*, 06:019, 2019.
 43. Matteo Martinelli, Erminia Calabrese, Francesco De Bernardis, Alessandro Melchiorri, Luca Pagano, and Roberto Scaramella. Constraining modified gravitational theories by weak lensing with euclid. *Physical Review D*, 83(2), Jan 2011.
 44. Jiro Matsumoto, Teppei Okumura, and Misao Sasaki. New measures to test modified gravity cosmologies. *Journal of Cosmology and Astroparticle Physics*, 2020(7), jul 2020.
 45. M. Moresco and F. Marulli. Cosmological constraints from a joint analysis of cosmic growth and expansion. *Monthly Notices of the Royal Astronomical Society: Letters*, 471(1):L82–L86, 2017.
 46. Seshadri Nadathur, Paul M. Carter, Will J. Percival, Hans A. Winther, and Julian E. Bautista. Beyond bao: Improving cosmological constraints from boss data with measurement of the void-galaxy cross-correlation. *Physical Review D*, 100(2), Jul 2019.
 47. Rafael C. Nunes. Structure formation in $f(T)$ gravity and a solution for H_0 tension. *JCAP*, 05:052, 2018.
 48. Rafael C. Nunes, Edésio M. Barboza, Jr., Everton M. C. Abreu, and Jorge Ananias Neto. Probing the cosmological viability of non-gaussian statistics. *JCAP*, 08:051, 2016.
 49. Rafael C. Nunes and Armando Bernui. BAO signatures in the 2-point angular correlations and the Hubble tension. *Eur. Phys. J. C*, 80(11):1025, 2020.
 50. Teppei Okumura, Chiaki Hikage, Tomonori Totani, Motonari Tonegawa, Hiroyuki Okada, Karl Glazebrook, Chris Blake, Pedro G. Ferreira, Surhud More, Atsushi Taruya, and et al. The subaru fmos galaxy redshift survey (fast-sound). iv. new constraint on gravity theory from redshift space distortions at $z \approx 1.4$. *Publications of the Astronomical Society of Japan*, 68(3):38, Apr 2016.
 51. Louis Perenon, Julien Bel, Roy Maartens, and Alvaro de la Cruz-Dombriz. Optimising growth of structure constraints on modified gravity. *JCAP*, 2019(6):020, June 2019.

52. Louis Perenon, Stéphane Ilíc, Roy Maartens, and Alvaro de la Cruz-Dombriz. Improvements in cosmological constraints from breaking growth degeneracy. *Astron. Astrophys.*, 642:A116, 2020.
53. Louis Perenon, Federico Piazza, Christian Marinoni, and Lam Hui. Phenomenology of dark energy: general features of large-scale perturbations. *JCAP*, 11:029, 2015.
54. A. Pezzotta, S. de la Torre, J. Bel, B. R. Granett, L. Guzzo, J. A. Peacock, B. Garilli, M. Scodreggio, M. Bolzonella, U. Abbas, and et al. The vimos public extragalactic redshift survey (vipers). *Astronomy & Astrophysics*, 604:A33, Jul 2017.
55. Adam G. Riess, Stefano Casertano, Wenlong Yuan, Lucas M. Macri, and Dan Scolnic. Large Magellanic Cloud Cepheid Standards Provide a 1% Foundation for the Determination of the Hubble Constant and Stronger Evidence for Physics beyond Λ CDM. *Astrophys. J.*, 876(1):85, 2019.
56. Thomas P. Sotiriou and Valerio Faraoni. $f(r)$ theories of gravity. *Reviews of Modern Physics*, 82(1):451–497, Mar 2010.
57. A. A. Starobinsky. Disappearing cosmological constant in $f(r)$ gravity. *JETP Letters*, 86(3):157–163, Oct 2007.
58. Michael A. Strauss and Jeffrey A. Willick. The density and peculiar velocity fields of nearby galaxies. *Physics Reports*, 261(5):271 – 431, 1995.
59. Ariel G. Sánchez, Francesco Montesano, Eyal A. Kazin, Eric Aubourg, Florian Beutler, Jon Brinkmann, Joel R. Brownstein, Antonio J. Cuesta, Kyle S. Dawson, Daniel J. Eisenstein, and et al. The clustering of galaxies in the sdss-iii baryon oscillation spectroscopic survey: cosmological implications of the full shape of the clustering wedges in the data release 10 and 11 galaxy samples. *Monthly Notices of the Royal Astronomical Society*, 440(3):2692–2713, Apr 2014.
60. Shinji Tsujikawa. Matter density perturbations and effective gravitational constant in modified gravity models of dark energy. *Physical Review D*, 76(2), Jul 2007.
61. Stephen J. Turnbull, Michael J. Hudson, Hume A. Feldman, Malcolm Hicken, Robert P. Kirshner, and Richard Watkins. Cosmic flows in the nearby universe from type ia supernovae. *Monthly Notices of the Royal Astronomical Society*, 420(1):447–454, Dec 2011.
62. A. M. Velasquez-Toribio and Júlio C. Fabris. The growth factor parametrization versus numerical solutions in flat and non-flat dark energy models. *The European Physical Journal C*, 80(12), Dec 2020.
63. Jan-Albert Viljoen, José Fonseca, and Roy Maartens. Constraining the growth rate by combining multiple future surveys. *Journal of Cosmology and Astroparticle Physics*, 2020(09):054–054, Sep 2020.
64. Michael J. Wilson. Geometric and growth rate tests of general relativity with recovered linear cosmological perturbations, 2016.
65. Yingjie Yang and Yungui Gong. The evidence of cosmic acceleration and observational constraints. *Journal of Cosmology and Astroparticle Physics*, 2020(06):059–059, jun 2020.
66. Gong-Bo Zhao, Yuting Wang, Shun Saito, Héctor Gil-Marín, Will J Percival, Dandan Wang, Chia-Hsun Chuang, Rossana Ruggeri, Eva-Maria Mueller, Fangzhou Zhu, and et al. The clustering of the sdss-iv extended baryon oscillation spectroscopic survey dr14 quasar sample: a tomographic measurement of cosmic structure growth and expansion rate based on optimal redshift weights. *Monthly Notices of the Royal Astronomical Society*, 482(3):3497–3513, Oct 2018.