Extended phase space thermodynamics of black holes: A study in Einstein's gravity and beyond

Krishnakanta Bhattacharya*

IUCAA, Post Bag 4, Pune University Campus, Ganeshkhind, Pune 411 007, Maharashtra, India.

December 3, 2021

Abstract

In the extended phase space approach, one can define thermodynamic pressure and volume that gives rise to the van der Waals type phase transition for black holes. For Einstein's GR, the expressions of these quantities are unanimously accepted. Of late, the van der Waals phase transition in black holes has been found in modified theories of gravity as well, such as the f(R)gravity and the scalar-tensor gravity. However, in the case of these modified theories of gravity, the expression of pressure (and, hence, volume) is not uniquely determined. In addition, for these modified theories, the extended phase space thermodynamics has not been studied extensively, especially in a covariant way. Since both the scalar-tensor and the f(R) gravity can be discussed in the two conformally connected frames (the Jordan and the Einstein frame respectively), the arbitrariness in the expression of pressure, will act upon the equivalence of the thermodynamic parameters in the two frames. We highlight these issues in the paper. In addition, in Einstein's gravity (GR), we obtain a general expression of the equilibrium state version of first law and the Smarr formula from the Einstein's equation for a general static and spherically symmetric (SSS) metric. Unlike the existing formalisms in literature which defines thermodynamic potential in order to express the first law, here we directly obtain the first law (and the Smarr formula) in GR in terms of the parameters present in the metric (such as mass, charge *etc.*). This study also shows how the extended phase space is formulated (by considering the cosmological constant as variable) and, also justifies why the cosmological constant plays the role of thermodynamic pressure in GR in extended phase space.

1 Introduction

The connection between general relativity and thermodynamics has been found almost fifty years ago from the works of Bekenstein and Hawking [1–3]. Till date, several interesting results are coming up with the course of time and further consolidate the connection between the two. Black hole (BH) phase transition is one of the most interesting results in black hole mechanics which was obtained long ago by Davies [4]. It was then identified that the critical point of the phase transition as one where the heat capacity shows an infinite discontinuity. The presence of phase transition and the critical phenomena was later found, more prominently for the AdS black holes. Firstly it was found by Hawking and Page [5], which is known as the Hawking-Page phase transition. Later, the P - V

^{*}krishnakanta@iucaa.in

criticality of conventional thermodynamics was found in BH mechanics, where the thermodynamic pressure (P) was defined in two different ways. It was first noticed in the non-extended phase space approach [6,7] where it was shown that the black hole undergoes a transition, which is similar to the van der Waals-Maxwell liquid-gas system, where the pressure was identified as the inverse of Hawking temperature. This approach was not accepted much due to the lack of proper canonical definition of pressure and volume. Later, in the extended phase space approach, the cosmological constant of the AdS black hole was treated as a variable and was found that the P - V criticality exists in the black hole thermodynamic system of the AdS black holes and the pressure (P) was identified in terms of the cosmological constant [8–14]. This extended phase space formalism was widely accepted and P - V criticality was later extensively studied with the identification of cosmological constant as the thermodynamic pressure (for general analysis in this topic, see [15–20]).

The extended phase space formalism, as discussed above, was mostly studied in Einstein's gravity (GR). Although, till date, GR is considered as the most viable theory to describe gravity, one cannot disregard the alternative theories of gravity for several obvious reasons. In addition, thermodynamic features of general relativity has been found in alternative gravity theories as well [21-28], which suggests that the thermodynamic property of gravity ranges well-beyond Einstein's GR. Interestingly, the P-V criticality has been found in scalar-tensor (ST) / Brans-Dicke and f(R) gravity as well [29–32]. However, there seems to have a disparity in the expression of the thermodynamic pressure. It is well known, in general, f(R) gravity can be studied equivalently as the scalar-tensor gravity with the identification $\phi = f'(R)$, where ϕ is the scalar field which is non-minimally coupled with the Ricci-scalar in the ST gravity and $f'(R) = \partial f(R)/\partial R$ in the f(R) gravity. Now, the expression of pressure has been considered as different in different papers. For example, in [29], the thermodynamic pressure is identified in terms of the cosmological constant multiplied with the factor f'(R) (or the conformal factor) and, on the other hand, in [30, 31] the cosmological constant itself is identified as the thermodynamic pressure in f(R) gravity and in Brans-Dicke (BD) gravity. Apart from this difference in the expression of thermodynamic pressure, there is another important aspect. As we have mentioned, both f(R) and BD gravity can be studied as a special case of scalar-tensor gravity and can be described in two conformally connected frames *i.e.*, Jordan and Einsten frame. In earlier works the equivalence of thermodynamic parameters, in the two conformally connected frames, has been established by us and other groups [19, 33-36]. But, those analysis were done in the non-extended phase space approach as the variation of the cosmological constant was not accounted. Therefore, the question arises how the thermodynamic parameters, defined in the extended phase space approach. are related in the two conformally connected frames. Furthermore, the question arises: whether this arbitrariness in the definition of pressure influences the equivalence of the thermodynamic parameters in the two frames.

In this work, we address these issues. Firstly, in Einstein gravity, we derive a novel approach and obtain the thermodynamic first law and the Smarr formula [37] from the Einstein's equation. The idea is, if a static and spherically symmetric solution of Einstein's equation depends upon several parameters (such as mass, charge, AdS radius *etc.*), using the presented approach, one can obtain the equilibrium state version of the first law and the Smarr formula in a general way. Thereby, we show that if the cosmological constant of the AdS black hole is considered a variable, it looks like thermodynamic pressure and its conjugate quantity appears to be the volume of the sphere, the radius of which is the horizon radius 1 . Thus, we show the formulation of extended phase space in GR. Later

¹The results (first law and the Smarr formula) obtained in this approach is nothing new, but the approach is new as far my knowledge. That is why, I call it as a novel approach instead of saying it as new findings.

we move on to discuss extended phase space thermodynamics in the conformally connected frames. Although there are debates regarding the physical equivalence of the two conformally connected frames [19, 28, 33–36, 38–58], it has been found that the thermodynamic parameters (of non-extended phase space) are equivalent in the two conformally connected frames [19, 33-36]. Therefore, following the extended phase space approach, here we study the connection of thermodynamic parameters in the two frames (especially pressure and volume). Firstly, we introduce the conformal frame of GR, which corresponds to a particular case of Jordan frame of ST gravity. Then, following Iyer-Wald formalism [59,60], we define thermodynamic quantities and obtain the first law considering cosmological constant as a variable. We then compare the thermodynamic parameters of the conformal frame (or the Jordan frame) with the thermodynamic parameters of GR (corresponding to the Einstein frame). We do this because, the thermodynamic parameters in GR is well defined. Therefore, it is simpler to compare with its conformal frame. After this, we generalize this result for the general scalar-tensor gravity in its two frames. Our analysis, not only proves the first law in the two frames, it provides the expressions of all thermodynamic parameters in a covariant way, especially thermodynamic volume which has not been shown for alternative theories of gravity. Also, we discuss that the arbitrariness in the expression of pressure indeed influences in the equivalence of the thermodynamic parameters.

The discussion of the paper is presented as follows. In the following section, we discuss our formalism to obtain first law and the Smarr formula in a general way for GR from the Einstein's equation. Here we also show how the identification of pressure in terms of cosmological constant gets justified. In section 3, we adopt Wald's formalism and compare the thermodynamic parameters of Einstein's gravity with its conformal frame. Finally, we generalize these results for ST gravity in section 4. We provide conclusion and outlook of our analysis in section 5.

Units and notations: In this article, we have adopted geometrized units and have set c, \hbar and G as unity. Quantities with a bar overhead (such as \bar{A}) correspond to the of Einstein's GR. Quantities with a tilde overhead (such as \tilde{A}) correspond to the Einstein frame of ST gravity. For Jordan frame, or for the conformal frame of Einstein's GR, we use plain notations (such as A).

2 Thermodynamic first law and Smarr-like formula from the spacetime metric in GR: A general analysis in spherically symmetric spacetime

It is well-known that Einstein's equation manifests itself as a thermodynamic law when it is projected on the black hole horizon [24–27, 61–63]. Here we have briefly mentioned two existing formalisms in literature to obtain thermodynamic first law from Einstein's equation for static and spherically symmetric spacetime. Then we discuss our general formalism of obtaining first law and the Smarr formula from the Einstein's equation. We show that, in the existing formalisms, the first law is defined in terms of thermodynamic potential. In our case, the first law appears as the change of the BH entropy in terms of the "virtual" change of the metric parameters (mass, charge etc.).

A general static and spherically symmetric metric is given as

$$d\bar{s}^{2} = -\bar{f}(\bar{r})d\bar{t}^{2} + \frac{d\bar{r}^{2}}{\bar{f}(\bar{r})} + \bar{r}^{2}(d\bar{\theta}^{2} + \sin^{2}\bar{\theta}d\bar{\phi}^{2}) , \qquad (1)$$

with $\bar{f}(\bar{r})$ being the general function of \bar{r} . This metric satisfies the radial Einstein's equation ($\bar{G}_{\bar{r}}^{\bar{r}}$ =

 $8\pi \bar{T}_{\bar{r}}^{\bar{r}}$), which is provided as

$$\frac{-1+\bar{f}(\bar{r})}{\bar{r}^2} + \frac{f'(\bar{r})}{\bar{r}} = 8\pi \bar{T}_{\bar{r}}^{\bar{r}} .$$
⁽²⁾

There are two routes to obtain the first law from the above Einstein's equation as described in literature. On the first route [24,25], the Einstein's equation (2) is cast on the horizon $\bar{r} = \bar{r}_H$ (where \bar{r}_H is obtained using $\bar{f}(\bar{r}_H) = 0$). Then both sides of (2) are multiplied with $(\bar{r}_H^2 d\bar{r}_H)/2$ to obtain

$$\bar{\Gamma}d\bar{S} - d\bar{E} = \bar{P}_{(r)}d\bar{V} .$$
(3)

The above equation is identified as the first law of a black hole, the horizon of which has undergone a virtual displacement from \bar{r}_H to $\bar{r}_H + d\bar{r}_H$. Here $\bar{P}_{(r)}$ is known as the radial pressure and \bar{V} is the geometric volume of a sphere of radius \bar{r}_H . The thermodynamic quantities are identified as

$$\bar{P}_{(r)} = \bar{T}_{\bar{r}}^{\bar{r}} \Big|_{\bar{r}_H} , \quad \bar{V} = \frac{4\pi}{3} \bar{r}_H^3 , \quad \bar{T} = \frac{f'(\bar{r}_H)}{4\pi} , \quad \bar{S} = \pi \bar{r}_H^2 , \quad \bar{E} = \frac{\bar{r}_H}{2} .$$
(4)

There is another route [62] to obtain thermodynamic law from the Einstein's equation (2) in the literature, which is as follows. The Eq. (2) can be written in terms of a generalized equation of state

$$\bar{P}_{(r)} = B(\bar{r}_H) + C(\bar{r}_H)\bar{T}$$
, (5)

where the expression of $\bar{P}_{(r)}$ and \bar{T} are given in (4). Therefore, taking the variation on both sides and multiplying it with \bar{V} leads to the thermodynamic relation

$$\bar{V}\delta\bar{P}_{(r)} = \bar{S}\delta\bar{T} + \delta\bar{G} , \qquad (6)$$

where,

$$\bar{G} = \int^{\bar{r}_{H}} d\bar{x} \bar{V}(\bar{x}) B'(\bar{x}) + \bar{T} \int^{\bar{r}_{H}} d\bar{x} \bar{V}(x) C'(\bar{x}) = \bar{P}_{(r)} \bar{V} - \bar{T} \bar{S} - \int^{\bar{r}_{H}} d\bar{x} \bar{V}'(\bar{x}) B(\bar{x}) , \bar{S} = \int^{\bar{r}_{H}} d\bar{x} \bar{V}'(\bar{x}) C(\bar{x}) .$$
(7)

In the present case (Einstein's gravity), $B(\bar{r}_H) = -(8\pi \bar{r}_H^2)^{-1}$ and $C(\bar{r}_H) = 1/(2\bar{r}_H)$, implying

$$\bar{G} = \frac{\bar{r}_H}{3} (1 - \pi \bar{r}_H \bar{T}) , \qquad (8)$$

and \bar{S} is given by quarter of the horizon area as given in (4). Thus, the thermodynamic quantities are identified as the same in the two different routes. The identified internal energy (\bar{E}) of the first route and the Gibbs function (\bar{G}) as defined in the second route are connected to each other by the Legendre transformation $\bar{E} = \bar{G} + \bar{T}\bar{S} - \bar{P}_{(r)}\bar{V}$.

The second route [62] is adopted (as it is mentioned there) in order to remove the ambiguity to distinguish the "heat" and "work" term of the first route [24,25] as both \bar{S} and \bar{V} are functions of \bar{r}_H only. However, in both the routes, the thermodynamic potentials (such as \bar{E} defined in the first route and \bar{G} in the second) are needed to be defined in order to match the derived relations with 1st law of

conventional thermodynamics. Note that the defined thermodynamic potentials, as defined in the two routes mentioned above, are not unique. For instance, for the Schwarzschild black hole, the internal energy (\bar{E}) , which is defined as half of the horizon radius, coincides with the mass of the black hole (\bar{M}) ; while for Reissner Nördstrom (RN) black hole, \bar{E} is not the same as \bar{M} . Furthermore, according to the no hair theorem, the parameters appearing in the black hole metric are more fundamental than the thermodynamic potentials, which are defined while obtaining the first law following these two routes. Moreover, as we have noted, the radial pressure is identified as $\bar{P}_{(r)} = \bar{T}_{\bar{r}}^{\bar{r}}(\bar{r}_H)$ in these methods. For a charged Reissner Nördstrom (RN) black hole in AdS spacetime, the radial pressure will be identified as $\bar{P}_{(r)} = \bar{T}_{\bar{r}}^{\bar{r}} = -\bar{Q}^2/(8\pi\bar{r}^4) - \bar{\Lambda}/8\pi$. However, for RN black hole in AdS space, the thermodynamic pressure is identified as $\bar{P} = -\bar{\Lambda}/8\pi$ which give rise to the $\bar{P} - \bar{V}$ criticality [8–14] and is consistent with the thermodynamics in the extended phase space. In other words, the radial pressure does not play the role of thermodynamic pressure for the $\bar{P} - \bar{V}$ criticality in the extended phase space.

Therefore, the natural question arises whether one can generally obtain the thermodynamic law (and the Smarr formula) of black holes in terms of the parameters (\overline{M} , \overline{Q} etc.) instead of defining any thermodynamic potential. More precisely, whether one can show the change of horizon area in terms of the change in parameters only. The formulation we developed in the following, requires Einstein's equation to be solved. However, the final form of the first law and the Smarr formula, requires only the spacetime metric and nothing else.

With the identification $\overline{T}_{\overline{r}}^{\overline{r}} = -\epsilon(\overline{r})/(8\pi)$, the Einstein's equation (2) leads to the general solution [64]

$$\bar{f}(\bar{r}) = 1 - \frac{\bar{r}_H}{\bar{r}} - \frac{1}{\bar{r}} \int_{\bar{r}_H}^{\bar{r}} \epsilon(x) x^2 dx = 1 - \frac{1}{\bar{r}} \int^{\bar{r}} \epsilon(x) x^2 dx - \frac{\bar{r}_H}{\bar{r}} \left[1 - \frac{1}{\bar{r}_H} \int^{\bar{r}_H} \epsilon(x) x^2 dx \right], \quad (9)$$

which implies

$$\bar{f}(\bar{r}) = g(\bar{r}) - \frac{C}{\bar{r}} , \qquad (10)$$

where C is the integration constant and

$$g(y) = 1 - \frac{1}{y} \int^y \epsilon(x) x^2 dx . \qquad (11)$$

The integration constant C can be identified as the mass of the black holes for the following reasons. The Birkhoff's theorem [65] says that the exterior solution of a static and spherically symmetric metric without any source is given by the Schwarzschild metric. From (11), we obtain $g(\bar{r}) = 1$ and $\bar{f}(\bar{r}) = 1 - C/\bar{r}$ for $\epsilon(\bar{r}) = 0$. Thus, we obtain the value of the integration constant as

$$C = 2M = \bar{r}_H g(\bar{r}_H) , \qquad (12)$$

and, thereby, we obtain

$$\bar{f}(\bar{r}) = g(\bar{r}) - \frac{2M}{\bar{r}} . \tag{13}$$

As we have mentioned earlier, for the spherically symmetric spacetime of the form (1), the Einstein's equation is first order differential equation. Therefore, in the solution (13), we have one

integration constant, which is defined as the mass parameter of BH. However, in the spacetime metric, we can have other parameters (such as \bar{Q}) coming from the component of energy-momentum tensor ϵ (such as, for RN black hole, $\epsilon(\bar{r}) = \bar{Q}^2/\bar{r}^4$). Thus, we can write $g(\bar{r}) \equiv g(\bar{r}, \bar{\mu}^i)$, where $\bar{\mu}^i$ are the parameters (other than the mass) that appear in the spacetime metric and the number of parameters are kept arbitrary. In addition, we express $\bar{f}(\bar{r})$ as $\bar{f}(\bar{r}, \bar{M}, \bar{\mu}^i)$.

Obtaining the first law

Let us now consider a small virtual change of parameters, leading to a small (virtual) change in the horizon radius $\delta \bar{r}_H^2$. Therefore, from (12), we obtain

$$2\delta\bar{M} = \left[g(\bar{r}_H, \bar{\mu}^i) + \frac{\partial g(\bar{r}_H, \bar{\mu}^i)}{\partial\bar{r}_H}\bar{r}_H\right]\delta\bar{r}_H + \bar{r}_H \frac{\partial g(\bar{r}_H, \bar{\mu}^i)}{\partial\bar{\mu}^i}\delta\bar{\mu}^i \tag{14}$$

Furthermore, from (10) we obtain

$$\bar{f}'(\bar{r}_H) = \frac{\partial \bar{f}(\bar{r})}{\partial \bar{r}}\Big|_{\bar{r}=\bar{r}_H} = \frac{\partial g(\bar{r}_H, \bar{\mu}^i)}{\partial \bar{r}_H} + \frac{g(\bar{r}_H, \bar{\mu}^i)}{\bar{r}_H} , \qquad (15)$$

In addition, from (13), we obtain

$$\frac{\partial g(\bar{r}_H, \bar{\mu}^i)}{\partial \bar{\mu}^i} = \frac{\partial \bar{f}(\bar{r}, \bar{M}, \bar{\mu}^i)}{\partial \bar{\mu}^i} \Big|_{\bar{r}=\bar{r}_H} \,. \tag{16}$$

Substituting (15) and (16) in (14), we finally obtain

$$\bar{T}\delta\bar{S} = \delta\bar{M} - \frac{\bar{r}_H}{2} \frac{\partial\bar{f}(\bar{r},\bar{M},\bar{\mu}^i)}{\partial\bar{\mu}^i}\Big|_{\bar{r}=\bar{r}_H} \delta\bar{\mu}^i \tag{17}$$

This is the explicit from the first law which has been generally obtained. As we have promised, here the change in entropy of BH is obtained in terms of the change in parameters only. No thermodynamic potentials are needed to be defined. The identification of temperature $(\bar{T} = \bar{f}'(\bar{r}_H)/4\pi)$ has been done in (17) via standard arguments of thermal quantum field theory and we have used the well-known result of BH entropy being quarter of horizon area. More importantly, we need not know about the source explicitly (unlike the two routes mentioned earlier) to obtain the first law. If the spacetime metric is provided, we can obtain the thermodynamic law from (17).

Generalized Smarr-like formula

Earlier, we have identified the mass of the black hole in terms of Eq. (12). We replace $g(\bar{r}_H, \bar{\mu}^i)$ there using Eq. (15) and obtain

$$\bar{M} = 2\bar{T}\bar{S} - \frac{\bar{r}_H^2}{2} \frac{\partial g(\bar{r}_H, \bar{\mu}^i)}{\partial \bar{r}_H} , \qquad (18)$$

where the expressions of \overline{T} and \overline{S} are mentioned earlier. The last term in Eq. (18) contributes when the source term (or the cosmological constant) is present. When there is no source (it corresponds

²by virtual change, we mean that the change in parameters and horizon radius are not happening by any physical process. We just compare two BH solutions with radii \bar{r}_H (mass \bar{M} and so on) and $\bar{r}_H + \delta \bar{r}_H$ (mass $\bar{M} + \delta \bar{M}$ and so on).

to the Schwartschild case), $g(\bar{r}_H, \bar{\mu}^i) = 1$ and we obtain $\bar{M} = 2\bar{T}\bar{S}$, which is the Smarr formula for Schwartzschild black hole.

From Eq. (13), we can define

$$g(\bar{r},\bar{\mu}^{i}) = \bar{f}(\bar{r},\bar{M},\bar{\mu}^{i})\Big|_{\bar{M}=0} .$$
(19)

Therefore, using the above relation (19) in (18), we obtain the Smarr-like formula as

$$\bar{M} = 2\bar{T}\bar{S} - \frac{\bar{r}_H^2}{2} \left(\frac{\partial\bar{f}(\bar{r},\bar{M},\bar{\mu}^i)}{\partial\bar{r}}\right)_{\bar{r}=\bar{r}_H,\bar{M}=0}.$$
(20)

Thus, we show that for a spherically symmetric spacetime, if the metric is known, we can obtain the Smarr formula directly using Eq. (20). Therefore, not only the first law, the Smarr formula can also be obtained from the Einstein's equation.

Identification of thermodynamic pressure in extended phase space

We now take a particular example to check whether the formulation, defined above, actually provides the first law for a known case. Let us consider charged spherically symmetric black hole solution with the cosmological constant. In that case, we know the metric solution (*i.e.* the RN black hole in AdS space), which is given as

$$\bar{f}(\bar{r},\bar{M},\bar{\mu}^{i}) = 1 - \frac{2\bar{M}}{\bar{r}} + \frac{\bar{Q}^{2}}{\bar{r}^{2}} - \frac{\bar{\Lambda}\bar{r}^{2}}{3}$$
(21)

Let us consider, that the horizon radius changes infinitesimally due to the change in \overline{M} , \overline{Q} and $\overline{\Lambda}$ (*i.e.* we consider the cosmological constant as a variable instead of being a true constant in the theory). Therefore, the first law, as provided by (17), is obtained as

$$\delta \bar{M} = \bar{T} \delta \bar{S} + \frac{\bar{Q}}{\bar{r}_H} \delta \bar{Q} - \frac{4\pi \bar{r}_H^3}{3} \delta \left(\frac{\bar{\Lambda}}{8\pi}\right) \,. \tag{22}$$

One can identify \bar{Q}/\bar{r}_H as the thermodynamic potential at the horizon $(\bar{\Phi}_H)$. $4\pi \bar{r}_H^3/3$ corresponds to the volume of a sphere with radius \bar{r}_H . Therefore, in AdS spacetime (spacetime with the value of cosmological constant being negative), if the cosmological constant is considered as a variable, one can identify the last term of (22) as $\bar{V}\delta\bar{P}$ and the first law looks like

$$\delta \bar{M} = \bar{T} \delta \bar{S} + \bar{\Phi}_H \delta \bar{Q} + \bar{V} \delta \bar{P} , \qquad (23)$$

where we have identified $\bar{P} = -\bar{\Lambda}/8\pi$. This identification of pressure (for GR) has been proved to be consistent in the literature. With these identifications the $\bar{P} - \bar{V}$ criticality has been obtained in BH thermodynamics [13,14]. It is noteworthy that, the BH mass (\bar{M}) can be identified as the enthalpy of the BH thermodynamic system [8–12] when $\bar{\Lambda}$ is considered as variable in the extended phase space approach.

One can obtain the Smarr formula using our formulation. From Eq. (20), we obtain the Smarr formula of RN-AdS black hole as

$$\bar{M} = 2\bar{T}\bar{S} + \bar{Q}\bar{\Phi}_H - 2\bar{P}\bar{V} , \qquad (24)$$

which agrees with the earlier results obtained in the literature [8, 13, 14]. However, the novelty in the present analysis is that we can obtain the Smarr formula directly from the metric.

Higher dimensional generalization of the first law and Smarr-like formula

The formulation mentioned above, can be generalized to higher dimension as well. A general spherically symmetric metric in arbitrary D spacetime dimension reads

$$d\bar{s}^2 = -\bar{f}(\bar{r})d\bar{t}^2 + \frac{d\bar{r}^2}{\bar{f}(\bar{r})} + \bar{r}^2 d\bar{\Omega}_{D-2}^2 .$$
(25)

For this metric, we obtain the first law and the Smarr-like formula in a general way (see the appendix A). The first law is given as

$$\delta \bar{M} = \bar{T} \delta \bar{S} + \frac{\bar{r}_H^{D-3}}{\omega_{D-2}} \frac{\partial \bar{f}(\bar{r}, \bar{M}, \bar{\mu}^i)}{\partial \bar{\mu}^i} \Big|_{\bar{r}=\bar{r}_H} \delta \bar{\mu}^i \tag{26}$$

The generalized Smarr-like formula in arbitrary D dimension is given as (see the appendix A)

$$(D-3)\bar{M} = (D-2)\bar{T}\bar{S} - \frac{\bar{r}_{H}^{D-2}}{\omega_{D-2}} \frac{\partial\bar{f}(\bar{r},\bar{M},\bar{\mu}^{i})}{\partial\bar{r}}\Big|_{\bar{r}=\bar{r}_{H},\bar{M}=0} .$$
 (27)

Again, the last term in (27) contributes when there is a source (or the cosmological constant). For, Schwartzschild metric in *D*-dimension we obtain the Smarr formula as $(D-3)\overline{M} = (D-2)\overline{T}\overline{S}$. Here, we have defined

$$\omega_n = \frac{16\pi}{nVol(S^n)} = \frac{8\Gamma\left(\frac{n+1}{2}\right)}{n\pi^{\frac{n-1}{2}}} , \qquad (28)$$

where $Vol(S^n)$ denotes the volume of a unit *n*-sphere $d\Omega_n^2$. In arbitrary dimension *D*, the expression of entropy is given as

$$\bar{S} = \frac{Vol(S^{D-2})\bar{r}_H^{D-2}}{4} \ . \tag{29}$$

RN-AdS black hole in D spacetime dimensions:

The spacetime metric of RN-AdS black hole in D dimension is given by the $f(\bar{r})$ as [66]

$$\bar{f}(\bar{r},\bar{M},\bar{\mu}^i) = 1 - \frac{\omega_n \bar{M}}{\bar{r}^{n-1}} + \frac{n\omega_n^2 \bar{Q}^2}{8(n-1)\bar{r}^{2n-2}} - \frac{2\bar{\Lambda}}{n(n+1)}\bar{r}^2 .$$
(30)

Here n = D - 2 corresponds to the number of angular coordinates. For this metric (30), we obtain the first law from Eq. (26), which is given by the Eq. (23). Thus, in the higher dimension, we get no change in the expression of the first law.

Here, $g(\bar{r}, \bar{\mu}^i)$ can be obtained as (see Eq. (68) of the appendix A)

Using the obtained Eq. (27) we obtain the Smarr formula as

$$(D-3)\bar{M} = (D-2)\bar{T}\bar{S} + (D-3)\bar{Q}\bar{\Phi}_H - 2\bar{P}\bar{V} , \qquad (31)$$

which agrees with the existing results in literature [8, 14]. However, as we have mentioned earlier, the importance of the present analysis lies in the fact that the results (1st law and the Smarr formula)

can be obtained from the metric itself. In *D*-dimension, the potential $(\bar{\Phi}_H)$ and the volume (\bar{V}) are defined as follows [66].

$$\bar{V} = \frac{Vol(S^n)}{n+1}\bar{r}_H^{n+1}$$
, and $\bar{\Phi}_H = \frac{n}{4(n-1)}\frac{\omega_n\bar{Q}}{\bar{r}_H^{n-1}}$. (32)

Thus, in the present section, we have generally obtained the first law and the Smarr formula. Then we have considered a charged black hole in AdS space. We have found, if the cosmological constant is considered as a variable, we obtain the extra pressure and volume parameters in BH thermodynamics. Thus, the phase space of the black hole gets extended and the pressure is identified in terms of the cosmological constant and volume is its conjugate quantity. The mass of the black hole plays the role of enthalpy in the extended phase space. For RN black hole, the thermodynamic volume surprisingly matches to the geometric volume of a sphere with radius being the horizon radius. But, this may not be true in general. We shall highlight this later in more detail in our subsequent discussions. Another goal of our paper is to find the connection of thermodynamic parameters in two conformally connected frames. The thermodynamic parameters in GR is known. Therefore, in the following section, we define a conformal frame of GR and draw the connection of the parameters in the two frames.

3 Extended phase space thermodynamics in the conformally connected frame: study of (in)equivalence of thermodynamic parameters

In the earlier section, we have shown that the Einstein's equation takes the form of the first law of black hole thermodynamics in the extended phase space. However, the van der Waals type phase transition (with the P - V criticality) is also observed in Brans-Dicke theory, f(R) theory etc. which has been mentioned earlier. These theories can be described in either Jordan frame or in Einstein frame, which are conformally connected to each other. The physical equivalence of two conformal frames is a longstanding debate in literature [19,28,33–36,38–58]. However, in our earlier work [19,33–36], we have shown that the thermodynamic parameters, like entropy, temperature, internal energy (which is given by the mass of the black hole), angular momentum etc. are conformally invariant. But, the earlier analysis has been performed in the non-extended phase space. On the contrary, in the extended phase space, we have another two thermodynamic quantities: pressure (P) and volume (V). Earlier, these two parameters has not been defined in a covariant way and has not been investigated whether these are conformally equivalent. The conformal equivalence of other thermodynamic quantities in nonextended phase space do not guarantee the equivalence of P and V in the two conformally connected frames. Moreover, the thermodynamics in extended phase space is quite different compared to the non-extended one. The mass of black holes plays the role of internal energy in the non-extended phase space, whereas the same plays the role of enthalpy in the extended case. Therefore, in this section, we make a detail analysis for that. For simplicity, in the beginning, we take Einstein's gravity (GR) and later we compare it with a frame, which is conformally connected to GR. This is done because, the expressions of thermodynamic parameters in GR is already known and well-accepted. Therefore, we can straightforwardly compare the parameters of GR with the defined thermodynamic parameters in the conformal frame. In the following section, we shall generalize this result for the scalar-tensor theory in its two frame (*i.e.* Jordan and the Einstein frame). The method, that we have adopted here, is the Noether charge formalism provided by Iyer and Wald [59,60], which provides a powerful way to derive the first law and to define thermodynamic parameters in a covariant way. We do not incorporate the (energy-momentum) source term for simplicity. However, one can always incorporate source, which will contribute to the hairs of the black hole.

The Einstein-Hilbert action with the cosmological constant is given as

$$\bar{\mathcal{A}} = \int \sqrt{-\bar{g}} \bar{L} d^4 x = \frac{1}{16\pi} \int \sqrt{-\bar{g}} (\bar{R} - 2\bar{\Lambda}) d^4 x .$$
(33)

The procedure of obtaining the thermodynamic first law for the above action (33) (in the extended phase space) can be found in our earlier work [16]. Therefore, instead of repeating it, we directly go to the conformal frame. We provide the following conformal transformation to the metric tensor \bar{g}_{ab} and start the analysis in the conformal frame.

$$\bar{g}_{ab} = \phi g_{ab} , \qquad (34)$$

where ϕ is the conformal factor. In the conformal frame (described by the metric-tensor g_{ab}), one can obtain the equivalent action, which is given as

$$\mathcal{A} = \int \sqrt{-g}L = \frac{1}{16\pi} \int \sqrt{-g} \Big[\phi R + \frac{3}{2\phi} (\nabla_i \phi) (\nabla^i \phi) - 3\Box \phi - 2\phi^n \Lambda \Big] , \qquad (35)$$

where $\Lambda = \phi^{2-n}\overline{\Lambda}$ (Note that here we have used n as the exponent and not the spacetime index. This has been followed throughout). One can ask why we do not write the last term inside the square bracket of Eq. (35) as $2\phi^2\overline{\Lambda}$ instead of $2\phi^n\Lambda$. For that, in the conformal frame, we define a variable Λ , which is the cosmological constant multiplied with the arbitrary power of the conformal factor (as $\Lambda = \phi^{2-n}\overline{\Lambda}$ and n is kept arbitrary). Doing this, we shall keep the generality in the expression of the thermodynamic pressure as we shall go on to define $P = -\Lambda/8\pi$. The reason for keeping this generality is the following. In literature, there are several works in which the pressure has been accounted as the cosmological constant multiplied with the arbitrary power of the conformal factor. For example, in ref [29], the thermodynamic pressure is identified as $P = -(\phi\overline{\Lambda})/8\pi$, which corresponds to the n = 1case of our analysis. Some other works [30,31] accounts $P = \overline{\Lambda}/8\pi$, which corresponds to n = 2 of our analysis. Therefore, for a general analysis we keep it as $\phi^n\Lambda$ in (35) and the pressure will be identified as $P = -\Lambda/8\pi$. Later, we will check the consequences for different values of n.

Here we plan to obtain the covariant thermodynamic description using the Wald's formulation. For that, we need to obtain the Noether current and potential due to the diffeomorphism invariance. We obtain the Noether current and potential in the following.

The variation of the above action (35) (in the extended phase space approach *i.e.* considering Λ as a variable) yields

$$\delta(\sqrt{-g}L) = \sqrt{-g} \Big[E_{ab} \delta g^{ab} + E_{(\phi)} \delta \phi - \frac{2\phi^n}{16\pi} \delta \Lambda + \nabla_a \Theta^a(q, \delta q) \Big] .$$
(36)

Here, $q = \{g_{ab}, \phi\}$ and the expressions of E_{ab} , $E_{(\phi)}$ and $\theta^a(q, \delta q)$ are provided as follows:

$$\begin{split} E_{ab} &= \frac{1}{16\pi} \Big[\phi G_{ab} + g_{ab} \nabla_i \nabla^i \phi - \nabla_a \nabla_b \phi + \frac{3}{2\phi} (\nabla_a \phi) (\nabla_b \phi) - \frac{3}{4\phi} g_{ab} (\nabla^i \phi) (\nabla_i \phi) + g_{ab} \phi^n \Lambda \Big] \ . \\ E_{(\phi)} &= \frac{1}{16\pi} \Big[R + \frac{3}{2\phi^2} (\nabla^i \phi) (\nabla_i \phi) - \frac{3}{\phi} \Box \phi - 2n\phi^{n-1} \Lambda \Big] \ . \end{split}$$
and

$$\theta^{a}(q,\delta q) = \frac{1}{16\pi} \Big[2\phi P^{ibad}(\nabla_{b}\delta g_{id}) - 2P^{iabd}(\nabla_{b}\phi)\delta g_{id} + \frac{3}{\phi}(\nabla^{a}\phi)\delta\phi - \frac{3}{2}g^{ij}(\nabla^{a}\phi)\delta g_{ij} - 3(\nabla_{b}\phi)\delta g^{ab} - 3\nabla^{a}(\delta\phi) \Big] .$$

$$(37)$$

Here, we define P^{abcd} as $P^{abcd} = (g^{ac}g^{bd} - g^{ad}g^{bc})/2$. We can obtain a general Bianchi-like identity in this case as follows

$$\nabla_b E^{ab} = -\frac{1}{2} E_{(\phi)}(\nabla^a \phi) + \frac{1}{16\pi} \phi^n (\nabla^a \Lambda) .$$
(38)

The above identity (38) will help us to obtain off-shell Noether current and potential due to the diffeomorphism invariance. Firstly, we want to obtain the change of the Lagrangian due to the diffeomorphism invariance $x^a \longrightarrow x^a + \xi^a$, which can be obtained by replacing arbitrary variation δ in (36) by Lie derivative (\pounds_{ξ}) . Thus, (36) yields

$$\nabla_a \left(\sqrt{-g} L \xi^a \right) = \sqrt{-g} \left(-2E^{ab} (\nabla_a \xi_b) + E_{(\phi)} \xi^a \nabla_a \phi - \frac{2\phi^n}{16\pi} \xi^a \nabla_a \phi + \nabla_a \theta^a (q, \pounds_{\xi} q) \right) .$$
(39)

To obtain Eq. (39), we have used $\pounds_{\xi}g^{ab} = -(\nabla^a\xi^b + \nabla^b\xi^a)$, $\pounds_{\xi}L = \xi^a\nabla_a L$, $\pounds_{\xi}\phi = \xi^a\nabla_a\phi$, $\pounds_{\xi}\Lambda = \xi^a\nabla_a\Lambda$ and so on. In (39), we use Bianchi like identity (given in (38)). This makes the RHS of (39) a total derivative term and one can show (39), as a whole, can be written as $\nabla_a J^a = 0$, where the conserved off-shell Noether current J^a is given as

$$J^a = L\xi^a + 2E^{ab}\xi_b - \theta^a(q, \pounds_{\xi}q) .$$

$$\tag{40}$$

The above Noether current (75) can be further expressed as a two-ranked anti-symmetric potential, also known as the Noether potential, which is given by the relation $J^a = \nabla_b J^{ab3}$, where

$$J^{ab}[\xi] = \frac{1}{16\pi} \left[\nabla^a(\phi\xi^b) - \nabla^b(\phi\xi^a) \right] \,. \tag{41}$$

So far, the Noether current and the potential is obtained off-shell, where the equation of motion is nor required to be used. In addition, the above expressions of the Noether current and potential are obtained for any arbitrary diffeomorphism. We shall follow the Wald's formulation to obtain the thermodynamic law in the extended phase space. In that case, it required to take the on-shell variation (or perturbation) of the Noether current with respect to the fields g_{ab} and ϕ (therefore, $\delta\xi^a = 0$ but $\delta\xi_a \neq 0$). In order to adopt Wald's formalism, we need to consider the arbitrary diffeomorphism as the Killing diffeomorphism. From here onward, ξ^a will be considered as the Killing vector. The onshell perturbation of the Noether current is given as

$$\delta(\sqrt{-g}J^a) = \xi^a \delta(\sqrt{-g}L) - \delta(\sqrt{-g}\theta^a(q, \pounds_{\xi}q))$$
$$= \sqrt{-g}\xi^a \nabla_b \theta^b(q, \delta q) - \delta(\sqrt{-g}\theta^a(q, \pounds_{\xi}q)) - \sqrt{-g}\frac{2\phi^n}{16\pi}\xi^a \delta\Lambda .$$
(42)

³see appendix A of [35] and consider $\omega(\phi) = -3/2$ and $V(\phi) = 2\phi^n \Lambda$ which mimics our system.

Upon using $\sqrt{-g}\xi^a \nabla_b \theta^b(q, \delta q) = \pounds_{\xi}(\sqrt{-g}\theta^a(q, \delta q)) + 2\sqrt{-g}\nabla_b \xi^{[a}\theta^{b]}(q, \delta q)$ in the above equation (42) (where $A^{[a}B^{b]} = (A^a B^b - A^b B^a)/2$), one obtains

$$\delta(\sqrt{-g}J^a) = -\omega^a + 2\sqrt{-g}\nabla_b\xi^{[a}\theta^{b]}(q,\delta q) - \sqrt{-g}\frac{2\phi^n}{16\pi}\xi^a\delta\Lambda , \qquad (43)$$

where,

$$\boldsymbol{\omega}^{a} = -\pounds_{\xi}(\sqrt{-g}\theta^{a}(q,\delta q)) + \delta(\sqrt{-g}\theta^{a}(q,\pounds_{\xi}q))$$
(44)

is identified as the symplectic Hamiltonian (for rigourous discussion see [59]). Variation of the total Hamiltonian is given as

$$\delta H = \int_{c} d\Sigma_{a} \frac{\omega^{a}}{\sqrt{-g}} = -\delta \int_{c} d\Sigma_{a} \nabla_{b} J^{ab} + 2 \int_{c} d\Sigma_{a} \nabla_{b} \xi^{[a} \theta^{b]}(q, \delta q) - \frac{\delta \Lambda}{8\pi} \int_{c} \phi^{n} d\Sigma_{a} \xi^{a} .$$
(45)

The above integration is performed on a three-dimensional Cauchy hypersurface, which has been symbolized by "c" in the subscript of the integral. Here, $d\Sigma_a = n_a \sqrt{h} d^3 x$ is the elemental surface area of the hypersurface, with n_a being the normal and h being the determinant of the induced metric of the surface. The first two terms of RHS of Eq. (45) can be written in terms of two-surface integral applying the Stoke's theorem. Since the spacetime contains a horizon, the two surface will not be a closed one. Instead, it will have two ends- one is the black hole horizon (\mathcal{H}) and the other one is the asymptotic infinity (∂c_{∞}). Following Wald [59,60], we also consider that the black hole horizon is a bifurcation Killing horizon, where the Killing normal $\xi^a = 0$. One can check, for the presence of the Killing vector in the spacetime, ω^a vanishes (*i.e.* H vanishes). Thereby, we obtain

$$-\frac{1}{2}\delta \int_{\mathcal{H}} d\Sigma_{ab} J^{ab}[\xi] + \frac{1}{2}\delta \int_{\partial c_{\infty}} d\Sigma_{ab} J^{ab}[\xi] - \frac{1}{16\pi} \int_{\partial c_{\infty}} d\Sigma_{ab} \xi^{[a}\theta^{b]}(q,\delta q) + \delta P \int_{c} \phi^{n} d\Sigma_{a} \xi^{a} = 0 , \quad (46)$$

where we identify $P = -\Lambda/8\pi$ and $d\Sigma_{ab} = \sqrt{\sigma}(\xi_a l_b - \xi_b l_a)$ is the elemental surface area of the two surfaces (\mathcal{H} and ∂c_{∞}), with l being the auxiliary null vector and σ being the determinant of the induced metric on the two surface. We consider the black hole metric to be stationary-axisymmetric, which possess the Killing vector $\boldsymbol{\xi} = \boldsymbol{\xi}_{(t)} + \Omega \boldsymbol{\xi}_{\phi}$, where $\boldsymbol{\xi}_{(t)}$ and $\boldsymbol{\xi}_{\phi}$ are the components of the Killing vector along the temporal and the azimuthal directions respectively and Ω is the angular velocity. Then the above relation (46) can be identified as the first law in the extended phase space, which is given as

$$\delta M = T\delta S + \Omega_H \delta J + V\delta P , \qquad (47)$$

where, following Wald's approach, one can identify mass (M), entropy (S) and the angular momentum (J) as

$$\delta S = \frac{\pi}{\kappa} \delta \int_{\mathcal{H}} d\Sigma_{ab} J^{ab} ;$$

$$\delta M = \frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\Sigma_{ab} J^{ab}) - 2d\Sigma_{ab} \xi^{[a} \Theta^{b]}(q, \delta q)] \Big|_{\xi = \xi_{(t)}} ;$$

$$\delta J = -\frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\Sigma_{ab} J^{ab}) - 2d\Sigma_{ab} \xi^{[a} \Theta^{b]}(q, \delta q)] \Big|_{\xi = \xi_{(\phi)}} .$$
(48)

In the above relations (48), κ is the surface gravity of the Killing horizon and the temperature is identified as $T = \kappa/2\pi$.

Let us now comment upon the thermodynamic volume (denoted as V) in the following. A lot of work has been done [67–74] in the context of thermodynamic volume in the extended phase space. In general, thermodynamic volume, is the conjugate quantity of P and may not be the same as the geometric volume. The expression of the thermodynamic volume is being provided in the following way. Note that $\phi^n \xi^a$ is divergenceless (*i.e.* $\nabla_a(\phi^n \xi^a) = 0$). Therefore, we can introduce a two-ranked anti-symmetric super-potential χ^{ab} as [8]

$$\phi^n \xi^a = \nabla_b(\chi^{ab}) \tag{49}$$

The potential χ^{ab} is not unique. One can add χ'^{ab} with χ^{ab} , where $\nabla_b(\chi'^{ab}) = 0$. In any case, the three integration of Eq. (46), can be written as

$$\int_{c} \phi^{n} d\Sigma_{a} \xi^{a} = \frac{1}{2} \Big[\int_{\mathcal{H}} d\Sigma_{ab} \chi^{ab} - \int_{\partial c_{\infty}} d\Sigma_{ab} \chi^{ab} \Big].$$
(50)

The last term, *i.e.* the integration on the asymptotic infinite, can provide a diverging contribution. Therefore, regularization procedure is required to be adopted (for the arguments in the similar line, please see [8, 74]). Using the regularization technique, as provided in literature [8, 74], we define the expression of the thermodynamic volume as

$$V = \frac{1}{2} \Big[\int_{\mathcal{H}} d\Sigma_{ab} \chi^{ab} - \int_{\partial c_{\infty}} d\Sigma_{ab} \Big(\chi^{ab} - \chi^{ab}_{(BG)} \Big) \Big], \tag{51}$$

where, the term containing $\chi^{ab}_{(BG)}$ is considered as the background contribution, which is incorporated in order to obliterate the divergence in the expression of thermodynamic volume. The similar expression of thermodynamic volume has been provided in the ref [8]. Moreover, as we see in the expression of the first law, as provided in Eq (47), the mass of the black hole plays the role of enthalpy instead of internal energy. Furthermore, the Wald's method is applicable only when there is a Killing symmetry in the spacetime and the BH horizon is the Killing horizon. However, for the two conformally connected frames, if there is a Killing vector in one frame, it will be a conformal Killing vector in the other frame and the BH horizon will be the conformal Killing horizon. However, if the conformal factor (here ϕ) is Lie-transported along the direction of the Killing vector (*i.e.* $\pounds_{\xi}\phi = 0$), the Killing vector in one frame will be the Killing vector in the other frame. Thus, in our analysis, we have considered ϕ to be Lie-transported along the direction of the Killing vector. This is a standard practice and have also been adopted in earlier works (such as [33]).

Equivalence of the thermodynamic parameters

Let us now see how the thermodynamic parameters of Einstein's gravity and the same in the conformally connected frames are related. As we have mentioned, using the same Iyer-Wald technique, the first law has been proved for Einstein-Hilbert action and the thermodynamic parameters have also been obtained [16]. In Einstein's gravity, the 1st law and the expression of the thermodynamic parameters are provided as follows.

$$\delta \bar{M} = \bar{T} \delta \bar{S} + \bar{\Omega}_H \delta \bar{J} + \bar{V} \delta \bar{P} , \qquad (52)$$

where the thermodynamic parameters are provided as

$$\delta \bar{S} = \frac{\pi}{\bar{\kappa}} \delta \int_{\mathcal{H}} d\bar{\Sigma}_{ab} \bar{J}^{ab} ;$$

$$\delta \bar{M} = \frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\bar{\Sigma}_{ab} \bar{J}^{ab}) - 2d\bar{\Sigma}_{ab} \bar{\xi}^{[a} \bar{\Theta}^{b]}(\bar{q}, \delta \bar{q})] \Big|_{\bar{\xi} = \bar{\xi}_{(t)}} ;$$

$$\delta \bar{J} = -\frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\bar{\Sigma}_{ab} \bar{J}^{ab}) - 2d\bar{\Sigma}_{ab} \bar{\xi}^{[a} \bar{\Theta}^{b]}(\bar{q}, \delta \bar{q})] \Big|_{\bar{\xi} = \bar{\xi}_{(\phi)}} .$$
(53)

In the above relation, q corresponds to the metric tensor describing the Einstein's gravity in its original frame (*i.e.* $q = \bar{g}_{ab}$). For GR, the Noether potential is given as $\bar{J}^{ab} = (\bar{\nabla}^a \bar{\xi}^b - \bar{\nabla}_b \bar{\xi}^a)/16\pi$ and the boundary term is given as $\bar{\Theta}^a(\bar{q}, \delta \bar{q}) = 2\bar{P}^{ibad}\bar{\nabla}_b \delta \bar{g}_{id}$ [64]. Here, we have assumed that the Killing vectors in the two frames coincide (*i.e.* $\bar{\xi}^a = \xi^a$, implying $\bar{\xi}_a = \phi \xi_a$). Thus, the surface gravity $\kappa = \sqrt{((\nabla_a \xi^b)(\nabla_b \xi^a))/2}$ and $\bar{\kappa} = \sqrt{((\bar{\nabla}_a \bar{\xi}^b)(\bar{\nabla}_b \bar{\xi}^a))/2}$ can be proved to be the same on the bifurcation 2-surface (where $\xi^a = \bar{\xi}^a = 0$). The proof of the zeroth law [36] suggests that the constancy of κ (or $\bar{\kappa}$) holds not only on the bifurcation surface, but also all over the BH horizon. Thus, $\kappa = \bar{\kappa}$ is valid all over the BH horizon. Which implies $\bar{T} = T$. The equivalence of the angular velocity can be proved along the same line of argument provided in [33]. One can prove that the Noether potential and the surface term are related in the two frames as

$$\bar{J}^{ab} = \frac{J^{ab}}{\phi^2} \tag{54}$$

and

$$\bar{\Theta}^a(\bar{q},\delta\bar{q}) = \frac{\Theta^a(q,\delta q)}{\phi^2} .$$
(55)

One can obtain the elemental surface area of the two surface is connected in the two frames as $d\bar{\Sigma}_{ab} = \phi^2 d\Sigma_{ab}$. Thus, one can prove the equivalence of the thermodynamic parameters as $\bar{M} = M$, $\bar{S} = S$ and $\bar{J} = J$ from the two equations (48) and (53). in the action (35), the term $3\Box\phi$ is a total derivative term, which could have been discarded from the action as the removal of a total derivative term does not affect the dynamics of the fields. However, if we had discarded $3\Box\phi$ term, the exact equivalence of mass, angular-momentum and entropy could not have been established (for more discussion along this line, please see [19, 35]). Thus, the $3\Box\phi$ is very important in proving the exact equivalence of the thermodynamic parameters.

Let us discuss upon the equivalence of pressure and volume. As we have identified $\Lambda = \phi^{2-n} \overline{\Lambda}$, we obtain

$$P = \phi^{2-n} \bar{P} \ . \tag{56}$$

Thus the expression of pressure is not equivalent in the two frames unless n = 2. The expression of volume is also equivalent iff n = 2. Thus, we can say, in the conformal frame for Einstein-Hilbert action, the thermodynamic quantities, such as mass, entropy, angular-momentum, temperature *etc.* are equivalent in the two frames and the equivalence of thermodynamic pressure and volume depends upon the choice of n. However, the first law can be proved for any arbitrary choice of n. Thus our analysis does not guarantee that the pressure and volume has to be conformally equivalent. Since the other parameters (such as mass, entropy *etc.*) are explicitly shown to be equivalent in the two

frames, one can expect the same. The formalism, mentioned above, is applicable irrespective of the equivalence of pressure and volume (*i.e.* irrespective of the choice of n).

In the next section, we show that the same results are valid for a more general case, *i.e.* for the scalar-tensor gravity which is described in the Jordan and the Einstein frame, which are conformally connected. The analysis presented in this section will provide us the hint for the general ST gravity, which is presented in the following section.

4 Extended phase space thermodynamics in the scalar-tensor gravity

In this section, we generalize our earlier results for the scalar-tensor gravity, which is described in two frames. In the original frame, which is known as the Jordan frame, the action is given as

$$\mathcal{A}^{(ST)} = \int d^4x \sqrt{-g} \frac{1}{16\pi} \Big(\phi R - \frac{\omega(\phi)}{\phi} g^{ab} \nabla_a \phi \nabla_b \phi - \mathcal{V}(\phi) \Big) .$$
(57)

In this frame, the scalar field ϕ is nonminimally coupled with the Ricci-scalar R. In addition, $\omega(\phi)$ is known as the Brans-Dicke parameter, which is considered as an arbitrary function of the scalar field ϕ . When $\omega(\phi)$ is constant, this theory boils down to the Brans-Dicke (BD) theory. Furthermore, we have the scalar field potential $\mathcal{V}(\phi)$, which can also contain the cosmological constant $\overline{\Lambda}$ (see ref [31,32] for instance), in extended phase space approach which will be regarded as a variable. Thus, from now onward, we consider $\mathcal{V}(\phi) \equiv \mathcal{V}(\phi, \overline{\Lambda})$. The nonminimal coupling in the original frame can be removed by a set of transformation, one is the conformal transformation and the other one is the rescaling of the scalar field. The transformation relations are given as

$$\tilde{g}_{ab} = \phi g_{ab},\tag{58}$$

and

$$\phi \to \tilde{\phi} \text{ with } d\tilde{\phi} = \sqrt{\frac{2\omega(\phi) + 3}{16\pi}} \frac{d\phi}{\phi} .$$
 (59)

With the help of above two transformation relations (58) and (59), one can write the equivalent form of the action (57) in the Einstein frame as

$$\tilde{\mathcal{A}} = \int d^4x \sqrt{-\tilde{g}} \tilde{\mathcal{L}} = \int d^4x \sqrt{-\tilde{g}} \Big[\frac{\tilde{R}}{16\pi} - \frac{1}{2} \tilde{g}^{ab} \tilde{\nabla}_a \tilde{\phi} \tilde{\nabla}_b \tilde{\phi} - \mathcal{U}(\tilde{\phi}, \bar{\Lambda}) \Big] , \qquad (60)$$

where $\mathcal{U}(\tilde{\phi}, \bar{\Lambda}) = \mathcal{V}(\phi, \bar{\Lambda})/(16\pi\phi^2)$. In literature, one finds that the action of ST gravity in the two frames are provided by Eqs (57) and (60). However, the two actions (57) and (60) are not exactly equivalent under the transformation relations (58) and (59). Instead the two actions (or the Lagrangians) differ by a total derivative term $(-3\Box\phi)$. If one neglects $-3\Box\phi$ term in the action the dynamics of the system remains invariant but, there remains an in-built inequivalence in the action, which results in the several inequivalences in the two frame; such as the violation of holographic principal in the Jordan frame, inequivalence of Noether current and potential in the two frames, which further lead to inequivalence of the thermodynamic parameters *etc.* (for more discussions in this regard, see [19,34]). For the specific choice of $\omega(\phi) = -3/2$ and $\mathcal{U}(\tilde{\phi}, \bar{\Lambda}) = \bar{\Lambda}/8\pi$, one recovers the Einstein's gravity with the cosmological constant term (where Eq. (60) boils down to the Eq. (33)). As we have mentioned earlier, in several works, the thermodynamic pressure for f(R) and BD theory has been considered as the cosmological constant multiplied with different powers of the the conformal factor (here ϕ) in different cases. Therefore, to maintain the generality in the expression of pressure, we consider the thermodynamic pressure in the Jordan frame is given as $P = -\Lambda/\pi$. Whereas, in Einstein frame we defined it as $\tilde{P} = -\bar{\Lambda}/8\pi$, where Λ and $\bar{\Lambda}$ are related by $\Lambda = \phi^{(n-2)}\Lambda^4$. Thus, incorporating $-3\Box\phi$ term and considering Λ as a variable instead of $\bar{\Lambda}$, the action of the Jordan frame is given as

$$\mathcal{A} = \int d^4x \sqrt{-g} \mathcal{L} = \int d^4x \sqrt{-g} \frac{1}{16\pi} \left(\phi R - \frac{\omega(\phi)}{\phi} g^{ab} \nabla_a \phi \nabla_b \phi - \mathcal{V}(\phi, \Lambda) - 3\Box \phi \right) \,. \tag{61}$$

From now onward, the action of the Jordan frame will be considered as \mathcal{A} (given in (61)) instead of $\mathcal{A}^{(ST)}$ (given in (57)) for the aforementioned reasons.

In both the frames, one can obtain the first law following the Wald's approach as shown in the previous section and can define the thermodynamic parameters as earlier. In order to avoid repeating similar calculations, in appendix B we have obtained the first law and have shown that Mass, entropy, angular momentum, angular velocity and temperature (*i.e.* the non-extended phase space variables) are conformally equivalent ⁵. The pressure in the Jordan frame is already identified as $P = -\Lambda/8\pi$ and, the thermodynamic volume is identified as

$$V = -\frac{1}{2} \int_{\mathcal{H}} \frac{\partial \mathcal{V}(\phi, \Lambda)}{\partial \Lambda} \sqrt{h} n_a \xi^a d^3 y + \frac{1}{2} \int_{\infty} \frac{\partial \mathcal{V}(\phi, \Lambda)}{\partial \Lambda} \sqrt{h} \Big[n_a \xi^a - (n_a \xi^a)_{BG} \Big] d^3 y , \qquad (62)$$

where n_a is the normal and h is the determinant of the induced metric of $t = \text{const. hypersurface}^6$

Similarly, in Einstein frame, the expression of pressure is given as $\tilde{P} = -\bar{\Lambda}/8\pi$ and the expression of thermodynamic volume is given as

$$\tilde{V} = -8\pi \int_{\mathcal{H}} \frac{\partial \mathcal{U}(\tilde{\phi}, \bar{\Lambda})}{\partial \bar{\Lambda}} \sqrt{\tilde{h}} \tilde{n}_a \tilde{\xi}^a d^3 y + 8\pi \int_{\infty} \frac{\partial \mathcal{U}(\tilde{\phi}, \bar{\Lambda})}{\partial \bar{\Lambda}} \sqrt{\tilde{h}} \Big[\tilde{n}_a \tilde{\xi}^a - (\tilde{n}_a \tilde{\xi}^a)_{BG} \Big] d^3 y , \qquad (63)$$

Again, here we have adopted the regularization prescription in order to ward off the divergence appearing in the expressions of volume as provided in (62) and (63). One can easily check and find that the thermodynamic pressure and volume will be equivalent in the two frames only when n = 2 and $\Lambda = \overline{\Lambda}$ (thus $P = \overline{P} = -\overline{\Lambda}/8\pi$).

We provide some important comments about the analysis done above. As we have mentioned earlier, it has been found that the expression of thermodynamic pressure for alternative theories of gravity, such as Brans-Dicke theory, f(R) theory etc., have been considered different in different literature while studying the P - V criticality. Whereas, the expression of thermodynamic pressure in Einstein's gravity is unanimously accepted. The f(R) theory and the Brans-Dicke theory can

⁴The thermodynamic pressure in the Einstein frame is considered as the same of GR due to the fact that in the limit $\omega(\phi) = -3/2$, Einstein frame action boils down to the Einstein-Hilbert action.

⁵In the context of ST gravity, the non-extended phase space variables are already defined and have shown equivalent in the two frames [19, 33–36]. However, the present discussion is valid for both non-extended phase space variables as well as extra variables (P and V) appearing in the extended phase space but, here we emphasize on P and V only.

⁶the above expression of volume (62) can be expressed as the integration of two surface (introducing Killingsuperpotential, as done in the section 3) when $\mathcal{V}(\phi, \Lambda)$ is $\mathcal{O}(\Lambda)$, as a result of which $\partial V(\phi, \Lambda)/\partial \Lambda$ is a function of ϕ only. For example, it can be done for the $\mathcal{V}(\phi, \Lambda)$ given in [31] (Eq. 13).

be studied as the special cases scalar-tensor gravity in the Jordan frame. On the other hand, one can roughly say that the GR can be considered as the special case of scalar-tensor theory in the Einstein frame. It has been found that the thermodynamic parameters of non-extended phase space are equivalent in the two conformally connected frames [19, 33–36]. Therefore, normally it can be expected that the extra two parameters in the extended phase space approach (*i.e.* P and V) should also be equivalent in the two frames. Nevertheless, the formalism described above to obtain the first law is valid irrespective of the equivalence of pressure and volume.

Thus, in our analysis, we have first considered the Einstein's gravity (GR) and have compared the thermodynamic parameters of GR (in the extended phase space approach) with that of its conformal frame. Here, we have adopted Wald's approach, which have been found useful not only to prove the first law but also to provide the expression of the thermodynamic parameters in a covariant manner and, thereby, establishing connection of the thermodynamic parameters in the two frames. Our analysis suggests, if thermodynamic pressure and volume have to be equivalent in the two frames (which is most likely as other thermodynamic parameters are shown to be equivalent) one always have to consider thermodynamic pressure in terms of cosmological constant only. Later, we have generalized this result for scalar-tensor gravity. The significance in our approach lies in the fact that we have provided all the expression of the thermodynamic parameters in a covariant way.

5 Conclusions and outlook

Extended phase space thermodynamics has been studied extensively under the framework of Einstein's GR, where the cosmological constant is considered as the thermodynamic pressure (P) and its conjugate quantity is considered as the thermodynamic volume (V). However, recently the P - Vcriticality has also been found for Brans-Dicke and for the f(R) gravity, which can be studied as a special case of the ST gravity and can be described in the two conformally connected frames. However, there remains a discrepancy in the expression of pressure in different literature. In some literature, the cosmological constant alone is considered as the pressure whereas, in another case, the cosmological constant, multiplied with conformal factor is identified as the thermodynamic pressure. In addition, the equivalence of the thermodynamic parameters, in the extended phase space approach, has not been studied for the conformally connected frames.

Motivated by these facts, we present the analysis of this paper. Firstly in GR, we provide a general way of obtaining the first law and the Smarr formula from Einstein's equation, which works both for the extended phase space and for the non-extended phase space. Generally, while proving the first law from the Einstein's equation, people define thermodynamic potential and obtain the equilibrium version of first law in terms of the virtual change of the defined thermodynamic potential. However, originally the equilibrium version of the first law is obtained in terms of the virtual change of the parameters present in the metric. We provide a general analysis along this route. Thereby we also show that in extended phase space thermodynamics, the pressure can be identified in terms of the cosmological constant. Later, we investigate the connection of thermodynamic parameters in the two conformally connected frames. Since, the thermodynamic parameters in Einstein's GR is already well-known and well-accepted, we introduce the conformal frame of GR and obtain the first law using Iyer-Wald formalism. Thereby, we also obtain the expression of thermodynamic parameters of extended phase space in a covariant manner, which we compare with the parameters is attainable only

when the pressure is identified in terms of cosmological constant only and not multiplied with the conformal factor. Finally, we have extended this analysis for the general scalar-tensor gravity in its two frames, where the similar conclusions are drawn.

Thus, in this work, the extended phase space thermodynamics has been thoroughly studied for two conformally connected frames to address the discrepancy in the literature. Most importantly, we have provided the expressions of thermodynamic parameters in a covariant way following standard existing formalisms of GR. In addition, we have shown the procedure to obtain equilibrium version of first law and the Smarr formula in a general way for a static and spherically symmetric black hole in GR. We hope the analysis provides some significant insights in the context of extended phase space thermodynamics. More on this topic is expected to be reported soon.

Acknowledgement: I am thankful to Late Prof. Thanu Padmanabhan for his comments on the draft of this article. Special thanks to Prof. Bibhas Ranjan Majhi for his comments on this article.

A Obtaining the first law and the Smarr-like formula in arbitrary dimension

For the metric (25), the Einstein's equation is given as

$$(D-3)\left(-1+\bar{f}(\bar{r})\right)+\bar{r}\bar{f}'(\bar{r})=\frac{16\pi}{D-2}\bar{r}^2\bar{T}_{\bar{r}}^{\bar{r}}.$$
(64)

The general solution of the above equation comes in the following form

$$\bar{f}(\bar{r},\bar{\mu}^{i}) = 1 - \left(\frac{\bar{r}_{H}}{\bar{r}}\right)^{D-3} + \frac{1}{\bar{r}^{D-3}} \int_{\bar{r}_{H}}^{\bar{r}} \frac{16\pi}{D-2} T_{\bar{r}}^{\bar{r}}(\bar{r},\bar{\mu}^{i}) \bar{r}^{D-2} d\bar{r} .$$
(65)

This expression can be written in the following form

$$\bar{f}(\bar{r},\bar{\mu}^{i}) = g(\bar{r},\bar{\mu}^{i}) - \left(\frac{\bar{r}_{H}}{\bar{r}}\right)^{D-3} g(\bar{r}_{H},\bar{\mu}^{i}) .$$
(66)

We identify

$$\omega_{D-2}\bar{M} = \bar{r}_H^{D-3} g(\bar{r}_H, \bar{\mu}^i) , \qquad (67)$$

and, thereby, we can write Eq. (66) as

$$\bar{f}(\bar{r}, \bar{M}, \bar{\mu}^i) = g(\bar{r}, \bar{\mu}^i) - \frac{\omega_{D-2}M}{\bar{r}^{D-3}} , \qquad (68)$$

where

$$g(x,\bar{\mu}^i) = 1 - \frac{1}{x^{D-3}} \int^x \frac{2}{D-2} \epsilon(x,\bar{\mu}^i) x^{D-2} dx .$$
(69)

Eq. (68) implies that $g(\bar{r},\bar{\mu}^i)=\bar{f}(\bar{r},\bar{M},\bar{\mu}^i)_{\bar{M}=0}$. From Eq. (66), we obtain

$$\bar{f}'(\bar{r}_H, \bar{M}, \bar{\mu}^i) = g'(\bar{r}_H, \bar{\mu}^i) + \frac{D-3}{\bar{r}_H}g(\bar{r}_H, \bar{\mu}^i) .$$
(70)

Replacing $g(\bar{r}_H, \bar{\mu}^i)$ of Eq. (67) using the above Eq. (70) we obtain Eq. (27). Let us now consider the horizon radius changes infinitesimally due to the change in parameters $(\bar{M} \text{ and } \bar{\mu}^i)$. In that case we have (from Eq. (67))

$$\omega_{D-2}\delta\bar{M} = \left[(D-3)\bar{r}_{H}^{D-4}g(\bar{r}_{H},\bar{\mu}^{i}) + \bar{r}_{H}^{D-3}g'(\bar{r}_{H},\bar{\mu}^{i}) \right]\delta\bar{r}_{H} + \bar{r}_{H}^{D-3}\frac{\partial g(\bar{r}_{H},\bar{\mu}^{i})}{\partial\bar{m}u^{i}}\delta\bar{\mu}^{i} .$$
(71)

Using equations (16) and (70) in (71), we obtain (26).

B Obtaining 1st law in ST gravity for both the frames in extended phase space

The variation of the action (61) yields

$$\delta(\sqrt{-g}\mathcal{L}) = \sqrt{-g} \Big[\mathcal{E}_{ab} \delta g^{ab} + \mathcal{E}_{(\phi)} \delta \phi - \frac{1}{16\pi} \frac{\partial \mathcal{V}(\phi, \Lambda)}{\partial \Lambda} \delta \Lambda + \nabla_a \Psi^a(q, \delta q) \Big] .$$
(72)

Here, $q = \{g_{ab}, \phi\}$ and the expressions of $\mathcal{E}_{ab}, \mathcal{E}_{(\phi)}$ and $\Psi^a(q, \delta q)$ are provided as follows:

$$\begin{aligned} \mathcal{E}_{ab} &= \frac{1}{16\pi} \Big[\phi G_{ab} + g_{ab} \nabla_i \nabla^i \phi - \nabla_a \nabla_b \phi - \frac{\omega(\phi)}{\phi} (\nabla_a \phi) (\nabla_b \phi) + \frac{\omega(\phi)}{2\phi} g_{ab} (\nabla^i \phi) (\nabla_i \phi) + \frac{1}{2} g_{ab} \mathcal{V}(\phi, \Lambda) \Big] \ . \\ \mathcal{E}_{(\phi)} &= \frac{1}{16\pi} \Big[R - \frac{\omega(\phi)}{\phi^2} (\nabla^i \phi) (\nabla_i \phi) + \frac{2\omega(\phi)}{\phi} \Box \phi - \frac{\partial \mathcal{V}(\phi, \Lambda)}{\partial \phi} + \frac{1}{\phi} \frac{d\omega(\phi)}{d\phi} (\nabla^i \phi) (\nabla_i \phi) \Big] \ . \\ &\text{and} \\ \Psi^a(q, \delta q) &= \frac{1}{16\pi} \Big[2\phi P^{ibad} (\nabla_b \delta g_{id}) - 2P^{iabd} (\nabla_b \phi) \delta g_{id} - 2\frac{\omega(\phi)}{\phi} (\nabla^a \phi) \delta \phi - \frac{3}{2} g^{ij} (\nabla^a \phi) \delta g_{ij} \Big] \end{aligned}$$

$$= \frac{1}{16\pi} \left[2\phi P^{ibaa} (\nabla_b \delta g_{id}) - 2P^{iaba} (\nabla_b \phi) \delta g_{id} - 2\frac{\nabla(\gamma)}{\phi} (\nabla^a \phi) \delta \phi - \frac{1}{2} g^{ij} (\nabla^a \phi) \delta g_{ij} - 3(\nabla_b \phi) \delta g^{ab} - 3\nabla^a (\delta \phi) \right] ,$$
(73)

where we have defined $P^{abcd} = (g^{ac}g^{bd} - g^{ad}g^{bc})/2$. The general Bianchi-like identity, in this case, is provided as

$$\nabla_b \mathcal{E}^{ab} = -\frac{1}{2} \mathcal{E}_{(\phi)}(\nabla^a \phi) + \frac{1}{32\pi} \frac{\partial \mathcal{V}(\phi, \Lambda)}{\partial \Lambda} (\nabla^a \Lambda) .$$
(74)

For diffeomorphism $(x^a \longrightarrow x^a + \xi^a)$ invariance, the arbitrary variation becomes Lie variation in (72). Then Substituting (74) in (72), one obtains the conserved offshell Noether current (\mathcal{J}^a) , which is given as

$$\mathcal{J}^a = \mathcal{L}\xi^a + 2\mathcal{E}^{ab}\xi_b - \Psi^a(q, \pounds_{\xi}q) .$$
(75)

This conserved Noether current can further be written as the divergence of two-ranked anti-symmetric potential, which is called Noether potential ($\mathcal{J}^a = \nabla_b \mathcal{J}^{ab}$), where the Noether potential is provided as⁷

$$\mathcal{J}^{ab}[\xi] = \frac{1}{16\pi} \Big[\nabla^a(\phi\xi^b) - \nabla^b(\phi\xi^a) \Big] .$$
(76)

 $^{^{7}}$ Note, that the expression of Noether current and potential is the same both for extended phase space approach and the non-extended one.

Using Wald's approach, as described in section 3, one can obtain the first law as $\delta M = T\delta S + \Omega_H \delta J + V\delta P$ with $T = \kappa/2\pi$ and

$$\delta S = \frac{\pi}{\kappa} \delta \int_{\mathcal{H}} d\Sigma_{ab} \mathcal{J}^{ab} ;$$

$$\delta M = \frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\Sigma_{ab} \mathcal{J}^{ab}) - 2d\Sigma_{ab} \xi^{[a} \Psi^{b]}(q, \delta q)] \Big|_{\xi = \xi_{(t)}} ;$$

$$\delta J = -\frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\Sigma_{ab} \mathcal{J}^{ab}) - 2d\Sigma_{ab} \xi^{[a} \Psi^{b]}(q, \delta q)] \Big|_{\xi = \xi_{(\phi)}} .$$
(77)

The expression of pressure and volume (adopting the regularization prescription) in provided in section 4.

Let us now quickly move forward to the Einstein frame. The variation of the Lagrangian, as provided in (60), yields

$$\delta(\sqrt{-\tilde{g}}\tilde{\mathcal{L}}) = \sqrt{-\tilde{g}}\tilde{\mathcal{E}}_{ab}\delta\tilde{g}^{ab} + \sqrt{-\tilde{g}}\tilde{\mathcal{E}}_{(\tilde{\phi})}\delta\tilde{\phi} - \sqrt{-\tilde{g}}\frac{\partial\mathcal{U}(\phi,\Lambda)}{\partial\bar{\Lambda}}\delta\bar{\Lambda} + \sqrt{-\tilde{g}}\tilde{\nabla}_{a}\tilde{\Psi}^{a}(\tilde{q},\delta\tilde{q}) , \qquad (78)$$

where $\tilde{q} \in {\tilde{g}_{ab}, \tilde{\phi}}$ and

$$\tilde{\mathcal{E}}_{ab} = \frac{G_{ab}}{16\pi} - \frac{1}{2} \tilde{\nabla}_a \tilde{\phi} \tilde{\nabla}_b \tilde{\phi} + \frac{1}{4} \tilde{g}_{ab} \tilde{\nabla}^i \tilde{\phi} \tilde{\nabla}_i \tilde{\phi} + \frac{1}{2} \tilde{g}_{ab} \mathcal{U}(\tilde{\phi}, \bar{\Lambda}) ;$$

$$\tilde{\mathcal{E}}_{(\tilde{\phi})} = \tilde{\nabla}_a \tilde{\nabla}^a \tilde{\phi} - \frac{\partial \mathcal{U}(\tilde{\phi}, \bar{\Lambda})}{\partial \tilde{\phi}} ;$$
and
$$\tilde{\Psi}^a(\tilde{q}, \delta \tilde{q}) = \frac{\delta \tilde{v}^a}{16\pi} - (\tilde{\nabla}^a \tilde{\phi}) \delta \tilde{\phi} .$$
(79)

where $\delta \tilde{v}^a = 2\tilde{p}^{ibad}\tilde{\nabla}_b\delta \tilde{g}_{id}$ and $\tilde{p}^{iabd} = (\tilde{g}^{ib}\tilde{g}^{ad} - \tilde{g}^{id}\tilde{g}^{ab})/2$. The, Bianchi-like identity is provided as

$$\tilde{\nabla}_{b}\tilde{E}^{ab} = -\frac{1}{2}(\tilde{\nabla}^{a}\tilde{\phi})\tilde{E}_{\phi} + \frac{1}{2}\frac{\partial\mathcal{U}(\phi,\Lambda)}{\partial\bar{\Lambda}}(\tilde{\nabla}^{a}\bar{\Lambda}) .$$
(80)

With these relations, one can obtain the conserved offshell Noether current in Einstein frame due to the diffeomorphism invariance $(\tilde{x}^a \longrightarrow \tilde{x}^a + \tilde{\xi}^a)$ as

$$\tilde{J}^a = \tilde{\mathcal{L}}\tilde{\xi}^a + 2\tilde{\mathcal{E}}^{ab}\tilde{\xi}_b - \tilde{\Psi}^a(\tilde{q}, \pounds_{\xi}\tilde{q}) .$$
(81)

and, the Noether potential is provided as

$$\tilde{J}^{ab} = \frac{1}{16\pi} (\tilde{\nabla}^a \tilde{\xi}^b - \tilde{\nabla}^b \tilde{\xi}^a) .$$
(82)

Again, following Wald's formalism, we obtain the first law as $\delta \tilde{M} = \tilde{T}\delta \tilde{S} + \tilde{\Omega}_H \delta \tilde{J} + \tilde{V}\delta \tilde{P}$ where $\tilde{T} = \tilde{\kappa}/2\pi$ and

$$\delta \tilde{S} = \frac{\pi}{\tilde{\kappa}} \delta \int_{\mathcal{H}} d\tilde{\Sigma}_{ab} \tilde{J}^{ab} ;$$

$$\delta \tilde{M} = \frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\tilde{\Sigma}_{ab} \tilde{J}^{ab}) - 2d\tilde{\Sigma}_{ab} \tilde{\xi}^{[a} \tilde{\Theta}^{b]}(\tilde{q}, \delta \tilde{q})] \Big|_{\tilde{\xi} = \tilde{\xi}_{(t)}} ;$$

$$\delta \tilde{J} = -\frac{1}{2} \int_{\partial c_{\infty}} [\delta(d\tilde{\Sigma}_{ab} \tilde{J}^{ab}) - 2d\tilde{\Sigma}_{ab} \tilde{\xi}^{[a} \tilde{\Theta}^{b]}(\tilde{q}, \delta \tilde{q})] \Big|_{\tilde{\xi} = \tilde{\xi}_{(\phi)}} .$$
(83)

In Einstein frame, the pressure and thermodynamic volume (with regularization prescription) is defined in section 4.

Equivalence of the thermodynamic parameters: One can obtain the Noether potentials are conformally connected as

$$\bar{\mathcal{J}}^{ab} = \frac{\mathcal{J}^{ab}}{\phi^2} \tag{84}$$

and

$$\tilde{\Psi}^a(\bar{q},\delta\bar{q}) = \frac{\Psi^a(q,\delta q)}{\phi^2} .$$
(85)

Also, one can obtain the elemental surface area of the two surface are connected in the two frames as $d\tilde{\Sigma}_{ab} = \phi^2 d\Sigma_{ab}$. With the help of the above two relations (84) and (85), one obtains the mass, charge and angular momentum are equivalent in Jordan and Einstein frame. The equivalence of the temperature and angular velocity can be proved applying the exactly same argument as presented in section 3, which we do not repeat here. The equivalence of pressure and volume is discussed in the section 4.

References

- [1] J. D. Bekenstein, "Black holes and entropy," Phys. Rev. D 7, 2333 (1973).
- [2] S. W. Hawking, "Particle Creation by Black Holes," Commun. Math. Phys. 43, 199 (1975)
 Erratum: [Commun. Math. Phys. 46, 206 (1976)].
- [3] J. M. Bardeen, B. Carter and S. W. Hawking, "The Four laws of black hole mechanics," Commun. Math. Phys. 31, 161 (1973).
- [4] P. C. W. Davies, "Thermodynamic Phase Transitions of Kerr-Newman Black Holes in De Sitter Space," Class. Quant. Grav. 6, 1909 (1989).
- [5] S. W. Hawking and D. N. Page, "Thermodynamics of Black Holes in anti-De Sitter Space," Commun. Math. Phys. 87, 577 (1983).
- [6] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, "Charged AdS black holes and catastrophic holography," Phys. Rev. D 60, 064018 (1999) [hep-th/9902170].
- [7] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, "Holography, thermodynamics and fluctuations of charged AdS black holes," Phys. Rev. D 60, 104026 (1999) [hep-th/9904197].
- [8] D. Kastor, S. Ray and J. Traschen, "Enthalpy and the Mechanics of AdS Black Holes," Class. Quant. Grav. 26, 195011 (2009) [arXiv:0904.2765 [hep-th]].
- B. P. Dolan, "The cosmological constant and the black hole equation of state," Class. Quant. Grav. 28, 125020 (2011) [arXiv:1008.5023 [gr-qc]].
- [10] B. P. Dolan, "Pressure and volume in the first law of black hole thermodynamics," Class. Quant. Grav. 28, 235017 (2011) [arXiv:1106.6260 [gr-qc]].

- [11] B. P. Dolan, "Compressibility of rotating black holes," Phys. Rev. D 84, 127503 (2011) [arXiv:1109.0198 [gr-qc]].
- [12] B. P. Dolan, "Where is the PdV term in the fist law of black hole thermodynamics?," arXiv:1209.1272 [gr-qc].
- [13] D. Kubiznak and R. B. Mann, "P-V criticality of charged AdS black holes," JHEP 1207, 033 (2012) [arXiv:1205.0559 [hep-th]].
- [14] D. Kubiznak, R. B. Mann and M. Teo, "Black hole chemistry: thermodynamics with Lambda," arXiv:1608.06147 [hep-th].
- [15] B. R. Majhi and S. Samanta, "P-V criticality of AdS black holes in a general framework," Phys. Lett. B 773, 203 (2017) [arXiv:1609.06224 [gr-qc]].
- [16] K. Bhattacharya and B. R. Majhi, "Thermogeometric description of the van der Waals like phase transition in AdS black holes," Phys. Rev. D 95, no.10, 104024 (2017) [arXiv:1702.07174 [gr-qc]].
- [17] K. Bhattacharya, B. R. Majhi and S. Samanta, "Van der Waals criticality in AdS black holes: a phenomenological study," Phys. Rev. D 96, no.8, 084037 (2017) [arXiv:1709.02650 [gr-qc]].
- [18] K. Bhattacharya and B. R. Majhi, "Thermogeometric study of van der Waals like phase transition in black holes: an alternative approach," Phys. Lett. B 802, 135224 (2020) [arXiv:1903.10370 [gr-qc]].
- [19] K. Bhattacharya, "Thermodynamic aspects and phase transition of black holes," http://gyan.iitg.ernet.in/bitstream/handle/123456789/1740/TH-2294_156121009.pdf?sequence=2&isAllowed
- [20] A. Dehyadegari, B. R. Majhi, A. Sheykhi and A. Montakhab, "Universality class of alternative phase space and Van der Waals criticality," Phys. Lett. B **791**, 30 (2019) [arXiv:1811.12308 [hep-th]].
- [21] C. Eling, R. Guedens and T. Jacobson, "Non-equilibrium thermodynamics of spacetime," Phys. Rev. Lett. 96, 121301 (2006) [arXiv:gr-qc/0602001 [gr-qc]].
- [22] E. Elizalde and P. J. Silva, "F(R) gravity equation of state," Phys. Rev. D 78, 061501 (2008) [arXiv:0804.3721 [hep-th]].
- [23] G. Chirco and S. Liberati, "Non-equilibrium Thermodynamics of Spacetime: The Role of Gravitational Dissipation," Phys. Rev. D 81, 024016 (2010) [arXiv:0909.4194 [gr-qc]].
- [24] T. Padmanabhan, "Classical and quantum thermodynamics of horizons in spherically symmetric space-times," Class. Quant. Grav. 19, 5387-5408 (2002) [arXiv:gr-qc/0204019 [gr-qc]].
- [25] T. Padmanabhan, "Gravity and the thermodynamics of horizons," Phys. Rept. 406, 49-125 (2005) [arXiv:gr-qc/0311036 [gr-qc]].
- [26] A. Paranjape, S. Sarkar and T. Padmanabhan, "Thermodynamic route to field equations in Lancos-Lovelock gravity," Phys. Rev. D 74, 104015 (2006) [arXiv:hep-th/0607240 [hep-th]].

- [27] D. Kothawala, S. Sarkar and T. Padmanabhan, "Einstein's equations as a thermodynamic identity: The Cases of stationary axisymmetric horizons and evolving spherically symmetric horizons," Phys. Lett. B 652, 338-342 (2007) [arXiv:gr-qc/0701002 [gr-qc]].
- [28] V. Faraoni, "Black hole entropy in scalar-tensor and f(R) gravity: An Overview," Entropy 12, 1246 (2010) [arXiv:1005.2327 [gr-qc]].
- [29] S. Chen, X. Liu, C. Liu and J. Jing, "P V criticality of AdS black hole in f(R) gravity," Chin. Phys. Lett. **30**, 060401 (2013) [arXiv:1301.3234 [gr-qc]].
- [30] A. Övgün, "P v criticality of a specific black hole in f(R) gravity coupled with Yang-Mills field," Adv. High Energy Phys. **2018**, 8153721 (2018) [arXiv:1710.06795 [gr-qc]].
- [31] S. H. Hendi and Z. Armanfard, "Extended phase space thermodynamics and P V criticality of charged black holes in Brans-Dicke theory," Gen. Rel. Grav. 47, no.10, 125 (2015) [arXiv:1503.07070 [gr-qc]].
- [32] S. H. Hendi, R. M. Tad, Z. Armanfard and M. S. Talezadeh, "Extended phase space thermodynamics and P–V criticality: Brans–Dicke–Born–Infeld vs. Einstein–Born–Infeld-dilaton black holes," Eur. Phys. J. C 76, no.5, 263 (2016) [arXiv:1511.02761 [gr-qc]].
- [33] J. i. Koga and K. i. Maeda, "Equivalence of black hole thermodynamics between a generalized theory of gravity and the Einstein theory," Phys. Rev. D 58, 064020 (1998) [arXiv:gr-qc/9803086 [gr-qc]].
- [34] K. Bhattacharya and B. R. Majhi, "Fresh look at the scalar-tensor theory of gravity in Jordan and Einstein frames from undiscussed standpoints," Phys. Rev. D 95, no.6, 064026 (2017) [arXiv:1702.07166 [gr-qc]].
- [35] K. Bhattacharya, A. Das and B. R. Majhi, "Noether and Abbott-Deser-Tekin conserved quantities in scalar-tensor theory of gravity both in Jordan and Einstein frames," Phys. Rev. D 97, no.12, 124013 (2018) [arXiv:1803.03771 [gr-qc]].
- [36] S. Dey, K. Bhattacharya and B. R. Majhi, "Thermodynamic structure of a generic null surface and the zeroth law in scalar-tensor theory," [arXiv:2105.07787 [gr-qc]].
- [37] L. Smarr, "Mass formula for Kerr black holes," Phys. Rev. Lett. 30, 71-73 (1973) [erratum: Phys. Rev. Lett. 30, 521-521 (1973)]
- [38] V. Faraoni and E. Gunzig, "Einstein frame or Jordan frame?," Int. J. Theor. Phys. 38, 217 (1999) [astro-ph/9910176].
- [39] A. Macias and A. Garcia, "Jordan frame or Einstein frame?," Gen. Rel. Grav. 33, 889 (2001).
- [40] V. Faraoni, E. Gunzig and P. Nardone, "Conformal transformations in classical gravitational theories and in cosmology," Fund. Cosmic Phys. 20, 121 (1999) [gr-qc/9811047].
- [41] V. Faraoni and S. Nadeau, "The (pseudo)issue of the conformal frame revisited," Phys. Rev. D 75, 023501 (2007) [gr-qc/0612075].
- [42] I. D. Saltas and M. Hindmarsh, "The dynamical equivalence of modified gravity revisited," Class. Quant. Grav. 28, 035002 (2011) [arXiv:1002.1710 [gr-qc]].

- [43] S. Capozziello, P. Martin-Moruno and C. Rubano, "Physical non-equivalence of the Jordan and Einstein frames," Phys. Lett. B 689, 117 (2010) [arXiv:1003.5394 [gr-qc]].
- [44] A. Padilla and V. Sivanesan, "Boundary Terms and Junction Conditions for Generalized Scalar-Tensor Theories," JHEP 1208, 122 (2012) [arXiv:1206.1258 [gr-qc]].
- [45] T. Jacobson and G. Kang, "Conformal invariance of black hole temperature," Class. Quant. Grav. 10, L201 (1993) [gr-qc/9307002].
- [46] G. Kang, "On black hole area in Brans-Dicke theory," Phys. Rev. D 54, 7483 (1996) [gr-qc/9606020].
- [47] S. Deser and B. Tekin, "Conformal Properties of Charges in Scalar-Tensor Gravities," Class. Quant. Grav. 23, 7479-7482 (2006) [arXiv:gr-qc/0609111 [gr-qc]].
- [48] M. H. Dehghani, J. Pakravan and S. H. Hendi, "Thermodynamics of charged rotating black branes in Brans-Dicke theory with quadratic scalar field potential," Phys. Rev. D 74, 104014 (2006) [hep-th/0608197].
- [49] A. Sheykhi and M. M. Yazdanpanah, "Thermodynamics of charged Brans-Dicke AdS black holes," Phys. Lett. B 679, 311 (2009) [arXiv:0904.1777 [hep-th]].
- [50] C. F. Steinwachs and A. Y. Kamenshchik, "One-loop divergences for gravity non-minimally coupled to a multiplet of scalar fields: calculation in the Jordan frame. I. The main results," Phys. Rev. D 84, 024026 (2011) [arXiv:1101.5047 [gr-qc]].
- [51] A. Y. Kamenshchik and C. F. Steinwachs, "Question of quantum equivalence between Jordan frame and Einstein frame," Phys. Rev. D 91, no. 8, 084033 (2015) [arXiv:1408.5769 [gr-qc]].
- [52] N. Banerjee and B. Majumder, "A question mark on the equivalence of Einstein and Jordan frames," Phys. Lett. B 754, 129 (2016) [arXiv:1601.06152 [gr-qc]].
- [53] S. Pandey and N. Banerjee, "Equivalence of Jordan and Einstein frames at the quantum level," arXiv:1610.00584 [gr-qc].
- [54] M. S. Ruf and C. F. Steinwachs, "Quantum equivalence of f(R) gravity and scalar-tensor theories," Phys. Rev. D 97, no. 4, 044050 (2018) [arXiv:1711.07486 [gr-qc]].
- [55] A. Karam, T. Pappas and K. Tamvakis, "Frame-dependence of higher-order inflationary observables in scalar-tensor theories," Phys. Rev. D 96, no. 6, 064036 (2017) [arXiv:1707.00984 [gr-qc]].
- [56] S. Bahamonde, S. D. Odintsov, V. K. Oikonomou and P. V. Tretyakov, "Deceleration versus acceleration universe in different frames of F(R) gravity," Phys. Lett. B **766**, 225 (2017) [arXiv:1701.02381 [gr-qc]].
- [57] A. Karam, A. Lykkas and K. Tamvakis, "Frame-invariant approach to higher-dimensional scalartensor gravity," Phys. Rev. D 97, no. 12, 124036 (2018) [arXiv:1803.04960 [gr-qc]].
- [58] K. Bhattacharya, B. R. Majhi and D. Singleton, "Fluid-gravity correspondence in the scalartensor theory of gravity: (in)equivalence of Einstein and Jordan frames," JHEP 07, 018 (2020) [arXiv:2002.04743 [hep-th]].

- [59] R. M. Wald, "Black hole entropy is the Noether charge," Phys. Rev. D 48, no.8, R3427-R3431 (1993) [arXiv:gr-qc/9307038 [gr-qc]].
- [60] V. Iyer and R. M. Wald, "Some properties of Noether charge and a proposal for dynamical black hole entropy," Phys. Rev. D 50, 846-864 (1994) [arXiv:gr-qc/9403028 [gr-qc]].
- [61] S. A. Hayward, "Unified first law of black hole dynamics and relativistic thermodynamics," Class. Quant. Grav. 15, 3147-3162 (1998) [arXiv:gr-qc/9710089 [gr-qc]].
- [62] D. Hansen, D. Kubiznak and R. Mann, "Horizon Thermodynamics from Einstein's Equation of State," Phys. Lett. B 771, 277-280 (2017) [arXiv:1610.03079 [gr-qc]].
- [63] K. Bhattacharya and B. R. Majhi, "Temperature and thermodynamic structure of Einstein's equations for a cosmological black hole," Phys. Rev. D 94, no.2, 024033 (2016) [arXiv:1602.07879 [gr-qc]].
- [64] T. Padmanabhan, "Gravitation, Foundations and Frontiers," Cambridge University Press (2013).
- [65] Garret Birkhoff, *Relativity and Modern Physics*, Harvard University Press, Cambridge, 1923.
- [66] R. G. Cai, "Cardy-Verlinde formula and thermodynamics of black holes in de Sitter spaces," Nucl. Phys. B 628, 375-386 (2002) [arXiv:hep-th/0112253 [hep-th]].
- [67] E. Caceres, P. H. Nguyen, and J. F. Pedraza, "Holographic entanglement entropy and the extended phase structure of STU black holes," JHEP 1509, 184 (2015) [arXiv:1507.06069 [hep-th]].
- [68] A. Karch and B. Robinson, "Holographic Black Hole Chemistry," JHEP 1512, 073 (2015) [arXiv:1510.02472 [hep-th]].
- [69] E. Caceres, P. H. Nguyen, and J. F. Pedraza, "Holographic Entanglement Chemistry," arXiv:1605.00595 [hep-th].
- [70] P. H. Nguyen, "An equal area law for holographic entanglement entropy of the AdS-RN black hole," JHEP 1512, 139 (2015) [arXiv:1508.01955 [hep-th]].
- [71] D. Kastor, S. Ray and J. Traschen, "Extended First Law for Entanglement Entropy in Lovelock Gravity," Entropy 18, no. 6, 212 (2016) [arXiv:1604.04468 [hep-th]].
- [72] D. Kastor, S. Ray and J. Traschen, "Chemical Potential in the First Law for Holographic Entanglement Entropy," JHEP 1411, 120 (2014) [arXiv:1409.3521 [hep-th]].
- [73] P. Pradhan, "Thermodynamic Products in Extended Phase Space," Int. J. Mod. Phys. D 26, 1750010 (2017) [arXiv:1603.07748 [gr-qc]].
- [74] J. Couch, W. Fischler and P. H. Nguyen, "Noether charge, black hole volume, and complexity," JHEP 03, 119 (2017) [arXiv:1610.02038 [hep-th]].
- [75] Y. Fujii, "Choosing a conformal frame in scalar tensor theories of gravity with a cosmological constant," Prog. Theor. Phys. 99, 599-622 (1998) [arXiv:gr-qc/9708010 [gr-qc]].