arXiv:2112.01295v1 [gr-qc] 2 Dec 2021

Exact solution for accretion onto a moving charged dilaton black hole

Rong-Jia Yang *,^{1, 2, 3, 4, †} Yinan Jia,¹ and Lei Jiao^{1, 5}

¹College of Physical Science and Technology, Hebei University, Baoding 071002, China

²Hebei Key Lab of Optic-Electronic Information and Materials, Hebei University, Baoding 071002, China

³National-Local Joint Engineering Laboratory of New Energy Photoelectric Devices, Hebei University, Baoding 071002, China

⁴Key Laboratory of High-pricision Computation and Application of Quantum

Field Theory of Hebei Province, Hebei University, Baoding 071002, China

⁵School of Physics and Astronomy, Sun Yat-Sen University, Zhuhai 519082, China

We present an analytic solution for accretion of a gaseous medium with adiabatic equation of state onto a charged dilaton black hole which moves at a constant velocity. We determine the fourvelocity of accreted flow and find that it possesses axial symmetry. We obtain the accretion rate which depends on the mass, the magnetic charge, and the dilation of black hole, meaning that these parameters take important roles in the process of accretion. The results may help us to get deeper understanding of the behavior of accreted flow near the event horizon of black hole.

PACS numbers:

I. INTRODUCTION

Accretion of matter onto a massive object is a long-standing interesting phenomenon in astrophysics. The pioneers works in this field were published in [1–5]. Since then accretion has been an extensively studied subject in the literature, see for example [6–23]. These studies mainly focused on accretion rate, critical radius, flow parameters, and so on. In fact, we can also use accretion to open up new possibilities. In [24], attempts were made to test the asymptotically safe scenario via the quantum correction to accretion onto a renormalization-group-improved Schwarzschild black hole; while in [25] accretion onto a Schwarzschild-like black hole was considered to constrain the parameter characterizing the breaking of Lorentz symmetry. It was shown in [26, 27] that the conditions under which the accretion is possible give limits on the ratio of mass to charge. Numerical analysis, in general, are needed to dealt with nonspherical accretion for either Newtonian or relativistic flow [28–31]. So finding analytical solutions is often a very valuable task. An exact fully relativistic solution describing stationary accretion of matter with adiabatic equation of state onto a moving Schwarzschild black hole was derived in [32]. An analytic nonspherical solutions were presented for dust shells onto a Schwarzschild black hole. Recently, an exact solution representing accretion of collisionless Vlasov gas onto a moving Schwarzschild black hole was given in [36] which based on the Hamiltonian formalism developed in [37] which allows for an analysis of more complex flows on the fixed Schwarzschild background.

Exact solutions are important to accurately and directly describe the behavior of accreted matters near a black hole. However, in general, the analytical solution for accreting process is very complex and difficult to obtain. So it is valuable to find analytical solutions for accretion onto different black holes. Here we aim to analysis the accreting process for a moving charged dilaton black hole which is a solution to low-energy string theory representing a static, spherically symmetric charged black hole [38, 39]. The results obtained here and in the literatures about analytic solutions may be helpful to understand the physical mechanism of accretion onto black holes.

In the following Section II, we briefly review the fundamental equations for accretion onto a moving black hole. In Section III, we hope to derive an exact solution for accretion onto a moving charged dilaton black hole. Finally, we will briefly summarize our results in section IV.

II. FUNDAMENTAL EQUATIONS

In this section, we briefly review some basic equations in the process of accrtion. We adopt the reference frame in which the black hole is rest while the homogeneous medium moves at a constant velocity at infinity. Assuming the

^{*} Corresponding author

[†]Electronic address: yangrongjia@tsinghua.org.cn

flow of matter is a perfect fluid, then the relativistic vorictity tensor is given by [32]

$$\omega_{\mu\nu} = P^{\alpha}_{\mu} P^{\beta}_{\nu} [(hu_{\alpha})_{;\beta} - (hu_{\beta})_{;\alpha}], \tag{1}$$

where u^{μ} is the component of four-velocity, $h \equiv (\rho + P)/n$ is the enthalpy, and $P^{\nu}_{\mu} = \delta^{\nu}_{\mu} + u_{\mu}u^{\nu}$ is the projection tensor, and the semicolon denotes the covariant derivative with respect to the coordinate. Throughout this paper, we use the units including c = G = 1, where c and G are the speed of light and Newtonian gravitational constant, respectively. We take the signs of components of metric tensor of Minkowski space-time as (-, +, +, +). Since the fluid is perfect, Euler's equation becomes

$$(hu_{\mu})_{;\alpha}u^{\alpha} + h_{,\mu} = 0,$$
 (2)

where the comma denotes the ordinary derivative with respect to the coordinate. From the Eqs. (1) and (2), one can obtain a simple expression for the vorticity

$$\omega_{\mu\nu} = [(hu_{\mu})_{;\nu} - (hu_{\nu})_{;\mu}]. \tag{3}$$

The constant velocity at which the black hole moves and the homogeneity of medium upstream imply that the vorticity is zero. And if the vorticity is zero on some initial hypersurface, it will be zero everywhere, like the case in Newtonian flow. Thus the quantity hu_{μ} can be expressed as the gradient of a potential [40]:

$$hu_{\mu} = \psi_{,\mu}.\tag{4}$$

If there no particles are destroyed or created, the equation of continuity for particle density n can be written as

$$(nu^{\alpha})_{:\alpha} = 0. \tag{5}$$

With Eq. (4), we can rewrite (5) in a differential form

$$[(n/h)\psi^{,\alpha}]_{;\alpha} = 0, (6)$$

where $\psi^{,\alpha}$ means the contravariant component of the vector $\psi_{,\alpha}$. In general, Eq. (6) is a nonlinear differential equation about ψ and its derivatives. However, if h is proportional to n, it will become a linear differential equation. Considering the simplest case, $P = \rho \propto n^2$, which implies that the speed of sound is equal to the speed of light and that the adiabatic index is equal to 2. The velocity of fluid must be subsonic everywhere, so no shock waves will arise. Then Eq. (6) now reads

$$(\psi^{,\alpha})_{;\alpha} = 0. \tag{7}$$

We must solve this differential equation under appropriate boundary conditions to obtain physical quantities of accretion which we desire. One of the physical quantities of accretion we want to get is the particle number accretion rate

$$\dot{N} = -\int_{S} nu^{i} \sqrt{-g} dS_{i} = -\int_{S} \psi_{,r} g^{rr} \sqrt{-g} d\Omega,$$
(8)

where S represents the boundary two-dimensional sphere centered on the black hole, r is the radial ordinate, and g^{rr} is the radius-radius component of the contravariant metric tensor, g denotes the determinant of the metric of the black hole under considering, and $d\Omega$ is the product of the differentials of angels.

Now, we consider the first boundary condition, i.e., the asymptotic boundary condition at large distances. Since the medium is homogeneous there, we can take $n_{\infty} = h_{\infty} = 1$ in appropriate units and restore n_{∞} in some final results. The asymptotic boundary condition in rectangular coordinates is

$$\psi = u_{\mu}x^{\mu} = -u_{\infty}^{0}t + \mathbf{u}_{\infty} \cdot \mathbf{x}.$$
(9)

In spherical coordinates, it becomes

$$\psi = -u_{\infty}^{0}t + u_{\infty}r[\cos\theta\cos\theta_{0} + \sin\theta\sin\theta_{0}\cos(\phi - \phi_{0})].$$
⁽¹⁰⁾

for $r \to \infty$. The asymptotic three-velocity vector, v_{∞} , can point in an arbitrary direction (θ_0, ϕ_0) . According to special relativity at infinity, there exists the following relation

$$u_{\infty}^{\mu} = (u_{\infty}^{0}, \mathbf{u}_{\infty}) = (1 - v_{\infty}^{2})^{-1/2} (1, \mathbf{v}_{\infty}).$$
(11)

Here we consider the accreted flow which moves into the black hole. Since the accreted flow is continuous at every point in space-time, and no disruption or infinity should exists in the flow, the physical quantities of the flow, such as the particle number density n and the enthalpy h, should be finite everywhere, including at the event horizon.

III. ACCRETION ONTO A MOVING CHARGED DILATON BLACK HOLE

In this section, we consider accretion onto a charged dilaton black hole which moves at a constant velocity through the medium. In heterotic string theory, the dilaton has a linear coupling to the square of electromagnetic field tensor, so every solution with nonzero electromagnetic field tensor must have a nonconstant dilaton. Thus the Reissner-Nordström solution, which describes charged black holes in general relativity, is not even an approximate solution in string theory. A solution for a statically charged black hole in the string theory was found to take the form [38, 39]

$$ds^{2} = -\left(1 - \frac{2M}{r}\right)dt^{2} + \left(1 - \frac{2M}{r}\right)^{-1}dr^{2} + r(r-a)(d\theta^{2} + \sin^{2}\theta d\phi^{2}),$$
(12)

where $a = Q^2 e^{2\phi_0}/M$ with ϕ_0 the asymptotic constant value of the dilaton, M the gravitational mass and Q the charge of black hole, respectively. The event horizon are localized at r = 2M. When r = a, the area of the sphere goes to zero and the surface is singular. The transition between black holes and naked singularities occurs at $\hat{Q} = Q_{\text{max}} \equiv \sqrt{2}e^{-\phi_0}M$. For $Q < Q_{\text{max}}$, the singularity is enclosed by the event horizon.

Considering the metric (12), Eq. (7) implies

$$-\frac{1}{1-\frac{2M}{r}}\frac{\partial^2\psi}{\partial t^2} + \frac{1}{r(r-a)}\frac{\partial}{\partial r}\left[r(r-a)\left(1-\frac{2M}{r}\right)\frac{\partial\psi}{\partial r}\right] + \frac{1}{r(r-a)}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\left(\sin\theta\frac{\partial\psi}{\partial\theta}\right) + \frac{1}{r(r-a)\sin^2\theta}\frac{\partial^2\psi}{\partial\phi^2} = 0.(13)$$

Taking into account the asymptotic boundary condition: $\psi = -u_{\infty}^{0}t + u_{\infty}r[\cos\theta\cos\theta_{0} + \sin\theta\sin\theta_{0}\cos(\phi - \phi_{0})]$ for $r \to \infty$, the general formula of ψ can be assumed to take form $\psi = -u_{\infty}^{0}t + u(r, \theta, \phi)$, which also satisfies the stationary flow condition that the gradient of ψ must be independent of time. Inserting ψ in the Eq. (13) with the general form, yields

$$\frac{\partial}{\partial r} \left[r(r-a) \left(1 - \frac{2M}{r} \right) \frac{\partial u}{\partial r} \right] + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial u}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 u}{\partial \phi^2} = 0.$$
(14)

This is the differential equation which the spatial component $u(r, \theta, \phi)$ of ψ should satisfy. If assuming u = $R(r)\Theta(\theta)\Phi(\phi)$, then the functions R(r), $\Theta(\theta)$, and $\Phi(\phi)$, respectively, satisfies the differential equations below

$$\frac{d^2\Phi}{d\phi^2} + m^2\Phi = 0,\tag{15}$$

$$\frac{1}{\sin\theta} \frac{d}{d\theta} \left(\sin\theta \frac{d\Theta}{d\theta} \right) - \frac{m^2 \Theta}{\sin^2\theta} + l(l+1)\Theta = 0, \tag{16}$$

and

$$\frac{d}{dr}\left[r(r-a)\left(1-\frac{2M}{r}\right)\frac{dR}{dr}\right] - l(l+1)R = 0.$$
(17)

Obviously, Eq. (17) is complex and difficult to solve directly. The key here is to introduce a new variable to simplify it. After some attempts, we found a variable, $\xi = \frac{2}{2M-a}r - \frac{2M+a}{2M-a}$, to simplify Eq. (17) as

$$(1-\xi^2)\frac{d^2R}{d\xi^2} - 2\xi\frac{dR}{d\xi} + l(l+1)R = 0,$$
(18)

which is Legendre equation. The general solutions for the differential equations (15), (16), and (18) can take the following forms, respectively

$$\Phi(\phi) = C\cos m\phi + D\sin m\phi \qquad m = 0, 1, 2, \cdots,$$
(19)

where C, and D are constants;

$$\Theta(\theta) = P_l^m(\cos\theta) \quad l, m = 0, 1, 2, \cdots,$$
⁽²⁰⁾

where $P_l^m(z)$ is the associated Legendre function; and

$$R(\xi) = AP_l(\xi) + BQ_l(\xi) \quad l = 0, 1, 2, \cdots,$$
(21)

where A and B are constants, $P_l(z)$ is the Legendre polynome, and $Q_l(z)$ is linearly independent from $P_l(z)$ and belongs to the second kind of Legendre function. Then the general formula of ψ for charged dilaton black hole (12) is

$$\psi = -u_{\infty}^{0}t + \sum_{l,m} [A_{lm}P_{l}(\xi) + B_{lm}Q_{l}(\xi)]Y_{lm}(\theta,\phi) \qquad l,m = 0, 1, 2, \cdots,$$
(22)

where A_{lm} , B_{lm} are constants needed to be determined from boundary conditions, $Y_{lm}(\theta, \phi)$ are spheric harmoics made up from $\Phi(\phi)$ and $\Theta(\theta)$. According to the boundary conditions, we can determine the specific form of Eq. (22). Since the particle number density is finite at the event horizon, we first obtain the four-velocities as

$$nu_t = -u_\infty^0,\tag{23}$$

$$nu_{r} = \frac{2}{2M - a} \sum_{l,m} \left[A_{lm} P_{l}^{'}(\xi) + B_{lm} Q_{l}^{'}(\xi) \right] Y_{lm}(\theta, \phi),$$
(24)

$$nu_{\theta} = \sum_{l,m} \left[A_{lm} P_l(\xi) + B_{lm} Q_l(\xi) \right] \frac{\partial Y_{lm}(\theta, \phi)}{\partial \theta}, \tag{25}$$

$$nu_{\phi} = \sum_{l,m} \left[A_{lm} P_l(\xi) + B_{lm} Q_l(\xi) \right] \frac{\partial Y_{lm}(\theta, \phi)}{\partial \phi}.$$
 (26)

where the prime denotes the derivative with respect to ξ . Then, substituting the corresponding quantities above in the normalization condition, we get

$$n^{2} = \frac{(nu_{t})^{2}}{1 - \frac{2M}{r}} - \left(1 - \frac{2M}{r}\right)(nu_{r})^{2} - \frac{(nu_{\theta})^{2}}{r(r-a)} - \frac{(nu_{\phi})^{2}}{r(r-a)\sin^{2}\theta}.$$
(27)

Thinking of the limit behaviour of the Legendre functions and the finiteness of n at the event horizon where $\xi = 1$, we have

$$n^{2} \to \frac{1}{1 - \frac{2M}{r}} \left\{ (u_{\infty}^{0})^{2} - \left[\frac{1}{4M} \sum_{l,m} B_{lm} Y_{lm}(\theta, \phi) \right]^{2} \right\}.$$
 (28)

Observing the right hand side of the equation above, we find that the denominator in the external part will tend to zero when closing to the event horizon, the internal part must be equal zero at the same time to ensure that the particle number density is finite at the event horizon. Therefore we find

$$B_{00}Y_{00} = 4Mu_{\infty}^0 \tag{29}$$

and all other B_{lm} s are zero. Therefore, Eq. (22) reduces to

$$\psi = -u_{\infty}^{0}t - 2Mu_{\infty}^{0}\ln\frac{r-2M}{r-a} + \sum_{l,m}A_{lm}P_{l}(\xi)Y_{lm}(\theta,\phi),$$
(30)

where A_{lm} can now be determined from the asymptotic boundary conditions in Eq. (11). Without loss of generality, we focus on the case of $\theta = 0$ which means the accretion flow at infinity moving towards the north pole of the coordinate system. Then the asymptotic boundary condition becomes

$$\psi = -u_{\infty}^{0}t + u_{\infty}r\cos\theta. \tag{31}$$

Comparing it with Eq. (30), yields

$$A_{10} = \frac{2M - a}{2} u_{\infty},$$
(32)

and all other A_{lm} s are zero. Finally, we find the solution ψ takes the form

$$\psi = -u_{\infty}^{0}t - 2Mu_{\infty}^{0}\ln\frac{r-2M}{r-a} + u_{\infty}\left(r - \frac{2M+a}{2}\right)\cos\theta.$$
(33)

Using Eq. (4) and the final expression of ψ in Eq. (33), the four-velocity of accreted flow is calculated as

$$nu_t = -u_\infty^0,\tag{34}$$

$$nu_r = -\frac{2Mu_{\infty}^0(2M-a)}{(r-a)(r-2M)} + u_{\infty}\cos\theta,$$
(35)

$$nu_{\theta} = -u_{\infty} \left(r - \frac{2M + a}{2} \right) \sin \theta, \tag{36}$$

$$nu_{\phi} = 0. \tag{37}$$

The velocity field is a function of the coordinates r and θ , meaning that it possesses axial symmetry. Substituting the components of four-velocity, Eqs. (34)-(37), into Eq. (27), we obtain the particle number density

$$n^{2} = (u_{\infty}^{0})^{2} \left\{ \frac{1}{1 - \frac{2M}{r}} \left[1 - \left[\frac{2M(2M-a)}{r(r-a)} \right]^{2} \right] \right\} - (u_{\infty})^{2} \left\{ \left(1 - \frac{2M}{r} \right) \cos^{2}\theta + \frac{r}{r-a} \left(1 - \frac{2M+a}{2r} \right)^{2} \sin^{2}\theta \right\} + (u_{\infty}^{0}u_{\infty}) \frac{4M(2M-a)}{r(r-a)} \cos\theta.$$
(38)

From this equation, we can judge that the particle number destiny is really finite at event horizon. By solving the corresponding equations, we can determine that the stagnation point at which the velocity is zero lies at $\theta = 0$ (directly downstream) and at radius

$$r = \frac{1}{2} \left\{ 2M + a + \left[(2M + a)^2 - 8M \left(a + \frac{a - 2M}{v_{\infty}} \right) \right]^{1/2} \right\}.$$
 (39)

With Eq. (8), we find the particle number accretion rate is given by

$$\dot{N} = 8\pi M (2M-a) u_{\infty}^0 n_{\infty},\tag{40}$$

where we have restored n_{∞} . The right hand side of Eq. (40) is just the area of the black hole multiplied by particle number destiny and Lorentz factor for the accreted flow at infinity. Since *a* depends on the magnetic charge and the dilaton, which means that these parameters play important parts in the process of accretion. If a = 0, all the results obtained here reduce to the case of Schwarzschild black hole.

IV. CONCLUSIONS AND DISCUSSIONS

We have obtained an analytic solution for accretion of a gaseous medium with a adiabatic equation of state $(P = \rho)$ onto a charged dilaton black hole which moves at a constant velocity. We have derived the four-velocity of accreted flow and found that it possesses axial symmetry. We have determined the particle number density and the location of the stagnation point. We have presented the accretion rate which depends on the the mass of black hole and the parameter *a* in string theory. Since *a* depends on the magnetic charge and the dilaton, which indicates that these parameters play important parts in the process of accretion onto the moving charged dilaton black hole. The results obtained here provides a possible observational effect to test the string theory and may help us to get deeper understanding of the behavior of accreted flow near the event horizon of black hole.

This work is supported in part by Hebei Provincial Natural Science Foundation of China (Grant No. A2014201068 and A2021201034).

- F. Hoyle and R. A. Lyttleton, "The effect of interstellar matter on climatic variation," in Mathematical Proceedings of the Cambridge Philosophical Society, vol. 35, pp. 405–415, Cambridge Univ Press, 1939.
- [2] R. A. Lyttleton and F. Hoyle, "The evolution of the stars," The Observatory, vol. 63, pp. 39–43, Feb. 1940.
- [3] H. Bondi and F. Hoyle, "On the mechanism of accretion by stars," Mon.Not.Roy.Astron.Soc., vol. 104, p. 273, 1944.
- [4] H. Bondi, "On spherically symmetrical accretion," Mon.Not.Roy.Astron.Soc., vol. 112, p. 195, 1952.
- [5] F. C. Michel, "Accretion of matter by condensed objects," Astrophysics and Space Science, vol. 15, no. 1, pp. 153–160, 1972.
- [6] M. Begelman, "Accretion of v > 5/3 gas by a schwarzschild black hole," Astronomy and Astrophysics, vol. 70, p. 583, 1978.
- [7] E. Malec, "Fluid accretion onto a spherical black hole: Relativistic description versus Bondi model," *Phys. Rev. D*, vol. 60, p. 104043, 1999.
- [8] E. Babichev, V. Dokuchaev, and Y. Eroshenko, "Black hole mass decreasing due to phantom energy accretion," *Phys. Rev. Lett.*, vol. 93, p. 021102, 2004.
- M. E. Rodrigues and E. L. B. Junior, "Spherical Accretion of Matter by Charged Black Holes on f(T) Gravity," Astrophys. Space Sci., vol. 363, no. 3, p. 43, 2018.
- [10] E. Contreras, A. Rincón, and J. M. Ramírez-Velasquez, "Relativistic dust accretion onto a scale-dependent polytropic black hole," *Eur. Phys. J. C*, vol. 79, no. 1, p. 53, 2019.
- [11] G. Abbas and A. Ditta, "Accretion onto a charged Kiselev black hole," Mod. Phys. Lett. A, vol. 33, no. 13, p. 1850070, 2018.
- [12] J. Zheng, R. Ye, J. Chen, and Y. Wang, "Accretion onto RN-AdS black hole surrounded by quintessence," Gen. Rel. Grav., vol. 51, no. 9, p. 123, 2019.
- [13] S. Yang, C. Liu, T. Zhu, L. Zhao, Q. Wu, K. Yang, and M. Jamil, "Spherical Accretion Flow onto General Parameterized Spherically Symmetric Black Hole Spacetimes," *Chin. Phys. C*, vol. 45, no. 1, p. 015102, 2021.
- [14] M. Umar Farooq, A. K. Ahmed, R.-J. Yang, and M. Jamil, "Accretion on high derivative asymptotically safe black holes," *Chin. Phys. C*, vol. 44, no. 6, p. 065102, 2020.
- [15] K. Nozari, M. Hajebrahimi, and S. Saghafi, "Quantum Corrections to the Accretion onto a Schwarzschild Black Hole in the Background of Quintessence," *Eur. Phys. J. C*, vol. 80, no. 12, p. 1208, 2020.
- [16] G. Panotopoulos, A. Rincon, and I. Lopes, "Accretion of matter and spectra of binary X-ray sources in massive gravity," Annals Phys., vol. 433, p. 168596, 2021.
- [17] S. Iftikhar, "Accretion onto some singularity-free black holes," Int. J. Mod. Phys. A, vol. 35, no. 13, p. 2050062, 2020.
- [18] C. Gao, X. Chen, V. Faraoni, and Y.-G. Shen, "Does the mass of a black hole decrease due to the accretion of phantom energy," *Phys. Rev.*, vol. D78, p. 024008, 2008.
- [19] A. J. John, S. G. Ghosh, and S. D. Maharaj, "Accretion onto a higher dimensional black hole," Phys. Rev. D, vol. 88, no. 10, p. 104005, 2013.
- [20] A. Ganguly, S. G. Ghosh, and S. D. Maharaj, "Accretion onto a black hole in a string cloud background," Phys. Rev. D, vol. 90, no. 6, p. 064037, 2014.
- [21] P. Mach and E. Malec, "Stability of relativistic Bondi accretion in Schwarzschild-(anti-)de Sitter spacetimes," Phys. Rev. D, vol. 88, no. 8, p. 084055, 2013.
- [22] G. M. Kremer and L. C. Mehret, "Post-Newtonian spherically symmetrical accretion," Phys. Rev. D, vol. 104, no. 2, p. 024056, 2021.
- [23] E. Tejeda and A. Aguayo-Ortiz, "Relativistic wind accretion on to a Schwarzschild black hole," Mon. Not. Roy. Astron. Soc., vol. 487, no. 3, pp. 3607–3617, 2019.
- [24] R. Yang, "Quantum gravity corrections to accretion onto a Schwarzschild black hole," Phys. Rev. D, vol. 92, no. 8, p. 084011, 2015.
- [25] R.-J. Yang, H. Gao, Y. Zheng, and Q. Wu, "Effects of Lorentz breaking on the accretion onto a Schwarzschild-like black hole," Commun. Theor. Phys., vol. 71, no. 5, pp. 568–572, 2019.
- [26] M. Jamil, M. A. Rashid, and A. Qadir, "Charged Black Holes in Phantom Cosmology," Eur. Phys. J. C, vol. 58, pp. 325– 329, 2008.
- [27] R. Yang, "Constraints from accretion onto a Tangherlini–Reissner–Nordstrom black hole," Eur. Phys. J. C, vol. 79, no. 4, p. 367, 2019.
- [28] P. Papadopoulos and J. A. Font, "Relativistic hydrodynamics around black holes and horizon adapted coordinate systems," *Phys. Rev. D*, vol. 58, p. 024005, 1998.
- [29] J. A. Font, J. M. Ibanez, and P. Papadopoulos, "Nonaxisymmetric relativistic Bondi-Hoyle accretion onto a Kerr black hole," Mon. Not. Roy. Astron. Soc., vol. 305, p. 920, 1999.
- [30] O. Zanotti, C. Roedig, L. Rezzolla, and L. Del Zanna, "General relativistic radiation hydrodynamics of accretion flows. I: Bondi-Hoyle accretion," Mon. Not. Roy. Astron. Soc., vol. 417, pp. 2899–2915, 2011.

- [31] F. D. Lora-Clavijo, A. Cruz-Osorio, and E. Moreno Méndez, "Relativistic Bondi-hoyle-lyttleton Accretion Onto a Rotating Black Hole: Density Gradients," Astrophys. J. Suppl., vol. 219, no. 2, p. 30, 2015.
- [32] L. I. Petrich, S. L. Shapiro, and S. A. Teukolsky, "Accretion onto a moving black hole: An exact solution," Phys. Rev. Lett., vol. 60, pp. 1781–1784, 1988.
- [33] L. Jiao and R.-J. Yang, "Accretion onto a moving Reissner-Nordström black hole," JCAP, vol. 09, p. 023, 2017.
- [34] Y. Liu and S. N. Zhang, "Exact solutions for shells collapsing towards a pre-existing black hole," Phys. Lett. B, vol. 679, pp. 88–94, 2009.
- [35] S.-X. Zhao and S.-N. Zhang, "Exact solutions for spherical gravitational collapse around a black hole: the effect of tangential pressure," Chin. Phys. C, vol. 42, no. 8, p. 085101, 2018.
- [36] P. Mach and A. Odrzywo, "Accretion of Dark Matter onto a Moving Schwarzschild Black Hole: An Exact Solution," Phys. Rev. Lett., vol. 126, no. 10, p. 101104, 2021.
- [37] P. Rioseco and O. Sarbach, "Accretion of a relativistic, collisionless kinetic gas into a Schwarzschild black hole," Class. Quant. Grav., vol. 34, no. 9, p. 095007, 2017.
- [38] G. W. Gibbons and K.-i. Maeda, "Black Holes and Membranes in Higher Dimensional Theories with Dilaton Fields," Nucl. Phys. B, vol. 298, pp. 741–775, 1988.
- [39] D. Garfinkle, G. T. Horowitz, and A. Strominger, "Charged black holes in string theory," Phys. Rev. D, vol. 43, p. 3140, 1991. [Erratum: Phys.Rev.D 45, 3888 (1992)].
- [40] V. Moncrief, "Stability of stationary, spherical accretion onto a Schwarzschild black hole," Astrophys. J., vol. 235, pp. 1038– 1046, Feb. 1980.