

ON MERGE TREES AND DISCRETE MORSE FUNCTIONS ON PATHS AND TREES

JULIAN BRÜGGEMANN

ABSTRACT. In this work we answer an open question asked by Johnson–Scoville. We show that each merge tree is represented by a discrete Morse function on a path. Furthermore, we present explicit constructions for two different but related kinds of discrete Morse functions on paths that induce any given merge tree. A refinement of the used methods allows us to define notions of equivalence of discrete Morse functions on trees which give rise to a bijection between equivalence classes of discrete Morse functions and isomorphism classes of certain labeled merge trees. We also compare our results to similar ones from the literature, in particular to work by Curry.

CONTENTS

1. Introduction	1
2. Conflict of Interest Statement	5
3. Preliminaries	5
3.1. Merge Trees	6
3.2. Generic Properties and Equivalences of DMFs	12
4. Construction of the Induced Index-Ordered DMF	21
4.1. The Index Morse Order	21
4.2. The Simplex Order	24
4.3. The Induced Index-Ordered DMF	29
5. The Sublevel-Connected DMF	31
6. Relationships Between Merge Trees and DMFs on Paths	33
7. Further Directions and Possible Applications	41
Appendix A. Step-by-Step Construction of the Induced Index-Ordered DMF	42
A.1. Step-by-Step Construction	43
A.2. Properties of the Step-by-Step Construction	44
A.3. The Constructed DMF Represents T	50
A.4. An Example for the Step-by-Step Construction	53
References	59

1. INTRODUCTION

Discrete Morse theory is a combinatorial version of the classical smooth Morse theory. It was originally developed by Forman in [For98].

In discrete Morse theory, topological properties of simplicial complexes X are analyzed by considering discrete Morse functions $f: X \rightarrow \mathbb{R}$. These topological properties can in turn be used to obtain cell decompositions of X with fewer cells. A good introduction to the topic is found in [For01].

Merge trees are used in Morse theory in order to keep track of the development of connected components of sublevel sets $X_a := f^{-1}(-\infty, a]$ of a given Morse function

$f: X \rightarrow \mathbb{R}$. Since the sublevel sets form a filtration of X , merge trees can be seen as a combinatorial description of the persistent connectivity of X . In particular, every branching in the induced merge tree $M(X, f)$ corresponds to a pair of connected components of a sublevel set $X_{a-\varepsilon}$ that merge to one connected component in a sublevel complex X_a of higher level.

Initially, merge trees were introduced to topological data analysis as an approximation to the Reeb graph, respectively contour tree, in [CSA03]. The Reeb graph is a graph that keeps track of the connected components of level sets of any given filtered manifold. In applications, the data set is often interpreted as a sampling of the graph of a function rather than a more general manifold, which is why the Reeb graph is often actually a tree, the so-called contour tree.

Among other implementations of techniques of smooth Morse theory, computational methods for the Reeb graph have originally been introduced in [SKK91] in order to handle surfaces embedded in 3D with the help of computers. Later on, several ways to compute and apply contour trees of data sets have been discussed in many articles, e.g. [KK94], [vKvOB⁺97], [TV98], and [CSA03]. Among other applications, merge trees have been used in visualization, e.g. in [OHWS13], [WBP07], [OHW⁺17], and [YWM⁺19]. Surveys about applications of merge trees and other concepts in visualization can be found in [HLH⁺16] and [LMW⁺16]. Furthermore, a certain version of chiral merge trees has been used in [Bar19] to analyze asymmetries of time series.

We focus on the more structural and theoretical side of merge trees, in particular the connection to discrete Morse theory. We consider a specific construction for merge trees induced by discrete Morse functions on trees which was introduced in [JS20]. We use this construction to gain a better understanding of the set of discrete Morse functions, the set of merge trees, and the relationship between the two.

Similar work has been done in [Cur19] for the relationship between Morse-like functions on the interval and a related version of merge trees. Furthermore, our work is in some sense similar to parts of [CDG⁺21], with the difference being that the authors of [CDG⁺21] consider the relationship between merge trees and their induced barcodes instead. It seems reasonable to adapt the following terms from both of these articles and say that we consider a different instance of the “fiber of the persistence map”, respectively a different instance of the “inverse problem” of the “persistence map”. These names refer to the fact that merge trees and barcodes are invariants of filtered spaces rather than just spaces. In this context, the name “persistent” became popular due to persistent homology as it appears in topological data analysis. Similar to the authors of [Cur19] and [CDG⁺21], we are interested in finding out what information is lost by considering our invariant at hand, the induced merge tree, instead of the given data, in our case a discrete Morse function, and how this information might be re-obtained. Such knowledge might be helpful to investigate the space of merge trees and the space of discrete Morse functions in future work. Furthermore, a good understanding of the fiber of the persistence map is useful for topological data analysis because it hints at features which might be lost due to the chosen invariant. Moreover, insights about the inverse problem might be helpful to enhance the chosen invariant in a way such that it preserves certain desired features of the data set.

We respond to an open question asked in [JS20] by showing that every merge tree is represented by a discrete Morse function (dMf). In particular, for any given merge

tree we construct a dMf on a path as a representative of the isomorphism class defined by said merge tree:

Theorem 6.5. *Let T be a merge tree. Then there is a path P such that $T \cong M(P, f_{io}) \cong M(P, f_{sc})$ holds as merge trees where f_{io} denotes the induced index-ordered dMf (Definition 4.21) and f_{sc} denotes the sublevel-connected dMf (Definition 5.5) on P .*

In particular, the discrete Morse function from Theorem 6.5 can be chosen to be index-ordered (Definition 3.1) or sublevel-connected (Definition 3.31).

The main tool for the construction is the corresponding Morse order (Definition 4.3), that is, the index Morse order (Definition 4.3) or the sublevel-connected Morse order (Definition 3.31) on the nodes of a given merge tree T . The index Morse order defines leaf nodes to be strictly less than inner nodes. Among leaf nodes and among inner nodes, the index Morse order is defined by using a twisted version of length-lexicographical order on the set of path words (Lemma 4.17) that correspond to the respective nodes. The path words are defined by the chirality of the nodes of the shortest path from the root to the corresponding node. For the sublevel-connected Morse order (Definition 5.1) we do not artificially distinguish between leaf nodes and inner nodes.

We use the index Morse order (Definition 4.3) to define the index Morse labeling (Definition 4.7) on the nodes of T . Together with the simplex order (Definition 4.11), which establishes a correspondence (Proposition 4.19) between the nodes of T and the simplices of a path P , the index Morse labeling defines the induced index-ordered discrete Morse function on said path P .

In Section 3.2 we introduce several kinds of equivalence relations on the sets of discrete Morse functions with only critical cells on paths and trees. These equivalence relations allow us to identify equivalence classes of discrete Morse functions with isomorphism classes of Morse labeled merge trees:

Theorem 6.4. *The induced labeled merge tree $M(-, -)$ and the induced dMf Φ define maps $M(-, -): DMF_P^{crit} \leftrightarrow MIT: \Phi$ that are inverse to each other in the sense that:*

- (1) *for a dMf (P, f) with only critical cells, the dMf $\Phi(M(P, f), \lambda_f)$ is symmetry-equivalent to (P, f) , and*
- (2) *for an Ml tree (T, λ) , the Ml tree $M(\Phi T, f_\lambda)$ is isomorphic to (T, λ) .*

Theorem 6.6. *The induced labeled merge tree $M(-, -)$ and the induced dMf Φ define maps $M(-, -): DMF_X^{crit} \leftrightarrow MIT: \Phi$ that are inverse to each other in the sense that:*

- (1) *for any dMf (X, f) with only critical cells, the dMf $\Phi(M(X, f), \lambda_f)$ is cm-equivalent to (X, f) , and*
- (2) *for any Ml tree (T, λ) , the Ml tree $M(\Phi T, f_\lambda)$ is isomorphic to (T, λ) .*

The construction of the discrete Morse function induced by a Morse labeling is similar to the construction of functions on the interval in [Cur19]. In particular, Theorem 6.4 is very similar to the result [Cur19, Prop 6.11]. The use of Morse labelings in this work basically plays the role of the function $\pi: T \rightarrow \mathbb{R}$ from [Cur19]. Moreover, the simplex order is almost the same as the use of chirality in [Cur19, Lem 6.4]. But in this work, Morse orders, and in turn Morse labelings, have to satisfy a certain compatibility with the chirality, that is, property (2) of Definition 3.17.

The notion of merge trees we use originates from [JS20, Def 5] and differs from the one used in [Cur19]: A priori, merge trees T in the sense of [JS20] do not carry a height function $T \rightarrow \mathbb{R}$ as part of their data. Instead, the two children of each node have a chirality assigned to them as part of the tree's data. This means that for any two child nodes of the same parent node, it is specified as part of data which is the right and which is the left child. This version of chirality is also canonically assigned to merge trees induced by discrete Morse functions. In contrast, the chirality of chiral merge trees in the sense of [Cur19] arises from a chosen orientation on the interval. We obtain a similar correspondence between the chirality of merge trees and orientations on paths using the simplex order, Definition 4.11. Apart from these differences, the notion of merge trees in the sense of [JS20] is closely related to the one from [Cur19]. Chirality in the sense of [JS20] is a specific version of the notion of chirality used in [Cur19]. In order to see this, we show that the construction of the merge tree $M(X, f)$ induced by a discrete Morse function f in the sense of [JS20] can be modified, see Proposition 3.20, to obtain a function $T \rightarrow \mathbb{R}$ from f , similarly to [Cur19]. This gives rise to the notion of Morse labelings, Definition 3.19, and Morse orders, Definition 4.3. It turns out that the use of chirality in [JS20] assumes a certain compatibility between Morse orders and the simplex order, whereas the use of chirality in [Cur19] does not. As a result, the induced merge tree in the sense of [Cur19] distinguishes between symmetry equivalences, Definition 3.43, whereas the induced merge tree in the sense of [JS20] identifies symmetry-equivalent discrete Morse functions with each other, see Proposition 3.49. We discuss this in a bit more detail at the end of Section 6.

In said discussion, we mention the notion of CMI trees, see Definition 6.9, which is as objects basically the same as the notion of merge trees from [CDG⁺21, Def 2.2]. However, the notion of combinatorial equivalence of labeled merge trees from [CDG⁺21, Def 2.6] corresponds to a non-chiral version of shuffle equivalence Definition 3.24 rather than an equivalence of the induced persistent set as in [Cur19, 5.1], which is more similar to an isomorphism of MI trees Definition 3.23. The chiral merge trees from [Bar19, Def 2.1] are as objects also very similar to the chiral merge trees from [Cur19] and, thus, differ similarly from our notion of labeled merge trees.

The aforementioned versions of labeled merge trees all have in common that their labelings need to be compatible with some other data inherent to the merge tree. In contrast to that, the labelings from [YWM⁺19] can be quite arbitrary and might even assign multiple labels to a single node. Hence, our MI trees a priori seem to be unrelated to the notion of labeled merge trees from [YWM⁺19].

The step-by-step approach (Appendix A) gives an alternative construction of the path P and the induced index-ordered discrete Morse function on P by induction over the nodes of T from highest to lowest with respect to the index Morse order.

Theorem A.9 shows that the two constructions of the induced index-ordered discrete Morse functions yield the same result.

We illustrate the step-by-step construction in an example in Appendix A.4.

Acknowledgements. The author would like to thank Benjamin Johnson and Nicholas A. Scoville for the suggested questions in their article [JS20] which led to this project. Furthermore, the author would like to thank the mathematical faculty of Ruhr-University of Bochum, especially the chair for topology, for the great scientific environment in which this endeavor was started. In addition to that the author thanks Max Planck

Institute for Mathematics for the great scientific environment in which this project was finished.

Most notably the author thanks his advisor, Viktoriya Ozornova, for her advice and the many helpful discussions which increased the quality of this article. Last but not least, the author thanks the anonymous referees for the helpful and detailed feedback.

2. CONFLICT OF INTEREST STATEMENT

The author states that there is no conflict of interest.

3. PRELIMINARIES

We consider discrete Morse functions (dMf) on trees. Recall that trees are finite acyclic simple graphs. Furthermore, simple graphs are 1-dimensional simplicial complexes. Where feasible, we introduce the preliminaries in the broader generality they are usually defined in, rather than in the lesser generality we actually need for this work. We adapt most notations and conventions from [JS20]. For simplicity, we assume all trees in this article to be non-empty. Similar to [JS20], we assume the dMfs to fulfill certain generic properties. In detail this means the following :

Definition 3.1. Let X be a simplicial complex. A map $f: X \rightarrow \mathbb{R}$ is called a **discrete Morse function (dMf)** if it fulfills the following properties for any pair of simplices $\sigma, \tau \in X$:

- (i) $\sigma \subseteq \tau \Rightarrow f(\sigma) \leq f(\tau)$ (weakly increasing)
- (ii) f is at most $2 - 1$
- (iii) $f(\sigma) = f(\tau) \Rightarrow (\sigma \subset \tau \vee \tau \subset \sigma)$ (matching)

Simplices on which f is $1 - 1$ are called **critical**. Simplices which belong to the preimage of the same value are called **matched**. The set of critical simplices is denoted by $\text{Cr}(f)$. Values of critical simplices under f are called **critical values** of f . A dMf is called **index-ordered** if for arbitrary critical simplices σ, τ the following holds: If $\dim(\sigma)$ is smaller than $\dim(\tau)$, then $f(\sigma) < f(\tau)$ holds.

Remark 3.2. The definition given above is not the most general definition of dMfs but rather assumes several generic properties. This means that any dMf in the sense of [For98] can be modified by a Forman equivalence (see [JS20, Def 4.1]) to fulfill these properties, that is, without changing the induced Morse matching. As usual in the context of dMfs, we write $f: X \rightarrow \mathbb{R}$ although the map f is actually defined on the face poset of X .

In [NTT18], a similar generic property is used to analyze flow paths induced by dMfs. The notion of faithful dMfs as defined in [NTT18, Def 2.9] is almost the same as index-ordered dMfs in this article. The only difference is that for index-ordered dMfs matched cells have the same value, whereas for faithful dMfs the values of the matched boundary simplices are higher than the values on the corresponding matched co-boundary simplices.

Furthermore, since simple graphs and in particular trees are examples of simplicial complexes, the definition of dMfs can be applied to them as well.

Remark 3.3. Because f maps $\text{Cr}(f)$ injectively into a totally ordered set, it induces a total order on $\text{Cr}(f)$. We refer to this induced order whenever we speak of simplices being **ordered by f** .

Notation 3.4. We use the following conventions regarding notation:

- We depict dMfs on graphs by labeling the graph with the values of the dMf.
- Let G be a graph and let v be a node of G . By $G[v]$ we denote the connected component of G which contains v .

Definition 3.5. • Let X be a simplicial complex, $f: X \rightarrow \mathbb{R}$ a dMf and $a \in \mathbb{R}$. The **sublevel complex** of level a , denoted by X_a^f , is defined by $X_a^f := \{\sigma \in X \mid f(\sigma) \leq a\}$. If the referred dMf f is clear from the context, we drop the superscript f from the notation.

- The ordered critical values $c_0 < c_1 < \dots < c_m$ induce a chain of sublevel complexes $X_{c_0}^f \subsetneq X_{c_1}^f \subsetneq \dots \subsetneq X_{c_m}^f$. Within this chain, we refer by $X_{c_i-\varepsilon}^f$ to the complex that immediately precedes X_{c_i} .

Remark 3.6. The given definition of sublevel complexes differs from the standard one used in the literature. We make use of the fact that a dMf f being weakly increasing implies that X_a^f as defined above is already a subcomplex of X . If we wanted to consider the general definition of dMfs as introduced in [For98], we would have to work with the smallest supercomplex of X_a^f in X instead. Taking the smallest supercomplex corresponds to additionally including all faces of simplices of X_a^f to make X_a^f a simplicial complex.

Lemma 3.7. *Let X be a finite simplicial complex and let $f: X \rightarrow \mathbb{R}$ be a dMf. Then f attains its minimum on a critical 0-simplex. Furthermore, the statement also holds for the restriction to any connected component of sublevel complexes.*

Proof. Because f is weakly increasing, no higher simplex τ can have a value which is strictly smaller than the value of f on any of τ 's boundary 0-simplices. Thus, the minimum is attained on a 0-simplex.

Let σ be a 0-simplex such that $f(\sigma)$ is the minimum of f . We prove by contradiction that σ is critical: If σ is not critical, then there is a 1-simplex τ such that σ is a boundary 0-simplex of τ and $f(\tau) = f(\sigma)$ holds. Let σ' be the other boundary 0-simplex of τ . Then $f(\sigma') < f(\tau) = f(\sigma)$ holds because f is by assumption weakly increasing and at most 2-1. This is a contradiction to f attaining its minimum on σ , so f attains its minimum on a critical 0-simplex.

Since all three properties of Definition 3.1 are inherited by arbitrary subcomplexes of X , the statement is also true for connected components of sublevel complexes. \square

Remark 3.8. The analogous statement for the maximum of a dMf f on arbitrary 1-simplices is false, as the following example shows:

$$\begin{array}{ccccccc} & & 3 & 3 & 0 & 2 & 1 \\ & & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

Here, the maximum is attained on a pair of matched simplices.

3.1. Merge Trees. We briefly recapture preliminaries about merge trees as they are explained in [JS20]. The basic idea is that merge trees keep track of the chronological development of the connected components of sublevel sets. We adapt the point of view of [JS20], that is, we consider them ‘upside down’. Thus, the children will appear above their parent node. Afterwards we introduce additional structure that dMfs induce on their corresponding merge trees and consider notions of equivalence which arise from that structure.

Definition 3.9 (Merge Tree). A *merge tree* is a full rooted chiral binary tree T . In detail this means that T is a rooted tree fulfilling the properties of being binary and full, and that T has the extra datum of being chiral:

full binary: Each node of T has either zero or two children.

chiral: Each child node in T carries the extra datum, the so-called *chirality*, of being a left or a right child.

Morphisms of merge trees are morphisms of rooted binary trees which are compatible with the chirality.


For rooted trees T we use the notions of *subtrees*, *ancestors* and *descendants* as they are commonly used in computer science.

Definition 3.10. For any node p of T , the *descendants* of p are defined inductively: A node c is a descendant of p if the parent node of c is a descendant of p or p itself.

A *subtree* of T is a subgraph of T that consists of exactly all of the descendants of some node p of T .

For a node p of T we call all nodes which lie on the shortest path between p and the root, including the root, the *ancestors* of p .

Notation 3.11. For an inner node c of T , we denote the left child of c with c_l and the right child of c with c_r . We illustrate this notation in the following example:



Remark 3.12. For full binary trees T with $i(T)$ inner nodes and $l(T)$ leaves it is a well-known result that $l(T) = i(T) + 1$ holds. It can be proved inductively.

Remark 3.13. The chirality of nodes will either be denoted by labels or indicated implicitly by embedding the merge tree on the page. Throughout the literature there are different notions of merge trees that are not always distinguished by name or notation. In this work, merge trees do not have explicit weights on edges. Moreover, merge trees in this work a priori do not carry a function to the real numbers. In that way, we distinguish between merge trees and Morse labeled merge trees which will be introduced in Definition 3.19. We use the chirality to obtain Morse labelings on unlabeled merge trees. This leads to Definition 4.3.

Construction 3.14 ([JS20, Thm 9]). Let X be a tree and let $f: X \rightarrow \mathbb{R}$ be a dMf. The *merge tree induced by f* , denoted by $M(X, f)$ is constructed as follows:

Let $c_0 < c_1 < \dots < c_m$ be the critical values of f that are assigned to 1-simplices. The associated merge tree $M(X, f)$ is constructed by induction over these critical values

in descending order. Furthermore, we label the nodes of $M(X, f)$ in order to refer to them later. The label of a node n will be denoted by $\lambda(n)$.

For the base case we begin by creating a node $M(c_m)$ which corresponds to the critical 1-simplex in X labeled c_m and setting its label $\lambda(M(c_m))$ to (c_m, L) .

For the inductive step, let $M(c_i)$ be a node of $M(X, f)$ that corresponds to a critical 1-simplex between two 0-simplices v and w . Define $\lambda_v := \max\{f(\sigma) \mid \sigma \in X_{c_i-\varepsilon}[v], \sigma \text{ critical}\}$ and $\lambda_w := \max\{f(\sigma) \mid \sigma \in X_{c_i-\varepsilon}[w], \sigma \text{ critical}\}$. Two child nodes of $M(c_i)$ are created, named n_{λ_v} and n_{λ_w} . Then label the new nodes $\lambda(n_{\lambda_v}) := \lambda_v$ and $\lambda(n_{\lambda_w}) := \lambda_w$. If $\min\{f(\sigma) \mid \sigma \in X_{c_i-\varepsilon}[v]\} < \min\{f(\sigma) \mid \sigma \in X_{c_i-\varepsilon}[w]\}$, we assign n_{λ_v} the same chirality (L or R) as $M(c_i)$ and give n_{λ_w} the opposite chirality. Continue the induction over the rest of the critical 1-simplices.

Remark 3.15. By construction, one of the following two cases holds for the labels λ_v and λ_w . They might be either critical values lower than c_i that are assigned to edges or critical values that are assigned to nodes. The two labels λ_v and λ_w do not necessarily belong to the same case.

Therefore, the nodes n_{λ_v} and n_{λ_w} will possibly be denoted as $M(c_j)$ and $M(c_k)$ for some $j, k < i$ in later steps of the induction. In particular, this means that the node which is considered in the first instance of the inductive step is c_m .

Remark 3.16. Although the induced merge tree as introduced above comes with a labeling, the labeling is not part of the data of the induced merge tree. This is one of the main differences between merge trees in [JS20] and the merge trees in [Cur19].

We consider one way to keep some information provided by the induced labeling on $M(X, f)$.

Definition 3.17. Let T be a merge tree. We call a total order \leq on the nodes of T a **Morse order** if it fulfills the following two properties for any subtree T' of T :

- (1) The restriction $\leq|_{T'}$ attains its maximum on the root of T' .
- (2) The restriction $\leq|_{T'}$ attains its minimum on the subtree with root p_l/p_r if L/R is the chirality of the root p of T' .

Moreover, we call a merge tree (T, \leq) together with a Morse order \leq a **Morse-ordered merge tree (Mo tree)**.

Remark 3.18. Assuming property (2) of Definition 3.17 for every subtree T' with root p of T is equivalent to either of the following:

- For any subtree T' with root p of T , the minimum of $\leq|_{T'}$ has the same chirality as p .
- For any subtree T' with root p of T , all nodes on the shortest path between p and the minimum of $\leq|_{T'}$ have the same chirality as p .

The equivalence can be proved by an inductive argument over all nodes of the shortest path between p and the minimum.

Because Morse orders \leq define in particular finite totally ordered sets, there are unique order-preserving isomorphisms $\lambda: (V(T), \leq) \xrightarrow{\cong} \{0, 1, \dots, i(T) + l(T) - 1\} \subseteq \mathbb{N}_0 \subset \mathbb{R}$ for each Morse order. Conversely, each injective labeling $\lambda: T \rightarrow \mathbb{R}$ induces an order \leq_λ on the nodes of T by usage of the total order on \mathbb{R} .

Definition 3.19. For Morse orders \leq we call the map $\lambda_{\leq}: (V(T), \leq) \rightarrow \{0, 1, \dots, i(T) + l(T) - 1\}$ the **Morse labeling induced by \leq** . We call an arbitrary labeling $\lambda: T \rightarrow \mathbb{R}$ a **Morse labeling** if it induces a Morse order on T .

We call a merge tree (T, λ) with a Morse labeling $\lambda: T \rightarrow \mathbb{R}$ a **Morse labeled merge tree (Ml tree)**.

For a Mo tree (T, \leq) we call the Ml tree (T, λ_{\leq}) the **Ml tree induced by (T, \leq)** .

Proposition 3.20. *Let $f: X \rightarrow \mathbb{R}$ be a dMf. The labeling which appears in Construction 3.14 induces a Morse order on $M(X, f)$. Hence, $M(X, f)$ canonically carries the structure of a Mo tree as well as an Ml tree.*

Proof. It is proved in [JS20, Thm 9] that $M(X, f)$ is a merge tree. We only have to prove that the labeling induces a Morse order. By Remark 3.3 the set $\text{Cr}(f)$ of critical values carries a total order induced by f . Since the critical values of f precisely define the labeling in Construction 3.14, the labeling induces a total order on the nodes of $M(X, f)$. It is only left to prove that this order is a Morse order.


In the construction, each inner node of $M(X, f)$ corresponds to a critical 1-simplex and is labeled with the critical value of said critical 1-simplex. Since parent nodes are created before their child nodes are, and since the critical values are considered from highest to lowest, property (1) of Definition 3.17 is fulfilled. The rule in the construction which decides the chirality of the child nodes is exactly the same as property (2) of Definition 3.17. Hence, it is fulfilled by construction. \square

We denote the canonical labeling of $M(X, f)$ by λ_f .

Remark 3.21. In the aforementioned proof, it becomes clear that both conditions of Definition 3.17 are necessary for a total order on $M(X, f)$ to be induced by a dMf f .

Morse orders are useful for the construction of dMfs that induce given merge trees T . We will see in Proposition 4.24 that for a total order on an arbitrary merge tree T condition (1) is sufficient for inducing a dMf in the sense of Definition 4.21 later on. Nonetheless, condition (2) is necessary to ensure that the induced dMf induces the given merge tree T .

Example 3.22. We consider the following example of a dMf $f: X \rightarrow \mathbb{R}$:




The critical values on edges are:


$$5 < 6 < 9 < 11 < 14$$

We now show the construction algorithm of $M(X, f)$ visually by depicting $X_{c_i - \varepsilon}$ on the left and the part of $M(X, f)$ that is created up to the step corresponding to c_i on the right.


Start:




$c_i = 14$:




$c_i = 11$:




$c_i = 9$:



$c_i = 6$:



$c_i = 5$:



There are no more critical edges left, so the construction of $M(X, f)$ is finished.

Since there are different notions of merge trees in the literature and since the merge trees in our setting carry a lot of structure, there are multiple possibilities of how to define equivalences of merge trees. In the remainder of this section, we define and discuss some versions of equivalence of merge trees.

Since merge trees are defined to be chiral rooted binary trees, the obvious notion for isomorphisms of merge trees is isomorphisms of chiral rooted binary trees. In detail, this means bijections between the sets of nodes and the sets of vertices which map the root to the root and are compatible with the chiral child relation. For Mo trees (Definition 3.17) we have more notions of equivalence.

Definition 3.23. Let (T, \leq) and (T', \leq') be Mo trees. An *isomorphism* of Mo trees $(T, \leq) \cong (T', \leq')$ is an order-preserving isomorphism of the underlying merge trees.

Let (T, λ) and (T', λ') be Ml trees. An *isomorphism* of Ml trees is an isomorphism of the underlying merge trees over \mathbb{R} , that is, an isomorphism of merge trees $\varphi: T \rightarrow T'$ such that $\lambda' \circ \varphi = \lambda$.

Definition 3.24. Let (T, λ) and (T', λ') be Ml trees. A *shuffle equivalence* $(\varphi, \psi): (T, \lambda) \rightarrow (T', \lambda')$ of Ml trees is a pair of an isomorphism of the underlying merge trees $\varphi: T \rightarrow T'$ and a bijection $\psi: \mathbb{R} \rightarrow \mathbb{R}$ such that

- $\psi \circ \lambda = \lambda' \circ \varphi$ holds,
- the restriction of ψ to values on leaves is order-preserving, and
- the restriction of ψ to values on inner nodes is order-preserving.

In the special case that the restriction $\psi|_{\text{im}(\lambda)}: \text{im}(\lambda) \rightarrow \text{im}(\lambda')$ is an order preserving bijection, we call (φ, ψ) an *order equivalence*.

A *shuffle equivalence* $(T, \leq) \rightarrow (T', \leq')$ between Mo trees is an isomorphism φ of the underlying merge trees such that

- the restriction of φ to leaf nodes is order-preserving, and
- the restriction of φ to inner nodes is order-preserving.

Remark 3.25. The name of shuffle equivalences hints at the fact that two given total orders, one total order on the leaves, and another total order on the inner nodes, might be combined to produce a total order on all nodes in different ways, a bit like shuffling cards. Shuffle equivalence checks if two Morse orders arise from the same underlying orders by different ways of shuffling. Nonetheless, the necessity of ranking ancestors

higher than descendants and being compatible with the chirality prevents arbitrary ways of shuffling two given orders on the leaves and inner nodes from producing Morse orders.

Shuffle equivalences induce by definition isomorphisms on the underlying merge trees. We now make the relationship between Mo trees and Ml trees precise.

Proposition 3.26. *The Morse labeling induced by a Morse order and the order induced by a Morse labeling define inverse bijections $\text{iMl}: \text{MoT}/\cong \xrightarrow{\cong} \text{MlT}/\sim: \text{iMo}$ where \sim denotes order equivalence.*

Proof. Let (T, \leq) be a Mo tree. It is immediate that (T, λ_{\leq}) has the property that λ_{\leq} induces \leq as its induced order on T . Thus, the composition $\text{iMo} \circ \text{iMl}$ is the identity on MoT .

Let (T, λ) be an Ml tree. By definition, the labeling λ induces a Morse order \leq_{λ} on T which makes (T, \leq_{λ}) a Mo tree. Since the induced Morse labeling $\lambda_{\leq_{\lambda}}$ by construction induces \leq_{λ} as its induced order, it follows that (T, λ) and $(T, \lambda_{\leq_{\lambda}})$ are order equivalent. \square

Corollary 3.27. *The induced Morse order iMo and the induced Morse labeling iMl induce a bijection between shuffle equivalences of Mo trees and shuffle equivalences of Ml trees.*

Proof. The assignment iMl maps shuffle equivalences of Mo trees to shuffle equivalences of Ml trees by Definition 3.24.

Let $\lambda: T \rightarrow \mathbb{R}$ and $\lambda': T' \rightarrow \mathbb{R}$ be shuffle-equivalent Morse labelings and let (φ, ψ) be the corresponding shuffle equivalence. Then by Definition 3.19 the map $\text{iMo}(\varphi, \psi) = \varphi: (T, \leq_{\lambda}) \rightarrow (T', \leq_{\lambda'})$ has the property that

- the restriction of φ to leaf nodes is order-preserving, and
- the restriction of φ to inner nodes is order-preserving.

Hence, $\varphi = \text{iMo}(\varphi, \psi)$ is a shuffle equivalence of Mo trees. \square

Remark 3.28. In particular, the aforementioned proposition and corollary mean that two Mo trees are isomorphic (respectively shuffle equivalent) if and only if the corresponding Ml trees are order equivalent (respectively shuffle equivalent) and vice versa.

3.2. Generic Properties and Equivalences of DMFs. In this subsection, we will take a closer look at generic properties that dMfs can be assumed to have. Furthermore we consider some notions of equivalences between dMfs.


A first example of a generic property is the notion of index-ordered dMfs as defined above. It is inspired by the eponymous notion from the smooth case. We use index-ordered dMfs in order to distinguish critical simplices by their dimension because merge trees have the same property: Critical 0-simplices appear as leaves whereas critical 1-simplices appear as inner nodes of the induced merge tree (see Construction 3.14). Therefore, index-ordered dMfs seem to be especially suitable for working with merge trees.

Nonetheless, index-ordered dMfs are not compatible with the structure of rooted subtrees. In detail, consider the following:

Remark 3.29. Let $f: X \rightarrow \mathbb{R}$ be an index-ordered dMf on a tree such that the following holds: The tree X has a critical 1-simplex σ such that the corresponding inner node

p in $M(X, f)$ has two inner nodes c and c' as children. Then the image of f on at least one of the connected components corresponding to c or c' is not an interval in $f(\text{Cr}(f))$.

Example 3.30. The following is a small example:



Neither the subtree with root labeled 4, nor the subtree with root labeled 5 is labeled with an interval in $f(\text{Cr}(f)) \subseteq \mathbb{R}$.


Nonetheless, we can assume compatibility with the structure of rooted subtrees as a property:

Definition 3.31. Let $f: X \rightarrow \mathbb{R}$ be a dMf. The function f is called **sublevel-connected** if for all critical 1-simplices v the set $f(X_{f(v)}[v])$ is an interval in $f(\text{Cr}(f))$.

Remark 3.32. Since both ‘index-ordered’ and ‘sublevel-connected’ are properties that only rely on the values of f on critical simplices, they can easily be arranged without changing the partial matching if the tree X is finite. It is also possible to do this without changing the induced merge tree. One way to obtain a sublevel-connected dMf would be to choose a collapsing order for the induced matching such that the respective connected components of sublevel sets correspond to intervals in said collapsing order.

Nonetheless, as seen in Remark 3.29, the two properties are in most cases mutually exclusive.

Example 3.33. Here is one possibility how to modify the dMf from the previous example in order to make it sublevel-connected without changing the induced merge tree:



Since dMfs can be modified to fulfill either of the two properties, one can always choose the one which is more convenient for the task at hand. Thus, we give two different constructions in this work, one for each property.

We recall that by Remark 3.3 each dMf $f: X \rightarrow \mathbb{R}$ induces an order on the 0-simplices of X and on the 1-simplices of X , respectively. This yields the following definition, which induces a merge-tree-invariant notion of equivalence between dMfs.

Definition 3.34. Let $f: X \rightarrow \mathbb{R}$ and $g: X \rightarrow \mathbb{R}$ be dMfs on a tree X . We call f and g **shuffle-equivalent** if they have the same critical simplices and if they induce the same order on the critical 0-simplices as well as the same order on the critical 1-simplices.

Let (X, f) and (X', f') be two dMfs on trees. A **shuffle equivalence** $(\varphi, \psi): f \rightarrow f'$ between f and f' consists of a simplicial map $\varphi: X \rightarrow X'$ and a bijection $\psi: \mathbb{R} \rightarrow \mathbb{R}$ such that

- $\psi \circ f = f' \circ \varphi$,
- $\varphi|_{\text{Cr}(f)}: \text{Cr}(f) \rightarrow \text{Cr}(f')$ is a bijection,
- the restriction of ψ to values on critical 0-simplices is order preserving, and
- the restriction of ψ to values on critical 1-simplices is order preserving.

In the special case that the restriction $\psi|_{\text{Cr}(f)}: \text{Cr}(f) \rightarrow \text{Cr}(f')$ is an order preserving bijection, we call (φ, ψ) an **order equivalence**.

Remark 3.35. It is immediate that shuffle equivalence of dMfs is an equivalence relation. The name is inspired analogously as for the eponymous notion for Mo trees and Ml trees. Shuffle equivalence checks if two dMfs arise from the same underlying orders by different ways of shuffling. Nonetheless not all ways of shuffling given orders on the critical 0-simplices and critical 1-simplices produce a dMf because dMfs have to be weakly increasing.

Furthermore, shuffle equivalence, and in particular order equivalence, also considers dMfs to be equivalent if they only differ by scaling because a different scaling does not change the induced orders on simplices.

We split the definition of shuffle equivalences in two steps to simplify the proofs of the following propositions. It is immediate that for two dMfs on the same tree X there is a shuffle equivalence between them if and only if they are shuffle equivalent.

Proposition 3.36. *Let X be a tree and let $f: X \rightarrow \mathbb{R}$ and $g: X \rightarrow \mathbb{R}$ be two shuffle-equivalent dMfs. Then $M(X, f)$ and $M(X, g)$ are isomorphic as merge trees.*

The following two lemmas will be helpful for the proof of the proposition:

Lemma 3.37. *Let X be a tree and let $f: X \rightarrow \mathbb{R}$ be a dMf such that X has at least one critical 1-simplex. Then the function $f|_{\text{Cr}(f)}$ attains its maximum on a critical 1-simplex.*

Proof. We prove the lemma by contradiction. Assume that $f|_{\text{Cr}(f)}$ attains its maximum on a critical 0-simplex σ . Let τ be a critical 1-simplex of X such that there is no other critical 1-simplex on the unique path between σ and τ . Since X is a tree, such a unique path always exists. Said unique path consists of a chain of simplices $\sigma \subset a_0 \supset a_1 \subset \dots \supset a_n \subset \tau$ such that all simplices a_i are non-critical. Then the sequence induces the sequence $f(\sigma) < f(a_0) = f(a_1) < \dots < f(a_n) < f(\tau)$. This is a contradiction to the assumption that $f|_{\text{Cr}(f)}$ attains its maximum on σ . \square

Remark 3.38. The chain of simplices used in the previous proof is an example of a gradient path, see [For98, Def 8.4].

Lemma 3.39. *Let X be a tree and let $f: X \rightarrow \mathbb{R}$ and $g: X \rightarrow \mathbb{R}$ be two shuffle-equivalent dMfs. We denote the critical 1-simplices of f and g by $c_0 < c_1 < \dots < c_n$*

where $<$ denotes the ordering induced by f or g , respectively. Then the connected components $X_{f(c_i)}^f[c_j]$ of sublevel complexes contain the same critical simplices as $X_{g(c_i)}^g[c_j]$ for all $j < i$.

Proof. Connected components of sublevel complexes are trees themselves. Furthermore, the restrictions of dMfs to subcomplexes are again dMfs because being a dMf only depends on properties that are inherited by arbitrary subcomplexes of X . It follows by Lemma 3.37 that for connected components $X_{f(c_i)}^f[c_j]/X_{g(c_i)}^g[c_j]$ which contain a critical 1-simplex c_j , the maps $f|_{X_{f(c_i)}^f[c_j]}$ and $g|_{X_{g(c_i)}^g[c_j]}$ attain their maxima on critical 1-simplices. It follows inductively that $f|_{X_{f(c_i)}^f[c_j]}$ and $g|_{X_{g(c_i)}^g[c_j]}$ attain their maxima in each step on the same critical 1-simplex, namely c_j , because both functions induce the same order on the critical 1-simplices. Thus, in each step in which a lower level is considered, the connected components of sublevel complexes are divided into smaller pieces at the same critical 1-simplex. It might happen that the connected components of sublevel complexes of lower levels differ by matched 1-simplices that have higher values than the next maximal critical value. However, by [For98, Lem 2.6] the homotopy type and, therefore, the connected components of sublevel complexes only change at critical levels. Because of that it follows inductively that critical simplices are located inside the same new connected components of sublevel sets of lower levels for both functions f and g . □

Proof of Proposition 3.36. We consider the construction of the induced merge tree (see Construction 3.14) and prove inductively the slightly stronger result that both functions yield isomorphic merge trees at every step of the construction. This implies that f and g induce isomorphic merge trees.

Since f and g impose the same order on the set of critical 1-simplices, the construction algorithm considers the same critical 1-simplices during the same steps for both functions. This already implies that the base case yields the same result for both functions. In particular, this means that the created root node corresponds to the same critical 1-simplex for both functions. Although the label of the created node might be different for the two functions, it does not affect the isomorphism type of the induced merge tree. This is because the labeling is not part of the data of merge trees.

For the inductive step, we observe that in every step of the construction we consider a connected component of sublevel complexes $X_{f(c_i)}^f[c_j]/X_{g(c_i)}^g[c_j]$ that contains at least one critical 1-simplex, namely the one with the highest remaining critical value c_j . Thus, by Lemma 3.39 in each step of the construction, the same critical simplices occur.

Assume we are at the step that considers the critical 1-simplex c_i . For the two new nodes which are created in the inductive step, two pieces of information are important for the isomorphism type of the induced merge tree, namely the chirality of the new nodes and which critical simplices the new nodes correspond to. The chirality of the new nodes affects the isomorphism type of the induced merge tree directly. The critical simplex corresponding to a child node c decides which connected component of the respective sublevel complex is used to build the subtree with root c and at which point said connected component will be subdivided next.

Both pieces of information are defined by the two connected components that belong to the boundary 0-simplices of c_i . The two child nodes correspond to the critical simplices with the highest critical values.

There are three cases:

- (1) Both connected components contain at least one critical 1-simplex c_j .
- (2) One connected component contains at least one critical 1-simplex c_j whereas the other one only contains one critical 0-simplex c .
- (3) Each of the two connected components contains only one critical 0-simplex c .

It follows by Lemma 3.37 that in case (1) the corresponding 1-simplices with the highest critical values are critical 1-simplices c_j . In case (2) the same is true for the connected component that contains at least one critical 1-simplex. For the connected components in case (2) and (3) that only contain one critical 0-simplex, respectively, it is true that the critical 0-simplex is the only critical simplex in its corresponding connected component. Thus, the new nodes correspond to the same critical simplices for f and for g because both functions induce the same order on 1-simplices and because connected components only correspond to critical 0-simplices if they are the only critical simplex left in the corresponding connected component. Furthermore, connected components $X_{f(c_i)}^f[c]/X_{g(c_i)}^g[c]$ that only contain one critical 0-simplex do not have any influence on the induced merge tree $M(X, f)/M(X, g)$ because their corresponding nodes have already been created during a step that considered a critical 1-simplex with a higher critical value and they are not considered in later steps of the construction.

The chirality of the new nodes depends on the minimal values on critical simplices of the two respective connected components $X_{f(c_i)}^f[c_j]/X_{g(c_i)}^g[c_j]$ or $X_{f(c_i)}^f[c]/X_{g(c_i)}^g[c]$. By Lemma 3.7, these minima belong to critical 0-simplices. By assumption, f and g induce the same order on the critical 0-simplices, so the same 0-simplex is minimal with respect to both functions. Thus, f and g assign the same chirality to the new nodes. □

Remark 3.40. The functions f and g might induce different order relations between 0-simplices and 1-simplices. Therefore, in sublevel complexes that appear during the same step of the construction there might be different connected components that contain only one critical 0-simplex each with respect to the two dMfs. However, those connected components that contain only one critical 0-simplex and no critical 1-simplices do not affect the isomorphism type of the induced merge tree. This is because those connected components correspond to leaves of the merge tree which have already been created during the step that considered the critical 1-simplex between said connected components and other connected components. Furthermore, those connected components do not appear in later steps of the construction of the induced merge tree because they do not contain any critical 1-simplices.


Proposition 3.41. *Shuffle equivalences of dMfs induce shuffle equivalences of the induced Ml trees. Moreover, order equivalences of dMfs induce order equivalences of the induced Ml trees.*

Proof. Let (X, f) and (X', f') be dMfs on trees and let (φ, ψ) be a shuffle equivalence between them. The proof that $M(X, f)$ is isomorphic to $M(X', f')$ as merge trees is almost the same as the proof for Proposition 3.36. The only difference is that instead

of id_X we now have a simplicial map $\varphi: X \rightarrow X'$ that induces a bijection on the critical simplices. Since φ commutes with f and f' , the map φ induces a bijection between the sublevel complexes of (X, f) and the sublevel complexes of (X', f') . Moreover, the map φ also induces a bijection between connected components of sublevel complexes because φ is simplicial. The rest of the proof that $M(X, f) \cong M(X', f')$ as merge trees is completely analogous to the proof of Proposition 3.36. In particular, the nodes of $M(X, f) \cong M(X', f')$ correspond to the critical simplices of X, X' , respectively, that are mapped to each other by φ .

It is only left to prove that the induced labelings induce the same Morse order on leaves and on inner nodes. Since the nodes of $M(X, f) \cong M(X', f')$ correspond to the simplices of X, X' , respectively, which are mapped to each other by φ , and since φ commutes with f, f' and ψ , the simplices of X and X' that are mapped to each other by φ differ in their labels as described by ψ . Since ψ is, by assumption, order preserving on values on critical 0-simplices and on values on critical 1-simplices, the two induced Morse orders coincide on leaves and on inner nodes. The same argument holds for the order equivalence case. \square

Remark 3.42. The criterion from Proposition 3.36 for merge tree equivalence is sufficient but not necessary, as the following example shows:



The two dMfs f and g induce inverse orders on the two 0-simplices. Nonetheless, f and g induce the same unlabeled merge tree.

This remark leads us to yet another kind of equivalence relation between dMfs that arises from symmetries of sublevel complexes. In order to make this notion of symmetry precise, we need some preparations.

Definition 3.43. Let $f: X \rightarrow \mathbb{R}$ be a dMf on a tree. For each non-empty connected component $X_c^f[v]$ of a sublevel complex X_c^f we denote by $\text{Aut}(X_c^f[v])$ the group of simplicial automorphisms of $X_c^f[v]$. For each $a \in \text{Aut}(X_c^f[v])$ there is an extension to a self-bijection $X \rightarrow X$ by the identity. The group $\widetilde{\text{Aut}}(X_c^f[v])$ is defined to be the group of said extensions of elements of $\text{Aut}(X_c^f[v])$ by the identity. We consider $\widetilde{\text{Aut}}(X_c^f[v])$ as a subgroup of the group of all self-bijections of X . The total order on $\text{Cr}(f)$ induced by f induces chains $\widetilde{\text{Aut}}(X_{c_0}^f[v]) \subset \widetilde{\text{Aut}}(X_{c_1}^f[v]) \subset \dots$ of inclusions of subgroups. Moreover, we have inclusions $\widetilde{\text{Aut}}(X_{c_i}^f[v]) \subset \widetilde{\text{Aut}}(X_{c_j}^f[v]) = \widetilde{\text{Aut}}(X_{c_j}^f[w]) \supset \widetilde{\text{Aut}}(X_{c_i}^f[w])$ if v and w are in different connected components of some sublevel complex $X_{c_i}^f$ that merge together in some other sublevel complex $X_{c_j}^f$ for $j > i$. We define the **sublevel automorphism group** of (X, f) , denoted by $\text{Aut}_{sl}(X, f)$, to be the subgroup generated by $\bigcup_{c \in \text{Cr}(f), v \in X} \widetilde{\text{Aut}}(X_c^f[v])$. We call the elements of $\text{Aut}_{sl}(X, f)$ **sublevel automorphisms**.

Remark 3.44. Even though sublevel automorphisms are built out of simplicial automorphisms of connected components $X_c^f[v]$ of sublevel complexes, they are in general not simplicial maps $X \rightarrow X$. To be precise, if a simplicial automorphism of $X_c^f[v]$ was used to construct a sublevel automorphism $a \in \text{Aut}_{sl}(X, f)$, then a will fail to be simplicial at the boundary of $X_c^f[v] \subset X$.

Proposition 3.45. *Let $f: X \rightarrow \mathbb{R}$ be a dMf on a tree and let $a \in \text{Aut}_{sl}(X, f)$ be a sublevel automorphism. Then $f * a$ defined by $f * a(\sigma) := f(a(\sigma))$ is a dMf on X . Moreover, this defines a right group action of $\text{Aut}_{sl}(X, f)$ on the set of dMfs on X .*

Proof. Since a is in particular a self-bijection of X , the map $f * a$ is distinct-valued and at most 2-1 because f is. Because a is an automorphism of a sublevel complex X_c^f , it preserves the face relations between cells of X_c^f . Thus, $f * a$ is weakly increasing on X_c^f because f is. The function $f * a$ is also weakly increasing at the boundary of X_c^f because X_c^f is a sublevel complex.

The compatibility of the group action follows directly by the associativity of the composition of maps. \square

Remark 3.46. Since automorphisms of simplicial complexes preserve the dimension of simplices, the action of $\text{Aut}_{sl}(X, f)$ on the set of dMfs on a tree X preserves the properties of being index-ordered or sublevel-connected.

Definition 3.47. Let $f: X \rightarrow \mathbb{R}$ and $g: X \rightarrow \mathbb{R}$ be dMfs on a tree X . We call f and g **sublevel-equivalent** if $\text{Cr}(f) = \text{Cr}(g)$ and $X_c^f \cong X_c^g$ for all $c \in \text{Cr}(f) = \text{Cr}(g)$. If additionally $g = f * a$ holds for a sublevel automorphism $a \in \text{Aut}_{sl}(X, f) = \text{Aut}_{sl}(X, g)$, then we call f and g **symmetry-equivalent**. We call the map a a **symmetry equivalence** from f to g .

We call two dMfs $f: X \rightarrow \mathbb{R}$ and $g: Y \rightarrow \mathbb{R}$ **symmetry-equivalent** if there is a simplicial isomorphism $\varphi: X \rightarrow Y$ such that f and $g \circ \varphi$ are symmetry-equivalent.

Example 3.48. We give a list of some symmetry-equivalent dMfs on a path with four vertices. Here we denote the sublevel equivalence induced by the reflection of the connected component of the sublevel complex of level k by $a_k[k]$:

$$\begin{array}{cccccccc}
 0 & 4 & 1 & 5 & 2 & 6 & 3 & \\
 \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \\
 & & & & & & \xrightarrow{a_6[6]} & \\
 & & & & & & & \bullet & 6 & 2 & 5 & 1 & 4 & 0 \\
 & & & & & & & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\
 a_5[5] & \rightarrow & 3 & 6 & 0 & 4 & 1 & 5 & 2 & a_4[4] & \rightarrow & 3 & 6 & 1 & 4 & 0 & 5 & 2 \\
 & & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & & & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet
 \end{array}$$

Proposition 3.49. *Let $f: X \rightarrow \mathbb{R}$ and $g: X \rightarrow \mathbb{R}$ be symmetry-equivalent dMfs on a tree X . Then $M(X, f)$ and $M(X, g)$ are isomorphic as Ml trees.*

Proof. It suffices to consider the case of a single sublevel automorphism $a \in \widetilde{\text{Aut}}(X_c^f[v])$. Since symmetry equivalences consist of compositions of sublevel automorphisms, the statement then follows by repeated application of the aforementioned case.

We consider how the application of a sublevel automorphism a affects the steps of the construction of the induced Ml tree, Construction 3.14. The sublevel automorphism a does not affect steps that do not remove any critical edge of $X_c^f[v]$ because a , by definition, acts identically outside of $X_c^f[v]$. Considering a critical edge of $X_c^f[v]$ in the constructing algorithm of the induced Ml tree corresponds to passing from a sublevel complex X_c^f in the sublevel filtration to a sublevel complex of a slightly lower level $X_{c-\varepsilon}^f$.

We observe that a is precisely a simplicial automorphism of $X_c^f[v]$. Since simplicial automorphisms preserve the filtration of the corresponding sublevel complex X_c^f into further sublevel complexes $X_{c-\varepsilon}^f$, the parts that $X_c^f[v]$ is divided into by passing to a slightly lower level $c - \varepsilon$ induce the same child nodes with the same chirality and label in the induced MI tree.

In conclusion, the construction yields the same results as without the application of a during all steps. \square


Remark 3.50. Sublevel automorphisms of dMfs on paths only consist of reflections of the corresponding connected component of a subcomplex. When such a connected component of a sublevel complex of a critical level c is considered during the construction of the induced MI tree, the reflection of the connected component only causes the two new parts that are obtained by considering a slightly lower level $c - \varepsilon$ to appear as their mirror images. In particular, the given dMf attains the same values on the two new parts as before. Hence, the two parts that appear are simplicially isomorphic to the ones that appear without application of the reflection.

Definition 3.51. Let (X, f) and (X', f') be dMfs on trees. A **component-merge equivalence (cm equivalence) of level a** is a bijection $\varphi: X \rightarrow X'$ such that one of the following, not necessarily exclusive, cases holds:

- (1) φ is a symmetry equivalence.
- (2) φ fulfills the following:
 - $f' \circ \varphi = f$,
 - φ induces a bijection between the sets of connected components of sublevel complexes such that each restriction $\varphi|_{X_{a-\varepsilon}[v]}: X_{a-\varepsilon}[v] \rightarrow X'_{a-\varepsilon}[\varphi(v)]$ is a cm equivalence of some level $b < a$, and
 - the edge $\sigma \in X$ with $f(\sigma) = a$ merges the two connected components $X_{a-\varepsilon}[v_1]$ and $X_{a-\varepsilon}[v_2]$ in $X_a[v_1] = X_a[v_2]$ if and only if the edge $\varphi(\sigma)$ merges the two connected components $X'_{a-\varepsilon}[\varphi(v_1)]$ and $X'_{a-\varepsilon}[\varphi(v_2)]$ in $X'_a[\varphi(v_1)] = X'_a[\varphi(v_2)]$.

If φ fulfills property (2) but not property (1), we call φ **non-trivial**.

Example 3.52. We give an example of two cm-equivalent dMfs on trees. The non-trivial cm equivalence from the left-hand-side to the right-hand-side consists of a symmetry equivalence of level 5 and the attachment of the edge labeled 6 between the vertices labeled 1 and 3 rather than 2 and 3. That is, it is a cm-equivalence of level 6.



Proposition 3.53. *CM equivalent dMfs on trees induce isomorphic MI trees.*

Proof. Let $\varphi: (X, f) \rightarrow (X', f')$ be a cm equivalence. By property (ii) of Definition 3.1, at most one non-trivial cm equivalence of level a can occur for any level a because there is at most one edge labeled a in (X, f) , (X', f') , respectively. Thus, we can decompose

any cm equivalence into a sequence $(\varphi_a)_a$ of non-trivial cm equivalences of decreasing levels such that each φ_a only changes the attachment of the single edge σ with $f(\sigma) = a$ and acts as a symmetry equivalence on the rest of path and dMf. It suffices to consider a single level a because the statement then follows by induction from highest to lowest over all levels a .

For such a non-trivial cm equivalence φ_a we consider the step of the construction of the induced Ml trees that considers the edge σ with $f(\sigma) = a$ and the edge $\varphi(\sigma)$. We inductively assume that φ induces an isomorphism of induced Ml trees everywhere outside the subtrees corresponding to the two connected components of $X_{a-\varepsilon}^f$ that are merged by the edge σ with $f(\sigma) = a$. That is, on the rest of $M(X, f)$ the map $M(\varphi)$ is a bijection compatible with the chiral child relation onto $M(X', f')$ except for the subtrees of $M(X', f')$ which correspond to the connected components of $X_{a-\varepsilon}^{f'}$ which are merged by the edge $\varphi(\sigma)$. Since the map φ is compatible with the dMfs and because it restricts to a cm equivalence $X_{a-\varepsilon}^f \rightarrow X_{a-\varepsilon}^{f'}$, the dMf f attains the same minima and maxima on the two relevant connected components of $X_{a-\varepsilon}^f$ as f' does on their counterparts of $X_{a-\varepsilon}^{f'}$ via φ . Since Construction 3.14 only considers which two connected components are merged by the considered edge, it makes no difference for the isomorphism type of the induced Ml trees that σ in general merges the two connected components of $X_{a-\varepsilon}^f$ at vertices that do not correspond via φ to the ones adjacent to $\varphi(\sigma)$ in $X_{a-\varepsilon}^{f'}$. Thus, the construction of the induced Ml tree produces nodes with the same chirality and label for both induced Ml trees in the steps that consider $\sigma, \varphi(\sigma)$, respectively. By assumption, the restriction $\varphi_{X_{a-\varepsilon}^f} : X_{a-\varepsilon}^f \rightarrow X_{a-\varepsilon}^{f'}$ is a symmetry equivalence, so the isomorphism of Ml trees extends to the subtrees that correspond to the respective connected components. \square

Proposition 3.54. *Let (X, f) be a dMf on a tree. There is a dMf on a path (P, f') such that (X, f) is cm-equivalent to (P, f') .*

Proof. Let σ be the minimal edge of X with respect to f such that σ is adjacent to a vertex v which is adjacent to more than 2 edges in $X_{f(\sigma)}$. Then the connected component $X_{f(\sigma)-\varepsilon}[v]$ is a path and the vertex v is adjacent to exactly two edges in $X_{f(\sigma)-\varepsilon}[v]$. We construct a cm equivalence of level $f(\sigma)$ by reattaching σ to one of the two vertices of $X_{f(\sigma)-\varepsilon}[v]$ that are only incident to one edge, respectively. If the other boundary vertex w of σ is incident to more than two edges in $X_{f(\sigma)-\varepsilon}[w]$, we apply the same procedure for w .

By induction over the level, we construct a sequence of cm equivalences of increasing level that starts at X and ends at P . The concatenation of all cm equivalences of that sequence yields the desired cm equivalence $(X, f) \rightarrow (P, f')$. \square

Remark 3.55. The way we defined cm equivalence makes it a generalization of symmetry equivalence. In fact, cm equivalences are the same as symmetry equivalences, i.e. they are always trivial, if we restrict ourselves to dMfs on paths: Without loss of generality, cm equivalences of some level a of a dMf on a path (P, f) describe all different possibilities of how to glue two paths together with a new edge in order to obtain a path again. This means that the edge labeled a can only be adjacent to the vertices that are adjacent to less than two edges, respectively, of the two old paths in $P_{a-\varepsilon}$. Thus, there are at most four possibilities for the two vertices which may be adjacent to the edge labeled a . All of these possibilities result in dMfs which are

related to each other by reflections of the original two paths in $P_{a-\varepsilon}$. Hence, they are all symmetry-equivalent to each other.

4. CONSTRUCTION OF THE INDUCED INDEX-ORDERED DMF

We address the inverse question: For any given merge tree T , is there a discrete Morse function f on a path P such that $M(P, f) \cong T$? We answer this question affirmatively by presenting an explicit construction of P and two possible choices for f . The path P is not unique in the sense that pairs of matched 0- and 1-simplices might be added without changing $M(P, f)$. Furthermore, f is not unique in the sense that different dMfs may induce the same critical cells and even dMfs with different critical cells may induce the same merge tree. We will construct P and two versions of f in a way such that all cells of P are critical with respect to f . One version of f will be index-ordered whereas the other version will be sublevel-connected.

The basic idea is to reverse-engineer the construction of the induced merge tree from Construction 3.14. Since we want all cells to be critical, the 1-simplices of P will be in bijection to the inner nodes of T . Moreover, the 0-simplices of P will be in bijection to the leaves of T . Since P is a path by assumption, its isomorphism type is completely determined by the nodes of T . Thus, if $i(T)$ is the number of inner nodes of T and $l(T)$ is the number of leaves of T , the path P consists of $i(T)$ 1-simplices and $l(T)$ 0-simplices. Because of that, we only have to find a dMf on P which induces T as its induced merge tree.

We start with the index-ordered case. There we have found two possible approaches: constructing P and f ‘all-at-once’, by constructing a Morse order on T that induces the desired dMf, or ‘step-by-step’, by induction on the nodes of T .

We choose the all-at-once approach because it openly displays the underlying structures we need. We present the step-by-step approach in Appendix A.1 as an alternative construction of the index-ordered dMf.

We define two different orders on T . First we define a Morse order on the nodes of T , which we call the ***index Morse order***. After that, we use the index Morse order to assign the natural numbers $0, \dots, i(T) + l(T) - 1 = 2 \cdot i(T)$ to the nodes of T in order to obtain a Morse labeling on T . Afterwards, we define the simplex order on the nodes of T , which we use to turn the labeling induced by the index Morse order into a dMf on P .

4.1. The Index Morse Order. To define the index Morse order, we first observe that every node a of T is uniquely determined by the shortest path from the root to a . We recall that the depth of T is the maximal length of any path in T that appears as the shortest path from the root to a leaf. Because T is chiral, we can identify such shortest paths with certain words:

Definition 4.1. Let T be a merge tree of depth n and let a be a node of T . The ***path word*** corresponding to a is a word $a_0 a_1 \dots a_n \in \{L, R, _ \}^{n+1}$ where $_$ denotes the empty letter. If a is of depth k , the letters $a_0 \dots a_k$ are given by the chirality of the nodes belonging to the shortest path from the root to a . The letters $a_{k+1} \dots a_n$ are then empty.

Remark 4.2. Let a, b be nodes of a merge tree T and let $a_0 a_1 \dots a_n$ be the path word corresponding to a and $b_0 b_1 \dots b_n$ be the path word corresponding to b . Then the

equation $a_0 = b_0 = L$ always holds because we consider paths that begin at the root. Because of $a_0 = b_0 = L$ and because we consider finite trees, there is always a maximal $k \in \mathbb{N}$ such that $a_i = b_i$ holds for all $i \leq k$. Furthermore, the last non-empty letter of a path word is always the chirality of the considered node.

We now define the index Morse order, which will produce an index-ordered dMf on P afterwards.

Definition 4.3. Let T be a merge tree. We define the *index Morse order* \leq_{io} on the nodes of T as follows:


Let a and b be arbitrary nodes of T . If a is a leaf node and b is an inner node, then we define $a \leq_{io} b$. If either both a and b are leaf nodes or both a and b are inner nodes, we consider the following:

Let $a_0 a_1 \dots a_n$ be the path word corresponding to a and $b_0 b_1 \dots b_n$ the path word corresponding to b . Furthermore, let $k \in \mathbb{N}$ be maximal such that $a_i = b_i$ for all $i \leq k$. If $a_k = b_k = L/R$ we define $a \leq_{io} b$ if and only if one of the following cases hold:

- a) $a_{k+1} = L$ and $b_{k+1} = R$ / $a_{k+1} = R$ and $b_{k+1} = L$
- b) $b_{k+1} = -$
- c) $a = b$ ($\Leftrightarrow k = n$)

The index Morse order is tailor-made to induce an index-ordered dMf later on. Nonetheless, we will see in Example 6.2 that it is in general not the only Morse order which induces an index-ordered dMf. In Section 5 we will introduce a different, perhaps more natural, Morse order that is more closely related to the sublevel filtration of the induced dMf. But for now we consider an example of the index Morse order and prove that \leq_{io} is actually a Morse order.

Example 4.4. We consider the following merge tree T :



The path words are written underneath their corresponding nodes. The index Morse order produces the following chain of inequalities where we denote the nodes by their corresponding path words:

$$LLL_- \preceq LLRR \preceq LLRL \preceq LRR_- \preceq LRL_- \preceq LLR_- \preceq LL-- \preceq LR-- \preceq L--$$

The inequalities from LLL_- to LRL_- arise from the path words of the leaf nodes. The inequality $LRL_- \preceq LLR_-$ holds because the node corresponding to LRL_- is a leaf node and the node corresponding to LLR_- is an inner node. The inequalities from LLR_- to $L--$ arise from the path words of the inner nodes.

Remark 4.5. By definition, the root node is always the maximal element of $(V(T), \leq_{io})$. Furthermore, the leftmost leaf node of T is always the minimal element of $(V(T), \leq_{io})$.

Proposition 4.6. *The index Morse order is a total order on $V(T)$. Furthermore, it is a Morse order.*

Proof.

Antisymmetry: Let $a, b \in V(T)$ such that $a \leq_{io} b$ and $b \leq_{io} a$. Then either both a and b are leaves or both a and b are inner nodes. Neither of the inequalities can be a result of case a) of Definition 4.3 because otherwise it would contradict the other inequality. Let $k \in \mathbb{N}$ be maximal such that $a_i = b_i$ holds for all $i \leq k$. If $a \leq_{io} b$ belongs to case b), then $b_{k+1} = -$ holds. However, if $b \leq_{io} a$ also belongs to case b), then $a_{k+1} = - = b_{k+1}$. This is a contradiction to the maximality of k . As a result $a_i = b_i$ for all i and $a = b$ holds.

Transitivity: Let a, b, c be nodes of T such that $a \leq_{io} b$ and $b \leq_{io} c$ holds. If one of the relations holds because one of the nodes is a leaf and the other one is an inner node, then a must be a leaf and c must be an inner node. In particular $a \leq_{io} c$ holds. Thus, consider the case that a, b and c are either all leaves or all inner nodes. Let $k, k' \in \mathbb{N}$ be maximal such that $a_i = b_i$ for all $i \leq k$ and $b_i = c_i$ for all $i \leq k'$. There are four cases:

$a = b$ or $b = c$:

If $a = b$ holds, then $a \leq_{io} c$ holds for the same reason as $b \leq_{io} c$ does. Analogously, if $b = c$ holds, then $a \leq_{io} c$ holds for the same reason as $a \leq_{io} b$ does.

$k < k'$:

In that case $a \leq_{io} c$ holds for the same reason that $a \leq_{io} b$ holds.

$k' < k$:

Then $a_i = b_i = c_i$ for $i \leq k'$. Furthermore, it is either $c_{k'+1} = c_{k+1} = -$ or $- \neq a_{k'+1} = b_{k'+1} \neq c_{k'+1} \neq -$ because k' was assumed to be maximal. In particular, $a \leq_{io} c$ holds because either case b) or case a) is fulfilled.


$k = k' < n$ If a, b and c are distinct nodes, the equality $k = k'$ is only possible if $c_{k+1} = -$ and $a_{k+1} = a_k = b_k = c_k \neq b_{k+1}$ holds. In any other configuration, the inequalities $a \leq_{io} b$ and $b \leq_{io} c$ would not hold. In consequence, $a \leq_{io} c$ holds by case b).

Totality: Let a and b be nodes of T . If one of a and b is a leaf and the other one is an inner node, then a and b are comparable by definition. Since T is in particular a rooted binary tree, the path words of a and b coincide at least at $a_0 = b_0 = L$. If there is a $j \in \mathbb{N}$ such that $a_j \neq b_j$, then a and b are comparable by case a) or b). If no such j exists, then $a = b$ holds, that is, they are comparable by case c).

The fact that \leq_{io} is a Morse order follows immediately from the definition: Case b) ensures condition (1) of Definition 3.17 and Case a) ensures condition (2) of Definition 3.17. The different treatment of leaves and inner nodes does not change this because it is defined in agreement with condition (1) and does not affect condition (2). \square

Definition 4.7. We call the Morse labeling $\lambda_{io}: (V(T), \leq_{io}) \rightarrow \{0, 1, \dots, i(T) + l(T) - 1\}$ induced by the index Morse order, see Definition 3.19, the **index Morse labeling** on T . That is, a node c of T is labeled with $\lambda_{io}(c)$.

Example 4.8. The index Morse order from Example 4.4 induces the following index Morse labeling:



4.2. The Simplex Order. We now define the simplex order on the nodes of T . The simplex order will tell us which nodes of T correspond to which simplices of P .

Remark 4.9. Let T be a merge tree and let a, b be nodes of T . Because T is in particular a rooted binary tree, there is a unique node p which is a common ancestor of a and b and has no descendants which are common ancestors of a and b .

Definition 4.10. We call the node p from Remark 4.9 the *youngest common ancestor of a and b* .

Definition 4.11. Let T be a merge tree. We define the *simplex order* \preceq on $V(T)$ as follows: For two nodes a and b of T we define $a \preceq b$ if and only if one of following mutually exclusive cases holds, where p denotes the youngest common ancestor of a and b :

- (1) a is a node of the subtree with root p_l and b is a node of the subtree with root p_r .
- (2) a is a node of the subtree with root b_l (in particular $b = p$).
- (3) b is a node of the subtree with root a_r (in particular $a = p$).
- (4) $a = b$.

Proposition 4.12. *The simplex order is a total order on the nodes of T .*

Before we prove the proposition, we remark the following:

Remark 4.13. Let T be a merge tree and let a, b, c be nodes of T such that b is an ancestor of c and a descendant of a . If c is a node of the subtree with root a_l/a_r , then b is also a node of the subtree with root a_l/a_r .

Proof of Proposition 4.12.

Antisymmetry: Let $a, b \in V(T)$ such that $a \preceq b$ and $b \preceq a$. We are going to prove that if the relation $a \preceq b$ corresponds to one of the cases (1),(2) or (3) from Definition 4.11, this contradicts the relation $b \preceq a$. It then follows that case (4) holds, that is, $a = b$.

1. If $a \preceq b$ corresponds to case (1), then a, b and p are three distinct nodes. In particular, $a \neq b$ holds, so case (4) is not fulfilled for $b \preceq a$. Furthermore, since $a \neq p \neq b$ holds, the cases (2) and (3) can not hold either for $b \preceq a$. Thus, the only possibility for the relation $b \preceq a$ would be that case (1) holds. But this would be a contradiction to case (1) holding for $a \preceq b$.
2. If $a \preceq b$ corresponds to case (2), then a is a node of the subtree with root b_l . In particular $a \neq b = p$. Thus, the cases (1) and (4) can obviously

not hold for $b \preceq a$. Additionally, case (2) can not hold for $b \preceq a$ because otherwise a would need to be an ancestor of b . Furthermore, a is not a node of the subtree with root b_r , so case (3) is also not possible for $b \preceq a$.


3. Case (3) works completely symmetric to case (2).

In conclusion, case (4) must hold. This means that $a = b$ holds.

Transitivity: Let $a, b, c \in V(T)$ such that $a \preceq b$ and $b \preceq c$ holds. If one of the relations belongs to case (4), there is nothing to prove. In order to prove transitivity we distinguish three cases depending on what case of Definition 4.11 the relation $b \preceq c$ belongs to:

$b \preceq c$ belongs to case (1): Let p be the youngest common ancestor of b and c . Then, by assumption, b is a node of the subtree with root p_l and c is a node of the subtree with root p_r . If the relation $a \preceq b$ belongs to case (2), a cannot be an ancestor of p and it follows by Remark 4.13 that a belongs to the subtree with root p_l , in particular, $a \preceq c$ holds. If $a \preceq b$ belongs to case (3), then a is either an ancestor of p or the other way around. If a is an ancestor of p , then, by Remark 4.13, p is a node of the subtree with root a_r . Thus, c is also contained in the subtree with root a_r and $a \preceq c$ holds by case (3). If p is an ancestor of a , then a belongs to the subtree with root p_l and $a \preceq c$ holds by case (1). The situation looks like one of the following:



$a \preceq b$ belongs to case (2):



$a \preceq b$ belongs to case (3):

a is an ancestor of p

p is an ancestor of a




In these pictures, the dotted lines indicate that there might be other nodes in between the two nodes which are connected by dotted lines.

If $a \preceq b$ belongs to case (1), we consider the youngest common ancestor p' of a and b . Then, by assumption, a is a node of the subtree with root p'_l and b is a node of the subtree with root p'_r . Since p and p' are both ancestors of b , either p is an ancestor of p' , or the other way around. If p is an ancestor of p' , then p' is contained in the subtree with root p_l by Remark 4.13. Hence, a is also contained in the subtree with root p_l . Furthermore, the node p is then the youngest common ancestor of a and


c as well as the youngest common ancestor of p' and c . This implies $a \preceq c$ because case (1) is fulfilled.

If p' is an ancestor of p , then p and c are contained in the subtree with root p' . Furthermore, the node p' is then the youngest common ancestor of a and c as well as the youngest common ancestor of a and p . It follows that $a \preceq c$ holds by case (1). The situation looks like this:

p is an ancestor of p' :




p' is an ancestor of p :




$b \preceq c$ belongs to case (2): The node b is a node of the subtree with root c_l . If the relation $a \preceq b$ belongs to case (2), then a is a node of the subtree with root c_l because a 's ancestor b is. Thus, $a \preceq c$ holds by case (2). If $a \preceq b$ belongs to case (3), then either a is an ancestor of c , or c an ancestor of a . If a is an ancestor of c , then, by Remark 4.13, the node c is contained in the subtree with root a_r . Thus, $a \preceq c$ holds by case (3). If c is an ancestor of a , then a is a node of the subtree with root c_l by Remark 4.13, in particular $a \preceq c$ holds by case (2). The situation looks like this:

$a \preceq b$ belongs to case (2):




$a \preceq b$ belongs to case (3):

a is an ancestor of c




c is an ancestor of a




If $a \preceq b$ belongs to case (1), we consider the youngest common ancestor p of a and b . Then a is a node of the subtree with root p_l and b is a node of the subtree with root p_r . Since p and c are both ancestors of b , it is either

p an ancestor of c or c an ancestor of p . If c is an ancestor of p , then a is a node of the subtree with root c_l because a 's ancestor p is by Remark 4.13. Therefore, $a \preceq c$ holds because of case (2). If p is an ancestor of c , then c has to be a node of the subtree with root p_r by Remark 4.13 and a is a node of the subtree with root p_l . Thus, $a \preceq c$ holds by case (1). The situation looks as follows:

c is an ancestor of p :



p is an ancestor of c :



$b \preceq c$ belongs to case (3): Again, the proof for case (3) works completely symmetrically to case (2).

Totality: Let a, b be nodes of T . Because T is by assumption a rooted binary tree, there is a common ancestor p of a and b . It follows that one of the cases 1-4 from Definition 4.11 must hold, so $a \preceq b$ or $b \preceq a$ holds. \square

We use the following definition to make the intuition of leaves being adjacent precise. This allows us to analyze the simplex order further.

Definition 4.14. Let T be a merge tree and let a and b be leaves of T . We call a and b **adjacent** if one of the following holds:

- (1) $a \preceq b$ and there is no leaf node c of T such that $a \preceq c \preceq b$ holds.
- (2) $b \preceq a$ and there is no leaf node c of T such that $b \preceq c \preceq a$ holds.

Lemma 4.15. *Subtrees of T form chains of cover relations in $(V(T), \preceq)$. In detail, this means the following:*

Let p be an inner node of T and let a/b be the leftmost/rightmost leaf of the subtree with root p . Then the nodes of the subtree with root p in T form the chain of cover relations $a \prec \dots \prec p \prec \dots \prec b$ in $(V(T), \preceq)$. Moreover, any two adjacent leaves of T have the property that the left one of the two adjacent leaves is covered by the youngest common ancestor of the two, whereas the right one covers the youngest common ancestor.

Proof. We prove the lemma inductively. Let p be a node of T such that both child nodes of p are leaves. It follows directly by Definition 4.11 that p_r covers p and p covers p_l . That is, the subtree with root p forms the chain of cover relations $a = p_l \prec p \prec p_r = b$.

If p is an arbitrary inner node, then by the inductive hypothesis the subtree with root p_l/p_r forms the chain of cover relations $a_1 \prec \dots \prec p_l \prec \dots \prec b_1/a_2 \prec \dots \prec p_r \prec \dots \prec b_2$ where a_1/a_2 is the leftmost and b_1/b_2 the rightmost leaf of the subtree

with root p_l/p_r . Since b_1/a_2 is a node of the subtree with root p_l/p_r , it follows by case (2)/(3) of Definition 4.11 that $b_1 \prec p/p \prec a_2$ holds. For all nodes c of T which are not nodes of the subtree with root p , the same case from Definition 4.11 holds for c and p as for c and b_1/a_2 . Thus, and because b_1/a_2 is maximal/minimal in the subtree with root p_l/p_r by the inductive assumption, there is no node c such that $b_1 \prec c \prec p/p \prec c \prec a_2$ holds. In conclusion, p covers b_1/a_2 covers p . \square

As mentioned before, we will use the simplex order to relate the nodes of T to the simplices of a path P . In order to do that, we now define a corresponding simplex order on the simplices of P .

Definition 4.16. Let P be a path. There are two 0-simplices p_0 and p_1 in P which belong only to one respective 1-simplex. For each simplex σ of P there is a unique shortest path γ_σ from p_0 to σ . We denote the length, that is, the number of simplices, of such a path γ_σ by $L(\gamma_\sigma)$. The **simplex order** on P is defined as follows: For two simplices σ and τ of P we define $\sigma \preceq \tau$ if and only if $L(\gamma_\sigma) \leq L(\gamma_\tau)$.

Lemma 4.17. *The simplex order on P is a total order on the simplices of P .*

Proof. The proof is straightforward and only uses that P is a path and that the integers are linearly ordered. \square

Remark 4.18. Connected subcomplexes of P correspond to chains of cover relations with respect to the simplex order. Furthermore, any 1-simplex covers its left boundary 0-simplex and is covered by its right boundary 0-simplex. If one visualizes P as being horizontally embedded in a plane such that p_0 is the leftmost 0-simplex of P and p_1 is the rightmost 0-simplex of P , then for simplices $s, s' \in P$ the relation $s \prec s'$ holds if and only if s is left of s' . This reminds us of the fact that the simplex order is only defined up to a choice of orientation.

Proposition 4.19. *Let T be a merge tree and let P be a path with $i(T)$ 1-simplices. Then we have an isomorphism $\phi: (P, \preceq) \xrightarrow{\cong} (V(T), \preceq)$ of totally ordered sets. The isomorphism ϕ only depends on the choice of p_0 and p_1 , that is, on a choice of orientation on P .*

Proof. It is immediate that $V(T)$ and P have the same number of elements. Since both sets are totally ordered, finite, and have the same number of elements, there is exactly one order-preserving bijection between them. Choosing p_0 and p_1 the other way around would reverse the total order on P . This would result in a different isomorphism according to the inverse order. \square

Before we continue with the definition of the index-ordered dMf, we consider how the simplex order can be used to classify the connected components of sublevel complexes of dMfs on paths (P, f) :

Proposition 4.20. *Let $f: P \rightarrow \mathbb{N}_0$ be a dMf on a path P . Then the connected components $P_c[v]$ of sublevel complexes P_c of P are precisely maximal sequences $\sigma := (s_0, \dots, v, \dots, s_k)$ of simplices of P such that $s_i \in P_c$ for all $i = 0, \dots, k$ and $s_0 \prec \dots \prec v \prec \dots \prec s_k$ is a chain of cover relations in (P, \preceq) .*

Proof. As mentioned in Remark 4.18, connected subcomplexes of P are given by chains σ of cover relations in (P, \preceq) . Thus, it follows directly that connected components of

sublevel complexes are maximal chains σ of cover relations in P that consist only of simplices of P_c .

For the other way around, it is clear that sequences σ of simplices of P_c form a subset of P_c . Because σ is supposed to be a sequence of cover relations in (P, \preceq) , it is connected by Remark 4.18. By the assumption that σ is maximal, its simplices form a maximal connected subcomplex, or in other words, a connected component of P_c . \square

4.3. The Induced Index-Ordered DMF. Now we explain how the simplex order can be used to construct dMfs on P from Morse orders on T :

Definition 4.21. Let T be a merge tree and P be a path such that the number of 1-simplices is $i(T)$ and let $\phi: (P, \preceq) \rightarrow (V(T), \preceq)$ be the isomorphism from Proposition 4.19.

For a Morse order \leq and its induced Morse labeling λ we define a map $f_\lambda: P \rightarrow \mathbb{N}_0$ by

$$\begin{array}{ccc} P & \xrightarrow{f_\lambda} & \mathbb{N}_0 \\ \downarrow \phi & \nearrow \lambda & \\ V(T) & & \end{array}$$

The map f_λ is then called the *dMf induced by the Morse order \leq* or the *dMf induced by the Morse labeling λ* .

In particular, the map $f_{io} := \lambda_{io} \circ \phi$ induced by the index Morse order is called the *induced index-ordered dMf*.

Remark 4.22. Although ϕ and λ are order-preserving maps with respect to the previously defined total orders, the map f_{io} does not respect the simplex order in general. Since f_{io} is supposed to be an index-ordered dMf, it does not need to respect the simplex order. Because the map f_{io} is supposed to be index-ordered, it rather needs to be compatible with face relation on P , which we will see to be true later on.

Example 4.23. The index Morse order from Example 4.4, respectively the index Morse labeling from Example 4.8, produces the following pair (P, f_{io}) :

$$\begin{array}{cccccccc} 0 & 6 & 2 & 5 & 1 & 8 & 4 & 7 & 3 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

Proposition 4.24. *For any given Morse order \leq on any merge tree T the dMf induced by \leq is a dMf that has only critical cells.*

Proof. Because the maps λ and ϕ are injective, the same is true for f_λ . In particular, f_λ is matching and all simplices of P are critical with respect to f .

We now prove that f_λ is weakly increasing: Let σ be a 1-simplex of P . By Remark 4.18 σ and its boundary 0-simplices form a chain of cover relations in (P, \preceq) . By Lemma 4.15, the boundary 0-simplices of σ correspond to adjacent leaves in T such that σ is their youngest common ancestor. By property (1) of Definition 3.17, it follows that σ has a higher Morse label than its boundary 0-simplices. Thus, the map f_λ is weakly increasing.

In conclusion, f_λ is a dMf with only critical cells. \square

Remark 4.25. The previous proposition proves that Morse orders \leq on merge trees T always induce dMfs f_λ . It is a priori unclear though whether the induced dMf f_λ induces the given merge tree T as its induced merge tree $M(P, f_\lambda)$. We prove this to be true in Theorem 6.5.

Furthermore, condition (1) from Definition 3.17 is necessary for f_λ to be a dMf, because a violation of (1) between an inner node and a leaf would result in a violation of f_λ being weakly increasing on the corresponding simplices.

Remark 4.26. It is clear that for a given merge tree T the path P from this section and the path P from Appendix A are isomorphic as paths since they both contain $i(T) + 1$ many 0-simplices and $i(T)$ many 1-simplices.

Before we continue with the sublevel-connected dMf, we consider how the simplex order can be used to improve our understanding of sublevel complexes of dMfs on paths (P, f_λ) and how they are related to subtrees of T . The condition for this approach to be applicable is that the dMf f_λ is induced by a Morse order \leq on T as in Definition 4.21, which we will see to be the general case later on. We will apply this approach to the sublevel-connected case in Section 5 where it will be of more importance.

Proposition 4.27. *Let $f_\lambda: P \rightarrow \mathbb{N}_0$ be a dMf on a path P that is induced by a Morse order \leq on T . Then the connected components $P_c[v]$ of sublevel complexes P_c of (P, f) induce subtrees of T via ϕ .*

Proof. It was shown in Proposition 4.20 that connected components of sublevel complexes $P_c[v]$ are maximal sequences $\sigma = (s_0, \dots, v, \dots, s_k)$ of simplices of P such that $s_i \in P_c$ holds for all $i = 0, \dots, k$ and $s_0 \prec \dots \prec v \prec \dots \prec s_k$ is a chain of cover relations in (P, \preceq) . It follows directly from the construction of ϕ in Proposition 4.19 that $\phi(\sigma) := \phi(s_0) \prec \dots \prec \phi(v) \prec \dots \prec \phi(s_n)$ is a maximal chain of cover relations in $(V(T), \preceq)$ such that $s_i \in P_c$ holds for all i .

Let λ be the Morse labeling induced by \leq . Since $f_\lambda = \lambda \circ \phi$ holds by construction of f_λ in Definition 4.21, this means that the condition for the simplices of σ to belong to a sublevel complex can be characterized as $\lambda(\phi(s_i)) \leq c$ for all i .

Since restrictions of total orders to subsets are again total orders, the chain $\phi(\sigma)$ has a maximal node $\phi(s_{max})$ with respect to the Morse labeling λ . Equivalently, $\phi(s_{max})$ is maximal among the nodes of $\phi(\sigma)$ with respect to the Morse order.

We now prove that the subtree with root $\phi(s_{max})$ is equal to $\phi(\sigma)$ by proving that the two sets of nodes are contained within each other.

Because the Morse labeling is induced by a Morse order, all descendants of $\phi(s_{max})$ in T have a lower Morse label than $\phi(s_{max})$ by property (1) of Definition 3.17. By Lemma 4.15 the subtree with root $\phi(s_{max})$ forms a chain of cover relations in $(V(T), \preceq)$. Thus, the subtree with root $\phi(s_{max})$ is contained in $\phi(\sigma)$ because s_{max} is an element of σ and $\phi(\sigma)$ is maximal such that $\lambda(\phi(s_i)) \leq c$ for all i .

We prove that $\phi(\sigma)$ is contained in the subtree with root $\phi(s_{max})$ by contradiction. We assume the existence of a node $\tau \in \phi(\sigma)$ which is not contained in the subtree with root $\phi(s_{max})$. Because the simplex order \preceq on T is a total order, either $\tau \preceq \phi(s_{max})$ or $\phi(s_{max}) \preceq \tau$ holds. Since $\phi(s_{max})$ holds the maximal Morse label among all nodes of $\phi(\sigma)$, no node of $\phi(\sigma)$ can be an ancestor of $\phi(s_{max})$. In particular, this holds for τ . The node τ can by assumption neither be a descendant nor an ancestor of $\phi(s_{max})$. Thus, there is a youngest common ancestor p of $\phi(s_{max})$ and τ which is responsible for

the comparison relation between the two in the simplex order. But then p would be a node of $\phi(\sigma)$ because $\phi(\sigma)$ is a chain of cover relations and either $\tau \preceq p \preceq \phi(s_{max})$ or $\phi(s_{max}) \preceq p \preceq \tau$ would hold. This contradicts the maximality of the Morse label of $\phi(s_{max})$ because p would have a higher Morse label than $\phi(s_{max})$ by property (1) of Definition 3.17.

Altogether we proved that $\phi(\sigma)$ is equal to the subtree with root $\phi(s_{max})$. \square

5. THE SUBLEVEL-CONNECTED DMF

As remarked in Section 3.2 it might sometimes be more convenient to work with sublevel-connected dMfs rather than with index-ordered dMfs. In this section we introduce a slightly different version of the Morse order from Definition 4.3 to construct a sublevel-connected dMf which is shuffle-equivalent to the induced index-ordered dMf and, hence, induces the same given merge tree.

Definition 5.1. Let T be a merge tree. We define the *sublevel-connected Morse order* \leq_{sc} on the nodes of T as follows:

Let a, b be arbitrary nodes of T . Let $a_0a_1 \dots a_n$ be the path word corresponding to a and $b_0b_1 \dots b_n$ the path word corresponding to b (see Definition 4.1). Furthermore, let $k \in \mathbb{N}$ be maximal such that $a_i = b_i$ for all $i \leq k$. If $a_k = b_k = L/R$ we define $a \leq_{sc} b$ if and only if one of the following cases hold:

- a) $a_{k+1} = L$ and $b_{k+1} = R/a_{k+1} = R$ and $b_{k+1} = L$
- b) $b_{k+1} = -$
- c) $a = b$

Remark 5.2. The only difference between the definition of the index Morse order and the definition of the sublevel-connected Morse order is that we do not treat leaves and inner nodes differently anymore. Thus, both orders induce the same order on inner nodes and the same order on leaves. However, the order relation between a leaf and an inner node is in general different then in the index Morse order.

Remark 5.3. The fact that the sublevel-connected Morse order is a Morse order can be proved the same way as the corresponding statement Proposition 4.6 for the index Morse order was proved. There are just fewer case distinctions to be made for the sublevel-connected Morse order.

Lemma 5.4. *Subtrees of T form intervals in $(V(T), \leq_{sc})$.*


Proof. Let T' be a subtree in T . The nodes of T' correspond to path words that begin with the same couple of letters. We denote the path words of nodes a of T' by $a_0a_1 \dots a_n$. Then there is a maximal $k \in \mathbb{N}$ such that $a_0 \dots a_k$ is the same for all nodes of T' . In particular, the path word $a_0 \dots a_k - \dots -$ corresponds to the root of T' .

Let b be a node of T that is not in T' and let $b_0 \dots b_n$ be the corresponding path word. Then $a_i \neq b_i$ holds for at least one $i \leq k$. Thus, the order relations between b and any node of T' are decided by the letters a_i and b_i for the minimal such i . In particular, the relation is the same for all nodes of T' . Thus, all nodes b which are not in T' are either greater than all nodes of T' or smaller than all nodes of T' . In conclusion, the nodes of T' form a chain of cover relations because \leq_{sc} is a total order. \square

The induced sublevel-connected labeling λ_{sc} and the induced sublevel-connected dMf f_{sc} are defined analogously to the index-ordered case:

Definition 5.5. Let T be a merge tree and P be a path such that the number of 1-simplices is $i(T)$. We call the Morse labeling $\lambda_{sc}: (V(T), \leq_{sc}) \rightarrow \{0, 1, \dots, i(T) + l(T) - 1\}$ induced by the sublevel-connected Morse order the **sublevel-connected Morse labeling** on T . The dMf $f_{sc} = \lambda_{sc} \circ \phi$ induced by λ_{sc} is called the **induced sublevel-connected dMf**.

Example 5.6. Let T be the merge tree from Example 4.4. The sublevel-connected Morse labeling on T and the induced sublevel-connected dMf are given below.



There are now two things left to prove: that the map f_{sc} is indeed a sublevel-connected dMf and that it induces the given merge tree T .

Proposition 5.7. *The induced sublevel-connected dMf f_{sc} is a sublevel-connected dMf that has only critical cells.*

Proof. As a dMf induced by a Morse order, the map f_{sc} is by Proposition 4.24 a dMf that has only critical cells. It is left to prove that f_{sc} is sublevel-connected:

By Proposition 4.27, the connected components of sublevel complexes of (P, f) induce subtrees of T via ϕ . By Lemma 5.4, subtrees of T form intervals in (T, \leq_{sc}) . The sublevel-connected Morse labeling λ_{sc} by definition maps intervals of (T, \leq_{sc}) to intervals of \mathbb{N}_0 . By concatenation of these arguments, it follows that $f_{sc} = \lambda_{sc} \circ \phi$ maps connected components of sublevel complexes to intervals of \mathbb{N}_0 , i.e. the map f_{sc} is sublevel-connected. \square

Theorem 5.8. *Let T be a merge tree and let P be a path such that the number of 1-simplices in P is $i(T)$. Then f_{io} and f_{sc} are shuffle-equivalent where f_{sc} is the induced sublevel-connected dMf and f_{io} is the induced index-ordered dMf. Thus, $M(P, f_{sc}) \cong M(P, f_{io})$ holds as merge trees.*

Proof. By Remark 5.2, the restrictions of the index Morse order and the sublevel-connected Morse order to the leaves of T are the same. Thus, the index Morse labeling and the sublevel-connected Morse labeling induce the same order on the leaves of T . Because the induced dMfs are constructed by the concatenation of λ with ϕ , the induced index-ordered dMf and the induced sublevel-connected dMf induce the same order on 0-simplices. It follows analogously that both induced dMfs induce the same order on the 1-simplices of P . That is, the dMfs f_{sc} and f_{io} are shuffle-equivalent.

Thus, $M(P, f_{sc}) \cong M(P, f_{io})$ follows by Proposition 3.36, where f_{io} is the induced index-ordered dMf defined in Definition 4.21. \square

6. RELATIONSHIPS BETWEEN MERGE TREES AND DMFS ON PATHS

In this section we want to take a look at the bigger picture again and consider how we can relate dMfs on paths and trees to merge trees. In order to do this in a structured way, we consider the different sets of dMfs and merge trees and relate them to each other by bijections that are compatible with the various notions of equivalence we introduced earlier. Afterwards, we use said bijections in order to prove that the aforementioned dMfs f_{io} , Definition 4.21, and f_{sc} , Proposition 5.7, both represent the given merge tree. We also relate the sets of dMfs and merge trees mentioned here to the setting of [Cur19].

Remark 6.1. In order to obtain bijections that are compatible with the various notions of equivalence, we restrict ourselves to the case of dMfs for which all simplices are critical. Since the induced merge tree does not take matched cells into account, we would otherwise need to define a notion of equivalence similar to Forman equivalence (see [JS20, Def 4.1]) of dMfs that in particular takes simple homotopy equivalences as well as combinatorial aspects of dMfs into account. This could be done by additional pre- and post-composition of the equivalences as we defined them with simple homotopy equivalences that are compatible with the given dMfs. For simplicity, we chose to leave this aspect out of this work.

We denote the set of merge trees up to isomorphism by Mer and the set of dMfs on paths with only critical simplices up to symmetry equivalence by DMF_P^{crit} . It follows by Proposition 3.49 that the assignment $M(-, -)$ is well-defined on DMF_P^{crit} . Furthermore, the construction of the induced dMf from Definition 4.21 extends to a map which we denote by Φ . Since ϕ is well defined up to a choice of orientation, the map Φ is in particular well defined up to symmetry equivalence. This leaves us with the following diagram:





FIGURE 1. Relationships Between Merge Trees and DMFs on Paths

The arrows induced by the index Morse order and the sublevel-connected Morse order are not left-inverse to the forget arrow as the following example shows. We have marked them in red because they are the only arrows that prevent the diagram from commuting completely. Nonetheless, it is obvious that the arrows induced by the two Morse orders are right-inverses of the forgetful arrow.

Example 6.2. Consider the following Ml trees and their corresponding Mo trees:



It is immediate that (T, λ) and (T', λ') are neither isomorphic nor shuffle equivalent. Thus, $(T', \lambda') \not\cong (T, \lambda) = (\text{forget}(T', \lambda'), \lambda_{io})$ holds. Nonetheless, the Ml tree (T', λ') induces an index-ordered dMf.

But we clearly have the following.

Remark 6.3. Let T, T' be isomorphic merge trees. Then $(T, \leq_{io}) \cong (T', \leq_{io})$ and $(T, \leq_{sc}) \cong (T', \leq_{sc})$ holds as Mo trees. Furthermore, we have $\text{forget}(T, \leq_{io}) \cong T \cong \text{forget}(T, \leq_{sc})$ as merge trees.

We have already seen in Proposition 3.26 that the maps iMo and iMl are inverse to each other in the sense that they are bijections compatible with isomorphisms, order equivalences and shuffle-equivalences. We now show that $M(-, -)$ and Φ are also inverse to each other up to the respective notions of equivalence.

Theorem 6.4. *The induced labeled merge tree $M(-, -)$ and the induced dMf Φ define maps $M(-, -): DMF_P^{crit} \leftrightarrow MlT: \Phi$ that are inverse to each other in the following sense:*

- (1) *For any dMf (P, f) with only critical cells, the dMf $\Phi(M(P, f), \lambda_f)$ is symmetry-equivalent to (P, f) , and*
- (2) *For any Ml tree (T, λ) , the Ml tree $M(\Phi T, f_\lambda)$ is isomorphic to (T, λ) .*

Proof. (1) Let (P, f) be a dMf on a path. We construct a symmetry equivalence between f and f_{λ_f} . It is given as follows: For any simplex σ of $\Phi M(P, f)$ there is exactly one simplex $\tilde{\sigma}$ of P such that $f(\tilde{\sigma}) = f_{\lambda_f}(\sigma)$. This induces a bijection $\varphi: P \rightarrow \Phi M(P, f)$ which is compatible with f, f_{λ_f} and $\text{id}_{\mathbb{R}}$ by definition. But in general, the map φ is not simplicial. This is because the simplex order on $M(P, f)$ might be different than the left/right relation on the corresponding simplices of P . In other words, the map $M(-): P \rightarrow M(P, f)$ is in general not compatible with the two different simplex orders. Nonetheless, the Morse labeling λ_f induced by f orders the nodes of $M(P, f)$ in the same order as their corresponding simplices of P . Thus, connected components of sublevel complexes of (P, f) still correspond to subtrees of $M(P, f)$. Since the induced merge tree assigns the chirality of child nodes according to which connected component carries the minimal value of f , at each inner node of $M(P, f)$ the chirality of the two child nodes is either assigned in accordance with the left/right relation on the corresponding sublevel complex, or it is the opposite. If it is the opposite, this can be corrected by application of the reflection of the corresponding sublevel complex, that is, by application of a sublevel equivalence. In consequence,

the difference between the right/left relation of the corresponding simplices in P and the simplex order only lies in symmetry equivalences of P . Thus, φ can be decomposed into a symmetry equivalence of P and a simplicial isomorphism $\tilde{\varphi}$. Hence, (φ, ψ) is a symmetry equivalence of dMfs on paths.

- (2) Let (T, λ) be an Ml tree. Let $c_0 < c_1 < \dots < c_n$ be the critical values of f_λ and let $\sigma_i \in \Phi T$ such that $f_\lambda(\sigma_i) = c_i$. We recall that the induced merge tree M defines in particular a bijection between the critical simplices of ΦT and the nodes of $M(\Phi T, f_\lambda)$. For any simplex $\sigma \in \Phi T$, we denote the node of $M(\Phi T, f_\lambda)$ that corresponds to σ by $M(\sigma)$. An isomorphism $\varphi: (T, \lambda) \rightarrow M(\Phi T, f_\lambda)$ is given by $\varphi := M \circ \phi^{-1}$. It is immediate that φ is a bijection because M and ϕ are. Furthermore, φ is by construction compatible with the respective Morse labelings. It is only left to show that φ is compatible with the chiral child relation and the respective roots.

Consider $\sigma_n \in \Phi T$. For both trees, the simplex σ_n corresponds to the root of the respective tree. In $M(\Phi T, f_\lambda)$ this is the case because σ_n carries the maximal value of f_λ . In (T, λ) this holds because $\phi(\sigma_n)$ holds the maximal Morse label $\lambda(\phi(\sigma_n)) = c_n$. Thus, the map φ maps the root of (T, λ) to the root of $M(\Phi T, f_\lambda)$.

Let σ_i be a simplex of ΦT . We now prove that φ is compatible with the chiral child relation, that is, that $\varphi(\phi(\sigma_i)_l) = M(\sigma_i)_l / \varphi(\phi(\sigma_i)_r) = M(\sigma_i)_r$ holds. If σ_i is a 0-simplex then there is nothing to show because then both $M(\sigma_i)$ and $\phi(\sigma_i)$ are leaves. Let σ_i be a critical edge with chirality L. The case for chirality R works symmetrically to the case with chirality L.

By Construction 3.14, the node $M(\sigma_i)_l / M(\sigma_i)_r$ corresponds to a critical simplex of the connected component of $\Phi T_{c_i-\varepsilon}$ that carries/does not carry the minimal value of f_λ on these two connected components. Furthermore, the node $M(\sigma_i)_l / M(\sigma_i)_r$ corresponds to the critical simplex that carries the maximal value of f_λ of the respective connected component of $\Phi T_{c_i-\varepsilon}$.

By application of Proposition 4.27, we see that the connected components $\Phi T_{c_i-\varepsilon}[M^{-1}(M(\sigma_i)_l)] / \Phi T_{c_i-\varepsilon}[M^{-1}(M(\sigma_i)_r)]$ induce subtrees of (T, λ) via ϕ . It follows that the node $\phi(\sigma_i)_l / \phi(\sigma_i)_r$ is contained in the subtree that corresponds to $\Phi T_{c_i-\varepsilon}[M^{-1}(M(\sigma_i)_l)] / \Phi T_{c_i-\varepsilon}[M^{-1}(M(\sigma_i)_r)]$ via ϕ because by (2) of Definition 3.17 the subtree with root $\phi(\sigma_i)_l$ does/ $\phi(\sigma_i)_r$ does not carry the minimal Morse label of the subtree with root $\phi(\sigma_i)$ in (T, λ) . Furthermore, the node $\phi(\sigma_i)_l / \phi(\sigma_i)_r$ corresponds to the simplex that carries the maximal value of f_λ of $\Phi T_{c_i-\varepsilon}[M^{-1}(M(\sigma_i)_l)] / \Phi T_{c_i-\varepsilon}[M^{-1}(M(\sigma_i)_r)]$ because by (1) of Definition 3.17 it carries the maximal Morse label on said subtree and because ϕ is order-preserving. Thus, $\varphi(\phi(\sigma_i)_l) = M(\sigma_i)_l / \varphi(\phi(\sigma_i)_r) = M(\sigma_i)_r$ holds. □

By considering Figure 1 we see that the difference between taking the induced merge tree of a dMf on a path is the same as taking its induced Ml tree and forgetting the Morse labeling. Thus, constructing a dMf that represents a given merge tree T is up to symmetry equivalence the same as choosing a Morse order on T . This leads us to the following:

Theorem 6.5. *Let T be a merge tree and let P be a path such that the number of 1-simplices in P is $i(T)$. Then $T \cong M(P, f_{io}) \cong M(P, f_{sc})$ holds as merge trees where f_{io}*

denotes the induced index-ordered dMf (Definition 4.21) and f_{sc} denotes the sublevel-connected dMf (Definition 5.5).

Proof. It is immediate that $M(P, f_{io}) \cong \text{forget}(\text{iMo}(M(P, f_{io}), \lambda_{f_{io}}))$ holds as merge trees, where on the left-hand-side $M(P, f_{io})$ denotes the induced merge tree and on the right-hand-side $M(P, f_{io})$ denotes the induced Ml tree. By Theorem 6.4, $(M(P, f_{io}), \lambda_{f_{io}}) \cong (T, \lambda_{io})$ holds as Ml trees. This implies $\text{iMo}(M(P, f_{io}), \lambda_{f_{io}}) \cong \text{iMo}(T, \lambda_{io})$ as Mo trees by Proposition 3.26. The statement follows because by Definition 3.23 isomorphisms of Mo trees are in particular isomorphisms of the underlying merge trees. Furthermore, $M(P, f_{io}) \cong M(P, f_{sc})$ holds by Theorem 5.8. \square

Using the notion of component-merge equivalence, Definition 3.51, we can extend Theorem 6.4 to a bijection between the set of dMf s with only critical simplices on trees up to cm equivalence DMF_X^{crit} and the set of Ml trees up to isomorphism MIT :

Theorem 6.6. *The induced labeled merge tree $M(-, -)$ and the induced dMf Φ define maps $M(-, -): DMF_X^{\text{crit}} \leftrightarrow MIT: \Phi$ that are inverse to each other in the sense that:*

- (1) *for any dMf (X, f) with only critical cells, the dMf $\Phi(M(X, f), \lambda_f)$ is cm -equivalent to (X, f) , and*
- (2) *for any Ml tree (T, λ) , the Ml tree $M(\Phi T, f_\lambda)$ is isomorphic to (T, λ) .*

Proof. The proof for statement (2) works exactly as in the proof for Theorem 6.4 because symmetry equivalences are in particular cm equivalences. For (1) we apply Proposition 3.54 to consider a representative of the cm equivalence class of (X, f) which is a dMf on a path (P, f') . By Proposition 3.53, the isomorphism type of the induced Ml tree does not depend on this choice. Thus, Theorem 6.4 implies that $\Phi(M(P, f'))$ is symmetry-equivalent to (P, f') . Since (P, f') is cm -equivalent to (X, f) , so is $\Phi(M(P, f'))$. \square

Corollary 6.7. *Let T be a merge tree. By Theorem 6.5 and Theorem 6.6 it follows that there are dMf s on trees (X, f) such that $M(X, f) \cong T$ as merge trees.*

Corollary 6.8. *Applying Theorem 6.6 together with Proposition 3.26 and Proposition 3.41 yields the result that there is a bijection $DMF_X^{\text{crit}}/\sim \cong MoT$ where \sim denotes order equivalence.*

We conclude this section by discussing our results and comparing them to the results of [Cur19]. Using the bijections appearing in Figure 1 and Theorem 6.6 we replaced the question of finding dMf s on paths or arbitrary trees that represent a given merge tree T by finding Morse orders on T instead. This argument can be used to replace the question of classifying merge equivalence classes of dMf s on trees by classifying merge equivalence classes of Mo trees. Example 6.2 tells us that, if a merge tree T has at least three leaves, there might be different Morse orders on T which are not shuffle-equivalent. That is, there are dMf s on trees (X, f) contained in the image of $\Phi \circ iMl$ which induce T as their merge tree but are neither isomorphic, nor symmetry-equivalent, nor cm -equivalent, nor shuffle-equivalent, nor a combination of the four to each other. A complete classification of merge-equivalence classes of dMf s on trees would hence need to classify Morse orders on merge trees as well. We found a non-empty shuffle equivalence class of Morse orders on any merge tree, defined by either the index Morse order or the sublevel-connected Morse order, since the two have been shown to be shuffle-equivalent and, hence, merge-equivalent in Theorem 5.8. This

allowed us to answer the aforementioned question of Johnson–Scoville affirmatively. That is, any merge tree is indeed represented by a dMf on a path, which is induced by a Morse order. Using a non-trivial cm equivalence, one can also find a dMf on a tree other than a path as a representative.

Furthermore, our results from Section 3.1, Section 3.2, and Section 6 allow us to structure the study of the set of merge equivalence classes of dMfs on trees using the following four notions of merge-invariant equivalences between dMfs on trees:

- Forman equivalence,
- symmetry equivalence,
- cm equivalence, and
- shuffle equivalence.

We discussed in Remark 6.1 how Forman equivalences could be considered together with the other notions of equivalence and why we left Forman equivalences out of this work. In Theorem 6.4 it becomes quite clear that passing to the induced Ml tree identifies symmetry-equivalent dMfs with each other up to isomorphism. Passing from the induced Ml tree to the induced Mo tree then identifies dMfs up to order equivalence with each other up to isomorphism of the underlying Mo tree. Last of all, passing from the induced Mo tree to the induced merge tree in particular identifies shuffle-equivalent dMfs with each other. This allows us to study the different equivalence classes separately for dMfs on paths. The more liberal notion of cm equivalence, Definition 3.51, allowed us to generalize Theorem 6.4 to dMfs on arbitrary trees as seen in Theorem 6.6.

In [Cur19], the author establishes a bijection between graph-equivalence ([Cur19, Def 6.1]) classes of Morse-like ([Cur19, Def 6.9]) continuous functions on the interval that attain minima at the boundary on the one hand and isomorphism classes of chiral merge trees ([Cur19, Def 5.3]) on the other hand.

At first glance it might seem likely that Ml trees in the sense of Definition 3.19 and chiral merge trees in the sense of [Cur19, Def 5.3] are directly related by geometric realization and considering the corresponding abstract simplicial complex but, as mentioned in the introduction, there is a subtle difference in the construction of the induced merge tree. To be precise, the two constructions only differ in the induced chirality. In [Cur19, Sec 5] the chirality is given by which of the two merging components is the left or right one with respect to the chosen orientation on the interval. In [JS20] the chirality is given such that drawing the induced merge tree is compatible with the elder rule: The component with the minimal value gets the same chirality as the merged component. This means that following the same chirality leads to the oldest component.

This convention leads to the necessity to assume property (2) of Definition 3.17: a certain compatibility between the Morse order and the chirality. The compatibility between Morse orders and the chirality implies, as seen in Proposition 3.49, that the induced Ml tree does not distinguish between symmetry-equivalent dMfs. If one defines induced Ml trees analogously to [Cur19], that is by inducing the chirality by a chosen orientation of the path, this notion of induced Ml trees would distinguish symmetry-equivalent dMfs on paths. Moreover, Ml trees induced by symmetry-equivalent dMfs would be related by sequences of reflections of subtrees. The definition could be as follows:

Definition 6.9. Let T be a merge tree. A *Curry Morse order* is a total order \leq on the nodes of T such that the maximal node of any subtree is the root of said subtree. A *Curry Morse labeling* on a merge tree T is a labeling λ on the nodes of T that induces a Curry Morse order on T .

A pair (T, \leq) of a merge tree with a Curry Morse order on it is called a *Curry Morse ordered merge tree (CMo tree)*. A pair (T, λ) of a merge tree with a Curry Morse labeling on it is called a *Curry Morse labeled merge tree (CML tree)*.


Let $f: P \rightarrow \mathbb{R}$ be a dMf where P is an oriented path. Then the CML tree induced by f is defined as follows:

Let $c_0 < c_1 < \dots < c_m$ be the critical values of f that are assigned to 1-simplices. The induced CML tree $M_C(P, f)$ is constructed by induction over these critical values in descending order. Furthermore, we label the nodes n of $M_C(P, f)$ with $\lambda(n)$.

For the start of the induction we begin by creating a node n_m which corresponds to the critical 1-simplex in P labeled c_m and setting its label $\lambda(n_m) := c_m$ and its chirality to L .

For the inductive step, let n_i be a node of $M(X, f)$ that corresponds to a critical 1-simplex between two 0-simplices v and w . Define $\lambda_v := \max\{f(\sigma) \mid \sigma \in P_{c_i-\varepsilon}[v], \sigma \text{ critical}\}$ and $\lambda_w := \max\{f(\sigma) \mid \sigma \in P_{c_i-\varepsilon}[w], \sigma \text{ critical}\}$. Two child nodes are created, named n_{λ_v} and n_{λ_w} . Then label the new nodes $\lambda(n_{\lambda_v}) := \lambda_v$ and $\lambda(n_{\lambda_w}) := \lambda_w$. If $P_{c_i-\varepsilon}[v]$ is left of $P_{c_i-\varepsilon}[w]$ with respect to the orientation on P , we assign n_{λ_v} the chirality L and give n_{λ_w} the opposite chirality. Continue the induction over the rest of the critical 1-simplices.

Example 6.10. We consider two of the symmetry-equivalent dMfs from Example 3.48 and see how the induced CML tree distinguishes them whereas the induced ML trees identifies them as one:



The notion of CML trees is related to the notion of chiral merge trees in the sense of [Cur19] by the interplay between abstract and geometrical simplicial complexes. In detail, the bijection is given as follows:

Construction 6.11. Let (T, λ) be a CML tree. We define a chiral merge tree $|(T, \lambda)|$ associated to (T, λ) as follows:

The compact rooted tree is given by the geometric realization $|T|$. We attach a distinguished edge e_∞ to the vertex which corresponds to the root of T in order to obtain a cell complex which we will by abuse of notation also refer to as $|T|$. The map $\pi: |T| \rightarrow \mathbb{R}$ is given by λ on vertices, and by a linear extension of λ on edges. To make the linear extensions on edges between parent nodes p and child nodes c unique, we assume that every edge in $|T|$ has length one and choose extensions of the form $\lambda(c) + t \cdot (\lambda(p) - \lambda(c))$, where $t \in [0, 1]$. The chirality on the edges in $|T|$ is given by the chirality of the child nodes adjacent to the corresponding edges in T .

For the other way around let $\pi: T \rightarrow \mathbb{R}$ be a chiral merge tree in the sense of [Cur19, Def 5.3]. We define a merge tree $\text{abs}(T)$ associated to T as follows:

We take the 0-skeleton T_0 as the vertex set and the 1-skeleton $(T \setminus \{e_\infty\})_1$ as the set of edges. We define the node which corresponds to v_∞ to be the root of $\text{abs}(T)$. The chiral child relation is induced by the chirality of the edges between the corresponding parent and child nodes in T . The labeling λ is directly induced by π . The proof that the two constructions are inverse to each other is straightforward.

Furthermore, there is a similar bijection between the two notions of Morse functions:

Construction 6.12. Let (P, f) be a dMf with only critical cells on an oriented path. We define a Morse-like function \tilde{f} on the interval which attains minima at the boundary as follows:

Let $k + 1$ be the number of 0-simplices in P . We denote the 0-simplices of P by n_0, n_1, \dots, n_k from left to right with respect to the given orientation. Then we define $\tilde{f}(\frac{i}{k}) := f(n_i)$ for $i \in \{0, 1, \dots, k\}$. We denote the 1-simplices of P by e_1, \dots, e_k , again according to the given orientation. Then we define $\tilde{f}(\frac{2i-1}{2k}) := f(e_i)$ for $i \in \{0, 1, \dots, k\}$. We define \tilde{f} on the rest of the interval as the linear extension. This makes \tilde{f} a distinct-valued PL function that attains minima at the boundary. Hence, \tilde{f} is Morse-like.

For the other way around let $f: I \rightarrow \mathbb{R}$ be a Morse-like function that attains minima at the boundary. We define a path P and a dMF f' on P as follows:

Let n_0, n_1, \dots, n_k be the local minima of f ordered by the orientation on I , so in particular $n_0 = 0$ and $n_k = 1$. We define P to be a path with $k+1$ simplices of dimension 0. We choose one of the 0-simplices which is a face of only one 1-simplex to be denoted by s_0 and the other such simplex by s_k . We denote the other 0-simplices such that their indices are in accordance with their position in the simplex order (Definition 4.16), making s_0 the minimal simplex with respect to the simplex order. The dMf f' is defined by $f'(s_i) := f(n_i)$ on 0-simplices. Let c_1, \dots, c_k be the local maxima of f , ordered in accordance to the orientation on I , and let e_1, \dots, e_k be the 1-simplices of P , ordered in accordance to the aforementioned simplex order on P . Then f' is defined by $f'(e_i) := f(c_i)$ on 1-simplices.

It is easy to check that the two given constructions are inverse to each other.

A theorem which, analogously to Theorem 6.4, defines a pair of inverse bijections $M_C(-, -): {}^+DMF_P^{\text{crit}} \leftrightarrow CMIT: \Phi$ can be proved similarly to the proof of Theorem 6.4. The difference is that the induced CMI tree keeps track of symmetry equivalences in the sense that symmetry equivalences of dMfs induce reflections at roots of subtrees on the induced CMI trees. The map Φ can be defined the same way as before. Alternatively, one could define the bijection Φ as the composition $\text{abs} \circ \Psi^{-1} \circ |_$ in the following diagram:

$$\begin{array}{ccccccc}
Mer & \xleftarrow{M(-, -)} & DMF_P^{\text{crit}} & \xleftarrow{o_{JS}} & +DMF_P^{\text{crit}} & \xleftarrow{|\cdot|} & \mathcal{M} \\
\downarrow \leq_{sc} & \uparrow \leq_{io} & \uparrow \text{forget} & \uparrow /_{symm} & \uparrow \Phi & \uparrow \text{abs} & \uparrow \Psi \\
MoT & \xleftarrow{iMl} & MIT & \xleftarrow{i} & CMIT & \xleftarrow{|\cdot|} & \mathcal{X} \\
& \xrightarrow{iMo} & & \xrightarrow{JS} & & \xrightarrow{\text{abs}} & \\
& & \uparrow \Phi & & \uparrow \Phi & & \uparrow \Psi^{-1}
\end{array}$$


FIGURE 2. Relationship to the continuous case

Here Ψ denotes the bijection between the set of Morse-like functions on the interval \mathcal{M} and the set of chiral merge trees \mathcal{X} from [Cur19, Cor 6.11]. The map i is the inclusion induced by considering Ml trees as CMI trees. Ml trees can be considered as a special kind of CMI trees because they only differ in the additional property (2) of Definition 3.17 which does not need to hold for CMI trees. The map $/_{symm}$ is the quotient map that identifies symmetry equivalent dMFs. The map o_{JS} is defined as the composition $\Phi \circ i \circ M(-, -)$. This definition coincides with choosing the representative of a dMf (P, f) with respect to symmetry equivalence such that the simplex order on P is compatible with f in the following way: At each critical edge e with $f(e) = c$ the connected component of $P_{c-\varepsilon}$ that corresponds to $M(e)_l/M(e)_r$ is left/right of the edge e with respect to the orientation induced by the simplex order. Here we recall that $M(e)$ denotes the inner node of $M(P, f)$ that corresponds to the critical edge e and that $M(e)_l/M(e)_r$ denotes the left/right child node of $M(e)$. In Example 6.10 we have $o_{JS}(P, g) = (P, f)$ where (P, f) is oriented from left to right.

The map JS is defined as $JS := M(-, -) \circ /_{symm} \circ \Phi$. This definition coincides with division by the equivalence relation generated by reflections of subtrees.

Example 6.13. We consider how a sequence of reflections of subtrees maps the CMI trees from Example 6.10 to each other. Here, we denote by a_λ the reflection of the

subtree with root labeled λ .



The proof that the two possible definitions for Φ coincide and that the diagram from Figure 2 commutes are straightforward.

As a result, the map Φ can be seen as a discrete version of the map Ψ from [Cur19, Cor 6.11]. Moreover, moving from the setting of [Cur19] to the setting of [JS20] basically means to divide by symmetry equivalences which allows the authors of [JS20] to generalize the construction of the induced merge tree to dMfs on trees. In order to extend our generalization Theorem 6.6 to the oriented case, one would need to find a notion of orientation-preserving cm equivalences which distinguishes symmetry-equivalent dMfs.

Furthermore, we conjecture that the correspondence between dMfs, Ml trees and Mo trees can be categorified. That is, there are categories of dMfs, Ml trees and Mo trees that have the aforementioned types of equivalences as isomorphisms and the aforementioned bijections define equivalences of categories.

7. FURTHER DIRECTIONS AND POSSIBLE APPLICATIONS

In this section we want to take a look at possible applications of our results. The structured overview of the different notions of equivalence of discrete Morse functions and their connections to each other might be useful to explore the space of discrete Morse functions on a given simplicial complex. Even though the construction of the induced merge tree from [JS20] does not easily extend to arbitrary simplicial complexes, the notions of equivalence from this article do. Hence, one could try to use e.g. the notion of symmetry equivalence to structure the space of discrete Morse functions in a nice way, i.e. into orbits of a groupoid action. In a second step, one could then try to define the space of merge trees as a quotient of the space of discrete Morse functions on some “large enough” simplicial complex. With such a construction, one could assemble

the results from this article and the results from [CDG⁺21] into an analysis of a larger instance of the persistence map.

Aside from this, one could try to generalize and enhance the construction of the induced merge tree from [JS20] to arbitrary simplicial complexes. Then, in a second step, one could try to use the induced merge tree to find possible cancellations of pairs of critical simplices. This way, the induced merge tree might be helpful to optimize discrete Morse functions.

Furthermore, one could study the possible Morse orders on a given merge tree. With a classification of all Morse orders on a given merge tree T , one could classify all discrete Morse functions on any given tree that induce T as the induced merge tree with the help of Definition 4.21 and cm equivalences.

Ultimately, the proofs of Theorem 6.6 and all the lemmas that lead to it describe exactly which information is lost when considering the induced Ml tree instead of the original dmf. This knowledge might be useful for applications in TDA because it tells the user which kind of features will not be seen by the induced merge tree, and hence, by the persistent zeroth homology and the barcode. Moreover, the knowledge about the exact lost information can be used to enhance the induced merge tree with extra structure such that it no longer disregards certain desired information.

APPENDIX A. STEP-BY-STEP CONSTRUCTION OF THE INDUCED INDEX-ORDERED DMF

The basic idea is to reconstruct T node by node, following the values of the index-ordered dmf f_{io} from higher values to lower values. This means that we build P and $f: P \rightarrow \mathbb{N}_0$ simplex by simplex according to the nodes of T .

The idea of reverse-engineering the construction of the induced merge tree is that each time a **left/right** node is being considered, the smallest value of its subtree is supposed to be in the **left/right** direction. Thus, the next highest value is supposed to be to the **right/left**. We always continue in the direction where the next highest value is supposed to be and add a new 1-simplex to P for each inner node we pass on the way. This corresponds precisely to considering the nodes of T from highest to lowest with respect to the index Morse order.


In each step of the construction we assign a label to the new 1-simplex of P such that it will induce a corresponding node in $M(P, f)$. The labels of the pre-existing simplices will be modified in an order preserving way such that the existing nodes of $M(P, f)$ will remain unchanged.

Before we actually start with the inductive construction, we need more notation:

Notation A.1. For an inner node c of T we denote the 1-simplex of P corresponding to c in the step-by-step construction by $s(c)$. It will be clear from the construction that this correspondence defines a bijection $s: V(T) \rightarrow P$ between the nodes of T and the simplices of P . The map s will turn out to be the inverse of the map ϕ from Proposition 4.19.

For the construction, we imagine P as being horizontally embedded in a plane. This way, every 1-simplex of P has a left and a right 0-simplex as its boundary. We denote the left boundary 0-simplex of a simplex $s(c)$ by $s_l(c)$ and the right boundary 0-simplex by $s_r(c)$. If we want to assign labels to all three, $s_l(c)$, $s(c)$ and $s_r(c)$ simultaneously, we write that we label $\bar{s}(c)$ with a triple (x, y, z) . That is, $s_l(c)$ is labeled with x , the


simplex $s(c)$ is labeled with y and $s_r(c)$ is labeled with z . Additionally, we recall that by Notation 3.11 we denote the left child of c with c_l and the right child with c_r . We illustrate this notation in the following example, where we label $\bar{s}(p)$ with (a, b, x) and $\bar{s}(c)$ with (x, y, z) :



For a node c of T we denote the path P that is obtained directly after considering c by P_c . The path that is obtained directly before c has been considered is denoted by $P_{c-\varepsilon}$. The path P_c is labeled by the map f_c and $P_{c-\varepsilon}$ is labeled by $f_{c-\varepsilon}$.

Analogously we denote the filled tree that contains all nodes of T that have been considered up until c by T_c . In particular, this means that c is also included in T_c . By the adjective ‘filled’ we mean that we artificially add leaves to T_c such that all considered nodes are inner nodes. We denote by $T_{c-\varepsilon}$ the filled tree that contains the nodes of T that have been considered up until c , except c . That is, T_c and $T_{c-\varepsilon}$ coincide except for that in $T_{c-\varepsilon}$ there is just a leaf where in T_c there is a subtree consisting of c and two leaves.

A.1. Step-by-Step Construction. We begin at the root ρ of T and start the construction of P and f with a 1-simplex which is labeled $(0, 2, 1)$. That is:



If the right child ρ_r of the root ρ is an inner node, we continue with ρ_r . Else, if the left child ρ_l of the root ρ is an inner node, we continue with ρ_l . Otherwise, we are already done.

For the inductive step, we continue as long as there are any inner nodes that have not been considered yet. Assume we are considering an inner node c that has not been considered yet. The inductive step depends on the chiralities of c and c ’s parent node p . That leaves us with four cases:

LR/RL First we suppose that c is a **right/left** child of a **left/right** parent node p . We attach a new 1-simplex $s(c)$ directly to the **right/left** of $s(p)$, that is, $s_l(c) = s_r(p)/s_r(c) = s_l(p)$. If there was a 1-simplex σ directly **right/left** of $s(p)$ beforehand, $s(c)$ is placed in between $s(p)$ and σ , that is, $\sigma_l = s_r(c)/\sigma_r = s_l(c)$. If $\bar{s}(p)$ was labeled (x, y, z) , we label $\bar{s}(c)$ with $(z + 1, y + 1, x + 1)/(z + 1, y + 1, x + 1)$. Then we increase the labels of all pre-existing 1-simplices by 2. Increase the labels of all pre-existing 0-simplices with labels λ such that $\lambda > z/\lambda > x$ by 1. We declare that c has been considered.

If c_l/c_r is an inner node, then proceed with c_l/c_r . Else, if c_r/c_l is an inner node, continue with c_r/c_l . Else, go back to p .

RR/LL Then we assume that c is a **right/left** child of a **right/left** parent node p . We attach a new 1-simplex $s(c)$ directly to the **right/left** of $s(p)$, that is, $s_l(c) = s_r(p)/s_r(c) = s_l(p)$. If there was a simplex σ directly **right/left** of $s(p)$ beforehand, $s(c)$ is placed in between, that is, $\sigma_l = s_r(c)/\sigma_r = s_l(c)$. Let (x, y, z) be the label of $\bar{s}(p)$. Define $m := \min\{\lambda | \lambda \text{ is the label of a simplex } s(d) \text{ for an inner node } d \text{ of the subtree with root } p \text{ in } T_{c-\varepsilon}\}$. We label $\bar{s}(c)$ with $(z+1, m+1, z)/(x, m+1, x+1)$. Then we increase the labels of all pre-existing 1-simplices by two and the labels of the pre-existing 0-simplices which had labels $\lambda > z/\lambda > x$ by 1. We declare that c has been considered.

If c_l/c_r is an inner node, then proceed with c_l/c_r . Else, if c_r/c_l is an inner node, continue with c_r/c_l . Else, go back to p .

If we are at a node c that already has been considered, there are two cases:

- L c is left node, so c_r has already been considered. If c_l is an inner node that has not been considered yet, proceed with c_l . Else, proceed with c 's parent node.
- R c is a right node, so c_l has already been considered. If c_r is an inner node that has not been considered yet, proceed with c_r . Else, proceed with c 's parent node.

As long as there are nodes of T that have not been considered yet, we keep following the described algorithm.

A.2. Properties of the Step-by-Step Construction. Before we take a closer look at the relationship between T and $M(P, f)$ we first notice that the algorithm visits every relevant node of T . We observe that leaves are irrelevant to the algorithm since their data is given implicitly by their parent nodes. Hence we only consider inner nodes.

Lemma A.2. *The algorithm visits all inner nodes of T from highest to lowest, corresponding to the index Morse order \leq_{io} .*

Proof. We consider the order in which the step-by-step algorithm visits the nodes of T and show that the node which is currently being considered always covers the node which is next in line. Since all leaf nodes are ranked lower than any inner node by the Morse order, we do not need to consider them here. In the entire proof, the covering relation is always meant with respect to the index Morse order.

Whenever an inner node p has been considered for the first time, the algorithm moves on to p 's child node c if c is an inner node with the other chirality than p 's. We now prove that in this case p covers c .

Because c has a different chirality than p , by case a) of Definition 4.3, the node c is ranked higher than any inner node of the subtree with p 's other child c' as root. Furthermore, by case b), the node c is also ranked higher than any other inner node of the subtree with root c . Additionally, $c \preceq p$ holds by case b) of Definition 4.3. Moreover, c is lower than any ancestor of p by case b). For any other inner node \tilde{c} of T which is neither an ancestor nor a descendant of p , the same relation between c and \tilde{c} holds as it does between p and \tilde{c} . Thus, the node p covers c in the index Morse order.

If c is a leaf, the step-by-step algorithm proceeds with the other child c' if it is an inner node. In that case one can prove that p covers c' with similar arguments as we have used to prove that p covers c in the previous case.

If both children of p are leaves, the algorithm goes back to the youngest ancestor \tilde{p} of p that has an inner node q' which has not yet been considered as a child. Let q be the other child of \tilde{p} . Because q was by assumption considered before q' , the node q has another chirality than \tilde{p} and q' has the same chirality as \tilde{p} . Thus, the node q' is considered to be less than p in the index Morse order because the corresponding path words differ after the position that corresponds to \tilde{p} . At that point, the path word corresponding to p continues with the other letter whereas the path word corresponding to q' has the same letter.

We now show that the node p is the smallest inner node in the subtree with root q : For the nodes on the shortest path between q and p it follows directly by case b) that p is smaller. In particular, $p \leq_{io} q$ holds by b). If there are other nodes in the subtree with root q which do not lie on the shortest path between q and p , it follows inductively that they have already been considered. Otherwise it would be a contradiction to the choice of \tilde{p} to be the youngest such ancestor.

Thus, it follows inductively that these nodes, together with the nodes on the shortest path between q and p , form a chain of cover relations with respect to the index Morse order.

Because p is smaller than q and the nodes that have been considered before q' are part of this chain of cover relations, p must be smaller than all nodes of the chain. In conclusion, the node p is the smallest node of the subtree with root q with respect to the index Morse order.

Additionally, all other inner nodes of the subtree with root q' are smaller than q' by case b) and it can be proved similarly as before that the relationship between any other inner node of T and q' is the same as with p .

In conclusion, the step-by-step algorithm considers the nodes in the descending index Morse order. That is, it follows a zig-zag path down to an inner node that has two leaves as children. Then the step-by-step algorithm alternates between going to the sibling, going down the sibling's subtree in a zig-zag, and going back up a level. This way, once a node c has been visited for the first time, the algorithm recursively considers all nodes of c 's subtree. Because the algorithm starts at the root, every node of T is considered. \square

We take a closer look at some properties of the step-by-step construction:

Lemma A.3. *The 1-simplex $s(c)$ has the smallest label among all 1-simplices of P_c . Furthermore, if \tilde{c} is the node that was considered directly before c , then the label of $s(\tilde{c})$ in P_c is exactly 1 higher than the label of $s(c)$.*

Proof. We prove the statement by induction over the nodes of T in the same order as they are considered by the construction algorithm: For the initial case, let $c = r$ be the root of T . Then T_r and (P_r, f_r) look like this:



The statement obviously holds. For the inductive step, we assume that for some node c of T the statement holds for the node \tilde{c} which was considered directly before c . In the cases **LR** and **RL**, the parent node p was considered directly before c , so $\tilde{c} = p$. By inductive hypothesis, this implies that $s(p)$ has the smallest label among 1-simplices in $P_{c-\varepsilon}$. Thus, by construction, the label of $s(c)$ in P_c is exactly 1 higher than the label of $s(p) = s(\tilde{c})$ in $P_{c-\varepsilon}$. In the cases **LL** and **RR** \tilde{c} is an inner node of the subtree with root p in $P_{c-\varepsilon}$. By construction, $s(c)$ gets a label exactly 1 higher than the label of $s(\tilde{c})$, because by inductive hypothesis $s(\tilde{c})$ has the smallest label among the simplices induced by nodes of the subtree with root p . In conclusion, $s(c)$ is assigned a label exactly 1 higher than the smallest value assigned to 1-simplices in $P_{c-\varepsilon}$. Because in every of the four cases the labels of all pre-existing 1-simplices are increased by 2, $s(c)$ has now the smallest label among all 1-simplices in P_c . \square

Remark A.4. If the given merge tree T has $i(T)$ inner nodes, it follows inductively from Lemma A.3 that the labels on the 1-simplices of P are $i(T) + 1, \dots, 2 \cdot (i(T))$. Furthermore, the order of the labels corresponds to the inverse order in which the corresponding nodes of T were considered.

We consider the relationship between the index Morse labeling and the function from the step-by-step approach:

Lemma A.5. *For every inner node c of T , the label on $s(c)$ as defined in the step-by-step construction is the same as the index Morse label $\lambda(c)$.*

Proof. It follows by Remark A.4 and Lemma A.2 that for an inner node c of T the label that the step-by-step construction assigns to the 1-simplex $s(c) \in P$ is the same as the index Morse label $\lambda(c)$ defined in Definition 4.7. \square

Lemma A.6. *For any leaf c in T , the label of $s(c)$ as constructed in the step-by-step construction coincides with the index Morse label $\lambda(c)$.*

Proof. Now we consider the labels on the leaves of T . The leaf nodes inherit the index Morse ordering from their parent nodes because their path words contain their parent's path words. In the step-by-step construction, the labels of the 0-simplices are constructed alongside their corresponding 1-simplices. As long as the algorithm continues on inner nodes of alternating chirality, it constructs new 0-simplices with increasing labels along the way. The leaves corresponding to these 0-simplices are increasing with respect to the index Morse order as the labels increase. This is because the alternating chirality of their parent nodes induces alternating letters in the corresponding path words.

Whenever the step-by-step algorithm visits a node c that has the same chirality as its parent node p , from now on the labels of all 0-simplices corresponding to the previously

considered descendants of p will be increased in every later step of the construction by $+1$ because their current labels are higher than the label of the 0-simplex corresponding to the leaf that was at c 's place in $T_{c-\varepsilon}$. Thus, all 0-simplices that are constructed in later steps of the construction will have lower labels than the ones corresponding to pre-existing descendants of p .

The leaves of T which are children of inner nodes that are considered later by the step-by-step construction algorithm are smaller with respect to the index Morse order because the same is true for their parent nodes. It can be proved inductively that the leaves of T are ordered by the index Morse order in the same order as by the labels of their corresponding 0-simplices. It follows by the same inductive argument that these labels are $\{0, 1, \dots, l(T) - 1 = i(T)\}$. \square

We now consider the relationship between the placement of simplices in the step-by-step construction and the simplex orders on P and T .

Lemma A.7. *For nodes a, b of T , the relation $a \prec b$ holds if and only if $s(a)$ is left of $s(b)$ in the sense of Appendix A.1.*

Proof.

\Rightarrow We consider the different cases from Definition 4.11 that the relation $a \prec b$ might correspond to. The colors in this proof are only used for conciseness and do not correspond to the colors used in Appendix A.1. First we assume that $a \prec b$ corresponds to case (1):

Let p be the youngest common ancestor of a and b . Then a/b is a node of the subtree with root p_l/p_r . In the construction algorithm from Appendix A.1, the simplex $s(p_l)/s(p_r)$ is placed **left/right** of $s(p)$. Since the algorithm from Appendix A.1 places new simplices directly next to the ones of the corresponding parent nodes, all simplices which correspond to nodes of the subtree with root p_l/p_r are placed **left/right** of $s(p)$. Thus, the simplex $s(a)/s(b)$ is placed **left/right** of $s(p)$, so in particular $s(a)$ is placed left of $s(b)$.

If $a \prec b$ corresponds to case (2)/(3), then a/b is a node of the subtree with root b_l/a_r . The statement follows by the same argument as for the nodes of the subtree with root p_l/p_r and p in case (1).

\Leftarrow Let $s(a)$ be left of $s(b)$ in P . Suppose $a \prec b$ does not hold. Then, by totality of \preceq , $b \preceq a$ must hold. If $a = b$ holds, then $s(a) = s(b)$ holds, which is a contradiction to the assumption. If $b \prec a$ holds, then $s(b)$ is left of $s(a)$, which is also a contradiction to the assumption. Thus, $a \prec b$ must hold. \square

Proposition A.8. *Let ϕ be the map from Proposition 4.19 where p_0 from Definition 4.16 was chosen to be the leftmost 0-simplex in the sense of the step-by-step construction. Then the assignment $s: T \rightarrow P$ that assigns each node c of T its corresponding simplex $s(c)$ as in Appendix A.1 is the inverse of ϕ as maps of totally ordered sets.*

Proof. Let σ and σ' be simplices of P . We show that $\sigma \prec \sigma'$ holds if and only if $s(\phi(\sigma)) \prec s(\phi(\sigma'))$ holds in P .

The relation $\sigma \prec \sigma'$ holds in P if and only if $\phi(\sigma) \prec \phi(\sigma')$ holds in T by Proposition 4.19. By Lemma A.7, the relation $\phi(\sigma) \prec \phi(\sigma')$ holds in T if and only if $s(\phi(\sigma))$ is left of $s(\phi(\sigma'))$ in P . By Remark 4.18, $s(\phi(\sigma))$ is left of $s(\phi(\sigma'))$ in P if and only if $s(\phi(\sigma)) \prec s(\phi(\sigma'))$ holds.

It follows now inductively that $s \circ \phi$ is the identity on P because P is a finite totally ordered set and $s \circ \phi$ is bijective and order-preserving as a composition of two order-preserving bijections. Thus, $s = \phi^{-1}$ holds. \square

Theorem A.9. *Let f be the dMf from the step-by-step approach and let f_{io} be the induced index-ordered dMf from the all-at-once approach (see Definition 4.21). Then $f = f_{io}$ holds.*

Proof. By Proposition A.8, the step-by-step construction assigns nodes c of T to simplices $s(c)$ such that $\phi(s(c)) = c$. By Lemma A.5 and Lemma A.6, the index Morse label $\lambda(c)$ is equal to the value of the Morse function $f(s(c))$ from Appendix A. In conclusion, by Definition 4.21, it follows that for any simplex s of P the equation $f_{io}(s) := \lambda(\phi(s)) := \lambda(c) = f(s(c)) = f(s)$ holds. \square

We present an alternative proof for the fact that the step-by-step construction produces an index-ordered dMf without using that we already know that f_{io} is an index-ordered dMf.

Lemma A.10. *If $\bar{s}(c)$ is labeled $(\alpha_c, \beta_c, \gamma_c)$ in P_c the following holds, depending on the chirality of c :*

$$\begin{aligned} L \quad & \gamma_c = \alpha_c + 1 \\ R \quad & \alpha_c = \gamma_c + 1 \end{aligned}$$

Furthermore, if $\bar{s}(c)$ is labeled (α, β, γ) in P , the following holds:

$$\begin{aligned} L \quad & \alpha < \gamma \\ R \quad & \gamma < \alpha \end{aligned}$$

Proof. Although the statement itself does not rely on the chirality of c 's parent node p , the proof does. In the cases **LL** and **RR**, the first statement follows directly by construction.

For the other two cases we observe that before a node c is considered during an **LR/RL** step, the preceding step is one of **L (the initial step)**, **LL**, **RL/RR**, **LR** and considers c 's parent node p . Thus, we prove the first statement in these cases inductively.

For the three base cases we observe that if the previous case was **L**, **LL/RR**, by construction, $\bar{s}(p)$ was labeled $(x, y, x + 1)$ for some $x, y \in \mathbb{N}/(z + 1, y, z)$ for some $y, z \in \mathbb{N}$. Thus, by construction, $\bar{s}(c)$ is labeled $(x + 2, y + 1, x + 1)/(z + 1, y + 1, z + 2)$.

For the inductive step it follows by inductive hypothesis that $\bar{s}(p)$ was labeled $(x, y, x + 1)$ for some $x, y \in \mathbb{N}/(z + 1, y, z)$ for some $y, z \in \mathbb{N}$. Thus, by construction, $\bar{s}(c)$ is labeled $(x + 2, y + 1, x + 1)/(z + 1, y + 1, z + 2)$. This concludes the proof of the first statement.

For the second statement we remark that the first statement implies that the desired inequalities for c hold in P_c . It is left to show that the following steps of the construction do not affect these inequalities.

If c has chirality **L/R** in T , the subtree with root c_r/c_l is considered completely before the subtree with root c_l/c_r is considered. While the subtree with root c_r/c_l is being considered, the label of $s_r(c)/s_l(c)$ increases during some steps, whereas the label of $s_l(c)/s_r(c)$ remains unchanged because inductively the label of 0-simplices corresponding to leaves of the subtree with root c_r/c_l is higher than the label of $s_l(c)/s_r(c)$. The increase of the label of $s_r(c)/s_l(c)$ only occurs when a step inserts a new 1-simplex which has $s_r(c)/s_l(c)$ as its **left/right** boundary simplex.

When the subtree with root c_l/c_r is being considered, the label of $s_l(c)/s_r(c)$ increases during some steps, but the label of $s_r(c)/s_l(c)$ increases during every step because inductively, the label of $s_r(c)/s_l(c)$ is higher than the label of 0-simplices corresponding to leaves of the subtree with root c_l/c_r . The increase of the label of $s_l(c)/s_r(c)$ only occurs when a step inserts a new 1-simplex which has $s_l(c)/s_r(c)$ as its right/left boundary simplex.

The inequalities are also not affected in later steps of the construction because the labels of 0-simplices corresponding to nodes of the subtree with root p in T form an interval in \mathbb{N} . This can be proven using the same argument as for the first statement of this lemma together with the observation that the subtree with root p in T is considered completely directly after p before any other node of T is considered.

In conclusion it follows inductively that the inequalities hold in P . \square

Lemma A.11. *The function f assigns strictly lower values to 0-simplices than to 1-simplices of P_c . Furthermore, the lowest label assigned to 1-simplices is exactly 1 higher than the highest label assigned to 0-simplices.*

Proof. The initial case is again straightforward. For the inductive step we assume again that for some node c of T the statement holds for the node \tilde{c} which was considered directly before c . Define $a := \max\{\lambda \in \mathbb{R} \mid \lambda \text{ is label of a 0-simplex of } P_{c-\varepsilon}\}$ and $b := \min\{\lambda \in \mathbb{R} \mid \lambda \text{ is label of a 1-simplex of } P_{c-\varepsilon}\}$ as well as $\alpha := \max\{\lambda \in \mathbb{R} \mid \lambda \text{ is label of a 0-simplex of } P_c\}$ and $\beta := \min\{\lambda \in \mathbb{R} \mid \lambda \text{ is label of a 1-simplex of } P_c\}$. By inductive hypothesis, $a + 1 = b$ holds and by Lemma A.3: $\beta = b + 1$. By construction, only labels that are at most 1 higher than pre-existing labels of 0-simplices in $P_{c-\varepsilon}$ are assigned to 0-simplices of P_c . Thus, $\alpha \leq a + 1$ holds. It follows that $\alpha \leq a + 1 = b < b + 1 = \beta$.

We now show the inductive step for the fact that $\alpha = a + 1$ holds. It is only left to show $\alpha \geq a + 1$. For that we again distinguish between the four cases depending on the chirality of c and p :

LR/RL If $\bar{s}(p)$ was labeled (x, y, z) in $P_{c-\varepsilon}$, $a \geq z/a \geq x$ holds. If $a = z/a = x$ holds, we consider that $s_l(c)/s_r(c)$ is labeled $z+1/x+1$ and since there was no label on 0-simplices larger than z/x beforehand, no other label on 0-simplices is increased. In conclusion, $\alpha \geq a + 1$. If $a > z/a > x$ holds, the label a is not assigned to a simplex of $s(p)$. Nonetheless the label a is increased by 1 because $a > z/a > x$ holds in $P_{c-\varepsilon}$. Thus, $\alpha \geq a + 1$ holds.

RR/LL If $\bar{s}(p)$ was labeled (x, y, z) in $P_{c-\varepsilon}$, by Lemma A.10 $x > z/z > x$ holds and in particular $a > z/a > x$. Thus, the label of any 0-simplex which was labeled a in $P_{c-\varepsilon}$ is increased by 1. In conclusion $\alpha \geq a + 1$ holds.

In all four cases the label a is increased by 1 during the step that considers c . Thus, $\alpha = a + 1$ holds which implies $\alpha = a + 1 = b < b + 1 = \beta \Rightarrow \beta = \alpha + 1$. \square

Theorem A.12. *The constructed function f is an index-ordered dMf that has only critical cells.*

Proof. We only have to show that f is injective. Lemma A.11 then implies that f is weakly increasing and index-ordered. The injectivity implies that all cells are critical.

We prove the injectivity of f inductively. The initial case again follows the same way as the initial step in Lemma A.3. The injectivity on 1-simplices follows by Lemma A.3 and Lemma A.11 and the fact that the map $+2: \mathbb{R} \rightarrow \mathbb{R}$ is a bijection. Thus, we only

have to show injectivity on 0-simplices. For the inductive step, we again distinguish between the four cases corresponding to the chirality of c and its parent node p :

LR/RL By construction, p was considered directly before c . If $\bar{s}(p)$ was labeled (x, y, z) in $P_{c-\varepsilon}$, Lemma A.10 implies that $z = x + 1/x = z + 1$ holds.

Thus, we can divide the 0-simplices of $P_{c-\varepsilon}$ into three disjoint types:

- a) 0-simplices that have labels lower than x/z ,
- b) the 0-simplices $s_l(p)$ and $s_r(p)$ which are labeled x and z , respectively, and
- c) 0-simplices with higher labels than z/x .

The 0-simplices of type a) keep their labels, so f_c is injective on these simplices in P_c . In P_c , the simplex $\bar{s}(p)$ is labeled $(x, y + 2, z + 1)/(x + 1, y + 2, z)$ and $\bar{s}(c)$ is labeled $(z + 1, y + 1, x + 1)/(z + 1, y + 1, x + 1)$. Because $z = x + 1/x = z + 1$ holds, it follows that f_c is injective on the simplices of type b). By construction, the 0-simplices of type c) are increased by 1, so they are assigned values larger than $z + 1/x + 1$. Because $+1: \mathbb{R} \rightarrow \mathbb{R}$ is a bijection, f_c is injective on the 0-simplices of type c) in P_c .

RR/LL We can divide the 0-simplices of $P_{c-\varepsilon}$ in three disjoint types:

- a) 0-simplices that have labels lower than z/x ,
- b) the 0-simplex with the label z/x , and
- c) 0-simplices with higher labels than z/x .

The 0-simplices of type a) keep their labels, so f_c is injective on these simplices in P_c . The 0-simplex that had the label z/x gets the label $z + 1/x + 1$ and the new 0-simplex gets the label z/x . By construction, the 0-simplices of type c) are assigned values larger than $z + 1/x + 1$. Because $+1: \mathbb{R} \rightarrow \mathbb{R}$ is a bijection, f_c is injective on the 0-simplices of type c) in P_c .

In both cases, the images of simplices of different types under f_c are disjoint. In conclusion, f_c is injective on all simplices of P_c , so it follows inductively that f is injective. \square

Remark A.13. In the preceding proof, we used the fact that the maps $+1: \mathbb{R} \rightarrow \mathbb{R}$ and $+2: \mathbb{R} \rightarrow \mathbb{R}$ are order-preserving bijections. We will use this fact later to show that the adjustment preserves the order in which the construction algorithm of the induced merge tree considers the 1-simplices of P .


A.3. The Constructed DMF Represents T .

Theorem A.14. *Let T be a merge tree. The step-by-step construction (Construction 3.14) yields a path P with a dMf $f: P \rightarrow \mathbb{R}$ such that $M(P, f) \cong T$ as merge trees.*

Proof. It was already shown in Theorem A.12 that f is an injective dMf. In particular, all simplices of P are critical with respect to f . Thus, the construction algorithm of the induced merge tree considers all 1-simplices of P .

We prove the theorem inductively: We prove that $M(P_c, f_c) \cong T_c$ holds throughout every step of the construction.

For the initial case, let r be the root of T . Then T_r , (P_r, f_r) and $M(P_r, f_r)$ look like this:




It is clear that $T_r \cong M(P_r, f_r)$ as merge trees.

By Lemma A.3 the order of the algorithm matches the order on nodes of T induced by the labels of their corresponding 1-simplices in P from highest to lowest.

We also recall that the step-by-step algorithm for the construction of the induced merge tree considers the critical 1-simplices of P ordered by label from highest to lowest label. In each step, the algorithm removes the critical 1-simplex from P which is being considered and creates two child nodes in $M(P, f)$ for the two resulting connective components. The child corresponding to the connective component with the lower minimal label gets the same chirality as its parent node.

We recall from Remark A.13 that in every step of the construction of P , after the creation of the new simplices is done, the adjustment of the pre-existing simplices is done in an order-preserving way. Thus, the construction algorithm of the induced merge tree considers the pre-existing 1-simplices of P_c in the same order as they appear in $P_{c-\varepsilon}$. By Lemma A.10, the new 1-simplex has the lowest label among 1-simplices, so it is considered last by the algorithm. For the inductive step, we distinguish between four cases corresponding to the chirality of c and its parent node p . Additionally, let $\bar{s}(p)$ be labeled (x, y, z) .


LR/RL Because there was a change in chirality from **left to right**/**right to left**, both children of p must be leaves in $T_{c-\varepsilon}$:




Due to the inductive hypothesis, $T_{c-\varepsilon} \cong M(P_{c-\varepsilon}, f_{c-\varepsilon})$ holds. As discussed above, the pre-existing 1- and 0-simplices are adjusted so that the construction algorithm of the induced merge tree considers the pre-existing edges of P_c in the same order as they appear in $P_{c-\varepsilon}$. Thus, the structure of $M(P_{c-\varepsilon}, f_{c-\varepsilon})$, except for the subtree of the node induced by $s(p)$, is preserved in $M(P_c, f_c)$.

By Lemmas A.10 and A.11, all labels of $\bar{s}(c)$ are greater than the label x of $s_l(p)/z$ of $s_r(p)$ in (P_c, f_c) . Thus, in $M(P_c, f_c)$ the simplex $s_l(p)/s_r(p)$ induces a leaf which is the left child of the node corresponding to $s(p)$. Furthermore, $\bar{s}(c)$ produces the subtree of the **right/left** child of the node corresponding to $s(p)$,

namely an inner node with two leaves as children. In conclusion $T_c \cong M(P_c, f_c)$ follows.



RR/LL Because there was no change of chirality, p_l/p_r might be an inner node which is the root of a non-trivial subtree in T_c : By Lemma A.10, the label z of $s_r(p)/x$ of $s_l(p)$ is smaller than the label of $s_l(p)/s_r(p)$ in $P_{c-\varepsilon}$. All labels of 0-simplices corresponding to nodes of the subtree with root p_l/p_r in $T_{c-\varepsilon}$ are inductively constructed from z/x . Therefore, by construction, these labels are bigger than z/x . Thus, p_r/p_l is a leaf in $T_{c-\varepsilon}$, whereas p_l/p_r might be the root of a non-trivial subtree:



Due to the inductive hypothesis, $T_{c-\varepsilon} \cong M(P_{c-\varepsilon}, f_{c-\varepsilon})$ holds. As discussed above, the pre-existing 1- and 0-simplices are adjusted so that the construction algorithm of the induced merge tree considers the pre-existing edges of P_c in the same order as they appear in $P_{c-\varepsilon}$. Thus, the structure of $M(P_{c-\varepsilon}, f_{c-\varepsilon})$, except possibly for the subtree of the node induced by $s(p)$, is preserved completely in $M(P_c, f_c)$.

By construction, the label $y+2$ of $s(p)$ is higher than all labels of simplices corresponding to other nodes of the subtree with root p in T_c . Additionally, the label $m+1$ of $s(c)$ is the smallest label among all labels of simplices corresponding to other nodes of the subtree with root p in T_c and the labels of 1-simplices belonging to this subtree form an interval in \mathbb{N} . Furthermore, all 1-simplices in P_c corresponding to nodes of the subtree with root p_l/p_r in T_c belong to a different connected component of $P_{c_{(y+2-\varepsilon)}}$ than $s(c)$. (We recall that $P_{c_{(y+2-\varepsilon)}}$ denotes the subcomplex of P_c consisting of simplices with labels lower than $y+2$.)


Thus, when $s(p)$ is being considered during the construction of the induced merge tree (M_c, f_c) , the connected component with $s(c)$ induces a **right/left**

child node of p , whereas the connected component corresponding to the subtree with root p_l/p_r induces a left/right child node.

Afterwards, the subtree with root induced by $s(p_l)/s(p_r)$ is preserved because the adjustment of the pre-existing simplices in the step that considered c is order preserving.


By construction, all labels of $\bar{s}(c)$ are smaller than the labels of simplices corresponding to the subtree with root p_l/p_r in T_c . Thus, $s(c)$ induces an inner node with two leaves as a right/left child of the node induced by $s(p)$ in $M(P_c, f_c)$.

In conclusion $T_c \cong M(P_c, f_c)$ follows.




It follows inductively that $T \cong M(P, f)$ holds. □

A.4. An Example for the Step-by-Step Construction. We apply the construction to an example that contains most of the interesting situations which may occur. We consider the following merge tree T :




In each step of the construction, we will depict $T \setminus T_c$, $T_{c-\epsilon} \cong M(P_{c-\epsilon}, f_{c-\epsilon})$, $T_c \setminus T_{c-\epsilon} \cong M(P_c, f_c) \setminus M(P_{c-\epsilon}, f_{c-\epsilon})$, $P_{c-\epsilon}$ and $P_c \setminus P_{c-\epsilon}$ using the corresponding colors. The choice of colors for this example does not correspond to the choice of colors for the case distinction above. Initially, the path P_c is empty.


Step 0 (L):




Step 1 (LR):




Step 2 (RL):




Step 3 (LR):




Step 4 (RL):




Step 5 (RR):






Remark A.15. Step 5 was the first step without a change of chirality. This implies that it was the first step where the labels of p_l and p_r differ by more than 1. Furthermore, it was the first step where labels of pre-existing 0-simplices needed to be adjusted. From now on, the adjustment of labels of pre-existing 0-simplices happens in every step.

Step 6 (RL):




Step 7 (LR):




Remark A.16. In step 8, we have an LL step where c is an inner node of the leftmost branch. This is the only kind of step where the labels on all 0-simplices are being adjusted.


Step 8 (LL):




Step 9 (LR):



Step 10 (RR):





This concludes the construction and we have $M(P, f) \cong T$.

REFERENCES

- [Bar19] Yuliy Baryshnikov. Time Series, Persistent Homology and Chirality. 2019. arXiv:1909.09846 [math.PR].
- [CDG⁺21] Justin Curry, Jordan DeSha, Adélie Garin, Kathryn Hess, Lida Kanari, and Brendan Mallery. From trees to barcodes and back again ii: Combinatorial and probabilistic aspects of a topological inverse problem. 07 2021. arXiv:2107.11212v2 [math.AT].
- [CSA03] Hamish Carr, Jack Snoeyink, and Ulrike Axen. Computing contour trees in all dimensions. *Computational Geometry: Theory and Applications*, pages 75–94, 2003.
- [Cur19] Justin Curry. The fiber of the persistence map for functions on the interval. *J Appl. and Comput. Topology*, pages 301–321, 02 2019.
- [For98] Robin Forman. Morse Theory for Cell Complexes. *Advances in Mathematics*, 134:90–145, 03 1998.
- [For01] Robin Forman. A User’s Guide To Discrete Morse Theory. *Sem. Lothar. Combin.*, 48, 12 2001.
- [HLH⁺16] Christian Heine, Heike Leitte, Mario Hlawitschka, Frederico Iuricich, Leila De Florian, Gerik Scheuermann, Hans Hagen, and Christoph Garth. A Survey of Topology-based Methods in Visualization. *Computer Graphics Forum*, 35:643–667, 07 2016.
- [JS20] Benjamin Johnson and Nicholas A. Scoville. Merge trees in discrete Morse theory. 2020. arXiv:2007.10272 [math.AT].
- [KK94] In So Kweon and Takeo Kanade. 1994.
- [LMW⁺16] Shusen Liu, Dan Maljovec, Bei Wang, Peer-Timo Bremer, and Valerio Pascucci. Visualizing High-Dimensional Data: Advances in the Past Decade. *IEEE Transactions on Visualization and Computer Graphics*, pages 1249–1268, 12 2016.
- [NTT18] Vidit Nanda, Dai Tamaki, and Kohei Tanaka. Discrete Morse theory and classifying spaces. *Advances in Mathematics*, 340:723–790, 10 2018.
- [OHW⁺17] Patrick Oesterling, Christian Heine, Gunther H. Weber, Dimitry Morozov, and Gerik Scheuermann. Computing and Visualizing Time-Varying Merge Trees for High-Dimensional Data. *Topological Methods in Data Analysis and Visualization*, pages 87–101, 2017.
- [OHWS13] Patrick Oesterling, Christian Heine, Gunther H. Weber, and Gerik Scheuermann. Visualizing nD Point Clouds as Topological Landscape Profiles to Guide Local Data Analysis. *IEEE Transactions on Visualization and Computer Graphics*, 19:514–526, 03 2013.
- [SKK91] Yoshihisa Shinagawa, Toshiyasu L. Kunii, and Yannick L. Kergosien. Surface Coding Based on Morse Theory. *IEEE Computer Graphics and Applications*, pages 66–78, 1991.
- [TV98] Sergey P. Tarasov and Michael N. Vyalyi. Construction of contour trees in 3D in $O(n \log n)$ steps. *SCG 98: Proceedings of the fourteenth annual symposium on Computational geometry*, pages 68–75, 06 1998.
- [vKvOB⁺97] Marc van Kreveld, René van Oostrum, Chandrajit Bajaj, Valerio Pascucci, and Dan Schikore. 08 1997.
- [WBP07] Gunther H. Weber, Peer-Timo Bremer, and Valerio Pascucci. Topological Landscapes: A Terrain Metaphor for Scientific Data. *IEEE Transactions on Visualization and Computer Graphics*, 13:1416–1423, 09 2007.
- [YWM⁺19] Lin Yan, Yusu Wang, Elizabeth Munch, Ellen Gasparovic, and Bei Wang. A Structural Average of Labeled Merge Trees for Uncertainty Visualization. *IEEE Transactions on Visualization and Computer Graphics*, 26:832–842, 08 2019.

(Julian Brüggemann) MAX PLANCK INSTITUTE FOR MATHEMATICS, BONN, GERMANY
 Email address: brueggemann@mpim-bonn.mpg.de