

Quadratic-in-spin interactions at the fifth post-Newtonian order probe new physics

Jung-Wook Kim,¹ Michèle Levi,^{1,2,*} and Zhewei Yin³

¹Queen Mary University of London, London E1 4NS, United Kingdom

²Niels Bohr Institute, University of Copenhagen, 2100 Copenhagen, Denmark

³Uppsala University, 75108 Uppsala, Sweden

(Dated: April 7, 2022)

We derive the observable binding energies and their relations to the angular momentum for the next-to-next-to-next-to-leading order (N^3LO) of all quadratic-in-spin interactions in post-Newtonian (PN) gravity. Our results are valid for all generic compact binaries, and enter at the fifth PN order for maximally-rotating compact objects, thus pushing the state of the art. This is accomplished through an extension of the effective field theory of spinning gravitating objects. We find a new operator that contributes to the observables, making the associated Wilson coefficient a new probe for candidate theories of gravity and for QCD.

The success of ground-based ground-breaking experiments in measuring gravitational waves (GWs) in recent years, since the first detection from a merger of a binary black hole [1] by the Advanced LIGO [2] and the Advanced VIRGO [3] collaboration has exceeded our expectations. By now there is an operational worldwide network of GW detectors of second generation technology, as the twin Advanced LIGO detectors in the US has been joined by the Advanced Virgo detector in Europe [3], and more recently also by the KAGRA detector in Japan [4]). These experiments have been reaching continually higher sensitivities, which yield more frequent GW detections, and the influx of GW data has been steeply growing [5–7].

Since 2017 these GW detections also include observations of neutron stars as individual components of the binary, either within binaries of neutron stars [8], or in mixed neutron star-black hole binaries [9]. The study of these various GW sources in their inspiral phase, that is when the compact objects in the binary are still orbiting in non-relativistic velocities, can be analytically carried out via the post-Newtonian (PN) approximation of General Relativity [10]. This study enables to make theoretical predictions for the gravitational waveforms, which in turn inform us not only on a wealth of astrophysical and cosmological scenarios that were previously unthought-of, but also serve as a unique means to test candidate theories of gravity and to learn on QCD in extreme conditions [11, 12]. For the latter, it is crucial to study spin and finite-size effects [13].

In recent years impressive progress has been made in pushing the state of the art in PN gravity. In particular, the point-mass interaction at the fifth PN (5PN) order within the conservative dynamics of the binary has been recently accomplished via a new exploitation of a combination of traditional GR methodologies [14–16], with crucial ingredients imported from gravitational self-force theory and the effective-one-body approach [17]. It was also completed via the effective field theory (EFT) approach [18, 19] in [20]. However, in order to attain the improved accuracy at the 5PN order, the spinning sec-

tors must also be completed. Accordingly, the work in [21, 22] for example, followed the footsteps of [14, 15], and was able to implement a similar approach with a hybrid of traditional GR methods to derive new results that are linear in the spin at the 4.5PN order, and for the restricted binary configuration of aligned spins, also for the bi-linear interaction between the two spins of the binary at the 5PN order.

It is critical to note however that while it is important and illustrative to target specific new PN results ad-hoc, via the capitalization of existing derivations from elementary methodologies, such as gravitational self-force theory in the aforementioned works, such an approach is essentially limited. It does not provide a conceptual framework to derive various results at other required sectors, nor an independent method to study self-force or PN theory, and we believe it is more prone to the propagation of intractable errors, due to its reliance on the combination of inputs dependent on the various ingredient methods.

In this work we complement the 5PN order accuracy by deriving all the interactions that are quadratic in the spins of the compact objects in the most generic settings. The derivation builds on the formulation of the EFT of spinning gravitating objects presented in [23], see also [24, 25], and its extensions to higher loop and spin orders in [26–35]. The formulation in [23], which provided the leading couplings to curvature to all orders in spin, also exclusively enabled the completion of the current state of the art at the 4PN order, as the next-to-next-to-leading order (NNLO) quadratic-in-spin sector [30, 31] is the only sector at this accuracy that was not derived as yet in another independent work. The evaluation of the sectors at the next-to-next-to-next-to-leading order (N^3LO) carried out in the present work is by far the most intricate and demanding undertaking in the spinning sectors to date. This would be intractable if not also for building on the infrastructure of `EFTofPNG` – a unique code for high-precision Feynman computation introduced in [25, 36].

The EFT of spinning gravitating objects. We build on the EFT of spinning gravitating objects introduced in

[30]. In order to obtain all the quadratic-in-spin interactions at this order we need to evaluate the two-particle effective action for the compact binary [18, 24]:

$$S_{\text{eff}} = S_g[g_{\mu\nu}] + \sum_{a=1}^2 S_{\text{pp}}(\lambda_a), \quad (1)$$

and carefully consider the effective action of the spinning particle, S_{pp} , localized on the worldline parametrized by λ_a for each of the two components of the binary.

Since we are concerned here with conservative dynamics, the propagators are taken as instantaneous, and it is useful to employ a Kaluza-Klein time-space decomposition of the field [37, 38], that was first studied and tested in the spin sectors in [39, 40]. The spatial dimension, d , must be kept generic throughout, as dimensional regularization is used to evaluate the effective action, with the modified minimal subtraction ($\overline{\text{MS}}$) prescription applied with the d -dimensional gravitational constant [33]

$$G_d \equiv G_N \left(\sqrt{4\pi e^{\gamma_E}} R_0 \right)^{d-3}, \quad (2)$$

in which $G_N \equiv G$ is Newton's gravitational constant in three-dimensional space, γ_E is Euler's constant, and R_0 is a fixed renormalization scale.

Since the quadratic-in-spin sector includes finite-size effects in addition to the minimal coupling of rotating objects to gravity, we need to take into account the following extended effective action for each of the spinning particles [23, 24]:

$$S_{\text{pp}}(\lambda) = \int d\lambda \left[-m\sqrt{u^2} - \frac{1}{2} \hat{S}_{\mu\nu} \hat{\Omega}^{\mu\nu} - \frac{\hat{S}^{\mu\nu} p_\nu}{p^2} \frac{Dp_\mu}{D\lambda} \right] + L_{\text{NMC}}[g_{\mu\nu}, u^\mu, S^\mu], \quad (3)$$

in which the non-minimal coupling due to the spin, L_{NMC} , is initially formulated in terms of the definite-parity classical analogue of the Pauli-Lubanski pseudovector, S_μ , as defined in [23, 28, 32]. We note that another formulation utilizing EFT techniques was approached in [41], where it was applied to lower orders.

The non-minimal coupling to all orders in the spin and linear in the curvature is given in the following compact form [23]:

$$L_{\text{NMC(R)}} = \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} \frac{C_{ES^{2n}}}{m^{2n-1}} D_{\mu_{2n}} \cdots D_{\mu_3} \frac{E_{\mu_1\mu_2}}{\sqrt{u^2}} S^{\mu_1} S^{\mu_2} \cdots S^{\mu_{2n-1}} S^{\mu_{2n}} + \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n+1)!} \frac{C_{BS^{2n+1}}}{m^{2n}} D_{\mu_{2n+1}} \cdots D_{\mu_3} \frac{B_{\mu_1\mu_2}}{\sqrt{u^2}} S^{\mu_1} S^{\mu_2} \cdots S^{\mu_{2n}} S^{\mu_{2n+1}}, \quad (4)$$

with the definite-parity electric and magnetic curvature components,

$$E_{\mu\nu} \equiv R_{\mu\alpha\nu\beta} u^\alpha u^\beta, \quad (5)$$

$$B_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\alpha\beta\gamma\mu} R^{\alpha\beta}{}_{\delta\nu} u^\gamma u^\delta, \quad (6)$$

and their covariant derivatives, D_μ . The only operator from this infinite series that contributes to our present sectors is the first electric term, which is preceded by a Wilson coefficient that corresponds to the quadrupolar deformation constant [42].

However at this high order the effective action of a spinning particle needs to be extended beyond linear order in the curvature, namely beyond (4). This extension was introduced in [34], following the symmetry considerations and the logic presented in [23], and from basic dimensional analysis and power-counting considerations [23, 24] only the following terms may enter at the 5PN

order to quadratic order in the spins:

$$L_{\text{NMC(R}^2)} = C_{E^2} \frac{E_{\alpha\beta} E^{\alpha\beta}}{\sqrt{u^2}^3} + C_{B^2} \frac{B_{\alpha\beta} B^{\alpha\beta}}{\sqrt{u^2}^3} + C_{E^2 S^2} S^\mu S^\nu \frac{E_{\mu\alpha} E_\nu^\alpha}{\sqrt{u^2}^3} + C_{B^2 S^2} S^\mu S^\nu \frac{B_{\mu\alpha} B_\nu^\alpha}{\sqrt{u^2}^3}. \quad (7)$$

These terms involve Wilson coefficients, that at this point absorb all numerical and mass factors, in contrast with those in (4). The terms on the second line are a new type of operators that would be relevant only for spinning objects, as opposed to those in the first line. Further scrutiny of the operators in question reveals that their dimensionless Wilson coefficients should be defined as

$$C_{[E/B]^2} \rightarrow \frac{1}{2} G^4 m^5 C_{[E/B]^2}, \quad (8)$$

$$C_{[E/B]^2 S^2} \rightarrow \frac{1}{2} G^2 m C_{[E/B]^2 S^2}. \quad (9)$$

Notice that these are the leading operators that exhibit such additional scaling in G in their coefficients, such that




FIG. 1. The unique Feynman graph at the NNNLO quadratic-in-spin sector, which originates from the new quadratic-curvature operator that is also quadratic in the spin, and is preceded by an unknown Wilson coefficient, see (7), (8). The new operator is represented here by the black square that is denoted by E^2 .

although they would start contributing through topologies of order G^2 , their actual contributions would be at order G^4 , and G^6 , respectively, unlike any PN contributions up to this order, including those in the quartic-in-spin sector at 5PN order [35], in which the order in G of the topologies has always been identical to the order of their actual contributions. One may suspect that these operators are redundant, and therefore may be trivially removed from the effective action. Yet, such redundancy is not identified at the level of the action, and in any case the physical observables will tell us if the operators are redundant. Notice also that interestingly the leading non-minimal couplings that are only linear in the spin will enter at higher orders.

From EFT formulation to observables. First we need to extend the relevant module of `EFTofPNG` in order to generate the Feynman rules for a generic number of spatial dimensions d at this order. This also requires a recursive implementation of the canonical gauge to fix the rotational variables of the spinning object that is defined in [23]. Notably also, the worldline operators include a coupling to the curvature up to quadratic order as noted, in addition to a proliferation of time derivatives on the worldline spin couplings, and from the bulk gravitational action. The graphs/contractions that contribute to these sectors are then generated through an extension of the relevant module of the code. All in all, we find that there are 1122 graphs that make up the N³LO quadratic-in-spin sectors, including the highest-loop graphs studied in the spinning sectors.

This large volume of complex graphs also carry a higher tensorial load due to the coupling to spins, and thus requires to further improve the projection of the integrals due to the high rank of their numerators [43], and to streamline new code to carry out algorithmic integration by parts (IBP) to reduce the integrals that show up to master integrals [44, 45] beyond what was needed for [33, 34]. In general, to verify the reliability of our new code extensions, we carried out independent development in parallel, and new results were compared only at crucial points along the elaborate derivation.

As to the operators that are quadratic in the curvature

in (7), an expansion of the operators reveals that only the electric-component operator that is quadratic in the spins enters at this order in our sectors. This operator gives rise to a unique Feynman graph of two-graviton exchange depicted in Fig. 1, whose value is given by

$$\text{Fig. 1} = -\frac{1}{2}C_{1(E^2S^2)} \frac{G^4 m_1 m_2^2}{r^6} \left[S_1^2 + 3(\vec{S}_1 \cdot \vec{n})^2 \right]. \quad (10)$$

Notice that this introduces a new Wilson coefficient into the effective action.

As noted the graphs are evaluated through dimensional regularization with the $\overline{\text{MS}}$ scheme, and similar to [33, 34] the values of the graphs contain poles in $\epsilon \equiv d - 3$ in conjunction with logarithms in r/R_0 . However, whereas in the piece with topologies of order G^4 only, which was evaluated in [34], all the poles and logarithms conspire to cancel out from the total sum, the total sum of the graphs over all the sector, does contain such divergent and logarithmic terms. These originate from one- and two-loop topologies at order G^2 and G^3 , respectively, at this PN order in our sectors, and thus they enter at these orders to the total effective action.

The growing load of higher-order time derivatives is also challenging, in particular in the spinning sectors, where they appear already at the leading spin effect. Summing up all the graphs we obtain an effective action that contains large pieces with higher-order time derivatives – up to the sixth order. We apply a rigorous procedure to remove these terms, using redefinitions of the position and spin variables, in analogy to field redefinitions that would be made to remove redundant operators, as noted in [27].

We extend the formulation of this procedure that was uniquely introduced for both the position and the rotational variables in [27]. Notice that this procedure is essential since such redefinitions need to be implemented beyond the linear level at high orders, in particular in those sectors that are beyond linear in the spins. The redefinitions of the rotational variables, the Lorentz matrices $\Delta\Lambda^{ij}$, and the spin ΔS^{ij} , that at this point are defined as Euclidean in the locally-flat frames, are parametrized by ω^{ij} , the anti-symmetric generator of rotations on the Lorentz matrices, that is

$$\Lambda^{ij} \equiv \Lambda^{ik} (e^\omega)^{kj}. \quad (11)$$

We then have here redefinitions of position that scale as $\Delta x \sim \epsilon^{-1}$, some of which are dependent in the spins, in addition to redefinitions of the rotational variables that also scale as $|\omega_{ij}| \sim \epsilon^{-1}$.

Now, in order to reduce the action all of the sectors at lower orders have to be treated consistently, i.e. from LO to N³LO of point-mass (S^0) through quadratic-in-spin (S^2) interactions. We get contributions to our new sectors from the full application of redefinitions in lower-order sectors. In particular, we recall that logarithmic terms, preceded by the dimensional parameter, ϵ ,

arise from the basic d -dimensional Fourier integral, see e.g. Eq. (4.35) in [24]. Therefore the LO point-mass, i.e. the Newtonian, and LO quadratic-in-spin sectors, should be considered to order ϵ^1 , as these will yield contributions to our sectors.

Altogether, after the full reduction of the action as described here the divergent and logarithmic terms vanish when going to observables, or alternatively we verified that they can also be removed at the level of the ac-

tion through the addition of a total time derivative. The transition to a Hamiltonian, and to various observables in simplified binary configurations, of e.g. circular orbits and aligned spins, is then straightforward. We created new modules that efficiently go from the unreduced effective actions to the gauge-invariant binding energies.

Let us present then the total binding energy due to the NNNLO quadratic-in-spin interactions, as the sum of the following independent pieces:

$$(e)_{S_2}^{\text{N}^3\text{LO}} = (e)_{S_1 S_2}^{\text{N}^3\text{LO}} + \left[(e)_{S_1^2}^{\text{N}^3\text{LO}} + (e)_{C_{1(\text{E}S^2)} S_1^2}^{\text{N}^3\text{LO}} + (e)_{C_{1(\text{E}^2 S^2)} S_1^2}^{\text{N}^3\text{LO}} + (1 \leftrightarrow 2) \right]. \quad (12)$$

As a function of the orbital frequency these various pieces are given by

$$(e)_{S_1 S_2}^{\text{N}^3\text{LO}}(x) = \tilde{S}_1 \tilde{S}_2 x^6 \nu \left[\frac{243}{16} - \left(\frac{2107}{16} - \frac{123}{32} \pi^2 \right) \nu + \frac{147}{8} \nu^2 + \frac{13}{16} \nu^3 \right], \quad (13)$$

$$(e)_{S_1^2}^{\text{N}^3\text{LO}}(x) = \tilde{S}_1^2 x^6 \nu \left[\left(\frac{1923}{112} \nu - \frac{9633}{112} \nu^2 + \frac{159}{16} \nu^3 \right) - q^{-1} \left(\frac{243}{16} - \left(\frac{795}{16} - \frac{189}{2048} \pi^2 \right) \nu + \frac{2727}{56} \nu^2 - \frac{153}{16} \nu^3 \right) \right], \quad (14)$$

$$(e)_{C_{1(\text{E}S^2)} S_1^2}^{\text{N}^3\text{LO}}(x) = C_{1(\text{E}S^2)} \tilde{S}_1^2 x^6 \nu \left[\left(\frac{789}{28} \nu - \frac{156}{7} \nu^2 + \frac{5}{8} \nu^3 \right) + q^{-1} \left(\frac{405}{32} - \left(\frac{2389}{32} - \frac{3747}{2048} \pi^2 \right) \nu - \frac{555}{56} \nu^2 + \frac{21}{32} \nu^3 \right) \right], \quad (15)$$

and

$$(e)_{C_{1(\text{E}^2 S^2)} S_1^2}^{\text{N}^3\text{LO}}(x) = C_{1(\text{E}^2 S^2)} \tilde{S}_1^2 x^6 \nu \left[-\frac{3}{2} \nu^2 (1 + q^{-1}) \right], \quad (16)$$

where all notations and definitions are identical to [27].

We note that (13) is in complete agreement with [22]. Importantly, we note that there is a new type of contribution in (16) due to the new operator that is quadratic in the curvature.

For the gauge-invariant relation of the binding energy to the angular momentum we find the following pieces:

$$(e)_{S_1 S_2}^{\text{N}^3\text{LO}}(\tilde{L}) = -\tilde{S}_1 \tilde{S}_2 \frac{\nu}{\tilde{L}^{12}} \left[\frac{102897}{16} - \left(\frac{31653}{32} - \frac{369}{32} \pi^2 \right) \nu - \frac{579}{16} \nu^2 + \frac{209}{64} \nu^3 \right], \quad (17)$$

$$(e)_{S_1^2}^{\text{N}^3\text{LO}}(\tilde{L}) = -\tilde{S}_1^2 \frac{\nu}{\tilde{L}^{12}} \left[\left(\frac{2117293}{896} \nu - \frac{142927}{448} \nu^2 + \frac{201}{128} \nu^3 \right) + q^{-1} \left(\frac{211653}{128} + \left(\frac{21211}{16} - \frac{63}{2048} \pi^2 \right) \nu - \frac{33535}{112} \nu^2 + \frac{3}{32} \nu^3 \right) \right], \quad (18)$$

$$(e)_{C_{1(\text{E}S^2)} S_1^2}^{\text{N}^3\text{LO}}(\tilde{L}) = -C_{1(\text{E}S^2)} \tilde{S}_1^2 \frac{\nu}{\tilde{L}^{12}} \left[\left(\frac{2593}{14} \nu - \frac{319}{28} \nu^2 + \frac{3}{8} \nu^3 \right) + q^{-1} \left(\frac{16065}{32} - \left(\frac{3061}{32} - \frac{11745}{2048} \pi^2 \right) \nu - \frac{313}{28} \nu^2 + \frac{17}{32} \nu^3 \right) \right], \quad (19)$$

and

$$(e)_{C_{1(\text{E}^2 S^2)} S_1^2}^{\text{N}^3\text{LO}}(\tilde{L}) = C_{1(\text{E}^2 S^2)} \tilde{S}_1^2 \frac{\nu}{\tilde{L}^{12}} \left[\frac{1}{2} \nu^2 (1 + q^{-1}) \right]. \quad (20)$$

State of the art and new physics. The EFT of spinning gravitating objects constitutes a self-contained

framework that allows for both formal and technical extension, which in turn enables to push the frontier of precision gravity. It addresses generic compact binaries, rather than only black holes, and provides a host of useful mathematical and observable quantities, which are not restricted to specific or simplified binary configurations, such as the center-of-mass frame or aligned-spins, and includes stringent self-consistency checks.

In this work the EFT of spinning gravitating objects also allows to uncover a new type of contribution to physical observables that shows up for rotating compact objects. Similarly to our new contribution, the spinless operators in the first line of (7), which are preceded by the Wilson coefficients C_{E^2} , C_{B^2} , would also naively contribute to the 5PN non-spinning interaction. It is exactly for this reason that it is crucial to undertake the matching of these Wilson coefficients, as they are bound to uncover further new physics. Indeed, recently there has been a surge of independent works that have been studying the rotational Love numbers of black holes and neutron stars [46–51], associated with C_{E^2} , C_{B^2} , with various new findings and insights. It remains for future work to uncover the new physics of gravity and QCD that is encapsulated in these Wilson coefficients.

Acknowledgments. We thank Johannes Blümlein and Alex Edison for pleasant discussions. J-WK was supported by the Science and Technology Facilities Council (STFC) consolidated grant ST/T000686/1 “*Amplitudes, Strings and Duality*”. ML received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant 847523, and is supported by the STFC Rutherford grant ST/V003895/1 “*Harnessing QFT for Gravity*”. ZY is supported by the Knut and Alice Wallenberg Foundation under grants KAW 2018.0116 and KAW 2018.0162.

* michele.levi@qmul.ac.uk

- [1] B. P. Abbott *et al.* (LIGO, VIRGO), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [2] J. Aasi *et al.* (LIGO Scientific), Advanced LIGO, *Class. Quant. Grav.* **32**, 074001 (2015), arXiv:1411.4547 [gr-qc].
- [3] F. Acernese *et al.* (VIRGO), Advanced Virgo: a second-generation interferometric gravitational wave detector, *Class. Quant. Grav.* **32**, 024001 (2015), arXiv:1408.3978 [gr-qc].
- [4] T. Akutsu *et al.* (KAGRA), Overview of KAGRA: Detector design and construction history, (2020), arXiv:2005.05574 [physics.ins-det].
- [5] B. P. Abbott *et al.* (LIGO Scientific, Virgo), GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs, *Phys. Rev.* **X9**, 031040 (2019), arXiv:1811.12907 [astro-ph.HE].
- [6] R. Abbott *et al.* (LIGO Scientific, Virgo), GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run, *Phys. Rev. X* **11**, 021053 (2021), arXiv:2010.14527 [gr-qc].
- [7] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run, (2021), arXiv:2111.03606 [gr-qc].
- [8] B. P. Abbott *et al.* (LIGO Scientific, Virgo), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.* **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [9] R. Abbott *et al.* (LIGO Scientific, KAGRA, VIRGO), Observation of Gravitational Waves from Two Neutron Star–Black Hole Coalescences, *Astrophys. J. Lett.* **915**, L5 (2021), arXiv:2106.15163 [astro-ph.HE].
- [10] L. Blanchet, Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries, *Living Rev. Rel.* **17**, 2 (2014), arXiv:1310.1528 [gr-qc].
- [11] B. P. a. o. Abbott (Virgo, LIGO Scientific), Tests of general relativity with GW150914, *Phys. Rev. Lett.* **116**, 221101 (2016), arXiv:1602.03841 [gr-qc].
- [12] B. P. a. o. Abbott (Virgo, LIGO Scientific), Properties of the Binary Black Hole Merger GW150914, *Phys. Rev. Lett.* **116**, 241102 (2016), arXiv:1602.03840 [gr-qc].
- [13] B. P. a. o. Abbott (Virgo, LIGO Scientific), Improved analysis of GW150914 using a fully spin-precessing waveform Model, *Phys. Rev.* **X6**, 041014 (2016), arXiv:1606.01210 [gr-qc].
- [14] D. Bini, T. Damour, and A. Gericco, Novel approach to binary dynamics: application to the fifth post-Newtonian level, *Phys. Rev. Lett.* **123**, 231104 (2019), arXiv:1909.02375 [gr-qc].
- [15] D. Bini, T. Damour, and A. Gericco, Binary dynamics at the fifth and fifth-and-a-half post-Newtonian orders, *Phys. Rev. D* **102**, 024062 (2020), arXiv:2003.11891 [gr-qc].
- [16] D. Bini, T. Damour, A. Gericco, S. Laporta, and P. Mastrolia, Gravitational dynamics at $O(G^6)$: perturbative gravitational scattering meets experimental mathematics, (2020), arXiv:2008.09389 [gr-qc].
- [17] A. Buonanno and T. Damour, Effective one-body approach to general relativistic two-body dynamics, *Phys. Rev.* **D59**, 084006 (1999), arXiv:gr-qc/9811091 [gr-qc].
- [18] W. D. Goldberger and I. Z. Rothstein, An Effective field theory of gravity for extended objects, *Phys. Rev.* **D73**, 104029 (2006), arXiv:hep-th/0409156 [hep-th].
- [19] W. D. Goldberger, Les Houches lectures on effective field theories and gravitational radiation, in *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime Les Houches, France, July 31-August 25, 2006* (2007) arXiv:hep-ph/0701129 [hep-ph].
- [20] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an

- effective field theory approach: potential contributions, Nucl. Phys. B **965**, 115352 (2021), arXiv:2010.13672 [gr-qc].
- [21] A. Antonelli, C. Kavanagh, M. Khalil, J. Steinhoff, and J. Vines, Gravitational spin-orbit coupling through third-subleading post-Newtonian order: from first-order self-force to arbitrary mass ratios, Phys. Rev. Lett. **125**, 011103 (2020), arXiv:2003.11391 [gr-qc].
- [22] A. Antonelli, C. Kavanagh, M. Khalil, J. Steinhoff, and J. Vines, Gravitational spin-orbit and aligned spin₁-spin₂ couplings through third-subleading post-Newtonian orders, Phys. Rev. D **102**, 124024 (2020), arXiv:2010.02018 [gr-qc].
- [23] M. Levi and J. Steinhoff, Spinning gravitating objects in the effective field theory in the post-Newtonian scheme, JHEP **09**, 219 (2015), arXiv:1501.04956 [gr-qc].
- [24] M. Levi, Effective Field Theories of Post-Newtonian Gravity: A comprehensive review, Rept. Prog. Phys. **83**, 075901 (2020), arXiv:1807.01699 [hep-th].
- [25] M. Levi and J. Steinhoff, EFTofPNG: A package for high precision computation with the Effective Field Theory of Post-Newtonian Gravity, Class. Quant. Grav. **34**, 244001 (2017), arXiv:1705.06309 [gr-qc].
- [26] M. Levi, Binary dynamics from spin1-spin2 coupling at fourth post-Newtonian order, Phys.Rev. **D85**, 064043 (2012), arXiv:1107.4322 [gr-qc].
- [27] M. Levi and J. Steinhoff, Equivalence of ADM Hamiltonian and Effective Field Theory approaches at next-to-next-to-leading order spin1-spin2 coupling of binary inspirals, JCAP **1412**, 003 (2014), arXiv:1408.5762 [gr-qc].
- [28] M. Levi and J. Steinhoff, Leading order finite size effects with spins for inspiralling compact binaries, JHEP **06**, 059 (2015), arXiv:1410.2601 [gr-qc].
- [29] M. Levi and J. Steinhoff, Next-to-next-to-leading order gravitational spin-orbit coupling via the effective field theory for spinning objects in the post-Newtonian scheme, JCAP **1601**, 011 (2016), arXiv:1506.05056 [gr-qc].
- [30] M. Levi and J. Steinhoff, Next-to-next-to-leading order gravitational spin-squared potential via the effective field theory for spinning objects in the post-Newtonian scheme, JCAP **1601**, 008 (2016), arXiv:1506.05794 [gr-qc].
- [31] M. Levi and J. Steinhoff, Complete conservative dynamics for inspiralling compact binaries with spins at the fourth post-Newtonian order, JCAP **09**, 029 (2021), arXiv:1607.04252 [gr-qc].
- [32] M. Levi, S. Mougiakakos, and M. Vieira, Gravitational cubic-in-spin interaction at the next-to-leading post-Newtonian order, JHEP **01**, 036 (2021), arXiv:1912.06276 [hep-th].
- [33] M. Levi, A. J. Mcleod, and M. Von Hippel, N³LO gravitational spin-orbit coupling at order G^4 , JHEP **07**, 115 (2021), arXiv:2003.02827 [hep-th].
- [34] M. Levi, A. J. Mcleod, and M. Von Hippel, N³LO gravitational quadratic-in-spin interactions at G^4 , JHEP **07**, 116 (2021), arXiv:2003.07890 [hep-th].
- [35] M. Levi and F. Teng, NLO gravitational quartic-in-spin interaction, JHEP **01**, 066 (2021), arXiv:2008.12280 [hep-th].
- [36] M. Levi, A public framework for Feynman calculations and post-Newtonian gravity, (2018), arXiv:1811.12401 [gr-qc].
- [37] B. Kol and M. Smolkin, Non-Relativistic Gravitation: From Newton to Einstein and Back, Class.Quant.Grav. **25**, 145011 (2008), arXiv:0712.4116 [hep-th].
- [38] B. Kol, M. Levi, and M. Smolkin, Comparing space+time decompositions in the post-Newtonian limit, Class.Quant.Grav. **28**, 145021 (2011), arXiv:1011.6024 [gr-qc].
- [39] M. Levi, Next to Leading Order gravitational Spin1-Spin2 coupling with Kaluza-Klein reduction, Phys.Rev. **D82**, 064029 (2010), arXiv:0802.1508 [gr-qc].
- [40] M. Levi, Next to Leading Order gravitational Spin-Orbit coupling in an Effective Field Theory approach, Phys.Rev. **D82**, 104004 (2010), arXiv:1006.4139 [gr-qc].
- [41] R. A. Porto, Post-Newtonian corrections to the motion of spinning bodies in NRGR, Phys.Rev. **D73**, 104031 (2006), arXiv:gr-qc/0511061 [gr-qc].
- [42] B. Barker and R. O'Connell, Gravitational Two-Body Problem with Arbitrary Masses, Spins, and Quadrupole Moments, Phys.Rev. **D12**, 329 (1975).
- [43] G. Passarino and M. J. G. Veltman, One Loop Corrections for $e^+ e^-$ Annihilation Into $\mu^+ \mu^-$ in the Weinberg Model, Nucl. Phys. **B160**, 151 (1979).
- [44] S. Laporta, High precision calculation of multiloop Feynman integrals by difference equations, Int. J. Mod. Phys. **A15**, 5087 (2000), arXiv:hep-ph/0102033 [hep-ph].
- [45] V. A. Smirnov, *Analytic tools for Feynman integrals*, Vol. 250 (2012).
- [46] A. Le Tiec and M. Casals, Spinning Black Holes Fall in Love, Phys. Rev. Lett. **126**, 131102 (2021), arXiv:2007.00214 [gr-qc].
- [47] A. Le Tiec, M. Casals, and E. Franzin, Tidal Love Numbers of Kerr Black Holes, Phys. Rev. D **103**, 084021 (2021), arXiv:2010.15795 [gr-qc].
- [48] H. S. Chia, Tidal deformation and dissipation of rotating black holes, Phys. Rev. D **104**, 024013 (2021), arXiv:2010.07300 [gr-qc].
- [49] P. Charalambous, S. Dubovsky, and M. M. Ivanov, On the Vanishing of Love Numbers for Kerr Black Holes, JHEP **05**, 038 (2021), arXiv:2102.08917 [hep-th].
- [50] P. Charalambous, S. Dubovsky, and M. M. Ivanov, Hidden Symmetry of Vanishing Love Numbers, Phys. Rev. Lett. **127**, 101101 (2021), arXiv:2103.01234 [hep-th].
- [51] G. Castro, L. Gualtieri, and P. Pani, Hidden symmetry between rotational tidal Love numbers of spinning neutron stars, Phys. Rev. D **104**, 044052 (2021), arXiv:2103.16595 [gr-qc].