

# Bootstrap percolation, probabilistic cellular automata and sharpness

Ivailo Hartarsky\*

CEREMADE, CNRS, Université Paris-Dauphine, PSL University  
75016 Paris, France

December 6, 2021

## Abstract

We establish new connections between percolation, bootstrap percolation, probabilistic cellular automata and deterministic ones. Surprisingly, by juggling with these in various directions, we effortlessly obtain a number of new results in these fields. In particular, we prove the sharpness of the phase transition of attractive absorbing probabilistic cellular automata, a class of bootstrap percolation models and kinetically constrained models. We further show how to recover a classical result of Toom on the stability of cellular automata w.r.t. noise and, inversely, how to deduce new results in bootstrap percolation universality from his work.

**MSC2020:** Primary 60K35; Secondary 37B15, 60C05, 82B43, 82C20

**Keywords:** probabilistic cellular automata, bootstrap percolation, kinetically constrained models, sharp phase transition, stability

## 1 Introduction

There are numerous links between probabilistic cellular automata (PCA) and percolation (see Section 1.1 for the definitions of our models of interest). In the case of additive PCA this link is very apparent, since they may be viewed as oriented percolation models (see e.g. [17]). Moreover, percolation is often used as a reference model for comparison in more complex cases (see e.g. [21]).

---

\*hartarsky@ceremade.dauphine.fr

Our first goal will be to import a recent technique [11] for proving the sharpness of phase transitions from percolation to the setting of attractive PCA. This allows us to establish that they all behave in the expected way throughout their ‘subcritical’ phase. This comes to complement a classical result of Bezuidenhout and Gray [6] showing that a certain ‘supercritical’ phase is also well-behaved.

Besides PCA, our other main motivation for pursuing this result comes from bootstrap percolation (BP). We establish a correspondence between the two, so as to deduce exponential decay above criticality previously conjectured for a class of BP models, as well as related kinetically constrained models (KCM), based on previous work of the author [16].

Finally, we show other uses of the correspondence between PCA and BP. Namely, it provides an equivalence between the non-triviality of the phase transition of certain BP models and the stability w.r.t. noise of certain deterministic cellular automata (CA). The former was studied recently in the framework of BP universality by Balister, Bollobás, Przykucki and Smith [3] and by Bollobás, Smith and Uzzell [8], while the latter was investigated over four decades ago by Toom [28] and subsequently by a number of authors [5, 9, 13, 14, 18]. Bridging their viewpoints yields results in both directions.

## 1.1 Models

### 1.1.1 General setting

**Convention** As it is common in set systems, we will denote points with lower case letters, sets of points with upper case ones (or with Greek lower case letters), families of such sets by capital calligraphic letters, systems of such families with capital script letters and in the rare case of classes of such systems, we will use capital fraktur letters. Here and below we use the words ‘set’, ‘family’, ‘system’ and ‘class’ as synonyms, but we will reserve their usage to the corresponding levels as much as possible.

Throughout, unless otherwise stated, we fix an integer dimension  $d \geq 1$  and range  $r \in [1, \infty)$ . We set  $R = ([-r, r]^d \times [-r, 0)) \cap \mathbb{Z}^{d+1}$ , so as to allow non-zero memory. Models *without memory* will be defined identically, taking  $R = ([-r, r]^d \times \{-1\}) \cap \mathbb{Z}^{d+1}$ . A *configuration* is any element of  $\Omega = \{0, 1\}^{\mathbb{Z}^d \times ([-r, 0) \cap \mathbb{Z})}$ . For any set  $X$ , we identify any  $\eta \in \{0, 1\}^X$  with a subset of  $X$  in the natural way. An *up-set* of a partially ordered set  $(P, \geq)$  is a subset  $U \subset P$  such that for any  $(p, u) \in P \times U$  such that  $p \geq u$  we have  $p \in U$ . We similarly define *down-sets* (which are the complements of up-sets). Let  $\mathcal{U}$  denote the system of all up-families of  $\Omega_R := \{0, 1\}^R$  equipped with the partial order  $\eta \geq \omega$  if  $\eta_x \geq \omega_x$  for all  $x \in R$  (that is, if  $\eta \supset \omega$ ). Note

that  $\emptyset \in \mathcal{U}$  and  $\Omega_R \in \mathcal{U}$ . We will further need to consider the class  $\mathfrak{D}$  of down-sets of the partially ordered system  $(\mathcal{U}, \supset)$ .

An *attractive PCA* will be defined by the rates  $v(\{\mathcal{U}\}) \in [0, 1]$  for  $\mathcal{U} \in \mathcal{U}$ . We require  $\sum_{\mathcal{U} \in \mathcal{U}} v(\{\mathcal{U}\}) = 1$  and view  $v$  as a probability measure on  $\mathcal{U}$ . We say simply *attractive CA*, if  $v$  is a Dirac measure. An attractive PCA is said to be *additive* if  $v(\{\mathcal{U}\}) = 0$  unless  $\mathcal{U}$  is generated by singletons, that is, there exists  $X \subset R$  (possibly empty) such that  $\mathcal{U} = \{Y \subset R : X \cap Y \neq \emptyset\}$ . We further say that it is *absorbing* if  $v(\{\Omega_R\}) = 0$ , which will be equivalent to saying that the  $\mathbf{0}$  configuration (which we identified with the set  $\emptyset$ ) is an absorbing state. Hence, attractive PCA identify with a finite dimensional simplex equipped with the standard topology (that is, the topology of weak convergence of the corresponding  $v$  measures) and similarly for additive, absorbing attractive and absorbing additive ones.

Given the rates  $v$ , we can construct the associated finite memory Markov chain on  $\{0, 1\}^{\mathbb{Z}^d}$  graphically as follows. In words, at each time step  $t \in \mathbb{Z}$ , the state of each site  $x \in \mathbb{Z}^d$  becomes 1 if and only if the restriction of the current configuration to  $R + (x, t)$  belongs to a randomly chosen up-family with law  $v$ . More formally, let  $(\mathcal{U}_{x,t})_{(x,t) \in \mathbb{Z}^{d+1}}$  be an i.i.d. random field of up-families with law  $v$ . Given the state of the PCA  $\eta$  at times  $t - r, \dots, t - 1$  (which form a configuration) and  $x \in \mathbb{Z}^d$ , we define

$$\eta_x(t) = \begin{cases} 1 & \text{if } \{(y, s) \in R : \eta_{x+y}(t+s) = 1\} \in \mathcal{U}_{x,t} \\ 0 & \text{otherwise.} \end{cases} \quad (1)$$

This defines the trajectory of the PCA, given the initial state and the up-family field. When we want to specify that the initial state is  $\omega \in \Omega$ , we write  $\eta^\omega$  for the corresponding process. When the PCA has no memory (i.e. for all  $\mathcal{U} \in \text{supp } v$  and  $U \in \mathcal{U}$  we have  $(U \cap [-r, r]^d \times \{-1\}) \in \mathcal{U}$ ), we may abusively write  $\eta^\omega$  with  $\omega \in \{0, 1\}^{\mathbb{Z}^d}$  and it is understood that this is the state of the process at time  $-1$ . We write  $\mathbb{P}_v$  for the law of the  $\mathcal{U}_{x,t}$  field, from which the process is constructed.

### 1.1.2 Examples

Let us now introduce a few relevant examples.

**Toom rule with death** The Toom rule [28, Example 1] is a deterministic CA in two dimensions, which updates the state of each site  $x \in \mathbb{Z}^2$  to the more common value (in  $\{0, 1\}$ ) among the current state of  $x$ ,  $x + (1, 0)$  and  $x + (0, 1)$ . We further subject this CA to a specific type of noise, obtaining a PCA that we will refer to as *Toom rule with death*. Namely, at each step

and each site independently with probability  $1 - p$ , instead of applying the previous rule, we directly set it to state 0.

With our notation this corresponds to  $d = 2$ ,  $r = 1$  and  $v$  charging only two up-families:  $v(\{\emptyset\}) = 1 - p$  (we will systematically put accolades to avoid confusing e.g. the singleton system consisting of the empty up-family appearing above and the empty system, which naturally verifies  $v(\emptyset) = 0$ ) and

$$v(\{\{X \subset R : |X \cap \{(0, 0, -1), (1, 0, -1), (0, 1, -1)\}| \geq 2\}\}) = p.$$

This PCA is attractive, but not additive. For  $p = 1$  it degenerates into the (deterministic) CA called Toom rule. In fact, Toom [28] studied random perturbations of attractive CA in much greater generality, but we will come back to this later.

**Generalised oriented site percolation** Fix  $X \subset R$  and  $p \in [0, 1]$ . We define GOSP to be the additive PCA with neighbourhood  $X$  by  $v(\{\emptyset\}) = 1 - p$  and  $v(\{\{Y \subset R : Y \cap X \neq \emptyset\}\}) = p$ . The name comes from the observation that  $\eta^{\{0\}}(t) \neq \mathbf{0}$  if and only if there is a path with steps in  $-X$  from 0 to some site of the form  $(x, t) \in \mathbb{Z}^{d+1}$ , using only sites  $(y, s) \in \mathbb{Z}^{d+1}$  such that  $\mathcal{U}_{y,s} \neq \emptyset$ . The standard oriented site percolation model is recovered by taking  $X = \{(-1, -1), (1, -1)\}$  in one dimension.

**Bootstrap percolation** A BP model is specified by an *update family*: a finite family  $\mathcal{X}$  of finite subsets of  $\mathbb{Z}^d \setminus \{0\}$ , both the sets and the family being possibly empty. At each time step the state of a site  $x \in \mathbb{Z}^d$  becomes 1 if it is already 1 or there exists  $X \in \mathcal{X}$  such that all elements of  $x + X$  are in state 1. In fact, this is just another way to parametrise the set of all attractive CA with no memory which are monotone in time.<sup>1</sup>

With our notation BP corresponds to taking as  $v$  the Dirac measure on the minimal up-family  $\mathcal{U} \subset \Omega_R$  such that  $\{X \times \{-1\} : X \in (\mathcal{X} \cup \{\{0\}\})\} \subset \mathcal{U}$ . If we drop the assumption that  $\{0\} \times \{-1\} \in \mathcal{U}$ , we retrieve the class of all attractive CA with no memory. The latter is essentially the class of models whose random perturbations were studied by Toom [28].

More generally, we define *inhomogeneous BP* by a measure  $\chi$  on update families. Then each site  $x \in \mathbb{Z}^d$  is assigned an i.i.d. update family  $\mathcal{X}_x$  with law  $\chi$  and at each step at site  $x$  we use the minimal up-family  $\mathcal{U}_x \subset \Omega_R$  such that  $\{X \times \{-1\} : X \in (\mathcal{X}_x \cup \{\{0\}\})\} \subset \mathcal{U}_x$  and define the evolution via

---

<sup>1</sup>This property is sometimes called *freezing* to distinguish from attractiveness, which is also a type of monotonicity.

Eq. (1)<sup>2</sup> with  $\mathcal{U}_{x,t} = \mathcal{U}_x$  for all  $t$ . Clearly, this is no longer a CA or a PCA, but rather what one would call *an inhomogeneous attractive CA*.

**PCA with death** Given an attractive PCA, we define its version with death by considering  $\tilde{v} = pv + (1-p)\delta_\emptyset$ , so that  $\tilde{v}$  defines another attractive PCA. In words, we run the original PCA with probability  $p$  and put state 0 with probability  $1-p$ , like we did for the Toom rule with death and GOSP.

**Kinetically constrained models** KCM are continuous time Markov processes with state space  $\{0,1\}^{\mathbb{Z}^d}$  informally defined as follows, given an *update family*: a finite family  $\mathcal{X}$  of finite subsets of  $\mathbb{Z}^d \setminus \{0\}$ , and a parameter  $q \in [0,1]$  (see [10]). Each site  $x \in \mathbb{Z}^d$  attempts to update at rate 1 to an independent Bernoulli state with parameter  $q$ , but is only allowed to do so if the configuration at some  $X \in \mathcal{X}$  is  $\mathbf{1}$ . Otherwise, the state of  $x$  cannot change until the above constraint becomes satisfied.

Superficially, KCM are not closely related to the attractive PCA that we study, as they are neither attractive, nor discrete time, nor synchronous. Nevertheless, we will see that our treatment entails new results for KCM as well.

## 1.2 The phase diagram of PCA

In 1994 Bezuidenhout and Gray [6] established the following fundamental result (see their work for a more formal statement).

**Theorem 1.1.** *Within the set of attractive PCA  $v$  without memory, those with  $v(\{\emptyset\}) > 0$  and  $\mathbb{P}_v(\forall t > 0, \eta^{\{0\}}(t) \neq \mathbf{0}) > 0$  is open.*

The main corollary of this is that the phase transition of survival from a single point for attractive PCA without memory but with positive death rate is continuous. Moreover, they showed that, within this ‘supercritical’ phase described in the theorem, one can perform renormalisation to highly supercritical oriented percolation. This entails a number of results and is a key step towards establishing that models in this phase are ‘well-behaved’ (see [17] for more detail on what we mean by this).

Our first main result is of a similar flavor and complements Theorem 1.1.

**Theorem 1.2.** *Let  $S$  be in the the set of attractive PCA such that  $\delta_{\mathbf{0}}$  is their unique invariant measure. Let  $\text{Int}$  and  $\bar{\cdot}$  be the interior and the closure*

---

<sup>2</sup>There may be issues defining this if  $\chi$  has infinite support. We will only consider finitely supported  $\chi$  measures in this work.

within the set of absorbing attractive PCA. Then for any  $\nu \in \text{Int}(S)$  there exist  $c, C > 0$  such that for all  $t > 0$  and finite  $A \subset \mathbb{Z}^d \times \{-r, \dots, -1\}$  it holds that

$$\mathbb{P}_\nu(\eta_0^{\mathbf{1}}(t) \neq \mathbf{0}) \leq C e^{-ct}, \quad (2)$$

$$\mathbb{P}_\nu(\eta^A(t) \neq \mathbf{0}) \leq C|A|e^{-ct}. \quad (3)$$

Moreover,  $S \subset \overline{\text{Int}(S)}$ .

This result can be informally rephrased as ‘the subcritical phase of attractive PCA is well-behaved.’ We leave the discussion of how to use this on concrete models to Section 1.3.

Unfortunately, the terms ‘supercritical’ for Theorem 1.1 and ‘subcritical’ for Theorem 1.2 are quite misleading. Indeed, it is known that there is an intermediate regime, which may naturally be called *cooperative survival* phase. More specifically, the Toom rule with death for small enough death rates exhibits such behaviour (see [28]) and more general examples will be discussed in Section 1.4.

### 1.3 Parametrised models and applications

In practice one is not interested in the set of *all* PCA, but rather has one specific PCA in mind, possibly with a parameter to tune. As it stands, Theorem 1.2 does not assert that for a specific model, as we vary the parameter, we will have a well-behaved subcritical phase immediately followed by a non-uniqueness one. Our next goal is to provide a simple sufficient criterion for this to happen. Indeed, hypotheses are needed, since the model of interest may glide along the boundary of the subcritical phase, in which case no sharpness of the phase transition is to be expected. To preclude this scenario we will require a few more definitions.

We say that a measure  $\nu$  on a partially ordered set  $(X, \geq)$  *stochastically dominates* another one,  $\nu$ , if for every down-set  $D$  of  $X$  we have  $\nu(D) \leq \nu(D)$ . A *parametrised model* is a continuously differentiable curve  $(\nu_p)_{p \in [0,1]}$  in the space of attractive PCA, so that  $\nu_p$  are probability measures on  $\mathcal{U}$ . Note that the function  $p \mapsto \nu(p)$  is nondecreasing for stochastic domination iff

$$\forall \mathcal{D} \in \mathfrak{D} \setminus \{\emptyset, \mathcal{U}\}, \forall p \in [0, 1], \nu'_p(\mathcal{D}) \leq 0. \quad (4)$$

If that is the case, we say that the parametrised model is *nondecreasing* and, if additionally  $\nu_p \neq \nu_q$  whenever  $p \neq q$ , we say it is *increasing*.

Morally, any increasing parametrised model should have a sharp phase transition in the sense of Theorem 1.2 (see Theorem 1.3 below). However, even for percolation some issues may arise if the model increases in

a ‘nonessential’ way. Furthermore, there has been some trouble establishing a necessary and sufficient condition for a modification being ‘essential’ [2, 4]. Circumventing these issues, we propose a sufficient criterion, which is satisfied in natural models.

We say the parametrised model is *strongly increasing* if there exists  $c > 0$  such that

$$\forall \mathcal{D} \in \mathfrak{D} \setminus \{\emptyset, \mathcal{U}\}, \forall p \in [0, 1], v'_p(\mathcal{D}) \leq -c(1 - v_p(\mathcal{D})), \quad (5)$$

where the derivative is w.r.t.  $p$ . For a nondecreasing parametrised model we define the *critical parameter*

$$p_c = \inf \{p \in [0, 1] : \delta_0 \text{ is not the unique invariant measure}\} \quad (6)$$

with  $\inf \emptyset = 1$ . Note that  $v_p(\{\emptyset\}) = 0$  iff  $\delta_1$  is invariant, while if  $v_p(\{\Omega_R\}) = 0$  iff  $\delta_0$  is invariant. We will say that a nondecreasing parametrised model has *exponential decay* if for any  $p < p_c$  there exist  $c, C > 0$  such that Eqs. (2) and (3) hold for all  $t > 0$  and finite  $A \subset \mathbb{Z}^d \times \{-r, \dots, -1\}$ .

**Theorem 1.3.** *Every strongly increasing parametrised model has exponential decay.*

We next state a few interesting applications to specific models. The first one recovers a classical result of Aizenman and Barsky [1, Theorem 7.3] (also see their section 8.1 for a discussion of additive PCA) and Menshikov [22], which also has other proofs [11, 12].

**Corollary 1.4.** *Every GOSP has exponential decay.*

The verification of Eq. (5) for GOSP is immediate, but we may also directly use Theorem 2.1 below instead of Theorem 1.3.

The next consequence makes further progress towards proving [16, Conjecture 8.1], generalising a question of Schonmann [24] from 1992, which remains open.

**Theorem 1.5.** *Consider a BP with update family  $\mathcal{X}$  contained in a half-space: there exists  $u \in \mathbb{R}^d$  such that  $\forall X \in \mathcal{X}, \forall x \in X, \langle x, u \rangle < 0$ . We use i.i.d. Bernoulli initial condition with parameter  $q$ , whose law we denote by  $\mu_q$  (BP being a CA, this is the only randomness). Then, setting*

$$q_c = \inf \left\{ q > 0 : \lim_{t \rightarrow \infty} \mu_q(\eta_0(t) \neq 1) = 0 \right\}. \quad (7)$$

we have

$$\forall q > q_c, \exists c, C > 0, \forall t > 0, \quad \mu_q(\eta_0(t) \neq 1) \leq Ce^{-ct}. \quad (8)$$

It should be noted that this application is not as direct, since BP has  $v(\{\emptyset\}) = 0$  by definition. Instead we will rely on the correspondence between BP and CA of Proposition 3.1. This correspondence presented in Section 3 is also at the root of the results discussed in Section 1.4.

Theorem 1.5 generalises directly to inhomogeneous BP as follows.

**Theorem 1.6.** *Consider an inhomogeneous BP with measure  $\chi$  supported on a finite set of update families contained in the same half-space (see Theorem 1.5). Denote by  $\mu_q$  the law of the i.i.d. update families with distribution  $\chi$  and i.i.d. initial condition with Bernoulli law of parameter  $q$ . Then Eq. (8) holds with  $q_c$  defined by Eq. (7).*

Moving on to KCM, the following is a direct consequence of Theorem 1.5 together with [16, Theorem 3.7].

**Corollary 1.7.** *Consider a KCM with update family  $\mathcal{X}$  contained in a half space (see Theorem 1.5). Then the spectral gap of its generator is positive for all  $q > q_c$  and 0 for all  $q < q_c$ , where  $q_c$  is the quantity defined in Eq. (7) for the BP with the same choice of update family  $\mathcal{X}$ .*

## 1.4 Bootstrap percolation and Toom perturbations of attractive cellular automata

Finally, we discuss consequences of the PCA representation of BP used to prove Theorem 1.5.

**Bootstrap percolation universality** We first need to introduce a few notions from BP universality. We will be led to restrict several of the following statements to  $d = 1$  or  $d = 2$ , as appropriate. However, let us emphasise that this is only because the rough BP universality has been established in these dimensions in [3, 8]. Yet, the higher dimensional theory was announced several years ago. When it becomes available, the results below will immediately generalise to any dimension.

We say that a BP update family  $\mathcal{X}$  is called *supercritical* if there exists a finite set  $A \subset \mathbb{Z}^d$  such that  $\bigcup_{t>0} \eta^A(t)$  is infinite. In  $d = 2$  (and, incidentally  $d = 1$ , since one-dimensional models naturally immerse in two dimensions) a more geometric characterisation is available based on the notion of stable direction. We say that  $u \in S^{d-1}$  (the unit sphere) is *unstable* if there exists  $X \in \mathcal{X}$  such that for all  $x \in X$  we have  $\langle u, x \rangle < 0$  and it is *stable otherwise*. It was proved in [8, Theorem 7.1, Definition 1.3] that in two dimensions  $\mathcal{X}$  is supercritical if and only if there exists an open hemisphere of unstable directions. Instead, we say that  $\mathcal{X}$  is *subcritical* if  $q_c > 0$  (recall Eq. (7)). It



was proved in [3, Corollary 2] that in two dimensions  $\mathcal{X}$  is subcritical if and only if every open hemisphere contains an open set of stable directions.

**Toom perturbations** In [28] random space-time perturbations of attractive CA with  $v(\{\emptyset\}) = 0$  (otherwise every configuration becomes  $\mathbf{0}$  after one step, since CA are deterministic) were considered, in order to assess whether the  $\mathbf{0}$  and  $\mathbf{1}$  states are stable. We will not define the exact noise used there, but using attractiveness to suppress ‘positive’ noise, it can be brought down to a noise stochastically dominated by a product measure with low parameter. For simplicity, we will work directly with the resulting product noise. Namely, we consider the CA with death  $\tilde{v}$ . We say that the CA  $v$  is an *eroder* if, starting from any configuration  $\omega$  such that  $\omega_{(x,s)} = 0$  for finitely many  $(x, s) \in \mathbb{Z}^d \times \{-r, \dots, -1\}$ , we have  $\eta^\omega(t) = \mathbf{1}$  for any  $t$  large enough depending on  $\omega$ .

**Interplay** We can recover the following result of Toom [28] directly from the BP–PCA correspondence and BP results.

**Theorem 1.8.** *In dimension  $d = 1$  any attractive CA is an eroder iff its version with death rate  $1 - p$  small enough has an invariant measure different from  $\delta_{\mathbf{0}}$ .*

It should be noted that proving this does not at all require the full power of [3, Theorem 1], but only the much easier partial result [3, Theorem 9] in view of Lemma 3.3 below. We believe this renormalisation approach to be simpler than the original proof of Toom, as well as its subsequent versions in [5, 13] based on somewhat involved Peierls arguments. Naturally, a carefully performed Peierls argument often gives a better quantitative bound on the critical parameter (see e.g. [25]), but this is seldom important. We also mention the more complex renormalisation approach of [9].

Inversely, the viewpoint of [28, Theorem 5] provides an interesting consequence for BP, recovering the main results of [3, 8] restricted to families contained in half-spaces, but extended to arbitrary dimension.

**Theorem 1.9.** *A BP in any dimension whose update family is contained in a half-space (recall Theorem 1.5) is either supercritical or subcritical (that is, either there are finite sets with infinite offspring or  $q_c > 0$ ).*

Although in two dimensions this can be checked easily (see Lemma 3.3 below) from geometric considerations based on the characterisations of supercritical and subcritical models from [3, 8], it does not seem to have been noticed. We should note that the half-space condition cannot be removed, as without it, already in two dimensions, another, *critical*, class emerges [8].

## 1.5 Organisation

The remainder of the paper is organised as follows. In Section 2 we prove our general results—Theorems 1.2 and 1.3 by adapting the randomised algorithm approach of Duminil-Copin, Raoufi and Tassion [11]. In Section 3 we introduce the correspondence between BP and CA and deduce Theorems 1.5, 1.6, 1.8 and 1.9.

## 2 Exponential decay

### 2.1 Linear parametristion

We first seek to prove the following preliminary result, from which Theorems 1.2 and 1.3 will be deduced in Section 2.2.

**Theorem 2.1.** *Let  $v$  be a rates measure and let  $v_p = pv + (1 - p)\delta_\emptyset$ . This nondecreasing parametrised model has exponential decay.*

The first step is a Russo formula adapted to our situation. We say that a space-time site  $(x_0, t_0)$  is *pivotal* for an event  $A$  and realisation of the  $\mathcal{U}_{x,t}$  variables, if  $A$  occurs for  $(\mathcal{U}_{x,t})_{(x,t) \in \mathbb{Z}^{d+1}}$ , but it does not occur if we replace  $\mathcal{U}_{x_0, t_0}$  by  $\emptyset$ .

**Lemma 2.2.** *Let  $v$  be a rates measure and let  $v_p = pv + (1 - p)\delta_\emptyset$ . Let  $A_n$  be the event that  $\eta_0^1(n) \neq 0$  and set  $\theta_n(p) = \mathbb{P}_{v_p}(A_n)$ . Then for all  $n \geq 0$  and  $p \in (0, 1]$*

$$\theta'_n(p) = \sum_{x,t} \mathbb{P}_{v_p}((x,t) \text{ is pivotal for } A_n).$$

*Proof.* Fix an i.i.d. random field  $\mathcal{U}_{x,t}^1$  with law  $v$  and i.i.d. random variables  $X_{x,t}$  uniform on  $[0, 1]$  and define

$$\mathcal{U}_{x,t}^p = \begin{cases} \mathcal{U}_{x,t}^1 & \text{if } X_{x,t} \leq p \\ \emptyset & \text{if } X_{x,t} > p. \end{cases}$$

Conditioning on the  $\mathcal{U}_{x,t}^1$ , we can then proceed as in the proof of the standard Russo formula (see e.g. [15, Sec. 2.4]). We then average over  $\mathcal{U}_{x,t}^1$  to obtain the desired equality.  $\square$

The next step is to adapt the Duminil-Copin–Raoufi–Tassion version of the O’Donnell–Saks–Schramm–Servedio decision tree result (see [11, 23] for background). It is important to note that the support of  $v$  is not necessarily binary or even totally ordered, so it is not clear how to adapt [11, Theorem

1.1]. We circumvent this problem by restricting our attention to events rather than real-valued observables.

**Lemma 2.3.** *Let  $n \in \mathbb{N}$  and  $A \subset \mathcal{U}^n$  be an increasing event (w.r.t. the pointwise inclusion order). Fix a randomised algorithm determining the occurrence of  $A$  and denote by  $\delta_i$  the probability that it reveals the value of the  $i$ -th up-family. Then*

$$\text{Var}_v(\mathbb{1}_A) \leq 2 \sum_{i=1}^n \delta_i \cdot \mathbb{P}_v(i \text{ is pivotal for } A)$$

where  $\text{Var}$  and  $\mathbb{P}$  are w.r.t. a product measure  $\bigotimes_{i=1}^n v$  on  $\mathcal{U}^n$ .

*Proof.* The result is proved like Theorem 1.1. of [11] (also see Remark 2.2. there). It suffices to replace their equation (5) by the fact that if  $\eta \in \mathcal{U}^n$  and  $\omega \in \mathcal{U}^n$  differ only at the  $i$ -th up-family, then

$$|\mathbb{1}_A(\eta) - \mathbb{1}_A(\omega)| \leq \mathbb{1}_{i \text{ is pivotal for } A}(\eta) + \mathbb{1}_{i \text{ is pivotal for } A}(\omega). \quad (9)$$

To see this, assume w.l.o.g. that  $\eta \notin A$  (if  $\mathbb{1}_A(\eta) = \mathbb{1}_A(\omega)$  there is nothing to prove). Then replacing  $\eta_i$  by  $\emptyset$  does not trigger the occurrence of  $A$ , as  $A$  is increasing. Hence, either  $\omega \notin A$  and the l.h.s. in Eq. (9) is 0, or  $i$  is pivotal for  $A$  and  $\omega$ , so the r.h.s. is 1.  $\square$

Our next task is to define a suitable randomised algorithm to which Lemma 2.3 will be applied.

**Lemma 2.4.** *Fix a  $p \in [0, 1]$  and a parametrised model. Recall  $A_n = \{\eta_0^1(n) \neq 0\}$  and  $\theta_n(p) = \mathbb{P}_p(A_n)$ . For each integer  $n \geq 1$  there exists a randomised algorithm determining the occurrence of  $A_n$  such that for every  $(x, t) \in \mathbb{Z}^{d+1}$  its revelation probability  $\delta_{x,t}$  satisfies*

$$\delta_{x,t} \leq \frac{2}{n} \sum_{i=0}^{n-1} \theta_i(p).$$

*Proof.* Clearly, it suffices to restrict our attention to the up-families in

$$S = \{(x, t) \in \mathbb{Z}^{d+1} : |x| \leq r \cdot (n - t), t \in [0, n]\}, \quad (10)$$

where  $r$  is the range of the process. The algorithm proceeds as follows (following [11, Lemma 3.2] and [16, Lemma 7.3]). Select a number  $k$  uniformly at random in  $\{1, \dots, n\}$ . Contrary to [16], we will explore forward in time, starting from time  $k$ . For  $t \in \{0, \dots, n - k\}$  denote by  $\omega^1(t)$  the configuration with initial condition  $\mathbf{1}$  and up-families  $\mathcal{U}'_{x,t} = \mathcal{U}_{x,t+k}$ , where  $\mathcal{U}_{x,k}$  are the

up-families in the graphical construction of  $\eta$ . We initialise the algorithm at  $t = 0$ . We explore the state of each  $\mathcal{U}_{x,t+k}$  for  $(x, t+k) \in S$  such that the currently explored up-families do not allow to conclude that  $\omega_x^1(t) = 0$  (so, at the first step that is all  $\mathcal{U}_{x,k}$  for  $(x, k) \in S$ ). Then we increment  $t$  and repeat the previous operation. Upon reaching  $t = n - k$ , if we are able to conclude that  $\omega_0^1(n - k) = 0$ , we terminate the algorithm (since we know by attractiveness that  $\eta_0^1(n) \leq \omega_0^1(n - k) = 0$ ). Otherwise we reveal all up-families in  $S$  to determine if  $A_n$  occurs and terminate.

Next observe that, conditionally on  $k$ , the probability of revealing  $U_{x,t}$  for  $(x, t) \in S$  is at most  $\theta_{n-k}(p) + \mathbb{1}_{t \geq k} \theta_{t-k}(p)$ , the first term bounding the probability that we reach  $t = n - k$  without having  $\omega_0^1(n - k) = 0$ . Averaging over the law of  $k$ , we obtain the desired conclusion.  $\square$

We are now ready to assemble the proof of Theorem 2.1.

*Proof of Theorem 2.1.* Fix  $v \neq \delta_\emptyset$  (otherwise the statement is trivial) and  $v_p = pv + (1-p)\delta_\emptyset$ . This clearly defines a nondecreasing parametrised model, since  $\mathcal{U} \supset \emptyset$  for all  $\mathcal{U} \in \mathcal{U}$ . Define  $A_n$ ,  $\theta_n(p)$  and  $S$  as above (see Lemma 2.4 and Eq. (10)). By Lemmas 2.3 and 2.4 we have

$$\theta_n(p)(1 - \theta_n(p)) \leq \frac{4}{n} \mathbb{E}_{v_p}[N] \sum_{i=0}^{n-1} \theta_i(p),$$

where  $N$  is the number of  $(x, t) \in S$  which are pivotal for  $A_n$ . Then, applying Lemma 2.2 we get

$$\theta'_n(p) \geq \frac{n\theta_n(p)(1 - \theta_n(p))}{4 \sum_{i=0}^{n-1} \theta_i(p)}.$$

Observing that  $1 - \theta_n(p) \geq v_p(\{\emptyset\}) = pv(\{\emptyset\}) + 1 - p$ , we may apply [11, Lemma 3.1] to conclude the proof of Eq. (2). The only subtlety arises in the neighbourhood of  $p = 1$  if  $v(\{\emptyset\}) = 0$ . Nevertheless, it suffices to apply [11, Lemma 3.1] on  $p \in [0, 1 - \varepsilon]$  for  $\varepsilon > 0$  arbitrarily small to obtain Eq. (2).

Finally, it remains to derive Eq. (3). Starting from a finite set  $A$ , the only sites which may be in state 1 at time  $t$  are those at distance at most  $rt$  from a site in  $A$ . Hence, taking the union bound of Eq. (2) over these sites and using attractiveness, we get

$$\mathbb{P}_{v_p}(\eta^A(t) \neq \mathbf{0}) \leq |A|rt \max_{x \in \mathbb{Z}^d} \mathbb{P}_{v_p}(\eta_x^A(t) \neq 0) \leq C|A|rte^{-ct} \leq C'|A|e^{-c't}$$

for a suitable choice of  $C', c' > 0$ , yielding Eq. (3).  $\square$

## 2.2 Parametrisations

With Theorem 2.1, we are in position to conclude the proof of Theorem 1.2.

*Proof of Theorem 1.2.* Let  $\nu$  be in the interior of the set of absorbing attractive PCA such that  $\delta_{\mathbf{0}}$  is their unique invariant measure, denoted by  $\text{Int}(S)$ . Clearly, we cannot have  $\nu(\{\emptyset\}) = 0$ , since in that case  $\delta_{\mathbf{1}}$  is invariant. Therefore, we may consider the linearly parametrised model  $\nu_p = p\nu + (1-p)\delta_{\emptyset}$  for  $p \in [0, 1 + \nu(\{\emptyset\})]$ . Since this parametrised model is absorbing and  $\nu_{\mathbf{1}} \in \text{Int}(S)$ , necessarily  $p_c > 1$ . But then Theorem 2.1 yields Eqs. (2) and (3) for  $\nu = \nu_{\mathbf{1}}$ .

Similarly, assume that  $\delta_{\mathbf{0}}$  is the unique invariant measure of the  $\nu$ -PCA, but not necessarily  $\nu \in \text{Int}(S)$ . Then Theorem 2.1 gives that  $p_c \geq 1$ , so that Eqs. (2) and (3) hold for any  $p < 1$ . Then Proposition 2.5 below allows us to conclude that  $\nu_p \in \text{Int}(S)$  for  $p \in [0, 1)$  and conclude that  $\nu \in \overline{\text{Int}(S)}$ .  $\square$

**Proposition 2.5.** *The set of attractive absorbing PCA for which there exist  $c, C > 0$  so that Eq. (2) holds for all  $t > 0$  is open.*

*Proof.* We use a standard renormalisation argument. Fix  $\nu$  such that Eq. (2) holds for some  $c, C > 0$  and let  $\varepsilon > 0$  small enough to be chosen later. Partition  $\mathbb{Z}^{d+1}$  into the boxes  $B_{x,t} = (xL, tL) + [0, L)^{d+1}$  for some large  $L$  to be chosen later. Let  $\mathcal{B}_{x,t}$  denote the event that the  $\nu$ -PCA with initial condition  $\mathbf{1}$  and up-family field translated by  $-L(x, t)$  is 1 for some  $(y, s) \in [0, L)^d \times [L-r, L)$ . By Eq. (2) and the union bound we have that for  $L$  large enough  $\mathbb{P}_{\nu}(\mathcal{B}_{x,t}) < \varepsilon$  for any  $(x, t) \in \mathbb{Z}^{d+1}$ . Moreover,  $\mathcal{B}_{x,t}$  only depends on the up-families  $\mathcal{U}_{y,s}$  for  $(y, s) \in B_{x',t'}$  with  $(x', t')$  at distance at most  $r$  from  $(x, t)$ .

Fix an attractive absorbing PCA  $\nu'$  sufficiently close to  $\nu$  (depending on  $\varepsilon$  and  $L$ ). Then we can couple the corresponding up-family fields  $\mathcal{U}_{y,s}$  and  $\mathcal{U}'_{y,s}$ , so that they differ with small probability and independently for each  $(y, s) \in \mathbb{Z}^{d+1}$ . We say that the box  $B_{x,t}$  is *bad* if  $\mathcal{B}_{x,t}$  occurs or there exists a site  $y, s$  at distance at most  $rL$  from  $B_{x,t}$  such that  $\mathcal{U}_{y,s} \neq \mathcal{U}'_{y,s}$ . By the well-known Liggett–Schonmann–Stacey theorem [19] the good boxes stochastically dominate a product measure with density of bad boxes approaching 1 if we choose  $\varepsilon$  small enough.

Finally, we consider oriented percolation of bad boxes with range  $r$  and observe that if a box does not belong to a bad connected component reaching time 0, then the state at its top boundary is necessarily  $\mathbf{0}$  in both the  $\nu$  and the  $\nu'$ -PCA with initial condition  $\mathbf{1}$ . Indeed, if the box  $B_{x,t}$  itself is not bad that is enough by definition of bad, boxes, while otherwise we can proceed by induction. Namely, we observe that if the state of all boxes  $B_{y,t-1}$  for

$\|x - y\|_\infty \leq r$  are good, then their top boundaries are in state  $\mathbf{0}$  and no absorbing PCA of range  $r$  can reach a nonzero state at the top of  $B_{x,t}$ , starting from that.

We can then conclude that Eq. (2) holds for  $\nu$  (and different  $c, C > 0$ ), since in highly subcritical independent percolation of range  $r$  the size of clusters has this exponential decay property (e.g. by Corollary 1.4).  $\square$

Our next goal is to prove Theorem 1.3. We will deduce this from the specific case, Theorem 2.1, and the following result.

**Lemma 2.6.** *Let  $(\nu_p)_{p \in [0,1]}$  be a strongly increasing parametrised model. Then there exists  $c' > 0$  such that for all  $p \in (0, 1]$  there exists  $\varepsilon > 0$  such that for all  $p' \in [1 - \varepsilon, 1]$  it holds that  $\nu_{p'} := p'\nu_p + (1 - p')\delta_\emptyset$  stochastically dominates  $\nu_{p-(1-p')/c'}$ .*

*Proof.* Notice that for any down-system  $\mathcal{D} \in \mathfrak{D} \setminus \{\emptyset, \mathcal{U}\}$  the desired inequality  $\nu_{p'}(\mathcal{D}) \leq \nu_{p-(1-p')/c'}(\mathcal{D})$  is an equality for  $p' = 1$ . Therefore, it suffices to show that the derivatives satisfy the inverse inequality:

$$\nu_p(\mathcal{D}) - 1 = \nu_{p'}(\mathcal{D}) \geq \frac{\partial \nu_{p-(1-p')/c'}}{\partial p'}(\mathcal{D}) = \frac{\nu'_{p-(1-p')/c'}(\mathcal{D})}{c'} \quad (11)$$

in the neighbourhood of  $p' = 1$ .

If  $\nu_p(\mathcal{D}) = 1$ , we may conclude directly by the fact that the parametrised model is nondecreasing. Otherwise, recalling Eq. (5) and setting  $c' = c/2$ , we have

$$\frac{\nu'_{p-(1-p')/c'}(\mathcal{D})}{c'} \leq 2 (\nu_{p-(1-p')/c'}(\mathcal{D}) - 1) \xrightarrow{p' \rightarrow 1} 2 (\nu_p(\mathcal{D}) - 1) < \nu_p(\mathcal{D}) - 1.$$

Hence, for  $p'$  sufficiently close to 1, Eq. (11) does hold for all  $\mathcal{D} \in \mathfrak{D} \setminus \{\emptyset, \mathcal{U}\}$ , since  $\mathfrak{D}$  is finite.  $\square$

*Proof of Theorem 1.3.* Consider a strongly increasing parametrised model  $(\nu_p)_{p \in [0,1]}$  and  $p \in (0, p_c)$  (if  $p_c = 0$  there is nothing to prove). Set  $\nu_{p'} = p'\nu_p + (1 - p')\delta_\emptyset$  for  $p' \in [1 - \varepsilon, 1]$ , where  $\varepsilon$  is from Lemma 2.6. Let  $p'_c$  denote the critical value of this parametrised model (defined like in Eq. (6) with the slight abuse that the parametrisation ranges over  $p' \in [1 - \varepsilon, 1]$  rather than  $p' \in [0, 1]$ ). Then Theorem 2.1 shows that Eqs. (2) and (3) hold for  $\nu_{p'}$  for  $p' < p'_c$ . By Lemma 2.6 and the fact that  $(\nu_p)_{p \in [0,1]}$  is nondecreasing, it suffices to prove that  $p'_c = 1$ . But this is clear, because the definition of  $p_c$  gives that  $\delta_0$  is the unique invariant measure for  $\nu_p = \nu_1$ .  $\square$

## 3 Bootstrap percolation and cellular automata

### 3.1 The correspondence

It was noticed already by Schonmann [24] that standard oriented site percolation, instead of a one-dimensional PCA or a two-dimensional percolation, can be viewed as a two-dimensional BP with an i.i.d. initial condition. This was exploited in [16] and extended to GOSP subsequently studied in [17]. We now show that this correspondence is in fact extends to all attractive PCA.<sup>3</sup>

**Proposition 3.1** (BP–CA correspondence). *Consider an attractive CA with  $v = \delta_{\mathcal{U}}$  for some  $\mathcal{U} \in \mathcal{U}$ . Let  $\tilde{\eta}$  be its version with death with rates measure  $\tilde{v} = pv + (1 - p)\delta_{\emptyset}$ . Let  $\mathcal{U}_{x,t}$  be the up-family field used to construct the  $\tilde{v}$ -PCA. Define the  $d + 1$ -dimensional up-family*

$$\mathcal{X} = \{R \setminus D : D \in \Omega_R \setminus \mathcal{U}\}. \quad (12)$$

*Consider the BP process  $\omega$  defined by the update family  $\mathcal{X}$  with initial condition  $(\mathbb{1}_{\mathcal{U}_{x,t}=\emptyset})_{(x,t) \in \mathbb{Z}^{d+1}}$ . Define its closure*

$$C = \{(x, t) \in \mathbb{Z}^{d+1} : \exists s \in \{0, 1, \dots\}, \omega_{(x,t)}(s) = 1\} \quad (13)$$

*and the configuration  $C_0 = (\mathbb{Z}^d \times \{-r, \dots, -1\}) \setminus C$ . Then for all  $t \geq 0$*

$$\tilde{\eta}^{C_0}(t) = \{x \in \mathbb{Z}^d : (x, t) \notin C\} \quad (14)$$

*and this is a version of the stationary  $\tilde{v}$ -PCA at its upper invariant measure, that is, the limit as  $t \rightarrow \infty$  of the law of  $\tilde{\eta}^1(t)$ . In particular,  $C = \mathbb{Z}^{d+1}$  a.s. if and only if  $\delta_{\mathbf{0}}$  is the unique invariant measure of the  $\tilde{v}$ -PCA.*

*Moreover, the map  $\mathcal{U} \leftrightarrow \mathcal{X}$  is a one-to-one correspondence between  $d$ -dimensional attractive CA and  $d + 1$ -dimensional BP with update family contained in the lower half-space  $\mathbb{Z}^d \times \{-1, -2, \dots\}$ .*

*Proof.* The bijectiveness follows from the fact that the *double complement* map  $\mathcal{U} \mapsto \{R \setminus D : D \in \Omega_R \setminus \mathcal{U}\}$  is an involution of the power set of  $\Omega_R$ , since the two complements commute and each of them is an involution.

Fix an attractive CA and its corresponding BP as in the statement. Let us first verify Eq. (14). By induction on  $t$ , it suffices to verify this for  $t = 0$ .

By definition  $(x, 0) \in C$  iff  $\mathcal{U}_{x,0} = \emptyset$  (this is the initial condition of  $\omega$ ) or there exists a minimal  $s > 0$  such that  $\omega_{(x,0)}(s) = 1$ . The latter, happens

---

<sup>3</sup>We refer the reader to [18] for a related correspondence between PCA and equilibrium statistical mechanics models.

iff  $R \setminus (\omega(s-1) - (x, 0)) \notin \mathcal{U}$  and  $\omega_{(x,0)}(s-1) = 0$ . Hence,  $(x, 0) \in C$  iff  $\mathcal{U}_{x,0} = \emptyset$  or  $R \cap (C_0 - (x, 0)) \notin \mathcal{U}$ . But this is equivalent to  $\tilde{\eta}_x^{C_0}(0) = 0$  by definition, so Eq. (14) indeed holds.

Thus,  $1 - \mathbb{1}_C$  is a trajectory of the  $\tilde{v}$ -PCA. Moreover, its law is clearly invariant by translation in  $\mathbb{Z}^{d+1}$ , so the process  $\tilde{\eta}_0^C$  is stationary. It remains to verify that this corresponds to the upper invariant measure. To see this it suffices to prove that

$$1 - \omega_{(x,s)}(t) \geq \tilde{\eta}_x^1(s) \quad (15)$$

for all  $(x, t, s) \in \mathbb{Z}^d \times \{1, 2, \dots\}^2$  such that  $s \geq rt$ , since  $1 - \mathbb{1}_C$  is the decreasing limit of  $1 - \omega(t)$ . Notice that  $1 - \omega_{(x,s)}(t)$  in fact only depends on the initial condition of  $\omega$  for  $(y, u) \in \mathbb{Z}^d \times \{0, \dots, s\}$ , since  $s \geq rt$ . Define the closure  $C'$  as in Eq. (13) with the process  $\omega'$  defined like  $\omega$  but with initial condition  $\mathbb{1}_{\mathcal{U}_{y,s}=\emptyset}$  for  $(y, u) \in \mathbb{Z}^d \times \{0, \dots, s\}$  and 0 elsewhere. Then in fact  $1 - \omega_{(x,s)}(t) = 1 - \omega'_{(x,s)}(t) \geq 1 - \mathbb{1}_{(x,s) \in C'} = \tilde{\eta}_x^1(s)$ . The last equality is Eq. (14) applied to a suitable choice of initial condition (which we may choose freely, since the relation is deterministic and not just a.s.). Thus, Eq. (15) is established and the proof is complete.  $\square$

Amusingly, since BP is an attractive CA, one can iterate this correspondence. As an example, consider the 0-dimensional CA with no memory given by the identity map  $0 \mapsto 0; 1 \mapsto 1$  (there are four 0-dimensional CA without memory—the identity, the constant 0, the constant 1 and a non-attractive one). It is clearly not an eroder, since 0 does not become 1. Its corresponding 1-dimensional BP model is East-BP, which makes the right neighbour of a 1 also 1. When applying the correspondence a second time, we obtain standard oriented site percolation in two dimensions (up to a linear transformation of the lattice).

**Proposition 3.2.** *The correspondence of Proposition 3.1 extends to a one-to-one mapping from the set of all  $d$ -dimensional attractive PCA (death may be integrated directly into  $v$ ) to  $d+1$ -dimensional inhomogeneous BP with  $\mathbf{0}$  initial condition and update families measure  $\chi$  supported on the set of update families contained in the lower half-space (deaths corresponding to  $\emptyset \in \mathcal{X}_x$ ). Namely,  $\chi$  is the image of  $v$  via the mapping  $\mathcal{U} \mapsto \mathcal{X}$  of Eq. (12) and vice versa. Then the same conclusions still hold (Eq. (14) and it being a stationary  $v$ -PCA at its upper invariant measure).*

The proof is identical to the one of Proposition 3.1 and therefore omitted.



### 3.2 Proof of Theorems 1.5 and 1.6

Fix an BP  $\omega$  with update family  $\mathcal{X}$  contained in a half-space. Up to an invertible linear transformation of the lattice, we may assume that this is the lower half-space and denote by  $v = \delta_{\mathcal{U}}$  the corresponding attractive CA via Proposition 3.1. The proposition gives us that  $q_c$  in Eq. (7) for  $\mathcal{X}$  is in fact equal to  $1 - p_c$  with  $p_c$  from Eq. (6) for the  $\tilde{v}$ -PCA with death rate  $1 - p$ . Theorem 2.1 applies to the  $\tilde{v}$ -PCA, so Eq. (2) holds for  $p < p_c$ .

Applying Eq. (14) to the initial condition equal to  $\mathbb{1}_{\mathcal{U}_{x,t}=\emptyset}$  for  $t \geq 0$  and 0 otherwise and denoting  $C'$  the corresponding closure, we get that  $\tilde{\eta}_x^1(t) = 1$  iff  $(x, t) \notin C'$ . Hence, for  $p < p_c$  there are  $c, C > 0$  such that for all  $(x, t) \in \mathbb{Z}^{d+1}$

$$\mathbb{P}_{\tilde{v}}((x, t) \notin C') = \mathbb{P}_{\tilde{v}}(\tilde{\eta}_x^1(t) \neq 0) \leq Ce^{-ct}.$$

Denote by  $\omega'$  the  $\mathcal{X}$ -BP with the above initial condition. Then,  $(x, t) \notin C'$  iff  $\omega'_{(x,t)}(t) = 0$ , since, by induction on  $t$ , the  $\omega'$  process becomes stationary in  $\mathbb{Z}^d \times \{0, \dots, t\}$  after  $t$  steps. Finally, since  $\omega'_{(x,t)}(t) \leq \omega_{(x,t)}(t)$  by attractiveness, the proof of Theorem 1.5 is concluded. Theorem 1.6 is proved identically in view of Proposition 3.2.

### 3.3 Proof of Theorems 1.8 and 1.9

Fix an attractive CA  $v = \delta_{\mathcal{U}}$  and let  $\mathcal{X}$  be its corresponding BP update family via Proposition 3.1. Consider a configuration  $\xi$  with finitely many 0s. Applying Eq. (14) to this initial condition and its closure  $C$ , we get that if  $\mathcal{U}$  is not an eroder, then  $\mathcal{X}$  is supercritical. Inversely, if  $\mathcal{X}$  is supercritical, considering the intersection of the closure of a finite initial set with infinite offspring and a horizontal strip of width  $r$  (which is clearly finite), we similarly obtain from Eq. (14) that  $\mathcal{U}$  is not an eroder.

Furthermore, Proposition 3.1 grants that  $\mathcal{X}$  is subcritical iff  $\mathcal{U}$  has at least two invariant measures when the death rate  $1 - p$  is small enough. Thus, Theorem 1.8 is equivalent to Theorem 1.9 restricted to two-dimensions and, inversely Theorem 1.8 without restrictions on the dimension is equivalent to Theorem 1.9.

Hence, Theorem 1.9 follows from Theorem 1.8, which is valid in all dimensions [28, Theorem 5]. Turning to Theorem 1.8 in one dimension, in order to avoid a circular reasoning, we recall from Section 1.4 that by [3, 8] it suffices to verify the following fact, which we prove for completeness (see [28, Theorem 6] or [27, Theorem 3]).

**Lemma 3.3.** *Fix a  $d$ -dimensional update family  $\mathcal{X}$  contained in a half-space. Then for any  $u \in S^{d-1}$  the set of stable directions  $v \in S^{d-1}$  such that  $\langle u, v \rangle <$*

$0\}$  is either empty or has a nonempty interior. Moreover, if  $d = 2$  and there is no open semicircle of unstable directions, then there exist two opposite directions in the interior of the set of stable directions.

*Proof.* Let  $e_1, \dots, e_d$  denote the canonical basis of  $\mathbb{R}^{d+1}$ . For concreteness let us assume that the family is contained in the upper half-space  $\{x \in \mathbb{R}^d : \langle x, e_d \rangle > 0\}$ . It is not hard to see from the definition (see [8, Remark 3.3]) that the set of stable directions can be written as a finite intersection of finite unions of closed hemispheres containing the direction  $e_d$  in their interior.

Clearly, any closed hemisphere contains the geodesic between any of its points and any point in its interior. Therefore, for any stable direction the geodesics to a neighbourhood of  $e_d$  consist of stable directions. Hence, the set of stable directions is the closure of its own interior. The general conclusion then follows immediately.

In two dimensions, matters are simpler. We already established that stable directions form a connected closed set, that is, a closed interval of  $S^1$ . Depending on whether it is smaller or larger than a semicircle, this gives the desired conclusion.  $\square$

To conclude, let us mention that, given opposite stable directions provided in Lemma 3.3, the renormalisation argument sketched in [3] proceeds as follows. Divide the plane into large rhombi in the usual way, so that their sides are close to being perpendicular to these directions. Then perform renormalisation, saying that the rhombus is good if it is initially in state  $\mathbf{0}$  for the  $\mathcal{X}$ -BP, which happens with high probability if the parameter  $q$  is small. Then it suffices to verify that an infinite oriented path of good rhombi will remain in state  $\mathbf{0}$  forever, which follows from the suitable choice of their geometry.

## 4 Further directions

To conclude, let us mention a few directions for further work.

Firstly, in view of Theorems 1.1 and 1.2, one would naturally like to know what happens in the cooperative survival phase. We are aware of no results in this direction. For instance one may expect the following to be true.

**Question 1.** *For an absorbing attractive PCA in the interior of the cooperative survival phase does one have exponential convergence to the upper invariant measure starting from the  $\mathbf{1}$  initial condition? Equivalently, in the corresponding inhomogeneous BP, does the truncated infection time have an*

exponential tail, that is, setting  $\tau_0 = \inf\{t > 0 : \eta_0(t) = 1\}$ , do we have

$$\limsup_{t \rightarrow \infty} \frac{\log \mathbb{P}(\tau_0 > t | \tau_0 < \infty)}{t} < 0?$$

For models, such as the Toom rule with death, for which one can prove that they are in the cooperative survival phase, this follows from the corresponding proof, but we rather ask for non-perturbative results valid throughout the phase.

It would also be very interesting to obtain an analogue of Theorem 1.2 for PCA with a unique invariant measure not necessarily equal to  $\delta_0$ . See [20, 26] for progress in this direction under other conditions.

Furthermore, it is natural to seek to extend Theorems 1.2 and 2.1 to continuous time absorbing attractive interacting particle systems with single spin flips. In the case of the contact process this is a well-known result of Bezuidenhout and Grimmett [7].

Finally, in the light of the BP–PCA correspondence of Propositions 3.1 and 3.2, can one transfer more interesting information between the two settings?

## Acknowledgements

This work is supported by ERC Starting Grant 680275 “MALIG.” We thank Irène Marcovici, Jan Swart, Réka Szabo, Siamak Taati and Cristina Toninelli for enlightening discussions. We also thank Réka for bringing important references to our attention.

## References

- [1] M. Aizenman and D. J. Barsky, *Sharpness of the phase transition in percolation models*, Comm. Math. Phys. **108** (1987), no. 3, 489–526 pp. MR874906
- [2] M. Aizenman and G. Grimmett, *Strict monotonicity for critical points in percolation and ferromagnetic models*, J. Stat. Phys. **63** (1991), no. 5-6, 817–835 pp. MR1116036
- [3] P. Balister, B. Bollobás, M. Przykucki, and P. Smith, *Subcritical  $U$ -bootstrap percolation models have non-trivial phase transitions*, Trans. Amer. Math. Soc. **368** (2016), no. 10, 7385–7411 pp. MR3471095
- [4] P. Balister, B. Bollobás, and O. Riordan, *Essential enhancements revisited*, arXiv e-prints (2014), available at arXiv:1402.0834.
- [5] P. Berman and J. Simon, *Investigations of fault-tolerant networks of computers*, Proceedings of the Twentieth Annual ACM Symposium on Theory of Computing, 1988, 66–77 pp.

- [6] C. Bezuidenhout and L. Gray, *Critical attractive spin systems*, Ann. Probab. **22** (1994), no. 3, 1160–1194 pp. MR1303641
- [7] C. Bezuidenhout and G. Grimmett, *Exponential decay for subcritical contact and percolation processes*, Ann. Probab. **19** (1991), no. 3, 984–1009 pp. MR1112404
- [8] B. Bollobás, P. Smith, and A. Uzzell, *Monotone cellular automata in a random environment*, Combin. Probab. Comput. **24** (2015), no. 4, 687–722 pp. MR3350030
- [9] M. Bramson and L. Gray, *A useful renormalization argument*, Random walks, Brownian motion, and interacting particle systems, 1991, 113–152 pp. MR1146444
- [10] N. Cancrini, F. Martinelli, C. Roberto, and C. Toninelli, *Kinetically constrained spin models*, Probab. Theory Related Fields **140** (2008), no. 3-4, 459–504 pp. MR2365481
- [11] H. Duminil-Copin, A. Raoufi, and V. Tassion, *Sharp phase transition for the random-cluster and Potts models via decision trees*, Ann. of Math. (2) **189** (2019), no. 1, 75–99 pp. MR3898174
- [12] H. Duminil-Copin and V. Tassion, *A new proof of the sharpness of the phase transition for Bernoulli percolation on  $\mathbb{Z}^d$* , Enseign. Math. **62** (2016), no. 1, 199–206 pp. MR3605816
- [13] P. Gács, *A new version of Toom’s proof*, arXiv e-prints (2021), available at arXiv:2105.05968.
- [14] L. F. Gray, *Toom’s stability theorem in continuous time*, Perplexing problems in probability, 1999, 331–353 pp. MR1703139
- [15] G. Grimmett, *Percolation*, Second edition, Grundlehren der mathematischen Wissenschaften, Springer, Berlin, Heidelberg, 1999. Originally published by Springer, New York (1989). MR1707339
- [16] I. Hartarsky,  *$U$ -bootstrap percolation: critical probability, exponential decay and applications*, Ann. Inst. Henri Poincaré Probab. Stat. **57** (2021), no. 3, 1255–1280 pp. MR4291442
- [17] I. Hartarsky and R. Szabó, *Generalised oriented site percolation*, Markov Process. Related Fields (To appear).
- [18] J. L. Lebowitz, C. Maes, and E. R. Speer, *Statistical mechanics of probabilistic cellular automata*, J. Stat. Phys. **59** (1990), no. 1-2, 117–170 pp. MR1049965
- [19] T. M. Liggett, R. H. Schonmann, and A. M. Stacey, *Domination by product measures*, Ann. Probab. **25** (1997), no. 1, 71–95 pp. MR1428500
- [20] P.-Y. Louis, *Ergodicity of PCA: equivalence between spatial and temporal mixing conditions*, Electron. Commun. Probab. **9** (2004), 119–131 pp. MR2108858
- [21] I. Marcovici, M. Sablik, and S. Taati, *Ergodicity of some classes of cellular automata subject to noise*, Electron. J. Probab. **24** (2019), Paper No. 41, 44 pp. MR3940771
- [22] M. V. Menshikov, *Coincidence of critical points in percolation problems*, Dokl. Akad. Nauk SSSR **288** (1986), no. 6, 1308–1311 pp. MR852458
- [23] R. O’Donnell, M. Saks, O. Schramm, and R. A. Servedio, *Every decision tree has an influential variable*, 46th Annual IEEE Symposium on Foundations of Computer Science, 2005, 31–39 pp.

- [24] R. H. Schonmann, *On the behavior of some cellular automata related to bootstrap percolation*, Ann. Probab. **20** (1992), no. 1, 174–193 pp. MR1143417
- [25] J. M. Swart, R. Szabó, and C. Toninelli, *Peierls bounds from Toom contours* (In preparation).
- [26] S. Taati, *Reversible cellular automata in presence of noise rapidly forget everything*, 27th IFIP WG 1.5 international workshop on cellular automata and discrete complex systems (AUTOMATA 2021), 2021, 3:1–3:15 pp.
- [27] A. Toom, *Cellular automata with errors: problems for students of probability*, Topics in contemporary probability and its applications, 1995, 117–157 pp. MR1410537
- [28] A. L. Toom, *Stable and attractive trajectories in multicomponent systems*, Multicomponent random systems, 1980, 549–575 pp. MR599548