A comparison of two possible nuclear effective field theory expansions around the one- and two-pion exchange potentials

Manuel Pavon Valderrama^{1, *}

¹School of Physics, Beihang University, Beijing 100191, China (Dated: December 6, 2021)

In the effective field theory formalism nuclear forces are organized as a low energy expansion. Usually the lowest order in this expansion corresponds to the non-perturbative iteration of the one-pion exchange potential and a few contact-range operators as to ensure renormalization group invariance. Here we consider an alternative expansion in which two-pion exchange becomes the lowest order finite range interaction and as such is expected to be iterated. A comparison of this new expansion with the standard one shows a better convergence pattern for a few selected partial waves $({}^{1}S_{0}, {}^{3}P_{0} \text{ and } {}^{3}S_{1} {}^{-3}D_{1})$ in two-nucleon scattering, in particular for the ${}^{1}S_{0}$ channel (though both expansions converge well). We briefly comment on the possible theoretical grounds for the expansion around two-pion exchange from the point of view of effective field theory.

I. INTRODUCTION

The theoretical derivation of the nuclear forces is still one of the most pressing and interesting open problems of nuclear physics [1]. Nowadays we expect any serious attempt of such a derivation to be grounded on quantum chromodynamics (QCD), the fundamental theory of strong interactions. Two strategies exists: a direct derivation in terms of lattice QCD [2], in which QCD is solved by resorting to brute force calculations, and an indirect one in which an effective field theory (EFT) [3–5] is formulated incorporating the low energy symmetries (most notably chiral symmetry) and degrees of freedom (pions, nucleons and possibly deltas) of QCD. This second strategy is in principle equivalent to QCD by virtue of being its renormalization group evolution at low energies.

Within the EFT formulation, we organize nuclear forces as a low energy expansion in terms of a ratio of scales, Q/M, where Q is a soft scale that can be usually identified with the pion mass ($m_{\pi} \sim 140 \text{ MeV}$) or the typical momenta of the nucleons within nuclei and M a hard scale that might correspond to the nucleon mass M_N or the chiral symmetry breaking scale $\Lambda_{\chi} = 4\pi f_{\pi} \sim 1 \text{ GeV}$. However, there is no agreement yet on how to organize the EFT expansion for nuclear forces, as shown by the existence of numerous power counting proposals [6–13], the eventual discovery of theoretical inconsistencies in a few countings [14–16] and daring and original [17, 18] (but sometimes contested [19]) reinterpretations of nuclear EFT to deal with these problems.

This predicament might be explained by a poor expansion parameter (Q/M): if the expansion parameter is not small enough, the relative importance of the terms in the expansion could be very different than what naive estimations suggest. This might happen even if the coefficients in the expansion are of order one. To give a

concrete example, we might consider the expansion of the matrix elements of a physical observable $\hat{\mathcal{O}}$:

$$\langle \hat{\mathcal{O}} \rangle = \sum_{\nu = \nu_{\min}}^{\infty} c_{\nu} x^{\nu} , \qquad (1)$$

where c_{ν} are coefficients, x the expansion parameter and $\nu \geq \nu_{\min}$ the order in the expansion. Provided |x| < 1and that the c_{ν} do not grow exponentially with the order ν , the previous expansion will eventually converge. However, there is significant leeway for mishaps in the previous expansion, particularly if we are limited to its first few terms. This might happen even if the $c_{\nu} = \mathcal{O}(1)$. For instance, it might happen that $|c_0| = 1/3$ and $|c_2| = 3$. The conventional expectation for the convergence of the EFT expansion will be $|c_0 x^0| > |c_2 x^2|$, but in this specific example this will only happen for x < 1/3. That is, if the convergence parameter is not small enough, it will be as if the EFT expansion is not working (even though it is). at least at low orders. Actually, this by itself does not invalidate the idea of the EFT expansion or its eventual convergence but it will certainly interfere with its practical implementation, particularly if we are not willing to calculate arbitrarily high orders for checking whether there is convergence of not.

Actually, something similar is probably happening in the two-nucleon sector. If we expand the finite-range part of the effective potential derived from chiral perturbation theory (i.e. the pion exchanges), we have

$$V_{\rm NN}^{\rm EFT}(r>0) = \sum_{\nu=0}^{\infty} V^{(\nu)}(r)$$

= $\underbrace{V^{(0)}(r)}_{V_{\rm OPE}} + \underbrace{V^{(2)}(r) + V^{(3)}(r)}_{V_{\rm TPE}} + \dots (2)$

where the superscript indicates the order of a contribution in naive dimensional analysis (NDA) and below them we specify whether this contribution is one-pionexchange (OPE) or (mostly ¹) two-pion-exchange (TPE),

^{*} mpavon@buaa.edu.cn

¹ The $\nu \geq 2$ pieces also contain small corrections to the OPE

which can be further subdivided into leading ($\nu = 2$) and subleading ($\nu = 3$) TPE, or TPE(L) and TPE(SL) for short. Naively, we expect the behavior of the Q^{ν} contributions to be

$$V^{(\nu)}(r) \sim \frac{4\pi}{M_N} \frac{1}{\left[\Lambda_{\rm NN}^{(\nu)} r\right]^{\nu+1}} \frac{1}{r^2} f^{(\nu)}(m_\pi r), \qquad (3)$$

where the characteristic scale $\Lambda_{\rm NN}^{(\nu)}$ is expected to be a hard scale, i.e. $\Lambda_{\rm NN}^{(\nu)} \sim M$ and with $f^{(\nu)}$ a function (such that $\lim_{x\to 0} x f^{(\nu)}(x) = 0$) that captures the features of the order ν contribution to the potential beyond its leading power-law behavior.

But, as it often happens, reality does not match expectations. If we consider the chiral limit $(m_{\pi} \rightarrow 0)$ for simplicity, the ¹S₀ and ³S₁ partial wave potentials can be expanded as follows

$$\lim_{m_{\pi}\to 0} V_{\rm NN}^{\rm EFT}(r) = -\frac{4\pi}{M_N} \left[\frac{1}{\Lambda_{\rm TPE(L)}^3} \frac{1}{r^5} + \frac{1}{\Lambda_{\rm TPE(SL)}^4} \frac{1}{r^6} \right] + \mathcal{O}\left(\left(\frac{Q}{M}\right)^4 \right), \tag{4}$$

where in this limit the OPE contribution vanishes for Swaves and the $1/r^5$ and $1/r^6$ contributions correspond to the $\nu = 2$ and 3 terms in the EFT expansion of the potential, i.e. TPE(L) and TPE(SL). From direct inspection of the potentials [20], we find that for the $c_3 = -3.4 \text{ GeV}$ and $c_3 = +3.4 \text{ GeV}$ choice of chiral couplings the actual numerical values of the leading and subleading TPE scales are

$$\Lambda_{\text{TPE(L)}}({}^{1}S_{0}) = +389 \,\text{MeV}\,,$$
 (5)

$$\Lambda_{\rm TPE(SL)}({}^{1}S_0) = +233\,{\rm MeV}\,,\tag{6}$$

$$\Lambda_{\rm TPE(L)}(^3S_1) = -370\,{\rm MeV}\,,\tag{7}$$

$$\Lambda_{\text{TPE(SL)}}(^{3}S_{1}) = +220 \,\text{MeV}\,,\tag{8}$$

which for the case of subleading TPE significantly differs from the NDA expectation of $\Lambda_{\text{TPE}} \sim (0.5 - 1.0) \text{ GeV}$.

Of course, the previous observation does not necessarily imply that the TPE potential has to be iterated (except maybe in the chiral limit). For physical pion masses, the TPE potential is suppressed by a $e^{-2m_{\pi}r}$ factor at large distances, which for sure attenuates its strength. Besides, if tensor OPE is present (as happens in the triplets), the TPE potential will be further attenuated owing to the distortions generated by the nonperturbative tensor OPE wave function. Be it as it may, it will be interesting to explore what happens when TPE is non-perturbative, particularly in comparison to what happen when it is a perturbation.

Actually, this idea was already explored in the past [16, 21], though usually not in the context of building a serious EFT expansion around it (except in [22]). In the Weinberg counting, scattering amplitudes are obtained from iterating the full effective potential with a finite cutoff. As has been often pointed out, this prescription does in general not generate renormalizable scattering amplitudes [12, 14], which thus might very well break the power counting assumptions of the effective potential from which they are calculated. Indeed, from analyzing the underlying power counting of the scattering amplitudes calculated in the Weinberg prescription by trial and error, it is easy to find that the Weinberg ${}^{1}S_{0}$ phase shifts are well reproduced if TPE (plus a contact operator to guarantee renormalizability) is LO and everything else is subleading (and perturbative) [16, 21]. Recently Ref. [22] has consider this expansion as a possible organizing principle for the ${}^{1}S_{0}$ singlet, leading to results which we agree with here.

For exploring the expansion around TPE, we will do as follows: first, we will consider the type of non-relativistic EFT expansion that is generated from an arbitrary choice of a leading order potential, which we will then particularize for the cases in which OPE and TPE are leading (Sect. II); after this, we will calculate the EFT expansion of the phase shifts for the two cases (Sect. III); and finally, we will discuss and compare these two EFTs, the possible justifications of the TPE expansions and its implications (Sect. IV).

II. POWER COUNTING AS A FUNCTION OF THE CHOICE OF A LEADING ORDER

Now we explore the possible form of the power countings generated from expanding around the cases in which OPE and TPE are non-perturbative. For concreteness we will refer to them as EFT(OPE) and EFT(TPE). Actually, EFT(OPE) has already been extensively discussed in the literature [23–28] and we will rely on preexisting analyses up to a certain extent. The expansion of interest here is EFT(TPE), which will require modifications to the power counting that are genuinely different from those of EFT(OPE).

For simplicity, we will consider a general EFT expansion which at LO requires the non-perturbative iteration of a piece of the effective potential, which in turns generates a LO wave function, $\Psi_{\rm LO}$, the form of which we assume to be known. The reason for adopting this approach is that the power counting of the contact-range potential is indeed determined by the power-law properties of $\Psi_{\rm LO}$ [29]. Then, by particularizing $\Psi_{\rm LO}$ to the EFT(OPE) and EFT(TPE) expansions we will deduce their power countings.

Before beginning, there are a few conventions we follow: the counting of the non-relativistic potential will be defined with respect to its scaling in momentum space,

potential, but for simplicity we will ignore them here.

e.g. a potential that has the form

$$V(\vec{q}) \propto \frac{1}{M^{2+n}Q^m} P_n(\vec{q}) f(\frac{\vec{q}}{Q}), \qquad (9)$$

where \vec{q} is the exchanged momentum, P_n a polynomial of order n and $f(\vec{x})$ some other function, will be counted as Q^{n-m} . When considering the iterations of the potential, $VG_0V \ldots G_0V$, the propagator G_0 between two instances of the potential is counted as Q. As a consequence, if a part of the potential is treated non-perturbatively, this is because we consider it to scale as Q^{-1} or LO. Also, the notation Q^{ν} is actually short-hand for $(Q/M)^{\nu}$.

We will assume the following power-law behavior for the LO wave function

$$\langle r|\Psi_{\rm LO}\rangle \sim r^{\alpha-1}$$
, (10)

where the exponent α can be calculated from the form of the LO potential. Following [29] we will consider the matrix elements of a contact-range interaction of the type

$$V_C(r) = \frac{C(R_c)}{4\pi R_c^2} \,\delta(r - R_c)\,, \tag{11}$$

which for concreteness we have regularized with a deltashell, where R_c is a cutoff radius. This contact will be sandwiched between the LO wave functions, leading to

$$\langle \Psi_{\rm LO} | V_C | \Psi_{\rm LO} \rangle \sim \frac{C(R_c)}{R_c^{2-2\alpha}} \,.$$
 (12)

If we demand renormalization group (RG) invariance for this matrix element

$$\frac{d}{dR_c} \left[\frac{C(R_c)}{R_c^{2-2\alpha}} \right] = 0 , \qquad (13)$$

then we realize that the running of $C(R_c)$ is given by

$$C(R_c) \propto R_c^{2\alpha - 2} \,. \tag{14}$$

If we evolve $C(R_c)$ from $MR_c \sim 1$ (where NDA applies and $C(R_c) \sim 1/M^2$ [30]) to $QR_c \sim 1$, we find that at low resolutions this coupling behaves as

$$C(R_c \sim 1/Q) \sim \frac{1}{M^2} \left(\frac{M}{Q}\right)^{2-2\alpha}.$$
 (15)

That is, if $\alpha > 1$ ($\alpha < 1$) then $C(R_c)$ will be demoted (promoted) to order $Q^{2\alpha-2}$. If we further expand the coupling in energy/momentum

$$C = C_0 + C_2 k^2 + C_4 k^4 + \dots, \qquad (16)$$

it is apparent that C_2, C_4, \ldots are further suppressed by factor of Q^2, Q^4, \ldots and scale as

$$C_2 \propto Q^{2\alpha}$$
, $C_4 \propto Q^{2\alpha+2}$ and so on. (17)

As a side note the previous argument can be crosschecked in two different ways: finiteness of the subleading order calculations [24, 25] and residual cutoff dependence [31–33]. If we consider the perturbative correction of a hypothetical, unaccounted for Q^{ν} contribution to the chiral potential in EFT, the matrix element will behave as [30]

$$\langle \Psi_{\rm LO} | V^{(\nu)} | \Psi_{\rm LO} \rangle \sim \int_{R_c}^{\infty} \frac{dr}{r^{3+\nu}} r^{2\alpha} \left(1 + k^2 r^2 + \dots \right),$$
(18)

where $V^{(\nu)}$ refers to a finite-range contribution of order ν to the EFT potential. From this, the lowest order at which we encounter a divergence in EFT is $Q^{2\alpha-2}$ (which can be absorbed by a recalibration of the scattering length), while at $Q^{2\alpha}$ we will find a divergence that requires a range correction. From the point of view of residual cutoff dependence, the LO phase shift converges as [34]

$$\frac{d}{dR_c}\delta_{\rm LO}(k;R_c) \propto k^3 R_c^{2\alpha+1} \,, \tag{19}$$

for $R_c \to 0$, which suggest that the range corrections enter $Q^{2\alpha+1}$ orders after LO (Q^{-1}) .

At this point it has to be stressed that two of the previous arguments (Eqs. (12) and (18)) are perturbative in nature and do not take into account that the LO wave function might already require a series of contactinteraction for its unambiguous determination. Indeed, this is what happens when the LO wave function is an attractive singular interaction [35, 36]. If this is the case, the necessary contact-range interactions will be automatically promoted to LO or Q^{-1} .

Now, all that remains is to determine the exponent α , which is a relatively well-known quantity:

- (i) For a LO regular potential, we have $\alpha = 0$.
- (ii) For a LO power-law singular potential of the type $1/r^n$ with $n \ge 2$, we have $\alpha = n/4$.

With this, the power counting of EFT(OPE) can be reproduced by taking into account that in the ${}^{1}S_{0}$ singlet $\alpha = 0$, while for the triplets in which tensor OPE is nonperturbative (e.g. ${}^{3}S_{1}$ - ${}^{3}D_{1}$ and ${}^{3}P_{0}$), $\alpha = 3/4$.

In contrast for EFT(TPE) we have $\alpha = 3/2$ for singlets and triplets alike. This implies in particular that the C_2 couplings, i.e. range corrections, are all demoted to Q^3 or N⁴LO. This demotion of the range corrections is probably the most characteristic feature of the power countings arising from expanding around singular interactions.

In Table I we briefly summarize the power counting of EFT(OPE) and EFT(TPE) as will be implemented and explored in this work. It includes a few simplifying assumptions, e.g. the fractional counting of C_2 with tensor OPE (i.e. $Q^{3/2}$ from $\alpha = 3/4$) has been approximated to Q^2 and a few couplings that are numerically small (the couplings that fix the E_1 and 3D_1 scattering lengths) have been demoted from $Q^{-1/2}$ to Q^2 .

The most important simplification in EFT(TPE) is that we have simply promoted the full $\nu \leq 3$ EFT potential to LO (instead of only promoting subleading TPE, which makes more sense in view of $\Lambda_{\text{TPE(SL)}}$, check

Contribution	NDA	EFT(OPE)	EFT(TPE)
V_{OPE}	Q^0	Q^{-1}	Q^{-1}
$V_{\rm TPE(L)}$	Q^2 Q^3	Q^2	Q^{-1}
$V_{\rm TPE(SL)}$	Q^3	Q^3	Q^{-1}
$C_0({}^1S_0)$	Q^0	Q^{-1}	Q^{-1}
$C_2({}^1S_0)$	Q^2	Q^0	Q^3
$C_4({}^1S_0)$	Q^4	Q^2	Q^5
$C_0({}^3S_1)$	Q^2	Q^{-1}	Q^{-1}
$C_0 \left(E_1 / {}^3 D_1 \right)$	Q^2	Q^2	Q^{-1}
$C_2 \left({}^3S_1 / E_1 / {}^3D_1 \right)$	Q^4	Q^2	Q^3
$C_0 ({}^3P_0)$	Q^0	Q^{-1}	Q^{-1}
$C_2 \left({}^3P_0\right)$	Q^2	Q^2	Q^3

TABLE I. Power counting for the different contributions to the effective potential in the EFT(OPE) and EFT(TPE) expansions as defined in this work and their comparison with NDA estimations. The EFT(OPE) counting has been extensively studied in Refs. [23–28, 30], where there exists differences in the details of how the different operators are counted (esteeming from the diverging assumptions made in each of the previous references). Here, while EFT(TPE) counting is purely driven by RG invariance, a few of the choices we have made in EFT(OPE) are practical in nature: in the ${}^{3}S_{1}$ - ${}^{3}D_{1}$ channel the couplings fixing the scattering volumes and hypervolumes in the E_{1} and ${}^{3}D_{1}$ partial waves ($C_{0}(E_{1}/{}^{3}D_{1}$ in the Table) is demoted to order Q^{2} , even though it should enter at order $Q^{-1/2}$ if we follow RGA [23]. For the singlet we follow Ref. [30].

Eqs. (5-8)). This might be justified from the observation that the contributions of OPE and leading TPE are numerically small. Yet, the actual reason why it is safe to promote them is that this choice does not change the RG evolution (RGE) of EFT(TPE): the power-law behavior of OPE and TPE(L) is unable to change the RGE of a theory in which TPE(SL) is treated non-perturbative. Besides, perturbative OPE is finite when sandwiched between the LO wave functions of EFT(TPE), while TPE(L) is renormalized by simply recalibrating the scattering length, i.e. by a subleading correction to C_0 .

III. COMPARISON BETWEEN THE TWO EXPANSIONS

Having decided the two power countings, the only thing left is to make the necessary calculations in the EFT(OPE) and EFT(TPE) expansions. The technical details are straightforward but tedious, where here we simply indicate the main features of the calculation:

(i) The LO wave functions and scattering amplitudes are obtained and renormalized as in Refs. [35–37], in which by using suitable boundary conditions it was shown the minimum number of couplings required for the different partial waves. In particular for the coupled channels (e.g. ${}^{3}S_{1} - {}^{3}D_{1}$) we have that

- (i.a) Non-perturbative OPE can be renormalized by fixing one of the scattering lengths (or their L-wave equivalents) [37].
- (i.b) Non-perturbative TPE can be renormalized by fixing the three scattering lengths (provided that the two eigenvalues of the coupled channel potentials are attractive, which is what happens in most partial waves) [35, 36].
- (ii) For the subleading order phase shifts, we basically follow the regularization used in Refs. [24, 25], which also shows that the perturbative corrections are indeed renormalizable after the inclusion of a series of couplings. A few differences with respect to Refs. [24, 25] are worth commenting:
 - (ii.a) For the ${}^{1}S_{0}$ partial wave in EFT(OPE), Refs. [24, 25] included both the C_{2} and C_{4} couplings at Q^{2} , which is the minimal requirement to render perturbative TPE finite and well-defined in the $R_{c} \rightarrow 0$ limit. However, finiteness is merely a subset of renormalizability and indeed RG invariance requires the C_{2} coupling to enter at order Q^{0} , which is what we do here, a choice that entails a series of iterations of the C_{2} coupling (which is computationally fastidious and has prompted expansions in which C_{2} is fully iterated at LO [38]). This is further justified by the fact that the LO calculation of the ${}^{1}S_{0}$ effective range only reproduces 50% of its value.
 - (ii.b) For the ${}^{3}S_{1}{}^{-3}D_{1}$ channel in EFT(OPE), Refs. [24, 25] use a total of six couplings for renormalizing TPE. However, Long and Yang proved that three couplings are enough [27] (just as happens in Weinberg's counting). We nonetheless will use six here, as this provides a more direct comparison between EFT(OPE) and EFT(TPE) at N⁴LO (as in both cases we will end up with six couplings).
 - (ii.c) This choice of six couplings generates a further complication for ${}^{3}S_{1}$ - ${}^{3}D_{1}$ in EFT(OPE): RGE arguments (either Eq. (13) particularized for $\alpha = 3/4$ or Ref. [23] for its original formulation) indicate that the couplings fixing the scattering lengths in the E_1 and 3D_1 channels enter at order $Q^{-1/2}$ or N^{1/2}LO. This is problematic as it will require an inordinate number of iterations of these couplings (far exceeding the required iterations of C_2 in the singlet). Yet, it is worth noticing that non-perturbative OPE actually reproduces relatively well these scattering lengths at LO [37] (in contrast with the ${}^{1}S_{0}$ case, in which this does not happen for the effective range), indicating that in most practical settings these couplings are not really required. For this reason we will simply demote the $C_0(E_1/{}^3S_1)$ couplings to Q^2 .

- (ii.d) In principle a similar problem appears in the ${}^{3}P_{0}$ partial wave, but in this case the scattering length can be reproduced at LO which automatically implies that the subleading corrections to $C_{0}({}^{3}P_{0})$ are trivial.
- (ii.e) Finally, in EFT(OPE) we further assume that the promotion of OPE from its naive scaling (Q^0) to LO does not affect the power counting of the TPE potential. This is in contrast to the approach of Long and Yang which also promotes the TPE potential by one order [26– 28], a choice with makes perfect sense in terms of scales once we look at Eqs. (5-8)).

Having explained the previous details, we might simply proceed to the calculation of the ${}^{1}S_{0}$, ${}^{3}P_{0}$ and ${}^{3}S_{1}$ - ${}^{3}D_{1}$ phase shifts in EFT(OPE) and EFT(TPE). For this, regardless of the expansion, we will calibrate the LO couplings as to reproduce the following values of the scattering length: $\alpha_0({}^1S_0) = -23.7 \,\text{fm}, \ \alpha_0({}^3P_0) = -2.65 \,\text{fm}^4,$ $\alpha_0({}^3S_1) = 5.42 \,\mathrm{fm}, \ \alpha_0(E_1) = 1.67, \mathrm{fm}^3 \ \mathrm{and} \ \alpha_0({}^3D_1) =$ 6.60 fm^5 , where the E_1 and 3D_1 scattering lengths are only required for EFT(TPE) and are defined for the Stapp-Ypsilantis-Metropolis (SYM, also known as nuclear bar) parametrization of the phase shifts [39] (where their detailed low energy behavior can be consulted in Ref. [40]). For the subleading order phase shifts, the couplings will be determined from fitting the Nijmegen II phase shifts [41] (expected to be equivalent to the ones extracted in the Nijmegen PWA [42] within errors) within the k = (100 - 200) MeV center-of-mass momentum window, except for the singlet channel where the momentum window changes with the order and includes lower momenta to ensure the correct scattering length (in particular we use k = (10 - 40) MeV, (10 - 80) MeV, (10-200) MeV and (10-200) MeV at LO, NLO, N²LO, $N^{3}LO$ and $N^{4}LO$). The explicit expressions for the effective OPE and TPE potential is taken from Ref. [20], where for the couplings $(g_A, f_{\pi}, m_{\pi}, d_{18}, c_1, c_3 \text{ and } c_4)$ we follow the same choices as in Ref. [25] and the recoil or $1/M_N$ corrections are included.

The results are shown in Fig. 1 for the ${}^{1}S_{0}$, ${}^{3}P_{0}$ and ${}^{3}S_{1}$ - ${}^{3}D_{1}$ partial waves, which uses the cutoff range $R_{c} = (0.5 - 1.0)$ fm to generate bands. The two expansions work relatively well for most partial waves, though EFT(TPE) works slightly better (particularly at higher momenta). Yet, the advantage of EFT(TPE) seems to concentrate mostly in the ${}^{1}S_{0}$ partial wave, for which it requires only two couplings (instead of four) at N⁴LO, with which it also displays a slightly better agreement with the Nijmegen II pseudodata. For ${}^{3}S_{1}$ - ${}^{3}D_{1}$ we have chosen a counting choice in EFT(OPE) with more parameters as strictly necessary, just to make calculations simpler. Thus the comparison made here does not take into account that EFT(OPE) can be improved as to require less couplings in deuteron channel.

IV. DISCUSSION

Here we have considered two possible EFT expansions of the nuclear force: a typical EFT expansion in which the LO is defined by the iteration of the OPE potential and an atypical one in which TPE is also iterated. In both cases contact-range operators are included as to guarantee RG invariance at each order, though a few liberties are taken to ease the computational burden (particularly in the ${}^{3}S_{1}$ - ${}^{3}D_{1}$ channel). The two expansions, which we have named EFT(OPE) and EFT(TPE), converge relatively well, though the atypical EFT(TPE) does a better job in the ${}^{1}S_{0}$ partial wave.

This begs the question: why is this the case? EFT(TPE) is in a sense a parody of what power counting should be, where contributions that should enter at Q^3 are promoted to Q^{-1} . In the absence of a good physical reason, EFT(TPE) should not be considered as a legitimate EFT expansion. The most apparent rationale of why EFT(TPE) might make sense lies in the slow convergence of the EFT expansion for nuclear physics, i.e. the poor separation of scales, which makes it plausible that higher order contributions might accidentally behave as lower order ones (check the discussion around Eq. (1) for further details). A cursory look at the S-wave potentials in the chiral limit, Eqs. (4-8), indicates that parts of TPE are indeed unexpectedly large.

In fact in the chiral limit tensor OPE, TPE(L) and TPE(SL) behave as pure $1/r^3$, $1/r^5$ and $1/r^6$ powerlaw infinite-range potentials (while spin-spin OPE vanishes). This was beautifully exploited by Birse [23] to determine the momenta below which tensor OPE can be treated perturbatively by applying previously known results from atomic physics (in particular, the failure of the secular perturbative expansion [43] for the $1/r^3$ potential, as calculated by Gao [44]). This suggested that tensor OPE is only perturbative for $k < 66 \,\mathrm{MeV}$ $(k < 182 \,\mathrm{MeV})$ in the ${}^{3}S_{1}$ - ${}^{3}D_{1}$ (${}^{3}P_{0}$) partial waves, at least in the chiral limit, explaining the previous observation that the Kaplan, Savage and Wise (KSW) counting [8, 9] (which treats all pion exchanges as perturbations) converges slowly and only for low momenta in the two-nucleon system [15] (check also Ref. [45] for a further confirmation of the limits of perturbative OPE).

It happens that Gao also analyzed the secular perturbative expansion of attractive $1/r^6$ interactions in Ref. [46], from which repeating Birse's arguments one obtains a critical momentum $k_{\rm crit} \simeq (150-200)$ MeV above which the perturbative treatment of subleading TPE will not converge in the chiral limit for the singlet. Taking into account that in the chiral limit the OPE vanishes in the singlet, it is sensible to assume that subleading TPE requires a non-perturbative treatment when $m_{\pi} \rightarrow 0$. In the real world significant deviations should be expected due to the finite pion mass effects and the OPE LO distortion. Indeed, the EFT(OPE) expansion shows good convergence properties up to N⁴LO, with no evident signs of a failure for $k > k_{\rm crit}$. However, the same was also true for the KSW expansion at NLO in the triplets, which later were shown to fail once OPE was iterated for the first time at N²LO [15]. Thus it might very well happen that the same fate awaits to the EFT(OPE) expansion once the first iteration of TPE(SL) enters at Q^7 / N⁸LO.

For triplet waves the situation is different, as TPE is distorted by tensor OPE. From this, it might be perfectly possible that its strength is screened by the longer-range OPE distortion, which is why EFT(OPE) works better in the triples when compared to the singlet. Actually, the results we show indicate that there is not a marked difference between the two expansions for the triplets. It should also be stressed that for EFT(TPE) we have made the simplification of promoting the full potential (including OPE) to LO. It might happen that leaving OPE as a NLO contribution, as its NDA scaling would suggest, might not lead to a converging expansion. If this were to be the case, the conclusion would be that EFT(TPE) is not a suitable expansion for the triplets. From an orthodox EFT perspective this would be good news, as this will reinforce the standard counting in which OPE drives the low energy physics of the deuteron, for instance. But to really confirm this hypothesis we would need to iterate tensor OPE up to relatively high orders in distorted wave perturbation theory, which is not exactly easy to do (particularly if we want to guarantee renormalizability at every step in the calculations). This would be nonetheless worth exploring in the future.

To summarize, the features of EFT(TPE) are intriguing: it is a really counterintuitive way of organizing nuclear EFT, yet it provides an expansion that converges

- R. Machleidt, Int. J. Mod. Phys. E26, 1730005 (2017), arXiv:1710.07215 [nucl-th].
- [2] S. R. Beane, W. Detmold, K. Orginos, and M. J. Savage, Prog. Part. Nucl. Phys. 66, 1 (2011), arXiv:1004.2935 [hep-lat].
- [3] P. F. Bedaque and U. van Kolck, Ann. Rev. Nucl. Part. Sci. 52, 339 (2002), nucl-th/0203055.
- [4] E. Epelbaum, H.-W. Hammer, and U.-G. Meissner, Rev. Mod. Phys. 81, 1773 (2009), arXiv:0811.1338 [nucl-th].
- [5] R. Machleidt and D. Entem, Phys.Rept. 503, 1 (2011), arXiv:1105.2919 [nucl-th].
- [6] S. Weinberg, Phys. Lett. **B251**, 288 (1990).
- [7] S. Weinberg, Nucl. Phys. B363, 3 (1991).
- [8] D. B. Kaplan, M. J. Savage, and M. B. Wise, Phys. Lett. B424, 390 (1998), nucl-th/9801034.
- [9] D. B. Kaplan, M. J. Savage, and M. B. Wise, Nucl. Phys. B534, 329 (1998), nucl-th/9802075.
- [10] J.-W. Chen, G. Rupak, and M. J. Savage, Nucl. Phys. A 653, 386 (1999), arXiv:nucl-th/9902056.
- [11] S. R. Beane, P. F. Bedaque, M. J. Savage, and U. van Kolck, Nucl. Phys. A700, 377 (2002), nucl-th/0104030.
- [12] A. Nogga, R. G. E. Timmermans, and U. van Kolck, Phys. Rev. C72, 054006 (2005), nucl-th/0506005.
- [13] S. R. Beane, D. B. Kaplan, and A. Vuorinen, Phys. Rev. C80, 011001 (2009), arXiv:0812.3938 [nucl-th].

- [14] D. B. Kaplan, M. J. Savage, and M. B. Wise, Nucl. Phys. B478, 629 (1996), nucl-th/9605002.
- [15] S. Fleming, T. Mehen, and I. W. Stewart, Nucl. Phys. A677, 313 (2000), nucl-th/9911001.
- [16] M. Pavon Valderrama, (2019), arXiv:1902.08172 [nucl-th].
- [17] E. Epelbaum and J. Gegelia, Eur. Phys. J. A41, 341 (2009), arXiv:0906.3822 [nucl-th].
- [18] E. Epelbaum, A. M. Gasparyan, J. Gegelia, and U.-G. Meißner, Eur. Phys. J. A54, 186 (2018), arXiv:1810.02646 [nucl-th].
- [19] M. Pavon Valderrama, Eur. Phys. J. A 55, 55 (2019), arXiv:1901.10398 [nucl-th].
- [20] M. C. M. Rentmeester, R. G. E. Timmermans, J. L. Friar, and J. J. de Swart, Phys. Rev. Lett. 82, 4992 (1999), nucl-th/9901054.
- [21] M. P. Valderrama, AIP Conf. Proc. 1322, 205, arXiv:1009.6100 [nucl-th].
- [22] C. Mishra, A. Ekström, G. Hagen, T. Papenbrock, and L. Platter, (2021), arXiv:2111.15515 [nucl-th].
- [23] M. C. Birse, Phys. Rev. C74, 014003 (2006), arXiv:nucl-th/0507077.
- [24] M. Valderrama, Phys.Rev. C83, 024003 (2011), arXiv:0912.0699 [nucl-th].
- [25] M. Pavon Valderrama, Phys.Rev. C84, 064002 (2011), arXiv:1108.0872 [nucl-th].

6

better than the standard way of organizing the EFT expansion. One might argue that EFT(TPE) is an ersatz Weinberg counting, but it is conceptually different: even though in both cases we are iterating a large chunk of the effective finite-range potential, this is not true for the contact-range operators, most of which are treated as perturbations. This detail makes EFT(TPE) renormalizable — the counting of the contacts is derived from RGE and the amplitudes have a well-defined $R_c \rightarrow 0$ limit — while the Weinberg counting does not always behave well in the hard cutoff limit (e.g. the ${}^{1}S_{0}$ channel is not renormalizable with TPE and two contacts, C_0 and C_2 [47], unless a very specific representation of the contact-range interaction is invoked [34]). If anything, the EFT(TPE) expansion provides more questions than answers and will require further theoretical effort to find its place within nuclear EFT.

ACKNOWLEDGMENTS

I would like to thank Ubirajara van Kolck for estimulating discussions on the topic of this manuscript and the IJCLab of Orsay, where part of this work was done, for its hospitality. This work is partly supported by the National Natural Science Foundation of China under Grants No. 11735003 and No. 11975041, the Fundamental Research Funds for the Central Universities and the Thousand Talents Plan for Young Professionals.

- [26] B. Long and C. Yang, Phys.Rev. C84, 057001 (2011), published version, arXiv:1108.0985 [nucl-th].
- [27] B. Long and C. J. Yang, Phys. Rev. C85, 034002 (2012), arXiv:1111.3993 [nucl-th].
- [28] B. Long and C. J. Yang, Phys. Rev. C86, 024001 (2012), arXiv:1202.4053 [nucl-th].
- [29] M. Pavón Valderrama and D. R. Phillips, Phys. Rev. Lett. **114**, 082502 (2015), arXiv:1407.0437 [nucl-th].
- [30] M. P. Valderrama, Int. J. Mod. Phys. E25, 1641007 (2016), arXiv:1604.01332 [nucl-th].
- [31] H. W. Grießhammer, Proceedings, 8th International Workshop on Chiral Dynamics (CD15): Pisa, Italy, June 29-July 3, 2015, PoS CD15, 104 (2016), arXiv:1511.00490 [nucl-th].
- [32] H. W. Griesshammer, Eur. Phys. J. A 56, 118 (2020), arXiv:2004.00411 [nucl-th].
- [33] D. Odell, M. P. Valderrama, and L. Platter, (2021), arXiv:2107.14250 [nucl-th].
- [34] M. Pavon Valderrama and E. R. Arriola, Annals Phys. 323, 1037 (2008), arXiv:0705.2952 [nucl-th].
- [35] M. Pavon Valderrama and E. R. Arriola, Phys. Rev. C74, 054001 (2006), arXiv:nucl-th/0506047.

- [36] M. Pavon Valderrama and E. Ruiz Arriola, Phys. Rev. C74, 064004 (2006), arXiv:nucl-th/0507075.
- [37] M. Pavon Valderrama and E. Ruiz Arriola, Phys. Rev. C72, 054002 (2005), nucl-th/0504067.
- [38] R. Peng, S. Lyu, S. König, and B. Long, (2021), arXiv:2112.00947 [nucl-th].
- [39] H. P. Stapp, T. J. Ypsilantis, and N. Metropolis, Phys. Rev. 105, 302 (1957).
- [40] M. P. Valderrama and E. R. Arriola, Phys. Rev. C72, 044007 (2005).
- [41] V. G. J. Stoks, R. A. M. Klomp, C. P. F. Terheggen, and J. J. de Swart, Phys. Rev. C49, 2950 (1994), nucl-th/9406039.
- [42] V. G. J. Stoks, R. A. M. Kompl, M. C. M. Rentmeester, and J. J. de Swart, Phys. Rev. C48, 792 (1993).
- [43] M. J. Cavagnero, Phys. Rev. A 50, 2841 (1994).
- [44] B. Gao, Phys. Rev. A59, 2778 (1999).
- [45] D. B. Kaplan, Phys. Rev. C 102, 034004 (2020), arXiv:1905.07485 [nucl-th].
- [46] B. Gao, Phys. Rev. A58, 1728 (1998).
- [47] D. R. Entem, E. Ruiz Arriola, M. Pavon Valderrama, and R. Machleidt, Phys. Rev. C77, 044006 (2008), arXiv:0709.2770 [nucl-th].




FIG. 1. Phase shifts of the ${}^{1}S_{0}$, ${}^{3}P_{0}$ and ${}^{3}S_{1}$ - ${}^{3}D_{1}$ (nuclear bar) partial waves in the two EFT expansions considered in this work, EFT(OPE) and EFT(TPE). EFT(OPE) is an expansion in which OPE is a non-perturbative LO effect and happens to be similar to the power counting described in Refs. [24, 25], except the ${}^{1}S_{0}$ partial wave in which EFT(OPE) includes all the necessary iterations of C_{2} , which enters at order Q^{0} in said expansion. EFT(TPE) is an expansion in which TPE is treated non-perturbatively, comprising the LO in the expansion (though for simplicity we also include OPE as part of the LO calculation as this will not significantly change the results). We follow the power counting described in Table I, where only the non-trivial orders (i.e. orders at which a new correction is included) are shown. The bands correspond to varying the cutoff in the $R_{c} = (0.5 - 1.0)$ fm range.