

PROJECTIONS ONTO HYPERBOLAS OR BILINEAR CONSTRAINT SETS IN HILBERT SPACES

Heinz H. Bauschke*, Manish Krishan Lal† and Xianfu Wang‡

December 1, 2021

Abstract

Sets of bilinear constraints are important in various machine learning models. Mathematically, they are hyperbolas in a product space. In this paper, we give a complete formula for projections onto sets of bilinear constraints or hyperbolas in a general Hilbert space.

2020 Mathematics Subject Classification: Primary 41A50, 90C26; Secondary 90C20, 46C05.

Keywords: bilinear constraint set, hyperbola, orthogonal projection, nonconvex minimization, prox-regularity.

1 Introduction

Throughout this paper, we assume that

$$X \text{ is a Hilbert space with inner product } \langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{R}, \quad (1)$$

*Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: heinz.bauschke@ubc.ca.

†Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: manish.krishanlal@ubc.ca.

‡Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: shawn.wang@ubc.ca.

and induced norm $\|\cdot\|$. In various learning models [9, 10], Elser utilizes projections onto the *bilinear constraint* set:

$$C_\gamma := \{(x, y) \in X \times X \mid \langle x, y \rangle = \gamma\} \quad (2)$$

where $\gamma \in \mathbb{R}$ is a fixed constant. Mathematically speaking, up to a rotation, this is just a hyperbola or quadratic surface in $X \times X$. Finding projections onto quadratic curves or surfaces algorithmically have many practical applications; see, e.g., [6, 7, 9, 10]. If $\gamma = 0$ in C_γ , then C_0 becomes the set ‘cross’ in $X \times X$. The projection formula for the ‘cross’ has been thoroughly investigated [2]. In [9, 10], although Elser has provided some results on projections onto C_γ only when $X = \mathbb{R}^n$, complete mathematical details are not presented.

The goal of this paper is to give a complete analysis of the projection onto the set C_γ with $\gamma \neq 0$, and extend the results to a general Hilbert space.

The remainder of the paper is organized as follows. Section 2 gives some general properties of bilinear constraint sets. In Section 3 we focus on full mathematical details for the existence and explicit formula of projections onto hyperbolas. Using results from Section 3, in Sections 4–5 we provide explicit formulas for projections onto sets of bilinear constraints. Our notation is standard and follows largely [1, 13].

2 General properties of C_γ

In an infinite-dimensional space, the existence of projection onto a set often requires the weak closedness of the set. Our first result says that although the set C_γ is norm closed it is not weakly closed in $X \times X$.

Proposition 2.1. The set C_γ is closed in the norm topology but not closed in the weak topology in $X \times X$. In fact, $\overline{C_\gamma}^{\text{weak}} = X \times X$.

Proof. Evidently, C_γ is norm closed. We show that $\overline{C_\gamma}^{\text{weak}} = X \times X$. Let $(x, y) \in X \times X$. Consider $S = \text{span}\{x, y\}$. The orthogonal decomposition theorem gives $X = S \oplus S^\perp$. Because X is infinite-dimensional and S is at most two-dimensional, S^\perp is infinite dimensional, so any orthonormal base of S^\perp must have a sequence $(e_n)_{n \in \mathbb{N}}$ which converges weakly to 0, i.e., $e_n \rightharpoonup 0$; see, e.g., [11]. Also $e_n \perp x$ and $e_n \perp y$. Set $\xi = \gamma - \langle x, y \rangle$. Then

$$\langle x - \xi e_n, y - e_n \rangle = \langle x, y \rangle - \langle x, e_n \rangle - \xi \langle e_n, y \rangle + \xi \langle e_n, e_n \rangle \quad (3)$$

$$= \langle x, y \rangle + \xi = \gamma, \quad (4)$$

so $(x - \xi e_n, y - e_n) \in C_\gamma$. Since $(x - \xi e_n, y - e_n) \rightharpoonup (x, y)$, we have $(x, y) \in \overline{C_\gamma}^{\text{weak}}$. Because $(x, y) \in X \times X$ was arbitrary, we conclude that $X \times X \subseteq \overline{C_\gamma}^{\text{weak}}$. ■

Proposition 2.1 indicates that finding P_{C_γ} , i.e., projections onto C_γ , might be complicated in a general Hilbert space. This seemingly difficult issue can be completely avoided by utilizing the structures of the optimization problem. See Subsection 3.1 below.

Our next result says that when $\gamma \neq 0$, locally around the set C_γ the projection onto the set is always single-valued.

Proposition 2.2. Let $\gamma \neq 0$, and $C_\gamma = \{(x, y) \in X \times X \mid \langle x, y \rangle = \gamma\}$. For every $(x, y) \in C_\gamma$, the the following hold:

- (i) C_γ is prox-regular at (x, y) .
- (ii) There exists a neighborhood of (x, y) on which the projection mapping onto C_γ is monotone and Lipschitz continuous.

Proof. (i): Write $\iota_{C_\gamma}(x, y) = \iota_{\{\gamma\}}(h(x, y))$ where $h(x, y) = \langle x, y \rangle$. Let $(x, y) \in C_\gamma$. Then $\nabla h(x, y) = (y, x) \neq (0, 0)$ because of $\gamma \neq 0$. Being a composition of a convex function $\iota_{\{\gamma\}}$ and a twice differentiable function h that is qualified at (x, y) , [3, Proposition 2.4] shows that ι_{C_γ} is prox-regular at (x, y) , so is C_γ .

(ii): Apply (i) and [3, Proposition 4.4]. ■

Observe that Proposition 2.2 also follow from [13, Proposition 13.32] and [13, Exercise 13.38] when $X = \mathbb{R}^n$; and that when $\gamma = 0$, C_0 is not prox-regular at $(0, 0)$. Although the projection exists locally around C_γ , it is still not clear for the global existence.

3 Projections onto hyperbolas

For ease of analysis, we start with

$$C_\gamma := \{(x, y) \in X \times X \mid h(x, y) := \langle x, y \rangle - \gamma = 0\}$$

where $\gamma > 0$. Our goal is to find the projection formula $P_{C_\gamma}(x_0, y_0)$ for every $(x_0, y_0) \in X \times X$. That is,

$$\text{minimize } f(x, y) := \|x - x_0\|^2 + \|y - y_0\|^2 \quad \text{subject to } (x, y) \in C_\gamma. \quad (\text{P})$$

3.1 Auxiliary problems and existence of projections

To determine the projection operator P_{C_γ} of the set C_γ , we shall introduce two equivalently reformulated problems. First, for every $(u_0, v_0) \in X \times X$, we solve the problem

$$\text{minimize } f_1(u, v) := \|u - u_0\|^2 + \|v - v_0\|^2 \quad \text{subject to } (u, v) \in \tilde{C}_1, \quad (\tilde{\text{P}}_1)$$

where $\tilde{C}_1 := \{(u, v) \in X \times X \mid h_1(u, v) := \|u\|^2 - \|v\|^2 - 2 = 0\}$. Next, for every $(\tilde{u}_0, \tilde{v}_0) \in X \times X$ we solve the problem

$$\text{minimize } f_\gamma(\tilde{u}, \tilde{v}) := \|\tilde{u} - \tilde{u}_0\|^2 + \|\tilde{v} - \tilde{v}_0\|^2 \quad \text{subject to } (\tilde{u}, \tilde{v}) \in \tilde{C}_\gamma, \quad (\tilde{P}_\gamma)$$

where $\tilde{C}_\gamma := \{(\tilde{u}, \tilde{v}) \in X \times X \mid h_\gamma(\tilde{u}, \tilde{v}) := \|\tilde{u}\|^2 - \|\tilde{v}\|^2 - 2\gamma = 0\}$. Both \tilde{C}_1 and \tilde{C}_γ are hyperbolas. \tilde{P}_1 and \tilde{P}_γ solve for projections $P_{\tilde{C}_1}$ and $P_{\tilde{C}_\gamma}$ respectively, and their connections to P_{C_γ} are given by the [Proposition 3.2](#) below. Recall

Definition 3.1 (Rotation with an angle ϕ). A change of coordinates $(u, v) \in X \times X$, by rotation through an angle ϕ , is defined by

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \cos \phi \text{Id} & -\sin \phi \text{Id} \\ \sin \phi \text{Id} & \cos \phi \text{Id} \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix}.$$

Put

$$A_\phi := \begin{bmatrix} \cos \phi \text{Id} & -\sin \phi \text{Id} \\ \sin \phi \text{Id} & \cos \phi \text{Id} \end{bmatrix}.$$

Then $A_\phi A_{-\phi} = \text{Id} = A_{-\phi} A_\phi$ and $A_\phi^{-1} = A_{-\phi}$. The relationships among P_{C_γ} , $P_{\tilde{C}_\gamma}$ and $P_{\tilde{C}_1}$ are summarized below.

Proposition 3.2. The following hold:

- (i) $P_{\tilde{C}_\gamma} = \sqrt{\gamma} P_{\tilde{C}_1} (\text{Id} / \sqrt{\gamma})$.
- (ii) $P_{C_\gamma} = A_{\pi/4} P_{\tilde{C}_\gamma} A_{-\pi/4}$.

Proof. (i): (\tilde{P}_1) is equivalent to (\tilde{P}_γ) by a change of variables of scaling $\text{Id} / \sqrt{\gamma}$. Indeed, using

$$\begin{bmatrix} u \\ v \end{bmatrix} = \frac{1}{\sqrt{\gamma}} \begin{bmatrix} \tilde{u} \\ \tilde{v} \end{bmatrix}, \text{ and } \begin{bmatrix} u_0 \\ v_0 \end{bmatrix} = \frac{1}{\sqrt{\gamma}} \begin{bmatrix} \tilde{u}_0 \\ \tilde{v}_0 \end{bmatrix}$$

$h_\gamma(\tilde{u}, \tilde{v}) = 0$ becomes $h_1(u, v) = 0$, and $f_\gamma(\tilde{u}, \tilde{v})$ becomes $\gamma f_1(u, v) = \gamma(\|u - u_0\|^2 + \|v - v_0\|^2)$.

(ii): (P) is equivalent to (\tilde{P}_γ) by a change of variables of rotation $A_{\pi/4}$. Indeed, with

$$\begin{bmatrix} x \\ y \end{bmatrix} = A_{\pi/4} \begin{bmatrix} \tilde{u} \\ \tilde{v} \end{bmatrix}, \text{ and } \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} = A_{\pi/4} \begin{bmatrix} \tilde{u}_0 \\ \tilde{v}_0 \end{bmatrix}$$

the objective $f(x, y) = \|x - x_0\|^2 + \|y - y_0\|^2 = \|(x, y) - (x_0, y_0)\|^2$ can be rewritten as

$$\begin{aligned} f(x, y) &= \left\| A_{\frac{\pi}{4}} \begin{bmatrix} \tilde{u} - \tilde{u}_0 \\ \tilde{v} - \tilde{v}_0 \end{bmatrix} \right\|^2 = \begin{bmatrix} \tilde{u} - \tilde{u}_0 \\ \tilde{v} - \tilde{v}_0 \end{bmatrix}^T A_{\frac{\pi}{4}}^T A_{\frac{\pi}{4}} \begin{bmatrix} \tilde{u} - \tilde{u}_0 \\ \tilde{v} - \tilde{v}_0 \end{bmatrix} \\ &= \|\tilde{u} - \tilde{u}_0\|^2 + \|\tilde{v} - \tilde{v}_0\|^2, \end{aligned}$$

and $h(x, y) = \langle x, y \rangle - \gamma = 0$ becomes $h_\gamma(\tilde{u}, \tilde{v}) = \|\tilde{u}\|^2 - \|\tilde{v}\|^2 - 2\gamma = 0$. \blacksquare

In view of Proposition 3.2(ii), to see that $P_{C_\gamma}(x, y) \neq \emptyset$ for every $(x, y) \in X \times X$, the following observation is crucial.

Proposition 3.3. For every $(\tilde{u}_0, \tilde{v}_0) \in X \times X$, the minimization problem

$$\text{minimize} \quad f_\gamma(\tilde{u}, \tilde{v}) = \|\tilde{u} - \tilde{u}_0\|^2 + \|\tilde{v} - \tilde{v}_0\|^2 \quad (5)$$

$$\text{subject to} \quad h_\gamma(\tilde{u}, \tilde{v}) = \|\tilde{u}\|^2 - \|\tilde{v}\|^2 - 2\gamma = 0 \quad (6)$$

always has a solution, i.e., $P_{\tilde{C}_\gamma}(\tilde{u}_0, \tilde{v}_0) \neq \emptyset$.

Proof. We shall illustrate only the case $\tilde{u}_0 \neq 0, \tilde{v}_0 \neq 0$, since the other cases are similar. We claim that the optimization problem is essentially 2-dimensional. To this end, we expand

$$f_\gamma(\tilde{u}, \tilde{v}) = \underbrace{\|\tilde{u}\|^2 - 2\langle \tilde{u}, \tilde{u}_0 \rangle + \|\tilde{u}_0\|^2}_{\|\tilde{u} - \tilde{u}_0\|^2} + \underbrace{\|\tilde{v}\|^2 - 2\langle \tilde{v}, \tilde{v}_0 \rangle + \|\tilde{v}_0\|^2}_{\|\tilde{v} - \tilde{v}_0\|^2}. \quad (7)$$

The constraint

$$h_\gamma(\tilde{u}, \tilde{v}) := \|\tilde{u}\|^2 - \|\tilde{v}\|^2 - 2\gamma = 0$$

means that only the norms $\|\tilde{u}\|$ and $\|\tilde{v}\|$ matter. With $\|\tilde{u}\|$ fixed, the Cauchy-Schwarz inequality in a Hilbert space, see, e.g., [11], shows that $\tilde{u} \mapsto \langle \tilde{u}, \tilde{u}_0 \rangle$ will be larger so that the first underlined part in f_γ will be smaller when \tilde{u} and \tilde{u}_0 are positively co-linear, i.e., $\tilde{u} = \alpha \tilde{u}_0$ for some $\alpha \geq 0$. Similarly, for fixed $\|\tilde{v}\|$ the second underlined part in f_γ will be smaller when $\tilde{v} = \beta \tilde{v}_0$ for some $\beta \geq 0$. It follows that the optimization problem given by (5)-(6) is equivalent to

$$\text{minimize} \quad g(\alpha, \beta) := (1 - \alpha)^2 \|\tilde{u}_0\|^2 + (1 - \beta)^2 \|\tilde{v}_0\|^2 \quad (8)$$

$$\text{subject to} \quad g_1(\alpha, \beta) := \alpha^2 \|\tilde{u}_0\|^2 - \beta^2 \|\tilde{v}_0\|^2 - 2\gamma = 0, \alpha \geq 0, \beta \geq 0. \quad (9)$$

Because $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuous and coercive, and $g_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuous, we conclude that the optimization problem given by (8)-(9) has a solution. \blacksquare

We are now ready for the investigation of projections onto $\tilde{C}_1, \tilde{C}_\gamma$ and C_γ .

3.2 Finding the projection $P_{\tilde{C}_1}$

Note that

$$\nabla f_1(u, v) = (2(u - u_0), 2(v - v_0)) \quad \text{and} \quad \nabla h_1(u, v) = (2u, -2v). \quad (10)$$

By [4, Proposition 4.1.1], every solution of (\tilde{P}_1) satisfies the necessarily conditions, i.e., the KKT system of (\tilde{P}_1) , given by

$$(1 + \lambda)u = u_0 \quad (11)$$

$$(1 - \lambda)v = v_0 \quad (12)$$

$$\|u\|^2 - \|v\|^2 - 2 = 0. \quad (13)$$

Lemma 3.4. Let $u_0 \neq 0, v_0 \neq 0$. If (u, v, λ) verifies (11)–(13) and (u, v) is an optimal solution of problem (\tilde{P}_1) , then the Lagrange multiplier λ satisfies $|\lambda| < 1$.

Proof. The constraint set

$$\|u\|^2 - \|v\|^2 = 2,$$

has a special structure: replacing u by $-u$, or v by $-v$ the constraint is still verified. Also consider

$$f(u, v) = \|u - u_0\|^2 + \|v - v_0\|^2 = \|u_0\|^2 - 2\langle u_0, u \rangle + \|u\|^2 + \|v_0\|^2 - 2\langle v_0, v \rangle + \|v\|^2,$$

Given u_0 and v_0 , for fixed $\|u\|$ and $\|v\|$, $f(u, v)$ becomes smaller if one choose $\langle u_0, u \rangle \geq 0$, and $\langle v_0, v \rangle \geq 0$. Indeed, one can do so by replacing u by $-u$ or v by $-v$ if needed. Now by (11) and (12),

$$(1 + \lambda)\langle u_0, u \rangle = \|u_0\|^2, \text{ and } (1 - \lambda)\langle v_0, v \rangle = \|v_0\|^2.$$

Because $u_0 \neq 0, v_0 \neq 0$, we have $\langle u_0, u \rangle > 0, \langle v_0, v \rangle > 0$ so that $1 + \lambda > 0, 1 - \lambda > 0$. Hence $|\lambda| < 1$. ■

Proposition 3.5. Let $u_0 \neq 0, v_0 \neq 0$. Define $p := \|u_0\|^2 - \|v_0\|^2$ and $q := \|u_0\|^2 + \|v_0\|^2$. Suppose that $u, v \in X$ and $\lambda \in \mathbb{R}$ verify (11)–(13) and (u, v) is an optimal solution to (\tilde{P}_1) . Then the following hold:

- (i) $u = \frac{u_0}{(1+\lambda)}, v = \frac{v_0}{(1-\lambda)}$, and $(1 - \lambda)^2\|u_0\|^2 - (1 + \lambda)^2\|v_0\|^2 = 2(1 - \lambda^2)^2$.
- (ii) The objective function has $f_1(u, v) = \lambda^2 \left(\frac{\|u_0\|^2}{(1+\lambda)^2} + \frac{\|v_0\|^2}{(1-\lambda)^2} \right)$.
- (iii) λ is the unique solution of

$$H(\lambda) := \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1 - \lambda^2)^2} - 1 = 0 \quad (14)$$

in $] -1, 1[$.

Proof. (i): Because $u_0 \neq 0, v_0 \neq 0$, we obtain $|\lambda| < 1$, $u = \frac{u_0}{(1+\lambda)}$ and $v = \frac{v_0}{(1-\lambda)}$ by Lemma 3.4 and (11)-(12). Then we have the following equivalences

$$\|u\|^2 - \|v\|^2 = 2 \Leftrightarrow \frac{\|u_0\|^2}{(1+\lambda)^2} - \frac{\|v_0\|^2}{(1-\lambda)^2} = 2 \quad (15)$$

$$\Leftrightarrow (1-\lambda)^2\|u_0\|^2 - (1+\lambda)^2\|v_0\|^2 = 2(1-\lambda)^2(1+\lambda)^2 \quad (16)$$

$$\Leftrightarrow (1-\lambda)^2\|u_0\|^2 - (1+\lambda)^2\|v_0\|^2 = 2(1-\lambda^2)^2. \quad (17)$$

(ii): Substitute $u = \frac{u_0}{(1+\lambda)}$ and $v = \frac{v_0}{(1-\lambda)}$ in f_1 .

(iii): By (i), we have

$$(1-\lambda)^2\|u_0\|^2 - (1+\lambda)^2\|v_0\|^2 = 2(1-\lambda^2)^2. \quad (18)$$

Since

$$(1-\lambda)^2\|u_0\|^2 - (1+\lambda)^2\|v_0\|^2 \quad (19)$$

$$= (1+\lambda^2-2\lambda)\|u_0\|^2 - (1+\lambda^2+2\lambda)\|v_0\|^2 \quad (20)$$

$$= \lambda^2(\|u_0\|^2 - \|v_0\|^2) + (\|u_0\|^2 - \|v_0\|^2) - 2\lambda(\|u_0\|^2 + \|v_0\|^2) \quad (21)$$

$$= \lambda^2 p + p - 2\lambda q, \quad (22)$$

using (22) on left side of (18), we obtain $\lambda^2 p + p - 2\lambda q = 2(1-\lambda^2)^2$, a univariate quartic equation in λ , equivalently,

$$H(\lambda) := \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1-\lambda^2)^2} - 1 = 0. \quad (23)$$

We show that (23) has a unique solution in $] -1, 1[$. Because $u_0 \neq 0, v_0 \neq 0$, we know that $q > |p|$, and using it along-with $|\lambda| < 1$ from Lemma 3.4, we get $p\lambda \leq |p||\lambda| \leq q|\lambda|$, and

$$H'(\lambda) = \frac{1}{(1-\lambda^2)^3} (-q(1+3\lambda^2) + p(\lambda^3+3\lambda)) \quad (24)$$

$$\leq \frac{1}{(1-\lambda^2)^3} (-q(1+3\lambda^2) + |p||\lambda|(\lambda^2+3)) \quad (25)$$

$$\leq \frac{1}{(1-\lambda^2)^3} (-q(1+3\lambda^2) + q|\lambda|(\lambda^2+3)) \quad (26)$$

$$= \frac{q}{(1-\lambda^2)^3} ((|\lambda|-1)^3) \quad (27)$$

$$= \frac{-q}{((1-|\lambda|)(1+|\lambda|))^3} ((1-|\lambda|)^3) \quad (28)$$

$$= \frac{-q}{(1+|\lambda|)^3}. \quad (29)$$

Since $q = \|u_0\|^2 + \|v_0\|^2 > 0$ and $(1 + |\lambda|)^3 > 0$, we have $H'(\lambda) < 0$ on $] -1, 1[$, which implies $H(\lambda)$ is strictly decreasing on $] -1, 1[$. Notice that $\lambda = \pm 1$ are vertical asymptotes because at $\lambda = \pm 1$ the numerator of $H(\lambda)$ is, $H(-1) = 2(p + q) = 4\|u_0\|^2 > 0$ and $H(1) = 2(p - q) = -4\|v_0\|^2 < 0$, so that $\lim_{\lambda \rightarrow 1^-} H(\lambda) = -\infty$ and $\lim_{\lambda \rightarrow -1^+} H(\lambda) = +\infty$. Since H is continuous, strictly decreasing, and its range is $] -\infty, \infty[$, we conclude that $H(\lambda) = 0$ has a unique zero in $] -1, 1[$. ■

Remark 3.6. As indicated in [9], an approximate solution to (14) can be found by the Bisection method, Newton's method, or a combined version. See also [5, 8].

Theorem 3.7. Let $u_0, v_0 \in X$, $u_0 \neq 0, v_0 \neq 0$. Then $P_{\tilde{C}_1}(u_0, v_0)$ is a singleton, and

$$P_{\tilde{C}_1}(u_0, v_0) = \left\{ \begin{bmatrix} \frac{u_0}{1+\lambda} \\ \frac{v_0}{1-\lambda} \end{bmatrix} \right\}$$

in which λ is the unique root of $H(\lambda) = 0$ in $] -1, 1[$ where

$$H(\lambda) = \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1 - \lambda^2)^2} - 1, \text{ and} \\ p = \|u_0\|^2 - \|v_0\|^2, \quad q = \|u_0\|^2 + \|v_0\|^2.$$

Proof. Apply Proposition 3.5. ■

Theorem 3.8. Let $u_0, v_0 \in X$ with either $u_0 = 0$ or $v_0 = 0$.

(i) When $u_0 = 0$, we have

$$P_{\tilde{C}_1}(0, v_0) = \left\{ \begin{bmatrix} u \\ \frac{v_0}{2} \end{bmatrix} \mid \|u\|^2 = 2 + \frac{\|v_0\|^2}{4} \right\}. \quad (30)$$

(ii) a) When $v_0 = 0$ and $\|u_0\| \geq 2\sqrt{2}$, we have

$$P_{\tilde{C}_1}(u_0, 0) = \left\{ \begin{bmatrix} \frac{u_0}{2} \\ v \end{bmatrix} \mid \|v\|^2 = \frac{\|u_0\|^2}{4} - 2 \right\}. \quad (31)$$

b) When $v_0 = 0$ and $0 < \|u_0\| < 2\sqrt{2}$, we have

$$P_{\tilde{C}_1}(u_0, 0) = \left\{ \begin{bmatrix} \sqrt{2} \frac{u_0}{\|u_0\|} \\ 0 \end{bmatrix} \right\}. \quad (32)$$

Proof. By (11)–(13), $(u, v) \in P_{\tilde{C}_1}(u_0, v_0)$ satisfies:

$$(1 + \lambda)u = u_0 \quad (33)$$

$$(1 - \lambda)v = v_0 \quad (34)$$

$$\|u\|^2 - \|v\|^2 = 2, \quad (35)$$

for some $\lambda \in \mathbb{R}$.

(i): $u_0 = 0, v_0 \neq 0$. (33) yields $(1 + \lambda)u = 0 \Rightarrow$ either $1 + \lambda = 0$ or $u = 0$. We consider two cases.

Case 1: $1 + \lambda = 0$, i.e., $\lambda = -1$. By (34), $2v = v_0 \Rightarrow v = \frac{v_0}{2}$ and then by (35), $\|u\|^2 - \|\frac{v_0}{2}\|^2 = 2 \Rightarrow \|u\|^2 = 2 + \frac{\|v_0\|^2}{4}$. So, $(u, v) = (u, \frac{v_0}{2})$ and $\|u\|^2 = 2 + \frac{\|v_0\|^2}{4}$. The objective function is

$$f(u, v) = \|u - u_0\|^2 + \|v - v_0\|^2 = \|u\|^2 + \left\|\frac{v_0}{2}\right\|^2 = 2 + \frac{\|v_0\|^2}{4} + \frac{\|v_0\|^2}{4} = 2 + \frac{\|v_0\|^2}{2}.$$

Case 2: $u = 0$. By (35), $(0 - \|v\|^2) = 2$, which is impossible. Combining both cases we have the formula (30).

(ii): $u_0 \neq 0$, but $v_0 = 0$, which implies $(1 - \lambda)v = 0$ by (34). We consider two cases:

Case 1: $1 - \lambda = 0$, i.e., $\lambda = 1$. By (33), $2u = u_0 \Rightarrow u = \frac{u_0}{2}$ and then by (35), $\|v\|^2 = \frac{\|u_0\|^2}{4} - 2 \Rightarrow \frac{\|u_0\|^2}{4} - 2 \geq 0 \Rightarrow \|u_0\|^2 \geq 8$. Thus, $(u, v) = (\frac{u_0}{2}, v)$ with $\|v\|^2 = \frac{\|u_0\|^2}{4} - 2$, where the objective is

$$f(u, v) = \|u - u_0\|^2 + \|v - v_0\|^2 = \frac{\|u_0\|^2}{4} + \frac{\|u_0\|^2}{4} - 2 = \frac{\|u_0\|^2}{2} - 2.$$

Note that Case 1 needs $\|u_0\| \geq 2\sqrt{2}$.

Case 2: $v = 0$. Then $\|u\|^2 = 2 \Rightarrow \|u\| = \sqrt{2}$ by (35). By (33), $|1 + \lambda|\|u\| = \|u_0\| \Rightarrow |1 + \lambda| = \frac{\|u_0\|}{\sqrt{2}} \Rightarrow 1 + \lambda = \pm \frac{\|u_0\|}{\sqrt{2}}$.

Subcase 1: $1 + \lambda = -\frac{\|u_0\|}{\sqrt{2}}$. By (33), $u = \frac{u_0}{(-\frac{\|u_0\|}{\sqrt{2}})} = -\frac{\sqrt{2}u_0}{\|u_0\|}$. Then

$$f(u, 0) = \|u - u_0\|^2 + \|0 - v_0\|^2 = \left\|-\frac{\sqrt{2}u_0}{\|u_0\|} - u_0\right\|^2 = (\|u_0\| + \sqrt{2})^2.$$

Subcase 2: $1 + \lambda = \frac{\|u_0\|}{\sqrt{2}}$. By (33), $u = \frac{\sqrt{2}u_0}{\|u_0\|}$. Then

$$f(u, 0) = \|u - u_0\|^2 + \|0 - v_0\|^2 = \left\|u_0 - \frac{\sqrt{2}u_0}{\|u_0\|}\right\|^2 = (\|u_0\| - \sqrt{2})^2.$$

Comparing Subcase 1 and Subcase 2, we obtain $|\|u_0\| - \sqrt{2}| < \|u_0\| + \sqrt{2}$ because $\|u_0\| \neq 0$. That is, Subcase 2 gives a smaller value at $(u, 0)$ with $u = \sqrt{2} \frac{u_0}{\|u_0\|}$. We need to compare it to Case 1 whenever it happens. Note that

$$\frac{\|u_0\|^2}{2} - 2 < (\|u_0\| - \sqrt{2})^2 \quad \text{whenever} \quad \|u_0\| \neq 2\sqrt{2}.$$

Indeed, we have

$$\frac{\|u_0\|^2}{2} - 2 < \|u_0\|^2 - 2\sqrt{2}\|u_0\| + 2 \iff \frac{\|u_0\|^2}{2} - 2\sqrt{2}\|u_0\| + 4 > 0 \iff (\|u_0\| - 2\sqrt{2})^2 > 0,$$

which always holds if $\|u_0\| \neq 2\sqrt{2}$. Hence, the nearest points are given by

$$\left(\frac{u_0}{2}, v\right) \quad \text{with} \quad \|v\|^2 = \frac{\|u_0\|^2}{4} - 2, \quad \text{when} \quad \|u_0\| \neq 2\sqrt{2} \quad (36)$$

$$\left(\frac{u_0}{2}, 0\right) = \left(\sqrt{2} \frac{u_0}{\|u_0\|}, 0\right), \quad \text{when} \quad \|u_0\| = 2\sqrt{2}. \quad (37)$$

However, Case 1 occurs only when $\|u_0\| \geq 2\sqrt{2}$. Hence, when $\|u_0\| \geq 2\sqrt{2}$ the nearest points are given by

$$\left(\frac{u_0}{2}, v\right) \quad \text{with} \quad \|v\|^2 = \frac{\|u_0\|^2}{4} - 2.$$

When $\|u_0\| < 2\sqrt{2}$, Case 1 is impossible. Then we only need to compare Subcase 1 and Subcase 2. Hence, the nearest point is

$$\left(\sqrt{2} \frac{u_0}{\|u_0\|}, 0\right) \quad \text{when} \quad \|u_0\| < 2\sqrt{2}.$$

Finally, since $\|u_0\| = 2\sqrt{2} \Rightarrow \|v\|^2 = 2 - 2 = 0$, (36) gives $(\frac{u_0}{2}, 0)$. Also $\frac{\sqrt{2}}{\|u_0\|} = \frac{1}{2}$, therefore $\sqrt{2} \frac{u_0}{\|u_0\|} = \frac{1}{2}u_0$. It follows that $(\frac{u_0}{2}, 0) = (\sqrt{2} \frac{u_0}{\|u_0\|}, 0)$ when $\|u_0\| = 2\sqrt{2}$, which implies that (37) can be obtained from (36) when $\|u_0\| = 2\sqrt{2}$. Hence formulas (31) and (32) hold. ■

3.3 Finding the projection $P_{\tilde{C}_\gamma}$

$P_{\tilde{C}_\gamma}$ can be found via $P_{\tilde{C}_1}$.

Theorem 3.9. Let $\gamma > 0, \tilde{u}_0, \tilde{v}_0 \in X$, and $\tilde{C}_\gamma = \{(\tilde{u}, \tilde{v}) \in X \times X \mid \|\tilde{u}\|^2 - \|\tilde{v}\|^2 = 2\gamma\}$. Then the following hold:

(i) When $\tilde{u}_0 \neq 0$ and $\tilde{v}_0 \neq 0$, we have

$$P_{\tilde{C}_\gamma}(\tilde{u}_0, \tilde{v}_0) = \left\{ \begin{bmatrix} \frac{\tilde{u}_0}{1+\lambda} \\ \frac{\tilde{v}_0}{1-\lambda} \end{bmatrix} \right\},$$

in which λ is the unique root of $H(\lambda) = 0$ in $] -1, 1[$, where

$$H(\lambda) = \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1 - \lambda^2)^2} - \gamma, \quad p = \|\tilde{u}_0\|^2 - \|\tilde{v}_0\|^2, \quad q = \|\tilde{u}_0\|^2 + \|\tilde{v}_0\|^2.$$

(ii) When $\tilde{u}_0 = 0$, we have

$$P_{\tilde{C}_\gamma}(0, \tilde{v}_0) = \left\{ \begin{bmatrix} \tilde{u} \\ \frac{\tilde{v}_0}{2} \end{bmatrix} \mid \|\tilde{u}\|^2 = 2\gamma + \frac{\|\tilde{v}_0\|^2}{4} \right\}. \quad (38)$$

(iii) a) When $\tilde{v}_0 = 0$ and $\|\tilde{u}_0\| \geq 2\sqrt{2\gamma}$, we have

$$P_{\tilde{C}_\gamma}(\tilde{u}_0, 0) = \left\{ \begin{bmatrix} \frac{\tilde{u}_0}{2} \\ \tilde{v} \end{bmatrix} \mid \|\tilde{v}\|^2 = \frac{\|\tilde{u}_0\|^2}{4} - 2\gamma \right\}. \quad (39)$$

b) When $\tilde{v}_0 = 0$ and $0 < \|\tilde{u}_0\| < 2\sqrt{2\gamma}$, we have

$$P_{\tilde{C}_\gamma}(\tilde{u}_0, 0) = \left\{ \begin{bmatrix} \sqrt{2\gamma} \frac{\tilde{u}_0}{\|\tilde{u}_0\|} \\ 0 \end{bmatrix} \right\}. \quad (40)$$

Proof. Proposition 3.2(i) states

$$P_{\tilde{C}_\gamma}(\tilde{u}_0, \tilde{v}_0) = \sqrt{\gamma} P_{\tilde{C}_1} \left(\frac{\tilde{u}_0}{\sqrt{\gamma}}, \frac{\tilde{v}_0}{\sqrt{\gamma}} \right).$$

Apply Theorem 3.7 and Theorem 3.8. ■

4 Projections onto bilinear constraint set C_γ when $\gamma > 0$

P_{C_γ} can be found via $P_{\tilde{C}_\gamma}$, which is the main result of this section.

Theorem 4.1. Let $\gamma > 0$, $x_0, y_0 \in X$, and $C_\gamma = \{(x, y) \in X \times X \mid \langle x, y \rangle = \gamma\}$. Then the following hold:

(i) When $x_0 \neq \pm y_0$, the projection is a singleton:

$$P_{C_\gamma}(x_0, y_0) = \left\{ \begin{bmatrix} \frac{x_0 - \lambda y_0}{1 - \lambda^2} \\ \frac{y_0 - \lambda x_0}{1 - \lambda^2} \end{bmatrix} \right\},$$

in which λ is the unique solution of $H(\lambda) = 0$ in $] -1, 1[$, where

$$H(\lambda) = \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1 - \lambda^2)^2} - \gamma, \quad p = 2\langle x_0, y_0 \rangle, \quad \text{and } q = \|x_0\|^2 + \|y_0\|^2.$$

(ii) When $x_0 = -y_0$, the projection is a set:

$$P_{C_\gamma}(x_0, -x_0) = \left\{ \begin{bmatrix} \frac{x_0}{2} + \frac{\tilde{u}}{\sqrt{2}} \\ -\frac{x_0}{2} + \frac{\tilde{u}}{\sqrt{2}} \end{bmatrix} \mid \|\tilde{u}\|^2 = 2\gamma + \frac{\|x_0\|^2}{2}, \quad \tilde{u} \in X \right\}.$$

(iii) a) When $x_0 = y_0$ and $\|x_0\| \geq 2\sqrt{\gamma}$, the projection is a set:

$$P_{C_\gamma}(x_0, x_0) = \left\{ \begin{bmatrix} \frac{x_0}{2} - \frac{\tilde{v}}{\sqrt{2}} \\ \frac{x_0}{2} + \frac{\tilde{v}}{\sqrt{2}} \end{bmatrix} \mid \|\tilde{v}\|^2 = \frac{\|x_0\|^2}{2} - 2\gamma, \quad \tilde{v} \in X \right\}.$$

b) When $x_0 = y_0$ and $0 < \|x_0\| < 2\sqrt{\gamma}$, the projection is a singleton:

$$P_{C_\gamma}(x_0, x_0) = \left\{ \sqrt{\gamma} \begin{bmatrix} \frac{x_0}{\|x_0\|} \\ \frac{x_0}{\|x_0\|} \end{bmatrix} \right\}.$$

Proof. Proposition 3.2(ii) states

$$P_{C_\gamma}(x_0, y_0) = A_{\frac{\pi}{4}} P_{\tilde{C}_\gamma} A_{-\frac{\pi}{4}}(x_0, y_0). \quad (41)$$

It suffices to apply Theorem 3.9. Indeed, with

$$\begin{bmatrix} \tilde{u}_0 \\ \tilde{v}_0 \end{bmatrix} = A_{-\pi/4} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} = \begin{bmatrix} \frac{x_0 + y_0}{\sqrt{2}} \\ \frac{-x_0 + y_0}{\sqrt{2}} \end{bmatrix}, \quad (42)$$

using (41) and (42) Theorem 3.9 gives:

(i): When $x_0 \neq \pm y_0$, we have $\tilde{u}_0 \neq 0, \tilde{v}_0 \neq 0$. By Theorem 3.9(i), we get $P_{\tilde{C}_\gamma}(\tilde{u}_0, \tilde{v}_0) = [\frac{\tilde{u}_0}{(1+\lambda)}, \frac{\tilde{v}_0}{(1-\lambda)}]^T$, so

$$\begin{aligned} P_{C_\gamma}(x_0, y_0) &= \begin{bmatrix} \frac{1}{\sqrt{2}} \text{Id} & -\frac{1}{\sqrt{2}} \text{Id} \\ \frac{1}{\sqrt{2}} \text{Id} & \frac{1}{\sqrt{2}} \text{Id} \end{bmatrix} \begin{bmatrix} \frac{1}{1+\lambda} \text{Id} & 0 \\ 0 & \frac{1}{1-\lambda} \text{Id} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} \text{Id} & \frac{1}{\sqrt{2}} \text{Id} \\ -\frac{1}{\sqrt{2}} \text{Id} & \frac{1}{\sqrt{2}} \text{Id} \end{bmatrix} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{1-\lambda^2} \text{Id} & \frac{-\lambda}{1-\lambda^2} \text{Id} \\ \frac{-\lambda}{1-\lambda^2} \text{Id} & \frac{1}{1-\lambda^2} \text{Id} \end{bmatrix} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix}. \end{aligned}$$

Also, $\|p\| = \|\tilde{u}_0\|^2 - \|\tilde{v}_0\|^2 = 2\langle x_0, y_0 \rangle$, and $\|q\| = \|\tilde{u}_0\|^2 + \|\tilde{v}_0\|^2 = \|x_0\|^2 + \|y_0\|^2$.

(ii): For $x_0 = -y_0$, we have $\tilde{u}_0 = 0, \tilde{v}_0 = -\sqrt{2}x_0$, so

$$\begin{aligned} P_{C_\gamma}(x_0, -x_0) &= A_{\frac{\pi}{4}} P_{\tilde{C}_\gamma}(0, -\sqrt{2}x_0) \\ &= \left\{ A_{\frac{\pi}{4}} \begin{bmatrix} \tilde{u} \\ \frac{-x_0}{\sqrt{2}} \end{bmatrix} \mid \|\tilde{u}\|^2 = 2\gamma + \frac{\|-\sqrt{2}x_0\|^2}{4} = 2\gamma + \frac{\|x_0\|^2}{2}, \tilde{u} \in X \right\}. \end{aligned}$$

(iii): When $x_0 \neq -y_0$, and $x_0 = y_0$, we have $x_0 \neq 0, (x_0, y_0) = (x_0, x_0)$, so that $\tilde{v}_0 = 0$ and $\tilde{u}_0 = \frac{x_0 + y_0}{\sqrt{2}} = \sqrt{2}x_0$. Then $\|\tilde{u}_0\| \geq 2\sqrt{2}\gamma \Leftrightarrow \|x_0\| \geq 2\sqrt{\gamma}$.

a) When $\|x_0\| \geq 2\sqrt{\gamma}$, we have

$$\begin{aligned} P_{C_\gamma}(x_0, x_0) &= A_{\frac{\pi}{4}} P_{\tilde{C}_\gamma}(\sqrt{2}x_0, 0) \\ &= A_{\frac{\pi}{4}} \left\{ \begin{bmatrix} \frac{x_0}{\sqrt{2}} \\ \tilde{v} \end{bmatrix} \mid \|\tilde{v}\|^2 = \frac{\|\sqrt{2}x_0\|^2}{4} - 2\gamma = \frac{\|x_0\|^2}{2} - 2\gamma, \tilde{v} \in X \right\}. \end{aligned}$$

b) When $0 < \|x_0\| < 2\sqrt{\gamma}$, we have $\|\tilde{u}_0\| < 2\sqrt{2\gamma}$, so

$$P_{C_\gamma}(x_0, x_0) = A_{\frac{\pi}{4}} P_{\tilde{C}_\gamma}(\sqrt{2}x_0, 0) = A_{\frac{\pi}{4}} \begin{bmatrix} \sqrt{2\gamma} \frac{\sqrt{2}x_0}{\|\sqrt{2}x_0\|} \\ 0 \end{bmatrix} = A_{\frac{\pi}{4}} \begin{bmatrix} \sqrt{2\gamma} \frac{x_0}{\|x_0\|} \\ 0 \end{bmatrix}.$$

■

5 Projections onto hyperbola \tilde{C}_γ and bilinear constraint set C_γ when $\gamma < 0$

Armed with the results in Sections 3 and 4, we can study $P_{\tilde{C}_\gamma}$ and P_{C_γ} when $\gamma < 0$. Define

$$T_1 : X \times X \rightarrow X \times X : (x, y) \mapsto (y, x), \text{ and}$$

$$T_2 : X \times X \rightarrow X \times X : (x, y) \mapsto (x, -y).$$

Theorem 5.1. Let $\gamma < 0, \tilde{u}_0, \tilde{v}_0 \in X$, and $\tilde{C}_\gamma = \{(u, v) \in X \times X \mid \|u\|^2 - \|v\|^2 = 2\gamma\}$. Then the following hold:

(i) When $\tilde{u}_0 \neq 0, \tilde{v}_0 \neq 0$, we have

$$P_{\tilde{C}_\gamma}(\tilde{u}_0, \tilde{v}_0) = \left\{ \begin{bmatrix} \frac{\tilde{u}_0}{1-\lambda} \\ \frac{\tilde{v}_0}{1+\lambda} \end{bmatrix} \right\}$$

in which λ is the unique root of $H(\lambda) = 0$ in $] -1, 1[$, where

$$H(\lambda) = \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1 - \lambda^2)^2} + \gamma, \quad p = \|\tilde{v}_0\|^2 - \|\tilde{u}_0\|^2, \quad \text{and } q = \|\tilde{u}_0\|^2 + \|\tilde{v}_0\|^2.$$

(ii) When $\tilde{v}_0 = 0$, we have

$$P_{\tilde{C}_\gamma}(\tilde{u}_0, 0) = \left\{ \begin{bmatrix} \frac{\tilde{u}_0}{2} \\ \tilde{v} \end{bmatrix} \mid \|\tilde{v}\|^2 = \frac{\|\tilde{u}_0\|^2}{4} - 2\gamma \right\}. \quad (43)$$

(iii) a) When $\tilde{u}_0 = 0$ and $\|\tilde{v}_0\| \geq 2\sqrt{2(-\gamma)}$, we have

$$P_{\tilde{C}_\gamma}(0, \tilde{v}_0) = \left\{ \begin{bmatrix} \tilde{u} \\ \frac{\tilde{v}_0}{2} \end{bmatrix} \mid \|\tilde{u}\|^2 = \frac{\|\tilde{v}_0\|^2}{4} + 2\gamma \right\}. \quad (44)$$

b) When $\tilde{u}_0 = 0$ and $0 < \|\tilde{v}_0\| < 2\sqrt{2(-\gamma)}$, we have

$$P_{\tilde{C}_\gamma}(0, \tilde{v}_0) = \left\{ \begin{bmatrix} 0 \\ \sqrt{2(-\gamma)} \frac{\tilde{v}_0}{\|\tilde{v}_0\|} \end{bmatrix} \right\}. \quad (45)$$

Proof. Since

$$\text{minimize } \|\tilde{u} - \tilde{u}_0\|^2 + \|\tilde{v} - \tilde{v}_0\|^2 \quad \text{subject to } \|\tilde{u}\|^2 - \|\tilde{v}\|^2 = 2\gamma$$

is equivalent to

$$\text{minimize } \|\tilde{v} - \tilde{v}_0\|^2 + \|\tilde{u} - \tilde{u}_0\|^2 \quad \text{subject to} \quad \|\tilde{v}\|^2 - \|\tilde{u}\|^2 = 2(-\gamma),$$

we have $P_{\tilde{C}_\gamma} = T_1 P_{\tilde{C}_{-\gamma}} T_1$. It suffices to apply [Theorem 3.9](#). ■

Theorem 5.2. Let $\gamma < 0$, $x_0, y_0 \in X$, and $C_\gamma = \{(x, y) \in X \times X \mid \langle x, y \rangle = \gamma\}$. Then the following hold:

(i) When $x_0 \neq \pm y_0$, the projection is a singleton:

$$P_{C_\gamma}(x_0, y_0) = \left\{ \begin{bmatrix} \frac{x_0 + \lambda y_0}{1 - \lambda^2} \\ \frac{y_0 + \lambda x_0}{1 - \lambda^2} \end{bmatrix} \right\}$$

in which λ is the unique solution of $H(\lambda) = 0$ in $] -1, 1[$, where

$$H(\lambda) = \frac{(\lambda^2 + 1)p - 2\lambda q}{2(1 - \lambda^2)^2} + \gamma, \quad p = -2\langle x_0, y_0 \rangle, \quad \text{and } q = \|x_0\|^2 + \|y_0\|^2.$$

(ii) When $x_0 = y_0$, the projection is a set:

$$P_{C_\gamma}(x_0, x_0) = \left\{ \begin{bmatrix} \frac{x_0}{2} + \frac{\tilde{u}}{\sqrt{2}} \\ \frac{x_0}{2} - \frac{\tilde{u}}{\sqrt{2}} \end{bmatrix} \mid \|\tilde{u}\|^2 = -2\gamma + \frac{\|x_0\|^2}{2}, \quad \tilde{u} \in X \right\}.$$

(iii) a) When $x_0 = -y_0$ and $\|x_0\| \geq 2\sqrt{-\gamma}$, the projection is a set:

$$P_{C_\gamma}(x_0, -x_0) = \left\{ \begin{bmatrix} \frac{x_0}{2} - \frac{\tilde{v}}{\sqrt{2}} \\ \frac{-x_0}{2} - \frac{\tilde{v}}{\sqrt{2}} \end{bmatrix} \mid \|\tilde{v}\|^2 = \frac{\|x_0\|^2}{2} + 2\gamma, \quad \tilde{v} \in X \right\}.$$

b) When $x_0 = -y_0$ and $0 < \|x_0\| < 2\sqrt{-\gamma}$, the projection is a singleton:

$$P_{C_1}(x_0, -x_0) = \left\{ \sqrt{-\gamma} \begin{bmatrix} \frac{x_0}{\|x_0\|} \\ \frac{-x_0}{\|x_0\|} \end{bmatrix} \right\}.$$

Proof. Since

$$\text{minimize } \|x - x_0\|^2 + \|y - y_0\|^2 \quad \text{subject to} \quad \langle x, y \rangle = \gamma$$

is equivalent to

$$\text{minimize } \|x - x_0\|^2 + \|z - (-y_0)\|^2 \quad \text{subject to} \quad \langle x, z \rangle = -\gamma,$$

we have $P_{C_\gamma} = T_2 P_{C_{-\gamma}} T_2$. It remains to apply [Theorem 4.1](#). ■

Acknowledgments

HHB and XW were supported by NSERC Discovery grants. MKL was supported by SERB-UBC fellowship and HHB and XW's NSERC Discovery grants. This work was presented at the SIAM Conference on Optimization in July 20–23, 2021 by MKL.

References

- [1] H.H. Bauschke and P.L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, second edition, Springer, 2017.
- [2] H.H. Bauschke, M. Krishan Lal, and X. Wang, The projection onto the cross, (2021), <https://arxiv.org/abs/2108.04382>.
- [3] F. Bernard and L. Thibault, Prox-regular functions in Hilbert spaces, *Journal of Mathematical Analysis and Applications* 303 (2005), 1–14.
- [4] D.P. Bertsekas, *Nonlinear Programming*, third edition, Athena Scientific, Belmont, Massachusetts, USA, 2016.
- [5] R.L. Burden, J.D. Faires, and A.M. Burden, *Numerical Analysis*, 10th edition, 2016.
- [6] N. Chernov and H. Ma, Least squares fitting of quadratic curves and surfaces, In *Computer Vision*, Editor S.R. Yoshida, Nova Science Publishers 2011; pp. 285–302.
- [7] N. Chernov and S. Wijewickrema, Algorithms for projecting points onto conics, *Journal of Computational and Applied Mathematics* 251 (2013), 8–21.
- [8] D. Eberly, *Robust and Error-free Geometric Computing*, CRC Press, 2021.
- [9] V. Elser, Learning without loss, *Fixed Point Theory and Algorithms for Sciences and Engineering* 2021, Paper No. 12, 51 pp.. <https://doi.org/10.1186/s13663-021-00697-1>
- [10] V. Elser, Matrix product constraints by projection methods, *Journal of Global Optimization* 68 (2017), 329–355.
- [11] E. Kreyszig, *Introductory Functional Analysis with Applications*, John Wiley & Sons, Inc., New York, 1989.
- [12] D.G. Luenberger, *Optimization by Vector Space Methods*, Wiley, 1969.
- [13] R.T. Rockafellar and R. J-B Wets, *Variational Analysis*, Springer-Verlag, New York, 1998.