

ON BOSE-EINSTEIN CONDENSATES IN THE THOMAS-FERMI REGIME

DANIELE DIMONTE AND EMANUELA L. GIACOMELLI

ABSTRACT. We study a system of N trapped bosons in the Thomas-Fermi regime with an interacting pair potential of the form $g_N N^{3\beta-1} V(N^\beta x)$, for some $\beta \in (0, 1/3)$ and g_N diverging as $N \rightarrow \infty$. We prove that there is complete Bose-Einstein condensation at the level of the ground state and, furthermore, that, if $\beta \in (0, 1/6)$, condensation is preserved by the time evolution.

CONTENTS

1. Introduction and main results	1
2. Ground state energy	8
2.1. Preliminary Estimates	8
2.2. Bound for \mathcal{N}_+	13
2.3. Proof of Theorem 1.1	14
3. Dynamics	14
3.1. Sobolev norms of the solutions	15
3.2. From the many-body problem to Hartree.	16
3.3. From Hartree to Gross-Pitaevskii	23
3.4. Proof of Theorem 1.6	24
Appendix A. Properties of the projectors p_j^φ, q_j^φ	25
References	26

1. INTRODUCTION AND MAIN RESULTS

In recent years, the analysis of quantum systems of interacting particles, in particular bosons, in suitable scaling regimes (mean-field, Gross-Pitaevskii, thermodynamic, etc.) has represented a very flourishing line of research in mathematical physics. Starting from the pioneering works [27, 31, 53], several results have been obtained along this research line about the ground state behavior of interacting bosons and Bose-Einstein condensation (BEC) [38, 40, 41, 42] in both the mean-field (see [28, 34, 35, 36, 49, 52] and references therein), in the Gross-Pitaevskii (GP) regimes (see [6, 7, 30, 43, 44, 45] and references therein) and in the thermodynamic limit (see [3, 25, 54] and references therein), as well as about the many-body dynamics in the mean-field (see [22, 33, 47] and references therein) and in the GP (see [8, 13, 23, 48] and references therein) settings. As we are going to see in more details later, in both these regimes, the effective behavior of the many-body system is shown to be suitably approximated studying an effective one-particle nonlinear problem.

In the typical setting, a large number N of bosonic particles is confined in a box or, more generally, by a trapping potential, and the bottom of the spectrum of the corresponding Hamiltonian, or its unitary evolution, is investigated as $N \rightarrow \infty$. The many-body Schrödinger operator has the form

$$H_N^{\text{trap}} := \sum_{j=1}^N (-\Delta_j + V_{\text{ext}}(x_j)) + g_N N^{3\beta-1} \sum_{1 \leq j < k \leq N} v(N^\beta(x_j - x_k)), \quad (1.1)$$

and it is supposed to act on $L_s^2(\mathbb{R}^{3N})$, which is the subspace of $L^2(\mathbb{R}^{3N})$ consisting of all functions which are symmetric with respect to permutations of the N particles in 3 dimensions. Here, V_{ext} is the trapping potential, which for simplicity is assumed to be homogeneous of order¹ s (see also Remark 1.2), i.e.,

$$V_{\text{ext}}(x) = \lambda|x|^s, \quad \text{with } s \geq 2 \text{ and } \lambda > 0, \quad (1.2)$$

while

$$v_N(x) := N^{3\beta}v(N^\beta x), \quad \text{with } \beta > 0, \quad (1.3)$$

is the pair interaction potential, which is suitably rescaled by the prefactor g_N/N . The parameter $\beta > 0$ varies in $[0, 1]$ and, with $g_N = g = \text{const.}$, identifies different scaling regimes: mean-field for $\beta = 0$, mean-field and zero-range for $\beta \in (0, 1)$ and GP for $\beta = 1$. In fact, β is also a measure of the *diluteness* of the system: indeed, denoting by $\bar{\rho}$ and R_{eff} the average density and the effective radius of the interaction potential, respectively, we have $\bar{\rho} \simeq N \|\rho_N^{\text{GP}}\|_\infty$ (see [16]), where ρ_N^{GP} is the density associated to the condensate wave function.

In this paper we will investigate the Thomas-Fermi (TF) limit, where $g_N \rightarrow \infty$. In the TF regime, one thus has (see Remark 1.3)

$$\bar{\rho}R_N^3 \propto \frac{N^{1-3\beta}}{g_N^{\frac{3}{s+3}}}$$

where $R_N \sim N^{-\beta}$ is the effective range of the interaction potential and $g_N^{\frac{1}{s+3}}$ is the size of the TF radius, i.e., the characteristic length scale of the condensate. Hence, since $g_N \rightarrow +\infty$ as $N \rightarrow \infty$, the above quantity certainly tends to 0 for $\beta > 1/3$, so that the gas is dilute and only binary scattering events are relevant; on the opposite, for $\beta < 1/3$, the quantity may diverge, depending on the how fast g_N tends to $+\infty$, so that any particle in the gas can interact with all the others, as in the typical mean-field picture.

The final outcome in terms of effective description is physically different but it is remarkable that, for any $\beta \in (0, 1]$, a one-particle Schrödinger operator with cubic nonlinearity is recovered as $N \rightarrow \infty$, giving rise to the GP functional

$$\mathcal{E}_{\text{GP}}^{\text{trap}}(u) := \langle u, hu \rangle + 4\pi g \int_{\mathbb{R}^3} dx |u(x)|^4. \quad (1.4)$$

in the stationary setting, where h is the one-body operator

$$h := -\Delta + V_{\text{ext}}, \quad (1.5)$$

and to the GP time-dependent equation

$$i\partial_t \psi_t = (-\Delta + 8\pi g |\psi_t|^2) \psi_t, \quad (1.6)$$

in the dynamical one: note the absence of the trapping potential in the above evolution equation, which is usually switched-off after the occurrence of BEC to observe the gas dynamics. Only the coefficient g of such a nonlinear term changes for different values of β : for $\beta \in (0, 1)$ the coefficient is proportional to the integral of the interaction potential (which has then to be assumed finite), while, for $\beta = 1$, its scattering length appears. In both cases the coefficient g is assumed to be constant as $N \rightarrow \infty$.

The physical motivation behind the choice of letting g_N grow with N is that in physical heuristics, the effective coefficient g is very often assumed to be large and the asymptotics $g \rightarrow +\infty$ considered, which goes under the name of *Thomas-Fermi regime*, because of a similarity at the level of the one-particle density with the Thomas-Fermi theory of electrons. The main reason behind such a choice is that in real experiments $g \propto aN$, a being the measured scattering length of the interaction potential, and therefore, even in presence of few thousands of atoms, the quantity might be quite large compared to the other characteristic lengths of the system.

Another important motivation to let $g = g_N$ diverge is to take into account other settings and, in particular, the thermodynamic limit, i.e., the regime in which $N, L \rightarrow \infty$, but $\rho := N/L^3$ is kept fixed: after rescaling the lengths this limit can be recast in the form above and recovered for $\beta = 1/3$, $g_N = N^{2/3}$. Finally, it is remarkable that a quite consistent literature is available about the ground state behavior of

¹The condition $s \geq 2$ is actually assumed only to simplify the presentation of the result, which holds true with different (more involved) estimates of the error terms also for $s \geq 0$.

GP functional in the asymptotic regime $g \rightarrow +\infty$, in particular in presence of rotation (see [17, 18] and references therein), when vortices might appear as a manifestation of the superfluidity of the system.

In this very same setting, the effective dynamics of vortices can also be studied [29] and shown to be driven by the trapping potential: if at initial time a trapped bosonic system is set in a condensate state with finitely many vortices (e.g., by putting it under sufficiently fast rotation) and then it is let to evolve (switching off the rotation), it is possible to follow the trajectories of vortex points and prove that they are determined by an ODE depending only on the external trap. We are going to comment further on this result below but we point out here that two key ingredients are needed in its proof: the system is assumed to be in the TF regime and the effective evolution equation for the condensate is given by the time-dependent GP equation. Here, we are precisely concerned with proving that such assumptions make sense. More precisely, we show that

- (1) in the TF regime (for $\beta < 1/3$, $g_N \ll N$) Bose-Einstein condensation takes place at the level of the ground state of a system of interacting bosons;
- (2) in the same TF limit, the effective time-evolution of a state with sufficiently low energy is governed by the time-dependent GP equation.

A similar regime has indeed been studied so far at the many-body level only in [2, 11, 16]. However, in [16] the focus is different: the object of the study is the interplay between the TF regime and the presence of a rapid rotation and its effect to the ground state energy is investigated to leading order. We comment later on the similarities between [2, 11] and our work.

Let us now provide some more details about the setting we are going to consider. The main object under investigation is the many-body Hamiltonian (1.1) and its time evolution, acting on bosonic (i.e., symmetric) wave functions in $L_s^2(\mathbb{R}^{3N})$. We restrict our analysis to interaction potentials satisfying the assumptions below.

Assumption 1. *The potential $v \neq 0$ is of positive type, i.e., $\hat{v} \geq 0$ and spherically symmetric. Furthermore, we assume that $|x|v \in L^1(\mathbb{R}^3)$, $v \in L^2(\mathbb{R}^2)$.*

Let us comment further about the characteristic length scales of the system. The most important one is indeed the range of the interaction potential $\propto N^{-\beta}$ and, if g_N is not too large, we claim that it is the only relevant one to be compared with the mean free path of the particles $\propto N^{-1/3}$. This yields the threshold at $\beta = 1/3$ mentioned above. Another important length scale is provided by the *scattering length* associated to v_N : for a potential v the scattering length is defined through the solution of the zero energy scattering equation, i.e.,

$$(-\Delta + \frac{1}{2}v) f = 0, \quad (1.7)$$

with the boundary condition $f(x) \rightarrow 1$ as $|x| \rightarrow \infty$. The scattering length a is then given by

$$8\pi a = \int_{\mathbb{R}^3} dx v(x) f(x). \quad (1.8)$$

If we denote by a_N the scattering length of the potential $g_N v_N(x) = g_N N^{3\beta-1} v(N^\beta x)$ then one can go through the Born approximation of a_N to write

$$a_N = \frac{g_N}{8\pi} \int_{\mathbb{R}^3} dx v_N(x) f_N(x) \sim \frac{g_N}{8\pi N} \int_{\mathbb{R}^3} dx v(x), \quad (1.9)$$

where f_N is the solution of the related scattering equation, i.e., $-\Delta f_N + (g_N/2)v_N f_N = 0$. In order for the expansion to make sense, one has to require that v is integrable and

$$g_N \ll N^{1-\beta}, \quad (1.10)$$

which we always do, but at the same this implies that

$$a_N \ll R_N \sim N^{-\beta}, \quad (1.11)$$

for any $\beta \in [0, 1)$. In other words, a_N is always much smaller than the range of the potential in the regime we are going to consider and plays no role in the picture. A transition is expected to occur when

$g_N \sim N^{1-\beta}$, for $\beta \geq 1/3$, which is precisely the case considered² in [11]. The last characteristic length of the problem is the *healing length* ξ_N , which is given by (see [24, Sect. 4.1])

$$\xi_N = \frac{1}{\sqrt{N \|\rho_N^{\text{GP}}\|_\infty}} \sim N^{-\frac{1}{2}} g_N^{-\frac{3}{2(s+3)}}. \quad (1.12)$$

Hence, one can readily check that under the conditions of next Theorem 1.1

$$a_N \ll \xi_N \ll R_N. \quad (1.13)$$

We denote by $E_0(N)$ the ground state energy of the system, i.e.,

$$E_0(N) := \inf \sigma(H_N^{\text{trap}}) \equiv \inf \{ \langle \psi, H_N^{\text{trap}} \psi \rangle \mid \psi \in L_s^2(\mathbb{R}^{3N}), \|\psi\|_{L_s^2(\mathbb{R}^{3N})} = 1 \}. \quad (1.14)$$

Moreover, $\Psi_0 \in L_s^2(\mathbb{R}^{3N})$ is the unique (up to an overall phase) ground state of H_N and we denote by γ_{Ψ_0} the associated reduced density matrix, i.e.,

$$\gamma_{\Psi_0} := \text{Tr}_{2, \dots, N} |\psi_0\rangle\langle\psi_0|, \quad (1.15)$$

where with $\text{Tr}_{2, \dots, N}$ we denoted the partial trace on all particles but one.

It is well known that in the mean-field regime ($\beta = 0$, $g_N = g$ constant), the ground state energy per particle is well approximated, in the limit $N \rightarrow \infty$, by the infimum of the Hartree functional, which approximately coincides with the restriction of the quadratic form associated to H_N to functions of the form $\psi = u^{\otimes N}$, with $u \in L^2(\mathbb{R}^3)$, $\|u\|_{L^2(\mathbb{R}^3)} = 1$, i.e.,

$$\mathcal{E}_H^{\text{trap}}(u) := \langle u, hu \rangle + \frac{1}{2} \langle u \otimes u, vu \otimes u \rangle. \quad (1.16)$$

More precisely, in [34], it is shown that

$$\lim_{N \rightarrow \infty} \frac{E_0(N)}{N} = \inf \{ \mathcal{E}_H^{\text{trap}}(u) \mid \|u\|_{L^2(\mathbb{R}^3)} = 1 \} =: E^{\text{H}}. \quad (1.17)$$

For trapped bosons in the mean field regime, it has been proven that the system exhibits BEC and its excitation spectrum can be approximated via the well-known Bogoliubov approximation [28], with corrections that can be explicitly identified [9, 10]. Analogous results are proven for a system of bosons in a box in the same regime [19, 36, 52].

Similarly, it is also known that in the GP regime ($\beta = 1$, $g_N = g$ constant), the ground state energy per particle can be approximated, in the limit $N \rightarrow \infty$, by the infimum of the GP effective energy (1.4). Indeed, in [44] it is proven that, in the limit $N \rightarrow \infty$, it holds (recall (1.4))

$$\lim_{N \rightarrow \infty} \frac{E_0(N)}{N} = \inf \{ \mathcal{E}_{\text{GP}}^{\text{trap}}(u) \mid \|u\|_{L^2(\mathbb{R}^3)} = 1 \} =: E_{\text{GP}}^{\text{trap}}. \quad (1.18)$$

The occurrence of BEC in this case has been proven [14, 43] as well as the validity of the Bogoliubov approximation [15, 45]. Similar results both on BEC and the excitation spectrum are available in (see [7, 30, 31] and references therein) for N interacting bosons in a box.

Concerning the intermediate regimes, we have to distinguish between two different cases. The first interesting regime is studied in [5] and it is characterized by taking $\beta \in (0, 1)$ and $g_N = g > 0$ small enough. In [5] it is proven the validity of the Bogoliubov theory in the aforementioned regime, studying both the ground state energy and the excitation spectrum of the system. More recently another regime has been investigated, which can be thought as an interpolation between the GP regime and the thermodynamic limit and it is described by $\beta \in (0, 1)$ small enough and $g_N = N^{1-\beta}$. In [2, 11] it is proven that, in such regime, the low-energy states exhibit BEC and it is studied the low-energy excitation spectrum. Note that, according to the discussion above on the characteristic length scales of the problem, this choice corresponds to a scattering length of the interaction of the same order of its effective range.

In the present work, on the other hand, we are mostly interested in the time-evolution of condensates in the TF regime ($g_N \gg 1$). For the sake of completeness, we also show that BEC occurs in the same setting, at least at the level of the ground state or of any approximate ground state, as it can be easily obtained

²In the notation of [11], $\beta = 1 - \kappa$ and $g_N = N^\kappa$, so that $g_N = N^{1-\beta}$.

by a suitable adaptation of the arguments in [28, 52] and with a much simpler proof (thanks to stronger assumption on β and g_N) than the ones in [2, 11]. More precisely, we prove that the ground state energy of (1.1), as long as $\beta < 1/3$, exhibits BEC in the (unique) minimizer of the GP functional, which in our setting reads as

$$\mathcal{E}_{\text{GP},N}^{\text{trap}}(\psi) := \langle \psi, h\psi \rangle + \frac{g_N}{2} (f v) \int_{\mathbb{R}^3} dx |\psi(x)|^4. \quad (1.19)$$

Theorem 1.1 (BEC in the ground state energy).

Let H_N be the Hamiltonian defined in (1.1) and let v_N be an interaction potential such that v satisfies Assumption 1. Assume that

$$\beta < \frac{1}{3}, \quad \text{and} \quad g_N \ll \min \left\{ N^{\frac{(1-3\beta)(s+3)}{s+5}}, N^{\frac{(s+3)\beta}{2(s+1)}} \right\}. \quad (1.20)$$

Let Ψ be any normalized many-body state such that $\langle \Psi, H_N \Psi \rangle \leq E_N + o(N)$ and let γ_Ψ be the associated reduced density matrix defined as in (1.15). Then,

$$\lim_{N \rightarrow \infty} \langle \varphi_N^{\text{GP}}, \gamma_\Psi \varphi_N^{\text{GP}} \rangle = 1 \quad (1.21)$$

where $\varphi_N^{\text{GP}} \in L^2(\mathbb{R}^3)$ is the unique (up to a phase) normalized minimizer of (1.19).

Remark 1.2 (Homogeneous trapping potential). For the sake of simplicity, we assumed that the trapping potential V_{ext} in (1.2) is homogeneous of order $s > 0$. As it is customary in the TF regime, such an assumption can be relaxed and replaced with the weaker condition that V_{ext} is asymptotically homogeneous, i.e., $V_{\text{ext}}(x) \sim |x|^s$, as $|x| \rightarrow +\infty$, because only the behavior at large $|x|$ matters to leading order. This however makes the estimates of the error terms as well as the conditions on the parameters more implicit.

Remark 1.3 (GP minimizer). We point out that, since the GP functional in (1.19) depends on N , the same holds for its minimizer φ_N^{GP} . Indeed, it is very well known that the density $|\varphi_N^{\text{GP}}|^2$ is pointwise close as $N \rightarrow \infty$ to the minimizer of the TF functional

$$\mathcal{E}_{\text{TF},g_N}^{\text{trap}}(\rho) = \int_{\mathbb{R}^3} dx \left\{ V_{\text{ext}}(x)\rho + \frac{g_N}{2} (f v)\rho^2 \right\} \quad (1.22)$$

among positive densities ρ normalized in L^1 . More precisely, one has (see, e.g., [41, Thm. 2.2])

$$\inf_{\|\psi\|_2=1} \mathcal{E}_{\text{GP},N}^{\text{trap}}(\psi) = \inf_{\rho \geq 0, \|\rho\|_1=1} \mathcal{E}_{\text{TF},g_N}^{\text{trap}}(\rho) (1 + o(1)) = g_N^{\frac{s}{s+3}} \inf_{\rho \geq 0, \|\rho\|_1=1} \mathcal{E}_{\text{TF},1}^{\text{trap}}(\rho) + O\left(g_N^{-\frac{2}{s+3}} \log g_N\right), \quad (1.23)$$

as $g_N \rightarrow +\infty$, and, denoting by $\rho_{g_N}^{\text{TF}}$ the unique TF minimizer,

$$g_N^{\frac{3}{s+3}} \rho_{g_N}^{\text{TF}}\left(g_N^{\frac{1}{s+3}} x\right) = \rho_1^{\text{TF}}(x), \quad \left\| g_N^{\frac{3}{s+3}} \left| \varphi_N^{\text{GP}}\left(g_N^{\frac{1}{s+3}} \cdot\right) \right|^2 - \rho_1^{\text{TF}} \right\|_\infty = o(1), \quad (1.24)$$

i.e., both the GP and TF minimizers spread over a length scale $g_N^{\frac{1}{s+3}}$ (TF radius) and tend pointwise to 0 as $N \rightarrow +\infty$.

Remark 1.4 (Excitation spectrum). In the present paper we do not investigate the excitation spectrum in the TF regime, but we remark that for a system of N bosons confined in a box, it is possible to study it with similar techniques as in [52]. The extension to the case of trapped bosons is however not completely trivial and beyond the scope of the present paper.

Remark 1.5 (Rotating systems). It would be interesting to generalize our result to rotating systems. It is indeed well-known that if the rotational velocity Ω_{rot} of the condensate is not too large, i.e., it is below a critical threshold $\Omega_c(N)$, the GP minimizer is unique. In this case, the proof of BEC is expected to hold with minor modifications. For faster rotation however the GP functional is known to have more than one minimizer. In that case the proof of BEC is more involved and one expects BEC in the mixed state given by convex combination of the GP minimizers (see [39, 44]).

As already anticipated above, after having proven that there is BEC in the ground state energy, a natural question that arises is whether BEC is preserved or not by time evolution. More explicitly, the question we would like to answer is whether we can find an *effective dynamics* for a single particle state that well describes the evolution of the condensed state. We consider a system of N trapped identical bosons, as described before, and we suppose that at the initial time the state is condensed. After a while, we remove the trap, thus the Hamiltonian of the system becomes

$$H_N = \sum_{j=1}^N (-\Delta_j) + g_N N^{3\beta-1} \sum_{1 \leq j < k \leq N} v(N^\beta(x_j - x_k)),$$

with domain $D(H_N) := L_s^2(\mathbb{R}^{3N}) \cap H^2(\mathbb{R}^{3N})$. The dynamics of the system is then encoded in the fact that the state at time t , $\psi_t \in L_s^2(\mathbb{R}^{3N})$, is a solution of the Schrödinger equation

$$\begin{cases} i\partial_t \psi_t = H_N \psi_t, \\ \psi_t|_{t=0} = \psi_0, \end{cases} \quad (1.25)$$

We prove that, up some conditions on β and g_N , we can find an effective equation, i.e., the time-dependent GP equation in the TF limit, which reads

$$i\partial_t \varphi_t = (-\Delta + g_N |\varphi_t|^2) \varphi_t \quad (1.26)$$

such that ψ_t converges to the solution of the effective problem. Although we need some technical assumptions on the solution ψ_t , we are able to provide an explicit rate of convergence.

In the case of dilute limits, this question already has some answers. For example, it is well known [21, 22, 26, 47, 51] that in the mean-field limit, if the initial state ψ_0 exhibits BEC on φ_0 , then ψ_t exhibits BEC on a family of single particle states $\{\varphi_t\}_{t \in \mathbb{R}}$ which solve the *nonlinear Hartree equation*:

$$i\partial_t \varphi_t = (-\Delta + v * |\varphi_t|) \varphi_t, \quad (1.27)$$

Moreover it is well known a similar result for the GP regime (see [4, 13, 22, 23, 48] and references therein). Meaning that if the initial datum exhibits BEC, then BEC is preserved also at later times on a family of single particle states $\{\varphi_t\}_{t \in \mathbb{R}}$ which solve the *nonlinear GP equation*:

$$i\partial_t \varphi_t = (-\Delta + g(\int v) |\varphi_t|^2) \varphi_t, \quad (1.28)$$

where g is a constant which is defined, in the limit as $N \rightarrow \infty$, as N times the scattering length of the system, i.e., $Na \rightarrow g$ as $N \rightarrow \infty$ (in our language it would correspond to $g_N \equiv g$).

As we already discussed for the ground state energy, also for the dynamical framework there are different intermediate regimes corresponding to $\beta \in (0, 1)$. Also in this case [48] if there is BEC at the initial state, there is still BEC in the many-body state of the system at later times on a family of single particle states $\{\varphi_t\}_{t \in \mathbb{R}}$ which solve the following nonlinear equation:

$$i\partial_t \varphi_t = (-\Delta + (\int v) |\varphi_t|^2) \varphi_t \quad (1.29)$$

Moreover, for the same regime, it is also available [11, 12] a norm approximation for the evolution of an initial state exhibiting BEC. This last nonlinear equation is rather similar to the GP equation, with the major difference being that instead of having the constant g here we find $\int v$. This is what one should expect since in this regime, N times the scattering length of the pair potential goes to $\int v$ in the limit $N \rightarrow \infty$.

Thus, if we now look at our setting in which $Na = \mathcal{O}(g_N)$, we can expect that the effective dynamics is a cubic nonlinear Schrödinger equation. Since the effective scattering length of our potential grows as g_N as $N \rightarrow \infty$, such an equation can not be independent of N .

Moreover, we have that, in the sense of distributions, $N^{3\beta} v(N^\beta x) \rightarrow (\int v) \delta_0(x)$ as $N \rightarrow \infty$. Using this, one could also see equation (1.29) as a limit for $N \rightarrow \infty$ of an Hartree equation as in (1.27) with the substitution $v \rightarrow N^{3\beta} v(N^\beta \cdot)$. Notice also that this sort-of intermediate equation is still N dependent, but has now the advantage of being a good candidate for the equation governing the dynamics of the single particle state for our system.

Our discussion above led us to guess that we can expect condensation at later times only on a state which is still N dependent. For this purpose we will need to consider two different differential equations. In analogy to the GP limit, from now on we define the GP equation as

$$i\partial_t \varphi_t^{\text{GP}} = \left(-\Delta + g_N (f v) |\varphi_t^{\text{GP}}|^2 \right) \varphi_t^{\text{GP}}. \quad (1.30)$$

As mentioned above, it will also be useful to consider an intermediate equation, similar to the Hartree equation we mentioned before in (1.27). From now on we refer to the Hartree equation as the equation given by

$$i\partial_t \varphi_t^{\text{H}} = \left(-\Delta + g_N v_N * |\varphi_t^{\text{H}}|^2 \right) \varphi_t^{\text{H}}, \quad (1.31)$$

where we recall $v_N(x) = N^{3\beta} v(N^\beta x)$. In the following we will need to work with the Hartree/GP energy functional. Since to study the dynamics we switch off the trap, to distinguish these functional to the ones introduced before, we define

$$\mathcal{E}_{\text{H}}^{\text{free}}(u) := \int_{\mathbb{R}^3} dx |\nabla u(x)|^2 + \frac{g_N}{2} \int_{\mathbb{R}^3} dx (v_N * |u|^2)(x) |u(x)|^2, \quad (1.32)$$

$$\mathcal{E}_{\text{GP}}^{\text{free}}(u) := \int_{\mathbb{R}^3} dx |\nabla u(x)|^2 + \frac{g_N}{2} (f v) \int_{\mathbb{R}^3} dx |u(x)|^4. \quad (1.33)$$

Theorem 1.6 (Dynamics). *Let $\beta \in (0, 1/6)$ and $\lambda \in (3\beta, 1 - 3\beta)$. Let ψ_t be the solution of (1.25) and φ_t^{GP} be the solution of (1.30) with initial data respectively ψ_0 and φ_0 , both with norm equal to 1. Suppose that v satisfies Assumption 1. It holds true that*

$$\begin{aligned} \left\| \gamma_{\psi_t}^{(1)} - |\varphi_t^{\text{GP}}\rangle\langle\varphi_t^{\text{GP}}| \right\| &\leq \sqrt{2} \left(N^{\frac{1-\lambda}{2}} \left\| \gamma_{\psi_0}^{(1)} - |\varphi_0\rangle\langle\varphi_0| \right\|^{\frac{1}{2}} + N^{\frac{3\beta-\lambda}{2}} \right) e^{C_v C_N(\varphi_0, t) g_N |t|} \\ &\quad + C \sqrt{g_N N^{-\beta}} \left(1 + \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N N^{-\beta} \|\varphi_0\|_{L^\infty}^2} \right) e^{C_v C_N(\varphi_0, t)^2 g_N |t|}, \end{aligned} \quad (1.34)$$

where the constant C_v only depends on the potential v and $C_N(\varphi_0, t)$ is given by

$$C_N(\varphi_0, t) := \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{C g_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4) |t|}. \quad (1.35)$$

Remark 1.7 (Time-scale of the GP approximation). The initial datum determines the maximal time-scale on which the GP approximation makes sense through the quantity $C_N(\varphi_0, t)$ and in particular the exponential appearing in (1.35). This in turn depends on the GP energy of the initial datum via (1.35), which, according to (1.23), can not be smaller than $C g_N^{s/(s+3)}$. Now, the exponential on the right hand side of (1.35) is bounded by a constant as long as

$$t < t_\star \sim g_N^{-\frac{3(s+2)}{s+3}}. \quad (1.36)$$

Note indeed that, assuming the H^2 norm of φ_0 to be bounded uniformly, the second summand in the exponent in (1.35) is much smaller than the first, i.e., of order $g_N^2 N^{-2\beta}$, which vanishes because of the factor $N^{-2\beta}$ (see also below). For a time shorter than t_\star now, $C_N(\varphi_0, t)$ can be bounded by a constant. Plugging this bound into (1.34) and using similar reasonment as above, one gets

$$\left\| \gamma_{\psi_t}^{(1)} - |\varphi_t^{\text{GP}}\rangle\langle\varphi_t^{\text{GP}}| \right\| \leq C \left(N^{\frac{1-\lambda}{2}} \left\| \gamma_{\psi_0}^{(1)} - |\varphi_0\rangle\langle\varphi_0| \right\|^{\frac{1}{2}} + N^{\frac{3\beta-\lambda}{2}} + g_N^{\frac{2s+3}{2(s+3)}} N^{-\frac{\beta}{2}} \right) \quad (1.37)$$

which tends to 0 as $N \rightarrow +\infty$ provided the initial datum is not too far from a condensate on φ_0 and assuming

$$g_N \ll N^{\frac{\beta(s+3)}{2s+3}}. \quad (1.38)$$

Remark 1.8 (Hartree vs GP approximation). Note that according to our result $\beta = 1/8$ is the threshold from Hartree to GP. Meaning that, if $\beta < 1/8$, then Hartree gives a better approximation than GP and the opposite happens if $\beta > 1/8$. However the interval of times we can cover in Theorem 1.6 is the same for both the Hartree and the GP approximations since the constant $C_N(\varphi_0, t)$ in (1.35) is appearing in both of them.

Remark 1.9 (Time-scale of the vortex dynamics). As anticipated, it would be interesting to compare our result with the findings of [29], concerning the effective dynamics of vortices. However, a first major difference with the setting considered there is the absence of the trapping potential, which is mostly due to a technical obstruction for the propagation in time of suitable bounds along the nonlinear dynamics. This difference makes a naive comparison (i.e., for $V_{\text{ext}} = 0$) not so meaningful, but in absence of an external trap, t_* in (1.36) is always much smaller than the characteristic time-scale of the vortex dynamics in [29], i.e., $t_{\text{vortex}} \sim g_N^{2/(s+3)} \log g_N$. However, if g_N is chosen to grow very slowly with respect to N , i.e. $g_N^3 \ll \log \log N$, then the time of the vortices can be achieved.

Acknowledgments. We are grateful to Michele Correggi for his feedback and several insightful discussions. We also thank Peter Pickl for useful comments and discussions. The support of the National Group of Mathematical Physics (GNFM–INdAM) through Progetto Giovani 2016 “Superfluidity and Superconductivity” and Progetto Giovani 2018 “Two-dimensional Phases” is acknowledged.

2. GROUND STATE ENERGY

The aim of this section is to prove Theorem 1.1. We start by recalling the GP functional in (1.19), which, under our hypothesis, admits a unique positive minimizer φ^{GP} satisfying the Euler-Lagrange equation

$$(-\Delta + V_{\text{ext}})\varphi^{\text{GP}} + g_N(fv)|\varphi^{\text{GP}}|^2\varphi^{\text{GP}} = \mu_{\text{GP}}\varphi^{\text{GP}}, \quad (2.1)$$

where

$$\mu_{\text{GP}} := E_{\text{GP},N}^{\text{trap}} + \frac{g_N}{2}(fv)\|\varphi^{\text{GP}}\|_{L^4(\mathbb{R}^3)}^4. \quad (2.2)$$

Because of the uniqueness of the solution to (2.2), φ^{GP} is also the ground state of the GP operator h^{GP} , defined as

$$h^{\text{GP}} := -\Delta + V_{\text{ext}} + g_N(fv)|\varphi^{\text{GP}}|^2 \quad (2.3)$$

One can find a complete set of normalized eigenfunctions $\{\varphi_n\}_{n \in \mathbb{N}}$ for the GP operator h^{GP} , where we identify $\varphi_0 \equiv \varphi^{\text{GP}}$. Given that the spectrum of h^{GP} is discrete, we can assume that the corresponding eigenvalues are ordered in such a way that $\mu_{\text{GP}} \equiv \mu_0 < \mu_1$ and $\mu_j \leq \mu_{j+1}$ for any $j \geq 1$. In what follows it is very important that the inequality $\mu_{\text{GP}} < \mu_1$ is strict, which is due to the fact that h^{GP} has a unique ground state (see also Proposition 2.1).

To prove Theorem 1.1 it is convenient to introduce an operator, which we call \mathcal{N}_+ , that counts the number of particles outside φ^{GP} , i.e., the number of excited particles. For short, we now set $P := |\varphi^{\text{GP}}\rangle\langle\varphi^{\text{GP}}|$ and $Q := 1 - P = \sum_{n \neq 0} |\varphi_n\rangle\langle\varphi_n|$. Furthermore, P_j and Q_j will denote copies of the operators P and Q acting on the j -th particle. The operator \mathcal{N}_+ can then be explicitly written as

$$\mathcal{N}_+ := \sum_{j=1}^N Q_j. \quad (2.4)$$

We now briefly explain the strategy of the proof of Theorem 1.1. In order to prove BEC of the ground state Ψ_0 of H_N in the one-particle state φ^{GP} , we prove in Proposition 2.5 an upper bound to $\langle\Psi_0, \mathcal{N}_+ \Psi_0\rangle$. This estimate in combination with a lower bound on the energy gap $\mu_{\text{GP}} - \mu_1$ yields the result.

2.1. Preliminary Estimates. In this section we prove a lower bound for the energy gap $\mu_{\text{GP}} - \mu_1$ and we estimate $\|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}$ and $\|\nabla\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}$ in terms of g_N .

2.1.1. Energy gap. Acting through the unitary transformation defined by

$$\varphi(x) = \varepsilon^{\frac{3}{s+2}}\psi\left(\varepsilon^{\frac{2}{s+2}}x\right), \quad \varepsilon := (g_N(fv))^{-\frac{s+2}{2(s+3)}} \quad (2.5)$$

one gets

$$\langle\varphi, h^{\text{GP}}\varphi\rangle = \varepsilon^{-\frac{2s}{s+2}}\langle\psi, (-\varepsilon^2\Delta + V_{\text{ext}} + |\psi_0^\varepsilon|^2)\psi\rangle =: \varepsilon^{-\frac{2s}{s+2}}\langle\psi, h_\varepsilon\psi\rangle, \quad (2.6)$$

where ψ_0^ε is the ground state energy of h_ε with energy μ_0^ε , which is explicitly given by

$$\varphi^{\text{GP}}(x) =: \varepsilon^{\frac{3}{s+2}}\psi_0^\varepsilon\left(\varepsilon^{\frac{2}{s+2}}x\right) \quad (2.7)$$

Analogously, we denote by $\mu_1^\varepsilon > \mu_0^\varepsilon$ the second lowest eigenvalue³ of h_ε , and by ψ_1^ε the associated normalized eigenstate. It easily follows from (2.6) that

$$\mu_1 - \mu_{\text{GP}} = \varepsilon^{-\frac{2s}{(s+2)}} (\mu_1^\varepsilon - \mu_0^\varepsilon). \quad (2.8)$$

To study the energy gap, it is then sufficient to find a bound for $\mu_1^\varepsilon - \mu_0^\varepsilon$, that is what we do in the next proposition.

Proposition 2.1 (Energy gap). *There exists a constant $C > 0$ depending only on v and V_{ext} such that*

$$\mu_1^\varepsilon - \mu_0^\varepsilon > C\varepsilon^2. \quad (2.9)$$

As a consequence, there exists a constant $C > 0$ depending only on v and V_{ext} such that

$$\mu_1 - \mu_{\text{GP}} > Cg_N^{-\frac{2}{s+3}}. \quad (2.10)$$

A key ingredient for the proof of Proposition 2.1 is an integral estimate for ψ_1^ε . This is the content of the next proposition, whose proof is very similar to [46, Proposition 3.5].

Proposition 2.2 (Integral decay estimates). *Let $A(x)$ be the Agmon distance associated to $V_{\text{ext}}(x) = k|x|^s$ ($s \geq 2$), i.e.,*

$$A(x) = \sqrt{k} \frac{|x|^{1+\frac{s}{2}}}{1+s/2}. \quad (2.11)$$

Then

$$\int_{\mathbb{R}^3} dx e^{\frac{A(x)}{\varepsilon^2}} |\psi_1^\varepsilon(x)|^2 \leq (1 + \mu_1^\varepsilon) e^{\frac{C}{\varepsilon^2}(1+\mu_1^\varepsilon)}, \quad (2.12)$$

where $C > 0$ is a positive constant which depends only on the trapping potential V_{ext} .

In the proof we use the following lemma.

Lemma 2.1. *Let $W(x) := V_{\text{ext}}(x) + |\psi_0^\varepsilon(x)|^2 - \mu_1^\varepsilon$ and let $\Phi(x) := A(x)/2\varepsilon^2$, where $A(x)$ is as in (2.11). It holds true that*

$$\int_{\mathbb{R}^3} dx (W(x) - \varepsilon^2 |\nabla \Phi(x)|^2) e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 \leq 0 \quad (2.13)$$

The proof of the lemma above is an adaptation of [46, Lemma 3.6].

Proof. We set

$$\Omega := \{x \in \mathbb{R}^3 \mid V_{\text{ext}}(x) - 4\mu_1^\varepsilon < 4\} \quad (2.14)$$

and

$$\Phi(x) := \frac{1}{2\varepsilon^2} A(x), \quad (2.15)$$

so that

$$|\nabla \Phi(x)|^2 = \frac{1}{4\varepsilon^2} V_{\text{ext}}(x). \quad (2.16)$$

We then have that for each $x \in \Omega^c \equiv \mathbb{R}^3 \setminus \Omega$,

$$(V_{\text{ext}}(x) + |\psi_0^\varepsilon(x)|^2 - \mu_1^\varepsilon) - \varepsilon^2 |\nabla \Phi(x)|^2 \geq 1. \quad (2.17)$$

In what follows we use $W(x)$ to denote the potential $W(x) := V_{\text{ext}}(x) + |\psi_0^\varepsilon(x)|^2 - \mu_1^\varepsilon$. By (2.17), we get

$$\begin{aligned} \int_{\Omega^c} dx e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 &\leq \int_{\Omega^c} dx (W(x) - \varepsilon^2 |\nabla \Phi(x)|^2) e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 \\ &\leq - \int_{\Omega} dx (W(x) - \varepsilon^2 |\nabla \Phi(x)|^2) e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2, \end{aligned} \quad (2.18)$$

where in the last inequality we used that

$$\int_{\mathbb{R}^3} dx (W(x) - \varepsilon^2 |\nabla \Phi(x)|^2) e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 \leq 0, \quad (2.19)$$

³Note that the strict inequality $\mu_1^\varepsilon > \mu_0^\varepsilon$ follows from the uniqueness of the ground state of h_ε .

which is what is stated in Lemma 2.1. Using now that $W(x) - \varepsilon^2 |\nabla \Phi|^2 \geq -\mu_1^\varepsilon$, from (2.18), we have

$$\int_{\Omega^c} dx e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 \leq \mu_1^\varepsilon \int_{\Omega} dx e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2, \quad (2.20)$$

which implies

$$\int_{\mathbb{R}^3} dx e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 \leq (\mu_1^\varepsilon + 1) \int_{\Omega} dx e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2. \quad (2.21)$$

To conclude, we bound the integral in the r.h.s. above. It is then enough to use that there exists a constant $C > 0$ which depends only on the trapping potential such that

$$\sup_{x \in \Omega} e^{\frac{\Phi(x)}{\varepsilon^2}} \leq e^{\frac{C}{\varepsilon^2}(1+\mu_1^\varepsilon)}. \quad (2.22)$$

We then get that

$$\int_{\mathbb{R}^3} dx e^{2\Phi(x)} |\psi_1^\varepsilon(x)|^2 \leq (\mu_1^\varepsilon + 1) e^{\frac{C}{\varepsilon^2}(1+\mu_1^\varepsilon)}. \quad (2.23)$$

□

Proposition 2.3 (Boundedness of μ_j^ε). *There exist a constants $C > 0$ depending only on v and V_{ext} such that, for $j = 0, 1$,*

$$0 < \mu_j^\varepsilon \leq C. \quad (2.24)$$

Proof. We first prove that μ_0^ε is bounded. It is trivial to see that the energy $\mathcal{E}^\varepsilon(\psi)$, where we set

$$\mathcal{E}^\varepsilon(\psi) := \langle \psi, (-\varepsilon^2 \Delta + V_{\text{ext}}) \psi \rangle + \frac{1}{2} \int_{\mathbb{R}^3} dx |\psi(x)|^4, \quad (2.25)$$

is uniformly bounded, by evaluating it on a suitable trial state. Therefore,

$$\mu_0^\varepsilon = \mathcal{E}^\varepsilon(\psi_0^\varepsilon) + \frac{1}{2} \|\psi_0^\varepsilon\|_4^4 \leq 2\mathcal{E}^\varepsilon(\psi_0^\varepsilon) \leq C. \quad (2.26)$$

We now bound μ_1^ε . First notice that given that V_{ext} is even, then ψ_0^ε is even. Now, by the min-max principle

$$\mu_1^\varepsilon = \inf_{\substack{\|\psi\|_{L^2(\mathbb{R}^3)}=1 \\ \psi \perp \psi_0^\varepsilon}} \langle \psi, h_\varepsilon \psi \rangle, \quad (2.27)$$

so that, for any normalized odd trial state $\psi_{\text{trial}} \perp \psi_0^\varepsilon$ we get

$$\mu_1^\varepsilon \leq \langle \psi_{\text{trial}}, h_\varepsilon \psi_{\text{trial}} \rangle \leq C(1 + \|\psi_0^\varepsilon\|_\infty^2). \quad (2.28)$$

Furthermore, a direct application of the maximum principle to the eigenvalue equation

$$\varepsilon^2 \Delta \psi_0^\varepsilon = (V_{\text{ext}} + |\psi_0^\varepsilon|^2 - \mu_0^\varepsilon) \psi_0^\varepsilon \quad (2.29)$$

provides an upper bound on $\|\psi_0^\varepsilon\|_\infty$: since by elliptic theory ψ_0^ε is smooth, at any maximum point $\bar{x} \in \mathbb{R}^3$, one has

$$0 \geq \Delta \psi_0^\varepsilon(\bar{x}) \geq (|\psi_0^\varepsilon(\bar{x})|^2 - \mu_{\text{GFP}}) \psi_0^\varepsilon(\bar{x}), \quad (2.30)$$

so that (recall that $\psi_0^\varepsilon > 0$)

$$|\psi_0^\varepsilon(\bar{x})|^2 \leq \mu_0^\varepsilon \leq C, \quad (2.31)$$

which in turn implies the uniform boundedness of μ_1^ε via (2.28). □

Proof of Proposition 2.1. We first prove (2.9) and then deduce from it the estimate (2.10).

Bound for $\mu_1^\varepsilon - \mu_0^\varepsilon$. Let $u(x) := \psi_1^\varepsilon(x)/\psi_0^\varepsilon(x)$, which is well posed since $\psi_0^\varepsilon(x) > 0$. We have that

$$\mu_1^\varepsilon = \mu_0^\varepsilon + \varepsilon^2 \int_{\mathbb{R}^3} dx |\nabla u(x)|^2 |\psi_0^\varepsilon(x)|^2,$$

since, by the eigenvalue equations for ψ_0^ε ,

$$\begin{aligned} \int_{\mathbb{R}^3} dx \varepsilon^2 |\nabla \psi_1^\varepsilon(x)|^2 &= \int_{\mathbb{R}^3} dx \varepsilon^2 |u(x) \nabla \psi_0^\varepsilon(x) + (\nabla u(x)) \psi_0^\varepsilon(x)|^2 \\ &= \int_{\mathbb{R}^3} dx \{ \varepsilon^2 |\nabla u(x)|^2 |\psi_0^\varepsilon(x)|^2 - \varepsilon^2 |u(x)|^2 \psi_0^\varepsilon(x) \Delta \psi_0^\varepsilon(x) \} \\ &= \int_{\mathbb{R}^3} dx \{ \varepsilon^2 |\nabla u(x)|^2 |\psi_0^\varepsilon(x)|^2 - V_{\text{ext}}(x) |u(x)|^2 |\psi_0^\varepsilon(x)|^2 \\ &\quad - |u(x)|^2 |\psi_0^\varepsilon(x)|^4 + \mu_0^\varepsilon |u(x)|^2 |\psi_0^\varepsilon(x)|^2 \}. \end{aligned} \quad (2.32)$$

To find a lower bound for $\mu_1^\varepsilon - \mu_0^\varepsilon$, we then use Poincaré's inequality. More precisely, let $B_R := \{x \in \mathbb{R}^3 \mid |x| < R\}$ for some $R > 0$ to be fixed later and let

$$u_R := \int_{B_R} u(x) |\psi_0^\varepsilon(x)|^2. \quad (2.33)$$

We can bound

$$\mu_1^\varepsilon - \mu_0^\varepsilon \geq C_R \varepsilon^2 \int_{B_R} dx |u(x) - u_R|^2 |\psi_0^\varepsilon(x)|^2,$$

by the weighted Poincaré's inequality (see [50] and references therein) and where $C_R > 0$ is a finite constant depending only on R . Now, recalling that $\psi_1^\varepsilon \perp \psi_0^\varepsilon$, we get

$$\begin{aligned} \int_{B_R} dx |u(x) - u_R|^2 |\psi_0^\varepsilon(x)|^2 &= \int_{B_R} dx |\psi_1^\varepsilon(x)|^2 + \left| \int_{B_R} dx \psi_0^\varepsilon(x) \psi_1^\varepsilon(x) \right|^2 \left(\int_{B_R} dx |\psi_0^\varepsilon(x)|^2 - 2 \right) \\ &= 1 - \int_{B_R^c} dx |\psi_1^\varepsilon(x)|^2 - \left| \int_{B_R^c} dx \psi_0^\varepsilon(x) \psi_1^\varepsilon(x) \right|^2 \left(1 + \int_{B_R^c} dx |\psi_0^\varepsilon(x)|^2 \right), \end{aligned} \quad (2.34)$$

where we also used that $\|\psi_j^\varepsilon\|_{L^2(\mathbb{R}^3)} = 1$ for $j = 0, 1$. Using now Cauchy-Schwarz, we obtain

$$\mu_1^\varepsilon - \mu_0^\varepsilon \geq C_R \varepsilon^2 \left(1 - 3 \int_{B_R^c} dx |\psi_1^\varepsilon(x)|^2 \right) \quad (2.35)$$

We then have to bound

$$\int_{B_R^c} dx |\psi_1^\varepsilon(x)|^2, \quad (2.36)$$

to do that we use Proposition 2.2. Indeed, from (2.12), we get that there exist two constants $C, C' > 0$ such that

$$\int_{B_R^c} dx |\psi_1^\varepsilon(x)|^2 \leq (1 + \mu_1^\varepsilon) e^{-\frac{C}{\varepsilon^2} (C' R^{1+\frac{s}{2}} - 1 - \mu_1^\varepsilon)}. \quad (2.37)$$

Given that μ_1^ε is uniformly bounded, there exists a constant $C'' > 0$ independent of ε such that $\mu_1^\varepsilon \leq C''$. We can then fix R so that

$$R^{1+\frac{s}{2}} > \frac{1 + C''}{C'} \geq \frac{1 + \mu_1^\varepsilon}{C'}, \quad (2.38)$$

to get

$$\mu_1^\varepsilon - \mu_0^\varepsilon > C \varepsilon^2. \quad (2.39)$$

Bound for $\mu_{\text{GP}} - \mu_1$. By (2.9) and (2.8)

$$\mu_{\text{GP}} - \mu_1 \geq C g_N^{-\frac{2}{s+3}}, \quad (2.40)$$

which implies the result. \square

2.1.2. *Estimates for $\|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}$, $\|\nabla\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}$.*

Proposition 2.4 (L^∞ bounds). *Let φ^{GP} be the normalized ground state of the operator h^{GP} defined in (2.3). There exists a positive constant $C > 0$ such that*

$$\|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \leq C g_N^{-\frac{3}{2(s+3)}}, \quad \|\nabla\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \leq C g_N^{\frac{2s-3}{2(s+3)}}. \quad (2.41)$$

Proof. Recall that φ^{GP} is such that $\langle \varphi^{\text{GP}}, h^{\text{GP}} \varphi^{\text{GP}} \rangle = \mu_{\text{GP}}$, it then follows that φ^{GP} satisfies the following variational equation

$$\Delta\varphi^{\text{GP}} = (V_{\text{ext}} + g_N(fv)|\varphi^{\text{GP}}|^2 - \mu_{\text{GP}}) \varphi^{\text{GP}}. \quad (2.42)$$

Proceeding similarly as in Proposition 2.3, one can prove that

$$|\varphi^{\text{GP}}(x_0)|^2 \leq \frac{\mu_{\text{GP}}}{g_N(fv)}. \quad (2.43)$$

From (2.8) together with (2.24), it follows that

$$\mu^{\text{GP}} \leq C g_N^{\frac{s}{s+3}}, \quad (2.44)$$

which immediately implies

$$\|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \leq C g_N^{-\frac{3}{2(s+3)}}, \quad s \geq 2. \quad (2.45)$$

We now estimate the L^∞ -norm of $\nabla\varphi^{\text{GP}}$. For short, we set

$$f(x) := V_{\text{ext}}(x)\varphi^{\text{GP}}(x) + g_N(fv)|\varphi^{\text{GP}}(x)|^2\varphi^{\text{GP}}(x) - \mu_{\text{GP}}\varphi^{\text{GP}}(x), \quad (2.46)$$

so that φ^{GP} satisfies $\Delta\varphi^{\text{GP}} = f$. The bound on the L^∞ -norm of $\nabla\varphi^{\text{GP}}$ can then be proven by standard elliptic estimates. More precisely, there exists a constant $\gamma > 0$ such that for all $x \in \mathbb{R}^3$ it holds

$$\|\nabla\varphi^{\text{GP}}\|_{L^\infty(B_{\rho/2}(x_0))} \leq \gamma\|f\|_{L^\infty(B_\rho(x_0))} + \frac{16\gamma}{\rho^4}\|\varphi^{\text{GP}}\|_{L^1(B_\rho(x_0))}, \quad (2.47)$$

where we denote by $B_r(x_0) := \{y \in \mathbb{R}^3 \mid |y - x_0| < r\}$. For a proof of (2.47) we refer to (Proposition 11.2, [20]). By Cauchy-Schwarz inequality, we get

$$|\nabla\varphi^{\text{GP}}(x_0)| \leq \|\nabla\varphi^{\text{GP}}\|_{L^\infty(B_{\rho/2}(x_0))} \leq C \left(\|f\|_{L^\infty(\mathbb{R}^3)} + \rho^{-\frac{5}{2}}\|\varphi^{\text{GP}}\|_{L^2(\mathbb{R}^3)} \right) \leq C\|f\|_{L^\infty(\mathbb{R}^3)}, \quad (2.48)$$

where in the last inequality we fix the radius ρ to be $\rho = \|f\|_{L^\infty(\mathbb{R}^3)}^{-2/5}$ and used that $\|\varphi^{\text{GP}}\|_{L^2(\mathbb{R}^3)} = 1$. We then have to estimate the L^∞ norm of f . To do that, we use the bounds (2.44), (2.45) and we get

$$\|f\|_{L^\infty(\mathbb{R}^3)} = \|V_{\text{ext}}(x)\varphi^{\text{GP}}(x) + g_N(fv)|\varphi^{\text{GP}}(x)|^2\varphi^{\text{GP}}(x) - \mu_{\text{GP}}\varphi^{\text{GP}}(x)\|_{L^\infty(\mathbb{R}^3)} \leq C g_N^{\frac{2s-3}{2(s+3)}} + \|V_{\text{ext}}\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}. \quad (2.49)$$

Using the exponential decay of φ^{GP} , which can be deduced from Proposition 2.2 together with the maximum principle for subharmonic functions and which implies that

$$|\varphi^{\text{GP}}(x)| \leq C g_N^{-\frac{3}{2(s+3)}} e^{-C g_N^{\frac{s+2}{s+3}} (|g_N^{-\frac{1}{s+3}} x| - C')} \quad \text{for } |x| \geq g_N^{\frac{1}{s+3}} R, \quad (2.50)$$

for some $R > 0$ big enough, it follows that

$$\|V_{\text{ext}}\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \leq C g_N^{\frac{2s-3}{2(s+3)}}. \quad (2.51)$$

Inserting (2.49) and (2.51) in (2.48), we get

$$\|\nabla\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \leq C g_N^{\frac{2s-3}{2(s+3)}}. \quad (2.52)$$

□

2.2. Bound for \mathcal{N}_+ . Before proving Theorem 1.1, we need a bound on \mathcal{N}_+ . This is the content of the next Proposition. The proof is very similar to the one of [28, Lemma 2]. In the proof we will use Lemma 2.2, which is proven below.

Proposition 2.5 (Bound for \mathcal{N}_+). *Let v satisfy Assumption 1. Then, as $N \rightarrow \infty$, there exists a constant C depending only on v such that*

$$\langle \Psi_0, \mathcal{N}_+ \Psi_0 \rangle \leq C \left(g_N^{\frac{s+5}{s+3}} N^{3\beta} + g_N^{\frac{2(s+1)}{s+3}} N^{1-\beta} \right). \quad (2.53)$$

Proof. We first prove an upper bound for $E_0(N)$. As a trial state we choose the one in which all the particles occupy the GP minimizer, i.e., $(\varphi^{\text{GP}})^{\otimes N}$, obtaining

$$\begin{aligned} E_0(N) &\leq N \langle \varphi^{\text{GP}}, (-\Delta + V_{\text{ext}}) \varphi^{\text{GP}} \rangle + g_N \frac{(N-1)}{2} \int_{\mathbb{R}^3} dx (v_N * |\varphi^{\text{GP}}|^2)(x) |\varphi^{\text{GP}}(x)|^2 \\ &\leq N \langle \varphi^{\text{GP}}, (-\Delta + V_{\text{ext}}) \varphi^{\text{GP}} \rangle + g_N (f v) \frac{N-1}{2} \int_{\mathbb{R}^3} dx |\varphi^{\text{GP}}(x)|^4 \\ &\quad + C g_N \frac{N-1}{N^\beta} \|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \|\nabla \varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \\ &\leq N E_{\text{GP},N}^{\text{trap}} + C g_N N^{1-\beta} \|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \|\nabla \varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}, \end{aligned} \quad (2.54)$$

where in the second inequality we used Lemma 2.2.

We now prove the lower bound. For any $m \in \mathbb{N}$ we set $\psi_m(x) := |\varphi^{\text{GP}}|^2(x) - \frac{1}{N} \sum_{j=1}^N \eta_m(x - x_j)$, where $\eta_m(x) \in C_c^\infty(\mathbb{R}^3)$ is a sequence of mollifiers, such that $\|\eta_m\|_2 = 1$ for any $m \in \mathbb{N}$ and $(v_N * \eta_m)(x) \rightarrow v_N(x)$ a.e. as $m \rightarrow \infty$. We then have

$$\begin{aligned} 0 &\leq \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \bar{\psi}_m(x) v_N(x-y) \psi_m(y) = \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy |\varphi^{\text{GP}}(x)|^2 v_N(x-y) |\varphi^{\text{GP}}(y)|^2 \\ &\quad - \frac{2}{N} \sum_{j=1}^N \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy |\varphi^{\text{GP}}(x)|^2 v_N(x-y) \eta_m(y-x_j) \\ &\quad + \frac{1}{N^2} \sum_{j,k=1}^N \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \eta_m(x-x_j) v_N(x-y) \eta_m(y-x_k). \end{aligned} \quad (2.55)$$

Taking the limit $m \rightarrow \infty$, by monotone convergence theorem, we get

$$0 \leq \int_{\mathbb{R}^3} dx (v_N * |\varphi^{\text{GP}}|^2)(x) |\varphi^{\text{GP}}(x)|^2 - \frac{2}{N} \sum_{j=1}^N (v_N * |\varphi^{\text{GP}}|^2)(x_j) + \frac{1}{N^2} \sum_{j,k=1}^N v_N(x_j - x_k), \quad (2.56)$$

which implies that

$$\frac{g_N}{N} \sum_{1 \leq j < k \leq N} v_N(x_j - x_k) \geq -\frac{g_N N}{2} \int_{\mathbb{R}^3} dx (v_N * |\varphi^{\text{GP}}|^2)(x) |\varphi^{\text{GP}}(x)|^2 + g_N \sum_{j=1}^N (v_N * |\varphi^{\text{GP}}|^2)(x_j) - g_N N^{3\beta} v(0). \quad (2.57)$$

From (2.57), we then get

$$\begin{aligned} H_N^{\text{trap}} &\geq \sum_{j=1}^N [-\Delta_j + V_{\text{ext}}(x_j) + g_N (v_N * |\varphi^{\text{GP}}|^2)(x_j)] \\ &\quad - \frac{g_N N}{2} \int_{\mathbb{R}^3} dx (v_N * |\varphi^{\text{GP}}|^2)(x) |\varphi^{\text{GP}}(x)|^2 - g_N N^{3\beta} v(0) \end{aligned}$$

Now, using Lemma 2.2 twice and exploiting the normalization of φ^{GP} , we can replace both convolutions, obtaining

$$H_N^{\text{trap}} - N E_{\text{GP},N}^{\text{trap}} \geq \sum_{j=1}^N (h_j^{\text{GP}} - \mu_{\text{GP}}) - g_N N^{3\beta} v(0) - C g_N N^{1-\beta} \|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \|\nabla \varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}. \quad (2.58)$$

Hence, recalling the eigenfunctions $\{\varphi_n\}_{n \in \mathbb{N}}$ of h^{GP} with eigenvalues $\{\mu_n\}_{n \in \mathbb{N}}$ (with $\varphi_0 \equiv \varphi^{\text{GP}}$, $\mu_0 \equiv \mu_{\text{GP}}$), we have that

$$\sum_{j=1}^N (h_j^{\text{GP}} - \mu_{\text{GP}}) = \sum_{j \geq 0} (\mu_j - \mu_0) |\varphi_j\rangle \langle \varphi_j| \geq (\mu_1 - \mu_0) \mathcal{N}_+ \geq 0 \quad (2.59)$$

that we plug into (2.58), so obtaining in combination with the upper bound (2.54)

$$(\mu_1 - \mu_{\text{GP}}) \langle \Psi, \mathcal{N}_+ \Psi \rangle \leq g_N N^{3\beta} v(0) + C g_N N^{1-\beta} \|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \|\nabla \varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}. \quad (2.60)$$

Applying Proposition 2.1 and Proposition 2.4, we finally get

$$C g_N^{-\frac{2}{s+3}} \langle \Psi_0, \mathcal{N}_+ \Psi_0 \rangle \leq (\mu_1 - \mu_{\text{GP}}) \langle \Psi_0, \mathcal{N}_+ \Psi_0 \rangle \leq C' \left(g_N N^{3\beta} + g_N^{\frac{2s}{s+3}} N^{1-\beta} \right). \quad (2.61)$$

for two positive constant C, C' . The result immediately follows. \square

We now prove a Lemma on the comparison between the Hartree and GP interactions.

Lemma 2.2 (Hartree and GP interactions). *Let v satisfy Assumption 1. Then, there exists a constant $C > 0$ such that*

$$\|v_N * |\varphi^{\text{GP}}|^2(x) - (f v) |\varphi^{\text{GP}}|^2(x)\|_{L^\infty(\mathbb{R}^3)} \leq \frac{C}{N^\beta} \|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \|\nabla \varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \quad (2.62)$$

Proof. A direct computation yields

$$\|v_N * |\varphi^{\text{GP}}|^2(x) - (f v) |\varphi^{\text{GP}}|^2(x)\|_{L^\infty(\mathbb{R}^3)} = \sup_{x \in \mathbb{R}^3} \left| \int_{\mathbb{R}^3} dy v(y) \left(|\varphi^{\text{GP}}(x - N^{-\beta} y)|^2 - |\varphi^{\text{GP}}|^2(x) \right) \right| \quad (2.63)$$

$$\begin{aligned} &= \sup_{x \in \mathbb{R}^3} \left| \int dy v(y) \int_0^1 d\lambda \frac{2y}{N^\beta} |\varphi^{\text{GP}}(x - \lambda N^{-\beta} y)| |\nabla \varphi^{\text{GP}}(x - \lambda N^{-\beta} y)| \right| \\ &\leq \frac{2}{N^\beta} \left(\int dy |y| v(y) \right) \|\varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \|\nabla \varphi^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}. \end{aligned} \quad (2.64)$$

\square

2.3. Proof of Theorem 1.1. By Proposition 2.5

$$\langle \Psi_0, \mathcal{N}_+ \Psi_0 \rangle \leq C \left(g_N^{\frac{s+5}{s+3}} N^{3\beta} + g_N^{\frac{2(s+1)}{s+3}} N^{1-\beta} \right), \quad (2.65)$$

so that

$$1 - \langle \varphi^{\text{GP}}, \gamma_{\Psi_0} \varphi^{\text{GP}} \rangle = \frac{1}{N} [N - \langle \Psi_0, a^*(\varphi^{\text{GP}}) a(\varphi^{\text{GP}}) \Psi_0 \rangle] = \frac{1}{N} \langle \Psi_0, \mathcal{N}_+ \Psi_0 \rangle \quad (2.66)$$

$$\leq C g_N^{\frac{s+5}{s+3}} N^{3\beta-1} + C g_N^{\frac{2(s+1)}{s+3}} N^{-\beta} \quad (2.67)$$

which completes the proof.

3. DYNAMICS

This section is devoted to the proof of Theorem 1.6. We summarize here the main steps.

- First, we use the techniques introduced in [47] to approximate the many-body state at time t , ψ_t , in terms of φ_t^{H} which we recall is the solution of the nonlinear Hartree equation introduced in (1.31):

$$i \partial_t \varphi_t^{\text{H}} = (-\Delta + g_N v_N * |\varphi_t^{\text{H}}|^2) \varphi_t^{\text{H}}$$

- Afterwards, we estimate the distance between φ_t^{H} and the solution φ_t^{GP} of the GP equation introduced in (1.30), i.e.,

$$i \partial_t \varphi_t^{\text{GP}} = (-\Delta + g_N (f v) |\varphi_t^{\text{GP}}|^2) \varphi_t^{\text{GP}}$$

3.1. Sobolev norms of the solutions. An important role in the proof of Theorem 1.6 is played the L^∞ norms of φ_t^{GP} and φ_t^{H} . We then adapt the proof of [4, Proposition 3.1 (ii)] to our framework to prove a bound for the H^2 -norm of φ^{GP} and φ^{H} , from which we deduce the estimates for the L^∞ -norms.

Proposition 3.1 (Sobolev bound for $\varphi_t^{\text{H}}, \varphi_t^{\text{GP}}$). *Let φ_t^{GP} and φ_t^{H} be solutions of, respectively, (1.30) and (1.31), both with initial datum $\varphi_0 \in H^n(\mathbb{R}^3)$. Assume that v satisfies Assumption 1.*

(i) *There exists a positive constant $C > 0$ which depends only on the interaction v such that*

$$\sup_{t \in \mathbb{R}} \|\varphi_t^{\text{GP}}\|_{H^1(\mathbb{R}^3)} \leq \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)}, \quad \sup_{t \in \mathbb{R}} \|\nabla \varphi_t^{\text{H}}\|_2 \leq C \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)} + C g_N N^{-\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^2 \quad (3.1)$$

(ii) *There exists a positive constant C such that for any $t \in \mathbb{R}$*

$$\|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)} \leq C \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{C g_N^2 \mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) |t|}, \quad (3.2)$$

$$\|\varphi_t^{\text{H}}\|_{H^2(\mathbb{R}^3)} \leq C \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{C g_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4) |t|}. \quad (3.3)$$

In the proof of Proposition 3.1, we use some Strichartz estimates. In particular we use that for any function $f \in L_t^\infty L_x^{\frac{6}{5}}$, one has

$$\sup_{t \in [0, T]} \left\| \int_0^t ds e^{i(t-s)\Delta} f(s, \cdot) \right\|_{L^2(\mathbb{R}^3)} \leq \sqrt{T} \sup_{t \in [0, T]} \|f(t, \cdot)\|_{L^{\frac{6}{5}}(\mathbb{R}^3)}. \quad (3.4)$$

For a proof of (3.4), see [32, Theorem 1.2].

Proof of Proposition 3.1. We start by showing the bounds in (3.1). The estimate for φ_t^{GP} directly follows from the fact that $\hat{v} \geq 0$ and that the energy is preserved by the GP dynamics. We now look at φ_t^{H} . Notice that, proceeding similarly as in Lemma 2.2 and using Assumption 1 we get

$$|\langle \varphi_0, (v_N * |\varphi_0|^2 - |\varphi_0|^2) \varphi_0 \rangle| \leq C g_N N^{-\beta} \|\nabla \varphi_0\|_{L^2(\mathbb{R}^3)} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^2 \quad (3.5)$$

Using now Cauchy-Schwartz and the fact that the energy is preserved by the Hartree dynamics, we can write

$$\|\nabla \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2 \leq \langle \varphi_0, (-\Delta + \frac{g_N}{2} v_N * |\varphi_0|^2) \varphi_0 \rangle \leq \mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + C \|\nabla \varphi_0\|_{L^2(\mathbb{R}^3)}^2 + C g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4. \quad (3.6)$$

We then get

$$\sup_{t \in \mathbb{R}} \|\nabla \varphi_t^{\text{H}}\|_2 \leq C \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)} + C g_N N^{-\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^2. \quad (3.7)$$

We now prove (3.3). Being φ_t^{GP} the solution of (1.30), we can write

$$\varphi_t^{\text{GP}} = e^{it\Delta} \varphi_{t_0}^{\text{GP}} - i g_N \int_{t_0}^t ds e^{i(t-s)\Delta} \left(|\varphi_t^{\text{GP}}|^2 \varphi_t^{\text{GP}} \right). \quad (3.8)$$

Consider now $\alpha \in \mathbb{N}^3$ with $|\alpha| = 2$; we use the notation $\partial^\alpha = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \partial_{x_3}^{\alpha_3}$. We have

$$\begin{aligned} \partial^\alpha \varphi_t^{\text{GP}} &= e^{it\Delta} \partial^\alpha \varphi_{t_0}^{\text{GP}} - i g_N \int_{t_0}^t ds e^{i(t-s)\Delta} \partial^\alpha \left(|\varphi_t^{\text{GP}}|^2 \varphi_t^{\text{GP}} \right) \\ &= e^{it\Delta} \partial^\alpha \varphi_{t_0}^{\text{GP}} - i g_N \int_{t_0}^t ds e^{i(t-s)\Delta} \sum_{\gamma \leq \beta \leq \alpha} \binom{\alpha}{\beta} \binom{\beta}{\gamma} \left(\partial^{(\alpha-\beta)} \overline{\varphi_t^{\text{GP}}} \partial^{(\beta-\gamma)} \varphi_t^{\text{GP}} \partial^\gamma \varphi_t^{\text{GP}} \right). \end{aligned} \quad (3.9)$$

By (3.4), we get

$$\begin{aligned} \sup_{t \in [t_0, t_0 + T]} \|\partial^\alpha \varphi_t^{\text{GP}}\|_{L^2(\mathbb{R}^3)} &\leq \|\partial^\alpha \varphi_{t_0}^{\text{GP}}\|_{L^2(\mathbb{R}^3)} \\ &+ g_N \sup_{t \in [t_0, t_0 + T]} \left\| \int_{t_0}^t ds e^{i(t-s)\Delta} \sum_{\gamma \leq \beta \leq \alpha} \binom{\alpha}{\beta} \binom{\beta}{\gamma} \left(\partial^{(\alpha-\beta)} \overline{\varphi_t^{\text{GP}}} \partial^{(\beta-\gamma)} \varphi_t^{\text{GP}} \partial^\gamma \varphi_t^{\text{GP}} \right) \right\|_{L^2(\mathbb{R}^3)} \\ &\leq \|\partial^\alpha \varphi_{t_0}^{\text{GP}}\|_{L^2(\mathbb{R}^3)} + 64g_N \sqrt{T} \sum_{\gamma \leq \beta \leq \alpha} \sup_{t \in [t_0, t_0 + T]} \left\| \partial^{(\alpha-\beta)} \varphi_t^{\text{GP}} \partial^{(\beta-\gamma)} \varphi_t^{\text{GP}} \partial^\gamma \varphi_t^{\text{GP}} \right\|_{L^{\frac{6}{5}}(\mathbb{R}^3)}. \end{aligned} \quad (3.10)$$

As in [4, Proposition 3.1], via Hölder inequality and Sobolev embeddings, we get that there exists a constant C (which depends only on the Sobolev embeddings) such that

$$\sup_{t \in [t_0, t_0 + T]} \|\partial^\alpha \varphi_t^{\text{GP}}\|_{L^2(\mathbb{R}^3)} \leq \|\partial^\alpha \varphi_{t_0}^{\text{GP}}\|_{L^2(\mathbb{R}^3)} + Cg_N \sqrt{T} \sup_{t \in [t_0, t_0 + T]} \|\varphi_t^{\text{GP}}\|_{H^1(\mathbb{R}^3)}^2 \|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)}. \quad (3.11)$$

This means, that up to increasing C and using (3.1), we get that for any $t \in [t_0, t_0 + T]$,

$$\|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)} \leq \|\varphi_{t_0}^{\text{GP}}\|_{H^2(\mathbb{R}^3)} + CEg_N \sqrt{T} \sup_{t \in [t_0, t_0 + T]} \|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)},$$

where we denoted by $E \equiv \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)}$. We can now fix T small enough so that $CEg_N \sqrt{T} = 1/2$, as a consequence we get for any $t \in [t_0, t_0 + T]$

$$\|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)} \leq 2 \|\varphi_{t_0}^{\text{GP}}\|_{H^2(\mathbb{R}^3)}. \quad (3.12)$$

Repeating the argument, we get that for any $t > 0$

$$\|\varphi_{t+T}^{\text{GP}}\|_{H^2(\mathbb{R}^3)} \leq 2 \|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)}. \quad (3.13)$$

Fix now $t > 0$ and let $k \in \mathbb{N}$ such that $t \in ((k-1)T, kT]$; then

$$\|\varphi_t^{\text{GP}}\|_{H^2(\mathbb{R}^3)} \leq 2 \|\varphi_{(k-1)T}^{\text{GP}}\|_{H^2(\mathbb{R}^3)} \leq 2^k \|\varphi_0\|_{H^2(\mathbb{R}^3)} \leq 2^{\frac{k}{T}+1} \|\varphi_0\|_{H^2(\mathbb{R}^3)} = 2^{4C^2 E^2 g_N^2 t+1} \|\varphi_0\|_{H^2(\mathbb{R}^3)}, \quad (3.14)$$

which implies (3.3).

The proof for (3.2) can be done in the same way. Proceeding as above, one gets

$$\begin{aligned} \sup_{t \in [t_0, t_0 + T]} \|\partial^\alpha \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)} &\leq \|\partial^\alpha \varphi_{t_0}^{\text{H}}\|_{L^2(\mathbb{R}^3)} \\ &+ g_N \sqrt{T} \sum_{\gamma \leq \beta \leq \alpha} \binom{\alpha}{\beta} \binom{\beta}{\gamma} \sup_{t \in [t_0, t_0 + T]} \left\| v_N * \left(\partial^{(\alpha-\beta)} \varphi_t^{\text{H}} \partial^{(\beta-\gamma)} \overline{\varphi_t^{\text{H}}} \right) \partial^\gamma \varphi_t^{\text{H}} \right\|_{L^{\frac{6}{5}}(\mathbb{R}^3)}. \end{aligned} \quad (3.15)$$

The only difference with respect to before is that to bound the interaction term one has to use Young's inequality for the convolution (see [37, Theorem 4.2]). \square

3.2. From the many-body problem to Hartree. In this section we want to compare the many-body problem with the Hartree approximation, the main result is in the proposition below.

Proposition 3.2 (From Many-Body to Hartree). *Let $\beta \in (0, \frac{1}{6})$ and $\lambda \in (3\beta, 1 - 3\beta)$. Let ψ_t and φ_t^{H} be the normalized solution of (1.25) and (1.31) with initial data ψ_0 and φ_0 , respectively. Suppose that v satisfies Assumption 1. We have*

$$\left\| |\varphi_t^{\text{H}} \rangle \langle \varphi_t^{\text{H}}| - \gamma_{\psi_t}^{(1)} \right\| \leq \sqrt{2} \left(N^{\frac{1-\lambda}{2}} \left\| |\varphi_0 \rangle \langle \varphi_0| - \gamma_{\psi_0}^{(1)} \right\|^{\frac{1}{2}} + N^{\frac{3\beta}{2}} \right) e^{C_v (C_N(\varphi_0, t) + C_N(\varphi_0, t)^2) g_N |t|}, \quad (3.16)$$

where C_v is a constant depending on the L^1 and the L^2 norms of v , and $C_N(\varphi_0, t)$ is given by

$$C_N(\varphi_0, t) = \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{Cg_N^2 \left(\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4 \right) |t|} \quad (3.17)$$

The key idea for the proof of Proposition 3.2 is to control the number of *bad particles* in the many-body system (i.e. the particles not in the state φ_t^H), we do that using methods introduced in [47].

We now fix the notation for the projectors on the spaces of good and bad particles. We collect and prove the main general properties of these operators in Appendix A.

Definition 3.1. Let $\varphi \in L^2(\mathbb{R}^3)$ and $\psi \in L^2_s(\mathbb{R}^{3N})$.

- (1) For any $1 \leq j \leq N$, the projectors $p_j^\varphi : L^2(\mathbb{R}^{3N}) \rightarrow L^2(\mathbb{R}^{3N})$ represent the probability of the j -th particle of the state ψ of being in the state φ , i.e.

$$\begin{aligned} p_j^\varphi \psi(x_1, \dots, x_N) &:= (|\varphi\rangle\langle\varphi|)_j \psi(x_1, \dots, x_N) \\ &\equiv \varphi(x_j) \int_{\mathbb{R}^3} dz \varphi^*(z) \psi(x_1, \dots, x_{j-1}, z, x_{j+1}, \dots, x_N). \end{aligned} \quad (3.18)$$

Analogously, $q_j^\varphi := 1 - p_j^\varphi$ represents the probability of the j -th particle of the state ψ not being in the state φ .

- (2) For any $0 \leq k \leq j \leq N$ we set

$$\mathcal{A}_k^j := \left\{ a := (a_1, a_2, \dots, a_j) \in \{0, 1\}^j : \sum_{l=1}^j a_l = k \right\} \quad (3.19)$$

and define the orthogonal projector $P_{j,k}^\varphi$ on $L^2(\mathbb{R}^{3N})$ as

$$P_{j,k}^\varphi := \sum_{a \in \mathcal{A}_k^j} \prod_{l=N-j+1}^N \left[(p_l^\varphi)^{1-a_l} (q_l^\varphi)^{a_l} \right]. \quad (3.20)$$

Note that the rank of $P_{j,k}^\varphi$ is the space of states that in the last j particles have exactly k that are not in the state φ (we sum over \mathcal{A}_k^j to make the state symmetric).

- (3) For any function $f : \{0, \dots, N\} \rightarrow \mathbb{R}$ we define the operator $\hat{f}^\varphi : L^2(\mathbb{R}^{3N}) \rightarrow L^2(\mathbb{R}^{3N})$ as

$$\hat{f}^\varphi := \sum_{k \in \mathbb{Z}} f(k) P_k^\varphi. \quad (3.21)$$

where $f(k) = 0$ for any $k \in \mathbb{Z} \setminus \{0, \dots, N\}$

Similarly, if $f_d(k) := f(k+d)$ for any $d, k \in \mathbb{Z}$, we define the operator $\hat{f}_d^\varphi : L^2(\mathbb{R}^{3N}) \rightarrow L^2(\mathbb{R}^{3N})$ associated to f_d as

$$\hat{f}_d^\varphi := \sum_{k \in \mathbb{Z}} f(k+d) P_k^\varphi. \quad (3.22)$$

Throughout the following Section, the hat $\hat{\cdot}$ will solely be used in the sense of Definition 3.1.

Following [47], we now introduce a functional which encodes the fraction of condensation within itself.

Definition 3.2. For any $\lambda \in (0, 1)$ we define the weight function $\mu^\lambda : \{0, \dots, N\} \rightarrow \mathbb{R}$ as

$$\mu^\lambda(k) := \begin{cases} \frac{k}{N^\lambda}, & \text{for } k \leq N^\lambda \\ 1, & \text{otherwise.} \end{cases} \quad (3.23)$$

Moreover, for any $N \in \mathbb{N}$, we define the functional $\alpha_N^\lambda : L^2_s(\mathbb{R}^{3N}) \times L^2(\mathbb{R}^{3N}) \rightarrow \mathbb{R}$ as⁴

$$\alpha_N^\lambda(\psi, \varphi) := \langle \psi, \hat{\mu}^{\lambda, \varphi} \psi \rangle = \left\| (\hat{\mu}^{\lambda, \varphi})^{1/2} \psi \right\|_{L^2_s(\mathbb{R}^{3N})}^2. \quad (3.24)$$

The proof of Proposition 3.2 is based on some Grönwall estimate on the quantity α_N^λ , which we prove in the next proposition.

⁴For ease of notation we will write $\hat{\mu}^{\lambda, \varphi}$ instead of $\widehat{\mu^\lambda}^\varphi$.

Proposition 3.3 (Grönwall estimate on α_N^λ). *Let $\beta \in (0, \frac{1}{6})$ and $\lambda \in (3\beta, 1 - 3\beta)$. Let ψ_t and φ_t^H be the normalized solution of (1.25) and (1.31) with initial data ψ_0 and φ_0 , respectively. Suppose that v satisfies Assumption 1; we then have*

$$\alpha_N^\lambda(\psi_t, \varphi_t^H) \leq \left(\alpha_N^\lambda(\psi_0, \varphi_0) + \frac{1}{N^{\lambda-3\beta}} \right) e^{C_v(C_N(\varphi_0, t) + C_N(\varphi_0, t)^2)g_N|t|}, \quad (3.25)$$

where C_v is a constant depending on the L^1 and the L^2 norms of v , and $C_N(\varphi_0, t)$ is given by

$$C_N(\varphi_0, t) = \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{C g_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4)} |t| \quad (3.26)$$

3.2.1. *Proof of Proposition 3.3.* To prove Proposition 3.3 we need some preliminary results which are stated in Lemma 3.1 and in Proposition 3.4. In the proof of these two intermediate results, we use some operator estimates proven in Appendix A. To simplify notation we will use the following definition.

Definition 3.3. We denote by $U_{j,k}$ the difference between the pair interaction and the mean-field interactions for two particles j and k , i.e.

$$U_{j,k} := (N-1)v_N(x_j - x_k) - Nv_N * |\varphi_t^H|^2(x_j) - Nv_N * |\varphi_t^H|^2(x_k). \quad (3.27)$$

Lemma 3.1. *Let $\lambda \in (0, 1)$. Let ψ_t and φ_t^H be the normalized solution of (1.25) and (1.31), respectively. Then we have,*

$$\partial_t \alpha_N^\lambda(\psi_t, \varphi_t^H) = \Gamma_N^\lambda(\psi_t, \varphi_t^H), \quad (3.28)$$

where $\Gamma_N^\lambda : L_s^2(\mathbb{R}^{3N}) \times L^2(\mathbb{R}^3) \rightarrow \mathbb{R}$ is defined as

$$\begin{aligned} \Gamma_N^\lambda(\psi, \varphi) := & 2g_N \text{Im} \left(\left\langle \psi, \left(\hat{\mu}^{\lambda, \varphi} - \hat{\mu}_1^{\lambda, \varphi} \right) p_1^\varphi p_2^\varphi U_{1,2} p_1^\varphi q_2^\varphi \psi \right\rangle \right) \\ & + g_N \text{Im} \left(\left\langle \psi, \left(\hat{\mu}^{\lambda, \varphi} - \hat{\mu}_2^{\lambda, \varphi} \right) p_1^\varphi p_2^\varphi U_{1,2} q_1^\varphi q_2^\varphi \psi \right\rangle \right) \\ & + 2g_N \text{Im} \left(\left\langle \psi, \left(\hat{\mu}^{\lambda, \varphi} - \hat{\mu}_1^{\lambda, \varphi} \right) p_1^\varphi q_2^\varphi U_{1,2} q_1^\varphi q_2^\varphi \psi \right\rangle \right). \end{aligned} \quad (3.29)$$

Proof. First notice that for any $1 \leq j \leq N$ if we differentiate $p_j^{\varphi_t^H}$ we get

$$\partial_t p_j^{\varphi_t^H} = (|\varphi_t^H \rangle \langle \partial_t \varphi_t^H|)_j + (|\partial_t \varphi_t^H \rangle \langle \varphi_t^H|)_j = i \left[p_j^{\varphi_t^H}, -\Delta_j + g_N v_N * |\varphi_t^H|^2(x_j) \right], \quad (3.30)$$

and similarly for $q_j^{\varphi_t^H}$

$$\partial_t q_j^{\varphi_t^H} = i \left[q_j^{\varphi_t^H}, -\Delta_j + g_N v_N * |\varphi_t^H|^2(x_j) \right]. \quad (3.31)$$

It is then useful to introduce H_H^φ defined as

$$H_H^\varphi := \sum_{j=1}^N \left[-\Delta_j + g_N v_N * |\varphi|^2(x_j) \right] \quad (3.32)$$

As a consequence, for any $1 \leq j \leq N$, we can write

$$\partial_t p_j^{\varphi_t^H} = i \left[p_j^{\varphi_t^H}, H_H^\varphi \right], \quad \partial_t q_j^{\varphi_t^H} = i \left[q_j^{\varphi_t^H}, H_H^\varphi \right]. \quad (3.33)$$

Given that for any $0 \leq k \leq j \leq N$ the operator $P_{j,k}^{\varphi_t^H}$ is just a linear combination of $p_i^{\varphi_t^H}$'s and $q_i^{\varphi_t^H}$'s, we deduce that for any weight $f : \{0, \dots, N\} \rightarrow \mathbb{R}^+$ we get

$$\partial_t f^{\varphi_t^H} = i \left[\partial_t f^{\varphi_t^H}, H_H^\varphi \right]. \quad (3.34)$$

In order to simplify the notation, for the rest of the proof we will drop the labels φ_t^H and λ in the formulas involving the weights and the effective many-body nonlinear operator. If now we differentiate $\partial_t \alpha_N^\lambda(\psi_t, \varphi_t^H)$, using (3.34) and the symmetry of ψ_t and $\hat{\mu}\psi_t$, we get

$$\partial_t \alpha_N^\lambda(\psi_t, \varphi_t^H) = -i \langle \psi_t, [\hat{\mu}, H_N - H_H] \psi_t \rangle = -i g_N \langle \psi_t, [\hat{\mu}, U_{1,2}] \psi_t \rangle = g_N \text{Im} \langle \langle \psi_t, \hat{\mu} U_{1,2} \psi_t \rangle \rangle. \quad (3.35)$$

We want to now decompose the action of the operator $\widehat{\mu} U_{1,2}$ in terms of p_1, p_2, q_1 and q_2 . To do so, recall that $P_{j,k}$ projects on the space of states in which in the last j particles there are exactly k particles that are not in the state φ_t^H ; we can then write

$$P_k \equiv P_{N,k} = p_1 p_2 P_{N-2,k} + (p_1 q_2 + q_1 p_2) P_{N-2,k-1} + q_1 q_2 P_{N-2,k-2}.$$

As a consequence, for any weight $f : \{0, \dots, N\} \rightarrow \mathbb{R}$ we can use the equality above to get

$$\begin{aligned} \widehat{f} &= \sum_{k \in \mathbb{Z}} f(k) P_k = \sum_{k \in \mathbb{Z}} f(k) (p_1 p_2 P_{N-2,k} + (p_1 q_2 + q_1 p_2) P_{N-2,k-1} + P_{N-2,k-2}) \\ &\quad - p_1 p_2 \sum_{k \in \mathbb{Z}} f(k+2) P_{N-2,k} - (p_1 q_2 + q_1 p_2) \sum_{k \in \mathbb{Z}} f(k+1) P_{N-2,k-1} \\ &= p_1 p_2 (\widehat{f} - \widehat{f}_2) + (p_1 q_2 + q_1 p_2) (\widehat{f} - \widehat{f}_1) + \sum_{k \in \mathbb{Z}} f(k) P_{N-2,k-2}. \end{aligned} \quad (3.36)$$

We can then insert (3.36) in (3.35) and we get

$$\begin{aligned} \partial_t \alpha_N^\lambda (\psi_t, \varphi_t^H) &= g_N \text{Im} (\langle \psi_t, (p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) + 2p_1 q_2 (\widehat{\mu} - \widehat{\mu}_1)) U_{1,2} \psi_t \rangle) \\ &= \sum_{\substack{\#_1 = p_1, q_1 \\ \#_2 = p_2, q_2}} g_N \text{Im} (\langle \psi_t, p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) U_{1,2} \#_1 \#_2 \psi_t \rangle) + g_N \text{Im} (\langle \psi_t, p_1 q_2 (\widehat{\mu} - \widehat{\mu}_1) U_{1,2} \#_1 \#_2 \psi_t \rangle), \end{aligned} \quad (3.37)$$

where we used that the last term in (3.36) is self-adjoint and commutes with $U_{1,2}$. It turns out that many of those terms are either identical or vanishing. In particular, we can use (A.3) to have that $p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) U_{1,2} p_1 p_2 = (p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) U_{1,2} p_1 p_2)^\dagger$, which implies that $g_N \text{Im} (\langle \psi_t, p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) U_{1,2} p_1 p_2 \psi_t \rangle) = 0$. The same holds true for the operators $p_1 q_2 (\widehat{\mu} - \widehat{\mu}_1) U_{1,2} p_1 q_2$ and $p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) U_{1,2} p_1 q_2$. Using again (A.3) we then get

$$\begin{aligned} \partial_t \alpha_N^\lambda (\psi_t, \varphi_t^H) &= 2g_N \text{Im} (\langle \psi_t, p_1 p_2 (\widehat{\mu} - \widehat{\mu}_1) U_{1,2} p_1 q_2 \psi_t \rangle) \\ &\quad + g_N \text{Im} (\langle \psi_t, p_1 p_2 (\widehat{\mu} - \widehat{\mu}_2) U_{1,2} q_1 q_2 \psi_t \rangle) \\ &\quad + 2g_N \text{Im} (\langle \psi_t, p_1 q_2 (\widehat{\mu} - \widehat{\mu}_1) U_{1,2} q_1 q_2 \psi_t \rangle), \end{aligned} \quad (3.38)$$

which concludes the proof. \square

The next step is to estimate separately the three terms appearing in $\Gamma_N^\lambda (\psi_t, \varphi_t^H)$ in terms of $\alpha_N^\lambda (\psi_t, \varphi_t^H)$ in order to prove a Grönwall-type estimate for $\alpha_N^\lambda (\psi_t, \varphi_t^H)$. This is the aim of the next Proposition.

Proposition 3.4. *Let $\lambda \in (0, 1)$, let ψ_t and φ_t be normalized solutions of (1.25) and (1.31), respectively. The following estimates hold true.*

$$\left| \text{Im} \left(\langle \psi_t, (\widehat{\mu}_1^{\lambda, \varphi_t^H} - \widehat{\mu}^{\lambda, \varphi_t^H}) p_1^{\varphi_t^H} p_2^{\varphi_t^H} U_{1,2} p_1^{\varphi_t^H} q_2^{\varphi_t^H} \psi_t \rangle \right) \right| \leq \frac{\|v\|_{L^1(\mathbb{R}^3)}}{N^{\frac{1+\lambda}{2}}} \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^{\frac{4}{3}} \sqrt{\alpha_N^\lambda (\psi_t, \varphi_t^H)}, \quad (3.39)$$

$$\left| \text{Im} \left(\langle \psi_t, (\widehat{\mu}_2^{\lambda, \varphi_t^H} - \widehat{\mu}^{\lambda, \varphi_t^H}) p_1^{\varphi_t^H} p_2^{\varphi_t^H} U_{1,2} q_1^{\varphi_t^H} q_2^{\varphi_t^H} \psi_t \rangle \right) \right| \leq \quad (3.40)$$

$$\leq 2 \max \left\{ \|v\|_{L^1(\mathbb{R})}, \|v\|_{L^2(\mathbb{R})} \right\} \max \left\{ \|\varphi_t^H\|_{L^4(\mathbb{R}^3)}, \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \right\}^2 \left(\alpha_N^\lambda (\psi_t, \varphi_t^H) + \frac{N^{3\beta-\lambda}}{2} \right), \quad (3.41)$$

$$\left| \text{Im} \left(\langle \psi_t, (\widehat{\mu}_1^{\lambda, \varphi_t^H} - \widehat{\mu}^{\lambda, \varphi_t^H}) p_1^{\varphi_t^H} q_2^{\varphi_t^H} U_{1,2} q_1^{\varphi_t^H} q_2^{\varphi_t^H} \psi_t \rangle \right) \right| \leq \quad (3.42)$$

$$\leq \frac{2 \max \left\{ \|v\|_{L^1(\mathbb{R}^3)}, \|v\|_{L^2(\mathbb{R}^3)} \right\}}{N^{\frac{1-\lambda}{2}}} \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \left(N^{\frac{3\beta}{2}} + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \right) \alpha_N^\lambda (\psi_t, \varphi_t^H). \quad (3.43)$$

Proof. In order to simplify the notation, during the proof we shall drop the labels λ and φ_t^H for short. To start we note that for any function we can rewrite the operator $p_1 v_N (x_1 - x_2) p_1$ in the following, more

convenient form:

$$p_1 v_N (x_1 - x_2) p_1 = (|\varphi_t^H \rangle \langle \varphi_t^H|)_1 v_N (x_1 - x_2) (|\varphi_t^H \rangle \langle \varphi_t^H|)_1 = (v_N * |\varphi_t^H|^2) (x_2) p_1. \quad (3.44)$$

Proof of (3.39). Using the fact $p_j q_j = q_j p_j = 0$ for any $1 \leq j \leq N$, by (3.44), we can rewrite

$$\begin{aligned} p_1 p_2 U_{1,2} p_1 q_2 &= p_1 p_2 [(N-1) v_N (x_1 - x_2) - N v_N * |\varphi_t^H|^2 (x_1) - N v_N * |\varphi_t^H|^2 (x_2)] p_1 q_2 \\ &= -N p_1 p_2 v_N * |\varphi_t^H|^2 (x_2) p_1 q_2 \\ &= -p_1 p_2 v_N * |\varphi_t^H|^2 (x_2) p_1 q_2. \end{aligned} \quad (3.45)$$

As a consequence, using (A.2) and (A.3), we are able to bound the left-hand term in (3.39) as

$$\begin{aligned} |\operatorname{Im} \langle \psi_t, (\hat{\mu}_1 - \hat{\mu}) p_1 p_2 v_N * |\varphi_t^H|^2 (x_2) p_1 q_2 \psi_t \rangle | &\leq \|v_N * |\varphi_t^H|^2 (x_2) p_2\| \|(\hat{\mu} - \hat{\mu}_{-1}) q_2 \psi_t\| \\ &\leq \|v_N * |\varphi_t^H|^2 (x_2) p_2\| \|(\hat{\mu} - \hat{\mu}_{-1}) \hat{\nu} \psi_t\|. \end{aligned} \quad (3.46)$$

We then bound separately these last two terms. To bound the first term, one can easily show that

$$\begin{aligned} \|v_N * |\varphi_t^H|^2 (x_2) p_2\|^2 &\leq \langle \varphi_t^H, (v_N * |\varphi_t^H|^2 (x_2))^2 \varphi_t^H \rangle \leq \|\varphi_t^H\|_{L^6(\mathbb{R}^3)}^2 \|v_N * |\varphi_t^H|^2 (x_2)\|_{L^3(\mathbb{R}^3)}^2 \\ &\leq \|\varphi_t^H\|_{L^6(\mathbb{R}^3)}^4 \|v\|_{L^1(\mathbb{R}^3)}^2 \leq \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^{\frac{4}{3}} \|v\|_{L^1(\mathbb{R}^3)}, \end{aligned} \quad (3.47)$$

To bound the second term we want to use the explicit definition of μ . In particular, we have that $|\mu(k) - \mu_{-1}(k)| \leq CN^\lambda$. Moreover, for any $k \leq N^\lambda$, we have

$$\left| (\mu(k) - \mu_{-1}(k))^2 \nu^2(k) \right| \leq \frac{k}{N^{1+2\lambda}} = \frac{1}{N^{1+\lambda}} \mu(k). \quad (3.48)$$

Given that $\mu(k) - \mu_{-1}(k) = 0$ if $k \geq N^\lambda + 1$, we get that

$$\|(\hat{\mu} - \hat{\mu}_{-1}) \hat{\nu} \psi_t\| = \left\langle \psi_t, (\hat{\mu} - \hat{\mu}_{-1})^2 \hat{\nu}^2 \psi_t \right\rangle^{\frac{1}{2}} \leq \frac{1}{N^{\frac{1+\lambda}{2}}} \langle \psi_t, \hat{\mu} \psi_t \rangle^{\frac{1}{2}} = \frac{1}{N^{\frac{1+\lambda}{2}}} \sqrt{\alpha_N^\lambda (\psi_t, \varphi_t^H)}. \quad (3.49)$$

The estimates (3.47) and (3.49) imply (3.39).

Proof of (3.40). To prove (3.40), we first look directly at the operator $p_1 p_2 U_{1,2} q_1 q_2$. We have

$$p_1 p_2 U_{1,2} q_1 q_2 = (N-1) p_1 p_2 v_N (x_1 - x_2) q_1 q_2. \quad (3.50)$$

We act similarly as before. In order to do so, notice first that the function $\mu_2 - \mu$ is positive and correspond still to a weight, so we can consider the square root of the operator $\hat{\mu}_2 - \hat{\mu}$ and using (A.3), we get

$$\begin{aligned} |\operatorname{Im} \langle \psi_t, (\hat{\mu}_2^{\lambda, \varphi_t^H} - \hat{\mu}^{\lambda, \varphi_t^H}) p_1^{\varphi_t^H} p_2^{\varphi_t^H} U_{1,2} q_1^{\varphi_t^H} q_2^{\varphi_t^H} \psi_t \rangle | &\leq (N-1) \left| \left\langle \psi_t, (\hat{\mu}_2 - \hat{\mu})^{\frac{1}{2}} p_1 p_2 v_N (x_1 - x_2) q_1 q_2 (\hat{\mu} - \hat{\mu}_{-2})^{\frac{1}{2}} \psi_t \right\rangle \right|. \end{aligned}$$

We now use that $v_N (x_1 - x_2)$ is nonzero only in a small region where $x_1 \approx x_2$, because of the quick decay of v_N . To exploit this fact is then convenient to symmetrize $(N-1) v_N (x_1 - x_2)$ replacing it with $(N-1) \sum_{k=2}^N v_N (x_1 - x_k)$ to get, using Cauchy-Schwarz,

$$\begin{aligned} (N-1) \left| \left\langle \psi_t, (\hat{\mu}_2 - \hat{\mu})^{\frac{1}{2}} p_1 p_2 v_N (x_1 - x_2) q_1 q_2 (\hat{\mu} - \hat{\mu}_{-2})^{\frac{1}{2}} \psi_t \right\rangle \right| &\leq \left\| \sum_{j=2}^N q_j v_N (x_1 - x_j) p_1 p_j (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\| \left\| (\hat{\mu} - \hat{\mu}_{-2})^{1/2} q_1 \psi_t \right\|. \end{aligned} \quad (3.51)$$

We now bound those terms separately. For the second term, we can again write explicitly the difference $\hat{\mu} - \hat{\mu}_{-2}$:

$$\hat{\mu}(k) - \hat{\mu}_{-2}(k) = \begin{cases} \frac{2}{N^\lambda} & \text{for } k \leq N^\lambda, \\ \frac{N^\lambda + 2 - k}{N^\lambda} & \text{for } N^\lambda < k \leq N^\lambda + 2, \\ 0 & \text{otherwise.} \end{cases} \quad (3.52)$$

and by (A.2) we then get

$$\left\| (\hat{\mu} - \hat{\mu}_{-2})^{1/2} q_1 \psi_t \right\| = \langle \psi_t, (\hat{\mu} - \hat{\mu}_{-2}) \hat{\nu}^2 \psi_t \rangle^{1/2} \leq \sqrt{\frac{2}{N}} \langle \psi_t, \hat{\mu} \psi_t \rangle^{1/2} = \sqrt{\frac{2}{N}} \sqrt{\alpha_N^\lambda (\psi_t, \varphi_t^H)}. \quad (3.53)$$

We now take into account the first term in the r.h.s of (3.51). Explicitely, it is

$$\begin{aligned} & \left\| \sum_{j=2}^N q_j v_N (x_1 - x_j) p_1 p_j (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\|^2 \\ &= \sum_{j=2}^N \sum_{\ell=2}^N \left\langle \psi_t, (\hat{\mu}_2 - \hat{\mu})^{1/2} p_1 p_j v_N (x_1 - x_j) q_j q_\ell v_N (x_1 - x_\ell) p_1 p_\ell (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\rangle, \end{aligned} \quad (3.54)$$

and now we bound it as a sum of two different terms, one in which $j = \ell$ and one for the case $j \neq \ell$. For the case $j = \ell$, by (A.9) and (A.10), we can write

$$\begin{aligned} & N^2 \left\langle \psi_t, (\hat{\mu}_2 - \hat{\mu})^{1/2} p_1 p_2 v_N (x_1 - x_2) q_2 q_3 v_N (x_1 - x_3) p_1 p_3 (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\rangle \\ & \leq N^2 \left\| \sqrt{|v_N (x_1 - x_2)|} \sqrt{|v_N (x_1 - x_3)|} p_1 p_3 (\hat{\mu}_2 - \hat{\mu})^{1/2} q_2 \psi_t \right\|^2 \\ & \leq N^2 \left\| \sqrt{|v_N (x_1 - x_2)|} p_2 \right\|^4 \left\| (\hat{\mu}_2 - \hat{\mu})^{1/2} q_2 \psi_t \right\|^2. \end{aligned}$$

Moreover, by (A.10), we get

$$\left\| \sqrt{|v_N (x_1 - x_2)|} p_2 \right\|^4 = \|p_2 |v_N (x_1 - x_2)| p_2\|^2 \leq \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^4 \|v\|_{L^1(\mathbb{R}^3)}^2,$$

and on the other hand proceeding similarly as in (3.53) we can estimate

$$\left\| (\hat{\mu}_2 - \hat{\mu})^{1/2} q_2 \psi_t \right\|^2 \leq \frac{2}{N} \alpha_N^\lambda (\psi_t, \varphi_t^H). \quad (3.55)$$

We then get

$$\begin{aligned} & \left| N^2 \left\langle \psi_t, (\hat{\mu}_2 - \hat{\mu})^{1/2} p_1 p_2 v_N (x_1 - x_2) q_2 q_3 v_N (x_1 - x_3) p_1 p_3 (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\rangle \right| \\ & \leq 2N \|v\|_{L^1(\mathbb{R}^3)}^2 \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^4 \alpha_N^\lambda (\psi_t, \varphi_t^H). \end{aligned} \quad (3.56)$$

We then look at the second term in the right hand side of (3.56): for this term we cannot bring any q close to the state ψ_t , and is therefore not possible to exploit the state ψ_t . We then bound every operator in it directly in norm using (A.11), we get

$$\begin{aligned} & N \left\langle \psi_t, (\hat{\mu}_2 - \hat{\mu})^{1/2} p_1 p_2 v_N (x_1 - x_2) q_2 v_N (x_1 - x_2) p_1 p_2 (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\rangle \\ & \leq N \|v_N (x_1 - x_2) p_1 p_2\|^2 \left\| (\hat{\mu}_2 - \hat{\mu})^{1/2} \right\|^2 \leq 2N^{1+3\beta-\lambda} \|\varphi_t^H\|_{L^4(\mathbb{R}^3)}^4 \|v\|_{L^2(\mathbb{R}^3)}^2. \end{aligned} \quad (3.57)$$

Combining (3.56) and (3.57), we can conclude that

$$\begin{aligned} & \left\| \sum_{j=2}^N q_j v_N (x_1 - x_j) p_1 p_j (\hat{\mu}_2 - \hat{\mu})^{1/2} \psi_t \right\| \\ & \leq \sqrt{2N \|v\|_{L^1(\mathbb{R}^3)}^2 \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^4 \alpha_N^\lambda (\psi_t, \varphi_t^H) + 2N^{1+3\beta-\lambda} \|\varphi_t^H\|_{L^4(\mathbb{R}^3)}^4 \|v\|_{L^2(\mathbb{R}^3)}^2} \end{aligned}$$

Moreover, the estimates (3.53) and (3.58), imply (3.40).

Proof of (3.42). Similarly as before, we note that

$$p_1 q_2 U_{1,2} q_1 q_2 = p_1 q_2 \left((N-1) v_N (x_1 - x_2) - N v_N * |\varphi_t^H|^2 (x_1) - N v_N * |\varphi_t^H|^2 (x_2) \right) q_1 q_2 \quad (3.58)$$

$$= p_1 q_2 \left((N-1) v_N (x_1 - x_2) - N v_N * |\varphi_t^H|^2 (x_1) \right) q_1 q_2. \quad (3.59)$$

We start the proof of (3.42) by writing

$$\begin{aligned} \left| \operatorname{Im} \left(\left\langle \psi_t, \left(\widehat{\mu}_1^{\lambda, \varphi_t^H} - \widehat{\mu}^{\lambda, \varphi_t^H} \right) p_1^{\varphi_t^H} q_2^{\varphi_t^H} U_{1,2} q_1^{\varphi_t^H} q_2^{\varphi_t^H} \psi_t \right\rangle \right) \right| &\leq \left\| (\widehat{\mu}_1 - \widehat{\mu})^{\frac{1}{2}} q_2 \psi_t \right\| \|U_{1,2} p_1\| \left\| (\widehat{\mu} - \widehat{\mu}_{-1})^{1/2} q_1 q_2 \psi_t \right\| \\ &\leq \sqrt{\frac{N}{N-1}} \left\| (\widehat{\mu}_1 - \widehat{\mu})^{\frac{1}{2}} \widehat{v} \psi_t \right\| \|U_{1,2} p_1\| \left\| (\widehat{\mu} - \widehat{\mu}_{-1})^{\frac{1}{2}} \widehat{v}^2 \psi_t \right\|. \end{aligned} \quad (3.60)$$

Proceeding as above we can estimate directly $(\widehat{\mu}_1 - \widehat{\mu}) \widehat{v}^2 \leq \frac{1}{N} \widehat{\mu}$ and $(\widehat{\mu} - \widehat{\mu}_{-1}) \widehat{v}^4 \leq \frac{1}{N^{2-\lambda}} \widehat{\mu}$. We now bound the potential term. We have, using (A.9),

$$\|U_{1,2} p_1\| \leq N \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \max \left\{ \|v\|_{L^1(\mathbb{R}^3)}, \|v\|_{L^2(\mathbb{R}^3)} \right\} \left(N^{\frac{3\beta}{2}} + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \right), \quad (3.61)$$

Inserting (3.61) in (3.60), we get (3.42). \square

Combining Lemma 3.1 and Proposition 3.4, we can prove Proposition 3.3.

Proof of Proposition 3.3. As stated before, the key idea of the proof is to exploit a Grönwall-type argument. In order to do that, we use Lemma 3.1 and Proposition 3.4 to write

$$\begin{aligned} \partial_t \alpha_N^\lambda(\psi_t, \varphi_t^H) &= \Gamma_N^\lambda(\psi_t, \varphi_t^H) \\ &\leq C_v g_N \left(\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \left[1 + \frac{1}{N^{\frac{1-\lambda-3\beta}{2}}} \right] + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^2 \left[1 + \frac{1}{N^{\frac{1-\lambda}{2}}} \right] \right) \alpha_N^\lambda(\psi_t, \varphi_t^H) \\ &\quad + C_v g_N \frac{\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^{\frac{4}{3}}}{N^{\frac{1+\lambda}{2}}} \sqrt{\alpha_N^\lambda(\psi_t, \varphi_t^H)} + \frac{C_v g_N}{N^{\lambda-3\beta}} \left(\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^2 \right). \end{aligned} \quad (3.62)$$

where C_v is a constant that depends only on the L^1 and L^2 norms of v . We can now do Cauchy-Schwarz in the next to the last term and discard some terms which are subleading for any $\lambda \in (0, 1 - 3\beta)$, to get

$$\begin{aligned} \partial_t \alpha_N^\lambda(\psi_t, \varphi_t^H) &\leq C_v g_N \left(\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^2 \right) \alpha_N^\lambda(\psi_t, \varphi_t^H) \\ &\quad + \frac{C_v g_N}{N^{1+\lambda}} \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^{\frac{5}{3}} + \frac{C_v g_N}{N^{\lambda-3\beta}} \left(\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^2 \right). \end{aligned} \quad (3.63)$$

Being $\lambda \in (3\beta, 1 - 3\beta)$, we get

$$\partial_t \alpha_N^\lambda(\psi_t, \varphi_t^H) \leq C_v g_N \left(\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} + \|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}^2 \right) \left(\alpha_N^\lambda(\psi_t, \varphi_t^H) + \frac{1}{N^{\lambda-3\beta}} \right) \quad (3.64)$$

To bound $\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)}$, we use Proposition 3.1. We find

$$\|\varphi_t^H\|_{L^\infty(\mathbb{R}^3)} \leq \|\varphi_t^H\|_{H^2(\mathbb{R}^3)} \leq \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{C g_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4) |t|} = C_N(\varphi_0, t). \quad (3.65)$$

By Grönwall, we can then conclude that

$$\alpha_N^\lambda(\psi_t, \varphi_t^H) \leq \left(\alpha_N^\lambda(\psi_0, \varphi_0) + \frac{1}{N^{\lambda-3\beta}} \right) e^{C_v (C_N(\varphi_0, t) + C_N(\varphi_0, t)^2) g_N |t|}. \quad (3.66)$$

\square

3.2.2. Proof of Proposition 3.2. The proof of Proposition 3.2 is splitted in two parts:

- First, we prove that

$$\left\| |\varphi_t^H \rangle \langle \varphi_t^H| - \gamma_{\psi_t}^{(1)} \right\|^2 \leq 2 \alpha_N^\lambda(\psi_t, \varphi_t^H) \quad (3.67)$$

- Secondly, we prove that for some $\lambda \in (0, 1)$, for any $\psi \in L_s^2(\mathbb{R}^{3N})$ with $\|\psi\|_{L_s^2(\mathbb{R}^{3N})} = 1$ and for any normalized $\varphi \in L^2(\mathbb{R}^3)$, one has

$$\alpha_N^\lambda(\psi, \varphi) \leq N^{1-\lambda} \left\| |\varphi \rangle \langle \varphi| - \gamma_\psi^{(1)} \right\|. \quad (3.68)$$

Combining the estimates in (3.67) and in (3.68) with the result in Proposition 3.3 we get the result.

Proof of Proposition 3.2. We start by proving (3.67). Being $|\varphi_t^H\rangle\langle\varphi_t^H| - \gamma_{\psi_t}^{(1)} \in \mathfrak{S}_2(L^2(\mathbb{R}^3))$, we have,

$$\left\| |\varphi_t^H\rangle\langle\varphi_t^H| - \gamma_{\psi_t}^{(1)} \right\| \leq \left\| |\varphi_t^H\rangle\langle\varphi_t^H| - \gamma_{\psi_t}^{(1)} \right\|_{\mathfrak{S}_2(L^2(\mathbb{R}^3))}. \quad (3.69)$$

Moreover, using (A.2), we can write

$$\begin{aligned} \left\| |\varphi_t^H\rangle\langle\varphi_t^H| - \gamma_{\psi_t}^{(1)} \right\|_{\mathfrak{S}_2(L^2(\mathbb{R}^3))}^2 &\leq 2 \left(1 - \text{tr} \left[|\varphi_t^H\rangle\langle\varphi_t^H| \gamma_{\psi_t}^{(1)} \right] \right) = 2 \left(1 - \left\| p_1^{\varphi_t^H} \psi_t \right\|_{L_s^2(\mathbb{R}^{3N})}^2 \right) \\ &= 2 \left\| q_1^{\varphi_t^H} \psi_t \right\|_{L_s^2(\mathbb{R}^{3N})}^2 = 2 \left\| \widehat{\mathcal{V}}^{\varphi_t^H} \psi_t \right\|_{L_s^2(\mathbb{R}^{3N})}^2 \end{aligned} \quad (3.70)$$

Given that $\nu^2 \leq \mu^\lambda$ we can write

$$\left\| |\varphi_t^H\rangle\langle\varphi_t^H| - \gamma_{\psi_t}^{(1)} \right\| \leq \sqrt{2} \left\| \widehat{\mathcal{V}}^{\varphi_t^H} \psi_t \right\|_{L_s^2(\mathbb{R}^{3N})} \leq \sqrt{2\alpha_N^\lambda(\psi_t, \varphi_t^H)} \quad (3.71)$$

Consider now $\alpha_N^\lambda(\psi, \varphi) = \langle \psi | \widehat{\mu}^{\lambda, \varphi} | \psi \rangle$, we have

$$\alpha_N^\lambda(\psi, \varphi) \leq N^{1-\lambda} \langle \psi, (\widehat{\mathcal{V}}^\varphi)^2 \psi \rangle = N^{1-\lambda} \|q_1^\varphi \psi\|_{L_s^2(\mathbb{R}^{3N})}^2 \leq N^{1-\lambda} \left\| |\varphi\rangle\langle\varphi| - \gamma_\psi^{(1)} \right\|, \quad (3.72)$$

where we used that $\mu^\lambda(k) \leq N^{1-\lambda}\nu^2$ and where the last inequality follows from the definition of q_1^φ . If we now apply (3.68) to $\psi_0 \in L^2(\mathbb{R}^{3N})$ and $\varphi \in L^2(\mathbb{R}^3)$, by and (3.67) and Proposition 3.3, we get

$$\left\| |\varphi_t^H\rangle\langle\varphi_t^H| - \gamma_{\psi_t}^{(1)} \right\|^2 \leq 2 \left(N^{1-\lambda} \left\| |\varphi_0\rangle\langle\varphi_0| - \gamma_{\psi_0}^{(1)} \right\| + \frac{1}{N^{\lambda-3\beta}} \right) e^{C_v(C_N(\varphi_0, t) + C_N(\varphi_0, t)^2)g_N|t|} \quad (3.73)$$

□

3.3. From Hartree to Gross-Pitaevskii. The main goal of this section is to prove the following proposition.

Proposition 3.5 (From Hartree to Gross-Pitaevskii). *Let $\beta \in (0, 1/3)$. Let φ_t^{GP} and φ_t^{H} be the normalized solution of (1.30) and (1.31), respectively, both with initial data $\varphi_0 \in L^2(\mathbb{R}^3)$. Suppose that v satisfies Assumption 1. We have*

$$\|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)} \leq C\sqrt{g_N} \frac{1 + \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N N^{-\beta} \|\varphi_0\|_{L^\infty}^2}}{N^{\frac{\beta}{2}}} e^{C_v C_N(\varphi_0, t)^2 g_N |t|}, \quad (3.74)$$

where C_v is a constant which depends only on v and $C_N(\varphi_0, t)$ is given by

$$C_N(\varphi_0, t) = \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{C g_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4) |t|} \quad (3.75)$$

Proof. We prove (3.74) using Grönwall estimate. To do that we want to bound the following quantity

$$\begin{aligned} \partial_t \|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2 &= 2\text{Re} \langle \varphi_t^{\text{GP}} - \varphi_t^{\text{H}}, \partial_t (\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}) \rangle = 2g_N \text{Im} \langle \varphi_t^{\text{H}}, (|\varphi_t^{\text{H}}|^2 - |\varphi_t^{\text{GP}}|^2) \varphi_t^{\text{GP}} \rangle \\ &\quad + 2g_N \text{Im} \langle \varphi_t^{\text{GP}}, (|\varphi_t^{\text{H}}|^2 - v_N * |\varphi_t^{\text{H}}|^2) \varphi_t^{\text{H}} \rangle. \end{aligned}$$

We now estimate the two quantities above separately. For the first one, we have

$$\begin{aligned} |2g_N \text{Im} \langle \varphi_t^{\text{H}}, (|\varphi_t^{\text{H}}|^2 - |\varphi_t^{\text{GP}}|^2) \varphi_t^{\text{GP}} \rangle| &\leq C g_N \|\varphi_t^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} (\|\varphi_t^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} + \|\varphi_t^{\text{H}}\|_{L^\infty(\mathbb{R}^3)}) \|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2 \\ &\leq C g_N (\|\varphi_t^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}^2 + \|\varphi_t^{\text{H}}\|_{L^\infty(\mathbb{R}^3)}^2) \|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2 \end{aligned} \quad (3.76)$$

We now estimate the second one, we have

$$\begin{aligned}
& |2g_N \text{Im} \langle \varphi_t^{\text{GP}}, (|\varphi_t^{\text{H}}|^2 - v_N * |\varphi_t^{\text{H}}|^2) \varphi_t^{\text{H}} \rangle| \\
& \leq 2g_N \int_{\mathbb{R}^3} dx |\varphi_t^{\text{GP}}(x)| |\varphi_t^{\text{H}}(x)| \int_{\mathbb{R}^3} dy v(y) \left(|\varphi_t^{\text{H}}(x)|^2 - \left| \varphi_t^{\text{H}} \left(x - \frac{y}{N^\beta} \right) \right|^2 \right) \\
& = 2g_N \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy v(y) |\varphi_t^{\text{GP}}(x)| |\varphi_t^{\text{H}}(x)| \int_0^1 d\lambda \frac{d}{d\lambda} \left| \varphi_t^{\text{H}} \left(x - \frac{\lambda y}{N^\beta} \right) \right|^2 \\
& \leq 2 \frac{g_N}{N^\beta} \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \int_0^1 d\lambda |y| v(y) |\varphi_t^{\text{GP}}(x)| |\varphi_t^{\text{H}}(x)| \left| \varphi_t^{\text{H}} \left(x - \frac{\lambda y}{N^\beta} \right) \right| \left| \nabla \varphi_t^{\text{H}} \left(x - \frac{\lambda y}{N^\beta} \right) \right|^2 \\
& \leq C \frac{g_N}{N^\beta} \|xv\|_{L^1(\mathbb{R}^3)} \|\varphi_t^{\text{H}}\|_{L^\infty(\mathbb{R}^3)}^2 (1 + \|\nabla \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2)
\end{aligned} \tag{3.77}$$

Then we find

$$\partial_t \|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2 \leq Cg_N \left(\|\varphi_t^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)}^2 + \|\varphi_t^{\text{H}}\|_{L^\infty(\mathbb{R}^3)}^2 \right) \left(\|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2 + \frac{1 + \|\nabla \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)}^2}{N^\beta} \right) \tag{3.79}$$

We can bound uniformly in time the L^∞ norms of φ_t^{GP} , φ_t^{H} , using the bound for the H^2 norms proved in Proposition 3.1, i.e.,

$$\|\varphi_t^{\text{GP}}\|_{L^\infty(\mathbb{R}^3)} \leq \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{Cg_N^2 \mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)|t|} \tag{3.80}$$

$$\|\varphi_t^{\text{H}}\|_{L^\infty(\mathbb{R}^3)} \leq C \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{Cg_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4) |t|} \tag{3.81}$$

Moreover, by Proposition 3.1, we also can bound uniformly in time

$$\|\nabla \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)} \leq C \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)} + Cg_N N^{-\beta} \|\varphi_0\|_{L^\infty}^2 \tag{3.82}$$

Thus, by Grönwall, we find

$$\|\varphi_t^{\text{GP}} - \varphi_t^{\text{H}}\|_{L^2(\mathbb{R}^3)} \leq C \sqrt{g_N} \frac{1 + \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)} + g_N N^{-\beta} \|\varphi_0\|_{L^\infty}^2}{N^{\frac{\beta}{2}}} e^{C_v C_N(\varphi_0, t)^2 g_N |t|}, \tag{3.83}$$

where C_v is a constant which depends on $\|xv\|_{L^1(\mathbb{R}^3)}$ and $C(\varphi_0)$ is defined as in (3.65), i.e.,

$$C_N(\varphi_0, t) = \|\varphi_0\|_{H^2(\mathbb{R}^3)} e^{Cg_N^2 (\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N^2 N^{-2\beta} \|\varphi_0\|_{L^\infty(\mathbb{R}^3)}^4) |t|} \tag{3.84}$$

□

3.4. Proof of Theorem 1.6. In this last section we combine the results of Proposition 3.3 and Proposition 3.5 to prove (1.34). We start by writing

$$\|\gamma_{\psi_t}^{(1)} - P_{\varphi_t^{\text{GP}}}\| \leq \|\gamma_{\psi_t}^{(1)} - P_{\varphi_t^{\text{H}}}\| + \|P_{\varphi_t^{\text{H}}} - P_{\varphi_t^{\text{GP}}}\| \leq \|\gamma_{\psi_t}^{(1)} - P_{\varphi_t^{\text{H}}}\| + 2 \|\varphi_t^{\text{H}} - \varphi_t^{\text{GP}}\|_{L^2(\mathbb{R}^3)}. \tag{3.85}$$

By Proposition 3.2, we get

$$\|\gamma_{\psi_t}^{(1)} - P_{\varphi_t^{\text{H}}}\| \leq \sqrt{2} \left(N^{\frac{1-\lambda}{2}} \|\gamma_{\psi_0}^{(1)} - P_{\varphi_0}\|^{\frac{1}{2}} + \frac{1}{N^{\frac{\lambda-3\beta}{2}}} \right) e^{C_v (C_N(\varphi_0, t) + C_N(\varphi_0, t)^2) g_N |t|}. \tag{3.86}$$

On the other hand, from Proposition 3.5 we have

$$2 \|\varphi_t^{\text{H}} - \varphi_t^{\text{GP}}\|_{L^2(\mathbb{R}^3)} \leq C \sqrt{g_N} \frac{1 + \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0)} + g_N N^{-\beta} \|\varphi_0\|_{L^\infty}^2}{N^{\frac{\beta}{2}}} e^{C_v C_N(\varphi_0, t)^2 g_N |t|}. \tag{3.87}$$

Combining the two previous results we then get

$$\begin{aligned} \left\| \gamma_{\psi_t}^{(1)} - P_{\varphi_t^{\text{GP}}} \right\| &\leq \sqrt{2} \left(N^{\frac{1-\lambda}{2}} \left\| \gamma_{\psi_0}^{(1)} - P_{\varphi_0} \right\|^{\frac{1}{2}} + \frac{1}{N^{\frac{\lambda-3\beta}{2}}} \right) e^{C_v(C_N(\varphi_0,t) + C_N(\varphi_0,t)^2)g_N|t|} \\ &\quad + C\sqrt{g_N} \frac{1 + \sqrt{\mathcal{E}_{\text{GP}}^{\text{free}}(\varphi_0) + g_N N^{-\beta}} \|\varphi_0\|_{L^\infty}^2}{N^{\frac{\beta}{2}}} e^{C_v C_N(\varphi_0,t)^2 g_N |t|}, \end{aligned} \quad (3.88)$$

where C_v now is a constant which depends on $\|v\|_{L^1(\mathbb{R}^3)}$, $\|v\|_{L^2(\mathbb{R}^3)}$ and $\|xv\|_{L^1(\mathbb{R}^3)}$.

APPENDIX A. PROPERTIES OF THE PROJECTORS p_j^φ , q_j^φ

Proposition A.1. *Let p_j^φ and q_j^φ be defined as in (3.18).*

(i) *Let $\nu : \{0, 1, \dots, N\} \rightarrow \mathbb{R}$ be given by $\nu(k) := \sqrt{\frac{k}{N}}$. Then the square of $\widehat{\nu}^\varphi$ is the fraction of particles not in the state φ , i.e.*

$$(\widehat{\nu}^\varphi)^2 = \frac{1}{N} \sum_{j=1}^N q_j^\varphi. \quad (A.1)$$

(ii) *For any $f : \{0, 1, \dots, N\} \rightarrow \mathbb{R}$ and any symmetric state $\psi \in L_s^2(\mathbb{R}^{3N})$*

$$\left\| \widehat{f}^\varphi q_1^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)} = \left\| \widehat{f}^\varphi \widehat{\nu}^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)}, \quad \left\| \widehat{f}^\varphi q_1^\varphi q_2^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)} \leq \sqrt{\frac{N}{N-1}} \left\| \widehat{f}^\varphi (\widehat{\nu}^\varphi)^2 \psi \right\|_{L_s^2(\mathbb{R}^3)}. \quad (A.2)$$

(iii) *For any $f : \{0, 1, \dots, N\} \rightarrow \mathbb{R}$, $v : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$ and $j, k = 0, 1, 2$, we have*

$$\widehat{f}^\varphi Q_j^\varphi v(x_1, x_2) Q_k^\varphi = Q_j^\varphi v(x_1, x_2) Q_k^\varphi \widehat{f}_{j-k}^\varphi, \quad (A.3)$$

where $Q_0^\varphi := p_1^\varphi p_2^\varphi$, $Q_1^\varphi := p_1^\varphi q_2^\varphi$ and $Q_2^\varphi := q_1^\varphi q_2^\varphi$.

Proof. To prove A.1 note that $\bigcup_{k=0}^N \mathcal{A}_k = \{0, 1\}^N$, so that $1 = \sum_{k=0}^N P_k^\varphi$. Using also $(q_j^\varphi)^2 = q_j^\varphi$ and $q_j^\varphi p_j^\varphi = 0$, we get

$$q_j^\varphi \prod_{l=1}^N \left[(p_l^\varphi)^{1-a_l} (q_l^\varphi)^{a_l} \right] = \begin{cases} \prod_{l=1}^N \left[(p_l^\varphi)^{1-a_l} (q_l^\varphi)^{a_l} \right] & \text{if } a_j = 1, \\ 0. & \text{else} \end{cases} = a_j \prod_{l=1}^N \left[(p_l^\varphi)^{1-a_l} (q_l^\varphi)^{a_l} \right]. \quad (A.4)$$

Given that for any $\mathbf{a} \in \mathcal{A}_k$ the sum of the a_j 's is k we deduce that

$$\frac{1}{N} \sum_{j=1}^N q_j^\varphi = \frac{1}{N} \sum_{j=1}^N q_j^\varphi \sum_{k=0}^N P_k^\varphi = \frac{1}{N} \sum_{k=0}^N \sum_{j=1}^N q_j^\varphi P_k^\varphi = \frac{1}{N} \sum_{k=0}^N k P_k^\varphi = \sum_{k=0}^N (\nu(k))^2 P_k^\varphi \quad (A.5)$$

and A.1 follows.

We now prove (A.2). We can use the symmetry of ψ and the previous point to get

$$\left\| \widehat{f}^\varphi \widehat{\nu}^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)}^2 = \left\langle \psi, \left(\widehat{f}^\varphi \right)^2 (\widehat{\nu}^\varphi)^2 \psi \right\rangle = \left\langle \psi, \left(\widehat{f}^\varphi \right)^2 q_1^\varphi \psi \right\rangle = \left\| \widehat{f}^\varphi q_1^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)}^2. \quad (A.6)$$

Similarly, for the second inequality in (A.2), we have

$$\left\| \widehat{f}^\varphi (\widehat{\nu}^\varphi)^2 \psi \right\|_{L_s^2(\mathbb{R}^3)}^2 = \frac{1}{N^2} \sum_{j,k=1}^N \left\langle \psi, \left(\widehat{f}^\varphi \right)^2 q_j^\varphi q_k^\varphi \psi \right\rangle = \frac{N-1}{N} \left\| \widehat{f}^\varphi q_1^\varphi q_2^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)}^2 + \frac{1}{N} \left\| \widehat{f}^\varphi \widehat{\nu}^\varphi \psi \right\|_{L_s^2(\mathbb{R}^3)}^2. \quad (A.7)$$

To prove (A.3) we use the definition of the Q_j 's and the previous points to get

$$\begin{aligned} \widehat{f}^\varphi Q_j^\varphi v(x_1, x_2) Q_k^\varphi &= \sum_{l \in \mathbb{Z}} f(l) P_l^\varphi Q_j^\varphi v(x_1, x_2) Q_k^\varphi = \sum_{l \in \mathbb{Z}} f(l) Q_j^\varphi v(x_1, x_2) Q_k^\varphi P_{N-2,l-j}^\varphi \\ &= \sum_{l \in \mathbb{Z}} f(l) Q_j^\varphi v(x_1, x_2) Q_k^\varphi P_{l-j+k}^\varphi = Q_j^\varphi v(x_1, x_2) Q_k^\varphi \widehat{f}_{j-k}^\varphi. \end{aligned} \quad (A.8)$$

□

Proposition A.2. *Let p_j^φ and q_j^φ be defined as in (3.18). Let $a, a' \in [1, +\infty]$ such that $1/a + 1/a' = 1$ and let $u \in L^{2a'}(\mathbb{R}^3)$, $\varphi \in L^{2a}(\mathbb{R}^3)$. It holds true that*

$$\|u(x_1 - x_2)p_1^\varphi\| \leq \|\varphi\|_{L^{2a}(\mathbb{R}^3)} \|u\|_{L^{2a'}(\mathbb{R}^3)}, \quad (\text{A.9})$$

$$\|p_1^\varphi u(x_1 - x_2)p_1^\varphi\| \leq \|\varphi\|_{L^{2a}(\mathbb{R}^3)}^2 \|u\|_{L^{a'}(\mathbb{R}^3)}. \quad (\text{A.10})$$

Moreover, if $a \in [1, 2]$, one has

$$\|u(x_1 - x_2)p_1^\varphi p_2^\varphi\| \leq \|\varphi\|_{L^{2a}(\mathbb{R}^3)}^2 \|u\|_{L^{a'}(\mathbb{R}^3)}. \quad (\text{A.11})$$

Proof. The estimate (A.9) directly follows from Hölder inequality,

$$\|u(x_1 - x_2)p_1^\varphi\|^2 = \sup_{\|\psi\|=1} \|u(x_1 - x_2)p_1^\varphi\psi\|^2 \leq \sup_{\|\psi\|=1} \|\psi\| \sup_{x_2 \in \mathbb{R}^3} \langle \varphi, u^2(\cdot - x_2)\varphi \rangle \leq \|\varphi\|_{L^{2a}(\mathbb{R}^3)}^2 \|u\|_{L^{2a'}(\mathbb{R}^3)}^2. \quad (\text{A.12})$$

To prove (A.10), we use (A.9). Indeed,

$$\|p_1^\varphi u(x_1 - x_2)p_1^\varphi\| \leq \left\| \sqrt{|u(x_1 - x_2)|} p_1^\varphi \right\|^2 \leq \|\varphi\|_{L^{2a}(\mathbb{R}^3)}^2 \|\sqrt{|u|}\|_{L^{2a'}(\mathbb{R}^3)}^2 \leq \|\varphi\|_{L^{2a}(\mathbb{R}^3)}^2 \|u\|_{L^{a'}(\mathbb{R}^3)}. \quad (\text{A.13})$$

Finally, the estimate (A.11) is a consequence of the Young inequality for the convolution (see [37, Theorem 4.2]). More precisely, we have

$$\|u(x_1 - x_2)p_1 p_2\|^2 = \sup_{\|\psi\|=1} \langle \psi, (|\varphi \times \varphi|)_1 (|\varphi \times \varphi|)_2 u^2(x_1 - x_2) (|\varphi \times \varphi|)_1 (|\varphi \times \varphi|)_2 \psi \rangle \quad (\text{A.14})$$

$$\leq \int_{\mathbb{R}^6} dx_1 dx_2 u^2(x_1 - x_2) |\varphi(x_1)|^2 |\varphi(x_2)|^2 = \left\langle |\varphi|^2, u^2 * |\varphi|^2 \right\rangle \quad (\text{A.15})$$

$$\leq \|\varphi\|_{L^{2p}(\mathbb{R}^3)}^2 \|\varphi\|_{L^{2q}(\mathbb{R}^3)}^2 \|u\|_{L^{2r}(\mathbb{R}^3)}^2, \quad (\text{A.16})$$

□

REFERENCES

- [1] Adams, R. A., Fournier, J. J. F. *Sobolev Spaces*, Elsevier (2003).
- [2] Adhikari A., Brennecke C., Schlein B., *Bose-Einstein Condensation Beyond the Gross-Pitaevskii Regime*, *Ann. Henri Poincaré* (2020).
- [3] Basti G., Cenatiempo S., Schlein B., *A new second order upper bound for the ground state energy of dilute Bose gases*, to appear in *Forum Math. Sigma*.
- [4] Benedikter N., de Oliveira G., Schlein B., *Quantitative Derivation of the Gross-Pitaevskii Equation*, *Communications on Pure and Applied Mathematics* 68.8 (2015), pp. 1399–1482.
- [5] Boccato C., Brennecke C., Cenatiempo S., Schlein S., *The excitation spectrum of Bose gases interacting through singular potentials*, *Eur. Math. Soc.*, **22**, 2331–2403 (2020).
- [6] Boccato C., Brennecke C., Cenatiempo S., Schlein B., *Bogoliubov Theory in the Gross-Pitaevskii Limit*, *Acta Mathematica* **222** (2019), no. 2, 219335.
- [7] Boccato, C., Brennecke, C., Cenatiempo, S., Schlein, B., *Optimal Rate for Bose-Einstein Condensation in the Gross-Pitaevskii Regime* *Comm. Math. Phys.* (2019).
- [8] Boccato, C., Cenatiempo S., Schlein, B., *Quantum Many-Body Fluctuations Around Nonlinear Schrödinger Dynamics*, *Annales Henri Poincaré* 18:113, 2017.
- [9] Bossmann, L., Petrat, S., Seiringer, R., *Asymptotic expansion of low-energy excitations for weakly interacting bosons*, *Forum of Mathematics, Sigma*, 9, E28, 2021.
- [10] Bossmann, L., Pavlović, N., Pickl, P., Soffer, A., *Higher Order Corrections to the Mean-Field Description of the Dynamics of Interacting Bosons*, *J Stat Phys* 178, 1362–1396 (2020).
- [11] Brennecke C., Caporaletti M., Schlein B., *Excitation Spectrum for Bose Gases beyond the Gross-Pitaevskii Regime*, arxiv:2104.13003.
- [12] Brennecke, C., Nam, P. T., Napiórkowski, M., Schlein, B., *Fluctuations of N -particle quantum dynamics around the nonlinear Schrödinger equation*, *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 36(5), 1201–1235, 2019.
- [13] Brennecke, C., Schlein, B. *Gross-Pitaevskii dynamics for Bose-Einstein condensates*, *Anal. PDE*, **12(6)**, 1513–1596 (2019).
- [14] Brennecke, C., Schlein, B., Schraven, S., *Bose-Einstein Condensation with Optimal Rate for Trapped Bosons in the Gross-Pitaevskii Regime*, arXiv:2102.11052.

- [15] Brennecke, C., Schlein, B., Schraven, S., *Bogoliubov Theory for Trapped Bosons in the Gross-Pitaevskii Regime*, arXiv:2108.11129.
- [16] Bru, J.B., Correggi, M., Pickl, P., Yngvason, J., *The TF Limit for Rapidly Rotating Bose Gases in Anharmonic Traps*, Commun. Math. Phys. 280, 517–544 (2008).
- [17] Correggi M., Pinsky F., Rougerie N., Yngvason J., *Critical rotational speeds for superfluids in homogeneous traps*, Journal of Mathematical Physics 53.9 (Sept. 2012), p. 095203.
- [18] Correggi M., Rougerie N., Yngvason J., *The Transition to a Giant Vortex Phase in a Fast Rotating Bose–Einstein Condensate*, Commun. Math. Phys. 303 (2011), 451–608.
- [19] Dereziński J., Napiórkowski M., *Excitation Spectrum of Interacting Bosons in the Mean-Field Infinite-Volume Limit*, Annales Henri Poincaré 15 (2014), 2409–2439.
- [20] DiBenedetto E., *Partial Differential Equations*, Springer Science+Business Media New York, (1995).
- [21] Elgart, A., Schlein, B. *Mean field dynamics of boson stars*, Communications on Pure and Applied Mathematics, **60(4)**, 500–545(2007).
- [22] Erdős, L., Schlein, B., Yau, H.-T. *Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems*, Inventiones Mathematicae, **167(3)**, 515–614 (2007).
- [23] Erdős, L., Schlein, B., Yau, H.-T. *Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensate*, Annals of Mathematics, **172(1)**, 291–370 (2010).
- [24] Fetter, A.L., Foot, C.J., *Bose Gas: Theory and Experiment*, editors: K. Levin, A.L. Fetter, D.M. Stamper-Kurn, *Contemporary Concepts of Condensed Matter Science*, Elsevier, Volume 5, 2012, 27–67.
- [25] Fournais, S., Solovej, J. P., *The energy of dilute Bose gases*, Annals of Mathematics, Vol. 192 (2020), 893–976.
- [26] Fröhlich, J., Knowles, A., Schwarz, S. *On the Mean-Field Limit of Bosons with Coulomb Two-Body Interaction* Communications in Mathematical Physics, **288(3)**, 1023–1059 (2009).
- [27] Ginibre, J., Velo, G., *The classical field limit of scattering theory for non-relativistic many-boson systems I. and II.* Comm. Math. Phys. 66 (1979), 37–76, and 68 (1979), 45–68.
- [28] Grech P., Seiringer R., *The Excitation Spectrum for Weakly Interacting Bosons in a Trap*, Communications in Mathematical Physics 322.2, pp. 559–591, (2013).
- [29] Jerrard, R.L., Smets, D., *Vortex dynamics for the two dimensional non homogeneous Gross-Pitaevskii equation*, Annali Scuola Norm. Sup. Pisa, vol. 14, no. 3, 729–766, (2015).
- [30] Hainzl, C., *Another proof of BEC in the GP-limit*, Journal of Mathematical Physics 62, 051901 (2021).
- [31] Hepp., K., *The classical limit for quantum mechanical correlation functions*, Comm. Math. Phys. 35 (1974), 265–277.
- [32] Keel, M., Tao, T. *Endpoint Strichartz estimates*, American Journal of Mathematics, **120(5)**, 955–980 (1998).
- [33] Knowles, A., Pickl, P., *Mean-Field Dynamics: Singular Potentials and Rate of Convergence*, Commun. Math. Phys. 298, 101–138 (2010).
- [34] Lewin, M., Nam, P. T., Rougerie, N., *Derivation of Hartree’s theory for generic mean- field Bose gases*, Adv. Math. 254 (2014), pp. 570–621.
- [35] Lewin, M., Nam, P. T., Rougerie, N., *The mean-field approximation and the non- linear Schrödinger functional for trapped Bose gases*, Trans. Amer. Math. Soc. 368 (2016), no. 9, 6131–6157.
- [36] Lewin, M., Nam P.T., Serfaty S., Solovej J. P., *Bogoliubov spectrum of interacting Bose gases*, Comm. Pure Appl. Math. 68 (2014), 3, 413 – 471.
- [37] Lieb E. H., Loss M., *Analysis: Second Edition. American Mathematical Society AMS*, (2001).
- [38] Lieb E. H., Seiringer R., *Proof of Bose-Einstein Condensation for Dilute Trapped Gases*, Phys. Rev. Lett. 88, 170409 – Published 16 April 2002.
- [39] Lieb, E. H., Seiringer, R., *Derivation of the Gross-Pitaevskii equation for rotating Bose gases*, Communications In Mathematical Physics, 264 (2), 505–537, (2006).
- [40] Lieb, E. H., Seiringer R., Solovej J. P., Yngvason, J., *The mathematics of the Bose gas and its condensation*, Oberwolfach Seminars 34 (2005), Birkhäuser Basel.
- [41] Lieb E. H., Seiringer, R., Yngvason, J., *Bosons in a trap: A rigorous derivation of the Gross-Pitaevskii energy functional*, Phys. Rev. A 61 (2000), 043602.
- [42] Lieb E. L., Yngvason, J., *Ground State Energy of the low density Bose Gas*, Phys. Rev. Lett. 80 (1998), 25042507.
- [43] Nam, P:T., Napiórkowski, M., Ricaud J., Triay, A., *Optimal rate of condensation for trapped bosons in the Gross-Pitaevskii regime*, Preprint arXiv:2001.04364.
- [44] Nam P.T., Rougerie N., Seiringer R., *Ground states of large bosonic systems: the Gross-Pitaevskii limit revisited*, Anal. PDE 9.2, pp. 459–485, (2016)
- [45] Nam, P.T., Triay, A., *Bogoliubov excitation spectrum of trapped Bose gases in the Gross-Pitaevskii regime*, Preprint arXiv:2106.11949.
- [46] Olgiati A., Rougerie N., *The Hartree functional in a double well*, arXiv:2004.14729
- [47] Pickl P., *A Simple Derivation of Mean Field Limits for Quantum Systems*, Letters in Mathematical Physics 97.2, pp. 151–164, (2011).
- [48] Pickl P., *Derivation of the time dependent Gross-Pitaevskii equation with external fields*, Reviews in Mathematical Physics 27.01, p. 1550003, (2015).

- [49] Pizzo, A., *Bose particles in a box III. A convergent expansion of the ground state of the Hamiltonian in the mean field limiting regime*, arXiv:1511.07026 [math-ph], Nov. 2015.
- [50] Rathmair, M., *On how Poincaré inequalities imply weighted ones*, Monatsh Math 188, 753–763 (2019).
- [51] Rodnianski, I., Schlein, B. *Quantum Fluctuations and Rate of Convergence Towards Mean Field Dynamics* Communications in Mathematical Physics, **291**(1), 31–61 (2009).
- [52] Seiringer, R., *The Excitation Spectrum for Weakly Interacting Bosons*, Commun. Math. Phys. **306**, 565-578 (2011).
- [53] Spohn, H., *Kinetic equations from hamiltonian dynamics: Markovian limits*, Rev. Mod. Phys. 52 (1980), 569–615.
- [54] Yau H.-T., Yin J., *The second order upper bound for the ground state energy of a Bose gas*, J. Stat. Phys. 136(3) (2009), 453–503.

UNIVERSITÄT BASEL, SPIEGELGASSE 1, CH-4051 BASEL, SWITZERLAND.

Email address: `daniele@dimonte.it`

DEPARTMENT OF MATHEMATICS, LUDWIG-MAXIMILIANS UNIVERSITÄT MÜNCHEN, THERESIENSTR., 39, 80333, MÜNCHEN, GERMANY.

Email address: `emanuela.giacomelli@math.lmu.de`