Bless and curse of smoothness and phase transitions in nonparametric regressions: a nonasymptotic perspective^{*}

Ying Zhu[†]

Abstract

When the regression function belongs to the *standard* smooth classes consisting of univariate functions with derivatives up to the $(\gamma+1)$ th order bounded by a common constant everywhere or a.e., it is well known that the minimax optimal rate of convergence in mean squared error (MSE) is $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$ when γ is finite and the sample size $n \to \infty$. From a nonasymptotic viewpoint that considers finite n, this paper shows that: for the *standard* Hölder and Sobolev classes, the minimax optimal rate is $\frac{\sigma^2(\gamma \vee 1)}{n} \gtrsim \left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$ when $\frac{n}{\sigma^2} \precsim (\gamma \vee 1)^{2\gamma+3}$ and $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}} \approx \frac{1}{2\gamma+3}$ $\frac{\sigma^2(\gamma \vee 1)}{n}$) when $\frac{n}{\sigma^2} \succeq (\gamma \vee 1)^{2\gamma+3}$. To establish these results, we derive upper and lower bounds on the covering and packing numbers for the *generalized* Hölder class where the *k*th ($k = 0, ..., \gamma$) derivative is bounded from above by a parameter R_k and the γ th derivative is $R_{\gamma+1}$ -Lipschitz (and also for the *generalized* ellipsoid class of smooth functions). Our bounds sharpen the classical metric entropy results for the standard classes, and give the general dependence on γ and R_k . By deriving the minimax optimal MSE rates under $R_k = 1, R_k \leq (k-1)!$ and $R_k = k!$ (with the latter two cases motivated in our introduction below) for the smooth classes with the help of our new entropy bounds, we show a couple of interesting results that cannot be shown with the existing entropy bounds in the literature. We further consider the Hölder class of *d*-variate functions. Our result suggests that the classical asymptotic rate $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+2+d}}$ could

be an underestimate of the MSE in finite samples.

Introduction 1

Estimation of an unknown smooth function f from the nonparametric regression model

$$y_i = f(x_i) + \epsilon_i, \ i = 1, \dots, n \tag{1}$$

has been a central object of study in statistics, numerical analysis and machine learning. The typical assumption about f is that it belongs to the standard Hölder class (or the standard Sobolev class) consisting of univariate functions with derivatives up to the $(\gamma + 1)$ th order bounded by a common constant everywhere or a.e.. When γ is finite and the sample size $n \to \infty$, it is well understood that the minimax optimal rate of convergence in mean squared error (MSE) is $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$, where n is the sample size and σ^2 is the variance of the noise term ϵ_i (e.g., [11, 12]). This classical result gives arise to the so called "bless of smoothness" and empirical researchers are often advised to exploit higher

^{*}AMS subject classifications: 41A99, 62G08, 94D99. Keywords: Hölder and ellipsoid classes of smooth functions; covering and packing numbers; nonasymptotic analysis; phase transition; minimax lower bound

[†]Assistant Professor at UC San Diego. yiz012@ucsd.edu.

degree smoothness assumptions if they are facing a small sample size. This suggestion is particularly common in economic applications where researchers need to perform subsample analyses and in these applications (e.g., studies on intergenerational mobility¹), n often ranges from tens to a couple of hundreds.

If n is small enough, higher smoothness could bring a curse. To see this, recall that any function f in the Hölder class, U_{r+1} [-1, 1], can be written as

$$f(x) = f(0) + \sum_{k=1}^{\gamma} \frac{x^k}{k!} f^{(k)}(0) + \frac{x^{\gamma}}{\gamma!} f^{(\gamma)}(z) - \frac{x^{\gamma}}{\gamma!} f^{(\gamma)}(0)$$
(2)

where z is some intermediate value between x and 0. Therefore, we can decompose \mathcal{U}_{r+1} into a polynomial subspace $\mathcal{U}_{r+1,1}$ associated with $f(0) + \sum_{k=1}^{\gamma} \frac{x^k}{k!} f^{(k)}(0)$ and a Hölder subspace $\mathcal{U}_{r+1,2}$ with $f^{(k)}(0) = 0$ for all $k = 0, ..., \gamma$. A similar decomposition also applies to the Sobolev class $\overline{\mathcal{S}}_{\gamma+1}$; in particular, $\overline{\mathcal{S}}_{\gamma+1} = \mathcal{U}_{r+1,1} + \mathcal{S}_{\gamma+1} := \{f_1 + f_2 : f_1 \in \mathcal{U}_{r+1,1}, f_2 \in \mathcal{S}_{\gamma+1}\}$, where the Sobolev subspace $\mathcal{S}_{\gamma+1}$ is imposed with the restrictions that $f^{(k)}(0) = 0$ for all $k \leq \gamma$ and $f^{(\gamma+1)}$ belongs to the space \mathcal{L}^2 . ² Higher smoothness increases the difficulty of estimating the polynomial component while decreases the difficulty of estimating the component in $\mathcal{U}_{r+1,2}$ and $\mathcal{S}_{\gamma+1}$ (as long as the magnitude of the derivatives is not too large). Understanding this trade-off in finite sample settings can lead to useful guidance for empirical researchers.

Curse of smoothness can also arise in another form through the derivatives. In noisy recovery of solutions to ordinary differential equations (ODE), researchers often use polynomials and spline bases to approximate the solutions (e.g., [5, 6, 7, 8, 9, 14]) to overcome computational challenges. As an example from studies of AIDs, Liang and Wu [5] use local polynomial regressions to estimate the ODE solutions y and their first derivatives y' from noisy measurements of plasma viral load and CD4+ T cell counts; then, the authors regress the estimates \hat{y}' on $f(\hat{y}; \theta)$ to obtain estimates of the parameters θ in the ODE model. Liang and Wu [5] discuss about the possible use of higher degree local polynomials for approximating the solutions, and doing so would require boundedness on the higher order derivatives of the solutions. Motivated by these statistical procedures for recovering ODEs in the literature, Zhu and Mirzaei [19] study how the smoothness of ODEs affects the smoothness of the underlying solutions. To illustrate, let us consider the autonomous ODE y'(x) = f(y(x)). Like other areas in nonparametric estimation, it can be desirable to only assume smoothness structures on f for hedging against misspecification of the functional form for f. Zhu and Mirzaei [19] show that: (i) If $|f^{(k)}(x)| \leq c_0$ for all x on the domain and $k = 0, ..., \gamma + 1$, then $|y^{(k+1)}(x)| \leq c_0^{k+1}k!$; (ii) the factorial bounds are attainable by the solutions to some ODE (e.g., $y' = e^{-y - \frac{1}{2}}$ and therefore tight. To this phenomenon, [19] gives the name "loss of smoothness".

Motivated by the "loss of smoothness" phenomenon, this paper generalizes the standard Hölder class in the literature to one where the kth $(k = 0, ..., \gamma)$ derivative of any member is bounded from above by a parameter R_k and the γ th derivative is $R_{\gamma+1}$ -Lipschitz (i.e., R_k is allowed to depend on k). We also generalize the standard ellipsoid class of smooth functions in a similar fashion by allowing its RKHS (reproducing kernel Hilbert space) radius to be bounded from above by $R_{\gamma+1}$; that is,

$$\mathcal{H}_{\gamma+1} = \left\{ f = \sum_{m=1}^{\infty} \theta_m \phi_m : \text{for } (\theta_m)_{m=1}^{\infty} \in \ell^2(\mathbb{N}) \text{ such that } \sum_{m=1}^{\infty} \frac{\theta_m^2}{\mu_m} \le R_{\gamma+1}^2 \right\}$$
(3)

¹The author is grateful to Professor Esfandiar Maasoumi at Emory University for pointing out this motivation.

²See [15] (Chapter 1) and [16] (Examples 12.17 and 12.29). Other decompositions of the Sobolev class are also possible, see, e.g. [18].

where $\ell^2(\mathbb{N}) := \{(\theta_m)_{m=1}^{\infty} | \sum_{m=1}^{\infty} \theta_m^2 < \infty\}, (\mu_m)_{m=1}^{\infty} \text{ and } (\phi_m)_{m=1}^{\infty} \text{ are the eigenvalues and eigenfunctions (that forms an orthonormal basis of <math>\mathcal{L}^2[0, 1]$), respectively, of an RKHS with $\mu_m = (cm)^{-2(\gamma+1)}$ for some positive constant c. The decay rate of the eigenvalues follows the standard assumption for $(\gamma + 1)$ -degree smooth functions in the literature (see, e.g., [10, 16, 18]) and $R_{\gamma+1} = 1$ in (3) gives the standard ellipsoid class of smooth functions in [16]. Moreover, (3) is equipped with the inner product $\langle h, g \rangle_{\mathcal{H}} = \sum_{m=1}^{\infty} \frac{\langle h, \phi_m \rangle \langle g, \phi_m \rangle}{\mu_m}$ where $\langle \cdot, \cdot \rangle$ is the inner product in $\mathcal{L}^2[0, 1]$. The ellipsoid class is a generalization of the Sobolev class.

In our nonasymptotic framework, n, γ and $\{R_k\}_{k=0}^{\gamma+1}$ are all finite. The objectives of this paper are to examine the impacts of γ and $\{R_k\}_{k=0}^{\gamma+1}$ on (i) the size of the generalized Hölder and ellipsoid classes of smooth functions, and on (ii) the mean squared error (MSE) convergence rates associated with (1) in finite sample settings. To accomplish the first objective, we establish upper and lower bounds on the covering and packing numbers of the generalized $\mathcal{U}_{r+1,1}, \mathcal{U}_{r+1,2}$ and $\mathcal{H}_{\gamma+1}$; see Table 1 for a summary of the results. With the help of our new entropy bounds, we then derive information theoretic lower bounds³ and matching upper bounds for the MSE under various R_k ; see Tables 2–3 for a summary of the results. In what follows, let us discuss the novelty of our results.

The lower bound $\max \{\underline{B}_1(\delta), \underline{B}_2\}$ and the upper bound $\overline{B}_1(\delta)$ in Table 1 are original. The (less original) bounds $\overline{B}_2(\delta)$ and $R^{*\frac{1}{\gamma+1}} \delta^{\frac{-1}{\gamma+1}}$ generalize the upper bounds associated with $\mathcal{U}_{\gamma+1,1}$ and $\mathcal{U}_{\gamma+1,2}$, respectively, in [3]. It is worth pointing out that $\overline{B}_2(\delta)$ holds for all $\delta \in (0, 1)$ (not just δ such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} < 0$) but is far from being tight when $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} \ge 0$. Obviously $\overline{B}_1(\delta) \preceq \overline{B}_2(\delta)$. When it comes to deriving the upper bounds for the MSE under large enough R_k (such as $R_k = (k-1)!$ or $R_k = k!$), $\overline{B}_1(\delta)$ will be very useful. In particular, [19] applies the counting argument⁴ in [3] to derive an upper bound for the covering number of $\mathcal{U}_{\gamma+1,1}$ under $R_k = (k-1)!$ and then derive an upper bound for the MSE. With the new bound $\overline{B}_1(\delta)$ developed in this paper, the upper bound in Table 3 for the MSE improves the one in [19] by a factor of $\gamma \log \gamma$.

Note that our lower bound for the generalized $\mathcal{U}_{\gamma+1,1}$ in Table 1 takes the maximum of two terms. The part \underline{B}_2 (a constant bound valid for all δ below a threshold detailed in Section 2) is useful for deriving the minimax lower bounds on the MSE if $R_k \leq (k-1)!$ for $k = 1, ..., \gamma$, while $\underline{B}_1(\delta)$ will be useful if $R_k = k!$. Taking the maximum of the lower bounds for $\mathcal{U}_{\gamma+1,1}$ and $\mathcal{U}_{\gamma+1,2}$ gives a lower bound on the log(δ -packing number) of $\mathcal{U}_{\gamma+1}$. The lower bound for $\mathcal{U}_{\gamma+1}$ in [3] (derived under the assumption that $R_k \approx 1$) is $\delta^{\frac{-1}{\gamma+1}}$, which does not take into account the contribution from $\mathcal{U}_{\gamma+1,1}$. Therefore, our lower bound sharpens the classical result for the standard Hölder class.

Compared to bounds $\underline{B}_1(\delta)$ and $\overline{B}_1(\delta)$ in Table 1, \underline{B}_2 is relatively straightforward. To establish $\underline{B}_1(\delta)$ and $\overline{B}_1(\delta)$, we discard the counting argument in [3] and instead, consider two classes (equivalent to $\mathcal{U}_{\gamma+1,1}$), each in the form of a $(\gamma + 1)$ –dimensional polyhedron. The lower bound $\underline{B}_1(\delta)$ is the more delicate part. In particular, for any $f \in \mathcal{U}_{\gamma+1,1}$ [-1, 1], we write $f(x) = \sum_{k=0}^{\gamma} \tilde{\theta}_k \phi_k(x)$, where $(\phi_k)_{k=0}^{\gamma}$ are the Legendre polynomials. The key step is to derive sharp nonasymptotic lower bounds for $(\phi_k)_{k=0}^{\gamma}$. We achieve this goal by exploiting the Legendre expansion of x^k .

Let us turn to the covering and packing numbers of the generalized ellipsoid class of smooth

³Following the literature on minimax optimality, we consider the standard setup of (1) where $\{x_i\}_{i=1}^n$ are independently and uniformly distributed on a bounded interval, and $\{\epsilon_i\}_{i=1}^n$ are independent $\mathcal{N}(0, \sigma^2)$.

⁴To derive an upper bound for the δ -covering number of $\mathcal{U}_{\gamma+1}$ (= $\mathcal{U}_{\gamma+1,1} + \mathcal{U}_{\gamma+1,2}$) under the assumption $R_k \simeq 1$, the proof in Kolmogorov and Tikhomirov [3] considers a grid of points $(x_{-s}, ..., x_{-1}, x_0, x_1, ..., x_s)$ on [-1, 1] where $s \simeq \delta^{\frac{-1}{\gamma+1}}$; the bound $(\frac{\gamma}{2}+1)\log \frac{1}{\delta}$ in [3] is obtained by counting the number of possible values of $\left(\left\lfloor \frac{f^{(k)}(x_0)}{\delta_k} \right\rfloor\right)_{k=0}^{\gamma}$ given the δ_k -covering accuracy for the kth derivative (where $\delta_0 = \delta$).

functions, $\mathcal{H}_{\gamma+1}$. Under the assumption $R_{\gamma+1} = 1$, the upper and lower bounds in Wainwright [16] (the last two inequalities on p.131) scale as $(\gamma \vee 1) \delta^{\frac{-1}{\gamma+1}}$ and $\delta^{\frac{-1}{\gamma+1}}$, respectively, while our upper and lower bounds in Table 1 have the same scaling $\delta^{\frac{-1}{\gamma+1}}$. We close the gap in [16] by finding the optimal "pivotal" eigenvalue that best balances the "estimation error" and the "approximation error" from truncating for a given resolution δ . More generally, for the case of $R_{\gamma+1} \preceq \gamma + 1$, we consider two different truncations, one giving the upper bound $\delta^{\frac{-1}{\gamma+1}}$ and the other giving the lower bound $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$. Note that $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}} \asymp \delta^{\frac{-1}{\gamma+1}}$ when $R_{\gamma+1} \asymp 1$. For the case of $R_{\gamma+1} \succeq \gamma + 1$, we use only one truncation to show that both the upper bound and the lower bound scale as $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$.

For the Hölder and ellipsoid classes, Tables 2–3 suggest that the "bless of smoothness" arises in two ways: (i) from the nonasymptotic viewpoint, when $\mathcal{U}_{\gamma+1}$ is imposed with the restrictions that $f^{(k)}(0) = 0$ for all $k \leq \gamma$ (hence, $\mathcal{U}_{\gamma+1} = \mathcal{U}_{\gamma+1,2}$), $(R^*)^{\frac{2}{2\gamma+3}} \approx 1$ for $\mathcal{U}_{\gamma+1,2}$, and $R_{\gamma+1}^{\frac{2}{2\gamma+3}} \approx 1$ for $\mathcal{H}_{\gamma+1}$, Table 2 shows that the minimax optimal rate for the MSE is $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$, which decreases in γ ; (ii) when γ is finite and $n \to \infty$, Tables 2–3 show that the minimax optimal rate for the MSE is $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$. Another way to think about (ii) in terms of Table 3 is, if the resolution δ is small enough (because of large enough n), the size of $\mathcal{U}_{\gamma+1,2}$ dominates the size of $\mathcal{U}_{\gamma+1,1}$; hence, in deriving the upper bound and the minimax lower bound for the MSE, one may simply take $\log(\delta-\text{covering number}) \approx \log(\delta-\text{packing number}) \approx \delta^{\frac{-1}{\gamma+1}}$ under the standard assumption $R_k = 1$. This practice is common in the literature.

In general, however, when n is small enough, higher smoothness can be a curse. To illustrate, note from Table 3 that even when $R_k = 1$ (the standard assumption), if $\gamma > 1$ and $\frac{n}{\sigma^2} \preceq \gamma^{2\gamma+3}$, the minimax optimal rate is $\frac{\sigma^2 \gamma}{n}$, which is greater than $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$. Obviously, this result also holds true for the Sobolev class $\overline{S}_{\gamma+1} = \mathcal{U}_{r+1,1} + S_{\gamma+1}$ mentioned at the beginning. In view of this result, exploiting higher degree smoothness assumptions in nonparametric regressions when n is small may not be a good idea, unless the researchers believe that $\mathcal{U}_{\gamma+1,1}$ has some sparsity or approximate sparsity structures (e.g., $f^{(k)}(0) = 0$ for most of $k \in \{0, ..., \gamma\}$). In these cases, for instance, when estimating an unknown smooth function, researchers should consider regularizing the polynomial component in $\mathcal{U}_{\gamma+1,1}$ (e.g., with the l_1 -penalty) besides regularizing the component in $\mathcal{H}_{\gamma+1}$ with the typical Sobolev penalty.

A couple of interesting facts are revealed by Tables 2–3. First, when $R_k = (k-1)!$, Table 2 shows that the optimal rates differ between $\mathcal{U}_{r+1,2}$ and $\mathcal{H}_{\gamma+1}$. Note that this difference cannot be revealed in the regime where γ is fixed and $n \to \infty$, and by the classical entropy bounds. Second, Table 3 shows that, as the scaling of R_k is increased from (k-1)! to k!, the minimax optimal rate is increased from $\frac{\sigma^2 \gamma}{n}$ to $\frac{\sigma^2 \gamma \log \gamma}{n}$ when the component in $\mathcal{U}_{r+1,1}$ dominates. Once $\frac{n}{\sigma^2} \succeq \gamma^{2\gamma+3}$ in the case of $R_k = (k-1)!$, and $\frac{n}{\sigma^2} \succeq (\gamma \log \gamma)^{2\gamma+3}$ in the case of $R_k = k!$, the optimal rate becomes $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$ as now the component in $\mathcal{U}_{r+1,2}$ dominates. The terms $(k!)_{k=0}^{\gamma}$ in (2) play a more important role on the size of $\mathcal{U}_{r+1,1}$ when R_k becomes large enough, which is why $\overline{B}_1(\delta)$ and $\underline{B}_1(\delta)$ in Table 1 are very useful for deriving the minimax optimal rate for the MSE under large R_k .

We want to point out that the implications of the results in this paper are not limited to nonparametric regressions. Many semiparametric estimators in statistics involve nonparametric regressions as an intermediate step; in addition, estimations of generalized additive models, partially

Table 1: Upper and lower bounds on the log(δ – covering number) and log(δ – packing number) of the generalized $\mathcal{U}_{r+1,1}$, $\mathcal{U}_{r+1,2}$ and $\mathcal{H}_{\gamma+1}$ in L_q –norm

	$\mathcal{U}_{\gamma+1,1} \ (q \in \{2, \infty\})$	$\mathcal{U}_{\gamma+1,2} \ (q \in \{2, \infty\})$	$\mathcal{H}_{\gamma+1} \ (q=2)$
Y2	$\begin{cases} \overline{B}_1\left(\delta\right) & \text{if } \min_{k \in \{0,,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} \geq 0\\ \overline{B}_2\left(\delta\right) & \text{otherwise} \end{cases}$	$R^* \frac{1}{\gamma+1} \delta^{\frac{-1}{\gamma+1}}$	$\begin{cases} R_{\gamma+1}^{\frac{1}{\gamma+1}}\delta^{\frac{-1}{\gamma+1}} & \text{if } R_{\gamma+1} \succsim \gamma+1\\ \delta^{\frac{-1}{\gamma+1}} & \text{if } R_{\gamma+1} \precsim \gamma+1 \end{cases}$
ž	$\max\left\{ \underline{B}_{1}\left(\delta\right),\underline{B}_{2}\right\}$	$\begin{cases} R^* \frac{1}{\gamma+1} \delta^{\frac{-1}{\gamma+1}} & \text{if } R_0 \succeq 1\\ (R^* R_0)^{\frac{1}{\gamma+1}} \delta^{\frac{-1}{\gamma+1}} & \text{if } R_0 \precsim 1 \end{cases}$	$R_{\gamma+1}^{\frac{1}{\gamma+1}}\delta^{\frac{-1}{\gamma+1}}$

where: $\overline{B}_1(\delta) = \sum_{k=0}^{\gamma} \log \frac{4(\gamma+1)R_k}{k!\delta}; \quad \overline{B}_2(\delta) = \left(\frac{\gamma}{2}+1\right) \log \frac{1}{\delta} + \sum_{k=0}^{\gamma} \log R_k; \quad \underline{B}_1(\delta) = \sum_{k=0}^{\gamma} \log \left(9^{-\gamma}\gamma^{-\gamma}\right) + \sum_{k=0}^{\gamma} \log \frac{C\sum_{m=0}^{\lfloor \gamma/2 \rfloor} R_{k+2m}}{\delta} \quad \text{(with } R_{k+2m} = 0 \text{ for } k + 2m > \gamma); \quad \underline{B}_2 = C'\gamma \text{ (valid for all } \delta \text{ below a threshold detailed in Section 2); } R^{\dagger} = \left(\sum_{k=0}^{\gamma} \frac{R_k}{k!}\right) \lor 1; \quad R^* = \left(\max_{k \in \{1, \dots, \gamma+1\}} \frac{R_k}{(k-1)!}\right) \lor 1; \quad C \text{ and } C' \text{ are positive universal constants that are: } \lesssim 1 \text{ and independent of } n, \gamma, \text{ and } \{R_k\}_{k=0}^{\gamma+1}.$ Bounds on the log of the covering or packing numbers of $\mathcal{U}_{\gamma+1}$ follow easily from the bounds in Table 1.

Table 2: Minimax optimal MSE rates of the generalized $\mathcal{U}_{r+1,2}$ and $\mathcal{H}_{\gamma+1}$

	$\mathcal{U}_{r+1,2}$	$\mathcal{H}_{\gamma+1}$
MSE	$\asymp (R^*)^{\frac{2}{2\gamma+3}} \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2\gamma+3}}$	$\asymp R_{\gamma+1}^{\frac{2}{2\gamma+3}} \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2\gamma+3}}$

where: $R^* = \left(\max_{k \in \{1,...,\gamma+1\}} \frac{R_k}{(k-1)!}\right) \vee 1$ and R_0 is bounded away from zero under the column for $\mathcal{U}_{r+1,2}$; $R_{\gamma+1}$ is bounded away from zero under the column for $\mathcal{H}_{r+1,2}$.

linear models, and single index models are all built upon nonparametric regressions. Our results also have important implications for these more complicated statistical procedures and models.

We conclude the paper by providing some insights about the Hölder class of d-variate functions, $\mathcal{U}_{\gamma+1}^d$. The nonasymptotic analysis for the higher dimensional problem is rather involved because of the additional interplay between the smoothness parameter γ and the dimension d. We are only able to provide some partial answers regarding $\mathcal{U}_{\gamma+1}^d$. Yet, these results suggest that the classical asymptotic minimax rate $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+2+d}}$ could be an underestimate of the MSE in finite sample settings.

1.1 Notation and definitions

Notation. Let $\lfloor x \rfloor$ denote the largest integer smaller than or equal to x. For two functions f(n) and g(n), let us write $f(n) \succeq g(n)$ if $f(n) \ge cg(n)$ for a universal constant $c \in (0, \infty)$; similarly, we write $f(n) \preceq g(n)$ if $f(n) \le c'g(n)$ for a universal constant $c' \in (0, \infty)$; and $f(n) \asymp g(n)$ if $f(n) \succeq g(n)$ and $f(n) \preceq g(n)$. Throughout this paper, we use various c and C letters to denote positive universal constants that are: $\preceq 1$ and independent of $n, \gamma, \{R_k\}_{k=0}^{\gamma+1}$ and d; these constants may vary from place to place. For a d-dimensional vector θ , the l_q -norm $|\theta|_q := \left(\sum_{j=1}^d |\theta_j|^q\right)^{1/q}$ if $1 \le q < \infty$ and $|\theta|_q := \max_{j \in \{1,\ldots,d\}} |\theta_j|$ if $q = \infty$. Let $\mathbb{B}_q^d(R) := \left\{\theta \in \mathbb{R}^d : |\theta|_q \le R\right\}$. For functions, the $\mathcal{L}^2(\mathbb{P}_n)$ -norm of the vector $f := \{f(x_i)\}_{i=1}^n$, denoted by $|f|_n$, is $\left[\frac{1}{n}\sum_{i=1}^n (f(x_i))^2\right]^{\frac{1}{2}}$; the supremum norm $|f - g|_\infty := \sup_{x \in [a, b]^d} |f(x) - g(x)|$ and the $L^2(\mathbb{P})$ norm $|f - g|_2 := \sqrt{\frac{1}{b-a}} \int_a^b [f(x) - g(x)]^2 dx$.

	$R_0 = 1, R_0$	$\mathcal{C}_k \le (k-1)! \forall k \le \gamma + 1$				
$\frac{n}{\sigma^2}$	$\precsim \gamma^{2\gamma+3}$	$\succsim \gamma^{2\gamma+3}$	$\precsim (\gamma \log \gamma)^{2\gamma+3}$	$\succeq (\gamma \log \gamma)^{2\gamma+3}$		
MSE	$\asymp \frac{\sigma^2 \gamma}{n}$	$\asymp \left(rac{\sigma^2}{n} ight)^{rac{2\gamma+2}{2\gamma+3}}$		$\asymp \left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$		
If $\gamma = 0$ or $\gamma = 1$, the minimax optimal rate is trivially $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+3}}$ as long as $\frac{n}{\sigma^2} \succeq 1$						

Table 3: Minimax optimal MSE rates of the generalized $\mathcal{U}_{r+1} (= \mathcal{U}_{r+1,1} + \mathcal{U}_{r+1,2}), \gamma > 1$

Definition (covering and packing numbers). Given a set Λ , a set $\{\eta^1, \eta^2, ..., \eta^N\} \subset \Lambda$ is a δ -cover of Λ in the metric ρ if for each $\eta \in \Lambda$, there exists some $i \in \{1, ..., N\}$ such that $\rho(\eta, \eta^i) \leq \delta$. The δ -covering number of Λ , denoted by $N_{\rho}(\delta, \Lambda)$, is the cardinality of the smallest δ -cover. A set $\{\eta^1, \eta^2, ..., \eta^M\} \subset \Lambda$ is a δ -packing of Λ in the metric ρ if for any distinct $i, j \in \{1, ..., M\}$, $\rho(\eta^i, \eta^j) > \delta$. The δ -packing number of Λ , denoted by $M_{\rho}(\delta, \Lambda)$, is the cardinality of the largest δ -packing. Throughout this paper, we use $N_q(\delta, \mathcal{F})$ and $M_q(\delta, \mathcal{F})$ to denote the δ -covering number and the δ -packing number, respectively, of a function class \mathcal{F} with respect to the function norm $|\cdot|_q$ where $q \in \{2, \infty\}$.

The following is a standard textbook result that summarizes the relationships between covering and packing numbers:

$$M_{\rho}(2\delta, \Lambda) \leq N_{\rho}(\delta, \Lambda) \leq M_{\rho}(\delta, \Lambda).$$

Given this sandwich result, a lower bound on the packing number gives a lower bound on the covering number, and vice versa; similarly, an upper bound on the covering number gives an upper bound on the packing number, and vice versa.

2 Hölder classes

Let $p = (p_j)_{j=1}^d$ and $P = \sum_{j=1}^d p_j$ where p_j s are non-negative integers; $x = (x_j)_{j=1}^d$ and $x^p = \prod_{j=1}^d x_j^{p_j}$. Write $D^p f(x) = \partial^P f / \partial x_1^{p_1} \cdots \partial x_d^{p_d}$.

For a non-negative integer γ , let $\mathcal{U}_{\gamma+1}\left((R_k)_{k=0}^{\gamma+1}, [-1, 1]^d\right)$ be the class of functions such that any function $f \in \mathcal{U}_{\gamma+1}\left((R_k)_{k=0}^{\gamma+1}, [-1, 1]^d\right)$ satisfies: (1) f is continuous on $[-1, 1]^d$, and all partial derivatives of f exist for all p with $P := \sum_{k=1}^d p_k \leq \gamma$; (2) $|D^p f(x)| \leq R_k$ for all p with P = k $(k = 0, ..., \gamma)$ and $x \in [-1, 1]^d$, where $D^0 f(x) = f(x)$; (3) $|D^p f(x) - D^p f(x')| \leq R_{\gamma+1} |x - x'|_{\infty}$ for all p with $P = \gamma$ and $x, x' \in [-1, 1]^d$.

Our focus in this section is on d = 1, where we use the shortform $\mathcal{U}_{\gamma+1}$. Section 4 consider a general d, where we use the shortform $\mathcal{U}_{\gamma+1}^d$.

In view of (2), we have the following decomposition:

$$\mathcal{U}_{\gamma+1} = \mathcal{U}_{\gamma+1,1} + \mathcal{U}_{\gamma+1,2} := \{ f_1 + f_2 : f_1 \in \mathcal{U}_{r+1,1}, f_2 \in \mathcal{U}_{\gamma+1,2} \}$$

where

$$\mathcal{U}_{\gamma+1,1} = \left\{ f(x) = \sum_{k=0}^{\gamma} \theta_k x^k : (\theta_k)_{k=0}^{\gamma} \in \mathcal{P}_{\gamma}, \ x \in [-1, 1] \right\}$$

with the $(\gamma + 1)$ –dimensional polyhedron

$$\mathcal{P}_{\gamma} = \left\{ (\theta_k)_{k=0}^{\gamma} \in \mathbb{R}^{\gamma+1} : \theta_k \in \left[\frac{-R_k}{k!}, \frac{R_k}{k!} \right] \right\}$$

and

$$\mathcal{U}_{\gamma+1,2} = \left\{ f \in \mathcal{U}_{\gamma+1} : f^{(k)}(0) = 0 \text{ for all } k \le \gamma \right\}.$$

2.1 Metric entropy bounds

Lemma 2.1. If δ is small enough such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} \ge 0$, we have

$$\log N_2\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \le \log N_\infty\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \le \underbrace{\sum_{k=0}^{\gamma} \log \frac{4\left(\gamma+1\right) R_k}{k!\delta}}_{\overline{B}_1(\delta)};\tag{4}$$

if δ is large enough such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} < 0$, we have

$$\log N_2\left(\delta, \,\mathcal{U}_{\gamma+1,1}\right) \le \log N_\infty\left(\delta, \,\mathcal{U}_{\gamma+1,1}\right) \le \underbrace{\left(\frac{\gamma}{2}+1\right)\log\frac{1}{\delta} + \sum_{k=0}^{\gamma}\log R_k}_{\overline{B}_2(\delta)}.\tag{5}$$

In terms of the lower bounds, we have

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \ge \log M_{2}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \ge \underline{B}_{1}\left(\delta\right)$$

where $\underline{B}_{1}(\delta) = \sum_{k=0}^{\gamma} \log \left(9^{-\gamma} \gamma^{-\gamma}\right) + \sum_{k=0}^{\gamma} \log \frac{C \sum_{m=0}^{\lfloor \gamma/2 \rfloor} R_{k+2m}}{\delta}$ (with $R_{k+2m} = 0$ for $k+2m > \gamma$) for some positive universal constant C. Let $R^{\dagger} = \left(\sum_{k=0}^{\gamma} \frac{R_{k}}{k!}\right) \vee 1$ and $\tilde{k} \in \arg \max_{k \in \{0,...,\gamma\}} \frac{R_{k}}{k!}$. If $\frac{cR_{\tilde{k}}}{k!R^{\dagger}\delta} \geq 2^{\gamma}$ and $3R^{\dagger}\delta \leq \frac{2R_{\tilde{k}}}{\tilde{k}!}$, we also have

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \geq \log M_{2}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \geq \underline{B}_{2} = C'\gamma$$

for some positive universal constant C'.

Remark. When $R_k \simeq 1$ for $k = 0, ..., \gamma$, $R^{\dagger} \simeq 1$; when $R_0 \simeq 1$ and $R_k \simeq (k-1)!$ for $k = 1, ..., \gamma$, $R^{\dagger} \simeq \log(\gamma \lor 2)$; when $R_k \simeq k!$ for all $k = 0, ..., \gamma$, $R^{\dagger} \simeq (\gamma \lor 1)$.

The proof for Lemma 2.1 is given in Section 5.1.

The upper bound $\overline{B}_1(\delta)$ and the lower bound max $\{\underline{B}_1(\delta), \underline{B}_2\}$ as well as their proofs are novel. Bound $\overline{B}_2(\delta)$ is the extension of the upper bound associated with $\mathcal{U}_{\gamma+1,1}$ in [3] and allows for general R_k . Obviously $\overline{B}_1(\delta) \preceq \overline{B}_2(\delta)$. It is worth mentioning that (5) holds for all $\delta \in (0, 1)$ (not just δ such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} < 0$) but is far from being tight when $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} \geq 0$. When it comes to deriving the upper bounds for the convergence rates concerning $\mathcal{U}_{\gamma+1}$ with large enough R_k , (4) will be very useful. In particular, if $R_k = (k-1)!$ or $R_k = k!$ for $k = 1, ..., \gamma$, bound (4) gives much sharper scalings (in terms of γ) than (5).

In terms of the packing numbers, \underline{B}_2 will be useful for deriving the minimax lower bounds for the convergence rates if $R_k \leq (k-1)!$, while $\underline{B}_1(\delta)$ will be useful if $R_k = k!$.

Lemma 2.2. Let
$$R^* = \left(\max_{k \in \{1,\dots,\gamma+1\}} \frac{R_k}{(k-1)!}\right) \vee 1$$
. We have
 $\log N_2\left(\delta, \mathcal{U}_{\gamma+1,2}\right) \leq \log N_\infty\left(\delta, \mathcal{U}_{\gamma+1,2}\right) \precsim R^* \frac{1}{\gamma+1} \delta^{\frac{-1}{\gamma+1}}$
 $if R^* \frac{k}{\gamma+1} \delta^{1-\frac{k}{\gamma+1}} \in (0, 1) \text{ for } k = 0, \dots, \gamma.$

We also have

$$\log M_{\infty} \left(\delta, \mathcal{U}_{\gamma+1,2}\right) \geq \log M_{2} \left(\delta, \mathcal{U}_{\gamma+1,2}\right) \succeq R^{*\frac{1}{\gamma+1}} \delta^{\frac{-1}{\gamma+1}}, \quad if \ R_{0} \succeq 1, \ \delta \in (0, 1); \\ \log M_{\infty} \left(\delta, \mathcal{U}_{\gamma+1,2}\right) \geq \log M_{2} \left(\delta, \mathcal{U}_{\gamma+1,2}\right) \succeq \left(R^{*} R_{0}\right)^{\frac{1}{\gamma+1}} \delta^{\frac{-1}{\gamma+1}}, \quad if \ R_{0} \precsim 1, \ \delta \in (0, 1).$$

The proof for Lemma 2.2 is given in Section 5.2.

Lemma 2.2 extends [3] to allow for general R_k . When $R_k \preceq k!$ for all $k = 1, ..., \gamma + 1$, $R^{*\frac{1}{\gamma+1}} \simeq 1$ and the bounds in Lemma 2.2 coincide with those associated with $\mathcal{U}_{\gamma+1,2}$ in [3]. If $R_k \succeq k!$ for all $k = 0, ..., \gamma+1$, $R^{*\frac{1}{\gamma+1}} \succeq 1$; for example, taking $R_k \succeq (k!)^2$ for all $k = 0, ..., \gamma+1$ yields $R^{*\frac{1}{\gamma+1}} \succeq \frac{\gamma}{e}$.

Theorem 2.1. Given Lemmas 2.1 and 2.2, we have

$$\log N_2 \left(2\delta, \mathcal{U}_{\gamma+1}\right) \le \log N_\infty \left(2\delta, \mathcal{U}_{\gamma+1}\right) \\ \le \begin{cases} \overline{B}_1 \left(\delta\right) + R^* \frac{1}{\gamma+1} \delta \frac{-1}{\gamma+1} & \text{if } \min_{k \in \{0, \dots, \gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} \ge 0 \\ \overline{B}_2 \left(\delta\right) + R^* \frac{1}{\gamma+1} \delta \frac{-1}{\gamma+1} & \text{if } \min_{k \in \{0, \dots, \gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} < 0 \end{cases}$$
(6)

and

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1}\right) \geq \log M_{2}\left(\delta, \mathcal{U}_{\gamma+1}\right) \succeq \begin{cases} \max\left\{\underline{B}_{1}\left(\delta\right), \underline{B}_{2}, R^{*\frac{1}{\gamma+1}}\delta^{\frac{-1}{\gamma+1}}\right\} & \text{if } R_{0} \succeq 1\\ \max\left\{\underline{B}_{1}\left(\delta\right), \underline{B}_{2}, \left(R^{*}R_{0}\right)^{\frac{1}{\gamma+1}}\delta^{\frac{-1}{\gamma+1}}\right\} & \text{if } R_{0} \preceq 1\end{cases}$$

where these bounds are subject to the conditions in Lemmas 2.1 and 2.2.

Theorem 2.1 follows easily from Lemmas 2.1 and 2.2; see the proof of Theorem 2.1 in Section 5.3.

2.2 Minimax optimality

Following the literature on minimax optimality, let us consider (1) where $\{x_i\}_{i=1}^n$ are independently and uniformly distributed on [-1, 1], and $\{\epsilon_i\}_{i=1}^n$ are independent $\mathcal{N}(0, \sigma^2)$, and for the achievability results, let us consider the least squares estimator

$$\hat{f} \in \arg\min_{\check{f}\in\mathcal{F}} \frac{1}{2n} \sum_{i=1}^{n} \left(y_i - \check{f}\left(x_i\right) \right)^2.$$
(7)

Theorem 2.2. Let $\mathcal{F} = \mathcal{U}_{\gamma+1,2}$ in (7). If $R_0 \succeq 1$ and $\frac{n}{\sigma^2} > (R^*)^{\frac{\gamma(2\gamma+3)+1}{\gamma+1}}$, we have

$$\mathbb{E}\left(\left|\hat{f}-f\right|_{2}^{2}\right) \lesssim r^{2} + \exp\left\{-cn\left(\sigma^{-2}\wedge 1\right)r^{2}\right\},$$
$$\min_{\tilde{f}}\max_{f\in\mathcal{U}_{\gamma+1,2}}\mathbb{E}\left(\left|\tilde{f}-f\right|_{2}^{2}\right) \gtrsim r^{2},$$

where $r^2 = (R^*)^{\frac{2}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$.

The proof for Theorem 2.2 is given in Section 5.4.

Theorem 2.3. Let $\mathcal{F} = \mathcal{U}_{\gamma+1}$ in (7). Suppose $R_0 \approx 1$ and $R_k \leq (k-1)!$ for $k = 1, ..., \gamma + 1$. If

$$1 < \frac{n}{\sigma^2} \precsim (\gamma \lor 1)^{2\gamma+3},\tag{8}$$

we have

$$\mathbb{E}\left(\left|\hat{f} - f\right|_{2}^{2}\right) \precsim r_{1}^{2} + \exp\left\{-cn\left(\sigma^{-2} \wedge 1\right)r_{1}^{2}\right\}$$

where $r_1^2 = \frac{\sigma^2(\gamma \vee 1)}{n}$, and $r_1^2 \succeq \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$ under (8). In addition to (8), if $n \succeq 4^{\gamma} \propto P^{\dagger 2}$ for any $\alpha > 1$

$$\frac{n}{\sigma^2} \succeq 4^{\gamma} \gamma R^{\dagger 2} \quad for \ any \ \gamma > 1, \tag{9}$$

we also have

$$\min_{\tilde{f}} \max_{f \in \mathcal{U}_{\gamma+1}} \mathbb{E}\left(\left|\tilde{f} - f\right|_2^2\right) \succeq \frac{\sigma^2 \gamma}{n} \quad for \ any \ \gamma > 1$$

On the other hand, if

$$\frac{n}{\sigma^2} \succeq (\gamma \lor 1)^{2\gamma+3}, \tag{10}$$

then we have

$$\mathbb{E}\left(\left|\hat{f}-f\right|_{2}^{2}\right) \lesssim r_{2}^{2} + \exp\left\{-cn\left(\sigma^{-2}\wedge 1\right)r_{2}^{2}\right\}$$
$$\min_{\tilde{f}}\max_{f\in\mathcal{U}_{\gamma+1}}\mathbb{E}\left(\left|\tilde{f}-f\right|_{2}^{2}\right) \gtrsim r_{2}^{2},$$

where $r_2^2 = \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$, and $r_2^2 \succeq \frac{\sigma^2(\gamma \lor 1)}{n}$ under (10).

Remark. Consider $\gamma > 1$. When $R_k = 1$ for $k = 0, ..., \gamma$, (9) becomes $\frac{n}{\sigma^2} \succeq 4^{\gamma} \gamma$; when $R_0 \asymp 1$ and $R_k = (k-1)!$ for $k = 1, ..., \gamma$, (9) is satisfied if $\frac{n}{\sigma^2} \succeq 4^{\gamma} \gamma (\log \gamma)^2$.

The proof for Theorem 2.3 is given in Section 5.5.

Theorem 2.4. Let $\mathcal{F} = \mathcal{U}_{\gamma+1}$ in (7). Suppose $R_k = k!$ for $k = 0, ..., \gamma + 1$. If

$$(\gamma \vee 1)^{2\gamma+1} < \frac{n}{\sigma^2} \precsim ((\gamma \vee 1) \log (\gamma \vee 2))^{2\gamma+3}, \tag{11}$$

we have

$$\mathbb{E}\left(\left|\hat{f} - f\right|_{2}^{2}\right) \precsim r_{1}^{2} + \exp\left\{-cn\left(\sigma^{-2} \wedge 1\right)r_{1}^{2}\right\}$$

where $r_1^2 = \frac{\sigma^2(\gamma \vee 1)\log(\gamma \vee 2)}{n}$, and $r_1^2 \succeq \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$ under (11). For any $\gamma > 1$, under the condition $\max\left(a^{2\gamma+1} - 4\gamma a \log a\right) \preceq \frac{n}{n} \preceq (a \log a)^{2\gamma+3}$

$$\max\left(\gamma^{2\gamma+1}, \, 4^{\gamma}\gamma\log\gamma\right) \precsim \frac{n}{\sigma^2} \precsim (\gamma\log\gamma)^{2\gamma+3},$$

we also have

$$\min_{\tilde{f}} \max_{f \in \mathcal{U}_{\gamma+1}} \mathbb{E}\left(\left|\tilde{f} - f\right|_2^2\right) \succeq \frac{\sigma^2 \gamma \log \gamma}{n} \quad for \ any \ \gamma > 1.$$

On the other hand, if

$$\frac{n}{\sigma^2} \succeq \left((\gamma \lor 1) \log \left(\gamma \lor 2 \right) \right)^{2\gamma+3},\tag{12}$$

then we have

$$\begin{split} \mathbb{E}\left(\left|\hat{f}-f\right|_{2}^{2}\right) & \precsim \quad r_{2}^{2}+\exp\left\{-cn\left(\sigma^{-2}\wedge 1\right)r_{2}^{2}\right\},\\ \min_{\tilde{f}}\max_{f\in\mathcal{U}_{\gamma+1}}\mathbb{E}\left(\left|\tilde{f}-f\right|_{2}^{2}\right) & \succsim \quad r_{2}^{2}, \end{split}$$

where $r_2^2 = \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$, and $r_2^2 \succeq \frac{\sigma^2(\gamma \lor 1)\log(\gamma \lor 2)}{n}$ under (12).

The proof for Theorem 2.4 is given in Section 5.6.

3 Ellipsoid classes

In this section, we turn to the generalized ellipsoid class of smooth functions, (3).

3.1 Metric entropy bounds

Theorem 3.1. If $R_{\gamma+1} \succeq \gamma + 1$, we have

$$\log N_2\left(\delta, \mathcal{H}_{\gamma+1}\right) \asymp \left(R_{\gamma+1}\delta^{-1}\right)^{\frac{1}{\gamma+1}}.$$

If $R_{\gamma+1} \preceq \gamma + 1$, we have

$$\log N_2\left(\delta, \mathcal{H}_{\gamma+1}\right) \precsim \delta^{\frac{-1}{\gamma+1}},\\ \log N_2\left(\delta, \mathcal{H}_{\gamma+1}\right) \succsim \left(R_{\gamma+1}\delta^{-1}\right)^{\frac{1}{\gamma+1}}.$$

The proof for Theorem 3.1 is given in Section 5.7.

When $R_{\gamma+1} = 1$, Theorem 3.1 sharpens the upper bound for $\log N_2(\delta, \mathcal{H}_{\gamma+1})$ in [16] from $(\gamma \vee 1) \delta^{-\frac{1}{\gamma+1}}$ to $\delta^{-\frac{1}{\gamma+1}}$. We discover the cause of the gap lies in that the "pivotal" eigenvalue (that balances the "estimation error" and the "approximation error" from truncating for a given resolution δ) in [16] is not optimal. We close the gap by finding the optimal "pivotal" eigenvalue.

More generally, for the case of $R_{\gamma+1} \preceq \gamma+1$, we consider two different truncations, one giving the upper bound $\delta^{\frac{-1}{\gamma+1}}$ and the other giving the lower bound $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$. Note that $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}} \approx \delta^{\frac{-1}{\gamma+1}}$ when $R_{\gamma+1} \simeq 1$. For the case of $R_{\gamma+1} \succeq \gamma+1$, we use only one truncation to show that both the upper bound and the lower bound scale as $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$.

3.2 Minimax optimality

Let us consider (1) where $\{x_i\}_{i=1}^n$ are independently and uniformly distributed on [0, 1], and $\{\epsilon_i\}_{i=1}^n$ are independent $\mathcal{N}(0, \sigma^2)$.

Theorem 3.2. Let $\mathcal{F} = \mathcal{H}_{\gamma+1}$ in (7). Suppose $R_{\gamma+1} \succeq 1$. Then we have

$$\mathbb{E}\left(\left|\hat{f} - f\right|_{2}^{2}\right) \lesssim r^{2} + \exp\left\{-cn\left(\sigma^{-2} \wedge 1\right)r^{2}\right\},$$
$$\min_{\tilde{f}} \max_{f \in \mathcal{H}_{\gamma+1}} \mathbb{E}\left(\left|\tilde{f} - f\right|_{2}^{2}\right) \gtrsim r^{2},$$
$$ere \ r^{2} = R_{\gamma+1}^{\frac{2}{2(\gamma+1)+1}} \left(\frac{\sigma^{2}}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}.$$

whe

The proof for Theorem 3.2 is given in Section 5.8.

Let us introduce the Sobolev class with the restrictions at zero:

$$\begin{aligned} \mathcal{S}_{\gamma+1} &:= \{ f: [0, 1] \to \mathbb{R} | f \text{ is } \gamma + 1 \text{ times differentiable a.e.,} \\ f^{(k)}(0) &= 0 \text{ for all } k \leq \gamma, \text{ and} \\ f^{(\gamma)} \text{ is absolutely continuous with} \\ \int_0^1 \left[f^{(\gamma+1)}(t) \right]^2 dt \leq R_{\gamma+1}^2 \}. \end{aligned}$$

Like the Hölder classes in Section 2, the Sobolev classes considered above can be extended to those without the conditions $f^{(k)}(0) = 0$ for all $k \leq \gamma$. When $R_k = 1$ for all $k = 0, ..., \gamma + 1$, based on Lemma 2.1 and Theorem 3.1, using arguments similar to those for Theorem 2.3, we would arrive at the same claim (as in Theorem 2.3) for $\overline{S}_{\gamma+1} = \mathcal{U}_{r+1,1} + S_{\gamma+1}$.

Conclusion and some insights about multivariate smooth func-4 tions

In the nonasymptotic framework, we have shown that higher smoothness can bring a curse unless the polynomial subspace has some sparsity or approximate sparsity structures. For the Hölder class of d-variate functions, there is an additional interplay between γ and d. We conclude the paper by providing some insights about the higher dimensional generalized Hölder class.

Given any function $f \in \mathcal{U}_{\gamma+1}^d$, we have

$$f(x) = \sum_{k=0}^{\gamma} \sum_{p:P=k} \frac{x^p D^p f(0)}{k!} + \sum_{p:P=\gamma} \frac{x^p D^p f(z)}{\gamma!} - \sum_{p:P=\gamma} \frac{x^p D^p f(0)}{\gamma!}$$

for some intermediate value z. Similar to Section 2, we have the following decomposition:

$$\mathcal{U}_{\gamma+1}^{d} = \mathcal{U}_{\gamma+1,1}^{d} + \mathcal{U}_{\gamma+1,2}^{d} := \left\{ f_1 + f_2 : f_1 \in \mathcal{U}_{r+1,1}^{d}, f_2 \in \mathcal{U}_{\gamma+1,2}^{d} \right\}$$

where

$$\mathcal{U}_{\gamma+1,1}^{d} = \left\{ f = \sum_{k=0}^{\gamma} \sum_{p:P=k} x^{p} \theta_{(p,k)} : \{\theta_{(p,k)}\}_{(p,k)} \in \mathcal{P}_{\Gamma}, \ x \in [-1, \ 1]^{d} \right\}$$

with the $\Gamma := \sum_{k=0}^{\gamma} \begin{pmatrix} d+k-1 \\ d-1 \end{pmatrix}$ –dimensional polyhedron

$$\mathcal{P}_{\Gamma} = \left\{ \{\theta_{(p,k)}\}_{(p,k)} \in \mathbb{R}^{\Gamma} : \text{for any given } k \in \{0, ..., \gamma\}, \ \theta_{(p,k)} \in \left[\frac{-R_k}{k!}, \frac{R_k}{k!}\right] \text{ for all } p \text{ with } P \le k \right\}$$

where $\theta = \{\theta_{(p,k)}\}_{(p,k)}$ denotes the collection of $\theta_{(p,k)}$ over all (p,k) configurations. And,

$$\mathcal{U}_{\gamma+1,2}^{d} = \left\{ f \in \mathcal{U}_{\gamma+1}^{d} : D^{p} f\left(0\right) = 0 \text{ for all } p \text{ with } P \leq k, \, k = 0, ..., \gamma \right\}$$

Lemma 4.1. Let $R^* = \left(\max_{k \in \{1, ..., \gamma+1\}} \frac{D_{k-1}^* R_k}{(k-1)!} \lor 1 \right)$. We have

$$\log N_2\left(\delta, \mathcal{U}^d_{\gamma+1,2}\right) \leq \log N_\infty\left(\delta, \mathcal{U}^d_{\gamma+1,2}\right) \precsim d^d R^* \frac{d}{\gamma+1} \delta^{\frac{-d}{\gamma+1}}, \tag{13}$$

$$if \ P^* \frac{k}{\alpha+1} \delta^{1-\frac{k}{\alpha+1}} \subset (0, 1) \quad \text{for } k = 0$$

$$y R^{\gamma+1} \delta^{-\gamma+1} \in (0, 1) \text{ for } k = 0, ..., \gamma;$$

$$\log M_{\infty} \left(\delta, \mathcal{U}_{\gamma+1,2}^{d}\right) \geq \log M_{2} \left(\delta, \mathcal{U}_{\gamma+1,2}^{d}\right) \succeq d^{d} R^{*\frac{d}{\gamma+1}} \delta^{\frac{-d}{\gamma+1}} \quad \text{if } \delta \in (0, 1).$$

$$(14)$$

Remark. With Lemma 4.1, we can easily establish the minimax optimal MSE rate for $\mathcal{U}_{\gamma+1,2}^d$, using arguments almost identical to those for Theorem 2.2.

The proof for Lemma 4.1 is given in Section 5.9.

Lemma 4.2. If δ is small enough such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)D_k^*R_k}{\delta k!} \geq 0$, we have

$$\log N_2\left(\delta, \mathcal{U}_{\gamma+1,1}^d\right) \le \log N_\infty\left(\delta, \mathcal{U}_{\gamma+1,1}^d\right) \le \sum_{k=0}^{\gamma} D_k^* \log \frac{4\left(\gamma+1\right) D_k^* R_k}{\delta k!}$$

where $D_k^* = \begin{pmatrix} d+k-1\\ d-1 \end{pmatrix}$; if δ is large enough such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)D_k^*R_k}{\delta k!} < 0$, we have

$$\log N_2\left(\delta, \mathcal{U}^d_{\gamma+1,1}\right) \le \log N_\infty\left(\delta, \mathcal{U}^d_{\gamma+1,1}\right) \precsim \left(\sum_{k=0}^{\gamma} D_k^*\right) \log \frac{1}{\delta} + \sum_{k=0}^{\gamma} D_k^* \log R_k.$$
(15)

Remark. As in Lemma 2.1, (15) holds for all $\delta \in (0, 1)$ (not just δ such that $\min_k \log \frac{4(\gamma+1)D_k^*R_k}{\delta k!} < 0$) but is too loose when δ is small enough.

Remark. A simple upper bound on $\sum_{k=0}^{\gamma} D_k^*$ is $\sum_{k=1}^{\gamma} d^k \approx d^{\gamma}$. Let us show a lower bound on $\sum_{k=0}^{\gamma} D_k^*$ for the case of $\gamma \geq 2d^2$ to illustrate how large $\frac{\sigma^2}{n} \sum_{k=0}^{\gamma} D_k^*$ can be. We can write $D_k^* = \frac{(k+d-1)!}{(d-1)!k!} = \prod_{j=1}^{d-1} \frac{k+j}{j}$. Because $\gamma \geq 2d^2$, we have

$$\sum_{k=0}^{\gamma} D_k^* = \left(\sum_{k=0}^{\gamma} \prod_{j=1}^{d-1} \frac{k+j}{j}\right) \ge \left(\sum_{k=d^2}^{\gamma} \prod_{j=1}^{d-1} \frac{k+j}{j}\right)$$
$$\ge \left(d^2 \prod_{j=1}^{d-1} \left(\frac{d^2}{j}+1\right)\right) \ge \left(d^2 \left(\frac{d^2}{d}+1\right)^{d-1}\right) \ge d^{d+1}$$

The proof for Lemma 4.2 is given in Section 5.10. In theory, our arguments for $\underline{B}_1(\delta)$ in Lemma 2.1 can be extended for analyzing the lower bound for $\log M_2(\delta, \mathcal{U}_{\gamma+1,1}^d)$. However, this extension is very intensive. Arguments similar to those for \underline{B}_2 in Lemma 2.1 will not lead to a useful bound for $\mathcal{U}_{\gamma+1,1}^d$. Despite the lack of lower bounds, we can still gain some insights from Lemma 4.2, as it implies

$$\mathbb{E}\left(\left|\hat{f} - f\right|_{2}^{2}\right) \precsim r^{2} + \exp\left\{-cn\left(\sigma^{-2} \land 1\right)r^{2}\right\}$$

where $r^2 = \frac{\sigma^2}{n} \sum_{k=0}^{\gamma} D_k^*$ and \hat{f} is the estimator in (7) with $\mathcal{F} = \mathcal{U}_{\gamma+1,1}^d$. The quantity $\sum_{k=0}^{\gamma} D_k^*$ is the higher dimensional analogue of $\gamma + 1$ and arises from the fact that a function in $\mathcal{U}_{\gamma+1,1}^d$ has D_k^* distinct kth partial derivatives. Therefore, there is a good reason to think the rate r^2 is minimax optimal.

Suppose $R_k = 1$ for all $k = 0, ..., \gamma + 1$. If d is small relative to γ and n, Lemma 4.1 implies that the minimax optimal rate concerning $\mathcal{U}_{\gamma+1,2}^d$ is roughly $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+2+d}}$, the classical rate for $\mathcal{U}_{\gamma+1}^d$ derived under the regime where γ and d are finite but $n \to \infty$. Observe that $\frac{\sigma^2}{n} \sum_{k=0}^{\gamma} D_k^* \gtrsim \left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+2+d}}$ whenever $\frac{n}{\sigma^2} \precsim \left(\sum_{k=0}^{\gamma} D_k^*\right)^{\frac{2\gamma+2+d}{d}}$, and $\frac{\sigma^2}{n} \sum_{k=0}^{\gamma} D_k^* \precsim \left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+2+d}}$ whenever $\frac{n}{\sigma^2} \precsim \left(\sum_{k=0}^{\gamma} D_k^*\right)^{\frac{2\gamma+2+d}{d}}$. Therefore, the classical asymptotic minimax rate $\left(\frac{\sigma^2}{n}\right)^{\frac{2\gamma+2}{2\gamma+2+d}}$ could be an underestimate of the MSE in finite sample settings where n is not large enough.

5 Proofs

5.1 Proof for Lemma 2.1

The upper bound. Recall the definition of $\mathcal{U}_{\gamma+1,1}$:

$$\mathcal{U}_{\gamma+1,1} = \left\{ f = \sum_{k=0}^{\gamma} \theta_k x^k : (\theta_k)_{k=0}^{\gamma} \in \mathcal{P}_{\gamma}, \, x \in [-1, \, 1] \right\}$$

with the $(\gamma + 1)$ –dimensional polyhedron

$$\mathcal{P}_{\gamma} = \left\{ (\theta_k)_{k=0}^{\gamma} \in \mathbb{R}^{\gamma+1} : \theta_k \in \left[\frac{-R_k}{k!}, \frac{R_k}{k!} \right] \right\}$$

where R_k is allowed to depend on $k \in \{0, ..., \gamma\}$ only. We first derive an upper bound for $N_{\infty}(\delta, \mathcal{U}_{\gamma+1,1})$. Because the $L^2(\mathbb{P})$ norm is no greater than the sup norm and a smallest δ -cover of $\mathcal{U}_{\gamma+1,1}$ with respect to the $|\cdot|_{\infty}$ norm also covers $\mathcal{U}_{\gamma+1,1}$ with respect to the $|\cdot|_2$ norm, we have

$$N_2(\delta, \mathcal{U}_{\gamma+1,1}) \leq N_\infty(\delta, \mathcal{U}_{\gamma+1,1})$$

To bound $\log N_{\infty}(\delta, \mathcal{U}_{\gamma+1,1})$ from above, note that for $f, f' \in \mathcal{U}_{\gamma+1,1}$, we have

$$\left|f - f'\right|_{\infty} \le \sum_{k=0}^{\gamma} \left|\theta_k - \theta'_k\right|$$

where $f' = \sum_{k=0}^{\gamma} \theta'_k x^k$ such that $\theta' \in \mathcal{P}_{\gamma}$. Therefore, the problem is reduced to bounding $N_1(\delta, \mathcal{P}_{\gamma})$.

Consider $(a_k)_{k=0}^{\gamma}$ such that $a_k > 0$ for every $k = 0, ..., \gamma$ and $\sum_{k=0}^{\gamma} a_k = 1$. To cover \mathcal{P}_{γ} within δ -precision, we find a smallest $a_k \delta$ -cover of $\left[\frac{-R_k}{k!}, \frac{R_k}{k!}\right]$ for each $k = 0, ..., \gamma$, $\left\{\theta_k^1, ..., \theta_k^{N_k}\right\}$, such that for any $\theta \in \mathcal{P}_{\gamma}$, there exists some $i_k \in \{1, ..., N_k\}$ with

$$\sum_{k=0}^{\gamma} \left| \theta_k - \theta_k^{i_k} \right| \le \delta.$$

As a consequence, we have

$$\log N_1(\delta, \mathcal{P}_{\gamma}) \le \sum_{k=0}^{\gamma} \log \frac{4R_k}{a_k k! \delta} = -\sum_{k=0}^{\gamma} \log a_k + \sum_{k=0}^{\gamma} \log \frac{4R_k}{k! \delta}.$$
 (16)

For $(a_k)_{k=0}^{\gamma}$ such that $\sum_{k=0}^{\gamma} a_k = 1$, the function

$$h(a_0, ..., a_\gamma) := -\sum_{k=0}^{\gamma} \log a_k = -\log\left(\prod_{k=0}^{\gamma} a_k\right)$$

is minimized at $a_k = \frac{1}{\gamma+1}$. Consequently, the minimum of $\sum_{k=0}^{\gamma} \log \frac{4R_k}{a_k k! \delta}$ equals $\sum_{k=0}^{\gamma} \log \frac{4(\gamma+1)R_k}{k! \delta}$ and we have

$$\log N_1(\delta, \mathcal{P}_{\gamma}) \le \sum_{k=0}^{\gamma} \log \frac{4(\gamma+1)R_k}{k!\delta}.$$

Therefore,

$$\log N_2\left(\delta, \,\mathcal{U}_{\gamma+1,1}\right) \le \log N_\infty\left(\delta, \,\mathcal{U}_{\gamma+1,1}\right) \le \sum_{k=0}^{\gamma} \log \frac{4\left(\gamma+1\right)R_k}{k!\delta}.\tag{17}$$

If δ is large enough such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)R_k}{k!\delta} < 0$, we can evoke the counting argument in [3] and obtain

$$\log N_2\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \le \log N_\infty\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \le \left(\frac{\gamma}{2} + 1\right) \log \frac{1}{\delta} + \sum_{k=0}^{\gamma} \log R_k.$$
(18)

The lower bound. We first derive a lower bound for $N_2(\delta, \mathcal{U}_{\gamma+1,1})$. Because the $L^2(\mathbb{P})$ norm is no greater than the sup norm and a largest δ -packing of $\mathcal{U}_{\gamma+1,1}$ with respect to the $|\cdot|_2$ norm also packs $\mathcal{U}_{\gamma+1,1}$ with respect to the $|\cdot|_{\infty}$ norm, we have

$$N_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \geq N_2\left(\delta, \mathcal{U}_{\gamma+1,1}\right).$$

Let $(\phi_k)_{k=0}^{\gamma}$ be the Legendre polynomials on [-1, 1]. For any function $f \in \mathcal{U}_{\gamma+1,1}$, we can write

$$f(x) = \sum_{k=0}^{\gamma} \tilde{\theta}_k \phi_k(x)$$
(19)

such that

$$\tilde{\theta}_k = \frac{(2k+1)}{2} \int_{-1}^1 f(x)\phi_k(x)dx.$$
(20)

The first step of our proof is to derive nonasymptotic lower bounds on $|\tilde{\theta}_k|$ in (19) by exploiting the Legendre expansion of x^k . In Lemma A.1 of Section 5.11, we carefully modify the argument in [2] to show that

$$\tilde{\theta}_k = \left(k + \frac{1}{2}\right) \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \frac{f^{(k+2m)}(0)}{2^{k+2m}m! \left(\frac{1}{2}\right)_{k+m+1}}$$

where $(a)_k = a(a+1)\cdots(a+k-1)$ is known as the Pochhammer symbol. Recall $|f^{(k)}(0)| \leq R_k$ for $k = 0, ..., \gamma$ and $f^{(k)}(0) = 0$ for $k > \gamma$. We can re-write

$$\mathcal{U}_{\gamma+1,1} = \left\{ f = \sum_{k=0}^{\gamma} \tilde{\theta}_k \phi_k\left(x\right) : \left(\theta_k\right)_{k=0}^{\gamma} \in \mathcal{P}_{\gamma}^L, \, x \in [-1, \, 1] \right\}$$

with the $(\gamma + 1)$ –dimensional polyhedron

$$\mathcal{P}_{\gamma}^{L} = \left\{ \left(\tilde{\theta}_{k} \right)_{k=0}^{\gamma} \in \mathbb{R}^{\gamma+1} : \tilde{\theta}_{k} \in \left[-\overline{R}_{k}, \overline{R}_{k} \right] \right\}$$

where $\overline{R}_k := \sum_{m=0}^{\lfloor \gamma/2 \rfloor} b_{k,m} R_{k+2m}$ and $b_{k,m} = \frac{\left(k + \frac{1}{2}\right)}{2^{k+2m} m! \left(\frac{1}{2}\right)_{k+m+1}}$. If we can bound \overline{R}_k from below by \underline{R}_k , then we have

$$\mathcal{P}_{\gamma}^{L} \supseteq \underline{\mathcal{P}}_{\gamma}^{L} = \left\{ \left(\tilde{\theta}_{k} \right)_{k=0}^{\gamma} \in \mathbb{R}^{\gamma+1} : \tilde{\theta}_{k} \in \left[-\underline{R}_{k}, \underline{R}_{k} \right] \right\}.$$
⁽²¹⁾

Let us derive \underline{R}_k . Because $f^{(l)}(0) = 0$ for $l > \gamma$,

$$\frac{f^{(k+2m)}(0)}{2^{k+2m}m!\left(\frac{1}{2}\right)_{k+m+1}} = 0 \qquad \text{if } k+2m > \gamma$$

There are at most $\gamma + 1$ terms that are multiplied in the product $m! \left(\frac{1}{2}\right)_{k+m+1}$. Note that $m \leq \frac{\gamma}{2} \leq \frac{3\gamma}{2} + 1$ and

$$\begin{pmatrix} \frac{1}{2} \\ _{k+m+1} \end{pmatrix}_{k+m+1} = \frac{1}{2} \frac{1+2}{2} \cdots \frac{1+2(k+m)}{2} \\ \leq \frac{2}{2} \frac{2+2}{2} \cdots \frac{2+2(k+m)}{2} \\ = (k+m+1)!$$

where $k + m + 1 \le \frac{3\gamma}{2} + 1$. Hence, we have

$$m! \left(\frac{1}{2}\right)_{k+m+1} \le m! \left(k+m+1\right)! \le 1 \cdot \left(\frac{3\gamma}{2}+1\right)^{\gamma} \le (3\gamma)^{\gamma}.$$

As a result, we have

$$\overline{R}_{k} = \sum_{m=0}^{\lfloor \gamma/2 \rfloor} b_{k,m} R_{k+2m} = \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \frac{\left(k + \frac{1}{2}\right) R_{k+2m}}{2^{k+2m} m! \left(\frac{1}{2}\right)_{k+m+1}}$$

$$\geq \left(k + \frac{1}{2}\right) 2^{-\gamma} 3^{-\gamma} \gamma^{-\gamma} \sum_{m=0}^{\lfloor \gamma/2 \rfloor} R_{k+2m}$$

$$\geq \frac{9^{-\gamma} \gamma^{-\gamma}}{2} \sum_{m=0}^{\lfloor \gamma/2 \rfloor} R_{k+2m} =: \underline{R}_{k}.$$
(22)

Note that for any $f, g \in \mathcal{U}_{\gamma+1,1}$ where $f(x) = \sum_{k=0}^{\gamma} \tilde{\theta}_k \phi_k(x)$ and $g(x) = \sum_{k=0}^{\gamma} \tilde{\theta}'_k \phi_k(x)$, we have

$$|f - g|_2^2 = \sum_{k=0}^{\gamma} \left[\sqrt{\frac{2}{2k+1}} \left(\tilde{\theta}_k - \tilde{\theta}'_k \right) \right]^2.$$
(23)

In view of (21) and (23), to construct a packing set of $\mathcal{U}_{\gamma+1,1}$ within δ -separation, we find a largest $\sqrt{\frac{2k+1}{2}} \frac{\delta}{\sqrt{\gamma+1}}$ -packing of $[-\underline{R}_k, \underline{R}_k]$ for each $k = 0, ..., \gamma$, $\{\tilde{\theta}_k^1, ..., \tilde{\theta}_k^{M_k}\}$, such that for any distinct $\theta_k^{i_k}$ and $\theta_k^{j_k}$ in the packing sets,

$$\sum_{k=0}^{\gamma} \left[\sqrt{\frac{2}{2k+1}} \left(\tilde{\theta}_k^{i_k} - \tilde{\theta}_k^{j_k} \right) \right]^2 > \delta^2.$$

Therefore,

$$\log M_2\left(\delta, \,\mathcal{U}_{\gamma+1,1}\right) \gtrsim \sum_{k=0}^{\gamma} \log \frac{\sqrt{2\left(\gamma+1\right)}\underline{R}_k}{\sqrt{2k+1}\delta}.$$
(24)

Bounds (24) and (22) together give

$$\log M_2\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \ge \sum_{k=0}^{\gamma} \log\left(9^{-\gamma}\gamma^{-\gamma}\right) + \sum_{k=0}^{\gamma} \log\frac{C\sum_{m=0}^{\lfloor \gamma/2 \rfloor} R_{k+2m}}{\delta}$$
(25)

for some positive universal constant C. Because the $L^{2}(\mathbb{P})$ norm is no greater than the sup norm, we have

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \geq \log M_{2}\left(\delta, \mathcal{U}_{\gamma+1,1}\right)$$

$$\geq \sum_{k=0}^{\gamma} \log\left(9^{-\gamma}\gamma^{-\gamma}\right) + \sum_{k=0}^{\gamma} \log\frac{C\sum_{m=0}^{\lfloor\gamma/2\rfloor} R_{k+2m}}{\delta} =: \underline{B}_{1}\left(\delta\right).$$

The following argument gives another useful bound for $\log M_2(\delta, \mathcal{U}_{\gamma+1,1})$. Let $R^{\dagger} := \left(\sum_{k=0}^{\gamma} \frac{R_k}{k!}\right) \vee 1$ and $\tilde{k} \in \arg \max_{k \in \{0,...,\gamma\}} \frac{R_k}{k!}$. We consider a $3R^{\dagger}\delta$ -grid of points on $\left[-\frac{R_{\tilde{k}}}{k!}, \frac{R_{\tilde{k}}}{k!}\right]$ (that is, each point is $3R^{\dagger}\delta$ apart) and denote the collection of these points by $\left(\theta_{\tilde{k}}^{*i}\right)_{i=1}^{M_0}$ where $M_0 = \frac{cR_{\tilde{k}}}{\tilde{k}!R^{\dagger}\delta}$. We choose δ such that $M_0 \geq 2^{\gamma}$ and $3R^{\dagger}\delta \leq \frac{2R_{\tilde{k}}}{\tilde{k}!}$. Let us fix $\theta_k^* \in \left[\frac{-R_k\delta}{k!}, \frac{R_k\delta}{k!}\right]$ for $k \in \{0, ..., \gamma\} \setminus \tilde{k}$ and define

$$f_{\lambda_i}^*(x) = \theta_{\tilde{k}}^{*i} + \sum_{k \in \{0, \dots, \gamma\} \setminus \tilde{k}} \lambda_{i,k} \theta_k^* x^k, \quad x \in [-1, 1]$$

where $(\lambda_{i,k})_{k \in \{0,...,\gamma\} \setminus \tilde{k}} =: \lambda_i \in \{0, 1\}^{\gamma}$ for all $i = 1, ..., 2^{\gamma}$. For any $\lambda_i, \lambda_j \in \{0, 1\}^{\gamma}$ such that $i \neq j$,

we have

$$\begin{split} \left| f_{\lambda_{i}}^{*} - f_{\lambda_{j}}^{*} \right|_{2} &= \left[\frac{1}{2} \int_{-1}^{1} \left[\theta_{\bar{k}}^{*i} - \theta_{\bar{k}}^{*j} + \sum_{k \in \{0, \dots, \gamma\} \setminus \bar{k}} \left(1 \left\{ \lambda_{i,k} \neq \lambda_{j,k} \right\} \theta_{\bar{k}}^{*} x^{k} \right) \right]^{2} dx \right]^{\frac{1}{2}} \\ &\geq \left[\frac{1}{\sqrt{2}} \left[\int_{0}^{1} \left[\theta_{\bar{k}}^{*i} - \theta_{\bar{k}}^{*j} + \sum_{k \in \{0, \dots, \gamma\} \setminus \bar{k}} \left(1 \left\{ \lambda_{i,k} \neq \lambda_{j,k} \right\} \theta_{\bar{k}}^{*} x^{k} \right) \right]^{2} dx \right]^{\frac{1}{2}} \\ &\geq \left[\frac{1}{\sqrt{2}} \int_{0}^{1} \left| \theta_{\bar{k}}^{*i} - \theta_{\bar{k}}^{*j} + \sum_{k \in \{0, \dots, \gamma\} \setminus \bar{k}} \left(1 \left\{ \lambda_{i,k} \neq \lambda_{j,k} \right\} \theta_{\bar{k}}^{*} x^{k} \right) \right| dx \\ &\geq \left[\frac{1}{\sqrt{2}} \left(3R^{\dagger} \delta - \delta \sum_{k=0}^{\gamma} \frac{R_{k}}{k!} \right) \\ &\geq \left[\frac{1}{\sqrt{2}} \left[\left(2\delta \sum_{k=0}^{\gamma} \frac{R_{k}}{k!} \right) \lor (2\delta) \right] \right] > \delta \end{split}$$

where the third line follows from the Jensen's inequality and the concavity of $\sqrt{\cdot}$ on (0, 1], and the fourth line follows from the triangle inequality. Hence, we have constructed a δ -packing set. The cardinality of this packing set is 2^{γ} . Consequently, we have

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \geq \log M_{2}\left(\delta, \mathcal{U}_{\gamma+1,1}\right) \succeq \gamma =: \underline{B}_{2}.$$

Remark. Note that the lower bound $\log M_2(\delta, \mathcal{U}_{\gamma+1,1}) \succeq \gamma$ holds for all δ such that $\frac{cR_{\tilde{k}}}{\tilde{k}!R^{\dagger}\delta} \geq 2^{\gamma}$ and $3R^{\dagger}\delta \leq \frac{2R_{\tilde{k}}}{\tilde{k}!}$.

5.2 Proof for Lemma 2.2

The upper bound. The following derivations generalize [3]. In particular, properly choosing the grid of points on [-1, 1] is the key modification here. Any function $f \in \mathcal{U}_{\gamma+1,2}$ can be written as

$$f(x + \Delta) = \underbrace{f(x) + \Delta f'(x) + \frac{\Delta^2}{2!} f''(x) + \dots + \frac{\Delta^{\gamma - 1}}{(\gamma - 1)!} f^{(\gamma - 1)}(x)}_{:=F_{\gamma - 1}(x)} + \frac{\Delta^{\gamma}}{\gamma!} f^{(\gamma)}(z)$$

where $x, x + \Delta \in (-1, 1)$ and z is some intermediate value. Let $REM_0(x + \Delta) := f(x + \Delta) - F_{\gamma-1}(x) - \frac{\Delta^{\gamma}}{\gamma!}f^{(\gamma)}(x)$ and note that

$$|REM_{0}(x+\Delta)| = \frac{|\Delta|^{\gamma}}{\gamma!} \left| f^{(\gamma)}(z) - f^{(\gamma)}(x) \right|$$

$$\leq \frac{|\Delta|^{\gamma+1}}{\gamma!} R_{\gamma+1}.$$
(26)

In other words,

$$f(x+\Delta) = \sum_{k=0}^{\gamma} \frac{\Delta^k}{k!} f^{(k)}(x) + REM_0(x+\Delta)$$

where $|REM_0(x + \Delta)| \leq \frac{|\Delta|^{\gamma+1}}{\gamma!} R_{\gamma+1}$. Similarly, any $f^{(i)} \in \mathcal{U}_{\gamma+1-i,2}$ for $1 \leq i \leq \gamma$ can be written as

$$f^{(i)}(x+\Delta) = \sum_{k=0}^{\gamma-i} \frac{\Delta^k}{k!} f^{(i+k)}(x) + REM_i(x+\Delta)$$
(27)

where $|REM_i(x + \Delta)| \leq \frac{|\Delta|^{\gamma+1-i}}{(\gamma-i)!} R_{\gamma+1-i}$.

For some $\delta_0, \ldots, \delta_{\gamma} > 0$, suppose that $|f^{(k)}(x) - g^{(k)}(x)| \leq \delta_k$ for $k = 0, \ldots, \gamma$, where $f, g \in \mathcal{U}_{\gamma+1,2}$. Then we have

$$|f(x+\Delta) - g(x-\Delta)| \le \sum_{k=0}^{\gamma} \frac{|\Delta|^k \delta_k}{k!} + 2\frac{|\Delta|^{\gamma+1}}{\gamma!} R_{\gamma+1}.$$

Let $\left(\max_{k \in \{1,\dots,\gamma+1\}} \frac{R_k}{(k-1)!}\right) \vee 1 =: R^*$. Consider $|\Delta| \leq (R^{*-1}\delta)^{\frac{1}{\gamma+1}}$ and $\delta_k = R^{*\frac{k}{\gamma+1}}\delta^{1-\frac{k}{\gamma+1}}$ for $k = 0, \dots, \gamma$ and δ such that $\delta_k \in (0, 1)$. Then,

$$|f(x+\Delta) - g(x-\Delta)| \leq \delta \sum_{k=0}^{\gamma} \left(R^* \frac{-k+k}{\gamma+1} \frac{1}{k!} \right) + 2R^* |\Delta|^{\gamma+1}$$
$$\leq \delta \sum_{k=0}^{\gamma} \frac{1}{k!} + 2\delta \leq 5\delta.$$
(28)

Let us consider the following $(R^{*-1}\delta)^{\frac{1}{\gamma+1}}$ –grid of points in [-1, 1]:

$$x_{-s} < x_{-s+1} \cdots < x_{-1} < x_0 < x_1 < \cdots < x_{s-1} < x_s$$

with

$$x_0 = 0$$
 and $s \preceq \left(R^{*-1}\delta\right)^{\frac{-1}{\gamma+1}}$.

It suffices to cover the kth derivatives of functions in $\mathcal{U}_{\gamma+1,2}$ within δ_k -precision at each grid point. Then by (28), we obtain a 5 δ -cover of $\mathcal{U}_{\gamma+1,2}$. Following the arguments in [3], bounding N_{∞} (δ , $\mathcal{U}_{\gamma+1,2}$) can be reduced to bounding the cardinality of

$$\Lambda = \left\{ \left(\left\lfloor \frac{f^{(k)}(x_i)}{\delta_k} \right\rfloor, -s \le i \le s, \ 0 \le k \le \gamma \right) : f \in \mathcal{U}_{\gamma+1,2} \right\}$$

with $\lfloor x \rfloor$ denoting the largest integer smaller than or equal to x. Starting with $x_0 = 0$, the number of possible values of the vector $\left(\left\lfloor \frac{f^{(k)}(x_0)}{\delta_k} \right\rfloor \right)_{k=0}^{\gamma}$ when f ranges over $\mathcal{U}_{\gamma+1,2}$ is 1. For i = 1, ..., s, given the value of $\left(\left\lfloor \frac{f^{(k)}(x_{i-1})}{\delta_k} \right\rfloor \right)_{k=0}^{\gamma}$, let us count the number of possible values of $\left(\left\lfloor \frac{f^{(k)}(x_i)}{\delta_k} \right\rfloor \right)_{k=0}^{\gamma}$. The counting for $\left(\left\lfloor \frac{f^{(k)}(x_{i-1})}{\delta_k} \right\rfloor \right)_{k=0}^{\gamma}$ is similar. For each $0 \le k \le \gamma$, let $B_{k,i-1} := \left\lfloor \frac{f^{(k)}(x_{i-1})}{\delta_k} \right\rfloor$. Observe that $B_{k,i-1}\delta_k \le f^{(k)}(x_0) < (B_{k,i-1}+1)\delta_k$.

Taking (27) with $x = x_{i-1}$ and $\Delta = x_i - x_{i-1}$ gives

$$f^{(i)}(x_i) - \sum_{k=0}^{\gamma-i} \frac{\Delta^k}{k!} f^{(i+k)}(x_{i-1}) \bigg| \le \frac{|\Delta|^{\gamma+1-i}}{(\gamma-i)!} R_{\gamma+1-i}$$

As a result,

$$\begin{aligned} \left| f^{(i)}(x_{i}) - \sum_{k=0}^{\gamma-i} \frac{\Delta^{k}}{k!} B_{i+k,i-1} \right| \\ &\leq \left| f^{(i)}(x_{i}) - \sum_{k=0}^{\gamma-i} \frac{\Delta^{k}}{k!} f^{(i+k)}(x_{i-1}) \right| + \left| \sum_{k=0}^{\gamma-i} \frac{\Delta^{k}}{k!} \left(f^{(i+k)}(x_{i-1}) - B_{i+k,i-1} \right) \right| \\ &\leq \frac{|\Delta|^{\gamma+1-i}}{(\gamma-i)!} R_{\gamma+1-i} + \sum_{k=0}^{\gamma-i} \frac{|\Delta|^{k}}{k!} \delta_{i+k} \\ &\leq \left(R^{*-1} \delta \right)^{1-\frac{i}{\gamma+1}} R^{*}_{\gamma+1-i} + \sum_{k=0}^{\gamma-i} \left[\frac{1}{k!} \left(R^{*-1} \delta \right)^{\frac{k}{\gamma+1}} R^{*\frac{i+k}{\gamma+1}} \delta^{1-\frac{i+k}{\gamma+1}} \right] \\ &\leq R^{*\frac{i}{\gamma+1}} \delta^{1-\frac{i}{\gamma+1}} + R^{*\frac{i}{\gamma+1}} \delta^{1-\frac{i}{\gamma+1}} \sum_{k=0}^{\gamma-i} \frac{1}{k!} \leq 4\delta_{i}. \end{aligned}$$

Hence, the number of possible values of $\left(\left\lfloor \frac{f^{(k)}(x_i)}{\delta_k} \right\rfloor\right)_{k=0}^{\gamma}$ is at most 4 given the value of $\left(\left\lfloor \frac{f^{(k)}(x_{i-1})}{\delta_k} \right\rfloor\right)_{k=0}^{\gamma}$. Consequently, we have

card (
$$\Lambda$$
) $\precsim 4^{2s} \precsim 16^{\left(R^{*-1}\delta\right)^{\frac{-1}{\gamma+1}}}$

which implies

$$\log N_2\left(\delta, \,\mathcal{U}_{\gamma+1,2}\right) \le \log N_\infty\left(\delta, \,\mathcal{U}_{\gamma+1,2}\right) \precsim R^* \frac{1}{\gamma+1} \delta^{\frac{-1}{\gamma+1}}.$$
(29)

The lower bound. In the derivation of the lower bound, [3] considers a $\delta^{\frac{1}{\gamma+1}}$ -grid of points

$$\cdots < \underline{a}_1 < \overline{a}_1 < \underline{a}_2 < \overline{a}_2 < \cdots < \underline{a}_{2s} < \overline{a}_{2s}$$

where $\overline{a}_i - \underline{a}_i = \delta^{\frac{1}{\gamma+1}}$ and $s \succeq \delta^{\frac{-1}{\gamma+1}}$. Recall that we have previously considered a $(R^{*-1}\delta)^{\frac{1}{\gamma+1}}$ -grid of points in [-1, 1] in the derivation of the upper bound for $\log N_{\infty}(\delta, \mathcal{U}_{\gamma+1,2})$. To obtain a lower bound for $\log M_{\infty}(\delta, \mathcal{U}_{\gamma+1,2})$ with the same scaling as our upper bound, the key modification we need is to replace the $\overline{a}_i - \underline{a}_i = \delta^{\frac{1}{\gamma+1}}$ with $\overline{a}_i - \underline{a}_i = (R^{*-1}\delta)^{\frac{1}{\gamma+1}}$ and $s \succeq \delta^{\frac{-1}{\gamma+1}}$ with $s \succeq R^{*\frac{1}{\gamma+1}}\delta^{\frac{-1}{\gamma+1}}$. The rest of the arguments are similar to those in [3]. In particular, let us consider

$$f_{\lambda}(x) = R^* \sum_{i=1}^{2s} \lambda_i \left(\overline{a}_i - \underline{a}_i\right)^{\gamma+1} h_0\left(\frac{x - \underline{a}_i}{\overline{a}_i - \underline{a}_i}\right)$$

where $\lambda_i \in \{0,1\}$ and $\lambda \in \{0,1\}^{2s}$, and h_0 is a function on \mathbb{R} satisfying: (1) h_0 restricted to [-1, 1] belongs to $\mathcal{U}_{\gamma+1,2}$; (2) $h_0(x) = 0$ for $x \notin (0, 1)$ and $h_0(x) > 0$ for $x \in (0, 1)$; (3) $h_0\left(\frac{1}{2}\right) = \max_{x \in [0,1]} h_0(x) = R_0$. As an example, we can take $h_0(x) = \begin{cases} 0 & x \notin (0, 1) \\ be^{\frac{-1}{x}}e^{\frac{-1}{(1-x)}} & x \in (0, 1) \end{cases}$ for some properly chosen constant *b* that can only depend on R_0 . Note that the functions $h(x) := R^* (\overline{a}_i - \underline{a}_i)^{\gamma+1} h_0\left(\frac{x-\underline{a}_i}{\overline{a}_i-\underline{a}_i}\right)$ and also $f_\lambda(x)$ belong to $\mathcal{U}_{\gamma+1,2}$ if $\delta \in (0, 1)$. For any distinct $\lambda, \lambda' \in \{0,1\}^{2s}$, we have

$$\left|f_{\lambda} - f_{\lambda'}\right|_{\infty} \ge R^* \left(\overline{a}_i - \underline{a}_i\right)^{\gamma+1} h_0\left(\frac{1}{2}\right) = R_0 \delta.$$

If $R_0 \succeq 1$, then $R_0 \delta \succeq \delta$ and

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,2}\right) \succeq R^{*\frac{1}{\gamma+1}} \delta^{\frac{-1}{\gamma+1}}$$

If $R_0 \preceq 1$, then we obtain

$$\log M_{\infty}\left(R_{0}\delta, \mathcal{U}_{\gamma+1,2}\right) \succeq R^{*\frac{1}{\gamma+1}}\delta^{\frac{-1}{\gamma+1}}$$

which implies that

$$\log M_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,2}\right) \succsim R^{*\frac{1}{\gamma+1}} \left(\frac{\delta}{R_0}\right)^{\frac{-1}{\gamma+1}}$$

Standard argument in the literature based on the Vasharmov-Gilbert Lemma further gives

$$\log M_2(\delta, \mathcal{U}_{\gamma+1,2}) \gtrsim \begin{cases} R^* \frac{1}{\gamma+1} \delta^{-1} & \text{if } R_0 \gtrsim 1\\ (R^* R_0)^{\frac{1}{\gamma+1}} \delta^{-1} & \text{if } R_0 \precsim 1 \end{cases}.$$
(30)

5.3 Proof for Theorem 2.1

To cover $\mathcal{U}_{\gamma+1}$ within 2δ -precision, we find a smallest δ -cover of $\mathcal{U}_{\gamma+1,1}$, $\{f_{1,1}, f_{1,2}, ..., f_{1,N_1}\}$, and a smallest δ -cover of $\mathcal{U}_{\gamma+1,2}$, $\{f_{2,1}, f_{2,2}, ..., f_{2,N_2}\}$. Given that any $f \in \mathcal{U}_{\gamma+1}$ can be expressed by $f = f_1 + f_2$ for some $f_1 \in \mathcal{U}_{\gamma+1,1}$ and $f_2 \in \mathcal{U}_{\gamma+1,2}$, there exist some $f_{1,i}$ and $f_{2,i'}$ from the covering sets such that

$$|f_1 + f_2 - f_{1,i} - f_{2,i}|_q \le |f_1 - f_{1,i}|_q + |f_2 - f_{2,i}|_q \le 2\delta, \ q \in \{2, \infty\}$$

Consequently, we obtain

$$\log N_q\left(2\delta, \mathcal{U}_{\gamma+1}\right) \le \log N_q\left(\delta, \mathcal{U}_{\gamma+1,1}\right) + \log N_q\left(\delta, \mathcal{U}_{\gamma+1,2}\right), \ q \in \{2, \infty\}.$$

In terms of log $M_q(\delta, \mathcal{U}_{\gamma+1})$, we have

$$\log M_q(\delta, \mathcal{U}_{\gamma+1}) \geq \max \left\{ \log M_q(\delta, \mathcal{U}_{\gamma+1,1}), \log M_q(\delta, \mathcal{U}_{\gamma+1,2}) \right\}, \ q \in \{2, \infty\}.$$

5.4 Proof for Theorem 2.2

The upper bound. In view of (7), the *basic inequality* gives

$$\frac{1}{n}\sum_{i=1}^{n} \left(\hat{f}(x_i) - f(x_i)\right)^2 \le \frac{2}{n}\sum_{i=1}^{n} \varepsilon_i \left(\hat{f}(x_i) - f(x_i)\right).$$
(31)

Let $\Omega(r; \bar{\mathcal{F}}) = \{f \in \bar{\mathcal{F}} : |f|_n \leq r\}$ with $\bar{\mathcal{F}} := \{g = g_1 - g_2 : g_1, g_2 \in \mathcal{F}\}$. We follow the standard recipe in the literature (see, e.g., [1, 4, 13, 16]) to bound the right-hand-side of (31). By Corollary 13.7 in [16], we solve for any $r \in (0, \sigma]$ such that

$$\frac{c}{\sqrt{n}} \int_{\frac{r^2}{4\sigma}}^{r} \sqrt{\log N_n(\delta, \ \Omega(r; \ \bar{\mathcal{F}}))} d\delta \le \frac{r^2}{\sigma}$$

where $N_n(\delta, \Omega(r; \bar{\mathcal{F}}))$ is the δ -covering number of the set $\Omega(r; \bar{\mathcal{F}})$ in the $|\cdot|_n$ norm. Note that

$$\frac{1}{\sqrt{n}} \int_{\frac{r^2}{4\sigma}}^{r} \sqrt{\log N_n(\delta, \Omega(r; \bar{\mathcal{F}}))} d\delta$$

$$\leq \frac{1}{\sqrt{n}} \int_0^r \sqrt{\log N_\infty(\delta, \bar{\mathcal{F}})} d\delta$$

$$\stackrel{\sim}{\sim} \underbrace{\frac{1}{\sqrt{n}} (R^*)^{\frac{1}{2\gamma+2}} r^{\frac{2\gamma+1}{2\gamma+2}}}_{T(r)}.$$

Setting $\sigma T(r) \asymp r^2$ yields

$$r^2 \asymp (R^*)^{\frac{2}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$$

By Theorem 14.12 in [16] and integrating the tail probability, we have

$$\mathbb{E}\left(\left|\hat{f}-f\right|_{2}^{2}\right) \precsim r^{2} + \exp\left\{-cn\left(\sigma^{-2}\wedge 1\right)r^{2}\right\}.$$

Note that in deriving T(r), we have used the upper bound in Lemma 2.2, $\log N_{\infty}(\delta, \mathcal{U}_{\gamma+1,2}) \preceq R^* \frac{1}{\gamma+1} \delta^{\frac{-1}{\gamma+1}}$, which is valid if $R^* \frac{k}{\gamma+1} \delta^{1-\frac{k}{\gamma+1}} \in (0, 1)$ for $k = 0, ..., \gamma$. These conditions are satisfied if

$$R^* \frac{\gamma}{\gamma+1} \left[(R^*)^{\frac{1}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{\gamma+1}{2(\gamma+1)+1}} \right]^{\frac{1}{\gamma+1}} = (R^*)^{\frac{\gamma(2\gamma+3)+1}{(2\gamma+3)(\gamma+1)}} \left(\frac{\sigma^2}{n}\right)^{\frac{1}{2(\gamma+1)+1}} < 1,$$

which is equivalent to

$$\frac{n}{\sigma^2} > \left(R^*\right)^{\frac{\gamma(2\gamma+3)+1}{\gamma+1}}.$$
(32)

The lower bound. The Yang and Barron version of Fano's inequality (see, e.g., [16], [17]) gives

$$\min_{\tilde{f}} \max_{f} \mathbb{E}\left(\left|\tilde{f} - f\right|_{2}^{2}\right) \ge \sup_{\eta,\epsilon} \frac{\delta^{2}}{4} \left(1 - \frac{\log 2 + \log N_{KL}\left(\epsilon, \mathcal{Q}_{\mathcal{U}_{2}}\right) + \epsilon^{2}}{\log M_{2}\left(\delta, \mathcal{U}_{\gamma+1,2}\right)}\right)$$
(33)

where $N_{KL}(\epsilon, \mathcal{Q}_{\mathcal{U}_2})$ denotes the ϵ -covering number of $\mathcal{U}_{\gamma+1,2}$ with respect to the square root of the KL-divergence. We denote the product distribution of $\{x_i\}_{i=1}^n$ by \mathbb{U} , and the distribution of y given $\{x_i\}_{i=1}^n$ by \mathbb{P}_j when the truth is f_j . Observe that

$$D_{KL} \left(\mathbb{P}_{j} \times \mathbb{U} \parallel \mathbb{P}_{k} \times \mathbb{U} \right) = \mathbb{E}_{X} \left[D_{KL} \left(\mathbb{P}_{j} \parallel \mathbb{P}_{k} \right) \right]$$
$$= \frac{n}{2\sigma^{2}} \left| \tilde{f} - f \right|_{2}^{2}$$

and consequently, under (32),

$$\log N_{KL}(\epsilon, \mathcal{Q}_{\mathcal{U}_2}) = \log N_2\left(\sqrt{\frac{2}{n}}\sigma\epsilon, \mathcal{U}_{\gamma+1,2}\right) \precsim \left(\frac{R^*\sqrt{n}}{\sigma\epsilon}\right)^{\frac{1}{\gamma+1}}.$$

Setting $\left(\frac{R^*\sqrt{n}}{\sigma\epsilon}\right)^{\frac{1}{\gamma+1}} \asymp \epsilon^2$ yields $\epsilon^2 \asymp \left(\frac{nR^{*2}}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}} =: \epsilon^{*2}$. Observe that setting

$$\delta \asymp (R^*)^{\frac{1}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{\gamma+1}{2(\gamma+1)+1}}$$

ensures

$$\left(R^*\delta^{-1}\right)^{\frac{1}{\gamma+1}} \asymp \left(R^*\right)^{\frac{2}{2\gamma+3}} \left(\frac{n}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}} \succeq \epsilon^{*2}.$$

Consequently, we have

$$1 - \frac{\log 2 + \log N_{KL}\left(\epsilon^*, \mathcal{Q}_{\mathcal{U}_2}\right) + \epsilon^{*2}}{\log M_2\left(\delta^*, \mathcal{U}_{\gamma+1,2}\right)} \ge \frac{1}{2}$$

and

$$\min_{\tilde{f}} \max_{f \in \mathcal{U}_{\gamma+1,2}} \mathbb{E}\left(\left|\tilde{f} - f\right|_2^2\right) \succeq (R^*)^{\frac{2}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$$

5.5 Proof for Theorem 2.3

The upper bound. Taking $R_0 \approx 1$ and $R_k \leq (k-1)!$ for $k = 1, ..., \gamma + 1$ in (6) yields

$$\log N_2\left(\delta, \,\mathcal{U}_{\gamma+1,1}\right) \precsim \left(\gamma \lor 1\right) \log \frac{1}{\delta} + \left(\frac{1}{\delta}\right)^{\frac{1}{\gamma+1}},\tag{34}$$

where we have used the fact that $R^* = 1$. Note that

$$\frac{1}{\sqrt{n}} \int_{\frac{r^2}{4\sigma}}^r \sqrt{\log N_n(\delta, \Omega(r; \bar{\mathcal{F}}))} d\delta$$

$$\leq \frac{1}{\sqrt{n}} \int_0^r \sqrt{\log N_\infty(\delta, \bar{\mathcal{F}})} d\delta$$

$$\asymp \underbrace{r\sqrt{\frac{\gamma \vee 1}{n}} + \frac{1}{\sqrt{n}} r^{\frac{2\gamma + 1}{2\gamma + 2}}}_{T(r)}$$

where the last line follows from (34). Note that when $R_0 \approx 1$ and $R_k \leq (k-1)!$ for $k = 1, ..., \gamma + 1$, (32) is reduced to $\frac{n}{\sigma^2} > 1$. Setting $\sigma T(r) \approx r^2$ yields

$$r^2 \simeq \max\left\{\frac{\sigma^2\left(\gamma \lor 1\right)}{n}, \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}\right\}.$$
 (35)

By Theorem 14.12 in [16] and integrating the tail probability, we have

$$\mathbb{E}\left(\left|\hat{f} - f\right|_{2}^{2}\right) \precsim r^{2} + \exp\left\{-cn\left(\sigma^{-2} \land 1\right)r^{2}\right\}$$

If $1 < \frac{n}{\sigma^2} \precsim (\gamma \lor 1)^{2(\gamma+1)+1}$, we have $r^2 \asymp \frac{\sigma^2(\gamma \lor 1)}{n}$. Otherwise, we have $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$.

The lower bound. The lower bound $\left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$ is well known and hence we only show the lower bound $\frac{\sigma^2(\gamma\vee 1)}{n}$ under the condition $1 < \frac{n}{\sigma^2} \preceq (\gamma \vee 1)^{2(\gamma+1)+1}$. If $\gamma = 0$ (or $\gamma = 1$), then as long as $\frac{n}{\sigma^2} \succeq 1$, we have $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{2}{3}}$ (respectively, $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{4}{5}}$) in the derivation of the upper bound. The minimax lower bound is trivially $\left(\frac{\sigma^2}{n}\right)^{\frac{2}{3}}$ (respectively, $\left(\frac{\sigma^2}{n}\right)^{\frac{4}{5}}$). Therefore, in what follows, we assume $\gamma > 1$.

Again, we use

$$\min_{\tilde{f}} \max_{f} \mathbb{E}\left(\left|\tilde{f} - f\right|_{2}^{2}\right) \ge \sup_{\eta,\epsilon} \frac{\delta^{2}}{4} \left(1 - \frac{\log 2 + \log N_{KL}\left(\epsilon, \mathcal{Q}_{\mathcal{U}}\right) + \epsilon^{2}}{\log M_{2}\left(\delta, \mathcal{U}_{\gamma+1}\right)}\right)$$
(36)

where $N_{KL}(\epsilon, \mathcal{Q}_{\mathcal{U}})$ denotes the ϵ -covering number of $\mathcal{U}_{\gamma+1}$ with respect to the square root of the KL-divergence. Similar to Section 5.4, under $\frac{n}{\sigma^2} > 1$, we have

$$\log N_{KL}(\epsilon, \mathcal{Q}_{\mathcal{U}}) = \log N_2 \left(\sqrt{\frac{2}{n}} \sigma \epsilon, \mathcal{U}_{\gamma+1} \right)$$
$$\lesssim \gamma \log \frac{\sqrt{n\gamma}}{\sigma \epsilon} - \sum_{k=0}^{\gamma} \log k! + \left(\frac{\sqrt{n}}{\sigma \epsilon} \right)^{\frac{1}{\gamma+1}}$$
$$\lesssim \left(\frac{\sqrt{n}}{\sigma \epsilon} \right)^{\frac{1}{\gamma+1}}$$

for any $\gamma \in \{0, ..., \beta\}$, where the last line follows by choosing a sufficiently small ϵ . Setting $\left(\frac{\sqrt{n}}{\sigma\epsilon}\right)^{\frac{1}{\gamma+1}} \simeq \epsilon^2$ yields $\epsilon^2 \simeq \left(\frac{n}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}} := \epsilon^{*2}$.

Next, we turn to $\log M_2(\delta, \mathcal{U}_{\gamma+1})$ in (36). For classes with $R_0 \simeq 1$ and $R_k \leq (k-1)!$ for $k = 1, ..., \gamma + 1$, it turns out the lower bound $\log M_2(\delta, \mathcal{U}_{\gamma+1,1}) \succeq \gamma$ will suffice. Recalling the definition of $\frac{R_{\tilde{k}}}{k!}$ in Section 5.1, we have $\frac{R_{\tilde{k}}}{k!} \simeq 1$ in the cases of $R_0 \simeq 1$ and $R_k \leq (k-1)!$. Also recall that the lower bound $\log M_2(\delta, \mathcal{U}_{\gamma+1,1}) \succeq \gamma$ in Section 5.1 holds for all δ such that $\frac{cR_{\tilde{k}}}{k!R^{\dagger}\delta} \geq 2^{\gamma}$ and $3R^{\dagger}\delta \leq \frac{2R_{\tilde{k}}}{k!}$. If

$$\gamma \gtrsim \epsilon^{*2} \asymp \left(\frac{n}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}},$$

$$\frac{n}{\sigma^2} \precsim \gamma^{2(\gamma+1)+1},$$
(37)

then we obtain

which is equivalent to

$$1 - \frac{\log 2 + \log N_{KL} \left(\epsilon^*, \mathcal{Q}_{\mathcal{U}}\right) + \epsilon^{*2}}{\log M_2 \left(\delta, \mathcal{U}_{\gamma+1}\right)} \ge \frac{1}{2}$$

for all δ such that $\delta \preceq \frac{2^{-\gamma}}{R^{\dagger}}$. Take $\delta^2 = \delta^{*2} \asymp \frac{\sigma^2 \gamma}{n}$ and observe that $\delta^* \preceq \frac{2^{-\gamma}}{R^{\dagger}}$ if

$$\frac{n}{\sigma^2} \gtrsim 2^{2\gamma} \gamma R^{\dagger 2} \tag{38}$$

where $R^{\dagger} \preceq \log \gamma$ in the cases of $R_0 \asymp 1$ and $R_k \leq (k-1)!$.

Thus, we have shown that under (37) and (38),

$$\min_{\tilde{f}} \max_{f \in \mathcal{U}_{\gamma+1}} \mathbb{E}\left(\left|\tilde{f} - f\right|_2^2\right) \succeq \frac{\sigma^2 \gamma}{n}$$

5.6 Proof for Theorem 2.4

In view of (17) and (29) (where $R^* \frac{1}{\gamma+1} \approx 1$), we have

$$\log N_2\left(\delta, \,\mathcal{U}_{\gamma+1}\right) \precsim \left(\gamma+1\right) \log \frac{\gamma+1}{\delta} + \left(\frac{1}{\delta}\right)^{\frac{1}{\gamma+1}}.\tag{39}$$

In view of (25) and (30) (where also $R_0 \approx 1$), for $\gamma > 1$, we have

$$\log M_2(\delta, \mathcal{U}_{\gamma+1}) \geq c' \left[(\gamma+1)\log\frac{1}{\delta} - \gamma^2 + \left(\frac{1}{\delta}\right)^{\frac{1}{\gamma+1}} \right] \\ - (\gamma+1)\log\gamma^{\gamma} + (\gamma+1)\log(\gamma-1)! \\ \approx \underbrace{\gamma\log\frac{1}{\delta}}_{T_1(\delta)} - \underbrace{\gamma^2}_{T_2} + \underbrace{\left(\frac{1}{\delta}\right)^{\frac{1}{\gamma+1}}}_{T_3(\delta)}.$$

The upper bound. In terms of the upper bound, we obtain

$$r^{2} \asymp \max\left\{\frac{\sigma^{2}\left(\gamma \lor 1\right)\log\left(\gamma \lor 2\right)}{n}, \left(\frac{\sigma^{2}}{n}\right)^{\frac{2\left(\gamma+1\right)}{2\left(\gamma+1\right)+1}}\right\}.$$
(40)

Note that when $R_k = k!$ for $k = 1, ..., \gamma + 1$, (32) is satisfied if $\frac{n}{\sigma^2} > (\gamma \lor 1)^{2\gamma+1}$. If $(\gamma \lor 1)^{2\gamma+1} < \frac{n}{\sigma^2} \precsim ((\gamma \lor 1) \log (\gamma \lor 2))^{2(\gamma+1)+1}$, we have $r^2 \asymp \frac{\sigma^2(\gamma \lor 1) \log(\gamma \lor 2)}{n}$. If $\frac{n}{\sigma^2} \succsim ((\gamma \lor 1) \log (\gamma \lor 2))^{2(\gamma+1)+1}$, we have $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$. Note that if $\gamma = 0$ or $\gamma = 1$, as long as $\frac{n}{\sigma^2} \succeq 1$, we have $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$.

The lower bound. Again, the lower bound $\left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$ is well known and hence we only show the lower bound $\frac{\sigma^2(\gamma \lor 1)\log(\gamma \lor 2)}{n}$ under the condition $(\gamma \lor 1)^{2\gamma+1} < \frac{n}{\sigma^2} \precsim ((\gamma \lor 1)\log(\gamma \lor 2))^{2(\gamma+1)+1}$. If $\gamma = 0$ (or $\gamma = 1$), then as long as $\frac{n}{\sigma^2} \succeq 1$, we have $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{2}{3}}$ (respectively, $r^2 \asymp \left(\frac{\sigma^2}{n}\right)^{\frac{4}{5}}$) in the derivation of the upper bound. The minimax lower bound is trivially $\left(\frac{\sigma^2}{n}\right)^{\frac{2}{3}}$ (respectively, $\left(\frac{\sigma^2}{n}\right)^{\frac{4}{5}}$). Therefore, in what follows, we assume $\gamma > 1$.

If $\frac{n}{\sigma^2} > \gamma^{2\gamma+1}$, we have $\log N_{KL}(\epsilon, \mathcal{Q}_{\mathcal{U}}) \precsim \left(\frac{\sqrt{n}}{\sigma\epsilon}\right)^{\frac{1}{\gamma+1}}$ for any $\gamma \in \{0, ..., \beta\}$ by choosing a sufficiently small ϵ . Setting $\left(\frac{\sqrt{n}}{\sigma\epsilon}\right)^{\frac{1}{\gamma+1}} \asymp \epsilon^2$ yields $\epsilon^2 \asymp \left(\frac{n}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}} := \epsilon^{*2}$. Whenever

$$\delta \precsim 2^{-\gamma},$$
 (41)

we have

$$T_1(\delta) - T_2 \succeq \gamma^2$$

Let us consider $\delta^{*2} \simeq \frac{\sigma^2 \gamma \log \gamma}{n}$. Observe that $\delta^* \preceq 2^{-\gamma}$ if

$$\frac{n}{\sigma^2} \gtrsim 2^{2\gamma} \gamma \log \gamma \tag{42}$$

and hence

$$T_1(\delta^*) - T_2 \succeq \gamma \log \gamma \quad \text{since } \gamma^2 \ge \gamma \log \gamma.$$
 (43)

In view of (43), if

$$\gamma \log \gamma \succeq \epsilon^{*2} \asymp \left(\frac{n}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}},$$

which is equivalent to

$$\frac{n}{\sigma^2} \precsim (\gamma \log \gamma)^{2(\gamma+1)+1}, \qquad (44)$$

then we obtain

$$1 - \frac{\log 2 + \log N_{KL}\left(\epsilon^*, \mathcal{Q}_{\mathcal{U}}\right) + \epsilon^{*2}}{\log M_2\left(\delta^*, \mathcal{U}_{\gamma+1}\right)} \ge \frac{1}{2}.$$

Therefore,

$$\min_{\tilde{f}} \max_{f \in \mathcal{U}_{\gamma+1}} \mathbb{E}\left(\left|\tilde{f} - f\right|_2^2\right) \succeq \frac{\sigma^2 \gamma \log \gamma}{n}.$$
(45)

5.7Proof for Theorem 3.1

In the special case of $R_{\gamma+1} = 1$, the argument below sharpens the upper bound for $\log N_2(\delta, \mathcal{H}_{\gamma+1})$ in [16] from $(\gamma \vee 1) \delta^{-\frac{1}{\gamma+1}}$ to $\delta^{-\frac{1}{\gamma+1}}$. We find the cause of the gap lies in that the "pivotal" eigenvalue (that balances the "estimation error" and the "approximation error" from truncating for a given resolution δ) in [16] is not optimal. We close the gap by finding the optimal "pivotal" eigenvalue.

More generally, for the case of $R_{\gamma+1} \preceq \gamma+1$, we consider two different truncations, one giving the upper bound $\delta^{\frac{-1}{\gamma+1}}$ and the other giving the lower bound $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$. Note that $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$ $\delta^{\frac{-1}{\gamma+1}}$ when $R_{\gamma+1} \approx 1$. For the case of $R_{\gamma+1} \succeq \gamma+1$, we use only one truncation to show that both the upper bound and the lower bound scale as $(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}}$. In view of (3), given $(\phi_m)_{m=1}^{\infty}$ and $(\mu_m)_{m=1}^{\infty}$, to compute $N_2(\delta, \mathcal{H}_{\gamma+1})$, it suffices to compute

 $N_2(\delta, \mathcal{E}_{\gamma+1})$ where

$$\mathcal{E}_{\gamma+1} = \left\{ (\theta_m)_{m=1}^{\infty} : \sum_{m=1}^{\infty} \frac{\theta_m^2}{\mu_m} \le R_{\gamma+1}^2, \, \mu_m = (cm)^{-2(\gamma+1)} \right\}.$$

Let us introduce the M-dimensional ellipsoid

 $\overline{\mathcal{E}}_{\gamma+1} = \left\{ (\theta_m)_{m=1}^M \text{ coincide with the first } M \text{ elements of } (\theta_m)_{m=1}^\infty \text{ in } \mathcal{E}_{\gamma+1} \right\}$

where $M (= M (\gamma + 1, \delta))$ is the smallest integer such that, for a given resolution $\delta > 0$ and weight $w_{\gamma+1}, w_{\gamma+1}^2 \delta^2 \ge \mu_M$. In other words, $\mu_m \ge w_{\gamma+1}^2 \delta^2$ for all indices $m \le M$. Consequently, we have: (1)

$$\mathbb{B}_{2}^{M}\left(w_{\gamma+1}R_{\gamma+1}\delta\right) \subseteq \overline{\mathcal{E}}_{\gamma+1};\tag{46}$$

(2)
$$\mu_{M-1} = (c(M-1))^{-2(\gamma+1)} > w_{\gamma+1}^2 \delta^2$$
 and $\mu_{M-1} = (c(M+1))^{-2(\gamma+1)} < w_{\gamma+1}^2 \delta^2$, which yield

$$M \asymp (w_{\gamma+1}\delta)^{-\frac{1}{\gamma+1}}.$$
(47)

Note that (46), (47), and the fact $\mathcal{E}_{\gamma+1} \supseteq \overline{\mathcal{E}}_{\gamma+1}$ give

$$\log N_2\left(\delta, \mathcal{E}_{\gamma+1}\right) \ge \log N_2\left(\delta, \overline{\mathcal{E}}_{\gamma+1}\right)$$

$$\succeq M \log\left(w_{\gamma+1}R_{\gamma+1}\right)$$

$$\approx \left(w_{\gamma+1}\delta\right)^{-\frac{1}{\gamma+1}}\log\left(w_{\gamma+1}R_{\gamma+1}\right).$$
(48)

In the following, let $A_1 + A_2 := \{a_1 + a_2 : a_1 \in A_1, a_2 \in A_2\}$ for sets A_1 and A_2 . For the upper bound, we have

$$N_{2}\left(\delta, \overline{\mathcal{E}}_{\gamma+1}\right) \leq \frac{\operatorname{vol}\left(\frac{2}{\delta}\overline{\mathcal{E}}_{\gamma+1} + \mathbb{B}_{2}^{M}\left(1\right)\right)}{\operatorname{vol}\left(\mathbb{B}_{2}^{M}\left(1\right)\right)}$$

$$\leq \left(\frac{2}{\delta}\right)^{M} \frac{\operatorname{vol}\left(\overline{\mathcal{E}}_{\gamma+1} + \mathbb{B}_{2}^{M}\left(\frac{\delta}{2}\right)\right)}{\operatorname{vol}\left(\mathbb{B}_{2}^{M}\left(1\right)\right)}$$

$$\leq \left(\frac{2}{\delta}\right)^{M} \max\left\{\frac{\operatorname{vol}\left(2\overline{\mathcal{E}}_{\gamma+1}\right)}{\operatorname{vol}\left(\mathbb{B}_{2}^{M}\left(1\right)\right)}, \frac{\operatorname{vol}\left(2\mathbb{B}_{2}^{M}\left(\frac{\delta}{2}\right)\right)}{\operatorname{vol}\left(\mathbb{B}_{2}^{M}\left(1\right)\right)}\right\}$$

$$\leq \max\left\{\left(\frac{4R_{\gamma+1}}{\delta}\right)^{M} \prod_{m=1}^{M} \sqrt{\mu_{m}}, 2^{M}\right\}$$
(49)

where the first inequality follows from the standard volumetric argument, and the last inequality follows from the standard result for the volume of ellipsoids. The fact $\mu_m = (cm)^{-2(\gamma+1)}$ and the elementary inequality $\sum_{m=1}^{M} \log m \ge M \log M - M$ give

$$\log\left[\left(\frac{4R_{\gamma+1}}{\delta}\right)^{M}\prod_{m=1}^{M}\sqrt{\mu_{m}}\right] \leq M\left(\log\left(4R_{\gamma+1}\right)+\gamma+1\right)+$$

$$M\left(\log\frac{1}{\delta}-(\gamma+1)\log\left(cM\right)\right)$$

$$= M\left(\log\left(4R_{\gamma+1}\right)+\gamma+1\right)+$$

$$M\left(\log\frac{1}{\delta}-(\gamma+1)\log\left(cM\right)+\log\frac{1}{w_{\gamma+1}}-\log\frac{1}{w_{\gamma+1}}\right)$$

$$\leq M\left(\log 4R_{\gamma+1}+\gamma+1\right)+M\log w_{\gamma+1}$$

$$\lesssim M\log\left(w_{\gamma+1}\left((\gamma+1)\vee R_{\gamma+1}\right)\right)$$
(50)

where we have used the fact $\mu_M = (cM)^{-2(\gamma+1)} \leq w_{\gamma+1}^2 \delta^2$ in the second inequality. Inequalities (47), (49) and (50) together yield

$$\log N_2\left(\delta, \overline{\mathcal{E}}_{\gamma+1}\right) \precsim (w_{\gamma+1}\delta)^{-\frac{1}{\gamma+1}} \max\left\{\log\left(w_{\gamma+1}\left((\gamma+1)\vee R_{\gamma+1}\right)\right), \log 2\right\}.$$

For any $\theta \in \mathcal{E}_{\gamma+1}$, note that for a given δ , we have

$$\sum_{m=M+1}^{\infty} \theta_m^2 \le \mu_M \sum_{m=M+1}^{\infty} \frac{\theta_m^2}{\mu_m} \le w_{\gamma+1}^2 R_{\gamma+1}^2 \delta^2.$$
(51)

To cover $\mathcal{E}_{\gamma+1}$ within $(1 + w_{\gamma+1}^2 R_{\gamma+1}^2)^{\frac{1}{2}} \delta$ -precision, we find a smallest δ -cover of $\overline{\mathcal{E}}_{\gamma+1}$, $\{\theta^1, ..., \theta^N\}$, such that for any $\theta \in \mathcal{E}_{\gamma+1}$, there exists some *i* from the covering set with

$$\left|\theta - \theta^{i}\right|_{2}^{2} \leq \sum_{m=1}^{M} \left(\theta_{m} - \theta_{m}^{i}\right)^{2} + w_{\gamma+1}^{2} R_{\gamma+1}^{2} \delta^{2} \leq \left(1 + w_{\gamma+1}^{2} R_{\gamma+1}^{2}\right) \delta^{2}$$

where we have used (51). Consequently, we have

$$\log N_2\left(\delta, \mathcal{E}_{\gamma+1}\right)$$

$$\lesssim \log N_2\left(\delta\left(1+w_{\gamma+1}^2 R_{\gamma+1}^2\right)^{\frac{-1}{2}}, \overline{\mathcal{E}}_{\gamma+1}\right)$$

$$\lesssim \left(w_{\gamma+1}\delta\left(1+w_{\gamma+1}^2 R_{\gamma+1}^2\right)^{\frac{-1}{2}}\right)^{-\frac{1}{\gamma+1}} \max\left\{\log\left(w_{\gamma+1}\left((\gamma+1)\vee R_{\gamma+1}\right)\right), \log 2\right\}.$$
(52)

Case 1: $R_{\gamma+1} \succeq \gamma + 1$. Setting $w_{\gamma+1} \simeq R_{\gamma+1}^{-1}$ in (48) and (52) solves

$$\left(w_{\gamma+1}\delta \left(1 + w_{\gamma+1}^2 R_{\gamma+1}^2 \right)^{\frac{-1}{2}} \right)^{-\frac{1}{\gamma+1}} \max \left\{ \log \left(w_{\gamma+1} \left((\gamma+1) \lor R_{\gamma+1} \right) \right), \log 2 \right\}$$

$$\approx (w_{\gamma+1}\delta)^{-\frac{1}{\gamma+1}} \log \left(w_{\gamma+1}R_{\gamma+1} \right)$$
 (53)

and gives

$$\log N_2\left(\delta, \, \mathcal{E}_{\gamma+1}\right) \asymp \left(R_{\gamma+1}\delta^{-1}\right)^{\frac{1}{\gamma+1}}.$$

Case 2: $R_{\gamma+1} \preceq \gamma + 1$. Setting $w_{\gamma+1} \asymp (\gamma+1)^{-1}$ in (52) gives

$$\log N_2\left(\delta, \mathcal{E}_{\gamma+1}\right) \precsim \delta^{\frac{-1}{\gamma+1}}.$$

Note that the lower bound obtained by setting $w_{\gamma+1} \approx (\gamma+1)^{-1}$ in (48) is not particularly useful. Instead, we consider a different truncation with $w_{\gamma+1} \approx R_{\gamma+1}^{-1}$. Then (48) with $w_{\gamma+1} \approx R_{\gamma+1}^{-1}$ gives

$$\log N_2\left(\delta, \, \mathcal{E}_{\gamma+1}\right) \succsim R_{\gamma+1}^{\frac{1}{\gamma+1}} \delta^{\frac{-1}{\gamma+1}}.$$

5.8 Proof for Theorem 3.2

The upper bound. For the upper bound associated with $\mathcal{H}_{\gamma+1}$, we have

$$r^{2} \asymp R_{\gamma+1}^{\frac{2}{2(\gamma+1)+1}} \left(\frac{\sigma^{2}}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}.$$
(54)

The lower bound. Setting $\left(\frac{R_{\gamma+1}\sqrt{n}}{\sigma\epsilon}\right)^{\frac{1}{\gamma+1}} \approx \epsilon^2$ yields $\epsilon^2 \approx \left(\frac{nR_{\gamma+1}^2}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}} =: \epsilon^{*2}$. Observe that setting

$$\delta \asymp R_{\gamma+1}^{\frac{1}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{\gamma+1}{2(\gamma+1)+1}}$$

ensures

$$(R_{\gamma+1}\delta^{-1})^{\frac{1}{\gamma+1}} \asymp R_{\gamma+1}^{\frac{2}{2\gamma+3}} \left(\frac{n}{\sigma^2}\right)^{\frac{1}{2(\gamma+1)+1}} \succeq \epsilon^{*2}.$$

Consequently, we have

$$1 - \frac{\log 2 + \log N_{KL}\left(\epsilon^*, \mathcal{Q}_{\mathcal{H}}\right) + \epsilon^{*2}}{\log M_2\left(\delta^*, \mathcal{H}_{\gamma+1}\right)} \ge \frac{1}{2}$$

and

$$\min_{\tilde{f}} \max_{f \in \mathcal{H}_{\gamma+1}} \mathbb{E}\left(\left|\tilde{f} - f\right|_2^2\right) \succeq R_{\gamma+1}^{\frac{2}{2(\gamma+1)+1}} \left(\frac{\sigma^2}{n}\right)^{\frac{2(\gamma+1)}{2(\gamma+1)+1}}$$

5.9 Proof for Lemma 4.1

Like in Section (5.2), the proper choice of the grid of points on each dimension of $[-1, 1]^d$ is the key in this case. Any function $f \in \mathcal{U}^d_{\gamma+1,2}$ can be written as

$$f(x + \Delta) = \sum_{k=0}^{\gamma} \sum_{p:P=k} \frac{\Delta^p D^p f(x)}{k!} + \sum_{\substack{p:P=\gamma \\ y! = P = \gamma}} \left[\frac{\Delta^p D^p f(z)}{\gamma!} - \frac{\Delta^p D^p f(x)}{\gamma!} \right]_{:=REM_0(x + \Delta)}$$

where $x, x + \Delta \in (-1, 1)^d$ and z is some intermediate value. For a given $k \in \{0, ..., \gamma\}$, recall $\operatorname{card}(\{p : P = k\}) = \begin{pmatrix} d+k-1\\ d-1 \end{pmatrix} = D_k^*$. Therefore, we have

$$|REM_0(x+\Delta)| \le \frac{D_{\gamma}^* R_{\gamma+1} |\Delta|_{\infty}^{\gamma+1}}{\gamma!}.$$
(55)

In a similar way, writing

$$D^{\tilde{p}}f(x+\Delta) = \sum_{k=0}^{\gamma-\tilde{P}} \sum_{p:P=k} \frac{\Delta^p D^{p+\tilde{p}}f(x)}{k!} + \sum_{\substack{p:P=\gamma-\tilde{P}\\ (\gamma-\tilde{P})! \\ :=REM_{\tilde{P}}(x+\Delta)}} \left[\frac{\Delta^p D^{p+\tilde{p}}f(\tilde{z})}{(\gamma-\tilde{P})!} - \frac{\Delta^p D^{p+\tilde{p}}f(x)}{(\gamma-\tilde{P})!} \right]$$

for $1 \leq \tilde{P} := \sum_{j=1}^{d} \tilde{p}_j \leq \gamma$ and $\tilde{p} = (\tilde{p}_j)_{j=1}^d$, we have

$$\left|REM_{\tilde{P}}(x+\Delta)\right| \leq \frac{D_{\gamma-\tilde{P}}^* R_{\gamma-\tilde{P}+1} \left|\Delta\right|_{\infty}^{\gamma+1-P}}{\left(\gamma-\tilde{P}\right)!}.$$
(56)

For some $\delta_0, \ldots, \delta_{\gamma} > 0$, suppose that $|D^p f(w) - D^p g(w)| \le \delta_k$ for all p with $P = k \in \{0, \ldots, \gamma\}$, where $f, g \in \mathcal{U}^d_{\gamma+1,2}$. Then we have

$$\begin{aligned} &|f(x+\Delta) - g(x+\Delta)| \\ &\leq \left| \sum_{k=0}^{\gamma} \sum_{p:P=k} \frac{\Delta^p}{k!} \left(D^p f\left(x\right) - D^p g\left(x\right) \right) \right| + 2 \frac{D_{\gamma}^* R_{\gamma+1} \left|\Delta\right|_{\infty}^{\gamma+1}}{\gamma!} \\ &\leq \left| \sum_{k=0}^{\gamma} \frac{D_k^* \left|\Delta\right|_{\infty}^k \delta_k}{k!} + 2 \frac{D_{\gamma}^* R_{\gamma+1} \left|\Delta\right|_{\infty}^{\gamma+1}}{\gamma!}. \end{aligned}$$

Let $\left(\max_{k \in \{1,\ldots,\gamma+1\}} \frac{D_{k-1}^* R_k}{(k-1)!} \vee 1\right) =: R^*$. Consider $|\Delta|_{\infty} \leq d^{-1} \left(R^{*-1}\delta\right)^{\frac{1}{\gamma+1}}$ and $\delta_k = R^* \frac{k}{\gamma+1} \delta^{1-\frac{k}{\gamma+1}}$ for $k = 0, \ldots, \gamma$ and δ such that $\delta_k \in (0, 1)$. Then,

$$f(x + \Delta) - g(x - \Delta)| \leq \delta \sum_{k=0}^{\gamma} \left(R^* \frac{-k+k}{\gamma+1} \frac{1}{k!} \right) + 2R^* |\Delta|_{\infty}^{\gamma+1}$$
$$\leq \delta \sum_{k=0}^{\gamma} \frac{1}{k!} + 2\delta \leq 5\delta$$
(57)

where we have used the fact that $D_k^* \leq d^k$. On each dimension of $[-1, 1]^d$, we consider a $d^{-1} \left(R^{*-1}\delta\right)^{\frac{1}{\gamma+1}}$ -grid of points. The rest of the arguments follow closely those in [3].

5.10 Proof for Lemma 4.2

For a given $k \in \{0, ..., \gamma\}$, let card $(\{p : P = k\}) = \begin{pmatrix} d+k-1 \\ d-1 \end{pmatrix} = \begin{pmatrix} d+k-1 \\ k \end{pmatrix} = D_k^*$. Recall the definition of $\mathcal{U}_{\gamma+1,1}^d$:

$$\mathcal{U}_{\gamma+1,1}^{d} = \left\{ f = \sum_{k=0}^{\gamma} \sum_{p:P=k} x^{p} \theta_{(p,k)} : \{\theta_{(p,k)}\}_{(p,k)} \in \mathcal{P}_{\Gamma}, \ x \in [-1, 1]^{d} \right\}$$

with the $\Gamma:=\sum_{k=0}^{\gamma}D_k^*-\text{dimensional polyhedron}$

$$\mathcal{P}_{\Gamma} = \left\{ \{\theta_{(p,k)}\}_{(p,k)} \in \mathbb{R}^{\Gamma} : \text{for any given } k, \, \{\theta_{(p,k)}\}_{p} \in \left[\frac{-R_{k}}{k!}, \, \frac{R_{k}}{k!}\right] \right\}$$

where $\theta = \{\theta_{(p,k)}\}_{(p,k)}$ denotes the collection of $\theta_{(p,k)}$ over all (p,k) configurations and $\{\theta_{(p,k)}\}_p$ denotes the collection of $\theta_{(p,k)}$ over all p configurations for a given $k \in \{0, ..., \gamma\}$.

To bound $\log N_{\infty}\left(\delta, \mathcal{U}_{\gamma+1,1}^d\right)$ from above, note that for $f, f' \in \mathcal{U}_{\gamma+1,1}^d$, we have

$$\left|f - f'\right|_{\infty} \le \sum_{k=0}^{\gamma} \sum_{p:P=k} \left|\theta_{(p,k)} - \theta'_{(p,k)}\right|$$

where $f' = \sum_{k=0}^{\gamma} \sum_{p:P=k} x^p \theta'_{(p,k)}$ such that $\theta' = \{\theta'_{(p,k)}\}_{(p,k)} \in \mathcal{P}_{\Gamma}$. Therefore, the problem is reduced to finding $N_1(\delta, \mathcal{P}_{\Gamma})$.

To cover \mathcal{P}_{Γ} within δ -precision, using arguments similar to those in Section 5.1, we find a smallest $\frac{\delta}{(\gamma+1)D_k^*}$ -cover of $\left[\frac{-R_k}{k!}, \frac{R_k}{k!}\right]$ for each $k = 0, ..., \gamma$, $\left\{\theta_k^1, ..., \theta_k^{N_k}\right\}$, such that for any $\theta \in \mathcal{P}_{\Gamma}$, there exists some $i_{(p,k)} \in \{1, ..., N_k\}$ with

$$\sum_{k=0}^{\gamma} \sum_{p:P=k} \left| \theta_{(p,k)} - \theta_k^{i_{(p,k)}} \right| \le \delta.$$

As a consequence, we have

$$\log N_1\left(\delta, \mathcal{P}_{\Gamma}\right) \le \sum_{k=0}^{\gamma} D_k^* \log \frac{4\left(\gamma+1\right) D_k^* R_k}{\delta k!}$$

and

$$\log N_2\left(\delta, \, \mathcal{U}^d_{\gamma+1,1}\right) \le \log N_\infty\left(\delta, \, \mathcal{U}^d_{\gamma+1,1}\right) \le \sum_{k=0}^{\gamma} D_k^* \log \frac{4\left(\gamma+1\right) D_k^* R_k}{\delta k!}$$

If δ is large enough such that $\min_{k \in \{0,...,\gamma\}} \log \frac{4(\gamma+1)D_k^*R_k}{\delta k!} < 0$, we use the counting argument in [3] to obtain

$$\log N_2\left(\delta, \, \mathcal{U}^d_{\gamma+1,1}\right) \leq \log N_\infty\left(\delta, \, \mathcal{U}^d_{\gamma+1,1}\right) \precsim \left(\sum_{k=0}^{\gamma} D_k^*\right) \log \frac{1}{\delta} + \sum_{k=0}^{\gamma} D_k^* \log R_k.$$

5.11 Lemma A.1 and its proof

Lemma A.1. Let $\{\phi_k\}_{k=1}^{\infty}$ be the Legendre polynomials on [-1, 1]. For any $f \in \mathcal{U}_{\gamma+1,1}[-1, 1]$, we have $f(x) = \sum_{k=0}^{\gamma} \tilde{\theta}_k \phi_k(x)$ such that

$$\tilde{\theta}_k = \left(k + \frac{1}{2}\right) \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \frac{f^{(k+2m)}(0)}{2^{k+2m}m! \left(\frac{1}{2}\right)_{k+m+1}}$$

where $(a)_k = a(a+1)\cdots(a+k-1)$ is known as the Pochhammer symbol.

Proof. To obtain the correct formula for finite sums, we carefully modify the derivations for Theorem 2 in [2] which concerns infinite sums. The Legendre expansion of x^k yields

$$\frac{x^k}{k!} = \frac{1}{2^k} \sum_{m=0}^{\lfloor k/2 \rfloor} \frac{k - 2m + \frac{1}{2}}{m! \left(\frac{1}{2}\right)_{k-m+1}} \phi_{k-2m}(x).$$
(58)

First, let us consider the case where γ is odd. Applying (58) gives

$$\begin{split} f(x) &= \sum_{k=0}^{\gamma} \frac{f^{(k)}(0)}{2^k} \sum_{m=0}^{\lfloor k/2 \rfloor} \frac{k-2m+\frac{1}{2}}{m! (\frac{1}{2})_{k-m+1}} \phi_{k-2m}(x) \\ &= \sum_{k=0}^{\lfloor \gamma/2 \rfloor} \frac{f^{(2k)}(0)}{2^{2k}} \sum_{m=0}^{k} \frac{2k-2m+\frac{1}{2}}{m! (\frac{1}{2})_{2k-m+1}} \phi_{2k-2m}(x) \quad (\text{even } k) \\ &+ \sum_{k=0}^{\lfloor \gamma/2 \rfloor} \frac{f^{(2k+1)}(0)}{2^{2k+1}} \sum_{m=0}^{k} \frac{2k-2m+\frac{3}{2}}{m! (\frac{1}{2})_{2k-m+2}} \phi_{2k-2m+1}(x) \quad (\text{odd } k) \quad (59) \\ &= \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \sum_{k=m}^{\lfloor \gamma/2 \rfloor} \frac{2k-2m+\frac{1}{2}}{m! (\frac{1}{2})_{2k-m+1}} \frac{f^{(2k)}(0)}{2^{2k}} \phi_{2k-2m}(x) \quad (\text{interchanging sums}) \\ &+ \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \sum_{k=m}^{\lfloor \gamma/2 \rfloor} \frac{2k-2m+\frac{3}{2}}{m! (\frac{1}{2})_{2k-m+1}} \frac{f^{(2k+1)}(0)}{2^{2k+1}} \phi_{2k-2m+1}(x) \\ &= \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \sum_{l=0}^{\lfloor \gamma/2 \rfloor} \frac{2l+\frac{1}{2}}{m! (\frac{1}{2})_{2l+m+1}} \frac{f^{(2l+2m)}(0)}{2^{2l+2m}} \phi_{2l}(x) \quad (\text{letting } l=k-m) \\ &+ \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \sum_{l=0}^{\lfloor \gamma/2 \rfloor} \frac{2l+\frac{3}{2}}{m! (\frac{1}{2})_{2l+m+2}} \frac{f^{(2l+2m+1)}(0)}{2^{2l+2m+1}} \phi_{2l+1}(x) \\ &= \sum_{l=0}^{\lfloor \gamma/2 \rfloor} \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \frac{2l+\frac{1}{2}}{m! (\frac{1}{2})_{2l+m+2}} \frac{f^{(2l+2m+1)}(0)}{2^{2l+2m}} \phi_{2l}(x) \quad (\text{interchanging sums}) \\ &+ \sum_{l=0}^{\lfloor \gamma/2 \rfloor} \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \frac{2l+\frac{3}{2}}{m! (\frac{1}{2})_{2l+m+2}} \frac{f^{(2l+2m+1)}(0)}{2^{2l+2m}} \phi_{2l}(x) \quad (\text{interchanging sums}) \\ &+ \sum_{l=0}^{\lfloor \gamma/2 \rfloor} \sum_{m=0}^{\lfloor \gamma/2 \rfloor} \frac{2l+\frac{3}{2}}{m! (\frac{1}{2})_{2l+m+2}} \frac{f^{(2l+2m+1)}(0)}{2^{2l+2m}} \phi_{2l+1}(x) \end{split}$$

which gives the claim in Lemma A.1.

For the case of even γ , note that the term in (59) takes the form

$$\sum_{k=0}^{\lfloor \gamma/2 \rfloor} \frac{f^{(2k+1)}(0)}{2^{2k+1}} \sum_{m=0}^{k} \frac{2k-2m+\frac{3}{2}}{m! \left(\frac{1}{2}\right)_{2k-m+2}} \phi_{2k-2m+1}(x)$$

$$= \sum_{k=0}^{\lfloor \gamma/2 \rfloor -1} \frac{f^{(2k+1)}(0)}{2^{2k+1}} \sum_{m=0}^{k} \frac{2k-2m+\frac{3}{2}}{m! \left(\frac{1}{2}\right)_{2k-m+2}} \phi_{2k-2m+1}(x)$$

$$+ \underbrace{\frac{f^{(\gamma+1)}(0)}{2^{\gamma+1}}}_{0} \sum_{m=0}^{\gamma/2} \frac{\gamma-2m+\frac{3}{2}}{m! \left(\frac{1}{2}\right)_{\gamma-m+2}} \phi_{\gamma-2m+1}(x)$$

and hence the previous derivations go through.

References

- P. Bartlett and S. Mendelson (2002). "Gaussian and Rademacher complexities: risk bounds and structural results". *Journal of Machine Learning Research*, 3, 463–482.
- [2] M. Cantero and A. Iserles (2012). "On rapid computation of expansions in ultrasperical polynomials". SIAM Journal on Numerical Analysis, 50(1), 307–327.
- [3] A. Kolmogorov and B. Tikhomirov (1959). "ε-entropy and ε-capacity of sets in functional spaces". Uspekhi Mat. Nauk., 86, 3-86. Appeared in English as American Mathematical Society Translations, 17, 277–364, 1961.
- [4] M. Ledoux and M. Talagrand (1991). Probability in Banach spaces: isoperimetry and processes. Springer, New York, NY.
- [5] H. Liang and H. Wu (2008). "Parameter estimation for differential equation models using a framework of measurement error in regression models". *Journal of the American Statistical Association*, 103:484, 1570–1583.
- [6] A. A. Poyton, M. S. Varziri, K. B. McAuley, P. J. McLellan, J. O. Ramsay (2006) "Parameter estimation in continuous-time dynamic models using principal differential analysis". *Computers* & Chemical Engineering, 30(4), 698–708.
- [7] J. O. Ramsay and B. W. Silverman (2005). Fitting differential equations to functional data: Principal differential analysis. In: Functional Data Analysis. Springer, New York, NY.
- [8] J. O. Ramsay (1996). "Principal differential analysis: Data reduction by differential operators". Journal of the Royal Statistical Society, Series B, 58, 495–508.
- [9] J. Ramsay, G. Hooker, D. Campbell and J. Cao (2007). "Parameter estimation for differential equations: A generalized smoothing approach". *Journal of the Royal Statistical Society*, 69, 741–796.
- [10] I. Steinwart and A. Christmann (2008). Support Vector Machines. Springer, New York.
- [11] C. J. Stone (1982). "Optimal global rates of convergence for nonparametric regression". Annals of Statistics, 10, 1040–1053.

- [12] A. B. Tsybakov (2009). Introduction to nonparametric estimation. New York, NY: Springer.
- [13] S. van de Geer (2000). Empirical Processes in M-Estimation. Cambridge University Press.
- [14] J. M. Varah (1982). "A spline least squares method for numerical parameter estimation in differential equations". SIAM Journal on Scientific and Statistical Computing, 3(1), 28–46.
- [15] G. Wahba (1990). Spline models for observational data. SIAM, CBMS-NSF Regional Conference Series in Applied Mathematics. Philadelphia, PA.
- [16] M. J. Wainwright (2019). High-dimensional statistics: A non-asymptotic viewpoint. Cambridge University Press.
- [17] Y. Yang and A. Barron (1999). "Information-theoretic determination of minimax rates of convergence". Annals of Statistics, 27(5), 1564–1599.
- [18] M. Yuan and T. Cai (2010). "A reproducing kernel Hilbert space approach to functional linear regression". Annals of Statistics, 38(6), 3412–3444.
- [19] Y. Zhu and M. Mirzaei (2021). "Classes of ODE solutions: smoothness, covering numbers, implications for noisy function fitting, and a 'loss of smoothness' phenomenon". https://sites.google.com/view/ying-zhu (also arXiv:2011.11371v3).