

# Dark Exciton Giant Rabi Oscillations with no External Magnetic Field

Vladimir Vargas-Calderón\* and Herbert Vinck-Posada  
*Grupo de Superconductividad y Nanotecnología, Departamento de Física  
Universidad Nacional de Colombia, 111321, Bogotá, Colombia*

J. M. Villas-Boas  
*Instituto de Física, Universidade Federal de Uberlândia, 38400-902, MG, Brazil*  
(Dated: December 8, 2021)

Multi-phonon physics is an emerging field that serves as a test bed for fundamental quantum physics and several applications in metrology, on-chip communication, among others. Quantum acoustic cavities or resonators are devices that are being used to study multi-phonon phenomena both theoretically and experimentally. In particular, we study a system consisting of a semiconductor quantum dot pumped by a driving laser, and coupled to an acoustic cavity. This kind of systems have proven to yield interesting multi-phonon phenomena, but the description of the quantum dot has been limited to a two-level system. This limitation restrains the complexity that a true semiconductor quantum dot can offer. Instead, in this work we consider a model where the quantum dot can have both bright and dark excitons, the latter being particularly useful due to their lower decoherence times, because they do not present spontaneous photon emission. In this setup, we demonstrate that by fine-tuning the driving laser frequency, one is able to realise giant Rabi oscillations between the vacuum state and a dark exciton state with  $N$ -phonon bundles. From this, we highlight two outstanding features: first, we are able to create dark states excitations in the quantum dot without the usual external magnetic field needed to do so; and second, in a dissipative scenario where the acoustic cavity and the quantum dot suffer from losses, the system acts as a phonon gun able to herald  $N$ -phonon bundles.

*Introduction*— Quantum vibrational modes of solids, described by phonons, have a great potential to be used in technological applications in metrology or quantum information processing [1–6]. Moreover, as the control of individual phonons continues to improve, the study of phonons and their interaction with other excitations in many-body quantum systems is also relevant for testing fundamental physics [7, 8]. Analogous to photons in quantum electrodynamics, phonons can be used to store, process, and transduce quantum information. The inclusion of phonons to the quantum toolbox gives a three-fold advantage: first, losses by radiation into the electromagnetic field vacuum are no longer present, as phonons can only propagate through some material medium (usually in solid-state devices); second, phonon energy scales are, in general, different from the optical energy scale, making phonons especially suited for on-chip communication [9, 10] in a variety of characteristic energies, from MHz to THz [11–16]; third, many experimental techniques developed by solid-state physicists [7, 16–18] become available for quantum information processing tasks with phonons [10].

Even though the majority of theoretical and experimental efforts have been devoted to single-phonon generation and control, many-phonon states are also required for highly non-classical sources, useful for quantum sensing and metrology [3], and quantum technologies such as quantum memories and transducers [4–6]. To this end, the recent work by Nori *et al.* [9] proposes a physical sys-

tem in quantum acoustodynamics composed of a semiconductor quantum dot (QD), modelled as a two-level system, coupled to the phonon mode of an acoustic cavity. The QD is coherently pumped by an external laser with a frequency that can be tuned to excite giant Rabi oscillations between a state that is mostly the vacuum state (no QD excitations and 0 phonons in the acoustic cavity), and a state that is composed of an exciton QD state, accompanied by  $N$  phonons in the acoustic cavity. The authors then show that dissipative channels allow the emission of  $N$ -phonon bundles, and analyse the quantum statistics of this emission, finding out that depending on dissipative and Hamiltonian parameters, they can realise a phonon laser or a phonon gun.

In this work we study a QD coherently pumped by an external laser and immersed in an acoustic single-mode cavity, but we take into account the richer exciton basis provided by the different spin alignments of the electron and hole that compose the exciton. Thus, the QD is described by a ground state, two bright exciton states, and two dark exciton states. The inclusion of this richer exciton basis allows the description of more complex interactions in the system, which allows the control of interesting phenomena. Therefore, the main contributions of our work, coming from the more complete exciton basis, are: dark excitons can be excited, taking advantage of the Bir-Pikus interaction [19], by fine-tuning the laser frequency without the need for an external magnetic field—which is both experimentally challenging as well as expensive [20, 21]; and giant-Rabi oscillations between a vacuum state and a dark exciton- $N$ -phonon state can be realised to herald  $N$ -phonon bundle emission.

---

\* vvargasc@unal.edu.co

*Model*— In this work we consider a pumped quantum dot embedded in an acoustic cavity described by the following Hamiltonian [9]:

$$H = H_{\text{QD}} + H_{\text{laser}} + H_{\text{cav}} + H_{\text{el-ph}}. \quad (1)$$

The bare quantum dot is described by a valence state  $|v\rangle$ , two bright exciton states  $|1\rangle$  and  $|2\rangle$  with anti-parallel electron-hole spins, and two dark exciton states  $|3\rangle$  and  $|4\rangle$  with parallel electron-hole spins. The corresponding Hamiltonian, taking into account the exchange interaction [22, 23], and making use of the ladder operators  $\sigma_{ij} = |i\rangle\langle j|$ , reads [20]

$$H_{\text{QD}} = \omega_X(\sigma_{11} + \sigma_{22}) + \omega_d(\sigma_{33} + \sigma_{44}) + \frac{\delta_1}{2}(\sigma_{12} + \sigma_{21}) + \frac{\delta_2}{2}(\sigma_{34} + \sigma_{43}), \quad (2)$$

$$H_{\text{el-ph}} = \left\{ g_{\text{bb}}[\sigma_{11} + \sigma_{22} + i(\sigma_{12} - \sigma_{21})] + \frac{g_{\text{bd}}}{\sqrt{2}}[(1+i)(\sigma_{13} + \sigma_{14}) + (1-i)(\sigma_{23} + \sigma_{24})] \right\} (b^\dagger + b) + \text{H.c.} \quad (3)$$

where  $g_{\text{bb}(\text{bd})}$  are bright-bright(bright-dark) exciton coupling rates through phonons.

*Results*— Such an electron-phonon coupling, accompanied by the optically driven Stokes process through the laser pumping, allows the excitation of giant Rabi oscillations between the vacuum state  $|\text{phonons} = 0\rangle \otimes |\text{QD} = v\rangle$  (for weak laser pumping) and an eigenstate of the Hamiltonian shown in eq. (1) [9]. In particular, one is able to target giant Rabi oscillations to states that have a large contribution from a state of -phonons and a symmetric dark exciton state  $|n, d_+\rangle$ . We show in fig. 1 two such states.




FIG. 1. Eigenstates of eq. (1) for weak electron-phonon coupling ( $g = g_{\text{bb}} = g_{\text{bd}} \approx 0.02\omega_b$ ) and pumping ( $\Omega_1 \approx 0.082\omega_b$ ,  $\Omega_2 = 0$ ). These values are used throughout the paper. The state  $|\lambda_{1(2)}\rangle$  is mostly composed by the state  $|2(3), d_+\rangle$ , where  $|d_\pm\rangle = (|3\rangle \pm |4\rangle)/\sqrt{2}$  is the dark exciton symmetric (+) or anti-symmetric (−) state.

where  $\omega_X$  is the bare bright exciton energy,  $\omega_d = \omega_X - \delta_0$  is the shifted bare dark exciton energy, and  $\delta_{1(2)}$  split the bright(dark) exciton energies. Further, the QD is driven by an external laser that pumps the bright exciton states through  $H_{\text{laser}} = \Omega_1(e^{-i\omega_L t}\sigma_{1v} + e^{i\omega_L t}\sigma_{v1}) + \Omega_2(e^{-i\omega_L t}\sigma_{2v} + e^{i\omega_L t}\sigma_{v2})$ . Here, the relative magnitudes of the laser amplitudes  $\Omega_1$  and  $\Omega_2$  depend on the laser polarisation [24]. The time dependence of the whole Hamiltonian is removed in the laser-frequency rotating frame via the unitary transformation  $U = \exp(i\omega_L t[\sigma_{11} + \sigma_{22} + \sigma_{33} + \sigma_{44}])$ . The acoustic cavity Hamiltonian accounts for the single phonon mode energy  $H_{\text{cav}} = \omega_b b^\dagger b$ , where  $b$  is the phonon annihilation operator. Throughout the paper we will use the values  $\delta_0 = 0.04\omega_b$ ,  $\delta_1 = 0.036\omega_b$  and  $\delta_2 = 0.01\omega_b$ , which match the values reported by Bayer *et al.* [23] for  $\omega_b = 5\text{meV}$ . Finally, we also consider the electron-phonon coupling described by the Bir-Pikus Hamiltonian [25]

Giant Rabi oscillations are achieved through a cascading effect [9], where the system transitions from the vacuum state  $|0, v\rangle$  to a QD bright exciton state (depending on the laser amplitudes), and then to a (mostly) dark exciton symmetric state with -phonons after the system is guided by the electron-phonon coupling mechanism. A couple of examples of such a giant Rabi oscillations can be found in fig. 2 for  $n = 2$  and  $n = 3$ . Both examples show that it is possible to tune the laser frequency to target giant Rabi oscillations between the vacuum state  $|0, v\rangle$ , and any eigenstate of eq. (1). It is worth noting that the states  $|n, d_-\rangle$ , i.e. with anti-symmetric dark exciton matter states, are eigenstates of eq. (1), but they are not accessible from non-dark anti-symmetric initial states, as can be easily corroborated by computing  $\langle m, \beta | H | n, d_-\rangle = 0$ , where  $|\beta\rangle$  is any state spanned by bright exciton states, and  $m$  is some number of phonons. However, in the presence of dissipation they can be accessed, as will be explored later.

The Hamiltonian analysis so far presented is only relevant to understand the underlying physics of giant Rabi oscillation production, but to study a realistic setup of the proposed system it is necessary to describe it as an open quantum system. Such a description allows understanding that giant Rabi oscillations can harvest  $N$ -phonon bundle excitations which can be emitted through dissipative channels. In the case of weak coupling to the environment, allowing the usual Born-Markov approximations [26], we consider four dissipative channels under the Gorini-Kossakowski-Sudarshan-Lindblad equa-




FIG. 2. Giant Rabi oscillations between the 0-phonon valence state (solid line) and the states shown in fig. 1 (dashed line). The dash-dotted line shows the evolution of the  $|n, d_+\rangle$  state for  $n = 2$  (top panel) and  $n = 3$  (bottom panel). The laser detuning is  $\Delta/\omega_b = (\omega_X - \omega_L)/\omega_b \approx -1.960$  for the top panel, and  $\approx -2.961$  for the bottom panel. The same coupling and pumping conditions from fig. 1 are used.

tion for the system's density operator [27, 28]

$$\begin{aligned} \frac{d\rho}{dt} = & i[\rho, H] + \kappa\mathcal{D}_b[\rho] + \gamma_b \sum_{j=1,2} \mathcal{D}_{\sigma_{vj}}[\rho] \\ & + \gamma_d \sum_{j=3,4} \mathcal{D}_{\sigma_{vj}}[\rho] + \gamma_\phi \sum_{j=1}^4 \mathcal{D}_{\sigma_{jj}}[\rho], \end{aligned} \quad (4)$$

where  $\mathcal{D}_A[\rho] = A\rho A^\dagger - \frac{1}{2}\rho A^\dagger A - \frac{1}{2}A^\dagger A\rho$  is the dissipator super-operator of the collapse operator  $A$ . The four dissipative channels considered in eq. (4) are: phonon escape from the acoustic nanocavity at a rate  $\kappa$  with a collapse operator  $b$  due to unwanted coupling with leaky modes [29, 30], spontaneous emission of the bright excitons at a rate  $\gamma_b$  with collapse operators  $\sigma_{vj}$  for  $j = 1, 2$ , effective spontaneous emission of the dark excitons (hole or electron spin flip followed by bright exciton spontaneous emission [31]) at a rate  $\gamma_d$  with collapse operators  $\sigma_{vj}$  for  $j = 3, 4$ , and pure dephasing of all exciton states at a rate  $\gamma_\phi$  with collapse operators  $\sigma_{jj}$  [32].

Solving eq. (4) using quantum trajectories [33, 34] exposes the  $N$ -phonon bundle nature of the excitations that are emitted: the  $N$ -phonon bundle behaves as a quasi-particle in the context of the dynamical process of emission [35]. Figure 3 depicts this concept by showing a single quantum trajectory that suffers a strongly-correlated phonon emission process. The initial state is the vacuum, as shown in fig. 3(b), right before a quantum jump that takes the system to a dark state with two phonons, as shown in fig. 3(c). Here the two-phonon state has been heralded. Now, dissipation makes the system undergo a quantum jump through the emission of a phonon, leaving the system in a dark state, but with only one phonon, as shown in fig. 3(d). Although the phonon-escape mechanism depletes the cavity phonon-

by-phonon, it is shortly after the emission of the first phonon that the second phonon is also emitted, leaving the system in a dark excitonic state with zero phonons, as shown in fig. 3(e). Finally, the QD emits a photon due to recombination, and the system is allowed to transit the same process once again. The two-phonon bundle emission process arises in all quantum trajectories, and becomes more common when the electron-phonon coupling  $g$  is increased, at the cost of realising giant Rabi oscillations between the vacuum and states that have a larger bright contribution. Nonetheless, we point out that for the presented parameters, one-phonon emission processes are more common [36].




FIG. 3. Evolution of a quantum trajectory due to Hamiltonian dynamics and dissipation-induced quantum jumps. (a) shows the occupation of the vacuum  $|0, v\rangle$  state, and of the  $N$ -phonon states, i.e. the occupation of  $\sum_i |N, i\rangle$ . (b)-(d) show the state composition of the system at the times pointed out by an arrow in panel (a). The laser detuning is the same as in the top panel of fig. 2. The dissipative parameters are  $\kappa \approx 7.89 \times 10^{-4}\omega_b$ ,  $\gamma_b \approx 1.869 \times 10^{-5}\omega_b$ ,  $\gamma_d = 0.1\gamma_b$  and  $\gamma_\phi = 4 \times 10^{-4}\omega_b$ . These parameters are used throughout the paper.

Moreover, as in the work by Bin *et al.* [9], a signature of strong correlations of the emitted phonons is found in the equal-time  $n$ -th order phonon correlation function  $g^{(n)}(\tau = 0) = \langle b^{\dagger n} b^n \rangle / (b^\dagger b)^n$  [37]. They noticed that near the resonances  $\Delta \approx -n\omega_b$  (for  $n \geq 2$ ), this corre-

lation function did not show the super-bunching peaks that would be expected for multi-phonon emission. Instead, there are dips in the middle of such peaks, as we show in fig. 4, signalling how the correlation function breaks down at specific resonances. In their case, only one dip corresponding to the resonance with the  $|n, c\rangle$  state was shown because their model was restricted only to one conduction  $|c\rangle$  exciton state, whereas in our case, we present three different dips corresponding to the resonances with the bright symmetric and anti-symmetric states, as well as the dark symmetric state. Nonetheless, as also pointed out by Fabrice, the breaking down of the usual equal-time correlation function is a signature of strong phonon-correlations that begin to form.




FIG. 4. Equal-time one- and two-phonon bundle correlation functions. At  $\Delta \approx -2\omega_b$  there are dips in the super-bunching peak of the usual correlation function at resonances corresponding to two-phonon states with bright (anti-)symmetric  $|b_{(-)+}\rangle$ , and with the dark symmetric state  $|d_+\rangle$ .

However, such correlation function is not adequate to investigate multi-phonon properties associated to the actual -phonon bundles that are excited. On the other hand, the equal-time  $m$ -th order -phonon bundle correlation function, defined as  $g_n^{(m)}(\tau = 0) =$

$\langle b^{\dagger nm} b^{nm} \rangle / \langle b^{\dagger n} b^n \rangle^m$  correctly describes these multi-phonon properties because it treats the -phonon bundle as a quasi-particle [35], with associated creation and annihilation operators  $b^{\dagger n}$  and  $b^n$ , respectively. In fig. 4 we show that this generalised correlation function reaches the sub-Poissonian anti-bunching regime at each of the resonances [38]. The resonance associated to the dark symmetric with two phonons state presents the lowest value of  $g_2^{(2)}$ , well within the anti-bunching regime. This shows that the Bir-Pikus mechanism allows the generation of robust anti-bunching behaviour when the laser frequency is tuned to target giant Rabi oscillations with the dark symmetric state, even more so than with the usually accessed bright states.

Finally, we examine the phonon emission spectrum (see fig. 5(c)) to show that the two-phonon emission processes can be frequency-resolved. We can interpret every single peak through the spectral theory of the Liouvillian superoperator  $\mathcal{L}$  that satisfies eq. (4), written as  $\frac{d\rho}{dt} = \mathcal{L}[\rho]$  [39–41]. A formal solution of this equation for a time-independent Liouvillian is  $\rho(t) = \sum_k e^{\Lambda_k t} \text{Tr}[\varrho_k \rho(0)] \varrho_k$ , where  $\Lambda_k$  are the complex eigenvalues of  $\mathcal{L}$  with corresponding eigenmatrices  $\varrho_k$ . The eigenvalues are associated to the emission peaks [42–44] as they show both the peak location  $\text{Im}\{\Lambda_k\}$  and the full width at half maximum  $-\text{Re}\{\Lambda_k\}$ . On the other hand, the eigenmatrices account for information about which states are involved in each transition [45]. These transitions can be systematically studied when they involve changes in the number of excitations through dissipative processes only [46–48]. In our case, such a theory cannot be used because of the presence of coherent pumping to the QD. Nonetheless, the examination of the eigenmatrix elements still shows which states are involved in the transitions, as displayed in fig. 5(a) and (b).

Figure 5(a) shows that the most prominent peak of the spectrum corresponds to transitions between dark states with one phonon and dark states with 0 phonons. It is worth noting that the magnitude of the matrix elements shown in fig. 5(a) and (b) do not indicate the contribution of those transitions to the emission peak. Instead, the excitation of the transitions depends upon the energy injected into the system, and the strength of the interactions that enable certain population transfers. Further, fig. 5(b) shows the allowed transitions for another peak, at a frequency  $\omega = 2\omega_b$ , which corresponds to transitions between dark states with two phonons decaying to dark states with no phonons, and also dark states with three phonons decaying to dark states with one phonon. We highlight that the emission peak corresponding to the two-phonon emission is much smaller than the one-phonon emission peak. However, the corresponding transitions can be differentiated due to the large energy difference in the spectrum. The associated side-bands on each major peak are due to transitions between dark and bright states accompanied by zero-, one- and two-phonon emission.

*Conclusions*— Dark excitons in QDs are more robust




FIG. 5. Phonon emission spectrum. Panels (a) and (b) show the elements  $|\langle\psi|\rho|\phi\rangle|^2$  of the Liouvillian eigenmatrices  $\rho$  that correspond to the peaks pointed in the phonon emission spectrum in panel (c). The eigenmatrices show that the peak (a) corresponds to a one phonon emission process, whereas the peak (b) corresponds to two-phonon emission processes. Eigenstates  $|\lambda_{10}\rangle$  and  $|\lambda_{11}\rangle$  of eq. (1) are superpositions of  $|0, v\rangle$  and  $|2, d_+\rangle$ , and eigenstates  $|\lambda_{15}\rangle$  and  $|\lambda_{16}\rangle$  are superpositions of  $|0, v\rangle$  and  $|3, d_+\rangle$ . The laser detuning is the same as in the top panel of fig. 2.

to decoherence than bright excitons, because they cannot couple to leaky optical modes. It is usually assumed that dark excitons are produced (and controlled) via external magnetic fields. However, in this work, we show that dark excitons in a QD coherently pumped by a laser and coupled with an acoustic cavity not only can be produced, but can also be targeted to realise giant Rabi oscillations between dark states with  $N$  phonons and the vacuum. These giant Rabi oscillations show a restricted cascading process that couples states with different phonon numbers, ultimately resulting in an effective coupling between the vacuum and a dark symmetric exciton state with  $N$ -phonons.

In the driven-dissipative scenario, other intermediate states are activated in the cascading process, culminating in  $N$ -phonon bundle emission, as shown by analysis of quantum trajectories. Furthermore, we show that  $N$ -phonon bundle correlation functions display quantum statistics corresponding to anti-bunching for the studied parameters, which is a desired property to realise  $N$ -phonon guns. Moreover, through the analysis of the emission spectrum, we characterise the acoustic transitions of the system, finding out that  $N$ -phonon bundle emission can be frequency resolved, which is an important feature to experimentally distinguish the phonon quasiparticles that escape the acoustic cavity.

Usually, high magnetic fields are needed to control the excitation of dark excitons, and producing these fields requires costly and large apparatuses that endanger the scalability possibilities of these systems. The system theoretically studied in this work provides an important alternative for quantum control in acoustic nanocavities without external magnetic fields to herald multi-phonon states through dark exciton manipulation.

- 
- [1] A. A. Balandin, Nanophononics: phonon engineering in nanostructures and nanodevices, *Journal of nanoscience and nanotechnology* **5**, 1015 (2005).
  - [2] R. Manenti, A. F. Kockum, A. Patterson, T. Behrle, J. Rahamim, G. Tancredi, F. Nori, and P. J. Leek, Circuit quantum acoustodynamics with surface acoustic waves, *Nature communications* **8**, 1 (2017).
  - [3] J. Zhang, M. Um, D. Lv, J.-N. Zhang, L.-M. Duan, and K. Kim, Noon states of nine quantized vibrations in two radial modes of a trapped ion, *Physical review letters* **121**, 160502 (2018).
  - [4] M. J. A. Schuetz, E. M. Kessler, G. Giedke, L. M. K. Vandersypen, M. D. Lukin, and J. I. Cirac, Universal quantum transducers based on surface acoustic waves, *Phys. Rev. X* **5**, 031031 (2015).
  - [5] A. Noguchi, R. Yamazaki, Y. Tabuchi, and Y. Nakamura, Qubit-assisted transduction for a detection of surface acoustic waves near the quantum limit, *Phys. Rev. Lett.* **119**, 180505 (2017).
  - [6] P. Arrangoiz-Arriola, E. A. Wollack, M. Pechal, J. D. Witmer, J. T. Hill, and A. H. Safavi-Naeini, Coupling a superconducting quantum circuit to a phononic crystal defect cavity, *Phys. Rev. X* **8**, 031007 (2018).
  - [7] A. Ask, M. Ekström, P. Delsing, and G. Johansson, Cavity-free vacuum-rabi splitting in circuit quantum acoustodynamics, *Phys. Rev. A* **99**, 013840 (2019).
  - [8] U. von Lüpke, Y. Yang, M. Bild, L. Michaud, M. Fadel, and Y. Chu, Parity measurement in the strong dispersive regime of circuit quantum acoustodynamics (2021), arXiv:2110.00263 [quant-ph].
  - [9] Q. Bin, X.-Y. Lü, F. P. Laussy, F. Nori, and Y. Wu, N-phonon bundle emission via the stokes process, *Physical review letters* **124**, 053601 (2020).
  - [10] K. Wan, S. Choi, I. H. Kim, N. Shutty, and P. Hayden, Fault-tolerant qubit from a constant number of components, *PRX Quantum* **2**, 040345 (2021).
  - [11] D. T. Bell and R. C. Li, Surface-acoustic-wave resonators, *Proceedings of the IEEE* **64**, 711 (1976).

- [12] E. M. Weig, R. H. Blick, T. Brandes, J. Kirschbaum, W. Wegscheider, M. Bichler, and J. P. Kotthaus, Single-electron-phonon interaction in a suspended quantum dot phonon cavity, *Phys. Rev. Lett.* **92**, 046804 (2004).
- [13] G. Rozas, M. F. P. Winter, B. Jusserand, A. Fainstein, B. Perrin, E. Semenova, and A. Lemaitre, Lifetime of the acoustic nanocavity modes, *Phys. Rev. Lett.* **102**, 015502 (2009).
- [14] O. O. Soykal, R. Ruskov, and C. Tahan, Sound-based analogue of cavity quantum electrodynamics in silicon, *Phys. Rev. Lett.* **107**, 235502 (2011).
- [15] A. Fainstein, N. D. Lanzillotti-Kimura, B. Jusserand, and B. Perrin, Strong optical-mechanical coupling in a vertical GaAs/AlAs microcavity for subterahertz phonons and near-infrared light, *Phys. Rev. Lett.* **110**, 037403 (2013).
- [16] P. Kharel, Y. Chu, M. Power, W. H. Renninger, R. J. Schoelkopf, and P. T. Rakich, Ultra-high-q phononic resonators on-chip at cryogenic temperatures, *Appl. Photonics* **3**, 066101 (2018).
- [17] Y. Xu, W. Fu, C.-I. Zou, Z. Shen, and H. X. Tang, High quality factor surface Fabry-Pérot cavity of acoustic waves, *Applied Physics Letters* **112**, 073505 (2018).
- [18] A. V. Akimov, C. L. Poysner, and A. J. Kent, Review of microwave electro-phononics in semiconductor nanostructures, *Semiconductor Science and Technology* **32**, 053003 (2017).
- [19] G. L. Bir, G. E. Pikus, *et al.*, *Symmetry and strain-induced effects in semiconductors*, Vol. 484 (Wiley New York, 1974).
- [20] C. A. Jiménez-Orjuela, H. Vinck-Posada, and J. M. Villas-Bôas, Dark excitons in a quantum-dot-cavity system under a tilted magnetic field, *Phys. Rev. B* **96**, 125303 (2017).
- [21] C. Adambukulam, V. Sewani, H. Stemp, S. Asaad, M. Madzik, A. Morello, and A. Laucht, An ultra-stable 1.5 T permanent magnet assembly for qubit experiments at cryogenic temperatures, *Review of Scientific Instruments* **92**, 085106 (2021).
- [22] S. Nomura, Y. Segawa, and T. Kobayashi, Confined excitons in a semiconductor quantum dot in a magnetic field, *Phys. Rev. B* **49**, 13571 (1994).
- [23] M. Bayer, G. Ortner, O. Stern, A. Kuther, A. A. Gorbunov, A. Forchel, P. Hawrylak, S. Fafard, K. Hinzer, T. L. Reinecke, S. N. Walck, J. P. Reithmaier, F. Kloppe, and F. Schäfer, Fine structure of neutral and charged excitons in self-assembled In(Ga)As/(Al)GaAs quantum dots, *Phys. Rev. B* **65**, 195315 (2002).
- [24] T. Belhadj, C.-M. Simon, T. Amand, P. Renucci, B. Chatel, O. Krebs, A. Lemaitre, P. Voisin, X. Marie, and B. Urbaszek, Controlling the polarization eigenstate of a quantum dot exciton with light, *Phys. Rev. Lett.* **103**, 086601 (2009).
- [25] K. Roszak, V. M. Axt, T. Kuhn, and P. Machnikowski, Exciton spin decay in quantum dots to bright and dark states, *Phys. Rev. B* **76**, 195324 (2007).
- [26] H.-P. Breuer, F. Petruccione, *et al.*, *The theory of open quantum systems* (Oxford University Press on Demand, 2002).
- [27] G. Lindblad, On the generators of quantum dynamical semigroups, *Communications in Mathematical Physics* **48**, 119 (1976).
- [28] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of  $n$ -level systems, *Journal of Mathematical Physics* **17**, 821 (1976).
- [29] J. Perea, D. Porrás, and C. Tejedor, Dynamics of the excitations of a quantum dot in a microcavity, *Physical Review B* **70**, 115304 (2004).
- [30] M. F. Pascual Winter, G. Rozas, A. Fainstein, B. Jusserand, B. Perrin, A. Huynh, P. O. Vaccaro, and S. Saravanan, Selective optical generation of coherent acoustic nanocavity modes, *Phys. Rev. Lett.* **98**, 265501 (2007).
- [31] S. Crooker, T. Barrick, J. Hollingsworth, and V. Klimov, Multiple temperature regimes of radiative decay in CdSe nanocrystal quantum dots: Intrinsic limits to the dark-exciton lifetime, *Applied Physics Letters* **82**, 2793 (2003).
- [32] T. Takagahara, Theory of exciton dephasing in semiconductor quantum dots, in *Semiconductor Quantum Dots* (Springer, 2002) pp. 353–388.
- [33] J. Dalibard, Y. Castin, and K. Mølmer, Wave-function approach to dissipative processes in quantum optics, *Phys. Rev. Lett.* **68**, 580 (1992).
- [34] K. Mølmer, Y. Castin, and J. Dalibard, Monte Carlo wave-function method in quantum optics, *JOSA B* **10**, 524 (1993).
- [35] C. S. Muñoz, E. Del Valle, A. G. Tudela, K. Müller, S. Lichtmannecker, M. Kaniber, C. Tejedor, J. Finley, and F. Laussy, Emitters of  $n$ -photon bundles, *Nature Photonics* **8**, 550 (2014).
- [36] Parameters can be tuned to get a larger amount of two-phonon emission processes. In this work, we tuned parameters with the tree-structured Parzen estimator implemented in Optuna [49] to minimise the generalised second-order correlation function at a value for the laser detuning that coupled the vacuum and the  $|2, d_+\rangle$  state through giant Rabi oscillations.
- [37] F. Dell’Anno, S. De Siena, and F. Illuminati, Multi-photon quantum optics and quantum state engineering, *Physics reports* **428**, 53 (2006).
- [38] X. T. Zou and L. Mandel, Photon-antibunching and sub-Poissonian photon statistics, *Phys. Rev. A* **41**, 475 (1990).
- [39] T. Petrosky and I. Prigogine, The Liouville space extension of quantum mechanics, in *Advances in Chemical Physics* (John Wiley & Sons, Ltd, 1996) pp. 1–120.
- [40] T. Petrosky, Complex Spectral Representation of the Liouvillian and Kinetic Theory in Nonequilibrium Physics, *Progress of Theoretical Physics* **123**, 395 (2010), <https://academic.oup.com/ptp/article-pdf/123/3/395/5308209/123-3-395.pdf>.
- [41] D. Manzano and P. Hurtado, Harnessing symmetry to control quantum transport, *Advances in Physics* **67**, 1 (2018), <https://doi.org/10.1080/00018732.2018.1519981>.
- [42] V. V. Albert and L. Jiang, Symmetries and conserved quantities in Lindblad master equations, *Phys. Rev. A* **89**, 022118 (2014).
- [43] N. Hatano and T. Petrosky, Eigenvalue problem of the Liouvillian of open quantum systems, *AIP Conference Proceedings* **1648**, 200005 (2015), <https://aip.scitation.org/doi/pdf/10.1063/1.4912483>.
- [44] V. Vargas-Calderón, Phonon-assisted tunnelling in double quantum dot molecules immersed in a microcavity, *Research Gate* 10.13140/RG.2.2.31486.23360/1 (2018).
- [45] B. A. Tay and T. Petrosky, Biorthonormal eigenbasis of a Markovian master equation for the quantum Brownian motion, *Journal of Mathematical Physics* **49**, 113301 (2008).
- [46] J. M. Torres, Closed-form solution of Lindblad master

- equations without gain, *Phys. Rev. A* **89**, 052133 (2014).
- [47] S. Echeverri-Arteaga, H. Vinck-Posada, and E. A. Gómez, Explanation of the quantum phenomenon of off-resonant cavity-mode emission, *Phys. Rev. A* **97**, 043815 (2018).
- [48] V. Vargas-Calderón and H. Vinck-Posada, Light emission properties in a double quantum dot molecule immersed in a cavity: Phonon-assisted tunneling, *Physics Letters A* **384**, 126076 (2020).
- [49] T. Akiba, S. Sano, T. Yanase, T. Ohta, and M. Koyama, Optuna: A next-generation hyperparameter optimization framework, in *Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (2019).