

CONSISTENCY RELATIONS OF RANK 2 CLUSTER SCATTERING DIAGRAMS OF AFFINE TYPE AND PENTAGON RELATION

KODAI MATSUSHITA

ABSTRACT. In this paper, we prove the consistency relations of rank 2 cluster scattering diagrams of affine type by using the pentagon relation.

1. INTRODUCTION

The cluster algebras are commutative algebras introduced by Fomin and Zelevinsky [FZ02]. The positivity of coefficients of F -polynomials and the sign-coherence of c -vectors were important conjectures. These were proved in [GHKK18] by using scattering diagram methods. Scattering diagrams for cluster algebras are characterized by the consistency relations in their structure groups G .

The structure group G of a given cluster scattering diagram is generated by the dilogarithm elements [GHKK18, Nak21]

$$(1.1) \quad \Psi[n] := \exp \left(\sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j^2} X_{jn} \right).$$

The precise definition of G is given in §2. These elements satisfy pentagon relations:

$$(1.2) \quad \Psi[n]^c \Psi[n']^c = \Psi[n]^c \Psi[n + n']^c \Psi[n]^c$$

where $\{n, n'\} = c^{-1}$.

In this paper, we prove the consistency relations of rank 2 cluster scattering diagrams of affine type, namely types $A_1^{(1)}$ and $A_2^{(2)}$. More precisely, we prove the following theorem. For simplicity, let $\begin{bmatrix} n_1 \\ n_2 \end{bmatrix} = \Psi[(n_1, n_2)]$.

Theorem 1. *The following relations holds:*

$$(1.3) \quad \begin{bmatrix} 0 \\ 1 \end{bmatrix}^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}^2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}^2 \begin{bmatrix} 2 \\ 1 \end{bmatrix}^2 \begin{bmatrix} 3 \\ 2 \end{bmatrix}^2 \cdots \prod_{j=0}^{\infty} \begin{bmatrix} 2^j \\ 2^j \end{bmatrix}^{2^{2-j}} \cdots \begin{bmatrix} 2 \\ 3 \end{bmatrix}^2 \begin{bmatrix} 1 \\ 2 \end{bmatrix}^2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}^2,$$

$$(1.4) \quad \begin{bmatrix} 0 \\ 1 \end{bmatrix}^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}^4 \begin{bmatrix} 3 \\ 4 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix}^4 \begin{bmatrix} 5 \\ 6 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix}^4 \cdots$$

$$\times \begin{bmatrix} 1 \\ 2 \end{bmatrix}^6 \prod_{j=1}^{\infty} \begin{bmatrix} 2^j \\ 2^{j+1} \end{bmatrix}^{2^{2-j}} \cdots \begin{bmatrix} 5 \\ 12 \end{bmatrix} \begin{bmatrix} 2 \\ 5 \end{bmatrix}^4 \begin{bmatrix} 3 \\ 8 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \end{bmatrix}^4 \begin{bmatrix} 1 \\ 4 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix}^4.$$

Moreover, these formulas can be reduced to trivial relations by iteration of the pentagon relations.

The relations (1.3) and (1.4) are the (unique) consistency relations of type $A_1^{(1)}$ and type $A_2^{(2)}$, respectively.

We say a product of dilogarithm elements is ordered, (resp. anti-ordered) if, for any adjacent pair $\begin{bmatrix} n_1 \\ n_2 \end{bmatrix} \begin{bmatrix} n'_1 \\ n'_2 \end{bmatrix}$, the inequality $n_1/n_2 \leq n'_1/n'_2$ (resp. $n_1/n_2 \leq n'_1/n'_2$) holds. The consistency relations of scattering diagrams in \mathbb{R}^2 have the form of

$$(1.5) \quad \text{“anti-ordered product”} = \text{“ordered product”}.$$

It was shown that the consistency relations are generated by the pentagon relation [Nak21], and the above theorem provides an simplest examples involving the *infinite* product.

We remark that the relations (1.3) first proved by [Rei10] by using quiver representations. Also, the relations (1.3) and (1.4) were proved by cluster mutation technique by [Rea20].

In §2, we introduce dilogarithm elements and the pentagon relations. In §3, we prove a generalization of the formula (1.3). In §4, we prove a generalization of the formula (1.4) by using a results of §3.

2. DILOGARITHM ELEMENTS AND PENTAGON RELATION

Let N be a rank 2 lattice with a skew-symmetric bilinear form

$$\{\cdot, \cdot\}: N \times N \longrightarrow \mathbb{Q}.$$

Let e_1, e_2 be a basis of N , and we define

$$N^+ := \{a_1 e_1 + a_2 e_2 \mid a_1, a_2 \in \mathbb{Z}_{\geq 0}, a_1 + a_2 > 0\}.$$

Let \mathbb{k} be a field of characteristic 0, and we define an N^+ -graded Lie algebra \mathfrak{g} over \mathbb{k} with generators X_n such that

$$\mathfrak{g} = \bigoplus_{n \in N^+} \mathfrak{g}_n, \quad \mathfrak{g}_n = \mathbb{k} X_n, \quad [X_n, X_{n'}] = \{n, n'\} X_{n+n'}.$$

Let $\mathcal{L} := \{L \subset N^+ \mid N^+ + L \subset L, \#(N^+ \setminus L) < \infty\}$. For $L \in \mathcal{L}$, we define a Lie algebra ideal $\mathfrak{g}^L := \bigoplus_{n \in L} \mathfrak{g}_n$ and the quotient of \mathfrak{g} by \mathfrak{g}^L

$$\mathfrak{g}_L := \mathfrak{g}/\mathfrak{g}^L = \bigoplus_{n \in N^+ \setminus L} \mathfrak{g}_n \quad (\text{as a vector space}).$$

Let G_L be a group with a set bijection

$$\exp_L: \mathfrak{g}_L \longrightarrow G_L$$

and the product is defined by a Baker-Campbell-Hausdorff (BCH) formula:
(2.1)

$$\exp_L(X) \exp_L(Y) = \exp_L\left(X + Y + \frac{1}{2}[X, Y] + \frac{1}{12}[X, [X, Y]] - \frac{1}{12}[Y, [X, Y]] + \cdots\right).$$

This product formula is well-defined because \mathfrak{g}_L is nilpotent.

For $L, L' \in \mathcal{L}$ such that $L \subset L'$, there exists the canonical Lie algebra homomorphism $\mathfrak{g}_L \rightarrow \mathfrak{g}'_L$, which induces the group homomorphism $G_L \rightarrow G'_L$. Thus, by the inverse limit we obtain a Lie algebra $\hat{\mathfrak{g}}$ and a group G :

$$\hat{\mathfrak{g}} := \varprojlim_{L \in \mathcal{L}} \mathfrak{g}_L, \quad G := \varprojlim_{L \in \mathcal{L}} G_L.$$

There is a set bijection

$$\exp: \hat{\mathfrak{g}} \rightarrow G, \quad (X_L)_{L \in \mathcal{L}} \mapsto (\exp_L(X_L))_{L \in \mathcal{L}}.$$

We use an infinite sum to express an element of $\hat{\mathfrak{g}}$.

We define important elements in G :

Definition 1 (Dilogarithm element). For any $n \in N^+$, define

$$[n] := \exp\left(\sum_{j>0} \frac{(-1)^{j+1}}{j^2} X_{jn}\right) \in G.$$

We call $[n]$ a *dilogarithm element* for n .

For $c \in \mathbb{Q}$ and $g = \exp(X) \in G$, we define $g^c := \exp(cX)$.

Proposition 1 (Pentagon relation [GHKK18], [Nak21]). *Let $n, n' \in N^+$. Then, the following relations hold in G :*

- (1) *If $\{n', n\} = 0$, then $[n'] [n] = [n] [n']$,*
- (2) *If $\{n', n\} = c^{-1}$ ($c \in \mathbb{Q} \setminus 0$), then*

$$[n']^c [n]^c = [n]^c [n + n']^c [n']^c \quad (\text{pentagon relation}).$$

3. PROOF OF FORMULA (1.3)

For a subset $I = \{i_1 < i_2 < i_3 < \cdots\}$ of \mathbb{Z} and a sequence $(a_i)_{i \in I}$ of elements of G , we write

$$\overrightarrow{\prod}_{i \in I} a_i := a_{i_1} a_{i_2} a_{i_3} \cdots, \quad \overleftarrow{\prod}_{i \in I} a_i := \cdots a_{i_3} a_{i_2} a_{i_1}.$$

For example, $\overrightarrow{\prod}_{i \geq 0} a_i = a_0 a_1 a_2 \cdots$ and $\overleftarrow{\prod}_{i \geq 0} a_i = \cdots a_2 a_1 a_0$.

The following is the main theorem of this section:

Theorem 2. *If $\{n', n\} = c^{-1}$ ($c \in \mathbb{Q} \setminus \{0\}$), then*

$$(3.1) \quad [n']^{2c} [n]^{2c} = \overrightarrow{\prod}_{p \geq 0} [n + p(n + n')]^{2c} \overrightarrow{\prod}_{p \geq 0} [2^p(n + n')]^{4c/2^p} \overleftarrow{\prod}_{p \geq 0} [n' + p(n + n')]^{2c}.$$

The case of $c = 1$, $n = [(1, 0)]$, $n' = [(0, 1)]$ is nothing but the formula (1.3).

To prove this theorem, we introduce some notations and lemmas.

Let $L \in \mathcal{L}$. For two elements g_1, g_2 of G , let us denote $g_1 \equiv g_2 \pmod{L}$ if their images in G_L are identical. For example, if $n \in N^+$ is in L , then $[n] \equiv \exp(0) = 1_G \pmod{L}$. By the definition of G , two elements g_1, g_2 of G are identical if and only if $g_1 \equiv g_2 \pmod{L}$ for all $L \in \mathcal{L}$.

Lemma 1. *If $\{n', n\} = c^{-1}$ ($c \in \mathbb{Q} \setminus \{0\}$), we obtain*

$$(3.2) \quad [n']^c [n]^{2c} = [n]^{2c} [2n + n']^c [n + n']^{2c} [n']^c,$$

$$(3.3) \quad [n']^{2c} [n]^c = [n]^c [n + n']^{2c} [n + 2n']^c [n']^{2c}.$$

Proof. The equality (3.2) can be proved by repeatedly applying the pentagon relation:

$$\begin{aligned} [n']^{2c} [n]^c &= [n']^c [n']^c [n]^c = [n']^c [n]^c [n + n']^c [n']^c \\ &= [n]^c [n + n']^c [n']^c [n + n']^c [n']^c = [n]^c [n + n']^{2c} [n + 2n']^c [n']^{2c}. \end{aligned}$$

Note that $\{n', n + n'\} = c^{-1}$, thus we can use the pentagon identity in the last equality.

If $\{n', n\} = c^{-1}$, then $\{n, n'\} = (-c)^{-1}$. By the equality (3.2),

$$(3.4) \quad [n]^{-c} [n']^{2(-c)} = [n']^{2(-c)} [n + 2n']^{-c} [n + n']^{2(-c)} [n]^{-c}.$$

Taking the inverse of both sides of (3.4), we obtain the equality (3.3). \square

The following is a key lemma:

Lemma 2. *Let l be a non-negative integer, and let $n, n' \in N^+$. If $\{n', n\} = c^{-1}$,*

$$(3.5) \quad [n']^{2c} \left(\overrightarrow{\prod}_{0 \leq p \leq l} [n + 2pn']^c \right) = [n]^c \left(\overrightarrow{\prod}_{1 \leq p \leq 2l+1} [n + pn']^{2c} \right) [n + (2l+2)n']^c [n']^{2c}$$

Proof. We will prove it by induction on l .

If $l = 0$, the equality (3.5) is nothing but (3.3).

Let $l > 0$. Suppose that the claim is true in the case of $l - 1$, then by the induction hypothesis,

$$\begin{aligned} &[n']^{2c} \left(\overrightarrow{\prod}_{0 \leq p \leq l} [n + 2pn']^c \right) \\ &= [n']^{2c} \left(\overrightarrow{\prod}_{0 \leq p \leq l-1} [n + 2pn']^c \right) [n + 2ln']^c \\ &= [n]^c \left(\overrightarrow{\prod}_{1 \leq p \leq 2l-1} [n + pn']^{2c} \right) [n + 2ln']^c [n']^{2c} [n + 2ln']^c \end{aligned}$$

$$= [n]^c \left(\overrightarrow{\prod}_{1 \leq p \leq 2l+1} [n + pn']^{2c} \right) [n + (2l+2)n']^c [n']^{2c}.$$

In the last equality, we use

$$[n']^{2c} [n + 2ln']^c = [n + 2ln']^c [n + (2l+1)n']^{2c} [n + (2l+2)n']^c [n']^{2c},$$

which is a specialization of (3.3). \square

Now we consider the limit of Lemma 2:

Lemma 3. *If $\{n', n\} = c^{-1}$, then*

$$(3.6) \quad [n']^{2c} \left(\overrightarrow{\prod}_{p \geq 0} [n + 2pn']^c \right) = [n]^c \left(\overrightarrow{\prod}_{p \geq 1} [n + pn']^{2c} \right) [n']^{2c},$$

$$(3.7) \quad \left(\overleftarrow{\prod}_{p \geq 0} [n' + 2pn]^c \right) [n]^{2c} = [n]^{2c} \left(\overleftarrow{\prod}_{p \geq 1} [n' + pn]^{2c} \right) [n']^c.$$

Proof. Let $L \in \mathcal{L}$. Then, by the cofiniteness of L , there exists some positive integer l such that $n + 2ln' \in L$. Then, by Lemma 2, we obtain

$$\begin{aligned} [n']^{2c} \left(\overrightarrow{\prod}_{p \geq 0} [n + 2pn']^c \right) &\equiv [n']^{2c} \left(\overrightarrow{\prod}_{0 \leq p \leq l} [n + 2pn']^c \right) \pmod{L} \\ &= [n]^c \left(\overrightarrow{\prod}_{1 \leq p \leq 2l+1} [n + pn']^{2c} \right) [n + (2l+2)n']^c [n']^{2c} \\ &\equiv [n]^c \left(\overrightarrow{\prod}_{p \geq 1} [n + pn']^{2c} \right) [n']^{2c} \pmod{L}. \end{aligned}$$

Thus, the equality (3.6) holds.

Since $\{n, n'\} = (-c)^{-1}$, by (3.6), we obtain

$$(3.8) \quad [n]^{-2c} \left(\overrightarrow{\prod}_{p \geq 0} [n' + 2pn]^{-c} \right) = [n']^{-c} \left(\overrightarrow{\prod}_{p \geq 1} [n' + pn]^{-c} \right) [n]^{-2c}.$$

The equality (3.7) is obtained by taking the inverse of the both sides of (3.8). \square

Proof of Theorem 2. For $L \in \mathcal{L}$ and $k \in \mathbb{Z}_{>0}$, let $P_L(k)$ be the following assertion: for $n, n' \in N^+$ and $c \in \mathbb{Q} \setminus \{0\}$, if $\{n', n\} = c^{-1}$ and $k(n+n') \in L$, then

$$[n']^{2c} [n]^{2c}$$

$$(3.9) \quad \begin{aligned} & \equiv \prod_{p \geq 0}^{\rightarrow} [n + p(n + n')]^{2c} \prod_{p \geq 0} [2^p(n + n')]^{4c/2^p} \prod_{p \geq 0}^{\leftarrow} [n' + p(n + n')]^{2c} \pmod{L}. \end{aligned}$$

For any $L \in \mathcal{L}$, there exists some positive integer k such that $k(n + n') \in L$. Thus, if $P_L(k)$ is true for any $k \in \mathbb{Z}_{>0}$ and $L \in \mathcal{L}$, then a relation (3.9) holds for any $L \in \mathcal{L}$, and Theorem 2 is proved. Fix $L \in \mathcal{L}$, and we prove $P_L(k)$ by induction on k .

If $k = 1$, the right hand side of (3.9) is equivalent to $[n]^{2c}[n']^{2c}$ because $ln + l'n' \in L$ for any $l, l' \in \mathbb{Z}_{\geq 1}$. Since

$$[n']^c[n]^c = [n]^c[n + n']^c[n']^c \equiv [n]^c[n']^c \pmod{L},$$

we obtain $[n']^{2c}[n]^{2c} \equiv [n']^{2c}[n']^{2c} \pmod{L}$.

Let $k \geq 2$, and we suppose a proposition $P_L(k-1)$ is true. By the equality (3.3),

$$\begin{aligned} [n']^{2c}[n]^{2c} &= ([n']^{2c}[n]^c)[n]^c \\ &= [n]^c[n + n']^{2c}[n + 2n']^c([n']^{2c}[n]^c) \\ &= [n]^c[n + n']^{2c}([n + 2n']^c[n]^c)[n + n']^{2c}[n + 2n']^c[n']^{2c}. \end{aligned}$$

Since $(k-1)(n + (n + 2n')) \in L$ and $\{n + 2n', n\} = (c/2)^{-1}$, by the induction hypothesis,

$$\begin{aligned} & [n + 2n']^c[n]^c \\ &= [n + 2n']^{(c/2) \cdot 2} [n]^{(c/2) \cdot 2} \\ & \stackrel{(3.9)}{\equiv} \prod_{p \geq 0}^{\rightarrow} [n + p(2n + 2n')]^{2 \cdot (c/2)} \prod_{p \geq 0} [2^p(2n + 2n')]^{4 \cdot (c/2)/2^p} \\ & \quad \times \prod_{p \geq 0}^{\leftarrow} [(n + 2n') + p(2n + 2n')]^{2 \cdot (c/2)} \pmod{L} \\ &= \prod_{p \geq 0}^{\rightarrow} [n + 2p(n + n')]^c \prod_{p \geq 1} [2^p(n + n')]^{4c/2^p} \prod_{p \geq 0}^{\leftarrow} [(n + 2n') + 2p(n + n')]^c. \end{aligned}$$

Since $\{n + n', n\} = c^{-1}$ and $\{n + 2n', n + n'\} = c^{-1}$, by Lemma 3,

$$\begin{aligned} [n']^{2c}[n]^{2c} &\equiv [n]^c[n + n']^{2c} \prod_{p \geq 0}^{\rightarrow} [n + 2p(n + n')]^c \prod_{p \geq 1} [2^p(n + n')]^{4c/2^p} \\ & \quad \times \prod_{p \geq 0}^{\leftarrow} [(n + 2n') + 2p(n + n')]^c \\ & \quad \times [n + n']^{2c}[n + 2n']^c[n']^{2c} \pmod{L} \\ &= [n]^c[n]^c \left(\prod_{p \geq 1}^{\rightarrow} [n + p(n + n')]^c \right) [n + n']^{2c} \prod_{p \geq 1} [2^p(n + n')]^{4c/2^p} \end{aligned}$$

$$\begin{aligned}
 & \times [n + n']^{2c} \prod_{p \geq 1}^{\leftarrow} [(n + 2n') + p(n + n')]^c \\
 & \times [n + 2n']^c [n + 2n']^c [n']^{2c} \\
 & = \prod_{p \geq 0}^{\rightarrow} [n + p(n + n')]^{2c} \prod_{p \geq 0} [2^p(n + n')]^{4c/2^p} \prod_{p \geq 0}^{\leftarrow} [n' + p(n + n')]^{2c}
 \end{aligned}$$

This completes the proof of Theorem 2. \square

4. PROOF OF FORMULA (1.4)

The formula (1.4) is the case of $c = -1$, $n = [(0, 1)]$, $n' = [(1, 0)]$ of the following theorem:

Theorem 3. *If $\{n', n\} = c^{-1}$ ($c \in \mathbb{Q}$), then we obtain*

$$\begin{aligned}
 [n']^c [n]^{4c} &= \prod_{p \geq 0}^{\rightarrow} ([(2p + 1)n + pn']^{4c} [(4p + 4)n + (2p + 1)n']^c) \\
 &\times [2n + n']^{2c} \prod_{p \geq 0} [2^p(2n + n')]^{4c/2^p} \\
 &\times \prod_{p \geq 0}^{\leftarrow} ([(2p + 1)n + (p + 1)n']^{4c} [4pn + (2p + 1)n']^c).
 \end{aligned}$$

To prove this theorem, we consider some lemmas.

Lemma 4. *If $\{n', n\} = c^{-1}$ ($c \in \mathbb{Q} \setminus \{0\}$), then we obtain*

$$\begin{aligned}
 & [n']^c \left(\prod_{0 \leq p \leq l}^{\rightarrow} [n + pn']^{2c} \right) \\
 &= [n]^{2c} \left(\prod_{1 \leq p \leq l}^{\rightarrow} ([2n + (2p - 1)n']^c [n + pn']^{4c}) \right) \\
 &\times [2n + (2l + 1)n']^c [n + (l + 1)n']^{2c} [n']^c
 \end{aligned}$$

Proof. We prove it by induction on l . The case of $l = 0$ is nothing less than the equality (3.2).

Let $l > 0$. Suppose that the claim is true in the case of $l - 1$, then

$$\begin{aligned}
 & [n']^c \left(\prod_{0 \leq p \leq l}^{\rightarrow} [n + pn']^{2c} \right) \\
 &= [n']^c \left(\prod_{0 \leq p \leq l-1}^{\rightarrow} [n + pn']^{2c} \right) [n + ln']^{2c}
 \end{aligned}$$

$$\begin{aligned}
&= [n]^{2c} \left(\prod_{1 \leq p \leq l-1}^{\rightarrow} ([2n + (2p-1)n']^c [n + pn']^{4c}) \right) \\
&\quad \times [2n + (2l-1)n']^c [n + ln']^{2c} [n']^c [n + ln']^{2c} \\
&= [n]^{2c} \left(\prod_{1 \leq p \leq l}^{\rightarrow} ([2n + (2p-1)n']^c [n + pn']^{4c}) \right) \\
&\quad \times [2n + (2l+1)n']^c [n + (l+1)n']^{2c} [n']^c
\end{aligned}$$

In the last equality, we use

$$[n']^c [n + ln']^{2c} = [n + ln']^{2c} [2n + (2l+1)n']^c [n + (l+1)n']^{2c} [n']^2,$$

which is a specialization of (3.2). \square

Now we consider the limit of Lemma 4:

Lemma 5. *If $\{n', n\} = c^{-1}$, then we obtain*

(4.1)

$$[n']^c \left(\prod_{p \geq 0}^{\rightarrow} [n + pn']^{2c} \right) = [n]^{2c} \left(\prod_{p \geq 1}^{\rightarrow} [2n + (2p-1)n']^c [n + pn']^{4c} \right) [n']^c,$$

(4.2)

$$\left(\prod_{p \geq 0}^{\leftarrow} [n' + pn]^{2c} \right) [n]^c = [n]^c \left(\prod_{p \geq 1}^{\leftarrow} [n' + pn]^{4c} [2n' + (2p-1)n]^c \right) [n']^{2c}.$$

Proof. Let $L \in \mathcal{L}$. Then, there exist some positive integer l such that $n + ln' \in L$. Then, by Lemma 4, we obtain

$$\begin{aligned}
&[n']^c \prod_{p \geq 0}^{\rightarrow} [n + pn']^{2c} \\
&\equiv [n']^c \left(\prod_{0 \leq p \leq l}^{\rightarrow} [n + pn']^{2c} \right) \pmod{L} \\
&= [n]^{2c} \left(\prod_{1 \leq p \leq l}^{\rightarrow} ([2n + (2p-1)n']^c [n + pn']^{4c}) \right) \\
&\quad \times [2n + (2l+1)n']^c [n + (l+1)n']^{2c} [n']^c \\
&\equiv [n]^{2c} \left(\prod_{p \geq 1}^{\rightarrow} [2n + (2p-1)n']^c [n + pn']^{4c} \right) [n']^c \pmod{L}
\end{aligned}$$

Thus, the equality (4.1) holds.

Since $\{n, n'\} = (-c)^{-1}$, by (4.1), we obtain

$$[n]^{-c} \left(\overrightarrow{\prod}_{p \geq 0} [n' + pn]^{-2c} \right) = [n']^{-2c} \left(\overrightarrow{\prod}_{p \geq 1} [2n' + (2p-1)n]^{-c} [n' + pn]^{-4c} \right) [n]^{-c}.$$

By taking the inverse of both sides of this equality, we have the equality (4.2). \square

Proof of Theorem 3. Using the pentagon relations, Theorem 2 and Lemma 5, we can calculate as follows:

$$\begin{aligned} & [n']^c [n]^{4c} \\ &= [n']^c [n]^{2c} [n]^{2c} \\ &= [n]^{2c} [2n + n']^c [n + n']^{2c} [n']^c [n]^{2c} \\ &= [n]^{2c} [2n + n']^c [n + n']^{2c} [n]^{2c} [2n + n']^c [n + n']^{2c} [n]^c \\ &\stackrel{(3.1)}{=} [n]^{2c} [2n + n']^c \\ &\quad \times \overrightarrow{\prod}_{p \geq 0} [n + p(2n + n')]^{2c} \\ &\quad \times \prod_{p \geq 0} [2^p (2n + n')]^{4c/2^p} \\ &\quad \times \overleftarrow{\prod}_{p \geq 0} [n + n' + p(2n + n')]^{2c} \\ &\quad \times [2n + n']^c [n + n']^{2c} [n']^c \\ &\stackrel{(4.1), (4.2)}{=} [n]^{2c} \times [n]^{2c} \left(\overrightarrow{\prod}_{p \geq 1} ([2n + (2p-1)(2n + n')]^c [n + p(2n + n')]^{4c}) \right) [2n + n']^c \\ &\quad \times \prod_{p \geq 0} [2^p (2n + n')]^{4c/2^p} \times [2n + n']^c \\ &\quad \times \left(\overleftarrow{\prod}_{p \geq 1} ([n + n'] + p(2n + n')]^{4c} [2(n + n') + (2p-1)(2n + n')]^c \right) \\ &\quad \times [n + n']^{2c} [n + n']^{2c} [n']^c \\ &= \overrightarrow{\prod}_{p \geq 0} ([n + p(2n + n')]^{4c} [2n + (2p+1)(2n + n')]^c) \\ &\quad \times [2n + n']^{2c} \prod_{p \geq 0} [2^p (2n + n')]^{4c/2^p} \\ &\quad \times \overleftarrow{\prod}_{p \geq 0} ([n + n'] + p(2n + n')]^{4c} [2(n + n') + (2p-1)(2n + n')]^c \end{aligned}$$

$$\begin{aligned}
&= \prod_{p \geq 0}^{\rightarrow} ([(2p+1)n + pn']^{4c} [(4p+4)n + (2p+1)n']^c) \\
&\quad \times [2n + n']^{2c} \prod_{p \geq 0} [2^p (2n + n')]^{4c/2^p} \\
&\quad \times \prod_{p \geq 0}^{\leftarrow} ([(2p+1)n + (p+1)n']^{4c} [4pn + (2p+1)n']^c)
\end{aligned}$$

This completes the proof. \square

REFERENCES

- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [GHKK18] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich. Canonical bases for cluster algebras. *Journal of the American Mathematical Society*, 31(2):497–608, 2018.
- [Nak21] Tomoki Nakanishi. Cluster algebras and scattering diagrams, part III. cluster scattering diagrams, 2021.
- [Rea20] Nathan Reading. A combinatorial approach to scattering diagrams. *Algebr. Comb.*, 3(3):603–636, 2020.
- [Rei10] Markus Reineke. Poisson automorphisms and quiver moduli. *Journal of the Institute of Mathematics of Jussieu*, 9(3):653–667, 2010.

(KODAI MATSUSHITA) GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY,
CHIKUSA-KU, NAGOYA, 464-8602 JAPAN
Email address: m19043g@math.nagoya-u.ac.jp