

Handling missing data when estimating causal effects with Targeted Maximum Likelihood Estimation

S. Ghazaleh Dashti^{1,2}, Katherine J. Lee^{1,2}, Julie A. Simpson³, Ian R. White⁴, John B. Carlin^{1,2}, Margarita Moreno-Betancur^{2,1}

1 Clinical Epidemiology and Biostatistics Unit, Murdoch Children's Research Institute, Melbourne, Victoria, Australia

2 Clinical Epidemiology and Biostatistics Unit, Department of Paediatrics, University of Melbourne, Melbourne, Victoria, Australia

3 Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, University of Melbourne, Melbourne, Victoria, Australia

4 MRC Clinical Trials Unit at UCL, University College London, London, UK

Corresponding author

S Ghazaleh Dashti, DDS, MPH, PhD

Clinical Epidemiology and Biostatistics Unit

Murdoch Children's Research Institute

Royal Children's Hospital, 50 Flemington Rd, Parkville VIC 3052

Email: ghazaleh.dashti@mcri.edu.au

Key words: missing data, causal inference, targeted maximum likelihood estimation, multiple imputation

Acknowledgements

This article used unit record data from the Victorian Adolescent Health Cohort Study. We thank the families that participated in the VAHCS, the study research team and the Principal Investigator, Professor George Patton. We also thank Anthony Charsley for his contributions to the coding in the preliminary stages of this work.

Funding

Ian White was supported by the Medical Research Council Programme MC_UU_00004/07. Katherine Lee was funded by a National Health and Medical Research Council (NHMRC) Career Development Fellowship (ID 1127984), Julie Simpson by a NHMRC Investigator Grant – Leadership (ID 1196068), and Margarita Moreno-Betancur by an NHMRC Investigator Grant (ID 2009572). The work was supported from the National Health and Medical Research Council Project Grant (ID 1166023). The Murdoch Children's Research Institute is supported by the Victorian Government's Operational Infrastructure Support Program.

Author Contributions

All authors participated in planning the simulation and case studies, interpretation of the results, and reviewing and revising the manuscript. SGD and MMB conceptualized and designed the study. SGD performed the simulation and case study analyses and led the writing of the manuscript.

Data availability statement

Data from the Victorian Adolescent Health Cohort Study (VAHCS) are not publicly available. Those interested in replicating these findings are welcome to contact the corresponding author, or the VAHCS team (<https://www.mcri.edu.au/research/projects/2000-stories/information-researchers>). Simulation study codes can be made available upon request to the corresponding author.

Conflicts of Interest

The authors declare no conflicts of interest.

Abstract

Targeted Maximum Likelihood Estimation (TMLE) is increasingly used for doubly robust causal inference, but how missing data should be handled when using TMLE with data-adaptive approaches is unclear. Based on Victorian Adolescent Health Cohort Study data, we conducted a simulation study to evaluate missing data methods when using TMLE. Methods compared were complete-case analysis, extended TMLE incorporating outcome-missingness model, a “missing covariate missing indicator” method, and five multiple imputation (MI) approaches using parametric or machine-learning imputation models. We considered simple (exposure/outcome generated from main-effects regressions) and complex (models also included interactions) scenarios, alongside two missingness mechanisms (different in whether the outcome influenced missingness in other variables). Parametric MI including interaction terms performed best across all settings in terms of bias and variance reduction. Non-MI methods had the largest variance and, like MI using machine-learning, had larger biases when the outcome influenced missingness in other variables. Parametric MI without interactions – the default in most MI software – had the largest bias in the complex scenario. When choosing a method to handle missing data in the context of TMLE with data-adaptive approaches, researchers must consider the missingness mechanism. Parametric MI that incorporates interactions and non-linearities appears to be the best approach.

Key words: causal inference, missing data, targeted maximum likelihood estimation, multiple imputation

Introduction

A key component of epidemiological research is causal inference from longitudinal cohort studies, where the objective is often to estimate the average causal effect (ACE) of an exposure on an outcome.(1-5) For a binary exposure ($X=1$ exposed; $X=0$ unexposed), the ACE is defined as the difference in the average potential outcome if everybody in the population were set to be exposed versus unexposed.(1-5) Under the assumptions of conditional exchangeability given a vector of measured confounders Z , consistency, and positivity, the ACE is identifiable from observable data by the g-formula $E[E(Y|X = 1, Z) - E(Y|X = 0, Z)]$, where Y is the outcome.(6)

Several singly robust approaches, including g-computation and propensity score methods, and doubly robust estimators, including Targeted Maximum Likelihood Estimation (TMLE), are available for ACE estimation in the absence of missing data. Here, we focus on TMLE, which combines models for the outcome and propensity score.(7-9) A detailed description of TMLE is available elsewhere.(7-9) Briefly, the first step is the same as in g-computation, where a model for the expected outcome conditional on exposure and confounders ($\hat{E}[Y|X, Z]$) is estimated and used to predict outcome for all records under exposure and no exposure. In g-computation these predictions are directly plugged into the g-formula to estimate the ACE. In TMLE, they are updated using information from the propensity score ($\hat{P}[X = 1|Z]$) (the targeting step), before being plugged into the g-formula.(7) The targeting step in TMLE ensures that the estimator is doubly robust, where only one of the two models (outcome or propensity model) needs to be consistently estimated to ensure consistent estimation of the ACE. Unlike with singly robust methods, data-adaptive approaches (e.g. machine learning methods) can be used for the exposure and outcome models in TMLE as long as the Donsker class condition (requiring that outcome and propensity score estimators do not heavily overfit the data) holds.(8-10) The possibility to use data-adaptive approaches provides a further avenue for reducing the risk of bias, with the Donsker condition enabling valid statistical inference. Given these desirable properties, interest in the application of TMLE for ACE estimation is growing in epidemiological studies (Figure 1).(11-80)

Missing data are ubiquitous in epidemiological studies and can lead to biased estimates and loss of precision if handled inappropriately.(81) Commonly used approaches for handling missing data in studies using TMLE for ACE estimation include multiple imputation (MI) (19, 23, 42, 44, 47, 53, 64, 78, 80), complete-case analysis (CCA) (13, 14, 17, 31, 41, 44, 62, 72, 73), extending TMLE to handle missing outcome data (39, 51, 74), and the missing covariate missing indicator (MCMI) approach for handling missing confounder data (20, 48, 63). No study has compared these missing data methods with each other in terms of bias and precision. Also, while a requirement for valid inference with MI is that the imputation should incorporate all relationships assumed to hold in the analysis method, (81) the optimal implementation of MI when using TMLE with data-adaptive methods for ACE estimation is unknown. Answers to these questions are key to

developing guidance for appropriate handling of missing data in the context of the growing use of TMLE in applied epidemiological research.

In the current paper, we seek to address this knowledge gap using a simulation study based on an illustrative example from the Victorian Adolescent Health Cohort Study (VAHCS). Our interest was to compare the performance of missing data methods readily available to applied researchers to inform current practice. We begin by introducing the VAHCS example, then describe methods for handling missing data with TMLE, present the simulation study we conducted to evaluate and compare the performance of these approaches, then illustrate the assessed approaches in the VAHCS example. We conclude with a general discussion.

Illustrative example

Our example was based on a previous investigation using data from VAHCS, a longitudinal cohort study of 1,943 participants (1,000 females), recruited at ages 14-15 years, between 1992-1993.(82) Data were collected from participants during their adolescence (waves one-six). Wave seven, conducted in 1998, was the young adulthood survey. Investigators aimed to estimate the ACE of self-reported frequent cannabis use in adolescent females (across waves two-six) on mental health in young adulthood (wave seven), measured using the Clinical Interview Schedule-Revised (CIS-R). We revisited this analysis using TMLE with data-adaptive approaches to adjust for the following confounders measured across waves two-six: parental divorce, antisocial behaviour, depression and anxiety, alcohol use, parental education.(83) **Table 1** shows descriptive statistics for analysis variables, and age at wave two, which is useful as an auxiliary variable in MI (a predictor of missing values but not included in the analysis method (81)). All variables had some degree of missingness, ranging from 0.1% (parental divorce) to 30.8% (frequent cannabis use).

Methods for handling missing data in TMLE

Approaches proposed for handling missing data when estimating the ACE using TMLE in studies like our example are described below.

Non-MI approaches

Complete-case analysis (CCA): records with missing data for any target analysis variable are deleted and only records with complete data are used.(84) This approach generally leads to loss of precision (85) and, depending on missingness mechanism, may inflict bias.(84)

Extended TMLE in the sample with complete exposure and confounders (Ext-TMLE+CEC): records with missing data for Z and X are deleted. Then $E[Y|X, Z]$ is estimated from records with complete data on Y , and the predictions of the outcome are updated in the targeting step using information from models for $P[X=1|Z]$ and $P[M_Y=0|X, Z]$, where M_Y is the missingness indicator for the outcome.(86) Updated predictions for the outcome under exposure and no exposure are obtained for all records, regardless of whether they have missing outcome.(86) As for the exposure and outcome models, the model for M_Y can be fitted using

data-adaptive approaches. With missingness only in the outcome, the extended TMLE method is unbiased under an extended exchangeability assumption ($Y^x \perp\!\!\!\perp M_Y \mid X, Z$ and $Y^x \perp\!\!\!\perp X \mid Z$ for $x = 0, 1$, where Y^x is the potential outcome when $X = x$ and $\perp\!\!\!\perp$ denotes independence).(87)

Extended TMLE plus missing covariate missing indicator (MCMI) approach (Ext-TMLE+MCMI): missing outcome data are handled using the extended TMLE approach, and missing confounder data by including missingness indicators for the incomplete confounders in the confounding adjustment set. Records with missing exposure data are excluded. In settings with complete exposure and outcome data, the MCMI approach can be expected to yield an unbiased estimate of the ACE under an extended exchangeability assumption ($Y^x \perp\!\!\!\perp X \mid Z, M_Z$ for $x = 0, 1$, where M_Z is the vector of missingness indicators for the incomplete confounders), and the assumption that the exposure or outcome depend on the confounder only when the confounder is observed.(88, 89) This assumption might be plausible in some settings, such as with electronic health record data where the decision to prescribe a medication might be influenced by family history of disease only when the clinician has the relevant information.

MI approaches

The MI fully conditional specification (FCS) framework (90) enables simultaneous handling of missing exposure, confounder, and outcome data. Under this approach, univariate imputation models are specified for each incomplete variable conditional on other variables and imputations are drawn sequentially until convergence.(90) The process is repeated multiple times to generate multiple completed datasets. Analysis is performed within each completed dataset and the results are pooled using Rubin's rules to obtain the final estimate and its standard error (SE).(90) For valid inference with MI, each univariate imputation model should be tailored to be compatible with the analysis method. To achieve this, all variables (i.e., exposure, outcome, confounders) and complexities such as interaction terms in the target analysis, and auxiliary variables should be included as predictors in each univariate imputation model.(90) There are various possible implementations of MI within the FCS framework:

Parametric MI with no interaction (MI-no int): each univariate imputation model is based on a regression model with main-effects terms only, which is the default in most MI software implementations. In the example and simulations, we consider main-effects logistic regression for the binary variables and predictive mean matching (PMM) for the continuous outcome based on a main-effects linear regression. In PMM, imputed values are drawn using the nearest observed value after fitting the regression (90) which makes it more robust to misspecification in the imputation model, for example in the presence of nonlinear associations.(91)

Parametric MI with two-way interactions (MI-2-way int): each univariate imputation models is based on a regression model as above, but including two-way exposure-outcome, exposure-confounder, confounder-outcome, and confounder-confounder interactions. Interaction terms are generated within each

cycle of the MI algorithm from current values of relevant variables involved in the interaction term (the so-called “passive” approach in *mice* in R).(92)

Parametric MI with two-, three-, and four-way interactions (MI-higher int): as above, but imputation models additionally include three- and four-way confounder-confounder interactions in the imputation models.

MI using classification and regression trees (MI-CART): instead of regression, for each variable with missing data a tree is fitted using a recursive partitioning technique, with all other variables as predictors. Each record belongs to a donor leaf, from which a randomly selected value for the variable is taken as the imputed value.(93) *MI-CART* (and *MI using random forest (MI-RF)*, see below) have been proposed to enable imputation that can more flexibly allow for interactions and non-linearities.(93)

MI using random forest (MI-RF): for each variable with missing data, multiple bootstrap samples are drawn and for each a separate tree is fitted. Each tree contributes a donor leaf, and a randomly selected value for the variable from all these donors is taken as the imputed value.(93)

All the above approaches can be implemented with the *mice* package in R (92).

Simulation study

To compare the performance of the described methods for handling missing data, we performed a simulation study based on the VAHCS example (**Figure 2**). We considered four scenarios. For each, we generated 2,000 datasets of 2,000 records.

Generating the complete data

We used parametric regression models to generate the variables, where the values of parameters in the models were determined by fitting similar models to the available data in VAHCS (unless stated otherwise). We considered two data generating scenarios (simple and complex), different in the confounder-confounder interaction terms involved in the data generation models. No exposure-confounder interaction terms were included (no effect modification). **Supplementary Table 1** provides the parameter values used for simulating the data and **Supplementary Table 2** the descriptive statistics of the variables in the simulated data.

For both scenarios, we generated a continuous auxiliary variable A (age at wave 2) and a set of confounders $Z = (Z_1$ (parental divorce), Z_2 (antisocial behaviour), Z_3 (depression and anxiety), Z_4 (alcohol use), Z_5 (parental education)). The models for generating these variables are (all binary variables coded 0/1 and $\text{logit}^{-1}(\cdot) = \exp(\cdot) / (1 + \exp(\cdot))$):

$$A \sim N(0,1)$$

$$\begin{aligned}
Z_1 &\sim \text{Binomial}(1, \text{logit}^{-1}(\alpha_0)) \\
Z_2 &\sim \text{Binomial}(1, \text{logit}^{-1}(\beta_0 + \beta_1 A)) \\
Z_3 &\sim \text{Binomial}(1, \text{logit}^{-1}(\gamma_0 + \gamma_1 A)) \\
Z_4 &\sim \text{Binomial}(1, \text{logit}^{-1}(\delta_0 + \delta_1 A)) \\
Z_5 &\sim \text{Binomial}(1, \text{logit}^{-1}(\zeta_0))
\end{aligned}$$

The scenarios differed in the exposure and outcome generation models:

Simple scenario: we used main-effects regression models to generate a binary exposure X (frequent cannabis use) and a continuous outcome Y (log-transformed, standardized CIS-R total score):

$$\begin{aligned}
X_{\text{simple}} &\sim \text{Binomial}(1, \text{logit}^{-1}(\eta_0 + \eta_1 Z_1 + \eta_2 Z_2 + \eta_3 Z_3 + \eta_4 Z_4 + \eta_5 Z_5 + \eta_6 A)) \\
Y_{\text{simple}} &\sim N(\theta_0 + \theta_1 X + \theta_2 Z_1 + \theta_3 Z_2 + \theta_4 Z_3 + \theta_5 Z_4 + \theta_6 Z_5, \text{sd} = 1)
\end{aligned}$$

Complex scenario: we used regression models that included confounder-confounder interactions (excluding interactions with Z_2 because of the low prevalence (15%)) to generate the exposure and outcome:

$$\begin{aligned}
X_{\text{complex}} &\sim \text{Binomial}(1, \text{logit}^{-1}(\eta_0^* + \eta_1 Z_1 + \eta_2 Z_2 + \eta_3 Z_3 + \eta_4 Z_4 + \eta_5 Z_5 + \eta_6 A + \eta_7 Z_1 Z_3 + \eta_8 Z_1 Z_4 \\
&\quad + \eta_9 Z_1 Z_5 + \eta_{10} Z_3 Z_4 + \eta_{11} Z_3 Z_5 + \eta_{12} Z_4 Z_5)) \\
Y_{\text{complex}} &\sim N(\theta_0^* + \theta_1 X + \theta_2 Z_1 + \theta_3 Z_2 + \theta_4 Z_3 + \theta_5 Z_4 + \theta_6 Z_5 + \theta_7 Z_1 Z_3 + \theta_8 Z_1 Z_4 + \theta_9 Z_1 Z_5 + \theta_{10} Z_3 Z_4 \\
&\quad + \theta_{11} Z_3 Z_5 + \theta_{12} Z_4 Z_5 + \theta_{13} Z_1 Z_3 Z_4 + \theta_{14} Z_1 Z_3 Z_5 + \theta_{15} Z_1 Z_4 Z_5 + \theta_{16} Z_3 Z_4 Z_5 \\
&\quad + \theta_{17} Z_1 Z_3 Z_4 Z_5, \text{sd} = 1)
\end{aligned}$$

We set the coefficients for the interaction terms approximately four times larger than values estimated in the VAHCS data. Under both outcome generation models, we set the coefficients for X (θ_1), which is the true value of the ACE to 0.2. For this effect and 2,000 records, the null hypothesis of no causal effect is formally rejected ($p < 0.05$) in approximately 80% of the simulated datasets.

Imposing missing data

We considered two missingness scenarios, A and B, depicted by m-DAGs (directed acyclic graphs) with missingness indicators for each variable with incomplete data as nodes (**Figure 3**). (85, 94)

To reflect the VAHCS example, we imposed missingness on Z_2 , Z_3 , Z_4 , X and Y, through generating missingness indicators M_{Z_2} , M_{Z_3} , M_{Z_4} , M_X and M_Y , coded 0 if the variable was observed and 1 if missing. We considered variables A, Z_1 , and Z_5 , which had a small proportion of missing values within VAHCS (<10% each, **Table 1**) as fully observed in the simulation study. The models used for generating the missingness indicators were:

$$\begin{aligned}
M_{Z_2} &\sim \text{Binomial}(1, \text{logit}^{-1}(\iota_0 + \iota_1 Z_1 + \iota_2 Z_5 + \iota_3 Z_2 + \iota_4 X + \iota_5 Y)) \\
M_{Z_3} &\sim \text{Binomial}(1, \text{logit}^{-1}(\kappa_0 + \kappa_1 Z_1 + \kappa_2 Z_5 + \kappa_3 Z_3 + \kappa_4 X + \kappa_5 Y + \kappa_6 M_{Z_2})) \\
M_{Z_4} &\sim \text{Binomial}(1, \text{logit}^{-1}(\lambda_0 + \lambda_1 Z_1 + \lambda_2 Z_5 + \lambda_3 Z_4 + \lambda_4 X + \lambda_5 Y + \lambda_6 M_{Z_2} + \lambda_7 M_{Z_3})) \\
M_X &\sim \text{Binomial}(1, \text{logit}^{-1}(\nu_0 + \nu_1 Z_1 + \nu_2 Z_5 + \nu_3 Z_2 + \nu_4 Z_3 + \nu_5 Z_4 + \nu_6 X + \nu_7 Y + \nu_8 M_{Z_2} + \nu_9 M_{Z_3} \\
&\quad + \nu_{10} M_{Z_4})) \\
M_Y &\sim \text{Binomial}(1, \text{logit}^{-1}(\xi_0 + \xi_1 Z_1 + \xi_2 Z_5 + \xi_3 Z_2 + \xi_4 Z_3 + \xi_5 Z_4 + \xi_6 X + \xi_7 Y + \xi_8 M_{Z_2} + \xi_9 M_{Z_3} \\
&\quad + \xi_{10} M_{Z_4} + \xi_{11} M_X))
\end{aligned}$$

For each missingness indicator, we set the coefficients for confounders without missing data (Z_1, Z_5), confounders with missing data (Z_2, Z_3, Z_4), and exposure to 0.9. We set the coefficients of the outcome to 0 for m-DAG A and 0.1 for m-DAG B. The missingness proportions were the same in both missingness scenarios and approximately the same as in the real VAHCS dataset, except for the outcome, which was increased to 20% (13% in VAHCS).

Analysis of the simulated data

The target analysis aimed to estimate the ACE of X on Y using TMLE with data-adaptive methods adjusting for Z_1 to Z_5 as confounders. We used the *TMLE* package in R (86). We fitted the exposure and outcome models using SuperLearner, following the current advice for selecting the candidate algorithms in the SuperLearner library, which is to include a range of parametric, semiparametric, and non-parametric methods.(95, 96) The selected methods were: mean (the average), glm (generalized linear model), glm.interaction (generalized linear model with 2-way interactions between all pairs of variables), bayesglm (Bayesian generalized linear model), gam (generalized additive model), glmnet (elastic net regression), earth (multivariate adaptive regression splines), rpart (recursive partitioning and regression trees), rpartPrune (recursive partitioning with pruning) and ranger (random forest). The SE for TMLE was obtained using the variance of the influence function.(9) This analysis was applied to each simulated incomplete dataset alongside each of the previously described missing data methods. For *Ext-TMLE+CEC* and *Ext-TMLE+MCMI* we used the same SuperLearner library for the outcome missingness model. We used the *mice* package in R to implement MI.(92) Due to computational constraints, for each MI approach, we generated five imputed datasets (see Discussion).(90) **Supplementary Tables 3 and 4** show the variables and interaction terms included in each imputation model for *MI-2-way int* and *MI-higher int*. We used the default settings of the *mice* package for the donor pool for PMM in parametric MI approaches and for the hyperparameters for *MI-CART* and *MI-RF*.(92)

Evaluation criteria

We compared the performance of the approaches for handling missing data by calculating the relative bias percent, the empirical standard errors (SE), and the percent error in average model-based SE relative to the empirical SE. For all, Monte-Carlo SEs were obtained.(97)

Analyses were performed in R version 3.6.1.(98)

Simulation Study Results

Relative bias

In the simple scenario, under m-DAG A, *CCA* and *Ext-TMLE+CEC* yielded small biases (<3%). *Ext-TMLE+MCMI* was more biased (10%) (**Figure 4**). These three approaches led to somewhat larger biases under m-DAG B (ranging from -13% to -16%). The three parametric MI approaches performed similarly to each other under both m-DAGs, yielding small biases (<|8%|). *MI-CART* performed similarly to

the parametric MI approaches under m-DAG A (relative bias -8%), but it had higher bias under m-DAG B (-20%). Of all the approaches, *MI-RF* had the highest bias under both m-DAGs (-32% m-DAG A, -42% m-DAG B).

Biases in the complex scenario, for both m-DAGs, were similar to the simple scenario, except that parametric MI without interaction had higher bias than in the simple scenario (33% m-DAG A, 23% m-DAG B)

Empirical standard error and relative error in model-based standard error

For each missing data method, the empirical SEs (**Figure 5**) were similar across the scenarios and m-DAGs. The SEs using *CCA* and *Ext-TMLE+CEC* were similar (0.13-0.16) and larger than *Ext-TMLE+MCMI* (0.12-0.14). Except for *MI-RF*, which had a lower empirical SE (0.7-0.8), the SEs obtained from the MI approaches were similar to each other (0.10-0.14) and to *Ext-TMLE+MCMI*.

The model SEs were underestimated using non-MI methods and overestimated using MI methods across all scenarios and m-DAGs (**Figure 6**). The errors were smaller under the simple than the complex scenario. Within each scenario, the performance among non-MI approaches was similar. The performance of the MI approaches was similar within each scenario, except *MI-RF*, which produced model SEs with considerably larger error.

Illustrative example results

We analysed the VAHCS example using the *tmle* package in R,(86) and applied the eight missing data methods described. Unlike in the simulations, a small proportion of participants had missing data for parental divorce and parental education (**Table 1**), which were handled here in the same way as missing data for the other confounders. Also, the auxiliary variable age had 9.3% missing data, which was multiply imputed in all the MI approaches. For the MI approaches, 100 imputations were performed. The obtained effect sizes were small, with *MI-no int* yielding somewhat larger effect size and non-MI methods and *MI-RF* smaller effect sizes (**Table 2**). The SEs for MI approaches were larger than the non-MI methods, which could be explained by the downward and upward biases in model SEs for non-MI and MI approaches, respectively, observed in our simulation study (**Figure 6**). For example, using the relative percent error in model SEs averaged over the four scenarios in the simulations, the corrected SEs in the case study would be 0.14 for *CCA*, 0.14 for *MI-no int* and 0.11 for *MI-RF*.

Discussion

We compared the performance of a range of methods for handling missing data when estimating the ACE using TMLE with data-adaptive approaches. We considered simple (exposure/outcome generated from main-effects regressions) and complex (models also included interactions) scenarios and two realistic

missingness mechanisms. Parametric MI approaches with interactions were the best performing approaches across all settings assessed in terms of bias and variance reduction. Non-MI methods had the largest variance and, along with MI using machine learning, had larger biases when the outcome influenced missingness in other variables. Parametric MI without interactions – the default in most MI software – yielded estimates with the highest bias in the complex scenarios.

Previous work by Moreno-Betancur et al illustrated that when the outcome variable did not influence missingness in any variable (e.g., our m-DAG A), the conditional distribution of the outcome could be consistently estimated from the available data using an appropriate procedure (the target parameter is said to be “recoverable”), and it can be expressed as the conditional distribution among the complete cases.(85) This was confirmed by findings in the present study where, under m-DAG A, the *CCA* and *Ext-TMLE+CEC* had small bias in both the simple and complex scenarios. The *Ext-TMLE+MCMI* approach, however, had higher bias. A key assumption under which the MCMI approach has been shown to be unbiased is when the exposure or outcome only depend on the confounder when the confounder is observed.(88, 89) We did not consider missingness scenarios where this held, because this assumption is implausible in a prospective cohort study, such as VAHCS, where the data are not being used for medical decision-making. Under m-DAG A, except for parametric MI without interactions in complex scenario and MI-RF in both scenarios, MI approaches also performed well in both simple and complex scenarios and improved precision compared with non-MI methods.

Also in reference (85), although the target parameter was conjectured to be non-recoverable when the outcome variable influenced missingness in variables other than the outcome (e.g., our m-DAG B), the regression-adjusted exposure-outcome association (which is equivalent to the ACE in the absence of effect modification) could still be estimated with negligible bias using MI in their simulation study.(85) Similarly, in our simulations, under the simple scenario, the use of parametric MI with no interactions generally led to small bias for both m-DAGs. For the complex scenario, where the imputation model with no interactions was misspecified and likely incompatible with the analysis method, MI with no interactions had high bias but including interaction terms in the imputation model improved the performance of parametric MI. In our simulation, the complex scenario was defined by the presence of confounder-confounder interaction terms in the exposure and outcome generating models, meaning that the MI models with interactions would be approximately correctly specified. Unfortunately, in practice, the data generating process underlying observational data is rarely fully understood. This limited knowledge, which itself is a motivation for using TMLE with data-adaptive approaches, makes defining parametric imputation models that are compatible with the analysis method challenging.

Within the FCS framework, recursive partitioning techniques, such as CART and RF, have been suggested as alternative approaches that could automatically incorporate interactions and non-linearities in the imputation process.(93) Two previous simulation studies have shown that MI using CART performs better

than parametric MI without interaction terms.(93, 99) However, in these studies, the target analysis was a correctly specified outcome regression model with interactions, and biases in estimates of the main effects were not that different following MI using CART or parametric MI without interaction. These studies imposed missingness in the outcome only (93) or outcome and covariates.(99) In both, missingness depended on fully observed variables. In the present study, *MI-CART* performed similarly to parametric MI approaches with interactions for m-DAG A but led to larger bias for m-DAG B. Bias in the ACE estimates following *MI-RF* was larger than *MI-CART*, in line with Dove et al.'s results, where it was speculated that this poorer performance might be because the tree building process for RF could miss interactions.(93).

In the present study, all non-MI approaches underestimated the model SE, which was not surprising. Standard TMLE variance estimation, which is based on the variance of the influence curve,(9) is valid if both the exposure and outcome models are consistently estimated and the Donsker class condition is satisfied.(10) It is, however, unclear if the latter is met when data-adaptive approaches are used for the exposure and outcome models.(100) This bias has been observed in other simulation studies,(96, 100, 101) and developing approaches to tackle it is an area of ongoing research. Additionally, Rubin's MI variance estimator is expected to perform poorly in the presence of incompatibility,(102) which might explain the overestimation of model SEs for the MI approaches. Incompatibility is the key challenge for using MI with TMLE with data-adaptive approaches, in terms of bias of point estimates as discussed previously, but even more so for bias in variance estimates. A promising alternative approach for obtaining SEs for MI in the presence of incompatibility has been recently proposed using the bootstrap,(102) but we did not explore this because of computational constraints.

Our simulation study was broadly based on VAHCS to emulate a realistic scenario. We evaluated the performance of missing data methods under realistic missingness mechanisms. We used m-DAGs instead of the more familiar framework to describe missingness mechanisms as missing completely at random (MCAR), missing at random (MAR), or missing not at random (MNAR) because with missingness in multiple variables, it is difficult to comprehend and substantively assess the meaning and plausibility of MAR assumption. Also, although it is possible to estimate key parameters unbiasedly if the MAR assumption holds, MAR is not necessary for unbiased estimation.(85) We did not consider missingness mechanisms where outcome influenced its own missingness, under which none of the approaches could be expected to perform well. For each MI approach, due to computational constraints we generated five completed datasets in the simulation study, which is fewer than we would do in practice.(90) We do not expect this to have affected the comparison between MI approaches, but it could have affected comparison of non-MI with MI methods. Our simulated data had a relatively simple structure across the assessed scenarios, and we assumed no effect modification. Extensions of our study could investigate the performance of these missing data methods for datasets with high-dimensional confounders, binary outcome, in the presence of effect modification, and more complex m-DAGs including longitudinal auxiliary variables.

Conclusion

We evaluated the performance of eight available approaches to handle missing data when estimating the ACE using TMLE with data-adaptive approaches under two data generation scenarios and two missingness mechanisms. Our results highlight the importance of considering the missingness mechanism when choosing a method to handle missing data. Parametric MI that incorporates interactions and non-linearities appears to be the best approach in the context of TMLE with data-adaptive approaches.

References

1. Greenland S, Robins JM, Pearl J. Confounding and collapsibility in causal inference. *Stat Sci* 1999;14(1):29-46.
2. Hernan MA. A definition of causal effect for epidemiological research. *J Epidemiol Commun H* 2004;58(4):265-71.
3. Hernan MA, Robins JM. Using Big Data to Emulate a Target Trial When a Randomized Trial Is Not Available. *Am J Epidemiol* 2016;183(8):758-64.
4. Rubin DB. Estimating Causal Effects of Treatments in Randomized and Nonrandomized Studies. *J Educ Psychol* 1974;66(5):688-701.
5. Rubin DB. Causal inference using potential outcomes: Design, modeling, decisions. *Journal of the American Statistical Association* 2005;100(469):322-31.
6. Hernan MA, Robins JM. *Causal Inference: What If*. Boca Raton: Chapman & Hall/CRC. 2020.
7. Schuler MS, Rose S. Targeted Maximum Likelihood Estimation for Causal Inference in Observational Studies. *Am J Epidemiol* 2017;185(1):65-73.
8. van der Laan MJ, Rubin DB. Targeted maximum likelihood learning. *Int J Biostat* 2006 2(1):Article 11.
9. van der Laan MJ, Rose S. *Targeted Learning: Causal Inference for Observational and Experimental Data*. New York, NY Springer-Verlag New York; 2011.
10. van der Laan MJ, Starmans RJCM. Entering the Era of Data Science: Targeted Learning and the Integration of Statistics and Computational Data Analysis. *Advances in Statistics* 2014;2014:502678.
11. Arnold B, Arana B, Mäusezahl D, et al. Evaluation of a pre-existing, 3-year household water treatment and handwashing intervention in rural Guatemala. *Int J Epidemiol* 2009;38(6):1651-61.
12. Rosenblum M, Deeks SG, van der Laan M, et al. The risk of virologic failure decreases with duration of HIV suppression, at greater than 50% adherence to antiretroviral therapy. *PloS one* 2009;4(9):e7196.
13. Mackey DC, Hubbard AE, Cawthon PM, et al. Usual physical activity and hip fracture in older men: an application of semiparametric methods to observational data. *Am J Epidemiol* 2011;173(5):578-86.
14. Padula AM, Mortimer K, Hubbard A, et al. Exposure to traffic-related air pollution during pregnancy and term low birth weight: estimation of causal associations in a semiparametric model. *Am J Epidemiol* 2012;176(9):815-24.
15. Wester CW, Stitelman OM, deGruttola V, et al. Effect modification by sex and baseline CD4+ cell count among adults receiving combination antiretroviral therapy in Botswana: results from a clinical trial. *AIDS Res Hum Retroviruses* 2012;28(9):981-8.
16. Legrand M, Pirrachio R, Rosa A, et al. Incidence, risk factors and prediction of post-operative acute kidney injury following cardiac surgery for active infective endocarditis: an observational study. *Crit Care* 2013;17(5):R220.
17. Decker AL, Hubbard A, Crespi CM, et al. Semiparametric Estimation of the Impacts of Longitudinal Interventions on Adolescent Obesity using Targeted Maximum-Likelihood: Accessible Estimation with the ltmle Package. *J Causal Inference* 2014;2(1):95-108.
18. Kotwani P, Balzer L, Kwarisiima D, et al. Evaluating linkage to care for hypertension after community-based screening in rural Uganda. *Trop Med Int Health* 2014;19(4):459-68.
19. Leslie HH, Karasek DA, Harris LF, et al. Cervical cancer precursors and hormonal contraceptive use in HIV-positive women: application of a causal model and semi-parametric estimation methods. *PloS one* 2014;9(6):e101090.
20. Ahern J, Karasek D, Luedtke AR, et al. Racial/Ethnic Differences in the Role of Childhood Adversities for Mental Disorders Among a Nationally Representative Sample of Adolescents. *Epidemiology* 2016;27(5):697-704.
21. Gianfrancesco MA, Balzer L, Taylor KE, et al. Genetic risk and longitudinal disease activity in systemic lupus erythematosus using targeted maximum likelihood estimation. *Genes Immun* 2016;17(6):358-62.
22. Hsu LI, Briggs F, Shao X, et al. Pathway Analysis of Genome-wide Association Study in Childhood Leukemia among Hispanics. *Cancer Epidemiol Biomarkers Prev* 2016;25(5):815-22.
23. Kreif N, Gruber S, Radice R, et al. Evaluating treatment effectiveness under model misspecification: A comparison of targeted maximum likelihood estimation with bias-corrected matching. *Stat Methods Med Res* 2016;25(5):2315-36.
24. Pearl M, Balzer L, Ahern J. Targeted Estimation of Marginal Absolute and Relative Associations in Case-Control Data: An Application in Social Epidemiology. *Epidemiology* 2016;27(4):512-7.
25. Salihi HM, Das R, Morton L, et al. Racial Differences in DNA-Methylation of CpG Sites Within Preterm-Promoting Genes and Gene Variants. *Matern Child Health J* 2016;20(8):1680-7.
26. Davis FM, Sutzko DC, Grey SF, et al. Predictors of surgical site infection after open lower extremity revascularization. *J Vasc Surg* 2017;65(6):1769-78.e3.
27. Herrera R, Berger U, von Ehrenstein OS, et al. Estimating the Causal Impact of Proximity to Gold and Copper Mines on Respiratory Diseases in Chilean Children: An Application of Targeted Maximum Likelihood Estimation. *Int J Environ Res Public Health* 2017;15(1).
28. Kreif N, Tran L, Grieve R, et al. Estimating the Comparative Effectiveness of Feeding Interventions in the Pediatric Intensive Care Unit: A Demonstration of Longitudinal Targeted Maximum Likelihood Estimation. *Am J Epidemiol* 2017;186(12):1370-9.
29. Skeem JL, Manchak S, Montoya L. Comparing Public Safety Outcomes for Traditional Probation vs Specialty Mental Health Probation. *JAMA Psychiatry* 2017;74(9):942-8.
30. Sukul D, Seth M, Dixon SR, et al. Contemporary Trends and Outcomes Associated With the Preprocedural Use of Oral P2Y12 Inhibitors in Patients Undergoing Percutaneous Coronary Intervention: Insights From the Blue Cross Blue Shield of Michigan Cardiovascular Consortium (BMC2). *J Invasive Cardiol* 2017;29(10):340-51.
31. Pearl M, Ahern J, Hubbard A, et al. Life-course neighbourhood opportunity and racial-ethnic disparities in risk of preterm birth. *Paediatr Perinat Epidemiol* 2018;32(5):412-9.

32. Platt JM, McLaughlin KA, Luedtke AR, et al. Targeted Estimation of the Relationship Between Childhood Adversity and Fluid Intelligence in a US Population Sample of Adolescents. *Am J Epidemiol* 2018;187(7):1456-66.
33. Skeem JL, Montoya L, Manchak SM. Comparing Costs of Traditional and Specialty Probation for People With Serious Mental Illness. *Psychiatr Serv* 2018;69(8):896-902.
34. Torres JM, Rudolph KE, Sofrygin O, et al. Longitudinal associations between having an adult child migrant and depressive symptoms among older adults in the Mexican Health and Aging Study. *Int J Epidemiol* 2018;47(5):1432-42.
35. Berkowitz SA, Basu S, Gundersen C, et al. State-Level and County-Level Estimates of Health Care Costs Associated with Food Insecurity. *Prev Chronic Dis* 2019;16:E90.
36. Gianfrancesco MA, Trupin L, Shiboski S, et al. Smoking Is Associated with Higher Disease Activity in Rheumatoid Arthritis: A Longitudinal Study Controlling for Time-varying Covariates. *J Rheumatol* 2019;46(4):370-5.
37. Izano MA, Sofrygin OA, Picciotto S, et al. Metalworking Fluids and Colon Cancer Risk: Longitudinal Targeted Minimum Loss-based Estimation. *Environ Epidemiol* 2019;3(1):e035.
38. Lim S, Tellez M, Ismail AI. Estimating a Dynamic Effect of Soda Intake on Pediatric Dental Caries Using Targeted Maximum Likelihood Estimation Method. *Caries Res* 2019;53(5):532-40.
39. Rodríguez-Molina D, Barth S, Herrera R, et al. An educational intervention to improve knowledge about prevention against occupational asthma and allergies using targeted maximum likelihood estimation. *Int Arch Occup Environ Health* 2019;92(5):629-38.
40. Torres JM, Rudolph KE, Sofrygin O, et al. Having an Adult Child in the United States, Physical Functioning, and Unmet Needs for Care Among Older Mexican Adults. *Epidemiology* 2019;30(4):553-60.
41. You Y, Doubova SV, Pinto-Masis D, et al. Application of machine learning methodology to assess the performance of DIABETIMSS program for patients with type 2 diabetes in family medicine clinics in Mexico. *BMC Med Inform Decis Mak* 2019;19(1):221.
42. Yu YH, Bodnar LM, Brooks MM, et al. Comparison of Parametric and Nonparametric Estimators for the Association Between Incident Prepregnancy Obesity and Stillbirth in a Population-Based Cohort Study. *Am J Epidemiol* 2019;188(7):1328-36.
43. Aiemjoy K, Aragie S, Wittberg DM, et al. Seroprevalence of antibodies against Chlamydia trachomatis and enteropathogens and distance to the nearest water source among young children in the Amhara Region of Ethiopia. *PLoS Negl Trop Dis* 2020;14(9):e0008647.
44. Bell-Gorrod H, Fox MP, Boule A, et al. The Impact of Delayed Switch to Second-Line Antiretroviral Therapy on Mortality, Depending on Failure Time Definition and CD4 Count at Failure. *Am J Epidemiol* 2020.
45. Bodnar LM, Cartus AR, Kirkpatrick SI, et al. Machine learning as a strategy to account for dietary synergy: an illustration based on dietary intake and adverse pregnancy outcomes. *Am J Clin Nutr* 2020;111(6):1235-43.
46. Clare PJ, Dobbins T, Bruno R, et al. The overall effect of parental supply of alcohol across adolescence on alcohol-related harms in early adulthood—a prospective cohort study. *Addiction* 2020;115(10):1833-43.
47. Decruyenaere A, Steen J, Colpaert K, et al. The obesity paradox in critically ill patients: a causal learning approach to a casual finding. *Crit Care* 2020;24(1):485.
48. Ehrlich SF, Neugebauer RS, Feng J, et al. Exercise During the First Trimester and Infant Size at Birth: Targeted Maximum Likelihood Estimation of the Causal Risk Difference. *Am J Epidemiol* 2020;189(2):133-45.
49. Kagawa RMC, Pear VA, Rudolph KE, et al. Distress level and daily functioning problems attributed to firearm victimization: sociodemographic-specific responses. *Ann Epidemiol* 2020;41:35-42.e3.
50. Kempker RR, Mikiashvili L, Zhao Y, et al. Clinical Outcomes Among Patients With Drug-resistant Tuberculosis Receiving Bedaquiline- or Delamanid-Containing Regimens. *Clin Infect Dis* 2020;71(9):2336-44.
51. Mozafar Saadati H, Mehrabi Y, Sabour S, et al. Estimating the effects of body mass index and central obesity on stroke in diabetics and non-diabetics using targeted maximum likelihood estimation: Atherosclerosis Risk in Communities study. *Obes Sci Pract* 2020;6(6):628-37.
52. Puryear SB, Balzer LB, Ayieko J, et al. Associations between alcohol use and HIV care cascade outcomes among adults undergoing population-based HIV testing in East Africa. *Aids* 2020;34(3):405-13.
53. Torres JM, Sofrygin O, Rudolph KE, et al. US Migration Status of Adult Children and Cognitive Decline Among Older Parents Who Remain in Mexico. *Am J Epidemiol* 2020;189(8):761-9.
54. Veit C, Herrera R, Weinmayr G, et al. Long-term effects of asthma medication on asthma symptoms: an application of the targeted maximum likelihood estimation. *BMC Med Res Methodol* 2020;20(1):307.
55. Westling T, Cowden C, Mwananyanda L, et al. Impact of chlorhexidine baths on suspected sepsis and bloodstream infections in hospitalized neonates in Zambia. *Int J Infect Dis* 2020;96:54-60.
56. Abdollahpour I, Nedjat S, Almasi-Hashiani A, et al. Estimating the Marginal Causal Effect and Potential Impact of Waterpipe Smoking on Risk of Multiple Sclerosis Using the Targeted Maximum Likelihood Estimation Method: A Large, Population-Based Incident Case-Control Study. *Am J Epidemiol* 2021;190(7):1332-40.
57. Almasi-Hashiani A, Nedjat S, Ghiasvand R, et al. The causal effect and impact of reproductive factors on breast cancer using super learner and targeted maximum likelihood estimation: a case-control study in Fars Province, Iran. *BMC Public Health* 2021;21(1):1219.
58. Amusa L, Zewotir T, North D, et al. Association of medical male circumcision and sexually transmitted infections in a population-based study using targeted maximum likelihood estimation. *BMC Public Health* 2021;21(1):1642.
59. Aris IM, Sarvet AL, Stensrud MJ, et al. Separating Algorithms From Questions and Causal Inference With Unmeasured Exposures: An Application to Birth Cohort Studies of Early Body Mass Index Rebound. *Am J Epidemiol* 2021;190(7):1414-23.

60. Beydoun HA, Beydoun MA, Huang S, et al. Hospitalization outcomes among brain metastasis patients receiving radiation therapy with or without stereotactic radiosurgery from the 2005-2014 Nationwide Inpatient Sample. *Sci Rep* 2021;11(1):19209.
61. Crouner JR, Marston WA, Freeman NLB, et al. The Society for Vascular Surgery Objective Performance Goals for Critical Limb Ischemia are attainable in select patients with ischemic wounds managed with wound care alone. *Ann Vasc Surg* 2021.
62. Dadi AF, Miller ER, Woodman RJ, et al. Effect of perinatal depression on risk of adverse infant health outcomes in mother-infant dyads in Gondar town: a causal analysis. *BMC Pregnancy Childbirth* 2021;21(1):255.
63. Ehrlich SF, Ferrara A, Hedderston MM, et al. Exercise During the First Trimester of Pregnancy and the Risks of Abnormal Screening and Gestational Diabetes Mellitus. *Diabetes Care* 2021;44(2):425-32.
64. Goel AR, Bruce HA, Williams N, et al. Long-Term Effects of Hearing Aids on Hearing Ability in Patients with Sensorineural Hearing Loss. *J Am Acad Audiol* 2021.
65. Hickey MD, Ayieko J, Owaraganise A, et al. Effect of a patient-centered hypertension delivery strategy on all-cause mortality: Secondary analysis of SEARCH, a community-randomized trial in rural Kenya and Uganda. *Plos Med* 2021;18(9):e1003803.
66. Ikeda T, Cooray U, Murakami M, et al. Maintaining Moderate or Vigorous Exercise Reduces the Risk of Low Back Pain at 4 Years of Follow-Up: Evidence From the English Longitudinal Study of Ageing. *J Pain* 2021.
67. Kahkoska AR, Abrahamson TJ, Alexander GC, et al. Association Between Glucagon-Like Peptide 1 Receptor Agonist and Sodium-Glucose Cotransporter 2 Inhibitor Use and COVID-19 Outcomes. *Diabetes Care* 2021;44(7):1564-72.
68. Kerschberger B, Boule A, Kuwengwa R, et al. The Impact of Same-Day Antiretroviral Therapy Initiation Under the World Health Organization Treat-All Policy. *Am J Epidemiol* 2021;190(8):1519-32.
69. Mozafar Saadati H, Sabour S, Mansournia MA, et al. Effect modification of general and central obesity by sex and age on cardiovascular outcomes: Targeted maximum likelihood estimation in the atherosclerosis risk in communities study. *Diabetes Metab Syndr* 2021;15(2):479-85.
70. Nardone A, Rudolph KE, Morello-Frosch R, et al. Redlines and Greenspace: The Relationship between Historical Redlining and 2010 Greenspace across the United States. *Environ Health Perspect* 2021;129(1):17006.
71. Nyabuti MN, Petersen ML, Bukusi EA, et al. Characteristics of HIV seroconverters in the setting of universal test and treat: Results from the SEARCH trial in rural Uganda and Kenya. *PloS one* 2021;16(2):e0243167.
72. Olivas-Martínez A, Cárdenas-Fragoso JL, Jiménez JV, et al. In-hospital mortality from severe COVID-19 in a tertiary care center in Mexico City; causes of death, risk factors and the impact of hospital saturation. *PloS one* 2021;16(2):e0245772.
73. Potter K, Masteller B, Balzer LB. Examining Obedience Training as a Physical Activity Intervention for Dog Owners: Findings from the Stealth Pet Obedience Training (SPOT) Pilot Study. *Int J Environ Res Public Health* 2021;18(3).
74. Rossides M, Kullberg S, Di Giuseppe D, et al. Infection risk in sarcoidosis patients treated with methotrexate compared to azathioprine: A retrospective 'target trial' emulated with Swedish real-world data. *Respirology* 2021;26(5):452-60.
75. Saadati HM, Sabour S, Mansournia MA, et al. The Direct Effect of Body Mass Index on Cardiovascular Outcomes among Participants Without Central Obesity by Targeted Maximum Likelihood Estimation. *Arq Bras Cardiol* 2021;116(5):879-86.
76. Shiba K, Torres JM, Daoud A, et al. Estimating the Impact of Sustained Social Participation on Depressive Symptoms in Older Adults. *Epidemiology* 2021;32(6):886-95.
77. Sun X, Wang L, Li H, et al. Identification of microenvironment related potential biomarkers of biochemical recurrence at 3 years after prostatectomy in prostate adenocarcinoma. *Aging (Albany NY)* 2021;13(12):16024-42.
78. Torres JM, Mitchell UA, Sofrygin O, et al. Associations between spousal caregiving and health among older adults in Mexico: A targeted estimation approach. *Int J Geriatr Psychiatry* 2021;36(5):775-83.
79. Wang L, Sun X, Jin C, et al. Identification of Tumor Microenvironment-Related Prognostic Biomarkers for Ovarian Serous Cancer 3-Year Mortality Using Targeted Maximum Likelihood Estimation: A TCGA Data Mining Study. *Front Genet* 2021;12:625145.
80. Zou R, El Marroun H, Cecil C, et al. Maternal folate levels during pregnancy and offspring brain development in late childhood. *Clin Nutr* 2021;40(5):3391-400.
81. Lee KJ, Tilling KM, Cornish RP, et al. Framework for the treatment and reporting of missing data in observational studies: The Treatment And Reporting of Missing data in Observational Studies framework. *J Clin Epidemiol* 2021;134:79-88.
82. Patton GC, Coffey C, Romaniuk H, et al. The prognosis of common mental disorders in adolescents: a 14-year prospective cohort study. *Lancet* 2014;383(9926):1404-11.
83. Patton GC, Coffey C, Carlin JB, et al. Cannabis use and mental health in young people: cohort study. *Bmj* 2002;325(7374):1195-8.
84. White IR, Carlin JB. Bias and efficiency of multiple imputation compared with complete-case analysis for missing covariate values. *Stat Med* 2010;29(28):2920-31.
85. Moreno-Betancur M, Lee KJ, Leacy FP, et al. Canonical Causal Diagrams to Guide the Treatment of Missing Data in Epidemiologic Studies. *Am J Epidemiol* 2018;187(12):2705-15.
86. Susan G, Mark JvdL. tmle : An R Package for Targeted Maximum Likelihood Estimation. *Journal of Statistical Software* 2012;51(13).
87. Díaz I, van der Laan MJ. Doubly robust inference for targeted minimum loss-based estimation in randomized trials with missing outcome data. *Stat Med* 2017;36(24):3807-19.
88. Blake HA, Leyrat C, Mansfield KE, et al. Propensity scores using missingness pattern information: a practical guide. *Stat Med* 2020;39(11):1641-57.

89. Blake HA, Leyrat C, Mansfield KE, et al. Estimating treatment effects with partially observed covariates using outcome regression with missing indicators. *Biom J* 2020;62(2):428-43.
90. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011;30(4):377-99.
91. Morris TP, White IR, Royston P. Tuning multiple imputation by predictive mean matching and local residual draws. *BMC Med Res Methodol* 2014;14:75.
92. van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software* 2011;45(3):1-67.
93. Doove LL, Van Buuren S, Dusseldorp E. Recursive partitioning for missing data imputation in the presence of interaction effects. *Computational Statistics and Data Analysis* 2014;72:92-104.
94. Mohan K, Pearl J. Graphical Models for Processing Missing Data. *Journal of the American Statistical Association* 2021;116(534):1023-37.
95. Naimi AI, Balzer LB. Stacked generalization: an introduction to super learning. *Eur J Epidemiol* 2018;33(5):459-64.
96. Naimi AI, Mishler AE, Kennedy EH. Challenges in Obtaining Valid Causal Effect Estimates with Machine Learning Algorithms. *arXiv e-prints* 2020.
97. Morris TP, White IR, Crowther MJ. Using simulation studies to evaluate statistical methods. *Stat Med* 2019;38(11):2074-102.
98. R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
99. Burgette LF, Reiter JP. Multiple imputation for missing data via sequential regression trees. *Am J Epidemiol* 2010;172(9):1070-6.
100. Balzer LB, Westling T. Demystifying Statistical Inference When Using Machine Learning in Causal Research. *Am J Epidemiol* 2021.
101. van der Laan MJ, Gruber S. Targeted Minimum Loss Based Estimation of an Intervention Specific Mean Outcome. U.C. Berkeley Division of Biostatistics Working Paper Series. Working Paper 290. 2011
102. Bartlett JW, Hughes RA. Bootstrap inference for multiple imputation under uncongeniality and misspecification. *Stat Methods Med Res* 2020;29(12):3533-46.

Table 1 – Description of variables used in the case study, their distribution and the proportion with missing data among the VAHCS female participants (n=1,000)

	Variable	Type	Grouping/unit	Notation	N (%*) coded 1 or mean (SD)	% with missing data
Confounder	Parental divorce	Binary	0=Not divorced/separated by wave 6 1= Divorced/separated by wave 6	Z1	221 (22.1)	0.1
	Antisocial behaviour	Binary	0=No across all waves 2 to 6 1=Yes at any wave 2 to 6	Z2	106 (14.6)	27.4
	Depression and anxiety	Binary	0=CIS-R score <12 across all waves 2 to 6 1=CIS-R score ≥12 at any wave 2 to 6	Z3	516 (59.9)	13.8
	Alcohol use	Binary	0=No across all waves 2 to 6 1=Yes at any wave 2 to 6	Z4	294 (37.2)	21.0
	Parental education	Binary	0=Did not complete high school by wave 6 1=Completed high school by wave 6	Z5	364 (37.7)	3.4
Exposure	Frequent cannabis use	Binary	0=Less than weekly use across all waves 2 to 6 1=At least weekly use at any wave 2 to 6	X	86 (12.4)	30.8
Outcome	CIS-R total score	Continuous	z-score, measured at wave 7	Y	0 (1)	13.4
Auxiliary variable	Age	Continuous	years, measured at wave 2	A	15.4 (0.4)	9.3
With any missing data						40.3

Abbreviations N number; SD standard deviation; CIS-R revised clinical interview schedule *Proportions reported among those with observed data for the variable


Table 2 – Estimated average causal effect (ACE) of frequent cannabis use during adolescence on CIS-R score (standardised z-score) using a TMLE approach under different missing data methods within the VAHCS case study

Missing data method	ACE (Difference in means)*	Standard error	95% confidence interval	Time to run
Complete-case	0.09	0.12	-0.14, 0.32	16.4 sec
Ext TMLE	0.12	0.11	-0.09, 0.33	11.2 sec
Ext TMLE+MCMI	0.13	0.13	-0.13, 0.39	21.7 sec
MI, no int	0.20	0.16	-0.11, 0.50	4.6 min
MI, 2-way int	0.16	0.17	-0.17, 0.49	5.8 min
MI, higher int	0.18	0.16	-0.13, 0.49	5.8 min
MI, CART	0.15	0.16	-0.16, 0.45	11.8 min
MI, RF	0.13	0.18	-0.21, 0.48	14.1 min

*Average causal effect estimated as the difference in the mean potential outcome under exposure and under no exposure


Abbreviations - Ext TML: Extended targeted maximum likelihood estimation (TMLE) approach; Ext TMLE+MCMI: extended TMLE plus missing covariate missing indicator (MCMI) approach; MI, no int (linear): parametric multiple imputation (MI) with no interaction – linear regression to impute missing outcome; MI no int: parametric MI with no interaction – predictive mean matching to impute missing outcome; MI, 2-way int: parametric MI with two-way interactions; MI higher int: parametric MI with two-, three-, and four-way interactions; MI, CART: MI using classification and regression trees; MI, RF: MI using random forest

Figure 1 – Number of studies applying targeted maximum likelihood estimation (TMLE) approach by year of publication between 2009 and November 2021 (total number of studies = 70)




Studies(11-80) were identified via a search of PubMed for papers that included “targeted maximum likelihood estimation (TMLE)” in their title or abstract. Abstracts were screened to identify studies whose focus was not on methodology and had applied TMLE to estimate causal effect of interest.

Figure 2 – Directed acyclic graph (DAG) used in data generation for the simulation study




Abbreviations: CIS-R revised clinical interview schedule

Figure 3 – Missingness directed acyclic graphs (m-DAGs) illustrating the missingness scenarios considered in the simulation study. Figure has been adapted from Moreno-Betancur et al. ⁽⁸⁵⁾




For simplicity of exposition, confounders without missing data (Z_1 and Z_5) are presented on a single node and confounders with missing data (Z_2, Z_3, Z_4) on another single node. Also, only one missingness indicator has been included for confounders with missing data (M_Z), coded as 1 when any of the variables Z_2, Z_3, Z_4 have missing data and as 0 when none has missing data.

Figure 4 – **Relative bias percent** (%) in ACE estimation (filled circles with error bars showing \pm Monte Carlo standard errors) using different missing data methods for simple and complex scenarios and m-DAGs A and B.




Abbreviations – *Ext-TMLE*: extended targeted maximum likelihood estimation (TMLE) approach; *Ext-TMLE+MCMI*: extended TMLE plus missing covariate missing indicator (MCMI) approach; *MI-no int (linear)*: parametric multiple imputation (MI) with no interaction – linear regression to impute missing outcome; *MI-no int*: parametric MI with no interaction – predictive mean matching to impute missing outcome; *MI-2-way int*: parametric MI with two-way interactions; *MI-higher int*: parametric MI with two-, three-, and four-way interactions; *MI-CART*: MI using classification and regression trees; *MI-RF*: MI using random forest

Figure 5 – **Empirical standard error** in ACE estimation (filled circles with error bars showing \pm Monte Carlo standard errors) using different missing data methods for simple and complex scenarios and m-DAGs A and B.



Abbreviations – *Ext-TMLE*: Extended targeted maximum likelihood estimation (TMLE) approach; *Ext-TMLE+MCMI*: extended TMLE plus missing covariate missing indicator (MCMI) approach; *MI-no int (linear)*: parametric multiple imputation (MI) with no interaction – linear regression to impute missing outcome; *MI-no int*: parametric MI with no interaction – predictive mean matching to impute missing outcome; *MI-2-way int*: parametric MI with two-way interactions; *MI-higher int*: parametric MI with two-, three-, and four-way interactions; *MI-CART*: MI using classification and regression trees; *MI-RF*: MI using random forest

Figure 6 – **Relative % error in model standard error** in ACE estimation (filled circles with error bars showing \pm Monte Carlo standard errors) using different missing data methods for simple and complex scenarios and m-DAGs A and B.



Abbreviations – *Ext-TMLE*: Eextended targeted maximum likelihood estimation (TMLE) approach; *Ext-TMLE+MCMI*: extended TMLE plus missing covariate missing indicator (MCMI) approach; *MI-no int (linear)*: parametric multiple imputation (MI) with no interaction – linear regression to impute missing outcome; *MI-no int*: parametric MI with no interaction – predictive mean matching to impute missing outcome; *MI-2-way int*: parametric MI with two-way interactions; *MI-higher int*: parametric MI with two-, three-, and four-way interactions; *MI-CART*: MI using classification and regression trees; *MI-RF*: MI using random fore

Supplementary Table 1 - Coefficient values used to simulate the variables and missingness indicators under the three assessed scenarios (simple, intermediate, and complex)

Model for	Regression coefficient of													
	Intercept	Z1	Z2	Z3	Z4	Z5	X	Y	A	MZ2**	MZ3**	MZ4**	MX**	
Complete data	Z1	-1.30												
	Z2	-1.90							0.40					
	Z3	0.40							0.70*					
	Z4	-0.60							-0.70*					
	Z5	-0.50												
	Simple scenario													
	X	-2.9*	1.30	1.90	0.40	0.20*	-0.30			0.70*				
	Y	-0.7*	0.10	0.40*	0.70	0.20*	0.30*	0.20*						
	Complex scenarios													
	X	-2.4*	1.30	1.90	0.40	0.20	-0.30							
Y	-0.70*	0.10	0.40*	0.70	0.20*	0.30*	0.20*							
DAG A	MZ2	-1.75	0.90	0.90		0.90	0.90							
	MZ3	-5.70	0.90		0.90	0.90	0.90			4.10				
	MZ4	-4.80	0.90			0.90	0.90	0.90		3.20	2.00			
	MX	-3.80	0.90	0.90	0.90	0.90	0.90	0.90		1.50	1.50	1.50		
	MY	-3.20	0.90	0.90	0.90	0.90	0.90	0.90		-0.60	0.10	0.10	0.10	
DAG B	MZ2	-1.75	0.90	0.90		0.90	0.90	0.10						
	MZ3	-5.95	0.90		0.90	0.90	0.90	0.10		4.30				
	MZ4	-4.80	0.90			0.90	0.90	0.90	0.10	3.50	1.30			
	MX	-3.80	0.90	0.90	0.90	0.90	0.90	0.90	0.10	1.50	1.50	1.50		
	MY	-3.20	0.90	0.90	0.90	0.90	0.90	0.90		-0.60	0.10	0.10	0.20	
* For the complex scenarios models for X and Y also included interactions as follows:														
	z1z3	z1z4	z1z5	z3z4	z3z5	z4z5	z1z3z4	z1z3z5	z1z4z5	z3z4z5	z1z3z4z5			
Complex scenario 1														
X	-1.60	-1.20	-0.50	-0.60	0.30	-1.50								
Y	-0.50	1.00	0.10	0.10	0.40	-0.10	-1.20	-1.00	-0.10	-0.40	1.70			
Complex scenario 2														
X	-3.20	-2.30	-1.00	-1.20	0.50	-2.90								
Y	-0.90	2.00	0.10	0.20	0.70	-0.20	-2.40	-2.00	-0.30	-0.80	3.40			

* These parameter values were modified from what is in the VAHCS data as follows: In generating confounders Z_3 and Z_4, we changed the coefficient value for A from what it was in VAHCS, so that it was a stronger auxiliary variable, and modified the intercepts so that the prevalence of the variables remained the same as in the VAHCS dataset. For the exposure model, we modified the coefficient value for A, so that it was a stronger auxiliary variable, and the coefficient value for Z_4, so that it was less strongly associated with X. We modified the intercept so that the prevalence of X was approximately 15% in the simulated data (12% in VAHCS) in all scenarios. For the outcome model, we modified the coefficient values for Z_2, Z_4, and Z_5, so that they were stronger confounders (the coefficient values for the confounders ranged from 0.1 (for Z_1) to 0.7 (for Z_3) in the simulation study). Under all outcome generation models, we set the coefficient value for X (θ_1), which is the true value of the ACE, to 0.2. We modified the intercept in the outcome model so that the mean of Y remained 0.

**As shown in the models provided in the manuscript, the regression model for generating M_(Z_3) included the missingness indicator M_(Z_2), the model for M_(Z_4) included M_(Z_2) and M_(Z_3), the model for M_X included M_(Z_2), M_(Z_3), and M_(Z_4), and the model for M_Y included all the preceding missingness indicators. We modified the coefficient values for these missingness indicators and the intercepts so that the missingness proportion for each variable and the overall proportions with missing data were the same across all missingness scenarios.

Supplementary Table 2– Description of variables in the simulated data (averaged over 2000 simulations)

	Variable	Type	Notation	%* coded 1 or mean (SD)	% with missing data
Confounder	Parental divorce	Binary	Z1	21	0
	Antisocial behaviour	Binary	Z2	14	30
	Depression and anxiety	Binary	Z3	59	15
	Alcohol use	Binary	Z4	37	20
	Parental education	Binary	Z5	38	0
Exposure	Frequent cannabis use	Binary	X	15	30
Outcome	CIS-R total score	Continuous	Y	0 (1)	20
With missing exposure or confounder data					40
With any missing data					50

Abbreviations SD standard deviation; CIS-R revised clinical interview schedule *Proportions reported in the simulated complete data

Supplementary Table 3 - The variables and interaction terms included in each imputation model for a multiple imputation approach that included all two-way interactions

Variable imputed	Variables included in imputation model																							
	A	Z1	Z2	Z3	Z4	Z5	X	Y	XY	XZ1	YZ1	XZ3	YZ3	XZ4	YZ4	XZ5	YZ5	Z1Z3	Z1Z4	Z1Z5	Z3Z4	Z3Z5	Z4Z5	
Z2	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Z3	1	1	1	0	1	1	1	1	1	1	1	0	0	1	1	1	1	0	1	1	0	0	1	1
Z4	1	1	1	1	0	1	1	1	1	1	1	1	1	0	0	1	1	1	0	1	0	1	0	1
X	1	1	1	1	1	1	0	1	0	0	1	0	1	0	1	0	1	1	1	1	1	1	1	1
Y	1	1	1	1	1	1	1	0	0	1	0	1	0	1	0	1	0	1	1	1	1	1	1	1

