QUANDLE COLORING QUIVERS OF (p, 2)-TORUS LINKS

JAGDEEP BASI AND CARMEN CAPRAU

ABSTRACT. A quandle coloring quiver is a quiver structure, introduced by Karina Cho and Sam Nelson, which is defined on the set of quandle colorings of an oriented knot or link by a finite quandle. We study quandle coloring quivers of (p, 2)-torus knots and links with respect to dihedral quandles.

1. INTRODUCTION

A quandle [7, 12] is a non-associate algebraic structure whose definition is motivated by knot theory, and it provides an axiomatization of the Reidemeister moves for link diagrams. The fundamental quandle of a link is a quandle that can be computed from any diagram of the link. Joyce [7] and Matveev [12] proved that the fundamental quandle is a stronger invariant of knots and links than the knot group. Quandles have been used to define other useful invariants of oriented links (see [1, 2, 3, 4, 5] and others). For example, the quandle counting invariant [4] is the size of the set of all homomorphisms from the fundamental quandle of an oriented link to a finite quandle. More recently, Cho and Nelson [3] introduced the notion of a quandle coloring quiver of an oriented link K with respect to a finite quandle \mathcal{X} , which is a directed graph that allows multiple edges and loops, and enhances the quandle counting invariant of K by \mathcal{X} .

In this paper, we study quandle coloring quivers of (p, 2)-torus links by dihedral quandles. The paper is organized as follows. In Sec. 2 we review some basic concepts about torus links and quandles. In Sec. 2.2 we also prove a couple of statements about the coloring space of (p, 2)-torus links by dihedral quandles, as they will be needed later in the paper. Sec. 3 is the heart of the paper; here we review the definition of quandle coloring quivers of oriented links with respect to finite quandles and study the quivers of (p, 2)-torus links with respect to dihedral quandles.

²⁰²⁰ Mathematics Subject Classification. 57K10, 57K12.

Key words and phrases. Quandles, quandle coloring quivers, torus knots and links. CC was partially supported by Simons Foundation grant #355640.

2. Preliminaries

In order to have a self-contained paper, we begin by reviewing relevant definitions and concepts.

2.1. **Basic definitions.** A knot or link K is called a (p,q)-torus knot or link if it lies, without any points of intersection, on the surface of a trivial torus in \mathbb{R}^3 . The integers p and q represent the number of times K wraps around the meridian and, respectively, the longitude of the torus. We denote a (p,q)-torus knot or link by $\mathcal{T}(p,q)$. It is known that $\mathcal{T}(p,q)$ is a knot if and only if gcd(p,q) = 1 and p and q are not equal to 1 simultaneously, and that it is a link with d components if and only if gcd(p,q) = d. Moreover, $\mathcal{T}(p,q)$ is the trivial knot if and only if either p or q is equal to 1 or -1. We use the term link to refer generically to both knots and links. It is also known that the (p,q)-torus link is isotopic to the (q,p)-torus link. The (-p, -q)-torus link is the (p,q)-torus link with reverse orientation, and torus links are invertible. For simplicity, we will assume that p,q > 0. The mirror image of $\mathcal{T}(p,q)$ is $\mathcal{T}(p,-q)$ and a non-trivial torus-knot is chiral.

It is known that the crossing number of a torus link with p, q > 0 is given by $c = \min((p-1)q, (q-1)p)$. The link $\mathcal{T}(p,q)$ can also be represented as the closure of the braid $(\sigma_1 \sigma_2 \dots \sigma_{p-1})^q$. Since $\mathcal{T}(p,q)$ is isotopic to $\mathcal{T}(q,p)$, this torus link has also a diagram which is the closure of the braid $(\sigma_1 \sigma_2 \dots \sigma_{q-1})^p$.

The simplest nontrivial torus knot is the trefoil knot, $\mathcal{T}(3,2)$, depicted in Fig. 1. For more details on torus knots and links, we refer the reader to the books [9, 10, 11, 13].




FIGURE 1. Oriented torus knot $\mathcal{T}(3,2)$

The term quandle is attributed to Joyce in his 1982 work [7] as an algebraic structure whose axioms are motivated by the Reidemeister moves.

A quandle [7, 8, 12] is a non-empty set \mathcal{X} together with a binary operation $\triangleright : \mathcal{X} \times \mathcal{X} \to \mathcal{X}$ satisfying the following axioms:

- (1) For all $x \in \mathcal{X}, x \triangleright x = x$.
- (2) For all $y \in \mathcal{X}$, the map $\beta_y : \mathcal{X} \to \mathcal{X}$ defined by $\beta_y(x) = x \triangleright y$ is invertible.

(3) For all $x, y, z \in \mathcal{X}$, $(x \triangleright y) \triangleright z = (x \triangleright z) \triangleright (y \triangleright z)$.

In this case, we use the notation $(\mathcal{X}, \triangleright)$ or $(\mathcal{X}, \triangleright_{\mathcal{X}})$, if we need to be specific about the quandle operation.

A quandle homomorphism [6] between two quandles $(\mathcal{X}, \triangleright_{\mathcal{X}})$ and $(\mathcal{Y}, \triangleright_{\mathcal{Y}})$ is a map $f: \mathcal{X} \to \mathcal{Y}$ such that $f(a \triangleright_{\mathcal{X}} b) = f(a) \triangleright_{\mathcal{Y}} f(b)$, for any $a, b \in \mathcal{X}$.

It is not hard to see that the composition of quandle homomorphisms is again a quandle homomorphism. We note that the second axiom of a quandle implies that for all $y \in \mathcal{X}$, the map β_y is an automorphism of \mathcal{X} . Moreover, by the first axiom of a quadle, the map β_y fixes y.

Example 1. Let $n \in \mathbb{Z}, n \geq 2, \mathcal{X} = \mathbb{Z}_n = \{0, 1, \dots, n-1\}$ and define $x \triangleright y = (2y - x) \mod n$, for all $x, y \in \mathbb{Z}_n$. That is, $x \triangleright y$ is the remainder of 2y - x upon division by n. Then \mathbb{Z}_n together with this operation \triangleright is a quandle, called the *dihedral quandle* (see [4, 14]).

Example 2. To every oriented link diagram of K, we can naturally associate its *fundamental quandle* (also referred to as the *knot quandle*), denoted by Q(K). Given a diagram D of K, its crossings divide D into arcs and we use a set of labels to label all of these arcs. The fundamental quandle Q(D) is the quandle generated by the set of arc-labels together with the crossing relations given by the operation \triangleright explained in Fig. 2.




FIGURE 2. Labelling a knot crossing in a fundamental quandle

The axioms of a quandle encode the Reidemeister moves, and thus the fundamental quandle is independent on the diagram D representing the oriented link K. Hence, Q(K) := Q(D) is a link invariant. It was shown independently by Joyce [7] and Matveev [12] that the fundamental quandle is a stronger invariant than the fundamental group of a knot or link. In fact, the fundamental quandle is a complete invariant, in the sense that if $Q(K_1)$ and $Q(K_2)$ are isomorphic quandles, then the (unoriented) links K_1 and K_2 are equivalent (up to chirality). However, determining whether or not two quandles are isomorphic is not an easy task.

Remark 3. We consider the fundamental quandle of a (p, 2)-torus link. By arranging the quandle relations of the fundamental quandle in a particular

way, we can present $Q(\mathcal{T}(p,2))$ in a recursive fashion. We know that a (p,2)-torus link can be represented as the closure of the braid σ_1^p , which is also a minimal diagram for the (p,2)-torus link. There are p crossings and p arcs labeled $\{x_1, x_2, \ldots, x_p\}$ in such a diagram of $\mathcal{T}(p,2)$. Considering the braid σ_1^p , we label its arcs starting from the bottom right with x_1 then moving to the bottom left arc to label it x_2 , and continuing up along the braid using labels x_3, x_4, \ldots, x_p , as shown in Fig. 3. We obtain the following presentation for the fundamental quandle of the (p, 2)-torus link:

$$Q(\mathcal{T}(p,2)) = \langle x_1, x_2, x_3, \dots, x_p \, | \, x_2 \triangleright x_1 = x_p, x_1 \triangleright x_p = x_{p-1}, \text{ and} \\ x_i \triangleright x_{i-1} = x_{i-2}, \text{ for all } 3 \le i \le p \rangle.$$




FIGURE 3. A diagram of a (p, 2)-torus link

When classifying links, it is useful to count the number of quandle homomorphisms from the fundamental quandle of a link to a fixed quandle. For example, Inoue [6] showed that the number of all quandle homomorphisms from the fundamental quandle of a link to an Alexander quandle [4, 6] has a module structure and that it is determined by the series of the Alexander polynomials of the link.

A quandle coloring of an oriented link K with respect to a finite quandle \mathcal{X} (also called an \mathcal{X} -quandle coloring) is given by a quandle homomorphism $\psi: Q(K) \to \mathcal{X}$. In this case, we refer to \mathcal{X} as the coloring quandle. We let $Hom(Q(K), \mathcal{X})$ be the collection of all \mathcal{X} -quandle colorings and we call it the coloring space of K with respect to \mathcal{X} . Finally, the cardinality $|Hom(Q(K), \mathcal{X})|$ of the coloring space is called the quandle counting invariant with respect to \mathcal{X} (see [3]).

When a quandle coloring consists of a single color, we use the term *trivial* coloring, and make use of the term *nontrivial* coloring otherwise.

2.2. The coloring space of (p, 2)-torus links with respect to dihedral quandles. In this paper we are concerned with (p, 2)-torus links, and we need to understand the coloring space of these links with respect to dihedral quandles. The following proposition and theorem are our first results.

Proposition 4. Let $\mathcal{T}(p, 2)$ be a torus link and \mathbb{Z}_n a dihedral quandle. Consider any two fixed and consecutive arcs of $\mathcal{T}(p, 2)$ with labels $x_1, x_2 \in Q(\mathcal{T}(p, 2))$. If $\psi \in Hom(Q(\mathcal{T}(p, 2)), \mathbb{Z}_n)$, then $p\psi(x_1) \equiv p\psi(x_2) \pmod{n}$.

Proof. By Remark 3, the fundamental quandle of the torus link $\mathcal{T}(p, 2)$ has the following presentation:

$$Q(\mathcal{T}(p,2)) = \langle x_1, x_2, x_3, \dots, x_p \,|\, x_2 \triangleright x_1 = x_p, x_1 \triangleright x_p = x_{p-1}, \text{ and} \\ x_i \triangleright x_{i-1} = x_{i-2}, \text{ for all } 3 \le i \le p \rangle.$$

For any \mathbb{Z}_n -quandle coloring $\psi \in Hom(Q(\mathcal{T}(p,2)), \mathbb{Z}_n)$ and $x_i, x_j \in Q(\mathcal{T}(p,2))$, we must have that $\psi(x_i \triangleright x_j) = \psi(x_i) \triangleright' \psi(x_j)$, where \triangleright and \triangleright' are the quandle operations for $Q(\mathcal{T}(p,2))$ and \mathbb{Z}_n , respectively. In particular,

$$\psi(x_p) = \psi(x_2 \triangleright x_1) = \psi(x_2) \triangleright' \psi(x_1) = (2\psi(x_1) - \psi(x_2)) \mod n,$$

$$\psi(x_{p-1}) = \psi(x_1 \triangleright x_p) = \psi(x_1) \triangleright' \psi(x_p) = (2\psi(x_p) - \psi(x_1)) \mod n.$$

Moreover, for all $3 \le i \le p$, the following holds:

$$\psi(x_{i-2}) = \psi(x_i \triangleright x_{i-1}) = \psi(x_i) \triangleright' \psi(x_{i-1}) = (2\psi(x_{i-1}) - \psi(x_i)) \mod n.$$

Using these statements inductively, we obtain:

$$\begin{split} \psi(x_2) &= (2\psi(x_3) - \psi(x_4)) \mod n \\ &= (2(2\psi(x_4) - \psi(x_5)) - \psi(x_4)) \mod n \\ &= (3\psi(x_4) - 2\psi(x_5)) \mod n \\ &= (3(2\psi(x_5) - \psi(x_6)) - 2\psi(x_5)) \mod n \\ &= (4\psi(x_5) - 3\psi(x_6)) \mod n \\ &\vdots \\ &= ((p-1)\psi(x_p) - (p-2)\psi(x_1)) \mod n \\ &= ((p-1)(2\psi(x_1) - \psi(x_2)) - (p-2)\psi(x_1)) \mod n \\ &= (p\psi(x_1) - (p-1)\psi(x_2)) \mod n. \end{split}$$

It follows that $p\psi(x_2) \equiv p\psi(x_1) \pmod{n}$.

We are now ready to prove the following theorem about the coloring space of (p, 2)-torus links with respect to dihedral quandles, which will be used extensively in Sec. 3.

Theorem 5. For a given torus link $\mathcal{T}(p, 2)$ and \mathbb{Z}_n a dihedral quandle, the following holds:

(i) If gcd(p,n) = 1, then $|Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)| = n$ and the coloring space $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$ is the collection of all trivial \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p,2)$.

(ii) If $gcd(p,n) = c \neq 1$, then $|Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)| = nc$, and the coloring space $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$ is the union of all n trivial quandle colorings together with a collection of n(c-1) nontrivial \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p,2)$.

Proof. Let $\psi \in Hom(Q(\mathcal{T}(p,2)), \mathbb{Z}_n)$. By Proposition 4,

$$p\psi(x_1) \equiv p\psi(x_2) \pmod{n}.$$

(i) If gcd(p, n) = 1, then the above congruence simplifies to

$$\psi(x_1) \equiv \psi(x_2) \pmod{n}$$

The latter can only occur when ψ maps x_1 and x_2 to the same color. Using the relations of $Q(\mathcal{T}(p,2))$, as described in Remark 3, this forces the quandle homomorphism ψ to color every arc in the diagram of $\mathcal{T}(p,2)$ the same. Since there are *n* quandle colorings to choose from, $|Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)| = n$, and $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$ is the collection of all trivial quandle colorings of $\mathcal{T}(p,2)$ with respect to \mathbb{Z}_n .

(ii) If $gcd(p, n) = c \neq 1$, then the congruence $p\psi(x_1) \equiv p\psi(x_2) \pmod{n}$ simplifies to the following:

$$\psi(x_1) \equiv \psi(x_2) \pmod{\frac{n}{c}}.$$

Then $\psi(x_2) \equiv \psi(x_1) + k\frac{n}{c} \pmod{n/c}$ also holds for all integers k, where $0 \leq k \leq c-1$. Hence, for each fixed color $\psi(x_1) \in \mathbb{Z}_n$, there are c possibilities for $\psi(x_2)$ in \mathbb{Z}_n such that $p\psi(x_1) \equiv p\psi(x_2) \pmod{n}$. Since there are n choices for $\psi(x_1) \in \mathbb{Z}_n$, we conclude that there are $nc \mathbb{Z}_n$ -quandle colorings of $\mathcal{T}(p, 2)$. Moreover, since there are n trivial quandle colorings with respect to \mathbb{Z}_n , the remaining $n(c-1) \mathbb{Z}_n$ -quandle colorings must be nontrivial quandle colorings of the link $\mathcal{T}(p, 2)$ with respect to \mathbb{Z}_n .

3. Quandle coloring quivers of (p, 2)-torus links with respect to dihedral quandles

A quandle enhancement is an invariant for oriented links that captures the quandle counting invariant and in general is a stronger invariant. For some examples of quandle enhacements, we refer the reader to [1, 2, 4, 5]. More recently, by representing quandle colorings as vertices in a graph, Cho and Nelson [3] imposed a combinatorial structure on the coloring space of a link that led to new quandle enhancements. For example, one of these

 $\mathbf{6}$

enhancements, the quandle coloring quiver, can distinguish links that have the same quandle counting invariant (see [3, Example 6]).

In what follows, we work with graphs that are allowed to have multiple edges and loops. To capture multiple edges in a graph G, we use a weight function $c: \{(x, y) | x, y \in V(G)\} \to \mathbb{N} \cup \{0\}$, where V(G) is the vertex set of G and (x, y) is any edge connecting vertices $x, y \in V(G)$. We denote the corresponding graph with a weight function c by (G, c). When c is a constant function c(x, y) = k, for all $x, y \in V(G)$ and some $k \in \mathbb{N}$, then we use the notation $c = \hat{k}$ for the weight function and (G, \hat{k}) for the corresponding graph.

We say that a graph is *complete*, provided that there is an edge between every pair of distinct vertices and a loop at each vertex; we denote a complete graph with n vertices by K_n . The graphs we work with in this paper are all directed, and we use the notation \overrightarrow{G} to denote a graph G with all edges directed both ways.

Given two graphs G_1 and G_2 with disjoint vertex sets and edge sets, the join of G_1 and G_2 , denoted $G_1 \nabla G_2$, is the graph with vertex set $V(G_1) \cup V(G_2)$ and edge set containing all edges in G_1 and G_2 , as well as all edges that connect vertices of G_1 with vertices of G_2 . Finally, if there is the same number of directed edges from each vertex of the graph G_2 to the graph G_1 with constant weight function \hat{k} and no directed edges from vertices of G_1 to vertices of G_2 , we use the notation $G_1 \nabla_{\hat{k}} G_2$ to denote such a join of graphs.

Let \mathcal{X} be a finite quandle and K an oriented link. For any set of quandle homomorphims $S \subset Hom(\mathcal{X}, \mathcal{X})$, the associated quandle coloring quiver [3], denoted $\mathcal{Q}_X^S(K)$, is the directed graph with a vertex v_f for every element $f \in Hom(Q(K), \mathcal{X})$ and an edge directed from the vertex v_f to the vertex v_g , whenever $g = \phi \circ f$, for some $\phi \in S$. When $S = Hom(\mathcal{X}, \mathcal{X})$, we denote the associated quiver by $\mathcal{Q}_{\mathcal{X}}(K)$, and call it the full \mathcal{X} -quandle coloring quiver of K.

It was explained in [3] that $\mathcal{Q}_X^S(K)$ is a link invariant, in the sense that the quandle coloring quivers associated to diagrams that are related by Reidemeister moves are isomorphic quivers.

Notation. We establish some notation for mappings. Consider a function $g: \mathcal{X} \to \mathcal{Y}$, where $\mathcal{X} = \{x_1, x_2, \ldots, x_n\}$ and $\mathcal{Y} = \{y_1, y_2, \ldots, y_m\}$. If $g(x_i) = y_{j_i}$, for each $x_i \in \mathcal{X}$, we will write g as $g_{y_{j_1}, y_{j_2}, y_{j_3}, \ldots, y_{j_n}}$ (omitting the commas when possible). This notation will simplify the formulas for various quandle homomorphisms considered in this paper.

Our goal is to study quandle coloring quivers of (p, 2)-torus links with respect to dihedral quandles \mathbb{Z}_n . Before we proceed, we need to prove the following result about the space $Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ of quandle automorphisms of the dihedral quandle \mathbb{Z}_n , where $n \geq 3$.

Proposition 6. Consider the dihedral quandle \mathbb{Z}_n , when $n \geq 3$. Then $Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ is a set of size n^2 of quandle homomorphisms $\phi_{\gamma} \colon \mathbb{Z}_n \to \mathbb{Z}_n$ that are defined recursively, where γ is any *n*-tuple $(\gamma_1, \gamma_2, \gamma_3, \ldots, \gamma_n) \in (\mathbb{Z}_n)^n$, such that

$$\phi_{\gamma} := \begin{cases} \phi_{\gamma}(0) = \gamma_1, \\ \phi_{\gamma}(1) = \gamma_2, \\ \phi_{\gamma}(k) = \phi_{\gamma}(k-2) \triangleright \phi_{\gamma}(k-1), & \text{for } 2 \le k \le n-1. \end{cases}$$

Proof. The operation of the dihedral quandle $(\mathbb{Z}_n, \triangleright)$ is given by $x \triangleright y = (2y - x) \mod n$, for any $x, y \in \mathbb{Z}_n$. Then, $k \triangleright (k + 1) = k + 2$, for all $0 \le k \le n-3$, where it follows that $(n-2) \triangleright (n-1) = 0$ and $(n-1) \triangleright 0 = 1$. Since any quandle homomorphism $\phi \in Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ must satisfy $\phi(x \triangleright y) = \phi(x) \triangleright \phi(y)$, for any $x, y \in \mathbb{Z}_n$ (using the same \triangleright), then ϕ must now satisfy $\phi(k) \triangleright \phi(k+1) = \phi(k \triangleright (k+1)) = \phi(k+2)$, for all $0 \le k \le n-3$, further yielding $\phi(n-2) \triangleright \phi(n-1) = \phi(0)$ and $\phi(n-1) \triangleright \phi(0) = \phi(1)$.

Hence, $\phi(0)$ and $\phi(1)$ completely determine ϕ , and we have the following recursive relation for the image of (0, 1, 2, ..., n-1) under ϕ :

$$(0,1,2,\ldots,n-1) \stackrel{\phi}{\mapsto} (\phi(0),\phi(1),\phi(0) \triangleright \phi(1),\phi(1) \triangleright \phi(2),\ldots,\phi(n-3) \triangleright \phi(n-2)).$$

Since there are *n* ways to assign each of $\phi(0)$ and $\phi(1)$ an element in \mathbb{Z}_n , we see that $|Hom(\mathbb{Z}_n, \mathbb{Z}_n)| = n^2$.

Remark 7. From the cyclic behavior of the dihedral quandle \mathbb{Z}_n and the proof of Proposition 6, we see that any homomorphism $\phi_{\gamma} \in Hom(\mathbb{Z}_n, \mathbb{Z}_n)$, where γ is an *n*-tuple $(\gamma_1, \gamma_2, \ldots, \gamma_n) \in (\mathbb{Z}_n)^n$, is completely determined by the images $\phi_{\gamma}(k) = \gamma_{k+1}$ and $\phi_{\gamma}(k+1) = \gamma_{k+2}$, where if k = n - 1 then k + 2 = 0 and if k = n then k + 1 = 0 and k + 2 = 1.

We also remark that using our notation, we can write $\phi_{\gamma} = \phi_{\gamma_1 \gamma_2 \dots \gamma_n}$.

Using Theorem 5 together with Proposition 6 and Corollary 7, we are ready to state and prove our main results regarding quivers.

Theorem 8. Given a torus link $\mathcal{T}(p, 2)$ and \mathbb{Z}_n a dihedral quandle, where gcd(p, n) = 1, the full \mathbb{Z}_n -quandle coloring quiver of $\mathcal{T}(p, 2)$ is the complete directed graph:

$$\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2)) = \left(\overleftrightarrow{K_n}, \hat{n}\right).$$

Proof. By Theorem 5, we know that $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$ is the collection of all *n* trivial \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p,2)$. Hence, the quandle coloring quiver $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2))$ has *n* vertices, one for each of the elements in the coloring space $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$. Let $\phi = \phi_{\gamma_1\gamma_2...\gamma_n} \in Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ be a quandle homomorphism given by: $0 \stackrel{\phi}{\mapsto} \gamma_1, 1 \stackrel{\phi}{\mapsto} \gamma_2, \ldots, (n-1) \stackrel{\phi}{\mapsto} \gamma_n$. In Proposition 6 we proved that $|Hom(\mathbb{Z}_n, \mathbb{Z}_n)| = n^2$, which was due to the fact that a quandle homomorphism $\phi_{\gamma_1\gamma_2...\gamma_n}$ is completely determined by the choices for γ_1 and γ_2 . In fact, by Remark 7, we know that $\phi_{\gamma_1\gamma_2...\gamma_n}$ is determined by the choices for any two consecutive values γ_k and γ_{k+1} , where if k = n then k + 1 = 1.

Let $\psi_i \in Hom(Q(\mathcal{T}(p,2)), \mathbb{Z}_n)$ denote the trivial \mathbb{Z}_n -quandle coloring of $\mathcal{T}(p,2)$ associated to some fixed $i \in \mathbb{Z}_n$ and let v_{ψ_i} be the corresponding vertex in the quandle coloring quiver $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2))$. Without loss of generality, by the above discussion, we can use that any map $\phi_{\gamma_1\gamma_2...\gamma_n}$ is completely determined by the choices for γ_i and γ_{i+1} .

Consider $\phi_{\gamma_1...\gamma_i i \gamma_{i+2}...\gamma_n} \in Hom(\mathbb{Z}_n, \mathbb{Z}_n)$. That is, $\gamma_{i+1} = i$, which means that $i \stackrel{\phi}{\mapsto} i$ under this map. There are *n* maps of the form $\phi_{\gamma_1...\gamma_i i \gamma_{i+2}...\gamma_n}$ (determined by the value of $\gamma_i \in \mathbb{Z}_n$) and they all satisfy that

$$\phi_{\gamma_1\dots\gamma_i i\gamma_{i+2}\dots\gamma_n} \circ \psi_i = \psi_i.$$

In addition, there are no other maps $\phi_{\gamma_1\gamma_2...\gamma_n}$ satisfying $\phi_{\gamma_1\gamma_2...\gamma_n} \circ \psi_i = \psi_i$. It follows that the quiver vertex v_{ψ_i} has n loops that represent ψ_i being fixed by those n maps.

Let $\psi_j \in Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$, where $j \in \mathbb{Z}_n$ is distinct from the fixed value of *i* above. For similar reasons as above, there are *n* quandle homomorphisms in $Hom(\mathbb{Z}_n,\mathbb{Z}_n)$ of the form $\phi_{\gamma_1\ldots\gamma_i j\gamma_{i+2}\ldots\gamma_n}$ that send $i \stackrel{\phi}{\mapsto} j$. Moreover,

$$\phi_{\gamma_1\dots\gamma_i j\gamma_{i+2}\dots\gamma_n} \circ \psi_i = \psi_j,$$

and there are no other maps $\phi_{\gamma_1\gamma_2...\gamma_n}$ satisfying $\phi_{\gamma_1\gamma_2...\gamma_n} \circ \psi_i = \psi_j$. Thus ψ_i is also mapped *n* times to each of the colorings ψ_j . This implies that there are *n* directed edges from the vertex v_{ψ_i} to each of the vertices v_{ψ_j} , for all $j \neq i$.

Hence, the full quandle coloring quiver of $\mathcal{T}(p,2)$ with respect to \mathbb{Z}_n is a graph with n vertices where every vertex has n directed edges from itself to all other vertices, including itself. Thus, $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2)) = \left(\overleftarrow{K_n}, \hat{n}\right)$, whenever gcd(p,n) = 1.

Theorem 9. Let $\mathcal{T}(p,2)$ be a torus link and \mathbb{Z}_n a dihedral quandle. If gcd(p,n) = c where c is prime, then the full quandle coloring quiver of $\mathcal{T}(p,2)$ with respect to \mathbb{Z}_n is the join of two complete directed graphs:

$$\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2)) = \left(\overleftarrow{K_n}, \hat{n}\right) \overleftarrow{\nabla}_{\hat{d}} \left(\overleftarrow{K_{n(c-1)}}, \hat{d}\right),$$

where d = n/c, and the two complete subgraphs correspond to the trivial and nontrivial, respectively, \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p, 2)$.

Proof. Using Theorem 5, the quandle coloring space $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$ is the union of all *n* trivial \mathbb{Z}_n -quandle colorings, together with n(c-1) nontrivial \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p,2)$. Therefore, the quandle coloring quiver $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2))$ contains *n* vertices corresponding to the trivial \mathbb{Z}_n -quandle colorings and n(c-1) vertices corresponding to the nontrivial \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p,2)$.

Let $\psi_{\sigma} = \psi_{\sigma_1 \sigma_2 \sigma_3 \dots \sigma_p}$ and $\psi_{\omega} = \psi_{\omega_1 \omega_2 \omega_3 \dots \omega_p}$ be any two quandle colorings in $Hom(Q(\mathcal{T}(p, 2)), \mathbb{Z}_n)$. Our notation means that $\psi_{\sigma}(x_i) = \sigma_i$ and $\psi_{\omega}(x_i) = \omega_i$, where $\sigma_i, \omega_i \in \mathbb{Z}_n$ and x_i is a generator of the fundamental quandle $Q(\mathcal{T}(p, 2))$.

If both ψ_{σ} and ψ_{ω} are trivial \mathbb{Z}_n -quandle colorings (possibly the same coloring), we know from the proof of Theorem 9 that there are *n* directed edges from the quiver vertex $v_{\psi_{\sigma}}$ to the vertex $v_{\psi_{\omega}}$ (including *n* directed loops from the vertex $v_{\psi_{\sigma}}$ to itself, if $\psi_{\sigma} = \psi_{\omega}$). It follows that there is a complete directed graph $\left(\overrightarrow{K_n}, \hat{n}\right)$ associated with the *n* trivial \mathbb{Z}_n -quandle colorings of $\mathcal{T}(p, 2)$, as a subgraph of the full quandle coloring quiver $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p, 2))$.

If ψ_{σ} is a nontrivial quandle coloring, then $\psi_{\sigma}(x_1) = \sigma_1 \neq \sigma_2 = \psi_{\sigma}(x_2)$, because equivalent consecutive colors in the subscript of ψ_{σ} would impose a trivial \mathbb{Z}_n -quandle coloring. By Proposition 4, for any $x_1, x_2 \in Q(\mathcal{T}(p, 2))$, a quandle coloring $\psi \in Hom(Q(\mathcal{T}(p, 2)), \mathbb{Z}_n)$ must satisfy the congruence $p\psi(x_1) \equiv p\psi(x_2) \pmod{n}$, which reduces to $\psi(x_1) \equiv \psi(x_2) \pmod{d}$, where $d = \frac{n}{c}$ and $c = \gcd(p, n), c \neq 1$. Since $\psi(x_1), \psi(x_2) \in \mathbb{Z}_n$, we have that for each fixed $\psi(x_1)$, there are c possibilities for $\psi(x_2)$ in \mathbb{Z}_n satisfying the above two congruences; specifically, $\psi(x_2) = \psi(x_1) + kd$, where k is an integer such that $0 \leq k \leq c-1$. Moreover, since $\psi_{\sigma}(x_1) \neq \psi_{\sigma}(x_2)$ for a nontrivial quandle coloring ψ_{σ} , then for each fixed $\psi_{\sigma}(x_1)$ there are c-1 possibilities for $\psi_{\sigma}(x_2)$ in \mathbb{Z}_n satisfying $p\psi_{\sigma}(x_1) \equiv p\psi_{\sigma}(x_2) \pmod{n}$.

Let ψ_{σ} be a fixed nontrivial quandle coloring in $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$. Then $\psi_{\sigma} = \psi_{\sigma_1(\sigma_1+kd)\sigma_3...\sigma_p}$, for some $1 \leq k \leq c-1$ and $\sigma_1, \sigma_3, \ldots, \sigma_p \in \mathbb{Z}_n$. Let $\psi_{\omega} = \psi_{\omega_1(\omega_1+hd)\omega_3...\omega_p}$ be any fixed coloring in $Hom(Q(\mathcal{T}(p,2)),\mathbb{Z}_n)$; that is, h is an integer such that $0 \leq h \leq c-1$, where $\omega_1, \omega_3, \ldots, \omega_p \in \mathbb{Z}_n$ and $h \neq 0$ if ψ_{ω} is a nontrivial quandle coloring.

Now suppose $\phi_{\gamma_1\gamma_2...\gamma_n} \in Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ satisfies $\phi_{\gamma_1\gamma_2...\gamma_n} \circ \psi_{\sigma} = \psi_{\omega}$. In particular, this means that

(1)
$$(\phi_{\gamma_1\gamma_2\dots\gamma_n} \circ \psi_{\sigma})(x_1) = \psi_{\omega}(x_1) = \omega_1$$

(2)
$$(\phi_{\gamma_1\gamma_2\dots\gamma_n} \circ \psi_{\sigma})(x_2) = \psi_{\omega}(x_2) = \omega_1 + hd.$$

Since $\psi_{\sigma}(x_1) = \sigma_1$ and $\psi_{\sigma}(x_2) = \sigma_1 + kd$, equations (1) and (2) imply that the following must hold:

$$\phi_{\gamma_1\gamma_2...\gamma_n}(\sigma_1) = \omega_1 \text{ and } \phi_{\gamma_1\gamma_2...\gamma_n}(\sigma_1 + kd) = \omega_1 + hd.$$

In Remark 7 we showed that a homomorphism $\phi_{\gamma_1\gamma_2...\gamma_n} \in Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ is determined by two consecutive values in its subscript. Moreover, by fixing the value of ω_1 , a homomorphism $\phi_{\gamma_1\gamma_2...\omega_1\tau...\gamma_n}$, with some $\tau \in \mathbb{Z}_n$, satisfying equation (1) yields:

$$\begin{split} \phi_{\gamma_1\gamma_2\dots\omega_1\tau\dots\gamma_n}(\sigma_1) &= \omega_1 \\ \phi_{\gamma_1\gamma_2\dots\omega_1\tau\dots\gamma_n}(\sigma_1+1) &= \tau \\ \phi_{\gamma_1\gamma_2\dots\omega_1\tau\dots\gamma_n}(\sigma_1+2) &= (2\tau-\omega_1) \bmod n \\ \phi_{\gamma_1\gamma_2\dots\omega_1\tau\dots\gamma_n}(\sigma_1+3) &= (3\tau-2\omega_1) \bmod n \\ &\vdots \\ \phi_{\gamma_1\gamma_2\dots\omega_1\tau\dots\gamma_n}(\sigma_1+kd) &= (kd\tau-(kd-1)\omega_1) \bmod n \end{split}$$

Hence, a homomorphism $\phi_{\gamma_1\gamma_2...\omega_1\tau...\gamma_n}$ satisfies both of the equations (1) and (2) if and only if

(3)
$$kd\tau - (kd-1)\omega_1 \equiv \omega_1 + hd \pmod{n}.$$

The latter congruence is equivalent to $kd\tau \equiv kd\omega_1 + hd \pmod{n}$, and since gcd(d, n) = d, this congruence reduces further to:

(4)
$$k\tau \equiv k\omega_1 + h \pmod{\frac{n}{d}}.$$

Note that k, ω_1 , and h are considered fixed in this congruence. Since $\frac{n}{d}$ is equal to the prime c and $1 \leq k \leq c-1$, we have that $gcd(k, \frac{n}{d}) = 1$ and thus the congruence in (4) has solutions τ . In particular, τ has a unique solution modulo $\frac{n}{d}$. It follows that for the congruence (3), there are d incongruent solutions modulo n for τ . Therefore, for a given nontrivial quandle coloring ψ_{σ} , there are d homomorphisms $\phi_{\gamma_1\gamma_2...\gamma_n}$ such that $\phi_{\gamma_1\gamma_2...\gamma_n} \circ \psi_{\sigma} = \psi_{\omega}$, for all (trivial and nontrivial) \mathbb{Z}_n -quandle colorings ψ_{ω} of $\mathcal{T}(p, 2)$.

The above reasoning shows that for each vertex $v_{\psi_{\sigma}}$ in the quiver corresponding to a nontrivial coloring, there are d directed edges to all of the vertices in the quiver, including to itself. It follows that the quiver $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2))$ contains as a subgraph, a complete directed graph $\left(\overleftarrow{K_{n(c-1)}}, \hat{d}\right)$ associated with the nontrivial quandle colorings that is joined, using the weight function \hat{d} , to the subgraph associated with trivial colorings:

$$\left(\overleftrightarrow{K_n}, \hat{n}\right) \overleftarrow{\nabla}_{\hat{d}} \left(\overleftrightarrow{K_{n(c-1)}}, \hat{d}\right).$$

Moreover, there is no map in $Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ that sends a fixed value in \mathbb{Z}_n to two distinct values in \mathbb{Z}_n . That is, there is no homomorphism $\phi_{\gamma_1\gamma_2...\gamma_n}$ in $Hom(\mathbb{Z}_n, \mathbb{Z}_n)$ that sends a trivial \mathbb{Z}_n -quandle coloring of $\mathcal{T}(p, 2)$ to a non-trivial \mathbb{Z}_n -quandle coloring of $\mathcal{T}(p, 2)$. Hence, the joining behavior cannot be reciprocated from $(\overleftarrow{K_n}, \widehat{n})$ to $(\overleftarrow{K_{n(c-1)}}, \widehat{d})$.

This completes the proof of $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2)) = (\overleftarrow{K_n}, \hat{n}) \overleftarrow{\nabla}_{\hat{d}} (\overleftarrow{K_{n(c-1)}}, \hat{d}),$ when gcd(p,n) = c and c is prime.

Corollary 10. For a given torus link $\mathcal{T}(p,2)$ and a prime n, we have:

• If $n \nmid p$, then $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2)) = \left(\overleftrightarrow{K_n}, \hat{n}\right)$. • If $n \mid p$, then $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p,2)) = \left(\overleftrightarrow{K_n}, \hat{n}\right) \overleftarrow{\nabla}_{\hat{1}} \left(\overleftrightarrow{K_{n(n-1)}}, \hat{1}\right)$.

Proof. The statements follow as particular cases from Theorems 8 and 9. Since n is prime, if $n \nmid p$, then gcd(n, p) = 1, and Theorem 8 implies the first statement of this corollary. If $n \mid p$, then gcd(n, p) = n where n is prime, and the application of Theorem 9 yields the second statement above. \Box

The following statement is a direct consequence of Corollary 10.

Corollary 11. Let n be a prime.

- If $n \nmid p_1$ and $n \nmid p_2$, then the quandle coloring quivers $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p_1, 2))$ and $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p_2, 2))$ are isomorphic.
- If $n \mid p_1$ and $n \mid p_2$, then the quandle coloring quivers $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p_1, 2))$ and $\mathcal{Q}_{\mathbb{Z}_n}(\mathcal{T}(p_2, 2))$ are isomorphic.

Example 12. As an example, we consider the full \mathbb{Z}_3 -quandle coloring quiver of the trefoil knot $\mathcal{T}(3,2)$. In this case, n = p = 3, and according to Theorem 9, the quiver $\mathcal{Q}_{\mathbb{Z}_3}(\mathcal{T}(3,2))$ has the following form:

$$\mathcal{Q}_{\mathbb{Z}_3}(\mathcal{T}(3,2)) = \left(\overleftrightarrow{K_3}, \hat{3}\right) \overleftarrow{\nabla}_{\hat{1}} \left(\overleftrightarrow{K_6}, \hat{1}\right).$$

In Fig. 4 on the left, we show the graph $(\overline{K_3}, \hat{3})$ associated with the trivial \mathbb{Z}_3 -quandle colorings of $\mathcal{T}(3, 2)$ and on the right of the figure, there is the graph $(\overline{K_6}, \hat{1})$ corresponding to the $n(c-1) = 3 \cdot 2$ nontrivial \mathbb{Z}_3 -quandle colorings of the trefoil knot. Both of these graphs are subgraphs of the full \mathbb{Z}_3 -quandle coloring quiver of the trefoil knot. In Fig. 5 we present the quandle coloring quiver $\mathcal{Q}_{\mathbb{Z}_3}(\mathcal{T}(3, 2))$ as the join of these subgraphs by the weight function $\hat{d} = \hat{1}: (\overline{K_3}, \hat{3}) \overleftarrow{\nabla}_{\hat{1}} (\overline{K_6}, \hat{1}).$

Remark 13. Taniguchi [15] studied quandle coloring quivers of links with respect to dihedral quandles, where the focus was on isomorphic quandle coloring quivers. Among other things, it was proved in [15, Theorem 3.3] that when the dihedral quandle \mathbb{Z}_n is of prime order, then the \mathbb{Z}_n -quandle coloring quivers of two links L_1 and L_2 are isomorphic if and only if the quandle coloring spaces of L_1 and L_2 with respect to the dihedral quandle \mathbb{Z}_n have the same size. That is, it was proved that if n is prime, then $\mathcal{Q}_{\mathbb{Z}_n}(L_1) \cong \mathcal{Q}_{\mathbb{Z}_n}(L_2)$ as quivers if and only if $|Hom(Q(L_1),\mathbb{Z}_n)| =$ $|Hom(Q(L_2),\mathbb{Z}_n)|$. Hence, for n prime, we can use Taniguchi's result in

12

combination with our results in Theorem 5 and Corollary 10, to say something about the full \mathbb{Z}_n -quandle coloring quiver of a certain link L. For this, we would need to have that $|Hom(Q(L), \mathbb{Z}_n)| = |Hom(Q(\mathcal{T}(p, 2)), \mathbb{Z}_n)|$, for a particular link L, a torus link $\mathcal{T}(p, 2)$, and a prime n. Specifically, if n is prime and $|Hom(Q(L), \mathbb{Z}_n)| = n$ for a certain link L, then $\mathcal{Q}_{\mathbb{Z}_n}(L) \cong$ $\left(\overrightarrow{K_n}, \hat{n}\right)$. Moreover, if n is prime and $|Hom(Q(L), \mathbb{Z}_n)| = n^2$, then $\mathcal{Q}_{\mathbb{Z}_n}(L) \cong$ $\left(\overrightarrow{K_n}, \hat{n}\right) \overleftarrow{\nabla}_{\hat{1}} \left(\overrightarrow{K_{n(n-1)}}, \hat{1}\right)$.




FIGURE 4. The subgraph components of the quiver $\mathcal{Q}_{\mathbb{Z}_3}(\mathcal{T}(3,2))$




FIGURE 5. The quandle coloring quiver $\mathcal{Q}_{\mathbb{Z}_3}(\mathcal{T}(3,2))$

References

- J. S. Carter, M. Elhamdadi, M. Graña and M. Saito, Cocycle knot invariants from quandle modules and generalized quandle homology, *Osaka Journal of Mathematics* 42, No. 3 (2005), 499-541.
- [2] J. S. Carter, D. Jelsovsky, S. Kamada, L. Langford and M. Saito, Quandle cohomology and state-sum invariants of knotted curves and surfaces, *Transactions of the American Mathematical Society* 355, No. 10 (2003), 3947-3989.
- [3] K. Cho and S. Nelson, Quandle coloring quivers, Journal of Knot Theory and Its Ramifications 28, No. 1, 1950001 (2019).
- [4] M. Elhamdadi and S. Nelson, Quandles: An introduction to the algebra of knots, Student Mathematical Library 74, American Mathematical Society, 2015.
- [5] G.R. Grindstaff and S. Nelson, Lie ideal enhacements of counting invariants, Osaka Journal of Mathematics 53, No. 4 (2016), 1015-1027.
- [6] A. Inoue, Quandle homomorphisms of knot quandles to Alexander quandles, Journal of Knot Theory and Its Ramifications 10, No. 6 (2001), 813-821.
- [7] D. Joyce, A classifying invariant of knots, the knot quandle, Journal of Pure and Applied Algebra 23, No. 1 (1982), 37-65.
- [8] S. Kamada, Knot invariants derived from quandles and racks, Geometry & Topology Monographs 4 (2002), 103-117.
- [9] A. Kawauchi, A survey of knot theory, *Birkchäuser*,1996.
- [10] W. B. R. Lickorish, An introduction to knot theory, Springer-Verlag, New York, 1997.
- [11] C. Livingston, Knot theory, The Mathematical Association of America, Washington DC, 1993.
- [12] S. V. Matveev, Distributive groupoids in knot theory, Mathematics of the USSR-Sbornik 47, No. 1 (1984), 73-83.
- [13] K. Murasugi, Knot theory and its applications, Birkchäuser, 1996.
- [14] M. Takasaki, Abstraction of symmetric transformations, Tohoku Mathematical Journal 49 (1943), 145-207.
- [15] Y. Taniguchi, Quandle coloring quivers of links using dihedral quandles, Journal of Knot Theory and Its Ramifications 30, No.2, 2150011 (2021).

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, FRESNO 5245 NORTH BACKER AVENUE, M/S PB108, CA 93740, USA

Email address: jbasi@protonmail.com

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, FRESNO 5245 NORTH BACKER AVENUE, M/S PB108, CA 93740, USA Email address: ccaprau@csufresno.edu