

THE SHIFT-DIMENSION OF MULTIPERSISTENCE MODULES

WOJCIECH CHACHÓLSKI, RENÉ CORBET, AND ANNA-LAURA SATTELBERGER

ABSTRACT. We present the shift-dimension of multipersistence modules and investigate its algebraic properties. This gives rise to a new invariant of multigraded modules over the multivariate polynomial ring arising from the hierarchical stabilization of the zeroth total multigraded Betti number. We give a fast algorithm for the computation of the shift-dimension of interval modules in the bivariate case. We construct multipersistence contours that are parameterized by multivariate functions and hence provide a large class of feature maps for machine learning tasks.

Acknowledgments. We are grateful to Mats Boij for insightful discussions. W.C. was partially supported by VR, the Wallenberg AI, Autonomous System and Software Program (WASP) funded by Knut and Alice Wallenberg Foundation, and MultipleMS funded by the European Union under the Horizon 2020 program, grant agreement 733161, and dBRAIN collaborative project at digital futures at KTH. R.C. got supported by the Brummer & Partners MathDataLab. A.-L.S. thanks the Brummer & Partners MathDataLab for its hospitality during her research stay at KTH Stockholm in 2021.

CONTENTS

1. Introduction	1
2. Measuring distances between multipersistence modules	3
3. The shift-dimension of multipersistence modules	7
4. The shift-dimension of multigraded $\mathbb{K}[x_1, \dots, x_r]$ -modules	16
5. Outlook to future work	18
References	19

1. INTRODUCTION

Topological data analysis (TDA) reveals structures of data by topological methods. One main tool is persistent homology. To the data, one associates a filtered simplicial complex. For every natural number n , one then takes the n -th homology with coefficients in a field \mathbb{K} ; the $(n$ -th) *persistence module* of the data. The persistent homology of a one-parameter filtration is algebraically well understood; by a basic structure theorem from algebra, the persistence module is uniquely determined by its barcode [18] in the discrete setting [14, 30] as well as in the real setting [3] under mild finiteness assumptions. Furthermore, the barcode representation of a persistence module is stable [12]. From the barcode, one reads the birth and death times of topological features. The case of several parameters, i.e., the study of multifiltered simplicial complexes and their homology as introduced by Carlsson and Zomorodian in [8] allows the extraction of finer information from the data. As was shown in [11], the homology

2020 *Mathematics Subject Classification.* 55N31, 16W50 (primary), 16G20, 68W30 (secondary).

Key words and phrases. Topological data analysis, multiparameter persistence, stable invariants, multigraded modules, persistence contours.

modules of a multifiltered simplicial complex can be obtained as the homology of a chain complex $F_1 \rightarrow F_2 \rightarrow F_3$ of free multigraded modules of finite rank by combinatorial techniques. This allows analysis of multipersistence modules using standard computer algebra software. However, the multiparameter case is algebraically intricate. In contrast to the case of a single parameter, as pointed out in [8], the respective moduli space is not zero-dimensional. Moreover, constructing stable and algorithmic invariants of multipersistence modules is challenging and is currently an active branch of research in topological data analysis.

We investigate a stable invariant of multipersistence modules that is based on the hierarchical stabilization of discrete invariants as introduced by Scomamario et al. in [27] and by Gäfvert and Chachólski in [17]. This construction requires the choice of a distance between multipersistence modules, and it turns a discrete invariant into a measurable real-valued function. The obtained invariant is continuous with respect to the chosen distance on multipersistence modules and with respect to the L_p distance on measurable real-valued functions. In this article, we focus on the invariant given by the zeroth total multigraded Betti number β_0 , often called *rank* of the multipersistence module in a TDA setup, defined as the minimal number of generators. Its hierarchical stabilization is commonly referred to as “stable rank” and was introduced in [17].

For the hierarchical stabilization, we also need to be able to construct distances on the space of tame persistence modules. A way to construct a big class of metrics other than the well-known interleaving distance [23] and matching distance [2, 21] is via so-called *persistence contours*. Persistence contours are rather well-studied in the one-parameter case. In [10], for example, the authors list various contours and show how classification, using the first one-persistence homology group, of point processes in the unit square (sampled with respect to different probability distributions) can be improved by choosing the right contour. For the case of several parameters, only a small number of examples of persistence contours is known besides the standard contours. The latter are parameterized by vectors. Persistence contours can be described by simple noise systems [17, 27]. Those are closely related to so-called “superlinear families” that were introduced independently in [6], a generalization of which can be found in [4]. Therein, no extensive class of parameterized such objects was given yet, but it is essential for machine learning tasks to have such a class at one’s disposal.

Multigraded modules over the polynomial ring $\mathbb{K}[x_1, \dots, x_r]$ with the standard multigrading can be considered naturally as multipersistence modules. In this article, we investigate the hierarchical stabilization of β_0 for multipersistence modules obtained in this way with respect to a broader family of contours. The obtained invariant is called *shift-dimension*.

The main contribution of this article is twofold. The first main contribution is the realization that the shift-dimension can be constructed and defined entirely algebraically without referring to the geometry of multipersistence modules induced by the choice of a distance. We give an explicit description of the shift-dimension and investigate some of its algebraic properties. We thereby establish a connection between topological data analysis and combinatorial commutative algebra. In general, the computation of the stable rank is algorithmic but NP-hard [17]. For the bivariate case, we present a linear-time algorithm for the computation of the shift-dimension of interval modules and therefore for the computation of the shift-dimension of quotients of monomial ideals.

The second main contribution lies in the construction of multipersistence contours. We present a class of contours that is parameterized not only by vectors, but by multivariate functions. In doing so, we provide a large class of multipersistence contours.

Related work. The investigation of invariants of multipersistence modules is currently a prominent topic in TDA. For two parameters, the Hilbert function and the rank invariant of [8] are implemented in RIVET [28] for an interactive data analysis. It has been used in applications such as [20]. Furthermore, minimal presentations can be computed efficiently for big data sets using RIVET [24] or `mpfree` [22]. Invariants such as the Hilbert series, associated primes, and local cohomology have been investigated [19], as well as the multirank function [29] and the compressed multiplicity [15].

Outline. This article is organized as follows. In Section 2, we review basic notions about multipersistence modules, how to measure distances between them, and the hierarchical stabilization of discrete invariants. We present several new contours for the multiparameter case. Section 3 introduces and studies the shift-dimension of tame persistence modules from an algebraic and combinatorial perspective. In particular, we focus on the investigation of (non-)additivity. We present a linear-time algorithm for the computation of the shift-dimension of interval modules in the bivariate case. In Section 4, we study the corresponding invariant of multigraded modules over the multivariate polynomial ring and investigate its algebraic properties. In Section 5, we give an outlook to open problems and future work.

2. MEASURING DISTANCES BETWEEN MULTIPERSISTENCE MODULES

2.1. Persistence modules. Let $(G, *)$ denote either $(\mathbb{N}^r, +)$ or $(\mathbb{R}_{\geq 0}^r, +)$. We consider G as a poset as follows. For $g_1, g_2 \in G$, $g_1 \leq g_2$ if this holds true component-wise. Let $\mathbb{K}[G]$ denote the monoid ring of G over the field \mathbb{K} with its natural G -grading. The category $\text{Mod}^{\text{gr}}(\mathbb{K}[G])$ of G -graded $\mathbb{K}[G]$ -modules is isomorphic to the thin category $\text{Fun}(G, \text{Vect}_{\mathbb{K}})$ of functors from the posetal category G to the category of \mathbb{K} -vector spaces $\text{Vect}_{\mathbb{K}}$ as follows. To each $M = \bigoplus_{g \in G} M_g \in \text{Ob}(\text{Mod}^{\text{gr}}(\mathbb{K}[G]))$, one associates the functor $(g \mapsto M_g) \in \text{Ob}(\text{Fun}(G, \text{Vect}_{\mathbb{K}}))$. Each graded morphism

$$\varphi = (\varphi_g)_{g \in G}: \bigoplus_{g \in G} M_g \rightarrow \bigoplus_{g \in G} N_g$$

naturally gives rise to a natural transformation $\{\varphi_g: M_g \rightarrow N_g\}_{g \in G}$ of the corresponding functors. The objects of those categories are called *(multi-)persistence modules*. We are going to switch seamlessly between both points of view. We denote by $\text{Tame}(G, \text{Vect}_{\mathbb{K}})$ the subcategory of $\text{Fun}(G, \text{Vect}_{\mathbb{K}})$ which corresponds to the subcategory $\text{Mod}_{\text{f.p.}}^{\text{gr}}(\mathbb{K}[G])$ of $\text{Mod}^{\text{gr}}(\mathbb{K}[G])$ consisting of finitely presented such modules, as investigated in [13]. In this article, we often consider *interval modules* which are defined as those elements of $\text{Fun}(G, \text{Vect}_{\mathbb{K}})$ that correspond to quotients of two monomial ideals. A subset $I \subseteq G$ is called *interval* in the poset (G, \leq) if for any $g, h \in I$ and any $f \in G$ the following two conditions hold:

- (i) If $g \leq f \leq h$, then $f \in I$.
- (ii) There exist $m \in \mathbb{N}$ and $g_1, \dots, g_{2m} \in I$ such that $g \leq g_1 \geq g_2 \leq \dots \geq g_{2m} \leq h$.

A *free* persistence module is a module of the form $F = \bigoplus_{g \in G} \mathbb{K}(g, \bullet)^{\beta_0(g)}$, where $\beta_0(g) \in \mathbb{N}$ and $\mathbb{K}(g, \bullet) \in \text{Fun}(G, \text{Vect}_{\mathbb{K}})$ denotes the functor

$$\mathbb{K}(g, h) := \begin{cases} \mathbb{K} & \text{if } g \leq h, \\ 0 & \text{if } g \not\leq h, \end{cases} \quad \mathbb{K}(g, h \leq f) := \begin{cases} \text{id}_{\mathbb{K}} & \text{if } g \leq h \leq f, \\ 0 & \text{otherwise.} \end{cases}$$

The natural number $\beta_0(F) := \sum_{g \in G} \beta_0(g)$ is called the *rank* of the free module F .

Definition 2.1. Let $M \in \text{Fun}((G, \leq), \text{Vect}_{\mathbb{K}})$ be a persistence module. The *total zeroth multigraded Betti number* of M , denoted $\beta_0(M)$ is the smallest possible rank of a free persistence module F such that there exists a surjection from F onto M .

Regarding M as an \mathbf{R}^r -graded $\mathbb{K}[\mathbf{R}^r]$ -module, the number $\beta_0(M)$ is the minimal number of homogeneous generators of M .

Caveat 2.2. In TDA literature, one often refers to β_0 as “rank”. This is not consistent with standard terminology in algebra. Yet, they do coincide for free modules.

2.2. Distances via persistence contours. It is crucial to be able to measure distances between persistence modules. In order to *learn* metrics in the sense of machine learning, one needs a large space of such metrics parameterized by simply describable objects. In this section, we recall a method to construct metrics on the category of tame functors, namely in terms of so-called *persistence contours*. Denote by \mathbf{R} the poset $\mathbb{R}_{\geq 0}$ and by \mathbf{R}_{∞}^r , where $r \in \mathbb{N}$, the poset obtained from \mathbf{R}^r by adding a single element ∞ such that $x \leq \infty$ for all $x \in \mathbf{R}^r$ and setting $x + \infty := \infty$ for all $x \in \mathbf{R}_{\infty}^r$.

Definition 2.3. A *(multi-)persistence contour* C is a functor $C: \mathbf{R}_{\infty}^r \times \mathbf{R} \rightarrow \mathbf{R}_{\infty}^r$ such that for any $x \in \mathbf{R}_{\infty}^r$ and $\varepsilon, \tau \in \mathbf{R}$, the following two conditions hold:

- (i) $x \leq C(x, \varepsilon)$ and
- (ii) $C(C(x, \varepsilon), \tau) \leq C(x, \varepsilon + \tau)$.

The second property in the definition indicates that a persistence contour fulfills a *lax* action only, i.e., the inequality is in general not an equality. For brevity, one often refers to persistence contours as just “contours”. One common example is the *standard contour in the direction of a vector* $v \in \mathbf{R}^r$ defined as $C(x, \varepsilon) := x + \varepsilon v$; i.e., one shifts x by the vector v scaled by ε . An immediate generalization of that contour is to travel along a curve instead of a line as explained in Example 2.9.

Following an immediate generalization of [10, Section 3], we now recall how persistence contours give rise to pseudometrics on the category of persistence modules.

Definition 2.4. Let C be a persistence contour, $M, N \in \text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}})$ tame persistence modules, and $\varepsilon \in [0, \infty)$. A morphism $f: M \rightarrow N$ is an ε -*equivalence* if for every $a \in [0, \infty)^r$ s.t. $C(a, \varepsilon) \neq \infty$, there is a \mathbb{K} -linear function $N(a) \rightarrow M(C(a, \varepsilon))$ making the following diagram commutative:

$$\begin{array}{ccc}
 M(a) & \xrightarrow{M(a \leq C(a, \varepsilon))} & M(C(a, \varepsilon)) \\
 \downarrow f(a) & \searrow \exists & \downarrow f(C(a, \varepsilon)) \\
 N(a) & \xrightarrow{N(a \leq C(a, \varepsilon))} & N(C(a, \varepsilon))
 \end{array}$$

Definition 2.5. Two tame functors M and N are called ε -*equivalent* (with respect to C) if there is a tame functor L and morphisms $f: M \rightarrow L \leftarrow N: g$ such that f is an ε_1 -equivalence, g is an ε_2 -equivalence, and $\varepsilon_1 + \varepsilon_2 \leq \varepsilon$.

Denote by $\mathcal{E} := \{\varepsilon \in [0, \infty) \mid M \text{ and } N \text{ are } \varepsilon\text{-equivalent}\}$. Then the function

$$d_C(M, N) := \begin{cases} \infty & \text{if } \mathcal{E} = \emptyset, \\ \inf(\mathcal{E}) & \text{otherwise} \end{cases}$$

is an extended pseudometric on $\text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}})$. There is an alternative way of thinking of contours. For that, we now recall the definition of a noise system from [27].

Definition 2.6. A *noise system* in $\text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}})$ is a collection $\{\mathcal{V}_\varepsilon\}_{\varepsilon \in \mathbf{R}}$ of sets of tame functors such that the following three conditions are satisfied:

- (i) For any ε , the zero functor is contained in \mathcal{V}_ε ,
- (ii) for all $0 \leq \tau < \varepsilon$, \mathcal{V}_τ is contained in \mathcal{V}_ε , and
- (iii) if $0 \rightarrow M \rightarrow L \rightarrow N \rightarrow 0$ is a short exact sequence in $\text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}})$, then
 - (1) if $L \in \mathcal{V}_\varepsilon$, then also $M, N \in \mathcal{V}_\varepsilon$ and
 - (2) if $M \in \mathcal{V}_\varepsilon$ and $N \in \mathcal{V}_\tau$, then $L \in \mathcal{V}_{\varepsilon+\tau}$.

Throughout this article, we will mainly focus on noise systems $\{\mathcal{V}_\varepsilon\}$ that are *closed under direct sums*, i.e., noise systems such that whenever $M, N \in \mathcal{V}_\varepsilon$, also their direct sum $M \oplus N$ is in \mathcal{V}_ε . Define $B(M, \tau)$ as

$$B(M, \tau) := \{U \mid U \text{ is a tame subfunctor of } M \text{ such that } M/U \in \mathcal{V}_\tau\}.$$

If $\{\mathcal{V}_\varepsilon\}$ is closed under direct sums, a minimal element in $B(M, \tau)$ is unique if it exists; this element is denoted by $M[\tau]$ and is called the τ -*shift* of M . Then $B(M, \tau)$ coincides with the set

$$\{U \mid U \text{ is a tame subfunctor of } M \text{ such that } M[\tau] \subseteq U \subseteq M\}.$$

Definition 2.7 ([17, Definition 8.2]). A noise system is called *simple* if

- (i) it is closed under direct sums,
- (ii) for any tame functor M and any $\tau \in \mathbf{R}$, $B(M, \tau)$ contains the minimal element $M[\tau]$, and
- (iii) $\beta_0(M[\tau]) \leq \beta_0(M)$ for any $\tau \in \mathbf{R}$.

The third condition inherently links simple noise systems to β_0 .

Proposition 2.8 ([17, Theorem 9.6]). For a persistence contour C , denote by $\mathcal{V}_{C, \varepsilon}$ the set of tame functors M such that $M(x \leq C(x, \varepsilon))$ is the zero-morphism whenever $C(x, \varepsilon) \neq \infty$. The function $C \mapsto \{\mathcal{V}_{C, \varepsilon}\}_\varepsilon$ is a bijection between the set of persistence contours and the set of simple noise systems.

Under this correspondence, for the standard contour in the direction of a vector v , $M[\tau \cdot v]$ is $(\tau \cdot v) * M$, where $*$ denotes the induced action of the monoid \mathbf{R}^r on the monoid ring $\mathbb{K}[\mathbf{R}^r]$.

2.3. Construction of new contours. In the setting of a single parameter, there are various contours suitable for concrete data-analytic tasks [10]. In this subsection, we construct several contours for the multiparameter case.

Example 2.9 (Curve contour). Consider a curve in \mathbb{R}^r that is given by a monotone non-decreasing function $I: \mathbf{R} \rightarrow \mathbb{R}^r$ for which $I(0)$ lies in one of the coordinate axes. Assume further that \mathbf{R}^r can be covered by translations of that curve parallel to $r-1$ coordinate axes such that every point in \mathbf{R}^r lies on precisely one of the curves. For $x \in \mathbf{R}_\infty^r$ and $\varepsilon \in [0, \infty)$, define $C(x, \varepsilon)$ to be the point that one obtains by traveling along the translated curve that runs ε -far through x ; i.e., if I_x denotes the translated curve running through x at ε_0 , we set $C(x, \varepsilon)$ to be $I_x(\varepsilon_0 + \varepsilon)$. \diamond

For some further examples of persistence contours in the one-parameter case, we refer to [10]. Among them is the contour of *distance type*. Fixing a vector, we can immediately generalize this contour to the multiparameter setting.

Example 2.10 (Distance type in a fixed direction). Let $r \in \mathbb{N}$, $x, v \in \mathbf{R}^r$, and f a non-negative (Lebesgue-)measurable function on \mathbf{R}^r . For $x \in \mathbf{R}_\infty^r$ and $\varepsilon \in [0, \infty)$, we

set $C(x, \varepsilon) := x + \delta(x, \varepsilon)v$ where $\delta(x, \varepsilon) \in \mathbb{R}_{\geq 0}$ is such that

$$\varepsilon = \int_{L_x^{x+\delta(x, \varepsilon)v}} f \, d\mathcal{L}^r.$$

If no such $\delta(x, \varepsilon)$ exists, we set $C(x, \varepsilon) := \infty$. Here, $L_x^{x+\delta(x, \varepsilon)v}$ denotes the set

$$L_x^{x+tv} := \{y \in \mathbf{R}^r \mid y \geq x, y \not\geq x + \delta(x, \varepsilon)v\},$$

to which we refer to as *L-shape (at x)*. \diamond

Remark 2.11. Note that the L-shapes in Example 2.10 can not be replaced by the rectangles of the same bounds; it would not define a lax action anymore. \diamond

Applying a one-parameter contour component-wise, as in the following example, yields a multiparameter contour.

Example 2.12 (Component-wise shift type). Let $r \in \mathbb{N}$, $f_1, \dots, f_r: [0, \infty) \rightarrow [0, \infty)$ be non-negative measurable functions and $x \in \mathbf{R}^r$. For each x_i , set y_i to be the infimum of all $\tilde{y}_i \in \mathbf{R}$ such that $x_i = \int_0^{\tilde{y}_i} f_i \, d\mathcal{L}^1$, if such \tilde{y}_i exists. Let m_1, \dots, m_r be super-additive non-negative functions. One obtains a contour by setting

$$C(x, \varepsilon) := \left(\begin{array}{c} \int_0^{y_1+m_1(\varepsilon)} f_1 \, d\mathcal{L}^1 \\ \vdots \\ \int_0^{y_r+m_r(\varepsilon)} f_r \, d\mathcal{L}^1 \end{array} \right)$$

if such y_1, \dots, y_r exist, and $C(x, \varepsilon) := \infty$ otherwise. \diamond

We now introduce a generalization of the contour of shift type that is based on multivariate functions. This yields a large class of persistence contours.

Example 2.13 (Multivariate shift type). Let $f_1, \dots, f_r: [0, \infty)^r \rightarrow [0, \infty)$ be non-negative measurable functions. For $y \in \mathbf{R}^r$, let $\text{Rec}(y)$ denote $[0, y_1] \times \dots \times [0, y_r] \subseteq \mathbf{R}^r$. For $x \in [0, \infty)^r$ define

$$B(x) := \left\{ y \mid \int_{\text{Rec}(y)} f_1 \, d\mathcal{L}^r \geq x_1, \dots, \int_{\text{Rec}(y)} f_r \, d\mathcal{L}^r \geq x_r \right\}$$

and $b(x) := \inf B(x)$, where \inf denotes the meet in the poset. Let $v \in \mathbf{R}^r$. One obtains a contour by setting

$$C(x, \varepsilon) := \sup \left(\left(\int_{\text{Rec}(b(x)+\varepsilon v)} f_1 \, d\mathcal{L}^r, \dots, \int_{\text{Rec}(b(x)+\varepsilon v)} f_r \, d\mathcal{L}^r \right), x \right)$$

where \sup denotes the join of the poset. If $B(x)$ is empty, set $C(x, \varepsilon) := \infty$. To verify that C is a contour, one can observe that, for every x and ε , the element $b(x) + \varepsilon v$ belongs to $B(C(x, \varepsilon))$. Consequently, $b(C(x, \varepsilon)) \leq b(x) + \varepsilon$, and hence

$$\int_{\text{Rec}(C(x, \varepsilon))+\tau v} f_i \, d\mathcal{L}^r \leq \int_{\text{Rec}(b(x)+\varepsilon v+\tau v)} f_i \, d\mathcal{L}^r$$

for every i . This inequality implies that condition (ii) of Definition 2.3 is fulfilled.

The computational bottleneck is to compute $b(x)$. This, however, can be done efficiently by an approximation scheme to arbitrary precision by studying the intersection of $B(x)$ with an equidistant grid and successively decreasing the grid length. \diamond

2.4. Hierarchical stabilization. In this section, we recall from [10, 17] the concept and properties of the *hierarchical stabilization*. This process turns discrete into stable invariants as follows. Let T be a set and $d: T \times T \rightarrow \mathbf{R} \cup \infty$ an extended pseudometric on T , where again \mathbf{R} denotes the poset of non-negative real numbers. Let $f: T \rightarrow \mathbb{N}$ be a discrete invariant and denote by \mathcal{M} the set of Lebesgue-measurable functions from $[0, \infty)$ to $[0, \infty)$, endowed with the interleaving distance as in [17, Section 3].

Definition 2.14. Fix an extended pseudometric d on T . The *hierarchical stabilization* of f , denoted \hat{f} , is defined to be

$$\hat{f}: T \rightarrow \mathcal{M}, \quad x \mapsto (\tau \mapsto \min \{f(y) \mid d(x, y) \leq \tau\}).$$

Hence, $\hat{f}(x) \in \mathcal{M}$ is the monotone, non-increasing measurable function that maps a real number τ to the minimal value that the discrete invariant f takes on a closed τ -neighborhood of x . Note that \hat{f} strongly depends on the chosen pseudometric on T .

Proposition 2.15. [10, Proposition 2.2] *For any choice of extended pseudometric on T , the function $\hat{f}: T \rightarrow \mathcal{M}$ is 1-Lipschitz.*

Therefore, the hierarchical stabilization turns discrete into stable invariants.

3. THE SHIFT-DIMENSION OF MULTIPERSISTENCE MODULES

In this section, we introduce the shift-dimension of persistence modules and investigate its algebraic properties. We particularly focus on (non-)additivity.

3.1. The shift-dimension and its origin.

Definition 3.1. Let $v \in \mathbf{R}^r$, M an \mathbf{R}^r -graded $\mathbb{K}[\mathbf{R}^r]$ -module. Elements m_1, \dots, m_k of M *v-generate* M if

$$v * (M / \langle m_1, \dots, m_k \rangle) = 0.$$

A *v-basis* of M is a set of *v-generators* m_1, \dots, m_k such that k is smallest possible. In this case, we call k the *shift-dimension* (or *v-dimension*) of M and denote it $\dim_v(M)$.

Throughout the article, we are going to stick to *homogeneous* elements when studying generators and *v-bases*. The following lemma justifies that convention.

Lemma 3.2. *If M is a tame persistence module, then for all $v \in \mathbf{R}^r$ there exists a *v-basis* consisting of finitely many homogeneous generators.*

Proof. This statement immediately follows from the fact that every finitely presented \mathbf{R}^r -graded $\mathbb{K}[\mathbf{R}^r]$ -module can be minimally generated by homogeneous elements. \square

The shift-dimension can be equally characterized as the smallest number of elements of M such that $v * M$ is contained in the submodule generated by those elements.

Remark 3.3. The 0-dimension of M is the minimal number of generators of M . For a fixed vector v , one may consider the family $\{\dim_{\tau v}(M)\}_{\tau \in \mathbb{R}_{\geq 0}}$ of shift-dimensions of a multipersistence module M . The non-increasing function $\mathbb{R}_{\geq 0} \rightarrow \mathbb{N}$ mapping $\tau \mapsto \dim_{\tau v}(M)$ is suitable as a feature map for machine learning algorithms. \diamond

In general, \mathbf{R}^r -graded $\mathbb{K}[\mathbf{R}^r]$ -modules are hard to handle. We will mostly restrict to the category of *finitely presented* \mathbf{R}^r -graded $\mathbb{K}[\mathbf{R}^r]$ -modules and thereby to the category $\text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}})$ of *tame* functors.

[illegible]

At first glance, the notion of the shift-dimension might seem rather artificial. But, in fact, it arises in a natural way; namely as the hierarchical stabilization of β_0 .

$$\widehat{\beta_{0v}}(M)(\delta) = \dim_{\delta v}(M).$$
$$\mathcal{V}_\delta := \{N \in \text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}}) \mid \forall_{u \in \mathbf{R}^r: C(u, \delta) < \infty} \ker(N(u \leq u + \delta \cdot v)) = N(u)\}$$
$$\widehat{\beta_{0,n}}(M)(\delta) = \min \{ \beta_0(U) \mid U \in B(M, \delta) \}.$$
$$B(M, \delta) = \{U \in \text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}}) \mid U \subseteq M, \delta v * (M/U) = 0\},$$

Note that $\hat{\beta}_{0,n}(M)$ is a monotone, non-increasing step function.

Remark 3.6. [17, Theorem 8.3] is closely related to [27, Proposition 11.1]. The definition we use implies uniqueness properties of the shift, cf. [17, Proposition 8.1]. Furthermore, in [17, 27], noise systems are described for the non-negative *rational* numbers instead of the non-negative real numbers. The corresponding arguments we use generalize to $\text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}})$. \diamond

Remark 3.7. As shown in [17, Section 11], computing the shift-dimension can be reduced to an NP-complete problem and hence is NP-hard itself: for a certain class of persistence modules, the computation of the shift-dimension is at least as hard as the RANK-3 problem [26], an optimization problem in linear algebra. \diamond

We obtain a truncated version of the shift-dimension as stabilization of the zeroth multigraded Betti number by changing the contour to be truncated. For C a contour and $\alpha \in \mathbf{R}^r$, denote by C_α the *truncation* of C at α . This is defined as

$$C_\alpha(x, \varepsilon) := \begin{cases} C(x, \varepsilon) & \text{if } \alpha \not\leq C(x, \varepsilon), \\ \infty & \text{otherwise.} \end{cases}$$

We call the corresponding noise system the α -truncated noise system.

Proposition 3.8. *Under the assumptions of Proposition 3.5 with α -truncated noise in the direction of a normalized vector v , β_0 stabilizes as follows with respect to the α -truncated standard contour in the direction of v :*

$$\widehat{\beta_{0v,\alpha}}(M)(\delta) = \dim_{\delta v}(M / \oplus_{\beta \geq \alpha} M_\beta).$$

Proof. The noise system associated to the truncated standard contour is given by

$$\mathcal{V}_{\delta,\alpha} := \{N \in \text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}}) \mid \forall u \in \mathbf{R}^r: C_\alpha(u, \delta) < \infty \implies \ker(N(u \leq u + \delta v)) = N(u)\}.$$

We denote the balls corresponding to this noise system by B_α . Applying [17, Theorem 8.3], we get

$$\widehat{\beta_{0v,\alpha}}(M)(\delta) = \min \{\beta_0(U) \mid U \in B_\alpha(M, \delta)\}.$$

We have $M/U \in \mathcal{V}_{\delta,\alpha}$ iff for all homogeneous $m \in M/U$ we have $\delta v * m = 0$ or $\alpha \leq \deg(\delta v * m)$. Hence,

$$B_\alpha(M, \delta) = \{U \in \text{Tame}(\mathbf{R}^r, \text{Vect}_{\mathbb{K}}) \mid U \subseteq M, \delta v * ((M / \oplus_{\beta \geq \alpha} M_\beta) / [U]) = 0\},$$

and thus the claim of the proposition follows. \square

Therefore, computing the stabilized β_0 with respect to a truncated contour corresponds to truncating the module at degree α .

3.2. Properties of the shift-dimension. We now state some fundamental properties of the shift-dimension. In particular, we investigate it regarding additivity.

On the level of epimorphisms, the following holds true.

Lemma 3.9. *Let $\varphi: M \twoheadrightarrow N$ be an epimorphism of tame persistence modules. Then the following holds true: $\dim_v(N) \leq \dim_v(M)$ for all $v \in \mathbf{R}^r$.*

Proof. Let $\{m_1, \dots, m_{\dim_v(M)}\}$ be a v -basis of M . Then $\varphi(m_1), \dots, \varphi(m_{\dim_v(M)})$ v -generate N and hence $\dim_v(N) \leq \dim_v(M)$. \square

For monomorphisms, we do not get a corresponding inequality in the reverse direction, as the following counterexample demonstrates.

Example 3.10. Let $v = (1, 1)$, M be the interval module generated in degrees $(0, 2)$ and $(2, 0)$, and N the free module generated by a single element in degree $(0, 0)$. Then $M \hookrightarrow N$ and $\dim_v(M) = 2 \not\leq 1 = \dim_v(N)$. \diamond

Lemma 3.11. *Let $v \in \mathbf{R}^r$ and $m \in M$. Then $\dim_v(M/\langle m \rangle)$ is contained in $\{\dim_v(M), \dim_v(M) - 1\}$, i.e., taking the quotient by a submodule that is generated by a single element can drop the shift-dimension by one at most.*

Proof. Assume there exist $m_1, \dots, m_{\dim_v(M)-2} \in M/\langle m \rangle$ that are a v -basis of $M/\langle m \rangle$. Then for any choice of representatives \tilde{m}_i of m_i , $m, \tilde{m}_1, \dots, \tilde{m}_{\dim_v(M)-2}$ v -generate M . It follows that $\dim_v(M) \leq \dim_v(M) - 1$, which is a contradiction. \square

In order to decide whether an element $m \in M$ can be extended to a v -basis $\{m, m_2, \dots, m_{\dim_v(M)}\}$ of M , one makes use of the following criterion.

Lemma 3.12. *Let $v \in \mathbf{R}^r$ and $m \in M$. There exists a v -basis of M containing m if and only if $\dim_v(M/\langle m \rangle) = \dim_v(M) - 1$.*

Proof. Let $\{m, m_2, \dots, m_{\dim_v(M)}\}$ be a v -basis of M . Then $[m_2], \dots, [m_{\dim_v(M)}]$ v -generate $M/\langle m \rangle$. Hence $\dim_v(M/\langle m \rangle) \leq \dim_v(M) - 1$ and therefore $\dim_v(M/\langle m \rangle) = \dim_v(M) - 1$ by Lemma 3.11. This proves the implication from left to right. Now let $\{m_1, \dots, m_{\dim_v(M)-1}\}$ be a v -basis of $M/\langle m \rangle$ and choose representatives $\tilde{m}_1, \dots, \tilde{m}_{\dim_v(M)-1} \in M$ for which $[\tilde{m}_i] = m_i$, $i = 1, \dots, \dim_v(M) - 1$. Then $m, \tilde{m}_1, \dots, \tilde{m}_{\dim_v(M)-1}$ v -generate M and hence already are a v -basis of M . \square

For short exact sequences, we prove the following two inequalities.

Theorem 3.13. *Let $0 \rightarrow M \xrightarrow{\varphi} L \xrightarrow{\psi} N \rightarrow 0$ be a short exact sequence of persistence modules. Then for all $v, w \in \mathbf{R}^r$, the following two inequalities hold:*

- (i) $\dim_{v+w}(L) \leq \dim_v(M) + \dim_w(N)$, and
- (ii) $\dim_v(L) \leq \dim_v(N) + \beta_0(M)$.

Proof. Choose a w -basis $\{n_1, \dots, n_a\}$ of N . For every i , choose $l_i \in L$ for which $\psi(l_i) = n_i$. Let $\{g_1, \dots, g_{\beta_0(L)}\}$ generate L . Then for all i there exist $\alpha_1, \dots, \alpha_a \in \mathbb{K}[\mathbf{R}^r]$ such that $w * g_i - \sum_{j=1}^a \alpha_j l_j \in \ker(\psi) = \text{im}(\varphi)$. Choose a v -basis $\{m_1, \dots, m_b\}$ of M . Thus, for all i , there exist $\gamma_1, \dots, \gamma_b \in \mathbb{K}[\mathbf{R}^r]$ such that $v * (w * g_i - \sum_{j=1}^a \alpha_j l_j) = \varphi(\sum_{k=1}^b \gamma_k m_k)$, which is equivalent to $(v + w)g_i = \sum_{j=1}^a (v * \alpha_j) l_j + \sum_{k=1}^b \gamma_k \varphi(m_k)$. Consequently $\dim_{v+w}(L) \leq b + a = \dim_v(M) + \dim_w(N)$, concluding the proof of (i). For the proof of statement (ii), let F be the free module generated by a v -basis of N . Since $\psi: L \twoheadrightarrow N$ is surjective, we obtain a morphism $F \rightarrow L$ and hence a morphism from $M \oplus F$ to L . Since $L/(M \oplus F) \cong N/F$, it follows that $\dim_v(L/(M \oplus F)) = 0$. Therefore, $\dim_v(L) \leq \beta_0(M \oplus F) = \beta_0(M) + \beta_0(F) = \beta_0(M) + \dim_v(N)$. \square

If $\dim_v(M) = \beta_0(M)$ or $\beta_0(M) = 1$, it follows that $\dim_v(L) \leq \dim_v(M) + \dim_v(N)$.

The shift-dimension is sub-additive, i.e., $\dim_v(M \oplus N) \leq \dim_v(M) + \dim_v(N)$. In general, it is not additive, as the following counterexample for L-shapes demonstrates.

Example 3.14 ($r = 2$). Consider the interval modules $M = \mathbb{K}((1, 0), \bullet) / \mathbb{K}((1, 2), \bullet)$ and $N = \mathbb{K}((0, 1), \bullet) / \mathbb{K}((2, 1), \bullet)$. Then $(1, 1) * (M \oplus N)$ is contained in the submodule of $M \oplus N$ that is generated by the element $(1, 1) \in (M \oplus N)_{(1,1)}$ in degree $(1, 1)$. It follows that $\dim_{(1,1)}(M \oplus N) = 1 \neq 1 + 1 = \dim_{(1,1)}(M) + \dim_{(1,1)}(N)$. \diamond

We now present some cases for which additivity of the shift-dimension does hold true.

Lemma 3.15. *If F and G are two free persistence modules of finite rank, then $\dim_v(F \oplus G) = \dim_v(F) + \dim_v(G)$ for all $v \in \mathbf{R}^r$.*

Proof. The statement follows from the fact that in sufficiently large degree, the elements of the v -basis need to generate $\mathbb{K}^{\beta_0(F) + \beta_0(G)}$. \square

Proposition 3.16. *Let M be an interval module generated by one element and F a free persistence module of rank one. Then $\dim_v(M \oplus F) = \dim_v(M) + 1$ for all $v \in \mathbf{R}^r$.*

Proof. For $v \in \mathbf{R}^r$ s.t. $\dim_v(M) = 0$, the statement is clear. Hence let $v \in \mathbf{R}^r$ be such that $\dim_v(M) = 1$. Then $\dim_v(M \oplus F) \leq \dim_v(M) + \dim_v(F) = 1 + 1 = 2$ and $\dim_v(M \oplus F) \geq \max\{\dim_v(M), \dim_v(F)\} = 1$. Assume $U = \langle (m, f) \rangle$ for some $(m, f) \in M \oplus F$. Then $v * ((M \oplus F)/U) \not\cong 0$, as we argue now. If $f = 0$, then for all $g \in F \setminus \{0\}$, the element $v * (0, g) \notin U$. If $f \neq 0$, then there exists $m \in M \setminus \{0\}$ such that $v * (m, 0) \notin U$, since by assumption there exists $m \in M$ for which $v * m \neq 0$. Hence $\dim_v(M \oplus F) > 1$, proving the claim. \square

Proposition 3.17. *Let $\dim_v(M) \leq 1$ and F a free persistence module of rank k . Then $\dim_v(M \oplus F) = \dim_v(M) + \dim_v(F)$.*

Proof. For $\dim_v(M) = 0$, the statement is clear. Now assume that $\dim_v(M) = 1$. Assume there exists a v -basis $\{(m_1, f_1), \dots, (m_k, f_k)\}$ of M of cardinality k . Since $\dim_v(F) = k$ and F is free, f_1, \dots, f_k need to be linearly independent. By assumption on the shift-dimension of M , there exists $m \in M$ s.t. $v * m \neq 0$. Because of the linear independence of f_1, \dots, f_k , the element $(v * m, 0)$ is not contained in the submodule generated by $(m_1, f_1), \dots, (m_k, f_k)$, in contradiction to the assumption. \square

3.3. A quantitative study of non-additivity. In this subsection, we investigate non-additivity of the shift-dimension in greater detail and give a measure for it. For a finite family of persistence modules $\{M_i\}_{i \in I}$ and a fixed, possibly learned $v \in \mathbf{R}^r$, we define the *locus of non-additivity* of the shift-dimension as

$$\text{Loc}_v(\{M_i\}_{i \in I}) := \left\{ \tau \in \mathbf{R} \mid \dim_{\tau v}(\oplus_{i \in I} M_i) \neq \sum_{i \in I} \dim_{\tau v}(M_i) \right\}.$$

For a quantitative study, we associate to $\{M_i\}_{i \in I}$ the L^p -distance between the functions $\sum_{i \in I} \dim_{\bullet}(M_i)$ and $\dim_{\bullet}(\oplus_{i \in I} M_i)$ of τ , i.e., for $1 \leq p < \infty$, we study

$$\text{err}_{v,p}(\{M_i\}_{i \in I}) := \left(\int_0^\infty \left(\left(\sum_{i \in I} \dim_{\tau v}(M_i) \right) - \dim_{\tau v} \left(\oplus_{i \in I} M_i \right) \right)^p d\tau \right)^{1/p}.$$

If this expression is sufficiently small, the sum of the shift-dimensions of the M_i yields a good approximation of the shift-dimension of their direct sum $\oplus M_i$. Revisiting and generalizing Example 3.10, we now undertake more quantitative investigations.

Example 3.18. Let $v \in \mathbb{R}_{>0}^r$. Let $M_1, M_2 \neq 0$ be interval modules generated by single elements \mathbf{g}_1 and \mathbf{g}_2 , resp., and quotiented out by single elements $\tilde{\mathbf{g}}_1$ and $\tilde{\mathbf{g}}_2$, resp. Assume further that $\text{lcm}(\deg(\mathbf{g}_1), \deg(\mathbf{g}_2)) < \deg(\tilde{\mathbf{g}}_2) \leq \deg(v * \mathbf{g}_1)$ and $\text{lcm}(\deg(\mathbf{g}_1), \deg(\mathbf{g}_2)) < \deg(\tilde{\mathbf{g}}_1) \leq \deg(v * \mathbf{g}_2)$ as well as $\deg(\tilde{\mathbf{g}}_1) \not\leq \deg(v * \mathbf{g}_1)$ and $\deg(\tilde{\mathbf{g}}_2) \not\leq \deg(v * \mathbf{g}_2)$. Then, $\dim_v(M_1 \oplus M_2) = 1 \neq 2 = \dim_v(M_1) + \dim_v(M_2)$, since the element $(1, 1) \in (M_1 \oplus M_2)_{\text{lcm}(\deg(\mathbf{g}_1), \deg(\mathbf{g}_2))}$ divides both $v * (\mathbf{g}_1, 0)$ and $v * (0, \mathbf{g}_2)$. Now, for all $\tau \in \text{Loc}_v(\{M_i\}_{i \in I})$ we get $\dim_{\tau v}(M_1 \oplus M_2) = 1 \neq 2 = \dim_{\tau v}(M_1) + \dim_{\tau v}(M_2)$. The length of the interval $\text{Loc}_v(\{M_i\}_{i \in I})$ yields $\text{err}_{v,1}(\{M_1, M_2\})$. \diamond

Let M_1, M_2 be interval modules as in the example above. In this case, the union of their underlying intervals is an interval. Denote by M the corresponding interval module. Then, whenever $\dim_{\tau v}(M_1 \oplus M_2) = 1$, we have $\dim_{\tau v}(M_1 \oplus M_2) = \dim_{\tau v}(M)$.

The following example demonstrates that the difference between $\dim_v(\oplus_{i \in I} M_i)$ and $\sum_{i \in I} \dim_v(M_i)$ can be arbitrarily large. In fact, the shift-dimension of a module can be 1 while the Hilbert function and the rank invariant attain arbitrarily large values.




FIGURE 1. Illustration of a persistence module as in Example 3.19 for $|I| = 2$. *Left:* Let $v = (4, 4)$. All shifted minimal generators have pairwise incomparable degrees but can be generated by the all-one vector in the degree of the greatest common divisor of their degrees. Hence, $\dim_v(\oplus_i(M_i)) = 1$. *Right:* The functions $\sum_i \dim_{\tau v}(M_i)$ (in lavender) and $\dim_{\tau v}(\oplus_i(M_i))$ (in blue) for $v = (1, 1)$. We have $\text{Loc}_v(\{M_i\}_{i \in I}) = [3, 4.3]$ and $\text{err}_{v,p} = (0.6 \cdot 2^p + 0.7 \cdot 3^p)^{1/p}$.

Example 3.19. Let M_1 and M_2 be as in Example 3.18. Define a finite collection of interval modules $\{M_i\}_{i \in I}$ such that each M_i has one generator \mathbf{g}_i and gets quotiented out by two elements $\tilde{\mathbf{g}}_i, \hat{\mathbf{g}}_i$ that are of the following form. The $\{\mathbf{g}_i\}_{i \in I}$ are assumed to have pairwise distinct degrees which lie on the straight line between $\deg(\mathbf{g}_1)$ and $\deg(\mathbf{g}_2)$. The degrees of $\tilde{\mathbf{g}}_i$ and $\hat{\mathbf{g}}_i$ are chosen such that $\deg(\tilde{\mathbf{g}}_i), \deg(\hat{\mathbf{g}}_i) \not\leq \deg(v * \mathbf{g}_i)$ and for all $j \neq i$ either $\deg(\tilde{\mathbf{g}}_i) \leq \deg(v * \mathbf{g}_j)$ or $\deg(\hat{\mathbf{g}}_i) \leq \deg(v * \mathbf{g}_j)$. Then, again, the all-one vector in degree $\text{lcm}(\deg(\mathbf{g}_1), \deg(\mathbf{g}_2))$ is a divisor of each $v * (0, \dots, \mathbf{g}_i, \dots, 0)$. Hence, $\dim_v(\oplus_{\{1,2\} \cup I} M_i) = 1 \neq 2 + |I| = \sum_{\{1,2\} \cup I} \dim_v(M_i)$. See Figure 1 for an illustration of one concrete example. \diamond

Note that $\text{Loc}_v(\{M_i\}_{i \in I})$ is in general not connected. Counterexamples can be constructed by taking direct sums of certain modules as in the examples above.

We give a further, more involved example for which additivity does not hold true.

Example 3.20 (Example 3.4 revisited). Let M_1 be the persistence module from Example 3.4 and M_2 be an interval module generated at degrees $(0, 5)$ and $(5, 1.5)$ and quotiented out at degrees $(2, 6)$ and $(9, 1.5)$. Let $v = (2, 1)$. Then,

$$\dim_{\tau v}(M_1) = \begin{cases} 5 & \text{if } 0 \leq \tau < 1, \\ 2 & \text{if } 1 \leq \tau < 2, \\ 0 & \text{if } 2 \leq \tau, \end{cases} \quad \dim_{\tau v}(M_2) = \begin{cases} 2 & \text{if } 0 \leq \tau < 1, \\ 1 & \text{if } 1 \leq \tau < 2, \\ 0 & \text{if } 2 \leq \tau, \end{cases}$$

and

$$\dim_{\tau v}(M_1 \oplus M_2) = \begin{cases} 7 & \text{if } 0 \leq \tau < 1, \\ 3 & \text{if } 1 \leq \tau < 1.5, \\ 2 & \text{if } 1.5 \leq \tau < 2, \\ 0 & \text{if } 2 \leq \tau. \end{cases}$$

We get $\text{Loc}_v(\{M_1, M_2\}) = [1.5, 2)$ and $\text{err}_{v,p}(\{M_1, M_2\}) = 0.5^{1/p}$. \diamond

Here, the non-triviality of the loci and L^p -errors of non-additivity is due to the existence of v -basis elements such that a suitable multiple of them yields a non-zero shifted generator in one direct summand and zero on the other direct summands.




FIGURE 2. In gray: the interval module M of Example 3.21. In lighter gray: $(4, 4) * M$. *Left:* In blue and lavender: the submodules generated by the bases \mathcal{B}_1 and \mathcal{B}_2 , resp. *Right:* Regions in which the basis elements can be exchanged, fixing $(8, 4)$ and $(4, 6)$, resp., following Lemma 3.22. Using this lemma again, both bases can be exchanged with $\mathcal{B}_3 := \{(4, 6), (6, 4)\}$. Those replacements do not preserve the indispensability relations, since $(4, 6)$ is indispensable only for $(0, 8)$ and $(6, 4)$ is indispensable only for $(11, 0)$.

3.4. Basis exchange properties. Unlike bases of vector spaces in linear algebra, v -bases do *not* give rise to a matroid if $r > 1$. The following example demonstrates that even for monomial ideals, neither the well-desired basis exchange property holds, nor that for every set of v -generators there exists a subset which is a v -basis of M .

Example 3.21. Let M be the interval module generated at degrees $(0, 8)$, $(4, 6)$, $(6, 4)$, $(8, 2)$, and $(11, 0)$. For $v = (4, 4)$, $\dim_v(M) = 2$. Both $\mathcal{B}_1 := \{(3.5, 11.5), (8, 4)\}$ and $\mathcal{B}_2 := \{(4, 12), (8, 4)\}$ are v -bases of M . The basis elements cannot be exchanged, as $(3.5, 11.5), (4, 6) \not\leq (4, 4) + (11, 0)$ and $(4, 12), (13, 4) \not\leq (4, 4) + (8, 2)$ (cf. Figure 2). Taking the shifts of all 5 minimal generators by v yields a set of v -generators of M . However, none of its proper subsets is a set of v -generators of M . \diamond

We now investigate how to modify elements of a given v -basis by multiplication with suitable elements of the monoid. Let $m \in M$ and \mathcal{B} be a v -basis of M . If for an element b of \mathcal{B} we have that $v * m \notin \langle \mathcal{B} \setminus \{b\} \rangle$, we call b *indispensable* for m . Otherwise, we call b *dispensable* for m .

Lemma 3.22. *Let \mathcal{B} be a v -basis of a persistence module M . Let $b \in \mathcal{B}$. Let $\{\mathbf{g}_i\}_{i \in I}$ be a minimal set of homogeneous generators of M . Then the following holds.*

- (i) *If there exists $\tilde{b} \in M$ and $r \in \mathbb{K}[\mathbf{R}^r]$ such that $r\tilde{b} = b$, then $(\mathcal{B} \setminus \{b\}) \cup \{\tilde{b}\}$ is a v -basis of M .*
- (ii) *Let $J \subseteq I$ be the set of all $j \in I$ s.t. b is indispensable for \mathbf{g}_j . Let $w \in \mathbf{R}^r$ s.t. $\deg(w * b) \leq \gcd(\{\deg(v * \mathbf{g}_j) \mid j \in J\})$. Then $(\mathcal{B} \setminus \{b\}) \cup \{w * b\}$ is a v -basis of M .*

Proof. Let \tilde{b} be as in the assumption of (i). Then $(\mathcal{B} \setminus \{b\}) \cup \{\tilde{b}\}$ is a set of v -generators as it generates $\langle \mathcal{B} \rangle$ by construction. Since it has the same cardinality as \mathcal{B} , it is indeed a v -basis. For the proof of the second statement, let w be as desired. We prove that for each $\mathbf{g} \in \{\mathbf{g}_i\}_{i \in I}$ we have $v * \mathbf{g} \in \langle (\mathcal{B} \setminus \{b\}) \cup \{w * b\} \rangle$. If b is dispensable for \mathbf{g} , then $v * \mathbf{g} \in \langle \mathcal{B} \setminus \{b\} \rangle \subseteq \langle (\mathcal{B} \setminus \{b\}) \cup \{w * b\} \rangle$. If b is indispensable for \mathbf{g} , then $v * \mathbf{g}$ can be expressed as linear combination of elements of $(\mathcal{B} \setminus \{b\}) \cup \{w * b\}$, since $\deg(w * b) \leq \deg(v * \mathbf{g})$. \square

Given $b \in \mathcal{B}$, there is—up to constants in the field \mathbb{K} —a unique maximal multiple of b by which one can replace it, but there might be several incomparable such choices for divisors of b . If one replaces a v -basis element by a multiple or by a divisor of it as described in the lemma above, it is important to note that the indispensability relations of the other v -basis elements with the minimal generators of M might change. Indispensability therefore encodes the combinatorial complexity of v -bases.

Corollary 3.23. *Let M be an interval module, \mathfrak{G} a minimal set of homogeneous generators of M , and \mathcal{B} a v -basis of M . Then the following statements are true.*

- (i) *There exists a v -basis consisting of minimal generators of M .*
- (ii) *If $b \in \mathcal{B}$ is indispensable for a $\mathfrak{g} \in \mathfrak{G}$, then $[v * \mathfrak{g}] = 0 \in M/\langle b \rangle$.*

Proof. Since M is an interval module, each homogeneous element is a multiple of some element of \mathfrak{G} . Using Lemma 3.22 iteratively, replace each element of \mathcal{B} by a respective element of \mathfrak{G} . If $b \in \mathcal{B}$ is indispensable for \mathfrak{g} , then $v * \mathfrak{g}$ is a multiple of b . \square

Note that the reverse of the second statement does not hold true.

3.5. Algorithm for interval modules. The software Topcat [16] of Gäfvert provides tools for calculating stable ranks for arbitrary persistence modules and, in principle, is based on searching through the entire $B(M, \delta)$ to look for an element with the smallest β_0 . This search can be done more efficiently for interval modules in the two-parameter case, and we now present its algorithm. An implementation in C++ of our algorithm is made available at https://github.com/recorb/shiftdim_stablerank.

Let $v = (v_1, v_2) \in \mathbb{R}_{>0}^2$, M be an interval module, and $\mathfrak{G} := \{\mathfrak{g}_i\}_{i \in I \subseteq \mathbb{R}}$ be the generating curve of M . Denote $\deg(\mathfrak{g}_i) = (a_i, b_i)$. W.l.o.g. assume that the generators are listed in the following total order: $a_i < a_j$ or $b_i > b_j$ whenever $i < j$.

Algorithm 3.24.

Step 1. Cluster $\{\mathfrak{g}_i\}_{i \in I_1}$ in increasing order as follows: Set $I_1 := I$, $k := 1$.

While there exists $i \in I_k$ such that $v * \mathfrak{g}_i \neq 0$, do:

- (a) Set i_k to be the maximum of the set $\{i \in I_k \mid \deg(\mathfrak{g}_i) \leq \deg(v * \mathfrak{g}_j) \text{ for all } j \in I_k \text{ s.t. } j < i \text{ and } v * \mathfrak{g}_j \neq 0\}$.
- (b) Define $I_{k+1} := I_k \setminus \{i \in I_k \mid \deg(\mathfrak{g}_{i_k}) \leq \deg(v * \mathfrak{g}_i)\}$.
- (c) Replace k by $k + 1$ and iterate.

Step 2. Set $\mathfrak{H} := \{\mathfrak{g}_{i_1}, \dots, \mathfrak{g}_{i_\ell}\}$ and $\ell := |\mathfrak{H}|$.

Step 3. Return ℓ and \mathfrak{H} .

Note that the computation of the i_k and I_{k+1} can be carried out geometrically: to obtain i_k , we project from $(\inf\{a_i \mid i \in I_k, v * \mathfrak{g}_i \neq 0\}, \sup\{b_i \mid i \in I_k, v * \mathfrak{g}_i \neq 0\}) + v$ down to the generating curve of M . For the construction of I_{k+1} , we project to the right from $\deg(\mathfrak{g}_{i_k}) - v$ to the generating curve and hit a coordinate (a, b) . We remove all generators whose second coordinate is not strictly greater than b from I_k to obtain I_{k+1} . See Figure 3 for an illustration of this geometric procedure.

Theorem 3.25. *Algorithm 3.24 computes $\dim_v(M)$ in at most $\lceil \frac{b_{i_1} - \inf\{b_i \mid i \in I\}}{v_2} \rceil$ iterations of Step 1.*

Proof. First note that by Corollary 3.23 (i), there is a subset of \mathfrak{G} that is a v -basis of M . Hence, a minimal subset \mathfrak{H} of \mathfrak{G} such that $v * \mathfrak{g}_i \in \langle \mathfrak{H} \rangle$ for all $i \in I$ is indeed a v -basis. If $\mathfrak{H} = \emptyset$, the statement is clear. Hence, let $\ell > 0$. Let us first argue that ℓ is finite. Since $v_1 \neq 0$, i_1 is well-defined. For all $0 < k < \ell$, $b_{i_k} - b_{i_{k+1}} \geq v_2 > 0$. Since $b_i \geq 0$ for all $i \in I$, it follows that ℓ is finite; in fact, $\ell < \lceil \frac{b_{i_1} - \inf\{b_i \mid i \in I\}}{v_2} \rceil < \lceil \frac{b_{i_1}}{v_2} \rceil$.




FIGURE 3. *Left:* Constructing a $(1.5, 1.5)$ -basis, using Algorithm 3.24, clustering from above. *Right:* Using the opposite version from below.

We now prove that $\ell = \dim_v(M)$ and \mathfrak{H} is a v -basis of M . We have $v*(M/\langle \mathfrak{H} \rangle) = 0$. Hence, $\dim_v(M) \leq \ell$. For the reverse direction, assume that $\ell > 0$ and let $\mathcal{B} := \{\mathbf{b}_j\}_{j \in J}$ be a v -basis of M . Applying Lemma 3.22 iteratively, we can w.l.o.g. assume that for each j , $\mathbf{b}_j = \mathbf{g}_i$ for some $i \in I$. Hence, w.l.o.g. \mathcal{B} is a subset of \mathfrak{G} that might differ from \mathfrak{H} . However, \mathcal{B} clusters I into $\dim_v(M)$ subsets; the cluster labeled by $j \in J$ consists of all $i \in I$ for which $\deg(\mathbf{b}_j) \leq (\deg v * \mathbf{g}_i)$. If two elements of I both are elements of one such cluster, then each element in between must be in the same cluster as well. By construction, for all $1 \leq k \leq \ell$ there is no $i \in I \setminus (\bigcup_{1 \leq j \leq k-1} I_j)$ such that there exists $\tilde{I}_k \supseteq I_k$ with $\tilde{I}_k \cap (\bigcup_{1 \leq j \leq k-1} I_j) = \emptyset$ and $\langle \{v * \mathbf{g}_j \mid j \in \tilde{I}_k\} \rangle \subseteq \langle \mathbf{g}_i \rangle$. Hence, the cluster given by \mathcal{B} which contains the smallest elements of I must be a subset of I_1 . By construction of I_2 , the second cluster given by \mathcal{B} can not be a proper superset of $I_1 \cup I_2$. Iteratively, we get that the cardinality of \mathcal{B} is at least ℓ . \square

Replacing the total order on the generators by the opposite total order yields another variant of the algorithm computing $\dim_v(M)$. This yields the upper bound $\ell \leq \lceil \frac{a_{i_1} - \inf\{a_i \mid i \in I\}}{v_1} \rceil \leq \lceil \frac{a_{i_1}}{v_1} \rceil$. The algorithm can be extended to the case $v_1 = 0$, $v_2 \neq 0$ if $\sup\{b_i \mid i \in I\} < \infty$ and—using the opposite total order—to the case $v_1 \neq 0$, $v_2 = 0$ if $\sup\{a_i \mid i \in I\} < \infty$. If in these cases the suprema are not finite, $\dim_v(M) = \infty$.

For the complexity of the algorithm, it remains to investigate the complexity of Steps 1(a) and 1(b). By the geometric construction described above, we need to compute infima, suprema, add and subtract a vector, project to a curve, and check if certain segments of a curve get shifted beyond another curve. In general, computing the shift-dimension is NP-hard, as was mentioned in Remark 3.7. We are now going to prove that the computation is linear for tame interval modules.

Corollary 3.26. *If M is a finitely presented interval module in the bivariate case, then $\dim_v(M)$ can be computed in $O(n)$ time, where $n := \beta_0(M)$.*

Proof. Each iteration of Step 1 boils down to carrying out two additions, determining three minima of a finite array of sorted data, and deciding whether $v * \mathbf{g}_j = 0$ only for a subset of minimal homogeneous generators. Deciding whether $v * \mathbf{g}_j = 0$ can again be done by finding the minimum of a finite array of sorted data. The total number of the latter operations can naively be bounded by $\beta_0(M)$ and can hence be done as a preprocessing step of $\beta_0(M)$ operations. All the individual operations can be done in $O(1)$ time. Running the algorithm from above or below, the number of iterations of Step 1 is bounded by $\lceil \frac{a_{i_1} - \min\{a_i \mid i \in I\}}{v_1} \rceil$ or $\lceil \frac{b_{i_1} - \min\{b_i \mid i \in I\}}{v_2} \rceil$, respectively. Since $\beta_0(M)$ is finite, it is an additional upper bound for the number of iterations of Step 1. \square

For some interval modules, such as monomial ideals, the module generated by \mathfrak{H} has an illustrative description: it is a staircase with the minimal possible number of steps that fits between M and $v * M$. This investigation might give additional ideas for a proof of an analogous algorithm in the setting of more than two parameters. Such a generalization would presumably increase the geometric complexity of the algorithm.

The insights of this algorithm in combination with a better structural understanding of the examples in Section 3.3 might give rise to an efficient way to compute the shift-dimension of direct sums of interval modules.

4. THE SHIFT-DIMENSION OF MULTIGRADED $\mathbb{K}[x_1, \dots, x_r]$ -MODULES

In this section, we explain how the shift-dimension for tame persistence modules translates to the monoid $G = \mathbb{N}^r$. Identifying the monoid ring $\mathbb{K}[\mathbb{N}^r]$ with the polynomial ring $\mathbb{K}[x_1, \dots, x_r]$ and hence translating $v * (\bullet)$ to multiplication by the monomial $x_1^{v_1} \cdots x_r^{v_r}$, the study of $\text{Mod}^{\text{gr}}(\mathbb{K}[G])$ is tantamount to the study of multigraded modules over the multivariate polynomial ring. Those modules play a fundamental role in algebraic geometry. We denote by $S := \mathbb{K}[x_1, \dots, x_r]$ the polynomial ring in r variables over a field \mathbb{K} with the standard \mathbb{N}^r -grading, i.e., the degree of x_i is the i -th standard unit vector $\mathbf{e}_i \in \mathbb{N}^r$. Motivated by the stabilization of β_0 with respect to the metric arising from the standard contour in the direction of a vector, we introduce the following invariant of multigraded S -modules.

Definition 4.1. Let M be an \mathbb{N}^r -graded S -module and $v \in \mathbb{N}^r$. Elements m_1, \dots, m_k of M *v-generate* M , if

$$x_1^{v_1} \cdots x_r^{v_r} \cdot M \subseteq \langle m_1, \dots, m_k \rangle.$$

A *v-basis* of M is a collection $\{m_1, \dots, m_k\}$ of elements of M that *v-generates* M and for which k is smallest possible. In this case, we call k the *v-dimension* of M and denote it by $\dim_v(M)$.

Again, for finitely generated persistence modules, we stick to *homogeneous* generators and *v-bases*. For fixed $v \in \mathbb{N}^r$, we abbreviate \dim_{nv} by \dim_n and refer to it as *n-dimension*, and similarly for “*n-basis*” and “*n-generators*”.

Remark 4.2. This definition generalizes to arbitrary rings R . Let M be an R -module and $r \in R$. Elements $\{m_1, \dots, m_k\}$ of M *r-generate* M if $rM \subseteq \langle m_1, \dots, m_k \rangle$. \diamond


Note that $\dim_0(M)$ is the minimal number of (homogeneous) generators of M . Moreover, $\dim_n(M)$ is zero if and only if $(x_1^{v_1} \cdots x_r^{v_r})^n M$ is the zero-module.

Remark 4.3. By Hilbert’s syzygy theorem, every finitely generated \mathbb{N}^r -graded S -module has a minimal free resolution F_\bullet of length at most r (see [25, Proposition 8.18]). The rank of F_i is the *i-th total multigraded Betti number* of the module. It would be intriguing to stabilize higher total multigraded Betti numbers as well. \diamond

The following proposition explains how to modify tame persistence modules to obtain a module in which every element can be extended to an *n-basis*. This construction works for the monoid ring $\mathbb{K}[\mathbb{N}^r]$, but not for $\mathbb{K}[\mathbf{R}^r]$.

Proposition 4.4. Let M be a finitely generated \mathbb{N}^r -graded S -module and $n \in \mathbb{N}$. Choose a non-zero element $m_1 \in M$ that is not contained in any *n-basis* of M and consider the quotient module $M/\langle m_1 \rangle$. Then choose a non-zero element $[m_2] \in M/\langle m_1 \rangle$ that is not contained in any *n-basis* of $M/\langle m_1 \rangle$, and repeat this process. There exists a natural number ℓ such that after ℓ iterations

$$M \longrightarrow M/\langle m_1 \rangle \longrightarrow M/\langle m_1, m_2 \rangle \longrightarrow \cdots \longrightarrow M/\langle m_1, \dots, m_\ell \rangle =: \widetilde{M},$$

FIGURE 4. Visualization of M , x_1x_2M , and $x_1^2x_2^2M$ for M from Example 4.9

one arrives at a module \widetilde{M} of the same n -dimension as M in which every non-zero element can be extended to an n -basis of \widetilde{M} .

Proof. Since M is finitely generated and S is Noetherian, the procedure described in the proposition stabilizes after a finite number of steps. \square

Note that the module \widetilde{M} can in general *not* be obtained by successively quotienting out non- n -basis-elements of M itself. For instance, consider $M = \langle x_1, x_2 \rangle$ and $v = \mathbb{1}$ the all-one vector. The element $m_1 := x_1^5x_2^3$ is not in any 2-basis of M and $m_2 := x_1^3x_2^5$ is not in a 2-basis of M , but $\{[m_2]\}$ is a 2-basis of the module $M/\langle m_1 \rangle$.

Remark 4.5. Recall that for two ideals I, J in a ring R , the *ideal quotient* of I by J $(I : J) := \{r \in R \mid rJ \subseteq I\} \triangleleft R$ is itself an ideal in R . Let $M = I \triangleleft \mathbb{K}[x_1, \dots, x_r]$ be a homogeneous ideal, $v \in \mathbb{N}^r$. Then for all $n \in \mathbb{N}$, the n -dimension of the ideal quotient $(I : (x_1^{v_1} \cdots x_r^{v_r})^n)$ bounds the minimal number of generators of I from below. \diamond

Example 4.6. Let M be the monomial ideal $\langle x_1^2, x_2^3, x_3^5 \rangle \triangleleft \mathbb{C}[x_1, x_2, x_3]$ and $v = \mathbb{1}$. For $n \in \{0, 1\}$, the three elements x_1^2, x_2^3, x_3^5 n -generate M . For $n \geq 2$, M is n -generated by the single element x_1^2 . Therefore, the sequence $(\dim_n(M))_{n \in \mathbb{N}}$ is $3, 3, 1, 1, 1, \dots$ \diamond

Example 4.7. Let $I = \langle x_1x_2^4, x_1^3x_2^2, x_1^5x_2 \rangle \triangleleft \mathbb{K}[x_1, x_2]$ and $v = \mathbb{1}$. Then I^2 is minimally generated by the 5 elements $x_1^2x_2^8, x_1^4x_2^6, x_1^6x_2^4, x_1^8x_2^3, x_1^{10}x_2^2$. The sequences of n -dimensions are $(\dim_n(I))_{n \in \mathbb{N}} = 3, 3, 1, 1, \dots$ and $(\dim_n(I^2))_{n \in \mathbb{N}} = 5, 5, 1, 1, \dots$ \diamond

Example 4.8 (Free multigraded module). Let $F = S(-a_1, \dots, -a_r) \cong S \cdot x_1^{a_1} \cdots x_r^{a_r}$ be a free multigraded module, where $a_1, \dots, a_r \in \mathbb{N}$. One reads that F is n -generated by the single element $x_1^{a_1+nv_1} \cdots x_r^{a_r+nv_r}$. Therefore $(\dim_n(F))_{n \in \mathbb{N}} = 1, 1, 1, \dots$ \diamond

Example 4.9 (Quotient of two monomial ideals). Let $M = \langle x_1^3x_2, x_1x_2^3 \rangle / \langle x_1^4x_2^4 \rangle$ and $v = \mathbb{1}$. Then $(\dim_n(M))_{n \in \mathbb{N}} = 2, 2, 1, 0, 0, \dots$. For a visualization, see Figure 4. \diamond

For $r = 1$, the shift-dimension is additive. For $r > 1$, this does in general *not* hold true, as the following counterexamples for $r = 2$ and $v = \mathbb{1}$ demonstrate.

Example 4.10 (Revisiting Example 3.14). Let $M = \langle x_1 \rangle / \langle x_1x_2^2 \rangle$ and $N = \langle x_2 \rangle / \langle x_1^2x_2 \rangle$. Then $x_1x_2(M \oplus N)$ is contained in the submodule $\langle (x_1x_2, x_1x_2) \rangle$ of $M \oplus N$. To see that, observe that $x_1(x_1x_2, x_1x_2) = x_1x_2(x_1, 0)$ and $x_2(x_1x_2, x_1x_2) = x_1x_2(0, x_2)$ in $M \oplus N$. It follows that $\dim_1(M \oplus N) = 1 \neq 1 + 1 = \dim_1(M) + \dim_1(N)$. \diamond

Example 4.11. Let $M_1 = \langle x_1^2 \rangle / \langle x_1^6, x_1^2x_2^4 \rangle$, $M_2 = \langle x_1x_2 \rangle / \langle x_1^5x_2, x_1x_2^5 \rangle$, and $M_3 = \langle x_2^2 \rangle / \langle x_1^4x_2^2, x_2^6 \rangle$. Then $\dim_3(M_i) = 1$ for $i = 1, 2, 3$ but $\dim_3(M_1 \oplus M_2 \oplus M_3) = 1$, since the singleton $\{(x_1^3x_2^3, x_1^3x_2^3, x_1^3x_2^3)\}$ is a 3-basis of the module $M_1 \oplus M_2 \oplus M_3$. \diamond

We now present some cases for which additivity of the shift-dimension does hold true. They all are immediate consequences of statements presented in Section 3.2. Here, we rephrase the statements for the discretized setting and omit the proofs.

Lemma 4.12. *If F and G are two free multigraded modules of finite rank, then $\dim_v(F \oplus G) = \dim_v(F) + \dim_v(G)$ for all $v \in \mathbb{N}^r$.* \square

Proposition 4.13. *Let $M = S/\langle x_1^{a_1} \cdots x_r^{a_r} \rangle$ and F a free multigraded module of rank one. Then $\dim_v(M \oplus F) = \dim_v(M) + \dim_v(F)$ for all v .* \square

Proposition 4.14. *Let $\dim_n(M) \leq 1$ and F a free multigraded module of rank k . Then $\dim_v(M \oplus F) = \dim_v(M) + k$.* \square

Theorem 4.15. *Let $0 \rightarrow M \xrightarrow{\varphi} L \xrightarrow{\psi} N \rightarrow 0$ be a short exact sequence of \mathbb{N}^r -graded S -modules. Then for all $v, w \in \mathbb{N}^r$, the following two inequalities hold:*

- (i) $\dim_{v+w}(L) \leq \dim_v(M) + \dim_w(N)$,
- (ii) $\dim_v(L) \leq \dim_v(N) + \beta_0(M)$. \square

If $\dim_v(M) = \beta_0(M)$ or $\beta_0(M) = 1$, it follows that $\dim_v(L) \leq \dim_v(M) + \dim_v(N)$.

5. OUTLOOK TO FUTURE WORK

Our article outlines several pathways for future research. On the one hand, it would be intriguing to stabilize discrete invariants other than the zeroth total multigraded Betti number, such as the rank of the highest syzygy module. The following question immediately arises: *Is the rank of the highest syzygy module “naturally” linked to a noise system—similar to the zeroth total multigraded Betti number being linked to the noise system in the direction of a vector in the sense of Definition 2.7?* A further challenging question is whether the stabilized Euler characteristic is the alternating sum of the stabilization of the ranks.

On the other hand, it would be worthwhile to investigate the stabilization of β_0 with respect to pseudometrics other than the one arising from the standard contour. For that, a first step is the description of the shift of the module. As described, for the standard contour in the direction of a vector, this is the module multiplied by the corresponding monomial. For other contours, there is no such algebraic description yet.

Another interesting direction would be to investigate the shift-dimension for (not necessarily graded) modules over arbitrary rings and to develop a geometric intuition.

We believe that the shift-dimension allows for the extraction of finer and new information of data. This makes it valuable for applications in the medical and life sciences, among others. In order to compute the shift-dimension of multipersistence modules of actual data arising in the sciences efficiently, one needs to extend the linear-time algorithm presented in this article to modules other than interval modules, such as direct sums of interval modules. Those indeed arise from data [15] and approximate persistence modules in the sense of [1], [5], and [9]. Using our algorithm, evaluating the shift-dimension summand-wise is already possible for this class of modules. It would be worthwhile to investigate if this additive version is stable, as was suggested to us by Ulrich Bauer. Furthermore, it would be interesting to find bounds for $\text{err}_{v,p}$ and to determine subclasses of modules for which this error is zero.

In the one-parameter case, two persistence modules M and N are known to be isomorphic if and only if $\widehat{\beta}_{0C}(M)$ and $\widehat{\beta}_{0C}(N)$ coincide for each persistence contour C . At present, it is not known if that statement holds true for the multiparameter case as well. As a first step in this direction, one might investigate whether one can distinguish non-isomorphic multigraded modules in the following sense: given two non-isomorphic modules M, N , can one find a noise system $(\mathcal{V}_\varepsilon)_{\varepsilon \in \mathbb{R}_{\geq 0}}$ such that $M \in \mathcal{V}_0$ but $N \notin \mathcal{V}_0$?

REFERENCES

- [1] H. Asashiba, M. Buchet, E. G. Escolar, K. Nakashima, and M. Yoshiwaki. On interval decomposability of 2D persistence modules. *Comput. Geom.*, 105–106:101879, 2022. 18
- [2] S. Biasotti, A. Cerri, P. Frosini, and D. Giorgi. A new algorithm for computing the 2-dimensional matching distance between size functions. *Pattern Recognition Letters*, 32(14):1735–1746, 2011. 2
- [3] M. B. Botnan and W. Crawley-Boevey. Decomposition of persistence modules. *Proc. Amer. Math. Soc.*, 148:4581–4596, 2020. 1
- [4] M. B. Botnan, J. Curry, and E. Munch. A relative theory of interleavings. Preprint arXiv:2004.14286, 2020. 2
- [5] M. B. Botnan, S. Oppermann, and S. Oudot. Signed barcodes for multi-parameter persistence via rank decompositions. In *Proceedings of the 38th International Symposium on Computational Geometry (SoCG 2022)*. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2022. 18
- [6] P. Bubenik, V. de Silva, and J. Scott. Metrics for generalized persistence modules. *Found. Comput. Math.*, 15:1501–1531, 2015. 2
- [7] M. Buchet and E. G. Escolar. Every 1D persistence module is a restriction of some indecomposable 2D persistence module. *J. Appl. and Comput. Topology*, 4:387–424, 2020. 8
- [8] G. Carlsson and A. Zomorodian. The theory of multidimensional persistence. *Discrete Comput. Geom.*, 42:71–93, 2009. 1, 2, 3
- [9] W. Chachólski, A. Guidolin, I. Ren, M. Scalamiero, and F. Tombari. Effective computation of relative homological invariants for functors over posets. Preprint arXiv:2209.05923, 2022. 18
- [10] W. Chachólski and H. Riihimäki. Metrics and stabilization in one parameter persistence. *SIAM J. Appl. Algebra Geom.*, 4(1):69–98, 2020. 2, 4, 5, 7
- [11] W. Chachólski, M. Scalamiero, and F. Vaccarino. Combinatorial presentation of multidimensional persistent homology. *J. Pure Appl. Algebra*, 221:1055–1075, 2017. 1
- [12] D. Cohen-Steiner, H. Edelsbrunner, and J. Harer. Stability of persistence diagrams. *Discrete Comput. Geom.*, 37:103–120, 2007. 1
- [13] R. Corbet and M. Kerber. The representation theorem of persistence revisited and generalized. *J. Appl. and Comput. Topology*, 2(1-2):1–31, 2018. 3
- [14] H. Edelsbrunner, D. Letscher, and A. J. Zomorodian. Topological persistence and simplification. *Discrete Comput. Geom.*, 28:511–533, 2002. 1
- [15] E. G. Escolar and Y. Hiraoka. Persistence modules on commutative ladders of finite type. *Discrete Comput. Geom.*, 55(1):100–157, 2016. 3, 18
- [16] O. Gäfvert. Topcat - a library for multiparameter persistence. Available at <https://github.com/olivergafvert/topcat>. 14
- [17] O. Gäfvert and W. Chachólski. Stable invariants for multidimensional persistence. Preprint arXiv:1703.03632, 2017. 2, 5, 7, 8, 9
- [18] R. Ghrist. Barcodes: the persistent topology of data. *Bull. Amer. Math. Soc.*, 45:61–75, 2008. 1
- [19] H. Harrington, N. Otter, H. Schenck, and U. Tillmann. Stratifying multiparameter persistent homology. *SIAM J. Appl. Algebra Geom.*, 3(3):439–471, 2017. 3
- [20] B. Keller, M. Lesnick, and T. L. Willke. Persistent Homology for Virtual Screening. Preprint ChemRxiv, 2018. 3
- [21] M. Kerber, M. Lesnick, and S. Oudot. Exact computation of the matching distance on 2-parameter persistence modules. In *Proceedings of the 35th International Symposium on Computational Geometry (SoCG 2019)*, pages 46:1–46:15, 2019. 2
- [22] M. Kerber and A. Rolle. Fast minimal presentations of bi-graded persistence modules. In *2021 Proceedings of the Symposium on Algorithm Engineering and Experiments (ALENEX)*, pages 207–220. SIAM, 2021. 3
- [23] M. Lesnick. The theory of the interleaving distance on multidimensional persistence modules. *Found. Comput. Math.*, 15(3):613–650, 2015. 2
- [24] M. Lesnick and M. Wright. Computing minimal presentations and Betti numbers of 2-parameter persistent homology. *SIAM J. Appl. Algebra Geom.*, 6(2), 2022. 3

- [25] E. Miller and B. Sturmfels. *Combinatorial Commutative Algebra*, volume 227 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005. 16
- [26] R. Peeters. Orthogonal representations over finite fields and the chromatic number of graphs. *Combinatorica*, 16:417–431, 1996. 9
- [27] M. Scalamiero, W. Chachólski, A. Lundman, R. Ramanujam, and S. Öberg. Multidimensional persistence and noise. *Found. Comput. Math*, 17(6):1367–1406, 2017. 2, 4, 8, 9
- [28] The RIVET Developers. RIVET: A program for the visualization and analysis of two-parameter persistent homology. Available at <https://github.com/rivetTDA/rivet>, 2020. 3
- [29] A. Thomas. *Invariants and Metrics for Multiparameter Persistent Homology*. PhD thesis, Duke University, 2019. 3
- [30] A. Zomorodian and G. Carlsson. Computing persistent homology. *Discrete Comput. Geom.*, 33(2):249–274, 2005. 1

WOJCIECH CHACHÓLSKI, DEPARTMENT OF MATHEMATICS, KTH ROYAL INSTITUTE OF TECHNOLOGY, LINDSTEDTSVÄGEN 25, 114 28 STOCKHOLM, SWEDEN
Email address: `wojtek@kth.se`

RENÉ CORBET, DEPARTMENT OF MATHEMATICS, KTH ROYAL INSTITUTE OF TECHNOLOGY, LINDSTEDTSVÄGEN 25, 114 28 STOCKHOLM, SWEDEN
Email address: `corbet@kth.se`

ANNA-LAURA SATTELBERGER, MAX-PLANCK-INSTITUT FÜR MATHEMATIK IN DEN NATURWISSENSCHAFTEN, INSELSTRASSE 22, 04103 LEIPZIG, GERMANY AND DEPARTMENT OF MATHEMATICS, KTH ROYAL INSTITUTE OF TECHNOLOGY, LINDSTEDTSVÄGEN 25, 114 28 STOCKHOLM, SWEDEN (*current*)
Email address: `alsat@kth.se`