

# A TODA BRACKET CONVERGENCE THEOREM FOR MULTIPLICATIVE SPECTRAL SEQUENCES

EVA BELMONT AND HANA JIA KONG

ABSTRACT. Moss’ theorem, which relates Massey products in the  $E_r$ -page of the classical Adams spectral sequence to Toda brackets of homotopy groups, is one of the main tools for calculating Adams differentials. Working in an arbitrary symmetric monoidal stable topological model category, we prove a general version of Moss’ theorem which applies to spectral sequences that arise from filtrations compatible with the monoidal structure. The theorem has broad applications, e.g. to the computation of the motivic slice and motivic Adams spectral sequences.

## 1. INTRODUCTION

In homological algebra, Massey products [Mas58] are operations that encode higher multiplicative structure on a differential graded algebra. Analogously in homotopy theory, Toda brackets (e.g. [Koc90, Tod62]) encode higher multiplicative structure on homotopy classes of maps, for example on the homotopy groups of a ring spectrum.

In a nice enough spectral sequence computing homotopy groups of a ring spectrum, the multiplicative structure on  $E_r$ -pages of the spectral sequence agrees with the multiplicative structure of the homotopy groups. A theorem by Moss [Mos70, Theorem 1.2] states that in the Adams spectral sequence, there is a correspondence of higher multiplicative structures as well: given a Massey product of permanent cycles in the  $E_r$ -page such that the corresponding Toda bracket is defined, and assuming a “crossing differential” hypothesis in the relevant degrees, the Massey product contains a permanent cycle that converges to an element of the Toda bracket. This correspondence of Toda brackets and Massey products is used heavily in determining the differentials of Adams spectral sequences; for modern work with classical references, see [Isa19, IWX20].

It is natural to ask if we can extend Moss’ theorem to other spectral sequences. Given a suitable multiplicative spectral sequence, the  $E_r$ -page is a differential graded algebra, and Massey products are naturally defined. On the other hand, the literature is less straightforward about the existence of Toda brackets in a symmetric monoidal stable model category, though such a construction is commonly assumed in practice. There are two main settings in which one considers Toda brackets:

- Given a sequence of maps  $Y \xrightarrow{\alpha''} X'' \xrightarrow{\alpha'} X' \xrightarrow{\alpha} X$  such that  $\alpha\alpha'$  and  $\alpha'\alpha''$  are null homotopic, one can define a *composition Toda bracket*  $\langle \alpha, \alpha', \alpha'' \rangle$ .
- Let  $S$  be the unit object, let  $X'$  be a left  $X''$ -module and let  $X'$  be a left  $X$ -module. Given classes  $\alpha'' : S \rightarrow X''$ ,  $\alpha' : S \rightarrow X'$ ,  $\alpha : S \rightarrow X$  we can define the products  $\alpha\alpha'$  and  $\alpha'\alpha''$ . If these products are zero then one may define a *multiplicative (or smash) Toda bracket*  $\langle \alpha, \alpha', \alpha'' \rangle$ .

The construction of both types of brackets in the stable homotopy category is classical; see [Koc80] for references and a comparison in the case  $Y = X'' = X' = S$ . There are also many general constructions of composition Toda brackets; see [CF17] for a summary and

comparison. In this paper, we set up 3-fold smash Toda brackets in the homotopy category of a general symmetric monoidal stable model category, and study the Moss convergence theorem in this generalized setting. Most of the material in this paper works for composition Toda brackets, but Proposition 4.10 requires the multiplicative setting, which is by far the more widely used one. Our result extends Moss' convergence theorem to any spectral sequence from a filtration with a compatible pairing. A related contribution of this paper is to reframe Moss' original work in a more modern framework.

Our motivation for writing this paper is for use in studying the motivic slice spectral sequences, which has emerged as a useful tool for computing motivic homotopy groups. In Section 5 we give examples of  $\mathbb{R}$ -motivic slice spectral sequence computations using our theorem. This theorem also applies to other spectral sequences. Recent computations of the  $\mathbb{C}$ -motivic Adams spectral sequence [Isa19, IWX20] assert but do not prove such a theorem is true in this context (see [Isa19, Theorem 3.1.1]); the present paper fills this gap in the literature. We feel certain the results here will also find other applications.

**1.1. Results and methods.** Let  $(\mathcal{C}, \wedge, S)$  be a symmetric monoidal stable topological model category and consider objects  $X(i) \in \mathcal{C}$  for  $i = 1, 2, 3$ . Suppose there are towers of fibrations

$$X(i) = X(i)_0 \longleftarrow X(i)_1 \longleftarrow X(i)_2 \longleftarrow \dots$$

and compatible pairings

$$\mu_1^{s,t} : X(2)_s \wedge X(1)_t \rightarrow X(1)_{s+t}, \quad \mu_2^{s,t} : X(3)_s \wedge X(2)_t \rightarrow X(2)_{s+t}, \quad \mu_3^{s,t} : X(3)_s \wedge X(1)_t \rightarrow X(1)_{s+t}.$$

Write  $\mu_i$  for  $\mu_i^{0,0}$  and assume that  $\mu_1\mu_2 = \mu_3\mu_1 : X(3) \wedge X(2) \wedge X(1) \rightarrow X(1)$ . The towers give rise to spectral sequences  $E_1(X(i)) = \pi_*(X(i)_f/X(i)_{f+1}) \implies \pi_*(X(i))$  where  $\pi_*(-)$  denotes  $\pi_*(\text{Map}_{\mathcal{C}}(S, -))$ .

**Theorem 1.1** (Theorem 4.11). *Fix  $r \geq 2$ . Suppose  $a_i \in E_r(X(i))$  for  $1 \leq i \leq 3$  are permanent cycles converging to elements  $\alpha_i \in \pi_*(X(i))$  such that  $0 = a_1a_2 = a_2a_3$  and  $0 = \alpha_1\alpha_2 = \alpha_2\alpha_3$ . Suppose the crossing differential hypothesis (see Definition 2.10) is satisfied at the  $E_{r-1}$ -page in the degrees of  $a_1a_2$  and  $a_2a_3$ . Then the Massey product  $\langle a_1, a_2, a_3 \rangle \subseteq E_r(X(1))$  contains a permanent cycle that converges to an element of the Toda bracket  $\langle \alpha_1, \alpha_2, \alpha_3 \rangle \subseteq \pi_*(X(1))$ .*

In the theorem, we assumed  $r \geq 2$  since in that case,  $E_r(X(i))$  is the homology of the differential graded algebra  $(E_{r-1}(X(i)), d_{r-1})$ . In the  $r = 1$  case, we consider Toda brackets on  $E_1(X(i))$  by regarding it as the homotopy groups of the associated graded objects  $X(i)_*/X(i)_{*+1}$  of the filtration on  $X(i)$ .

**Theorem 1.2** (Theorem 4.16). *Assume the setup of Theorem 4.11 with  $r = 1$ . Then the Toda bracket  $\langle a_1, a_2, a_3 \rangle \subseteq \pi_*(X(1)_*/X(1)_{*+1}) = E_1(X(1))$  contains a permanent cycle that converges to an element of the Toda bracket  $\langle \alpha_1, \alpha_2, \alpha_3 \rangle \subseteq \pi_*(X(1))$ .*

In fact, the  $r = 1$  case is not as different as it seems: in Section 2.2 we identify  $E_r(X(i))$  for  $r \geq 1$  with a quotient of  $\pi_*(X(i)_*/X(i)_{*+r})$ . Our strategy consists of the comparison of three kinds of brackets:

- (1) Massey products of classes in  $E_r(X(i))$  defined in terms of classes in  $\pi_*(X(i)_*/X(i)_{*+r})$ ,
- (2) Toda brackets of classes in  $\pi_*(X(i)_*)$ ,
- (3) Toda brackets of classes in  $\pi_*(X(i))$ .

We show that any element in (2) represents an element in (1), and detects an element of (3). This implies our main theorem. The crossing differential hypothesis (see Definition 2.10, Assumption 3.3) is necessary for constructing (2). In practice, the crossing differential hypothesis is satisfied in many cases.

1.2. **Outline.** In Section 2, we introduce notation for the spectral sequence associated to a tower of fibrations and give a reformulation of the  $E_r$ -page that will be used in the rest of the paper. In Section 3 we give notation for the setup of the main theorem, namely a pairing of spectral sequences satisfying certain properties along with certain elements that will form the relevant Toda brackets and Massey products. We define 3-fold Toda brackets and Massey products in our general setting. In Section 4 we prove our versions of Moss' theorem (Theorems 4.11 and 4.16). In Section 5 we give some example applications of these theorems to specific computations in the  $\mathbb{R}$ -motivic slice spectral sequence.

The reader who wishes to jump to the precise statements of the theorems should read Section 3.2 and then skip to Section 4.2.

1.3. **Acknowledgements.** We would like to thank Dan Isaksen for helpful conversations and comments, and Peter May for helpful comments on our manuscript. The second author was supported by the National Science Foundation under Grant No. DMS-1926686.

## 2. THE SPECTRAL SEQUENCE ASSOCIATED TO A TOWER

2.1. **The spectral sequence for a tower of fibrations.** We work in a stable topological model category  $\mathcal{C}$  and its homotopy category  $h\mathcal{C}$ . Let  $X$  and  $Y$  be two objects in  $\mathcal{C}$ . (Starting in Section 3.2 we will assume that  $Y$  is the unit object.) Suppose we have a tower of fibrations:

$$(2.1) \quad \begin{array}{ccccccc} X = X_0 & \xleftarrow{i} & X_1 & \xleftarrow{\quad} & \dots & \xleftarrow{i} & X_f & \xleftarrow{i} & X_{f+1} & \xleftarrow{i} & \dots \\ & & \searrow \kappa & & & & \searrow \kappa & & & & \\ & p \downarrow & & & & & p \downarrow & & & & \\ & X_0/X_1 & & & & & X_f/X_{f+1} & & & & \end{array}$$

Applying  $\pi_* \text{Map}_{\mathcal{C}}(Y, -)$ , we have a spectral sequence

$$E_1^{n,f}(Y, X) = \pi_n \text{Map}_{\mathcal{C}}(Y, X_f/X_{f+1}) \Rightarrow \pi_n \text{Map}_{\mathcal{C}}(Y, X).$$

We will use the bigrading convention  $(n, f)$  where  $n$  is the stem and  $f$  is the filtration.

**Notation 2.2.** When the object  $Y$  is clear, we will drop it from the notation, writing

$$\pi_n(X) := \pi_n \text{Map}_{\mathcal{C}}(Y, X)$$

$$E_r^{n,f}(X) := E_r^{n,f}(Y, X).$$

If the stem  $n$  is clear or unimportant, we will also drop it from the notation, writing  $E_r^f := E_r^{*,f}$ . For ease of notation, we set the following conventions: let

$$X_{s,r} := X_s/X_{s+r},$$

$$X_{s,r,t} := X_{s,r}/X_{s+t,r}$$

where  $X_{s,r}/X_{s+t,r}$  is the cofiber of the map  $X_{s,r} \rightarrow X_{s+t,r}$  which arises as in the following diagram of cofiber sequences.

$$(2.3) \quad \begin{array}{ccccc} X_{s+t+r} & \longrightarrow & X_{s+t} & \longrightarrow & X_{s+t,r} \\ \downarrow & & \downarrow & & \downarrow \\ X_{s+r} & \longrightarrow & X_s & \longrightarrow & X_{s,r} \\ \downarrow & & \downarrow & & \downarrow \\ X_{s+r,t} & \longrightarrow & X_{s,t} & \longrightarrow & X_{s,r,t} \end{array}$$

The following lemma is standard, but we use the statement to define the map  $\zeta$ .


**Lemma 2.4.** *If  $1 \leq t \leq r$ , there is a homotopy equivalence*

$$(\xi, \xi') : X_{s,r,t} \xrightarrow{\cong} X_{s,t} \oplus \Sigma X_{s+r,t}.$$

*Under this equivalence, the connecting homomorphism  $\kappa : X_{s,r,t} \rightarrow \Sigma X_{s+t,r}$  is the composite  $X_{s,t} \xrightarrow{\kappa'} \Sigma X_{s+t} \xrightarrow{p} \Sigma X_{s+t,r}$  on the first component and the map  $\Sigma X_{s+r,t} \xrightarrow{i} \Sigma X_{s+t,r}$  induced by  $X_{s+r} \rightarrow X_{s+t}$  on the second component.*

*Proof.* Note that when  $t \leq r$ , the first map in the third row of in Diagram 2.3 is zero. The result follows.  $\blacksquare$

We present the following figure as a mnemonic to help keep track of the indexing.



**2.2. Cycles, boundaries, and the crossing differential condition.** By definition,  $E_r^{*,f}(X)$  is a subquotient of  $E_1^{*,f}(X) = \pi_*(X_{f,1})$ . We will reformulate this in terms of elements of  $\pi_*(X_{f,r})$ . We start with a minor reformulation of the usual notions.

**Definition 2.5.** Let

$$Z_r^{*,f} = \{a \in \pi_*(X_{f,1}) : \kappa(a) \in \pi_{*-1}(X_{f+1}) \text{ lifts to } \pi_{*-1}(X_{f+r})\}, \text{ and}$$

$$B_r^{*,f} = \begin{cases} \{a \in \pi_*(X_{f,1}) : a \text{ lifts to some } \alpha \in \ker(\pi_*(X_f) \rightarrow \pi_*(X_{f-r+1}))\} & f \geq r-1, \\ 0 & f < r-1. \end{cases}$$

Say that an element of  $Z_r^{*,*}$  is a  $d_{r-1}$ -cycle, and say that an element of  $B_r^{*,*}$  is a  $d_{r-1}$ -boundary. Note that there is an inclusion  $B_r^{*,f} \subseteq Z_r^{*,f}$  since  $\kappa(a) = 0$  for  $a \in B_r^{*,f}$ . Denote the quotient map  $Z_r^{*,*} \rightarrow Z_r^{*,*}/B_r^{*,*}$  by  $[-]_r$ .

**Remark 2.6.** This is not the usual definition of cycles and boundaries, but we claim that

$$E_r^{*,f}(X) \cong Z_r^{*,f}/B_r^{*,f}.$$

A cycle in the usual sense is an equivalence class of elements  $a \in E_1(X)$  in the kernel of  $d_i$  for all  $i \leq r-1$ , modulo  $d_i$ -boundaries for  $i < r-1$ . Our  $Z_r^{*,*}$  consists of all such elements  $a$  (note that  $\kappa(a)$  lifts to  $\pi_{*-1}(X_{f+r})$  if and only if its image in  $\pi_{*-1}(X_{f+r-1,1})$  is zero). Our  $B_r^{*,*}$  consists of all  $d_i$ -boundaries for  $i \leq r-1$ : a differential  $d_i(b) = a$  means that there is a class  $\tilde{a} \in \pi_*(X_f)$  that projects to  $a \in \pi_*(X_{f,1})$  and whose image in  $\pi_*(X_{f-i+1})$  equals  $\kappa(b)$ ; hence the image of  $\tilde{a}$  in  $\pi_*(X_{f-i})$  is zero.

We begin with an easy lemma summarizing some useful properties about these definitions.

**Lemma 2.7.** *We have*

$$(2.8) \quad Z_r^{*,f} = \{a \in \pi_*(X_{f,1}) : a \text{ lifts to } \pi_*(X_{f,r})\}.$$

*That is, a class  $a \in \pi_*(X_{f,1})$  supports no nontrivial differentials of length  $\leq r$  if and only if  $a$  lifts to  $\pi_*(X_{f,r})$ .*

If  $\alpha \in \pi_*(X_f)$  maps to  $a \in \pi_*(X_{f,r})$  and  $a \neq 0$ , we say that  $a$  is an  $E_\infty$ -representative of  $\alpha$ . We say that  $a \in \pi_*(X_{f,r})$  is a permanent cycle if  $a$  lifts to  $\pi_*(X_f)$ .

*Proof.* Consider the following diagram of cofiber sequences.

$$\begin{array}{ccccc}
X_f & \longrightarrow & X_{f,r} & \xrightarrow{\kappa} & \Sigma X_{f+r} \\
\downarrow & & \downarrow & & \downarrow \\
X_f & \longrightarrow & X_{f,1} & \xrightarrow{\kappa} & \Sigma X_{f+1} \\
\downarrow & & \downarrow & & \downarrow \\
* & \longrightarrow & \Sigma X_{f+1,r-1} & \xrightarrow{=} & \Sigma X_{f+1,r-1}
\end{array}$$

If  $\kappa(a)$  lifts to  $\pi_{*-1}(X_{f+r})$ , then it has zero image in  $\pi_{*-1}(X_{f+1,r-1})$ , and hence  $a$  has zero image in  $\pi_*(X_{f+1,r-1})$ , implying the desired lift. The other direction is clear.  $\blacksquare$

**Lemma 2.9.** *Let*

$$\tilde{Z}_r^{*,f} := \text{coker}(\pi_*(X_{f+1,r-1}) \rightarrow \pi_*(X_{f,r})).$$

*Then the map  $\pi_*(X_{f,r}) \rightarrow \pi_*(X_{f,1})$  factors through an isomorphism*

$$\pi_*(X_{f,r}) \twoheadrightarrow \tilde{Z}_r^{*,f} \xrightarrow{\cong} Z_r^{*,f} \hookrightarrow \pi_*(X_{f,1}).$$

*Moreover, the composition*

$$\tilde{d}_r : X_{f,r} \xrightarrow{\kappa} \Sigma X_{f+r} \xrightarrow{p} \Sigma X_{f+r,r}$$

*is compatible with the differential  $d_r : E_r^{*,f} \rightarrow E_r^{*-1,f+r}$  under the identification  $Z_r \cong \tilde{Z}_r$ .*

Here  $\kappa$  is the boundary map in the cofiber sequence  $X_f \rightarrow X_r \rightarrow X_{f,r}$ .

*Proof.* By Lemma 2.7, we have that the map  $u : \pi_*(X_{f,r}) \rightarrow \pi_*(X_{f,1})$  factors through a surjection to  $Z_r^{*,f}$ , and it suffices to show that  $u(a) = 0$  if and only if  $a$  is in the image of  $\pi_*(X_{f+1,r-1})$ . This follows from the cofiber sequence  $X_{f+1,r-1} \rightarrow X_{f,r} \rightarrow X_{f,1}$ .

Let  $\tilde{d}_r$  denote the composition  $\pi_*(X_{f,r}) \xrightarrow{\kappa} \pi_{*-1}(X_{f+r}) \xrightarrow{p} \pi_{*-1}(X_{f+r,r})$ . We wish to show that the diagram

$$\begin{array}{ccc}
\pi_*(X_{f,r}) & \xrightarrow{\tilde{d}_r} & \pi_{*-1}(X_{f+r,r}) \\
\downarrow & & \downarrow \\
Z_r^{*,f} & & Z_r^{*-1,f+r} \\
\downarrow [-]_r & & \downarrow [-]_r \\
E_r^{*,f} & \xrightarrow{d_r} & E_r^{*-1,f+r}
\end{array}$$

commutes. Suppose we have a differential  $d_r([b]_r) = [a]_r$  where  $b \in Z_r^{*,f} \subseteq \pi_*(X_{f,r})$  is a representative of  $[b]_r$  that has a lift  $\tilde{b} \in \pi_*(X_{f,r})$ . To show that  $\tilde{d}_r(\tilde{b})$  maps to  $[a]_r$  along the right vertical map, we must show that  $\tilde{d}_r(\tilde{b})$  lifts to an element  $\tilde{a} \in \pi_{*-1}(X_{f+r})$  whose image in  $\pi_{*-1}(X_{f+1,r-1})$  is  $\kappa(b)$ .

$$\begin{array}{ccc}
X_{f,r} & \xrightarrow{\kappa} & \Sigma X_{f+r} & \xrightarrow{p} & \Sigma X_{f+r,r} & & \tilde{b} & \longrightarrow & \tilde{a} & \longrightarrow & \tilde{d}_r(\tilde{b}) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
X_{f,1} & \xrightarrow{\kappa} & \Sigma X_{f+1} & & & & b & \longrightarrow & \kappa(b) & & 
\end{array}$$


The diagrams above show that one may take  $\tilde{a} = \kappa(\tilde{b})$ .  $\blacksquare$

If an equivalence class in  $\tilde{Z}_r^{*,f+r} = \text{coker}(\pi_*(X_{f+r+1,r-1}) \rightarrow \pi_*(X_{f+r,r}))$  contains an element in the image of  $\tilde{d}_r : \pi_*(X_{f,r}) \rightarrow \pi_*(X_{f+r,r})$ , not every other representative need be in the image. This is a problem, because we would like to take a chosen representative of a boundary in  $\tilde{Z}_r^{*,f+r}$  and produce an explicit representative of the  $d_r$ -preimage. Proposition 2.12 says that this problem does not occur if we assume a criterion called the crossing differentials hypothesis.

**Definition 2.10.** Say that a spectral sequence  $\{E_*(X)\}$  satisfies the crossing differential hypothesis at the  $E_r$ -page in degree  $(n, f)$  if

$$(2.11) \quad \text{every element in } E_{f-m+1}^{n+1,m}(X) \text{ is a permanent cycle for } 0 \leq m \leq f - r - 1.$$

In the context of a differential  $d_r(x) = y$  for  $x \in E_r^{n+1,f-r}(X)$ ,  $y \in E_r^{n,f}(X)$ , the crossing differentials hypothesis is satisfied in the degree of  $y$  if there are no differentials from stem  $n+1$  to stem  $n$  with source filtration  $< f-r$  and target filtration  $> f$ .



**Proposition 2.12** ([Mos70, Proposition 6.3]). Suppose  $a \in \pi_*(X_{f,1})$  is a  $d_r$ -boundary, and suppose  $\alpha \in \pi_*(X_f)$  is any lift of  $a$  such that  $\alpha \in \ker(\pi_*(X_f) \rightarrow \pi_*(X))$ . If the crossing differential hypothesis at the  $E_r$ -page is satisfied in the degree of  $a$ , then the map  $\pi_*(X_f) \rightarrow \pi_*(X_{f-r})$  sends  $\alpha$  to zero; i.e. the composite  $Y \xrightarrow{\alpha} X_f \xrightarrow{i} X_{f-r}$  is null homotopic.

Proposition 2.12 is used in Lemma 3.4, where we have a specified lift (given as a product) to  $\pi_*(X_f)$  of a class in  $\pi_*(X_{f,1})$ ; Proposition 2.12 guarantees that the chosen lift becomes zero in  $\pi_*(X_{f-r})$ , whereas a priori we only know that there exists a lift with this property.

**2.3. Source of a differential.** In Section 2.2 we described the  $E_r$ -page as a quotient  $Z_r/B_r$ , where  $Z_r$  is isomorphic to a quotient  $\tilde{Z}_r$  of  $\pi_*(X_{f,r})$  (Lemma 2.9). We had a map  $\tilde{d}_r : \pi_*(X_{f,r}) \rightarrow \pi_{*-1}(X_{f+r,r})$  compatible with the differential. While  $\tilde{d}_r$  need not extend to a map  $\tilde{Z}_r \rightarrow \tilde{Z}_r$ , we continue to develop basic spectral sequence machinery using representatives in  $\pi_*(X_{f,r})$  instead of  $\pi_*(X_{f,1})$ . In the next lemma, given  $\alpha \in \pi_*(X_{f,r+1})$  representing a class in the image of  $\tilde{d}_r$ , we give an explicit construction of the source of this differential up to higher filtration, given some additional assumptions related to Proposition 2.12.

**Lemma 2.13.** Suppose  $a \in \pi_*(X_{f,1})$  is a  $d_r$ -boundary and  $\alpha \in \pi_*(X_f)$  is a lift in  $\ker(\pi_*(X_f) \rightarrow \pi_*(X_{f-r}))$ . Then a lift  $\beta$  exists in (2.14), and for any choice of  $\beta$ , we have

$$\tilde{d}_r(\xi\beta) \equiv p(\alpha) \text{ in } \tilde{Z}_r^{*,f}$$

where  $p$  denotes  $\pi_*(X_{f,r+1}) \rightarrow \pi_*(X_{f,r})$ . In other words,  $[\xi\beta]_r$  supports the  $d_r$ -differential that hits  $[a]_r$ .

$$(2.14) \quad \begin{array}{ccc} & \Sigma^{-1}X_{f-r,r+1,r} & \xrightarrow{\xi} \Sigma^{-1}X_{f-r,r} \\ & \beta \nearrow & \downarrow \kappa_{r+1} \\ Y & \xrightarrow{\alpha} & X_{f,r+1} \\ & \searrow \text{null} & \downarrow i \\ & & X_{f-r,r+1} \\ & & \downarrow p \\ & & X_{f-r,r+1,r} \end{array}$$

*Proof.* In this proof we write  $\kappa_{r+1}$  for the boundary map  $\Sigma^{-1}X_{f-r,r+1,r} \rightarrow X_{f,r+1}$  to distinguish it from the boundary map of the cofiber sequence  $X_f \rightarrow X_{f-r} \xrightarrow{\kappa} X_{f,r}$ . A lift  $\beta$  exists because the column in the middle is a cofiber sequence.

Using Lemma 2.4 we may write  $\beta = (\beta_1, \beta_2)$  where  $\beta_1 \in \pi_*(\Sigma^{-1}X_{f-r,r})$  and  $\beta_2 \in \pi_*(X_{f+1,r})$ ; note that  $\beta_1 = \xi\beta$ . Since the diagram

$$(2.15) \quad \begin{array}{ccccc} & \Sigma^{-1}X_{f-r,r+1,r} & \longleftarrow & \Sigma^{-1}X_{f-r,r} & \\ & \kappa_{r+1} \downarrow & & \downarrow \kappa & \\ X_{f,r} & \xleftarrow{p} & X_{f,r+1} & \xleftarrow{p'} & X_f \end{array}$$

commutes, we have  $\tilde{d}_r(\xi\beta) \stackrel{\text{def}}{=} pp'\kappa(\xi\beta) \stackrel{(2.15)}{=} p\kappa_{r+1}(\beta_1)$ . By Lemma 2.4,  $\kappa_{r+1}\beta_2 \equiv 0$  in  $\tilde{Z}_r^{*,f}$ . Since  $\alpha = \kappa_{r+1}\beta$ , the result follows.  $\blacksquare$

### 3. TODA BRACKET AND MASSEY PRODUCT SETUP

Let  $(\mathcal{C}, \wedge, S)$  be a symmetric monoidal stable topological model category.

**3.1. Pairings.** So far, we have only considered a single filtered object  $\{X_f\}_{f \geq 0}$ . The main subject of this paper concerns (higher) multiplicative structures on these filtrations and their associated spectral sequences. The basic notion is of an associative pairing of filtered objects, which in turn allows us to multiply homotopy classes.

**Definition 3.1.** Given a pairing  $\mu : X' \wedge X \rightarrow X$  and classes  $\alpha \in [\Sigma^n S, X]$ ,  $\alpha' \in [\Sigma^m S, X]$ , the product  $\alpha\alpha'$  is defined as the composition  $\Sigma^{m+n} S \simeq \Sigma^m S \wedge \Sigma^n S \xrightarrow{\alpha' \wedge \alpha} X' \wedge X \xrightarrow{\mu} X$ .

**Remark 3.2.** In the rest of the paper, we will make the simplifying assumption that all maps have degree zero, i.e. have source  $S$ . The only change required for nonzero degrees concerns the insertion of minus signs. These are the same as in the classical case, which is well known; see for example [Rav86, §1.4] for the signs in the definition of Massey products on a differential graded algebra.

If there is a corresponding pairing  $E_r(X') \otimes E_r(X) \xrightarrow{\mu} E_r(X)$  then one can define the product of classes  $a' \in E_r(X')$  and  $a \in E_r(X)$  to be the image along the pairing.



**3.2. Setup.** Now we fix notation that will be used in the rest of the paper.

Let  $X, X',$  and  $X''$  be objects in  $\mathcal{C}$  with associative pairings  $\mu'' : X'' \wedge X \rightarrow X, \mu' : X'' \wedge X' \rightarrow X',$  and  $\mu : X' \wedge X \rightarrow X$  such that  $\mu\mu' = \mu''\mu$  as maps  $X'' \wedge X' \wedge X \rightarrow X.$

Suppose there are maps  $\alpha'' : S \rightarrow X'', \alpha' : S \rightarrow X', \alpha : S \rightarrow X$  with null homotopic compositions  $\alpha\alpha' \simeq 0 \simeq \alpha'\alpha''.$  We assume that  $X, X',$  and  $X''$  each have a filtration as in (2.1) such that the pairings are compatible with these filtrations (i.e. there are maps  $\mu : X'_n \wedge X_m \rightarrow X_{n+m}$  and similarly for  $\mu', \mu''$ ). One can show that this gives rise to pairings of spectral sequences, e.g.  $E_r(X') \otimes E_r(X) \rightarrow E_r(X).$

Furthermore, suppose that  $\alpha, \alpha',$  and  $\alpha''$  are detected by nonzero permanent cycles  $a \in E_r^{*,f-f'}(X), a' \in E_r^{*,f'-f''}(X'),$  and  $a'' \in E_r^{*,f''}(X''),$  respectively, such that  $aa'$  and  $a'a''$  are hit by  $d_r.$

Then we have analogous filtered versions of  $\alpha, \alpha', \alpha'',$  e.g.  $\alpha : S \rightarrow X_{f-f'},$  and compositions  $\alpha'\alpha'' : S \rightarrow X'_{f'}, \alpha\alpha' : S \rightarrow X_{f-f''}.$  We will use the symbol  $\alpha$  to refer to one of three maps, depending on context: the original map  $\alpha : S \rightarrow X,$  the filtered version  $S \rightarrow X_{f-f'},$  or the induced map  $S \rightarrow X_{f-f',r}.$

Beware that we need not have  $\alpha'\alpha'' = \alpha\alpha' = 0$  in the filtered context; compare with Lemma 3.4 below.

**Assumption 3.3.** In the rest of the paper, assume that  $E_r(X)$  satisfies the crossing differentials hypothesis (Definition 2.10) at the  $E_r$ -page in the degree of  $aa'$  and  $E_r(X')$  satisfies the crossing differentials hypothesis at the  $E_r$ -page in the degree of  $a'a''.$

**Lemma 3.4.** *Given Assumption 3.3, the following compositions are nullhomotopic.*

$$\begin{aligned} S &\xrightarrow{\alpha\alpha'} X_{f-f''} \rightarrow X_{f-f''-r}, & S &\xrightarrow{\alpha\alpha'} X_{f-f'',r+1} \rightarrow X_{f-f''-r,r+1}, \\ S &\xrightarrow{\alpha'\alpha''} X'_{f'-f''} \rightarrow X'_{f'-f''-r}, & S &\xrightarrow{\alpha'\alpha''} X'_{f',r+1} \xrightarrow{i} X'_{f'-r,r+1}. \end{aligned}$$

*Proof.* Note that  $\alpha\alpha' : S \rightarrow X' \rightarrow X$  factors as  $\alpha\alpha' : S \rightarrow X'_{f'-f''} \rightarrow X_{f-f''} \rightarrow X.$  Since this is null and  $\alpha\alpha' : S \rightarrow X'_{f'-f''} \rightarrow X_{f-f''}$  lifts  $aa'$  which is also assumed to be null, Proposition 2.12 implies that  $S \rightarrow X_{f-f''} \rightarrow X_{f-f''-r}$  is null. This is the first statement. The second statement follows from composing with the map  $X_{f-f''-r} \rightarrow X_{f-f''-r,r+1}.$  The remaining two statements are similar.  $\blacksquare$

**3.3. Toda brackets.** We define the multiplicative Toda bracket  $\langle \alpha, \alpha', \alpha'' \rangle$  following [Koc90, Definitions 2.2.1, 2.2.2]. Let  $CX$  denote the cofiber of  $X \xrightarrow{\text{id}} X.$  Let  $F$  and  $G$  be two nullhomotopies  $\alpha'\alpha'' \xrightarrow{F} 0$  and  $\alpha\alpha' \xrightarrow{G} 0.$  Equivalently, we have maps  $F : CS \rightarrow X'$  and  $G : CS \rightarrow X$  extending  $\alpha'\alpha''$  and  $\alpha\alpha',$  respectively.

**Definition 3.5.** With the above notation, the *Toda bracket*  $\langle \alpha, \alpha', \alpha'' \rangle$  is the set of maps

$$\Sigma S \simeq CS \amalg_S CS \xrightarrow{(\mu \circ (X' \wedge \alpha) \circ F, \mu'' \circ (X'' \wedge G) \circ C\alpha'')} X.$$

It is helpful to represent this using the following diagram.

$$\begin{array}{ccccccc} CS & & & & & & \\ & \nearrow F & & & & & \\ S & \xrightarrow{\alpha''} & X'' & \xrightarrow{\mu' \circ (X'' \wedge \alpha')} & X' & \xrightarrow{\mu \circ (X' \wedge \alpha)} & X \\ & \downarrow & \downarrow & & & & \\ CS & \xrightarrow{C\alpha''} & CX'' & \xrightarrow{\mu'' \circ (X'' \wedge G)} & & & \end{array}$$



### 3.4. Massey products.

**Definition 3.6.** Define the Massey product  $\langle a, a', a'' \rangle$  as follows:

$$\langle a, a', a'' \rangle = \{[ab' - b'a''] : d_r(b') = a'a'' \text{ and } d_r(b) = aa'\} \subseteq E_{r+1}(X).$$

The reader may notice two subtle differences from the usual definition (e.g. see [Rav86, §A.1.4]) of the Massey product. First, we have defined  $a, a'$ , and  $a''$  as cycles in the  $E_r$ -page, and are silently identifying them with their image in the  $E_{r+1}$ -page. Second, for the missing signs, recall (Remark 3.2) that we are simplifying notation by assuming that all classes have degree zero.

Note that  $b \in E_r(X)$  and  $b' \in E_r(X')$ , so  $ab' - b'a''$  is defined as an element of  $E_r(X)$  using the multiplication pairing; the basic setup of Massey products implies that it survives to an element in  $E_{r+1}(X)$ . In the next lemma, which relies on Assumption 3.3, we explain how to relate this to the notion of cycles in Section 2.2.

**Lemma 3.7.** Let  $\beta^+$  be any choice of lift in Figure 1 and  $\beta^-$  be any lift in Figure 2. Let  $\mu^+ = \alpha\beta^+$  and  $\mu^- = \beta^-\alpha''$ . Then there exists a lift  $D : S \rightarrow \Sigma^{-1}X_{f-r,r+1}$  as in Figure 4 such that  $pD = \mu^+ - \mu^-$ . The map  $D$  represents an element of  $\langle a, a', a'' \rangle \subseteq E_{r+1}(X)$ .

In the diagrams below, all columns are cofiber sequences. We make some simplifications to the notation such as labeling  $\mu \circ (X'_{f',r+1} \wedge \alpha)$  as  $\alpha$ . Recall that the products  $\alpha\beta^+$  and  $\beta^-\alpha''$  are defined as  $\mu \circ (X'_{f',r+1} \wedge \alpha) \circ \beta^+$  and  $\mu'' \circ (X''_{f'',r+1}) \circ \alpha''$ .




FIGURE 1.




FIGURE 2.




FIGURE 3.




FIGURE 4.

*Proof.* The marked nullhomotopies are such because of Lemma 3.4. First note that the lifts  $\beta^+$  and  $\beta^-$  exist because the vertical chains of maps form cofiber sequences. By construction we have  $\kappa\mu^+ = \alpha\alpha'\alpha'' = \kappa\mu^-$ , and so the lift  $D$  exists in Figure 4. By Lemma 2.13, we have  $d_r[\zeta\beta^+]_r = a'a''$  and  $d_r[\zeta\beta^-]_r = aa'$ . Thus  $\alpha(\zeta\beta^+) - (\zeta\beta^-)\alpha''$  represents an element of  $\langle a, a', a'' \rangle$ .

To relate this to  $\zeta(\mu^+ - \mu^-) = \zeta(\alpha\beta^+) - \zeta(\beta^-\alpha'')$ , we claim that  $\zeta(\alpha\beta^+) = \alpha(\zeta\beta^+)$ ; this follows from the commutativity of the following diagram.

$$\begin{array}{ccc} S & \xrightarrow{\beta^+} & \Sigma^{-1}X'_{f'-r,r+1,r} & \xrightarrow{\xi} & \Sigma^{-1}X'_{f'-r,r} \\ & & \downarrow \mu \circ (X'_{f'-r,r+1,r} \wedge \alpha) & & \downarrow \mu \circ (X'_{f'-r,r} \wedge \alpha) \\ & & \Sigma^{-1}X_{f-r,r+1,r} & \xrightarrow{\xi} & \Sigma^{-1}X_{f-r,r} \end{array}$$

To conclude, use the top right square in Figure 4 to see that  $D$  is a lift of  $\zeta(\mu^+ - \mu^-)$ , so  $D$  represents the same class in  $E_{r+1}$ .  $\blacksquare$

#### 4. COMPARISON STATEMENTS

We assume the notation in Section 3.2. The goal of this section is to prove Theorems 4.11 and 4.16, which compare the Toda bracket  $\langle \alpha, \alpha', \alpha'' \rangle$  with the Massey product  $\langle a, a', a'' \rangle$  via a third bracket which we define in Definition 4.5.

**4.1. Difference of maps.** We use the following relative definition of the difference of maps, generalizing the usual definition using the pinch map followed by the fold map  $\nabla$ . If  $q$  is  $\text{id}_Y$ , we recover the difference of two maps  $\Sigma T \rightarrow Y$ .

**Definition 4.1.** Suppose we have two maps of cofiber sequences as follows.

$$(4.2) \quad \begin{array}{ccccc} T & \longrightarrow & CT & \longrightarrow & \Sigma T \\ \downarrow h & & \downarrow f^+ & & \downarrow g^+ \\ W & \xrightarrow{j} & Y & \xrightarrow{q} & Cj \end{array} \quad \begin{array}{ccccc} T & \longrightarrow & CT & \longrightarrow & \Sigma T \\ \downarrow h & & \downarrow f^- & & \downarrow g^- \\ W & \xrightarrow{j} & Y & \xrightarrow{q} & Cj \end{array}$$

Then  $g^+ - g^- : \Sigma T \rightarrow Cj$  is the composition

$$CT \amalg_T CT \xrightarrow{\text{pinch}} CT \amalg_T CT \amalg_T CT \xrightarrow{(f^+, C(h), f^-)} Y \amalg_W CW \amalg_W Y \xrightarrow{\nabla} Y \amalg_W CW.$$

The following lemma is the crucial step comparing the Massey product constructed in Lemma 3.7, which is a difference of maps, and the definition of the Toda bracket (Definition 3.5), which is defined by gluing two maps  $CS \rightarrow X$  which agree as maps  $S \rightarrow X$ .

**Lemma 4.3.** *Suppose we have two maps of cofiber sequences (4.2). Let  $f : \Sigma T \rightarrow Y$  be the induced map on the homotopy pushout  $CT \amalg_T CT \xrightarrow{(f^+, f^-)} Y$ . Then  $g^+ - g^- = qf$ .*

*Proof.* The difference map comes from a map of diagrams

$$(4.4) \quad \begin{array}{ccc} T \rightarrow CT & \xrightarrow{f^+, f^-} & CW \rightarrow CW \amalg_W Y \\ \downarrow & \downarrow f^- & \downarrow \\ CT \xrightarrow{f^+} Y & & Y \amalg_W CW \rightarrow Y \amalg_W CW \end{array}$$

where the map on bottom left corners is  $CT \hookrightarrow CT \amalg_T CT \xrightarrow{f^+} CW \amalg_W Y$  and the map on top right corners is  $CT \hookrightarrow CT \amalg_T CT \xrightarrow{f^-} CW \amalg_W Y$ . In each square in (4.4) there is a map from the homotopy pushout to the bottom right corner, and these commute to form

$$\begin{array}{ccc} CT \amalg_T CT & \longrightarrow & Y \amalg_W CW \amalg_W Y \\ f \downarrow & & \downarrow \nabla \\ Y & \xrightarrow{q} & Y \amalg_W CW \end{array}$$

where the top right composition is  $g^+ - g^-$ . ■

**4.2. Comparison statements.** We use the setup in Section 3.2 for  $r \geq 1$ , including Assumption 3.3. Lemma 3.4 allows us to define the following Toda brackets.

**Definition 4.5.**

- Let  $\langle \alpha, \alpha', \alpha'' \rangle \subseteq \pi_*(X)$  denote the multiplicative Toda bracket of the classes  $S \xrightarrow{\alpha''} X''$ ,  $S \xrightarrow{\alpha'} X'$ , and  $S \xrightarrow{\alpha} X$ .
- Let  $\langle \alpha, \alpha', \alpha'' \rangle_*$  denote the multiplicative Toda bracket of the classes  $S \xrightarrow{\alpha''} X''_{f''}$ ,  $S \xrightarrow{\alpha'} X'_{f'}$ , and  $S \xrightarrow{\alpha} X_{f-f}$ .

**Remark 4.6.** These are subsets of the composition Toda brackets associated to the compositions  $S \xrightarrow{\alpha''} X'' \xrightarrow{\alpha'} X' \xrightarrow{\alpha} X$  and  $S \xrightarrow{\alpha''} X''_{f''} \xrightarrow{\alpha'} X'_{f'} \xrightarrow{\alpha} X_{f-f}$ , respectively.

**Proposition 4.7.** For every element  $\zeta : \Sigma S \rightarrow X_{f-r}$  of  $\langle \alpha, \alpha', \alpha'' \rangle_*$ , the image of  $\zeta$  in  $\pi_{*+1}(X_{f-r,r+1})$  represents an element in the Massey product  $\langle a, a', a'' \rangle$ .

*Proof.* Let  $F, G$  be any choice of nullhomotopies for the multiplicative Toda bracket  $\langle \alpha, \alpha', \alpha'' \rangle_*$  (defined as in Definitions 3.5, 4.5).

We use Lemma 3.7 to construct an element of the Massey product; it suffices to construct maps  $\beta^+, \beta^-$  in a way that is compatible with  $F, G$ . Consider the following diagrams, made out of maps of vertical cofiber sequences.

$$\begin{array}{ccc}
 S \xrightarrow{\tilde{\beta}^+} \Sigma^{-1} X'_{f'-r,r} \longrightarrow \Sigma^{-1} X'_{f'-r,r+1,r} & & S \xrightarrow{\tilde{\beta}^-} \Sigma^{-1} X_{f-f''-r,r} \longrightarrow \Sigma^{-1} X_{f-f''-r,r+1,r} \\
 \parallel \downarrow \kappa & & \parallel \downarrow \kappa \\
 S \xrightarrow{\alpha' \alpha''} X'_{f'} \longrightarrow X'_{f',r+1} & & S \xrightarrow{\alpha \alpha'} X_{f-f''} \longrightarrow X_{f-f'',r+1} \\
 \downarrow & & \downarrow \\
 CS \xrightarrow{F} X'_{f'-r} \longrightarrow X'_{f'-r,r+1} & & CS \xrightarrow{G} X_{f-f''-r} \longrightarrow X_{f-f''-r,r+1}
 \end{array}$$

We claim that  $\beta^+$  in Figure 1 can be taken to be the composition across the top row of the first diagram above: indeed, composing with  $\kappa$  yields  $\alpha'' \alpha'$ . Similarly, we take  $\beta^-$  in Figure 3 to be the composition across the top row of the second diagram.

That is, we have a diagram of cofiber sequences as follows.

$$(4.8) \quad
 \begin{array}{ccccc}
 S & \longrightarrow & X_f & \longrightarrow & X_{f,r+1} \\
 \downarrow & & \downarrow i & & \downarrow \\
 CS & \xrightarrow[\alpha F]{G \alpha''} & X_{f-r} & \longrightarrow & X_{f-r,r+1} \\
 \downarrow & & \downarrow p & & \downarrow \\
 \Sigma S & \xrightarrow[\Sigma \tilde{\beta}^- \alpha'']{\Sigma \alpha \tilde{\beta}^+} & X_{f-r,r} & \longrightarrow & X_{f-r,r+1,r}
 \end{array}$$

In the notation of Lemma 3.7 with the choices of  $\beta^+, \beta^-$  above, we have that

$$\Sigma(\mu^+ - \mu^-) = \Sigma(\alpha \beta^+ - \beta^- \alpha'') : \Sigma S \rightarrow X_{f-r,r+1,r}$$

is the difference of the two compositions across the third row, and so any lift  $D : \Sigma S \rightarrow X_{f-r,r+1}$  of this difference is an element of the Massey product.

Since the Toda bracket  $\zeta$  is the map

$$\Sigma S \simeq CS \amalg_S CS \xrightarrow{(\alpha F, G\alpha'')} X_{f-r},$$

by taking  $T$  to be  $S$  and the bottom cofiber sequence to be  $X_f \xrightarrow{i} X_{f-r} \xrightarrow{p} X_{f-r,r}$  in Lemma 4.3, we have

$$\Sigma(\alpha\tilde{\beta}^+ - \tilde{\beta}^-\alpha'') = p\zeta.$$

Therefore the image of  $\zeta$  along  $\pi_{*+1}(X_{f-r}) \rightarrow \pi_{*+1}(X_{f-r,r+1})$  is a lift and thus an element of the Massey product.  $\blacksquare$

**Remark 4.9.** If  $\alpha'' : \Sigma^n S \rightarrow X''_{f''}$  has degree  $n$ , then  $G\alpha'' : \Sigma^n S \wedge CS \rightarrow X_{f-r}$  is homotopic but not equal to the map  $C(\Sigma^n S) \rightarrow X_{f-r}$  required in Definition 3.5, and the induced pushout maps  $\Sigma^n S \wedge \Sigma S \rightarrow X_{f-r,r}$  and  $\Sigma^{n+1} S \rightarrow X_{f-r,r}$  as in (4.8) differ by a factor of  $(-1)^n$ . This explains why Toda brackets have signs corresponding to those in Massey products [Rav86, Definition A1.4.1].

**Proposition 4.10.** *For every element  $\Sigma S \rightarrow X_{f-r}$  in  $\langle \alpha, \alpha', \alpha'' \rangle_*$ , composing with the map  $X_{f-r} \rightarrow X$  yields an element of  $\langle \alpha, \alpha', \alpha'' \rangle$ .*

*Proof.* An element of  $\langle \alpha, \alpha', \alpha'' \rangle_*$  is defined using nullhomotopies  $F_* : CS \rightarrow X'_{f'-r}$  factoring through  $\alpha'\alpha''$  and  $G_* : CS \rightarrow X_{f-f''-r}$  factoring through  $\alpha\alpha'$ . Let  $F : CS \rightarrow X'_{f'-r} \rightarrow X'$  and  $G : CS \rightarrow X_{f-f''-r} \rightarrow X$  denote the compositions. Then the pair  $(F, G)$  defines an element in  $\langle \alpha, \alpha', \alpha'' \rangle$ .  $\blacksquare$

Now we obtain the main theorem.

**Theorem 4.11.** *Suppose  $\alpha\alpha' = 0 = \alpha'\alpha''$  and  $aa' = 0 = a'a''$  where  $a \in E_r^{f-f'}(X)$  is an  $E_r$ -representative of  $\alpha \in \pi_*(X)$ , and similarly for  $a'$  and  $a''$ . Given Assumption 3.3, there exists an element of the Massey product  $\langle a, a', a'' \rangle$  that is a permanent cycle converging to an element of  $\langle \alpha, \alpha', \alpha'' \rangle$ .*

*Proof.* Combine Proposition 4.7 and Proposition 4.10.  $\blacksquare$

**4.3. Toda brackets in the  $E_1$ -page.** In the previous section, we worked with  $E_r$ -page classes  $a, a', a''$  for  $r \geq 1$  such that  $aa' = 0 = a'a''$  in  $E_{r+1}(X)$ . In this section, we consider  $\alpha, \alpha', \alpha''$  with  $\alpha\alpha' = 0 = \alpha'\alpha''$  that are detected by  $a, a', a''$  in  $E_1^{f-f'}(X)$ ,  $E_1^{f'-f''}(X')$ , and  $E_1^{f''}(X'')$ , respectively, such that  $aa' = 0 = a'a''$  in  $E_1(X)$ .

In this case, we cannot define a corresponding Massey product. Instead, we recall that  $E_1^f(X) = \pi_*(X_{f,1})$  by definition, and our setup allows us to consider Toda brackets of such homotopy elements. We still make Assumption 3.3 at the  $E_0$ -page in the degrees of  $aa'$  and  $a'a''$ , which does not actually require a definition of  $E_0$ -pages; this is restated below.

**Assumption 4.12.** Let  $n_1$  be the stem of  $aa'$  and let  $n_2$  be the stem of  $a'a''$ . Assume that every element in  $E_{f-f''-m+1}^{n_1+1,m}$  for  $0 \leq m \leq f-f''-1$  and  $E_{f'-m+1}^{n_2+1,m}$  for  $0 \leq m \leq f'-1$  is a permanent cycle.

The goal of this section is to prove Theorem 4.16, which is the analogue of Theorem 4.11 where the Massey product is replaced by a Toda bracket of homotopy classes representing  $E_1$ -page elements. More precisely, let  $\langle a, a', a'' \rangle_{*,*}$  denote the multiplicative Toda bracket of the classes  $S \xrightarrow{a''} X''_{f'',1}$ ,  $S \xrightarrow{a'} X'_{f'-f'',1}$ , and  $S \xrightarrow{a} X_{f-f',1}$ . This is a subset of the composition Toda bracket associated to

$$S \xrightarrow{a''} X''_{f'',1} \xrightarrow{a'} X'_{f',1} \xrightarrow{a} X_{f,1}.$$

Proposition 2.12 does not make sense in this case; the replacement is Lemma 4.13.

**Lemma 4.13.** *Suppose  $\zeta \in \pi_n(X_f)$  is a class whose images in both  $\pi_n(X)$  and  $\pi_n(X_{f,1})$  are zero. Suppose  $E_r(X)$  satisfies the crossing differentials hypothesis at the  $E_0$ -page in degree  $(n, f)$ . Then  $\zeta = 0$ .*

*Proof.* If not, then there exists  $i \leq f$  such that the image of  $\zeta$  in  $\pi_n(X_i)$  is nonzero but its image in  $\pi_n(X_{i-1})$  is zero. This means that there exists  $z \in \pi_{n+1}(X_{i-1,1})$  such that  $\kappa(z)$  lifts to  $\zeta \in \pi_n(X_f)$ . Moreover, since  $p(\zeta) \in \pi_n(X_{f,1})$  is zero,  $\zeta$  lifts to a class in  $\pi_n(X_{f+1})$ . That is,  $z$  supports a nontrivial differential of length  $\geq 2$ , contradicting the crossing differentials hypothesis.  $\blacksquare$

**Proposition 4.14.** *The compositions  $S \rightarrow X''_{f''} \rightarrow X'_{f'}$  and  $S \rightarrow X'_{f'-f''} \rightarrow X_{f-f'}$  are null. For every element  $\Sigma S \rightarrow X_f$  in  $\langle \alpha, \alpha', \alpha'' \rangle_{*,*}$ , composing with the map  $X_f \rightarrow X$  yields an element of  $\langle \alpha, \alpha', \alpha'' \rangle$ .*

*Proof.* This is analogous to Proposition 4.10, using Lemma 4.13 instead of Proposition 2.12.  $\blacksquare$

**Proposition 4.15.** *Given Assumption 4.12, every class in  $\langle \alpha, \alpha', \alpha'' \rangle_*$  projects to a class in  $\langle a, a', a'' \rangle_{*,*}$ .*

*Proof.* Given nullhomotopies  $F_* : CS \rightarrow X'_{f'}$  and  $G_* : CS \rightarrow X_f$  as in Definition 3.5 applied to  $\langle \alpha, \alpha', \alpha'' \rangle$ , compose to get nullhomotopies  $F_{*,*} : CS \rightarrow X'_{f'} \rightarrow X'_{f',1}$  and  $G_{*,*} : CS \rightarrow X_f,1$  which in turn define a compatible element of  $\langle a, a', a'' \rangle_{*,*}$ .  $\blacksquare$

**Theorem 4.16.** *Suppose  $\alpha\alpha' = 0 = \alpha'\alpha''$  and  $aa' = 0 = a'a''$  where  $a \in \pi_*(X_{f-f',1})$  is an  $E_1$  representative of  $\alpha \in \pi_*(X)$ , and similarly for  $a'$  and  $a''$ . Given Assumption 4.12, there is a permanent cycle in  $\langle a, a', a'' \rangle_{*,*}$  converging to an element of  $\langle \alpha, \alpha', \alpha'' \rangle$ .*

*Proof.* Combine Propositions 4.15 and 4.14.  $\blacksquare$

## 5. EXAMPLES

Our motivation for writing this paper is to study Toda brackets in the  $\mathbb{R}$ -motivic slice spectral sequence for the sphere at  $p = 2$ . This is the spectral sequence associated to Voevodsky's effective slice filtration: given an effective motivic spectrum  $X$ , there is a tower

$$\begin{array}{ccccccc} X = f_0X & \longleftarrow & f_1X & \longleftarrow & f_2X & \cdots & \\ \downarrow & & \downarrow & & & & \\ s_0X & & s_1X & & \cdots & & \end{array}$$

which gives rise to the slice spectral sequence. By [Pel11], if  $X$  is a ring spectrum, we get a multiplicative pairing of spectral sequences  $E_r(X) \otimes E_r(X) \rightarrow E_r(X)$ . The quotients  $s_qX$  are called *slices*. See [Lev13, RSØ19] for more details.

Røndigs, Spitzweck, and Østvær [RSØ19] calculated the slices of the motivic sphere spectrum:

$$(5.1) \quad s_*(S) = H\mathbb{Z} \otimes \text{Ext}_{MU_*MU}^{*,*}(MU_*, MU_*)$$

where the notation means that a copy of  $\mathbb{Z}/2^n$  in the Ext term corresponds to an (appropriately degree-shifted)  $H\mathbb{Z}/2^n$  summand, where  $H\mathbb{Z}/2^n$  is the motivic Eilenberg–MacLane spectrum. In the identification (5.1), (higher) multiplicative data about the Adams–Novikov  $E_2$ -term  $\text{Ext}_{MU_*MU}^{*,*}(MU_*, MU_*)$  carries over to the slice  $E_1$ -page  $\pi_{*,*}(s_*(S))$ . In practice, we work one prime at a time, and focus on the  $p = 2$  case. Combining results of [RSØ19] and Hu, Kriz, and Ormsby [HKO11], the 2-completed slice spectral sequence for the  $\mathbb{R}$ -motivic sphere converges to the homotopy groups of the 2-completed sphere, which we denote by  $\pi_{*,*}^{\mathbb{R}}(S)$ .

**Example 5.2.** There is a strictly defined Adams–Novikov Massey product  $\langle \alpha_1, 2, \alpha_1 \rangle = 2\alpha_{2/2}$ . The class  $\alpha_1$  is a permanent cycle converging to  $\eta \in \pi_{1,1}^{\mathbb{R}}(S)$  and  $\alpha_{2/2}$  is a permanent cycle converging to  $\nu \in \pi_{3,2}^{\mathbb{R}}(S)$ . We have  $2 \cdot \alpha_1 = 0$  in the slice  $E_1$ -page but  $2 \cdot \eta$  is nonzero in  $\pi_{*,*}^{\mathbb{R}}(S)$ . However, there is a class  $\omega = 2 + \rho\eta$  detected by 2 in the slice spectral sequence (as  $\rho\eta$  has higher slice filtration) such that  $\omega\eta = 0 = \omega\rho$ . Hence there is a Toda bracket  $\langle \alpha_1, 2, \alpha_1 \rangle$  in the slice  $E_1$ -page containing  $2\alpha_{2/2}$  and there is a corresponding Toda bracket  $\langle \eta, \omega, \eta \rangle$  in homotopy that is also strictly defined. Then Theorem 4.16 implies that  $2\alpha_{2/2}$  is a permanent cycle in the  $\mathbb{R}$ -motivic slice spectral sequence converging to  $\omega\nu$  which is contained in the homotopy Toda bracket  $\langle \eta, \omega, \eta \rangle$ .

While this example is in low enough stems that the Toda bracket argument is not needed, this argument applies to a wealth of other Adams–Novikov Massey products.

**Example 5.3.** In  $\pi_{*,*}^{\mathbb{R}}(HZ/2^n)$ , there is a strictly defined Toda bracket  $\langle \rho, 2, 2^{n-1} \rangle = 2^{n-1}\tau$ , which can be seen using the spectral sequence associated to the cofiber sequence of motivic Eilenberg-MacLane spectra

$$HZ/2 \rightarrow HZ/2^n \rightarrow HZ/2^{n-1}.$$

This is another source of slice  $E_1$ -page Toda brackets: if  $x \in \text{Ext}_{BP_*BP}^{*,*}(BP_*, BP_*)$  generates a copy of  $\mathbb{Z}/2^n$ , then one may consider the  $E_1$ -page Toda bracket  $\langle \rho, 2, 2^{n-1}x \rangle \ni 2^{n-1}\tau x$ . For example, there is a strictly defined  $E_1$ -page Toda bracket  $\langle \rho, 2, \alpha_1 \rangle = \tau\alpha_1$ . Using the notation in Example 5.2, Theorem 4.16 implies that the  $\pi_{*,*}^{\mathbb{R}}(S)$  Toda bracket  $\langle \rho, \omega, \eta \rangle$  contains  $\tau\eta$ .

**Example 5.4.** There are also higher slice Massey products which converge to Toda brackets. For example, the slice differential  $d_1(\tau^2) = \rho^2\tau\alpha_1$  implies that  $\langle 2, \rho^2, \tau\alpha_1 \rangle$  is strictly defined as a slice  $E_2$ -page Massey product consisting of  $2\tau^2$ . The products  $\omega \cdot \rho^2$  and  $\rho^2 \cdot \tau\eta$  are zero in homotopy (see Example 5.2), and so the Toda bracket  $\langle \omega, \rho^2, \tau\eta \rangle$  in  $\pi_{*,*}^{\mathbb{R}}(S)$  is defined and contains  $\omega\tau^2$ . Then we may deduce the hidden multiplication  $\eta \cdot \omega\tau^2$  using a shuffle argument as follows. We have

$$\eta \cdot \langle \omega, \rho^2, \tau\eta \rangle = \langle \eta, \omega, \rho^2 \rangle \tau\eta.$$

By Example 5.3, the right hand side contains  $\rho(\tau\eta)^2$ . Moreover, the right hand side is strictly defined:  $\langle \eta, \omega, \rho^2 \rangle$  contains indeterminacy generated by  $\rho^3\nu$ , but this is annihilated by  $\tau\eta$ , since  $\rho^2\tau\eta = 0$ . Thus we have

$$\eta \cdot \omega\tau^2 = \rho(\tau\eta)^2.$$

## REFERENCES

- [CF17] J. Daniel Christensen and Martin Frankland. Higher Toda brackets and the Adams spectral sequence in triangulated categories. *Algebr. Geom. Topol.*, 17(5):2687–2735, 2017.
- [HKO11] P. Hu, I. Kriz, and Kyle Ormsby. Convergence of the motivic Adams spectral sequence. *Journal of K-theory: K-theory and its Applications to Algebra, Geometry, and Topology*, 7:573 – 596, 06 2011.
- [Isa19] Daniel Isaksen. Stable stems. *Memoirs of the American Mathematical Society*, 262(1269):0–0, nov 2019.
- [IWX20] Daniel C. Isaksen, Guozhen Wang, and Zhouli Xu. More stable stems, 2020.
- [Koc80] Stanley O. Kochman. Uniqueness of Massey products on the stable homotopy of spheres. *Canadian J. Math.*, 32(3):576–589, 1980.
- [Koc90] Stanley O. Kochman. *Stable homotopy groups of spheres*, volume 1423 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1990. A computer-assisted approach.
- [Lev13] Marc Levine. Convergence of Voevodsky’s slice tower. *Doc. Math.*, 18:907–941, 2013.
- [Mas58] W. S. Massey. Some higher order cohomology operations. In *Symposium internacional de topología algebraica International symposium on algebraic topology*, pages 145–154. Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958.
- [Mos70] R. Michael F. Moss. Secondary compositions and the Adams spectral sequence. *Math. Z.*, 115:283–310, 1970.

- [Pel11] Pablo Pelaez. *Multiplicative properties of the slice filtration*. Number 335 in Astérisque. Société mathématique de France, 2011.
- [Rav86] Douglas C. Ravenel. *Complex cobordism and stable homotopy groups of spheres*, volume 121 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1986.
- [RSØ19] Oliver Röndigs, Markus Spitzweck, and Paul Arne Østvær. The first stable homotopy groups of motivic spheres. *Ann. of Math. (2)*, 189(1):1–74, 2019.
- [Tod62] Hirosi Toda. *Composition methods in homotopy groups of spheres*. Annals of Mathematics Studies, No. 49. Princeton University Press, Princeton, N.J., 1962.