

The $21N$ -moment multi-temperature coefficients for Zhdanov closure

M. Raghunathan and Y. Marandet

Aix-Marseille Univ., CNRS, PIIM, Marseille, France

H. Bufferand, G. Ciraolo, Ph. Ghendrih, and P. Tamain

IRFM-CEA, F-13108 Saint-Paul-Lez-Durance, France

E. Serre

Aix-Marseille Univ., CNRS, M2P2, Marseille, France

We provide the $21N$ -moment multi-temperature collision coefficients for the Boltzmann collision operator using the Sonine-Hermite polynomial ansatz in the style of Zhdanov et al. First, we outline the general derivation method. Then, we provide the collision coefficients in the most general form in terms of the Chapman-Cowling integrals for any potential of interaction for the $21N$ -moment approximation. Then, we provide the collision coefficients for the specific case of the approximated Coulomb potential cross-section with Debye cutoff. These coefficients help in order to find easy implementation in current SOL/edge fluid packages which currently implement the $21N$ -moment single-temperature coefficients. This provides the completion of a missing link in the current literature.

I. INTRODUCTION

Understanding impurity dynamics in the scrape-off layer (SOL) and edge of tokamaks remains a key factor in improving the performance of current and next-step machines such as WEST, JET and ITER. For example, reducing the peak heat fluxes on plasma facing components to manageable levels relies on a large extent on impurity radiation, which in turn depends on the impurity density, flow, and temperature. Generally, the SOL/edge is modeled as a high-collisionality regime, and the plasma is generally simulated with fluid packages, normally coupled to kinetic neutrals, e.g. Soledge3x-EIRENE[1]. Generally, the collision terms used in the fluid packages are found by averaging over the kinetic equation with different constants of motion. However, the procedure of such averaging typically leaves the system of equations obtained unclosed. Closing the system of equations is the procedure for obtaining these unclosed quantities, such as the collisional friction and thermal forces, in terms of the plasmadynamical moments such as density, flow velocity and temperature.

One such method of closure is called the Zhdanov closure[2, 3], which involves using Grad's method[4] with a Sonine-Hermite polynomial ansatz for the distribution functions and the moments. On averaging over the Boltzmann kinetic equation with the moments, and assuming low Knudsen number, one finds prescription for the friction and thermal forces in terms of the flow velocities and the temperature gradients. The advantage of this closure is that it does not depend on any trace density assumptions or low temperature difference assumptions, as opposed to earlier closures such as Braginskii's[5]. This is important for modelling impurity dynamics because impurities may be present as a significant fraction of the background plasma species, or, in case of deuterium-tritium plasmas, tritium, being a main ion species, cannot be modelled as trace over an existing deuterium-electron plasma. The Zhdanov closure is already implemented, though only using single-temperature collision coefficients at the plasma common temperature, in SOL/edge simulators such as Soledge3x-EIRENE[1, 6, 7], SOLPS[8–10], B2-EIRENE[11], and EDGE2D[12].

Generally, the depth of the closure is represented by the number of moments, i.e. $5N$ -moment, $13N$ -moment, $21N$ -moment or $29N$ -moment, where N represents the number of species. In Refs. [13] and [14], we found that generally, the multi-temperature $21N$ -moment scheme exhibits values of transport coefficients and friction/thermal forces about 20-40% different from the single-temperature scheme, and furthermore, is adequate for obtaining converged values of friction and thermal forces for fusion-relevant scenarios. However, the collision coefficients for the $21N$ -moment multi-temperature Zhdanov closure scheme have not been provided explicitly in any literature. Thus, in this article, we provide the $21N$ -moment multi-temperature collisional coefficients derived from the Boltzmann collision operator to fill the existing gap. We hope that the availability of these coefficients will lead to a straightforward implementation in current SOL/edge fluid packages.

The article is organized as follows. Sections II and III give an overview of the moment averaging process over the collision operator using the Sonine-Hermite polynomial ansatz, describes all the relevant terminology, and derives expressions for the collision coefficients. In Section IV, we provide the values of the multi-temperature collision coefficients for the $21N$ -moment scheme in the most general form in terms of Chapman-Cowling integrals, which are applicable to any potential of interaction. In Section V, we provide the values of the $21N$ -moment collision coefficients for the Coulomb potential with Debye cutoff approximation. With these values, one should be able to implement Zhdanov closure for the multi-temperature case. In Section VI, we summarize the article.

II. THE BOLTZMANN EQUATION FOR MULTI-SPECIES PLASMA

The Boltzmann equation for the distribution function for species α , f_α in the frame of the peculiar velocity of species α , $\mathbf{c}_\alpha = \mathbf{v}_\alpha - \mathbf{u}$, is given by

$$\frac{df_\alpha}{dt} + \mathbf{c}_\alpha \cdot \nabla f_\alpha + \frac{1}{m_\alpha} \mathbf{F}_\alpha^* \cdot \nabla_{\mathbf{c}_\alpha} f_\alpha - c_{\alpha s} \frac{\partial f_\alpha}{\partial c_{\alpha r}} \frac{\partial u_r}{\partial x_s} = \sum_\beta J_{\alpha\beta}, \quad (1)$$

where the common plasma flow velocity \mathbf{u} is given by

$$\rho \mathbf{u} = \sum_\alpha \rho_\alpha \mathbf{u}_\alpha, \quad \rho = \sum_\alpha \rho_\alpha, \quad (2)$$

where ρ represents the mass density. The d/dt represents full time derivative given by $d/dt = \partial/\partial t + \mathbf{u} \cdot \nabla$, and where the force term \mathbf{F}_α and $d\mathbf{u}/dt$ are combined to write the relative force in the moving frame given by $\mathbf{F}_\alpha^* = \mathbf{F}_\alpha - m_\alpha d\mathbf{u}/dt$.

The LHS is referred to as the ‘‘free-streaming term’’, and the RHS is the collisional contribution between species α and every other species of the system. The collisional RHS is represented by the Boltzmann collision integral,

$$J_{\alpha\beta} = \iint (f'_\alpha f'_{1\beta} - f_\alpha f_{1\beta}) g \sigma_{\alpha\beta}(g, \chi) d\Omega d\mathbf{c}_{1\beta}, \quad (3)$$

where $\alpha, 1\beta$ refer to species of the two particles colliding, subscript l refers to properties after the collision, g is the relative velocity between the colliding particles, $\sigma_{\alpha\beta}$ is the collision cross section, and Ω is the solid angle in which the collision occurs.

Now, for any quantity ψ_α depending purely on species α , one can average over the RHS of Eq. 1 which attains the following form

$$R_\alpha = \sum_\beta \int \psi_\alpha J_{\alpha\beta} d\mathbf{v}_\alpha = \sum_\beta \iiint \psi_\alpha (f'_\alpha f'_{1\beta} - f_\alpha f_{1\beta}) g \sigma_{\alpha\beta}(g, \chi) d\Omega d\mathbf{c}_\alpha d\mathbf{c}_{1\beta}. \quad (4)$$

For elastic collisions, the moment averaged collision operator can be transformed into

$$R_\alpha = \sum_\beta \iiint (\psi'_\alpha - \psi_\alpha) f_\alpha f_{1\beta} g \sigma_{\alpha\beta}(g, \chi) d\Omega d\mathbf{c}_\alpha d\mathbf{c}_{1\beta}, \quad (5)$$

since the distribution functions for any given species are purely a function of the species peculiar velocity.

We now describe the modification of Grad’s method[4] as used by Zhdanov[15] in his previous papers. In the ansatz for the solution of the Boltzmann equation, it is assumed that the solution f_α is already near thermodynamic equilibrium for species α , $f_\alpha^{(0)}$ as follows

$$\begin{aligned} f_\alpha^{(0)} &= n_\alpha \left(\frac{m_\alpha}{2\pi k T_\alpha} \right)^{3/2} \exp\left(-\frac{m_\alpha c_\alpha^2}{2k T_\alpha} \right) \\ &= n_\alpha \left(\frac{\gamma_\alpha}{2\pi} \right)^{3/2} \exp\left(-\frac{\gamma_\alpha c_\alpha^2}{2} \right), \end{aligned} \quad (6)$$

where $\gamma_\alpha = m_\alpha/kT_\alpha$, n_α is the number density, and T_α the temperature of species α . In order to solve the Boltzmann equation (1), Zhdanov and Yushmanov choose an ansatz of the form

$$f_\alpha = f_\alpha^{(0)} \sum_{m,n} 2^{2n} m_\alpha^{-2} \gamma_\alpha^{2n+m} \tau_{mn} b_{\alpha r_1 \dots r_m}^{mn} G_{\alpha r_1 \dots r_m}^{mn}, \quad (7)$$

where

$$G_\alpha^{mn}(\mathbf{c}_\alpha, \gamma_\alpha) = (-1)^n n! m_\alpha \gamma_\alpha^{-(n+m/2)} S_{m+1/2}^n \left(\frac{\gamma_\alpha}{2} \mathbf{c}_\alpha^2 \right) P^{(m)}(\gamma_\alpha^{1/2} \mathbf{c}_\alpha). \quad (8)$$

Here, $S_{m+1/2}^n$ are the Sonine polynomials, given by,

$$S_{m+1/2}^n \left(\frac{\gamma_\alpha}{2} \mathbf{c}_\alpha^2 \right) = \sum_{p=0}^n \left(-\frac{\gamma_\alpha}{2} \mathbf{c}_\alpha^2 \right)^p \frac{(m+n+1/2)!}{p!(n-p)!(m+p+1/2)!}.$$

Further, $P^{(m)}$ are the irreducible projection of the tensorial monomial $\mathbf{c}_\alpha^m = c_{\alpha r_1} \dots c_{\alpha r_m}$, derived by the following recurrence relation

$$P^{(m+1)}(\gamma_\alpha^{1/2} \mathbf{c}_\alpha) = \gamma_\alpha^{1/2} \mathbf{c}_\alpha P^{(m)} - \gamma_\alpha^{1/2} \frac{c_\alpha^2}{2m+1} \frac{\partial P^{(m)}}{\partial \mathbf{c}_\alpha},$$

with $P^{(0)} = 1$. The constant τ_{mn} arises as a result of internal contractions between b_α^{mn} and G_α^{mn} , and is given by

$$\tau_{mn} = \frac{(2m+1)!(m+n)!}{n!(m!)^2(2m+2n+1)!}.$$

The forms mentioned in Refs. [4] and [15] are cosmetically different because of the choice to use full factorial representations of functions and because of summing over full indices rather than over half-indices, but they are equivalent. The coefficients b_α^{mn} are calculated as

$$n_\alpha b_\alpha^{mn} = \int G_\alpha^{mn} f_\alpha d\mathbf{c}_\alpha. \quad (9)$$

Some values of G_α^{mn} are as follows

$$\begin{aligned} G_\alpha^{00} &= m_\alpha, \quad G_\alpha^{10} = m_\alpha \mathbf{c}_\alpha, \quad G_\alpha^{01} = \frac{m_\alpha}{2} \left(c_\alpha^2 - \frac{3}{\gamma_\alpha} \right), \\ G_\alpha^{11} &= \frac{m_\alpha}{2} \mathbf{c}_\alpha \left(c_\alpha^2 - \frac{5}{\gamma_\alpha} \right), \quad G_\alpha^{20} = m_\alpha \left(\mathbf{c}_\alpha \mathbf{c}_\alpha - \frac{1}{3} U c_\alpha^2 \right) \\ G_\alpha^{12} &= \frac{m_\alpha}{4} \mathbf{c}_\alpha (c_\alpha^4 - 14\gamma_\alpha^{-1} c_\alpha^2 + 35\gamma_\alpha^{-12}), \\ G_\alpha^{21} &= \frac{m_\alpha}{2} (c_\alpha^2 - 7\gamma_\alpha^{-1}) \left(\mathbf{c}_\alpha \mathbf{c}_\alpha - \frac{1}{3} U c_\alpha^2 \right), \end{aligned}$$

and the corresponding b_α^{mn} are given by

$$\begin{aligned} n_\alpha b_\alpha^{00} &= \rho_\alpha, \quad n_\alpha b_\alpha^{10} = \rho_\alpha \mathbf{w}_\alpha, \quad n_\alpha b_\alpha^{01} = 0, \\ n_\alpha b_\alpha^{11} &= \mathbf{h}_\alpha, \quad n_\alpha b_\alpha^{20} = \pi_\alpha, \\ n_\alpha b_\alpha^{12} &= \mathbf{r}_\alpha, \quad n_\alpha b_\alpha^{21} = \sigma_\alpha. \end{aligned}$$

Here, b_α^{00} , b_α^{01} and b_α^{10} represent the intuitive hydrodynamical moments density, diffusion velocity and temperature ρ_α , $\mathbf{w}_\alpha = \mathbf{u}_\alpha - \mathbf{u}$, and T_α . the higher moments b_α^{11} and b_α^{20} represent the thermodynamically privileged moments (as per Balescu's nomenclature[16]), the heat flux \mathbf{h}_α and the divergence free pressure-stress tensor π_α , which are called "privileged" because they contribute to the entropy production. The higher-order moments b_α^{12} and b_α^{21} are non-privileged moments \mathbf{r}_α and σ_α , which do not have a clear physical meaning, however which may contribute to the accuracy of moment equations in terms of representing the Boltzmann equation. As one can notice, these are all moments of ranks less than 2. Moments of rank-0 are scalar, like density ρ_α and temperature T_α , are constitute N variables each. Moments rank-1 are vectorial moments, like momentum $m_\alpha \mathbf{w}_\alpha$ and heat-flux \mathbf{h}_α , and contribute $3N$ variables each. Moments of rank-2 are tensorial in nature, like the stress-tensor π_α and σ_α , and contribute $5N$ variables each (and not $9N$, since they are symmetric and traceless). In principle, one can construct a $5N$ -moment system with just the hydrodynamical moments, a $13N$ -system with including the thermodynamically privileged moments, and a $21N$ -system including \mathbf{r}_α and σ_α .

III. DERIVATION OF THE RIGHT HAND SIDE OF THE BOLTZMANN EQUATION

In the Boltzmann collision integral $J_{\alpha\beta}$, it is possible to choose a distribution function which takes the form

$$f_\alpha = f_\alpha^{(0)}(1 + \Phi_\alpha),$$

which essentially represents the ansatz as a perturbed Maxwellian. The linearized moment-averaged collision operator Eq. (5) can then be written as

$$R_{\alpha\beta} = \int \psi_\alpha J_{\alpha\beta} d\mathbf{c}_\alpha \iint f_\alpha^{(0)} f_\beta^{(0)} (\psi'_\alpha - \psi_\alpha) (1 + \Phi_\alpha + \Phi_\beta) g \sigma_{\alpha\beta}(g, \chi) d\Omega d\mathbf{c}_\alpha d\mathbf{c}_{1\beta},$$

This allows us to decompose the moment-average into sums of smaller terms, which is useful analytically. This also is similar to the properties exhibited by some other linearized operators such as the linearized Landau operator[16, 17].

On substituting the Sonine-Hermite polynomial ansatz from Eq. (7) for the distribution functions f , and set the moment $\psi = G_\alpha^{mn}$ from Eq. (8), we obtain

$$R_{\alpha\beta}^{mnkl} = \iiint f_\alpha^{(0)} f_\beta^{(0)} \{G_\alpha^{mn}(\mathbf{c}'_\alpha) - G_\alpha^{mn}(\mathbf{c}_\alpha)\} \{1 + 2^{2l} \gamma_\alpha^{2l+k} m_\alpha^{-2} \tau_{kl} G_\alpha^{kl}(\mathbf{c}_\alpha) b_\alpha^{kl} + 2^{2l} \gamma_\beta^{2l+k} m_\beta^{-2} \tau_{kl} G_\beta^{kl}(\mathbf{c}_\beta) b_\beta^{kl}\} g \sigma_{\alpha\beta}(g, \chi) d\Omega d\mathbf{c}_\alpha d\mathbf{c}_\beta,$$

where $R_{\alpha\beta}^{mnkl}$ represents the part of $R_{\alpha\beta}^{mn}$ averaging over the kl term of the ansatz Eq. (7). Noting that $G^{kl} b^{kl}$ is an inner product, we now substitute the definition of G^{mn} , and use the following integral identity [18, 19]

$$\int P^{(m)}(P^{(k)} : W) G(v) dv = \frac{W}{2m+1} \delta_{km} \int P^{(m)} : P^{(m)} G(v) dv, \quad (10)$$

where W is symmetric and traceless tensor of rank k not a function of \mathbf{v} . Furthermore, we define a ‘‘bracket’’ integrals of the following form

$$n_\alpha n_\beta [F, G] = \iiint f_\alpha^{(0)} f_\beta^{(0)} G(F' - F) g \sigma_{\alpha\beta}(g, \chi) d\Omega d\mathbf{c}_\alpha d\mathbf{c}_\beta, \quad (11)$$

through which we can contract over index k in $R_{\alpha\beta}^{mnkl}$ and write it as $R_{\alpha\beta}^{mnl}$, such that

$$R_{\alpha\beta}^{mnl} = (1 - \delta_{m0} \delta_{l0}) (A_{\alpha\beta}^{mnl} b_\alpha^{ml} + B_{\alpha\beta}^{mnl} b_\beta^{ml}) + \delta_{m0} \delta_{l0} C_{\alpha\beta}^{mnl}, \quad (12)$$

where $A_{\alpha\beta}^{mnl}$, $B_{\alpha\beta}^{mnl}$ and $C_{\alpha\beta}^{mnl}$ are given by

$$\begin{aligned} A_{\alpha\beta}^{mnl} &= Q_{\alpha\beta}^{mnl} \gamma_\alpha^{l-n} \left[S_{m+1/2}^n(W_\alpha^2) P^{(m)}(\mathbf{W}_\alpha), S_{m+1/2}^l(W_\alpha^2) P^{(m)}(\mathbf{W}_\alpha) \right] \\ B_{\alpha\beta}^{mnl} &= Q_{\alpha\beta}^{mnl} \frac{\gamma_\beta^{l+m/2}}{\gamma_\alpha^{n+m/2}} \frac{m_\alpha}{m_\beta} \left[S_{m+1/2}^n(W_\alpha^2) P^{(m)}(\mathbf{W}_\alpha), S_{m+1/2}^l(W_\beta^2) P^{(m)}(\mathbf{W}_\beta) \right] \\ C_{\alpha\beta}^{mnl} &= (-1)^n n! \gamma_\alpha^{-(n+m/2)} m_\alpha n_\alpha n_\beta \left[S_{1/2}^n(W_\alpha^2), 1 \right], \end{aligned} \quad (13)$$

where

$$Q_{\alpha\beta}^{mnl} = (-1)^{n+l} 2^{2l+m} \frac{(2m)!(m+l)!n!}{(m!)^2(2m+2l+1)!} n_\alpha n_\beta. \quad (14)$$

This moment-averaged collision operator is valid for any difference of masses or temperatures of the colliding species. The general expressions for the collision coefficients $A_{\alpha\beta}^{mnl}$, $B_{\alpha\beta}^{mnl}$, and $C_{\alpha\beta}^{mnl}$ take the following general forms

$$A_{\alpha\beta}^{mnl}, B_{\alpha\beta}^{mnl}, C_{\alpha\beta}^{mnl} \sim \sum_{rl} \mathcal{A}_{\alpha\alpha}^{pqr, m} \Omega_{\alpha\beta}^{lr}, \sum_{rl} \mathcal{A}_{\alpha\beta}^{pqr, m} \Omega_{\alpha\beta}^{lr}, \sum_{rl} \mathcal{A}^{pqr, m} \Omega_{\alpha\beta}^{lr}$$

respectively. The \mathcal{A} -coefficients are expressed as functions of mass and temperature ratios of the species α and β . The terms $\Omega_{\alpha\beta}^{lr}$ are the effective cross-section moment integrals of Chapman and Cowling (henceforth referred to as the ‘‘Chapman-Cowling integrals’’) which are dependent on the potential of interaction between species α and β . The exact derivations of the generalized coefficients $\mathcal{A}_{\alpha\beta}^{pqr, m}$ and $\mathcal{A}_{\alpha\alpha}^{pqr, m}$ for different bracket integrals up to rank-2, with all steps supplied for verification purposes, are provided in Ref. 13.

The Chapman-Cowling integral is written in the following form for our case

$$\Omega_{\alpha\beta}^{lr} = \left(\frac{2\pi}{\gamma_{\alpha\beta}} \right)^{1/2} \int_0^\infty \exp(-\zeta^2) \zeta^{2r+3} \phi_{\alpha\beta}^{(l)} d\zeta, \quad (15)$$

$$\phi_{\alpha\beta}^{(l)} = \int_0^\infty (1 - \cos^l \chi) \sigma_{\alpha\beta}(g, \chi) \sin \chi d\chi, \quad (16)$$

where $\zeta = (\gamma_{\alpha\beta}/2)^{1/2} g$.

IV. MULTI-TEMPERATURE COLLISION COEFFICIENTS IN TERMS OF THE CHAPMAN-COWLING INTEGRALS

In this section, the following notation is used in addition to the ones introduced in the previous sections,

$$\eta_{\alpha\beta} = \frac{m_\alpha}{m_\beta}, \text{ and } \theta_{\alpha\beta} = \frac{T_\alpha}{T_\beta}, \quad (17)$$

for the simplicity of representing the collision coefficients. The rank-1 coefficients up to $p, q \leq 1$ and rank-2 coefficients for $p, q = 0$ are exactly the same as the ones provided by Zhdanov et al in Chapter 4 of Ref. [15], represented slightly differently. For the case of equal temperatures, i.e. $\theta_{\alpha\beta} = 1$, they are exactly the same as the transport coefficients provided in Chapter 8 of Ref. [15] (with minor corrections which can be found in Appendix D of Ref. [13]). As mentioned earlier, the coefficients in this section are valid for any potential of interaction, and we provide the specific ones for the Coulomb potential in the next section.

A. Value of $C_{\alpha\beta}^{010}$ for $p, q = 0$

$$p = 1, q = 0 : \mathcal{A}_{\alpha\beta}^{1011,0} \Omega_{\alpha\beta}^{11}$$

$$\mathcal{A}_{\alpha\beta}^{1011,0} = -\frac{16k\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha}}{(\eta_{\alpha\beta} + 1)^2\theta_{\alpha\beta}} \quad (18)$$

B. Values of $B_{\alpha\beta}^{1pq}$ for $p, q \leq 2$

$$p = 0, q = 0 : \mathcal{A}_{\alpha\beta}^{0011,1} \Omega_{\alpha\beta}^{11}$$

$$\mathcal{A}_{\alpha\beta}^{0011,1} = \frac{16\eta_{\alpha\beta}n_{\alpha}n_{\beta}}{3\eta_{\alpha\beta} + 3} \quad (19)$$

$$p = 0, q = 1 : \mathcal{A}_{\alpha\beta}^{0111,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{0121,1} \Omega_{\alpha\beta}^{12}$$

$$\mathcal{A}_{\alpha\beta}^{0111,1} = -\frac{16\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta}}{3k(\eta_{\alpha\beta} + 1)T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})}$$

$$\mathcal{A}_{\alpha\beta}^{0121,1} = \frac{32\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta}}{15k(\eta_{\alpha\beta} + 1)T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (20)$$

$$p = 1, q = 0 : \mathcal{A}_{\alpha\beta}^{1011,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{1021,1} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{1022,1} \Omega_{\alpha\beta}^{22}$$

$$\mathcal{A}_{\alpha\beta}^{1011,1} = -\frac{40k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}(3\theta_{\alpha\beta} - 2) + \theta_{\alpha\beta})}{3(\eta_{\alpha\beta} + 1)^2 m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})}$$

$$\mathcal{A}_{\alpha\beta}^{1021,1} = \frac{16k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}^2(3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2)}{3(\eta_{\alpha\beta} + 1)^3 \theta_{\alpha\beta} m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})}$$

$$\mathcal{A}_{\alpha\beta}^{1022,1} = \frac{32k\eta_{\alpha\beta}^2(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha}}{3(\eta_{\alpha\beta} + 1)^3 \theta_{\alpha\beta} m_{\alpha}} \quad (21)$$

$$p = 1, q = 1 : \mathcal{A}_{\alpha\beta}^{1111,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{1121,1} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{1131,1} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{1122,1} \Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{1132,1} \Omega_{\alpha\beta}^{23}$$

$$\mathcal{A}_{\alpha\beta}^{1111,1} = \frac{8\eta_{\alpha\beta}\theta_{\alpha\beta}n_{\alpha}n_{\beta}(\eta_{\alpha\beta}(21\theta_{\alpha\beta} - 10) + 11\theta_{\alpha\beta})}{3(\eta_{\alpha\beta} + 1)^2(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2}$$

$$\mathcal{A}_{\alpha\beta}^{1121,1} = -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta}(\eta_{\alpha\beta}^2(21\theta_{\alpha\beta}^2 - 26\theta_{\alpha\beta} + 10) + 2\eta_{\alpha\beta}\theta_{\alpha\beta}(8\theta_{\alpha\beta} - 3) + 5\theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2}$$

$$\mathcal{A}_{\alpha\beta}^{1131,1} = \frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta}(\eta_{\alpha\beta}^2(3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2}$$

$$\mathcal{A}_{\alpha\beta}^{1122,1} = -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta}(\eta_{\alpha\beta}(7\theta_{\alpha\beta} - 5) + 2\theta_{\alpha\beta})}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})}$$

$$\mathcal{A}_{\alpha\beta}^{1132,1} = \frac{64\eta_{\alpha\beta}^2(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (22)$$

$$\begin{aligned}
p = 0, q = 2 : \mathcal{A}_{\alpha\beta}^{0211,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{0221,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{0231,1}\Omega_{\alpha\beta}^{13} \\
\mathcal{A}_{\alpha\beta}^{0211,1} = \frac{8\eta_{\alpha\beta}^3 m_{\beta}^2 n_{\alpha} n_{\beta}}{3k^2 (\eta_{\alpha\beta} + 1) T_{\beta}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{0221,1} = -\frac{32\eta_{\alpha\beta}^3 m_{\beta}^2 n_{\alpha} n_{\beta}}{15k^2 (\eta_{\alpha\beta} + 1) T_{\beta}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{0231,1} = \frac{32\eta_{\alpha\beta}^3 m_{\beta}^2 n_{\alpha} n_{\beta}}{105k^2 (\eta_{\alpha\beta} + 1) T_{\beta}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (23)
\end{aligned}$$

$$\begin{aligned}
p = 2, q = 0 : \mathcal{A}_{\alpha\beta}^{2011,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{2021,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{2031,1}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{2022,1}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{2032,1}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\beta}^{2033,1}\Omega_{\alpha\beta}^{33} \\
\mathcal{A}_{\alpha\beta}^{2011,1} = \frac{140k^2 \eta_{\alpha\beta} \theta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}^2 (\eta_{\alpha\beta} (5\theta_{\alpha\beta} - 4) + \theta_{\alpha\beta})}{3 (\eta_{\alpha\beta} + 1)^2 m_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2021,1} = -\frac{112k^2 \eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}^2 \left(\eta_{\alpha\beta}^3 (5\theta_{\alpha\beta}^3 - 12\theta_{\alpha\beta}^2 + 12\theta_{\alpha\beta} - 4) + 3\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^3 + 3\eta_{\alpha\beta} \theta_{\alpha\beta}^3 + \theta_{\alpha\beta}^3 \right)}{3 (\eta_{\alpha\beta} + 1)^4 \theta_{\alpha\beta} m_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2031,1} = \frac{16k^2 \eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}^2 \left(\eta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right) \left(\eta_{\alpha\beta}^2 (5\theta_{\alpha\beta}^2 - 10\theta_{\alpha\beta} + 6) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{3 (\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta}^2 m_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2022,1} = -\frac{224k^2 \eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha}^2 (\eta_{\alpha\beta} (2\theta_{\alpha\beta} - 1) + \theta_{\alpha\beta})}{3 (\eta_{\alpha\beta} + 1)^4 \theta_{\alpha\beta} m_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\beta}^{2032,1} = \frac{64k^2 \eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha}^2 \left(\eta_{\alpha\beta}^2 (2\theta_{\alpha\beta}^2 - 4\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{3 (\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta}^2 m_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\beta}^{2033,1} = \frac{64k^2 \eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1)^2 n_{\alpha} n_{\beta} T_{\alpha}^2}{3 (\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta}^2 m_{\alpha}^2} \quad (24)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 2 : \mathcal{A}_{\alpha\beta}^{1211,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{1221,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{1231,1}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{1241,1}\Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\beta}^{1222,1}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{1232,1}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\beta}^{1242,1}\Omega_{\alpha\beta}^{24} \\
\mathcal{A}_{\alpha\beta}^{1211,1} = -\frac{4\eta_{\alpha\beta}^2 \theta_{\alpha\beta} m_{\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (27\theta_{\alpha\beta} - 10) + 17\theta_{\alpha\beta})}{3k (\eta_{\alpha\beta} + 1)^2 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1221,1} = \frac{8\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta} \left(\eta_{\alpha\beta}^2 (81\theta_{\alpha\beta}^2 - 74\theta_{\alpha\beta} + 20) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} (44\theta_{\alpha\beta} - 17) + 27\theta_{\alpha\beta}^2 \right)}{15k (\eta_{\alpha\beta} + 1)^3 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1231,1} = -\frac{16\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta} \left(\eta_{\alpha\beta}^2 (81\theta_{\alpha\beta}^2 - 118\theta_{\alpha\beta} + 56) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} (22\theta_{\alpha\beta} - 3) + 19\theta_{\alpha\beta}^2 \right)}{105k (\eta_{\alpha\beta} + 1)^3 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1241,1} = \frac{32\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta} \left(\eta_{\alpha\beta}^2 (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{105k (\eta_{\alpha\beta} + 1)^3 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1222,1} = \frac{16\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (9\theta_{\alpha\beta} - 5) + 4\theta_{\alpha\beta})}{15k (\eta_{\alpha\beta} + 1)^3 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{1232,1} = -\frac{64\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (9\theta_{\alpha\beta} - 7) + 2\theta_{\alpha\beta})}{105k (\eta_{\alpha\beta} + 1)^3 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{1242,1} = \frac{64\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1) m_{\beta} n_{\alpha} n_{\beta}}{105k (\eta_{\alpha\beta} + 1)^3 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (25)
\end{aligned}$$

$$\begin{aligned}
p = 2, q = 1 : & \mathcal{A}_{\alpha\beta}^{2111,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{2121,1} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{2131,1} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{2141,1} \Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\beta}^{2122,1} \Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{2132,1} \Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\beta}^{2142,1} \Omega_{\alpha\beta}^{24} \\
& + \mathcal{A}_{\alpha\beta}^{2133,1} \Omega_{\alpha\beta}^{33} + \mathcal{A}_{\alpha\beta}^{2143,1} \Omega_{\alpha\beta}^{34} \\
\mathcal{A}_{\alpha\beta}^{2111,1} = & - \frac{28k\eta_{\alpha\beta}\theta_{\alpha\beta}^2 n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta} (45\theta_{\alpha\beta} - 28) + 17\theta_{\alpha\beta})}{3(\eta_{\alpha\beta} + 1)^2 m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{2121,1} = & \frac{56k\eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^4 m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} (\eta_{\alpha\beta}^3 (135\theta_{\alpha\beta}^3 - 236\theta_{\alpha\beta}^2 + 168\theta_{\alpha\beta} - 40) + \eta_{\alpha\beta}^2 \theta_{\alpha\beta} (169\theta_{\alpha\beta}^2 - 136\theta_{\alpha\beta} + 48) \\
& + \eta_{\alpha\beta} \theta_{\alpha\beta}^2 (101\theta_{\alpha\beta} - 20) + 27\theta_{\alpha\beta}^3) \\
\mathcal{A}_{\alpha\beta}^{2131,1} = & - \frac{16k\eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} (\eta_{\alpha\beta}^4 (135\theta_{\alpha\beta}^4 - 388\theta_{\alpha\beta}^3 + 492\theta_{\alpha\beta}^2 - 280\theta_{\alpha\beta} + 60) \\
& + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (38\theta_{\alpha\beta}^3 - 45\theta_{\alpha\beta}^2 + 36\theta_{\alpha\beta} - 10) + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (23\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 2) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (20\theta_{\alpha\beta} - 1) + 19\theta_{\alpha\beta}^4) \\
\mathcal{A}_{\alpha\beta}^{2141,1} = & \frac{32k\eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} (\eta_{\alpha\beta}^4 (5\theta_{\alpha\beta}^4 - 20\theta_{\alpha\beta}^3 + 36\theta_{\alpha\beta}^2 - 32\theta_{\alpha\beta} + 12) + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) \\
& + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 + \theta_{\alpha\beta}^4) \\
\mathcal{A}_{\alpha\beta}^{2122,1} = & \frac{224k\eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta}^2 (18\theta_{\alpha\beta}^2 - 21\theta_{\alpha\beta} + 5) + \eta_{\alpha\beta} \theta_{\alpha\beta} (15\theta_{\alpha\beta} - 11) + 2\theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^4 m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2132,1} = & - \frac{64k\eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta}^3 (36\theta_{\alpha\beta}^3 - 75\theta_{\alpha\beta}^2 + 56\theta_{\alpha\beta} - 15) + \eta_{\alpha\beta}^2 \theta_{\alpha\beta} (33\theta_{\alpha\beta}^2 - 38\theta_{\alpha\beta} + 11) + 2\eta_{\alpha\beta} \theta_{\alpha\beta}^2 (7\theta_{\alpha\beta} - 4) + 2\theta_{\alpha\beta}^3)}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2142,1} = & \frac{128k\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta}^2 (2\theta_{\alpha\beta}^2 - 4\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2133,1} = & - \frac{64k\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta} (9\theta_{\alpha\beta} - 5) + 4\theta_{\alpha\beta})}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\beta}^{2143,1} = & \frac{128k\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1)^2 n_{\alpha} n_{\beta} T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (26)
\end{aligned}$$

$$\begin{aligned}
p = 2, q = 2 : & \mathcal{A}_{\alpha\beta}^{2211,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{2221,1} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{2231,1} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{2241,1} \Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\beta}^{2251,1} \Omega_{\alpha\beta}^{15} + \mathcal{A}_{\alpha\beta}^{2222,1} \Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{2232,1} \Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\beta}^{2242,1} \Omega_{\alpha\beta}^{24} \\
& + \mathcal{A}_{\alpha\beta}^{2252,1} \Omega_{\alpha\beta}^{25} + \mathcal{A}_{\alpha\beta}^{2233,1} \Omega_{\alpha\beta}^{33} + \mathcal{A}_{\alpha\beta}^{2243,1} \Omega_{\alpha\beta}^{34} + \mathcal{A}_{\alpha\beta}^{2253,1} \Omega_{\alpha\beta}^{35} \\
& \mathcal{A}_{\alpha\beta}^{2211,1} = \frac{6\eta_{\alpha\beta} \theta_{\alpha\beta}^3 n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (55\theta_{\alpha\beta} - 28) + 27\theta_{\alpha\beta})}{(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \\
\mathcal{A}_{\alpha\beta}^{2221,1} = & -\frac{16\eta_{\alpha\beta} \theta_{\alpha\beta} n_{\alpha} n_{\beta}}{15(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} (\eta_{\alpha\beta}^3 (495\theta_{\alpha\beta}^3 - 684\theta_{\alpha\beta}^2 + 378\theta_{\alpha\beta} - 70) + 3\eta_{\alpha\beta}^2 \theta_{\alpha\beta} (267\theta_{\alpha\beta}^2 - 204\theta_{\alpha\beta} + 56) \\
& + 3\eta_{\alpha\beta} \theta_{\alpha\beta}^2 (165\theta_{\alpha\beta} - 46) + 119\theta_{\alpha\beta}^3) \\
\mathcal{A}_{\alpha\beta}^{2231,1} = & \frac{16\eta_{\alpha\beta} n_{\alpha} n_{\beta}}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} (\eta_{\alpha\beta}^4 (1485\theta_{\alpha\beta}^4 - 3348\theta_{\alpha\beta}^3 + 3294\theta_{\alpha\beta}^2 - 1400\theta_{\alpha\beta} + 210) \\
& + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (648\theta_{\alpha\beta}^3 - 864\theta_{\alpha\beta}^2 + 597\theta_{\alpha\beta} - 140) + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (360\theta_{\alpha\beta}^2 - 178\theta_{\alpha\beta} + 59) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (271\theta_{\alpha\beta} - 30) + 241\theta_{\alpha\beta}^4) \\
\mathcal{A}_{\alpha\beta}^{2241,1} = & -\frac{64\eta_{\alpha\beta} n_{\alpha} n_{\beta}}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} (\eta_{\alpha\beta}^4 (55\theta_{\alpha\beta}^4 - 172\theta_{\alpha\beta}^3 + 240\theta_{\alpha\beta}^2 - 158\theta_{\alpha\beta} + 42) \\
& + 2\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (240\theta_{\alpha\beta}^3 - 180\theta_{\alpha\beta}^2 + 3\theta_{\alpha\beta} + 5) + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (90\theta_{\alpha\beta}^2 - 50\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (13\theta_{\alpha\beta} + 1) + 7\theta_{\alpha\beta}^4) \\
\mathcal{A}_{\alpha\beta}^{2251,1} = & \frac{32\eta_{\alpha\beta} n_{\alpha} n_{\beta}}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} (\eta_{\alpha\beta}^4 (5\theta_{\alpha\beta}^4 - 20\theta_{\alpha\beta}^3 + 36\theta_{\alpha\beta}^2 - 32\theta_{\alpha\beta} + 12) + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) \\
& + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 + \theta_{\alpha\beta}^4) \\
\mathcal{A}_{\alpha\beta}^{2222,1} = & -\frac{16\eta_{\alpha\beta} \theta_{\alpha\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta}^2 (198\theta_{\alpha\beta}^2 - 189\theta_{\alpha\beta} + 35) + \eta_{\alpha\beta} \theta_{\alpha\beta} (207\theta_{\alpha\beta} - 119) + 44\theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{2232,1} = & \frac{32\eta_{\alpha\beta} n_{\alpha} n_{\beta}}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} (\eta_{\alpha\beta}^3 (594\theta_{\alpha\beta}^3 - 972\theta_{\alpha\beta}^2 + 539\theta_{\alpha\beta} - 105) + 2\eta_{\alpha\beta}^2 \theta_{\alpha\beta} (405\theta_{\alpha\beta}^2 - 433\theta_{\alpha\beta} + 112) \\
& + \eta_{\alpha\beta} \theta_{\alpha\beta}^2 (377\theta_{\alpha\beta} - 209) + 56\theta_{\alpha\beta}^3) \\
\mathcal{A}_{\alpha\beta}^{2242,1} = & -\frac{64\eta_{\alpha\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta}^3 (66\theta_{\alpha\beta}^3 - 153\theta_{\alpha\beta}^2 + 133\theta_{\alpha\beta} - 42) + \eta_{\alpha\beta}^2 \theta_{\alpha\beta} (45\theta_{\alpha\beta}^2 - 40\theta_{\alpha\beta} + 7) + \eta_{\alpha\beta} \theta_{\alpha\beta}^2 (25\theta_{\alpha\beta} - 13) + 4\theta_{\alpha\beta}^3)}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{2252,1} = & \frac{128\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} (\eta_{\alpha\beta}^2 (2\theta_{\alpha\beta}^2 - 4\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2)}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{2233,1} = & \frac{32\eta_{\alpha\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta}^2 (99\theta_{\alpha\beta}^2 - 126\theta_{\alpha\beta} + 35) + 8\eta_{\alpha\beta} \theta_{\alpha\beta} (9\theta_{\alpha\beta} - 7) + 8\theta_{\alpha\beta}^2)}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{2243,1} = & -\frac{128\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (11\theta_{\alpha\beta} - 7) + 4\theta_{\alpha\beta})}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{2253,1} = \frac{128\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1)^2 n_{\alpha} n_{\beta}}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (27)
\end{aligned}$$

C. Values of $A_{\alpha\beta}^{1pq}$ for $p, q \leq 2$

$$p = 0, q = 0 : \mathcal{A}_{\alpha\alpha}^{0011,1} \Omega_{\alpha\beta}^{11}$$

$$\mathcal{A}_{\alpha\alpha}^{0011,1} = -\frac{16n_{\alpha} n_{\beta}}{3\eta_{\alpha\beta} + 3} \quad (28)$$

$$p = 0, q = 1 : \mathcal{A}_{\alpha\alpha}^{0111,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{0121,1} \Omega_{\alpha\beta}^{12}$$

$$\mathcal{A}_{\alpha\alpha}^{0111,1} = \frac{16\theta_{\alpha\beta} m_{\alpha} n_{\alpha} n_{\beta}}{3k(\eta_{\alpha\beta} + 1) T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})}$$

$$\mathcal{A}_{\alpha\alpha}^{0121,1} = -\frac{32\theta_{\alpha\beta} m_{\alpha} n_{\alpha} n_{\beta}}{15k(\eta_{\alpha\beta} + 1) T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (29)$$

$$\begin{aligned}
p = 1, q = 0 : \mathcal{A}_{\alpha\alpha}^{1011,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{1021,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{1022,1}\Omega_{\alpha\beta}^{22} \\
\mathcal{A}_{\alpha\alpha}^{1011,1} = \frac{40kn_{\alpha}n_{\beta}T_{\alpha} \left(-2\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}\theta_{\alpha\beta}^2 + \theta_{\alpha\beta}^2 \right)}{3(\eta_{\alpha\beta} + 1)^2\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{1021,1} = -\frac{16kn_{\alpha}n_{\beta}T_{\alpha} \left(\eta_{\alpha\beta}^2 (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{3(\eta_{\alpha\beta} + 1)^3\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{1022,1} = -\frac{32k\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha}}{3(\eta_{\alpha\beta} + 1)^3\theta_{\alpha\beta}m_{\alpha}} \quad (30)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 1 : \mathcal{A}_{\alpha\alpha}^{1111,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{1121,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{1131,1}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{1122,1}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{1132,1}\Omega_{\alpha\beta}^{23} \\
\mathcal{A}_{\alpha\alpha}^{1111,1} = -\frac{8n_{\alpha}n_{\beta} \left(-2\eta_{\alpha\beta}^2 (5\theta_{\alpha\beta} - 8) + 5\eta_{\alpha\beta}\theta_{\alpha\beta}^2 + 6\eta_{\alpha\beta}^3 + 5\theta_{\alpha\beta}^2 \right)}{3(\eta_{\alpha\beta} + 1)^2(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{1121,1} = -\frac{32n_{\alpha}n_{\beta} \left(11\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}^2 (-10\theta_{\alpha\beta}^2 + 26\theta_{\alpha\beta} - 21) - 5\eta_{\alpha\beta}\theta_{\alpha\beta} (\theta_{\alpha\beta} + 1) - 5\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{1131,1} = -\frac{32n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{1122,1} = -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} (2\eta_{\alpha\beta} - 5\theta_{\alpha\beta} + 7)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{1132,1} = -\frac{64\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (31)
\end{aligned}$$

$$\begin{aligned}
p = 0, q = 2 : \mathcal{A}_{\alpha\alpha}^{0211,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{0221,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{0231,1}\Omega_{\alpha\beta}^{13} \\
\mathcal{A}_{\alpha\alpha}^{0211,1} = -\frac{8\theta_{\alpha\beta}^2 m_{\alpha}^2 n_{\alpha} n_{\beta}}{3k^2 (\eta_{\alpha\beta} + 1) T_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{0221,1} = \frac{32\theta_{\alpha\beta}^2 m_{\alpha}^2 n_{\alpha} n_{\beta}}{15k^2 (\eta_{\alpha\beta} + 1) T_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{0231,1} = -\frac{32\theta_{\alpha\beta}^2 m_{\alpha}^2 n_{\alpha} n_{\beta}}{105k^2 (\eta_{\alpha\beta} + 1) T_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (32)
\end{aligned}$$

$$\begin{aligned}
p = 2, q = 0 : & \mathcal{A}_{\alpha\alpha}^{2011,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{2021,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{2031,1}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{2022,1}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{2032,1}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\alpha}^{2033,1}\Omega_{\alpha\beta}^{33} \\
& \mathcal{A}_{\alpha\alpha}^{2011,1} = \frac{140k^2 n_\alpha n_\beta T_\alpha^2 \left(4\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) - \eta_{\alpha\beta} \theta_{\alpha\beta}^2 - \theta_{\alpha\beta}^2 \right)}{3 (\eta_{\alpha\beta} + 1)^2 m_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{2021,1} = \frac{112k^2 n_\alpha n_\beta T_\alpha^2}{3 (\eta_{\alpha\beta} + 1)^4 \theta_{\alpha\beta}^2 m_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \left(-2\eta_{\alpha\beta}^4 (\theta_{\alpha\beta}^3 - 3\theta_{\alpha\beta}^2 + 4\theta_{\alpha\beta} - 2) + \eta_{\alpha\beta}^3 \theta_{\alpha\beta} (3\theta_{\alpha\beta}^3 - 6\theta_{\alpha\beta}^2 + 4) \right. \\
& \quad \left. + 3\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + \eta_{\alpha\beta} \theta_{\alpha\beta}^3 (\theta_{\alpha\beta} + 2) + \theta_{\alpha\beta}^4 \right) \\
& \mathcal{A}_{\alpha\alpha}^{2031,1} = -\frac{16k^2 n_\alpha n_\beta T_\alpha^2}{3 (\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta}^2 m_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \left(\eta_{\alpha\beta}^4 (5\theta_{\alpha\beta}^4 - 20\theta_{\alpha\beta}^3 + 36\theta_{\alpha\beta}^2 - 32\theta_{\alpha\beta} + 12) \right. \\
& \quad \left. + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 + \theta_{\alpha\beta}^4 \right) \\
& \mathcal{A}_{\alpha\alpha}^{2022,1} = \frac{224k^2 \eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) n_\alpha n_\beta T_\alpha^2 \left(\eta_{\alpha\beta}^2 (-\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta} \theta_{\alpha\beta}^2 + \theta_{\alpha\beta}^2 \right)}{3 (\eta_{\alpha\beta} + 1)^4 \theta_{\alpha\beta}^2 m_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\alpha}^{2032,1} = -\frac{64k^2 \eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) n_\alpha n_\beta T_\alpha^2 \left(\eta_{\alpha\beta}^2 (2\theta_{\alpha\beta}^2 - 4\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{3 (\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta}^2 m_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\alpha}^{2033,1} = -\frac{64k^2 \eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1)^2 n_\alpha n_\beta T_\alpha^2}{3 (\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta}^2 m_\alpha^2} \quad (33)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 2 : & \mathcal{A}_{\alpha\alpha}^{1211,1}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{1221,1}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{1231,1}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{1241,1}\Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\alpha}^{1222,1}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{1232,1}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\alpha}^{1242,1}\Omega_{\alpha\beta}^{24} \\
& \mathcal{A}_{\alpha\alpha}^{1211,1} = \frac{4\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta \left(\eta_{\alpha\beta}^2 (22 - 10\theta_{\alpha\beta}) + 5\eta_{\alpha\beta} \theta_{\alpha\beta}^2 + 12\eta_{\alpha\beta}^3 + 5\theta_{\alpha\beta}^2 \right)}{3k (\eta_{\alpha\beta} + 1)^2 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1221,1} = -\frac{8\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta \left(\eta_{\alpha\beta}^3 (68 - 44\theta_{\alpha\beta}) + \eta_{\alpha\beta}^2 (25\theta_{\alpha\beta}^2 - 74\theta_{\alpha\beta} + 76) + 10\eta_{\alpha\beta} \theta_{\alpha\beta} (2\theta_{\alpha\beta} + 1) + 12\eta_{\alpha\beta}^4 + 15\theta_{\alpha\beta}^2 \right)}{15k (\eta_{\alpha\beta} + 1)^3 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1231,1} = \frac{16\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta \left(-34\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}^2 (47\theta_{\alpha\beta}^2 - 118\theta_{\alpha\beta} + 90) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} (5\theta_{\alpha\beta} + 14) + 19\theta_{\alpha\beta}^2 \right)}{105k (\eta_{\alpha\beta} + 1)^3 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1241,1} = -\frac{32\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta \left(\eta_{\alpha\beta}^2 (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{105k (\eta_{\alpha\beta} + 1)^3 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1222,1} = \frac{16\eta_{\alpha\beta} \theta_{\alpha\beta} m_\alpha n_\alpha n_\beta (4\eta_{\alpha\beta} - 5\theta_{\alpha\beta} + 9)}{15k (\eta_{\alpha\beta} + 1)^3 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1232,1} = -\frac{64\eta_{\alpha\beta} \theta_{\alpha\beta} m_\alpha n_\alpha n_\beta (2\eta_{\alpha\beta} - 7\theta_{\alpha\beta} + 9)}{105k (\eta_{\alpha\beta} + 1)^3 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1242,1} = -\frac{64\eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) \theta_{\alpha\beta} m_\alpha n_\alpha n_\beta}{105k (\eta_{\alpha\beta} + 1)^3 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (34)
\end{aligned}$$

$$\begin{aligned}
p = 2, q = 1: & \mathcal{A}_{\alpha\alpha}^{2111,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{2121,1} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{2131,1} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{2141,1} \Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\alpha}^{2122,1} \Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{2132,1} \Omega_{\alpha\beta}^{23} \\
& + \mathcal{A}_{\alpha\alpha}^{2142,1} \Omega_{\alpha\beta}^{24} + \mathcal{A}_{\alpha\alpha}^{2133,1} \Omega_{\alpha\beta}^{33} + \mathcal{A}_{\alpha\alpha}^{2143,1} \Omega_{\alpha\beta}^{34} \\
\mathcal{A}_{\alpha\alpha}^{2111,1} = & -\frac{28kn_{\alpha}n_{\beta}T_{\alpha} \left(8\eta_{\alpha\beta}^4 (\theta_{\alpha\beta} - 1) - 12\eta_{\alpha\beta}^3 \theta_{\alpha\beta}^2 + 4\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (5\theta_{\alpha\beta} - 8) - 5\eta_{\alpha\beta} \theta_{\alpha\beta}^4 - 5\theta_{\alpha\beta}^4 \right)}{3(\eta_{\alpha\beta} + 1)^2 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\alpha}^{2121,1} = & -\frac{56kn_{\alpha}n_{\beta}T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^4 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(12\eta_{\alpha\beta}^5 (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) - 4\eta_{\alpha\beta}^4 (16\theta_{\alpha\beta}^3 - 35\theta_{\alpha\beta}^2 + 32\theta_{\alpha\beta} - 22) \right. \\
& \left. + \eta_{\alpha\beta}^3 \theta_{\alpha\beta} (35\theta_{\alpha\beta}^3 - 148\theta_{\alpha\beta}^2 + 100\theta_{\alpha\beta} + 64) + \eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (45\theta_{\alpha\beta}^2 - 104\theta_{\alpha\beta} + 116) + 5\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (5\theta_{\alpha\beta} + 4) + 15\theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2131,1} = & -\frac{16kn_{\alpha}n_{\beta}T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(4\eta_{\alpha\beta}^5 (17\theta_{\alpha\beta}^3 - 51\theta_{\alpha\beta}^2 + 64\theta_{\alpha\beta} - 30) \right. \\
& \left. + \eta_{\alpha\beta}^4 (-67\theta_{\alpha\beta}^4 + 252\theta_{\alpha\beta}^3 - 336\theta_{\alpha\beta}^2 + 312\theta_{\alpha\beta} - 180) - 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (21\theta_{\alpha\beta}^3 - 33\theta_{\alpha\beta}^2 - 15\theta_{\alpha\beta} + 46) \right. \\
& \left. - 2\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (43\theta_{\alpha\beta}^2 - 70\theta_{\alpha\beta} + 84) - 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (7\theta_{\alpha\beta} + 12) - 19\theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2141,1} = & -\frac{32kn_{\alpha}n_{\beta}T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(\eta_{\alpha\beta}^4 (5\theta_{\alpha\beta}^4 - 20\theta_{\alpha\beta}^3 + 36\theta_{\alpha\beta}^2 - 32\theta_{\alpha\beta} + 12) \right. \\
& \left. + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 + \theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2122,1} = & -\frac{224k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha} \left(6\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}^2 (-7\theta_{\alpha\beta}^2 + 16\theta_{\alpha\beta} - 11) + \eta_{\alpha\beta} \theta_{\alpha\beta}^2 (5\theta_{\alpha\beta} - 9) + \theta_{\alpha\beta}^2 (5\theta_{\alpha\beta} - 7) \right)}{15(\eta_{\alpha\beta} + 1)^4 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{2132,1} = & -\frac{64k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \left(\eta_{\alpha\beta}^3 (19\theta_{\alpha\beta}^2 - 38\theta_{\alpha\beta} + 21) + \eta_{\alpha\beta}^2 (-17\theta_{\alpha\beta}^3 + 56\theta_{\alpha\beta}^2 - 69\theta_{\alpha\beta} + 36) \right. \\
& \left. + 2\eta_{\alpha\beta} \theta_{\alpha\beta} (-7\theta_{\alpha\beta}^2 + 3\theta_{\alpha\beta} + 7) + 2(7 - 6\theta_{\alpha\beta}) \theta_{\alpha\beta}^2 \right) \\
\mathcal{A}_{\alpha\alpha}^{2142,1} = & -\frac{128k\eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha} \left(\eta_{\alpha\beta}^2 (2\theta_{\alpha\beta}^2 - 4\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{2133,1} = & -\frac{64k\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha} (4\eta_{\alpha\beta} - 5\theta_{\alpha\beta} + 9)}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{2143,1} = & -\frac{128k\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1)^2 n_{\alpha} n_{\beta} T_{\alpha}}{15(\eta_{\alpha\beta} + 1)^5 \theta_{\alpha\beta} m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (35)
\end{aligned}$$

$$\begin{aligned}
p = 2, q = 2 : & \mathcal{A}_{\alpha\alpha}^{2211,1} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{2221,1} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{2231,1} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{2241,1} \Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\alpha}^{2251,1} \Omega_{\alpha\beta}^{15} + \mathcal{A}_{\alpha\alpha}^{2222,1} \Omega_{\alpha\beta}^{22} \\
& + \mathcal{A}_{\alpha\alpha}^{2232,1} \Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\alpha}^{2242,1} \Omega_{\alpha\beta}^{24} + \mathcal{A}_{\alpha\alpha}^{2252,1} \Omega_{\alpha\beta}^{25} + \mathcal{A}_{\alpha\alpha}^{2233,1} \Omega_{\alpha\beta}^{33} + \mathcal{A}_{\alpha\alpha}^{2243,1} \Omega_{\alpha\beta}^{34} + \mathcal{A}_{\alpha\alpha}^{2253,1} \Omega_{\alpha\beta}^{35} \\
\mathcal{A}_{\alpha\alpha}^{2211,1} = & - \frac{2n_\alpha n_\beta \left(-8\eta_{\alpha\beta}^4 (14\theta_{\alpha\beta} - 19) + 168\eta_{\alpha\beta}^3 \theta_{\alpha\beta}^2 - 28\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (5\theta_{\alpha\beta} - 11) + 35\eta_{\alpha\beta} \theta_{\alpha\beta}^4 + 40\eta_{\alpha\beta}^5 + 35\theta_{\alpha\beta}^4 \right)}{3(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \\
\mathcal{A}_{\alpha\alpha}^{2221,1} = & \frac{16n_\alpha n_\beta}{15(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \left(-68\eta_{\alpha\beta}^6 (\theta_{\alpha\beta} - 1) + 4\eta_{\alpha\beta}^5 (42\theta_{\alpha\beta}^2 - 97\theta_{\alpha\beta} + 76) \right. \\
& + \eta_{\alpha\beta}^4 (-189\theta_{\alpha\beta}^3 + 511\theta_{\alpha\beta}^2 - 376\theta_{\alpha\beta} + 306) + 7\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (10\theta_{\alpha\beta}^3 - 59\theta_{\alpha\beta}^2 + 68\theta_{\alpha\beta} + 22) + 7\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (15\theta_{\alpha\beta}^2 - 37\theta_{\alpha\beta} + 49) \\
& \left. + 35\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (2\theta_{\alpha\beta} + 1) + 35\theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2231,1} = & - \frac{16n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \left(12\eta_{\alpha\beta}^6 (31\theta_{\alpha\beta}^2 - 62\theta_{\alpha\beta} + 40) + \eta_{\alpha\beta}^5 (-714\theta_{\alpha\beta}^3 + 2410\theta_{\alpha\beta}^2 - 3064\theta_{\alpha\beta} + 1800) \right. \\
& + \eta_{\alpha\beta}^4 (399\theta_{\alpha\beta}^4 - 2128\theta_{\alpha\beta}^3 + 2916\theta_{\alpha\beta}^2 - 1936\theta_{\alpha\beta} + 1530) + 2\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (329\theta_{\alpha\beta}^3 - 924\theta_{\alpha\beta}^2 + 465\theta_{\alpha\beta} + 612) \\
& \left. + 2\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (301\theta_{\alpha\beta}^2 - 504\theta_{\alpha\beta} + 656) + 266\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (\theta_{\alpha\beta} + 1) + 133\theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2241,1} = & - \frac{64n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \left(\eta_{\alpha\beta}^5 (27\theta_{\alpha\beta}^3 - 81\theta_{\alpha\beta}^2 + 100\theta_{\alpha\beta} - 46) + \eta_{\alpha\beta}^4 (-28\theta_{\alpha\beta}^4 + 118\theta_{\alpha\beta}^3 - 183\theta_{\alpha\beta}^2 + 174\theta_{\alpha\beta} - 88) \right. \\
& \left. + \eta_{\alpha\beta}^3 \theta_{\alpha\beta} (-21\theta_{\alpha\beta}^3 + 12\theta_{\alpha\beta}^2 + 75\theta_{\alpha\beta} - 94) + \eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (-35\theta_{\alpha\beta}^2 + 68\theta_{\alpha\beta} - 75) - 7\eta_{\alpha\beta} \theta_{\alpha\beta}^3 (\theta_{\alpha\beta} + 3) - 7\theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2251,1} = & - \frac{32n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \left(\eta_{\alpha\beta}^4 (5\theta_{\alpha\beta}^4 - 20\theta_{\alpha\beta}^3 + 36\theta_{\alpha\beta}^2 - 32\theta_{\alpha\beta} + 12) + 4\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (3\theta_{\alpha\beta}^2 - 6\theta_{\alpha\beta} + 4) \right. \\
& \left. + 6\eta_{\alpha\beta}^2 \theta_{\alpha\beta}^2 (\theta_{\alpha\beta}^2 - 2\theta_{\alpha\beta} + 2) + 4\eta_{\alpha\beta} \theta_{\alpha\beta}^3 + \theta_{\alpha\beta}^4 \right) \\
\mathcal{A}_{\alpha\alpha}^{2222,1} = & - \frac{16\eta_{\alpha\beta} n_\alpha n_\beta \left(\eta_{\alpha\beta}^3 (116 - 84\theta_{\alpha\beta}) + \eta_{\alpha\beta}^2 (63\theta_{\alpha\beta}^2 - 154\theta_{\alpha\beta} + 135) + 7\eta_{\alpha\beta} (13 - 5\theta_{\alpha\beta}) \theta_{\alpha\beta}^2 + 16\eta_{\alpha\beta}^4 + 7(9 - 5\theta_{\alpha\beta}) \theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\alpha}^{2232,1} = & - \frac{32\eta_{\alpha\beta} n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(132\eta_{\alpha\beta}^4 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}^3 (-294\theta_{\alpha\beta}^2 + 796\theta_{\alpha\beta} - 558) \right. \\
& \left. + \eta_{\alpha\beta}^2 (168\theta_{\alpha\beta}^3 - 658\theta_{\alpha\beta}^2 + 853\theta_{\alpha\beta} - 531) + 14\eta_{\alpha\beta} \theta_{\alpha\beta} (14\theta_{\alpha\beta}^2 - 17\theta_{\alpha\beta} - 9) + 7\theta_{\alpha\beta}^2 (19\theta_{\alpha\beta} - 27) \right) \\
\mathcal{A}_{\alpha\alpha}^{2242,1} = & - \frac{64\eta_{\alpha\beta} n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(\eta_{\alpha\beta}^3 (31\theta_{\alpha\beta}^2 - 62\theta_{\alpha\beta} + 35) + \eta_{\alpha\beta}^2 (-35\theta_{\alpha\beta}^3 + 122\theta_{\alpha\beta}^2 - 152\theta_{\alpha\beta} + 77) \right. \\
& \left. - 2\eta_{\alpha\beta} \theta_{\alpha\beta} (7\theta_{\alpha\beta}^2 + 5\theta_{\alpha\beta} - 18) + (25 - 21\theta_{\alpha\beta}) \theta_{\alpha\beta}^2 \right) \\
\mathcal{A}_{\alpha\alpha}^{2252,1} = & - \frac{128\eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) n_\alpha n_\beta \left(\eta_{\alpha\beta}^2 (2\theta_{\alpha\beta}^2 - 4\theta_{\alpha\beta} + 3) + 2\eta_{\alpha\beta} \theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\alpha}^{2233,1} = & - \frac{32\eta_{\alpha\beta}^2 n_\alpha n_\beta \left(-8\eta_{\alpha\beta} (7\theta_{\alpha\beta} - 9) + 8\eta_{\alpha\beta}^2 + 35\theta_{\alpha\beta}^2 - 126\theta_{\alpha\beta} + 99 \right)}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{2243,1} = & - \frac{128\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_\alpha n_\beta (4\eta_{\alpha\beta} - 7\theta_{\alpha\beta} + 11)}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{2253,1} = & - \frac{128\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1)^2 n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^5 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (36)
\end{aligned}$$

D. Values of $B_{\alpha\beta}^{2pq}$ for $p, q \leq 1$

$$\begin{aligned}
p = 0, q = 0 : & \mathcal{A}_{\alpha\beta}^{0011,2} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{0021,2} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{0022,2} \Omega_{\alpha\beta}^{22} \\
& \mathcal{A}_{\alpha\beta}^{0011,2} = \frac{32\eta_{\alpha\beta}\theta_{\alpha\beta}n_{\alpha}n_{\beta}}{3(\eta_{\alpha\beta} + 1)(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{0021,2} = -\frac{64\eta_{\alpha\beta}^2(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}}{15(\eta_{\alpha\beta} + 1)^2(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\beta}^{0022,2} = -\frac{16\eta_{\alpha\beta}n_{\alpha}n_{\beta}}{5(\eta_{\alpha\beta} + 1)^2} \quad (37)
\end{aligned}$$

$$\begin{aligned}
p = 0, q = 1 : & \mathcal{A}_{\alpha\beta}^{0111,2} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{0121,2} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{0131,2} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{0122,2} \Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{0123,2} \Omega_{\alpha\beta}^{23} \\
& \mathcal{A}_{\alpha\beta}^{0111,2} = -\frac{32\eta_{\alpha\beta}^2\theta_{\alpha\beta}m_{\beta}n_{\alpha}n_{\beta}}{3k(\eta_{\alpha\beta} + 1)T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{0121,2} = \frac{64\eta_{\alpha\beta}^2m_{\beta}n_{\alpha}n_{\beta}(\eta_{\alpha\beta}(2\theta_{\alpha\beta} - 1) + \theta_{\alpha\beta})}{15k(\eta_{\alpha\beta} + 1)^2T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{0131,2} = -\frac{128\eta_{\alpha\beta}^3(\theta_{\alpha\beta} - 1)m_{\beta}n_{\alpha}n_{\beta}}{105k(\eta_{\alpha\beta} + 1)^2T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{0122,2} = \frac{16\eta_{\alpha\beta}^2m_{\beta}n_{\alpha}n_{\beta}}{5k(\eta_{\alpha\beta} + 1)^2T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\beta}^{0123,2} = -\frac{32\eta_{\alpha\beta}^2m_{\beta}n_{\alpha}n_{\beta}}{35k(\eta_{\alpha\beta} + 1)^2T_{\beta}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (38)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 0 : & \mathcal{A}_{\alpha\beta}^{1011,2} \Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{1021,2} \Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{1031,2} \Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{1022,2} \Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{1023,2} \Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\beta}^{1033,2} \Omega_{\alpha\beta}^{33} \\
& \mathcal{A}_{\alpha\beta}^{1011,2} = -\frac{112k\eta_{\alpha\beta}\theta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}(2\theta_{\alpha\beta} - 1) + \theta_{\alpha\beta})}{3(\eta_{\alpha\beta} + 1)^2m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{1021,2} = \frac{224k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}^2(4\theta_{\alpha\beta}^2 - 7\theta_{\alpha\beta} + 4) + \eta_{\alpha\beta}\theta_{\alpha\beta}(\theta_{\alpha\beta} + 1) + \theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^3m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{1031,2} = -\frac{32k\eta_{\alpha\beta}^2(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}^2(4\theta_{\alpha\beta}^2 - 8\theta_{\alpha\beta} + 5) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^4\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\beta}^{1022,2} = \frac{56k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}(11\theta_{\alpha\beta} - 8) + 3\theta_{\alpha\beta})}{15(\eta_{\alpha\beta} + 1)^3m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\beta}^{1032,2} = -\frac{16k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha}(\eta_{\alpha\beta}^2(11\theta_{\alpha\beta}^2 - 22\theta_{\alpha\beta} + 14) + 6\eta_{\alpha\beta}\theta_{\alpha\beta} + 3\theta_{\alpha\beta}^2)}{15(\eta_{\alpha\beta} + 1)^4\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\beta}^{1033,2} = -\frac{32k\eta_{\alpha\beta}^2(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha}}{5(\eta_{\alpha\beta} + 1)^4\theta_{\alpha\beta}m_{\alpha}} \quad (39)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 1 : & \mathcal{A}_{\alpha\beta}^{1111,2}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\beta}^{1121,2}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\beta}^{1131,2}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\beta}^{1141,2}\Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\beta}^{1122,2}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\beta}^{1132,2}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\beta}^{1142,2}\Omega_{\alpha\beta}^{24} \\
& + \mathcal{A}_{\alpha\beta}^{1133,2}\Omega_{\alpha\beta}^{33} + \mathcal{A}_{\alpha\beta}^{1143,2}\Omega_{\alpha\beta}^{34} \\
\mathcal{A}_{\alpha\beta}^{1111,2} = & \frac{16\eta_{\alpha\beta}\theta_{\alpha\beta}^2 n_{\alpha}n_{\beta} (\eta_{\alpha\beta} (18\theta_{\alpha\beta} - 7) + 11\theta_{\alpha\beta})}{3(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1121,2} = & -\frac{64\eta_{\alpha\beta}\theta_{\alpha\beta}n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (27\theta_{\alpha\beta}^2 - 34\theta_{\alpha\beta} + 14) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} (10\theta_{\alpha\beta} - 3) + 7\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1131,2} = & \frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^3 (108\theta_{\alpha\beta}^3 - 230\theta_{\alpha\beta}^2 + 173\theta_{\alpha\beta} - 35) + 2\eta_{\alpha\beta}^2\theta_{\alpha\beta} (47\theta_{\alpha\beta}^2 - 57\theta_{\alpha\beta} + 34) + \eta_{\alpha\beta}\theta_{\alpha\beta}^2 (37\theta_{\alpha\beta} + 11) + 16\theta_{\alpha\beta}^3 \right)}{105(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1141,2} = & -\frac{64\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (4\theta_{\alpha\beta}^2 - 8\theta_{\alpha\beta} + 5) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
\mathcal{A}_{\alpha\beta}^{1122,2} = & -\frac{8\eta_{\alpha\beta}\theta_{\alpha\beta}n_{\alpha}n_{\beta} (\eta_{\alpha\beta} (99\theta_{\alpha\beta} - 56) + 43\theta_{\alpha\beta})}{15(\eta_{\alpha\beta} + 1)^3 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{1132,2} = & \frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (99\theta_{\alpha\beta}^2 - 127\theta_{\alpha\beta} + 49) + \eta_{\alpha\beta}\theta_{\alpha\beta} (71\theta_{\alpha\beta} - 29) + 21\theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{1142,2} = & -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (11\theta_{\alpha\beta}^2 - 22\theta_{\alpha\beta} + 14) + 6\eta_{\alpha\beta}\theta_{\alpha\beta} + 3\theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\beta}^{1133,2} = & \frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} (\eta_{\alpha\beta} (9\theta_{\alpha\beta} - 7) + 2\theta_{\alpha\beta})}{35(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\beta}^{1143,2} = & -\frac{64\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) n_{\alpha}n_{\beta}}{35(\eta_{\alpha\beta} + 1)^4 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (40)
\end{aligned}$$

E. Values for $A_{\alpha\beta}^{2pq}$ for $p, q \leq 1$

$$\begin{aligned}
p = 0, q = 0 : & \mathcal{A}_{\alpha\alpha}^{0011,2}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{0021,2}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{0022,2}\Omega_{\alpha\beta}^{22} \\
\mathcal{A}_{\alpha\alpha}^{0011,2} = & -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta}}{3(\eta_{\alpha\beta} + 1) (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{0021,2} = & -\frac{64\eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) n_{\alpha}n_{\beta}}{15(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{0022,2} = & -\frac{16n_{\alpha}n_{\beta}}{5(\eta_{\alpha\beta} + 1)^2} \quad (41)
\end{aligned}$$

$$\begin{aligned}
p = 0, q = 1 : & \mathcal{A}_{\alpha\alpha}^{0111,2}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{0121,2}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{0131,2}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{0122,2}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{0123,2}\Omega_{\alpha\beta}^{23} \\
\mathcal{A}_{\alpha\alpha}^{0111,2} = & \frac{32\eta_{\alpha\beta}\theta_{\alpha\beta}m_{\alpha}n_{\alpha}n_{\beta}}{3k(\eta_{\alpha\beta} + 1) T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{0121,2} = & -\frac{64\eta_{\alpha\beta}\theta_{\alpha\beta}m_{\alpha}n_{\alpha}n_{\beta} (\eta_{\alpha\beta} - \theta_{\alpha\beta} + 2)}{15k(\eta_{\alpha\beta} + 1)^2 T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{0131,2} = & -\frac{128\eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) \theta_{\alpha\beta}m_{\alpha}n_{\alpha}n_{\beta}}{105k(\eta_{\alpha\beta} + 1)^2 T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
\mathcal{A}_{\alpha\alpha}^{0122,2} = & \frac{16\theta_{\alpha\beta}m_{\alpha}n_{\alpha}n_{\beta}}{5k(\eta_{\alpha\beta} + 1)^2 T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
\mathcal{A}_{\alpha\alpha}^{0132,2} = & -\frac{32\theta_{\alpha\beta}m_{\alpha}n_{\alpha}n_{\beta}}{35k(\eta_{\alpha\beta} + 1)^2 T_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (42)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 0 : & \mathcal{A}_{\alpha\alpha}^{1011,2}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{1021,2}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{1031,2}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{1022,2}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{1023,2}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\alpha}^{1033,2}\Omega_{\alpha\beta}^{33} \\
& \mathcal{A}_{\alpha\alpha}^{1011,2} = \frac{112k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha} \left(\eta_{\alpha\beta}^2 (-\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}\theta_{\alpha\beta}^2 + \theta_{\alpha\beta}^2 \right)}{3(\eta_{\alpha\beta} + 1)^2\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1021,2} = \frac{224k\eta_{\alpha\beta}n_{\alpha}n_{\beta}T_{\alpha} \left(\eta_{\alpha\beta}^2 (-3\theta_{\alpha\beta}^2 + 6\theta_{\alpha\beta} - 4) + \eta_{\alpha\beta}\theta_{\alpha\beta} (\theta_{\alpha\beta}^2 - \theta_{\alpha\beta} - 2) + (\theta_{\alpha\beta} - 2)\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1031,2} = -\frac{32k\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha} \left(\eta_{\alpha\beta}^2 (4\theta_{\alpha\beta}^2 - 8\theta_{\alpha\beta} + 5) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^4\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1022,2} = -\frac{56kn_{\alpha}n_{\beta}T_{\alpha} \left(8\eta_{\alpha\beta}^2(\theta_{\alpha\beta} - 1) - 3\eta_{\alpha\beta}\theta_{\alpha\beta}^2 - 3\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\alpha}^{1032,2} = -\frac{16kn_{\alpha}n_{\beta}T_{\alpha} \left(\eta_{\alpha\beta}^2 (11\theta_{\alpha\beta}^2 - 22\theta_{\alpha\beta} + 14) + 6\eta_{\alpha\beta}\theta_{\alpha\beta} + 3\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^4\theta_{\alpha\beta}m_{\alpha}(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\alpha}^{1033,2} = -\frac{32k\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}T_{\alpha}}{5(\eta_{\alpha\beta} + 1)^4\theta_{\alpha\beta}m_{\alpha}} \quad (43)
\end{aligned}$$

$$\begin{aligned}
p = 1, q = 1 : & \mathcal{A}_{\alpha\alpha}^{1111,2}\Omega_{\alpha\beta}^{11} + \mathcal{A}_{\alpha\alpha}^{1121,2}\Omega_{\alpha\beta}^{12} + \mathcal{A}_{\alpha\alpha}^{1131,2}\Omega_{\alpha\beta}^{13} + \mathcal{A}_{\alpha\alpha}^{1141,2}\Omega_{\alpha\beta}^{14} + \mathcal{A}_{\alpha\alpha}^{1122,2}\Omega_{\alpha\beta}^{22} + \mathcal{A}_{\alpha\alpha}^{1132,2}\Omega_{\alpha\beta}^{23} + \mathcal{A}_{\alpha\alpha}^{1142,2}\Omega_{\alpha\beta}^{24} \\
& \quad + \mathcal{A}_{\alpha\alpha}^{1133,2}\Omega_{\alpha\beta}^{33} + \mathcal{A}_{\alpha\alpha}^{1143,2}\Omega_{\alpha\beta}^{34} \\
& \mathcal{A}_{\alpha\alpha}^{1111,2} = -\frac{16\eta_{\alpha\beta}n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (11 - 7\theta_{\alpha\beta}) + 7\eta_{\alpha\beta}\theta_{\alpha\beta}^2 + 4\eta_{\alpha\beta}^3 + 7\theta_{\alpha\beta}^2 \right)}{3(\eta_{\alpha\beta} + 1)^2(\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1121,2} = -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} \left(19\eta_{\alpha\beta}^3(\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}^2 (-28\theta_{\alpha\beta}^2 + 61\theta_{\alpha\beta} - 47) + 7\eta_{\alpha\beta}\theta_{\alpha\beta} (\theta_{\alpha\beta}^2 - 3\theta_{\alpha\beta} - 2) + 7(\theta_{\alpha\beta} - 3)\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1131,2} = -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta}}{105(\eta_{\alpha\beta} + 1)^4(\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(2\eta_{\alpha\beta}^3 (33\theta_{\alpha\beta}^2 - 66\theta_{\alpha\beta} + 41) + \eta_{\alpha\beta}^2 (-42\theta_{\alpha\beta}^3 + 164\theta_{\alpha\beta}^2 - 191\theta_{\alpha\beta} + 117) \right. \\
& \quad \left. + 2\eta_{\alpha\beta}\theta_{\alpha\beta} (-14\theta_{\alpha\beta}^2 + 15\theta_{\alpha\beta} + 23) + (37 - 21\theta_{\alpha\beta})\theta_{\alpha\beta}^2 \right) \\
& \mathcal{A}_{\alpha\alpha}^{1141,2} = -\frac{64\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (4\theta_{\alpha\beta}^2 - 8\theta_{\alpha\beta} + 5) + 2\eta_{\alpha\beta}\theta_{\alpha\beta} + \theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^4(\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \\
& \mathcal{A}_{\alpha\alpha}^{1122,2} = -\frac{8n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (78 - 56\theta_{\alpha\beta}) + 21\eta_{\alpha\beta}\theta_{\alpha\beta}^2 + 22\eta_{\alpha\beta}^3 + 21\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^3(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1132,2} = -\frac{32n_{\alpha}n_{\beta} \left(50\eta_{\alpha\beta}^3(\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta}^2 (-49\theta_{\alpha\beta}^2 + 127\theta_{\alpha\beta} - 99) - 21\eta_{\alpha\beta}\theta_{\alpha\beta}(\theta_{\alpha\beta} + 1) - 21\theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^4(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1142,2} = -\frac{32n_{\alpha}n_{\beta} \left(\eta_{\alpha\beta}^2 (11\theta_{\alpha\beta}^2 - 22\theta_{\alpha\beta} + 14) + 6\eta_{\alpha\beta}\theta_{\alpha\beta} + 3\theta_{\alpha\beta}^2 \right)}{105(\eta_{\alpha\beta} + 1)^4(\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \\
& \mathcal{A}_{\alpha\alpha}^{1133,2} = -\frac{32\eta_{\alpha\beta}n_{\alpha}n_{\beta} (2\eta_{\alpha\beta} - 7\theta_{\alpha\beta} + 9)}{35(\eta_{\alpha\beta} + 1)^4(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \\
& \mathcal{A}_{\alpha\alpha}^{1143,2} = -\frac{64\eta_{\alpha\beta}(\theta_{\alpha\beta} - 1)n_{\alpha}n_{\beta}}{35(\eta_{\alpha\beta} + 1)^4(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (44)
\end{aligned}$$

V. THE 21N-MOMENT COEFFICIENTS WITH THE COULOMB POTENTIAL

Following Zhdanov[15], we choose the value of the Chapman-Cowling integrals $\Omega_{\alpha\beta}^{lr}$ for Coulomb potential with Debye cutoff as follows

$$\Omega_{\alpha\beta}^{lr} = \sqrt{\pi}l(r-1)! \left(\frac{2\pi}{\gamma_{\alpha\beta}} \right)^{1/2} \frac{\Delta_{\alpha\beta}^2}{4} \ln \Lambda_{\alpha\beta}, \quad (45)$$

where the Coulomb logarithm $\ln \Lambda_{\alpha\beta}$ is given by

$$\Lambda_{\alpha\beta} = \frac{12\lambda_D}{\Delta_{\alpha\beta}}, \quad (46)$$

where the Debye length λ_D is given by

$$\lambda_D^{-2} = \sum_{\alpha} \frac{n_{\alpha} Z_{\alpha}^2 e^2}{\epsilon_0 k T_{\alpha}},$$

and the mean distance of closest approach is given by

$$\Delta_{\alpha\beta} = \frac{|Z_{\alpha} Z_{\beta}| e^2}{4\pi\epsilon_0} \frac{\gamma_{\alpha\beta}}{\mu_{\alpha\beta}} \quad (47)$$

where $\gamma_{\alpha\beta}$ and $\mu_{\alpha\beta}$ are the reduced- γ ($= \gamma_{\alpha}\gamma_{\beta}/(\gamma_{\alpha} + \gamma_{\beta})$) and reduced mass respectively.

Although, this form of cross section is not fully accurate (there are some corrections to the Coulomb logarithm which can be found in Refs.[13, 20, 21]), it is a reasonable approximation to the order of the Coulomb logarithm within the 21N-moment scheme, and they allow us to simplify the coefficients in the previous section significantly. Furthermore, we define a $\lambda_{\alpha\beta, coulomb}$ such that

$$\lambda_{\alpha\beta, coulomb} = \sqrt{\pi} \left(\frac{2\pi}{\gamma_{\alpha\beta}} \right)^{1/2} \frac{\Delta_{\alpha\beta}^2}{4} \ln \Lambda_{\alpha\beta}. \quad (48)$$

This value of $\lambda_{\alpha\beta, coulomb}$ will remain a common factor in all Coulomb collision coefficients. We now present the coefficients divided by $\lambda_{\alpha\beta, coulomb}$ for clarity, as follows.

A. Value of $C_{\alpha\beta}^{010}/\lambda_{\alpha\beta, coulomb}$ for $p, q = 0$

$$p = 1, q = 0 : - \frac{16k\eta_{\alpha\beta} (\theta_{\alpha\beta} - 1) n_{\alpha} n_{\beta} T_{\alpha}}{(\eta_{\alpha\beta} + 1)^2 \theta_{\alpha\beta}} \quad (49)$$

B. Values of $B_{\alpha\beta}^{1pq}/\lambda_{\alpha\beta, coulomb}$ for $p, q \leq 2$

$$p = 0, q = 0 : \frac{16\eta_{\alpha\beta} n_{\alpha} n_{\beta}}{3\eta_{\alpha\beta} + 3} \quad (50)$$

$$p = 0, q = 1 : - \frac{16\eta_{\alpha\beta}^2 m_{\beta} n_{\alpha} n_{\beta}}{5k (\eta_{\alpha\beta} + 1) T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (51)$$

$$p = 1, q = 0 : - \frac{8k\eta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta} (3\theta_{\alpha\beta} - 2) + \theta_{\alpha\beta})}{(\eta_{\alpha\beta} + 1)^2 m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (52)$$

$$p = 1, q = 1 : \frac{24\eta_{\alpha\beta} \theta_{\alpha\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (5\theta_{\alpha\beta} - 2) + 3\theta_{\alpha\beta})}{5 (\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (53)$$

$$p = 0, q = 2 : \frac{8\eta_{\alpha\beta}^3 m_{\beta}^2 n_{\alpha} n_{\beta}}{7k^2 (\eta_{\alpha\beta} + 1) T_{\beta}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (54)$$

$$p = 2, q = 0 : \frac{20k^2 \eta_{\alpha\beta} \theta_{\alpha\beta} n_{\alpha} n_{\beta} T_{\alpha}^2 (\eta_{\alpha\beta} (5\theta_{\alpha\beta} - 4) + \theta_{\alpha\beta})}{(\eta_{\alpha\beta} + 1)^2 m_{\alpha}^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (55)$$

$$p = 1, q = 2 : - \frac{12\eta_{\alpha\beta}^2 \theta_{\alpha\beta} m_{\beta} n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (7\theta_{\alpha\beta} - 2) + 5\theta_{\alpha\beta})}{7k (\eta_{\alpha\beta} + 1)^2 T_{\beta} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \quad (56)$$

$$p = 2, q = 1 : - \frac{20k\eta_{\alpha\beta} \theta_{\alpha\beta}^2 n_{\alpha} n_{\beta} T_{\alpha} (\eta_{\alpha\beta} (7\theta_{\alpha\beta} - 4) + 3\theta_{\alpha\beta})}{(\eta_{\alpha\beta} + 1)^2 m_{\alpha} (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \quad (57)$$

$$p = 2, q = 2 : \frac{10\eta_{\alpha\beta} \theta_{\alpha\beta}^3 n_{\alpha} n_{\beta} (\eta_{\alpha\beta} (9\theta_{\alpha\beta} - 4) + 5\theta_{\alpha\beta})}{(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \quad (58)$$

C. Values of $A_{\alpha\beta}^{1pq}/\lambda_{\alpha\beta,coulomb}$ for $p, q \leq 2$

$$p = 0, q = 0 : -\frac{16n_\alpha n_\beta}{3\eta_{\alpha\beta} + 3} \quad (59)$$

$$p = 0, q = 1 : \frac{16\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta}{5k(\eta_{\alpha\beta} + 1)T_\alpha(\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (60)$$

$$p = 1, q = 0 : -\frac{8kn_\alpha n_\beta T_\alpha \left(10\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta} (\theta_{\alpha\beta} - 4) \theta_{\alpha\beta} - 3\theta_{\alpha\beta}^2 \right)}{3(\eta_{\alpha\beta} + 1)^2 \theta_{\alpha\beta} m_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (61)$$

$$p = 1, q = 1 : -\frac{8n_\alpha n_\beta \left(\eta_{\alpha\beta}^2 (52 - 6\theta_{\alpha\beta}) + \eta_{\alpha\beta} \theta_{\alpha\beta} (9\theta_{\alpha\beta} + 20) + 30\eta_{\alpha\beta}^3 + 13\theta_{\alpha\beta}^2 \right)}{15(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (62)$$

$$p = 0, q = 2 : -\frac{8\theta_{\alpha\beta}^2 m_\alpha^2 n_\alpha n_\beta}{7k^2 (\eta_{\alpha\beta} + 1) T_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (63)$$

$$p = 2, q = 0 : \frac{4k^2 n_\alpha n_\beta T_\alpha^2 \left(28\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 1) + \eta_{\alpha\beta} \theta_{\alpha\beta} (3\theta_{\alpha\beta} - 8) - 5\theta_{\alpha\beta}^2 \right)}{(\eta_{\alpha\beta} + 1)^2 m_\alpha^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (64)$$

$$p = 1, q = 2 : \frac{4\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta \left(-2\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 59) + \eta_{\alpha\beta} \theta_{\alpha\beta} (19\theta_{\alpha\beta} + 36) + 84\eta_{\alpha\beta}^3 + 23\theta_{\alpha\beta}^2 \right)}{35k(\eta_{\alpha\beta} + 1)^2 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \quad (65)$$

$$p = 2, q = 1 : -\frac{4kn_\alpha n_\beta T_\alpha}{15(\eta_{\alpha\beta} + 1)^2 \theta_{\alpha\beta} m_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \left(280\eta_{\alpha\beta}^4 (\theta_{\alpha\beta} - 1) + 84\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 4) \theta_{\alpha\beta} \right. \\ \left. + 348\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 2) \theta_{\alpha\beta}^2 + \eta_{\alpha\beta} \theta_{\alpha\beta}^3 (19\theta_{\alpha\beta} - 184) - 69\theta_{\alpha\beta}^4 \right) \quad (66)$$

$$p = 2, q = 2 : -\frac{2n_\alpha n_\beta}{105(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^4} \left(-56\eta_{\alpha\beta}^4 (2\theta_{\alpha\beta} - 59) + 8\eta_{\alpha\beta}^3 \theta_{\alpha\beta} (303\theta_{\alpha\beta} + 380) \right. \\ \left. + 4\eta_{\alpha\beta}^2 (1329 - 139\theta_{\alpha\beta}) \theta_{\alpha\beta}^2 + \eta_{\alpha\beta} \theta_{\alpha\beta}^3 (233\theta_{\alpha\beta} + 1288) + 1400\eta_{\alpha\beta}^5 + 433\theta_{\alpha\beta}^4 \right) \quad (67)$$

D. Values of $B_{\alpha\beta}^{2pq}/\lambda_{\alpha\beta,coulomb}$ for $p, q \leq 1$

$$p = 0, q = 0 : \frac{32\eta_{\alpha\beta} n_\alpha n_\beta (\eta_{\alpha\beta} (3\theta_{\alpha\beta} - 1) + 2\theta_{\alpha\beta})}{15(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (68)$$

$$p = 0, q = 1 : -\frac{32\eta_{\alpha\beta}^2 m_\beta n_\alpha n_\beta (\eta_{\alpha\beta} (5\theta_{\alpha\beta} - 1) + 4\theta_{\alpha\beta})}{35k(\eta_{\alpha\beta} + 1)^2 T_\beta (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (69)$$

$$p = 1, q = 0 : -\frac{32k\eta_{\alpha\beta} \theta_{\alpha\beta} n_\alpha n_\beta T_\alpha (\eta_{\alpha\beta} (5\theta_{\alpha\beta} - 3) + 2\theta_{\alpha\beta})}{5(\eta_{\alpha\beta} + 1)^2 m_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (70)$$

$$p = 1, q = 1 : \frac{32\eta_{\alpha\beta} \theta_{\alpha\beta}^2 n_\alpha n_\beta (\eta_{\alpha\beta} (7\theta_{\alpha\beta} - 3) + 4\theta_{\alpha\beta})}{7(\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \quad (71)$$

E. Values for $A_{\alpha\beta}^{2pq}/\lambda_{\alpha\beta,coulomb}$ for $p, q \leq 1$

$$p = 0, q = 0 : - \frac{32n_\alpha n_\beta \left(2\eta_{\alpha\beta} (\theta_{\alpha\beta} + 3) + 5\eta_{\alpha\beta}^2 + 3\theta_{\alpha\beta} \right)}{15 (\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})} \quad (72)$$

$$p = 0, q = 1 : \frac{32\theta_{\alpha\beta} m_\alpha n_\alpha n_\beta \left(2\eta_{\alpha\beta} (\theta_{\alpha\beta} + 4) + 7\eta_{\alpha\beta}^2 + 3\theta_{\alpha\beta} \right)}{35k (\eta_{\alpha\beta} + 1)^2 T_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (73)$$

$$p = 1, q = 0 : - \frac{16kn_\alpha n_\beta T_\alpha \left(35\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} - 1) + 7\eta_{\alpha\beta}^2 (\theta_{\alpha\beta} - 4) \theta_{\alpha\beta} + 2\eta_{\alpha\beta} (\theta_{\alpha\beta} - 16) \theta_{\alpha\beta}^2 - 9\theta_{\alpha\beta}^3 \right)}{15 (\eta_{\alpha\beta} + 1)^2 \theta_{\alpha\beta} m_\alpha (\eta_{\alpha\beta} + \theta_{\alpha\beta})^2} \quad (74)$$

$$p = 1, q = 1 : - \frac{16n_\alpha n_\beta \left(21\eta_{\alpha\beta}^3 (\theta_{\alpha\beta} + 13) + 3\eta_{\alpha\beta}^2 \theta_{\alpha\beta} (39\theta_{\alpha\beta} + 74) + 2\eta_{\alpha\beta} \theta_{\alpha\beta}^2 (13\theta_{\alpha\beta} + 105) + 140\eta_{\alpha\beta}^4 + 51\theta_{\alpha\beta}^3 \right)}{105 (\eta_{\alpha\beta} + 1)^2 (\eta_{\alpha\beta} + \theta_{\alpha\beta})^3} \quad (75)$$

VI. SUMMARY

In this article, we have provided an overview of the calculation of the collisional coefficients, and then provided the collisional coefficients for the Boltzmann collision operator for the $21N$ -moment multi-temperature case, both generally and specifically for the Coulomb potential with Debye cutoff. This fills an important gap in existing literature. Furthermore, we hope that these coefficients will lead to an easy implementation in existing state-of-the-art plasma fluid packages.

-
- [1] Bufferand H, et al. Three-dimensional modelling of edge multi-component plasma taking into account realistic wall geometry. *Nuclear Materials and Energy*. 2019 Jan;18:82–86.
 - [2] Zhdanov V, Kagan Y, Sazykin A. Effect of viscous transfer of momentum on diffusion in a gas mixture. *Sov Phys JETP*. 1962;15:596–602.
 - [3] Zhdanov VM, Yushmanov PN. Diffusion and heat transfer in a multicomponent completely ionized plasma. *Journal of Applied Mechanics and Technical Physics*. 1980 Jul;21(4):453–461.
 - [4] Grad H. Asymptotic Theory of the Boltzmann Equation. *The Physics of Fluids*. 1963 Feb;6(2):147–181.
 - [5] Braginskii SI. Transport Processes in a Plasma. *Reviews of Plasma Physics*. 1965 Jan;1:205.
 - [6] Bufferand H, Bensiali B, Bucalossi J, Ciraolo G, Genesio P, Ghendrih P, et al. Near wall plasma simulation using penalization technique with the transport code SolEdge2D-Eirene. *Journal of Nuclear Materials*. 2013 Jul;438:S445–S448.
 - [7] Bufferand H, Balbin J, Baschetti S, Bucalossi J, Ciraolo G, Gallo A, et al. Implementation of multi-component Zhdanov closure in SOLEDGE3X. Submitted to *Plasma Physics and Controlled Fusion*. 2021 Jul; Available from: <https://hal.archives-ouvertes.fr/hal-03243371>.
 - [8] Rozhansky V, et al. Momentum balance for impurities in SOLPS transport code. *Journal of Nuclear Materials*. 2015 Aug;463:477–479.
 - [9] Sytova E, et al. Impact of a new general form of friction and thermal forces on SOLPS-ITER modelling results. *Contributions to Plasma Physics*. 2018;58(6-8):622–628.
 - [10] Makarov SO, Coster DP, Rozhansky VA, Stepanenko AA, Zhdanov VM, Kaveeva EG, et al. Equations and improved coefficients for parallel transport in multicomponent collisional plasmas: Method and application for tokamak modeling. *Physics of Plasmas*. 2021 Jun;28(6):062308. Publisher: American Institute of Physics. Available from: <https://aip.scitation.org/doi/full/10.1063/5.0047618>.
 - [11] Fichtmüller M, et al. Multi-Species Developments in the EDGE2D Code. *Contributions to Plasma Physics*. 1998;38(1-2):284–289.
 - [12] Bergmann A, et al. Implementation into B2 of a 21-moment description for the parallel transport. *Contributions to Plasma Physics*. 1996;36(2-3):192–196.
 - [13] Raghunathan M, Marandet Y, Bufferand H, Ciraolo G, Ghendrih P, Tamain P, et al. Generalized collisional fluid theory for multi-component, multi-temperature plasma using the linearized Boltzmann collision operator for scrape-off layer/edge applications. *Plasma Physics and Controlled Fusion*. 2021 May;63(6):064005. Publisher: IOP Publishing. Available from: <https://doi.org/10.1088/1361-6587/abf670>.
 - [14] Raghunathan M, Marandet Y, Bufferand H, Ciraolo G, Ghendrih P, Tamain P, et al. Multi-temperature Generalized Zhdanov Closure for Scrape-Off Layer/Edge Applications. Submitted to *Plasma Physics and Controlled Fusion* (arXiv:210708940 [physics]). 2021 Jul; Available from: <http://arxiv.org/abs/2107.08940>.
 - [15] Zhdanov VM. *Transport processes in multicomponent plasma*. London: Taylor and Francis; 2002.
 - [16] Balescu R. *Transport processes in plasmas*. vol. 1. Amsterdam (Netherlands): North-Holland; 1988.
 - [17] Helander P, Sigmar DJ. *Collisional transport in magnetized plasmas*. vol. 4. Cambridge University Press; 2005.

- [18] Ji JY, Held ED. Exact linearized Coulomb collision operator in the moment expansion. *Physics of Plasmas*. 2006 Oct;13(10):102103.
- [19] Weinert U. Spherical tensor representation. *Archive for Rational Mechanics and Analysis*. 1980 Aug;74(2):165–196.
- [20] Liboff RL. Transport Coefficients Determined Using the Shielded Coulomb Potential. *The Physics of Fluids*. 1959 Jan;2(1):40–46. Publisher: American Institute of Physics.
- [21] Capitelli M, Bruno D, Laricchiuta A. Transport Cross Sections: Classical and Quantum Approaches. In: Capitelli M, Bruno D, Laricchiuta A, editors. *Fundamental Aspects of Plasma Chemical Physics: Transport*. Springer Series on Atomic, Optical, and Plasma Physics. New York, NY: Springer; 2013. p. 57–98.