

# The rational cohomology groups of the classifying spaces of Kac-Moody groups\*

Zhao Xu-an, zhaoxa@bnu.edu.cn

Gao Hongzhu, hzgao@bnu.edu.cn

School of Mathematical Sciences, Beijing Normal University

Key Laboratory of Mathematics and Complex Systems

Ministry of Education, China, Beijing 100875

Ruan Yangyang, ruanyy@amss.ac.cn

Institute of Mathematics, Chinese Academy of Sciences

China, Beijing 100190

## Abstract

In this paper, we compute the rational cohomology groups of the classifying space of a simply connected Kac-Moody group of infinite type. The fundamental principle is “from finite to infinite”. That is, for a Kac-Moody group  $G(A)$  of infinite type, the input data for computation are the rational cohomology of classifying spaces of parabolic subgroups of  $G(A)$  (which are of finite type), and the homomorphisms induced by inclusions of these subgroups. In some special cases, we can further determine the cohomology rings. Our method also applies to study the mod  $p$  cohomology of the classifying spaces of Kac-Moody groups.

## 1 Introduction

Let  $A = (a_{ij})$  be an  $n \times n$  integer matrix which satisfies:

- (1) For each  $i$ ,  $a_{ii} = 2$ ;
- (2) For  $i \neq j$ ,  $a_{ij} \leq 0$ ;
- (3) If  $a_{ij} = 0$ , then  $a_{ji} = 0$ .

then  $A$  is called a Cartan matrix.

Let  $h$  be the rational linear space with basis  $\Pi^\vee = \{\alpha_1^\vee, \alpha_2^\vee, \dots, \alpha_n^\vee\}$ , and denote the dual basis of  $\Pi^\vee$  in the dual space  $h^*$  by  $\{\omega_1, \omega_2, \dots, \omega_n\}$ , that is,  $\omega_i(\alpha_j^\vee) = \delta_{ij}$  for  $1 \leq i, j \leq n$ . Let  $\Pi = \{\alpha_1, \dots, \alpha_n\} \subset h^*$  be given by  $\alpha_j(\alpha_i^\vee) = a_{ij}$  for all  $i, j$ , then  $\alpha_j = \sum_{i=1}^n a_{ij}\omega_i$ .  $\Pi$  and  $\Pi^\vee$  are called the simple root system and simple co-root system associated to the Cartan matrix  $A$ .  $\alpha_i, \alpha_i^\vee, \omega_i, 1 \leq i \leq n$  are called respectively the simple roots, simple co-roots and fundamental dominant weights.

For each Cartan matrix  $A$ , there is a Lie algebra  $g(A)$  associated to  $A$  called Kac-Moody Lie algebra, see Kac[6][7] and Moody[14].

---

\*The authors are supported by National Science Foundation of China, 12071034.

The Kac-Moody Lie algebra  $g(A)$  is generated by elements  $\alpha_i^\vee, e_i, f_i, 1 \leq i \leq n$  over  $\mathbb{C}$ , with the defining relations:

- (1)  $[\alpha_i^\vee, \alpha_j^\vee] = 0, 1 \leq i, j \leq n;$
- (2)  $[e_i, f_j] = \delta_{ij} \alpha_i^\vee, 1 \leq i, j \leq n;$
- (3)  $[\alpha_i^\vee, e_j] = a_{ij} e_j, [\alpha_i^\vee, f_j] = -a_{ij} f_j, 1 \leq i, j \leq n;$
- (4)  $\text{ad}(e_i)^{-a_{ij}+1}(e_j) = 0, 1 \leq i \neq j \leq n;$
- (5)  $\text{ad}(f_i)^{-a_{ij}+1}(f_j) = 0, 1 \leq i \neq j \leq n.$

Kac and Peterson[8][9][10] constructed the Kac-Moody group  $G(A)$  with Lie algebra  $g(A)$ .

A Cartan matrix  $A$  is indecomposable if  $A$  can not be written as a direct sum of two Cartan matrices  $A_1$  and  $A_2$ . Indecomposable Cartan matrices, their associated Kac-Moody Lie algebras, or Kac-Moody groups are divided into three types.

- (1) Finite type, if  $A$  is positive definite. In this case,  $G(A)$  is just a simply connected complex semisimple Lie group.
- (2) Affine type, if  $A$  is positive semi-definite and has rank  $n - 1$ .
- (3) Indefinite type, otherwise.

The last two types of Cartan matrices or Kac-Moody groups are also called of infinite type. A decomposable Kac-Moody group is isomorphic to a product of Kac-Moody groups of these three types.

The Weyl group  $W(A)$  associated to a Cartan matrix  $A$  is the group generated by the Weyl reflections  $\sigma_i : h^* \rightarrow h^*$  with respect to  $\alpha_i^\vee, 1 \leq i \leq n$ , where  $\sigma_i(\alpha) = \alpha - \alpha(\alpha_i^\vee)\alpha_i$ . The action of  $\sigma_i$  on the fundamental dominant weights is given by  $\sigma_i(\omega_j) = \omega_j - \omega_j(\alpha_i^\vee)\alpha_i = \omega_j - \delta_{ij}\alpha_i$ . For details see Kac[7].

Let  $S$  be the set  $\{1, 2, \dots, n\}$ . For each proper subset  $I \subset S$ , the matrix  $A_I = (a_{ij})_{i,j \in I}$  is also a Cartan matrix. And there is a parabolic subalgebra  $g_I(A) \subseteq g(A)$  generated by  $e_i, i \in I$  and  $\alpha_j^\vee, f_j, 1 \leq j \leq n$ . Corresponding to  $g_I(A)$ , there is a connected parabolic subgroup  $G_I(A)$  of  $G(A)$ . For  $I = S$ ,  $G_I(A) = G(A)$ . And for  $I = \emptyset$ ,  $G_I(A) = B(A)$ , the Borel subgroup of  $G(A)$ .

Let  $C(A)$  be the category whose objects are proper subsets  $I$  of  $S$  such that  $A_I$  are of finite type, and morphisms are the inclusions of these proper subsets. There is a functor  $F : C(A) \rightarrow Top$  which sends  $I$  to the classifying space  $BG_I(A)$  and sends  $I \subset J$  to the map  $BG_I(A) \rightarrow BG_J(A)$ . By a theorem of Kichiloo[12][13], there is

**Theorem.** For a Cartan matrix  $A$  of infinite type, the homotopy type of  $BG(A)$  is the homotopy colimit of the functor  $F$ , i.e.  $BG(A) \simeq \text{hocolimit} F$ .

This theorem enables us to compute the cohomology of  $BG(A)$  from the cohomology of  $BG_I(A)$  for those  $A_I$  of finite type. It gives a principle that “from finite to infinite”. By Bousfield and Kan[2], there is a multiplicative spectral sequence with  $E_2$ -term given by the derived limits of the functor  $H^* \circ F$  which converges to the cohomology of  $BG(A)$ . This spectral sequence is used in Kitchiloo[13] to study the cohomology of the classifying spaces of Kac-Moody groups. But the  $E_2$ -term of Bousfield-Kan spectral sequence is hard to compute. So in this paper we introduce a homotopy simplification procedure. In fact we show that there are spaces  $X; X_{I_1}, X_{I_2}, \dots, X_{I_d}$  for certain  $d \geq 2$  such that  $X = X_{I_1} \cup X_{I_2} \cup \dots \cup X_{I_d}$ ,  $X$  has the same homotopy type as  $BG(A)$ , and for  $1 \leq i \leq d$ ,  $X_{I_i}$  has the same homotopy type as  $BG_{I_i}(A)$ . These spaces  $X$  and  $X_{I_i}, 1 \leq i \leq d$  are constructed explicitly in section 2. As a consequence we can use Mayer-Vietoris spectral sequence

of  $X = X_{I_1} \cup X_{I_2} \cup \cdots \cup X_{I_d}$  to compute the cohomology groups of  $BG(A)$ .

The contents of this paper are as follows. In section 2, we simplify the homotopy colimit of the classifying space  $BG(A)$  and construct the spaces  $X; X_{I_1}, X_{I_2}, \cdots, X_{I_d}$ . In section 3 we use the Mayer-Vietoris spectral sequence to compute the rational cohomology groups of  $BG(A)$ . We will explain how to compute the  $E_2$ -term of this spectral sequence. And we prove that for rational coefficients the spectral sequence collapses at  $E_2$ -term. So it gives the cohomology groups of  $BG(A)$ . In section 4 we give an algorithm to compute the cohomology groups and list the computation results for the cases of  $n = 3$  and 4. In section 5 we give some results about the rational polynomial invariants of Weyl groups  $W(A)$ , which appear naturally in the M-V spectral sequence. In section 6 we discuss some applications of our results and give some conjectures on the cohomology of classifying spaces of Kac-Moody groups.

## 2 The simplification of the homotopy colimit

Let us recall the definition of homotopy colimit. See Dugger[4] for details.

Let  $C$  be a small category and  $F : C \rightarrow Top$  be a functor from  $C$  to the category of topological spaces. Then we have a simplicial space  $\text{srep}(F)$  which is called the simplicial replacement of  $F$  with  $\text{srep}(F)_n = \coprod_{i_0 \rightarrow i_1 \rightarrow \cdots \rightarrow i_n} F(i_0)$ , the coproduct ranges over chains of composable morphisms in  $C$ . The face and degeneracy maps are defined as follows. If  $\sigma : i_0 \rightarrow i_1 \rightarrow \cdots \rightarrow i_n$  is a chain, then by deleting  $i_j, j \geq 0$  we obtain a chain  $\sigma(j)$  of  $n - 1$  composable maps. When  $j > 0$ , the map  $d_j : \text{srep}(F)_n \rightarrow \text{srep}(F)_{n-1}$  sends the summand  $F(i_0)$  corresponding to the chain  $\sigma$  to the identical copy of  $F(i_0)$  in  $\text{srep}(F)_{n-1}$  indexed by  $\sigma(j)$ . When  $j = 0$ ,  $d_0 : \text{srep}(F)_n \rightarrow \text{srep}(F)_{n-1}$  sends the summand  $F(i_0)$  corresponding to the chain  $\sigma$  to the summand  $F(i_1)$  corresponding to  $\sigma(0)$  and the map we use here is  $F(i_0) \rightarrow F(i_1)$  given by the first map  $i_0 \rightarrow i_1$  in  $\sigma$ . For the degeneracy maps each  $s_j$  sends the summand  $F(i_0)$  corresponding to the chain  $\sigma$  to the identical summand  $F(i_0)$  corresponding to the chain in which one has inserted the identity map  $i_j \rightarrow i_j$ . The homotopy colimit of the functor  $F : C \rightarrow Top$  is defined as the geometric realization of its simplicial replacement  $\text{srep}(F)$ .

For a simplicial complex  $L$  with vertex set  $S = \{1, 2, \cdots, n\}$ , there is a small category  $C(L)$  with objects the simplices of  $L$ . And the morphisms of  $C(L)$  are given by inclusions of simplices. We called the category  $C(L)$  the simplicial category associated to  $L$ . In a simplicial category  $C(L)$  all the automorphisms are identities, and for two objects there is at most one morphism between them. Given a Cartan matrix  $A$ , the category  $C(A)$  defined in the introduction is a simplicial category since if  $A_J$  is of finite type, then for any  $I \subset J$ ,  $A_I$  is also of finite type. And for each  $i \in S$ ,  $\{i\} \in C(A)$  because the rank 1 Cartan matrix is of finite type. Therefore for a Kac-Moody group  $G(A)$ , it is convenient to regard the category  $C(A)$  as a proper simplicial sub-complex of the  $n - 1$ -simplex  $\Delta_{n-1}$  with vertex set  $S = \{1, 2, \cdots, n\}$ , and regard an object  $I \in C(A)$  as a simplex.

Let  $C$  be the simplicial category associated to a simplicial complex  $L$  and  $F : C \rightarrow Top$  be a functor, For an object  $I \in C$ , let  $C_I$  be the full subcategory of  $C$  whose objects are  $I$  and faces of  $I$ . We define  $X_I = \text{hocolimit} F|_{C_I}$ , the homotopy colimit of  $F$  restricted to  $C_I$ . Since  $C_I$  has a terminal object  $I$  we know  $X_I \simeq F(I)$ . The inclusion  $I \subset J$  induces the map  $i_{IJ} : X_I \rightarrow X_J$ . As a summary, we have

**Lemma 2.1.** *1. For  $I \subset J$ , the map  $i_{IJ} : X_I \rightarrow X_J$  is an inclusion(In fact a cofibration).*

2. For each  $I \in C$ ,  $X_I$  is a simplicial subspace of  $X = \text{hocolimit} F$ .
3.  $X_I \cap X_J = X_{I \cap J}$  for  $I, J \in C$ .

This lemma can be checked by the definition of homotopy colimits. It can be regarded as a homotopy replacement result for the functor  $F$ . By applying it to the functor  $F : C(A) \rightarrow \text{Top}$ , we can replace the map  $BG_I(A) \rightarrow BG_J(A)$  by a cofibration  $X_I \subset X_J$ , and the classifying space  $BG_I(A)$  by  $X_I$ . Then the homotopy colimit of  $F$  is replaced by a union of subspaces  $X_I$  in  $X$ .


We call a simplex  $\sigma$  of a simplicial complex  $L$  a maximal simplex if it is not a face of any other simplex. It is obvious that the geometric realizations of a simplicial complex is the union of geometric realization of its maximal simplices. For a finite simplicial category  $C = C(L)$ , the homotopy colimit of a functor  $F : C \rightarrow \text{Top}$  has a simple description.

**Theorem 2.1.** *Let  $L$  be a simplicial complex contained in  $\Delta_{n-1}$  and  $I_1, I_2, \dots, I_d$  be the set of maximal simplices of  $L$ , then for a functor  $F : C(L) \rightarrow \text{Top}$ , we have  $X = \text{hocolimit} F = \bigcup_{1 \leq i \leq d} X_{I_i}$ .*

Combining Theorem 2.1 with Kitchloo's theorem, we have

**Corollary 2.1.** *Let  $A$  be a Cartan matrix of infinite type,  $I_1, I_2, \dots, I_d \in C(A)$  be the maximal objects of  $C(A)$ , then  $BG(A) \simeq \bigcup_{1 \leq i \leq d} X_{I_i}$ .*

**Example 2.1.** *For a Cartan matrix  $A$  of rank 3, up to permutation of 1, 2, 3, the simplicial category  $C(A)$  corresponding to  $A$  is one of the following 4 types.*



In the four cases the maximal objects of  $C(A)$  and the homotopy type of  $BG(A)$  are

- (1).  $\{1\}, \{2\}, \{3\}$ ,  $BG(A) \simeq (X_1 \cup_{X_\emptyset} X_2) \cup_{X_\emptyset} X_3$ ;
- (2).  $\{1, 2\}, \{3\}$ ,  $BG(A) \simeq X_{12} \cup_{X_\emptyset} X_3$ ;
- (3).  $\{1, 2\}, \{1, 3\}$ ,  $BG(A) \simeq X_{12} \cup_{X_1} X_{13}$ ;
- (4).  $\{1, 2\}, \{1, 3\}, \{2, 3\}$ ,  $BG(A) \simeq (X_{12} \cup_{X_1} X_{13}) \cup_{X_2 \cup_{X_\emptyset} X_3} X_{23}$ .

### 3 Mayer-Vietoris spectral sequence

In section 2 we write the homotopy type of the classifying space  $BG(A)$  as a union of subspaces  $X_{I_1}, X_{I_2}, \dots, X_{I_d}$  which are homotopy equivalent to  $BG_{I_1}(A), BG_{I_2}(A), \dots, BG_{I_d}(A)$  respectively. Now we can use the Mayer-Vietoris exact sequences to compute the cohomology groups of  $BG(A)$  inductively. The computation procedure can be organized into an M-V spectral sequence. The spectral sequence  $E_r, r \geq 0$  is constructed from the bi-complex  $E_0^{r,s} = \bigoplus_{i_1 < i_2 < \dots < i_{r+1}} S_{\mathcal{U}}^s(X_{I_{i_1}} \cap X_{I_{i_2}} \cap \dots \cap X_{I_{i_{r+1}}})$

$\cdots \cap X_{I_{i_{r+1}}}$ ) with horizontal differential  $\delta$  and vertical differential  $d$ . The  $E_1$ -term and  $E_2$ -term of this spectral sequence are given by  $H^d(E_0)$  and  $H^\delta E_1 = H^\delta H^d(E_0)$ . This spectral sequence converges to  $H^*(X)$ . See Bott and Tu[3] for details. By applying it to  $X_{I_1} \cup X_{I_2} \cup \cdots \cup X_{I_d}$ , we have

**Theorem 3.1.** *For arbitrary abelian group  $G$ , there is an M-V spectral sequence with  $E_1^{r,s} = \bigoplus_{i_1 < i_2 < \cdots < i_{r+1}} H^s(X_{I_{i_1}} \cap X_{I_{i_2}} \cap \cdots \cap X_{I_{i_{r+1}}}; G)$ , which converges to the cohomology  $H^*(BG(A); G)$ .*

As a consequence, with coefficient group a field  $k$ , we have  $H^*(BG(A); k) \cong E_\infty^{0,*} \oplus E_\infty^{1,*-1} \oplus \cdots \oplus E_\infty^{d-1,0}$ . From the expression of  $E_1^{r,s}$  it is obvious that for  $r \geq d$ ,  $E_1^{r,s} = 0$ . So the spectral sequence is concentrated on the range of  $0 \leq r \leq d-1$ . Let  $E_\infty^r = \bigoplus_{s=0}^\infty E_\infty^{r,s}$  for  $r \geq 0$ , then  $E_\infty^r = 0$

for  $r \geq d$  and we have  $H^*(BG(A); k) \cong \bigoplus_{r=0}^{d-1} \Sigma^r E_\infty^r$  as a graded  $k$ -vector space, where  $\Sigma^r$  lifts the degree of a graded vector space by  $r$ .

In the following all cohomology groups are of rational coefficients.

The input data to compute the rational cohomology  $H^*(BG(A))$  come from the functor  $H^* \circ F : C(A) \rightarrow \mathbb{Q}Alg$  which sends  $I \in C(A)$  to  $H^*(BG_I(A))$  and sends a morphism  $I \subset J$  to  $H^*(BG_J(A)) \rightarrow H^*(BG_I(A))$ . The ring  $H^*(BG_I(A))$  and homomorphism  $H^*(BG_J(A)) \rightarrow H^*(BG_I(A))$  can be computed by the following theorem of Borel, see [1].

**Theorem 3.2.** *Let  $K$  be a connected compact Lie group with Weyl group  $W(K)$ , and  $T$  be a maximal torus subgroup of  $K$ , then the cohomology homomorphism  $Bi^* : H^*(BK; \mathbb{Q}) \rightarrow H^*(BT; \mathbb{Q})$  induced by the inclusion  $T \rightarrow K$  is injective. And its image is exactly the invariants  $H^*(BT; \mathbb{Q})^{W(K)}$  of Weyl group.*

For a parabolic subgroup  $G_I(A)$  of finite type, the unitary form of  $G_I(A)$  is a compact Lie subgroup  $K_I(A)$  of the unitary form  $K(A)$  of  $G(A)$ . And the inclusion  $K_I(A) \subset G_I(A)$  is a homotopy equivalence. Hence for the computation of the cohomology we can replace the group  $G_I(A)$  by its unitary form  $K_I(A)$ . Since the Weyl group of  $G_I(A)$ (or  $K_I(A)$ ) is  $W_I(A)$ , which is generated by Weyl reflection  $\sigma_i, i \in I$ , the ring  $H^*(BG_I(A)) \cong H^*(BK_I(A))$  can be identified with  $H^*(BT)^{W_I(A)}$  by Borel's theorem. For convenience we write  $P$  for the polynomial algebra  $H^*(BT)$ , which is naturally identified with  $\mathbb{Q}[w_1, \dots, w_n]$ , where  $w_i, 1 \leq i \leq n$  are the fundamental dominant weights. And for each  $I \subset S$  we write  $P_I$  for the polynomial subalgebra  $H^*(BT)^{W_I(A)}$ . If we identify  $H^*(BG_I(A))$  with  $P_I$ , then the homomorphism  $Bi_{I,J}^* : H^*(BG_J(A)) \rightarrow H^*(BG_I(A))$  for  $I \subset J$  can be identified with the inclusion  $P_J = H^*(BT)^{W_J(A)} \subset H^*(BT)^{W_I(A)} = P_I$ . An easy but useful fact is that  $P_{I_1} \cap P_{I_2} = P_{I_1 \cup I_2}$ .


Let  $E_1^r = \bigoplus_{s=0}^\infty E_1^{r,s}$  for  $0 \leq r \leq d-1$ . From the above discussion we have

$$E_1^r \cong \bigoplus_{s=0}^\infty \bigoplus_{i_1 < i_2 < \cdots < i_{r+1}} H^s(X_{I_{i_1}} \cap X_{I_{i_2}} \cap \cdots \cap X_{I_{i_{r+1}}}; \mathbb{Q}) \cong \bigoplus_{i_1 < i_2 < \cdots < i_{r+1}} P_{I_{i_1 \cup I_{i_2} \cup \cdots \cup I_{i_{r+1}}}}.$$

The  $E_2$ -term of the M-V spectral sequence can be computed from  $E_1$ -term as the homology of the following chain complex  $0 \rightarrow E_1^0 \xrightarrow{\delta_0} E_1^1 \xrightarrow{\delta_1} E_1^2 \xrightarrow{\delta_2} \cdots \xrightarrow{\delta_{d-2}} E_1^{d-1} \rightarrow 0$ .

The following is an example.

**Example 3.1.** For the case  $X = X_{I_1} \cup X_{I_2} \cup X_{I_3} \cup X_{I_4}$  with  $d = 4$ , the chain complex can be depicted as follows.



In this diagram we omit the sign  $\oplus$  between the adjacent items in each row. For the description of the differential see Bott and Tu[3] or Example 4.1 in the next section.

For  $I \in C(A)$ , by Borel's theorem  $P_I$  is concentrated on the even degree. So we have  $E_1^{r,s} = \{0\}$  for  $s$  odd. Then we have  $E_k^{r,s} = \{0\}$  for  $s$  odd,  $k \geq 1$ . We need this observation to prove the following theorem.

**Theorem 3.3.** *The rational M-V spectral sequence for  $BG(A)$  collapses at  $E_2$ -term.*

**Proof:** By Kitchloo[13], let  $l$  be a prime number so that Weyl group  $W(A)$  contains no elements of order  $l$ , then there is an unstable Adams map  $\psi$  on  $BG(A)$ , and it induces the Adams map  $\psi_I$  on  $BG_I(A)$  for all  $I \in C(A)$ . The cohomology endomorphism  $\psi_I^*$  acting on  $H^{2q}(BG_I(A))$  for  $G_I(A)$  of finite type is the multiplication by  $l^q$ . The map  $\psi$  induces an endomorphism on the M-V spectral sequence. For even integer  $2k$ ,  $d_{2k}^{r,s} = 0$  since the domain or the range of  $d_{2k}^{p,q}$  is trivial by the observation  $E_{2k}^{r,s} = \{0\}$  for  $s$  odd,  $k \geq 1$ . For odd integer  $2k-1$ , if  $x \in E_{2k-1}^{r,2s}$ , since  $\psi^*$  is commutative with the differential of the M-V spectral sequence, we have  $d_{2k-1}\psi^*(x) = \psi^*d_{2k-1}(x)$ , hence we get  $l^s d_{2k-1}(x) = l^{s-k+1} d_{2k-1}(x)$ . As a result if  $k > 1$ , we have  $d_{2k-1}x = 0$ . This shows that we always have  $d_k(x) = 0$ , for  $k \geq 2$ . Therefore the M-V spectral sequence collapses.

**Remark 3.1.** *There are similar M-V spectral sequences for all coefficient groups other than  $\mathbb{Q}$  which converge to the associated cohomology group  $H^*(BG(A))$ . But in general they do not collapse at  $E_2$ -term.*

The explicit computation of the  $E_2$ -term of the rational M-V spectral sequence will be discussed in the next section.

## 4 The computation of rational cohomology groups of $BG(A)$

In this section we compute the rational cohomology groups of the classifying space  $BG(A)$ . Since for rational coefficient the corresponding M-V spectral sequence collapses at  $E_2$ -term, the  $E_2$ -term of the spectral sequence gives the rational cohomology groups of  $BG(A)$ .

The following is an example of explicit computation.

**Example 4.1.** We demonstrate the computation procedure for the  $M$ - $V$  spectral sequence of  $X = X_{I_1} \cup X_{I_2} \cup X_{I_3} \cup X_{I_4}$ . The cochain complex  $E_1 : 0 \rightarrow E_1^0 \xrightarrow{\delta_0} E_1^1 \xrightarrow{\delta_1} E_1^2 \xrightarrow{\delta_2} \dots \xrightarrow{\delta_{d-2}} E_1^{d-1} \rightarrow 0$  is illustrated as in Example 3.1. Here we compute the cohomology  $E_2 = H^*(E_1) = \bigoplus_{r=0}^{d-1} H^r(E_1)$ .

**Step 1:** It is easy to see that  $H^0(E_1) \cong P_{I_1} \cap P_{I_2} \cap P_{I_3} \cap P_{I_4} = P_{I_1 \cup I_2 \cup I_3 \cup I_4}$ .

**Step 2:** For  $H^1(E_1)$ , the element in  $\ker \delta_1$  has the form  $(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})$  with  $v_{ij} \in P_{I_i \cap I_j}$ ,  $1 \leq i < j \leq 4$  and satisfies

$$v_{12} - v_{13} + v_{23} = 0. \quad (1)$$

$$v_{12} - v_{14} + v_{24} = 0. \quad (2)$$

$$v_{13} - v_{14} + v_{34} = 0. \quad (3)$$

$$v_{23} - v_{24} + v_{34} = 0. \quad (4)$$

The image of  $\delta_0$  is generated by the elements of the forms  $\delta_0(v_i)$  with  $v_i \in P_{I_i}$ ,  $1 \leq i \leq 4$ , which is represented by the row vectors of the following matrix.

$$\begin{pmatrix} 12 & 13 & 14 & 23 & 24 & 34 \\ v_1 & v_1 & v_1 & 0 & 0 & 0 \\ -v_2 & 0 & 0 & -v_2 & -v_2 & 0 \\ 0 & -v_3 & 0 & -v_3 & 0 & v_3 \\ 0 & 0 & 0 & -v_4 & -v_4 & -v_4 \end{pmatrix}$$

By the equations  $v_{12} - v_{13} + v_{23} = 0$  and  $v_{12} - v_{14} + v_{24} = 0$ , we have  $v_{12} \in P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) \cap (P_{I_1 \cap I_4} + P_{I_2 \cap I_4})$ . We compute  $H^1(E_1)$  by constructing an epimorphism

$$\begin{aligned} \phi_1 : H_1(E_1) &\rightarrow P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) \cap (P_{I_1 \cap I_4} + P_{I_2 \cap I_4}) - (P_{I_1} + P_{I_2}) \\ &[(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})] \rightarrow v_{12} + P_{I_1} + P_{I_2}. \end{aligned}$$

Here we use  $A - B$  to denote the quotient of  $A$  with respect to subspace  $B$  for convenience. It is easy to see that  $\phi_1$  is well defined. And as a rational vector space  $H_1(E_1) \cong \text{im} \phi_1 \oplus \ker \phi_1$ .

Now we compute  $\ker \phi_1$ . For  $[(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})] \in \ker \phi_1$ , we must have  $v_{12} \in P_{I_1} + P_{I_2}$ . Therefore there exist  $v_1 \in P_{I_1}, v_2 \in P_{I_2}$  such that  $v_{12} = v_1 - v_2$ . Substituting  $v_{12}$  with  $v_1 - v_2$  in Equations 1 and 2, we get  $v_{13} - v_1 = v_{23} - v_2$  and  $v_{14} - v_1 = v_{24} - v_2$ . This means that  $v_{13} - v_1 \in P_{I_1 \cap I_3} \cap P_{I_2 \cap I_3} = P_{(I_1 \cup I_2) \cap I_3}$  and  $v_{14} - v_1 \in P_{I_1 \cap I_4} \cap P_{I_2 \cap I_4} = P_{(I_1 \cup I_2) \cap I_4}$ . By Equation 3,  $v_{13} - v_{14} + v_{34} = 0$ , we get  $(v_{13} - v_1) - (v_{14} - v_1) + v_{34} = 0$ . So  $v_{13} - v_1 \in P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4}$ . Therefore  $v_{13} - v_1 \in P_{(I_1 \cup I_2) \cap I_3} \cap (P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4})$ .

We define a homomorphism

$$\begin{aligned} \phi_2 : \ker \phi_1 &\rightarrow P_{(I_1 \cup I_2) \cap I_3} \cap (P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4}) - (P_{I_1 \cup I_2} + P_{I_3}) \\ &[(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})] \rightarrow v_{13} - v_1 + P_{I_1 \cup I_2} + P_{I_3}. \end{aligned}$$

It is easy to check that this homomorphism is well defined. Note that we need the  $P_{I_1 \cup I_2}$ -item to eliminate the effect of indeterminacy of  $v_1$ , and the  $P_{I_3}$ -item to send  $[(0, -v_3, 0, -v_3, 0, v_3)] \in \ker \delta_0$  to 0. By definition  $\phi_2$  is an epimorphism and we have  $\ker \phi_1 \cong \text{im} \phi_2 \oplus \ker \phi_2$ .

For  $[(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})] \in \ker \phi_2$ , we already have  $v_1 \in P_{I_1}$  and  $v_2 \in P_{I_2}$  such that  $v_{12} = v_1 - v_2$  for  $\ker \phi_2 \subset \ker \phi_1$ , and there are  $\alpha \in P_{I_1 \cup I_2}$  and  $v_3 \in P_{I_3}$  such that  $v_{13} - v_1 = \alpha - v_3$ . So we have  $v_{12} = (v_1 + \alpha) - (v_2 + \alpha)$  and  $v_{13} = (v_1 + \alpha) - v_3$ . This means that if we choose  $v'_1 = v_1 + \alpha$  and  $v'_2 = v_2 + \alpha$  instead of  $v_1, v_2$  in the previous step, then we will have  $v_{12} = v'_1 - v'_2$  and  $v_{13} = v'_1 - v_3$ . For simplicity we still use  $v_1, v_2$  to denote the newly chosen elements  $v'_1, v'_2$ , then  $\alpha$  disappears and we have  $v_{12} = v_1 - v_2$  and  $v_{13} = v_1 - v_3$ . Combining them with Equation 1, we can solve  $v_{23} = v_2 - v_3$ . By Equations 2, 3 and 4, we get  $v_{14} - v_1 = v_{24} - v_2 = v_{34} - v_3$ . These equations imply that  $v_{14} - v_1 \in P_{I_1 \cap I_4} \cap P_{I_2 \cap I_4} \cap P_{I_3 \cap I_4} = P_{(I_1 \cup I_2 \cup I_3) \cap I_4}$ .

Now we define an epimorphism

$$\begin{aligned} \phi_3 : \ker \phi_2 &\rightarrow P_{(I_1 \cup I_2 \cup I_3) \cap I_4} - (P_{I_1 \cup I_2 \cup I_3} + P_{I_4}) \\ [(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})] &\rightarrow v_{14} - v_1 + P_{I_1 \cup I_2 \cup I_3} + P_{I_4}. \end{aligned}$$

For  $[(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})] \in \ker \phi_3$ , we have  $v_{12} = v_1 - v_2, v_{13} = v_1 - v_3, v_{23} = v_2 - v_3$  and  $v_{14} - v_1 = \beta - v_4$  for certain  $v_i \in P_{I_i}, 1 \leq i \leq 4$  and  $\beta \in P_{I_1 \cup I_2 \cup I_3}$ . If we replace  $v_1, v_2, v_3$  in the previous step by  $v'_1 = v_1 + \beta, v'_2 = v_2 + \beta, v'_3 = v_3 + \beta$ , then  $\beta$  disappears, and we have  $v_{14} - v_1 = -v_4$ . As a result we get  $v_{14} = v_1 - v_4, v_{24} = v_2 - v_4$  and  $v_{34} = v_3 - v_4$  from the Equations 2, 3 and 4. Hence  $(v_{12}, v_{13}, v_{14}, v_{23}, v_{24}, v_{34})$  is in  $\text{im} \delta_0$  and  $\ker \phi_3 = 0$ . Hence  $\ker \phi_2 \cong \text{im} \phi_3 \oplus \ker \phi_3 \cong \text{im} \phi_3$ .

So we have

$$\begin{aligned} H^1(E_1) &\cong \text{im} \phi_1 \oplus \text{im} \phi_2 \oplus \text{im} \phi_3 \\ &\cong P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) \cap (P_{I_1 \cap I_4} + P_{I_2 \cap I_4}) - (P_{I_1} + P_{I_2}) \\ &\oplus P_{(I_1 \cup I_2) \cap I_3} \cap (P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4}) - (P_{I_1 \cup I_2} + P_{I_3}) \oplus P_{(I_1 \cup I_2 \cup I_3) \cap I_4} - (P_{I_1 \cup I_2 \cup I_3} + P_{I_4}) \end{aligned}$$

**Step 3:** For  $H^2(E_1)$ , the element in  $\ker \delta_2$  has the form  $(v_{123}, v_{124}, v_{134}, v_{234})$  with  $v_{ijk} \in P_{I_i \cap I_j \cap I_k}, 1 \leq i < j < k \leq 4$  and satisfies

$$v_{123} - v_{124} + v_{134} - v_{234} = 0. \quad (5)$$

The image of  $\delta_1$  is generated by all the elements of the forms  $\delta_1(v_{ij})$  with  $v_{ij} \in P_{I_i \cap I_j}, 1 \leq i, j \leq 4$ .

$$\begin{pmatrix} 123 & 124 & 134 & 234 \\ v_{12} & v_{12} & 0 & 0 \\ -v_{13} & 0 & v_{13} & 0 \\ 0 & -v_{14} & -v_{14} & 0 \\ v_{23} & 0 & 0 & v_{23} \\ 0 & v_{24} & 0 & -v_{24} \\ 0 & 0 & v_{34} & v_{34} \end{pmatrix}$$

We compute  $H^2(E_1)$  by constructing an epimorphism

$$\begin{aligned} \psi_1 : H^2(E_1) &\rightarrow P_{I_1 \cap I_2 \cap I_3} \cap (P_{I_1 \cap I_2 \cap I_4} + P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) - (P_{I_1 \cap I_2} + P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) \\ [(v_{123}, v_{124}, v_{134}, v_{234})] &\rightarrow v_{123} + P_{I_1 \cap I_2} + P_{I_1 \cap I_3} + P_{I_2 \cap I_3} \end{aligned}$$

Then as a vector space  $H^2(E_1) \cong \text{im} \psi_1 \oplus \ker \psi_1$ .



For  $[(v_{123}, v_{124}, v_{134}, v_{234})] \in \ker \psi_1$ , there are elements  $v_{12} \in P_{I_1 \cap I_2}$ ,  $v_{13} \in P_{I_1 \cap I_3}$  and  $v_{23} \in P_{I_2 \cap I_3}$  such that  $v_{123} = v_{12} - v_{13} + v_{23}$ . Combining with Equation 5, we get  $(v_{124} - v_{12}) - (v_{134} - v_{13}) + (v_{234} - v_{23}) = 0$ . This implies that  $v_{124} - v_{12} \in P_{I_1 \cap I_2 \cap I_4} \cap (P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4})$ .

We compute  $\ker \psi_1$  by constructing a homomorphism

$$\begin{aligned} \psi_2 : \ker \psi_1 &\rightarrow P_{I_1 \cap I_2 \cap I_4} \cap (P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) - (P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) + P_{I_1 \cap I_4} + P_{I_2 \cap I_4}) \\ &[(v_{123}, v_{124}, v_{134}, v_{234})] \rightarrow v_{124} - v_{12} + P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) + P_{I_1 \cap I_4} + P_{I_2 \cap I_4} \end{aligned}$$

It is easy to see that the homomorphism  $\psi_2$  is well defined. Note that we add the item  $P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3})$  to eliminate the effect of indeterminacy of  $v_{12}$ . By definition,  $\psi_2$  is an epimorphism. So as a vector space  $\ker \psi_1 \cong \text{im} \psi_2 \oplus \ker \psi_2$ .

For  $[(v_{123}, v_{124}, v_{134}, v_{234})] \in \ker \psi_2$ , a similar discussion as above shows that there are elements  $v_{12} \in P_{I_1 \cap I_2}$ ,  $v_{13} \in P_{I_1 \cap I_3}$  and  $v_{23} \in P_{I_2 \cap I_3}$  with  $v_{123} = v_{12} - v_{13} + v_{23}$ . And there are  $\gamma \in P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3})$ ,  $v_{14} \in P_{I_1 \cap I_4}$  and  $v_{24} \in P_{I_2 \cap I_4}$  such that  $v_{124} - v_{12} = \gamma - v_{14} + v_{24}$ . Similarly we can assume  $\gamma = 0$  if we choose  $v_{12}$  and  $v_{13}$  carefully in the previous step ( $v'_{12} = v_{12} + \gamma$  and  $v'_{13} = v_{13} + \gamma$ ). Combining with Equation 5, we get  $v_{134} - v_{13} + v_{14} = v_{234} - v_{23} + v_{24}$ . This shows that  $v_{134} - v_{13} + v_{14} \in P_{I_1 \cap I_3 \cap I_4} \cap P_{I_2 \cap I_3 \cap I_4} = P_{(I_1 \cup I_2) \cap I_3 \cap I_4}$ .

Now we can construct an epimorphism

$$\begin{aligned} \psi_3 : \ker \psi_2 &\rightarrow P_{(I_1 \cup I_2) \cap I_3 \cap I_4} - (P_{(I_1 \cup I_2) \cap I_3} + P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4}) \\ &[(v_{123}, v_{124}, v_{134}, v_{234})] \mapsto v_{134} - v_{13} + v_{14} + P_{(I_1 \cup I_2) \cap I_3} + P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4} \end{aligned}$$

We add the items  $P_{(I_1 \cup I_2) \cap I_3}$  and  $P_{(I_1 \cup I_2) \cap I_4}$  to eliminate the effect of indeterminacy of  $v_{13}$  and  $v_{14}$ . For  $[(v_{123}, v_{124}, v_{134}, v_{234})] \in \ker \psi_3$  we have  $v_{123} = v_{12} - v_{13} + v_{23}$ ,  $v_{124} = v_{12} - v_{14} + v_{24}$  and  $v_{134} = v_{13} - v_{14} + \epsilon + \theta + v_{34}$ . If we choose  $v'_{13} = v_{13} + \epsilon$ ,  $v'_{14} = v_{14} + \epsilon$  and  $v'_{23} = v_{23} + \theta$ ,  $v'_{24} = v_{24} + \theta$  in the previous step, then we have  $v_{134} = v_{13} - v_{14} + v_{34}$  and  $v_{234} = v_{23} - v_{24} + v_{34}$ . This shows that  $\ker \psi_3 = 0$ .

Similar to the result for  $H^1(E_1)$ , we get

$$\begin{aligned} H^2(E_1) &\cong \text{im} \psi_1 \oplus \text{im} \psi_2 \oplus \text{im} \psi_3 \\ &\cong P_{(I_1 \cup I_2) \cap I_3 \cap I_4} - (P_{(I_1 \cup I_2) \cap I_3} + P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4}) \\ &\oplus P_{I_1 \cap I_2 \cap I_4} \cap (P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) - (P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) + P_{I_1 \cap I_4} + P_{I_2 \cap I_4}) \\ &\oplus P_{I_1 \cap I_2 \cap I_3} \cap (P_{I_1 \cap I_2 \cap I_4} + P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) - (P_{I_1 \cap I_2} + P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) \end{aligned}$$

**Step 4:** For  $H^3(E_1)$ , it is easy to see that

$$H^3(E_1) \cong P_{I_1 \cap I_2 \cap I_3 \cap I_4} - (P_{I_1 \cap I_2 \cap I_3} + P_{I_1 \cap I_2 \cap I_4} + P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}).$$

As a result, the rational cohomology group  $H^*(BG(A))$  is isomorphic to  $\Sigma^3 H^3(E_1) \oplus \Sigma^2 H^2(E_1) \oplus \Sigma H^1(E_1) \oplus H^0(E_1)$ .

**Remark 4.1.** We will get different expressions of  $H^*(BG(A))$  if we change the order of indices  $I_1, I_2, \dots, I_d$ .

The computation for the general case  $X = X_{I_1} \cup X_{I_2} \cup \dots \cup X_{I_d}$  is very complicate. We have the following inductive algorithm to write down the final results.

**Step 1:** We begin with the cohomology group of  $X_1 = X_{I_1}$  which is isomorphic to  $P_{I_1}$  as a graded abelian group.

**Step 2:** We compute the cohomology group of  $X_2 = X_{I_1} \cup X_{I_2}$ . The intersection  $X_{I_1} \cap X_{I_2}$  has the cohomology groups  $P_{I_1 \cap I_2}$ . We need the following form of the M-V exact sequence which was used in Zhao and Gao[16]

**Lemma 4.1.** *Let  $X = X_1 \cup_{X_0} X_2$  be the push-out of the diagram  $X_1 \xleftarrow{j_1} X_0 \xrightarrow{j_2} X_2$ . The homomorphism  $j : H^*(X_1) \oplus H^*(X_2) \rightarrow H^*(X_0)$  is given by  $j(u, v) = j_1^*(u) - j_2^*(v)$ . If  $X_1, X_2$  are deformation retracts of some open subspaces of  $X$ , then there exists a short exact sequence*

$$0 \rightarrow \Sigma \text{coker} j \rightarrow H^*(X) \rightarrow \ker j \rightarrow 0.$$

Applying this lemma to  $X_{I_1} \cup X_{I_2}$ , there are two homomorphisms  $j_1 : H^*(X_{I_1}) \rightarrow H^*(X_{I_1 \cap I_2})$ , and  $j_2 : H^*(X_{I_2}) \rightarrow H^*(X_{I_1 \cap I_2})$ . Then  $\ker j \cong P_{I_1 \cup I_2}$  and  $\text{coker} j \cong P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2})$ . And we get

$$H^*(X_1 \cup X_2) \cong \Sigma \text{coker} j \oplus \ker j \cong \Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2})) \oplus P_{I_1 \cup I_2}.$$

We denote the two items in the expression by  $F_1$  and  $F_2$ . Then  $F_1 = \Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2}))$  has the form  $\Sigma(A - B)$  and  $F_2$  has the form  $A'$ .

**Step 3:** We compute the cohomology group of  $X_3 = X_{I_1} \cup X_{I_2} \cup X_{I_3}$ . Note that  $X_3 = X_2 \cup X_{I_3}$  and the intersection of  $X_2$  and  $X_{I_3}$  is  $\bar{X}_2 = (X_{I_1} \cup X_{I_2}) \cap X_{I_3} = X_{I_1 \cap I_3} \cup X_{I_2 \cap I_3}$ . By applying the result of Step 2 to  $\bar{I}_1 = I_1 \cap I_3$  and  $\bar{I}_2 = I_2 \cap I_3$ , we get

$$H^*(X_{I_1 \cap I_3} \cup X_{I_2 \cap I_3}) \cong \Sigma \text{coker} \bar{j} \oplus \ker \bar{j} \cong \Sigma(P_{(I_1 \cap I_3) \cap (I_2 \cap I_3)} - (P_{I_1 \cap I_3} + P_{I_2 \cap I_3})) \oplus P_{(I_1 \cap I_3) \cup (I_2 \cap I_3)}.$$

And we have two homomorphisms  $j_1 : H^*(X_{I_1} \cup X_{I_2}) \rightarrow H^*(X_{(I_1 \cap I_3) \cup (I_2 \cap I_3)})$  and  $j_2 : H^*(X_{I_3}) \rightarrow H^*(X_{(I_1 \cap I_3) \cup (I_2 \cap I_3)})$ . These homomorphisms can be depicted in the following diagram.

$$\begin{array}{ccccc} H^*(X_{I_1} \cup X_{I_2}) & \cong & \Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2})) & \oplus & P_{I_1 \cup I_2} \\ \downarrow j_1 & & \downarrow j_1^{\text{coker}} & & \downarrow j_1^{\text{ker}} \\ H^*(X_{(I_1 \cap I_3) \cup (I_2 \cap I_3)}) & \cong & \Sigma P_{(I_1 \cap I_3) \cap (I_2 \cap I_3)} - (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) & \oplus & P_{(I_1 \cap I_3) \cup (I_2 \cap I_3)} \\ j_2 \uparrow & & j_2^{\text{coker}} \uparrow & & j_2^{\text{ker}} \uparrow \\ H^*(X_{I_3}) & \cong & 0 & \oplus & P_{I_3} \end{array}$$

In the M-V exact sequence we combine the homomorphisms  $j_1, j_2$  to give the homomorphism  $j$  and  $H^*(X_{I_1} \cup X_{I_2} \cup X_{I_3}) \cong \Sigma \text{coker} j \oplus \ker j$ . If we assume that the homomorphisms  $j_1$  and  $j_2$  are decomposed into sum of its summand, then from the M-V exact sequence we get  $H^*(X_{I_1} \cup X_{I_2} \cup X_{I_3}) \cong \Sigma \text{coker} j_1^{\text{coker}} \oplus \Sigma \text{coker} j_2^{\text{ker}} \oplus \ker j_1^{\text{coker}} \oplus \ker j_2^{\text{ker}}$ . Detailed computation gives the following rule to obtain  $H^*(X_3)$  from  $H^*(X_2)$ .

i) We write the cohomology of  $\bar{X}_2 = X_{I_1 \cap I_3} \cup X_{I_2 \cap I_3}$  below the cohomology of  $X_2 = X_{I_1} \cup X_{I_2}$  and we get

$$\begin{array}{c} \Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2})) \quad \oplus \quad P_{I_1 \cup I_2} \\ \Sigma(P_{I_1 \cap I_2 \cap I_3} - (P_{I_1 \cap I_3} + P_{I_2 \cap I_3})) \oplus P_{(I_1 \cup I_2) \cap I_3} \end{array}$$

The result for  $\bar{X}_2$  is obtained from the result for  $X_2$  by replacing  $I_1$  and  $I_2$  to  $\bar{I}_1 = I_1 \cap I_3$  and  $\bar{I}_2 = I_2 \cap I_3$ . The two expressions contain four direct sum items, that is  $\Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2}))$ ,  $P_{I_1 \cup I_2}$ ,  $\Sigma(P_{I_1 \cap I_2 \cap I_3} - (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}))$  and  $P_{(I_1 \cup I_2) \cap I_3}$ .

ii) Each of these items will be transformed into an item in the expression of  $H^*(X_3)$ . The two corresponding items  $\Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2}))$  and  $\Sigma(P_{I_1 \cap I_2 \cap I_3} - (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}))$  have the forms

as  $\Sigma(A - B)$  and  $\Sigma(\bar{A} - \bar{B})$  with  $B \subset A, \bar{B} \subset \bar{A}$  and  $A \subset \bar{A}, B \subset \bar{B}$ . They will be transformed into two new items  $\Sigma(A \cap \bar{B} - B)$  and  $\Sigma^2(\bar{A} - (\bar{B} + A))$ . The two items  $P_{I_1 \cup I_2}$  and  $P_{(I_1 \cup I_2) \cap I_3}$  have the forms  $A'$  and  $\bar{A}'$  with  $A' \subset \bar{A}'$ , which are transformed into the two items  $\Sigma(\bar{A}' - (A' + P_{I_3}))$  and  $A' \cap P_{I_3}$ . We write the four items as  $F_1 \oplus F_2 \oplus F_3 \oplus F_4$ , and require  $F_4$  to be the unique item without grade lifting (no  $\Sigma$  appears).

And we continue this process until **Step**  $k$ .

Suppose we have finished the step  $k$  and obtained the cohomology group  $H^*(X_k) = H^*(X_{I_1} \cup X_{I_2} \cup \dots \cup X_{I_k})$  and it has the form of  $F_1 \oplus F_2 \oplus \dots \oplus F_{2^{k-1}-1} \oplus F_{2^{k-1}}$ . And we require that there is no  $\Sigma$  in  $F_{2^{k-1}}$ .

**Step**  $k + 1$ : now we write  $H^*(\bar{X}_k) = H^*(X_{I_1 \cap I_{k+1}} \cup X_{I_2 \cap I_{k+1}} \cup \dots \cup X_{I_k \cap I_{k+1}})$  below the cohomology groups  $H^*(X_k)$  and we get

$$\begin{aligned} &F_1 \oplus F_2 \oplus \dots \oplus F_{2^{k-1}-1} \oplus F_{2^{k-1}} \\ &\bar{F}_1 \oplus \bar{F}_2 \oplus \dots \oplus \bar{F}_{2^{k-1}-1} \oplus \bar{F}_{2^{k-1}} \end{aligned}$$

Through the procedure similar to that in step 3 we know that the two items  $F_i$  and  $\bar{F}_i, 1 \leq i \leq 2^{k-1} - 1$  have the forms of  $\Sigma^r(A - B)$  and  $\Sigma^r(\bar{A} - \bar{B})$  with  $B \subset A, \bar{B} \subset \bar{A}$  and  $A \subset \bar{A}, B \subset \bar{B}$ , which will be transformed into two new items  $\Sigma^r(A \cap \bar{B} - B)$  and  $\Sigma^{r+1}(\bar{A} - (\bar{B} + A))$  in  $H^*(X_{k+1})$ . The last two items  $F_{2^{k-1}}$  and  $\bar{F}_{2^{k-1}}$  have the forms  $A'$  and  $\bar{A}'$  with  $A' \subset \bar{A}'$ , which are transformed into the two items  $\Sigma(\bar{A}' - (A' + P_{I_{k+1}}))$  and  $A' \cap P_{I_{k+1}}$ .

The algorithm ends when  $k = d - 1$ .

The following is an example illustrating the algorithm for  $d = 4$ .

**Example 4.2. Step 1.**  $X_{I_1}$ , input  $P_{I_1}$

**Step 2.**  $X_{I_1} \cup X_{I_2}$

We write the cohomology of  $X_1$  and  $\bar{X}_1$ .

$$P_{I_1}$$

$$P_{I_1 \cap I_2}$$

And we get  $H^*(X_2) \cong \Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2})) \oplus P_{I_1 \cup I_2}$

**Step 3.**  $X_{I_1} \cup X_{I_2} \cup X_{I_3}$

$$(X_{I_1} \cup X_{I_2}) \cap X_{I_3} = (X_{I_1} \cap X_{I_3}) \cup (X_{I_2} \cap X_{I_3}) = X_{I_1 \cap I_3} \cup X_{I_2 \cap I_3}$$

We write the cohomology of  $X_2$  and  $\bar{X}_2$ .

$$\Sigma(P_{I_1 \cap I_2} - (P_{I_1} + P_{I_2})) \oplus P_{I_1 \cup I_2}$$

$$\Sigma(P_{I_1 \cap I_2 \cap I_3} - (P_{I_1 \cap I_3} + P_{I_2 \cap I_3})) \oplus P_{(I_1 \cup I_2) \cap I_3}$$

And we get  $H^*(X_3) \cong \Sigma^2(P_{I_1 \cap I_2 \cap I_3} - (P_{I_1 \cap I_2} + P_{I_1 \cap I_3} + P_{I_2 \cap I_3})) \oplus \Sigma(P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) - (P_{I_1} + P_{I_2})) \oplus \Sigma(P_{(I_1 \cup I_2) \cap I_3} - (P_{I_1 \cup I_2} + P_{I_3})) \oplus P_{I_1 \cup I_2 \cup I_3}$

**Step 4.**  $X_{I_1} \cup X_{I_2} \cup X_{I_3} \cup X_{I_4}$

$$(X_{I_1} \cup X_{I_2} \cup X_{I_3}) \cap X_{I_4} = X_{I_1 \cap I_4} \cup X_{I_2 \cap I_4} \cup X_{I_3 \cap I_4}$$

We write the cohomology of  $X_3$  and  $\bar{X}_3$ .

$$\begin{aligned} &\Sigma^2(P_{I_1 \cap I_2 \cap I_3} - (P_{I_1 \cap I_2} + P_{I_1 \cap I_3} + P_{I_2 \cap I_3})) \oplus \Sigma(P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) - (P_{I_1} + P_{I_2})) \oplus \\ &\Sigma(P_{(I_1 \cup I_2) \cap I_3} - (P_{I_1 \cup I_2} + P_{I_3})) \oplus P_{I_1 \cup I_2 \cup I_3} \end{aligned}$$

$$\Sigma^2(P_{I_1 \cap I_2 \cap I_3 \cap I_4} - (P_{I_1 \cap I_2 \cap I_4} + P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4})) \oplus \Sigma(P_{I_1 \cap I_2 \cap I_4} \cap (P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) -$$

$$(P_{I_1 \cap I_4} + P_{I_2 \cap I_4}) \oplus \Sigma(P_{(I_1 \cup I_2) \cap I_3 \cap I_4} - (P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4})) \oplus P_{(I_1 \cup I_2 \cup I_3) \cap I_4}$$

And we get

$$\begin{aligned} H^*(X_4) \cong & \Sigma^3(P_{I_1 \cap I_2 \cap I_3 \cap I_4} - (P_{I_1 \cap I_2 \cap I_3} + P_{I_1 \cap I_2 \cap I_4} + P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4})) \\ & \oplus \Sigma^2(P_{(I_1 \cup I_2) \cap I_3 \cap I_4} - (P_{(I_1 \cup I_2) \cap I_3} + P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4})) \\ & \oplus \Sigma^2(P_{I_1 \cap I_2 \cap I_4} \cap (P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) - (P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) + P_{I_1 \cap I_4} + P_{I_2 \cap I_4})) \\ & \oplus \Sigma^2(P_{I_1 \cap I_2 \cap I_3} \cap (P_{I_1 \cap I_2 \cap I_4} + P_{I_1 \cap I_3 \cap I_4} + P_{I_2 \cap I_3 \cap I_4}) - (P_{I_1 \cap I_2} + P_{I_1 \cap I_3} + P_{I_2 \cap I_3})) \\ & \oplus \Sigma(P_{(I_1 \cup I_2 \cup I_3) \cap I_4} - (P_{I_1 \cup I_2 \cup I_3} + P_{I_4})) \\ & \oplus \Sigma(P_{(I_1 \cup I_2) \cap I_3} \cap (P_{(I_1 \cup I_2) \cap I_4} + P_{I_3 \cap I_4}) - (P_{I_1 \cup I_2} + P_{I_3})) \\ & \oplus \Sigma(P_{I_1 \cap I_2} \cap (P_{I_1 \cap I_3} + P_{I_2 \cap I_3}) \cap (P_{I_1 \cap I_4} + P_{I_2 \cap I_4}) - (P_{I_1} + P_{I_2})) \\ & \oplus P_{I_1 \cup I_2 \cup I_3 \cup I_4} \end{aligned}$$

By classifying the maximal objects in  $C(A)$  and using the above algorithm, we obtain the results for rational cohomology groups of the classifying spaces of Kac-Moody groups of infinite type with  $n = 3$  and 4.

**Example 4.3.** *The rational cohomology groups of the classifying space  $BG(A)$  for  $n = 3$  are listed case by case as below.*

1.  $\{1\}, \{2\}, \{3\}$   
 $\Sigma(P - (P_1 + P_2)) \oplus \Sigma(P - (P_{12} + P_3)) \oplus P_{123}$
2.  $\{1, 2\}, \{3\}$   
 $\Sigma(P - (P_{12} + P_3)) \oplus P_{123}$
3.  $\{1, 2\}, \{1, 3\}$   
 $\Sigma(P_1 - (P_{12} + P_{13})) \oplus P_{123}$
4.  $\{1, 2\}, \{1, 3\}, \{2, 3\}$   
 $\Sigma^2(P - (P_1 + P_2 + P_3)) \oplus \Sigma(P_1 \cap (P_2 + P_3) - (P_{12} + P_{13})) \oplus P_{123}$

**Example 4.4.** *The rational cohomology groups of the classifying space  $BG(A)$  for  $n = 4$  are listed case by case as below.*

1.  $\{1\}, \{2\}, \{3\}, \{4\},$   
 $\Sigma(P - (P_1 + P_2)) \oplus \Sigma(P - (P_{12} + P_3)) \oplus \Sigma(P - (P_{123} + P_4)) \oplus P_{1234}$
2.  $\{1, 2\}, \{3\}, \{4\},$   
 $\Sigma(P - (P_{12} + P_3)) \oplus \Sigma(P - (P_{123} + P_4)) \oplus P_{1234}$
3.  $\{1, 2\}, \{3, 4\},$   
 $\Sigma(P - (P_{12} + P_{34})) \oplus P_{1234}$
4.  $\{1, 2\}, \{1, 3\}, \{4\}$   
 $\Sigma(P_1 - (P_{12} + P_{13})) \oplus \Sigma(P - (P_{123} + P_4)) \oplus P_{1234}$
5.  $\{1, 2\}, \{1, 3\}, \{1, 4\}$   
 $\Sigma(P_1 - (P_{12} + P_{13})) \oplus \Sigma(P_1 - (P_{123} + P_{14})) \oplus P_{1234}$
6.  $\{1, 2\}, \{1, 3\}, \{3, 4\},$   
 $\Sigma(P_1 - (P_{12} + P_{13})) \oplus \Sigma(P_3 - (P_{123} + P_{34})) \oplus P_{1234}$
7.  $\{1, 2\}, \{1, 3\}, \{2, 3\}, \{4\}$

- $\Sigma^2(P - (P_1 + P_2 + P_3)) \oplus \Sigma(P_1 \cap (P_2 + P_3) - (P_{12} + P_{13})) \oplus \Sigma(P - (P_{123} + P_4)) \oplus P_{1234}$   
8.  $\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}$   
 $\Sigma^2(P - (P_1 + P_2 + P_3)) \oplus \Sigma(P_1 \cap (P_2 + P_3) - (P_{12} + P_{13})) \oplus \Sigma(P_1 - (P_{123} + P_{14})) \oplus P_{1234}$   
9.  $\{1, 2\}, \{1, 3\}, \{2, 4\}, \{3, 4\}$   
 $\Sigma^2(P - (P_2 + P_3 + P_4)) \oplus \Sigma(P_1 - (P_{12} + P_{13})) \oplus \Sigma(P_2 \cap (P_3 + P_4) - (P_{123} + P_{24})) \oplus P_{1234}$   
10.  $\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}$   
 $\Sigma^2(P - (P_1 + P_2 + P_3)) \oplus \Sigma^2(P - (P_1 + P_2 + P_4)) \oplus \Sigma(P_1 \cap (P_2 + P_4) - (P_{123} + P_{14})) \oplus \Sigma(P_1 \cap (P_2 + P_3) - (P_{12} + P_{13})) \oplus P_{1234}$   
11.  $\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}$   
 $\Sigma^2(P - (P_1 + P_2 + P_3)) \oplus \Sigma^2(P - (P_1 + P_2 + P_4)) \oplus \Sigma^2(P - (P_1 \cap (P_2 + P_4) + P_3 + P_4)) \oplus \Sigma(P_1 \cap (P_2 + P_4) \cap (P_3 + P_4) - (P_{123} + P_{14})) \oplus \Sigma(P_1 \cap (P_2 + P_3) - (P_{12} + P_{13})) \oplus P_{1234}$   
12.  $\{1, 2, 3\}, \{4\}$   
 $\Sigma(P - (P_{123} + P_4)) \oplus P_{1234}$   
13.  $\{1, 2, 3\}, \{1, 4\}$   
 $\Sigma(P_1 - (P_{123} + P_{14})) \oplus P_{1234}$   
14.  $\{1, 2, 3\}, \{1, 4\}, \{2, 4\},$   
 $\Sigma^2(P - (P_1 + P_2 + P_4)) \oplus \Sigma(P_1 \cap (P_2 + P_4) - (P_{123} + P_{14})) \oplus P_{1234}$   
15.  $\{1, 2, 3\}, \{1, 4\}, \{2, 4\}, \{3, 4\}$   
 $\Sigma^2(P - (P_1 + P_2 + P_4)) \oplus \Sigma^2(P - (P_1 \cap (P_2 + P_4) + P_3 + P_4)) \oplus \Sigma((P_1 \cap (P_2 + P_4) \cap (P_3 + P_4) - (P_{123} + P_{14})) \oplus P_{1234}$   
16.  $\{1, 2, 3\}, \{1, 2, 4\}$   
 $\Sigma(P_{12} - (P_{123} + P_{124})) \oplus P_{1234}$   
17.  $\{1, 2, 3\}, \{1, 2, 4\}, \{3, 4\}$   
 $\Sigma^2(P - (P_{12} + P_3 + P_4)) \oplus \Sigma(P_{12} \cap (P_3 + P_4) - (P_{123} + P_{124})) \oplus P_{1234}$   
18.  $\{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 4\}$   
 $\Sigma^2(P_1 - (P_{12} + P_{13} + P_{14})) \oplus \Sigma(P_{12} \cap (P_{13} + P_{14}) - (P_{123} + P_{124})) \oplus P_{1234}$   
19.  $\{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 4\}, \{2, 3, 4\}$   
 $\Sigma^3(P - (P_1 + P_2 + P_3 + P_4)) \oplus \Sigma^2(P_1 \cap (P_2 + P_3 + P_4) - (P_{12} + P_{13} + P_{14})) \oplus \Sigma^2(P_2 \cap (P_3 + P_4) - (P_{12} \cap (P_{13} + P_{14}) + P_{23} + P_{24})) \oplus \Sigma(P_{12} \cap (P_{13} + P_{14}) \cap (P_{23} + P_{24}) - (P_{123} + P_{124})) \oplus P_{1234}$

By the above algorithm there are always  $2^{d-1}$  items in the cohomology of  $H^*(BG(A))$ . But some items are zero for trivial reason. We delete them from the lists. We have done the similar computations for  $n = 5$  with the help of Zhang Yukun, Luo Yajing, Gong Tao, Wang Ran, Li Yingxin, Qu Qingrui, Liang Peng and Gu Peiqi. The result is too lengthy to be listed here.

## 5 The polynomial invariants of Weyl groups and their subgroups

To compute the rational cohomology of the classifying space  $BG(A)$ , it is vital to know the ring of polynomial invariants  $P_I$  for  $I \subset S$  (see the expressions in Examples 4.3 and 4.4). In some cases the computation of  $P_I$  for  $I \subset S$  can be reduced to the computation of rational polynomial invariants of Weyl group  $W(A_I)$  for the Cartan matrix  $A_I$ .

Each decomposable  $n \times n$  Cartan matrix  $A$  can be written, up to order of the elements in  $S$ , as a sum  $A = \bigoplus_{i=1}^k A_{S_i}$  of sub-matrices with  $S$  the disjoint union of  $S_i$ . And the corresponding Kac-Moody groups can be decomposed into  $G(A) = \prod_{i=1}^k G(A_{S_i})$ . For each  $I \subset S$  the parabolic subgroup  $G_I(A)$  has a decomposition  $G_I(A) = \prod_{i=1}^k G_{I \cap S_i}(A_i)$ . And the Weyl group of  $G_I(A)$  is isomorphic to  $\prod_{i=1}^k W_{I \cap S_i}(A_i)$ . It is obvious that the ring of invariants  $P_I$  is isomorphic to  $\bigotimes_{i=1}^k P_{I \cap S_i}$ . So the computation of  $P_I$  can be reduced to the indecomposable case.

Let  $A = (a_{ij})_{n \times n}$  be a Cartan matrix. The Weyl reflections  $\sigma_1, \sigma_2, \dots, \sigma_n$  satisfy

$$\sigma_i(w_j) = \begin{cases} w_j - \alpha_j, & \text{for } i = j; \\ w_j, & \text{for } i \neq j. \end{cases}$$

Thus we have

$$(\sigma_1(w_1), \sigma_2(w_2), \dots, \sigma_n(w_n)) = (w_1, w_2, \dots, w_n)(I_n - A).$$

We write  $(w_1, w_2, \dots, w_n)$  as  $w$ . For  $S = I \sqcup J$ , we write  $w$  as  $(w_I, w_J)$ , where  $w_I = (w_1, w_2, \dots, w_{|I|})$  and  $w_J = (w_{|I|+1}, w_{|I|+2}, \dots, w_n)$ . We have

$$(\sigma_1(w_1), \sigma_2(w_2), \dots, \sigma_{|I|}(w_{|I|})) = (w_1, w_2, \dots, w_n) \begin{pmatrix} I_{|I|} - A_I \\ -A_{JI} \end{pmatrix}$$

and  $\sigma_i(w_j) = w_j$  for all  $i \in I, j \in J$ , where  $A_{JI}$  is the submatrix  $A_{JI} = (a_{ji})_{j \in J, i \in I}$  of  $A$ .

By choosing a  $|J| \times |I|$  matrix  $C$  with rational elements, we get another basis for the rational linear space of weights,

$$w' = (w'_I, w'_J) = (w_I, w_J) \begin{pmatrix} I_{|I|} & 0 \\ C & I_{|J|} \end{pmatrix}.$$

By a direct computation we get

$$(\sigma_1(w'_1), \sigma_2(w'_2), \dots, \sigma_{|I|}(w'_{|I|})) = (w_1, w_2, \dots, w_n) \begin{pmatrix} I_{|I|} - A_I \\ C - A_{JI} \end{pmatrix} = (w'_1, w'_2, \dots, w'_n) \begin{pmatrix} I_{|I|} - A_I \\ CA_I - A_{JI} \end{pmatrix}$$

If the rank of matrix  $A_I$  is the same as that of  $\begin{pmatrix} A_I \\ A_{JI} \end{pmatrix}$ , then the rational matrix  $C$  can be chosen such that  $CA_I = A_{JI}$ . As a consequence we get

$$(\sigma_1(w'_1), \sigma_2(w'_2), \dots, \sigma_{|I|}(w'_{|I|})) = (w'_1, w'_2, \dots, w'_{|I|})(I_{|I|} - A_I).$$

This implies that the linear action of  $W_I(A)$  on the  $\mathbb{Q}$ -linear space spanned by  $w'_1, w'_2, \dots, w'_{|I|}$  is isomorphic to the action of  $W(A_I)$  on the  $\mathbb{Q}$ -linear space of weights for  $G(A_I)$ . Combining with the fact that  $W_I(A)$  acts trivially on  $w_j, j \in J$ , we prove the following proposition.

**Proposition 5.1.** *Let  $A$  be an  $n \times n$  Cartan matrix and the matrix equation  $XA_I = A_{JI}$  has a rational solution  $C$ , then the ring of invariants  $P_I$  of  $W_I(A)$  is isomorphic to the tensor product of the ring of invariants of the Weyl group  $W(A_I)$  acting on  $\mathbb{Q}$ -linear space of weights for  $G(A_I)$  and the polynomial ring  $\mathbb{Q}[w_{|I|+1}, w_{|I|+2}, \dots, w_n]$ .*

Particularly the proposition is true if  $\det A_I \neq 0$ .

Even one knows all the rings of invariants  $P_I$ , it is still difficult to compute the rational cohomology groups of  $BG(A)$  explicitly. There are expressions like  $P - (P_1 + P_2 + P_3)$  and  $P_1 \cap (P_2 + P_4) \cap (P_3 + P_4) - (P_{123} + P_{14})$  for which we even can not compute their Poincaré series.

In [5] Foley claimed that for a  $3 \times 3$  Cartan matrix  $A$  of infinite type,  $P - (P_1 + P_2 + P_3) = 0$ . This result hints the following conjecture.

**Conjecture 5.1.** *For an  $n \times n$  Cartan matrix  $A$  of infinite type,  $P = P_1 + P_2 + \cdots + P_n$ .*

## 6 Some applications and conjectures

The computation for the cohomology groups of  $BG(A)$ (or  $BK(A)$ ) has several applications.

**Theorem 6.1.** *For a Kac-Moody group  $K(A)$  of infinite type, the image of the Borel homomorphism  $Bi^* : H^*(BK(A); \mathbb{Q}) \rightarrow H^*(BT, \mathbb{Q})$  is the ring of invariants  $P^{W(A)}$ .*

**Proof:** We have  $BK(A) \sim X_{I_1} \cup X_{I_2} \cup \cdots \cup X_{I_d}$ . We can also write  $BT$  as  $BT \cup BT \cup \cdots \cup BT$  and there is a homomorphism  $f$  between the M-V spectral sequences of  $H^*(BK(A))$  and  $H^*(BT)$ . Borel's theorem asserts that the homomorphism  $H^*(X_{I_i}) \rightarrow H^*(BT)$  has image the ring of invariants  $P^{W_{I_i}(A)}$ . Since  $S = \bigcup_{i=1}^d I_i$ , the image of  $Bi^*$  is isomorphic to  $\bigcap_{i=1}^d P^{W_{I_i}(A)} = P^{W(A)}$ .

The following result is proved in Ruan and Zhao[15].

**Theorem 6.2.** *For a rank 3 Kac-Moody group  $K(A)$  of infinite type, the integral cohomology  $H^*(BK(A); \mathbb{Z})$  has  $p$ -torsion elements for any prime number  $p$ .*

Based on theorem 6.2 we can give the following conjecture.

**Conjecture 6.1.** *For a Kac-Moody group  $K(A)$  of infinite type,  $H^*(BG(A); \mathbb{Z})$  has  $p$ -torsion elements for any prime  $p$ .*

In [16] we show that for a generic Kac-Moody group, the rational cohomology ring  $H^*(BG(A); \mathbb{Q})$  is infinitely generated. So it is natural to give the following conjecture.

**Conjecture 6.2.** *For a Cartan matrix  $A = (a_{ij})_{n \times n}$  of indefinite type with  $n \geq 3$ , the rational cohomology ring  $H^*(BG(A); \mathbb{Q})$  is infinitely generated.*

**Remark 6.1.** *For Conjecture 6.2 it is known that for Cartan matrix  $A = (a_{ij})_{n \times n}$  of finite type, affine type or  $n \leq 2$ , the rational cohomology ring  $H^*(BG(A); \mathbb{Q})$  is finitely generated.*

The computation in this paper can be used to attack the above conjectures.

## References

- [1] A. Borel, Sur la cohomologie des espaces fibres principaux et des espaces homogenes de groupes de Lie compacts, Ann.math. 57(1953), 115-207.

- [2] A. Bousfield, D. Kan, Homotopy limits, completions and localizations. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972.
- [3] R. Bott, and L. W. Tu, Differential forms in algebraic topology, Graduate Texts in Mathematics, 82. New York-Berlin, Springer-Verlag, 1982.
- [4] D. Dugger, A primer on homotopy colimits. <http://math.uoregon.edu/~ddugger/hocolim.pdf>, 2008, preprint.
- [5] J. D. Foley, Recognizing nullhomotopic maps into the classifying space of a Kac-Moody group, 2015, arXiv:1502.06520.
- [6] V. Kac, Simple irreducible graded Lie algebras of finite growth. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 32, 1968, 1323-1367.
- [7] V. Kac, Infinite Dimensional Lie Algebras, Cambridge University Press, 1982.
- [8] V. Kac; D. Peterson, Regular functions on certain infinite-dimensional groups. *Arithmetic and geometry*, Vol. II, 141-166, *Progr. Math.*, 36, Birkhäuser Boston, Boston, MA, 1983.
- [9] V. Kac; D. Peterson, Defining relations of certain infinite-dimensional groups. *The mathematical heritage of élie Cartan (Lyon, 1984)*. *Astérisque* 1985, Numéro Hors Série, 165-208.
- [10] V. Kac, Constructing groups associated to infinite-dimensional Lie algebras. *Infinite dimensional groups with applications (Berkeley, Calif., 1984)*, 167-216, *Math. Sci. Res. Inst. Publ.*, 4, Springer, New York, 1985.
- [11] N. Kitchloo, Topology of Kac-Moody groups, PhD thesis, MIT, 1998.
- [12] C. Broto and N. Kitchloo, Classifying spaces of Kac-Moody groups, *Math. Z.* 240 (2002), no. 3, 621-649.
- [13] N. Kitchloo, On some applications of unstable Adams operations to the topology of Kac-Moody groups, *Proc. Amer. Math. Soc.*, 2017, 145(2), 915-924.
- [14] R. V. Moody, A new class of Lie algebras, *J. Algebra*, Vol 10, 1968, 211-230.
- [15] Ruan Yangyang, Zhao Xu-an, Cohomology of classifying spaces of rank 3 Kac-Moody groups, preprint.
- [16] Zhao Xu-an, Gao Hongzhu, The rational cohomology Hopf algebra of a generic Kac-Moody group, *Pacific J. Math.*, Vol 305(2), 2020, 757-766.