Estimation based on nearest neighbor matching: from density ratio to average treatment effect

Zhexiao Lin^{*}, Peng Ding[†], and Fang Han[‡]

December 28, 2021

Abstract

Nearest neighbor (NN) matching as a tool to align data sampled from different groups is both conceptually natural and practically well-used. In a landmark paper, Abadie and Imbens (2006) provided the first large-sample analysis of NN matching under, however, a crucial assumption that the number of NNs, M, is fixed. This manuscript reveals something new out of their study and shows that, once allowing M to diverge with the sample size, an intrinsic statistic in their analysis actually constitutes a consistent estimator of the density ratio. Furthermore, through selecting a suitable M, this statistic can attain the minimax lower bound of estimation over a Lipschitz density function class. Consequently, with a diverging M, the NN matching provably yields a doubly robust estimator of the average treatment effect and is semiparametrically efficient if the density functions are sufficiently smooth and the outcome model is appropriately specified. It can thus be viewed as a precursor of double machine learning estimators.

Keywords: graph-based statistics, stochastic geometry, double robustness, double machine learning, propensity score.

1 Introduction

With observations from different groups, matching methods (Greenwood, 1945; Chapin, 1947) aim to balance them through minimizing group differences in observed covariates. Such methods have proven their usefulness for causal inference in various disciplines, including economics (Imbens, 2004), epidemiology (Brookhart et al., 2006), political science (Ho et al., 2007; Sekhon, 2008), sociology (Morgan and Harding, 2006), and statistics (Cochran and Rubin, 1973; Rubin, 2006; Rosenbaum, 2010).

Among all the matching methods, nearest neighbor (NN) matching (Rubin, 1973) is likely the most well-used and easiest to implement approach. In addition, it is computationally attractive as the time complexity of locating NNs is low. In the simplest treatment-control study, NN matching assigns each treatment (control) individual to M control (treatment) individuals with the smallest distance to it. In this regard, two natural questions arise. First, how do we select the number of

^{*}Department of Statistics, University of Washington, Seattle, WA 98195, USA; e-mail: zxlin@uw.edu

[†]Department of Statistics, University of California, Berkeley, CA 94720, USA; email: pengdingpku@berkeley.edu

[‡]Department of Statistics, University of Washington, Seattle, WA 98195, USA; e-mail: fanghan@uw.edu

matches, *M*? This is referred to in the literature as ratio matching, and is both important and delicate, well-known to be related to the bias-variance tradeoff in nonparametic statistics (Smith, 1997; Rubin and Thomas, 2000). Second, how do we perform large-sample statistical inference for NN matching methods? Such an analysis is usually nonstandard and thus believed to be mathematically challenging. Indeed, it was long-lacking in the literature until Abadie and Imbens (2006).

To answer the above two questions, in a series of ingenious papers, Abadie and Imbens (2006, 2008, 2011, 2012) established large-sample properties of M-NN matching for estimating the average treatment effect (ATE). These results are, however, only valid under a crucial assumption that, in ratio matching, M is fixed. The according message is then mixed. As a matter of fact, Abadie and Imbens (2006) argued — which we quote here — that the ATE estimator based on M-NN matching with a fixed M is both asymptotically biased and statistically inefficient, namely, it "does not achieve the semiparametric efficiency bound as calculated by Hahn (1998)". While bias correction is now feasible to alleviate the first issue (Abadie and Imbens, 2011), the lack of efficiency seems fundamental.

This manuscript revisits the study of Abadie and Imbens (2006) from a new perspective, bridging M-NN matching to density ratio estimation (Nguyen et al., 2010; Sugiyama et al., 2012) as well as double robustness (Scharfstein et al., 1999; Bang and Robins, 2005). To this end, our analysis stresses, in ratio matching, the importance of forcing M to diverge with the sample size n in order to achieve statistical efficiency. Our claim is thus aligned with similar ones in various other random graph-based inference problems (Wald and Wolfowitz, 1940; Friedman and Rafsky, 1979; Henze, 1988; Liu and Singh, 1993; Henze and Penrose, 1999; Berrett et al., 2019; Bhattacharya, 2019), which also include a series of results made by some of the authors in this paper (Shi et al., 2021a,b; Lin and Han, 2021).

The contributions of this manuscript are mainly two-fold. First, we show that an intrinsic statistic that plays a central role in the analysis of Abadie and Imbens (2006), $K_M(x)$ (Abadie and Imbens (2006, Page 240); to be defined in (2.2) of Section 2), actually gives rise to a consistent density ratio estimator in the two-sample setting. Even more interestingly, from the angle of density ratio estimation, this NN matching-based estimator is to our knowledge the first one that simultaneously satisfies the following three properties.

- (P1) Conceptually one-step: it directly estimates the density ratio with no need to estimate individual densities, and is thus in line with Vapnik's rule that "[w]hen solving a problem of interest, one should not solve a more general problem as an intermediate step" (Vapnik, 2006, on the top of Page 477).
- (P2) Computationally of low complexity: it is of a sub-quadratic (and nearly linear when M is small) time complexity via a careful algorithmic formulation based on k-d trees (cf. Algorithms 1-2 and Theorem 3.1 ahead), and thus in many scientific applications is computationally more attractive than its optimization-based alternatives (Lima et al., 2008; Kremer et al., 2015; Borgeaud et al., 2021).
- (P3) Statistically *rate-optimal*: it is information-theoretically efficient in terms of achieving an upper bound of estimation accuracy that matches the corresponding minimax lower bound

over a class of Lipschitz density functions (cf. the set of results in Section 4 ahead).

This estimator itself is accordingly an appealing alternative to existing density ratio estimators. Moreover, it is potentially useful in many data analysis problems (e.g., f-divergence estimation, classification, importance sampling, and etc.) where density ratio estimation plays a pivotal role.

Getting back to the original ATE estimation problem, our second contribution is to bridge the above insights to the bias-corrected matching-based estimator proposed in Abadie and Imbens (2011) as well as the double robustness and double machine learning framework introduced in Scharfstein et al. (1999), Bang and Robins (2005), and Chernozhukov et al. (2018). In fact, their bias-corrected estimator can be formulated as

$$\widehat{\tau}_M^{\rm bc} = \widehat{\tau}^{\rm reg} + \frac{1}{n} \Big[\sum_{i=1,D_i=1}^n \Big(1 + \frac{K_M^1(i)}{M} \Big) \widehat{R}_i - \sum_{i=1,D_i=0}^n \Big(1 + \frac{K_M^0(i)}{M} \Big) \widehat{R}_i \Big]$$

(notation to be introduced in Section 5), where $1 + K_M^1(\cdot)/M$ and $1 + K_M^0(\cdot)/M$ approach the inverse of the propensity score e(x) and 1 - e(x), respectively. One could then leverage the general double robustness and double machine learning theory to validate the following two claims of (a double machine learning version of) $\hat{\tau}_M^{\text{bc}}$.

- (1) Consistency: as long as either the density (propensity score) functions satisfy certain conditions or the outcome (regression) model is correctly specified, $M \log n/n \to 0$, and $M \to \infty$ as $n \to \infty$, $\hat{\tau}_M^{\rm bc}$ converges in probability to the population ATE, denoted as τ .
- (2) Semiparametric efficiency: if the density functions are sufficiently smooth, the outcome model is appropriately specified, and M scales with n at an proper rate, then (a sample-splitting and cross-fitting version of) $\hat{\tau}_{M}^{\rm bc}$ is an asymptotically normal estimator of τ with the asymptotic variance attaining the semiparametric efficiency lower bound (Hahn, 1998). Furthermore, a simple consistent estimator of the asymptotic variance is available.

Our results thus complement those made in Abadie and Imbens (2006, 2011), rendering necessary confidence for practitioners to implement NN matching for inferring the ATE. In addition, although Abadie and Imbens (2006, Theorem 5) hints at the necessity of allowing M to diverge for gaining efficiency, we provide rigorous theory for their conjecture.

Technically speaking, our analysis hinges on those M's that grow with n. Existing results in NN matching literature, including Abadie and Imbens (2006), Abadie and Imbens (2008), Abadie and Imbens (2011), and Abadie and Imbens (2012), are then limited as they are all focused on a fixed M. Instead, we take a different route to establish nonasymptotic moment bounds of $K_M(x)$, where there is more room for M to move around; of note, similar ideas were also pursued in Lin and Han (2021) by the authors to analyze rank-based statistics. Detailed explanation of our theoretical analysis, however, has to be left to latter sections.

Paper organization. The rest of this manuscript is organized as follows. In Sections 2-4 we introduce the method, computation, and theory of density ratio estimation via NN matching. In detail, Section 2 introduces the statistical setup and the matching-based estimator of density ratio. Section 3 introduces the algorithms to implement the matching-based estimator constructed on

the k-d tree structure. Section 4 delivers the main theory, quantifying both pointwise and global approximation accuracy. Built on the previous three sections, Section 5 formally elaborates on the double robustness and semiparametric efficiency of the bias-corrected NN matching-based estimator of the ATE. More applications of the proposed matching-based estimator will be covered in Section 6, with proofs of the main results relegated to Section 7 and the rest put in the appendix. Additional results about NN matching that cannot be incorporated in the double robustness and double machine learning framework are exhibited in the supplement.

Notation. For any integers $n, d \ge 1$, let $[n] := \{1, 2, ..., n\}$, n! be the factorial of n, and \mathbb{R}^d be the d-dimensional real space. A set consisting of distinct elements $x_1, ..., x_n$ is written as either $\{x_1, ..., x_n\}$ or $\{x_i\}_{i=1}^n$, and its cardinality is written by $|\{x_i\}_{i=1}^n|$. The corresponding sequence is denoted $[x_1, ..., x_n]$ or $[x_i]_{i=1}^n$. The notation $\mathbb{1}(\cdot)$ is saved for the indicator function. For any $a, b \in \mathbb{R}$, write $a \lor b = \max\{a, b\}$ and $a \land b = \min\{a, b\}$. For any two real sequences $\{a_n\}$ and $\{b_n\}$, write $a_n \lesssim b_n$ (or equivalently, $b_n \gtrsim a_n$ or $a_n = O(b_n)$) if there exists a universal constant C > 0 such that $a_n/b_n \leq C$ for all sufficiently large n, and write $a_n \prec b_n$ (or equivalently, $b_n \succ a_n$ and $a_n = o(b_n)$) if $a_n/b_n \to 0$ as n goes to infinity. We write $a_n \asymp b_n$ if both $a_n \lesssim b_n$ and $b_n \lesssim a_n$ holds. We use $\stackrel{\mathsf{d}}{\longrightarrow}$ and $\stackrel{\mathsf{P}}{\longrightarrow}$ to denote convergence in distribution and in probability, respectively. For any sequence of random variables $\{X_n\}$, write $X_n = o_P(1)$ if $X_n \stackrel{\mathsf{P}}{\longrightarrow} 0$. For any random variable Z, P_Z represents its law. Denote the closed ball in \mathbb{R}^d centered at x with radius δ by $B_{x,\delta}$. In the sequel, let c, C, C', C'', C''', C''', ... be generic positive constants whose actual values may change at different locations.

2 Density ratio estimation I: method

From this section to Section 4, we consider X, Z to be two general random vectors in \mathbb{R}^d that are defined on the same probability space, with d to be a fixed positive integer.

Let ν_0 and ν_1 represent the probability measures of X and Z, respectively. Assume ν_0 and ν_1 are absolutely continuous with respect to the Lebesgue measure λ on \mathbb{R}^d equipped with the Euclidean norm $\|\cdot\|$; denote the corresponding densities (Radon-Nikodym derivatives) by f_0 and f_1 . Assume further that ν_1 is absolutely continuous with respect to ν_0 and write the corresponding density ratio, f_1/f_0 , as r; we set 0/0 = 0 by default.

Assume X_1, \ldots, X_{N_0} are N_0 independent copies of X, Z_1, \ldots, Z_{N_1} are N_1 independent copies of Z, and $[X_i]_{i=1}^{N_0}$ and $[Z_j]_{j=1}^{N_1}$ are mutually independent. The problem of estimating the density ratio r based on $\{X_1, \ldots, X_{N_0}, Z_1, \ldots, Z_{N_1}\}$ is fundamental in economics (Cunningham, 2021), information theory (Cover and Thomas, 2006), machine learning (Sugiyama et al., 2012), statistics (Imbens and Rubin, 2015), and other fields.

In density ratio estimation, NN-based estimators are advocated before due to its computational efficiency; cf. Lima et al. (2008), Póczos and Schneider (2011), Kremer et al. (2015), Noshad et al. (2017), Berrett et al. (2019), Zhao and Lai (2020b), among many others. In this manuscript, based on Abadie and Imbens (2006, 2008, 2011, 2012)'s NN matching framework, we unveil a new density ratio estimator based on NN matching. To this end, some necessary notation is introduced first.

Definition 2.1 (NN matching). For any $x, z \in \mathbb{R}^d$ and $M \in [N_0]$,

(i) let $\mathcal{X}_{(M)}(\cdot) : \mathbb{R}^d \to \{X_i\}_{i=1}^{N_0}$ be the mapping that returns the value of the input z's *M*-th NN in $\{X_i\}_{i=1}^{N_0}$, i.e., the value of $x \in \{X_i\}_{i=1}^{N_0}$ such that

$$\sum_{i=1}^{N_0} \mathbb{1}\left(\|X_i - z\| \le \|x - z\| \right) = M;$$
(2.1)

(ii) let $K_M(\cdot) : \mathbb{R}^d \to [\![N_1]\!]$ be the mapping that returns the number of matched times of x, i.e.,

$$K_M(x) = K_M\left(x; \{X_i\}_{i=1}^{N_0}, \{Z_j\}_{j=1}^{N_1}\right) := \sum_{j=1}^{N_1} \mathbb{1}\left(\|x - Z_j\| \le \|\mathcal{X}_{(M)}(Z_j) - Z_j\|\right);$$
(2.2)

(iii) let $A_M(\cdot) : \mathbb{R}^d \to \mathcal{B}(\mathbb{R}^d)$ be the corresponding mapping from \mathbb{R}^d to the class of all Borel sets in \mathbb{R}^d so that

$$A_M(x) = A_M\left(x; \{X_i\}_{i=1}^{N_0}\right) := \left\{z \in \mathbb{R}^d : \|x - z\| \le \|\mathcal{X}_{(M)}(z) - z\|\right\},\tag{2.3}$$

returns the catchment area of x in the setting of (ii);

(iv) for $i \in [N_0]$, let $K_M(i)$ and $A_M(i)$ be shorthands of $K_M(X_i)$ and $A_M(X_i)$, respectively.

Remark 2.1. Of note, since ν_0 is absolutely continuous with respect to the Lebesgue measure, a solution to (2.1) exists and is unique. Abadie and Imbens (2006, Pages 240 and 260) introduced the terms $K_M(\cdot)$ and $A_M(\cdot)$ to analyze the asymptotic behavior of their NN matching-based ATE estimator. We also adopt the terminology "catchment area" in Definition 2.1(iii) to align with them.

The following proposition formally links $K_M(\cdot)$ to $A_M(\cdot)$, was shown and used in the proof of Abadie and Imbens (2006, Lemma 3), and is stated here for aiding understanding.

Proposition 2.1 (Abadie and Imbens (2006)). For any $x \in \mathbb{R}^d$, we have

$$K_M(x) = \sum_{j=1}^{N_1} \mathbb{1}\Big(Z_j \in A_M(x)\Big).$$

Remark 2.2 (Relation between $A_M(i)$'s and Voronoi tessellation when M = 1). It is easy to verify that, due to the absolute continuity of ν_0 , $[A_1(i)]_{i=1}^{N_0}$ are almost surely disjoint except for a Lebesgue measure zero area, and partition \mathbb{R}^d into N_0 polygons. Furthermore, one can verify that $\{A_1(i)\}_{i=1}^{N_0}$ are exactly the Voronoi tessellation defined in Voronoi (1908), which plays a vital role in stochastic and computational geometry. In this case, each element $A_1(i)$ is the Voronoi cell from the definition of (2.3).

With these notation and concepts, we are now ready to introduce the following density ratio estimator based on NN matching.

Definition 2.2 (NN matching-based density ratio estimator). For any $M \in [N_0]$ and $x \in \mathbb{R}^d$, we define the following estimator of r(x),

$$\widehat{r}_M(x) = \widehat{r}_M\left(x; \{X_i\}_{i=1}^{N_0}, \{Z_j\}_{j=1}^{N_1}\right) := \frac{N_0}{N_1} \frac{K_M(x)}{M}.$$
(2.4)

The estimator $\hat{r}_M(\cdot)$ is by construction a one-step estimator, and thus automatically satisfies Property (P1) in Introduction. In the following two sections, we will show that $\hat{r}_M(\cdot)$ also satisfies Properties (P2) and (P3).

3 Density ratio estimation II: computation

This section discusses implementation of and establishes Property (P2) for the proposed estimator $\hat{r}_M(\cdot)$. To this end, we separately discuss two cases:

Case I: estimating only the values of $\hat{r}_M(\cdot)$ at the observed data points X_1, \ldots, X_{N_0} ;

Case II: estimating the values of $\hat{r}_M(\cdot)$ at both the observed data points X_1, \ldots, X_{N_0} and n new points $x_1, \ldots, x_n \in \mathbb{R}^d$.

Case I. In many applications, we are only interested in a functional of density ratios at observed sample points, i.e., the values of $\Phi(\{r(X_i)\}_{i=1}^{N_0})$ for some given functions Φ defined on \mathbb{R}^{N_0} . Check, e.g., (6.1) and – in a slightly different but symmetric form – (5.2) ahead for such examples on Kullback-Leibler (KL) divergence and ATE estimation. To this end, it is natural to consider the plug-in estimator $\Phi(\{\hat{r}_M(X_i)\}_{i=1}^{N_0})$, for which it suffices to compute the values of $\{\hat{r}_M(X_i)\}_{i=1}^{N_0}$.

Built on the k-d tree structure (Bentley, 1975) for tracking NNs, Algorithm 1 below outlines an easy to implement algorithm to simultaneously compute all the values of $\{\hat{r}_M(X_i)\}_{i=1}^{N_0}$. This algorithm could be regarded as a direct extension of the celebrated Friedman-Bentley-Finkel algorithm (Friedman et al., 1977) to the NN matching setting.

Algorithm 1: Density ratio estimators at sample points.

Input: $\{X_i\}_{i=1}^{N_0}, \{Z_j\}_{j=1}^{N_1}, \text{ and } M.$ Output: $\{\widehat{r}_M(X_i)\}_{i=1}^{N_0}$. Build a k-d tree using $\{X_i\}_{i=1}^{N_0}$; for $j = 1 : N_1$ do $\begin{bmatrix} \text{Search the } M\text{-NNs of } Z_j \text{ in } \{X_i\}_{i=1}^{N_0} \text{ using the } k\text{-d tree}; \\ \text{Store the indices of the } M\text{-NNs of } Z_j \text{ as } S_j; \\ \text{Count and store the number of occurrence in } \bigcup_{j=1}^{N_1} S_j \text{ for each element in } \llbracket N_0 \rrbracket$, which is then $\{K_M(i)\}_{i=1}^{N_0}$; Obtain $\{\widehat{r}_M(X_i)\}_{i=1}^{N_0}$ based on (2.4).

Case II. Suppose we are interested in estimating density ratios at both the observed and n new points in \mathbb{R}^d . A naive algorithm is then to insert each new point into observed points and perform Algorithm 1 in order. However, this algorithm is not ideal as the corresponding time complexity would be n times the complexity of Algorithm 1, which could be computationally heavy with a large number of new points.

Instead, we develop a more sophisticated implementation. Let the new points be $\{x_i\}_{i=1}^n$. Algorithm 2 computes all the values of $\{\widehat{r}_M(x_i)\}_{i=1}^n$ as well as $\{\widehat{r}_M(X_i)\}_{i=1}^{N_0}$. The key message delivered

Algorithm 2: Density ratio estimators at both sample and new points.

Input: $\{X_i\}_{i=1}^{N_0}, \{Z_j\}_{j=1}^{N_1}, M$, and new points $\{x_i\}_{i=1}^n$. Output: $\{\hat{r}_M(X_i)\}_{i=1}^{N_0}$ and $\{\hat{r}_M(x_i)\}_{i=1}^n$. Build a k-d tree using $\{X_i\}_{i=1}^{N_0} \cup \{x_i\}_{i=1}^n$; for $j = 1 : N_1$ do Set S_j and S'_j be two empty sets; $m \leftarrow 1$; while $|S_j| < M$ do Search the m-th NN of Z_j in $\{X_i\}_{i=1}^{N_0} \cup \{x_i\}_{i=1}^n$; if the m-th NN of Z_j is $in \{X_i\}_{i=1}^{N_0} \cup \{x_i\}_{i=1}^n$; if the m-th NN of Z_j is $in \{X_i\}_{i=1}^{N_0}$ then \mid add the index into S_j ; else \lfloor add the index into S'_j ; $m \leftarrow m + 1$; Store the indices sets S_j and S'_j ; Count and store the number of occurrence in $\bigcup_{j=1}^{N_1} S_j$ for each element in $[N_0]$, which is then $\{K_M(i)\}_{i=1}^{N_0}$. Count and store the number of occurrence in $\bigcup_{j=1}^{N_1} S'_j$ for each element in [n], which is then $\{K_M(x_i)\}_{i=1}^n$;

Obtain $\{\widehat{r}_M(X_i)\}_{i=1}^{N_0}$ and $\{\widehat{r}_M(x_i)\}_{i=1}^n$ based on (2.4).

here is that, compared to the aforementioned naive implementation, in Algorithm 2 we only need to construct one single k-d tree; the matching elements are then categorized to two different sets, corresponding to those with regard to X_i 's and x_i 's, separately. Such an implementation is thus intuitively much more efficient.

The following is the main theorem of this section, elaborating on the computational advantage of the proposed estimator.

Theorem 3.1. (1) The average time complexity of Algorithm 1 to compute all the values of $\{\hat{r}_M(X_i)\}_{i=1}^{N_0}$ is

$$O\Big((d+N_1M/N_0)N_0\log N_0\Big).$$

(2) Assume $[x_i]_{i=1}^n$ are independent and identically distributed (i.i.d.) following ν_0 and are independent of $[X_i]_{i=1}^{N_0}$. Then the average time complexity of Algorithm 2 to compute all the values of $\{\widehat{r}_M(x_i)\}_{i=1}^n$ and $\{\widehat{r}_M(X_i)\}_{i=1}^{N_0}$ is

$$O((d + N_1 M/N_0)(N_0 + n)\log(N_0 + n))$$

Remark 3.1 (Comparison to non-NN-based estimators). Assuming $N_0 \simeq N_1 \simeq N$, it is worth noting that optimization-based methods are commonly of a time complexity $O(N^2)$ if not worse (Noshad et al., 2017). They are thus less appealing in terms of handling gigantic data as was argued in, e.g., astronomy (Lima et al., 2008; Kremer et al., 2015) and big text analysis (Borgeaud et al., 2021) applications. **Remark 3.2** (Comparison to the two-step NN-based density ratio estimator). Regarding Case I, a direct calculation yields that the time complexity of the simple two-step NN-based method, which separately estimates f_1 and f_0 based on individual *M*-NN density estimators, is

$$O(dN_0 \log N_0 + dN_1 \log N_1 + N_0 M \log N_0 + N_0 M \log N_1).$$

It is thus of the same order as Algorithm 1 when $N_1 \simeq N_0$, while computationally heavier when $N_1 \prec N_0$. Regarding Case II, the time complexity of the simple two-step NN-based method is

$$O(dN_0 \log N_0 + dN_1 \log N_1 + (N_0 + n)M \log N_0 + (N_0 + n)M \log N_1)$$

Thus, if n is of less or equal order of N_0 , it is of the same order when $N_1 \simeq N_0$, while computationally heavier than Algorithm 2 when $N_1 \prec N_0$.

Remark 3.3 (Comparison to the one-step NN-based density ratio estimator in Noshad et al. (2017)). It is worth noting that, in order to estimate *f*-divergence measures, Noshad et al. (2017) constructed another one-step NN-based estimator admitting the following simple form,

$$\widehat{r}'_M(x) = \frac{N_0}{N_1} \frac{\mathcal{M}_i}{\mathcal{N}_i + 1},$$

where \mathcal{N}_i and \mathcal{M}_i are the numbers of points in $\{X_i\}_{i=1}^{N_0}$ and $\{Z_i\}_{i=1}^{N_1}$ among the *M* NNs of *x*; cf. Noshad et al. (2017, Equ. (20)). For Case I, its time complexity is

$$O(d(N_0 + N_1)\log(N_0 + N_1) + N_0M\log(N_0 + N_1));$$

while for Case II it is

$$O(d(N_0 + N_1)\log(N_0 + N_1) + (N_0 + n)M\log(N_0 + N_1))$$

Both are at the same order as the naive NN-based one, but unlike the naive approach, this estimator is indeed one-step. However, it is still theoretically unclear if this estimator is statistically efficient; see Remark 4.4 ahead for more details.

4 Density ratio estimation III: theory

This section introduces the theory for density ratio estimation based on NN matching. To this end, before establishing detailed theoretical properties (e.g., consistency and the rate of convergence) for $\hat{r}_M(\cdot)$, we first exhibit a lemma elaborating on the (asymptotic) L^p moments of $\nu_1(A_M(x))$, the ν_1 -measure of the catchment area. This novel result did not appear in Abadie and Imbens's analysis. It is also of independent interest in stochastic and computational geometry in light of Remark 2.2.

Lemma 4.1 (Asymptotic L^p moments of catchment areas's ν_1 -measure). Assuming $M \log N_0/N_0 \rightarrow 0$ as $N_0 \rightarrow \infty$, we have

$$\lim_{N_0 \to \infty} \frac{N_0}{M} \mathbf{E} \left[\nu_1 \left(A_M(x) \right) \right] = r(x)$$

holds for ν_0 -almost all x. If we further assume $M \to \infty$, then for any positive integer p, we have

$$\lim_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbf{E} \left[\nu_1^p \left(A_M(x)\right)\right] = \left[r(x)\right]^p$$

holds for ν_0 -almost all x.

Remark 4.1 (Relation to the measure of Voronoi cells). When M = 1 and $\nu_0 = \nu_1$, the measure of catchment areas reduces to the measure of Voronoi cells as pointed out in Remark 2.2. Interestingly, in the stochastic geometry literature, Devroye et al. (2017) studied a related problem of bounding the moments of the measure of Voronoi cells (cf. Theorem 2.1 therein). Setting M = 1 and $\nu_0 = \nu_1$ in the first part of Lemma 4.1 and recalling Remark 2.2, we can derive their Theorem 2.1(i). On the other hand, Devroye et al. (2017, Theorem 2.1(ii)) showed that, as $\nu_0 = \nu_1$, p = 2, and $d \leq 3$, unlike $(M^{-1}N_0)^2 \mathbb{E}[\nu_1^2(A_M(x))]$, $N_0^2 \mathbb{E}[\nu_1^2(A_1(x))]$ does not converge to 1; cf. Devroye et al. (2017, Section 4.2). This supports the necessity of forcing $M \to \infty$ for stabilizing the moments of $\hat{r}_M(\cdot)$.

4.1 Consistency

We first establish the pointwise consistency of the estimator $\hat{r}_M(x)$ for estimating r(x). This requires nearly no assumption on ν_0, ν_1 except for those made at the beginning of Section 2, in line with similar observations made in NN-based density estimation (Biau and Devroye, 2015, Theorem 3.1).

Theorem 4.1 (Pointwise consistency). Assume $M \log N_0/N_0 \rightarrow 0$.

(i) (Asymptotic unbiasedness) For ν_0 -almost all x, we have

$$\lim_{N_0 \to \infty} \mathbf{E} \big[\widehat{r}_M(x) \big] = r(x).$$

(ii) (Pointwise L_p consistency) Let p be any positive integer and assume $MN_1/N_0 \to \infty$ and $M \to \infty$ as $N_0 \to \infty$. Then for ν_0 -almost all x, we have

$$\lim_{N_0 \to \infty} \mathbf{E} \left[|\widehat{r}_M(x) - r(x)|^p \right] = 0.$$

For evaluating the global consistency of the estimator, on the other hand, it is necessary to introduce the following (global) L_p risk:

$$L_{p} \text{ risk} := \mathbb{E} \left[\left| \hat{r}_{M}(X) - r(X) \right|^{p} \mid X_{1}, \dots, X_{N_{0}}, Z_{1}, \dots, Z_{N_{1}} \right] \\ = \int_{\mathbb{R}^{d}} \left| \hat{r}_{M}(x) - r(x) \right|^{p} f_{0}(x) \mathrm{d}x,$$
(4.1)

where X is a copy drawn from ν_0 that is independent of the data. For the L_p risk consistency of the estimator, we impose conditions on ν_0 and ν_1 further as follows.

Denote the supports of ν_0 and ν_1 by S_0 and S_1 , respectively. For any set $S \subset \mathbb{R}^d$, denote the diameter of S by

$$\operatorname{diam}(S) := \sup_{x,z \in S} \left\| x - z \right\|.$$

Assumption 4.1. (i) ν_0, ν_1 are two probability measures on \mathbb{R}^d , both are absolutely continuous with respect to λ , and ν_1 is absolutely continuous with respect to ν_0 .

- (ii) There exists a constant R > 0 such that diam $(S_0) \leq R$.
- (iii) There exist two constants $f_L, f_U > 0$ such that for any $x \in S_0$ and $z \in S_1, f_L \leq f_0(x) \leq f_U$ and $f_1(z) \leq f_U$.

(iv) There exists a constant $a \in (0,1)$ such that for any $\delta \in (0, \operatorname{diam}(S_0)]$ and $z \in S_1$,

$$\lambda(B_{z,\delta} \cap S_0) \ge a\lambda(B_{z,\delta}),$$

recalling that $B_{z,\delta}$ represents the closed ball in \mathbb{R}^d with center at z and radius δ .

Remark 4.2. Assumption 4.1 is standard in the literature for establishing the global consistency of density ratio estimators. The regularity conditions on the support ensure that the angle of the support is not too sharp, which trivially hold for any *d*-dimensional cube. These conditions were also enforced in Nguyen et al. (2010, Theorem 1), Sugiyama et al. (2008, Assumption 1), Kpotufe (2017, Definition 1), among many others.

We then establish the L_p risk consistency of the estimator via the Hardy–Littlewood maximal inequality (Stein, 2016); cf. Lemma 7.2 ahead. Of note, this inequality was used in Han et al. (2020) in a relative manner in order to study the information-theoretical limit of entropy estimation.

Theorem 4.2 (L_p risk consistency). Assume the pair of ν_0, ν_1 satisfies Assumption 4.1. Let p be any positive integer. Assume further that $M \log N_0/N_0 \rightarrow 0$, $MN_1/N_0 \rightarrow \infty$, and $M \rightarrow \infty$ as $N_0 \rightarrow \infty$. We then have

$$\lim_{N_0 \to \infty} \mathbf{E} \Big[\int_{\mathbb{R}^d} \left| \widehat{r}_M(x) - r(x) \right|^p f_0(x) \mathrm{d}x \Big] = 0.$$

As a direct corollary of Theorem 4.2, one can obtain the limit of any finite moment of $\nu_1(A_M(\cdot))$ with a random center. This could be regarded as a global extension of Lemma 4.1.

Corollary 4.1. Assume the same conditions as in Theorem 4.2. We then have

$$\lim_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p(A_M(W))\right] = \mathbf{E}\left(\left[r(W)\right]^p\right),$$

where W follows an arbitrary distribution that is absolutely continuous with respect to ν_0 and has density bounded upper and below by two positive constants. In particular, it holds when W is drawn from ν_0 .

4.2 Rates of Convergence

In this section we establish the rates of convergence for $\hat{r}(x)$ under both pointwise and global measures. We first consider the pointwise mean square error (MSE) convergence rate and show that $\hat{r}_M(\cdot)$ is minimax optimal in that regard. In the sequel, we fix an $x \in \mathbb{R}^d$ and consider the following local assumption on (ν_0, ν_1) .

Assumption 4.2 (Local assumption). (i) ν_0, ν_1 are two probability measures on \mathbb{R}^d , both are absolutely continuous with respect to λ , and ν_1 is absolutely continuous with respect to ν_0 .

- (ii) There exist two constants $f_L, f_U > 0$ such that $f_0(x) \ge f_L$ and $f_1(x) \le f_U$.
- (iii) There exists a constant $\delta > 0$ such that for any $z \in B_{x,\delta}$,

$$|f_0(x) - f_0(z)| \lor |f_1(x) - f_1(z)| \le L ||x - z||,$$

for some constants L > 0.

Define the following probability class

$$\mathcal{P}_{x,p}(f_L, f_U, L, d, \delta) := \Big\{ (\nu_0, \nu_1) : \text{Assumption 4.2 holds} \Big\}.$$

The following theorem establishes the uniform pointwise convergence rate of $\hat{r}_M(\cdot)$.

Theorem 4.3 (Pointwise rates of convergence). Assume $M \log N_0/N_0 \to 0$ and $M/\log N_0 \to \infty$, and consider a sufficiently large N_0 .

(i) Asymptotic bias:

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{x,p}(f_L,f_U,L,d,\delta)} \left| \mathbb{E}\big[\widehat{r}_M(x)\big] - r(x) \right| \le C\Big(\frac{M}{N_0}\Big)^{1/d},$$

where C > 0 is a constant only depending on f_L, f_U, L, d .

Further assume $MN_1/N_0 \rightarrow \infty$.

(ii) Asymptotic variance:

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{x,\mathbf{p}}(f_L,f_U,L,d,\delta)} \operatorname{Var}\left[\widehat{r}_M(x)\right] \le C' \left[\left(\frac{1}{M}\right) + \left(\frac{N_0}{MN_1}\right) \right],$$

where C' > 0 is a constant only depending on f_L, f_U .

(iii) Asymptotic MSE:

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{x,p}(f_L,f_U,L,d,\delta)} \mathbb{E}\Big[\widehat{r}_M(x) - r(x)\Big]^2 \le C''\Big[\Big(\frac{M}{N_0}\Big)^{2/d} + \Big(\frac{1}{M}\Big) + \Big(\frac{N_0}{MN_1}\Big)\Big],$$

where C'' > 0 is a constant only depending on f_L, f_U, L, d .

Further assume $N_1^{-\frac{d}{2+d}} \log N_0 \to 0.$

(iv) Fix
$$\alpha > 0$$
 and take $M = \alpha \cdot \{N_0^{\frac{2}{2+d}} \lor (N_0 N_1^{-\frac{2}{2+d}})\}$. We have

$$\sup_{\substack{(\nu_0,\nu_1) \in \mathcal{P}_{x,p}(f_L, f_U, L, d, \delta)}} \mathbb{E}[\hat{r}_M(x) - r(x)]^2 \le C''' (N_0 \land N_1)^{-\frac{2}{2+d}}, \tag{4.2}$$

where C''' > 0 is a constant only depending on f_L, f_U, L, d, α .

The final rate of convergence in Theorem 4.3, Equ. (4.2) matches the established minimax lower bound in Lipschitz density function estimation (Tsybakov, 2009, Section 2). By some simple manipulation, the argument in Tsybakov (2009, Exercise 2.8) directly extends to density ratio as the latter is a harder statistical problem (Kpotufe, 2017, Remark 3). This is formally stated in the following proposition.

Proposition 4.1 (Pointwise MSE minimax lower bound). For all sufficiently large N_0 ,

$$\inf_{\tilde{r}} \sup_{(\nu_0,\nu_1)\in\mathcal{P}_{x,p}(f_L,f_U,L,d,\delta)} \mathbb{E}\big[\tilde{r}(x) - r(x)\big]^2 \ge c(N_0 \wedge N_1)^{-\frac{2}{2+d}}$$

where c > 0 is a constant only depending on f_L , f_U , L, d and the infimum is taken over all measurable functions.

We then move on to the case of a global risk and study the rates of convergence in this regard. To this end, a global assumption on (ν_0, ν_1) is given below.

- **Assumption 4.3** (Global assumption). (i) ν_0, ν_1 are two probability measures on \mathbb{R}^d , both are absolutely continuous with respect to λ , and ν_1 is absolutely continuous with respect to ν_0 .
 - (ii) There exists a constant R > 0 such that diam $(S_0) \leq R$.
- (iii) There exist two constants $f_L, f_U > 0$ such that for any $x \in S_0$ and $z \in S_1, f_L \leq f_0(x) \leq f_U$ and $f_1(z) \leq f_U$.
- (iv) There exists a constant $a \in (0, 1)$ such that for any $\delta \in (0, \operatorname{diam}(S_0)]$ and any $z \in S_1$,

$$\lambda(B_{z,\delta} \cap S_0) \ge a\lambda(B_{z,\delta}).$$

- (v) There exists a constant H > 0 such that the surface area (Hausdorff measure, Evans and Garzepy (2018, Section 3.3)) of S_1 is bounded by H.
- (vi) There exists a constant L > 0 such that for any $x, z \in S_1$,

$$|f_0(x) - f_0(z)| \lor |f_1(x) - f_1(z)| \le L ||x - z||.$$

Remark 4.3. Assumption 4.3 is standard in the literature for establishing the global risk of density ratio estimators; similar assumptions were made in Zhao and Lai (2020a, Assumption 1) and Zhao and Lai (2020b, Assumption 1). Note that the regularity conditions on the support automatically hold for d-dimensional cubes, and the restriction on the surface area is added to control the boundary effect on NN-based methods.

Define the following probability class

$$\mathcal{P}_{g}(f_L, f_U, L, d, a, H, R) := \left\{ (\nu_0, \nu_1) : \text{Assumption 4.3 holds} \right\}.$$

$$(4.3)$$

The next theorem establishes the uniform rate of convergence of $\hat{r}(\cdot)$ within the above probability class and under the L_1 risk. This rate is further matched by a minimax lower bound derived in Theorem 1 of Zhao and Lai (2020a) using similar arguments as in the pointwise case.

Theorem 4.4 (Global rates of convergence under the L_1 risk). Assume $M \log N_0/N_0 \to 0$, $M/\log N_0 \to \infty$, $MN_1/N_0 \to \infty$, and consider a sufficiently large N_0 .

(i) We have the following uniform upper bound,

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{g}(f_L,f_U,L,d,a,H,R)} \mathbb{E}\Big[\int_{\mathbb{R}^d} \Big|\widehat{r}_M(x) - r(x)\Big| f_0(x) \mathrm{d}x\Big] \le C\Big[\Big(\frac{M}{N_0}\Big)^{1/d} + \Big(\frac{1}{M}\Big)^{1/2} + \Big(\frac{N_0}{MN_1}\Big)^{1/2}\Big],$$

where C > 0 is a constant only depending on f_L, f_U, a, H, L, d .

(ii) Further assume $N_1^{-\frac{d}{2+d}} \log N_0 \to 0$, fix $\alpha > 0$, and take $M = \alpha \cdot \{N_0^{\frac{2}{2+d}} \lor (N_0 N_1^{-\frac{d}{2+d}})\}$. We then have

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{g}(f_L,f_U,L,d,a,H,R)} \mathbb{E}\Big[\int_{\mathbb{R}^d} \Big|\widehat{r}_M(x) - r(x)\Big| f_0(x) \mathrm{d}x\Big] \le C'(N_0 \wedge N_1)^{-\frac{1}{2+d}},$$

where C' > 0 is a constant only depending on $f_L, f_U, a, H, L, d, \alpha$.

Proposition 4.2 (Global minimax lower bound under the L_1 risk). If a is sufficiently small and H, R are sufficiently large, then for all sufficiently large N_0 ,

$$\inf_{\widetilde{r}} \sup_{(\nu_0,\nu_1)\in\mathcal{P}_{g}(f_L,f_U,L,d,a,H,R)} \mathbb{E}\Big[\int_{\mathbb{R}^d} \left|\widetilde{r}(x) - r(x)\right| f_0(x) \mathrm{d}x\Big] \ge c(N_0 \wedge N_1)^{-\frac{1}{2+d}}$$

where c > 0 is a constant only depending on f_L , f_U , L, d and the infimum is taken over all measurable functions.

Remark 4.4 (Comparison to the one-step estimator in Noshad et al. (2017)). The estimator introduced in Remark 3.3 by Noshad et al. (2017) is, to our knowledge, the only alternative density ratio estimator in the literature that is able to attain both the properties (P1) and (P2). However, the arguments in Noshad et al. (2017, Section III) can only yield the bound

$$\mathbf{E}\left[\widehat{r}'_{M}(x) - r(x)\right]^{2} \lesssim \left(\frac{M}{N_{0}}\right)^{1/d} + \left(\frac{1}{M}\right)$$

for $(\nu_0, \nu_1) \in \mathcal{P}_{x,p}(f_L, f_U, L, d, \delta)$. This is via Equ. (21) therein, de-poissonizing the estimator, and further assuming N_1/N_0 converges to a positive constant. The above bound is strictly looser than the bound $(M/N_0)^{2/d} + M^{-1}$ for $\hat{r}_M(\cdot)$ shown in Theorem 4.3. However, it seems mathematically challenging to improve their analysis and accordingly, unlike $\hat{r}_M(\cdot)$, it is still theoretically unclear if the estimator $\hat{r}'_M(x)$ is a statistically efficient density ratio estimator.

5 Revisiting the matching-based estimator of the ATE

This section studies the bias-corrected NN matching-based estimator of the ATE, proposed in Abadie and Imbens (2011) to correct the asymptotic bias of the original matching-based estimator derived by Abadie and Imbens (2006). To this end, we leverage the new insights obtained in the last three sections, and bridge the study to both the classic double robustness framework (Scharfstein et al., 1999; Bang and Robins, 2005) and the modern double machine learning framework (Chernozhukov et al., 2018).

We first introduce the setup for the NN matching-based estimator and its bias-corrected version. Following Abadie and Imbens (2006), let $[(X_i, D_i, Y_i)]_{i=1}^n$ be *n* independent independent copies of (X, D, Y), where $D \in \{0, 1\}$ is a binary variable, $X \in \mathbb{R}^d$ represents the individual covariates, assumed to be absolute continuous admitting a density f_X , and $Y \in \mathbb{R}$ stands for the outcome variable.

For each unit $i \in [n]$, we observe $D_i = 1$ if in the treated group and $D_i = 0$ if in the control group. Let $n_0 := \sum_{i=1}^n (1 - D_i)$ and $n_1 := \sum_{i=1}^n D_i$ be the numbers of control and treated units, respectively. Under the potential outcome framework (Rubin, 1974), the unit *i* has two potential outcomes, $Y_i(1)$ and $Y_i(0)$, but we observe only one of them:

$$Y_i = \begin{cases} Y_i(0), & \text{if } D_i = 0, \\ Y_i(1), & \text{if } D_i = 1. \end{cases}$$

The goal is to estimate the following population ATE,

$$\tau := \mathbf{E}\Big[Y_i(1) - Y_i(0)\Big],$$

based on the observations $\{(X_i, D_i, Y_i)\}_{i=1}^n$. To estimate ATE, we consider its empirical counterpart

$$\widehat{\tau}_M := \frac{1}{n} \sum_{i=1}^n \left[\widehat{Y}_i(1) - \widehat{Y}_i(0) \right],$$

where $\widehat{Y}_i(0)$ and $\widehat{Y}_i(1)$ are the imputed outcomes of $Y_i(0)$ and $Y_i(1)$. With a fixed M, the matchingbased estimator in Abadie and Imbens (2006) imputes the missing potential outcomes by

$$\widehat{Y}_{i}(0) := \begin{cases} Y_{i}, & \text{if } D_{i} = 0, \\ \frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{0}(i)} Y_{j}, & \text{if } D_{i} = 1, \end{cases} \text{ and } \widehat{Y}_{i}(1) := \begin{cases} \frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{1}(i)} Y_{j}, & \text{if } D_{i} = 0, \\ Y_{i}, & \text{if } D_{i} = 1. \end{cases}$$

Here for $\omega \in \{0,1\}$, $\mathcal{J}_M^{\omega}(i)$ represents the index set of *M*-NNs of X_i in $\{X_j : D_j = \omega\}_{j=1}^n$, i.e., the set of all indices $j \in [n]$ such that $D_j = \omega$ and

$$\sum_{\ell=1, D_{\ell}=\omega}^{n} \mathbb{1}\left(\|X_{\ell} - X_{i}\| \le \|X_{j} - X_{i}\| \right) \le M$$

Let $K_M^{\omega}(i)$ be the number of matched times for unit *i* such that $D_i = \omega$, i.e.,

$$K_M^{\omega}(i) := \sum_{j=1, D_j=1-\omega}^n \mathbb{1}\left(i \in \mathcal{J}_M^{\omega}(j)\right).$$

With the above notation and concepts, the matching-based estimator in Abadie and Imbens (2006) can be written as

$$\widehat{\tau}_M = \frac{1}{n} \Big[\sum_{i=1, D_i=1}^n \Big(1 + \frac{K_M^1(i)}{M} \Big) Y_i - \sum_{i=1, D_i=0}^n \Big(1 + \frac{K_M^0(i)}{M} \Big) Y_i \Big].$$
(5.1)

However, when d > 1, the bias term of $\hat{\tau}_M$ is asymptotically non-negligible (Abadie and Imbens, 2006). To fix this, Abadie and Imbens (2011) proposed the following bias-corrected version for $\hat{\tau}_M$. In detail, let $\hat{\mu}_0(x)$ and $\hat{\mu}_1(x)$ be mappings from \mathbb{R}^d to \mathbb{R} that estimate the conditional means of the outcomes

 $\mu_0(x) := \mathrm{E}[Y \,|\, X = x, D = 0] \ \, \text{and} \ \, \mu_1(x) := \mathrm{E}[Y \,|\, X = x, D = 1],$

respectively, with the corresponding residuals

$$\widehat{R}_i := Y_i - \widehat{\mu}_{D_i}(X_i), \quad i \in \llbracket n \rrbracket.$$

The estimator based on the outcomes regression is

$$\widehat{\tau}^{\operatorname{reg}} := n^{-1} \sum_{i=1}^{n} \Big[\widehat{\mu}_1(X_i) - \widehat{\mu}_0(X_i) \Big].$$

One is then ready to check that the bias-corrected matching-based estimator in Abadie and Imbens (2011) has the following equivalent form, summarized as a lemma.

Lemma 5.1. The bias-corrected matching-based estimator in Abadie and Imbens (2011) can be rewritten as

$$\widehat{\tau}_{M}^{\rm bc} = \widehat{\tau}^{\rm reg} + \frac{1}{n} \Big[\sum_{i=1, D_i=1}^{n} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \widehat{R}_{i} - \sum_{i=1, D_i=0}^{n} \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \widehat{R}_{i} \Big].$$
(5.2)

Equ. (5.2) is related to doubly robust estimators. To compare them, we review the general double robustness framework. In detail, we first have some outcome models and residuals defined in the same way as above, and then let

$$\widehat{e}(x): \mathbb{R}^d \to \mathbb{R}$$

be a genetic estimator of the propensity score

$$e(x) := P(D = 1 | X = x).$$

The doubly robust estimator in Scharfstein et al. (1999) and Bang and Robins (2005) could then be formulated as

$$\hat{\tau}^{\rm dr} = \hat{\tau}^{\rm reg} + \frac{1}{n} \Big[\sum_{i=1, D_i=1}^n \frac{1}{\hat{e}(X_i)} \hat{R}_i - \sum_{i=1, D_i=0}^n \frac{1}{1 - \hat{e}(X_i)} \hat{R}_i \Big].$$
(5.3)

Notice that conditional on $\mathbf{D} := (D_1, \ldots, D_n)$, $[X_i : D_i = \omega]_{i=1}^n$ are n_{ω} i.i.d. random variables sampled from the distribution of $X \mid D = \omega$, and the two groups of sample points,

$$[X_i: D_i = 0]_{i=1}^n$$
 and $[X_i: D_i = 1]_{i=1}^n$,

are mutually independent. Denote the density of $X | D = \omega$ by $f_{X | D = \omega}$. From the construction of $K_M^0(i), K_M^1(i)$ and results in Section 4, under the density assumptions and an appropriate choice of M, conditional on D,

$$\frac{n_0}{n_1} \frac{K_M^0(i)}{M}$$
 and $\frac{n_1}{n_0} \frac{K_M^1(i)}{M}$

are consistent estimators of

$$\frac{f_{X \mid D=1}(X_i)}{f_{X \mid D=0}(X_i)}$$
 and $\frac{f_{X \mid D=0}(X_i)}{f_{X \mid D=1}(X_i)}$,

respectively. Noting further that n_1/n_0 converges almost surely to P(D = 1)/P(D = 0) by the law of large numbers, the following two statistics

$$\frac{K_M^0(i)}{M}$$
 and $\frac{K_M^1(i)}{M}$

are then consistent estimators of

$$\frac{e(X_i)}{1 - e(X_i)} \quad \text{and} \quad \frac{1 - e(X_i)}{e(X_i)},$$

respectively. Thus, in view of (5.3), the bias-corrected matching-based estimator $\hat{\tau}_M^{\text{bc}}$ in (5.2) is actually a doubly robust estimator of τ , and accordingly, should also enjoy all the desirable properties of doubly robust estimators. This novel insight into Abadie and Imbens (2011)'s bias-corrected matching estimator allows us to establish its asymptotic properties with a diverging M, which is the main topic in the rest of this section. It is worth mentioning here that Abadie and Imbens never pointed out the relation between matching-based ATE estimators and the double robustness framework, and there is no study in the double robustness literature that has analyzed NN matchingbased estimators.

To formally state the properties, we first leverage the results of Chernozhukov et al. (2018). In the sequel, let $U_{\omega} := Y(\omega) - \mu_{\omega}(X)$ for $\omega \in \{0, 1\}$ and X be the support of X. **Assumption 5.1.** (i) For almost all $x \in \mathbb{X}$, D is independent of (Y(0), Y(1)) conditional on X = x, and there exists some constant $\eta > 0$ such that $\eta < P(D = 1 | X = x) < 1 - \eta$.

- (ii) $[(X_i, D_i, Y_i)]_{i=1}^n$ are i.i.d. following the joint distribution of (X, D, Y).
- (iii) $\mathbb{E}[U_{\omega}^2 | X = x]$ is uniformly bounded for almost all $x \in \mathbb{X}$ and $\omega \in \{0, 1\}$.
- (iv) $E[\mu_{\omega}^2]$ is bounded for $\omega \in \{0, 1\}$.

Assumption 5.2. (i) $E[U_{\omega}^2]$ is bounded away from zero for $\omega \in \{0, 1\}$.

(ii) There exists some constants $\kappa > 0$ such that $E[|Y|^{2+\kappa}]$ is bounded.

Assumption 5.3. For $\omega \in \{0,1\}$, there exists a deterministic function $\bar{\mu}_{\omega}(\cdot) : \mathbb{R}^d \to \mathbb{R}$ such that $\mathrm{E}[\bar{\mu}^2_{\omega}(X)]$ is bounded and the estimator $\hat{\mu}_{\omega}(x)$ satisfies

$$\|\widehat{\mu}_{\omega} - \overline{\mu}_{\omega}\|_{\infty} = o_{\mathrm{P}}(1),$$

where $\|\cdot\|_{\infty}$ denotes the function L_{∞} norm.

Assumption 5.4. For $\omega \in \{0,1\}$, the estimator $\widehat{\mu}_{\omega}(x)$ satisfies

$$\|\widehat{\mu}_{\omega} - \mu_{\omega}\|_{\infty} = o_{\mathrm{P}}(1)$$

Assumption 5.5. For $\omega \in \{0, 1\}$, the estimator $\widehat{\mu}_{\omega}(x)$ satisfies

$$\|\widehat{\mu}_{\omega} - \mu_{\omega}\|_{\infty} = o_{\mathrm{P}}(n^{-d/(4+2d)})$$

Remark 5.1. Assumption 5.1(i) is the unconfoundedness and overlap assumptions, and is often referred to as the strong ignorability condition (Rosenbaum and Rubin, 1983). Assumption 5.2 corresponds to Assumption 5.1 in Chernozhukov et al. (2018), and are similar to Assumption 4 in Abadie and Imbens (2006). Assumption 5.3 allows for the misspecification of the outcome models; for example, if $\hat{\mu}_{\omega} = \bar{\mu}_{\omega} = 0$, $\hat{\tau}_{M}^{\rm bc}$ then reduces to $\hat{\tau}_{M}$. Assumption 5.4 assumes that the outcome models are correctly specified. Assumption 5.5 assumes approximation accuracy of the outcome model. Abadie and Imbens (2011) uses the power series approximation (Newey, 1997) to estimate the outcome model, which under some classic nonparametric statistics assumptions automatically satisfy Assumption 5.5 (cf. Lemma A.1 in Abadie and Imbens (2011)).

Remark 5.2. In Assumption 5.5, we assume an approximation rate under L_{∞} norm. This is different from the L_2 norm put in Assumption 5.1 in Chernozhukov et al. (2018), but can be handled with some trivial modification to the proof of Chernozhukov et al. (2018, Theorem 5.1) since one can replace the Cauchy–Schwarz inequality by the L_1 - L_{∞} Hölder's inequality. Theorem 4.4 can then be applied directly.

Lastly, we review the semiparametric efficiency lower bound for estimating ATE (Hahn, 1998):

$$\sigma^{2} := \mathbf{E} \Big[\mu_{1}(X) - \mu_{0}(X) + \frac{D(Y - \mu_{1}(X))}{e(X)} - \frac{(1 - D)(Y - \mu_{0}(X))}{1 - e(X)} - \tau \Big]^{2}.$$

Theorem 4.2 above and standard results on the double machine learning estimators (cf. Chernozhukov et al. (2018, Theorem 5.1)) then imply the following theorem, showing that, once allowing M to diverge at an appropriate rate, $\hat{\tau}_M^{\rm bc}$ in (5.2) constitutes a doubly robust estimator of τ . In addition, the counterpart of $\hat{\tau}_M^{\rm bc}$ via sample splitting and cross fitting (Chernozhukov et al., 2018, Definition 3.1), denoted by $\tilde{\tau}_{M,K}^{\text{bc}}$ with $K \geq 2$ representing a fixed number of partitions, is semiparametrically efficient. Check the beginning of Theorem 5.1's proof for a formal definition of $\tilde{\tau}_{M,K}^{\text{bc}}$.

Theorem 5.1. (i) (Double robustness) On one hand, if the distribution of (X, D, Y) satisfies Assumptions 5.1, 5.3, and either $(P_{X|D=0}, P_{X|D=1})$ or $(P_{X|D=1}, P_{X|D=0})$ satisfies Assumption 4.1, then if $M \log n/n \to 0$ and $M \to \infty$ as $n \to \infty$,

 $\widehat{\tau}_M^{\mathrm{bc}} - \tau \xrightarrow{\mathbf{p}} 0 \quad \text{and} \quad \widetilde{\tau}_{M,K}^{\mathrm{bc}} - \tau \xrightarrow{\mathbf{p}} 0.$

On the other hand, if the distribution of (X, D, Y) satisfies Assumptions 5.1 and 5.4, then

$$\widehat{\tau}_{M}^{\mathrm{bc}} - \tau \xrightarrow{\mathsf{p}} 0 \quad \text{and} \quad \widetilde{\tau}_{M,K}^{\mathrm{bc}} - \tau \xrightarrow{\mathsf{p}} 0.$$

(ii) (Semiparametric efficiency of $\tilde{\tau}_{M,K}^{\text{bc}}$) Assume the distribution of (X, D, Y) satisfies Assumptions 5.1, 5.2, 5.5 and either $(P_{X|D=0}, P_{X|D=1})$ or $(P_{X|D=1}, P_{X|D=0})$ satisfies Assumption 4.3. Then if we pick $M = \alpha n^{\frac{2}{2+d}}$ for some constant $\alpha > 0$,

$$\sqrt{n}(\widetilde{\tau}_{M,K}^{\mathrm{bc}}-\tau) \xrightarrow{\mathsf{d}} N(0,\sigma^2).$$

In addition, letting

$$\widehat{\sigma}^2 := \frac{1}{n} \sum_{i=1}^n \Big[\widehat{\mu}_1(X_i) - \widehat{\mu}_0(X_i) + D_i \Big(1 + \frac{K_M^1(i)}{M} \Big) \widehat{R}_i - (1 - D_i) \Big(1 + \frac{K_M^0(i)}{M} \Big) \widehat{R}_i - \widehat{\tau}_M^{\mathrm{bc}} \Big]^2,$$

we have $\widehat{\sigma}^2 \xrightarrow{\mathbf{p}} \sigma^2$.

Remark 5.3. Both $\hat{\tau}_M^{\text{bc}}$ and $\tilde{\tau}_{M,K}^{\text{bc}}$ directly estimate 1/e(X) and 1/(1-e(X)) using $1+K_M^1(X)/M$ and $1+K_M^0(X)/M$, respectively. This is slightly different from the setup in Chernozhukov et al. (2018), which considered plug-in estimators based on an estimate of e(X). Accordingly, some minor modifications are needed for employing their Theorem 5.1, and are completed in the proof of Theorem 5.1(ii).

Remark 5.4. To be in line with the double robustness terminology, we can call Assumptions 4.1 and 4.3 in Theorem 5.1 the "density (or propensity) model assumptions" and Assumptions 5.4 and 5.5 the "outcome (or regression) model assumptions". Note also that in Theorem 5.1 we require additional smoothness (Lipschitz) conditions on the density functions. This is stronger than the corresponding conditions made in Abadie and Imbens (2011) but is needed for us to prove the semiparametric efficiency of $\tilde{\tau}_{M,K}^{\rm bc}$'s based on the double machine learning framework. On the other hand, this requirement could be removed for $\hat{\tau}_M^{\rm bc}$ via a more careful treatment based on the particular structure of the matching-based estimators; cf. Theorem 5.2 and Remark 5.6 ahead.

The analysis of $\hat{\tau}_M^{\text{bc}}$ itself, on the other hand, cannot be directly incorporated in the double machine learning and its analysis has to be done independently. This is given in the next theorem based on the following two sets of assumptions.

Assumption 5.6. (i) $E[U_{\omega}^2 | X = x]$ is uniformly bounded away from zero for almost all $x \in X$ and $\omega \in \{0, 1\}$.

- (ii) There exists some constants $\kappa > 0$ such that $E[|U_{\omega}|^{2+\kappa} | X = x]$ is uniformly bounded for almost all $x \in X$ and $\omega \in \{0, 1\}$.
- (iii) $\max_{t \in \Lambda_{\lfloor d/2 \rfloor + 1}} \|\partial^t \mu_\omega\|_{\infty}$ is bounded, where for any positive integer k, Λ_k is the set of all ddimensional vectors of nonnegative integers $t = (t_1, \ldots, t_d)$ such that $\sum_{i=1}^d t_i = k$ and $\lfloor \cdot \rfloor$ stands for the floor function.

Assumption 5.7. There exists some constant $\varepsilon > 0$ such that for $\omega \in \{0,1\}$, the estimator $\hat{\mu}_{\omega}(x)$ satisfies

$$\max_{t \in \Lambda_{\lfloor d/2 \rfloor + 1}} \|\partial^t \widehat{\mu}_{\omega}\|_{\infty} = O_{\mathcal{P}}(1) \quad \text{and} \quad \max_{\ell \in \llbracket \lfloor d/2 \rfloor \rrbracket} \max_{t \in \Lambda_{\ell}} \|\partial^t \widehat{\mu}_{\omega} - \partial^t \mu_{\omega}\|_{\infty} = O_{\mathcal{P}}(n^{-\gamma_{\ell}}),$$

with some constants γ_{ℓ} 's satisfying $\gamma_{\ell} > \frac{1}{2} - \frac{\ell}{d} + \varepsilon$ for $\ell = 1, 2, \dots, \lfloor d/2 \rfloor$.

Theorem 5.2 (Semiparametric efficiency of $\hat{\tau}_M^{\text{bc}}$). Assume the distribution of (X, D, Y) satisfies Assumptions 5.1, 5.6, 5.7 and either $(P_{X \mid D=0}, P_{X \mid D=1})$ or $(P_{X \mid D=1}, P_{X \mid D=0})$ satisfies Assumption 4.1. Define

$$\gamma := \min_{\ell \in \llbracket \lfloor d/2 \rfloor \rrbracket} \left\{ \left[1 - \left(\frac{1}{2} - \gamma_{\ell} + \varepsilon \right) \frac{d}{\ell} \right] \land \left[1 - \left(\frac{1}{2} + \varepsilon \right) \frac{d}{\lfloor d/2 \rfloor + 1} \right] \right\}$$

recall that γ_{ℓ} 's and the constant ε were introduced in Assumption 5.7. Then, if $M \to \infty$ as $n \to \infty$ and $M \leq n^{\gamma}$,

$$\sqrt{n}(\widehat{\tau}_M^{\mathrm{bc}} - \tau) \stackrel{\mathsf{d}}{\longrightarrow} N(0, \sigma^2).$$

If in addition Assumption 5.4 holds, we have $\widehat{\sigma}^2 \xrightarrow{\mathsf{p}} \sigma^2$.

Remark 5.5. Assumption 5.6 is comparable to Assumption A.4 and the assumptions of Theorem 2 in Abadie and Imbens (2011). Compared to the assumptions of Theorem 2 in Abadie and Imbens (2011), Assumption 5.6(iii) is weaker in the sense that we only require a finite order of smoothness. Assumption 5.7 again assumes the approximation accuracy of the outcome model, with lower convergence rates required for higher order derivatives of the outcome model. We note that under some smoothness conditions on the outcome model as made in Abadie and Imbens (2011), Assumption 5.7 is satisfied using power series approximation (Abadie and Imbens, 2011, Lemma A.1).

Remark 5.6. Due to the intrinsic structure of bias-corrected matching-based estimators, unlike a direct use of Hölder's inequality and assuming Assumption 5.5 for double machine learning estimators, we instead assume Assumption 5.7 on the approximation accuracy of the derivatives of the outcome models. There are additionally two differences between Theorem 5.1(ii) and Theorem 5.2. First, unlike in Theorem 5.1(ii) where M needs to grow polynomially fast with n, in Theorem 5.2 we only require M to (i) diverge not so fast for controlling the difference of matching units; and (ii) diverge to infinity (no matter how slowly it is) for achieving semiparametric efficiency. The sets of assumptions in Theorems 5.1(ii) and 5.2 both render semiparametric efficiency for bias-corrected matching-based estimators. Second, in Theorem 5.2 we only require Assumption 4.1 to hold for the density model. This is again weaker than the Lipschitz-type conditions (Assumption 4.3) assumed in Theorem 5.1(ii) but is in line with the observations made in Abadie and Imbens (2006) and Abadie and Imbens (2011).

As d = 1, by picking $\hat{\mu}_{\omega} = 0$ for $\omega \in \{0, 1\}$, Assumption 5.7 is automatically satisfied and the bias-corrected estimator $\hat{\tau}_M^{\text{bc}}$ reduces to the original estimator $\hat{\tau}_M$ studied in Abadie and Imbens (2006). Theorem 5.2 then directly implies the following corollary, which corresponds to Abadie and Imbens (2006, Corollary 1) but with one key difference that $M \to \infty$ here.

Corollary 5.1 (Semiparametric efficiency of $\hat{\tau}_M$ when d = 1). Assume d = 1, the distribution of (X, D, Y) satisfies Assumptions 5.1, 5.6, and either $(P_{X \mid D=0}, P_{X \mid D=1})$ or $(P_{X \mid D=1}, P_{X \mid D=0})$ satisfies Assumption 4.1. If $M \to \infty$ as $n \to \infty$ and $M \lesssim n^{\frac{1}{2}-\varepsilon}$ for some $\varepsilon > 0$, then we have

$$\sqrt{n}(\widehat{\tau}_M - \tau) \stackrel{\mathsf{d}}{\longrightarrow} N(0, \sigma^2).$$

Remark 5.7. By picking $\hat{\mu}_{\omega} = 0$ for $\hat{\tau}_M$, Assumption 5.4 is no longer satisfied. Accordingly, in Corollary 5.1, $\hat{\sigma}^2$ may not be a consistent estimator of σ^2 without additional assumptions. However, by decomposing σ^2 into the form of Theorem 1 in Hahn (1998), one could still estimate σ^2 via a similar and direct way as what is outlined in Section 4 in Abadie and Imbens (2006). We do not pursue this track in detail here as the case of d = 1 without Assumption 5.4 is beyond the main scope of this manuscript.

Remark 5.8. There are three additional problems in Abadie and Imbens (2006, 2012). First, estimation of the average treatment effect on the treated (ATT) can be incorporated in the double robustness and double machine learning framework (Theorem 5.1) and matching framework (Theorem 5.2) in the same way. Second, asymptotic normality (with an additional asymptotic bias term) of $\hat{\tau}_M$ in general d can be established as Theorem 5.2. Third, unbalanced designs with $n_0 \succ n_1$ cannot be incorporated in the double robustness and double machine learning framework. In the supplement, we shall study the above unbalanced design case as M is forced to diverge with n and establish the corresponding properties.

6 Discussion

The success of NN matching in estimating ATE raises natural question about possible extension to other functional estimation problems. In what follows we present some partial results along this direction for the KL divergence estimation, which is a topic of interest in many applications; cf. the references at the beginning of Zhao and Lai (2020b).

In detail, let's consider the same two-sample setting as Sections 2-4. Denote $\phi(x) := x \log x$ with the convention that $\phi(0) = 0$. We consider estimating the KL divergence between ν_1 and ν_0 , written as

$$D_{\phi}(\nu_1 \| \nu_0) := \int \phi(r(x)) f_0(x) \mathrm{d}x,$$

based on the data points $\{X_i\}_{i=1}^{N_0}, \{Z_j\}_{j=1}^{N_1}$. To estimate $D_{\phi}(\nu_1 \| \nu_0)$, it is natural to consider the following plug-in estimator,

$$\widehat{D}_{\phi} = \widehat{D}_{\phi} \Big(\{X_i\}_{i=1}^{N_0}, \{Z_j\}_{j=1}^{N_1} \Big) := \frac{1}{N_0} \sum_{i=1}^{N_0} \phi \Big(\widehat{r}_M(X_i) \Big),$$
(6.1)

where $\hat{r}_M(\cdot)$ is the NN matching-based density ratio estimator introduced in Section 2.

Define the following probability class

$$\mathcal{P}_{\mathrm{KL}}(f_L, f_U, L, d, a, H, R, f'_U) := \left\{ (\nu_0, \nu_1) : \text{Assumption 4.3 holds} \\ \text{and } \mathrm{E}_{X \sim \nu_0} \{ [\log(f_1(X))]^4 \} \le f'_U \right\}.$$
(6.2)

The following results establish the MSE convergence rate of \hat{D}_{ϕ} and show that the estimator is minimax optimal over $\mathcal{P}_{\mathrm{KL}}(f_L, f_U, L, d, a, H, R, f'_U)$. Under the condition that $N_0 \gtrsim N_1$, this rate is further (up to some log *n*-terms) minimax optimal in view of Han et al. (2020, Theorem 7) and similar arguments as used in Section 4.

Theorem 6.1 (Rates of convergence, KL divergence estimation). (i) Assume $M \log N_0/N_0 \to 0$, $M/\log N_0 \to \infty$, $MN_1/(N_0 \log^2 N_1) \to \infty$, $N_1^{-\frac{d}{1+d}} \log N_0 \to 0$. Then for all sufficiently large N_0 ,

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{\mathrm{KL}}(f_L,f_U,L,d,a,H,R,f'_U)} \mathbb{E}\Big[\widehat{D}_{\phi} - D_{\phi}(\nu_1 \| \nu_0)\Big]^2 \le C\Big[\frac{1}{N_0} + \frac{1}{N_1} + \Big(\frac{M}{N_0}\Big)^{2/d} + \Big(\frac{1}{M}\Big)^2 + \Big(\frac{N_0}{MN_1}\Big)^2\Big],$$

where $C > 0$ is a constant only depending on f_L , f_U , a, H, L, d, f'_U .

(*ii*) Fix $\alpha > 0$, and take $M = \alpha(N_0^{\frac{1}{1+d}}) \vee (N_0 N_1^{-\frac{d}{1+d}})$. Then for all sufficiently large N_0 ,

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{\mathrm{KL}}(f_L,f_U,L,d,R,a,H,f_U')} \mathrm{E}\Big[\widehat{D}_{\phi} - D_{\phi}(\nu_1 \| \nu_0)\Big]^2 \le C'(N_0 \wedge N_1)^{-\frac{2}{1+d}},$$

where C' > 0 is a constant only depending on $f_L, f_U, a, H, L, d, f'_U, \alpha$.

Proposition 6.1 (MSE minimax rate, KL divergence estimation). If a is sufficiently small and L, R, H, f'_U are sufficiently large, then for all sufficiently large N_0 ,

$$\inf_{\widetilde{D}_{\phi}} \sup_{(\nu_{0},\nu_{1})\in\mathcal{P}_{\mathrm{KL}}(f_{L},f_{U},L,d,a,H,R,f_{U}')} \mathbb{E}\Big[\widetilde{D}_{\phi} - D_{\phi}(\nu_{1}\|\nu_{0})\Big]^{2} \ge c\Big[\Big(\frac{1}{N_{1}}\Big)^{\frac{2}{1+d}}\Big(\frac{1}{\log N_{1}}\Big)^{\frac{2+4d}{1+d}} + \frac{1}{N_{1}}\Big],$$

where c > 0 is a constant only depending on f_L , f_U , d and the infimum is taken over all measurable functions.

7 Proofs of the main results

Additional notation. We use X and Z to represent $(X_1, X_2, \ldots, X_{N_0})$ and $(Z_1, Z_2, \ldots, Z_{N_1})$, respectively. Let U(0, 1) denote the uniform distribution on [0, 1]. In the sequel, let $U \sim U(0, 1)$ and $U_{(M)}$ be the M-th order statistic of N_0 independent random variables from U(0, 1), assumed to be mutually independent and both independent of (X, Z). It is well known that $U_{(M)}$ follows the beta distribution $\text{Beta}(M, N_0 + 1 - M)$. Let $\text{Bin}(\cdot, \cdot)$ denote the binomial distribution. Let $L_1(\mathbb{R}^d)$ denote the space of all functions $f : \mathbb{R}^d \to \mathbb{R}$ such that $\int |f(x)| dx < \infty$. For any $x \in \mathbb{R}^d$ and function $f : \mathbb{R}^d \to \mathbb{R}$, we say x is a Lebesgue point of f if

$$\lim_{\delta \to 0^+} \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f(x) - f(y)| \mathrm{d}y = 0.$$

7.1 Proof of Lemma 4.1

Proof of Lemma 4.1. From the Lebesgue differentiation theorem, for any $f \in L_1(\mathbb{R}^d)$, for λ -almost all x, x is a Lebesgue point of f. Then for ν_0 -almost all x, we have $f_0(x) > 0$ and x is a Lebesgue point of f_0 and f_1 from the absolute continuity of ν_0 and ν_1 . We then only need to consider those $x \in \mathbb{R}^d$ such that $f_0(x) > 0$ and x is a Lebesgue point of f_0 and f_1 .

We first introduce a lemma about the Lebesgue point.

Lemma 7.1. Let ν be a probability measure on \mathbb{R}^d admitting a density f with respect to the Lebesgue measure. Let $x \in \mathbb{R}^d$ be a Lebesgue point of f. We then have, for any $\epsilon \in (0,1)$, there exists $\delta = \delta_x > 0$ such that for any $z \in \mathbb{R}^d$ such that $||z - x|| \leq \delta$, we have

$$\left|\frac{\nu(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f(x)\right| \le \epsilon, \quad \left|\frac{\nu(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f(x)\right| \le \epsilon.$$

Part I. This part proves the first claim. We separate the proof of Part I into two cases based on the value of $f_1(x)$.

Case I.1. $f_1(x) > 0$. Since x is a Lebesgue point of ν_0 and ν_1 , by Lemma 7.1, for any $\epsilon \in (0, 1)$, there exists some $\delta = \delta_x > 0$ such that for any $z \in \mathbb{R}^d$ with $||z - x|| \leq \delta$, we have

$$\begin{aligned} \left| \frac{\nu_0(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_0(x) \right| &\leq \epsilon f_0(x), \quad \left| \frac{\nu_0(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_0(x) \right| \leq \epsilon f_0(x), \\ \left| \frac{\nu_1(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_1(x) \right| &\leq \epsilon f_1(x), \quad \left| \frac{\nu_1(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_1(x) \right| \leq \epsilon f_1(x). \end{aligned}$$

Accordingly, if $||z - x|| \leq \delta$, we have

$$\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} \le \frac{\nu_0(B_{z,\|x-z\|})}{\lambda(B_{z,\|x-z\|})} \frac{\lambda(B_{x,\|x-z\|})}{\nu_1(B_{x,\|x-z\|})} \le \frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)}.$$

Since $\lambda(B_{z,||x-z||}) = \lambda(B_{x,||x-z||})$, we then have

$$\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} \le \frac{\nu_0(B_{z,\|x-z\|})}{\nu_1(B_{x,\|x-z\|})} \le \frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)}.$$
(7.1)

On the other hand, for any $z \in \mathbb{R}^d$ such that $||z - x|| > \delta$,

$$\nu_0(B_{z,\|z-x\|}) \ge \nu_0(B_{z^*,\delta}) \ge (1-\epsilon)f_0(x)\lambda(B_{z^*,\delta}) = (1-\epsilon)f_0(x)\lambda(B_{0,\delta}),$$

where z^* is the intersection point of the surface of $B_{x,\delta}$ and the line connecting z and x.

Let $\eta_N = 4 \log(N_0/M)$. Since $M \log N_0/N_0 \to 0$, we can take N_0 large enough so that

$$\eta_N \frac{M}{N_0} = 4 \frac{M}{N_0} \log\left(\frac{N_0}{M}\right) < (1-\epsilon) f_0(x) \lambda(B_{0,\delta}).$$

Then for any $z \in \mathbb{R}^d$ such that $\nu_0(B_{z,||z-x||}) \leq \eta_N M/N_0$, we have $||z-x|| \leq \delta$ since otherwise it would contradict the selection of N_0 .

Let Z be a copy from ν_1 independent of the data. Then

$$E[\nu_1(A_M(x))] = P(Z \in A_M(x))$$

=P(||x - Z|| \le ||\mathcal{X}_{(M)}(Z) - Z||) = P(\nu_0(B_{Z,||x-Z||}) \le \nu_0(B_{Z,||\mathcal{X}_{(M)}(Z) - Z||})). (7.2)

For any given $z \in \mathbb{R}^d$, $[\nu_0(B_{z,||X_i-z||})]_{i=1}^{N_0}$ are i.i.d. from U(0,1) since $[X_i]_{i=1}^{N_0}$ are i.i.d. from ν_0 and we use the probability integral transform. Then $\nu_0(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||})$ has the same distribution as $U_{(M)}$ and is independent of Z.

Upper bound. With a slight abuse of notation, we note $W = \nu_0(B_{Z,||x-Z||})$. We then have, from (7.1) and (7.2),

$$E[\nu_{1}(A_{M}(x))] = P\left(W \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|})\right)$$

$$\leq P\left(W \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \le \eta_{N}\frac{M}{N_{0}}\right) + P\left(\nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) > \eta_{N}\frac{M}{N_{0}}\right)$$

$$= P\left(W \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \le \eta_{N}\frac{M}{N_{0}}, \|Z-x\| \le \delta\right) + P\left(U_{(M)} > \eta_{N}\frac{M}{N_{0}}\right)$$

$$\leq P\left(\nu_{0}(B_{Z,\|x-Z\|}) \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \|Z-x\| \le \delta\right) + P\left(U_{(M)} > \eta_{N}\frac{M}{N_{0}}\right)$$

$$\leq P\left(\frac{1-\epsilon}{1+\epsilon}\frac{f_{0}(x)}{f_{1}(x)}\nu_{1}(B_{x,\|x-Z\|}) \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \|Z-x\| \le \delta\right) + P\left(U_{(M)} > \eta_{N}\frac{M}{N_{0}}\right)$$

$$\leq P\left(\frac{1-\epsilon}{1+\epsilon}\frac{f_{0}(x)}{f_{1}(x)}\nu_{1}(B_{x,\|x-Z\|}) \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|})\right) + P\left(U_{(M)} > \eta_{N}\frac{M}{N_{0}}\right)$$

$$= P\left(\frac{1-\epsilon}{1+\epsilon}\frac{f_{0}(x)}{f_{1}(x)}U \le U_{(M)}\right) + P\left(U_{(M)} > \eta_{N}\frac{M}{N_{0}}\right).$$
(7.3)

For the second term in (7.3), notice that $\eta_N \to \infty$ as $N_0 \to \infty$. Then from the Chernoff bound and for N_0 sufficiently large, we have

$$\frac{N_0}{M} P\left(U_{(M)} > \eta_N \frac{M}{N_0}\right) = \frac{N_0}{M} P\left(Bin\left(N_0, \eta_N \frac{M}{N_0}\right) \le M\right)$$
$$\le \frac{N_0}{M} \exp\left((1 + \log \eta_N - \eta_N)M\right) \le \frac{N_0}{M} \exp\left(-\frac{1}{2}\eta_N M\right) = \left(\frac{N_0}{M}\right)^{1-2M}.$$

Since $M/N_0 \to 0$ and $M \ge 1$, we then obtain

$$\lim_{N_0 \to \infty} \frac{N_0}{M} \mathcal{P}\left(U_{(M)} > \eta_N \frac{M}{N_0}\right) = 0.$$
(7.4)

For the first term in (7.3), we have

$$\frac{N_0}{M} \operatorname{P}\left(\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} U \leq U_{(M)}\right) = \frac{N_0}{M} \int_0^1 \operatorname{P}\left(U_{(M)} \geq \frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} t\right) \mathrm{d}t$$

$$= \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \int_0^{\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} \frac{N_0}{M}} \operatorname{P}\left(U_{(M)} \geq \frac{M}{N_0} t\right) \mathrm{d}t \leq \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \int_0^\infty \operatorname{P}\left(\frac{N_0}{M} U_{(M)} \geq t\right) \mathrm{d}t$$

$$= \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \frac{N_0}{M} \operatorname{E}[U_{(M)}] = \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \frac{N_0}{N_0+1}.$$
(7.5)

We then obtain

$$\limsup_{N_0 \to \infty} \frac{N_0}{M} \mathbb{P}\left(\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} U \le U_{(M)}\right) \le \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)}.$$
(7.6)

Plugging (7.4) and (7.6) to (7.3) then yields

$$\limsup_{N_0 \to \infty} \frac{N_0}{M} \mathbb{E} \left[\nu_1 \left(A_M(x) \right) \right] \le \frac{1 + \epsilon}{1 - \epsilon} \frac{f_1(x)}{f_0(x)}.$$
(7.7)

Lower bound. We have, from (7.1) and (7.2),

$$E\left[\nu_{1}(A_{M}(x))\right] = P\left(W \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|})\right)$$

$$\geq P\left(W \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \le \eta_{N}\frac{M}{N_{0}}\right)$$

$$= P\left(W \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \le \eta_{N}\frac{M}{N_{0}}, \|Z-x\| \le \delta\right)$$

$$\geq P\left(\frac{1+\epsilon}{1-\epsilon}\frac{f_{0}(x)}{f_{1}(x)}\nu_{1}(B_{x,\|x-Z\|}) \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \le \eta_{N}\frac{M}{N_{0}}, \|Z-x\| \le \delta\right)$$

$$= P\left(\frac{1+\epsilon}{1-\epsilon}\frac{f_{0}(x)}{f_{1}(x)}\nu_{1}(B_{x,\|x-Z\|}) \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \le \eta_{N}\frac{M}{N_{0}}\right)$$

$$\geq P\left(\frac{1+\epsilon}{1-\epsilon}\frac{f_{0}(x)}{f_{1}(x)}\nu_{1}(B_{x,\|x-Z\|}) \le \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|})\right) - P\left(\nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) > \eta_{N}\frac{M}{N_{0}}\right)$$

$$= P\left(\frac{1+\epsilon}{1-\epsilon}\frac{f_{0}(x)}{f_{1}(x)}U \le U_{(M)}\right) - P\left(U_{(M)} > \eta_{N}\frac{M}{N_{0}}\right).$$
(7.8)

The second last equality is from the fact that for $||Z - x|| > \delta$,

$$\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} \nu_1(B_{x,\|x-Z\|}) \ge \frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} \nu_1(B_{x,\delta})$$
$$\ge \frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} f_1(x)(1-\epsilon)\lambda(B_{0,\delta}) = (1+\epsilon)f_0(x)\lambda(B_{0,\delta}) > \eta_N \frac{M}{N_0},$$

and then that $\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} \nu_1(B_{x,\|x-Z\|}) \leq \eta_N \frac{M}{N_0}$ implies $\|Z - x\| \leq \delta$. For the first term in (7.8), we have

$$\frac{N_0}{M} \mathbf{P}\Big(\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} U \le U_{(M)}\Big) = \frac{N_0}{M} \int_0^1 \mathbf{P}\Big(U_{(M)} \ge \frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} t\Big) \mathrm{d}t$$
$$= \frac{1-\epsilon}{1+\epsilon} \frac{f_1(x)}{f_0(x)} \int_0^{\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} \frac{N_0}{M}} \mathbf{P}\Big(U_{(M)} \ge \frac{M}{N_0} t\Big) \mathrm{d}t.$$

If $\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} \ge 1$, since $U_{(M)} \in [0,1]$, we have

$$\frac{N_0}{M} \mathbf{P}\Big(\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} U \le U_{(M)}\Big) = \frac{1-\epsilon}{1+\epsilon} \frac{f_1(x)}{f_0(x)} \frac{N_0}{M} \mathbf{E}[U_{(M)}] = \frac{1-\epsilon}{1+\epsilon} \frac{f_1(x)}{f_0(x)} \frac{N_0}{N_0+1}.$$

Then

$$\lim_{N_0 \to \infty} \frac{N_0}{M} \mathbb{P}\left(\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} U \le U_{(M)}\right) = \frac{1-\epsilon}{1+\epsilon} \frac{f_1(x)}{f_0(x)}.$$

If $\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} < 1$, from the Chernoff bound,

$$\begin{split} &\int_{\frac{1+\epsilon}{1-\epsilon}\frac{f_0(x)}{f_1(x)}\frac{N_0}{M}}^{\frac{N_0}{M}} \mathbf{P}\Big(U_{(M)} \ge \frac{M}{N_0}t\Big) \mathrm{d}t \\ \le &\Big[1 - \frac{1+\epsilon}{1-\epsilon}\frac{f_0(x)}{f_1(x)}\Big]\frac{N_0}{M}\mathbf{P}\Big(U_{(M)} \ge \frac{1+\epsilon}{1-\epsilon}\frac{f_0(x)}{f_1(x)}\Big) \\ \le &\Big[1 - \frac{1+\epsilon}{1-\epsilon}\frac{f_0(x)}{f_1(x)}\Big]\frac{N_0}{M}\exp\Big[M - \frac{1+\epsilon}{1-\epsilon}\frac{f_0(x)}{f_1(x)}N_0 - M\log M + M\log\Big(\frac{1+\epsilon}{1-\epsilon}\frac{f_0(x)}{f_1(x)}N_0\Big)\Big]. \end{split}$$

Since $f_0(x) > 0$ and $M \log N_0/N_0 \to 0$, we obtain

$$\lim_{N_0 \to \infty} \int_{\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} \frac{N_0}{M}}^{\frac{N_0}{M}} \mathcal{P}\left(U_{(M)} \ge \frac{M}{N_0} t\right) \mathrm{d}t = 0.$$

Then

$$\lim_{N_0 \to \infty} \frac{N_0}{M} \mathbb{P}\left(\frac{1+\epsilon}{1-\epsilon} \frac{f_0(x)}{f_1(x)} U \le U_{(M)}\right) = \frac{1-\epsilon}{1+\epsilon} \frac{f_1(x)}{f_0(x)}.$$

Plugging the above argument along with (7.4) to (7.8) yields

$$\liminf_{N_0 \to \infty} \frac{N_0}{M} \mathbb{E} \left[\nu_1 \left(A_M(x) \right) \right] \ge \frac{1 - \epsilon}{1 + \epsilon} \frac{f_1(x)}{f_0(x)}.$$
(7.9)

Lastly, combining (7.7) with (7.9) and noticing that ϵ is arbitrary, we obtain

$$\lim_{N_0 \to \infty} \frac{N_0}{M} \mathbb{E} \left[\nu_1 \left(A_M(x) \right) \right] = \frac{f_1(x)}{f_0(x)} = r(x).$$
(7.10)

Case I.2. $f_1(x) = 0$. Again, for any $\epsilon \in (0, 1)$, by Lemma 7.1, there exists some $\delta = \delta_x > 0$ such that for any $z \in \mathbb{R}^d$ with $||z - x|| \leq \delta$, we have

$$\left|\frac{\nu_0(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_0(x)\right| \le \epsilon f_0(x), \quad \left|\frac{\nu_1(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})}\right| \le \epsilon$$

Recall that $W = \nu_0(B_{Z,||x-Z||})$. Then if $||Z - x|| \le \delta$, we have

$$Z \ge (1-\epsilon)f_0(x)\lambda(B_{Z,\|x-Z\|}) = (1-\epsilon)f_0(x)\lambda(B_{x,\|x-Z\|}) \ge \frac{1-\epsilon}{\epsilon}f_0(x)\nu_1(B_{x,\|x-Z\|}).$$

Proceeding in the same way as (7.3), we obtain

$$E[\nu_1(A_M(x))]$$

$$\leq P\Big(W \leq \nu_0(B_{Z,\|\mathcal{X}_{(M)}(Z) - Z\|}), \nu_0(B_{Z,\|\mathcal{X}_{(M)}(Z) - Z\|}) \leq \eta_N \frac{M}{N_0}, \|Z - x\| \leq \delta\Big) + P\Big(U_{(M)} > \eta_N \frac{M}{N_0}\Big)$$

$$\leq P\Big(\frac{1 - \epsilon}{\epsilon} f_0(x)U \leq U_{(M)}\Big) + P\Big(U_{(M)} > \eta_N \frac{M}{N_0}\Big).$$

For the first term above,

$$\frac{N_0}{M} \operatorname{P}\left(\frac{1-\epsilon}{\epsilon} f_0(x)U \le U_{(M)}\right) = \frac{N_0}{M} \int_0^1 \operatorname{P}\left(U_{(M)} \ge \frac{1-\epsilon}{\epsilon} f_0(x)t\right) \mathrm{d}t$$
$$= \frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)} \int_0^{\frac{1-\epsilon}{\epsilon} f_0(x)\frac{N_0}{M}} \operatorname{P}\left(U_{(M)} \ge \frac{M}{N_0}t\right) \mathrm{d}t \le \frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)} \int_0^\infty \operatorname{P}\left(\frac{N_0}{M}U_{(M)} \ge t\right) \mathrm{d}t$$
$$= \frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)} \frac{N_0}{M} \operatorname{E}[U_{(M)}] = \frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)} \frac{N_0}{N_0+1}.$$

By (7.4) and noticing ϵ is arbitrary, one has

$$\lim_{N_0 \to \infty} \frac{N_0}{M} \mathbb{E} \left[\nu_1 (A_M(x)) \right] = 0 = r(x).$$
(7.11)

Combining (7.10) and (7.11) completes the proof of the first claim.

Part II. This part proves the second claim. We also separate the proof of Part II into two cases

based on the value of $f_1(x)$.

Case II.1. $f_1(x) > 0$. Again, for any $\epsilon \in (0, 1)$, we take δ in the same way as in Case I.1. Let $\eta_N = \eta_{N,p} = 4p \log(N_0/M)$. We also take N_0 sufficiently large so that

$$\eta_N \frac{M}{N_0} = 4p \frac{M}{N_0} \log\left(\frac{N_0}{M}\right) < (1-\epsilon) f_0(x) \lambda(B_{0,\delta}).$$

Let $\widetilde{Z}_1, \ldots, \widetilde{Z}_p$ be p independent copies that are drawn from ν_1 independent of the data. Then

$$E[\nu_{1}^{p}(A_{M}(x))] = P(\tilde{Z}_{1}, \dots, \tilde{Z}_{p} \in A_{M}(x))$$

=P($||x - \tilde{Z}_{1}|| \le ||\mathcal{X}_{(M)}(\tilde{Z}_{1}) - \tilde{Z}_{1}||, \dots, ||x - \tilde{Z}_{p}|| \le ||\mathcal{X}_{(M)}(\tilde{Z}_{p}) - \tilde{Z}_{p}||)$
=P($\nu_{0}(B_{\tilde{Z}_{1},||x - \tilde{Z}_{1}||) \le \nu_{0}(B_{\tilde{Z}_{1},||\mathcal{X}_{(M)}(\tilde{Z}_{1}) - \tilde{Z}_{1}||), \dots, \nu_{0}(B_{\tilde{Z}_{p},||x - \tilde{Z}_{p}||) \le \nu_{0}(B_{\tilde{Z}_{p},||\mathcal{X}_{(M)}(\tilde{Z}_{p}) - \tilde{Z}_{p}||)).$

Let $W_k = \nu_0(B_{\widetilde{Z}_k, \|x - \widetilde{Z}_k\|})$ and $V_k = \nu_0(B_{\widetilde{Z}_k, \|\mathcal{X}_{(M)}(\widetilde{Z}_k) - \widetilde{Z}_k\|})$ for any $k \in [\![p]\!]$. Then $[W_k]_{k=1}^p$ are i.i.d. since $[\widetilde{Z}_k]_{k=1}^p$ are i.i.d. For any $k \in [\![p]\!]$ and $\widetilde{Z}_k \in \mathbb{R}^d$ given, $V_k | \widetilde{Z}_k$ has the same distribution as $U_{(M)}$. Then for any $k \in [\![p]\!]$, V_k has the same distribution as $U_{(M)}$, and V_k is independent of \widetilde{Z}_k .

Let $W_{\max} = \max_{k \in \llbracket p \rrbracket} W_k$ and $V_{\max} = \max_{k \in \llbracket p \rrbracket} V_k$. Then

$$E[\nu_1^p(A_M(x))] = P\left(W_1 \le V_1, \dots, W_p \le V_p\right) \le P\left(W_{\max} \le V_{\max}\right)$$
$$\le P\left(W_{\max} \le V_{\max}, V_{\max} \le \eta_N \frac{M}{N_0}\right) + P\left(V_{\max} > \eta_N \frac{M}{N_0}\right)$$
(7.12)

For the second term in (7.12),

$$\mathbf{P}\left(V_{\max} > \eta_N \frac{M}{N_0}\right) \le \sum_{k=1}^p \mathbf{P}\left(V_k > \eta_N \frac{M}{N_0}\right) = p\mathbf{P}\left(U_{(M)} > \eta_N \frac{M}{N_0}\right).$$

Proceeding as (7.4),

$$\left(\frac{N_0}{M}\right)^p \mathcal{P}\left(U_{(M)} > \eta_N \frac{M}{N_0}\right) \le \left(\frac{N_0}{M}\right)^p \exp\left(-\frac{1}{2}\eta_N M\right) = \left(\frac{N_0}{M}\right)^{p(1-2M)}$$

We then obtain

$$\lim_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathcal{P}\left(V_{\max} > \eta_N \frac{M}{N_0}\right) = 0.$$
(7.13)

For the first term in (7.12), notice that $[\nu_1(B_{x,\|\widetilde{Z}_k-x\|})]_{k=1}^p$ are i.i.d. from U(0,1) since $[\widetilde{Z}_k]_{k=1}^p$ are i.i.d. We then have

$$\left(\frac{N_0}{M}\right)^p \mathbf{P}\left(W_{\max} \leq V_{\max}, V_{\max} \leq \eta_N \frac{M}{N_0}\right)$$

$$= \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(W_{\max} \leq V_{\max}, V_{\max} \leq \eta_N \frac{M}{N_0}, \max_{k \in \llbracket p \rrbracket} \|\widetilde{Z}_k - x\| \leq \delta\right)$$

$$\leq \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\widetilde{Z}_k - x\|}) \leq V_{\max}, V_{\max} \leq \eta_N \frac{M}{N_0}, \max_{k \in \llbracket p \rrbracket} \|\widetilde{Z}_k - x\| \leq \delta\right)$$

$$\leq \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)} \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\widetilde{Z}_k - x\|}) \leq V_{\max}\right)$$

$$= \left(\frac{N_0}{M}\right)^p \int_0^1 pt^{p-1} \mathbf{P}\left(V_{\max} \geq \frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{f_1(x)}t \mid \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\widetilde{Z}_k - x\|}) = t\right) \mathrm{d}t$$

$$\begin{split} &= p\Big(\frac{1+\epsilon}{1-\epsilon}\frac{f_1(x)}{f_0(x)}\Big)^p \int_0^{\frac{1-\epsilon}{1+\epsilon}\frac{f_0(x)}{f_1(x)}\frac{N_0}{M}} t^{p-1}\mathbf{P}\Big(V_{\max} \ge \frac{M}{N_0}t\Big| \max_{k\in \llbracket p \rrbracket} \nu_1(B_{x,\Vert \tilde{Z}_k - x\Vert}) = \frac{1+\epsilon}{1-\epsilon}\frac{f_1(x)}{f_0(x)}\frac{M}{N_0}t\Big) \mathrm{d}t \\ &= p\Big(\frac{1+\epsilon}{1-\epsilon}\frac{f_1(x)}{f_0(x)}\Big)^p \Big[\int_0^1 t^{p-1}\mathbf{P}\Big(V_{\max} \ge \frac{M}{N_0}t\Big| \max_{k\in \llbracket p \rrbracket} \nu_1(B_{x,\Vert \tilde{Z}_k - x\Vert}) = \frac{1+\epsilon}{1-\epsilon}\frac{f_1(x)}{f_0(x)}\frac{M}{N_0}t\Big) \mathrm{d}t \\ &+ \int_1^{\frac{1-\epsilon}{1+\epsilon}\frac{f_0(x)}{f_1(x)}\frac{N_0}{M}} t^{p-1}\mathbf{P}\Big(V_{\max} \ge \frac{M}{N_0}t\Big| \max_{k\in \llbracket p \rrbracket} \nu_1(B_{x,\Vert \tilde{Z}_k - x\Vert}) = \frac{1+\epsilon}{1-\epsilon}\frac{f_1(x)}{f_0(x)}\frac{M}{N_0}t\Big) \mathrm{d}t \Big]. \end{split}$$

For the first term,

$$\int_0^1 t^{p-1} \mathbf{P}\Big(V_{\max} \ge \frac{M}{N_0} t \Big| \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \Vert \widetilde{Z}_k - x \Vert}) = \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \frac{M}{N_0} t\Big) \mathrm{d}t \le \int_0^1 t^{p-1} \mathrm{d}t = \frac{1}{p}.$$

For the second term, using the Chernoff bound, conditional on $\widetilde{Z} = (\widetilde{Z}_1, \ldots, \widetilde{Z}_p)$,

$$\begin{split} &\int_{1}^{\frac{1-\epsilon}{1+\epsilon}\frac{f_{0}(x)}{M}} t^{p-1} \mathbf{P}\Big(V_{\max} \ge \frac{M}{N_{0}}t \,\Big|\, \widetilde{\mathbf{Z}}\Big) \mathrm{d}t \\ &\le \int_{1}^{\infty} t^{p-1} \mathbf{P}\Big(V_{\max} \ge \frac{M}{N_{0}}t \,\Big|\, \widetilde{\mathbf{Z}}\Big) \mathrm{d}t = \int_{0}^{\infty} (1+t)^{p-1} \mathbf{P}\Big(V_{\max} \ge \frac{M}{N_{0}}(1+t) \,\Big|\, \widetilde{\mathbf{Z}}\Big) \mathrm{d}t \\ &\le \int_{0}^{\infty} (1+t)^{p-1} \Big[\sum_{k=1}^{p} \mathbf{P}\Big(V_{k} \ge \frac{M}{N_{0}}(1+t) \,\Big|\, \widetilde{\mathbf{Z}}\Big)\Big] \mathrm{d}t = p \int_{0}^{\infty} (1+t)^{p-1} \mathbf{P}\Big(U_{(M)} \ge \frac{M}{N_{0}}(1+t)\Big) \mathrm{d}t \\ &\le p \int_{0}^{\infty} (1+t)^{p-1} (1+t)^{M} \exp(-tM) \mathrm{d}t = p e^{M} \int_{1}^{\infty} t^{M+p-1} \exp(-tM) \mathrm{d}t \\ &\le p e^{M} \int_{0}^{\infty} t^{M+p-1} \exp(-tM) \mathrm{d}t = \frac{p e^{M}}{M^{M+p}} \Gamma(M+p) = \frac{p e^{M}}{M^{M+p}} (M+1)^{p-1} \Gamma(M+1)(1+o(1)) \\ &= \frac{p e^{M}}{M^{M+p}} (M+1)^{p-1} \sqrt{2\pi M} \frac{M^{M}}{e^{M}} (1+o(1)) = \sqrt{2\pi} p M^{-1/2} \Big(1+\frac{1}{M}\Big)^{p-1} (1+o(1)), \end{split}$$

where the last three steps are from Stirling's approximation using $M \to \infty$.

Then

$$\lim_{N_0 \to \infty} p \Big(\frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \Big)^p \int_1^{\frac{1-\epsilon}{1+\epsilon} \frac{f_0(x)}{N_1} \frac{N_0}{M}} t^{p-1} \mathbf{P} \Big(V_{\max} \ge \frac{M}{N_0} t \Big| \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\widetilde{Z}_k - x\|}) = \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)} \frac{M}{N_0} t \Big) \mathrm{d}t = 0,$$

and then we obtain

$$\limsup_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbb{P}\left(W_{\max} \le V_{\max}, V_{\max} \le \eta_N \frac{M}{N_0}\right) \le \left(\frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)}\right)^p.$$
(7.14)

Plugging (7.13) and (7.14) into (7.12) yields

$$\limsup_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p(A_M(x))\right] \le \left(\frac{1+\epsilon}{1-\epsilon}\frac{f_1(x)}{f_0(x)}\right)^p = \left(\frac{1+\epsilon}{1-\epsilon}r(x)\right)^p.$$
(7.15)

Lastly, using Hölder's inequality,

$$\left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p(A_M(x))\right] \ge \left[\frac{N_0}{M}\mathbf{E}\left[\nu_1(A_M(x))\right]\right]^p.$$

Employing the first claim, we have

$$\liminf_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p(A_M(x))\right] \ge \left[r(x)\right]^p.$$
(7.16)

Combining (7.15) with (7.16) and noting that ϵ is arbitrary, we obtain

$$\lim_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbb{E}\left[\nu_1^p \left(A_M(x)\right)\right] = \left[r(x)\right]^p.$$
(7.17)

Case II.2. $f_1(x) = 0$. For any $\epsilon \in (0, 1)$, we take δ in the same way as in the proof of Case I.2 and take η_N as in the proof of Case II.1.

By (7.12),

$$\left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p(A_M(x))\right] \le \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(W_{\max} \le V_{\max}, V_{\max} \le \eta_N \frac{M}{N_0}\right) + \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(V_{\max} > \eta_N \frac{M}{N_0}\right).$$

For the first term,

For the first term,

$$\begin{split} & \left(\frac{N_0}{M}\right)^p \mathbf{P}\Big(W_{\max} \leq V_{\max}, V_{\max} \leq \eta_N \frac{M}{N_0}\Big) \\ &= \left(\frac{N_0}{M}\right)^p \mathbf{P}\Big(W_{\max} \leq V_{\max}, V_{\max} \leq \eta_N \frac{M}{N_0}, \max_{k \in \llbracket p \rrbracket} \|Z_k - x\| \leq \delta\Big) \\ &\leq \left(\frac{N_0}{M}\right)^p \mathbf{P}\Big(\frac{1-\epsilon}{\epsilon} f_0(x) \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\tilde{Z}_k - x\|}) \leq V_{\max}, V_{\max} \leq \eta_N \frac{M}{N_0}, \max_{k \in \llbracket p \rrbracket} \|Z_k - x\| \leq \delta\Big) \\ &\leq \left(\frac{N_0}{M}\right)^p \mathbf{P}\Big(\frac{1-\epsilon}{\epsilon} f_0(x) \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\tilde{Z}_k - x\|}) \leq V_{\max}\Big) \\ &= \left(\frac{N_0}{M}\right)^p \int_0^1 pt^{p-1} \mathbf{P}\Big(V_{\max} \geq \frac{1-\epsilon}{\epsilon} f_0(x)t\Big| \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\tilde{Z}_k - x\|}) = t\Big) \mathrm{d}t \\ &= p\Big(\frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)}\Big)^p \int_0^{\frac{1-\epsilon}{\epsilon} f_0(x)\frac{N_0}{M}} t^{p-1} \mathbf{P}\Big(V_{\max} \geq \frac{M}{N_0}t\Big| \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\tilde{Z}_k - x\|}) = t\Big) \mathrm{d}t \\ &= p\Big(\frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)}\Big)^p \Big[\int_0^1 t^{p-1} \mathbf{P}\Big(V_{\max} \geq \frac{M}{N_0}t\Big| \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\tilde{Z}_k - x\|}) = t\Big) \mathrm{d}t \\ &+ \int_1^{\frac{1-\epsilon}{\epsilon} f_0(x)\frac{N_0}{M}} t^{p-1} \mathbf{P}\Big(V_{\max} \geq \frac{M}{N_0}t\Big| \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \|\tilde{Z}_k - x\|}) = t\Big) \mathrm{d}t\Big]. \end{split}$$

Then proceeding in the same way as (7.14), we have

$$\limsup_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbb{P}\left(W_{\max} \le V_{\max}, V_{\max} \le \eta_N \frac{M}{N_0}\right) \le \left(\frac{\epsilon}{1-\epsilon} \frac{1}{f_0(x)}\right)^p.$$

Lastly, using (7.13) and noting again that ϵ is arbitrary, one obtains

$$\lim_{N_0 \to \infty} \left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p(A_M(x))\right] = 0 = [r(x)]^p.$$
(7.18)
7.18) then completes the proof of the second claim.

Combining (7.17) and (7.18) then completes the proof of the second claim.

7.2 Proof of Theorem 4.1

Proof of Theorem 4.1(i). By (2.4) and that $[Z_j]_{j=1}^{N_1}$ are i.i.d,

$$\mathbf{E}[\hat{r}_{M}(x)] = \mathbf{E}\Big[\frac{N_{0}}{N_{1}}\frac{K_{M}(x)}{M}\Big] = \frac{N_{0}}{N_{1}M}\mathbf{E}\Big[\sum_{j=1}^{N_{1}}\mathbb{1}(Z_{j} \in A_{M}(x))\Big] = \frac{N_{0}}{M}\mathbf{E}\big[\nu_{1}(A_{M}(x))\big].$$

Employing Lemma 4.1 then completes the proof.

Proof of Theorem 4.1(ii). By Hölder's inequality, it suffices to consider the case when p is even. Noticing that x^p is convex for p > 1 and x > 0, one has

$$\mathbf{E}\left[\left|\widehat{r}_{M}(x) - r(x)\right|^{p}\right] \leq 2^{p-1} \left(\mathbf{E}\left[\left|\widehat{r}_{M}(x) - \mathbf{E}[\widehat{r}_{M}(x) \mid \boldsymbol{X}]\right|^{p}\right] + \mathbf{E}\left[\left|\mathbf{E}[\widehat{r}_{M}(x) \mid \boldsymbol{X}] - r(x)\right|^{p}\right]\right).$$
(7.19)

For the second term in (7.19), Lemma 4.1 implies

$$\lim_{N_0 \to \infty} \mathbb{E}\left[\left|\mathbb{E}[\hat{r}_M(x) \mid \boldsymbol{X}] - r(x)\right|^p\right] = \lim_{N_0 \to \infty} \mathbb{E}\left[\left|\frac{N_0}{M}\nu_1(A_M(x)) - r(x)\right|^p\right] = 0$$
(7.20)

by expanding the product term.

For the first term in (7.19), noticing that $[Z_j]_{j=1}^{N_1}$ are i.i.d, we have $K_M(x) | \mathbf{X} \sim Bin(N_1, \nu_1(A_M(x)))$. Using Lemma 4.1 and $MN_1/N_0 \to \infty$, for any positive integers p and q, we have

$$\lim_{N_0 \to \infty} \left(\frac{N_0}{N_1 M}\right)^p \mathbb{E}[N_1^p \nu_1^p(A_M(x))] = [r(x)]^p,$$
$$\lim_{N_0 \to \infty} \left(\frac{N_0}{N_1 M}\right)^p \left(\frac{N_0}{M}\right)^q \mathbb{E}[N_1^p \nu_1^{p+q}(A_M(x))] = [r(x)]^p,$$

and then $\mathbb{E}[N_1^p \nu_1^p(A_M(x))]$ is the dominated term among $[\mathbb{E}[N_1^k \nu_1^{k+q}(A_M(x))]]_{k \le p,q \ge 0}$.

To complete the proof, for any positive integer c and $Z \sim \text{Bin}(n, p')$, let $\mu_c := \text{E}[(Z - \text{E}[Z])^c]$ be the *c*-th central moment. According to Romanovsky (1923), we have

$$\mu_{c+1} = p'(1-p') \left(nc\mu_{c-1} + \frac{\mathrm{d}\mu_c}{\mathrm{d}p'} \right).$$

Then for even p, we obtain

$$\mathbb{E}\Big[\Big(K_M(x) - N_1\nu_1(A_M(x))\Big)^p\Big] \lesssim \mathbb{E}[N_1\nu_1(A_M(x))]^{p/2} \lesssim \Big(\frac{N_1M}{N_0}\Big)^{p/2}.$$

The first term in (7.19) then satisfies

$$\mathbf{E}\Big[\Big|\widehat{r}_M(x) - \mathbf{E}[\widehat{r}_M(x) \mid \boldsymbol{X}]\Big|^p\Big] = \Big(\frac{N_0}{N_1M}\Big)^p \mathbf{E}\Big[\Big(K_M(x) - N_1\nu_1(A_M(x))\Big)^p\Big] \lesssim \Big(\frac{N_0}{N_1M}\Big)^{p/2}.$$

Since $MN_1/N_0 \to \infty$, we obtain

$$\lim_{N_0 \to \infty} \mathbf{E} \left[\left| \widehat{r}_M(x) - \mathbf{E} [\widehat{r}_M(x) \,|\, \boldsymbol{X}] \right|^p \right] = 0.$$
(7.21)

Plugging (7.20) and (7.21) into (7.19) then completes the proof.

7.3 Proof of Theorem 4.2

Proof of Theorem 4.2. We first cite the Hardy–Littlewood maximal inequality.

Lemma 7.2 (Hardy–Littlewood maximal inequality (Stein, 2016)). For any locally integrable function $f : \mathbb{R}^d \to \mathbb{R}$, define

$$\mathsf{M}f(x) := \sup_{\delta > 0} \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f(z)| \mathrm{d}z.$$

Then for $d \ge 1$, there exists a constant $C_d > 0$ only depending on d such that for all t > 0 and $f \in L_1(\mathbb{R}^d)$, we have

$$\lambda(\{x: \mathsf{M}f(x) > t\}) < \frac{C_d}{t} \|f\|_{L_1},$$

where $\|\cdot\|_{L_1}$ stands for the function L_1 norm.

Let $\epsilon > 0$ be given. We assume $\epsilon \leq f_L$. From Assumption 4.1, S_0 and S_1 are bounded, then ν_0 and ν_1 are compactly supported. Since $f_0, f_1 \in L_1$ and the class of continuous functions are dense in the class of compactly supported L_1 functions from simple use of Lusin's theorem, we can find g_0, g_1 such that g_0, g_1 are continuous and $||f_0 - g_0||_{L_1} \leq \epsilon^3$ and $||f_1 - g_1||_{L_1} \leq \epsilon^3$.

Since g_0, g_1 are continuous with compact supports, they are uniformly continuous, that is, there exists $\delta > 0$ such that for any $x, z \in \mathbb{R}^d$ and $||z - x|| \leq \delta$, we have

$$|g_0(x) - g_0(z)| \le \frac{\epsilon^2}{3}$$
 and $|g_1(x) - g_1(z)| \le \frac{\epsilon^2}{3}$

For any $x \in \mathbb{R}^d$, we have

$$\frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(x) - f_0(z)| dz
\leq \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(x) - g_0(x)| dz + \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |g_0(x) - g_0(z)| dz + \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(z) - g_0(z)| dz
= |f_0(x) - g_0(x)| + \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |g_0(x) - g_0(z)| dz + \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(z) - g_0(z)| dz.$$
(7.22)

For the first term in (7.22), using Markov's inequality, one has

$$\lambda\Big(\Big\{x: \Big|f_0(x) - g_0(x)\Big| > \frac{\epsilon^2}{3}\Big\}\Big) \le \frac{3}{\epsilon^2} ||f_0 - g_0||_{L_1} \le 3\epsilon.$$
(7.23)

For the second term in (7.22), by the selection of δ ,

$$\frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |g_0(x) - g_0(z)| dz \le \max_{z \in B_{x,\delta}} |g_0(x) - g_0(z)| \le \frac{\epsilon^2}{3}.$$
(7.24)

For the third term,

$$\frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(z) - g_0(z)| \mathrm{d}z \le \sup_{\delta > 0} \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(z) - g_0(z)| \mathrm{d}z = \mathsf{M}(f_0 - g_0)(x).$$

Lemma 7.2 then yields

$$\lambda\Big(\Big\{x: \mathsf{M}(f_0 - g_0)(x) > \frac{\epsilon^2}{3}\Big\}\Big) < \frac{3C_d}{\epsilon^2} \|f_0 - g_0\|_{L_1} \le 3C_d\epsilon.$$
(7.25)

We can establish similar results for f_1, g_1 .

Let

$$A_{1} := \left\{ x : \left| f_{0}(x) - g_{0}(x) \right| > \frac{\epsilon^{2}}{3} \right\} \bigcup \left\{ x : \left| f_{1}(x) - g_{1}(x) \right| > \frac{\epsilon^{2}}{3} \right\} \bigcup \left\{ x : \mathsf{M}(f_{0} - g_{0})(x) > \frac{\epsilon^{2}}{3} \right\} \bigcup \left\{ x : \mathsf{M}(f_{1} - g_{1})(x) > \frac{\epsilon^{2}}{3} \right\}.$$

Plugging (7.23), (7.24), (7.25) into (7.22), for any $x \notin A_1$ and $||z - x|| \le \delta$, we have

$$\frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(x) - f_0(z)| \mathrm{d}z \le \epsilon^2, \quad \frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_1(x) - f_1(z)| \mathrm{d}z \le \epsilon^2,$$

and

$$\lambda(A_1) \le 6(C_d + 1)\epsilon.$$

Let $A_2 := \{x : f_1(x) \le \epsilon\}$. We then separate the proof into three cases. In the following, it suffices to consider $f_0(x) > 0$ due to the definition of L_p risk in (4.1).

Case I. $x \notin A_1 \cup A_2$.

According to $\epsilon \leq f_L$ and by the definition of A_2 , for any $x \notin A_1 \cup A_2$ and $||z - x|| \leq \delta$,

$$\frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_0(x) - f_0(y)| dz \le \epsilon^2 \le \epsilon f_L \le \epsilon f_0(x),$$
$$\frac{1}{\lambda(B_{x,\delta})} \int_{B_{x,\delta}} |f_1(x) - f_1(y)| dz \le \epsilon^2 \le \epsilon f_1(x).$$

We then obtain

$$\begin{aligned} \left| \frac{\nu_0(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_0(x) \right| &\leq \epsilon f_0(x), \quad \left| \frac{\nu_0(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_0(x) \right| \leq \epsilon f_0(x), \\ \left| \frac{\nu_1(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_1(x) \right| &\leq \epsilon f_1(x), \quad \left| \frac{\nu_1(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_1(x) \right| \leq \epsilon f_1(x). \end{aligned}$$

Let $\eta_N = \eta_{N,p} = 4p \log(N_0/M)$. We also take N_0 large enough so that

$$\eta_N \frac{M}{N_0} = 4p \frac{M}{N_0} \log\left(\frac{N_0}{M}\right) < (1 - \epsilon) f_L \lambda(B_{0,\delta}).$$

Then for any $x \in \mathbb{R}^d$ such that $f_0(x) > 0$, we have

$$\eta_N \frac{M}{N_0} < (1-\epsilon) f_0(x) \lambda(B_{0,\delta})$$

Proceeding as in the proof of Case II.1 of Lemma 4.1 and also Theorem 4.1 by using Fubini's theorem, since ϵ is arbitrary, we obtain

$$\lim_{N_0 \to \infty} \mathbb{E} \left[\int_{\mathbb{R}^d} \left| \widehat{r}_M(x) - r(x) \right|^p f_0(x) \mathbb{1}(x \notin A_1 \cup A_2) \mathrm{d}x \right] = 0.$$
(7.26)

Case II. $x \in A_2 \setminus A_1$.

In this case, we have

$$\frac{\nu_0(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_0(x) \Big| \le \epsilon f_0(x), \quad \Big| \frac{\nu_0(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_0(x) \Big| \le \epsilon f_0(x),$$
$$\frac{\nu_1(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_1(x) \Big| \le \epsilon^2, \quad \Big| \frac{\nu_1(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_1(x) \Big| \le \epsilon^2.$$

Take η_N and take N_0 sufficiently large as in Case I above. Proceeding as the proof of Case II.2 of Lemma 4.1 and also Theorem 4.1 by using Fubini's theorem, since ϵ is arbitrary, we obtain

$$\lim_{N_0 \to \infty} \mathbb{E} \Big[\int_{\mathbb{R}^d} \left| \widehat{r}_M(x) - r(x) \right|^p f_0(x) \mathbb{1}(x \in A_2 \setminus A_1) \mathrm{d}x \Big] = 0.$$
(7.27)

Case III. $x \in A_1$.

In this case, for any $x \in A_1$ and $z \in S_1$,

$$\nu_0(B_{z,\|z-x\|}) \ge f_L \lambda(B_{z,\|z-x\|} \cap S_0) \ge a f_L \lambda(B_{z,\|z-x\|}) \ge \frac{a f_L}{f_U} \nu_1(B_{x,\|z-x\|}).$$

Then for any $x \in A_1$, from (7.12) and in the same way as (7.14),

$$\left(\frac{N_0}{M}\right)^p \mathbf{E}\left[\nu_1^p \left(A_M(x)\right)\right] = \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(W_1 \le V_1, \dots, W_p \le V_p\right) \le \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(W_{\max} \le V_{\max}\right)$$
$$\le \left(\frac{N_0}{M}\right)^p \mathbf{P}\left(\frac{af_L}{f_U} \max_{k \in \llbracket p \rrbracket} \nu_1(B_{x, \lVert \widetilde{Z}_k - x \rVert}) \le V_{\max}\right) \le \left(\frac{f_U}{af_L}\right)^p (1 + o(1)) = O(1).$$

Proceeding as in the proof of Theorem 4.1, and due to the boundedness assumptions on f_0 and f_1 , for any $x \in A_1$ and p even,

$$\mathbf{E}\left[\left|\widehat{r}_{M}(x)-r(x)\right|^{p}\right] \lesssim \mathbf{E}\left[\left|\widehat{r}_{M}(x)-\mathbf{E}[\widehat{r}_{M}(x) \mid \boldsymbol{X}]\right|^{p}\right] + \mathbf{E}\left[\left(\mathbf{E}[\widehat{r}_{M}(x) \mid \boldsymbol{X}]\right)^{p}\right] + \left|r(x)\right|^{p} \lesssim 1.$$

Then

$$\mathbb{E}\Big[\int_{\mathbb{R}^d} \left| \widehat{r}_M(x) - r(x) \right|^p f_0(x) \mathbb{1}(x \in A_1) \mathrm{d}x \Big] \lesssim f_U \lambda(A_1) \lesssim \epsilon.$$

Since ϵ is arbitrary, one has

$$\lim_{N_0 \to \infty} \mathbb{E} \left[\int_{\mathbb{R}^d} \left| \widehat{r}_M(x) - r(x) \right|^p f_0(x) \mathbb{1}(x \in A_1) \mathrm{d}x \right] = 0.$$
(7.28)
(7.27), and (7.28) completes the proof.

Combining (7.26), (7.27), and (7.28) completes the proof.

7.4 Proof of Theorem 4.3

We only have to prove the first two claims as the rest are trivial.

Proof of Theorem 4.3(i). For any $z \in \mathbb{R}^d$ such that $||z-x|| \leq \delta/2$, since $B_{z,||z-x||} \subset B_{x,2||z-x||} \subset B_{x,\delta}$, we have

$$\begin{aligned} \left| \frac{\nu_0(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f_0(x) \right| &\leq \frac{1}{\lambda(B_{z,\|z-x\|})} \int_{B_{z,\|z-x\|}} |f_0(y) - f_0(x)| \mathrm{d}y \leq 2L \|z - x\|, \\ \left| \frac{\nu_1(B_{x,\|z-x\|})}{\lambda(B_{x,\|z-x\|})} - f_1(x) \right| &\leq \frac{1}{\lambda(B_{x,\|z-x\|})} \int_{B_{x,\|z-x\|}} |f_1(z) - f_1(x)| \mathrm{d}z \leq L \|z - x\|. \end{aligned}$$

Consider any $\delta_N > 0$ such that $\delta_N \leq \delta/2$. If $||z - x|| \leq \delta_N$ and $f_0(x) > 2L\delta_N$, then

$$\frac{f_0(x) - 2L\delta_N}{f_1(x) + L\delta_N} \le \frac{\nu_0(B_{z,\|x-z\|})}{\lambda(B_{z,\|x-z\|})} \frac{\lambda(B_{x,\|x-z\|})}{\nu_1(B_{x,\|x-z\|})}$$

If further $f_1(x) > L\delta_N$, then

$$\frac{\nu_0(B_{z,\|x-z\|})}{\lambda(B_{z,\|x-z\|})} \frac{\lambda(B_{x,\|x-z\|})}{\nu_1(B_{x,\|x-z\|})} \le \frac{f_0(x) + 2L\delta_N}{f_1(x) - L\delta_N}$$

On the other hand, if $||z - x|| \ge \delta_N$ and $f_0(x) > 2L\delta_N$,

$$\nu_0(B_{z,\|z-x\|}) \ge (f_0(x) - 2L\delta_N)\lambda(B_{0,\delta_N}) = (f_0(x) - 2L\delta_N)V_d\delta_N^d,$$

where V_d is the Lebesgue measure of the unit ball on \mathbb{R}^d .

Let $\delta_N = (\frac{4}{f_L V_d})^{1/d} (\frac{M}{N_0})^{1/d}$. Since $M/N_0 \to 0$, we have $\delta_N \to 0$ as $N_0 \to \infty$. Taking N_0 large enough so that $\delta_N < f_L/(4L)$ and $\delta_N \le \delta/2$, then

$$2LV_d\delta_N^{d+1} = \frac{M}{N_0}\frac{8L}{f_L}\delta_N < 2\frac{M}{N_0}.$$

Then for any $(\nu_0, \nu_1) \in \mathcal{P}_{x,p}(f_L, f_U, L, d, \delta)$,

$$(f_0(x) - 2L\delta_N)V_d\delta_N^d > 4\frac{f_0(x)}{f_L}\frac{M}{N_0} - 2\frac{M}{N_0} \ge 2\frac{M}{N_0}.$$

With a slight abuse of notation, let $W := \nu_0(B_{Z,||x-Z||})$. Then $W \leq 2\frac{M}{N_0}$ implies that $||Z - x|| \leq \delta_N$. Depending on the value of $f_1(x)$, the proof is separated into two cases.

Case I. $f_1(x) > L\delta_N$.

Upper bound. Proceeding similar to (7.3), we have

$$E[\widehat{r}_{M}(x)] = \frac{N_{0}}{M} E[\nu_{1}(A_{M}(x))] = \frac{N_{0}}{M} P\left(W \le \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||})\right) \\
 \le \frac{N_{0}}{M} P\left(W \le \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||}), \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||}) \le 2\frac{M}{N_{0}}\right) + \frac{N_{0}}{M} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \\
 = \frac{N_{0}}{M} P\left(W \le \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||}), \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||}) \le 2\frac{M}{N_{0}}, ||Z - x|| \le \delta_{N}\right) + \frac{N_{0}}{M} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \\
 \le \frac{N_{0}}{M} P\left(W \le \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||}), ||Z - x|| \le \delta_{N}\right) + \frac{N_{0}}{M} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \\
 \le \frac{N_{0}}{M} P\left(\frac{f_{0}(x) - 2L\delta_{N}}{f_{1}(x) + L\delta_{N}}\nu_{1}(B_{x,||x-Z||}) \le \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||}), ||Z - x|| \le \delta_{N}\right) + \frac{N_{0}}{M} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \\
 \le \frac{N_{0}}{M} P\left(\frac{f_{0}(x) - 2L\delta_{N}}{f_{1}(x) + L\delta_{N}}\nu_{1}(B_{x,||x-Z||}) \le \nu_{0}(B_{Z,||\mathcal{X}_{(M)}(Z)-Z||})\right) + \frac{N_{0}}{M} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \\
 = \frac{N_{0}}{M} P\left(\frac{f_{0}(x) - 2L\delta_{N}}{f_{1}(x) + L\delta_{N}}U \le U_{(M)}\right) + \frac{N_{0}}{M} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right).$$

$$(7.29)$$

For the second term in (7.29), since $M/\log N_0 \to \infty$, for any $\gamma > 0$,

$$\frac{N_0}{M} P\left(U_{(M)} > 2\frac{M}{N_0}\right) = \frac{N_0}{M} P\left(Bin\left(N_0, 2\frac{M}{N_0}\right) \le M\right) \\
\le \frac{N_0}{M} \exp\left((\log 2 - 1)M\right) = \frac{N_0}{M} N_0^{-(1 - \log 2)M/\log N_0} \prec N_0^{-\gamma}.$$
(7.30)

For the first term in (7.29), proceeding as (7.5), we obtain

$$\frac{N_0}{M} P\Big(\frac{f_0(x) - 2L\delta_N}{f_1(x) + L\delta_N} U \le U_{(M)}\Big) \le \frac{f_1(x) + L\delta_N}{f_0(x) - 2L\delta_N} \frac{N_0}{N_0 + 1}.$$

Then we obtain

$$\mathbf{E}[\hat{r}_M(x)] \le \frac{f_1(x) + L\delta_N}{f_0(x) - 2L\delta_N} \frac{N_0}{N_0 + 1} + o(N_0^{-\gamma}).$$
(7.31)

Lower bound. Proceeding similar to (7.8), we have

$$\begin{split} \mathbf{E}[\widehat{r}_{M}(x)] &= \frac{N_{0}}{M} \mathbf{E}[\nu_{1}(A_{M}(x))] = \frac{N_{0}}{M} \mathbf{P}\left(W \leq \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|})\right) \\ &\geq \frac{N_{0}}{M} \mathbf{P}\left(W \leq \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \leq 2\frac{M}{N_{0}}\right) \\ &= \frac{N_{0}}{M} \mathbf{P}\left(W \leq \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \leq 2\frac{M}{N_{0}}, \|Z-x\| \leq \delta_{N}\right) \\ &\geq \frac{N_{0}}{M} \mathbf{P}\left(\frac{f_{0}(x) + 2L\delta_{N}}{f_{1}(x) - L\delta_{N}} \nu_{1}(B_{x,\|x-Z\|}) \leq \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|}) \leq 2\frac{M}{N_{0}}, \|Z-x\| \leq \delta_{N}\right) \end{split}$$

$$\begin{split} &= \frac{N_0}{M} \mathbb{P}\Big(\frac{f_0(x) + 2L\delta_N}{f_1(x) - L\delta_N} \nu_1(B_{x, \|x - Z\|}) \le \nu_0(B_{Z, \|\mathcal{X}_{(M)}(Z) - Z\|}), \nu_0(B_{Z, \|\mathcal{X}_{(M)}(Z) - Z\|}) \le 2\frac{M}{N_0} \\ &\ge \frac{N_0}{M} \mathbb{P}\Big(\frac{f_0(x) + 2L\delta_N}{f_1(x) - L\delta_N} U \le U_{(M)}\Big) - \frac{N_0}{M} \mathbb{P}\Big(U_{(M)} > 2\frac{M}{N_0}\Big) \\ &= \frac{f_1(x) - L\delta_N}{f_0(x) + 2L\delta_N} \int_0^{\frac{f_0(x) + 2L\delta_N}{f_1(x) - L\delta_N} \frac{N_0}{M}} \mathbb{P}\Big(U_{(M)} \ge \frac{M}{N_0}t\Big) \mathrm{d}t - \frac{N_0}{M} \mathbb{P}\Big(U_{(M)} > 2\frac{M}{N_0}\Big). \end{split}$$

Consider the first term. If $\frac{f_0(x)+2L\delta_N}{f_1(x)-L\delta_N} \ge 1$, then

$$\frac{f_1(x) - L\delta_N}{f_0(x) + 2L\delta_N} \int_0^{\frac{f_0(x) + 2L\delta_N}{f_1(x) - L\delta_N} \frac{N_0}{M}} P\Big(U_{(M)} \ge \frac{M}{N_0}t\Big) dt = \frac{f_1(x) - L\delta_N}{f_0(x) + 2L\delta_N} \frac{N_0}{N_0 + 1}.$$

If $\frac{f_0(x)+2L\delta_N}{f_1(x)-L\delta_N} < 1$, using the Chernoff bound, for any $\gamma > 0$,

$$\int_{\frac{f_0(x)+2L\delta_N}{f_1(x)-L\delta_N}\frac{N_0}{M}}^{\frac{N_0}{M}} P\Big(U_{(M)} \ge \frac{M}{N_0}t\Big) dt \le \int_{\frac{f_0(x)}{f_1(x)}\frac{N_0}{M}}^{\frac{N_0}{M}} P\Big(U_{(M)} \ge \frac{M}{N_0}t\Big) dt \le \int_{\frac{f_L}{f_U}\frac{N_0}{M}}^{\frac{N_0}{M}} P\Big(U_{(M)} \ge \frac{M}{N_0}t\Big) dt \le \left[1 - \frac{f_L}{f_U}\right] \frac{N_0}{M} P\Big(U_{(M)} \ge \frac{f_L}{f_U}\Big) \le \left[1 - \frac{f_L}{f_U}\right] \frac{N_0}{M} \exp\left[M - \frac{f_L}{f_U}N_0 - M\log M + M\log\left(\frac{f_L}{f_U}N_0\right)\right] \\ \prec N_0^{-\gamma}.$$

The last step is due to $M \log N_0/N_0 \rightarrow 0$. Recalling (7.30), we then obtain

$$\mathbf{E}\left[\widehat{r}_{M}(x)\right] \geq \frac{f_{1}(x) - L\delta_{N}}{f_{0}(x) + 2L\delta_{N}} \frac{N_{0}}{N_{0} + 1} - o(N_{0}^{-\gamma}).$$
(7.32)

Combining (7.31) and (7.32), and taking N_0 large enough so that $L\delta_N \leq f_U \wedge (f_L/4)$, one obtains $\left| \mathbf{E}[\hat{r}_M(x)] - r(x) \right| \le \left| \frac{f_1(x) + L\delta_N}{f_0(x) - 2L\delta_N} \frac{N_0}{N_0 + 1} - \frac{f_1(x)}{f_0(x)} \right| \bigvee \left| \frac{f_1(x) - L\delta_N}{f_0(x) + 2L\delta_N} \frac{N_0}{N_0 + 1} - \frac{f_1(x)}{f_0(x)} \right| + o(N_0^{-\gamma})$ $\leq \frac{f_0(x)L\delta_N + 2f_1(x)L\delta_N}{f_0(x)(f_0(x) - 2L\delta_N)} + \frac{1}{N_0 + 1}\frac{f_1(x) + L\delta_N}{f_0(x) - 2L\delta_N} + o(N_0^{-\gamma})$ $\leq \left(\frac{2}{f_L} + \frac{4f_U}{f_L^2}\right) L\delta_N + \frac{4f_U}{f_L} \frac{1}{N_0 + 1} + o(N_0^{-\gamma}).$

By the selection of δ_N and that the right hand side does not depend on x, we complete the proof for this case.

Case II. $f_1(x) \leq L\delta_N$.

The upper bound (7.31) in Case I still holds for this case. Accordingly, taking N_0 large enough so that $L\delta_N \leq f_L/4$, one has

$$\begin{aligned} \left| \mathbf{E}[\hat{r}_{M}(x)] - r(x) \right| &\leq \mathbf{E}[\hat{r}_{M}(x)] + r(x) \\ &\leq \frac{f_{1}(x) + L\delta_{N}}{f_{0}(x) - 2L\delta_{N}} \frac{N_{0}}{N_{0} + 1} + \frac{f_{1}(x)}{f_{0}(x)} + o(N_{0}^{-\gamma}) \leq \frac{4}{f_{L}}L\delta_{N} + \frac{1}{f_{L}}L\delta_{N} + o(N_{0}^{-\gamma}). \end{aligned}$$
complete the whole proof.

We thus complete the whole proof.

Proof of Theorem 4.3(ii). By the law of total variance,

$$\operatorname{Var}\left[\widehat{r}_{M}(x)\right] = \operatorname{E}\left[\operatorname{Var}\left[\widehat{r}_{M}(x) \mid \boldsymbol{X}\right]\right] + \operatorname{Var}\left[\operatorname{E}\left[\widehat{r}_{M}(x) \mid \boldsymbol{X}\right]\right].$$
(7.33)

For the first term in (7.33), let Z be a copy drawn from ν_1 independently of the data. Then, since $[Z_j]_{j=1}^{N_1}$ are i.i.d,

where C > 0 is a constant only depending on f_L, f_U . The last step is due to (7.31).

For the second term in (7.33), notice that

$$\operatorname{Var}\left[\operatorname{E}\left[\widehat{r}_{M}(x) \mid \boldsymbol{X}\right]\right] = \operatorname{Var}\left[\operatorname{E}\left[\frac{N_{0}}{N_{1}M}K_{M}(x) \mid \boldsymbol{X}\right]\right] = \left(\frac{N_{0}}{M}\right)^{2}\operatorname{Var}\left[\nu_{1}\left(A_{M}(x)\right)\right].$$

Recalling that $W = \nu_0(B_{Z,||x-Z||})$, we have the following lemma about the density of W near x.

Lemma 7.3. Denote the density of W by f_W . Then for any $(\nu_0, \nu_1) \in \mathcal{P}_{x,p}(f_L, f_U, L, d, \delta)$,

$$f_W(0) = r(x).$$

Furthermore, for any $\epsilon > 0$ and N_0 sufficiently large, we have for all $0 \le w \le 2M/N_0$,

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{\mathbf{P}}(f_L,f_U,\delta,L,d)} f_W(w) \le (1+\epsilon)\frac{f_U}{f_L}.$$

Due to Lemma 7.3, we can take N_0 sufficiently large so that for any $0 \le w \le 2M/N_0$,

$$\sup_{(\nu_0,\nu_1)\in\mathcal{P}_{\mathbf{p}}(f_L,f_U,\delta,L,d)} f_W(w) \le 2\frac{f_U}{f_L}.$$

Let Z, \widetilde{Z} be two independent copies from ν_1 that are further independent of the data. Let $W = \nu_0(B_{Z,\|x-Z\|})$ and $\widetilde{W} = \nu_0(B_{\widetilde{Z},\|x-\widetilde{Z}\|})$. Let $V = \nu_0(B_{Z,\|\mathcal{X}_{(M)}(Z)-Z\|})$ and $\widetilde{V} = \nu_0(B_{\widetilde{Z},\|\mathcal{X}_{(M)}(\widetilde{Z})-\widetilde{Z}\|})$. We then have

$$\operatorname{Var}\left[\nu_{1}\left(A_{M}(x)\right)\right] = \operatorname{E}\left[\nu_{1}^{2}\left(A_{M}(x)\right)\right] - \left(\operatorname{E}\left[\nu_{1}\left(A_{M}(x)\right)\right]\right)^{2}$$

$$= \operatorname{P}\left(Z \in A_{M}(x), \widetilde{Z} \in A_{M}(x)\right) - \operatorname{P}\left(Z \in A_{M}(x)\right)\operatorname{P}\left(\widetilde{Z} \in A_{M}(x)\right)$$

$$= \operatorname{P}\left(\|x - Z\| \leq \|\mathcal{X}_{(M)}(Z) - Z\|, \|x - \widetilde{Z}\| \leq \|\mathcal{X}_{(M)}(\widetilde{Z}) - \widetilde{Z}\|\right)$$

$$- \operatorname{P}\left(\|x - Z\| \leq \|\mathcal{X}_{(M)}(Z) - Z\|\right)\operatorname{P}\left(\|x - \widetilde{Z}\| \leq \|\mathcal{X}_{(M)}(\widetilde{Z}) - \widetilde{Z}\|\right)$$

$$= \operatorname{P}\left(W \leq V, \widetilde{W} \leq \widetilde{V}\right) - \operatorname{P}\left(W \leq V\right)\operatorname{P}\left(\widetilde{W} \leq \widetilde{V}\right).$$

Due to the independence between Z and \widetilde{Z} , W and \widetilde{W} are independent. Notice that V | Z have the same distribution as $U_{(M)}$ for any $Z \in \mathbb{R}^d$, then V and Z are independent, so are \widetilde{V} and \widetilde{Z} .

Let's expand the variance further as

$$\operatorname{Var}\left[\nu_1\big(A_M(x)\big)\right]$$

$$= \left[P\left(W \le V, \widetilde{W} \le \widetilde{V}, W \le 2\frac{M}{N_0}, \widetilde{W} \le 2\frac{M}{N_0} \right) - P\left(W \le V, W \le 2\frac{M}{N_0} \right) P\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0} \right) \right] \\ + \left[P\left(W \le V, \widetilde{W} \le \widetilde{V} \right) - P\left(W \le V, \widetilde{W} \le \widetilde{V}, W \le 2\frac{M}{N_0}, \widetilde{W} \le 2\frac{M}{N_0} \right) \right] \\ - \left[P\left(W \le V \right) P\left(\widetilde{W} \le \widetilde{V} \right) - P\left(W \le V, W \le 2\frac{M}{N_0} \right) P\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0} \right) \right].$$
(7.35)

For the first term in (7.35), we have the following lemma.

Lemma 7.4. We have

$$\left(\frac{N_0}{M}\right)^2 \left[\mathbf{P}\left(W \le V, \widetilde{W} \le \widetilde{V}, W \le 2\frac{M}{N_0}, \widetilde{W} \le 2\frac{M}{N_0} \right) - \mathbf{P}\left(W \le V, W \le 2\frac{M}{N_0}\right) \mathbf{P}\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0}\right) \right] \\ \le C \frac{1}{M},$$

where C > 0 is a constant only depending on f_L, f_U .

For the second term in (7.35),

$$\begin{split} & P\left(W \leq V, \widetilde{W} \leq \widetilde{V}\right) - P\left(W \leq V, \widetilde{W} \leq \widetilde{V}, W \leq 2\frac{M}{N_0}, \widetilde{W} \leq 2\frac{M}{N_0}\right) \\ & \leq P\left(W \leq V, \widetilde{W} \leq \widetilde{V}, W > 2\frac{M}{N_0}\right) + P\left(W \leq V, \widetilde{W} \leq \widetilde{V}, \widetilde{W} > 2\frac{M}{N_0}\right) \\ & \leq P\left(V > 2\frac{M}{N_0}\right) + P\left(\widetilde{V} > 2\frac{M}{N_0}\right) \\ & = 2P\left(U_{(M)} > 2\frac{M}{N_0}\right). \end{split}$$

Using the Chernoff bound and $M/\log N_0 \to \infty$, for any $\gamma > 0$,

$$\left(\frac{N_0}{M}\right)^2 \mathbf{P}\left(U_{(M)} > 2\frac{M}{N_0}\right) \le \left(\frac{N_0}{M}\right)^2 \exp\left[-(1-\log 2)M\right] \prec N_0^{-\gamma},$$

We then have

$$\left(\frac{N_{0}}{M}\right)^{2} \left[P\left(W \leq V, \widetilde{W} \leq \widetilde{V}\right) - P\left(W \leq V, \widetilde{W} \leq \widetilde{V}, W \leq 2\frac{M}{N_{0}}, \widetilde{W} \leq 2\frac{M}{N_{0}}\right) \right]
- \left(\frac{N_{0}}{M}\right)^{2} \left[P\left(W \leq V\right) P\left(\widetilde{W} \leq \widetilde{V}\right) - P\left(W \leq V, W \leq 2\frac{M}{N_{0}}\right) P\left(\widetilde{W} \leq \widetilde{V}, \widetilde{W} \leq 2\frac{M}{N_{0}}\right) \right]
\leq 2 \left(\frac{N_{0}}{M}\right)^{2} P\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \prec N_{0}^{-\gamma}.$$
(7.36)

For the third term in (7.35), it is easy to see

$$\left[\mathbf{P}\left(W \le V\right) \mathbf{P}\left(\widetilde{W} \le \widetilde{V}\right) - \mathbf{P}\left(W \le V, W \le 2\frac{M}{N_0}\right) \mathbf{P}\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0}\right) \right] \ge 0.$$

Plugging Lemma 7.4 and (7.36) into (7.35), we obtain

$$\left(\frac{N_0}{M}\right)^2 \operatorname{Var}\left[\nu_1(A_M(x))\right] \lesssim C\frac{1}{M},\tag{7.37}$$

where C > 0 is a constant only depending on f_L, f_U .

Plugging (7.34) and (7.37) into (7.33) completes the proof.

7.5 Proof of Theorem 4.4

Proof of Theorem 4.4. We only have to prove the first claim as the second is trivial.

Take $\delta_N = (\frac{4}{f_L V_d})^{1/d} (\frac{M}{N_0})^{1/d}$ as in the proof of Theorem 4.3(i). Take $\delta'_N = (\frac{2}{af_L V_d})^{1/d} (\frac{M}{N_0})^{1/d}$. For any $x \in \mathbb{R}^d$, denote the distance of x to the boundary of S_1 by $\Delta(x)$, i.e., $\Delta(x) = \inf_{z \in \partial S_1} ||z - x||$.

Depending on the position of x and the value of $\Delta(x)$, we separate the proof into three cases. Case I. $x \in S_1$ and $\Delta(x) > 2\delta_N$.

In this case, since $\Delta(x) > 2\delta_N$, for any $||z - x|| \le \delta_N$, we have $B_{z,||z-x||} \subset S_1$. From the smoothness conditions on f_0 and f_1 , similar to the proof of Theorem 4.3, we have

$$E\left[\int_{\mathbb{R}^{d}} \left| \hat{r}_{M}(x) - r(x) \right| f_{0}(x) \mathbb{1}(x \in S_{1}, \Delta(x) > 2\delta_{N}) dx \right] \\
 = \int_{\mathbb{R}^{d}} E\left[\left| \hat{r}_{M}(x) - r(x) \right| \right] f_{0}(x) \mathbb{1}(x \in S_{1}, \Delta(x) > 2\delta_{N}) dx \\
 \leq \int_{\mathbb{R}^{d}} \left(E\left[\hat{r}_{M}(x) - r(x) \right]^{2} \right)^{1/2} f_{0}(x) \mathbb{1}(x \in S_{1}, \Delta(x) > 2\delta_{N}) dx \\
 \leq C\left[\left(\frac{M}{N_{0}} \right)^{1/d} + \left(\frac{1}{M} \right)^{1/2} + \left(\frac{N_{0}}{MN_{1}} \right)^{1/2} \right] \int_{\mathbb{R}^{d}} f_{0}(x) \mathbb{1}(x \in S_{1}, \Delta(x) > 2\delta_{N}) dx \\
 \leq C\left[\left(\frac{M}{N_{0}} \right)^{1/d} + \left(\frac{1}{M} \right)^{1/2} + \left(\frac{N_{0}}{MN_{1}} \right)^{1/2} \right] \int_{\mathbb{R}^{d}} f_{0}(x) dx \\
 = C\left[\left(\frac{M}{N_{0}} \right)^{1/d} + \left(\frac{1}{M} \right)^{1/2} + \left(\frac{N_{0}}{MN_{1}} \right)^{1/2} \right],$$
 (7.38)

where the constant C > 0 only depends on f_L, f_U, L, d .

Case II. $x \in S_0 \setminus S_1$ and $\Delta(x) > \delta'_N$.

In this case, r(x) = 0 and for any $z \in S_1$,

$$\nu_0(B_{z,\|z-x\|}) \ge f_L \lambda(B_{z,\|z-x\|} \cap S_0) \ge a f_L \lambda(B_{z,\|z-x\|}) > a f_L V_d \delta_N'^d \ge 2\frac{M}{N_0}$$

Then for any $\gamma > 0$,

$$\mathbb{E}\left[\left|\widehat{r}_{M}(x) - r(x)\right|\right] = \mathbb{E}\left[\widehat{r}_{M}(x)\right] = \frac{N_{0}}{M} \mathbb{E}\left[\nu_{1}\left(A_{M}(x)\right)\right]$$
$$= \frac{N_{0}}{M} \mathbb{P}\left(W \le \nu_{0}(B_{Z, \|\mathcal{X}_{(M)}(Z) - Z\|})\right) \le \frac{N_{0}}{M} \mathbb{P}\left(U_{(M)} > 2\frac{M}{N_{0}}\right) \prec N_{0}^{-\gamma}.$$

One then obtains

Case III. $x \in S_0$ and $\Delta(x) \leq (2\delta_N) \vee \delta'_N$. In this case, for any $z \in S_1$,

$$\nu_0(B_{z,\|z-x\|}) \ge f_L \lambda(B_{z,\|z-x\|} \cap S_0) \ge a f_L \lambda(B_{z,\|z-x\|}) \ge \frac{a f_L}{f_U} \nu_1(B_{x,\|z-x\|}).$$
Accordingly,

$$\begin{split} & \mathbf{E}\Big[\Big|\widehat{r}_{M}(x) - r(x)\Big|\Big] \leq \mathbf{E}\Big[\widehat{r}_{M}(x)\Big] + r(x) = \frac{N_{0}}{M}\mathbf{P}\Big(W \leq \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z) - Z\|})\Big) + r(x) \\ & \leq \frac{N_{0}}{M}\mathbf{P}\Big(\frac{af_{L}}{f_{U}}\nu_{1}(B_{x,\|x-Z\|}) \leq \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z) - Z\|}), \nu_{0}(B_{Z,\|\mathcal{X}_{(M)}(Z) - Z\|}) \leq 2\frac{M}{N_{0}}\Big) + r(x) \\ & \leq \frac{N_{0}}{M}\mathbf{P}\Big(\frac{af_{L}}{f_{U}}U \leq U_{(M)}\Big) + r(x) = \frac{f_{U}}{af_{L}}(1 + o(1)) + \frac{f_{U}}{f_{L}}. \end{split}$$

From the definition of δ_N, δ'_N and $M/N_0 \to 0$, we have $\delta_N, \delta'_N \to 0$ as $N_0 \to \infty$. Since the surface area of S_1 is bounded by H, we have

$$\lambda(\{x: \Delta(x) \le (2\delta_N) \lor \delta'_N\}) \lesssim H\{(2\delta_N) \lor \delta'_N\}$$

Then we obtain

where the constant C > 0 only depends on f_L, f_U, a, H, d .

Combining (7.38), (7.39), (7.40) completes the proof.

A Proofs of the rest results

A.1 Proofs of the results in Section 3

A.1.1 Proof of Theorem 3.1

Proof of Theorem 3.1. We consider the complexities of two algorithms separately.

Algorithm 1.

The worst-case computation complexity of building a balanced k-d tree is $O(dN_0 \log N_0)$ (cf. Brown (2015)) since the size of the k-d tree is N_0 .

The average computation complexity of searching a NN is $O(\log N_0)$ from Friedman et al. (1977), and then the average computation complexity of search *M*-NNs in $\{X_i\}_{i=1}^{N_0}$ for all $\{Z_j\}_{j=1}^{N_1}$ is $O(MN_1 \log N_0)$.

Notice that $|S_j| = M$ for any $j \in [N_1]$ and then $|\bigcup_{j=1}^{N_1} S_j| \leq N_1 M$. Since the elements of each S_j are in $[N_0]$, the largest integer in $\bigcup_{j=1}^{N_1} S_j$ is N_0 . Then the computation complexity of counting step is $O(N_1M + N_0)$ due to the counting sort algorithm (Cormen et al., 2009, Section 8.2).

Combining the above three steps completes the proof for Algorithm 1.

Algorithm 2.

The computation complexity of building a k-d tree is $O(d(N_0+n)\log(N_0+n))$ from Algorithm 1 since the size of the k-d tree is $N_0 + n$.

For the searching step, for each $j \in [N_1]$, the number of NNs to be searched is

$$M + \sum_{i=1}^{n} \mathbb{1}(\|x_i - Z_j\| \le \|\mathcal{X}_{(M)}(Z_j) - Z_j\|).$$

Then from (2.2), the total number of NNs searched for all $j \in [N_1]$ is

$$\sum_{j=1}^{N_1} \left(M + \sum_{i=1}^n \mathbb{1}\left(\|x_i - Z_j\| \le \|\mathcal{X}_{(M)}(Z_j) - Z_j\| \right) \right) = N_1 M + \sum_{i=1}^n K_M(x_i).$$

Let X, Y be two independent copies from ν_0, ν_1 , respectively. Since $[Z_j]_{j=1}^{N_1}$ are i.i.d. and $[X_i]_{i=1}^{N_0} \bigcup [x_i]_{i=1}^n$ are i.i.d, we have

$$E\Big[\sum_{i=1}^{n} K_M(x_i)\Big] = nE[K_M(X)] = N_1 nE[\nu_1(A_M(X))] = N_1 n\frac{M}{N_0 + 1},$$

where the last equation is due to

$$E[\nu_1(A_M(X))] = P(||X - Y|| \le ||\mathcal{X}_{(M)}(Y) - Y||) = P(U \le U_{(M)}) = \frac{M}{N_0 + 1},$$

by using the probability integral transform. Then the average computation complexity for the searching step is $O(\frac{N_1M}{N_0}(N_0+n)\log(N_0+n))$

For the counting step, the computation complexity for counting $\bigcup_{j=1}^{N_1} S_j$ is $O(N_0 + N_1M)$ since the cardinality of $\bigcup_{j=1}^{N_1} S_j$ is N_1M and the largest integer is N_0 . The average computation complexity for counting $\bigcup_{j=1}^{N_1} S'_j$ is $O(\frac{N_1M}{N_0}n+n)$ since the average cardinality of $\bigcup_{j=1}^{N_1} S'_j$ is $\frac{N_1M}{N_0}n$ and the largest integer is n.

Combining the above three steps completes the proof for Algorithm 2.

A.2 Proofs of the results in Section 4

A.2.1 Proof of Corollary 4.1

Proof of Corollary 4.1. Corollary 4.1 can be established following the same way as that of Theorem 4.2 but with less effort since we only have to show

$$\lim_{N_0 \to \infty} \mathbb{E} \left[\int_{\mathbb{R}^d} \left| \mathbb{E} \left[\widehat{r}_M(x) \, \middle| \, \mathbf{X} \right] - r(x) \right|^p f_0(x) \mathrm{d}x \right] = 0.$$

In detail, denote the Radon-Nikodym derivative of the probability measure of W with respect to ν_0 by r_W . We then have

$$\begin{split} &\lim_{N_0 \to \infty} \operatorname{E} \left[\left| \frac{N_0}{M} \nu_1 \left(A_M(W) \right) - r(W) \right|^p \right] = \lim_{N_0 \to \infty} \operatorname{E} \left[\int_{\mathbb{R}^d} \left| \frac{N_0}{M} \nu_1 \left(A_M(x) \right) - r(x) \right|^p r_W(x) d\nu_0(x) \right] \\ &= \lim_{N_0 \to \infty} \operatorname{E} \left[\int_{\mathbb{R}^d} \left| \frac{N_0}{M} \nu_1 \left(A_M(x) \right) - r(x) \right|^p r_W(x) f_0(x) dx \right] \\ &\lesssim \lim_{N_0 \to \infty} \operatorname{E} \left[\int_{\mathbb{R}^d} \left| \frac{N_0}{M} \nu_1 \left(A_M(x) \right) - r(x) \right|^p f_0(x) dx \right] \end{split}$$

$$= \limsup_{N_0 \to \infty} \mathbb{E} \Big[\int_{\mathbb{R}^d} \Big| \mathbb{E} \Big[\widehat{r}_M(x) \, \Big| \, \mathbf{X} \Big] - r(x) \Big|^p f_0(x) \mathrm{d}x \Big] = 0$$

Here the last line has intrinsically been established in the proof of Theorem 4.2.

Noticing that $E[r(W)]^p$ is bounded under Assumption 4.1, the proof is thus complete.

A.2.2 Proof of Proposition 4.1

Proof of Proposition 4.1. We take ν_0 and ν_1 to share the same support, and assume x to be the origin of \mathbb{R}^d without loss of generality.

When $N_1 \leq N_0$, we take ν_0 to be the uniform distribution with density f_L on $[-f_L^{-1/d}/2, f_L^{-1/d}/2]^d$. Then the MSE is lower bounded by the density estimation over Lipchitz class with N_1 samples.

When $N_0 \leq N_1$, we take ν_1 to be the uniform distribution with density f_U on $[-f_U^{-1/d}/2, f_U^{-1/d}/2]^d$. Notice that $1/f_0$ is also local Lipchitz from the lower boundness condition and local Lipchitz condition on f_0 . Then the MSE is lower bounded by the density estimation over Lipchitz class with N_0 samples.

We then complete the proof by combining the above two lower bounds and then using the famous minimax lower bound in Lipschitz density estimation (Tsybakov, 2009, Exercise 2.8), \Box

A.2.3 Proof of Proposition 4.2

Proof of Proposition 4.2. We take ν_0 and ν_1 to be of the same support.

When $N_1 \leq N_0$, we take ν_0 to be the uniform distribution with density f_L on $[-f_L^{-1/d}/2, f_L^{-1/d}/2]^d$. Then the L_1 risk is lower bounded by the L_1 risk over support of density estimation over Lipchitz class with N_1 samples.

When $N_0 \leq N_1$, we take ν_1 to be the uniform distribution with density f_U on $[-f_U^{-1/d}/2, f_U^{-1/d}/2]^d$. Notice $1/f_0$ is also Lipchitz from the lower boundness condition and Lipchitz condition on f_0 . From the lower boundness condition on f_0 , the L_1 risk is lower bounded by the L_1 risk over support of density estimation over Lipchitz class with N_0 samples.

We then complete the proof by combining the above two lower bounds and then using then the minimax lower bound of L_1 risk for density estimation over Lipchitz class (Zhao and Lai, 2020a, Theorem 1).

A.3 Proofs of the results in Section 5

We first formally define $\tilde{\tau}_{M,K}^{\text{bc}}$. Assume *n* is divisible by *K* for simplicity. Let $[I_k]_{k=1}^K$ be a *K*-fold random partition of [n], with each of size equal to n' = n/K. For each $k \in [K]$ and $\omega \in \{0, 1\}$, construct $K_{M,k}^{\omega}(\cdot), \hat{\mu}_{\omega,k}(\cdot)$ using data $[(X_i, D_i, Y_i)]_{i=1, i \notin I_k}^n$. Then define

$$\begin{aligned} \breve{\tau}_{M,k}^{\text{bc}} &= \frac{1}{n'} \sum_{i=1,i\in I_k}^n \left[\widehat{\mu}_{1,k}(X_i) - \widehat{\mu}_{0,k}(X_i) \right] \\ &+ \frac{1}{n'} \Big[\sum_{i=1,i\in I_k, D_i=1}^n \Big(1 + \frac{K_{M,k}^1(X_i)}{M} \Big) \Big(Y_i - \widehat{\mu}_{1,k}(X_i) \Big) - \sum_{i=1,i\in I_k, D_i=0}^n \Big(1 + \frac{K_{M,k}^0(X_i)}{M} \Big) \Big(Y_i - \widehat{\mu}_{0,k}(X_i) \Big) \Big] \end{aligned}$$

and

$$\widetilde{\tau}_{M,K}^{\mathrm{bc}} := \frac{1}{K} \sum_{k=1}^{K} \widecheck{\tau}_{M,k}^{\mathrm{bc}}.$$

A.3.1 Proof of Lemma 5.1

Proof of Lemma 5.1. From Equ.(3) in Abadie and Imbens (2011), the bias-corrected estimator imputes the missing potential outcomes by

$$\widehat{Y}_{i}^{\text{bc}}(0) := \begin{cases} Y_{i}, & \text{if } D_{i} = 0, \\ \frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{0}(i)} (Y_{j} + \widehat{\mu}_{0}(X_{i}) - \widehat{\mu}_{0}(X_{j})), & \text{if } D_{i} = 1, \end{cases}$$

and

$$\widehat{Y}_{i}^{\text{bc}}(1) := \begin{cases} \frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{1}(i)} (Y_{j} + \widehat{\mu}_{1}(X_{i}) - \widehat{\mu}_{1}(X_{j})), & \text{if } D_{i} = 0, \\ Y_{i}, & \text{if } D_{i} = 1. \end{cases}$$

The bias-corrected estimator is defined as:

$$\widehat{\tau}_M^{\mathrm{bc}} := \frac{1}{n} \sum_{i=1}^n \left[\widehat{Y}_i^{\mathrm{bc}}(1) - \widehat{Y}_i^{\mathrm{bc}}(0) \right].$$

Then by simple algebra, we have

$$\begin{aligned} \widehat{\tau}_{M}^{\rm bc} &= \frac{1}{n} \sum_{i=1}^{n} \left[\widehat{Y}_{i}^{\rm bc}(1) - \widehat{Y}_{i}^{\rm bc}(0) \right] \\ &= \frac{1}{n} \sum_{i=1,D_{i}=1}^{n} \left[Y_{i} - \frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{0}(i)} (Y_{j} + \widehat{\mu}_{0}(X_{i}) - \widehat{\mu}_{0}(X_{j})) \right] + \frac{1}{n} \sum_{i=1,D_{i}=0}^{n} \left[\frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{1}(i)} (Y_{j} + \widehat{\mu}_{1}(X_{i}) - \widehat{\mu}_{1}(X_{j})) - Y_{i} \right] \\ &= \frac{1}{n} \sum_{i=1,D_{i}=1}^{n} \left[\widehat{R}_{i} + \widehat{\mu}_{1}(X_{i}) - \widehat{\mu}_{0}(X_{i}) - \frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{0}(i)} \widehat{R}_{j} \right] + \frac{1}{n} \sum_{i=1,D_{i}=0}^{n} \left[\frac{1}{M} \sum_{j \in \mathcal{J}_{M}^{1}(i)} \widehat{R}_{j} - \widehat{R}_{i} + \widehat{\mu}_{1}(X_{i}) - \widehat{\mu}_{0}(X_{i}) \right] \\ &= \frac{1}{n} \sum_{i=1}^{n} \left[\widehat{\mu}_{1}(X_{i}) - \widehat{\mu}_{0}(X_{i}) \right] + \frac{1}{n} \left[\sum_{i=1,D_{i}=1}^{n} \left(1 + \frac{K_{M}^{1}(i)}{M} \right) \widehat{R}_{i} - \sum_{i=1,D_{i}=0}^{n} \left(1 + \frac{K_{M}^{0}(i)}{M} \right) \widehat{R}_{i} \right]. \end{aligned}$$
This completes the proof.

This completes the proof.

Proof of Theorem 5.1 A.3.2

Proof of Theorem 5.1(i). Part I. Suppose the density function is sufficiently smooth. For any $i \in [n], \text{ let } \bar{R}_i := Y_i - \bar{\mu}_{D_i}(X_i). \text{ From } (5.2),$

$$\hat{\tau}_{M}^{\text{bc}} = \hat{\tau}^{\text{reg}} + \frac{1}{n} \Big[\sum_{i=1,D_{i}=1}^{n} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \hat{R}_{i} - \sum_{i=1,D_{i}=0}^{n} \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \hat{R}_{i} \Big]$$
$$= \hat{\tau}^{\text{reg}} + \frac{1}{n} \Big[\sum_{i=1}^{n} D_{i} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \hat{R}_{i} - \sum_{i=1}^{n} (1 - D_{i}) \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \hat{R}_{i} \Big]$$

$$= \frac{1}{n} \sum_{i=1}^{n} \left[\widehat{\mu}_{1}(X_{i}) - \overline{\mu}_{1}(X_{i}) \right] - \frac{1}{n} \sum_{i=1}^{n} \left[\widehat{\mu}_{0}(X_{i}) - \overline{\mu}_{0}(X_{i}) \right] \\ + \frac{1}{n} \left[\sum_{i=1}^{n} D_{i} \left(1 + \frac{K_{M}^{1}(i)}{M} \right) \left(\overline{\mu}_{1}(X_{i}) - \widehat{\mu}_{1}(X_{i}) \right) - \sum_{i=1}^{n} (1 - D_{i}) \left(1 + \frac{K_{M}^{0}(i)}{M} \right) \left(\overline{\mu}_{0}(X_{i}) - \widehat{\mu}_{0}(X_{i}) \right) \right] \\ + \frac{1}{n} \left[\sum_{i=1}^{n} D_{i} \left(1 + \frac{K_{M}^{1}(i)}{M} - \frac{1}{e(X_{i})} \right) \overline{R}_{i} - \sum_{i=1}^{n} (1 - D_{i}) \left(1 + \frac{K_{M}^{0}(i)}{M} - \frac{1}{1 - e(X_{i})} \right) \overline{R}_{i} \right] \\ + \frac{1}{n} \left[\sum_{i=1}^{n} \left(1 - \frac{D_{i}}{e(X_{i})} \right) \overline{\mu}_{1}(X_{i}) - \sum_{i=1}^{n} \left(1 - \frac{1 - D_{i}}{1 - e(X_{i})} \right) \overline{\mu}_{0}(X_{i}) \right] \\ + \frac{1}{n} \left[\sum_{i=1}^{n} \frac{D_{i}}{e(X_{i})} Y_{i} - \sum_{i=1}^{n} \frac{1 - D_{i}}{1 - e(X_{i})} Y_{i} \right].$$
(A.1)

For each term, we only establish the first half part, and the second half can be established in the same way.

For the first term in (A.1),

$$\left|\frac{1}{n}\sum_{i=1}^{n} \left[\widehat{\mu}_{1}(X_{i}) - \overline{\mu}_{1}(X_{i})\right]\right| \leq \|\widehat{\mu}_{1} - \overline{\mu}_{1}\|_{\infty} = o_{\mathrm{P}}(1).$$

Then

$$\frac{1}{n}\sum_{i=1}^{n} \left[\widehat{\mu}_{1}(X_{i}) - \overline{\mu}_{1}(X_{i})\right] - \frac{1}{n}\sum_{i=1}^{n} \left[\widehat{\mu}_{0}(X_{i}) - \overline{\mu}_{0}(X_{i})\right] = o_{\mathrm{P}}(1).$$
(A.2)

For the second term in (A.1),

$$\left|\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)\left(\bar{\mu}_{1}(X_{i})-\hat{\mu}_{1}(X_{i})\right)\right|$$

$$\leq \|\hat{\mu}_{1}-\bar{\mu}_{1}\|_{\infty}\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)=\|\hat{\mu}_{1}-\bar{\mu}_{1}\|_{\infty}=o_{\mathrm{P}}(1),$$

where the last step is due to $\sum_{i=1}^{n} D_i K_M^1(i) = n_0 M$. We then have

$$\frac{1}{n} \Big[\sum_{i=1}^{n} D_i \Big(1 + \frac{K_M^1(i)}{M} \Big) \Big(\bar{\mu}_1(X_i) - \hat{\mu}_1(X_i) \Big) - \sum_{i=1}^{n} (1 - D_i) \Big(1 + \frac{K_M^0(i)}{M} \Big) \Big(\bar{\mu}_0(X_i) - \hat{\mu}_0(X_i) \Big) \Big] = o_{\mathrm{P}}(1).$$
(A.3)

For the third term in (A.1), by Theorem 4.2,

$$\begin{split} & \mathbf{E}\Big[\Big|\frac{1}{n}\sum_{i=1}^{n}D_{i}\Big(1+\frac{K_{M}^{1}(i)}{M}-\frac{1}{e(X_{i})}\Big)\bar{R}_{i}\Big|\Big] \leq \mathbf{E}\Big[\Big|D_{i}\Big(1+\frac{K_{M}^{1}(i)}{M}-\frac{1}{e(X_{i})}\Big)\bar{R}_{i}\Big|\Big] \\ & \leq \mathbf{E}\Big[1+\frac{K_{M}^{1}(i)}{M}-\frac{1}{e(X_{i})}\Big]^{2}\mathbf{E}[D_{i}\bar{R}_{i}]^{2}=\mathbf{E}\Big[1+\frac{K_{M}^{1}(i)}{M}-\frac{1}{e(X_{i})}\Big]^{2}\mathbf{E}[D_{i}(Y_{i}(1)-\bar{\mu}_{1}(X_{i}))^{2}] \\ & \leq \mathbf{E}\Big[1+\frac{K_{M}^{1}(i)}{M}-\frac{1}{e(X_{i})}\Big]^{2}\mathbf{E}\Big[\sigma_{1}^{2}(X_{i})+(\mu_{1}(X_{i})-\bar{\mu}_{1}(X_{i}))^{2}\Big]=o(1). \end{split}$$

We then obtain

$$\frac{1}{n} \Big[\sum_{i=1}^{n} D_i \Big(1 + \frac{K_M^1(i)}{M} - \frac{1}{e(X_i)} \Big) \bar{R}_i - \sum_{i=1}^{n} (1 - D_i) \Big(1 + \frac{K_M^0(i)}{M} - \frac{1}{1 - e(X_i)} \Big) \bar{R}_i \Big] = o_{\mathrm{P}}(1). \quad (A.4)$$

For the fourth term in (A.1), notice that

$$\operatorname{E}\left[\frac{1}{n}\sum_{i=1}^{n}\left(1-\frac{D_{i}}{e(X_{i})}\right)\bar{\mu}_{1}(X_{i}) \mid \boldsymbol{X}\right] = 0$$

Then

$$\operatorname{Var}\left[\frac{1}{n}\sum_{i=1}^{n}\left(1-\frac{D_{i}}{e(X_{i})}\right)\bar{\mu}_{1}(X_{i})\right] = \operatorname{E}\left[\operatorname{Var}\left[\frac{1}{n}\sum_{i=1}^{n}\left(1-\frac{D_{i}}{e(X_{i})}\right)\bar{\mu}_{1}(X_{i}) \middle| \mathbf{X}\right]\right]$$
$$=\frac{1}{n}\operatorname{E}\left[\bar{\mu}_{1}^{2}(X_{i})\left(\frac{1}{e(X_{i})}-1\right)\right] = O(n^{-1}).$$

Then

$$\frac{1}{n} \Big[\sum_{i=1}^{n} \Big(1 - \frac{D_i}{e(X_i)} \Big) \bar{\mu}_1(X_i) - \sum_{i=1}^{n} \Big(1 - \frac{1 - D_i}{1 - e(X_i)} \Big) \bar{\mu}_0(X_i) \Big] = o_{\mathbf{P}}(1).$$
(A.5)

For the fifth term in (A.1), notice that $E[Y^2]$ are bounded and $[(X_i, D_i, Y_i)]_{i=1}^n$ are i.i.d.. Using the weak law of large numbers (Durrett, 2019, Theorem 2.2.3) yields

$$\frac{1}{n} \left[\sum_{i=1}^{n} \frac{D_i}{e(X_i)} Y_i - \sum_{i=1}^{n} \frac{1 - D_i}{1 - e(X_i)} Y_i \right] \xrightarrow{\mathsf{p}} \mathbf{E} \left[Y_i(1) - Y_i(0) \right] = \tau.$$
(A.6)

Plugging (A.2), (A.3), (A.4), (A.5), (A.6) into (A.1) completes the proof.

Part II. Suppose the outcome model is correct. By (5.2),

$$\begin{aligned} \widehat{\tau}_{M}^{\text{bc}} &= \widehat{\tau}^{\text{reg}} + \frac{1}{n} \Big[\sum_{i=1,D_{i}=1}^{n} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \widehat{R}_{i} - \sum_{i=1,D_{i}=0}^{n} \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \widehat{R}_{i} \Big] \\ &= \widehat{\tau}^{\text{reg}} + \frac{1}{n} \Big[\sum_{i=1}^{n} D_{i} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \widehat{R}_{i} - \sum_{i=1}^{n} (1 - D_{i}) \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \widehat{R}_{i} \Big] \\ &= \frac{1}{n} \sum_{i=1}^{n} \Big[\widehat{\mu}_{1}(X_{i}) - \mu_{1}(X_{i}) \Big] - \frac{1}{n} \sum_{i=1}^{n} \Big[\widehat{\mu}_{0}(X_{i}) - \mu_{0}(X_{i}) \Big] \\ &+ \frac{1}{n} \Big[\sum_{i=1}^{n} D_{i} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \Big(\mu_{1}(X_{i}) - \widehat{\mu}_{1}(X_{i}) \Big) - \sum_{i=1}^{n} (1 - D_{i}) \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \Big(\mu_{0}(X_{i}) - \widehat{\mu}_{0}(X_{i}) \Big) \Big] \\ &+ \frac{1}{n} \Big[\sum_{i=1}^{n} D_{i} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \Big(Y_{i} - \mu_{1}(X_{i}) \Big) - \sum_{i=1}^{n} (1 - D_{i}) \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \Big(Y_{i} - \mu_{0}(X_{i}) \Big) \Big] \\ &+ \frac{1}{n} \sum_{i=1}^{n} \Big[\mu_{1}(X_{i}) - \mu_{0}(X_{i}) \Big]. \end{aligned}$$
(A.7)

For the first term in (A.7),

$$\left|\frac{1}{n}\sum_{i=1}^{n}\left[\widehat{\mu}_{1}(X_{i})-\mu_{1}(X_{i})\right]\right| \leq \|\widehat{\mu}_{1}-\mu_{1}\|_{\infty}=o_{\mathrm{P}}(1).$$

Then

$$\frac{1}{n}\sum_{i=1}^{n} \left[\widehat{\mu}_{1}(X_{i}) - \mu_{1}(X_{i})\right] - \frac{1}{n}\sum_{i=1}^{n} \left[\widehat{\mu}_{0}(X_{i}) - \mu_{0}(X_{i})\right] = o_{\mathrm{P}}(1).$$
(A.8)

For the second term in (A.7),

$$\left|\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)\left(\mu_{1}(X_{i})-\widehat{\mu}_{1}(X_{i})\right)\right|$$

$$\leq \|\widehat{\mu}_{1}-\mu_{1}\|_{\infty}\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)=\|\widehat{\mu}_{1}-\mu_{1}\|_{\infty}=o_{\mathrm{P}}(1).$$

where the last step is due to $\sum_{i=1}^{n} D_i K_M^1(i) = n_0 M$. We then have

$$\frac{1}{n} \Big[\sum_{i=1}^{n} D_i \Big(1 + \frac{K_M^1(i)}{M} \Big) \Big(\mu_1(X_i) - \widehat{\mu}_1(X_i) \Big) - \sum_{i=1}^{n} (1 - D_i) \Big(1 + \frac{K_M^0(i)}{M} \Big) \Big(\mu_0(X_i) - \widehat{\mu}_0(X_i) \Big) \Big] = o_{\mathrm{P}}(1).$$
(A.9)

For the third term in (A.7), noticing that $K_M^1(\cdot)$ is a function of \boldsymbol{X} and \boldsymbol{D} , one could obtain

$$\operatorname{E}\left[\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)\left(Y_{i}-\mu_{1}(X_{i})\right) \mid \boldsymbol{X},\boldsymbol{D}\right]=0.$$

Accordingly, we have

$$\operatorname{Var}\left[\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)\left(Y_{i}-\mu_{1}(X_{i})\right)\right]$$
$$=\operatorname{E}\left[\operatorname{Var}\left[\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)\left(Y_{i}-\mu_{1}(X_{i})\right)\right] \middle| \boldsymbol{X}, \boldsymbol{D}\right]$$
$$=\frac{1}{n}\operatorname{E}\left[D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)^{2}\sigma_{1}^{2}(X_{i})\right] \leq \frac{1}{n}\operatorname{E}\left[D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)^{2}\right] \|\sigma_{1}^{2}\|_{\infty},$$

where $\sigma_1^2(x) = \mathbf{E}[U_1^2 | X = x]$ for $x \in \mathbb{X}$.

Conditional on \mathbf{D} , for any $j, \ell \in [\![n]\!]$ and $j \neq \ell$ such that $D_j = D_\ell = 0$, for $i \in [\![n]\!]$ such that $D_i = 1$, it is easy to check $\mathrm{E}[K_M^1(i) \mid \mathbf{D}] = Mn_0/n_1$ and $\mathrm{E}[(K_M^1(i))^2 \mid \mathbf{D}] \asymp M^2$. Then

$$\operatorname{Var}\left[\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(1+\frac{K_{M}^{1}(i)}{M}\right)\left(Y_{i}-\mu_{1}(X_{i})\right)\right]=O(n^{-1})$$

We then obtain

$$\frac{1}{n} \Big[\sum_{i=1}^{n} D_i \Big(1 + \frac{K_M^1(i)}{M} \Big) \Big(Y_i - \mu_1(X_i) \Big) - \sum_{i=1}^{n} (1 - D_i) \Big(1 + \frac{K_M^0(i)}{M} \Big) \Big(Y_i - \mu_0(X_i) \Big) \Big] = o_{\mathrm{P}}(1).$$
(A.10)

For the fourth term in (A.7), notice that $E[\mu_{\omega}^2]$ is bounded for $\omega \in \{0, 1\}$. Using the weak law of large number, we obtain

$$\frac{1}{n}\sum_{i=1}^{n}\left[\mu_1(X_i) - \mu_0(X_i)\right] \xrightarrow{\mathbf{p}} \mathbf{E}\left[\mu_1(X_i) - \mu_0(X_i)\right] = \tau.$$
(A.11)

Plugging (A.8), (A.9), (A.10), (A.11) into (A.7) completes the proof.

Analogous analysis can be performed on $\check{\tau}_{M,k}^{\mathrm{bc}}$ for any $k \in [\![K]\!]$. Then the results apply to $\tilde{\tau}_{M,K}^{\mathrm{bc}}$ automatically since K is fixed.

Proof of Theorem 5.1(ii). From Definition 3.1 in Chernozhukov et al. (2018), $\tilde{\tau}_{M,K}^{\text{bc}}$ is the double machine learning estimator. We then follow the proof of Theorem 5.1 (recalling Remark 5.3) and use the notations in Chernozhukov et al. (2018), essentially checking Assumption 3.1 and 3.2 therein. In the following the notation in Chernozhukov et al. (2018) is adopted.

For estimating the ATE, from Equ. (5.3) in Chernozhukov et al. (2018), the score is

$$\psi(X, D, Y; \tilde{\tau}, \tilde{\zeta}) := \tilde{\mu}_1(X) - \tilde{\mu}_0(X) + \frac{D(Y - \tilde{\mu}_1(X))}{\tilde{e}(X)} - \frac{(1 - D)(Y - \tilde{\mu}_0(X))}{1 - \tilde{e}(X)} - \tilde{\tau},$$

where $\tilde{\zeta}(x) = (\tilde{\mu}_0(x), \tilde{\mu}_1(x), \tilde{\rho}_0(x), \tilde{\rho}_1(x))$ are the nuisance parameters by letting $\tilde{\rho}_0(x) = 1/(1-\tilde{e}(x))$ and $\tilde{\rho}_1(x) = 1/\tilde{e}(x)$. Let $\rho_0(x) = 1/(1-e(x))$ and $\rho_1(x) = 1/e(x)$. Then the true value is $\zeta(x) = (\mu_0(x), \mu_1(x), \rho_0(x), \rho_1(x))$.

We can then write the score as

$$\psi(X, D, Y; \widetilde{\tau}, \zeta) = \widetilde{\mu}_1(X) - \widetilde{\mu}_0(X) + D(Y - \widetilde{\mu}_1(X))\widetilde{\rho}_1(X) - (1 - D)(Y - \widetilde{\mu}_0(X))\widetilde{\rho}_0(X) - \widetilde{\tau}.$$

For any p > 0, let $||f||_p := ||f(X, D, Y)||_p = (\int |f(\omega)|^p dP_{(X, D, Y)}(\omega))^{1/p}$. For the κ in Assumption 5.1, let $q = 2 + \kappa/2$, $q_1 = 2 + \kappa$ and q_2 such that $q^{-1} = q_1^{-1} + q_2^{-1}$. Let \mathcal{T}_n be the set consisting of all $\widetilde{\zeta}$ such that for $\omega \in \{0, 1\}$,

$$\|\widetilde{\mu}_{\omega} - \mu_{\omega}\|_{\infty} = o(n^{-d/(4+2d)}), \quad \|\widetilde{\rho}_{\omega} - \rho_{\omega}\|_{1} = O(n^{-1/(d+2)}), \quad \|\widetilde{\rho}_{\omega} - \rho_{\omega}\|_{q_{2}} = o(1).$$

Then the selection of \mathcal{T}_n satisfies Assumption 3.2(a) in Chernozhukov et al. (2018) from Assumption 5.5, Theorem 4.4, and Theorem 4.2, respectively.

For step 1, we verify the Neyman orthogonality. It is easy to see $E\psi(X, D, Y; \tau, \zeta) = 0$. For any $\tilde{\zeta} \in \mathcal{T}_n$, the Gateaux derivative in the direction $\tilde{\zeta} - \zeta$ is

$$\begin{aligned} \partial_{\tilde{\zeta}} & \mathrm{E}\psi(X, D, Y; \tau, \zeta)[\tilde{\zeta} - \zeta] = \mathrm{E}[\tilde{\mu}_1(X) - \mu_1(X)] - \mathrm{E}[\tilde{\mu}_0(X) - \mu_0(X)] \\ & - \mathrm{E}[D(\tilde{\mu}_1(X) - \mu_1(X))\rho_1(X)] + \mathrm{E}[(1 - D)(\tilde{\mu}_0(X) - \mu_0(X))\rho_0(X)] \\ & + \mathrm{E}[D(Y - \mu_1(X))(\tilde{\rho}_1(X) - \rho_1(X))] - \mathrm{E}[(1 - D)(Y - \mu_0(X))(\tilde{\rho}_0(X) - \rho_0(X))] \end{aligned}$$

It is easy to check the above quantity is zero, which completes this step.

Step 2 and step 3 can be directly applied.

For step 4, we can establish in the same way that for $\omega \in \{0, 1\}$, $\|\mu_{\omega}\|_{q_1} = O(1)$ from $\|Y\|_{q_1} = O(1)$, and $\tau = O(1)$. Then from Hölder's inequality and $\|\rho_{\omega}\|_{\infty}$ is bounded for $\omega \in \{0, 1\}$,

$$\begin{split} \|\psi(X,D,Y;\tau,\widetilde{\zeta})\|_{q} &= \|\widetilde{\mu}_{1}(X) - \widetilde{\mu}_{0}(X) + D(Y - \widetilde{\mu}_{1}(X))\widetilde{\rho}_{1}(X) - (1 - D)(Y - \widetilde{\mu}_{0}(X))\widetilde{\rho}_{0}(X) - \tau\|_{q} \\ \leq &\|\widetilde{\mu}_{1}(X)\|_{q} + \|\widetilde{\mu}_{0}(X)\|_{q} + \|(Y - \widetilde{\mu}_{1}(X))\widetilde{\rho}_{1}(X)\|_{q} + \|(Y - \widetilde{\mu}_{0}(X))\widetilde{\rho}_{0}(X)\|_{q} + \tau \\ \leq &\|\mu_{1}\|_{q} + \|\widetilde{\mu}_{1} - \mu_{1}\|_{\infty} + \|\mu_{0}\|_{q} + \|\widetilde{\mu}_{0} - \mu_{0}\|_{\infty} + (\|Y\|_{q_{1}} + \|\mu_{1}\|_{q_{1}} + \|\widetilde{\mu}_{1} - \mu_{1}\|_{\infty})\|\widetilde{\rho}_{1}\|_{q_{2}} \\ &+ (\|Y\|_{q_{1}} + \|\mu_{0}\|_{q_{1}} + \|\widetilde{\mu}_{0} - \mu_{0}\|_{\infty})\|\widetilde{\rho}_{0}\|_{q_{2}} + \tau = O(1). \end{split}$$

The last step is from the definition of \mathcal{T}_n . Then we complete this step.

For step 5, from Hölder's inequality,

$$\begin{split} &\|\psi(X,D,Y;\tau,\zeta) - \psi(X,D,Y;\tau,\zeta)\|_{2} \\ \leq &\|\widetilde{\mu}_{1} - \mu_{1}\|_{2} + \|\widetilde{\mu}_{0} - \mu_{0}\|_{2} + \|D(Y - \widetilde{\mu}_{1}(X))\widetilde{\rho}_{1}(X) - D(Y - \mu_{1}(X))\rho_{1}(X)\|_{2} \\ &+ \|(1 - D)(Y - \widetilde{\mu}_{0}(X))\widetilde{\rho}_{0}(X) - (1 - D)(Y - \mu_{0}(X))\rho_{0}(X)\|_{2} \\ \leq &\|\widetilde{\mu}_{1} - \mu_{1}\|_{2} + \|\widetilde{\mu}_{0} - \mu_{0}\|_{2} + \|(Y - \mu_{1}(X))(\widetilde{\rho}_{1} - \rho_{1})\|_{2} + \|(\widetilde{\mu}_{1} - \mu_{1})\widetilde{\rho}_{1}\|_{2} \\ &+ \|(Y - \mu_{0}(X))(\widetilde{\rho}_{0} - \rho_{0})\|_{2} + \|(\widetilde{\mu}_{0} - \mu_{0})\widetilde{\rho}_{0}\|_{2} \\ \leq &\|\widetilde{\mu}_{1} - \mu_{1}\|_{\infty} + \|\widetilde{\mu}_{0} - \mu_{0}\|_{\infty} + (\|Y\|_{q_{1}} + \|\mu_{1}(X)\|_{q_{1}})\|\widetilde{\rho}_{1} - \rho_{1}\|_{q_{2}} + \|\widetilde{\mu}_{1} - \mu_{1}\|_{\infty}\|\widetilde{\rho}_{1}\|_{2} \\ &+ (\|Y\|_{q_{1}} + \|\mu_{0}(X)\|_{q_{1}})\|\widetilde{\rho}_{0} - \rho_{0}\|_{q_{2}} + \|\widetilde{\mu}_{0} - \mu_{0}\|_{\infty}\|\widetilde{\rho}_{0}\|_{2} = o(1). \end{split}$$

The last step is due to the definition of \mathcal{T}_n .

Notice that for any $t \in (0, 1)$,

$$\begin{split} &\partial_t^2 \mathcal{E}\psi(X, D, Y; \tau, \zeta + t(\widetilde{\zeta} - \zeta)) \\ &= -2\Big(\mathcal{E}[D(\widetilde{\mu}_1(X) - \mu_1(X))(\widetilde{\rho}_1(X) - \rho_1(X))] - \mathcal{E}[(1 - D)(\widetilde{\mu}_0(X) - \mu_0(X))(\widetilde{\rho}_0(X) - \rho_0(X))]\Big). \end{split}$$
 Then from the definition of \mathcal{T}_n ,

 $|\partial_t^2 \mathcal{E}\psi(X, D, Y; \tau, \zeta + t(\tilde{\zeta} - \zeta))| \le 2[\|\tilde{\mu}_1 - \mu_1\|_{\infty} \|\tilde{\rho}_1 - \rho_1\|_1 + \|\tilde{\mu}_0 - \mu_0\|_{\infty} \|\tilde{\rho}_0 - \rho_0\|_1] = o(n^{-1/2}).$ We then complete this step and thus complete the proof.

Proof of Theorem 5.1(ii), consistency of $\hat{\sigma}^2$. We decompose $\hat{\sigma}^2$ as

$$\begin{split} \widehat{\sigma}^2 &- \sigma^2 = \frac{1}{n} \sum_{i=1}^n \left[\widehat{\mu}_1(X_i) - \widehat{\mu}_0(X_i) + D_i \left(1 + \frac{K_M^1(i)}{M} \right) \widehat{R}_i - (1 - D_i) \left(1 + \frac{K_M^0(i)}{M} \right) \widehat{R}_i - \widehat{\tau}_M^{\rm bc} \right]^2 \\ &= \left\{ \frac{1}{n} \sum_{i=1}^n \left[\widehat{\mu}_1(X_i) - \widehat{\mu}_0(X_i) + D_i \left(1 + \frac{K_M^1(i)}{M} \right) \widehat{R}_i - (1 - D_i) \left(1 + \frac{K_M^0(i)}{M} \right) \widehat{R}_i - \widehat{\tau}_M^{\rm bc} \right]^2 \\ &- \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + D_i \left(1 + \frac{K_M^1(i)}{M} \right) (Y_i - \mu_1(X_i)) - (1 - D_i) \left(1 + \frac{K_M^0(i)}{M} \right) (Y_i - \mu_0(X_i)) - \widehat{\tau}_M^{\rm bc} \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + D_i \left(1 + \frac{K_M^1(i)}{M} \right) (Y_i - \mu_1(X_i)) - (1 - D_i) \left(1 + \frac{K_M^0(i)}{M} \right) (Y_i - \mu_0(X_i)) - \widehat{\tau}_M^{\rm bc} \right]^2 \right\} \\ &- \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \widehat{\tau}_M^{\rm bc} \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \widehat{\tau}_M^{\rm bc} \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \widehat{\tau}_M^{\rm bc} \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \tau \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \tau \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \tau \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \tau \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \tau \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i)} (Y_i - \mu_0(X_i)) - \tau \right]^2 \right\} \\ &+ \left\{ \frac{1}{n} \sum_{i=1}^n \left[\mu_1(X_i) - \mu_0(X_i) + \frac{D_i}{e(X_i)} (Y_i - \mu_1(X_i)) - \frac{1 - D_i}{1 - e(X_i$$

 $=:T_1 + T_2 + T_3 + T_4.$

By Assumption 5.4, Assumption 5.1, Theorem 4.2, as well as the fact that $\hat{\tau}_M^{\text{bc}} = O_P(1)$, we have $T_1 = o_P(1)$. By Assumption 5.1, Theorem 4.2, and $\hat{\tau}_M^{\text{bc}} = O_P(1)$, we have $T_2 = o_P(1)$. By Assumption 5.1 and $\hat{\tau}_M^{\text{bc}} - \tau = o_P(1)$, we have $T_3 = o_P(1)$. By the fact that $[(X_i, D_i, Y_i)]_{i=1}^n$ are i.i.d., Assumption 5.1 and the weak law of large numbers, we have $T_4 = o_P(1)$. Combining the above four facts together then completes the proof.

A.3.3 Proof of Theorem 5.2

Proof of Theorem 5.2. For $\omega \in \{0, 1\}$, let $j_M^{\omega}(i)$ represent the index of Mth-NN of X_i in $\{X_j : D_j = \omega\}_{j=1}^n$, i.e., the index $j \in [n]$ such that $D_j = \omega$ and

$$\sum_{\ell=1, D_{\ell}=\omega}^{n} \mathbb{1}\left(\|X_{\ell} - X_{i}\| \le \|X_{j} - X_{i}\| \right) = M.$$

With a little abuse of notation, let $\epsilon_i = Y_i - \mu_{D_i}(X_i)$ for any $i \in [n]$. Checking the definition of $\hat{\tau}_M^{\rm bc}$ in (5.2), we decompose $\hat{\tau}_M^{\rm bc}$ to be

$$\begin{split} \hat{\tau}_{M}^{\text{bc}} &= \hat{\tau}^{\text{reg}} + \frac{1}{n} \Big[\sum_{i=1,D_{i}=1}^{n} \Big(1 + \frac{K_{M}^{1}(i)}{M} \Big) \hat{R}_{i} - \sum_{i=1,D_{i}=0}^{n} \Big(1 + \frac{K_{M}^{0}(i)}{M} \Big) \hat{R}_{i} \Big] \\ &= \frac{1}{n} \sum_{i=1}^{n} \Big[\hat{\mu}_{1}(X_{i}) - \hat{\mu}_{0}(X_{i}) \Big] + \frac{1}{n} \sum_{i=1}^{n} (2D_{i} - 1) \Big(1 + \frac{K_{M}^{D_{i}}(i)}{M} \Big) \Big(Y_{i} - \hat{\mu}_{D_{i}}(X_{i}) \Big) \\ &= \frac{1}{n} \sum_{i=1}^{n} \Big[\mu_{1}(X_{i}) - \mu_{0}(X_{i}) \Big] + \frac{1}{n} \sum_{i=1}^{n} (2D_{i} - 1) \Big(1 + \frac{K_{M}^{D_{i}}(i)}{M} \Big) \hat{\epsilon}_{i} \\ &+ \frac{1}{n} \sum_{i=1}^{n} (2D_{i} - 1) \Big[\frac{1}{M} \sum_{m=1}^{M} \Big(\mu_{1-D_{i}}(X_{i}) - \mu_{1-D_{i}}(X_{j_{m}^{1-D_{i}}(i)}) \Big) \Big] \\ &- \frac{1}{n} \sum_{i=1}^{n} (2D_{i} - 1) \Big[\frac{1}{M} \sum_{m=1}^{M} \Big(\hat{\mu}_{1-D_{i}}(X_{i}) - \hat{\mu}_{1-D_{i}}(X_{j_{m}^{1-D_{i}}(i)}) \Big) \Big] \\ &:= \bar{\tau}(\mathbf{X}) + E_{M} + B_{M} - \hat{B}_{M}. \end{split}$$

We have the following central limit theorem on $\bar{\tau}(\mathbf{X}) + E_M$.

Lemma A.1.

$$\sqrt{n}\sigma^{-1/2}\Big(\bar{\tau}(\boldsymbol{X})+E_M-\tau\Big) \stackrel{\mathsf{d}}{\longrightarrow} N\Big(0,1\Big).$$

For the bias term $B_M - \hat{B}_M$, define

$$U_{m,i} := X_{j_m^{1-D_i}(i)} - X_i \quad \text{for any } i \in \llbracket n \rrbracket \text{ and } m \in \llbracket M \rrbracket.$$

We then have the following lemma bounding the moments of $U_{M,i}$.

Lemma A.2. Let p be any positive integer. Then there exists a constant $C_p > 0$ only depending on

p such that for any $i \in [n]$,

$$\mathbf{E}\left[\left\|U_{M,i}\right\|^{p} \mid \boldsymbol{D}\right] \leq C_{p}\left(\frac{M}{n_{1-D_{i}}}\right)^{p/d}.$$

In light of the smoothness conditions on μ_{ω} and approximation conditions on $\hat{\mu}_{\omega}$ for $\omega \in \{0, 1\}$, we can establish the following lemma using Lemma A.2.

Lemma A.3.

$$\sqrt{n} \Big(B_M - \widehat{B}_M \Big) \stackrel{\mathsf{p}}{\longrightarrow} 0.$$

Combining Lemma A.1 and Lemma A.3 completes the proof. The proof of the consistency of the variance estimator is the same as Theorem 5.1(ii) since only Assumption 5.1 and Assumption 5.4 are needed.

A.3.4 Proof of Lemma A.1

Proof of Lemma A.1. For any $x \in \mathbb{X}$, define $\sigma_{\omega}^2(x) = \mathbb{E}[[Y(\omega) - \mu_{\omega}(X)]^2 | X = x]$ for $\omega \in \{0, 1\}$. Let

$$V^{\tau} := \mathbf{E} \Big[\mu_1(X) - \mu_0(X) - \tau \Big]^2 \quad \text{and} \quad V^E := \frac{1}{n} \sum_{i=1}^n \Big(1 + \frac{K_M^{D_i}(i)}{M} \Big)^2 \sigma_{D_i}^2(X_i).$$

From the standard central limit theorem, we have

$$\sqrt{n} \Big(\bar{\tau}(\boldsymbol{X}) - \tau \Big) \stackrel{\mathsf{d}}{\longrightarrow} N \Big(0, V^{\tau} \Big).$$
 (A.12)

Let $E_{M,i} = (2D_i - 1)(1 + K_M^{D_i}(i)/M)\epsilon_i$ for any $i \in [n]$. Conditional on $\mathbf{X}, \mathbf{D}, [E_{M,i}]_{i=1}^n$ are independent. Notice that $\mathbb{E}[E_{M,i} | \mathbf{X}, \mathbf{D}] = 0$ and $\sum_{i=1}^n \operatorname{Var}[E_{M,i} | \mathbf{X}, \mathbf{D}] = nV^E$. To apply the Lindeberg-Feller central limit theorem (Billingsley, 1995, Theorem 27.2), it suffices to verify that: for a given (\mathbf{X}, \mathbf{D}) ,

$$\frac{1}{nV^E} \sum_{i=1}^{n} \mathbb{E}\left[\left(E_{M,i}\right)^2 \mathbb{1}\left(|E_{M,i}| > \delta\sqrt{nV^E}\right) \mid \boldsymbol{X}, \boldsymbol{D}\right] \to 0,$$

for all $\delta > 0$.

Let $C_{\sigma} := \sup_{x \in \mathbb{X}, \omega \in \{0,1\}} \{ \mathbb{E}[|U_{\omega}|^{2+\kappa} | X = x] \lor \mathbb{E}[U_{\omega}^2 | X = x] \} < \infty$. Let $p_1 := 1 + \kappa/2$ and p_2 be the constant such that $p_1^{-1} + p_2^{-1} = 1$. From Hölder's inequality and Markov's inequality,

$$\frac{1}{nV^{E}} \sum_{i=1}^{n} \mathbb{E}\left[\left(E_{M,i}\right)^{2} \mathbb{1}\left(|E_{M,i}| > \delta\sqrt{nV^{E}}\right) \mid \boldsymbol{X}, \boldsymbol{D}\right] \\
\leq \frac{1}{nV^{E}} \sum_{i=1}^{n} \left(\mathbb{E}\left[\left|E_{M,i}\right|^{2+\kappa} \mid \boldsymbol{X}, \boldsymbol{D}\right]\right)^{1/p_{1}} \left(\mathbb{P}\left(|E_{M,i}| > \delta\sqrt{nV^{E}} \mid \boldsymbol{X}, \boldsymbol{D}\right)\right)^{1/p_{2}} \\
\leq \frac{1}{nV^{E}} \sum_{i=1}^{n} \left(\mathbb{E}\left[\left|E_{M,i}\right|^{2+\kappa} \mid \boldsymbol{X}, \boldsymbol{D}\right]\right)^{1/p_{1}} \left(\frac{1}{\delta^{2}nV^{E}} \mathbb{E}\left[\left(E_{M,i}\right)^{2} \mid \boldsymbol{X}, \boldsymbol{D}\right]\right)^{1/p_{2}} \\
\leq \frac{C_{\sigma}}{nV^{E}} \left(\frac{1}{\delta^{2}nV^{E}}\right)^{1/p_{2}} \sum_{i=1}^{n} \left(1 + \frac{K_{M}^{D_{i}}(i)}{M}\right)^{2(1+1/p_{2})}.$$

Notice that $E[1 + K_M^{D_i}(i)/M]^{2(1+1/p_2)} < \infty$ from Theorem 4.2. Let $c_{\sigma} = \inf_{x \in \mathbb{X}, \omega \in \{0,1\}} E[U_{\omega}^2 | X = x] > 0$. From the definition of V^E , we have $V^E \ge c_{\sigma}$ for almost all X, D. Then

$$\mathbf{E}\left[\frac{1}{nV^E}\sum_{i=1}^{n}\mathbf{E}\left[\left(E_{M,i}\right)^2\mathbb{1}\left(|E_{M,i}| > \delta\sqrt{nV^E}\right) \mid \boldsymbol{X}, \boldsymbol{D}\right]\right] = O(n^{-1/p_2}) = o(1).$$

We thus obtain

$$\frac{1}{nV^E} \sum_{i=1}^{n} \mathbb{E}\left[\left(E_{M,i}\right)^2 \mathbb{1}\left(|E_{M,i}| > \delta\sqrt{nV^E}\right) \mid \boldsymbol{X}, \boldsymbol{D}\right] = o_{\mathrm{P}}(1).$$

Employing the Lindeberg-Feller central limit theorem then yields

$$\sqrt{n}(V^E)^{-1/2}E_M = \left(nV^E\right)^{-1/2}\sum_{i=1}^n E_{M,i} \xrightarrow{\mathsf{d}} N(0,1).$$
 (A.13)

Noticing that $\sqrt{n}(\bar{\tau}(\mathbf{X}) - \tau)$ and $\sqrt{n}(V^E)^{-1/2}E_M$ are asymptotically independent, leveraging the same argument as made in Abadie and Imbens (2006, Proof of Theorem 4, Page 267) and then combining (A.12) and (A.13) reaches

$$\sqrt{n} \left(V^{\tau} + V^{E} \right)^{-1/2} \left(\bar{\tau}(\boldsymbol{X}) + E_{M} - \tau \right) \xrightarrow{\mathsf{d}} N(0, 1).$$
(A.14)

We decompose V^E as:

$$V^{E} = \frac{1}{n} \sum_{i=1,D_{i}=1}^{n} \left(1 + \frac{K_{M}^{1}(i)}{M}\right)^{2} \sigma_{1}^{2}(X_{i}) + \frac{1}{n} \sum_{i=1,D_{i}=0}^{n} \left(1 + \frac{K_{M}^{0}(i)}{M}\right)^{2} \sigma_{0}^{2}(X_{i})$$

$$= \left[\frac{1}{n} \sum_{i=1,D_{i}=1}^{n} \left(\frac{1}{e(X_{i})}\right)^{2} \sigma_{1}^{2}(X_{i}) + \frac{1}{n} \sum_{i=1,D_{i}=0}^{n} \left(\frac{1}{1 - e(X_{i})}\right)^{2} \sigma_{0}^{2}(X_{i})\right]$$

$$+ \frac{1}{n} \sum_{i=1,D_{i}=1}^{n} \left[\left(1 + \frac{K_{M}^{1}(i)}{M}\right)^{2} - \left(\frac{1}{e(X_{i})}\right)^{2}\right] \sigma_{1}^{2}(X_{i})$$

$$+ \frac{1}{n} \sum_{i=1,D_{i}=0}^{n} \left[\left(\frac{1}{1 - e(X_{i})}\right)^{2} - \left(1 + \frac{K_{M}^{0}(i)}{M}\right)^{2}\right] \sigma_{0}^{2}(X_{i}). \quad (A.15)$$

For the first term in (A.15), notice that $[(X_i, D_i, Y_i)]_{i=1}^n$ are i.i.d. and $\mathbb{E}[D_i(e(X_i))^{-2}\sigma_1^2(X_i)], \mathbb{E}[(1-D_i)(1-e(X_i))^{-2}\sigma_0^2(X_i)] < \infty$. Using the weak law of large numbers (Durrett, 2019, Theorem 2.2.14), we have

$$\frac{1}{n} \sum_{i=1,D_i=1}^n \left(\frac{1}{e(X_i)}\right)^2 \sigma_1^2(X_i) + \frac{1}{n} \sum_{i=1,D_i=0}^n \left(\frac{1}{1-e(X_i)}\right)^2 \sigma_0^2(X_i) \xrightarrow{\mathsf{p}} \mathbf{E}\Big[\frac{\sigma_1^2(X)}{e(X)} + \frac{\sigma_0^2(X)}{1-e(X)}\Big].$$

For the second term in (A.15), using the Cauchy–Schwarz inequality,

$$E\left|\frac{1}{n}\sum_{i=1,D_{i}=1}^{n} \left[\left(1+\frac{K_{M}^{1}(i)}{M}\right)^{2}-\left(\frac{1}{e(X_{i})}\right)^{2}\right]\sigma_{1}^{2}(X_{i})\right| \\ \leq C_{\sigma} E\left[D_{i}\left|\left(1+\frac{K_{M}^{1}(i)}{M}\right)^{2}-\left(\frac{1}{e(X_{i})}\right)^{2}\right|\right] = C_{\sigma} E\left[D_{i} E\left[\left|\left(1+\frac{K_{M}^{1}(i)}{M}\right)^{2}-\left(\frac{1}{e(X_{i})}\right)^{2}\right|\right| \boldsymbol{D}\right]\right]$$

$$\leq C_{\sigma} \mathbb{E} \Big[D_i \Big(\mathbb{E} \Big[\Big(\frac{K_M^1(i)}{M} - \frac{1 - e(X_i)}{e(X_i)} \Big)^2 \Big| \mathbf{D} \Big] \mathbb{E} \Big[\Big(2 + \frac{K_M^1(i)}{M} + \frac{1 - e(X_i)}{e(X_i)} \Big)^2 \Big| \mathbf{D} \Big] \Big)^{1/2} \Big] = o(1),$$

where the last step is due to Theorem 4.2. Then we obtain

$$\frac{1}{n}\sum_{i=1,D_i=1}^n \left[\left(1 + \frac{K_M^1(i)}{M}\right)^2 - \left(\frac{1}{e(X_i)}\right)^2 \right] \sigma_1^2(X_i) \stackrel{\mathsf{p}}{\longrightarrow} 0.$$

For the third term in (A.15), we can establish in the same way that

$$\frac{1}{n} \sum_{i=1,D_i=0}^{n} \left[\left(\frac{1}{1-e(X_i)} \right)^2 - \left(1 + \frac{K_M^0(i)}{M} \right)^2 \right] \sigma_0^2(X_i) \stackrel{\mathsf{p}}{\longrightarrow} 0.$$

Then from (A.15),

$$V^E \xrightarrow{\mathbf{p}} \mathbf{E}\Big[\frac{\sigma_1^2(X)}{e(X)} + \frac{\sigma_0^2(X)}{1 - e(X)}\Big].$$

Using Slutsky's lemma (van der Vaart, 1998, Theorem 2.8) and then the definition of σ^2 , we complete the proof by (A.14).

A.3.5 Proof of Lemma A.2

Proof of Lemma A.2. From Assumption 4.1 and Assumption 5.1, let $R := \text{diam}(\mathbb{X}) < \infty$ and $f_L := \inf_{x \in \mathbb{X}, \omega \in \{0,1\}} f_{\omega}(x) > 0$. For any $x \in \mathbb{X}, \omega \in \{0,1\}$ and $u \leq R$, from Assumption 4.1,

$$\nu_{\omega}(B_{x,u} \cap \mathbb{X}) \ge f_L \lambda(B_{x,u} \cap \mathbb{X}) \ge f_L a \lambda(B_{x,u}) = f_L a V_d u^d.$$

Let $c_0 = f_L a V_d$. For any $i \in [n]$, $x \in \mathbb{X}$, $M \leq n_{1-D_i}$, if $0 \leq u \leq R n_{1-D_i}^{1/d}$, we have

$$P\left(\left\|X_{j}-X_{i}\right\| \geq un_{1-D_{i}}^{-1/d} \mid \boldsymbol{D}, X_{i}=x, j=j_{M}^{1-D_{i}}(i)\right)$$

$$\leq P\left(Bin\left(n_{1-D_{i}}, \nu_{1-D_{i}}\left(B_{x,un_{1-D_{i}}^{-1/d}} \cap \mathbb{X}\right)\right) \leq M \mid \boldsymbol{D}\right)$$

$$\leq P\left(Bin\left(n_{1-D_{i}}, c_{0}\left(un_{1-D_{i}}^{-1/d}\right)^{d}\right) \leq M \mid \boldsymbol{D}\right)$$

$$= P\left(Bin\left(n_{1-D_{i}}, c_{0}u^{d}n_{1-D_{i}}^{-1}\right) \leq M \mid \boldsymbol{D}\right).$$

Using the Chernoff bound, if $M < c_0 u^d$, then

$$\mathbb{P}\left(\operatorname{Bin}\left(n_{1-D_{i}}, c_{0}u^{d}n_{1-D_{i}}^{-1}\right) \leq M \mid \boldsymbol{D}\right) \leq \exp\left(M - c_{0}u^{d} + M\log\left(\frac{c_{0}u^{a}}{M}\right)\right).$$

.1

Notice that the above upper bound does not depend on x. We then obtain

$$\begin{aligned} & \operatorname{P}\left(\left\|X_{j} - X_{i}\right\| \geq u n_{1-D_{i}}^{-1/d} \left| \boldsymbol{D}, j = j_{M}^{1-D_{i}}(i)\right) \\ & \leq \mathbb{1}\left(M < c_{0} u^{d}\right) \exp\left(M - c_{0} u^{d} + M \log\left(\frac{c_{0} u^{d}}{M}\right)\right) + \mathbb{1}\left(M \geq c_{0} u^{d}\right). \end{aligned}$$

On the other hand, if $u > Rn_{1-D_i}^{1/d}$, then the probability is zero from the definition of R. Accordingly, the above bound holds for any $u \ge 0$.

For any $i \in [n]$, we thus have

$$n_{1-D_{i}}^{p/d} \mathbb{E}\left[\left\|U_{M,i}\right\|^{p} \middle| \mathbf{D}\right] = p \int_{0}^{\infty} \mathbb{P}\left(\left\|X_{j} - X_{i}\right\| \ge u n_{1-D_{i}}^{-1/d} \middle| \mathbf{D}, j = j_{M}^{1-D_{i}}(i)\right) u^{p-1} du$$

$$\le p \int_{0}^{\infty} \left[\mathbb{1}\left(M < c_{0}u^{d}\right) \exp\left(M - c_{0}u^{d} + M \log\left(\frac{c_{0}u^{d}}{M}\right)\right) + \mathbb{1}\left(M \ge c_{0}u^{d}\right)\right] u^{p-1} du$$

$$= p c_{0}^{-p/d} d^{-1} \left[\int_{M}^{\infty} \left(\frac{e}{M}\right)^{M} t^{M+\frac{p}{d}-1} e^{-t} dt + \int_{0}^{M} t^{\frac{p}{d}-1} dt\right],$$
(A.16)

where the last step is through taking $t = c_0 u^d$.

For the first term in (A.16), from Stirling's formula and $M \to \infty$,

$$\int_{M}^{\infty} \left(\frac{e}{M}\right)^{M} t^{M+\frac{p}{d}-1} e^{-t} dt \leq \int_{0}^{\infty} \left(\frac{e}{M}\right)^{M} t^{M+\frac{p}{d}-1} e^{-t} dt = \left(\frac{e}{M}\right)^{M} \Gamma\left(M+\frac{p}{d}\right)$$
$$\sim \left(\frac{e}{M}\right)^{M} (M+1)^{\frac{p}{d}-1} \Gamma(M+1) \sim \left(\frac{e}{M}\right)^{M} (M+1)^{\frac{p}{d}-1} \sqrt{2\pi M} \left(\frac{M}{e}\right)^{M} \sim \sqrt{2\pi} M^{\frac{p}{d}-\frac{1}{2}},$$

where \sim means asymptotic convergence.

For the second term in (A.16),

$$\int_0^M t^{\frac{p}{d}-1} \mathrm{d}t = \frac{d}{p} M^{\frac{p}{d}}.$$

Combining the above two bounds then complete the proof.

A.3.6 Proof of Lemma A.3

Proof of Lemma A.3. We decompose $B_M - \widehat{B}_M$ as

$$|B_{M} - \widehat{B}_{M}|$$

$$= \left|\frac{1}{n}\sum_{i=1}^{n} (2D_{i} - 1) \left[\frac{1}{M}\sum_{m=1}^{M} \left(\mu_{1-D_{i}}(X_{i}) - \mu_{1-D_{i}}(X_{j_{m}^{1-D_{i}}(i)}) - \widehat{\mu}_{1-D_{i}}(X_{i}) + \widehat{\mu}_{1-D_{i}}(X_{j_{m}^{1-D_{i}}(i)})\right)\right]\right|$$

$$\leq \max_{i \in \llbracket n \rrbracket, m \in \llbracket M \rrbracket} \left|\mu_{1-D_{i}}(X_{i}) - \mu_{1-D_{i}}(X_{j_{m}^{1-D_{i}}(i)}) - \widehat{\mu}_{1-D_{i}}(X_{i}) + \widehat{\mu}_{1-D_{i}}(X_{j_{m}^{1-D_{i}}(i)})\right|$$

$$\leq \max_{i \in \llbracket n \rrbracket, m \in \llbracket M \rrbracket, \omega \in \{0,1\}} \left|\mu_{\omega}(X_{i}) - \mu_{\omega}(X_{j_{m}^{1-D_{i}}(i)}) - \widehat{\mu}_{\omega}(X_{i}) + \widehat{\mu}_{\omega}(X_{j_{m}^{1-D_{i}}(i)})\right|. \tag{A.17}$$

Let $k = \lfloor d/2 \rfloor + 1$. For any $\omega \in \{0, 1\}$, from Taylor expansion to kth order,

$$\left|\mu_{\omega}(X_{j_{m}^{1-D_{i}}(i)}) - \mu_{\omega}(X_{i}) - \sum_{\ell=1}^{k-1} \frac{1}{\ell!} \sum_{t \in \Lambda_{\ell}} \partial^{t} \mu_{\omega}(X_{i}) U_{m,i}^{t}\right| \leq \max_{t \in \Lambda_{k}} \|\partial^{t} \mu_{\omega}\|_{\infty} \frac{1}{k!} \sum_{t \in \Lambda_{k}} \|U_{m,i}\|^{k}.$$
(A.18)

In the same way,

$$\left|\widehat{\mu}_{\omega}(X_{j_{m}^{1-D_{i}}(i)}) - \widehat{\mu}_{\omega}(X_{i}) - \sum_{\ell=1}^{k-1} \frac{1}{\ell!} \sum_{t \in \Lambda_{\ell}} \partial^{t} \widehat{\mu}_{\omega}(X_{i}) U_{m,i}^{t} \right| \leq \max_{t \in \Lambda_{k}} \|\partial^{t} \widehat{\mu}_{\omega}\|_{\infty} \frac{1}{k!} \sum_{t \in \Lambda_{k}} \|U_{m,i}\|^{k}.$$
(A.19)

We also have

$$\left|\sum_{\ell=1}^{k-1} \frac{1}{\ell!} \sum_{t \in \Lambda_{\ell}} (\partial^t \widehat{\mu}_{\omega}(X_i) - \partial^t \mu_{\omega}(X_i)) U_{m,i}^t \right| \le \sum_{\ell=1}^{k-1} \max_{t \in \Lambda_{\ell}} \|\partial^t \widehat{\mu}_{\omega} - \partial^t \mu_{\omega}\|_{\infty} \frac{1}{\ell!} \sum_{t \in \Lambda_{\ell}} \|U_{m,i}\|^{\ell}.$$
(A.20)

Notice that $||U_{M,i}|| = \max_{m \in \llbracket M \rrbracket} ||U_{m,i}||$ for any $i \in \llbracket n \rrbracket, \omega \in \{0, 1\}$. Then for any $\omega \in \{0, 1\}$, plugging (A.18), (A.19), (A.20) into (A.17), we obtain

$$|B_M - \widehat{B}_M| \lesssim (1 + O_{\mathrm{P}}(1)) \max_{i \in \llbracket n \rrbracket} \|U_{M,i}\|^k + \max_{\ell \in \llbracket k - 1 \rrbracket} \max_{t \in \Lambda_\ell} \|\partial^t \widehat{\mu}_\omega - \partial^t \mu_\omega\|_\infty \max_{i \in \llbracket n \rrbracket} \|U_{M,i}\|^\ell.$$

From Lemma A.2, all moments of $(n/M)^{p/d} ||U_{M,i}||^p$ are bounded. Then for any positive integer p and any $\varepsilon > 0$, we have

$$\max_{i \in [n]} \|U_{M,i}\|^p = o_{\mathrm{P}}\left(n^{\varepsilon} \left(\frac{M}{n}\right)^{p/d}\right)$$

We then obtain

$$B_M - \widehat{B}_M = o_{\mathrm{P}}\left(n^{\varepsilon} \left(\frac{M}{n}\right)^{k/d}\right) + \max_{\ell \in [k-1]} o_{\mathrm{P}}\left(n^{-\gamma_{\ell}} n^{\varepsilon} \left(\frac{M}{n}\right)^{\ell/d}\right)$$

The proof is thus complete by noticing the definition of γ and $M \lesssim n^{\gamma}$.

A.4 Proofs of the results in Section 6

Before starting the proof, we first introduce some necessary notation. Given two samples $\{X_i\}_{i=1}^{N_0}$ and $\{Z_j\}_{j=1}^{N_1}$, for any $M \in [\![N_0]\!]$ and $\ell \in [\![N_1]\!]$, define $j_M(\ell)$ to be the index of the *M*-th NN of Z_ℓ in $\{X_i\}_{i=1}^{N_0}$. In other words, $j_M(\ell)$ is the index of $k \in [\![N_0]\!]$ such that

$$\sum_{i=1}^{N_0} \mathbb{1}(\|X_i - Z_\ell\| \le \|X_k - Z_\ell\|) = M.$$

Further define $\mathcal{J}_M(\ell)$ to be the set of indices of all *M*-NNs of Z_ℓ in $\{X_i\}_{i=1}^{N_0}$, i.e.,

$$\mathcal{J}_M(\ell) := \{ j_m(\ell) : m \in \llbracket M \rrbracket \}.$$

A.4.1 Proof of Theorem 6.1

Proof of Theorem 6.1. Before starting the proof, we first introduce the following lemma about the truncation level for f_1 to attain the desirable estimation accuracy.

Lemma A.4. There exists a constant $C_0 > 0$ only depending on f_L, f_U, L, d such that for any $x \in \mathbb{R}^d$ with

$$f_1(x) \ge C_0(\frac{M}{N_0})^{1/d}$$
 and $\Delta(x) \ge C_0(\frac{M}{N_0})^{1/d}$ (A.21)

we have, for all N_0 sufficiently large,

- (i) $\left| \mathbb{E} \left[\widehat{r}_M(x) \right] r(x) \right| \leq \frac{1}{2} r(x);$
- (*ii*) Var $\left[\mathbb{E} \left[\widehat{r}_M(x) \mid \boldsymbol{X} \right] \right] \leq \frac{32}{M} [r(x)]^2;$
- (iii) with probability one,

$$\frac{N_0}{M}\nu_1(A_M(x)) \ge C_1 r(x),$$

where $C_1 > 0$ is a constant only depending on f_L, f_U, d .

Let X be drawn from ν_0 independent of the data. We then have

$$D_{\phi}(\nu_1 \| \nu_0) = \int \phi(r(x)) f_0(x) dx = \mathbb{E}[\phi(r(X))].$$

Let \mathcal{E} be the set of all $x \in \mathbb{R}^d$ satisfying (A.21) in Lemma A.4. We then decompose \widehat{D}_{ϕ} as

$$\begin{split} \widehat{D}_{\phi} - D_{\phi}(\nu_{1} \| \nu_{0}) &= \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \phi(\widehat{r}_{M}(x)) - \mathbf{E}[\phi(r(X))] \\ &= \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \left[\phi\Big(\widehat{r}_{M}(X_{i})\Big) - \phi\Big(\mathbf{E}[\widehat{r}_{M}(X_{i}) \mid \mathbf{X}]\Big) \right] \\ &+ \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \left[\phi\Big(\mathbf{E}[\widehat{r}_{M}(X_{i}) \mid \mathbf{X}]\Big) - \phi\Big(\mathbf{E}[\widehat{r}_{M}(X_{i}) \mid X_{i}]\Big) \right] \mathbb{1}\Big(X_{i} \in \mathcal{E}\Big) \\ &+ \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \left[\phi\Big(\mathbf{E}[\widehat{r}_{M}(X_{i}) \mid \mathbf{X}]\Big) - \phi\Big(r(X_{i})\Big) \right] \mathbb{1}\Big(X_{i} \in \mathcal{E}\Big) \\ &+ \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \left[\phi\Big(\mathbf{E}[\widehat{r}_{M}(X_{i}) \mid \mathbf{X}]\Big) - \phi\Big(r(X_{i})\Big) \right] \mathbb{1}\Big(X_{i} \notin \mathcal{E}\Big) \\ &+ \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \phi\Big(r(X_{i})\Big) - \mathbf{E}[\phi(r(X))] \\ &=: T_{1} + T_{2} + T_{3} + T_{4} + T_{5}. \end{split}$$

By the Cauchy–Schwarz inequality,

$$\mathbf{E}\Big[\widehat{D}_{\phi} - D_{\phi}(\nu_1 \| \nu_0)\Big]^2 \le 5\Big(\mathbf{E}[T_1]^2 + \mathbf{E}[T_2]^2 + \mathbf{E}[T_3]^2 + \mathbf{E}[T_4]^2 + \mathbf{E}[T_5]^2\Big).$$

We then consider $E[T_1]^2$, $E[T_2]^2$, $E[T_3]^2$, $E[T_4]^2$, $E[T_5]^2$ separately. We have the following lemmas. Lemma A.5. It holds that

$$\mathbf{E}[T_1]^2 \le C \Big[\frac{1}{N_1} + \Big(\frac{N_0}{N_1 M} \Big)^2 \Big],$$

where C > 0 is a constant only depending on f_L, f_U .

Lemma A.6. It holds that

$$\mathbf{E}[T_2]^2 \le C \Big[\frac{1}{N_0} + \left(\frac{1}{M}\right)^2 \Big],$$

where C > 0 is a constant only depending on f_L, f_U, d, f'_U .

Lemma A.7. It holds that

$$\mathbf{E}[T_3]^2 \le C\left(\frac{M}{N_0}\right)^{2/d},$$

where C > 0 is a constant only depending on f_L, f_U, L, d, f'_U .

Lemma A.8. It holds that

$$\mathbf{E}[T_4]^2 \le C\left(\frac{M}{N_0}\right)^{2/d},$$

where C > 0 is a constant only depending on f_L, f_U, a, H, d, f'_U .

Notice that T_5 is a linear combination of the function of i.i.d. random variables with mean zero. From the boundedness assumption on r, it is easy to establish

$$E[T_5]^2 \le C \frac{1}{N_0},$$
 (A.22)

where C > 0 is a constant only depending on f_L, f_U .

Combining Lemma A.5, Lemma A.6, Lemma A.7, Lemma A.8, and (A.22) completes the proof.

A.4.2 Proof of Lemma A.4

Proof of Lemma A.4. For Part (i), recall that in the proof of Theorem 4.3(i), we take $\delta_N = (\frac{4}{f_L V_d})^{1/d} (\frac{M}{N_0})^{1/d}$. Take $C_0 > L(\frac{4}{f_L V_d})^{1/d}$. Then $f_1(x) \ge C_0(\frac{M}{N_0})^{1/d} > L\delta_N$. Take $C_0 > 2(\frac{4}{f_L V_d})^{1/d}$. Then $\Delta(x) > 2\delta_N$. Then for any $z \in \mathbb{R}^d$ such that $||z - x|| \le \delta_N$, we have $B_{z,||x-z||} \subset S_1$. We take N_0 large enough so that $\delta_N < f_L/(4L)$. Then from the proof of Theorem 4.3(i), $W \le 2\frac{M}{N_0}$ implies $||Z - x|| \le \delta_N$. From Case I in the proof of Theorem 4.3(i), we use (7.31) and (7.32), and take N_0 large enough so that $L\delta_N \le f_U \land (f_L/4)$. We then have

$$\begin{aligned} \left| \mathbf{E} \left[\widehat{r}_{M}(x) \right] - r(x) \right| &\leq \left| \frac{f_{1}(x) + L\delta_{N}}{f_{0}(x) - 2L\delta_{N}} \frac{N_{0}}{N_{0} + 1} - \frac{f_{1}(x)}{f_{0}(x)} \right| \bigvee \left| \frac{f_{1}(x) - L\delta_{N}}{f_{0}(x) + 2L\delta_{N}} \frac{N_{0}}{N_{0} + 1} - \frac{f_{1}(x)}{f_{0}(x)} \right| + o(N_{0}^{-\gamma}) \\ &\leq \frac{f_{0}(x)L\delta_{N} + 2f_{1}(x)L\delta_{N}}{f_{0}(x)(f_{0}(x) - 2L\delta_{N})} + \frac{1}{N_{0} + 1} \frac{f_{1}(x) + L\delta_{N}}{f_{0}(x) - 2L\delta_{N}} + o(N_{0}^{-\gamma}) \\ &\leq \frac{2}{f_{L}}L\delta_{N} + r(x)\frac{4}{f_{L}}L\delta_{N} + \frac{4f_{U}}{f_{L}}\frac{1}{N_{0} + 1} + o(N_{0}^{-\gamma}). \end{aligned}$$

Taking $C_0 > L \frac{4}{f_L} (\frac{4}{f_L V_d})^{1/d}$, Part (i) then holds when N_0 is sufficiently large.

For Part (ii), we take C_0 large enough as in Part (i). Then $W \leq 2\frac{M}{N_0}$ implies $||Z - x|| \leq \delta_N$. For any $\epsilon > 0$, we take C_0 and N_0 sufficiently large such that for any $||z - x|| \leq 2\delta_N$, we have $|f_1(z) - f_1(x)| \leq \epsilon C_0(\frac{M}{N_0})^{1/d}$ and $|f_0(z) - f_0(x)| \leq \epsilon f_L$, which is achievable from the Lipschitz condition. Then $|f_1(z) - f_1(x)| \leq \epsilon f_1(x)$ and $|f_0(z) - f_0(x)| \leq \epsilon f_0(x)$. For any $x \in \mathbb{R}^d$ such that $f_1(x) \geq C_0(\frac{M}{N_0})^{1/d}$ and $\Delta(x) \geq C_0(\frac{M}{N_0})^{1/d}$, for N_0 sufficiently large, we have for all $0 \leq w \leq w + t \leq 2M/N_0$,

$$P\left(w \le W \le w + t\right) = \nu_1\left(\left\{z \in \mathbb{R}^d : \nu_0(B_{z,\|x-z\|}) \in [w, w+t]\right\}\right)$$

$$\leq \frac{f_1(x)(1+\epsilon)}{f_0(x)(1-\epsilon)}\nu_0\left(\left\{z \in \mathbb{R}^d : \nu_0(B_{z,\|x-z\|}) \in [w, w+t]\right\}\right),$$

and then

$$\limsup_{t \to 0} t^{-1} \mathbf{P} \left(w \le W \le w + t \right) \le r(x) \frac{1 + \epsilon}{1 - \epsilon} (1 + \epsilon).$$

Proceeding in the same way as the proof of Theorem 4.3(ii) and Lemma 7.4, we prove Part (ii).

On Part (iii), we cite a result on lower bounding the M-NN distance without proof.

Lemma A.9. (*Biau and Devroye, 2015, Theorem 4.1*) Let $X_1, \ldots, X_{N_0} \in \mathbb{R}^d$ be N_0 independent copies from a probability measure with density f_0 . Assume that f_0 is uniformly bounded by a universal

constant f_U . If $M/\log N_0 \to \infty$ as $N_0 \to \infty$, then with probability one, for all N_0 large enough,

$$\inf_{x \in \mathbb{R}^d} \left\| \mathcal{X}_{(M)}(x) - x \right\| \ge C \left(\frac{M}{N_0} \right)^{1/d},$$

where C > 0 is a constant only depending on f_U and d.

Notice that for $z \in \mathbb{R}^d$, if $||z - x|| \leq \frac{1}{2} ||\mathcal{X}_{(M)}(x) - x||$, then $z \in A_M(x)$. From Lemma A.9, with probability one, for all N_0 sufficiently large, we have

$$\frac{N_0}{M}\nu_1(A_M(x)) \ge \frac{N_0}{M}\nu_1(B_{x,\|\mathcal{X}_{(M)}(x) - x\|/2}) \ge \frac{N_0}{M}\nu_1(B_{x,C(M/N_0)^{1/d}/2}),$$

where C > 0 is the constant in Lemma A.9. Taking C_0 large enough so that $C_0 > C/2$ and $C_0 \ge LC$, then $\Delta(x) > C(M/N_0)^{1/d}/2$ and for any $||z - x|| \le C(M/N_0)^{1/d}/2$, we have $f_1(z) \ge \frac{1}{2}f_1(x)$. Then

$$\frac{N_0}{M}\nu_1(A_M(x)) \ge \frac{N_0}{M} \frac{f_1(x)}{2} V_d \left(\frac{C}{2} \left(\frac{M}{N_0}\right)^{\frac{1}{d}}\right)^d = 2^{-(d+1)} C^d V_d f_1(x) \ge 2^{-(d+1)} C^d V_d f_L r(x).$$

We then complete the proof of Part (iii).

A.4.3 Proof of Lemma A.5

Proof of Lemma A.5. Notice that

$$\mathbf{E}[T_1]^2 = \mathbf{E}\left[\mathbf{E}[T_1^2 \mid \boldsymbol{X}]\right] = \mathbf{E}\left[\left(\mathbf{E}[T_1 \mid \boldsymbol{X}]\right)^2 + \operatorname{Var}[T_1 \mid \boldsymbol{X}]\right].$$

We consider the above two terms separately.

Step I. $E[T_1 | X]$.

Let g(x) be a given twice continuously differentiable function on $[0, \infty)$. Equation 2.5 in Totik (1994) gives

$$|g(t) - g(x) - g'(x)(t - x)| \le 4 \frac{(t - x)^2}{x} ||xg''(x)||_{\infty},$$
(A.23)

for all $x, t \in [0, \infty)$.

From (A.23),

$$\begin{split} |\mathbf{E}[T_{1} | \mathbf{X}]| &= \left| \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \mathbf{E} \left[\phi \left(\widehat{r}_{M}(X_{i}) \right) - \phi \left(\mathbf{E}[\widehat{r}_{M}(X_{i}) | \mathbf{X}] \right) \Big| \mathbf{X} \right] \right| \\ &\leq 2 \| \phi - g \|_{\infty} + \left| \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \mathbf{E} \left[g \left(\widehat{r}_{M}(X_{i}) \right) - g \left(\mathbf{E}[\widehat{r}_{M}(X_{i}) | \mathbf{X}] \right) \Big| \mathbf{X} \right] \right| \\ &\leq 2 \| \phi - g \|_{\infty} + 4 \| x g''(x) \|_{\infty} \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \mathbf{E} \left[\frac{1}{\mathbf{E}[\widehat{r}_{M}(X_{i}) | \mathbf{X}]} \left(\widehat{r}_{M}(X_{i}) - \mathbf{E}[\widehat{r}_{M}(X_{i}) | \mathbf{X}] \right)^{2} \Big| \mathbf{X} \right] \\ &= 2 \| \phi - g \|_{\infty} + 4 \| x g''(x) \|_{\infty} \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \frac{\mathrm{Var}[\widehat{r}_{M}(X_{i}) | \mathbf{X}]}{\mathbf{E}[\widehat{r}_{M}(X_{i}) | \mathbf{X}]}. \end{split}$$

From the definition of $\widehat{r}_M(X_i)$,

$$\widehat{r}_M(X_i) = \frac{N_0}{N_1} \frac{K_M(i)}{M} = \frac{N_0}{N_1 M} \sum_{j=1}^{N_1} \mathbb{1} \left(Z_j \in A_M(i) \right).$$

Then conditional on \boldsymbol{X} ,

$$\frac{N_1M}{N_0}\widehat{r}_M(X_i) \,\Big|\, \boldsymbol{X} \sim \operatorname{Bin}(N_1, A_M(i)).$$

It is easy to calculate

$$E[\hat{r}_M(X_i) | \mathbf{X}] = \frac{N_0}{M} \nu_1(A_M(i)), \quad Var[\hat{r}_M(X_i) | \mathbf{X}] = \frac{N_0^2}{N_1 M^2} \nu_1(A_M(i)) \left[1 - \nu_1(A_M(i))\right].$$

Then

$$|\mathbf{E}[T_1 \mid \mathbf{X}]| \le 2\|\phi - g\|_{\infty} + 4\|xg''(x)\|_{\infty} \frac{1}{N_0} \sum_{i=1}^{N_0} \frac{N_0}{N_1 M} \left[1 - \nu_1 \left(A_M(i)\right)\right]$$
$$\le 2\|\phi - g\|_{\infty} + 4\frac{N_0}{N_1 M} \|xg''(x)\|_{\infty}.$$

We then cite the equivalence theorem.

Lemma A.10. (*Ditzian and Totik, 2012, Theorem 2.1.1*) For any real-valued function ϕ on $[0, \infty)$ and t > 0, there exists a universal constant C > 0 such that

$$\inf_{g \in C^2[0,\infty)} \left(\|\phi - g\|_{\infty} + t^2 \|xg''(x)\|_{\infty} \right) \le C\omega_{\varphi}^2(\phi, t)_{\infty},$$

where $\omega_{\varphi}^2(\phi, t)_{\infty}$ is the second-order modulus of smoothness with $\varphi(x) = \sqrt{x}$, and $C^2[0, \infty)$ is the class of all twice continuously differentiable functions on $[0, \infty)$.

From Example 3.4.2 in Ditzian and Totik (2012), when $\phi(x) = x \log x$, we have $\omega_{\varphi}^2(\phi, t)_{\infty} \approx t^2$. Then from Lemma A.10, we obtain

$$|\mathbf{E}[T_1 \mid \boldsymbol{X}]| \lesssim \frac{N_0}{N_1 M}.\tag{A.24}$$

Step II. $\operatorname{Var}[T_1 \mid \boldsymbol{X}]$. Conditional on \boldsymbol{X} ,

$$\operatorname{Var}[T_1 \mid \boldsymbol{X}] = \operatorname{Var}\left[\frac{1}{N_0} \sum_{i=1}^{N_0} \left[\phi\left(\widehat{r}_M(X_i)\right) - \phi\left(\operatorname{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\right)\right] \mid \boldsymbol{X}\right]$$
$$= \operatorname{Var}\left[\frac{1}{N_0} \sum_{i=1}^{N_0} \phi\left(\widehat{r}_M(X_i)\right) \mid \boldsymbol{X}\right].$$
(A.25)

We then apply Efron-Stein inequality (Boucheron et al., 2013, Theorem 3.1). Recall $\mathbf{Z} = (Z_1, \ldots, Z_{N_1})$ and for any $\ell \in [N_1]$, define

$$\boldsymbol{Z}_{\ell} := (Z_1, \ldots, Z_{\ell-1}, \widetilde{Z}_{\ell}, Z_{\ell+1}, \ldots, Z_{N_1}),$$

where $[\widetilde{Z}_{\ell}]_{\ell=1}^{N_1}$ are independent copies of $[Z_{\ell}]_{\ell=1}^{N_1}$.

Fix $\ell \in [N_1]$. Recall that $\mathcal{J}_M(\cdot)$ are defined based on X and Z. We then define $\mathcal{J}'_M(\cdot)$ in the same way but based on X and Z_{ℓ} .

Let \hat{r}'_M be the density ratio estimator based on X and Z_ℓ . Then for any $i \in [N_0]$,

$$\widehat{r}_M(X_i) = \frac{N_0}{N_1 M} \sum_{j=1}^{N_1} \mathbb{1}\left(i \in \mathcal{J}_M(j)\right), \quad \widehat{r}'_M(X_i) = \frac{N_0}{N_1 M} \sum_{j=1}^{N_1} \mathbb{1}\left(i \in \mathcal{J}'_M(j)\right).$$

Notice that for $j \in [\![N_1]\!]$ and $j \neq \ell$, $\mathcal{J}_M(j) = \mathcal{J}'_M(j)$ since both $\mathcal{J}_M(\cdot)$ and $\mathcal{J}'_M(\cdot)$ are based on \mathbf{X} . Then if $i \in \mathcal{J}_M(\ell) \setminus \mathcal{J}'_M(\ell)$ or $i \in \mathcal{J}'_M(\ell) \setminus \mathcal{J}_M(\ell)$, we have $|\hat{r}_M(X_i) - \hat{r}'_M(X_i)| = \frac{N_0}{N_1M}$. If $i \in \mathcal{J}_M(\ell) \cap \mathcal{J}'_M(\ell)$ or $i \notin \mathcal{J}_M(\ell) \cup \mathcal{J}'_M(\ell)$, we have $\hat{r}_M(X_i) = \hat{r}'_M(X_i)$. As a result,

$$\begin{split} & \operatorname{E}\left[\left[\frac{1}{N_{0}}\sum_{i=1}^{N_{0}}\left(\phi\left(\widehat{r}_{M}(X_{i})\right)-\phi\left(\widehat{r}_{M}'(X_{i})\right)\right)\right]^{2} \middle| \mathbf{X}\right] \\ \leq & \operatorname{E}\left[\left[\frac{1}{N_{0}}\sum_{i=1}^{N_{0}}\left|\phi\left(\widehat{r}_{M}(X_{i})\right)-\phi\left(\widehat{r}_{M}'(X_{i})\right)\right|\right]^{2} \middle| \mathbf{X}\right] \\ = & \operatorname{E}\left[\left[\frac{1}{N_{0}}\sum_{i\in(\mathcal{J}_{M}(\ell)\setminus\mathcal{J}_{M}'(\ell)\cup\mathcal{J}_{M}'(\ell)\setminus\mathcal{J}_{M}(\ell))}^{N_{0}}\left|\phi\left(\widehat{r}_{M}(X_{i})\right)-\phi\left(\widehat{r}_{M}'(X_{i})\right)\right|\right]^{2} \middle| \mathbf{X}\right] \\ \leq & \frac{1}{N_{0}^{2}}\operatorname{E}\left[\left|\left(\mathcal{J}_{M}(\ell)\setminus\mathcal{J}_{M}'(\ell)\right)\cup\left(\mathcal{J}_{M}'(\ell)\setminus\mathcal{J}_{M}(\ell)\right)-\phi\left(\widehat{r}_{M}'(X_{i})\right)\right]^{2} \middle| \mathbf{X}\right] \\ & \sum_{i\in(\mathcal{J}_{M}(\ell)\setminus\mathcal{J}_{M}'(\ell))\cup(\mathcal{J}_{M}'(\ell)\setminus\mathcal{J}_{M}(\ell))}^{N_{0}}\left[\phi\left(\widehat{r}_{M}(X_{i})\right)-\phi\left(\widehat{r}_{M}'(X_{i})\right)\right]^{2} \middle| \mathbf{X}\right] \\ \leq & \frac{2M}{N_{0}^{2}}\operatorname{E}\left[\sum_{i\in(\mathcal{J}_{M}(\ell)\setminus\mathcal{J}_{M}'(\ell))\cup(\mathcal{J}_{M}'(\ell)\setminus\mathcal{J}_{M}(\ell))}^{N_{0}}\left[\phi\left(\widehat{r}_{M}(X_{i})\right)-\phi\left(\widehat{r}_{M}'(X_{i})\right)\right]^{2} \middle| \mathbf{X}\right], \end{split}$$

where for any set A, |A| is the cardinality of A. The last two steps are due to the Cauchy–Schwarz inequality and the fact that $|\mathcal{J}_M(\ell)| \leq M, |\mathcal{J}'_M(\ell)| \leq M$. Thus

$$|(\mathcal{J}_M(\ell) \setminus \mathcal{J}'_M(\ell)) \bigcup (\mathcal{J}'_M(\ell) \setminus \mathcal{J}_M(\ell))| \le 2M.$$

Since $[X_i]_{i=1}^{N_0}$ i.i.d., we have

where the last step is from symmetry.

Let

$$\widehat{r}_M^{-\ell}(X_i) = \frac{N_0}{N_1 M} \sum_{j=1, j \neq \ell}^{N_1} \mathbb{1}\left(i \in \mathcal{J}_M(j)\right).$$

Then $\widehat{r}_M(X_i) = \widehat{r}_M^{-\ell}(X_i) + \frac{N_0}{N_1M} \mathbb{1}\left(i \in \mathcal{J}_M(\ell)\right), \widehat{r}'_M(X_i) = \widehat{r}_M^{-\ell}(X_i) + \frac{N_0}{N_1M} \mathbb{1}\left(i \in \mathcal{J}'_M(\ell)\right).$ Notice that

 $\widehat{r}_M^{-\ell}(X_i)$ is a function of \boldsymbol{X} and $[Z_j]_{j=1,j\neq\ell}^{N_0}$. Since $[Z_j]_{j=1}^{N_1}$ are i.i.d. and independent with \widetilde{Z}_ℓ ,

For the second term in (A.27),

$$\mathbf{E}\left[\mathbb{1}\left(i\in\mathcal{J}_{M}(\ell)\setminus\mathcal{J}_{M}'(\ell)\right)\,\middle|\,\mathbf{X}\right]\leq\mathbf{E}\left[\mathbb{1}\left(i\in\mathcal{J}_{M}(\ell)\right)\,\middle|\,\mathbf{X}\right]=\nu_{1}\left(A_{M}(i)\right).\tag{A.28}$$

For the first term in (A.27), notice that $\frac{N_1M}{N_0} \hat{r}_M^{-\ell}(X_i) \sim \operatorname{Bin}\left(N_1 - 1, \nu_1(A_M(i))\right)$. Then using the Chernoff bound, we obtain

$$P\left(\frac{N_1M}{N_0}\hat{r}_M^{-\ell}(X_i) \le \frac{1}{2}(N_1 - 1)\nu_1(A_M(i))\right) \le \exp\left(-\frac{1}{2}(1 - \log 2)(N_1 - 1)\nu_1(A_M(i))\right), \\
 P\left(\frac{N_1M}{N_0}\hat{r}_M^{-\ell}(X_i) \ge 2(N_1 - 1)\nu_1(A_M(i))\right) \le \exp\left(-(2\log 2 - 1)(N_1 - 1)\nu_1(A_M(i))\right).$$

If $\nu_1(A_M(i)) \ge C_{\gamma} \frac{\log N_1}{N_1 - 1}$, then for any $\gamma > 0$, there exists a constant $C_{\gamma} > 0$ only depending on γ such that, with probability at least $1 - N_1^{-\gamma}$, we have

$$\frac{1}{2}\frac{N_0}{M}\left(1 - \frac{1}{N_1}\right)\nu_1\left(A_M(i)\right) \le \hat{r}_M^{-\ell}(X_i) \le 2\frac{N_0}{M}\left(1 - \frac{1}{N_1}\right)\nu_1\left(A_M(i)\right)$$

Notice that $\frac{N_0}{N_1M} \prec \frac{N_0}{M}\nu_1(A_M(i))$ when $\nu_1(A_M(i)) \geq C_{\gamma} \frac{\log N_1}{N_1-1}$. Then for N_0 sufficiently large, with probability at least $1 - N_1^{-\gamma}$, the following sandwich inequality holds:

$$\frac{1}{M} \frac{N_0}{M} \nu_1 (A_M(i)) \le \hat{r}_M^{-\ell}(X_i) \le \hat{r}_M^{-\ell}(X_i) + \frac{N_0}{N_1 M} \le 4 \frac{N_0}{M} \nu_1 (A_M(i)).$$
(A.29)

If $\nu_1(A_M(i)) \leq C_{\gamma} \frac{\log N_1}{N_1 - 1}$, then by the Chernoff bound,

$$P\left(\frac{N_1M}{N_0}\widehat{r}_M^{-\ell}(X_i) \ge 2C_{\gamma}\log N_1\right)$$

$$\leq \exp\left(2C_{\gamma}\log N_1 - (N_1 - 1)\nu_1(A_M(i)) - 2C_{\gamma}\log N_1\log\left(\frac{2C_{\gamma}\log N_1}{(N_1 - 1)\nu_1(A_M(i))}\right)\right)$$

$$\leq \exp\left(-(2\log 2 - 1)C_{\gamma}\log N_1\right) \le N_1^{-\gamma}.$$

Then for N_0 sufficiently large, with probability at least $1 - N_1^{-\gamma}$,

$$\widehat{r}_{M}^{-\ell}(X_{i}) \le 2C_{\gamma} \frac{N_{0} \log N_{1}}{N_{1}M}.$$
(A.30)

Then for any $\epsilon \in (0, 1)$,

where $C_{\epsilon} > 0$ is a constant only depending on ϵ and C > 0 is a universal constant. The first term

is due to (A.30) and the inequality

$$|x\log x - y\log y| \le C_{\epsilon}|x - y|^{1-\epsilon},$$

whenever $x, y \in [0, 1]$ and $|x - y| \leq \frac{1}{2}$. The second term is from $\phi(x) = x \log x$ and $\widehat{r}_M^{-\ell}(X_i) + \frac{N_0}{N_1M} \leq \frac{N_0}{M}$. The third term is from $\phi'(x) = 1 + \log x$ and the mean value theorem by using the sandwich inequality (A.29).

Plugging (A.28) and (A.31) into (A.26) and tracing back to (A.27) yields

$$E\left[\frac{1}{N_0}\sum_{i=1}^{N_0} \left(\phi\left(\hat{r}_M(X_i)\right) - \phi\left(\hat{r}'_M(X_i)\right)\right)\right]^2 \\ \leq \frac{4M}{N_0} E\left[C_{\epsilon}\left(\frac{N_0}{N_1M}\right)^{2(1-\epsilon)} \nu_1\left(A_M(i)\right) \mathbb{1}\left(\nu_1\left(A_M(i)\right) \le C_{\gamma}\frac{\log N_1}{N_1}\right) + \frac{2}{N_1^{\gamma}} \left(2\frac{N_0}{M}\log\frac{N_0}{M}\right)^2 \nu_1\left(A_M(i)\right) \\ + C\left[1 + \left(\log\left(\frac{N_0}{M}\nu_1\left(A_M(i)\right)\right)\right)^2\right] \left(\frac{N_0}{N_1M}\right)^2 \nu_1\left(A_M(i)\right) \mathbb{1}\left(\nu_1\left(A_M(i)\right) \ge C_{\gamma}\frac{\log N_1}{N_1}\right)\right].$$
(A.32)

For the first term in (A.32), since $MN_1/(N_0 \log^2 N_1) \to \infty$,

$$C_{\epsilon} \frac{M}{N_0} \left(\frac{N_0}{N_1 M}\right)^{2(1-\epsilon)} \nu_1 \left(A_M(i)\right) \mathbb{1} \left(\nu_1 \left(A_M(i)\right) \le C_{\gamma} \frac{\log N_1}{N_1}\right) \\ \le C_{\epsilon} C_{\gamma} \frac{M}{N_0} \left(\frac{N_0}{N_1 M}\right)^{2(1-\epsilon)} \frac{\log N_1}{N_1} = C_{\epsilon} C_{\gamma} \frac{1}{N_1^2} \left(\frac{N_0}{N_1 M}\right)^{1-2\epsilon} \log N_1 \prec \frac{1}{N_1^2},$$
(A.33)

where we take ϵ sufficiently small.

For the second term in (A.32), using Corollary 4.1, E[r(X)] = 1 and $N_1^{-\frac{d}{1+d}} \log N_0 \to 0$,

$$\frac{M}{N_0} \frac{1}{N_1^{\gamma}} \left(\frac{N_0}{M} \log \frac{N_0}{M}\right)^2 \mathbf{E} \left[\nu_1 \left(A_M(i)\right)\right] = \frac{1}{N_1^{\gamma}} \left(\log \frac{N_0}{M}\right)^2 \mathbf{E} \left[\frac{N_0}{M} \nu_1 \left(A_M(i)\right)\right]$$
$$\lesssim \frac{1}{N_1^{\gamma}} \left(\log \frac{N_0}{M}\right)^2 \prec \frac{1}{N_1^2}, \tag{A.34}$$

where we take γ sufficiently large.

For the third term in (A.32), using Corollary 4.1,

$$\frac{M}{N_{0}} \mathbb{E}\left[\left[1 + \left(\log\left(\frac{N_{0}}{M}\nu_{1}(A_{M}(i))\right)\right)^{2}\right]\left(\frac{N_{0}}{N_{1}M}\right)^{2}\nu_{1}\left(A_{M}(i)\right)\mathbb{1}\left(\nu_{1}(A_{M}(i)) \geq C_{\gamma}\frac{\log N_{1}}{N_{1}}\right)\right] \\
\leq \frac{M}{N_{0}} \mathbb{E}\left[\left[1 + \left(\log\left(\frac{N_{0}}{M}\nu_{1}(A_{M}(i))\right)\right)^{2}\right]\left(\frac{N_{0}}{N_{1}M}\right)^{2}\nu_{1}\left(A_{M}(i)\right)\right] \\
= \frac{1}{N_{1}^{2}} \mathbb{E}\left[\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right) + \left(\log\left(\frac{N_{0}}{M}\nu_{1}(A_{M}(i))\right)\right)^{2}\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right)\right] \\
\leq \frac{1}{N_{1}^{2}} \mathbb{E}\left[\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right) + \left(\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right)\right)^{2} \lor 4e^{-2}\right] \lesssim C\frac{1}{N_{1}^{2}},$$
(A.35)

where C > 0 is a constant only depending on f_L , f_U . The last step is from $(\log x)^2 \le x$ when $x \ge 1$ and the maximum of $(\log x)^2 x$ is $4e^{-2}$ when $x \in [0, 1]$.

Plugging (A.33), (A.34), (A.35) into (A.32) yields

$$\mathbf{E}\Big[\frac{1}{N_0}\sum_{i=1}^{N_0}\Big(\phi\Big(\widehat{r}_M(X_i)\Big)-\phi\Big(\widehat{r}'_M(X_i)\Big)\Big)\Big]^2 \lesssim C\frac{1}{N_1^2},$$

where C > 0 is a constant only depending on f_L, f_U .

By Efron-Stein inequality and (A.25),

$$E\left[\operatorname{Var}[T_1 \mid \boldsymbol{X}]\right] = E\left[\operatorname{Var}\left[\frac{1}{N_0}\sum_{i=1}^{N_0}\phi(\hat{r}_M(X_i)) \mid \boldsymbol{X}\right]\right]$$

$$\lesssim N_1 E\left[\frac{1}{N_0}\sum_{i=1}^{N_0}\left(\phi(\hat{r}_M(X_i)) - \phi(\hat{r}'_M(X_i))\right)\right]^2 \lesssim C\frac{1}{N_1},$$
(A.36)

where C > 0 is a constant only depending on f_L, f_U .

Combining (A.24) and (A.36) completes the proof.

A.4.4 Proof of Lemma A.6

Proof of Lemma A.6. From (A.23),

$$\begin{split} |\mathbf{E}[T_{2} | \mathbf{X}]| &= \Big| \mathbf{E} \Big[\frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \Big[\phi \Big(\mathbf{E}[\hat{r}_{M}(X_{i}) | \mathbf{X}] \Big) - \phi \Big(\mathbf{E}[\hat{r}_{M}(X_{i}) | X_{i}] \Big) \Big] \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \Big] \Big| \\ &\leq 2 \| \phi - g \|_{\infty} + \Big| \mathbf{E} \Big[\frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \Big[g \Big(\mathbf{E}[\hat{r}_{M}(X_{i}) | \mathbf{X}] \Big) - g \Big(\mathbf{E}[\hat{r}_{M}(X_{i}) | X_{i}] \Big) \Big] \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \Big] \Big| \\ &\leq 2 \| \phi - g \|_{\infty} + 4 \| x g''(x) \|_{\infty} \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \mathbf{E} \Big[\frac{(\mathbf{E}[\hat{r}_{M}(X_{i}) | \mathbf{X}] - \mathbf{E}[\hat{r}_{M}(X_{i}) | X_{i}])^{2}}{\mathbf{E}[\hat{r}_{M}(X_{i}) | X_{i}]} \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \Big] \\ &= 2 \| \phi - g \|_{\infty} + 4 \| x g''(x) \|_{\infty} \frac{1}{N_{0}} \sum_{i=1}^{N_{0}} \mathbf{E} \Big[\frac{\mathrm{Var}[\mathbf{E}[\hat{r}_{M}(X_{i}) | \mathbf{X}] | X_{i}]}{\mathbf{E}[\hat{r}_{M}(X_{i}) | X_{i}]} \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \Big]. \end{split}$$

Using Lemma A.4,

$$\begin{aligned} |\mathbf{E}[T_2 \mid \mathbf{X}]| &\leq 2 \|\phi - g\|_{\infty} + 4 \|xg''(x)\|_{\infty} \frac{1}{N_0} \sum_{i=1}^{N_0} \mathbf{E}\Big[\frac{\frac{32}{M}[r(X_i)]^2}{\frac{1}{2}r(X_i)} \mathbb{1}\Big(X_i \in \mathcal{E}\Big)\Big] \\ &\leq 2 \|\phi - g\|_{\infty} + 4 \|xg''(x)\|_{\infty} \frac{1}{N_0} \sum_{i=1}^{N_0} \frac{64}{M} \mathbf{E}[r(X_i)] \\ &= 2 \|\phi - g\|_{\infty} + \frac{256}{M} \|xg''(x)\|_{\infty}. \end{aligned}$$

Using Lemma A.10, we obtain

$$|\mathbf{E}[T_2 \mid \boldsymbol{X}]| \lesssim C \frac{1}{M},\tag{A.37}$$

where C > 0 is a constant only depending on f_L, f_U .

For the variance, we have

$$\operatorname{Var}[T_2] = \operatorname{Var}\left[\frac{1}{N_0}\sum_{i=1}^{N_0} \left[\phi\left(\operatorname{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(\operatorname{E}[\widehat{r}_M(X_i) \mid X_i]\right)\right] \mathbb{1}\left(X_i \in \mathcal{E}\right)\right].$$

To apply Efron-Stein inequality, recall $\boldsymbol{X} = (X_1, \dots, X_{N_0})$ and for any $\ell \in [\![N_0]\!]$,

$$\boldsymbol{X}_{\ell} = (X_1, \dots, X_{\ell-1}, \widetilde{X}_{\ell}, X_{\ell+1}, \dots, X_{N_0}),$$

where $[\widetilde{X}_{\ell}]_{\ell=1}^{N_0}$ are independent copies of $[X_{\ell}]_{\ell=1}^{N_0}$.

Fix $\ell \in [N_0]$. Recall that $A_M(\cdot)$ are defined based on \mathbf{X} . We then define $A'_M(\cdot)$ in the same way but based on \mathbf{X}_ℓ , and define $A^\ell_M(\cdot)$ based on $(X_1, \ldots, X_{\ell-1}, X_{\ell+1}, \ldots, X_{N_0})$.

(1) If $i = \ell$,

$$\begin{split} & \Big| \Big[\phi\Big(\mathrm{E}[\widehat{r}_{M}(X_{\ell}) \mid \boldsymbol{X}] \Big) - \phi\Big(\mathrm{E}[\widehat{r}_{M}(X_{\ell}) \mid X_{\ell}] \Big) \Big] \mathbb{1} \Big(X_{\ell} \in \mathcal{E} \Big) \\ & - \Big[\phi\Big(\mathrm{E}[\widehat{r}_{M}(\widetilde{X}_{\ell}) \mid \boldsymbol{X}_{\ell}] \Big) - \phi\Big(\mathrm{E}[\widehat{r}_{M}(\widetilde{X}_{\ell}) \mid \widetilde{X}_{\ell}] \Big) \Big] \mathbb{1} \Big(\widetilde{X}_{\ell} \in \mathcal{E} \Big) \Big| \\ \leq & \Big| \phi\Big(\mathrm{E}[\widehat{r}_{M}(X_{\ell}) \mid \boldsymbol{X}] \Big) - \phi\Big(\mathrm{E}[\widehat{r}_{M}(X_{\ell}) \mid X_{\ell}] \Big) \Big| \mathbb{1} \Big(X_{\ell} \in \mathcal{E} \Big) \\ & + \Big| \phi\Big(\mathrm{E}[\widehat{r}_{M}(\widetilde{X}_{\ell}) \mid \boldsymbol{X}_{\ell}] \Big) - \phi\Big(\mathrm{E}[\widehat{r}_{M}(\widetilde{X}_{\ell}) \mid \widetilde{X}_{\ell}] \Big) \Big| \mathbb{1} \Big(\widetilde{X}_{\ell} \in \mathcal{E} \Big). \end{split}$$

(2) If $i \neq \ell$,

$$\begin{aligned} & \left| \left[\phi \Big(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}] \Big) - \phi \Big(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid X_{i}] \Big) \right] \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \\ & - \left[\phi \Big(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}_{\ell}] \Big) - \phi \Big(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid X_{i}] \Big) \right] \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \right| \\ & = \left| \phi \Big(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}] \Big) - \phi \Big(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}_{\ell}] \Big) \Big| \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \\ & = \left| \phi \Big(\frac{N_{0}}{M} \nu_{1} \big(A_{M}(i) \big) \Big) - \phi \Big(\frac{N_{0}}{M} \nu_{1} \big(A'_{M}(i) \big) \Big) \Big| \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \\ & \leq \left| \phi \Big(\frac{N_{0}}{M} \nu_{1} \big(A_{M}(i) \big) \Big) - \phi \Big(\frac{N_{0}}{M} \nu_{1} \big(A'_{M}(i) \big) \Big) \Big| \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big) \\ & + \left| \phi \Big(\frac{N_{0}}{M} \nu_{1} \big(A'_{M}(i) \big) \Big) - \phi \Big(\frac{N_{0}}{M} \nu_{1} \big(A'_{M}(i) \big) \Big) \Big| \mathbb{1} \Big(X_{i} \in \mathcal{E} \Big). \end{aligned}$$

Since $[X_\ell]_{\ell=1}^{N_0}$ and $[\widetilde{X}_\ell]_{\ell=1}^{N_0}$ are exchangeable, using Efron-Stein inequality, we have

$$\operatorname{Var}[T_{2}] \lesssim \frac{1}{N_{0}} \operatorname{E}\left[\left|\phi\left(\operatorname{E}[\widehat{r}_{M}(X_{\ell}) \mid \boldsymbol{X}]\right) - \phi\left(\operatorname{E}[\widehat{r}_{M}(X_{\ell}) \mid X_{\ell}]\right)\right| \mathbb{1}\left(X_{\ell} \in \mathcal{E}\right)\right]$$
$$+ \sum_{i=1, i \neq \ell}^{N_{0}} \left|\phi\left(\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right)\right) - \phi\left(\frac{N_{0}}{M}\nu_{1}\left(A_{M}^{\ell}(i)\right)\right)\right| \mathbb{1}\left(X_{i} \in \mathcal{E}\right)\right]^{2}$$
$$\lesssim \frac{1}{N_{0}} \operatorname{E}\left[\left[\phi^{2}\left(\operatorname{E}[\widehat{r}_{M}(X_{\ell}) \mid \boldsymbol{X}]\right) + \phi^{2}\left(\operatorname{E}[\widehat{r}_{M}(X_{\ell}) \mid X_{\ell}]\right)\right] \mathbb{1}\left(X_{\ell} \in \mathcal{E}\right)\right]$$
$$+ \frac{1}{N_{0}} \operatorname{E}\left[\sum_{i=1, i \neq \ell}^{N_{0}} \left|\phi\left(\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right)\right) - \phi\left(\frac{N_{0}}{M}\nu_{1}\left(A_{M}^{\ell}(i)\right)\right)\right| \mathbb{1}\left(X_{i} \in \mathcal{E}\right)\right]^{2}.$$

For the first term, from Corollary 4.1, r(x) is bounded and $\phi^2(x) \le x^3$ when $x \ge 1$, we obtain

$$\mathbf{E}\left[\left[\phi^{2}\left(\mathbf{E}[\widehat{r}_{M}(X_{\ell}) \mid \boldsymbol{X}]\right) + \phi^{2}\left(\mathbf{E}[\widehat{r}_{M}(X_{\ell}) \mid X_{\ell}]\right)\right] \mathbb{1}\left(X_{\ell} \in \mathcal{E}\right)\right] \lesssim C,$$

where C > 0 is a constant only depending on f_L, f_U .

For the second term, notice that $A_M(i) \subset A_M^{\ell}(i)$ and

$$A_M^{\ell}(i) \setminus A_M(i) = \{ z \in \mathbb{R}^d : j_{M+1}(z) = i, \ell \in \mathcal{J}_M(z) \}$$

Then for an independent copy $Z \sim \nu_1$,

$$\frac{N_0}{M}\nu_1\big(A_M^\ell(i)\big) - \frac{N_0}{M}\nu_1\big(A_M(i)\big) = \frac{N_0}{M} \mathcal{P}\Big(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \mid \mathbf{X}\Big) \le \frac{N_0}{M}\nu_1\big(A_M(\ell)\big).$$

By the mean value theorem, we have

$$\sum_{i=1,i\neq\ell}^{N_0} \left| \phi\left(\frac{N_0}{M}\nu_1(A_M(i))\right) - \phi\left(\frac{N_0}{M}\nu_1(A_M^\ell(i))\right) \right| \mathbb{1}\left(X_i \in \mathcal{E}\right) \\ = \sum_{i=1,i\neq\ell}^{N_0} \left| 1 + \log \xi_i \right| \left[\frac{N_0}{M} P\left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \mid \mathbf{X}\right) \right] \mathbb{1}\left(X_i \in \mathcal{E}\right),$$

where ξ_i is between $\frac{N_0}{M}\nu_1(A_M(i))$ and $\frac{N_0}{M}\nu_1(A_M^\ell(i))$. When $\xi_i \leq 1$, $|\log \xi_i| \leq -\log \left(\frac{N_0}{M}\nu_1(A_M(i))\right) \mathbb{1}\left(\frac{N_0}{M}\nu_1(A_M(i)) \leq 1\right)$. When $\xi_i \geq 1$, $|\log \xi_i| \leq 1$. $\log\left(\frac{N_0}{M}\nu_1(A_M^{\ell}(i))\right) \leq \left(\frac{N_0}{M}\nu_1(A_M(i)) + \frac{N_0}{M}\nu_1(A_M(\ell))\right).$ Then $\left|1 + \log \xi_i\right| \le 1 - \log \left(\frac{N_0}{M} \nu_1(A_M(i))\right) \mathbb{1}\left(\frac{N_0}{M} \nu_1(A_M(i)) \le 1\right) + \left(\frac{N_0}{M} \nu_1(A_M(i)) + \frac{N_0}{M} \nu_1(A_M(\ell))\right).$

Notice that events $\{j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z)\}$ are disjoint for $i \neq \ell$. We then have

$$\sum_{i=1,i\neq\ell}^{N_0} \left[1 + \frac{N_0}{M} \nu_1 (A_M(\ell)) \right] \left[\frac{N_0}{M} P \left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \mid \mathbf{X} \right) \right] \mathbb{1} \left(X_i \in \mathcal{E} \right)$$

$$\leq \sum_{i=1,i\neq\ell}^{N_0} \left[1 + \frac{N_0}{M} \nu_1 (A_M(\ell)) \right] \left[\frac{N_0}{M} P \left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \mid \mathbf{X} \right) \right]$$

$$= \left[1 + \frac{N_0}{M} \nu_1 (A_M(\ell)) \right] \left[\frac{N_0}{M} P \left(\ell \in \mathcal{J}_M(Z) \mid \mathbf{X} \right) \right]$$

$$= \left[1 + \frac{N_0}{M} \nu_1 (A_M(\ell)) \right] \frac{N_0}{M} \nu_1 (A_M(\ell)).$$

From Corollary 4.1,

$$\mathbf{E}\Big[\sum_{i=1,i\neq\ell}^{N_0} \Big[1 + \frac{N_0}{M}\nu_1\big(A_M(\ell)\big)\Big]\Big[\frac{N_0}{M}\mathbf{P}\Big(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \,\Big|\, \mathbf{X}\Big)\Big]\mathbb{1}\Big(X_i \in \mathcal{E}\Big)\Big]^2 \lesssim C, \quad (A.38)$$

where C > 0 is a constant only depending on f_L, f_U .

Let $\Xi_{\ell} \mid \mathbf{X}$ follows the same distribution as $Z \mid \mathbf{X}, \{Z \in A_M(\ell)\}$ for $Z \sim \nu_1$. Then

$$\begin{split} &\sum_{i=1,i\neq\ell}^{N_0} \frac{N_0}{M} \nu_1 \left(A_M(i) \right) \left[\frac{N_0}{M} \mathbf{P} \left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \mid \mathbf{X} \right) \right] \mathbb{1} \left(X_i \in \mathcal{E} \right) \\ &\leq \sum_{i=1,i\neq\ell}^{N_0} \frac{N_0}{M} \nu_1 \left(A_M(i) \right) \left[\frac{N_0}{M} \mathbf{P} \left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \mid \mathbf{X} \right) \right] \\ &= \sum_{i=1,i\neq\ell}^{N_0} \mathbf{E} \left[\left(\frac{N_0}{M} \right)^2 \nu_1 \left(A_M(i) \right) \mathbb{1} \left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \right) \mid \mathbf{X} \right] \\ &= \left(\frac{N_0}{M} \right)^2 \mathbf{E} \left[\sum_{i=1,i\neq\ell}^{N_0} \nu_1 \left(A_M(j_{M+1}(Z)) \right) \mathbb{1} \left(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \right) \mid \mathbf{X} \right] \\ &= \left(\frac{N_0}{M} \right)^2 \mathbf{E} \left[\nu_1 \left(A_M(j_{M+1}(Z)) \right) \mathbb{1} \left(\ell \in \mathcal{J}_M(Z) \right) \mid \mathbf{X} \right] \end{split}$$

$$= \left(\frac{N_0}{M}\right)^2 \mathbf{E} \left[\nu_1 \left(A_M(j_{M+1}(\Xi_\ell)) \right) \, \middle| \, \mathbf{X} \right] \nu_1 \left(A_M(\ell) \right)$$

Then

$$\begin{split} & \operatorname{E}\Big[\sum_{i=1,i\neq\ell}^{N_{0}}\frac{N_{0}}{M}\nu_{1}\left(A_{M}(i)\right)\Big[\frac{N_{0}}{M}\operatorname{P}\left(j_{M+1}(Z)=i,\ell\in\mathcal{J}_{M}(Z)\,\middle|\,\mathbf{X}\right)\Big]\mathbb{1}\left(X_{i}\in\mathcal{E}\right)\Big]^{2} \\ &\leq \left(\frac{N_{0}}{M}\right)^{4}\operatorname{E}\Big[\operatorname{E}\Big[\nu_{1}\left(A_{M}(j_{M+1}(Z))\right)\mathbb{1}\left(\ell\in\mathcal{J}_{M}(Z)\right)\,\middle|\,\mathbf{X}\Big]\Big]^{2} \\ &= \left(\frac{N_{0}}{M}\right)^{4}\operatorname{E}\Big[\operatorname{E}\Big[\nu_{1}\left(A_{M}(j_{M+1}(\Xi_{\ell}))\right)\,\middle|\,\mathbf{X}\Big]\nu_{1}\left(A_{M}(\ell)\right)\Big]^{2} \\ &\leq \left(\frac{N_{0}}{M}\right)^{4}\left(\operatorname{E}\Big[\nu_{1}\left(A_{M}(j_{M+1}(\Xi_{\ell}))\right)\Big]^{4}\operatorname{E}\Big[\nu_{1}\left(A_{M}(\ell)\right)\Big]^{4}\right)^{1/2} \\ &= \left(\frac{N_{0}}{M}\right)^{4}\left(\operatorname{E}\Big[\nu_{1}\left(A_{M}(i)\right)\Big]^{4}\operatorname{E}\Big[\nu_{1}\left(A_{M}(\ell)\right)\Big]^{4}\right)^{1/2} \end{split}$$

for some $i \neq \ell$. The last step is due to the fact that $[X_i]_{i=1,i\neq\ell}^{N_0}$ are exchangeable and $\ell \in \mathcal{J}_M(Z)$ implies $j_{M+1}(Z) \neq \ell$.

By Corollary 4.1, we obtain

$$\mathbf{E}\Big[\sum_{i=1,i\neq\ell}^{N_0} \frac{N_0}{M} \nu_1\big(A_M(i)\big) \Big[\frac{N_0}{M} \mathbf{P}\Big(j_{M+1}(Z) = i, \ell \in \mathcal{J}_M(Z) \,\Big|\, \mathbf{X}\Big)\Big] \mathbb{1}\Big(X_i \in \mathcal{E}\Big)\Big]^2 \lesssim C, \qquad (A.39)$$

where C > 0 is a constant only depending on f_L, f_U .

Using the same trick, by Lemma A.4 and Corollary 4.1, we have

$$\begin{split} & \operatorname{E}\Big[\sum_{i=1,i\neq\ell}^{N_{0}}\log\Big(\frac{N_{0}}{M}\nu_{1}\big(A_{M}(i)\big)\Big)\mathbb{1}\Big(\frac{N_{0}}{M}\nu_{1}\big(A_{M}(i)\big) \leq 1\Big)\Big[\frac{N_{0}}{M}\operatorname{P}\Big(j_{M+1}(Z) = i, \ell \in \mathcal{J}_{M}(Z) \,\Big|\,\mathbf{X}\Big)\Big]\mathbb{1}\Big(X_{i} \in \mathcal{E}\Big)\Big]^{2} \\ & \leq \Big(\frac{N_{0}}{M}\Big)^{2}\Big(\operatorname{E}\Big[\log^{4}\Big(\frac{N_{0}}{M}\nu_{1}\big(A_{M}(i)\big)\Big)\mathbb{1}\Big(\frac{N_{0}}{M}\nu_{1}\big(A_{M}(i)\big) \leq 1\Big)\mathbb{1}\Big(X_{i} \in \mathcal{E}\Big)\Big]\operatorname{E}\Big[\nu_{1}^{4}\big(A_{M}(\ell)\big)\Big]\Big)^{1/2} \\ & \leq \Big(\frac{N_{0}}{M}\Big)^{2}\Big(\operatorname{E}\Big[\log^{4}\Big(C_{1}r(X_{i})\Big)\mathbb{1}\Big(\frac{N_{0}}{M}\nu_{1}\big(A_{M}(i)\big) \leq 1\Big)\mathbb{1}\Big(X_{i} \in \mathcal{E}\Big)\Big]\operatorname{E}\Big[\nu_{1}^{4}\big(A_{M}(\ell)\big)\Big]\Big)^{1/2} \\ & \leq \Big(\frac{N_{0}}{M}\Big)^{2}\Big(\operatorname{E}\Big[\log^{4}\Big(C_{1}r(X_{i})\Big)\Big]\operatorname{E}\Big[\nu_{1}^{4}\big(A_{M}(\ell)\big)\Big]\Big)^{1/2} \\ & = \Big[\operatorname{E}\Big[\log^{4}\Big(C_{1}r(X_{i})\Big)\Big]\operatorname{E}\Big[\Big(\frac{N_{0}}{M}\Big)^{4}\nu_{1}^{4}\big(A_{M}(\ell)\big)\Big]\Big)^{1/2} \lesssim C, \end{split}$$

where C > 0 is a constant only depending on f_L, f_U, d, f'_U .

A.4.5 Proof of Lemma A.7

Proof of Lemma A.7. We have

$$|T_3| = \left|\frac{1}{N_0}\sum_{i=1}^{N_0} \left[\phi\left(\mathbb{E}[\widehat{r}_M(X_i) \mid X_i]\right) - \phi\left(r(X_i)\right)\right] \mathbb{1}\left(X_i \in \mathcal{E}\right)\right]$$
$$\leq \frac{1}{N_0}\sum_{i=1}^{N_0} \left|\phi\left(\mathbb{E}[\widehat{r}_M(X_i) \mid X_i]\right) - \phi\left(r(X_i)\right)\right| \mathbb{1}\left(X_i \in \mathcal{E}\right).$$

From Lemma A.4, if $X_i \in \mathcal{E}$, then for N_0 sufficiently large,

$$\frac{1}{2}r(X_i) \le \operatorname{E}[\widehat{r}_M(X_i) \,|\, X_i] \le \frac{3}{2}r(X_i).$$

From the mean value theorem and the proof of Theorem 4.3(i),

$$\begin{aligned} |T_3| \lesssim \frac{1}{N_0} \sum_{i=1}^{N_0} \left[1 + \left| \log r(X_i) \right| \right] \left| \mathbb{E}[\hat{r}_M(X_i) \mid X_i] - r(X_i) \right| \mathbb{1} \left(X_i \in \mathcal{E} \right) \\ \lesssim C \left(\frac{M}{N_0} \right)^{1/d} \frac{1}{N_0} \sum_{i=1}^{N_0} \left[1 + \left| \log r(X_i) \right| \right] \mathbb{1} \left(X_i \in \mathcal{E} \right), \end{aligned}$$

where C > 0 is a constant only depending on f_L, f_U, L, d, f'_U .

Then we obtain

$$\operatorname{E}[|T_3|] \le C\left(\frac{M}{N_0}\right)^{1/d},\tag{A.40}$$

where C > 0 is a constant only depending on f_L, f_U, L, d, f'_U .

From the i.i.d.-ness of $[X_i]_{i=1}^{N_0}$,

$$\operatorname{Var}[T_{3}] = \operatorname{Var}\left[\frac{1}{N_{0}}\sum_{i=1}^{N_{0}}\left[\phi\left(\operatorname{E}[\widehat{r}_{M}(X_{i}) \mid X_{i}]\right) - \phi\left(r(X_{i})\right)\right]\mathbb{1}\left(X_{i} \in \mathcal{E}\right)\right]$$
$$= \frac{1}{N_{0}}\operatorname{Var}\left[\left[\phi\left(\operatorname{E}[\widehat{r}_{M}(X_{i}) \mid X_{i}]\right) - \phi\left(r(X_{i})\right)\right]\mathbb{1}\left(X_{i} \in \mathcal{E}\right)\right]$$
$$\leq \frac{1}{N_{0}}\operatorname{E}\left[\left[\phi\left(\operatorname{E}[\widehat{r}_{M}(X_{i}) \mid X_{i}]\right) - \phi\left(r(X_{i})\right)\right]^{2}\mathbb{1}\left(X_{i} \in \mathcal{E}\right)\right]$$
$$\leq \frac{1}{N_{0}}C\left(\frac{M}{N_{0}}\right)^{2/d},$$
(A.41)

where C > 0 is a constant only depending on f_L, f_U, L, d, f'_U . The last step is due to the mean value theorem.

Combining (A.40) and (A.41) completes the proof.

A.4.6 Proof of Lemma A.8

Proof of Lemma A.8. We seperate $S_0 \setminus \mathcal{E}$ into three sets $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$, where $\mathcal{E}_1 = \{x \in S_0 : 0 < f_1(x) \le C_0(\frac{M}{N_0})^{1/d}\}$, $\mathcal{E}_2 = \{x \in \mathbb{R}^d : f_1(x) = 0, \Delta(x) > \delta'_N\}$, $\mathcal{E}_3 = \{x \in S_0 : f_1(x) = 0, \Delta(x) \le \delta'_N\} \cup \{x \in S_0 : f_1(x) > C_0(\frac{M}{N_0})^{1/d}, \Delta(x) \le C_0(\frac{M}{N_0})^{1/d}\}$, where C_0 is the constant in Lemma A.4, and δ'_N is defined in the proof of Theorem 4.4.

For \mathcal{E}_1 , from the Cauchy–Schwarz inquality,

$$\mathbb{E}\left[\frac{1}{N_0}\sum_{i=1}^{N_0} \left[\phi\left(\mathbb{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(r(X_i)\right)\right] \mathbb{1}\left(X_i \in \mathcal{E}_1\right)\right]^2 \\
 \leq \mathbb{E}\left[\left[\phi\left(\mathbb{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(r(X_i)\right)\right]^2 \mathbb{1}\left(X_i \in \mathcal{E}_1\right)\right]$$

We take N_0 sufficiently large such that $C_0(\frac{M}{N_0})^{1/d} \leq f_L e^{-1} \wedge 1$. Then for any $x \in \mathcal{E}_1$, $r(x) \leq e^{-1}$ and $f_1(x) \leq 1$.

If $E[\hat{r}_M(X_i) | \mathbf{X}] \leq r(X_i)$, then

$$\begin{split} & \operatorname{E}\left[\left[\phi\left(\operatorname{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}]\right) - \phi\left(r(X_{i})\right)\right]^{2} \mathbb{1}\left(X_{i} \in \mathcal{E}_{1}\right)\right] \\ & \leq \operatorname{E}\left[\phi^{2}\left(r(X_{i})\right) \mathbb{1}\left(X_{i} \in \mathcal{E}_{1}\right)\right] \leq \phi^{2}\left(f_{L}^{-1}C_{0}\left(\frac{M}{N_{0}}\right)^{1/d}\right) \operatorname{P}\left(0 < f_{1}(X_{i}) \leq C_{0}\left(\frac{M}{N_{0}}\right)^{1/d}\right) \\ & \lesssim C\left(\frac{M}{N_{0}}\right)^{2/d} \log^{2}\left(\frac{N_{0}}{M}\right) \operatorname{P}\left(0 < f_{1}(X_{i}) \leq C_{0}\left(\frac{M}{N_{0}}\right)^{1/d}\right) \\ & \leq C\left(\frac{M}{N_{0}}\right)^{2/d} \log^{2}\left(\frac{N_{0}}{M}\right) \operatorname{P}\left(\left|\log f_{1}(X_{i})\right| \geq \left|\frac{1}{d}\log\left(\frac{N_{0}}{M}\right) - \log C_{0}\right|\right) \\ & \leq C\left(\frac{M}{N_{0}}\right)^{2/d} \log^{2}\left(\frac{N_{0}}{M}\right)\left(\frac{1}{d}\log\left(\frac{N_{0}}{M}\right) - \log C_{0}\right)^{-4} \operatorname{E}\left[\log^{4}\left(f_{1}(X_{i})\right)\right] \\ & \prec C\left(\frac{M}{N_{0}}\right)^{2/d}, \end{split}$$

where C > 0 is a constant only depending on f_L, f_U, L, d .

If $E[\hat{r}_M(X_i) | \mathbf{X}] > r(X_i)$, then from mean value theorem,

$$\begin{split} & \operatorname{E}\left[\left[\phi\left(\operatorname{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}]\right) - \phi\left(r(X_{i})\right)\right]^{2} \mathbb{1}\left(X_{i} \in \mathcal{E}_{1}\right)\right] \\ & \leq \operatorname{E}\left[\left(1 + \log\left(r(X_{i})\right)\right)^{2}\left(\operatorname{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}] - r(X_{i})\right)^{2} \mathbb{1}\left(X_{i} \in \mathcal{E}_{1}\right)\right] \\ & \lesssim C\left(\frac{M}{N_{0}}\right)^{2/d} \operatorname{E}\left[\left(1 + \log\left(r(X_{i})\right)\right)^{2} \mathbb{1}\left(X_{i} \in \mathcal{E}_{1}\right)\right] \\ & \lesssim C\left(\frac{M}{N_{0}}\right)^{2/d} \left(1 + \operatorname{E}\left[\log^{4}\left(f_{1}(X_{i})\right)\right]\right) \lesssim C\left(\frac{M}{N_{0}}\right)^{2/d}, \end{split}$$

where C > 0 is a constant only depending on f_L, f_U, L, d, f'_U .

We then obtain

$$\mathbf{E}\Big[\frac{1}{N_0}\sum_{i=1}^{N_0}\Big[\phi\Big(\mathbf{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\Big) - \phi\Big(r(X_i)\Big)\Big]\mathbb{1}\Big(X_i \in \mathcal{E}_1\Big)\Big]^2 \lesssim C\Big(\frac{M}{N_0}\Big)^{2/d},$$

where C > 0 is a constant only depending on f_L, f_U, L, d, f'_U . For \mathcal{E}_2 , notice that $(\log x)^2 < \frac{1}{x} + x$. In the same way as case II of the proof of Theorem 4.4, for any $\gamma > 0$, we have

$$E\left[\frac{1}{N_0}\sum_{i=1}^{N_0} \left[\phi\left(E[\hat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(r(X_i)\right)\right] \mathbb{1}\left(X_i \in \mathcal{E}_2\right)\right]^2$$

$$\leq E\left[\left[\phi\left(E[\hat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(r(X_i)\right)\right]^2 \mathbb{1}\left(X_i \in \mathcal{E}_2\right)\right]$$

$$= E\left[\phi^2\left(E[\hat{r}_M(X_i) \mid \boldsymbol{X}]\right) \mathbb{1}\left(X_i \in \mathcal{E}_2\right)\right] \prec N_0^{-\gamma}.$$

For \mathcal{E}_3 ,

$$E\left[\frac{1}{N_0}\sum_{i=1}^{N_0} \left[\phi\left(E[\hat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(r(X_i)\right)\right] \mathbb{1}\left(X_i \in \mathcal{E}_3\right)\right]^2 \\
 = \frac{1}{N_0} E\left[\left[\phi\left(E[\hat{r}_M(X_i) \mid \boldsymbol{X}]\right) - \phi\left(r(X_i)\right)\right]^2 \mathbb{1}\left(X_i \in \mathcal{E}_3\right)\right] + \left(1 - \frac{1}{N_0}\right)$$

$$\mathbb{E}\Big[\Big[\phi\Big(\mathbb{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\Big) - \phi\Big(r(X_i)\Big)\Big]\Big[\phi\Big(\mathbb{E}[\widehat{r}_M(X_j) \mid \boldsymbol{X}]\Big) - \phi\Big(r(X_j)\Big)\Big]\mathbb{1}\Big(X_i \in \mathcal{E}_3\Big)\mathbb{1}\Big(X_j \in \mathcal{E}_3\Big)\Big].$$
Notice that

Notice that

$$\left|\phi\left(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}]\right)\right| + \left|\phi\left(r(X_{i})\right)\right| \leq \left(\mathrm{E}[\widehat{r}_{M}(X_{i}) \mid \boldsymbol{X}]\right)^{2} + r(X_{i})^{2} + 2.$$

Employing the same approach as in the proof of case III in Theorem 4.4, for any $X_i, X_j \in \mathcal{E}_3$, one can establish

$$\mathbf{E}\left[\left(\mathbf{E}[\hat{r}_M(X_i) \mid \boldsymbol{X}]\right)^4 \mid X_i, X_j\right] \lesssim C,$$

where C > 0 is a constant only depending on f_L, f_U, a .

Then from the independence of X_i and X_j ,

$$\mathbf{E}\Big[\frac{1}{N_0}\sum_{i=1}^{N_0}\Big[\phi\Big(\mathbf{E}[\widehat{r}_M(X_i) \mid \boldsymbol{X}]\Big) - \phi\Big(r(X_i)\Big)\Big]\mathbb{1}\Big(X_i \in \mathcal{E}_3\Big)\Big]^2 \lesssim C\Big(\frac{M}{N_0}\Big)^{2/d},$$

where the constant C > 0 only depending on f_L, f_U, a, H, d .

A.4.7 Proof of Proposition 6.1

Proof of Proposition 6.1. We take ν_0 to be the uniform distribution of an arbitrary support such that the density is f_L . Notice that in this case, the KL divergence estimation is reduced to the estimation of differential entropy with N_1 samples. We then complete the proof using the minimax lower bound in bounded support Lipschitz entropy estimation without the assumption that the density is bounded away from zero (Han et al., 2020, Theorem 7).

A.5 Proofs of results in Section 7

A.5.1 Proof of Lemma 7.1

Proof of Lemma 7.1. The first inequality is directly from the definition of Lebesgue points. The second inequality follows by

$$\begin{split} & \left| \frac{\nu(B_{z,\|z-x\|})}{\lambda(B_{z,\|z-x\|})} - f(x) \right| \leq \frac{1}{\lambda(B_{z,\|z-x\|})} \int_{B_{z,\|z-x\|}} |f(y) - f(x)| \mathrm{d}y \\ \leq & \frac{1}{\lambda(B_{z,\|z-x\|})} \int_{B_{x,2\|z-x\|}} |f(y) - f(x)| \mathrm{d}y = \frac{\lambda(B_{x,2\|z-x\|})}{\lambda(B_{z,\|z-x\|})} \frac{1}{\lambda(B_{x,2\|z-x\|})} \int_{B_{x,2\|z-x\|}} |f(y) - f(x)| \mathrm{d}y \\ = & 2^d \frac{1}{\lambda(B_{x,2\|z-x\|})} \int_{B_{x,2\|z-x\|}} |f(y) - f(x)| \mathrm{d}y, \end{split}$$

and then the definition of Lebesgue points.

A.5.2 Proof of Lemma 7.3

Proof of Lemma 7.3. Fix any $(\nu_0, \nu_1) \in \mathcal{P}_{x,p}(f_L, f_U, L, d, \delta)$.

We first prove the first claim. First consider $f_1(x) > 0$. For any $\epsilon > 0$, there exists $\delta' > 0$ such that for any $z \in \mathbb{R}^d$ satisfying $||z - x|| \leq \delta'$, we have $|f_0(z) - f_0(x)| \leq \epsilon f_0(x)$ and $|f_1(z) - f_0(x)| \leq \epsilon f_0(x)$

 $|f_1(x)| \leq \epsilon f_1(x)$ from the local Lipschitz assumption. We take w > 0 sufficiently small such that $w < (1-\epsilon)f_0(x)\lambda(B_{0,\delta'})$. Then $W \leq w$ implies $||x-Z|| \leq \delta'$. Then for w > 0 sufficiently small,

$$P(W \le w) = P\left(W \le w, \|x - Z\| \le \delta'\right) \le P\left(\frac{1 - \epsilon}{1 + \epsilon} \frac{f_0(x)}{f_1(x)} \nu_1(B_{x,\|x - Z\|}) \le w\right) = \frac{1 + \epsilon}{1 - \epsilon} \frac{f_1(x)}{f_0(x)} w_2$$

and

$$P(W \le w) = P\left(W \le w, \|x - Z\| \le \delta'\right) \ge P\left(\frac{1 + \epsilon}{1 - \epsilon} \frac{f_0(x)}{f_1(x)} \nu_1(B_{x,\|x - Z\|}) \le w, \|x - Z\| \le \delta\right)$$
$$= P\left(\frac{1 + \epsilon}{1 - \epsilon} \frac{f_0(x)}{f_1(x)} \nu_1(B_{x,\|x - Z\|}) \le w\right) = \frac{1 - \epsilon}{1 + \epsilon} \frac{f_1(x)}{f_0(x)} w.$$

Then we have

$$\frac{1-\epsilon}{1+\epsilon} \frac{f_1(x)}{f_0(x)} \le \liminf_{w \to 0} w^{-1} \mathbf{P}(W \le w) \le \limsup_{w \to 0} w^{-1} \mathbf{P}(W \le w) \le \frac{1+\epsilon}{1-\epsilon} \frac{f_1(x)}{f_0(x)}$$

Since ϵ is arbitrary, we obtain

$$f_W(0) = \lim_{w \to 0} w^{-1} \mathbf{P}(W \le w) = \frac{f_1(x)}{f_0(x)} = r(x).$$

The case for $f_1(x) = 0$ can be established in the same way. This completes the proof of the first claim.

For the second claim, for any $0 < \epsilon < f_L$, there exists $\delta' > 0$ such that for any $z \in \mathbb{R}^d$ satisfying $||z-x|| \le \delta'$, we have $|f_0(z) - f_0(x)| \le \epsilon$ and $|f_1(z) - f_1(x)| \le \epsilon$ from the local Lipschitz assumption. We take N_0 sufficiently large such that $2\frac{M}{N_0} < (f_L - \epsilon)\lambda(B_{0,\delta'})$. Then for any $0 < w \le 2\frac{M}{N_0}$, we have $w < (f_L - \epsilon)\lambda(B_{0,\delta'})$. We take t > 0 such that $w + t < (f_L - \epsilon)\lambda(B_{0,\delta'})$. Then for any $(\nu_0, \nu_1) \in \mathcal{P}_{x,p}(f_L, f_U, L, d, \delta)$,

$$P\left(w \le W \le w+t\right) = \nu_1\left(\left\{z \in \mathbb{R}^d : \nu_0(B_{z,\|x-z\|}) \in [w,w+t]\right\}\right)$$
$$\leq \frac{f_1(x) + \epsilon}{f_0(x) - \epsilon}\nu_0\left(\left\{z \in \mathbb{R}^d : \nu_0(B_{z,\|x-z\|}) \in [w,w+t]\right\}\right).$$

Notice that f_0 is lower bounded by f_L . Then for N_0 sufficiently large,

$$\limsup_{t \to 0} t^{-1} \mathbf{P} \left(w \le W \le w + t \right) \le \frac{f_1(x) + \epsilon}{f_0(x) - \epsilon} (1 + \epsilon).$$

This then completes the proof.

A.5.3 Proof of Lemma 7.4

Proof of Lemma 7.4. Due to the i.i.d.-ness of Z and \tilde{Z} ,

$$\left(\frac{N_0}{M}\right)^2 \left[P\left(W \le V, \widetilde{W} \le \widetilde{V}, W \le 2\frac{M}{N_0}, \widetilde{W} \le 2\frac{M}{N_0}\right) - P\left(W \le V, W \le 2\frac{M}{N_0}\right) P\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0}\right) \right]$$

$$= \left(\frac{N_0}{M}\right)^2 \int_0^{2\frac{M}{N_0}} \int_0^{2\frac{M}{N_0}} \left[P\left(V \ge w_1, \widetilde{V} \ge w_2\right) - P\left(V \ge w_1\right) P\left(\widetilde{V} \ge w_2\right) \right] f_W(w_1) f_W(w_2) dw_1 dw_2$$

$$\le 4 \left(\frac{f_U}{f_L}\right)^2 \left(\frac{N_0}{M}\right)^2 \int_0^{2\frac{M}{N_0}} \int_0^{2\frac{M}{N_0}} \left| P\left(V \ge w_1, \widetilde{V} \ge w_2\right) - P\left(V \ge w_1\right) P\left(\widetilde{V} \ge w_2\right) \right| dw_1 dw_2$$

$$=4\left(\frac{f_U}{f_L}\right)^2 \int_{-1}^{1} \int_{-1}^{1} \left| P\left(V \ge \frac{M}{N_0}(1+t_1), \widetilde{V} \ge \frac{M}{N_0}(1+t_2)\right) - P\left(V \ge \frac{M}{N_0}(1+t_1)\right) P\left(\widetilde{V} \ge \frac{M}{N_0}(1+t_2)\right) \right| dt_1 dt_2$$

where the last step is from taking $w_1 = \frac{M}{N_0}(1+t_1)$ and $w_2 = \frac{M}{N_0}(1+t_2)$. Let

$$S(t_1, t_2) := \left| \mathbf{P} \left(V \ge \frac{M}{N_0} (1 + t_1), \widetilde{V} \ge \frac{M}{N_0} (1 + t_2) \right) - \mathbf{P} \left(V \ge \frac{M}{N_0} (1 + t_1) \right) \mathbf{P} \left(\widetilde{V} \ge \frac{M}{N_0} (1 + t_2) \right) \right|.$$

If $t_1 \ge t_2 \ge 0$,

$$S(t_1, t_2) \le P\left(V \ge \frac{M}{N_0}(1+t_1)\right) = P\left(U_{(M)} \ge \frac{M}{N_0}(1+t_1)\right).$$

If $t_2 \ge t_1 \ge 0$,

$$S(t_1, t_2) \le P\Big(\widetilde{V} \ge \frac{M}{N_0}(1+t_2)\Big) = P\Big(U_{(M)} \ge \frac{M}{N_0}(1+t_2)\Big).$$

Then for $t_1, t_2 \ge 0$,

$$\begin{split} S(t_1, t_2) &\leq \mathsf{P}\Big(U_{(M)} \geq \frac{M}{N_0}(1 + t_1 \lor t_2)\Big). \\ \text{If } t_1 \leq t_2 \leq 0 \text{ and } \mathsf{P}\Big(V \geq \frac{M}{N_0}(1 + t_1), \widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) \geq \mathsf{P}\Big(V \geq \frac{M}{N_0}(1 + t_1)\Big)\mathsf{P}\Big(\widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big), \\ S(t_1, t_2) &\leq \mathsf{P}\Big(\widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) - \mathsf{P}\Big(V \geq \frac{M}{N_0}(1 + t_1)\Big)\mathsf{P}\Big(\widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) \\ &= \mathsf{P}\Big(V \leq \frac{M}{N_0}(1 + t_1)\Big)\mathsf{P}\Big(\widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) \leq \mathsf{P}\Big(V \leq \frac{M}{N_0}(1 + t_1)\Big) = \mathsf{P}\Big(U_{(M)} \leq \frac{M}{N_0}(1 + t_1)\Big). \\ \text{If } t_1 \leq t_2 \leq 0 \text{ and } \mathsf{P}\Big(V \geq \frac{M}{N_0}(1 + t_1), \widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) \leq \mathsf{P}\Big(V \geq \frac{M}{N_0}(1 + t_1)\Big)\mathsf{P}\Big(\widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big), \\ S(t_1, t_2) \leq \mathsf{P}\Big(\widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) - \mathsf{P}\Big(V \geq \frac{M}{N_0}(1 + t_1), \widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) \\ &= \mathsf{P}\Big(V \leq \frac{M}{N_0}(1 + t_1), \widetilde{V} \geq \frac{M}{N_0}(1 + t_2)\Big) \leq \mathsf{P}\Big(V \leq \frac{M}{N_0}(1 + t_1)\Big) = \mathsf{P}\Big(U_{(M)} \leq \frac{M}{N_0}(1 + t_1)\Big). \\ \text{If } t_1 \leq t_2 \leq 0 \text{ and } \mathsf{P}(V \geq \frac{M}{N_0}(1 + t_2)\Big) \leq \mathsf{P}\Big(V \leq \frac{M}{N_0}(1 + t_1)\Big) = \mathsf{P}\Big(U_{(M)} \leq \frac{M}{N_0}(1 + t_1)\Big). \\ \end{array}$$

If $t_2 \leq t_1 \leq 0$, we can establish in the same way that

$$S(t_1, t_2) \le P\Big(U_{(M)} \le \frac{M}{N_0}(1+t_2)\Big).$$

Then for $t_1, t_2 \leq 0$,

$$S(t_1, t_2) \le P\Big(U_{(M)} \le \frac{M}{N_0}(1 + t_1 \wedge t_2)\Big).$$

For $t_1 \ge 0 \ge t_2$, if $t_1 + t_2 \ge 0$,

$$S(t_1, t_2) \le P\Big(U_{(M)} \ge \frac{M}{N_0}(1+t_1)\Big),$$

and if $t_1 + t_2 \le 0$,

$$S(t_1, t_2) \le P\Big(U_{(M)} \le \frac{M}{N_0}(1+t_2)\Big).$$

Then

$$\left(\frac{N_0}{M}\right)^2 \left[\mathbf{P}\left(W \le V, \widetilde{W} \le \widetilde{V}, W \le 2\frac{M}{N_0}, \widetilde{W} \le 2\frac{M}{N_0} \right) - \mathbf{P}\left(W \le V, W \le 2\frac{M}{N_0}\right) \mathbf{P}\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0} \right) \right]$$

$$\leq 4 \left(\frac{f_U}{f_L}\right)^2 \int_{-1}^{1} \int_{-1}^{1} S(t_1, t_2) dt_1 dt_2$$

$$= 4 \left(\frac{f_U}{f_L}\right)^2 \left[\int_{0}^{1} \int_{0}^{1} S(t_1, t_2) dt_1 dt_2 + \int_{-1}^{0} \int_{-1}^{0} S(t_1, t_2) dt_1 dt_2$$

$$+ \int_{-1}^{0} \int_{0}^{1} S(t_1, t_2) dt_1 dt_2 + \int_{0}^{1} \int_{-1}^{0} S(t_1, t_2) dt_1 dt_2\right]$$

$$= 4 \left(\frac{f_U}{f_L}\right)^2 \left[\int_{0}^{1} \int_{0}^{1} S(t_1, t_2) dt_1 dt_2 + \int_{-1}^{0} \int_{-1}^{0} S(t_1, t_2) dt_1 dt_2 + 2 \int_{0}^{1} \int_{-1}^{0} S(t_1, t_2) dt_1 dt_2\right], \quad (A.42)$$
where the last step is from the symmetry of $S(t_1, t_2)$

where the last step is from the symmetry of $S(t_1, t_2)$.

For the first term in (A.42), by the symmetry of $S(t_1, t_2)$ and the Chernoff bound,

$$\begin{split} &\int_{0}^{1} \int_{0}^{1} S(t_{1}, t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} \leq \int_{0}^{\infty} \int_{0}^{\infty} S(t_{1}, t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} = 2 \int_{0}^{\infty} \int_{0}^{\infty} S(t_{1}, t_{2}) \mathbb{1}(t_{1} \geq t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} \\ \leq & 2 \int_{0}^{\infty} \int_{0}^{\infty} \mathrm{P}\Big(U_{(M)} \geq \frac{M}{N_{0}} (1 + t_{1} \lor t_{2})\Big) \mathbb{1}(t_{1} \geq t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} = 2 \int_{0}^{\infty} t \mathrm{P}\Big(U_{(M)} \geq \frac{M}{N_{0}} (1 + t)\Big) \mathrm{d}t \\ \leq & 2 \int_{0}^{\infty} t(1 + t)^{M} e^{-Mt} \mathrm{d}t. \end{split}$$

Notice that

$$\begin{split} &\int_{0}^{\infty} t(1+t)^{M} e^{-Mt} dt = \int_{0}^{\infty} (1+t)^{M+1} e^{-Mt} dt - \int_{0}^{\infty} (1+t)^{M} e^{-Mt} dt \\ &= -\frac{1}{M} \Big(-1 - (M+1) \int_{0}^{\infty} (1+t)^{M} e^{-Mt} dt \Big) - \int_{0}^{\infty} (1+t)^{M} e^{-Mt} dt \\ &= \frac{1}{M} + \frac{1}{M} \int_{0}^{\infty} (1+t)^{M} e^{-Mt} dt = \frac{1}{M} + \frac{e^{M}}{M} \int_{1}^{\infty} t^{M} e^{-Mt} dt \le \frac{1}{M} + \frac{e^{M}}{M} \int_{0}^{\infty} t^{M} e^{-Mt} dt \\ &= \frac{1}{M} + \frac{e^{M}}{M} \frac{1}{M^{M+1}} \int_{0}^{\infty} t^{M} e^{-t} dt = \frac{1}{M} + \frac{e^{M}}{M^{M+2}} \Gamma(M+1) = \frac{1}{M} + \frac{e^{M}}{M^{M+2}} \sqrt{2\pi M} \Big(\frac{M}{e} \Big)^{M} (1+o(1)) \\ &= \frac{1}{M} (1+o(1)), \end{split}$$
(A.43)

where the second last inequality from Stirling's approximation since $M \to \infty$. We then obtain

$$\int_{0}^{1} \int_{0}^{1} S(t_1, t_2) \mathrm{d}t_1 \mathrm{d}t_2 \le \frac{2}{M} (1 + o(1)). \tag{A.44}$$

For the second term in (A.42),

$$\begin{split} &\int_{-1}^{0} \int_{-1}^{0} S(t_{1}, t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} = 2 \int_{-1}^{0} \int_{-1}^{0} S(t_{1}, t_{2}) \mathbb{1}(t_{1} \leq t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} \leq 2 \int_{-1}^{0} \int_{-1}^{0} S(t_{1}, t_{2}) \mathbb{1}(t_{1} \leq t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} \\ \leq 2 \int_{-1}^{0} \int_{-1}^{0} \mathbb{P}\Big(U_{(M)} \leq \frac{M}{N_{0}} (1 + t_{1} \wedge t_{2})\Big) \mathbb{1}(t_{1} \leq t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} = 2 \int_{-1}^{0} (-t) \mathbb{P}\Big(U_{(M)} \leq \frac{M}{N_{0}} (1 + t)\Big) \mathrm{d}t \\ = 2 \int_{0}^{1} t \mathbb{P}\Big(U_{(M)} \leq \frac{M}{N_{0}} (1 - t)\Big) \mathrm{d}t \leq 2 \int_{0}^{1} t (1 - t)^{M} e^{Mt} \mathrm{d}t. \end{split}$$
 Notice that

Notice that

$$\int_0^1 t(1-t)^M e^{Mt} dt = -\int_0^1 (1-t)^{M+1} e^{Mt} dt + \int_0^1 (1-t)^M e^{Mt} dt$$

$$= -\frac{1}{M} \Big(-1 + (M+1) \int_0^1 (1-t)^M e^{Mt} dt \Big) + \int_0^1 (1-t)^M e^{Mt} dt \le \frac{1}{M}.$$
 (A.45)

We then obtain

$$\int_{-1}^{0} \int_{-1}^{0} S(t_1, t_2) \mathrm{d}t_1 \mathrm{d}t_2 \le \frac{2}{M}.$$
(A.46)

For the third term in (A.42),

$$\begin{split} &\int_{0}^{1} \int_{-1}^{0} S(t_{1}, t_{2}) \mathrm{d}t_{1} \mathrm{d}t_{2} \\ &= \int_{0}^{1} \int_{-t_{1}}^{0} \mathrm{P}\Big(U_{(M)} \geq \frac{M}{N_{0}}(1+t_{1})\Big) \mathrm{d}t_{1} \mathrm{d}t_{2} + \int_{0}^{1} \int_{-1}^{-t_{1}} \mathrm{P}\Big(U_{(M)} \leq \frac{M}{N_{0}}(1+t_{2})\Big) \mathrm{d}t_{1} \mathrm{d}t_{2} \\ &= \int_{0}^{1} t \mathrm{P}\Big(U_{(M)} \geq \frac{M}{N_{0}}(1+t)\Big) \mathrm{d}t + \int_{-1}^{0} (-t) \mathrm{P}\Big(U_{(M)} \leq \frac{M}{N_{0}}(1+t)\Big) \mathrm{d}t \\ &\leq \int_{0}^{\infty} t \mathrm{P}\Big(U_{(M)} \geq \frac{M}{N_{0}}(1+t)\Big) \mathrm{d}t + \int_{-1}^{0} (-t) \mathrm{P}\Big(U_{(M)} \leq \frac{M}{N_{0}}(1+t)\Big) \mathrm{d}t \\ &\leq \frac{1}{M}(1+o(1)) + \frac{1}{M} = \frac{2}{M}(1+o(1)), \end{split}$$

where the last step is from (A.43) and (A.45).

We then obtain

$$\int_{0}^{1} \int_{-1}^{0} S(t_1, t_2) dt_1 dt_2 \le \frac{2}{M} (1 + o(1)).$$
(A.47)

Plugging (A.44), (A.46), (A.47) into (A.42) yields

$$\left(\frac{N_0}{M}\right)^2 \left[\mathbf{P}\left(W \le V, \widetilde{W} \le \widetilde{V}, W \le 2\frac{M}{N_0}, \widetilde{W} \le 2\frac{M}{N_0} \right) - \mathbf{P}\left(W \le V, W \le 2\frac{M}{N_0}\right) \mathbf{P}\left(\widetilde{W} \le \widetilde{V}, \widetilde{W} \le 2\frac{M}{N_0}\right) \right]$$

$$\le 32 \left(\frac{f_U}{f_L}\right)^2 \frac{1}{M} (1 + o(1)),$$
(A.48)

and thus completes the proof.

References

- Abadie, A. and Imbens, G. W. (2006). Large sample properties of matching estimators for average treatment effects. *Econometrica*, 74(1):235–267.
- Abadie, A. and Imbens, G. W. (2008). On the failure of the bootstrap for matching estimators. *Econometrica*, 76(6):1537–1557.
- Abadie, A. and Imbens, G. W. (2011). Bias-corrected matching estimators for average treatment effects. *Journal of Business and Economic Statistics*, 29(1):1–11.
- Abadie, A. and Imbens, G. W. (2012). A martingale representation for matching estimators. Journal of the American Statistical Association, 107(498):833–843.
- Bang, H. and Robins, J. M. (2005). Doubly robust estimation in missing data and causal inference models. *Biometrics*, 61(4):962–973.

- Bentley, J. L. (1975). Multidimensional binary search trees used for associative searching. Communications of the ACM, 18(9):509–517.
- Berrett, T. B., Samworth, R. J., and Yuan, M. (2019). Efficient multivariate entropy estimation via k-nearest neighbour distances. The Annals of Statistics, 47(1):288–318.
- Bhattacharya, B. B. (2019). A general asymptotic framework for distribution-free graph-based two-sample tests. *Journal of the Royal Statistical Society. Series B*, 81(3):575–602.
- Biau, G. and Devroye, L. (2015). Lectures on the nearest neighbor method. Springer Series in the Data Sciences. Springer, Cham.
- Billingsley, P. (1995). Probability and measure (3rd Edition). Wiley Series in Probability and Mathematical Statistics. John Wiley and Sons, Inc., New York, NY.
- Borgeaud, S., Mensch, A., Hoffmann, J., Cai, T., Rutherford, E., Millican, K., Driessche, G. v. d., Lespiau, J.-B., Damoc, B., Clark, A., et al. (2021). Improving language models by retrieving from trillions of tokens. arXiv preprint arXiv:2112.04426.
- Boucheron, S., Lugosi, G., and Massart, P. (2013). Concentration inequalities: A nonasymptotic theory of independence. Oxford University Press.
- Brookhart, M. A., Schneeweiss, S., Rothman, K. J., Glynn, R. J., Avorn, J., and Stürmer, T. (2006). Variable selection for propensity score models. *American Journal of Epidemiology*, 163(12):1149–1156.
- Brown, R. A. (2015). Building k-d tree in O(knlogn) time. Journal of Computer Graphics Techniques, 4(1):50–68.
- Chapin, F. S. (1947). Experimental designs in sociological research. Harper and Brothers.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68.
- Cochran, W. G. and Rubin, D. B. (1973). Controlling bias in observational studies: A review. Sankhyā: The Indian Journal of Statistics, Series A, 35(4):417–446.
- Cormen, T. H., Leiserson, C. E., Rivest, R. L., and Stein, C. (2009). Introduction to algorithms. MIT press.
- Cover, T. M. and Thomas, J. (2006). *Elements of information theory (2nd Edition)*. John Wiley and Sons.
- Cunningham, S. (2021). Causal inference: the mixtape. Yale University Press.
- Devroye, L., Györfi, L., Lugosi, G., and Walk, H. (2017). On the measure of Voronoi cells. *Journal* of Applied Probability, 54(2):394–408.

Ditzian, Z. and Totik, V. (2012). Moduli of smoothness. Springer.

- Durrett, R. (2019). Probability: theory and examples (5th Edition). Cambridge University Press.
- Evans, L. C. and Garzepy, R. F. (2018). Measure theory and fine properties of functions. Routledge.
- Friedman, J. H., Bentley, J. L., and Finkel, R. A. (1977). An algorithm for finding best matches in logarithmic expected time. ACM Transactions on Mathematical Software, 3(3):209–226.
- Friedman, J. H. and Rafsky, L. C. (1979). Multivariate generalizations of the Wald-Wolfowitz and Smirnov two-sample tests. *The Annals of Statistics*, 7(4):697–717.
- Greenwood, E. (1945). *Experimental sociology*. Columbia University Press.
- Hahn, J. (1998). On the role of the propensity score in efficient semiparametric estimation of average treatment effects. *Econometrica*, 66(2):315–331.
- Han, Y., Jiao, J., Weissman, T., and Wu, Y. (2020). Optimal rates of entropy estimation over Lipschitz balls. *The Annals of Statistics*, 48(6):3228–3250.
- Henze, N. (1988). A multivariate two-sample test based on the number of nearest neighbor type coincidences. The Annals of Statistics, 16(2):772–783.
- Henze, N. and Penrose, M. D. (1999). On the multivariate runs test. *The Annals of Statistics*, 27(1):290–298.
- Ho, D. E., Imai, K., King, G., and Stuart, E. A. (2007). Matching as nonparametric preprocessing for reducing model dependence in parametric causal inference. *Political Analysis*, 15(3):199–236.
- Imbens, G. W. (2004). Nonparametric estimation of average treatment effects under exogeneity: A review. Review of Economics and Statistics, 86(1):4–29.
- Imbens, G. W. and Rubin, D. B. (2015). *Causal inference in statistics, social, and biomedical sciences.* Cambridge University Press.
- Kpotufe, S. (2017). Lipschitz density-ratios, structured data, and data-driven tuning. In 2017 International Conference on Artificial Intelligence and Statistics, pages 1320–1328. PMLR.
- Kremer, J., Gieseke, F., Pedersen, K. S., and Igel, C. (2015). Nearest neighbor density ratio estimation for large-scale applications in astronomy. *Astronomy and Computing*, 12:67–72.
- Lima, M., Cunha, C. E., Oyaizu, H., Frieman, J., Lin, H., and Sheldon, E. S. (2008). Estimating the redshift distribution of photometric galaxy samples. *Monthly Notices of the Royal Astronomical Society*, 390(1):118–130.
- Lin, Z. and Han, F. (2021). On boosting the power of Chatterjee's rank correlation. arXiv preprint arXiv:2108.06828.

- Liu, R. Y. and Singh, K. (1993). A quality index based on data depth and multivariate rank tests. Journal of the American Statistical Association, 88(421):252–260.
- Morgan, S. L. and Harding, D. J. (2006). Matching estimators of causal effects: Prospects and pitfalls in theory and practice. *Sociological Methods and Research*, 35(1):3–60.
- Newey, W. K. (1997). Convergence rates and asymptotic normality for series estimators. *Journal* of *Econometrics*, 79(1):147–168.
- Nguyen, X., Wainwright, M. J., and Jordan, M. I. (2010). Estimating divergence functionals and the likelihood ratio by convex risk minimization. *IEEE Transactions on Information Theory*, 56(11):5847–5861.
- Noshad, M., Moon, K. R., Sekeh, S. Y., and Hero, A. O. (2017). Direct estimation of information divergence using nearest neighbor ratios. In 2017 IEEE International Symposium on Information Theory (ISIT), pages 903–907.
- Póczos, B. and Schneider, J. (2011). On the estimation of alpha-divergences. In 2011 International Conference on Artificial Intelligence and Statistics, pages 609–617.
- Romanovsky, V. (1923). Note on the moments of a binomial $(p+q)^n$ about its mean. *Biometrika*, 15(3/4):410–412.
- Rosenbaum, P. R. (2010). Design of observational studies. Springer.
- Rosenbaum, P. R. and Rubin, D. B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55.
- Rubin, D. B. (1973). Matching to remove bias in observational studies. *Biometrics*, 29(1):159–183.
- Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology*, 66(5):688–701.
- Rubin, D. B. (2006). Matched sampling for causal effects. Cambridge University Press.
- Rubin, D. B. and Thomas, N. (2000). Combining propensity score matching with additional adjustments for prognostic covariates. *Journal of the American Statistical Association*, 95(450):573–585.
- Scharfstein, D. O., Rotnitzky, A., and Robins, J. M. (1999). Adjusting for nonignorable dropout using semiparametric nonresponse models. *Journal of the American Statistical Association*, 94(448):1096–1120.
- Sekhon, J. S. (2008). Multivariate and propensity score matching software with automated balance optimization: the matching package for R. Journal of Statistical Software, 42(7):1–52.
- Shi, H., Drton, M., and Han, F. (2021a). On Azadkia-Chatterjee's conditional dependence coefficient. Technical report, Department of Statistics, University of Washington.
- Shi, H., Drton, M., and Han, F. (2021+b). On the power of Chatterjee's rank correlation. *Biometrika*. (in press).
- Smith, H. L. (1997). Matching with multiple controls to estimate treatment effects in observational studies. Sociological Methodology, 27(1):325–353.
- Stein, E. M. (2016). Singular integrals and differentiability properties of functions. Princeton University Press.
- Sugiyama, M., Suzuki, T., and Kanamori, T. (2012). Density ratio estimation in machine learning. Cambridge University Press.
- Sugiyama, M., Suzuki, T., Nakajima, S., Kashima, H., von Bünau, P., and Kawanabe, M. (2008). Direct importance estimation for covariate shift adaptation. Annals of the Institute of Statistical Mathematics, 60(4):699–746.
- Totik, V. (1994). Approximation by Bernstein polynomials. *American Journal of Mathematics*, 116(4):995–1018.
- Tsybakov, A. B. (2009). Introduction to nonparametric estimation. Springer.
- van der Vaart, A. W. (1998). Asymptotic statistics. Cambridge University Press.
- Vapnik, V. (2006). Estimation of dependences based on empirical data. Springer.
- Voronoi, G. (1908). Nouvelles applications des paramètres continus à la théorie des formes quadratiques. deuxième mémoire. recherches sur les parallélloèdres primitifs. Journal für die reine und angewandte Mathematik (Crelles Journal), 1908(134):198–287.
- Wald, A. and Wolfowitz, J. (1940). On a test whether two samples are from the same population. Annals of Mathematical Statistics, 11(2):147–162.
- Zhao, P. and Lai, L. (2020a). Analysis of KNN density estimation. arXiv preprint arXiv:2010.00438.
- Zhao, P. and Lai, L. (2020b). Minimax optimal estimation of KL divergence for continuous distributions. *IEEE Transactions on Information Theory*, 66(12):7787–7811.