

WEIERSTRASS SEMIGROUPS FROM CYCLIC COVERS OF HYPERELLIPTIC CURVES

ETHAN COTTERILL, NATHAN PFLUEGER, AND NAIZHEN ZHANG

ABSTRACT. The *Weierstrass semigroup* of pole orders of meromorphic functions in a point p of a smooth algebraic curve C is a classical object of study; a celebrated problem of Hurwitz is to characterize which semigroups $S \subset \mathbb{N}$ with finite complement are *realizable* as Weierstrass semigroups $S = S(C, p)$. In this note, we establish realizability results for cyclic covers $\pi : (C, p) \rightarrow (B, q)$ of hyperelliptic targets B marked in hyperelliptic Weierstrass points; and we show that realizability is dictated by the behavior under j -fold multiplication of certain divisor classes in hyperelliptic Jacobians naturally associated to our cyclic covers, as j ranges over all natural numbers.

CONTENTS

1. Introduction	1
1.1. Roadmap	2
2. Background	4
3. Weierstrass-realizability for $2N$ -semigroups	12
4. The multiplication profile of a point on a hyperelliptic Jacobian	22
5. Explicit realizability constructions	27
Appendix A. Separability results for univariate polynomials	30
Appendix B. Flynn’s models	32
References	35

1. INTRODUCTION

Hurwitz spaces of covers $C \rightarrow B$ of algebraic curves of fixed genus have long played an important role in algebraic geometry and number theory. *Cyclic* covers are distinguished by their simplicity; indeed, the geometry of a cyclic cover is controlled by its branch locus on the target curve B . It is natural to ask, in particular, about the behavior of *Weierstrass semigroups* in ramification points of a cyclic cover; this is a cyclic cover version of Hurwitz’ celebrated *realization problem* for Weierstrass semigroups. Resolving it, however, is untenable without introducing further hypotheses. In this paper, we address the realization problem for ramification points p of *marked* cyclic covers $(C, p) \rightarrow (B, q)$ whose targets are hyperelliptic curves marked in hyperelliptic Weierstrass points. In this context it turns out that realizability is controlled by the multiplicative behavior of divisor classes in the Jacobian of B that are naturally extracted from the branch divisors of our covers.

During much of the preparation of this work, Naizhen Zhang was supported by the DFG Priority Programme 2026 “Geometry at infinity”.

To relate the realizability of semigroups at totally-ramified points of covers to multiplication in the Jacobians of hyperelliptic targets, we make essential use of Mumford’s *reduced* presentation for divisors on a hyperelliptic curve; see Section 2.3. Specifically, we always reduce with respect to the fixed Weierstrass point q , and use presentations that split as an effective divisor minus a multiple of q . As the behavior of arbitrary degree-zero divisors under multiplication is extremely complicated, however, we focus primarily on those elements $[p - q]$ of the Jacobian that belong to image of (B, q) under the Abel–Jacobi map. In particular, the study of *torsion* points in hyperelliptic Jacobians that arise as Abel–Jacobi images is an active area of inquiry in its own right, with links to Diophantine solutions of generalized Pell equations (see [17, 18]) and the dynamics of billiards in ellipsoids (see [11, 6]). Here we introduce *multiplication profiles* for points of odd hyperelliptic Jacobians marked in specified Weierstrass points q , which characterize the sizes of the supports of their j -fold multiples’ q -reductions for every positive integer j ; we give a complete characterization of those multiplication profiles associated with (the branch divisors of) cyclic triple covers; and we produce a list of numerical conditions that need to be satisfied (and which we conjecture to be sufficient) for a sequence of integers to arise from some point class $[p - q]$.

We then focus in more detail on cyclic covers whose underlying branch divisors represent Abel–Jacobi images of torsion classes. When the covering degree N is sufficiently small relative to γ (roughly, whenever $N \leq 2\gamma + 8$) we use earlier work of Flynn to build examples of marked hyperelliptic curves (B, q) equipped with points p whose associated multiplication profiles exhaust all conjectural possibilities. We also construct an explicit quasi-projective parameter space R_N for marked hyperelliptic curves equipped with N -torsion points, and we make an explicit conjecture about the stratification of R_N according to multiplication profile whenever $N \leq 2\gamma + 6$.

1.1. Roadmap. The material following this introduction is organized as follows. In Section 2, we recall the basic theory of cyclic covers, and of reduced divisors on hyperelliptic curves. Any cyclic cover is specified by *covering data* on the target variety; and for *simple* cyclic covers, our primary focus here, covering data consists of the branch divisor D , together with a line bundle L for which $L^{\otimes N} \cong \mathcal{O}(D)$. Lemma 2.13 computes the pushforward to the target of twists of the structure sheaf of the domain curve (C, p) of a simple cyclic cover by multiples of a ramification point p , and as such determines how the Weierstrass semigroup in p is controlled cohomologically on the hyperelliptic target (B, q) . On the other hand, Mumford showed that every divisor class on an odd hyperelliptic curve (B, q) admits a representative that is *semi-reduced* with respect to q , and which may be codified as a pair of univariate polynomials; and Cantor explained how to further algorithmically obtain a q -*reduced* pair of polynomials. See Lemmas 2.17 and 2.18, respectively. Cantor’s algorithm allows us to immediately derive an explicit characterization of those odd hyperelliptic curves $y^2 = f(x)$ with torsion points of orders $2\gamma + i$, $i = 1, 2, 3$, where $\deg f(x) = 2\gamma + 1$; see Corollary 2.20. In light of Zarhin’s [19, Cor. 2.8], these are the lowest possible orders greater than 2.

In Section 3, we begin our investigation of realizability for Weierstrass semigroups arising from cyclic covers of hyperelliptic curves (B, q) in earnest. To avoid an additional layer of cohomological complexity, we assume that the genus g of our

domain curve C is always at least $(2N-1)$ times the genus γ of the target B . As our marked points of reference q are always Weierstrass points, numerical semigroups S that may arise from our N -fold cyclic covers are all of multiplicity $2N$. As such, S has a *standard basis* (e_0, \dots, e_{2N-1}) comprised of nonzero elements of S that are minimal in their respective equivalence classes modulo $2N$. Lemma 3.9 establishes that the semigroup with standard basis (e_0, \dots, e_{2N-1}) is realizable if and only if there exists a pair (L, D) with $L^{\otimes N} \cong \mathcal{O}(D)$ for which L^{-j} has a meromorphic section with exact pole order $a_j := \min(e_j, e_{N+j})$ for every $j = 1, \dots, N-1$.¹

In Section 3.2, we obtain explicit numerical restrictions on the set of admissible tuples (a_1, \dots, a_N) underlying Weierstrass-realizable semigroups S . Theorem 3.12 gives a restriction based on the cohomological Lemma 2.13, while further inequalities singled out by Morrison and Pinkham [13] arise from the additive structure of S . Taken together, these inequalities define what we call the *feasible set* $F(N)$ of $(N-1)$ -tuples (a_1, \dots, a_{N-1}) potentially associated to cyclic covers. As we explain in Example 3.14, the fact that every element of $F(2)$ indexes a realizable semigroup is a straightforward consequence of the existence of q -reduced representatives in each divisor class; so the smallest “interesting” covering degree is $N = 3$. Lemma 3.16 reframes the realizability of the semigroup indexed by (a_1, \dots, a_{N-1}) as a statement about the q -reduced representations of integer multiples of a single effective divisor E_1 fixed at the outset.

Accordingly, we turn our attention to the effective part E_j of the q -reduced representative of each multiple jE_1 , $j \in \mathbb{N}$. Their degrees comprise the *multiplication profile* of the cyclic cover determined by $L = \mathcal{O}(E_1)$ and its N -th tensor power. A given multiplication profile $(d_j)_{j \in \mathbb{N}}$ is realized by a degree- N cyclic cover of (B, q) precisely when $\Theta_{d_1} \cap \bigcap_{j=2}^{N-1} m_j^{-1} \Theta_{d_j}$ is nonempty, where m_j denotes j -fold multiplication on the Jacobian of B and Θ_{d_j} denotes the image of the d_j -th Abel–Jacobi map with base point q . Moreover, the multiplication profiles given by $d_j = 0$ for every j and $d_j = \gamma$ for every j are always associated with realizable semigroups, and correspond to extremal elements in $F(N)$. Lemma 3.20 establishes basic properties of multiplication profiles, which we in turn use in Section 3.4 to classify all realizable semigroups arising from cyclic covers with either elliptic targets or covering degree 4 and target genus 2. In Theorem 3.22, we classify all realizable semigroups arising from cyclic triple covers via an inductive divisor-theoretic analysis on (B, q) .

In Section 4, we initiate a systematic investigation of multiplication profiles of *point classes* $[p - q]$ on hyperelliptic Jacobians. Lemma 4.4, Corollary 4.5, Lemma 4.6, and Zarhin’s nonexistence theorem [19, Cor. 2.8] for torsion points of orders $2 < N < 2\gamma - 1$ impose the main restrictions on these sequences in general; our expectation that there are no additional restrictions is formalized in Conjecture 4.9. Our main tools for tackling this conjecture are Lemma 4.11, which characterizes the rational function $p_j(x) + q_j(x)y$ on $B : y^2 = f(x)$ whose associated divisor represents d_j ; and its consequence Lemma 4.14, which establishes that profile entries d_j less than γ are codified by the vanishing of certain explicit determinants D_j in the \mathbb{P}^{N-1} of coefficients of degree at most $N-1$ of power series $\sqrt{f(x)}$. As we explain in the discussion following Lemma 4.14, every multiplication profile of a point class is uniquely characterized by the vanishing or non-vanishing of the determinants D_j .

¹We assume thereafter that the branch divisor D is a reduced scheme.

We put these tools to work in Section 4.2, in which we study the multiplication profiles of torsion point classes. The multiplication profile of an N -torsion point class is uniquely determined by the restricted sequence of d_j for j between $\gamma + 2$ and $\lfloor \frac{N}{2} \rfloor$, and in this range it is uniquely determined by the vanishing or non-vanishing of the corresponding D_j ; see Proposition 4.16. Similarly, N -torsionness is codified by a determinantal condition in \mathbb{P}^{N-1} ; accordingly, we are led to define a parameter space R_N of separable polynomials in the coefficients of degree at most $N - 1$ of power series in x that satisfy the determinantal conditions imposed by N -torsionness and quadric *polynomiality relations* that ensure that the associated power series square to polynomials. The space R_N , in turn, is stratified by multiplication profiles defined by additional determinantal conditions D_j . Strikingly, all of the determinantal conditions operative here are defined by maximal minors of Hankel matrices. Conjecture 4.18 gives an explicit prediction about the irreducible decomposition of R_N in the first two nontrivial cases of $N = 2\gamma + 4$ and $N = 2\gamma + 6$. In these cases, each irreducible component selects for a particular multiplication profile.

Finally, in Section 5 we leverage Mumford presentations, Cantor's reduction algorithm, and earlier work of Flynn [9] to produce explicit models of hyperelliptic curves with torsion orders $N \leq 2\gamma + 9$; whenever $N \leq 2\gamma + 7$, our models realize all possible multiplication profiles. In Example 5.11, we combinatorially analyze those sequences of numbers eligible to be multiplication profiles of N -torsion point classes, and show that in each respective parity, the number of eligible sequences is a Fibonacci number. Appendix A is a collection of separability results for polynomials that are needed in order to certify that our models are smooth, while Appendix B is a detailed treatment of (our adaptations of) Flynn's models.

1.2. Conventions. In this paper, we work over an algebraically closed field k of characteristic zero. A variety over k will always mean an integral, separated scheme of finite type over k unless otherwise specified.

1.3. Acknowledgements. We thank Joe Harris for illuminating conversations. Fernando Torres invited the first author to speak in Campinas on this topic in early 2020, and sadly left us before this project's completion. We dedicate this paper to his memory.

2. BACKGROUND

2.1. Cyclic covers of algebraic curves. In this section, we review the basic theory of cyclic covers of smooth projective algebraic curves.

Definition 2.1 (Cf. [15], [16]). Given a smooth variety Y and a finite abelian group G , a G -cover of Y is a finite, surjective morphism $\pi : X \rightarrow Y$ for which X is normal, $K(X)/K(Y)$ is Galois, and $\text{Gal}(K(X)/K(Y)) = G$.

Definition 2.1 has a number of immediate consequences. First of all, finiteness of π implies that it is affine, and therefore $X \cong \underline{\text{spec}}(\pi_*\mathcal{O}_X)$. Secondly, G acts on X and finiteness of G implies $Y = X/G$; moreover, G acts transitively on the fibers of π . Thirdly, whenever π is flat (e.g., when π is a finite map between smooth projective curves), $\pi_*\mathcal{O}_X$ is a locally-free sheaf on Y carrying the structure of an \mathcal{O}_Y -algebra with a G -action. It is therefore useful to work in the category of G -sheaves, whose basic features we now recall.

Definition 2.2 (See [3], Sec 4). Given a finite group G acting on the left of a scheme X by automorphisms, a G -sheaf is a quasicohent \mathcal{O}_X -module E together with a set of isomorphisms $\{\lambda_g^E : E \rightarrow g^*E \mid g \in G\}$ for which

- (1) $\lambda_{1_G}^E = \text{Id}_E$; and
- (2) $\lambda_{g_1 g_2}^E = g_2^*(\lambda_{g_1}^E) \circ \lambda_{g_2}^E$.

Given G -sheaves E and F , $g \in G$ acts on $\text{Hom}_{\mathcal{O}_X}(E, F)$ by sending f to $(\lambda_g^F)^{-1} \circ g^*f \circ \lambda_g^E$. We let $\text{G-Hom}_{\mathcal{O}_X}(E, F)$ denote the subspace of module morphisms invariant under this action.

One basic property of relevance to us is that the adjointness of pull-backs and pushforwards extends to the category of G -sheaves:

Lemma 2.3. [See *loc. cit.*] Given a G -equivariant morphism of schemes $f : X \rightarrow Y$ and G -sheaves E and F on Y and X resp., there are natural isomorphisms

$$\text{G-Hom}_{\mathcal{O}_X}(f^*E, F) \cong \text{G-Hom}_{\mathcal{O}_X}(E, f_*F).$$

In our study of degree- N cyclic covers $\pi : C \rightarrow B$ of curves, G will always be the cyclic Galois group $\text{Gal}(K(C)/K(B)) \cong \mu_N = \langle \xi \mid \xi^N = 1 \rangle$, which acts trivially on B and permutes points in the fibers of π . In particular, π is G -equivariant and for any G -line bundle L on C (resp. any line bundle M on B), L and π_*L (resp. M and π^*M) are G -sheaves on C (resp. B) and B (resp. C). In this context, Lemma 2.3 yields

$$(1) \quad \text{G-Hom}_{\mathcal{O}_C}(\pi^*M, L) \cong \text{G-Hom}_{\mathcal{O}_B}(M, \pi_*L).$$

According to definition 2.2, L being a G -line bundle requires that $L \cong g^*L$ for every $g \in G$. This holds in the following case, on which we will focus hereafter.

Example 2.4. Let $\pi : C \rightarrow B$ be a degree- N cyclic cover of smooth projective curves equipped with a totally ramified point $p \in C$. Then $\mathcal{O}_C(mp)$ is μ_N -invariant, as p is fixed by the action of the Galois group. As a result, $\pi_*\mathcal{O}_C(mp)$ is a locally-free G -sheaf on B and hence decomposes as a direct sum of line bundles

$$\pi_*\mathcal{O}_C(mp) = \bigoplus_{i=0}^{N-1} L_i$$

where $L_i = \ker(\lambda_\xi^{\pi_*\mathcal{O}_C(mp)} - \eta^i \cdot \text{Id})$, for some fixed primitive N -th root of unity η . Given our assumptions on the ground field as well as on the extension $K(C)/K(B)$, we have $K(C) = K(B)[x]/(x^N - \theta)$ for some $\theta \in K(B)$. In particular, $K(C)$ decomposes as $K(C) = \bigoplus_{i=0}^{N-1} (K(B) \cdot x^i)$, where $K(B) \cdot x^i$ is the eigenspace with eigenvalue η^i for the action of our pre-chosen generator ξ of the Galois group. It follows that $L_i(V) = \mathcal{O}_C(mp)(\pi^{-1}(V)) \cap (K(B) \cdot x^i)$ for every open set $V \subset B$.

2.2. Covering data. As we have already seen, every degree- N cyclic cover $\pi : C \rightarrow B$ of smooth projective curves induces a decomposition $\pi_*\mathcal{O}_C = \bigoplus_{i=0}^{N-1} L_i$, where $L_0 = \mathcal{O}_B$.

Example 2.5. Applying duality for finite morphisms of noetherian schemes ([12], Ex.III.6.10(b)), together with the fact that $\pi^!\omega_B \cong \omega_C$, it follows that $\pi_*\omega_C$ also explicitly decomposes as a direct sum of line bundles on B :

$$\pi_*\omega_C \cong (\pi_*\mathcal{O}_C)^{-1} \otimes \omega_B \cong \left(\bigoplus_{i=0}^{N-1} L_i^{-1} \right) \otimes \omega_B.$$

See [13, Theorem 3.5].

On the other hand, the general theory of abelian covers developed by Pardini ([15]) explains how to build covers out of certain geometric data on the target variety. In this subsection, we briefly review the main ingredients of Pardini's theory in the context most relevant to us.

Definition 2.6. ([15, Thm. 2.1]) Given a finite abelian group G , let \hat{G} (resp., \mathfrak{C}) denote its character group (resp., the set of cyclic subgroups of G). For every $H \in \mathfrak{C}$, let S_H denote a generating set for the character group of H . Given a smooth projective variety Y , a collection of G -building data on Y consists of the following:

- (1) For every $\chi \in \hat{G}$, an invertible sheaf L_χ on Y ; and
- (2) for every $H \in \mathfrak{C}$ and $\psi \in S_H$, a divisor $D_{H,\psi}$ on Y .

These satisfy the linear equivalence relations

$$(2) \quad L_{\chi_1} \otimes L_{\chi_2} \cong L_{\chi_1 \chi_2} \otimes \mathcal{O}_Y \left(\sum_{H \in \mathfrak{C}} \sum_{\psi \in S_H} \epsilon_{\chi_1, \chi_2}^{H, \psi} D_{H, \psi} \right)$$

for every $\chi_1, \chi_2 \in \hat{G}$, where the constant $\epsilon_{\chi_1, \chi_2}^{H, \psi}$ is determined by requiring that

$$\epsilon_{\chi_1, \chi_2}^{H, \psi} = \begin{cases} 0 & \text{if } i_1 + i_2 < |H| \\ 1 & \text{otherwise} \end{cases}$$

where $\chi_j|_H = \psi^{i_j}$ and $0 \leq i_j \leq |H| - 1$, $j \in \{1, 2\}$.

Remark 2.7. Starting from a given abelian cover $\pi : X \rightarrow Y$, we may recover building data for π as follows: we extract L_χ from the eigensheaf decomposition of $\pi_* \mathcal{O}_X$; while $D_{H,\psi}$ is the sum of irreducible components E of the branch divisor D of π for which the stabilizer group of any irreducible component of $\pi^{-1}(E)$ is H , and the induced action of H on the cotangent space at the generic point of such components is given by ψ .

Conversely, a collection of covering data as in Definition 2.6 canonically prescribes a morphism $\pi : X = \underline{\text{spec}}(\mathcal{O}_Y \oplus \bigoplus_\chi L_\chi^{-1}) \rightarrow Y$, where the algebra structure on $\mathcal{O}_Y \oplus \bigoplus_\chi L_\chi^{-1}$ is defined using the data $\{D_{H,\psi}\}_{H \in \mathfrak{C}, \psi \in S_H}$; see [15, Thm 2.1] for further details. This morphism satisfies all conditions in Definition 2.1, except that X may fail to be normal. Thus, in general one needs to pass to the the normalization X^ν of the previously-constructed variety X in order to obtain an abelian cover in the sense of Definition 2.1. In particular, the building data one starts with may not be equal to the ‘‘final’’ building data of the cover $\pi^\nu : X^\nu \rightarrow Y$: the line bundles L_χ are in general not the eigensheaves appearing in the decomposition of $\pi_* \mathcal{O}_X$, and $D = \sum D_{H,\psi}$ is not the branch divisor of π^ν .

For cyclic covers, the required building data reduces to the specification of a line bundle L on the target B together with effective divisors $\{D_{H,\psi}\}_{H \in \mathfrak{C}, \psi \in S_H}$ for which

$$|G| \cdot L \cong \mathcal{O}_B \left(\sum_{H, \psi} [G : H] r_{H, \psi} D_{H, \psi} \right)$$

where $\chi|_H = \psi^{r_{H, \psi}}$, for a pre-chosen generator χ of \hat{G} ; see [15, Eq. 2.18]. In the more special case of *simple* cyclic covers, the building data reduces to a pair (L, D) for which $L^{\otimes N} \cong \mathcal{O}_B(D)$, where $N = |G|$; see [15, Eq. 2.19].

Hereafter we will focus on simple cyclic covers of smooth projective curves, each of which is determined by a pair (L, D) for which $L^{\otimes N} \cong \mathcal{O}_B(D)$. More concretely, there is an \mathcal{O}_Y -algebra structure on $\bigoplus_{i=0}^{N-1} L^{-i}$ given by the natural morphisms

$$L^{k-N} \cong L^k \otimes \mathcal{O}_B(-D) \rightarrow L^k \otimes \mathcal{O}_B \cong L^k.$$

Just as in Remark 2.7, $\text{spec}(\bigoplus_{i=0}^{N-1} L^{-i})$ is not always normal and in general one needs to normalize; however, in this instance the outcome is well-known:

Lemma 2.8. *Let D be an effective divisor, L be an invertible sheaf on a smooth projective curve B , such that $L^{\otimes N} \cong \mathcal{O}_B(D)$. There is a degree- N cyclic cover $\pi : C \rightarrow B$ of smooth projective curves for which*

$$\pi_* \mathcal{O}_C \cong \bigoplus_{i=0}^{N-1} L^{-i} \left(\left[\frac{iD}{N} \right] \right)$$

which is the normalization of the cover

$$\text{spec} \left(\bigoplus_{i=0}^{N-1} L^{-i} \right) \rightarrow B.$$

Here $[\cdot]$ denotes the integral part of a \mathbb{Q} -Cartier divisor. The cover (2.8) is sometimes called the N -th extraction of D (see [8]).

Proof. This is a special case of [8, Lem. 1.8]. \square

Remark 2.9. We shall refer to pairs (L, D) such that $L^{\otimes N} \cong \mathcal{O}_B(D)$ on curves as *covering data*, so as to distinguish from *building data* in the sense of Pardini.

Unlike with building data, there is no natural way to assign covering data to a given cyclic cover:

Theorem 2.10 (Thm. 3.3, [13]). *Associated with each collection of covering data $(L, D = \sum a_k q_k)$, $1 \leq a_k < N$ for which $L^{\otimes N} \cong \mathcal{O}_B(D)$, there is a degree- N cyclic cover $\pi : C \rightarrow B$ branched over D_{red} , where $\pi^{-1}(q_k)$ consists of $\gcd(a_k, N)$ many points. Conversely, any degree- N cyclic cover $\pi : C \rightarrow B$ may be constructed from some covering data $(L, D = \sum a_k q_k)$ using the method of Lemma 2.8; the same cover may also be constructed using the covering data $(L^{(i)}, D^{(i)})$, where $i < N$, $\gcd(i, N) = 1$ and*

$$L^{(i)} = L^i \left(- \left[\frac{iD}{N} \right] \right), D^{(i)} = \sum a_k^{(i)} q_k$$

with $a_k^{(i)} \equiv ia_k \pmod{N}$ and $1 \leq a_k^{(i)} < N$.

Proof. The construction of a cyclic cover from covering data (L, D) is the content of Lemma 2.8. It is easy to see that π is branched over D_{red} , as $\pi_* \mathcal{O}_C|_{B \setminus D_{\text{red}}} \cong \mathcal{O}_{B \setminus D_{\text{red}}}^{\oplus N}$. The fact that $\pi^{-1}(q_k)$ consists of $\gcd(a_k, N)$ many points follows from the fact that $K(C)/K(B)$ is a Kummer extension of function fields.²

Conversely, given any degree- N cyclic cover $\pi : C \rightarrow B$, we have $\pi_* \mathcal{O}_C \cong \mathcal{O}_Y \oplus \bigoplus_{\chi} L_{\chi}$. Choose $L = L_{\chi}^{-1}$ for some χ generating $\widehat{\mu}_N$, together with a section s of $L^{\otimes N}$, and let $D = \text{div}(s)$; then up to isomorphism of covers, (L, D) is a collection

²Recall also that in this case all ramification points over a given point q_k in the branch locus have the same ramification index $N_k = \frac{N}{\gcd(a_k, N)}$.

of covering data for π . Moreover, whenever $\gcd(i, N) = 1$ and $j \in [1, N]$ is such that $i \cdot j \equiv 1 \pmod{N}$, we have $(L^{(i)})^{(j)} = L$ and $(D^{(i)})^{(j)} = D$. Thus, $(L^{(i)}, D^{(i)})$ determines a collection of covering data for π whenever (L, D) does. \square

Remark 2.11. In particular, π is totally ramified at all ramification points whenever N is prime.

Now suppose $D = \sum a_k q_k$ and that p is a ramification point of π with image $\pi(p) = q_k$. Set $N_k := \frac{N}{\gcd(a_k, N)}$, $a'_k := \frac{a_k}{\gcd(a_k, N)}$. Recall that $K(C) = K(B)[x]/(x^N - \theta)$; so $x^n = \theta$. As $v_{q_k}(\theta) = a_k$, it follows that $v_p(\theta) = a_k N_k$ and therefore $v_p(x) = a'_k$.

On the other hand, recall that $L^{-i}([\frac{iD}{N}])$ is the eigensheaf associated to η^i , where η is our pre-chosen primitive N -th root; so $\pi^* L^{-i}([\frac{iD}{N}])$ is the invertible sub-sheaf of \mathcal{O}_C locally generated by some multiple of x^i . In particular, we deduce the following:

Lemma 2.12. *Suppose that $\pi(p) = q_1$. The image of $(\pi^* L^{-i}([\frac{iD}{N}]))_p$ in $\mathcal{O}_{C,p}$ is \mathfrak{m}_p^b , where b is the unique integer in $[0, N_1 - 1]$ congruent to ia'_1 modulo N_1 .*

Proof. Elements in $(\pi^* L^{-i}([\frac{iD}{N}]))_p$ are represented by functions of the form $f x^i$, where $f \in K(B)$ is regular near p . Let t_{q_1} be a uniformizer at $q_1 \in B$. The desired conclusion follows from the facts that $v_p(\pi^* t_{q_1}) = N_1$ and $v_p(x^i) = i \cdot a'_1$. \square

Lemma 2.13. *Suppose $\pi : C \rightarrow B$ is a cyclic cover constructed out of the covering data $(L, D = \sum a_i q_i)$ for which $L^{\otimes N} \cong \mathcal{O}_B(D)$. Fix $p \in C$ such that $q_1 = \pi(p)$, and suppose π is totally ramified at p . Then for every non-negative integer m , we have*

$$(3) \quad \pi_* \mathcal{O}_C(mp) = \bigoplus_{i=0}^{N-1} (L^{(i)})^{-1} \left(\left\lfloor \frac{m}{N} + \left\langle \frac{ia_1}{N} \right\rangle \right\rfloor q_1 \right)$$

where $L^{(i)} = L^i(-[\frac{iD}{N}])$, for $i = 0, \dots, N-1$, and $\langle c \rangle$ denotes the fractional part of a positive rational number c .

Proof. As discussed in Example 2.4, $\pi_* \mathcal{O}_C(mp)$ decomposes as a direct sum of line bundles $\pi_* \mathcal{O}_C(mp) = \bigoplus_{i=0}^{N-1} L_i$, where L_i is the eigensheaf on which a fixed generator ξ of μ_N acts by multiplication-by- η^i , for some preselected primitive N -th root of unity η .

Now let $U = B \setminus \{q_1\}$. As $\mathcal{O}_C(mp)|_{\pi^{-1}(U)} = \mathcal{O}_C|_{\pi^{-1}(U)}$, we have $\pi_* \mathcal{O}_C(mp)|_U = \pi_* \mathcal{O}_C|_U = \bigoplus_{i=0}^{N-1} (L^{(i)})^{-1}|_U$. This means, in particular, that

$$\pi_* \mathcal{O}_C(mp) = \bigoplus_{i=0}^{N-1} (L^{(i)})^{-1}(n_i q_1)$$

for certain integers n_i , and it remains to determine these. For this purpose, we use the natural isomorphism of Equation (1), which yields

$$\mu_N\text{-Hom } \mathcal{O}_B((L^{(i)})^{-1}(n_i q_1), \pi_* \mathcal{O}_C(mp)) \cong \mu_N\text{-}h_{\mathcal{O}_C}((\pi^*(L^{(i)})^{-1})(n_i N p), \mathcal{O}_C(mp)).$$

The upshot is that n_i is the maximal integer n for which there is a natural inclusion $\pi^*(L^{(i)})^{-1}(n_i q_1) \hookrightarrow \mathcal{O}_C(mp)$ of μ_N -sheaves.

According to Lemma 2.12, the image of $(\pi^* L^{(i)})^{-1}_p$ is \mathfrak{m}_p^b , where b is the unique integer between 1 and $N-1$ for which $b \equiv i \cdot a_1 \pmod{N}$. It follows that the image

of the natural μ_N -morphism $\pi^*(L^{(i)})^{-1} \rightarrow \mathcal{O}_C$ factors through $\mathcal{O}_C(-bp)$ (and not $\mathcal{O}_C(-(b+1)p)$). Hence, the natural inclusion $\pi^*(L^{(i)})^{-1}(nq_1) \hookrightarrow \mathcal{O}_C(mp)$ exists if and only if $nN - b \leq m$, i.e. $n \leq \frac{m+b}{N}$. As $b = ia_1 - \lfloor \frac{ia_1}{N} \rfloor N$, the desired result follows. \square

2.3. Reduced divisors on a hyperelliptic curve. One result that we will frequently use hereafter is the unique *reduced* presentation of divisor classes in a hyperelliptic Jacobian due to Mumford. This should not be confused with the usual notion of reducedness for a scheme.

Definition 2.14. Let B be a hyperelliptic curve of genus γ , and fix a Weierstrass point $q \in B$. A divisor D of degree zero on B is *reduced* (resp., *semi-reduced*) with respect to q whenever the following conditions (resp., the first two listed) are satisfied:

- (1) $D = \sum_{i=1}^n p_i - nq$, where p_i are all distinct from q .
- (2) For every $p \in B$, D is not supported simultaneously at p and $\iota(p)$, where ι is the hyperelliptic involution.
- (3) $n \leq \gamma$.

Lemma 2.15 ([14]). *Let B be a hyperelliptic curve of genus γ , and let $q \in B$ be a Weierstrass point. Each divisor class in $\text{Jac}(B)$ contains a unique representative that is reduced with respect to q .*

We shall use the uniqueness of reduced representatives in the following form:

Corollary 2.16. *Every effective divisor $D = \sum_{i=1}^k p_i$ on B for which $D - kq$ is reduced satisfies $h^0(\mathcal{O}_B(D)) = 1$ and $h^0(\mathcal{O}_B(D - q)) = 0$.*

Proof. For the sake of argument, suppose $h^0(\mathcal{O}_B(D)) > 1$; then there exists a non-constant function f whose pole divisor is bounded above by D . Up to reducing k , we may assume the pole divisor of f is precisely D . Thus, $D \sim D'$, where D' is also effective. Displacing multiples of q to the left-hand side of the linear equivalence if necessary (and then subtracting appropriate multiples of q on both sides), we obtain two distinct reduced divisors that are linearly equivalent, a contradiction. The same argument also yields the second assertion. \square

Following Mumford, every semi-reduced divisor on an odd hyperelliptic curve may be presented in a unique way by a pair of univariate polynomials:

Lemma 2.17. *Let B be a genus γ hyperelliptic curve with equation $y^2 = f(x)$, where $f(x)$ is a monic and separable polynomial of degree $2\gamma+1$. Let $D = \sum_{i=1}^s p_i - sq$ be a divisor on B that is semi-reduced with respect to q .³ Then there exists a unique pair of polynomials $u(x), v(x) \in k[x]$ satisfying the following conditions:*

- (1) u is monic and $\deg u(x) = s$;
- (2) $\deg v(x) < \deg u(x)$;
- (3) $u(x) \mid (f(x) - v^2(x))$; and
- (4) $p_i = (a_i, v(a_i))$, where a_1, \dots, a_s are the roots of $u(x)$.

On the other hand, Cantor's algorithm ([4]) explains how to compute the *reduced* presentation of a semi-reduced divisor:

³We always assume in practice that q is the unique point over ∞ .

Lemma 2.18. *Keeping the same setup as in Lemma 2.17, suppose that $(u(x), v(x))$ represents the semi-reduced divisor D . Set $u_1(x) = \frac{f(x) - v^2(x)}{u(x)}$, renormalizing the highest-order coefficient of $u_1(x)$ to obtain a monic polynomial if necessary; and choose $v_1(x) \equiv -v(x) \pmod{u_1(x)}$ such that $\deg v_1 < \deg u_1$. Then $D' = \sum_j (p'_j - q) \sim D$, where $p'_j = (b_j, v_1(b_j))$ and the b_j range over all roots of $u_1(x)$. Moreover, D' is semi-reduced with respect to q .*

Notice that $\deg u_1(x) = \max\{2\gamma + 1, 2 \deg v(x)\} - \deg u(x) < \deg u(x)$ and that $v_1(x)$ may be chosen so that $\deg v_1(x) < \deg u_1(x)$. Moreover, $u_1 | (f - v_1^2)$. After iterating Cantor's algorithm finitely many times, we arrive at a pair $(u_n(x), v_n(x))$ that represents a q -reduced divisor. In particular, the algorithm terminates when $\deg u_i \leq \gamma$.

Now say that applying Cantor's algorithm T times to a Mumford presentation $(u_0(x), v_0(x))$ of a divisor that is semi-reduced with respect to q yields a pair $(u_T(x), v_T(x))$ that represents a q -reduced divisor. Then

$$u_i(x) \sim \frac{f(x) - v_{i-1}^2(x)}{u_{i-1}(x)}$$

for every $i = 1, \dots, T$.⁴ In particular, we have

$$u_T(x) \sim \frac{f(x) - v_{T-1}^2(x)}{u_{T-1}(x)} \sim \frac{f(x) - v_{T-1}^2(x)}{f(x) - v_{T-2}^2(x)} u_{T-2}(x).$$

By iteratively replacing $u_i(x)$ by $\frac{f(x) - v_{i-1}^2(x)}{u_{i-1}(x)}$, we see that

(4)

$$\begin{aligned} u_T(x) \prod_{k=1}^{\frac{T}{2}} (f(x) - v_{2k-2}^2(x)) &\sim u_0(x) \prod_{k=1}^{\frac{T}{2}} (f(x) - v_{2k-1}^2(x)) \text{ when } T \text{ is even; and} \\ u_T(x) u_0(x) \prod_{k=1}^{\lfloor \frac{T}{2} \rfloor} (f(x) - v_{2k-1}^2(x)) &\sim \prod_{k=0}^{\lfloor \frac{T}{2} \rfloor} (f(x) - v_{2k}^2(x)) \text{ when } T \text{ is odd.} \end{aligned}$$

We will use Cantor's algorithm to analyze the existence of torsion points of prescribed order on B , as follows. Suppose $N \cdot (p - q) \sim 0$, where $N \geq 2\gamma + 1$; and let $p = (a, b)$ be a point for which $f(a) \neq 0$. Starting with the pair $(u_0(x), v_0(x)) = ((x - a)^N, v(x))$ that represents $N \cdot (p - q)$ and applying the algorithm repeatedly, we eventually arrive at $(u_T(x), v_T(x)) = (1, 0)$. Indeed, if $u_n(x)$ is not constant, then $(u_T(x), v_T(x))$ represents a nonzero divisor of the form $\sum ((a_i, b_i) - q)$, where the a_i range over all roots of $u_n(x)$ and $b_i = v_n(a_i)$; in particular, the representative may be further reduced.

When (a, b) is torsion of order $N \geq 2\gamma + 1$, the equations (4) specialize to

$$(5) \quad (x - a)^N \prod_{k=1}^{\frac{T}{2}} (f(x) - v_{2k-1}^2(x)) \sim \prod_{k=0}^{\frac{T}{2}-1} (f(x) - v_{2k}^2(x))$$

⁴Here we use \sim to denote equality up to a scalar.

and

$$(6) \quad (x-a)^N \prod_{k=1}^{\lfloor \frac{T}{2} \rfloor} (f(x) - v_{2k-1}^2(x)) \sim \prod_{k=0}^{\lfloor \frac{T}{2} \rfloor} (f(x) - v_{2k}^2(x))$$

respectively. The degrees of the v_i satisfy recursively-defined inequalities; namely, we have $\deg v_i < \sum_{k=0}^i (-1)^{i-k} R_k$, where the integers R_k are prescribed by $R_k = \max\{2\gamma + 1, 2 \deg v_{k-1}\}$, subject to $R_0 = N$.

Lemma 2.19. *Suppose that $p - q$ is an N -torsion, $N \geq 2\gamma + 1$. Starting with the Mumford presentation $((x-a)^N, v(x))$ of $N \cdot (p-q)$, Cantor's algorithm terminates in at most T steps, where $T = \lceil \frac{N-2\gamma}{2} \rceil$.*

Proof. It suffices to see that $\deg u_i \equiv \deg u_{i+1} \pmod{2}$ whenever $2 \deg v_i(x) > 2\gamma + 1$; otherwise the algorithm terminates after the subsequent step. \square

Corollary 2.20. *Suppose $\gamma \geq 2$, and that $f(x)$ is a monic separable polynomial of degree $2\gamma + 1$. The hyperelliptic curve $y^2 = f(x)$ admits an N -torsion point with*

- (1) $N = 2\gamma + 1$ if and only if $f(x) = (x-a)^{2\gamma+1} + v_0^2(x)$ with $\deg v_0(x) \leq \gamma$;
- (2) $N = 2\gamma + 2$ if and only if $f(x) = (x-a)^{2\gamma+2} + v_0^2(x)$ with $\deg v_0(x) = \gamma + 1$;
- (3) $N = 2\gamma + 3$ if and only if $f(x) = \frac{v_0^2(x) - (x-a)^{2\gamma+3} \cdot v_1^2(x)}{1 - (x-a)^{2\gamma+3}}$ with $\deg v_0(x) = 2\gamma + 2$ and $\deg v_1(x) \leq \gamma$.

Proof. In the first two cases, the algorithm terminates in one step; while in the last case, it must terminate in two steps. Applying Equations 5 and 6, respectively, we derive the shape of $f(x)$ in each of these three cases.

More precisely, Cantor's algorithm implies that $N \cdot (p - q) \sim 0$ in each of these cases. On the other hand, the fact that B has no N' -torsion points for any $3 \leq N' \leq 2\gamma$ ([19, Corollary 2.8]) implies $p - q$ cannot have order smaller than N . \square

Remark 2.21. The case of $(2\gamma + 1)$ -torsion points is covered by [1, Theorem 1].

Whenever $N \geq 2\gamma + 4$, it is no longer possible to give a uniform characterization of odd-degree hyperelliptic curves with N -torsion points, as *a priori* there are various distinct ways in which the reduction algorithm might terminate. We nevertheless have the following existence result:

Example 2.22. Let $1 \leq s \leq \gamma$, and set $f(x) = \frac{-1}{2}[(x^{\gamma+s} - x^{\gamma-s+1} + 1)^2 - x^{2(\gamma+s)}]$, and let $p = (0, \sqrt{\frac{-1}{2}})$. Then, $f(x)$ is a monic, separable polynomial of degree $2\gamma + 1$, and $(p - q)$ is a $(2\gamma + 2s)$ -torsion point of the hyperelliptic curve $y^2 = f(x)$.

Indeed, the fact that f is monic is clear. On the other hand, to show that $f(x) = \frac{1}{2}(2x^{\gamma+s} - x^{\gamma-s+1} + 1)(x^{\gamma-s+1} - 1)$ is separable it suffices to show that $g(x) = 2x^{\gamma+s} - x^{\gamma-s+1} + 1$ is separable, as f clearly does not vanish at $(\gamma - s + 1)$ -th roots of unity; and the separability of $g(x)$ may be checked directly.⁵

Applying the reduction algorithm to $(x^{2\gamma+2s}, \sqrt{\frac{-1}{2}}(x^{\gamma+s} - x^{\gamma-s+1} + 1))$, we see that $(2\gamma + 2s) \cdot (p - q) \sim 0$. Since $\gamma + s \leq 2\gamma$, by [19, Corollary 2.8] we conclude $(p - q)$ has order $2\gamma + 2s$.

⁵Indeed, $g(x) = g'(x) = 0$ would imply that $x^{\gamma-s+1} = \frac{\gamma+s}{2s-1}$ and $x^{2s-1} = \frac{\gamma-s+1}{2\gamma+2s}$, which is impossible.

3. WEIERSTRASS-REALIZABILITY FOR $2N$ -SEMIGROUPS

Definition 3.1. A numerical semigroup S is *Weierstrass-realizable* whenever there exists a pointed smooth projective curve (C, p) for which $S = S(C, p)$ is the Weierstrass semigroup of C at p .

In investigating the Weierstrass-realizability of a given semigroup S , it is often to useful to work modulo the multiplicity M of S . Hereafter, we refer to numerical semigroups of multiplicity M as *M -semigroups*.

Definition 3.2. Given an M -semigroup S , we set

$$e_i = e_i(S) := \min\{s \in S \mid s \equiv i \pmod{M}\}.$$

The *standard basis* of S is the M -dimensional vector (M, e_1, \dots, e_{M-1}) .

Our strategy. We will certify the Weierstrass-realizability of prescribed $2N$ -semigroups S by constructing cyclic covers of degree- N of hyperelliptic curves and computing the Weierstrass semigroups at totally ramified points lying over Weierstrass points of the targets. We focus on cyclic covers of the following form.

Situation 3.3. Let $\pi : C = \text{spec}(\bigoplus_{i=0}^{N-1} (L^{(i)})^{-1}) \rightarrow B$ be a cyclic Galois cover of degree N between smooth projective curves, where B is a hyperelliptic curve of genus γ and $g(C) = g$. Here L is a line bundle on B for which $L^{\otimes N} \cong \mathcal{O}_B(D)$ (for some fixed choice of isomorphism) where D is an effective divisor of the form $aq + \sum a_i q_i$, and $L^{(i)} = L^i(-[\frac{iD}{N}])$. Fix $p \in C$ such that $\pi(p) = q$, where q is a Weierstrass point of B and π is totally ramified at p .

According to Lemma 2.13, a given integer m belongs to $S(C, p)$ if and only if

$$(7) \quad \sum_{i=0}^{N-1} h^0\left(\left(L^{(i)}\right)^{-1} \left(\left[\frac{m}{N} + \left\langle \frac{ia}{N} \right\rangle\right] q\right)\right) \neq \sum_{i=0}^{N-1} h^0\left(\left(L^{(i)}\right)^{-1} \left(\left[\frac{m-1}{N} + \left\langle \frac{ia}{N} \right\rangle\right] q\right)\right).$$

The inequality holds if and only if there exists at least one i such that $s(m, i) := \frac{m}{N} + \langle \frac{ia}{N} \rangle$ is an integer and $h^0((L^{(i)})^{-1}(s(m, i)q)) \neq h^0((L^{(i)})^{-1}((s(m, i) - 1)q))$.

By definition, $s(m, i) \in \mathbb{Z}$ if and only if $m + ia \equiv 0 \pmod{N}$. Since π is totally ramified at p , by Theorem 2.10 $\gcd(a, N) = 1$, and hence there is *precisely* one i such that $s(m, i)$ is an integer. In this case, we shall write s_m for $s(m, i)$.

Definition 3.4. Given any line bundle L on C , we set

$$S_L(q) := \{n \in \mathbb{Z} \mid h^0(L(nq)) \neq h^0(L((n-1)q))\}$$

and refer to it the *Weierstrass set* at q of L .

Given a $2N$ -semigroup S with standard basis $(2N, e_1, \dots, e_{2N-1})$, our goal is to decide whether $S = S(C, p)$, for some pair (C, p) . To do so, we will adapt the inequality (7) to the case of $m = e_i$. It is worth noting that specifying a standard basis $(2N, e_1, \dots, e_{2N-1})$ is equivalent to fixing a *gap vector* $(0, v_1, \dots, v_{2N-1})$, in which v_i denotes the number of elements in $\mathbb{N} \setminus S$ with $2N$ -residue equal to i , for every $i = 1, \dots, 2N - 1$. Indeed, since e_i is a minimal representative of the i -th residue class, we have $v_i = \lfloor \frac{e_i}{2N} \rfloor$. But the value of every sum $v_i + v_{N+i}$ is fixed by Theorem 3.12; so the standard basis is determined by $\min\{e_1, e_{N+1}\}, \dots, \min\{e_{N-1}, e_{2N-1}\}$, and consequently hereafter we only specify these elements of the standard basis when determining a semigroup.

The upshot of the preceding discussion is the following criterion for the Weierstrass-realizability of $2N$ -semigroups:

Lemma 3.5. *Let S be a $2N$ -semigroup with standard basis $(2N, e_1, \dots, e_{2N-1})$. Then S is Weierstrass-realizable by a pointed cyclic cover $\pi : (C, p) \rightarrow (B, q)$ as in Situation 3.3 if and only if there exists an invertible sheaf L on B and an effective divisor $D = aq + \sum a_k q_k$ for which $L^{\otimes N} \cong \mathcal{O}_B(D)$, $\gcd(a, N) = 1$ and*

$$h^0((L^{(i)})^{-1}(s_{\bar{e}_j}q)) = 1 \text{ and } h^0((L^{(i)})^{-1}((s_{\bar{e}_j} - 1)q)) = 0$$

for $j = 1, \dots, N-1$, where $\bar{e}_j = \min\{e_j, e_{j+N}\}$, $\bar{e}_j + ia \equiv 0 \pmod{N}$ and $L^{(i)} = L^i(-\lfloor \frac{iD}{N} \rfloor)$; moreover, $\max\{e_j, e_{j+N}\} = \min\{n \in S_{(L^{(i)})^{-1}}(q) \mid n - s_{\bar{e}_j} \equiv 1 \pmod{2}\}$.

Proof of Lemma 3.5. We have $S = S(C, p)$ if and only if $e_j = \min\{n \in S(C, p) \mid n \equiv j \pmod{2N}\}$. This in turn is equivalent to $s_{\bar{e}_j}$ being the minimal element of $S_{(L^{(i)})^{-1}}(q)$, which is cohomologically characterized by the given conditions. Notice also that $s_{e_j} = \frac{e_j}{N} + \langle \frac{ia}{N} \rangle$ and $s_{e_{j+N}} = \frac{e_{j+N}}{N} + \langle \frac{ia}{N} \rangle$ are integers of different parity, which yields the last item. \square

It is easy to see that if $L^{\otimes N} \cong \mathcal{O}_B(D)$ and D is effective on B , then $\deg(L^{-1}) < 0$ and hence that $S_{L^{-n}}(q) \subset \mathbb{Z}_{>0}$ for every $n \geq 1$.⁶

3.1. Weierstrass-realizability via cyclic covers with reduced branch locus.

A particularly simple class of cyclic covers is comprised of those for which the covering data (L, D) includes a *reduced* effective divisor D . Hereafter, we focus on those $2N$ -semigroups that are realizable by pointed cyclic covers $\pi : (C, p) \rightarrow (B, q)$ constructed out of such covering data. More precisely, we study the Weierstrass-realizability of $2N$ -semigroups via degree- N cyclic covers $\pi : C \rightarrow B$ of hyperelliptic curves satisfying the following hypotheses:

Situation 3.6. (1) *The cover π satisfies the conditions of Situation 3.3; and*
 (2) *the underlying (branch) divisor D is reduced (and hence $L^{(i)} = L^i$).*

In this case, the cover π is totally ramified at every point in the ramification locus. Let g (resp., γ) denote the genus of the source (resp., target curve); we have $g = 1 + N(\gamma - 1) + \frac{1}{2} \deg(D)(N - 1)$ by the Riemann–Hurwitz formula. *Hereafter, we always assume $g \geq (2N - 1)\gamma$.*

Remark 3.7. A $2N$ -semigroup S realized by a pointed cyclic cover $\pi : C \rightarrow B$ as in Situation 3.6 with target B of genus γ is (N, γ) -hyperelliptic in the sense of [5]; that is, $\mathbb{N} \setminus S$ contains precisely γ multiples of N , all of which are strictly less than $2N\gamma$. In particular, semigroups S_1 and S_2 associated with distinct pairs (N_i, γ_i) , $i = 1, 2$ are always distinct.

Notation 3.8. Given $n \geq 1$, let $a_n(L, q)$ denote the minimal element of $S_{L^{-n}}(q)$ strictly greater than zero.

⁶In fact, it is also true that $S_{(L^{(i)})^{-1}}(q) \subset \mathbb{Z}_{>0}$, as

$$\deg((L^{(i)})^{-1}) = -i \cdot \deg(L) + \deg\left(\mathcal{O}_B\left(\left\lfloor \frac{iD}{N} \right\rfloor\right)\right)$$

$\deg(D) = N \cdot \deg(L)$ and we have assumed from the outset that D is *not* of the form $D = N \cdot D'$, where D' is effective.

Note that $s_{\overline{e_j}} = \frac{\overline{e_j} - j}{N} + 1$ for every $j = 1, \dots, N - 1$. Accordingly, the cohomological condition in Lemma 3.5 simplifies to the requirement that

$$(8) \quad h^0(L^{-(N-j)}(s_{\overline{e_j}}q)) = 1, h^0(L^{-(N-j)}((s_{\overline{e_j}} - 1)q)) = 0, 1 \leq j \leq N - 1.$$

Conversely, assume we are given an invertible sheaf L and a reduced effective divisor D for which $L^{\otimes N} \cong \mathcal{O}_B(D)$, it is easy to see that $Na_j(L, q) - (N - j) \in S(C, p)$ for every $j = 1, \dots, N - 1$. Suppose now that S is a $2N$ -semigroup with standard basis $(2N, e_1, \dots, e_{2N-1})$ and that $S = S(C, p)$. Because of the minimality of the integers $a_j(L, q)$ in their respective Weierstrass sets, we conclude that

$$Na_{N-j}(L, q) - (N - j) = \overline{e_j} = \min\{e_j, e_{j+N}\}$$

or equivalently, that $a_{N-j}(L, q) = s_{\overline{e_j}}$. In particular, $(a_1(L, q), \dots, a_{N-1}(L, q))$ determines $S = S(C, p)$.

Accordingly, in the current setting Lemma 3.5 specializes to the following statement:

Lemma 3.9. *The $2N$ -semigroups that are Weierstrass-realizable by pointed cyclic covers of (B, q) as described in Situation 3.6 correspond bijectively to tuples of positive integers (a_1, \dots, a_{N-1}) for which there exist a line bundle L and a reduced effective divisor D on B such that*

- (1) $L^{\otimes N} \cong \mathcal{O}_B(D)$; and
- (2) $h^0(L^{-j}(a_jq)) = 1, h^0(L^{-j}((a_j - 1)q)) = 0$, for $j = 1, \dots, N - 1$.

Notation 3.10. Set $d_{g,\gamma,N} = \frac{(2g-2)-N(2\gamma-2)}{N(N-1)}$.

Riemann-Hurwitz implies that the cover $\pi : (C, p) \rightarrow (B, q)$ is totally ramified over precisely $Nd_{g,\gamma,N}$ distinct points. In particular, we have $\deg(L) = d_{g,\gamma,N}$. Our additional numerical assumption $g \geq (2N - 1)\gamma$ ensures that $L^{\otimes N}$ is very ample, and thus by Bertini's theorem the associated complete linear series contains a reduced divisor D . The upshot is that in constructing cyclic covers that realize Weierstrass semigroups, we may focus on verifying the cohomological conditions of Lemma 3.9. This point is illustrated in the next example.

Example 3.11. (A first example.) Suppose $L = \mathcal{O}_B(d_{g,\gamma,N}q)$. Note that $(L^{\otimes N})(-q)$ is itself very ample; accordingly, we fix a general reduced divisor D' in $|(L^{\otimes N})(-q)|$ not supported at q and use $D = D' + q \in |L^{\otimes N}|$ to build a cyclic cover as in Situation 3.6. In so doing we realize the $2N$ -semigroup with standard basis

$$e_0 = 2N \text{ and } \overline{e_j} = \min\{e_j, e_{N+j}\} = (Nd_{g,\gamma,N} - 1)(N - j) \text{ for every } j = 1, \dots, N - 1.$$

This semigroup was studied in [5]; hereafter we will refer to it as being of *Carvalho-Torres type*.

Example 3.11 illustrates our general strategy towards realizing a given numerical semigroup S : we seek a divisor Y on (B, q) and a reduced effective divisor D on B such that $D \in |NY|$ and $(\mathcal{O}_B(Y), D)$ provides a cyclic cover $\pi : (C, p) \rightarrow (B, q)$ for which $S = S(C, p)$. We may reformulate our realizability problem as follows:

Question 1. *Let S be a $2N$ -semigroup with standard basis $(2N, e_1, \dots, e_{2N-1})$, and set $a_j = \frac{\min\{e_{N-j}, e_{2N-j}\} + j}{N}$. For which tuples of positive integers (a_1, \dots, a_{N-1}) does there exist a divisor Y of degree $d_{g,\gamma,N}$, not necessarily effective, such that for every $1 \leq j \leq N - 1$, we have*

$$\begin{cases} h^0(\mathcal{O}_B(-(N-j)Y + a_jq)) = 1 \text{ and} \\ h^0(\mathcal{O}_B(-(N-j)Y + (a_j - 1)q)) = 0? \end{cases}$$

Hereafter, we shall refer to the $2N$ -semigroup H whose standard basis is determined by (a_1, \dots, a_{N-1}) as above as *the $2N$ -semigroup indexed by (a_1, \dots, a_{N-1})* .

3.2. Restrictions on Weierstrass semigroups arising from cyclic covers.

Let $\mathcal{L}(a, b) := L^{\otimes -a}(b \cdot q)$. We begin by explicitly relating the cohomology of $\mathcal{O}(m \cdot p)$ on C to that of $\mathcal{L}(a, \lfloor \frac{m+a}{N} \rfloor)$ on B . To this end, note first that

$$h^0(C, m \cdot p) - h^0(C, (m-1) \cdot p) = h^0\left(B, \mathcal{L}\left(a, \left\lfloor \frac{m+a+1}{N} \right\rfloor\right)\right) - h^0\left(B, \mathcal{L}\left(a, \left\lfloor \frac{m+a}{N} \right\rfloor\right)\right)$$

precisely when N divides $(m+a+1)$. It follows that

$$h^0(C, (Nb-a)p) - h^0(C, (Nb-a-1)p) = h^0(B, \mathcal{L}(a, b)) - h^0(B, \mathcal{L}(a, b-1))$$

and, consequently, that

$$h^0(B, \mathcal{L}(a, b)) - h^0(B, \mathcal{L}(a, b-1)) = \delta(Nb-a \in S(C, p))$$

where $\delta(\cdot)$ denotes Kronecker's delta. As $h^0(B, \mathcal{L}(a, -1)) = 0$, we deduce that

$$(9) \quad \begin{aligned} h^0(B, \mathcal{L}(a, b)) &= \sum_{i=0}^n (h^0(B, \mathcal{L}(a, i)) - h^0(B, \mathcal{L}(a, i-1))) \\ &= \#\{m \in S(C, p) : m \leq Nb-a \text{ and } m \equiv -a \pmod{N}\}. \end{aligned}$$

The following result shows that (9) uniquely determines the *gap vector* computing the number of elements of $\mathbb{N} \setminus S(C, p)$ in each N -residue class between 0 and $N-1$.

Theorem 3.12. *We have*

$$\#\{\text{gaps} \equiv -a \pmod{N}\} = \gamma - 1 + \delta(a=0) + ad_{g, \gamma, N}$$

for every $a = 0, 1, \dots, N-1$.

Proof. Riemann–Roch implies that

$$(10) \quad h^0(B, \mathcal{L}(a, b)) = b + 1 - \gamma - ad_{g, \gamma, N}.$$

whenever $b \gg 0$. On the other hand, (9) implies that

$$\begin{aligned} h^0(B, \mathcal{L}(a, b)) &= \#\{m \leq Nb-a : m \geq 0 \text{ and } m \equiv -a \pmod{N}\} - \#\{\text{gaps} \equiv -a \pmod{N}\} \\ &= \left\lfloor \frac{Nb-a}{N} \right\rfloor + 1 - (\#\text{gaps} \equiv -a \pmod{N}) \\ &= b + \left\lfloor \frac{-a}{N} \right\rfloor + 1 - (\#\text{gaps} \equiv -a \pmod{N}). \end{aligned}$$

Comparing the latter equality with (10) yields the desired result. \square

Additional restrictions on (a_1, \dots, a_{N-1}) arise from the additive structure of S ; indeed, according to [13, (1.2)], for every $i, j \in \{0, 1, \dots, 2N-1\}$ we have

$$(11) \quad v_i + v_j \geq v_{i+j} \text{ if } i+j < 2N, \text{ while } 1 + v_i + v_j \geq v_{i+j-2N} \text{ if } i+j > 2N$$

where v_i denotes the cardinality of the set of gaps $\{\ell \in \mathbb{N} \setminus S : \ell \equiv i \pmod{2N}\}$ with $2N$ -residues equal to i . Here $v_i = \lfloor \frac{e_i}{2N} \rfloor$. We will refer to the inequalities (11) as the *Morrison–Pinkham inequalities*; they characterize those vectors (e_0, \dots, e_{2N-1}) that arise as standard bases of numerical semigroups (and in particular, are independent of conditions arising from the construction of cyclic covers). A useful reformulation of (11) is the requirement that (e_0, \dots, e_{2N-1}) satisfy

$$(12) \quad e_j + e_k \geq e_{(j+k) \bmod 2N}$$

for every $j, k \in \{0, 1, \dots, 2N - 1\}$. On the other hand, Theorem 3.12 implies that standard bases (e_0, \dots, e_{2N-1}) of $2N$ -semigroups that arise from our cyclic covers built out of reduced branch divisors as in Situation 3.6 satisfy

$$(13) \quad e_{N-i} + e_{2N-i} = iNd_{g,\gamma,N} + (2\gamma + 1)N - 2i$$

for every $i = 1, \dots, N - 1$. Applying the Morrison–Pinkham inequalities (12) with $j = N, k = N - i$ in tandem with the cyclic cover restriction (13), we deduce that

$$iNd_{g,\gamma,N} - i \leq e_{N-i}, e_{2N-i} \leq iNd_{g,\gamma,N} + (2\gamma + 1)N - i$$

and consequently that

$$iNd_{g,\gamma,N} \leq \min(e_{N-i}, e_{2N-i}) + i \leq iNd_{g,\gamma,N} + \gamma$$

for every $i = 1, \dots, N - 1$. In other words, $a_i \in [id_{g,\gamma,N}, id_{g,\gamma,N} + \gamma]$ for every i .

3.3. Weierstrass-realizability for feasible $2N$ -semigroups.

Definition 3.13. For $N \geq 2$, we set $\tilde{F}(N) = \left(\prod_{j=1}^{N-1} [jd_{g,\gamma,N}, jd_{g,\gamma,N} + \gamma] \right) \cap \mathbb{Z}^{N-1}$. Similarly we set

$$F(N) = \{(a_1, \dots, a_{N-1}) \in \tilde{F}(N) \mid a_{i+j} \leq a_i + a_j \text{ for every } 1 \leq i, j, i+j \leq N-1\}.$$

We refer to $F(N)$ as the N -th *feasible set*.

Example 3.14. Say $N = 2$. The feasible set $F(2)$ is an interval of length γ , and each of the $\gamma + 1$ possible values $a_1 = d_{g,\gamma,2} + j$, $j = 0, \dots, \gamma$ indexes a 4-semigroup that is realizable as the Weierstrass semigroup of a (totally ramified point of a) double cover of a hyperelliptic curve. Indeed, according to Corollary 2.16, the existence of a line bundle L that satisfies the cohomological conditions of Lemma 3.9 is a straightforward consequence of the existence of q -reduced divisors in every divisor class of any given pointed hyperelliptic curve (B, q) , which in turn is the content of Lemma 2.15.

Example 3.15. Say $N = 3$. This time, we have

$$(14) \quad F(3) := \{(a_1, a_2) \in ([d_{g,\gamma,3}, d_{g,\gamma,3} + \gamma] \times [2d_{g,\gamma,3}, 2d_{g,\gamma,3} + \gamma]) \cap \mathbb{Z}^2 \mid a_2 \leq 2a_1\}$$

and for each $j = 1, 2$, the cohomological conditions

$$h^0(\mathcal{O}_B(-jY + a_jq)) = 1, h^0(\mathcal{O}_B(-jY + (a_j - 1)q)) = 0$$

of Lemma 3.9 determine an (at-most) unique effective divisor $E_j \sim a_jq - jY$ on B , which may or may not exist. Note that $\deg(E_j) = a_j - j \cdot \deg(Y) = a_j - jd_{g,\gamma,3}$.

With the two preceding examples in mind, we reformulate our realizability criterion once again:

Lemma 3.16. *The $2N$ -semigroup indexed by (a_1, \dots, a_{N-1}) is Weierstrass-realizable via a cyclic cover as in Situation 3.6 if and only if there exists an $(N - 1)$ -tuple of effective divisors $(E_1, \dots, E_{N-1}) \in \prod_{j=1}^{N-1} \text{Sym}^{a_j - jd_{g,\gamma,N}} B$ for which*

- (1) $E_j - (a_j - jd_{g,\gamma,N})q$ is q -reduced in the sense of Definition 2.14; and
- (2) $j(E_1 - a_1q) \sim E_j - a_jq$

for every $j = 1, \dots, N - 1$.

Proof. Given E_1, \dots, E_{N-1} satisfying (1) and (2), let $L = \mathcal{O}_B(-E_1 + a_1q)$; it then follows from (3) that $L^j \cong \mathcal{O}_B(a_jq - E_j)$. It also follows from (1) and Corollary 2.16 that

$$h^0(L^{-j}(a_jq)) = h^0(\mathcal{O}_B(E_j)) = 1 \text{ and } h^0(L^{-j}((a_j - 1)q)) = h^0(\mathcal{O}_B(E_j - q)) = 0.$$

Recall that we assume g is large with respect to γ , so that a reduced representative D in $|L^{\otimes N}|$ exists. Applying Lemma 3.9, we conclude that the $2N$ -semigroup indexed by (a_1, \dots, a_{N-1}) is Weierstrass-realizable.

Conversely, suppose S is Weierstrass-realizable via a cyclic cover as in Situation 3.6. According to Lemma 3.9, we have $h^0(L^{-j}(a_jq)) = 1$ and $h^0(L^{-j}((a_j - 1)q)) = 0$, for $j = 1, \dots, N - 1$. This means that there exist effective divisors E_1, \dots, E_{N-1} for which $\mathcal{O}_B(E_j) \cong L^{-j}(a_jq)$ for every j . In particular, $\deg(E_j) = a_j - jd_{g,\gamma,N} \leq \gamma$. Consequently $E_j - (a_j - jd_{g,\gamma,N})q$ fails to be reduced if and only if one of the following two situations occurs:

- (1) E_j is supported at q ; or else
- (2) E_j is supported at p and $\iota(p)$, where ι is the hyperelliptic involution on B .

The fact that $h^0(L^{-j}((a_j - 1)q)) = h^0(\mathcal{O}_B(E_j)(-q)) = 0$ rules out the first case; while the fact that $h^0(L^{-j}(a_jq)) = h^0(\mathcal{O}_B(E_j)) = 1$ rules out the second. Finally, since $\mathcal{O}_B(E_j) \cong L^{-j}(a_jq)$ for every j , item (2) in the statement of the lemma clearly holds. This concludes the proof. \square

Lemma 3.16 shows that the problem of constructing pointed cyclic covers of a hyperelliptic curve with prescribed Weierstrass semigroup is fundamentally about the behavior of multiplication maps on a hyperelliptic Jacobian.

Notation 3.17. Let $u_d : B^d \rightarrow \text{Jac}(B) : \sum p_i \mapsto [\sum p_i - dq]$ be the d -th Abel-Jacobi map. Let Θ_d denote the image of u_d , and set $\Theta_d^0 = \Theta_d \setminus \Theta_{d-1}$.

Notation 3.18. Let $m_j : \text{Jac}(B) \rightarrow \text{Jac}(B)$ denote the multiplication-by- j map.

Lemma 3.16 establishes that the $2N$ -semigroup indexed by (a_1, \dots, a_{N-1}) is realizable via a cover as in Situation 3.6 if and only if $\bigcap_{j=1}^{N-1} m_j^{-1}(\Theta_{a_j - jd_{g,\gamma,N}}^0)$ is non-empty.

Remark 3.19. It is worth noting that more refined information is known about set-theoretic intersections of the form $(m_2^{-1}\Theta_d) \cap \Theta_{d'}$. For example, when $\gamma > 1$, $m_2^{-1}\Theta_1 \cap \Theta_1$ consists of the u_1 -images of all Weierstrass points in $\text{Jac}(B)$, and $m_2^{-1}\Theta_1 \cap \Theta_{\gamma-1} \subset \text{Jac}(B)[2]$ (see [19, Thm 2.5]).

For the sake of simplicity, hereafter we set $\epsilon_j = a_j - jd_{g,\gamma,N}$ and use the vector $(\epsilon_1, \dots, \epsilon_{N-1})$ to indicate the semigroup indexed by (a_1, \dots, a_{N-1}) .

Lemma 3.20. *Suppose the $2N$ -semigroup indexed by $(\epsilon_1, \dots, \epsilon_{N-1})$ is realizable via a cyclic cover as in Situation 3.6. Then the following hold:*

- (1) *The set $\{j \mid \epsilon_j = 0\}$ is either empty or consists of all multiples of a given integer n in $\{1, \dots, N - 1\}$ and $\epsilon_{j+n} = \epsilon_j$ for every j .*
- (2) *When $\gamma > 1$ and $\epsilon_1 = 1$, we have $\epsilon_2, \epsilon_3 \neq 1$ unless $(\epsilon_1, \epsilon_2, \epsilon_3) = (1, 0, 1)$.*
- (3) *When $\gamma > 1$ and $\epsilon_1 = 1$, we have $\epsilon_j \neq 0$ for every $j \leq 2\gamma$ unless j is even and $\epsilon_2 = 0$.*
- (4) *The semigroup indexed by (γ, \dots, γ) is realizable.*
- (5) *If $\epsilon_i = 0$, then $\epsilon_j = \epsilon_{i-j}$ for every index j strictly less than i .*

- Proof.* (1) Without loss of generality, suppose that the index set $\{j \mid \epsilon_j = 0\}$ is nonempty; let n be the smallest such index. Then $E_1 - \epsilon_1 q$ defines an order- n element of $\text{Jac}(B)$, and the conclusion follows from the requirement that $j(E_1 - \epsilon_1 q) \sim E_j - \epsilon_j q$.
- (2) The fact that $m_2^{-1}\Theta_1 \cap \Theta_1$ consists exclusively of Weierstrass points implies that $\epsilon_2 \neq 1$; on the other hand, we have $H^0(\mathcal{O}(2q + p)) = H^0(\mathcal{O}(2q))$ for every $p \in B$.
- (3) Follows from the fact that B has no points of order $j \in \{3, \dots, 2\gamma\}$ ([19, Cor. 2.8]).
- (4) Immediate, as the preimages $m_j^{-1}(\Theta_j^0)$ are non-empty open subsets of $\text{Jac}(B)$.
- (5) Let $x = E_1 - \epsilon_1 q$. If $\epsilon_i = 0$, then $i \cdot x = 0$, which in turn implies $j \cdot x = -(i - j) \cdot x$. The item follows from the fact that every Θ_j is invariant under multiplication by (-1) : the inverse of $\sum_{k=1}^j p_k - jq$ is $\sum_{k=1}^j \iota(p_k) - jq$. \square

3.4. Realizability for small target genera.

$\gamma = \mathbf{1}$. In this case, the coordinates ϵ_i of every candidate vector $(\epsilon_1, \dots, \epsilon_{N-1})$ satisfy $\epsilon_i + \epsilon_j \geq \epsilon_{i+j}$ for every pair of indices $i, j \leq \frac{N-1}{2}$. Furthermore, we have $m_j^{-1}(\Theta_1^0) = B \setminus B[j]$ and $m_j^{-1}(\Theta_0) = B[j]$. Candidate ϵ -vectors that are realizable by cyclic covers as in Situation 3.6 are as follows:

- (1) $(\epsilon_1, \dots, \epsilon_{N-1}) = \mathbf{0}$. The trivial ϵ -vector indexes a Carvalho–Torres semigroup, which is realizable; see Example 3.11.
- (2) For some fixed i , we have $\epsilon_{ki} = 0$ for every $k = 1, \dots, \lfloor \frac{N-1}{i} \rfloor$; and $\epsilon_j = 1$ for all other indices j . To realize the corresponding semigroup by a cyclic cover as in Situation 3.6, choose $E_1 = p - q$, where $p \in B$ is any point (with Abel–Jacobi image) of order i .
- (3) $(\epsilon_1, \dots, \epsilon_{N-1}) = \mathbf{1}$. Realization follows trivially for topological reasons in this subcase; see Lemma 3.20(4).

$\gamma = \mathbf{2}, \mathbf{N} = \mathbf{4}$. Realizable ϵ -vectors in lexicographic order are as follows:

- (1) $(\epsilon_1, \dots, \epsilon_3) = \mathbf{0}$. See Example 3.11.
- (2) $(\epsilon_1, \dots, \epsilon_3) = (1, 0, 1)$. In this subcase, choose $E_1 = p - q$, where p defines a point of order 2.
- (3) $(\epsilon_1, \dots, \epsilon_3) = (1, 2, 2)$. Choose $E_1 = p - q$, where $p - q$ is non-torsion, and apply Lemma 3.20(2).
- (4) $(\epsilon_1, \dots, \epsilon_3) = (2, 0, 2)$. Choose $E_1 \in \text{Jac}(B)[2] \setminus \Theta_1$.
- (5) $(\epsilon_1, \dots, \epsilon_3) = (2, 1, 1)$. Start with any B that admits a 5-torsion point p ($[2]$). Then $E_1 = 2(p - q)$ is q -reduced and $[2E_1] = [3E_1] = [p - q]$.
- (6) $(\epsilon_1, \dots, \epsilon_3) = (2, 1, 2)$. According to Lemma 3.18 (2), we have $m_2(\Theta_1) \cap \Theta_1 = \Theta_0$, and therefore $m_2^{-1}(\Theta_1^0) \subset \Theta_2^0$. Now choose $E_1 \in m_2^{-1}(\Theta_1^0)$ such that $[2E_1] = [p - q]$, where p is any Weierstrass point distinct from q . Then $[3E_1] = [p - q] + E_1 \neq [s - q]$, as otherwise we would have $[2E_1] = [2s - 2q]$, which is absurd.
- (7) $(\epsilon_1, \dots, \epsilon_3) = (2, 2, 0)$. Choose $E_1 \in \text{Jac}(B)[3]$ and apply Lemma 3.20(5).
- (8) $(\epsilon_1, \dots, \epsilon_3) = (2, 2, 1)$. To prove realizability in this subcase, we prove the following two claims:
 - $m_2^{-1}(\Theta_1)$ does not contain $m_3^{-1}(\Theta_1)$; and
 - $m_3^{-1}(\Theta_1)$ intersects Θ_1 in at most finitely many points.

For the first, suppose $x \in \text{Jac}(B)[3]$ is nonzero; then $x, 2x \in \Theta_2^0$ by items (3) and (5) of Lemma 3.20. It follows that $x \notin m_2^{-1}(\Theta_1)$.

For the second, let $p \in B$ be a general point; then $2p - 2q$ is reduced. If $3(p - q) \sim R - q$, we have $2(p - q) \sim \iota(p) + R - 2q$, which would violate the uniqueness of reduced presentations. Since Θ_1 is irreducible, $m_3(\Theta_1)$ intersects Θ_1 in at most finitely points and the claim follows.

The upshot of these two claims is that there is an irreducible component Z of $m_3^{-1}(\Theta_1)$ that intersects $m_2^{-1}(\Theta_1)$ in at most finitely many points.

Choosing a general element $E_1 \in Z$, we have $[2E_1] \in \Theta_2^0$ and $[3E_1] \in \Theta_1^0$.

(9) $(\epsilon_1, \dots, \epsilon_3) = \mathbf{2}$. See Lemma 3.20(4).

Realizability when $\gamma = 2$ and $N \geq 5$ is already significantly more delicate; we return to it in Section 4.

3.5. Realizability in small covering degree.

Example 3.21. Whenever $1 \leq k \leq \frac{\gamma}{N-1}$, $\sum_{i=1}^k p_i - kq$ is reduced and no p_i is a Weierstrass point on B , the first $(N-1)$ multiples $j(\sum_{i=1}^k p_i - kq)$, $1 \leq j \leq N-1$ are also reduced.

Accordingly, suppose that $1 \leq k \leq \frac{\gamma}{N-1}$, $\sum_{i=1}^k p_i - kq$ is reduced and no p_i is a Weierstrass point, and set $L = \mathcal{O}_B((d_{g,\gamma,N} + k)q - \sum_{i=1}^k p_i)$. Corollary 2.16 now implies that

$$h^0(L^{-j}((jd_{g,\gamma,N} + jk)q)) = 1, h^0(L^{-j}((jd_{g,\gamma,N} + jk - 1)q)) = 0$$

for every $j = 1, \dots, N-1$. It follows that the $2N$ -semigroup indexed by $(d_{g,\gamma,N} + k, 2(d_{g,\gamma,N} + k), \dots, (N-1)(d_{g,\gamma,N} + k))$ is realizable, for every $k = 1, \dots, \frac{\gamma}{N-1}$.

N=2. As explained in Example 3.14, the realizability problem is trivial in this case and every 4-semigroup is realizable.

N=3. In this case, we obtain the following result:

Theorem 3.22. *The 6-semigroup with associated a -vector (a_1, a_2) is Weierstrass-realizable by a construction described as in Situation 3.6 if and only if either*

- (1) $(a_1, a_2) \in F(3)$ and a_2 is even; or
- (2) $(a_1, a_2) \in F(3)$, $4d_{g,\gamma,3} + 2\gamma - 2a_1 + 1 \leq a_2 \leq 2a_1$, and a_2 is odd.

We split the argument into two cases, depending upon the parity of a_2 .

Case one: a_2 is even.

Proposition 3.23. *The 6-semigroups indexed by $(a_1 = d_{g,\gamma,3} + k, a_2 = 2d_{g,\gamma,3} + 2m) \in F(3)$ are all Weierstrass-realizable.*

Proof. Note that the feasible range of m is $0 \leq m \leq \frac{\gamma}{2}$ and the corresponding feasible range of k , given m , is $m \leq k \leq \gamma$.

Now set $Z = \sum_{i=1}^m p_i$ be an effective divisor as in Example 3.21 that underlies a cyclic cover realizing the 6-semigroup indexed by $(d_{g,\gamma,3} + m, 2d_{g,\gamma,3} + 2m)$; and suppose that $m < k \leq \gamma$. Let $E_1 = Z + \sum_{i=1}^{k-m} q_i$, where the q_i are distinct Weierstrass points other than q . Note that $E_1 - kq$ is a reduced divisor. Set $L = \mathcal{O}_B((d_{g,\gamma,3} + k)q - E_1)$; Corollary 2.16 implies that

$$h^0(L^{-1}((d_{g,\gamma,3} + k)q)) = h^0(\mathcal{O}_B(E_1)) = 1, h^0(L^{-1}((d_{g,\gamma,3} + k - 1)q)) = 0$$

and also

$$h^0(L^{-2}((2d_{g,\gamma,3} + 2m)q)) = h^0(\mathcal{O}_B(2Z)) = 1, h^0(L^{-1}((2d_{g,\gamma,3} + 2m - 1)q)) = 0.$$

Our proof is now complete. \square

Case two: a_2 is odd. When $a_2 = 2d_{g,\gamma,3} + 2m + 1$ is odd, we have further restrictions on the feasible range of a beyond what is given in Equation (14):

Lemma 3.24. *Every 6-semigroup indexed by $(a_1, 2d_{g,\gamma,3} + 2m + 1) \in F(3)$ that is Weierstrass realizable by a cyclic cover as in Situation 3.6 satisfies $a_1 \geq d_{g,\gamma,3} + (\gamma - m)$.*

Proof. According to Lemma 3.16, any cyclic cover that realizes $(a_1, 2d_{g,\gamma,3} + 2m + 1)$ is associated with a pair of effective divisors E_1, E_2 for which the corresponding degree-0 divisors $E_i - \deg(E_i) \cdot q$, $i = 1, 2$ are reduced. Moreover, the condition $2[E_1 - \epsilon_1 q] \sim E_2 - (2m + 1)q$ implies that

$$[2\epsilon_1 - (2m + 1)]q \sim 2E_1 - E_2 \sim 2E_1 + \iota(E_2) - 2(2m + 1)q.^7$$

Thus $[2\epsilon_1 + (2m + 1)]q \sim 2E_1 + \iota(E_2)$. Since there is no further cancellation on the two sides, we conclude that there exists a rational function on B with (odd) pole order $2\epsilon_1 + (2m + 1)$ at q . Since q is a Weierstrass point, $2\epsilon_1 + (2m + 1) \geq 2\gamma + 1$ must hold; unwinding yields $a_1 \geq d_{g,\gamma,3} + (\gamma - m)$. \square

Next, we shall show that for every $(a_1, a_2) \in F(3)$ such that $a_1 \geq d_{g,\gamma,3} + (\gamma - m)$ and $a_2 = 2d_{g,\gamma,3} + 2m + 1$, the 6-semigroup indexed by (a_1, a_2) is Weierstrass-realizable via a triple cover described in Situation 3.6. In other words, the previous lemma characterizes the only obstruction to realizability when $N = 3$. We start with a concrete example.

Example 3.25. The realizability problem for the semigroup indexed by $(d_{g,\gamma,3} + \gamma, 2d_{g,\gamma,3} + 2m + 1) \in F(3)$ is solvable by induction on m .

For the base case $m = 0$, we use the fact that $m_2^{-1}\Theta_1 \cap \Theta_{\gamma-1} \subset \text{Jac}(B)[2]$ ([19, Thm 2.5]). Given a general point $q' \in B$, it follows that there is an effective divisor E_1 for which $E_1 - \gamma q$ is reduced and $[2(E_1 - \gamma q)] = [q' - q]$. In particular, the pair (E_1, q') resolves the realizability problem for $(a_1, a_2) = (d_{g,\gamma,3} + \gamma, 2d_{g,\gamma,3} + 1)$.

Now let (E_1, E_2) be a pair of effective divisors as in Lemma 3.16 that resolves the realizability problem for $(a_1, a_2) = (d_{g,\gamma,3} + \gamma, 2d_{g,\gamma,3} + 2m - 1)$; in particular, we have $\deg(E_1) = \gamma$ and $\deg(E_2) = 2m - 1$. Given a general point $q' \in B$, let $D_2 = E_2 + 2q'$. The fact that q' is general guarantees that $D_2 - (2m + 1)q$ is reduced, and trivially we have

$$D_2 - (2m + 1)q \sim 2[E_1 + q' - (\gamma + 1)q].$$

As $\deg(E_1 + q') = \gamma + 1$, we have $E_1 + q' \sim D_1 + q$ for some effective divisor D_1 of degree γ . Thus $D_2 - (2m + 1)q \sim 2(D_1 - \gamma q)$.

We now check that D_1 is not supported at q . Indeed, otherwise we would have

$$E_1 \sim D' - q' + nq \sim D' + \iota(q') + (n - 2)q$$

where D' is effective and not supported at q , and $n \geq 2$. That is,

$$E_1 - \gamma q \sim D' + \iota(q') - [\gamma - (n - 2)]q.$$

⁷Recall here that $\epsilon_j = a_j - jd_{g,\gamma,3}$.

By uniqueness of q -reduced representatives (Lemma 2.15), the latter equality is only possible if $n = 2$ and $E_1 = D' + \iota(q')$, a possibility that is precluded because q' is general.

Finally, we check that $h^0(\mathcal{O}_B(D_1)) = 1$. Given the previous paragraph, either $D_1 - \gamma q$ is reduced, in which case we are done; or else D_1 is simultaneously supported at p and $\iota(p)$ for some $p \in B$. The latter possibility would also violate the uniqueness of q -reduced presentation for $E_1 - \gamma q$, however, in view of $E_1 + q' \sim D_1 + q$. By Lemma 3.16, we are done.

In fact, the same argument leads to a general solution to the realizability problem for all $(a_1, a_2) \in F(3)$ for which $\epsilon_2 = 2m + 1$ is odd and $a_1 \geq d_{g,\gamma,3} + (\gamma - m)$:

Proposition 3.26. *Every a -vector $(d_{g,\gamma,3} + k, 2d_{g,\gamma,3} + 2m + 1)$ for which $k \geq \gamma - m$ and $2m + 1 \leq \gamma$ indexes a 6-semigroup realizable by a cyclic triple cover as in Situation 3.6.*

Proof. According to Lemma 3.16, any solution to the semigroup-realizability problem indexed by $(d_{g,\gamma,3} + k, 2d_{g,\gamma,3} + 2m + 1)$ is presented by a pair of effective divisors $(E_{k,m}^1, E_{k,m}^2)$. In Example 3.25, every pair $(E_{\gamma,m}^1, E_{\gamma,m}^2)$ is such that

- (1) $E_{\gamma,0}^2 = q'$, where q' is a general point on B ; and
- (2) $E_{\gamma,m}^1 + q \sim E_{\gamma,m-1}^1 + q_m$, $E_{\gamma,m}^2 = E_{\gamma,m-1}^2 + 2q_m$, where q_m is general.

In particular, we have $\text{Supp}(E_{\gamma,0}^1) - \text{Supp}(E_{\gamma,0}^2) \neq \emptyset$, as otherwise $(2\gamma - 1)(q' - q) \sim 0$, which would violate the generality of q' . In fact $q' \notin \text{Supp}(E_{\gamma,0}^1)$, as otherwise

$$2E' + q' \sim (2\gamma - 1)q$$

where $E' = E_{\gamma,0}^1 - q'$, which would imply the existence of a rational function on B with pole order $2\gamma - 1$ at q . This is clearly impossible. A similar argument shows that $\text{Supp}(E_{\gamma,0}^1) - \text{Supp}(E_{\gamma,0}^2)$ contains no Weierstrass point of B .

Now suppose that $p \in \text{Supp}(E_{\gamma,0}^1) - \text{Supp}(E_{\gamma,0}^2)$, and set

$$E_{\gamma-1,1}^1 = E_{\gamma,0}^1 - p, E_{\gamma-1,1}^2 = E_{\gamma,0}^2 + 2\iota(p).$$

Thanks to our choice of p , the divisors $E_{\gamma-1,1}^1 - (\gamma - 1)q$, $E_{\gamma-1,1}^2 - 3q$ are q -reduced, and

$$2[E_{\gamma-1,1}^1 - (\gamma - 1)q] \sim E_{\gamma,0}^2 - 2p + 2q \sim E_{\gamma-1,1}^2 - 3q.$$

By Lemma 3.16 again, we obtain a solution for the realizability problem for the semigroup indexed by $(d_{g,\gamma,3} + \gamma - 1, 2d_{g,\gamma,3} + 3)$. Notice that $\text{Supp}(E_{\gamma,1}^1) - \text{Supp}(E_{\gamma,1}^2)$ is nonempty and contains no Weierstrass point; so by induction, we obtain solutions for the realizability problem for those 6-semigroups indexed by $(d_{g,\gamma,3} + \gamma - m, 2d_{g,\gamma,3} + 2m + 1)$ for which $0 \leq m < \frac{\gamma}{2}$.

The general statement now follows from a simple construction. For $m = 1, \dots, \frac{\gamma}{2}$, let $(E_{\gamma-m,m}^1, E_{\gamma-m,m}^2)$ be a solution to the realizability problem for 6-semigroups indexed by $(d_{g,\gamma,3} + \gamma - m, 2d_{g,\gamma,3} + 2m + 1)$. For any m' between 1 and m , let $q_1, \dots, q_{m'}$ be distinct Weierstrass points on B not in $\text{Supp}(E_{\gamma-m,m}^1) \cup \{q\}$.⁸ Let

⁸Such points exist because $\deg(E_{\gamma-m,m}^1) = \gamma - m$, and there are $2\gamma + 2$ distinct Weierstrass points on B .

$E_{\gamma-m+m',m}^1 = E_{\gamma-m,m}^1 + \sum_{k=1}^{m'} q_k$. It follows easily from our choice of $q_1, \dots, q_{m'}$ that $E_{\gamma-m+m',m}^1 - (\gamma - m + m')q$ is q -reduced and that

$$2(E_{\gamma-m+m',m}^1 - (\gamma - m + m')q) \sim 2(E_{\gamma-m,m}^1 - (\gamma - m)q) \sim E_{\gamma-m,m}^2 - (2m + 1)q.$$

It follows that $(E_{\gamma-m+m',m}^1, E_{\gamma-m,m}^2)$ underlies a cyclic cover that realizes the 6-semigroup indexed by $(d_{g,\gamma,3} + \gamma - m + m', 2d_{g,\gamma,3} + 2m + 1)$. \square

4. THE MULTIPLICATION PROFILE OF A POINT ON A HYPERELLIPTIC JACOBIAN

Definition 4.1. Given $x \in \Theta_d^0 \subset \text{Jac}(B)$, let d_j denote the unique integer for which $m_j(x) \in \Theta_{d_j}^0$ for every positive integer j . We refer to the integer sequence $m(x) := (d_1 = d, d_2, d_3, \dots)$ as the *multiplication profile* of x . We also refer to the subsequence $m(x)_{N-1} := (d_j)_{j=1}^{N-1}$ as the $(N-1)$ -th *truncated multiplication profile* of x .

Remark 4.2. When $\gamma = 1$, possible multiplication profiles are as follows: $m(x) = (0, 0, \dots)$ when x is the neutral element; $m(x) = (1, 1, \dots)$ when x is non-torsion; and otherwise $m(x)$ is of some finite order n , in which case the multiplication profile is determined by its truncation $m_n(x) = (1, \dots, 1, 0)$. Throughout the remainder of this section, we assume that $\gamma \geq 2$.

Hereafter, we will focus on *point classes* that arise from Abel-Jacobi maps of degree one.

Notation 4.3. For a fixed point $p \neq q$, let $R_N(p)$ denote the q -reduced representative of $[N(p - q)]$.

Lemma 4.4. *For every index j , we have $d_j > d_{j+1}$ if and only if $\iota(p) \in \text{Supp}(R_j(p))$ for every $p \neq q$.*

Proof. Note that $R_j(p) \sim R_{j+1}(p) + (\iota(p) - q)$. As $d_{j+1} < d_N$, it follows that $R_j(p) = R_{j+1}(p) + (\iota(p) - q)$. Conversely, suppose $R_j(p) = E + (\iota(p) - q)$, where $E \in \Theta_{d_j-1}^0$. Then E is q -reduced, and $R_{j+1}(p) = E$. \square

Corollary 4.5. *We have $|d_j - d_{j+1}| \leq 1$ for every j ; and $d_j \neq d_{j+1}$ whenever $d_j < \gamma$.*

Proof. According to Lemma 4.4, we have $d_{j+1} = d_j - 1$ whenever $d_{j+1} < d_j$. If $d_{j+1} > d_j$, then because $R_j(p) \sim R_{j+1}(p) + (\iota(p) - q)$ is not q -reduced, we must have $R_j(p) = R_{j+1}(p) - (p - q)$, and hence $d_{j+1} = d_j + 1$. On the other hand, if $d_j < \gamma$, then either $\iota(p) \in \text{Supp}(R_j(p))$ and $d_{j+1} < d_j$; or else $\iota(p) \notin \text{Supp}(R_j(p))$ and $d_{j+1} > d_j$. \square

Lemma 4.6. *Fix $p \neq q$. There is no index j for which $d_{j+1} = d_j + 1 = d_{j+2} + 1$, unless $d_j = d_2 = 0$.*

Proof. The equalities $d_{j+1} = d_j + 1 = d_{j+2} + 1$ force $R_{j+1}(p) = R_j(p) + (p - q) = R_{j+2}(p) + (\iota(p) - q)$, in which case $p = \iota(p)$ necessarily. \square

In light of the above, we now develop two instructive examples.

Example 4.7. Whenever $x = [p - q]$ is a general point of Θ_1 , Lemma 4.6 implies that

$$m(x)_{\gamma+1} = (1, 2, \dots, \gamma, \gamma).$$

On the other hand, whenever $x \in \Theta_1$ is 2-torsion, we have $m(x) = (1, 0, 1, 0, \dots)$.

Finally, whenever $x \in \Theta_1$ is $(2\gamma + 1)$ -torsion, we have $m(x)_{2\gamma+1} = (1, 2, \dots, \gamma, \gamma, \gamma - 1, \dots, 1, 0)$, and $m(x)_{2\gamma+1}$ uniquely determines $m(x)$.

The above gives a complete solution for Weierstrass-realizability by cyclic covers whenever $2 \leq N \leq \gamma + 2$ and $a_1 = 1 + d_{g,\gamma,N}$.

Example 4.8. Let a be a general element of k^\times , $1 \leq n \leq \gamma + 1$, and $f(x) = (x - a)^{\gamma+n}x^{\gamma-n+1} + 1$. Working over the hyperelliptic curve $B : y^2 = f(x)$, let $p = (a, 1)$ and $p' = (0, 1)$. Note that $\iota(p) = (a, -1)$ and $\iota(p') = (0, -1)$.

Now $\text{div}(y - 1) = (\gamma + n)p + (\gamma - n + 1)p' - (2\gamma + 1)q$. In particular, we have $[(\gamma + n)(p - q)] = [(\gamma - n + 1)(\iota(p') - q)]$, and it follows that

$$[(\gamma + m)(p - q)] = [(\gamma - n + 1)(\iota(p') - q) + (m - n)(p - q)].$$

Note that $(\gamma - n + 1)(\iota(p') - q) + (m - n)(p - q)$ is q -reduced for every $m = n, \dots, 2n - 1$. We claim that

$$m([p - q])_{\gamma+2n-1} = (1, 2, \dots, \gamma, \gamma, \gamma - 1, \dots, \gamma - n + 1, \gamma - n + 2, \dots, \gamma).$$

Indeed, given our previous discussion, it only remains to show that $d_{\gamma+1+j} = \gamma - j$ for every $j = 1, \dots, n - 2$. This follows, in turn, from the fact that $|d_k - d_{k+1}| \leq 1$: letting R_k denote the q -reduced presentation of $m_k([p - q])$, we have $R_k \sim R_{k+1} + \iota(p) - q$.

Notice that the case in which $[p - q]$ is $(2\gamma + 1)$ -torsion is covered by this example.

4.1. Multiplication profiles of point classes. We begin with a general speculation about the multiplication profile of an element of Θ_1^0 .

Conjecture 4.9. Any sequence $(d_i)_{i=1}^\infty$ of non-negative integers that satisfies the following conditions is realized as the multiplication profile of an element of $[p - q] \in \Theta_1^0(B) \subset \text{Jac}(B)$, for some hyperelliptic curve B of genus γ :

- (1) $d_1 = 1$, and $0 \leq d_j \leq \gamma$ for every j ;
- (2) $d_i + d_j \geq d_{i+j}$;
- (3) $|d_j - d_{j+1}| \leq 1$ and if $d_j < \gamma$, then $d_j \neq d_{j+1}$;
- (4) there is no j for which $d_{j+1} = d_j + 1 = d_{j+2} + 1$, unless $d_j = d_2 = 0$;
- (5) the first index N for which $d_N = 0$ is either $N = 2$ or else $N \geq 2\gamma + 1$, in which case $d_{sN+r} = d_r$ and $d_{N-r} = d_r$ for every $1 \leq r < N$ and $s \geq 0$.

Note that these are all *necessary* conditions. Indeed, items (1), (2) follow from the definition of d_j ; (3) is the content of Corollary 4.5; (4) is Lemma 4.6; while (5) follows from the definition of torsion order and the non-existence of torsion points of small orders ([19, Corollary 2.8]).

To tackle this conjecture, we analyze how to read the multiplication profile of a specific point $p - q$ from the affine equation $y^2 = f(x)$ of the curve B . Without loss of generality, we assume $f(0) \neq 0$ and $p = (0, \sqrt{f(0)})$, where $\sqrt{f(0)}$ is a fixed square root of $f(0)$.

Definition 4.10. Given an effective divisor D on B of degree d , let $p(x) - q(x)y$ be a rational function whose associated principal divisor is $D - dq$. We shall refer to d as the *support length* of this function.

The j -th entry d_j of $m([p - q])$ is codified by a rational function $p_j(x) - q_j(x)y \in K(B)$ that has minimal support length d among all functions that vanish in p to order at least j and are regular away from q . For this minimizing value of d , we have $d_j = d - j$; moreover, the pair (p_j, q_j) is well-defined up to a scalar multiple. Hereafter, we let $\sum_{i=0}^{\infty} a_i x^i$ denote the image of y in $\widehat{\mathcal{O}}_{\mathbb{P}^1, 0} = k[[x]]$ under the natural map $\mathcal{O}_{B, p} \rightarrow \mathcal{O}_{\mathbb{P}^1, 0} \rightarrow \widehat{\mathcal{O}}_{\mathbb{P}^1, 0}$ induced by the hyperelliptic structure map $B \rightarrow \mathbb{P}^1$.

Lemma 4.11. *Up to a scalar multiple, (p_j, q_j) is the unique pair of polynomials $(p(x), q(x))$ for which $x^j | (p - (\sum a_i x^i) \cdot q)$ that minimizes the quantity*

$$\max\{2 \deg(p), 2\gamma + 1 + 2 \deg(q)\}.$$

Proof. The map $\mathcal{O}_{B, p} \rightarrow \mathcal{O}_{\mathbb{P}^1, 0}$ induced by the hyperelliptic structure map sends m_p to $m_0 = (x)$; so the rational function $p(x) - q(x)y \in K(B)$ vanishes in p to order at least j if and only if x^j divides $p(x) - (\sum a_i x^i) \cdot q(x)$. On the other hand, $p(x) - q(x)y$ plainly has support length $\max\{2 \deg(p), 2\gamma + 1 + 2 \deg(q)\}$. \square

Example 4.12. The following are immediate consequences of Lemma 4.11:

- (1) For every $j = 1, \dots, \gamma$, we have $(p_j(x), q_j(x)) = (x^j, 0)$.
- (2) $(p_{\gamma+1}(x), q_{\gamma+1}(x)) = (\sum_{i=0}^{\gamma} a_i x^i, 1)$.
- (3) $(p_{\gamma+2}(x), q_{\gamma+2}(x)) = (\sum_{i=0}^{\gamma+1} a_i x^i, 1)$.
- (4) Suppose $[p - q]$ is N -torsion. Then $\deg p_N(x) = \frac{N}{2}$ and $\deg q_N(x) \leq \frac{N - (2\gamma + 2)}{2}$ whenever N is even; while $\deg p_N(x) \leq \frac{N-1}{2}$ and $\deg q_N(x) = \frac{N - (2\gamma + 1)}{2}$ whenever N is odd.

Definition 4.13. We say that d_{j_0} is a *critical entry* of a multiplication profile (d_j) whenever $d_{j_0} < \min(d_{j_0-1}, d_{j_0+1})$.

Whenever d_j is a critical entry, $\text{div}(p_j(x) - q_j(x)y)$ is not supported at $\iota(p)$. This means that x divides neither p_j nor q_j . Furthermore, the fact that $\text{div}(p_j(x) - q_j(x)y)$ has minimal support length means that p_j and q_j cannot share other linear terms; so $\gcd(p_j, q_j) = 1$. Now set $i = j + d_j - \gamma$. We then have $p_i = p_{i+1} = \dots = p_j$ and $q_i = q_{i+1} = \dots = q_j$, while $p_{j+k} = x^k p_j$ and $q_{j+k} = x^k q_j$ for $k = 1, \dots, \gamma - d_j$.

Lemma 4.14. *Fix $n \geq \gamma + 2$. When $n + \gamma$ is odd (resp. even), we have $d_n < \gamma$ if and only if $D_n = 0$, where*

$$D_n := \begin{vmatrix} a_{\gamma+2} & a_{\gamma+3} & \dots & a_{\frac{n+\gamma+1}{2}} \\ a_{\gamma+3} & a_{\gamma+4} & \dots & a_{\frac{n+\gamma+3}{2}} \\ \vdots & \vdots & & \vdots \\ a_{\frac{n+\gamma+1}{2}} & a_{\frac{n+\gamma+3}{2}} & \dots & a_{n-1} \end{vmatrix} \quad (\text{resp. } D_n := \begin{vmatrix} a_{\gamma+1} & a_{\gamma+2} & \dots & a_{\frac{n+\gamma}{2}} \\ a_{\gamma+2} & a_{\gamma+3} & \dots & a_{\frac{n+\gamma+2}{2}} \\ \vdots & \vdots & & \vdots \\ a_{\frac{n+\gamma}{2}} & a_{\frac{n+\gamma+2}{2}} & \dots & a_{n-1} \end{vmatrix}).$$

Proof. The idea is to calculate p_n and q_n explicitly. Since the argument in each of the two cases is nearly identical, we only discuss the case in which $n + \gamma$ is odd. To begin, recall that the support length of $p_n(x) - q_n(x)y$ is $\max\{2 \deg p_n, 2\gamma + 1 + 2 \deg q_n\}$. For this quantity to be smaller than $\gamma + n$, we must have $\deg q_n < \frac{n - \gamma - 1}{2}$. Now set $q_n(x) = \sum c_i x^i$. Then $(\sum a_i x^i) q_n(x) = \sum e_k x^k$, where $e_k = \sum_{i+j=k} c_i a_j$. Note that $p_n(x)$ is merely an appropriate truncation of $(\sum a_i x^i) q_n(x)$. Clearly we also require $\deg p_n < \frac{n + \gamma}{2}$; it follows that $e_{\frac{n+\gamma+1}{2}} = \dots = e_{n-1} = 0$. By definition

of the e_k , the linear system of equations

$$\begin{bmatrix} a_{\gamma+2} & a_{\gamma+3} & \cdots & a_{\frac{n+\gamma+1}{2}} \\ a_{\gamma+3} & a_{\gamma+4} & \cdots & a_{\frac{n+\gamma+3}{2}} \\ \vdots & \vdots & & \vdots \\ a_{\frac{n+\gamma+1}{2}} & a_{\frac{n+\gamma+3}{2}} & \cdots & a_{n-1} \end{bmatrix} \begin{bmatrix} x_{\frac{n-\gamma-3}{2}} \\ x_{\frac{n-\gamma-1}{2}} \\ \vdots \\ x_0 \end{bmatrix} = 0$$

has a non-zero solution, and the result follows. \square

Remark 4.15. An upshot of Lemma 4.14 is that $d_{\gamma+2} = \gamma - 1$ if and only if $a_{\gamma+1} = 0$; and $d_{\gamma+3} < \gamma$ if and only if $a_{\gamma+2} = 0$.

From our previous discussion about restrictions on multiplication profiles, it is clear that whenever there are indices $j > i$ for which $d_i = d_j = \gamma$ and $d_k < \gamma$ for all k strictly between i and j , we in fact have

$$(d_i, d_{i+1}, \dots, d_{j-1}, d_j) = (\gamma, \gamma-1, \dots, \gamma - \frac{j-i}{2} + 1, \gamma - \frac{j-i}{2}, \gamma - \frac{j-i}{2} + 1, \dots, \gamma-1, \gamma).$$

Consequently, every multiplication profile is uniquely characterized by the vanishing or non-vanishing of the corresponding determinants D_n .

4.2. Multiplication profiles of torsion points. Suppose $x = [p - q] \in \Theta_1^0$ is an N -torsion point. Its multiplication profile $m(x) = (d_j)$ is then determined by $(d_{\gamma+2}, \dots, d_{\lfloor \frac{N}{2} \rfloor})$. Moreover, the following rules apply:

- (1) $d_j = 0$ if and only if $N|j$;
- (2) $d_{\gamma+2} \in \{\gamma - 1, \gamma\}$;
- (3) $d_i + d_j \geq d_{i+j}$;
- (4) there is no j for which $(d_j, d_{j+1}, d_{j+2}) = (a, a + 1, a)$ for any a ; and
- (5) there is no j for which $(d_j, d_{j+1}) = (a, a)$ with $a < \gamma$.

In particular, we have either $d_{\frac{N}{2}} = d_{\frac{N}{2}-1} = \gamma$ or $d_{\frac{N}{2}} = d_{\frac{N}{2}-1} - 1$ when N is even; and $d_{\frac{N-1}{2}} = \gamma$ when N is odd.

Now suppose that $N \geq 2\gamma + 1$, and that $y^2 = f(x)$ is the affine equation of a hyperelliptic curve with an N -torsion divisor $p - q$, where $p = (0, \sqrt{f(0)})$ and $\sum a_i x^i$ is a chosen square root of $f(x)$ inside $k[[x]]$; then as explained previously, there exist polynomials $p_N(x), q_N(x)$ for which x^N divides $p_N(x) - (\sum a_i x^i)q_N(x)$. Conversely, if our aim is to construct $f(x)$ so that the associated hyperelliptic curve has a divisor of particular torsion order, one possible approach involves specifying an approximating power series $\sum a_i x^i$ with certain properties. This approach serves our purposes well, as we may control $m([p - q])$ by imposing conditions on the coefficients a_i .

Proposition 4.16. *Given $p, q, f(x)$, and $\sum a_i x^i$ as before, suppose that $[p - q]$ has order precisely N . Let $(d_j)_{j=1}^{\infty}$ be a sequence satisfying the five necessary conditions stated in the beginning of this subsection. Then $m([p - q]) = (d_j)$ if and only if $D_j = 0$ for precisely those j between $\gamma + 1$ and $\lfloor \frac{N}{2} \rfloor$ for which $d_j < \gamma$.*

Proof. From our earlier discussion, it is clear that the sequence $(d_{\gamma+1}, \dots, d_{\lfloor \frac{N}{2} \rfloor})$ determines the full multiplication profile in this particular case. The desired conclusion now follows from Lemma 4.14 and the discussion following it. \square

We are now ready to give a complete list of conditions that characterize those power series $\sum a_i x^i$ that give rise to a two-pointed hyperelliptic curve (B, p, q) for which the order of $[p - q]$ is precisely N and $m([p - q]) = (d_j)_{j=1}^\infty$. A key observation is that for every fixed value of N , finding a power series that satisfies these conditions is a *finitely-determined* problem. Namely, every coefficient a_j with $j \geq N$ is uniquely determined by a_0, \dots, a_{N-1} ; so the set of admissible power series is a sublocus of the projective space \mathbb{P}^{N-1} in these coordinates.

Conditions for admissibility:

- (Polynomiality and degree): For every $k \geq 2\gamma + 2$, we have $\sum_{i+j=k} a_i a_j = 0$, while $\sum_{i+j=2\gamma+1} a_i a_j \neq 0$.
- (Separability): The Sylvester determinant $S(f)(a_0, \dots, a_{N-1})$ is nonzero (see Notation A.1), where $f(x) = \sum_{k=0}^{2\gamma+1} (\sum_{i+j=k} a_i a_j) x^k$.
- (Torsionness) Whenever N is even (resp. odd), the Hankel matrix

$$(15) \quad M_N := \begin{bmatrix} a_{\gamma+2} & a_{\gamma+3} & \cdots & a_{\frac{N}{2}+1} \\ a_{\gamma+3} & a_{\gamma+4} & \cdots & a_{\frac{N}{2}+2} \\ \vdots & \vdots & & \vdots \\ a_{\frac{N}{2}+\gamma} & a_{\frac{N}{2}+\gamma+1} & \cdots & a_{N-1} \end{bmatrix} \quad (\text{resp., } M_N := \begin{bmatrix} a_{\gamma+1} & a_{\gamma+2} & \cdots & a_{\frac{N+1}{2}} \\ a_{\gamma+2} & a_{\gamma+3} & \cdots & a_{\frac{N+3}{2}} \\ \vdots & \vdots & & \vdots \\ a_{\frac{N-1}{2}+\gamma} & a_{\frac{N+1}{2}+\gamma} & \cdots & a_{N-1} \end{bmatrix})$$

has less-than-maximal rank.

- (Profile-specific conditions): See Proposition 4.16.

Definition 4.17. Given $N \geq 2\gamma + 3$, let \tilde{R}_N denote the closed subset of \mathbb{P}^{N-1} given by the intersection of the closed subset defined by the homogeneous equations $\sum_{i+j=k} x_i x_j = 0$ for $k = 2\gamma + 2, \dots, N - 1$ with the closed subset given by the vanishing of maximal minors of the $(\frac{N}{2} - 1) \times (\frac{N}{2} - \gamma)$ Hankel matrix M_N . Let R_N denote the locally-closed subset given by the intersection of \tilde{R}_N with the open loci $(\sum_{i+j=2\gamma+1} x_i x_j \neq 0)$ and $(S(f)(x_0, \dots, x_{N-1}) \neq 0)$ (see Notation A.1), where $f(x) = \sum_{k=0}^{2\gamma+1} (\sum_{i+j=k} x_i x_j) x^k$.

We think of R_N as a parameter space for square roots of separable polynomials $f(x)$ of degree $2\gamma + 1$ whose corresponding hyperelliptic curves admit N -torsion points $p = (0, \sqrt{f(0)})$. Note that the definition of f guarantees that $a_0 \neq 0$.

4.2.1. *Conjectural classification of multiplication profiles for torsion orders $N \leq 2\gamma + 6$.* According to [7, Proposition 4.3], whenever N is even (resp. odd) the maximal minors of M_N cut out a closed subscheme of \mathbb{P}^{N-1} that is a cone over an integral, normal closed subscheme W of $\mathbb{P}_{(x_{\gamma+2}, \dots, x_{N-1})}^{N-\gamma-2}$ (resp. $\mathbb{P}_{(x_{\gamma+1}, \dots, x_{N-1})}^{N-\gamma-1}$) of codimension γ . In particular, the ideal I generated by the maximal minors of M_N is a homogeneous prime ideal generated by some homogeneous elements of degree $\lfloor \frac{N+1}{2} \rfloor - \gamma$.

In light of [19, Cor. 2.8], we may assume that $N \geq 2\gamma + 1$ without loss of generality. Moreover, as explained in Examples 5.1, 5.2, and 5.5 below, whenever $N \in \{2\gamma + 1, 2\gamma + 2, 2\gamma + 3, 2\gamma + 5\}$ a single multiplication profile is possible; so the interesting cases are those for which $N = 2\gamma + 4$ or $N = 2\gamma + 6$. On the other hand, Example 5.4 (resp., Example 5.7) establishes that there are two (resp., three) possible multiplication profiles whenever $N = 2\gamma + 4$ (resp., $N = 2\gamma + 6$).

Experimental computer evidence suggests that each of these profiles determines a distinguished component of the corresponding parameter space R_N .

Conjecture 4.18. *Suppose that either $N = 2\gamma + 4$ or $N = 2\gamma + 6$. Each irreducible component V_i of the parameter space \tilde{R}_N is cut out by a prime ideal I_i generated by the polynomiality quadrics $\sum_{i=0}^{N-1} x_i x_{k-i}$, $k = 2\gamma + 2, \dots, N$ and maximal minors of the Hankel matrix M_N in (15), together with a distinguished set of auxiliary generators. More precisely, we have*

$$\tilde{R}_{2\gamma+4} = V_1 \cup V_2$$

where $I_1 = I(V_1)$ is generated by the polynomiality quadrics, 2×2 Hankel minors, and the quadrics $2x_{\lfloor \frac{\gamma+1+j}{2} \rfloor} x_{\lfloor \frac{\gamma+1+j}{2} \rfloor + 1} - x_{\gamma+1} x_j$, $j = \gamma + 3, \dots, 2\gamma + 3$; while $I_2 = I(V_2)$ is generated by the polynomiality quadrics, maximal Hankel minors, and $x_{\gamma+2}$. Similarly, we have a decomposition

$$\tilde{R}_{2\gamma+6} = V_1 \cup V_2 \cup V_3.$$

This time, $I_1 = I(V_1)$ is generated by the polynomiality quadrics, 3×3 Hankel minors, and the cubics

$$-2x_{\gamma+3} D_{\ell_1, \ell_2}^{(1,2)} + x_{\gamma+2} D_{\ell_1, \ell_2}^{(1,3)}, (\ell_1, \ell_2) \in \binom{[\gamma+2]}{2}$$

in which $D_{\ell_1, \ell_2}^{(j_1, j_2)}$ denotes the 2×2 determinant associated to rows (ℓ_1, ℓ_2) and columns (j_1, j_2) of $M_{2\gamma+6}$. The ideal $I_2 = I(V_2)$ is generated by the polynomiality quadrics, 3×3 Hankel minors, the cubics

$$2x_{\gamma+3} D_{\ell_1, \ell_2}^{(1,2)} + x_{\gamma+1} D_{\ell_1, \ell_2}^{(2,3)}$$

in which $(\ell_1, \ell_2) \in \binom{[\gamma+2]}{2}$ indexes a pair of rows with $\ell_i \geq 2$, $i = 1, 2$, the cubics

$$2x_{\gamma+3}(x_{\gamma+1}x_{j+2} - x_{\gamma+3}x_j) - x_{\gamma+1}(x_{\gamma+4}x_{j+1} + x_{\gamma+3}x_{j+2}), j = \gamma + 3, \dots, 2\gamma + 3$$

and $x_{\gamma+2}$. The ideal $I_3 = I(V_3)$ is generated by the polynomiality quadrics, 3×3 Hankel minors, and x_j , $j = \gamma + 1, \gamma + 2$.

Remark 4.19. Note that Conjecture 4.18 is coherent with the numerology of Examples 5.4 and 5.7, respectively. Indeed, the order of the components we have chosen is in increasing order of speciality here, as it is there. The conjecture predicts that a general point of V_i corresponds to an N -torsion point whose multiplication profile is (uniquely determined by) the i -th sequence listed in the corresponding example.

5. EXPLICIT REALIZABILITY CONSTRUCTIONS

Example 5.1. Let $x = [p - q]$ be a 2-torsion (resp. $(2\gamma + 1)$ -torsion, $(2\gamma + 3)$ -torsion) point. Then $m(x)$ is determined by $m(x)_2 = (1, 0)$ (resp. $m(x)_{2\gamma+1} = (1, 2, \dots, \gamma, \gamma, \dots, 1, 0)$, $m(x)_{2\gamma+3} = (1, 2, \dots, \gamma, \gamma, \gamma, \gamma, \dots, 1, 0)$).

Example 5.2. Here we compute the multiplication profiles of those torsion points constructed in Example 2.22.

Reducing the Mumford presentation $(x^{\gamma+s}, v(x) = \sqrt{\frac{-1}{2}}(-x^{\gamma-s+1} + 1))$ using Cantor's algorithm, we obtain $\frac{f(x) - v^2(x)}{x^{\gamma+s}} = c(x^{\gamma-s+1} - 1)$. It follows that $(\gamma + s) \cdot$

$(p - q) \sim \sum((\xi_j, 0) - q)$, where ξ_j runs through all $(\gamma - s + 1)$ -th roots of unity. In particular, $d_{\gamma+s} = \gamma - s + 1$ and the multiplication profile of $p - q$ is determined by

$$m(p - q)_{2\gamma+2s} = (1, \dots, \gamma, \gamma, \gamma - 1, \dots, \gamma - s + 1, \dots, \gamma, \gamma, \dots, 1, 0).$$

Note that any $(2\gamma + 2)$ -torsion point must have the latter profile, with $s = 1$.

The following observation is obvious, yet useful in practice:

Lemma 5.3. *For every even number $N \geq 2\gamma + 1$, $[p - q]$ has order N if and only if $\frac{N}{2}(p - q)$ has order 2.*

On the other hand, elements of $\text{Jac}(B)[2]$ have q -reduced presentations of the form $\sum_{i=1}^s q_i - sq$, where q_i are pairwise-distinct Weierstrass points distinct from q . Finally, up to linear change of coordinates, we may always assume that the x -coordinate of any fixed torsion point is 0. Together these three observations often enable us to simplify the equations involved in determining the multiplication profiles of torsion points of even order.

Example 5.4. Fix $s, r \geq 0$ such that $s + r = \gamma + 1$. For a general choice of $(a_1, \dots, a_s, b_1, \dots, b_r)$, Lemma A.2 implies that $x^{2\gamma+2-s} - \prod_{i=1}^s (x - a_i) \cdot \prod_{j=1}^r (x - b_j)^2$ is separable of degree $2\gamma + 1 - s$.⁹ Letting $c = (\sum a_i + 2\sum b_j)^{-1}$, it follows that

$$f(x) = c \cdot \prod_{i=1}^s (x - a_i) [x^{2\gamma+2-s} - \prod_{i=1}^s (x - a_i) \cdot \prod_{j=1}^r (x - b_j)^2]$$

is a monic, separable polynomial of degree $2\gamma + 1$. Now let

$$v(x) = \sqrt{-c} \prod_{i=1}^s (x - a_i) \cdot \prod_{j=1}^r (x - b_j), \text{ so that } f(x) - v^2(x) = c \cdot \prod_{i=1}^s (x - a_i) \cdot x^{2\gamma+2-s}.$$

Letting $p = (0, v(0))$, we see that $(2\gamma + 2 - s)(p - q) \sim \sum_{i=1}^s ((a_i, 0) - q)$ is a 2-torsion point of the Jacobian of $B : y^2 = f(x)$.

When $s = \gamma$, it follows that $x = p - q$ is of order $2\gamma + 4$ and its multiplication profile is determined by $m_{2\gamma+4}(x) = (1, 2, \dots, \gamma, \gamma, \gamma, \gamma, \gamma - 1, \dots, 1, 0)$. This profile and that of Example 5.2 (with $s = 2$) are the only two possible profiles for a $(2\gamma + 4)$ -torsion element in Θ_1^0 on some hyperelliptic Jacobian.

More generally, for every $s = 1, \dots, \gamma$, $x = p - q$ is of order $4\gamma + 4 - 2s$ and its multiplication profile is determined by

$$m_{4\gamma+4-2s}(x) = (1, 2, \dots, \gamma, \gamma, \gamma - 1, \dots, s, s + 1, \dots, \gamma, \gamma, \gamma, \dots, 1, 0).$$

Example 5.5. Say that $x = p - q$ is a $(2\gamma + 5)$ -torsion point; then $m(x)$ is determined by

$$m(x)_{2\gamma+5} = (1, \dots, \gamma, \gamma, \gamma, \gamma, \gamma, \gamma, \dots, 0).$$

Indeed, according to Example 4.7, every d_j is already determined, except for $d_{\gamma+2} = d_{\gamma+3}$. On the other hand, notice that $2 \cdot (\gamma + 2) \cdot x = [u(p) - q] \in \Theta_1^0$; it follows from [19, Theorem 2.5] that $d_{\gamma+2} = d_{\gamma+3} = \gamma$. (Alternatively, use condition (4) in Conjecture 4.9.)

⁹The generality assumption guarantees that $e_1(a, \dots, a_s, b_1, b_1, \dots, b_r, b_r) \neq 0$, and the conclusion on the degree of the difference follows.

Example 5.6. Fix $1 \leq s \leq \gamma - 1$, and let $N = 2\gamma + 1 - s$. By Lemma A.4, $g(x) = cx^N + \prod_{i=1}^s (x - a_i) \cdot (x - b)^2$ is separable of degree N , for general choices of a_1, \dots, a_s, b, c . Let $a_{s+1}, \dots, a_{2\gamma+1}$ denote the roots of g ; then $a_1, \dots, a_{2\gamma+1}$ are all distinct. Let

$$f(x) = \prod_{i=1}^{2\gamma+1} (x - a_i) \text{ and } v(x) = \sqrt{c^{-1}} \prod_{i=1}^s (x - a_i)(x - b).$$

Set $p = (0, v(0))$. Reducing $(x^N, v(x))$ one time using Cantor's algorithm, we conclude that $N(p - q) \sim \sum_{i=1}^s ((a_i, 0) - q)$. Consequently, $x = p - q$ has order $4\gamma + 2 - 2s$ and $m(x)$ is determined by

$$m(x)_{4\gamma+2-2s} = (1, \dots, \gamma, \gamma, \gamma - 1, \dots, s, s + 1, \dots, \gamma, \gamma, \gamma - 1, \dots, 0).$$

This is the multiplication profile of a $(4\gamma + 2 - 2s)$ -torsion point in Θ_1 for which $d_{2\gamma+1-s}$ is minimized.

Example 5.7. Let $N = 2\gamma + 6$. By symmetry, candidate multiplication profiles associated with torsion points of this order are determined by either

- (1) $(d_{\gamma+2}, d_{\gamma+3}) = (\gamma, \gamma)$;
- (2) $(d_{\gamma+2}, d_{\gamma+3}) = (\gamma, \gamma - 1)$; or
- (3) $(d_{\gamma+2}, d_{\gamma+3}) = (\gamma - 1, \gamma - 2)$.

The second (resp., third) possibility is realized in Ex. 5.4 (resp., Ex. 5.6) with $s = \gamma - 1$ (resp., $s = \gamma - 2$). We claim the first possibility is also realizable. Indeed, set $s = \gamma + 1$ and $\ell = 3$; and apply Lemma A.3 to obtain a separable polynomial of degree γ of the form $g(x) = x^{\gamma+3} - \prod_{i=1}^s (x - a_i) \cdot (x - b)^2 := c \cdot \prod_{i=\gamma+2}^{2\gamma+1} (x - a_i)$. The generality of the a_i together with the fundamental theorem of symmetric polynomials guarantees that the constants $a_1, \dots, a_{\gamma+1}, b$ are all non-zero; and it's easy to see that $a_1, \dots, a_{2\gamma+1}$ are pairwise distinct, provided that $a_1, \dots, a_{\gamma+1}$ are. In the case at hand, to ensure that the a_i , $i = 1, \dots, \gamma + 1$ are pairwise distinct, we use the fact that $4e_2(x_1, \dots, x_{\gamma+3}) - 3e_1^2(x_1, \dots, x_{\gamma+3}) = 0$ admits a solution $(a_1, \dots, a_{\gamma+3})$ whose entries are pairwise distinct. Setting $b = -\frac{1}{2}(\sum a_i)$, we obtain $e_1(a_1, \dots, a_{\gamma+3}, b, b) = e_2(a_1, \dots, a_{\gamma+3}, b, b) = 0$.

Now let $f(x) = \prod_{i=1}^{2\gamma+1} (x - a_i)$, $u_0(x) = x^{\gamma+3}$, $v_0(x) = \sqrt{-c} \prod_{i=1}^{\gamma+1} (x - a_i) \cdot (x - b)$. Apply the reduction algorithm once to the pair (u_0, v_0) , we obtain $u_1(x) = \prod_{i=1}^{\gamma+1} (x - a_i)$. In particular, we have $(\gamma + 3) \cdot (p - q) \sim \sum_{i=1}^{\gamma+1} (q_i - q)$, where $p = (0, v(0))$ and $q_i = (a_i, 0)$. We conclude that $x = p - q$ has order $2\gamma + 6$ and $\text{div}(y) = \sum_{i=1}^{2\gamma+1} (q_i - q)$; so $m(x)$ matches the first candidate profile on our list.

Example 5.8. Let $N = 2\gamma + 7$. By symmetry, candidate multiplication profiles are indexed by

- (1) $(d_{\gamma+2}, d_{\gamma+3}) = (\gamma, \gamma)$; and
- (2) $(d_{\gamma+2}, d_{\gamma+3}) = (\gamma - 1, \gamma)$.

Both profiles are realized in Appendix B. For the first (resp., second), we require that $\gamma \geq 3$ (resp., $\gamma \geq 4$).

Example 5.9. When $N = 2\gamma + 8$, candidate profiles are indexed by

- (1) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma, \gamma, \gamma)$;
- (2) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma, \gamma - 1, \gamma - 2)$;
- (3) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma, \gamma, \gamma - 1)$;

- (4) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma - 1, \gamma - 2, \gamma - 3)$; and
 (5) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma - 1, \gamma, \gamma)$.

Of these, the second is realized in Ex. 5.4, setting $s = \gamma - 2$. The third may be realized using a similar construction applying Lemma A.3 as used to realize the first listed profile in Ex. 5.7. The fourth is realized in Ex. 5.6, setting $s = \gamma - 3$.

According to Lemma 4.14, to realize the first profile we need to find a power series $\sum a_i x^i$ in $R_{2\gamma+8}$ for which $a_{\gamma+1} \neq 0$, $a_{\gamma+2} \neq 0$, and $a_{\gamma+1}a_{\gamma+3} - a_{\gamma+2}^2 \neq 0$. Similarly, in order to realize the fifth profile, the required conditions are $a_{\gamma+1} = 0$ and $a_{\gamma+2} \neq 0$.

Example 5.10. When $N = 2\gamma + 9$, candidate profiles are indexed by

- (1) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma, \gamma, \gamma)$;
 (2) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma, \gamma - 1, \gamma)$; and
 (3) $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma - 1, \gamma, \gamma)$.

The first profile is realized in Appendix B for every $\gamma \geq 4$.

Example 5.11. The classification of indexing sequences for candidate multiplication profiles associated with torsion points of arbitrary even order $2\gamma + 2k$ is similar to that of Example 5.9; an upshot is that the number N_k of candidate profiles associated with torsion order $2\gamma + 2k$ is the $(k + 1)$ -th Fibonacci number for every k . Indeed, we claim that

$$(16) \quad N_k = \sum_{i=1}^{k-2} N_i + 2$$

for every positive integer $k \geq 3$. To see this, note that indexing sequences $(d_{\gamma+2}, d_{\gamma+3}, \dots, d_{\gamma+k})$ are of four possible types: (γ, γ, \dots) , $(\gamma, \gamma - 1, \dots)$, $(\gamma - 1, \gamma, \dots)$, and $(\gamma - 1, \gamma - 2, \dots)$. In the first case, the remaining $(k - 3)$ entries comprise an arbitrary profile associated with a torsion order of index $(k - 2)$, as manifested by the instance of N_{k-2} on the right-hand side of (16). Similarly, in the third case, $d_{\gamma+4} = \gamma$ is forced, and the remaining $(k - 4)$ entries comprise an arbitrary profile associated with a torsion order of index $(k - 3)$; these account for the instance of N_{k-3} . In the second and fourth cases, we have the following dichotomy: either the sequence is strictly decreasing (here there are precisely two possibilities, which accounts for the instance of 2 in (16)) or it eventually reverses course and starts to increase. In the latter situation, the sequence starts increasing after j steps, where $2 \leq j \leq \lfloor \frac{k}{2} \rfloor - 1$, as symmetry requires that this first ascent end in γ, γ ; and by varying the choice of j we obtain the remaining terms on the right-hand side of (16).

On the other hand, the number \tilde{N}_k of candidate multiplication profiles associated with torsion points of arbitrary *odd* order $2\gamma + 2k - 1$, $k \geq 1$ is equal to the k -th Fibonacci number. Indeed, an analysis similar to that of the previous paragraph yields $\tilde{N}_k = \sum_{i=1}^{k-2} \tilde{N}_i + 1$ for every $k \geq 3$.

APPENDIX A. SEPARABILITY RESULTS FOR UNIVARIATE POLYNOMIALS

A classical formula of Sylvester establishes that a univariate polynomial $f(x) \in k[x]$ has a multiple root if and only if the *resultant* of (f, f') is zero. Here we derive some separability criteria for univariate polynomials of a particular form based on

an analysis of the corresponding resultants, in order to justify the constructions carried out in the body of the paper.

Notation A.1. Let $S(f)$ denote the Sylvester polynomial of $f(x) = \sum_{i=0}^n a_i x^i$; that is, the determinant of the following $(2n-1) \times (2n-1)$ matrix:

$$\begin{bmatrix} a_n & 0 & \dots & 0 & na_n & 0 & \dots & 0 \\ a_{n-1} & a_n & \dots & 0 & (n-1)a_{n-1} & na_n & \dots & 0 \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ a_1 & a_2 & \dots & a_{n-1} & a_1 & 2a_2 & \dots & na_n \\ a_0 & a_1 & \dots & a_{n-2} & 0 & a_1 & \dots & (n-1)a_{n-1} \\ 0 & a_0 & \dots & a_{n-3} & 0 & 0 & \dots & (n-2)a_{n-2} \\ \vdots & \vdots & & \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & a_0 & 0 & 0 & \dots & a_1 \end{bmatrix}.$$

Lemma A.2. Let $(a_1, \dots, a_s, b_1, \dots, b_r)$ be a general point of $\mathbb{A}^{s+r}(k)$ and $N = s + 2r$. Then $f(x) = x^N - \prod_{i=1}^s (x - a_i) \cdot \prod_{j=1}^r (x - b_j)^2$ is a separable polynomial.

Proof. Let e_i denote the i -th elementary symmetric polynomial in N variables, and let $\bar{e}_i = e_i(-a_1, \dots, -a_s, -b_1, -b_1, \dots, -b_r, -b_r)$. Then $f(x) = -\sum_{i=1}^N \bar{e}_i x^{N-i}$. By definition, the Sylvester polynomial $S(f)$ of $f(x)$ is homogeneous of degree $2N - 3$ in $(\bar{e}_1, \dots, \bar{e}_N)$, since $\deg(F) = N - 1$.

Now let $F : \mathbb{A}^N \rightarrow \mathbb{A}^N$ denote the morphism

$$(a_1, \dots, a_s, b_1, \dots, b_{2r}) \mapsto (\bar{e}_i)_{i=1}^N = (e_i(-a_1, \dots, -a_s, -b_1, \dots, -b_{2r}))_{i=1}^N$$

that sends the roots of a degree- N polynomial to the coefficients of its expansion. Pulling $S(f)$ back along F , we obtain a polynomial S that is homogeneous in $a_1, \dots, a_s, b_1, \dots, b_{2r}$ of degree $N^2 - 2$.

To conclude, it suffices to show that S restricts to a non-zero function on the linear subspace defined by $b_{2i-1} - b_{2i} = 0$, for $i = 1, \dots, r$. For this purpose, it further suffices to show that S restricts to a non-zero function on the linear subspace defined by $b_1 = \dots = b_r = 1$. In fact, the polynomial $x^N - (x-1)^N$ is separable, with roots $\frac{\xi^i}{\xi^i - 1}$ ($i = 1, \dots, N-1$), where ξ is a primitive N -th root of unity; but this means exactly that $S(1, \dots, 1) \neq 0$. \square

Using basic properties of symmetric polynomials, we obtain a strengthened form of Lemma A.2:

Lemma A.3. Suppose $N = s + 2$ and that (a_1, \dots, a_s, b, b) is a general point of the algebraic subset of $\mathbb{A}^N(k)$ defined by $e_1(x_1, \dots, x_N) = \dots = e_{\ell-1}(x_1, \dots, x_N) = 0$ and $x_{s+2} - x_{s+1} = 0$, where $1 \leq \ell < N$.¹⁰ Then $f(x) = x^N - \prod_{i=1}^s (x - a_i) \cdot (x - b)^2$ is separable.

¹⁰When $\ell = 1$, we interpret these to be the hypotheses of Lemma A.2.

Proof. Under the given assumptions, we have $f(x) = -\sum_{i=\ell}^N \overline{e_i} x^{N-i}$; the corresponding Sylvester polynomial $S(f)$ is¹¹

$$- \begin{vmatrix} \overline{e_\ell} & 0 & \dots & 0 & 0 & 0 & \dots & 0 \\ \overline{e_{\ell+1}} & \overline{e_\ell} & \dots & 0 & -\overline{e_{\ell+1}} & 0 & \dots & 0 \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ \overline{e_{N-2}} & \overline{e_{N-3}} & \dots & \overline{e_\ell} & -(N-\ell-2)\overline{e_{N-2}} & -(N-\ell-3)\overline{e_{N-3}} & \dots & 0 \\ \overline{e_{N-1}} & \overline{e_{N-2}} & \dots & \overline{e_{\ell+1}} & -(N-\ell-1)\overline{e_{N-1}} & -(N-\ell-2)\overline{e_{N-2}} & \dots & (N-\ell)\overline{e_\ell} \\ \overline{e_N} & \overline{e_{N-1}} & \dots & \overline{e_{\ell+2}} & -(N-\ell)\overline{e_N} & -(N-\ell-1)\overline{e_{N-1}} & \dots & (N-\ell-1)\overline{e_{\ell+1}} \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \overline{e_N} & 0 & 0 & \dots & \overline{e_{N-1}} \end{vmatrix}.$$

In particular, it is homogeneous of degree $2N - 2\ell - 1$ in $\overline{e_1}, \dots, \overline{e_N}$, and symmetric and homogeneous of degree $N^2 - N - \ell^2$ in the affine coordinates x_1, \dots, x_N of \mathbb{A}^N .

Our aim is to show that the pull-back of $S(f)$ to the algebraic set Z defined by $e_i(x_1, \dots, x_N) = 0$ ($i = 1, \dots, \ell - 1$) and $x_{s+2} - x_{s+1} = 0$ is not identically zero. Indeed, if $S(f)$ vanishes along Z , then because $S(f)$ is symmetric it must vanish on the algebraic set defined by $e_i(x_1, \dots, x_N) = 0$, $i = 1, \dots, \ell - 1$ and $V_N^2 = 0$, where $V_N(x_1, \dots, x_N) = \prod_{i < j} (x_i - x_j)$ is the Vandermonde polynomial in N variables. Thus, letting $I = \langle e_1, \dots, e_{\ell-1}, V_N^2 \rangle$, we have $S(f) \in \sqrt{I}$, i.e., $S(f)^m = P_N(0, \dots, 0, e_\ell, \dots, e_N) \cdot h(e_\ell, \dots, e_N)$ for some $m \geq 1$, where $V_N^2 = P_N(e_1, \dots, e_N)$. However, $S(f)$ is irreducible in the variables e_1, \dots, e_N ; see [10, Example 1.1.4]. So the vanishing of $S(f)$ along Z would imply that $P_N(0, \dots, 0, e_\ell, \dots, e_N) = S(f)^{m'}$ for some m' . This last possibility, however, is impossible as $P_N(0, \dots, 0, e_\ell, \dots, e_N)$ is *inhomogeneous* in e_ℓ, \dots, e_N . Indeed, via Newton's method we may express $V_n^2(x_1, \dots, x_n)$ as a polynomial in elementary symmetric functions $\sum_{\lambda \vdash n(n-1)} c_\lambda^{(n)} e_\lambda$, where $e_\lambda = \prod e_{\lambda_i}$ and $c_{n^{n-1}}^{(n)} \neq 0$.¹² The fact that the coefficients $c_{n^{n-1}}^{(n)}$ are nonzero follows via induction using the relation $V_n^2 = P_{n-1}(e_1, \dots, e_{n-1})e_{n-1}^2 + e_n h_n$. \square

The next lemma builds on the same idea, and its proof is even simpler.

Lemma A.4. *Fix $s \geq 1$ and $N > s+2r$. For general choices of $(a_1, \dots, a_s, b_1, \dots, b_r) \in \mathbb{A}^{s+r}(k)$ and $c \in k$, the polynomial $f(x) = c \cdot x^N + \prod_{i=1}^s (x - a_i) \prod_{j=1}^r (x - b_j)^2$ is separable.*

Proof. For a fixed choice of $a_1, \dots, a_s, b_1, \dots, b_r$, $S(f)$ is a polynomial in c of degree N whose leading term is $N^N e^{N-1} c^N$, where $e = \prod a_i \cdot \prod b_j^2$. (As the a_i and b_j are general, we have $e \neq 0$.) Hence $S(f) \neq 0$ for a general value of c . \square

APPENDIX B. FLYNN'S MODELS

In [9], Flynn developed effective methods for producing rational torsion points of large orders on special families of hyperelliptic curves. In this section, we use Flynn's models in order to certify the realizability of certain multiplication profiles.

¹¹Starting from the definition, add $-(N - \ell - 1)$ times the i th column to the $(N - \ell - 2 + i)$ th column for $i = 1, \dots, N - \ell - 2$.

¹²This is a consequence of the fundamental theorem of symmetric polynomials.

B.1. Torsion orders that grow linearly with respect to the genus. In *loc. cit.* Flynn recursively defined sequences of polynomials $\phi_r(x), \theta_r(x)$ given by

$$\theta_{r+1}(x) = (x+2)\theta_r(x) + 2(x+1)\phi_r(x), \phi_{r+1}(x) = 2\theta_r(x) + (x+2)\phi_r(x)$$

subject to $\theta_1(x) = \phi_1(x) = 1$; and studied genus- γ hyperelliptic curves with presentations

$$(17) \quad y^2 + (\phi_r(x)y - \theta_r(x)x^\gamma)w(x) = x^{2\gamma+1} + x^{2\gamma}$$

where $w(x)$ is a polynomial of degree at most $\gamma - 2r + 1$ and $w(0) \notin \{0, -4^{2-r}\}$. Flynn showed that for $p = (0, 0)$ and q equal to the point over infinity, $p - q$ is a torsion element in the Jacobian of (17) of order $2g + 2r - 1$ ([9, Result 1(a)]).

Our goal is to determine the multiplication profile of $p - q$. By symmetry, this is tantamount to computing $d_{\gamma+i}$, for $i = 2, \dots, r$ (though sometimes the profile will be determined by a smaller subset of the d_j). When i is small, the task further reduces to calculating the degree of $v(x)$, where $(x^{\gamma+i}, v(x))$ is the Mumford presentation for $(\gamma + i) \cdot (p - q)$. For example, when $i = 2$, we have $d_{\gamma+2} = \gamma$ if $\deg(v) = \gamma + 1$ and $d_{\gamma+2} = \gamma - 1$ if $\deg(v) \leq \gamma$. Indeed, the reduction algorithm terminates in one step in both cases and $d_{\gamma+2} = \deg(\frac{v^2 + vhw - f}{x^{\gamma+2}})$. Similarly, let $(x^{\gamma+3}, v(x))$ be the Mumford presentation of $(\gamma + 3)(p - q)$. Then $d_{\gamma+3} = \gamma$ if $\deg(v) = \gamma + 2$ and $d_{\gamma+3} \leq \gamma - 1$ otherwise; either way, the reduction algorithm terminates in two steps.

Notation B.1. Given a polynomial $f(x)$, let $f(x)_i$ denote the x^i -coefficient of its expansion.

To compute the degree of $v(x)$ in these cases, we first solve the equation

$$Z^2 + \phi_r(x)w(x)Z - \theta_r(x)w(x)x^\gamma = 0, \quad i = 2, 3$$

over $k[[x]]$, and then appropriately truncate the result. Since the discriminant of the left-hand side is $\phi_r^2 w^2 + 4\theta_r w x^\gamma$, it has a square root of the form $\phi_r(x)w(x) + \sum_{i=\gamma}^{\infty} a_i x^i$. This, in turn, leads to the following equations:

$$\begin{cases} 2(\phi_r w)_0 \cdot a_\gamma = 4(\theta_r w)_0 \\ 2(\phi_r w)_1 \cdot a_\gamma + 2(\phi_r w)_0 \cdot a_{\gamma+1} = 4(\theta_r w)_1 \\ 2(\phi_r w)_2 \cdot a_\gamma + 2(\phi_r w)_1 \cdot a_{\gamma+1} + 2(\phi_r w)_0 \cdot a_{\gamma+2} = 4(\theta_r w)_2 \\ \dots \end{cases}$$

Using the inductive definitions of ϕ and θ , respectively, we deduce that

$$(\phi_r)_0 = (\theta_r)_0 = 4^{r-1} \text{ and } (\theta_{r+1})_k - (\phi_{r+1})_k = (\theta_r)_{k-1} + (\phi_r)_{k-1}, 1 \leq k \leq r.$$

Thus, $a_\gamma = 2$, $a_{\gamma+1} = \frac{2((\theta_r w)_1 - (\phi_r w)_1)}{(\phi_r w)_0} = \frac{2((\theta_{r-1} w)_0 + (\phi_{r-1} w)_0)}{(\phi_r w)_0} = 1$, and

$$2(\phi_r)_1(w)_0 + 8(\phi_{r-1})_0(w)_1 + 2(\phi_r w)_0 a_{\gamma+2} = 4[(\theta_r w)_2 - (\phi_r w)_2].$$

In view of the fact that

$$(\theta_r w)_2 - (\phi_r w)_2 = [(\theta_r)_2 - (\phi_r)_2]w_0 + [(\theta_r)_1 - (\phi_r)_1]w_1 = [(\theta_{r-1})_1 + (\phi_{r-1})_1]w_0 + [(\theta_{r-1})_0 + (\phi_{r-1})_0]w_1$$

it follows that

$$2(\phi_r)_1(w)_0 + 2(\phi_r w)_0 a_{\gamma+2} = 4[(\theta_{r-1})_1 + (\phi_{r-1})_1]w_0.$$

Finally, applying the relation $(\phi_r)_1 = 2(\theta_{r-1})_1 + 2(\phi_{r-1})_1 + (\phi_{r-1})_0$, we deduce that $2(\phi_{r-1} w)_0 + 2(\phi_r w)_0 a_{\gamma+2} = 0$, so $a_{\gamma+2} = -\frac{1}{4}$. Since we are merely interested

in realizing multiplication profiles, we may fix $w(x) = 1$ without loss of generality (and thereby remove the restriction placed on γ in [9]). Letting $r = 4, 5$ yields the following corollary:

Corollary B.2. *For $r = 4$ and 5 , let $p = (0, 0)$ be the point defined as above. When $r = 4$ and $\gamma \geq 3$ (resp. $r = 5$ and $\gamma \geq 4$), $p - q$ has order $2\gamma + 7$ (resp. $2\gamma + 9$) and $m(p - q)$ is the multiplication profile determined by $(d_{\gamma+2}, d_{\gamma+3}) = (\gamma, \gamma)$ (resp. $(d_{\gamma+2}, d_{\gamma+3}, d_{\gamma+4}) = (\gamma, \gamma, \gamma)$).*

B.2. A variant of Flynn's construction. Assume $\gamma \geq 4$. Let B denote the hyperelliptic affinely presented by $y^2 + (x^3 + 4)y = x^{2\gamma+1} + x^{2\gamma-2} - (3x^3 + 4)x^{\gamma-1}$. Let $p = (0, 0)$ and q be the point over infinity. We claim that the principal divisor associated to the rational function $(3x^3 + 4)x^{\gamma-1} + (x^3 + 4)y$ on B is $(2\gamma + 7)(p - q)$.

To begin, note that when $(3x^3 + 4)x^{\gamma-1} + (x^3 + 4)y = 0$, we have

$$(3x^3 + 4)^2 x^{2\gamma-2} - (x^3 + 4)^2 y^2 = (3x^3 + 4)^2 x^{2\gamma-2} - (x^3 + 4)^2 (x^{2\gamma+1} + x^{2\gamma-2}) = -x^{2\gamma+7} = 0.$$

As a result, the given function vanishes only at $p = (0, 0)$, and its associated principal divisor is $\text{div}((3x^3 + 4)x^{\gamma-1} + (x^3 + 4)y) = (2\gamma + 7)(p - q)$.

Now $\iota(p) = (0, -4)$; so p is not a Weierstrass point. Furthermore $2\gamma + 7$ is not divisible by $2\gamma + i$, $1 \leq i \leq 6$ when $\gamma \geq 2$; so $p - q$ is of order exactly $2\gamma + 7$.

In order to determine $m(p - q)$, we solve the quadratic equation $Z^2 + (x^3 + 4)Z - 16x^{\gamma-1} = 0$ over $k[[x]]$.¹³ A square root of the discriminant is of the form $x^3 + 4 + \sum_{i \geq \gamma-1} a_i x^i$, which leads to the following equations:

$$\begin{cases} 2 \cdot 4 \cdot a_{\gamma-1} = 4 \cdot (-4) \\ 2 \cdot 0 \cdot a_{\gamma-1} + 2 \cdot 4 \cdot a_{\gamma} = 0 \\ 2 \cdot 0 \cdot a_{\gamma-1} + 2 \cdot 0 \cdot a_{\gamma} + 2 \cdot 4 \cdot a_{\gamma+1} = 0 \\ \dots \end{cases}$$

Hence, $a_{\gamma-1} = -2$ and $a_{\gamma} = a_{\gamma+1} = 0$. It follows that there is some $v(x)$ that solves the congruence

$$Z^2 + (x^3 + 4)Z - [x^{2\gamma+1} + x^{2\gamma-2} - (3x^3 + 4)x^{\gamma-1}] \equiv 0 \pmod{x^{\gamma+2}}$$

and for which $\deg v(x) = \gamma - 1$. So $m(p - q)$ is the second profile in Ex. 5.8.

Remark B.3. Technically speaking, one should check that the discriminant of $(x^3 + 4) - 4[(3x^3 + 4)^2 x^{2\gamma-2} - (x^{2\gamma+1} + x^{2\gamma-2})]$ is non-zero in order to guarantee the smoothness of the affine curve. However, as the curve B is non-singular at every point involved in the putative linear equivalence relations, it suffices to replace B by its normalization, if necessary, to certify the existence of a torsion element with the desired multiplication profile.¹⁴

¹³This is where we require that $\gamma \geq 4$.

¹⁴More explicitly, let $F(x, y) = y^2 + (x^3 + 4)y + (3x^3 + 4)x^{\gamma-1} - x^{2\gamma+1} - x^{2\gamma-2}$. We have $\frac{\partial F}{\partial y}(0, 0) = 4$, so the curve cut out by $F(x, y)$ is non-singular at p . The associated principal divisor of $-x^{\gamma-1} - y$ yields the linear equivalence relation $(\gamma + 2)(p - q) \sim \sum_{i=1}^{\gamma-1} (r_i - q)$, where $r_i = (\xi_i, -\xi_i^3 - 2)$ and the ξ_i run through all roots of $x^{\gamma-1} - 2 = 0$. It is easy to see $\frac{\partial F}{\partial y}(r_i) = -\xi_i^3 \neq 0$.

REFERENCES

- [1] B. Bekker and Y. Zarhin, *Torsion points of order $2g + 1$ on odd degree hyperelliptic curves of genus g* , Trans. Amer. Math. Soc. **373** (2020), no. 11, 8059–8094.
- [2] J. Boxall, D. Grant, and F. Leprévost, *5-torsion points on curves of genus 2*, J. London Math. Soc. **64** (2001), no. 1, 29–43.
- [3] T. Bridgeland, A. King, and M. Reid, *The McKay correspondence as an equivalence of derived categories*, J. Amer. Math. Soc. **14** (2001), no. 3, 535–554.
- [4] D. Cantor, *Computing in the Jacobian of a hyperelliptic curve*, Math. Comp. **48** (1987), 95–101.
- [5] C. Carvalho and F. Torres, *On numerical semigroups related to covering of curves*, Semigroup Forum **67** (2003), no. 3, 344–354.
- [6] V. Dragović and M. Radnović, *Poncelet porisms and beyond*, Birkhäuser, 2011.
- [7] D. Eisenbud, *Linear sections of determinantal varieties*, Amer. J. Math. **110** (1988), no. 3, 541–575.
- [8] H. Esnault and E. Viehweg, *Revêtements cycliques*, in “Algebraic threefolds”, 241–250, Springer, 1982.
- [9] E.V. Flynn, *Sequences of rational torsions on abelian varieties*, Invent. Math. **106** (1991), no. 1, 433–442.
- [10] I.M. Gelfand, M. Kapranov, and A. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Springer, 2008.
- [11] P. Griffiths and J. Harris, *On Cayley’s explicit solution to Poncelet’s porism*, Enseign. Math. **24** (1978), no. 1-2, 31–40.
- [12] R. Hartshorne, *Algebraic geometry*, Springer, 1977.
- [13] I. Morrison and H. Pinkham, *Galois Weierstrass points and Hurwitz characters*, Annals Math. **124** (1986), no. 3, 591–625.
- [14] D. Mumford, *Tata lectures on Theta II, Progress in Math.* **43**, 1984.
- [15] R. Pardini, *Abelian covers of algebraic varieties*, J. Reine Angew. Math. **417** (1991), 191–214.
- [16] R. Pardini, *On the period map for abelian covers of projective varieties*, Ann. Sc. Norm. Sup. Pisa **26** (1998), no. 4, 719–735.
- [17] J.-P. Serre, *Distribution asymptotique des valeurs propres des endomorphismes de Frobenius (d’après Abel, Chebyshev, Robinson, . . .)*, Bourbaki **70** (2018), no. 1146, 1–32.
- [18] U. Zannier, *Hyperelliptic continued fractions and generalized Jacobians*, to appear in Amer. J. Math.
- [19] Y. Zarhin, *Division by 2 on odd-degree hyperelliptic curves and their Jacobians*, Izv. Math. **83** (2019), no. 3, 501–520.

(Ethan Cotterill) INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL FLUMINENSE, RUA PROF WALDEMAR DE FREITAS, S/N, CAMPUS DO GRAGOATÁ, CEP 24.210-201, NITERÓI, RJ, BRAZIL

Email address: cotterill.ethan@gmail.com

(Nathan Pflueger) DEPARTMENT OF MATHEMATICS AND STATISTICS, AMHERST COLLEGE, AMHERST, MA 01002

Email address: npflueger@amherst.edu

(Naizhen Zhang) INSTITUT FÜR DIFFERENTIALGEOMETRIE, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER, GERMANY

Email address: naizhen.zhang@math.uni-hannover.de