arXiv:2201.00085v1 [quant-ph] 1 Jan 2022

Strong quantum nonlocality for unextendible product bases in heterogeneous systems

Fei Shi,^{1,*} Mao-Sheng Li,^{2,3,†} Lin Chen,^{4,5,‡} and Xiande Zhang^{6,§}

¹School of Cyber Security, University of Science and Technology of China, Hefei, 230026, People's Republic of China

²Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China

³Department of Physics, University of Science and Technology of China, Hefei 230026, China

⁴LMIB(Beihang University), Ministry of Education,

and School of Mathematical Sciences, Beihang University, Beijing 100191, China

⁵International Research Institute for Multidisciplinary Science, Beihang University, Beijing 100191, China

⁶School of Mathematical Sciences, University of Science and Technology of China, Hefei, 230026, People's Republic of China

A set of multipartite orthogonal product states is strongly nonlocal if it is locally irreducible in every bipartition, which shows the phenomenon of strong quantum nonlocality without entanglement. It is known that unextendible product bases (UPBs) can show the phenomenon of quantum nonlocality without entanglement. Thus it is interesting to investigate the strong quantum nonlocality for UPBs. Most of the UPBs with the minimum size cannot demonstrate strong quantum nonlocality. In this paper, we construct a series of UPBs with different large sizes in $d_A \otimes d_B \otimes d_C$ and $d_A \otimes d_B \otimes d_C \otimes d_D$ for $d_A, d_B, d_C, d_D \ge 3$, and we also show that these UPBs have strong quantum nonlocality, which answers an open question given by Halder *et al.* [Phys. Rev. Lett. **122**, 040403 (2019)] and Yuan *et al.* [Phys. Rev. A **102**, 042228 (2020)] for any possible three and four-partite systems. Furthermore, we propose an entanglement-assisted protocol to locally discriminate the UPB in $3 \otimes 3 \otimes 4$, and it consumes less entanglement resource than the teleportation-based protocol. Our results build the connection between strong quantum nonlocality and UPBs.

Keywords: strong quantum nonlocality, unextendible product bases, entanglement-assisted discrimination

I. INTRODUCTION

It is known that a set of nonorthogonal states cannot be perfectly distinguished, and a set of mutually orthogonal states can be always distinguished by performing a global measurement [1]. However, when the composite quantum system is distributed among several spatially separated parties, it is not always possible to distinguish the states by performing local operations and classical communications (LOCC). These states are said to be locally indistinguishable. The local indistinguishability plays an important role in quantum data hiding $\left[2-5\right]$ and quantum secret sharing [6-8]. Bennett *et al.* first showed the phenomenon of quantum nonlocality without entanglement by constructing a locally indistinguishable orthogonal product basis in a two qutrit system [9]. Consequently, the nonlocality here (or the local distinguishability based nonlocality that we call) is very different from the most well-known form of quantum nonlocality also known as Bell nonlocality [10, 11] which can only arise from entangled states. This leads us to explore this kind of nonlocality. After that, locally indistinguishable sets have been widely investigated [12–30]. Unextendible product bases (UPBs) are special kinds of locally indistinguishable sets [31, 32]. A UPB for a multipartite quantum system is an incomplete orthogonal product basis whose complementary subspace contains no product state. The UPBs can be used to construct positive-partial-transpose (PPT) entangled states and Peres sets [31, 33]. It is also connected to bound entangled states, fermionic systems, Bell inequalities without quantum violation [12, 31, 34– 37]. Most of the constructions for UPBs are about UPBs with the minimum size [9, 12, 38–41]. It is interesting to investigate UPBs with large sizes in multipartite systems.

Recently, Halder *et al.* proposed a strong form of nonlocality based on the concept of local irreducibility of quantum states [42]. An orthogonal product set (OPS) is locally irreducible means that it is not possible to eliminate one or more states from the set by orthogonality-preserving local measurements. Under this definition, a locally irreducible set must be a locally indistinguishable set, while the converse is not true in general. Therefore, constructing locally irreducible set of orthogonal quantum states is an efficient way to show the local distinguishability based nonlocality. An OPS is said to be *strongly nonlocal* if it is locally irreducible in every bipartition. For further study this kind of nonlocality, it is interesting to investigate the locally irreducibility and the strong quantum nonlocality for OPSs. Halder *et al.* constructed two strongly nonlocal OPSs in $3 \otimes 3 \otimes 3$ and $4 \otimes 4 \otimes 4$ respectively, which shows the phenomenon of strong quantum nonlocality without entanglement [42]. In addition, there were several results about

 $^{\dagger} li.mao sheng.math@gmail.com$

^{*}shifei@mail.ustc.edu.cn

[‡]linchen@buaa.edu.cn

[§]Corresponding author: drzhangx@ustc.edu.cn

strongly nonlocal OPSs. A strongly nonlocal OPS in $d \otimes d \otimes d$, $d \otimes d \otimes (d+1)$, $3 \otimes 3 \otimes 3 \otimes 3$ and $4 \otimes 4 \otimes 4 \otimes 4$ for $d \geq 3$ was given in [43]. The authors in [44] constructed a strongly nonlocal OPS in $d_A \otimes d_B \otimes d_C$, $d_A \otimes d_B \otimes d_C \otimes d_D$, and $d_A \otimes d_B \otimes d_C \otimes d_D \otimes d_E$ for $d_A, d_B, d_C, d_D, d_E \geq 3$. Both of Refs. [42, 43] propose an open question: whether one can find strongly nonlocal UPBs? Specially, Ref. [43] indicates that most of the previous UPBs with the minimum size cannot be used for building strongly nonlocal UPBs. In [45], the authors show that a strongly nonlocal UPB with large size in $d \otimes d \otimes d$ exists for $d \geq 3$, where the UPB was constructed from [46]. However, we still do not know whether there exist strongly nonlocal UPBs in multipartite systems with non-equal local dimensions. So it is interesting to consider strongly nonlocal UPBs in general systems, like any possible three and four-partite systems. Further, some strongly nonlocal orthogonal entangled sets were shown in [47, 48]

When a set of orthogonal states is not locally distinguishable, entanglement can be used as a resource for distinguishability of such states. This is called the entanglement-assisted discrimination, which was first proposed by Cohen. In [49], Cohen showed that the tile UPB in $3 \otimes 3$ can be perfectly distinguished by using a two-qubit maximally entangled state. Since then, entanglement-assisted discrimination has attracted a lot of interest [47, 49–57]. Since a strongly nonlocal UPB is locally indistinguishable in any bipartition, a perfect discrimination of this set needs a resource state that must be entangled in all bipartitions. In case of the teleportation-based protocol [58], any set of orthogonal states in $m \otimes n$ ($m \leq n$) can be perfectly distinguished by using an $m \otimes m$ maximally entangled state. Then the teleportation-based protocol can perfectly distinguish the strongly nonlocal UPB in $3 \otimes 3 \otimes 4$ by using $3 \otimes 3$ maximally entangled states shared between any two pairs. Since entanglement is a costly resource in quantum information, it is important to find a protocol using cheaper resources.

In this paper, we focus on the construction of strongly nonlocal UPBs in any possible three and four-partite systems. The construction of strongly nonlocal UPBs is more difficult than the construction of strongly nonlocal OPSs, since it is not easy to check that an OPS is a UPB usually. Thus, new method is required. By using the relation between OPSs and grid representations, we successfully construct a series of UPBs with different large sizes in three and four-partite systems. That is, for any $0 \le s \le \lfloor \frac{d_A-3}{2} \rfloor$, there exists a UPB of size $d_A d_B d_C - 8(s+1)$ in $d_A \otimes d_B \otimes d_C$ and a UPB of size $d_A d_B d_C d_D - 16(s+1)$ in $d_A \otimes d_B \otimes d_C \otimes d_D$ for $3 \le d_A \le d_B \le d_C \le d_D$. We also show that these UPBs are strongly nonlocal by using the techniques from [45]. Further, we propose an entanglement-assisted protocol for local discrimination of the strongly nonlocal UPB in $3 \otimes 3 \otimes 4$, which consumes less entanglement resource than the teleportation-based protocol.

The rest of this paper is organized as follows. In Sec. II, we introduce preliminary knowledge. In Sec. III, we construct tripartite UPBs and show that these UPBs are strongly nonlocal. In Theorem 3 of supplementary material [59], we show a series of strongly nonlocal UPBs in four-partite systems. In Sec. IV, we consider the entanglement-assisted discrimination for the strongly nonlocal UPB in $3 \otimes 3 \otimes 4$. Finally, we conclude in Sec. V.

II. PRELIMINARY

In this section, we introduce the preliminary knowledge and facts. Throughout this paper, we do not normalize states and operators for simplicity, and we consider only pure states and positive operator-valued measure (POVM) measurements. For any positive integer $n \ge 1$, we denote \mathbb{Z}_n as the set $\{0, 1, \dots, n-1\}$. Assume that $\{|i\rangle\}_{i\in\mathbb{Z}_m}$ is the computational basis of an *m*-dimensional Hilbert space. A bipartite state $|\psi\rangle$ in $m \otimes n$ can be expressed by

$$|\psi\rangle = \sum_{i \in \mathbb{Z}_m} \sum_{j \in \mathbb{Z}_n} a_{i,j} |i\rangle_A |j\rangle_B.$$
(1)

Then $|\psi\rangle$ corresponds to an $m \times n$ matrix,

$$M = (a_{i,j})_{i \in \mathbb{Z}_m, j \in \mathbb{Z}_n}.$$
(2)

Note that $|\psi\rangle$ is a product state if and only if rank(M) = 1. Assume that $|\psi_i\rangle$ in $m \otimes n$ corresponds to an $m \times n$ matrix M_i for i = 1, 2, then the inner product

$$\langle \psi_1 | \psi_2 \rangle = \operatorname{Tr}(M_1^{\dagger} M_2). \tag{3}$$

An unextendible product basis (UPB) for a multipartite quantum system is an incomplete orthogonal product basis whose complementary subspace contains no product state. For example, the SHIFTS UPB in $2 \otimes 2 \otimes 2$ is as follows [31],

$$\begin{split} \psi_0 \rangle &= |0\rangle_A |1\rangle_B |+\rangle_C, \quad |\psi_1\rangle &= |1\rangle_A |+\rangle_B |0\rangle_C, \\ \psi_2 \rangle &= |+\rangle_A |0\rangle_B |1\rangle_C, \quad |\psi_3\rangle &= |-\rangle_A |-\rangle_B |-\rangle_C, \end{split}$$

where $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$.

A local measurement performed to distinguish a set of multipartite mutually orthogonal states is called an *orthogonality-preserving local measurement*, if the postmeasurement states keep being mutually orthogonal. Specially, a measurement is *trivial* if all the POVM elements are proportional to the identity operator. In [42], the authors proposed the concept of strong quantum nonlocality. An orthogonal product set (OPS) is said to be *strongly nonlocal* if it is locally irreducible in every bipartition. Note that an OPS is a *locally irreducible set* means that it is not possible to eliminate one or more states from the set by orthogonality-preserving local measurements. There exists an efficient way to check whether an OPS is strongly nonlocal [45]. Assume that an OPS $\{|\psi\rangle\} \subset \bigotimes_{i=1}^{n} \mathcal{H}_{A_i}$. Let $B_1 = \{A_2A_3 \ldots A_n\}, B_2 = \{A_3 \ldots A_nA_1\}, B_3 = \{A_4 \ldots A_nA_1A_2\}, \ldots, B_n = \{A_1 \ldots A_{n-1}\}$. If the party B_i can only perform a trivial orthogonality-preserving POVM for any $1 \leq i \leq n$, then the OPS $\{|\psi\rangle\}$ is strongly nonlocal.

In this paper, we show a series of strongly nonlocal UPBs in $d_A \otimes d_B \otimes d_C$ and $d_A \otimes d_B \otimes d_C \otimes d_D$ for any $d_A, d_B, d_C, d_D \geq 3$ respectively. Without loss of generality, we always assume that $3 \leq d_A \leq d_B \leq d_C \leq d_D$. In [44], the authors gave a decomposition for the outermost layer of 3,4-dimensional hypercubes, and our construction of UPBs in this paper is inspired by this decomposition. Since any OPS in $2 \otimes n$ is locally reducible [12, 33], a strongly nonlocal UPB in $d_A \otimes d_B \otimes d_C$ and $d_A \otimes d_B \otimes d_C \otimes d_D$ must satisfy $d_A, d_B, d_C, d_D \geq 3$. Our construction achieves the minimum quantum system necessary for the existence of such a UPB.

Next, we introduce two basic lemmas from [45], which are useful for showing strong quantum nonlocality. Let \mathcal{H}_n be an *n*-dimensional Hilbert space. Assume that the computational basis of \mathcal{H}_n is $\{|0\rangle, |1\rangle, \dots, |n-1\rangle\}$. For any operator E on \mathcal{H}_n , we denote the matrix E as the matrix representation of the operator E under the computational basis. In general, we do not distinguish the operator E and the matrix E. Given any $n \times n$ matrix $E := \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} a_{i,j} |i\rangle\langle j|$, for $\mathcal{S}, \mathcal{T} \subseteq \{|0\rangle, |1\rangle, \dots, |n-1\rangle\}$, we define

$${}_{\mathcal{S}}E_{\mathcal{T}} := \sum_{|s\rangle \in \mathcal{S}} \sum_{|t\rangle \in \mathcal{T}} a_{s,t} |s\rangle \langle t|.$$

It means that ${}_{\mathcal{S}}E_{\mathcal{T}}$ is a submatrix of E with row coordinates \mathcal{S} and column coordinates \mathcal{T} . In the case $\mathcal{S} = \mathcal{T}$, we denote $E_{\mathcal{S}} := {}_{\mathcal{S}}E_{\mathcal{S}}$ for simplicity. Given a set $\mathcal{S} \subseteq \{|0\rangle, |1\rangle, \cdots, |n-1\rangle\}$, an orthogonal set $\{|\psi_i\rangle\}_{i\in\mathbb{Z}_s}$ is spanned by \mathcal{S} , if for any $i \in \mathbb{Z}_s, |\psi_i\rangle$ is a linear combination of the states from \mathcal{S} .

Lemma 1 (Block Zeros Lemma [45]) Let an $n \times n$ matrix $E = (a_{i,j})_{i,j \in \mathbb{Z}_n}$ be the matrix representation of an operator $E = M^{\dagger}M$ under the basis $\mathcal{B} = \{|0\rangle, |1\rangle, \ldots, |n-1\rangle\}$. Given two nonempty disjoint subsets \mathcal{S} and \mathcal{T} of \mathcal{B} , assume that $\{|\psi_i\rangle\}_{i\in\mathbb{Z}_s}, \{|\phi_j\rangle\}_{j\in\mathbb{Z}_t}$ are two orthogonal sets spanned by \mathcal{S} and \mathcal{T} respectively, where $s = |\mathcal{S}|$, and $t = |\mathcal{T}|$. If $\langle \psi_i | E | \phi_i \rangle = 0$ for any $i \in \mathbb{Z}_s$ and $j \in \mathbb{Z}_t$, then ${}_{\mathcal{S}}E_{\mathcal{T}} = \mathbf{0}$ and ${}_{\mathcal{T}}E_{\mathcal{S}} = \mathbf{0}$.

Lemma 2 (Block Trivial Lemma [45]) Let an $n \times n$ matrix $E = (a_{i,j})_{i,j \in \mathbb{Z}_n}$ be the matrix representation of an operator $E = M^{\dagger}M$ under the basis $\mathcal{B} = \{|0\rangle, |1\rangle, \ldots, |n-1\rangle\}$. Given a nonempty subset $\mathcal{S} := \{|u_0\rangle, |u_1\rangle, \ldots, |u_{s-1}\rangle\}$ of \mathcal{B} , let $\{|\psi_j\rangle\}_{j \in \mathbb{Z}_s}$ be an orthogonal set spanned by \mathcal{S} . Assume that $\langle \psi_i | E | \psi_j \rangle = 0$ for any $i \neq j \in \mathbb{Z}_s$. If there exists a state $|u_t\rangle \in \mathcal{S}$, such that $\{|u_t\rangle\} E_{\mathcal{S} \setminus \{|u_t\rangle\}} = \mathbf{0}$ and $\langle u_t | \psi_j \rangle \neq 0$ for any $j \in \mathbb{Z}_s$, then $E_{\mathcal{S}} \propto \mathbb{I}_{\mathcal{S}}$.

III. STRONGLY NONLOCAL TRIPARTITE UPBS

In this section, we construct strongly nonlocal tripartite UPBs. In Example 3, we show a UPB in $3 \otimes 3 \otimes 4$. Then we generalize this UPB to the space $d_A \otimes d_B \otimes d_C$ in Proposition 4, and prove its strong quantum nonlocality in Proposition 5. In fact, we show a series of strongly nonlocal UPBs of different sizes in $d_A \otimes d_B \otimes d_C$ in Theorem 6. Let $w_n = e^{\frac{2\pi\sqrt{-1}}{n}}$ be the *n*-th unit root, and let sum(*M*) be the sum of all entries of the matrix *M*.

First, we consider a simple example in $3 \otimes 3 \otimes 4$. Let

$$\begin{aligned} \mathcal{A}_{1} &:= \{ |\psi_{1}(i,k)\rangle = |\xi_{i}\rangle_{A}|0\rangle_{B}|\eta_{k}\rangle_{C} : (i,k) \in \mathbb{Z}_{2} \times \mathbb{Z}_{3} \setminus \{(0,0)\} \}, \\ \mathcal{A}_{2} &:= \{ |\psi_{2}(i,j)\rangle = |\xi_{i}\rangle_{A}|\eta_{j}\rangle_{B}|3\rangle_{C} : (i,j) \in \mathbb{Z}_{2} \times \mathbb{Z}_{2} \setminus \{(0,0)\} \}, \\ \mathcal{A}_{3} &:= \{ |\psi_{3}(j,k)\rangle = |2\rangle_{A}|\xi_{j}\rangle_{B}|\eta_{k}\rangle_{C} : (j,k) \in \mathbb{Z}_{2} \times \mathbb{Z}_{3} \setminus \{(0,0)\} \}, \\ \mathcal{A}_{4} &:= \{ |\psi_{4}\rangle = |2\rangle_{A}|2\rangle_{B}|3\rangle_{C} \}, \\ \mathcal{B}_{1} &:= \{ |\phi_{1}(i,k)\rangle = |\eta_{i}\rangle_{A}|2\rangle_{B}|\xi_{k}\rangle_{C} : (i,k) \in \mathbb{Z}_{2} \times \mathbb{Z}_{3} \setminus \{(0,0)\} \}, \\ \mathcal{B}_{2} &:= \{ |\phi_{2}(i,j)\rangle = |\eta_{i}\rangle_{A}|\xi_{j}\rangle_{B}|0\rangle_{C} : (i,j) \in \mathbb{Z}_{2} \times \mathbb{Z}_{2} \setminus \{(0,0)\} \}, \\ \mathcal{B}_{3} &:= \{ |\phi_{3}(j,k)\rangle = |0\rangle_{A}|\eta_{j}\rangle_{B}|\xi_{k}\rangle_{C} : (j,k) \in \mathbb{Z}_{2} \times \mathbb{Z}_{3} \setminus \{(0,0)\} \}, \\ \mathcal{B}_{4} &:= \{ |\phi_{4}\rangle = |0\rangle_{A}|0\rangle_{B}|0\rangle_{C} \}, \\ \mathcal{F} &:= \{ |\varphi(k)\rangle = |1\rangle_{A}|1\rangle_{B}|\beta_{k}\rangle_{C} : k \in \mathbb{Z}_{2} \setminus \{0\} \}, \end{aligned}$$

$$|S\rangle := \left(\sum_{i=0}^{2} |i\rangle\right)_{A} \left(\sum_{j=0}^{2} |j\rangle\right)_{B} \left(\sum_{k=0}^{3} |k\rangle\right)_{C},$$

where $|\eta_s\rangle_X = |0\rangle_X + (-1)^s |1\rangle_X$, $|\xi_s\rangle_X = |1\rangle_X + (-1)^s |2\rangle_X$ for $s \in \mathbb{Z}_2$, $X \in \{A, B\}$, $|\eta_s\rangle_C = \sum_{t=0}^2 w_3^{st} |t\rangle_C$, $|\xi_s\rangle_C = \sum_{t=0}^3 |t\rangle_C$

 $\sum_{t=0}^{2} w_{3}^{st} |t+1\rangle_{C} \text{ for } s \in \mathbb{Z}_{3} \text{ and } |\beta_{s}\rangle_{C} = |1\rangle_{C} + (-1)^{s}|2\rangle_{C} \text{ for } s \in \mathbb{Z}_{2}.$ The state $|S\rangle$ is called a "stopper" state. It is easy to see that $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$ is an OPS, and $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$ is an OPS, and $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$ is an OPS, and $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$ is an OPS, and $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$ is a OPS, and $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{i_{i=1}^{3} \{|\psi_{i}(0,0)\rangle, |\phi_{i}(0,0)\rangle\} \cup \{|\psi_{4}\rangle\} \cup \{|\phi_{4}\rangle\} \cup \{|\varphi(0)\rangle\}$ is a complete orthogonal basis in $\mathbb{C}^{3} \otimes \mathbb{C}^{3} \otimes \mathbb{C}^{4}$. The nine subsets $\mathcal{A}_i, \mathcal{B}_i$ (i = 1, 2, 3, 4) and \mathcal{F} in A|BC bipartition correspond to the nine blocks of 3×12 grid in Fig. 1. For example, \mathcal{A}_1 corresponds to the 2 × 3 grid {(1,2) × (00,01,02)}. Moreover, \mathcal{A}_i is symmetrical to \mathcal{B}_i for $1 \le i \le 4$. If we delete $\mathcal{A}_4, \mathcal{B}_4$, and add the "stopper" state $|S\rangle$, we can obtain a UPB in $3 \otimes 3 \otimes 4$.



FIG. 1: The corresponding 3×12 grid of $\mathcal{A}_i, \mathcal{B}_i$ (i = 1, 2, 3, 4) and \mathcal{F} (Eq. (4)) in A|BC bipartition. For example, \mathcal{A}_1 corresponds to the 2 × 3 grid $\{(1,2) \times (00,01,02)\}$. Moreover, \mathcal{A}_i is symmetrical to \mathcal{B}_i for $1 \leq i \leq 4$.

Example 3 In $3 \otimes 3 \otimes 4$, the set of states $\bigcup_{i=1}^{3} (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F} \cup \{|S\}\}$ given by Eq. (4) is a UPB of size 28.

Let \mathcal{H} be the space spanned by the states in $\bigcup_{i=1}^{3} (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F} \cup \{|S\rangle\}$. For any state $|\psi\rangle \in \mathcal{H}^{\perp}$, we only Proof. need to show that $|\psi\rangle$ must be an entangled state. We prove it by contradiction. Assume there exists a product state $|\psi\rangle \in \mathcal{H}^{\perp}$. Let \mathcal{H}_1 be the space spanned by the states in $\bigcup_{i=1}^3 (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F}$. Since $\mathcal{H}_1 \subset \mathcal{H}$, we have $\mathcal{H}^{\perp} \subset \mathcal{H}_1^{\perp}$. Moreover,

$$\mathcal{H}_{1}^{1} = \operatorname{span}\{|\psi_{1}(0,0)\rangle, |\psi_{2}(0,0)\rangle, |\psi_{3}(0,0)\rangle, |\psi_{4}\rangle, |\phi_{1}(0,0)\rangle, |\phi_{2}(0,0)\rangle, |\phi_{3}(0,0)\rangle, |\phi_{4}\rangle, |\varphi(0)\rangle\}.$$

Then $|\psi\rangle$ can be expressed by

$$|\psi\rangle = a_0|\psi_1(0,0)\rangle + b_0|\psi_2(0,0)\rangle + c_0|\psi_3(0,0)\rangle + d_0|\psi_4\rangle + a_1|\phi_1(0,0)\rangle + b_1|\phi_2(0,0)\rangle + c_1|\phi_3(0,0)\rangle + d_1|\phi_4\rangle + e|\varphi(0)\rangle,$$

where $a_0, b_0, c_0, d_0, a_1, b_1, c_1, d_1, e \in \mathbb{C}$. By assumption, $|\psi\rangle$ is a product state, and $\langle S|\psi\rangle = 0$.

Next, we consider the matrix form of $|\psi\rangle$ in A|BC bipartition. It corresponds to the 3×12 matrix M in A|BC bipartition, where

$$M = \begin{bmatrix} a_0 & a_0 & a_0 & b_0 & b_0 & c_0 & c_0 & c_0 & c_0 & c_0 & d_0 \\ a_0 & a_0 & a_0 & b_0 & b_0 & e & e & b_1 & b_1 & a_1 & a_1 \\ d_1 & c_1 & c_1 & c_1 & c_1 & c_1 & b_1 & b_1 & a_1 & a_1 \\ 0 & 00 & 01 & 02 & 03 & 13 & 11 & 12 & 10 & 20 & 21 & 22 & 23 \\ & & & BC & & & & \\ \end{bmatrix}$$

and

$$\operatorname{rank}(M) = 1. \tag{5}$$

Note that M has a similar structure as Fig. 1. It means that x_0 is symmetrical to x_1 for $x \in \{a, b, c, d\}$. The state $|S\rangle$ corresponds to the matrix S in A|BC bipartition, where

Every element of M has coordinate (A, BC). For example, d_0 has coordinate (2, 23). If we consider AB|C bipartition, then we can rearrange the first row of M to a 3×4 matrix, denoted by M_2 through (AB, C) coordinates. For example, d_0 has coordinate (22,3) in M_2 . Similarly, we can rearrange the last row of M to a 3×4 matrix, denoted by M_0 through (AB, C) coordinates. Thus we obtain M_2 and M_0 , where

$$M_{2} = \begin{bmatrix} c_{0} & c_{0} & c_{0} & d_{0} \\ c_{0} & c_{0} & c_{0} & b_{0} \\ a_{0} & a_{0} & a_{0} & b_{0} \end{bmatrix} \begin{bmatrix} 22 \\ 21 \\ AB \\ 20 \end{bmatrix}, \qquad M_{0} = \begin{bmatrix} b_{1} & a_{1} & a_{1} \\ b_{1} & c_{1} & c_{1} & c_{1} \\ d_{1} & c_{1} & c_{1} & c_{1} \end{bmatrix} \begin{bmatrix} 02 \\ 01 \\ AB \\ 00 \end{bmatrix} AB.$$

Since $|\psi\rangle$ is separable in AB|C bipartition, we must have

$$\operatorname{rank} M_2 = 0 \quad \text{or} \quad 1, \tag{7}$$

$$\operatorname{rank} M_0 = 0 \quad \text{or} \quad 1. \tag{8}$$

Assume $a_0 \neq 0$. Since rank(M) = 1, we have $c_1 = d_1$, and $c_0 = e = b_1 = a_1 = d_0$.

- (i) If $c_0 = 0$, then $b_0 = 0$ by Eq. (7), and $c_1 = 0$ by Eq. (8), which is impossible by Eq. (6).
- (ii) If $c_0 \neq 0$, then $c_0 = a_0 = b_0$ by Eq. (7), and $c_0 = c_1$ by Eq. (8), which is impossible by Eq. (6).
- Thus we have $a_0 = 0$. By the symmetry of M, we must have $a_1 = 0$.

Assume $b_0 \neq 0$. Since rank(M) = 1, we have $c_1 = d_1 = 0$, and $c_0 = e = b_1 = a_1 = d_0 = 0$. This is impossible by Eq. (6). So we must have $b_0 = b_1 = 0$.

Assume $d_1 \neq 0$. Since rank(M) = 1, we have $e = c_0 = d_0 = 0$. By Eq. (8), we obtain $c_1 = 0$. This is impossible by Eq. (6). So we must have $d_0 = d_1 = 0$.

Assume $c_1 \neq 0$. Since rank(M) = 1, we have $e = c_0 = 0$. This is impossible by Eq. (6). So we have $c_0 = c_1 = 0$. Since sum(M) = 0, we must have e = 0, which contradicts rank(M) = 1.

Thus $|\psi\rangle$ must be an entangled state, and the set of states $\bigcup_{i=1}^{3} (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F} \cup \{|S\rangle\}$ is a UPB.

Next, we generalize the above UPB to the space $d_A \otimes d_B \otimes d_C$. Let

$$\begin{aligned}
\mathcal{A}_{1} &:= \{|\xi_{i}\rangle_{A}|0\rangle_{B}|\eta_{k}\rangle_{C}: (i,k) \in \mathbb{Z}_{d_{A}-1} \times \mathbb{Z}_{d_{C}-1} \setminus \{(0,0)\}\}, \\
\mathcal{A}_{2} &:= \{|\xi_{i}\rangle_{A}|\eta_{j}\rangle_{B}|d_{C}-1\rangle_{C}: (i,j) \in \mathbb{Z}_{d_{A}-1} \times \mathbb{Z}_{d_{B}-1} \setminus \{(0,0)\}\}, \\
\mathcal{A}_{3} &:= \{|d_{A}-1\rangle_{A}|\xi_{j}\rangle_{B}|\eta_{k}\rangle_{C}: (j,k) \in \mathbb{Z}_{d_{B}-1} \times \mathbb{Z}_{d_{C}-1} \setminus \{(0,0)\}\}, \\
\mathcal{A}_{4} &:= \{|d_{A}-1\rangle_{A}|d_{B}-1\rangle_{B}|d_{C}-1\rangle_{C}\}, \\
\mathcal{B}_{1} &:= \{|\eta_{i}\rangle_{A}|d_{B}-1\rangle_{B}|\xi_{k}\rangle_{C}: (i,k) \in \mathbb{Z}_{d_{A}-1} \times \mathbb{Z}_{d_{C}-1} \setminus \{(0,0)\}\}, \\
\mathcal{B}_{2} &:= \{|\eta_{i}\rangle_{A}|\xi_{j}\rangle_{B}|0\rangle_{C}: (i,j) \in \mathbb{Z}_{d_{A}-1} \times \mathbb{Z}_{d_{B}-1} \setminus \{(0,0)\}\}, \\
\mathcal{B}_{3} &:= \{|0\rangle_{A}|\eta_{j}\rangle_{B}|\xi_{k}\rangle_{C}: (j,k) \in \mathbb{Z}_{d_{B}-1} \times \mathbb{Z}_{d_{C}-1} \setminus \{(0,0)\}\}, \\
\mathcal{B}_{4} &:= \{|0\rangle_{A}|0\rangle_{B}|0\rangle_{C}\}, \\
\mathcal{F} &:= \{|\beta_{i}\rangle_{A}|\beta_{j}\rangle_{B}|\beta_{k}\rangle_{C}: (i,j,k) \in \mathbb{Z}_{d_{A}-2} \times \mathbb{Z}_{d_{B}-2} \times \mathbb{Z}_{d_{C}-2} \setminus \{(0,0,0)\}\}, \\
|S\rangle &:= \left(\sum_{i=0}^{d_{A}-1}|i\rangle\right)_{A} \left(\sum_{j=0}^{d_{B}-1}|j\rangle\right)_{B} \left(\sum_{k=0}^{d_{C}-1}|k\rangle\right)_{C},
\end{aligned}$$

where $|\eta_s\rangle_X = \sum_{t=0}^{d_X-2} w_{d_X-1}^{st} |t\rangle_X$, and $|\xi_s\rangle_X = \sum_{t=0}^{d_X-2} w_{d_X-1}^{st} |t+1\rangle_X$ for $s \in \mathbb{Z}_{d_X-1}$, and $X \in \{A, B, C\}, |\beta_s\rangle_X = \sum_{t=0}^{d_X-2} w_{d_X-1}^{st} |t+1\rangle_X$

 $\sum_{t=0}^{d_X-3} w_{d_X-2}^{st} |t+1\rangle_X \text{ for } s \in \mathbb{Z}_{d_X-2}, \text{ and } X \in \{A, B, C\}.$ Note that $\{|\eta_s\rangle_X\}_{s\in\mathbb{Z}_{d_X-1}}, \{|\xi_s\rangle_X\}_{s\in\mathbb{Z}_{d_X-1}}, \text{ and } \{|\beta_s\rangle_X\}_{s\in\mathbb{Z}_{d_X-2}}$ are three orthogonal sets, $X \in \{A, B, C\}$, which are spanned by $\{|t\rangle_X\}_{t=0}^{d_X-2}, \{|t\rangle_X\}_{t=1}^{d_X-1}, \text{ and } \{|t\rangle_X\}_{t=1}^{d_X-2}$, respectively. This extends the definitions of states in $3 \otimes 3 \otimes 4$ in Eq. (4). The nine subsets $\mathcal{A}_i, \mathcal{B}_i$ (i = 1, 2, 3, 4) and \mathcal{F} in A|BC bipartition correspond to the nine blocks of $d_A \times d_B d_C$ grid in Fig. 2. Then we have the following proposition.



FIG. 2: The corresponding $d_A \times d_B d_C$ grid of $\mathcal{A}_i, \mathcal{B}_i$ (i = 1, 2, 3, 4) and \mathcal{F} (Eq.(9)) in A|BC bipartition. Moreover, \mathcal{A}_i is symmetrical to \mathcal{B}_i for $1 \le i \le 4$.

Proposition 4 In $d_A \otimes d_B \otimes d_C$, $3 \leq d_A \leq d_B \leq d_C$, the set of states $\bigcup_{i=1}^3 (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F} \cup \{|S\}\}$ given by Eq. (9) is a UPB of size $d_A d_B d_C - 8$.

The proof of Proposition 4 is given in Appendix A. Next, we consider the strong quantum nonlocality for UPBs.

Proposition 5 In $d_A \otimes d_B \otimes d_C$, $3 \leq d_A \leq d_B \leq d_C$, the UPB $\cup_{i=1}^3 (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F} \cup \{|S\}\}$ given by Eq. (9) is strongly nonlocal.

Proof. Denote $\mathcal{U} := \bigcup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$. Let B and C come together to perform a joint orthogonality-preserving POVM $\{E = M^{\dagger}M\}$, where $E = (a_{ij,k\ell})_{i,k \in \mathbb{Z}_{d_{B}}, j,\ell \in \mathbb{Z}_{d_{C}}}$. Then the postmeasurement states $\{\mathbb{I}_{A} \otimes M | \psi\rangle : |\psi\rangle \in \mathcal{U}\}$ should keep being mutually orthogonal. Assume that $|\psi_{1}\rangle_{A}|\psi_{2}\rangle_{B}|\psi_{3}\rangle_{C}, |\varphi_{1}\rangle_{A}|\varphi_{2}\rangle_{B}|\varphi_{3}\rangle_{C} \in \mathcal{U}$. Then

$${}_{A}\langle\psi_{1}|_{B}\langle\psi_{2}|_{C}\langle\psi_{3}|\mathbb{I}_{A}\otimes E|\varphi_{1}\rangle_{A}|\varphi_{2}\rangle_{B}|\varphi_{3}\rangle_{C} = \langle\psi_{1}|\varphi_{1}\rangle_{A}({}_{B}\langle\psi_{2}|_{C}\langle\psi_{3}|E|\varphi_{2}\rangle_{B}|\varphi_{3}\rangle_{C}) = 0.$$
(10)

If $\langle \psi_1 | \varphi_1 \rangle_A \neq 0$, then $B_i \langle \psi_2 | C_i \langle \psi_3 | E | \varphi_2 \rangle_B | \varphi_3 \rangle_C = 0$. By using this property, we need to show that $E \propto \mathbb{I}$. If we can show that $E \propto \mathbb{I}$, then it means that BC can only perform a trivial orthogonality-preserving POVM. Since the nine subsets $\mathcal{A}_i, \mathcal{B}_i$ $(i = 1, 2, 3, 4), \mathcal{F}$ in any bipartition of $\{A | BC, C | AB, B | CA\}$ correspond to a similar grid as Fig. 2, this implies that any of the party $\{BC, AB, CA\}$ can only perform a trivial orthogonality-preserving POVM. Then we obtain that the UPB \mathcal{U} is strongly nonlocal. More details for showing $E \propto \mathbb{I}$ are given in Appendix B.

Note that the states in \mathcal{A}_i or \mathcal{B}_i (i = 1, 2, 3, 4) in Eq. (9) are defined by the outermost layer of a $d_A \times d_B \times d_C$ cube, and the states in \mathcal{F} are just defined by all inner cells. By observing this, we can construct more strongly nonlocal UPBs in $d_A \otimes d_B \otimes d_C$ by continuing to decompose \mathcal{F} in Fig. 2 by the similar tiling method. Suppose we are on the *n*-th layer from outside to inside, $0 \le n \le \lfloor \frac{d_A - 3}{2} \rfloor$. Let $X_n := d_X - 2n$ for $X \in \{A, B, C\}$. Then we can define the following states,

$$\begin{aligned} \mathcal{A}_{1}^{(n)} &:= \{|\xi_{i}^{(n)}\rangle_{A}|n\rangle_{B}|\eta_{k}^{(n)}\rangle_{C} : (i,k) \in \mathbb{Z}_{A_{n}-1} \times \mathbb{Z}_{C_{n}-1} \setminus \{(0,0)\}\}, \\ \mathcal{A}_{2}^{(n)} &:= \{|\xi_{i}^{(n)}\rangle_{A}|\eta_{j}^{(n)}\rangle_{B}|d_{C}-1-n\rangle_{C} : (i,j) \in \mathbb{Z}_{A_{n}-1} \times \mathbb{Z}_{B_{n}-1} \setminus \{(0,0)\}\}, \\ \mathcal{A}_{3}^{(n)} &:= \{|d_{A}-1-n\rangle_{A}|\xi_{j}^{(n)}\rangle_{B}|\eta_{k}^{(n)}\rangle_{C} : (j,k) \in \mathbb{Z}_{B_{n}-1} \times \mathbb{Z}_{C_{n}-1} \setminus \{(0,0)\}\}, \\ \mathcal{A}_{4}^{(n)} &:= \{|d_{A}-1-n\rangle_{A}|d_{B}-1-n\rangle_{B}|d_{C}-1-n\rangle_{C}\}, \\ \mathcal{B}_{1}^{(n)} &:= \{|\eta_{i}^{(n)}\rangle_{A}|d_{B}-1-n\rangle_{B}|\xi_{k}^{(n)}\rangle_{C} : (i,k) \in \mathbb{Z}_{A_{n}-1} \times \mathbb{Z}_{C_{n}-1} \setminus \{(0,0)\}\}, \\ \mathcal{B}_{2}^{(n)} &:= \{|\eta_{i}^{(n)}\rangle_{A}|\xi_{j}^{(n)}\rangle_{B}|n\rangle_{C} : (i,j) \in \mathbb{Z}_{A_{n}-1} \times \mathbb{Z}_{B_{n}-1} \setminus \{(0,0)\}\}, \\ \mathcal{B}_{3}^{(n)} &:= \{|n\rangle_{A}|\eta_{j}^{(n)}\rangle_{B}|\xi_{k}^{(n)}\rangle_{C} : (j,k) \in \mathbb{Z}_{B_{n}-1} \times \mathbb{Z}_{C_{n}-1} \setminus \{(0,0)\}\}, \\ \mathcal{F}^{(n)} &:= \{|\beta_{i}^{(n)}\rangle_{A}|\beta_{j}^{(n)}\rangle_{B}|\beta_{k}^{(n)}\rangle_{C} : (i,j,k) \in \mathbb{Z}_{A_{n}-2} \times \mathbb{Z}_{B_{n}-2} \times \mathbb{Z}_{C_{n}-2} \setminus \{(0,0,0)\}\}, \\ |S\rangle &= \left(\sum_{i=0}^{d_{A}-1}|i\rangle\right)_{A} \left(\sum_{j=0}^{d_{B}-1}|j\rangle\right)_{B} \left(\sum_{k=0}^{d_{C}-1}|k\rangle\right)_{C}, \end{aligned}$$

where $|\eta_s^{(n)}\rangle_X = \sum_{t=n}^{X_n+n-2} w_{X_n-1}^{s(t-n)} |t\rangle_X$, and $|\xi_s^{(n)}\rangle_X = \sum_{t=n}^{X_n+n-2} w_{X_n-1}^{s(t-n)} |t+1\rangle_X$, for $s \in \mathbb{Z}_{X_n-1}$, and $X \in \{A, B, C\}$, $|\beta_s^{(n)}\rangle_X = \sum_{t=n}^{X_n+n-3} w_{X_n-2}^{s(t-n)} |t+1\rangle_X$ for $s \in \mathbb{Z}_{X_n-2}$, and $X \in \{A, B, C\}$.

Note that $\{|\eta_s^{(n)}\rangle_X\}_{s\in\mathbb{Z}_{X_n-1}}, \{|\xi_s^{(n)}\rangle_X\}_{s\in\mathbb{Z}_{X_n-1}}, \text{ and } \{|\beta_s^{(n)}\rangle_X\}_{s\in\mathbb{Z}_{X_n-2}}$ are three orthogonal sets, $X \in \{A, B, C\}$, which are spanned by $\{|t\rangle_X\}_{t=n}^{X_n+n-2}, \{|t\rangle_X\}_{t=n+1}^{X_n+n-1}, \text{ and } \{|t\rangle_X\}_{t=n+1}^{X_n+n-2}$, respectively. This extends the definition of states in Eq. (9) from n = 0 to general n. Specially, $\mathcal{A}_i^{(0)}, \mathcal{B}_i^{(0)}$ $(i = 1, 2, 3, 4), \mathcal{F}^{(0)}$ are $\mathcal{A}_i, \mathcal{B}_i$ $(i = 1, 2, 3, 4), \mathcal{F}$ of Eq. (9) exactly, which correspond to Fig. 2 in A|BC bipartition. Next, when $d_A \ge 5, \mathcal{A}_i^{(0)}, \mathcal{B}_i^{(0)}, \mathcal{A}_i^{(1)}, \mathcal{B}_i^{(1)}$ $(i = 1, 2, 3, 4), \mathcal{F}^{(1)}$ correspond to Fig. 3 in A|BC bipartition. Then we have the following theorem.



FIG. 3: The corresponding $d_A \times d_B d_C$ grid of $\mathcal{A}_i^{(0)}, \mathcal{B}_i^{(0)}, \mathcal{A}_i^{(1)}, \mathcal{B}_i^{(1)}$ $(i = 1, 2, 3, 4), \mathcal{F}^{(1)}$ (Eq. (11)) in A|BC bipartition, where $d_A \ge 5$.

Theorem 6 In $d_A \otimes d_B \otimes d_C$, $3 \leq d_A \leq d_B \leq d_C$, for any $0 \leq n \leq \lfloor \frac{d_A - 3}{2} \rfloor$, the set

$$\mathcal{U}_n := \bigcup_{t=0}^n (\bigcup_{i=1}^3 (\mathcal{A}_i^{(t)} \cup \mathcal{B}_i^{(t)})) \cup \mathcal{F}^{(n)} \cup \{|S\rangle\}$$

given by Eq. (11) is a strongly nonlocal UPB of size $d_A d_B d_C - 8(n+1)$.

The proof of Theorem 6 is given in Appendix C. By now, we have shown strongly nonlocal UPBs exist in any threepartite systems. For four-partite systems, strongly nonlocal UPBs are shown to exist in Supplementary material [59] by using the similar method. In [44], the authors gave a decomposition of the 5-dimensional hypercube, which may be used for constructing strongly nonlocal UPBs in any five-partite systems. However, it requires more calculations. We leave this as an open question.

IV. ENTANGLEMENT-ASSISTED DISCRIMINATION

In this section, we investigate the local discrimination of the strongly nonlocal UPB in $3 \otimes 3 \otimes 4$ in Example 3 by using entanglement as a resource. In a protocol of local quantum state discrimination, a multipatite quantum system is prepared with a state which is secretly chosen from a known set, and the purpose is to determine in which state the system is by using LOCC only. If the set of states is locally indistinguishable, then additional entanglement resources can assist for perfect discrimination. This process is called entanglement-assisted discrimination [49].



FIG. 4: Entanglement-assisted discrimination. A strongly nonlocal tripartite UPB is shared by Alice, Bob and Charlie. Their aim is to distinguish these states in the UPB by using LOCC only. For perfect discrimination of the UPB, an amount p of the maximally entangled state $|\Phi(d_1)\rangle_{A,B}$ is shared by Alice and Bob; an amount q of the maximally entangled state $|\Phi(d_2)\rangle_{A,C}$ is shared by Alice and Charlie, and an amount r of the maximally entangled state $|\Phi(d_2)\rangle_{A,C}$ is shared by Alice and Charlie, and an amount r of the maximally entangled state $|\Phi(d_2)\rangle_{B,C}$ is shared by Bob and Charlie, where $|\Phi(d)\rangle_{X,Y} = \sum_{k=0}^{d-1} |k\rangle_X |k\rangle_Y$ for $X, Y \in \{A, B, C\}$. It consumes $p \log_2 d_1 + q \log_2 d_2 + r \log_2 d_3$ ebits of entanglement resource.

Assume a strongly nonlocal tripartite UPB is shared by Alice, Bobs and Charlie. Since a strongly nonlocal UPB is locally indistinguishable in any bipartition, a perfect discrimination of this set needs a resource state that must be entangled in all bipartitions. The configuration of entanglement resources can be described by $\{(p, |\Phi(d_1)\rangle)_{A,B}, (q, |\Phi(d_2)\rangle)_{A,C}, (r, |\Phi(d_3)\rangle)_{B,C}\}$ [47, 55]. Here, $(p, |\Phi(d_1)\rangle)_{A,B}$ means that an amount p of the maximally entangled state $|\Phi(d_1)\rangle$ is shared by Alice and Bob; $(q, |\Phi(d_2)\rangle)_{A,C}$ means that an amount q of the maximally entangled state $|\Phi(d_2)\rangle$ is shared by Alice and Charlie; $(r, |\Phi(d_3)\rangle)_{B,C}$ means that an amount r of the maximally entangled state $|\Phi(d_3)\rangle$ is shared by Alice and Charlie; $(r, |\Phi(d_3)\rangle)_{B,C}$ means that an amount r of the maximally entangled state $|\Phi(d_3)\rangle$ is shared by Alice and Charlie, where $|\Phi(d)\rangle_{X,Y} = \sum_{k=0}^{d-1} |k\rangle_X |k\rangle_Y$ for $X, Y \in \{A, B, C\}$. It consumes $p \log_2 d_1 + q \log_2 d_2 + r \log_2 d_3$ ebits of entanglement resource. See Fig. 4 for this configuration of entanglement-assisted discrimination. Now, we give a discrimination protocol for the strongly nonlocal UPB given by Eq. (4).

Proposition 7 In $3 \otimes 3 \otimes 4$, the strongly nonlocal UPB $\cup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$ given by Eq. (4) can be locally distinguished by using $\{(1, |\Phi(2)\rangle)_{A,B}, (0, |\Phi(d_{2})\rangle)_{A,C}, (1, |\Phi(2)\rangle)_{B,C}\}$ for any positive d_{2} , which consumes 2 ebits of entanglement resource.

The proof of Proposition 7 is given in Appendix D. If we only use the teleportation-based protocol [55, 58], the strongly nonlocal UPB given by Eq. (4) can be locally distinguished by using $\{(1, |\Phi(3)\rangle)_{A,B}, (0, |\Phi(d_2)\rangle)_{A,C}, (1, |\Phi(3)\rangle)_{B,C}\}$, which consumes $2\log_2 3$ ebits of entanglement resource. Thus the protocol in Proposition 7 consumes less entanglement resource than the teleportation-based protocol.

V. CONCLUSION AND DISCUSSION

In this work, we have constructed a series of strongly nonlocal UPBs of different sizes in any possible three and four-partite systems. We have also proposed an entanglement-assisted protocol for local discrimination of the strongly nonlocal UPB in $3 \otimes 3 \otimes 4$. All of our UPBs of large sizes can be used to construct PPT entangled states with small rank. For example, in Proposition 4, for a normalized UPB $\{|\psi_i\rangle\}_{i=1}^{d_A d_B d_C - 8}$ in $d_A \otimes d_B \otimes d_C$ for $3 \leq d_A \leq d_B \leq d_C$, the mixed state

$$\rho = \frac{1}{8} \left(\mathbb{I} - \sum_{i=1}^{d_A d_B d_C - 8} |\psi_i\rangle \langle \psi_i | \right)$$

is a PPT entangled state with rank 8. Further our UPBs of large sizes can be also used for low-rank noisy bound entangled states which satisfy the range criterion [60]. Bipartite noisy bound entangled states that satisfy the range criterion were shown in [61]. There are some open questions left. Whether UPBs with the minimum size can show strong quantum nonlocality? Whether one can generalize our constructions on $d_1 \otimes d_2 \otimes \cdots \otimes d_n$ for $n \ge 5$?

Acknowledgments

The authors are very grateful to the reviewers for providing many useful suggestions which have greatly improved the presentation of our paper. F.S. and X.Z. were supported by the NSFC under Grants No. 11771419 and No. 12171452, the Anhui Initiative in Quantum Information Technologies under Grant No. AHY150200, and the National Key Research and Development Program of China 2020YFA0713100. M.-S.L. was supported by NSFC (Grants No. 12005092) and the China Postdoctoral Science Foundation (2020M681996). L.C. was supported by the NNSF of China (Grant No. 11871089), and the Fundamental Research Funds for the Central Universities (Grant No. ZG216S2005).

Appendix A: The proof of Proposition 4

Proof. For the same discussion as Example 3, we can assume that $|\psi\rangle$ is a product state in the complementary space of the space spanned by the states in $\bigcup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$. Then we consider the matrix form of $|\psi\rangle$ in A|BC and AB|C bipartitions. First, we consider A|BC bipartition. We have

and

$$\operatorname{rank}(M) = 1, \quad \operatorname{sum}(M) = 0. \tag{A2}$$

Then we consider AB|C bipartition. We can rearrange the first row of M to the $d_B \times d_C$ matrix $M_{(d_A-1)}$ through (AB, C) coordinates of M, and rearrange the last row of M to the $d_B \times d_C$ matrix M_0 through (AB, C) coordinates of M, where

$$M_{(d_{A}-1)} = \begin{bmatrix} c_{0} & c_{0} & \cdots & c_{0} & d_{0} \\ c_{0} & c_{0} & \cdots & c_{0} & b_{0} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ c_{0} & c_{0} & \cdots & c_{0} & b_{0} \\ a_{0} & a_{0} & \cdots & a_{0} & b_{0} \end{bmatrix}, \quad \operatorname{rank}(M_{(d_{A}-1)}) = 0 \text{ or } 1,$$
(A3)

and

$$M_{0} = \begin{bmatrix} b_{1} & a_{1} & \cdots & a_{1} & a_{1} \\ b_{1} & c_{1} & \cdots & c_{1} & c_{1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ b_{1} & c_{1} & \cdots & c_{1} & c_{1} \\ d_{1} & c_{1} & \cdots & c_{1} & c_{1} \end{bmatrix}, \quad \operatorname{rank}(M_{0}) = 0 \text{ or } 1.$$
(A4)

For the same proof as Example 3, we can show that M do not exist by Eqs. (A2), (A3) and (A4). Thus $|\psi\rangle$ must be an entangled state, and $\bigcup_{i=1}^{3} (\mathcal{A}_i \cup \mathcal{B}_i) \cup \mathcal{F} \cup \{|S\rangle\}$ is a UPB.

Appendix B: The proof of Proposition 5

Proof. First of all, we need to introduce some notations which have been introduced by [45]. Let $S = \{|\psi_1\rangle_A |\psi_2\rangle_B |\psi_3\rangle\}$ be a tripartite orthogonal product set. Define

$$\mathcal{S}(|\psi\rangle_A) := \{ |\psi_2\rangle_B |\psi_3\rangle_C : \ |\psi\rangle_A |\psi_2\rangle_B |\psi_3\rangle_C \in \mathcal{S} \}.$$

Moreover, define $\mathcal{S}^{(A)} = \{|j\rangle_B | k\rangle_C : j, k \in \mathbb{Z}_n\}$ as the support of $\mathcal{S}(|\psi\rangle_A)$ which spans $\mathcal{S}(|\psi\rangle_A)$. For example, in Eq. (4), $\mathcal{A}_1 := \{|\xi_i\rangle_A | 0\rangle_B | \eta_j\rangle_C : (i,j) \neq (0,0) \in \mathbb{Z}_2 \times \mathbb{Z}_3\}$. Then $\mathcal{A}_1(|\xi_1\rangle_A) = \{|0\rangle_A | \eta_j\rangle_C\}_{j\in\mathbb{Z}_3}$, $\mathcal{A}_1^{(A)} = \{|0\rangle_B | 0\rangle_C, |0\rangle_B | 1\rangle_C, |0\rangle_B | 2\rangle_C\}$, and $\mathcal{A}_1(|\xi_1\rangle_A)$ is spanned by $\mathcal{A}_1^{(A)}$. Actually, $\mathcal{A}_i^{(A)}, \mathcal{B}_i^{(A)}$ $(i = 1, 2, 3, 4), \mathcal{F}^{(A)}$ in Eq. (4) can be easily observed by Fig. 1. They are the projection sets of $\mathcal{A}_i, \mathcal{B}_i$ $(i = 1, 2, 3, 4), \mathcal{F}$ in *BC* party in Fig. 1. Now, we give three steps for the proof.

Step 1 Since $\langle \xi_1 | \eta_1 \rangle_A \neq 0$, applying Block Zeros Lemma to any two elements of $\{\mathcal{A}_1(|\xi_1\rangle_A), \mathcal{A}_2(|\xi_1\rangle_A), \mathcal{B}_2(|\eta_1\rangle_A), \mathcal{B}_1(|\eta_1\rangle_A)\}$, we obtain

$$\mathcal{A}_{i}^{(A)} E_{\mathcal{A}_{j}^{(A)}} = \mathbf{0}, \ \mathcal{A}_{i}^{(A)} E_{\mathcal{B}_{k}^{(A)}} = \mathbf{0}, \ \mathcal{B}_{k}^{(A)} E_{\mathcal{B}_{\ell}^{(A)}} = \mathbf{0}, \ \mathcal{B}_{k}^{(A)} E_{\mathcal{A}_{i}^{(A)}} = \mathbf{0},$$
(B1)

for $1 \leq i \neq j \leq 2, 1 \leq k \neq \ell \leq 2$. Note that if $d_A = 3$, $|\beta_i\rangle_A$ must be $|\beta_0\rangle_A$ in \mathcal{F} . So we can not apply Block Zeros Lemma to $\mathcal{F}(|\beta_1\rangle_A)$ and $\mathcal{A}_1(|\xi_1\rangle_A)$. In order to obtain $_{\mathcal{F}^{(A)}}E_{\mathcal{A}_1^{(A)}} = \mathbf{0}$, we consider $\mathcal{A}_3(|d_A - 1\rangle_A)$ and $\mathcal{A}_1(|\xi_1\rangle_A)$. Then for $(j,k) \in \mathbb{Z}_{d_B-1} \times \mathbb{Z}_{d_C-1} \setminus \{(0,0)\}$, and $i \in \mathbb{Z}_{d_C-1}$, we have

$${}_{B}\langle\xi_{j}|_{C}\langle\eta_{k}|E|0\rangle_{B}|\eta_{i}\rangle_{C} = {}_{B}\left(\sum_{t_{1}=0}^{d_{B}-2} w_{d_{B}-1}^{-jt_{1}}\langle t_{1}+1|\right)_{C}\left(\sum_{t_{2}=0}^{d_{C}-2} w_{d_{C}-1}^{-kt_{2}}\langle t_{2}|\right)E|0\rangle_{B}\left(\sum_{t_{3}=0}^{d_{C}-2} w_{d_{C}-1}^{it_{3}}|t_{3}\rangle\right)_{C} = 0.$$
(B2)

We have shown that $_B\langle j+1|_C\langle 0|E|0\rangle_B|i\rangle_C = 0$ for $j \in \mathbb{Z}_{d_B-1}$, $i \in \mathbb{Z}_{d_C-1}$, and $_B\langle d_B-1|_C\langle k+1|E|0\rangle_B|i\rangle_C = 0$ for $k, i \in \mathbb{Z}_{d_C-1}$ by Eq. (B1). Then Eq. (B2) can be expressed by

$${}_{B}\left(\sum_{t_{1}=0}^{d_{B}-3} w_{d_{B}-1}^{-jt_{1}} \langle t_{1}+1|\right) {}_{C}\left(\sum_{t_{2}=1}^{d_{C}-2} w_{d_{C}-1}^{-kt_{2}} \langle t_{2}|\right) {}_{E}|0\rangle_{B}\left(\sum_{t_{3}=0}^{d_{C}-2} w_{d_{C}-1}^{it_{3}}|t_{3}\rangle\right)_{C}=0,$$

10

i.e.

$$\sum_{t_1=0}^{d_B-3} \sum_{t_2=1}^{d_C-2} \sum_{t_3=0}^{d_C-2} w_{d_B-1}^{-jt_1} w_{d_C-1}^{-kt_2} w_{d_C-1B}^{it_3} \langle t_1+1|_C \langle t_2|E|0\rangle_B |t_3\rangle_C = 0,$$

for any $0 \le j \le d_B - 3$, $1 \le k \le d_C - 2$, $0 \le i \le d_C - 2$. It means that

$$[H_1^{\dagger} \otimes H_2^{\dagger} \otimes H_3]X = \mathbf{0}$$

where

$$H_{1} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & w_{d_{B}-1} & \cdots & w_{d_{B}-1}^{(d_{B}-3)} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & w_{d_{B}-1}^{(d_{B}-3)} & \cdots & w_{d_{B}-1}^{(d_{B}-3)^{2}} \end{pmatrix}, \quad H_{2} = \begin{pmatrix} w_{d_{C}-1} & w_{d_{C}-1}^{2} & \cdots & w_{d_{C}-1}^{(d_{C}-2)} \\ w_{d_{C}-1}^{2} & w_{d_{C}-1}^{4} & \cdots & w_{d_{C}-1}^{2} \\ \vdots & \vdots & \ddots & \vdots \\ w_{d_{C}-1}^{(d_{C}-2)} & w_{d_{C}-1}^{2(d_{C}-2)} & \cdots & w_{d_{C}-1}^{(d_{C}-2)^{2}} \end{pmatrix}$$

 $H_3 = (w_{d_C-1}^{ij})_{i,j \in \mathbb{Z}_{d_C-1}}$ and X is a column vector,

$$X = ({}_{B}\langle t_{1} + 1 | {}_{C}\langle t_{2} | E | 0 \rangle_{B} | t_{3} \rangle_{C})_{\{0 \le t_{1} \le d_{B} - 3, \ 1 \le t_{2} \le d_{C} - 2, \ 0 \le t_{3} \le d_{C} - 2\}}.$$

Since H_1, H_2, H_3 are all full-rank matrices, it implies that $H_1^{\dagger} \otimes H_2^{\dagger} \otimes H_3$ is a full-rank matrix. Then $X = \mathbf{0}$, i.e.

$$_{B}\langle t_{1}+1|_{C}\langle t_{2}|E|0\rangle_{B}|t_{3}\rangle_{C}=0, \text{ for } 0\leq t_{1}\leq d_{B}-3, \ 1\leq t_{2}\leq d_{C}-2, \ 0\leq t_{3}\leq d_{C}-2.$$

It also means that

$$_{\mathcal{F}^{(A)}}E_{\mathcal{A}_{1}^{(A)}}=\mathbf{0}. \tag{B3}$$

By using $\mathcal{A}_3(|d_A-1\rangle_A)$ and $\mathcal{A}_2(|\xi_1\rangle_A)$, we can also show that

$$\mathcal{F}^{(A)}E_{\mathcal{A}_{2}^{(A)}} = \mathbf{0} \tag{B4}$$

by the similar discussion as above. Further, by the symmetry of Fig. 2, we can also obtain that

$$_{\mathcal{F}^{(A)}}E_{\mathcal{B}_{1}^{(A)}}=\mathbf{0}, \quad _{\mathcal{F}^{(A)}}E_{\mathcal{B}_{2}^{(A)}}=\mathbf{0}.$$
(B5)

Thus, by Eqs. (B1), (B3), (B4) and (B5), E is a block diagonal matrix. It can be expressed by

$$E = E_{\mathcal{A}_1^{(A)}} \oplus E_{\mathcal{A}_2^{(A)}} \oplus E_{\mathcal{F}^{(A)}} \oplus E_{\mathcal{B}_2^{(A)}} \oplus E_{\mathcal{B}_1^{(A)}}.$$
(B6)

Step 2 Considering $|S\rangle$ and $\{|\beta_0\rangle_A|\beta_j\rangle_B|\beta_k\rangle_C\}_{(j,k)\in\mathbb{Z}_{d_B-2}\times\mathbb{Z}_{d_C-2}\setminus\{(0,0)\}}\subset \mathcal{F}$, where $d_C \geq 4$, then by Eq. (B6), we have

$${}_{B}\left(\sum_{i_{1}=0}^{d_{B}-1}\langle i_{1}|\right){}_{C}\left(\sum_{i_{2}=0}^{d_{C}-1}\langle i_{2}|\right){}_{E}|\beta_{j}\rangle_{B}|\beta_{k}\rangle_{C} = {}_{B}\left(\sum_{i_{1}=1}^{d_{B}-2}\langle i_{1}|\right){}_{C}\left(\sum_{i_{2}=1}^{d_{C}-2}\langle i_{2}|\right){}_{E}|\beta_{j}\rangle_{B}|\beta_{k}\rangle_{C} = 0,$$

for $(j,k) \in \mathbb{Z}_{d_B-2} \times \mathbb{Z}_{d_C-2} \setminus \{(0,0)\}$. Moreover, we have

$$\left(\sum_{i_1=1}^{d_B-2} |i_1\rangle\right)_B \left(\sum_{i_2=1}^{d_C-2} |i_2\rangle\right)_C = |\beta_0\rangle_B |\beta_0\rangle_C.$$

Therefore, by using the states in $\{|S\rangle\} \cup \{|\beta_0\rangle_A |\beta_j\rangle_B |\beta_k\rangle_C\}_{(j,k)\in\mathbb{Z}_{d_B-2}\times\mathbb{Z}_{d_C-2}} \setminus \{(0,0)\}$, we obtain

$${}_{B}\langle\beta_{i}|_{C}\langle\beta_{j}|E|\beta_{k}\rangle_{B}|\beta_{\ell}\rangle_{C} = 0, \quad \text{for } (i,j) \neq (k,\ell) \in \mathbb{Z}_{d_{B}-2} \times \mathbb{Z}_{d_{C}-2}.$$
(B7)

By Eq. (B7), there exists a real number $e_{s,t}$ $(E^{\dagger} = E)$ for $(s,t) \in \mathbb{Z}_{d_B-2} \times \mathbb{Z}_{d_C-2}$, such that

$$E_{\mathcal{F}^{(A)}} = \sum_{s=0}^{d_B-3} \sum_{t=0}^{d_C-3} e_{s,t} |\beta_s\rangle_B \langle\beta_s| \otimes |\beta_t\rangle_C \langle\beta_t|.$$
(B8)

Note that Eq. (B8) also holds for $d_C = 3$ (in this case, $E_{\mathcal{F}^{(A)}} = e_{0,0}|\beta_0\rangle_B\langle\beta_0| \otimes |\beta_0\rangle_C\langle\beta_0|$, and $|\beta_0\rangle_X = |1\rangle_X$ for $X \in \{B, C\}$). Next, by using the states in $\mathcal{A}_1(|\xi_1\rangle_A)$, we have

$${}_{B}\langle 0|_{C}\langle \eta_{i}|E|0\rangle_{B}|\eta_{j}\rangle_{C}=0, \text{ for } i\neq j\in\mathbb{Z}_{d_{C}-1}.$$

Then there exists a real number a_s for $s \in \mathbb{Z}_{d_C-1}$ such that

$$E_{\mathcal{A}_{1}^{(A)}} = \sum_{s=0}^{d_{C}-2} a_{s} |0\rangle_{B} \langle 0| \otimes |\eta_{s}\rangle_{C} \langle \eta_{s}|$$

In the same way, there exist real numbers $a_s, b_s, c_t, d_t, e_{s,t}$ such that the operator

$$E = \sum_{s=0}^{d_C-2} a_s |0\rangle_B \langle 0| \otimes |\eta_s\rangle_C \langle \eta_s| + \sum_{s=0}^{d_B-2} b_s |\eta_s\rangle_B \langle \eta_s| \otimes |d_C - 1\rangle_C \langle d_C - 1| + \sum_{s=0}^{d_B-3} \sum_{t=0}^{d_C-3} e_{s,t} |\beta_s\rangle_B \langle \beta_s| \otimes |\beta_t\rangle_C \langle \beta_t|$$

$$+ \sum_{t=0}^{d_B-2} c_t |\xi_t\rangle_B \langle \xi_t| \otimes |0\rangle_C \langle 0| + \sum_{t=0}^{d_C-2} d_t |d_B - 1\rangle_B \langle d_B - 1| \otimes |\xi_t\rangle_C \langle \xi_t|.$$
(B9)

By using those states $\{|0\rangle_A |\eta_i\rangle_B |\xi_j\rangle_C\}_{(i,j)\in\mathbb{Z}_{d_B-1}\times\mathbb{Z}_{d_C-1}\setminus\{(0,0)\}} = \mathcal{B}_3$, we can show that

$${}_{B}\langle \eta_{k}|_{C}\langle \xi_{\ell}|E|\eta_{i}\rangle_{B}|\xi_{j}\rangle_{C}=0, \quad \text{for } (k,\ell)\neq (i,j)\in\mathbb{Z}_{d_{B}-1}\times\mathbb{Z}_{d_{C}-1}\setminus\{(0,0)\}$$

Assume $k = 0, \ell \neq 0, i \neq 0, j = 0$. By Eq. (B9), we have

$$0 =_{B} \langle \eta_{0} |_{C} \langle \xi_{\ell} | E | \eta_{i} \rangle_{B} | \xi_{0} \rangle_{C}$$

$$= \sum_{s=0}^{d_{C}-2} a_{s} \langle \eta_{0} | 0 \rangle_{B} \langle 0 | \eta_{i} \rangle_{B} \langle \xi_{\ell} | \eta_{s} \rangle_{C} \langle \eta_{s} | \xi_{0} \rangle_{C} + \sum_{s=0}^{d_{B}-3} \sum_{t=0}^{d_{C}-3} e_{s,t} \langle \eta_{0} | \beta_{s} \rangle_{B} \langle \beta_{s} | \eta_{i} \rangle_{B} \langle \xi_{\ell} | \beta_{t} \rangle_{C} \langle \beta_{t} | \xi_{0} \rangle_{C}$$

$$= \sum_{s=0}^{d_{C}-2} a_{s} \langle \xi_{\ell} | \eta_{s} \rangle_{C} \langle \eta_{s} | \xi_{0} \rangle_{C} + w_{d_{C}-1}^{\ell} (d_{B}-2) (d_{C}-2) e_{0,0}.$$
(B10)

There are three cases for the terms in the summation of the last equality.

(a) If s = 0, then

$$\langle \xi_{\ell} | \eta_0 \rangle_C \langle \eta_0 | \xi_0 \rangle_C = (d_C - 2) \sum_{n=1}^{d_C - 2} w_{d_C - 1}^{-(n-1)\ell} = -(d_C - 2) w_{d_C - 1}^{\ell}$$

(b) If $s = \ell$, then

$$\langle \xi_{\ell} | \eta_{\ell} \rangle_C \langle \eta_{\ell} | \xi_0 \rangle_C = -\sum_{n=1}^{d_C - 2} w_{d_C - 1}^{\ell} = -(d_C - 2) w_{d_C - 1}^{\ell}$$

(c) If $s \neq 0, \ell$, then

$$\langle \xi_{\ell} | \eta_s \rangle_C \langle \eta_s | \xi_0 \rangle_C = -\sum_{n=1}^{d_C-2} w_{d_C-1}^{ns-(n-1)\ell} = -w_{d_C-1}^{\ell} \sum_{n=1}^{d_C-2} w_{d_C-1}^{(s-\ell)n} = w_{d_C-1}^{\ell}.$$

Thus by Eq. (B10), we have

$$\sum_{s=0}^{d_C-2} a_s - (d_C - 1)(a_0 + a_\ell) + (d_B - 2)(d_C - 2)e_{0,0} = 0.$$
 (B11)

Since $\ell \neq 0 \in \mathbb{Z}_{d_C-1}$, we must have $a_1 = a_2 = \cdots = a_{d_C-2}$. Then Eq. (B11) can be expressed by

$$-(d_C - 2)a_0 - a_1 + (d_B - 2)(d_C - 2)e_{0,0} = 0.$$
(B12)

Further, by using the states $|S\rangle$, $|0\rangle_A |\eta_1\rangle_B |\xi_0\rangle \in \mathcal{B}_3$ and Eq. (B9) we obtain

$${}_{B}\left(\sum_{i_{1}=0}^{d_{B}-1}\langle i_{1}|\right){}_{C}\left(\sum_{i_{2}=0}^{d_{B}-1}\langle i_{2}|\right){}_{E}|\eta_{1}\rangle_{B}|\xi_{0}\rangle_{C} = (d_{C}-1)(d_{C}-2)a_{0} - (d_{B}-2)(d_{C}-2)^{2}e_{0,0} = 0,$$

i,e.

$$(d_C - 1)a_0 - (d_B - 2)(d_C - 2)e_{0,0} = 0.$$
(B13)

Then, by Eqs. (B12) and (B13), it implies $a_0 = a_1$. Thus $a_0 = a_1 = \ldots = a_{d_C-2} = k$. It means that the operator

$$E_{\mathcal{A}_1^{(A)}} = k \sum_{s=0}^{d_C-2} |0\rangle_B \langle 0| \otimes |\eta_s\rangle_C \langle \eta_s|,$$

which is equivalent to

$$E_{\mathcal{A}_{1}^{(A)}} = k \mathbb{I}_{\mathcal{A}_{1}^{(A)}}.$$
(B14)

Step 3 Considering $|S\rangle$ and $\{|0\rangle_A |\eta_i\rangle_B |\xi_j\rangle_C\}_{(i,j)\in\mathbb{Z}_{d_B-1}\times\mathbb{Z}_{d_C-1}\setminus\{(0,0)\}} = \mathcal{B}_3$. By using Eqs. (B6) and (B14), we have

$${}_{B}\left(\sum_{i_{1}=0}^{d_{B}-1}\langle i_{1}|\right){}_{C}\left(\sum_{i_{2}=0}^{d_{C}-1}\langle i_{2}|\right){}_{E}|\eta_{i}\rangle_{B}|\xi_{j}\rangle_{C}={}_{B}\left(\sum_{i_{1}=0}^{d_{B}-2}\langle i_{1}|\right){}_{C}\left(\sum_{i_{2}=1}^{d_{C}-1}\langle i_{2}|\right){}_{E}|\eta_{i}\rangle_{B}|\xi_{j}\rangle_{C}=0.$$

Moreover, we have

$$\left(\sum_{i_1=0}^{d_B-2} |i_1\rangle\right)_B \left(\sum_{i_2=1}^{d_C-1} |i_2\rangle\right)_C = |\eta_0\rangle_B |\xi_0\rangle_C.$$

Therefore, by using the states $\{|S\rangle\} \cup \{|0\rangle_A |\eta_i\rangle_B |\xi_j\rangle_C\}_{(i,j)\in\mathbb{Z}_{d_B-1}\times\mathbb{Z}_{d_C-1}\setminus\{(0,0)\}}$, we have

$${}_{B}\langle \eta_{k}|_{C}\langle \xi_{\ell}|E|\eta_{i}\rangle_{B}|\xi_{j}\rangle_{C}=0, \quad \text{for } (k,\ell)\neq (i,j)\in\mathbb{Z}_{d_{B}-1}\times\mathbb{Z}_{d_{C}-1}.$$

For any $|j\rangle_B|k\rangle_C \in \mathcal{A}_1^{(A)} \cap \mathcal{B}_3^{(A)}$, we have $_{\{|j\rangle_B|k\rangle_C\}} E_{\mathcal{B}_3^{(A)} \setminus \{|j\rangle_B|k\rangle_C\}} = \mathbf{0}$ by Eqs. (B6) and (B14). Applying Block Trivial Lemma to $\{|\eta_i\rangle_B|\xi_j\rangle_C\}_{(i,j)\in\mathbb{Z}_{d_B-1}\times\mathbb{Z}_{d_C-1}}$, we have

$$E_{\mathcal{B}_{3}^{(A)}} = k_{1} \mathbb{I}_{\mathcal{B}_{3}^{(A)}}.$$
 (B15)

Since $\mathcal{A}_1^{(A)} \cap \mathcal{B}_3^{(A)} \neq \emptyset$, it implies $k = k_1$. Thus, by Eqs. (B14) and (B15), we obtain

$$E_{\mathcal{A}_1^{(A)} \cup \mathcal{B}_3^{(A)}} = k \mathbb{I}_{\mathcal{A}_1^{(A)} \cup \mathcal{B}_3^{(A)}}.$$

By the symmetry of Fig. 2, we can obtain $E = k\mathbb{I}$. Thus, E is trivial.

Appendix C: The proof of Theorem 6

Proof. First, we need to show that \mathcal{U} is a UPB. For the same discussion as Proposition 4, we can obtain the matrices,

$$M^{(d_A)} = \begin{bmatrix} a_0 & a_0 & \cdots & a_0 & b_0 & \cdots & b_0 & c_0 & \cdots & c_0 & c_0 & \cdots & c_0 & d_0 \\ a_0 & a_0 & \cdots & a_0 & b_0 & \cdots & b_0 & & & & b_1 & \cdots & b_1 & a_1 & \cdots & a_1 & a_1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & & & & \\ a_0 & a_0 & \cdots & a_0 & b_0 & \cdots & b_0 & & & & & b_1 & \cdots & b_1 & a_1 & \cdots & a_1 & a_1 \\ d_1 & c_1 & \cdots & c_1 & c_1 & \cdots & c_1 & c_1 & \cdots & c_1 & b_1 & \cdots & b_1 & a_1 & \cdots & a_1 & a_1 \end{bmatrix},$$

where $M^{(d_A-2s)}$ is similar to M of Eq. (A1) in Proposition 4. Next, we have

$$\operatorname{rank}(M^{(d_A)}) = 1, \quad \operatorname{sum}(M^{(d_A)}) = 0.$$
 (C1)

Then we consider AB|C bipartition. We can rearrange the first row of $M^{(d_A)}$ to the $d_B \times d_C$ matrix $M_{(d_A-1)}$ through (AB, C) coordinates of $M^{(d_A)}$, and rearrange the last row of $M^{(d_A)}$ to the $d_B \times d_C$ matrix M_0 through (AB, C) coordinates of $M^{(d_A)}$, where

$$M_{(d_A-1)} = \begin{bmatrix} c_0 & c_0 & \cdots & c_0 & d_0 \\ c_0 & c_0 & \cdots & c_0 & b_0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ c_0 & c_0 & \cdots & c_0 & b_0 \\ a_0 & a_0 & \cdots & a_0 & b_0 \end{bmatrix}, \quad \operatorname{rank}(M_{(d_A-1)}) = 0 \quad \text{or} \quad 1,$$
(C2)

and

$$M_{0} = \begin{bmatrix} b_{1} & a_{1} & \cdots & a_{1} & a_{1} \\ b_{1} & c_{1} & \cdots & c_{1} & c_{1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ b_{1} & c_{1} & \cdots & c_{1} & c_{1} \\ d_{1} & c_{1} & \cdots & c_{1} & c_{1} \end{bmatrix}, \quad \operatorname{rank}(M_{0}) = 0 \text{ or } 1.$$
(C3)

For the same proof of Example 3, we can show that $a_0 = a_1 = b_0 = b_1 = c_0 = c_1 = d_0 = d_1 = 0$ by Eqs. (C1), (C2), and (C3). Then we obtain that

$$\operatorname{rank}(M^{(d_A-2)}) = 1, \quad \operatorname{sum}(M^{(d_A-2)}) = 0.$$
 (C4)

We can rearrange the second row of $M^{(d_A)}$ to the $d_B \times d_C$ matrix $M_{(d_A-2)}$ through (AB, C) coordinates of $M^{(d_A)}$, and rearrange the last but two row of $M^{(d_A)}$ to the $d_B \times d_C$ matrix M_1 through (AB, C) coordinates of $M^{(d_A)}$, where

$$M_{(d_{A}-2)} = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & c_{0}^{(1)} & c_{0}^{(1)} & \cdots & c_{0}^{(1)} & d_{0}^{(1)} & 0 \\ 0 & c_{0}^{(1)} & c_{0}^{(1)} & \cdots & c_{0}^{(1)} & b_{0}^{(1)} & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & c_{0}^{(1)} & c_{0}^{(1)} & \cdots & c_{0}^{(1)} & b_{0}^{(1)} & 0 \\ 0 & a_{0}^{(1)} & a_{0}^{(1)} & \cdots & a_{0}^{(1)} & b_{0}^{(1)} & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix} , \quad \operatorname{rank}(M_{d_{A}-2}) = 0 \text{ or } 1, \tag{C5}$$

14

and

$$M_{1} = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & b_{0}^{(1)} & a_{0}^{(1)} & \cdots & a_{0}^{(1)} & a_{0}^{(1)} & 0 \\ 0 & b_{0}^{(1)} & c_{0}^{(1)} & \cdots & c_{0}^{(1)} & c_{0}^{(1)} & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & b_{0}^{(1)} & c_{0}^{(1)} & \cdots & c_{0}^{(1)} & c_{0}^{(1)} & 0 \\ 0 & d_{0}^{(1)} & c_{0}^{(1)} & \cdots & c_{0}^{(1)} & c_{0}^{(1)} & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix}, \quad \text{rank}(M_{1}) = 0 \text{ or } 1.$$
(C6)

Similarly, we can also show that $a_0^{(1)} = a_1^{(1)} = b_0^{(1)} = b_1^{(1)} = c_0^{(1)} = c_1^{(1)} = d_0^{(1)} = d_1^{(1)} = 0$ by Eqs. (C4), (C5), and (C6). Repeating this process s times, then we obtain that

$$\operatorname{rank}(M^{(d_A-2s)}) = 1, \quad \operatorname{sum}(M^{(d_A-2s)}) = 0.$$
 (C7)

Since $M^{(d_A-2s)}$ is similar to M of Eq. (A1) in Proposition 4, and we can show that $a_0^{(s)} = a_1^{(s)} = b_0^{(s)} = b_1^{(s)} = c_0^{(s)} = c_1^{(s)} = d_0^{(s)} = d_1^{(s)} = d_1^{(s)} = e^{(s)} = 0$ by the proof of Proposition 4. Thus we obtain that $M^{(d_A)}$ is a zero matrix, and it is impossible for rank $(M^{(d_A)}) = 1$.

We can obtain that \mathcal{U} is strongly nonlocal by induction on t along with Proposition 5.

Appendix D: The proof of Proposition 7

Proof. We denote $|\Phi(2)\rangle_{A,B}$ as $|\Phi(2)\rangle_{a,b_1}$, and $|\Phi(2)\rangle_{B,C}$ as $|\Phi(2)\rangle_{b_2,c}$. Assume that $|\Phi(2)\rangle_{a,b_1}$ is distributed between Alice and Bob, and $|\Phi(2)\rangle_{b_{2,c}}$ is distributed between Bob and Charlie. The initial states are

$$|\psi\rangle_{A,B,C}(|0\rangle_{a}|0\rangle_{b_{1}}+|1\rangle_{a}|1\rangle_{b_{1}})(|0\rangle_{b_{2}}|0\rangle_{c}+|1\rangle_{b_{2}}|1\rangle_{c}),$$
(D1)

for $|\psi\rangle_{A,B,C} \in \bigcup_{i=1}^{3} (\mathcal{A}_{i} \cup \mathcal{B}_{i}) \cup \mathcal{F} \cup \{|S\rangle\}$, where *a* is the ancillary system of Alice, *b*₁ and *b*₂ are the ancillary systems of Bob, and *c* is the ancillary system of Charlie. Denote $P[|i\rangle_{\bigstar}] := |i\rangle\langle i|_{\bigstar}$, $P[(|i_1\rangle, |i_2\rangle, \dots, i_r)_{\bigstar}; (|j_1\rangle, |j_2\rangle, \dots, |j_s\rangle)_{\bigstar}; (|k_1\rangle, |k_2\rangle, \dots, |k_t\rangle)_{\diamondsuit}] := (|i_1\rangle\langle i_1| + |i_2\rangle\langle i_2| + \dots + |i_r\rangle\langle i_r|)_{\bigstar} \otimes (|j_1\rangle\langle j_1| + |j_2\rangle\langle j_2| + \dots + |j_s\rangle\langle j_s|)_{\bigstar} \otimes (|k_1\rangle\langle k_1| + |k_2\rangle\langle k_2| + \dots + |k_t\rangle\langle k_t|)_{\diamondsuit}.$ Now the discrimination protocol proceeds as follows.

Step 1. Alice performs the measurement $\{M_1 := P[(|0\rangle, |1\rangle)_A; |0\rangle_a] + P[|2\rangle_A; |1\rangle_a], \overline{M_1} := \mathbb{I} - M_1\}$. Charlie performs the measurement $\{L_1 := P[(|1\rangle, |2\rangle, |3\rangle)_C; |0\rangle_c] + P[|0\rangle_C; |1\rangle_c], L_1 := \mathbb{I} - L_1\}$. If M_1 and L_1 clicks (it means that the operators M_1 and L_1 act on the states in Eq. (D1)), the resulting postmeasurement states are

$$\begin{split} |\psi_{1}(0,1)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|1\rangle_{b}|1\rangle_{c} + |2\rangle_{A}|0\rangle_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|1\rangle_{c} \\ + |1\rangle_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{c}|2\rangle_{c}|2\rangle_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{1}(0,2)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|1\rangle_{b}|1\rangle_{c} + |2\rangle_{A}|0\rangle_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|1\rangle_{c} \\ + |1\rangle_{A}|0\rangle_{B}(w_{3}^{2}|1\rangle + w_{3}|2\rangle)_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{c}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|1\rangle_{c} \\ + |1\rangle_{A}|0\rangle_{B}(w_{3}^{2}|1\rangle + w_{3}|2\rangle)_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|2\rangle_{C} \\ + |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|1\rangle_{b}|1\rangle_{c} - |2\rangle_{A}|0\rangle_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{1}(1,0)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|1\rangle_{b}|0\rangle_{c}|0\rangle_{c} - |2\rangle_{A}|0\rangle_{B}(|1\rangle + |2\rangle)_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{1}(1,1)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|1\rangle_{b}|0\rangle_{c}|0\rangle_{c} - |2\rangle_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{1}(1,2)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{c}|0\rangle_{c} - |2\rangle_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{1}(1,2)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{b}|0\rangle_{c}|0\rangle_{c} - |2\rangle_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{1}(1,2)\rangle \rightarrow |1\rangle_{A}|0\rangle_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{b}|0\rangle_{c}|0\rangle_{c} - |2\rangle_{A}|0\rangle_{B}(w_{3}^{2}|1\rangle + w_{3}|2\rangle)_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{2}(1,0)\rangle \rightarrow |1\rangle_{A}|0\rangle - |1\rangle)_{B}|3\rangle_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{b}|0\rangle_{c}|2\rangle_{c}|2\rangle_{A}|0\rangle - |1\rangle)_{B}|3\rangle_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{2}(1,1)\rangle \rightarrow |1\rangle_{A}|0\rangle - |1\rangle)_{B}|3\rangle_{C}|0\rangle_{a}|0\rangle_{b}|0\rangle_{b}|0\rangle_{c}|2\rangle_{c}|2\rangle_{A}|0\rangle - |1\rangle)_{B}|3\rangle_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{3}(1,0)\rangle \rightarrow |2\rangle_{A}|1\rangle + |2\rangle)_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|1\rangle_{c}|1\rangle_{c}|2\rangle_{A}|1\rangle + |2\rangle)_{B}(w_{3}^{2}|1\rangle + w_{3}^{2}|2\rangle_{C}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{3}(1,0)\rangle \rightarrow |2\rangle_{A}|1\rangle + |2\rangle)_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b}|1\rangle_{b}|1\rangle_{c}|1\rangle_{c}|2\rangle_{A}|1\rangle - |2\rangle_{B}|0\rangle_{c}|1\rangle_{a}|1\rangle_{b}|0\rangle_{b}|0\rangle_{c}, \\ |\psi_{3}(1,0)\rangle \rightarrow |2\rangle_{$$

$$\begin{split} |\psi_{3}(1,2)\rangle \rightarrow |2\rangle_{A}(|1\rangle - |2\rangle)_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c} + |2\rangle_{A}(|1\rangle - |2\rangle)_{B}(w_{3}^{2}|1\rangle + w_{3}|2\rangle)_{C}|1\rangle_{a}|1\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(0,1)\rangle \rightarrow (|0\rangle + |1\rangle)_{A}|2\rangle_{B}(|1\rangle + w_{3}|2\rangle + w_{3}^{2}|3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(1,0)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}|2\rangle_{B}(|1\rangle + |2\rangle + |3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(1,1)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}|2\rangle_{B}(|1\rangle + w_{3}^{2}|2\rangle + w_{3}^{2}|3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(1,2)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}|2\rangle_{B}(|1\rangle + w_{3}^{2}|2\rangle + w_{3}^{2}|3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(1,2)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}|2\rangle_{B}(|1\rangle + w_{3}^{2}|2\rangle + w_{3}^{2}|3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{2}(0,1)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}(|1\rangle - |2\rangle)_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c}, \\ |\phi_{2}(1,0)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}(|1\rangle + |2\rangle)_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c}, \\ |\phi_{2}(1,1)\rangle \rightarrow (|0\rangle - |1\rangle)_{A}(|1\rangle + |2\rangle)_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{3}(0,1)\rangle \rightarrow |0\rangle_{A}(|0\rangle + |1\rangle)_{B}(|1\rangle + w_{3}^{2}|2\rangle + w_{3}^{2}|3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{3}(1,0)\rangle \rightarrow |0\rangle_{A}(|0\rangle - |1\rangle)_{B}(|1\rangle + |2\rangle + |3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{3}(1,1)\rangle \rightarrow |0\rangle_{A}(|0\rangle - |1\rangle)_{B}(|1\rangle + w_{3}^{2}|2\rangle + w_{3}^{2}|3\rangle_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{3}(1,2)\rangle \rightarrow |0\rangle_{A}(|0\rangle - |1\rangle)_{B}(|1\rangle + w_{3}^{2}|2\rangle + w_{3}^{2}|3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(1\rangle) \rightarrow |1\rangle_{A}|1\rangle_{B}(|1\rangle - |2\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi_{1}(1\rangle) \rightarrow |1\rangle_{A}(|0\rangle - |1\rangle)_{B}(|1\rangle + |2\rangle)_{B}|0\rangle_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\phi(1\rangle) \rightarrow |1\rangle_{A}|1\rangle_{B}(|1\rangle - |2\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|1\rangle_{c} + |2\rangle_{A}(|0\rangle + |1\rangle + |2\rangle)_{B}|0\rangle_{C}|1\rangle_{a}|1\rangle_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c} + \\ (|0\rangle + |1\rangle)_{A}(|0\rangle + |1\rangle + |2\rangle)_{B}(|1\rangle + |2\rangle + |3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\psi(1\rangle) \rightarrow |1\rangle_{A}|0\rangle + |1\rangle + |2\rangle)_{B}(|1\rangle + |2\rangle + |3\rangle)_{C}|0\rangle_{a}|0\rangle_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c} .$$

Step 2. Bob performs the measurement $\{M_{2,1} := P[|2\rangle_B; |0\rangle_{b_1}; |0\rangle_{b_2}], M_{2,2} := P[(|1\rangle - |2\rangle)_B; |0\rangle_{b_1}; |1\rangle_{b_2}], M_{2,3} := P[(|1\rangle + |2\rangle)_B; |0\rangle_{b_1}; |1\rangle_{b_2}], \overline{M_2} := \mathbb{I} - \sum_{j=1}^3 M_{2,j}\}$. If $M_{2,1}$ clicks (it means that the operator $M_{2,1}$ acts on the states in Eq. (D2)), then the postmeasurement states are $\{|\phi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, and $|S\rangle \to (|0\rangle + |1\rangle)_A |2\rangle_B(|1\rangle + |2\rangle + |3\rangle)_C |0\rangle_a |0\rangle_{b_1} |0\rangle_{b_2} |0\rangle_c$. Then Alice performs the measurement $\{M_{2,1,1} := P[(|0\rangle - |1\rangle)_A], \overline{M_{2,1,1}} = \mathbb{I} - M_{2,1,1}\}$. If $M_{2,1,1}$ clicks (it means that the operator $M_{2,1,1}$ acts on the states $\{|\phi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$ and $|S\rangle$), then the postmeasurement states are $\{|\phi_1(1,j)\rangle\}_{j\in\mathbb{Z}_3}$, which are locally distinguishable; if $\overline{M_{2,1,1}}$ clicks, then the postmeasurement states are $\{|\phi_1(0,j)\rangle\}_{j\neq0\in\mathbb{Z}_3}$ and $|S\rangle$ that are locally distinguishable. Next, if $M_{2,2}$ clicks, then the postmeasurement states are $\{|\phi_2(i,1)\rangle\}_{i\in\mathbb{Z}_2}$ that are locally distinguishable; if $M_{2,3}$ clicks, then the postmeasurement states are $\{|\phi_2(i,1)\rangle\}_{i\in\mathbb{Z}_2}$ that are locally distinguishable; if $M_{2,3}$ clicks, then the postmeasurement states are $\{|\phi_1(i,j)\rangle\}_{i\in\mathbb{Z}_2}$ that are locally distinguishable; if $M_{2,3}$ clicks, then the postmeasurement states are $\{|\phi_1(i,j)\rangle\}_{i\in\mathbb{Z}_2}$ that are locally distinguishable; if $M_{2,3}$ clicks, then the postmeasurement states are $\{|\phi_2(i,1)\rangle\}_{i\in\mathbb{Z}_2}$ that are locally distinguishable; if $M_{2,3}$ clicks, then the postmeasurement states are $\{|\phi_1(i,j)\rangle\}_{i\in\mathbb{Z}_2}$, $\|\phi_1(i,j)\rangle|_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_2(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_3(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_1(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_1(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_2(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_3(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_1(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_2(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{|\psi_3(i,j)\rangle\}_{i,j\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$, $\{$

 $\begin{array}{l} Step \ 3. & \text{Alice performs the measurement } \{M_3 := P[|0\rangle_A], \overline{M_3} = \mathbb{I} - M_3\}. & \text{If } M_3 \text{ clicks } \\ (\text{it means that the operator } M_3 \text{ acts on the states } \{|\psi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \ \{|\psi_2(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_2}, \\ \{|\psi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \ \{|\phi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \ |\varphi(1)\rangle \text{ and } |S\rangle), \text{ then the postmeasurement states are } \\ \{|\phi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3} \text{ and } |S\rangle \rightarrow |0\rangle_A |0\rangle_B |0\rangle_C |0\rangle_a |0\rangle_{b_1} |1\rangle_{b_2} |1\rangle_c + |0\rangle_A (|0\rangle + |1\rangle)_B (|1\rangle + |2\rangle + |3\rangle)_C |0\rangle_a |0\rangle_{b_1} |0\rangle_{b_2} |0\rangle_c, \\ \text{which are locally distinguishable; if } \overline{M_3} \text{ clicks, then the postmeasurement states are } \\ \{\psi_2(i,j)\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_2}, \ \{|\psi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \ |\varphi(1)\rangle, \text{ and } |S\rangle \rightarrow |1\rangle_A |0\rangle_B |0\rangle_C |0\rangle_a |0\rangle_{b_1} |1\rangle_{b_2} |1\rangle_c + |2\rangle_A (|0\rangle + |1\rangle_A (|0\rangle + |1\rangle)_B (|1\rangle + |2\rangle + |3\rangle)_C |0\rangle_a |0\rangle_{b_1} |0\rangle_{b_2} |0\rangle_c + |2\rangle_A (|0\rangle + |1\rangle + |2\rangle)_B (|1\rangle + |2\rangle + |3\rangle_B |0\rangle_C |1\rangle_a |1\rangle_{b_1} |1\rangle_{b_2} |1\rangle_c + |1\rangle_A (|0\rangle + |1\rangle)_B (|1\rangle + |2\rangle + |3\rangle)_C |0\rangle_a |0\rangle_{b_1} |0\rangle_{b_2} |0\rangle_c + |2\rangle_A (|0\rangle + |1\rangle + |2\rangle)_B (|1\rangle + |2\rangle + |3\rangle)_C |1\rangle_a |1\rangle_{b_1} |0\rangle_{b_2} |0\rangle_c. \\ \end{array}$

Step 4. Charlie performs the measurement $\{M_4 := P[|3\rangle_C], \overline{M_4} = \mathbb{I} - M_4\}$. If M_4 clicks (it means that the operator M_4 acts on the states $\{|\psi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \{|\psi_2(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_2}, \{|\psi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, |\varphi(1)\rangle$ and $|S\rangle$), then then the postmeasurement states are $\{\psi_2(i,j)\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_2}, \text{ and } |S\rangle \rightarrow |1\rangle_A(|0\rangle + |1\rangle)_B|3\rangle_C|0\rangle_a|0\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c + |2\rangle_A(|0\rangle + |1\rangle + |2\rangle)_B|3\rangle_C|1\rangle_a|1\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c$, which are locally distinguishable; if $\overline{M_4}$ clicks, then the postmeasurement states are $\{|\psi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \{|\psi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, |\varphi(1)\rangle$ and $|S\rangle \rightarrow |1\rangle_A|0\rangle_B|0\rangle_C|0\rangle_a|0\rangle_{b_1}|1\rangle_{b_2}|1\rangle_c+|2\rangle_A(|0\rangle+|1\rangle+|2\rangle)_B|0\rangle_C|1\rangle_a|1\rangle_{b_1}|1\rangle_{b_2}|1\rangle_c+|1\rangle_A(|0\rangle+|1\rangle)_B(|1\rangle+|2\rangle)_C|0\rangle_a|0\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c + |2\rangle_A(|0\rangle+|1\rangle+|2\rangle)_C|1\rangle_a|1\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c.$

Step 5. Bob performs the measurement $\{\overline{M}_5 := P[|0\rangle_B], \overline{M}_5 = \mathbb{I} - M_5\}$. If M_5 clicks (it means that the operator M_5 acts on the states $\{|\psi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, \{|\psi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, |\varphi(1)\rangle$ and $|S\rangle$), then the postmeasurement states are $\{|\psi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$ and $|S\rangle \rightarrow |1\rangle_A|0\rangle_B|0\rangle_C|0\rangle_a|0\rangle_b_1|1\rangle_b_2|1\rangle_c + |2\rangle_A|0\rangle_B|0\rangle_C|1\rangle_a|1\rangle_{b_1}|1\rangle_{b_2}|1\rangle_c + |1\rangle_A|0\rangle_B(|1\rangle + |2\rangle)_C|0\rangle_a|0\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c + |2\rangle_A|0\rangle_B(|1\rangle + |2\rangle)_C|0\rangle_a|0\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c + |2\rangle_A|0\rangle_B(|1\rangle + |2\rangle)_C|1\rangle_a|1\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c$. Then, Alice performs the measurement $\{M_{5,1} := P[(|0\rangle + |1\rangle)_a], \overline{M}_{5,1} = P[(|0\rangle - |1\rangle)_a\}$. Bob performs the measurement $\{M_{5,2} := P[(|0\rangle + |1\rangle)_{b_1}], \overline{M}_{5,2} = P[(|0\rangle + |1\rangle)_{b_1}|0\rangle_{b_2}|0\rangle_c$.

 $P[(|0\rangle - |1\rangle)_{b_1}]$. If $M_{5,1}$ and $M_{5,2}$ clicks (It means that the operators $M_{5,1}$ and $M_{5,2}$ act on the states $\{|\psi_1(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$ and $|S\rangle$), then the postmeasurement states are

$$\begin{split} |\psi_{1}(0,1)\rangle \rightarrow (|1\rangle + |2\rangle)_{A}|0\rangle_{B}|0\rangle_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c} \\ + (|1\rangle + |2\rangle)_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\psi_{1}(0,2)\rangle \rightarrow (|1\rangle + |2\rangle)_{A}|0\rangle_{B}|0\rangle_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c} \\ + (|1\rangle + |2\rangle)_{A}|0\rangle_{B}(w_{3}^{2}|1\rangle + w_{3}|2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\psi_{1}(1,0)\rangle \rightarrow (|1\rangle - |2\rangle)_{A}|0\rangle_{B}|0\rangle_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c} \\ + (|1\rangle - |2\rangle)_{A}|0\rangle_{B}(|1\rangle + |2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\psi_{1}(1,1)\rangle \rightarrow (|1\rangle - |2\rangle)_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\psi_{1}(1,2)\rangle \rightarrow (|1\rangle - |2\rangle)_{A}|0\rangle_{B}(w_{3}|1\rangle + w_{3}^{2}|2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |\psi_{1}(1,2)\rangle \rightarrow (|1\rangle - |2\rangle)_{A}|0\rangle_{B}(w_{3}^{2}|1\rangle + w_{3}|2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \\ |S\rangle \rightarrow (|1\rangle + |2\rangle)_{A}|0\rangle_{B}|0\rangle_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|1\rangle_{b_{2}}|1\rangle_{c} \\ + (|1\rangle + |2\rangle)_{A}|0\rangle_{B}(|1\rangle + |2\rangle)_{C}(|0\rangle + |1\rangle)_{a}(|0\rangle + |1\rangle)_{b_{1}}|0\rangle_{b_{2}}|0\rangle_{c}, \end{aligned}$$

which can be easily locally distinguished. All other cases obtain a similar protocol. If $\overline{M_5}$ clicks, then the postmeasurement states are $\{|\psi_3(i,j)\rangle\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, |\varphi(1)\rangle$ and $|S\rangle \rightarrow |2\rangle_A(|1\rangle+|2\rangle)_B|0\rangle_C|1\rangle_a|1\rangle_{b_1}|1\rangle_{b_2}|1\rangle_c+|1\rangle_A|1\rangle_B(|1\rangle+|2\rangle)_C|0\rangle_a|0\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c+|2\rangle_A(|1\rangle+|2\rangle)_C|1\rangle_a|1\rangle_{b_1}|0\rangle_{b_2}|0\rangle_c$. Step 6. Alice performs the means that $\{M_6 := P[[0\rangle_A], \overline{M_6} = 1 - M_6\}$. If M_6 clicks (it means that the

Step 6. Alice performs the measurement $\{M_6 := P[|0\rangle_A], M_6 = \mathbb{I} - M_6\}$. If M_6 clicks (it means that the operator M_6 acts on the states $\{|\psi_3(i,j)\rangle_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}, |\varphi(1)\rangle$ and $|S\rangle$), then the postmeasurement states are $\{\psi_3(i,j)\}_{(i,j)\neq(0,0)\in\mathbb{Z}_2\times\mathbb{Z}_3}$ and $|S\rangle$ that are locally distinguishable. If $\overline{M_6}$ clicks, then the postmeasurement state is $|\varphi(1)\rangle$.

All other cases in step 1 obtain a similar protocol. This completes the proof.

- M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information. Cambridge University Press, Cambridge, UK, 2004.
- [2] B. M. Terhal, D. P. Divincenzo, and D. Leung, "Hiding bits in bell states." Phys. Rev. Lett., vol. 86, no. 25, pp. 5807–5810, 2001.
- [3] D. P. Divincenzo, D. Leung, and B. M. Terhal, "Quantum data hiding," IEEE Trans. Inf. Theory, vol. 48, no. 3, pp. 580–598, 2002.
- [4] T. Eggeling and R. F. Werner, "Hiding classical data in multipartite quantum states," Phys. Rev. Lett., vol. 89, no. 9, p. 097905, 2002.
- [5] W. Matthews, S. Wehner, and A. Winter, "Distinguishability of quantum states under restricted families of measurements with an application to quantum data hiding," *Commun. Math. Phys.*, vol. 291, no. 3, pp. p.813–843, 2009.
- [6] D. Markham and B. C. Sanders, "Graph states for quantum secret sharing," Phys. Rev. A, vol. 78, no. 4, pp. 144–144, 2008.
- [7] M. Hillery, V. Buzek, and A. Berthiaume, "Quantum secret sharing," Phys. Rev. A, vol. 59, p. 1829, 1999.
- [8] R. Rahaman and M. G. Parker, "Quantum scheme for secret sharing based on local distinguishability," Phys. Rev. A, vol. 91, p. 022330, 2015.
- [9] C. H. Bennett, D. P. Divincenzo, C. A. Fuchs, T. Mor, E. M. Rains, P. W. Shor, J. A. Smolin, and W. K. Wootters, "Quantum nonlocality without entanglement," *Phys. Rev. A*, vol. 59, no. 2, pp. 1070–1091, 1999.
- [10] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, "Quantum entanglement," *Reviews of modern physics*, vol. 81, no. 2, p. 865, 2009.
- [11] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, "Bell nonlocality," *Reviews of Modern Physics*, vol. 86, no. 2, p. 419, 2014.
- [12] D. P. Divincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. M. Terhal, "Unextendible product bases, uncompletable product bases and bound entanglement," *Commun. Math. Phys.*, vol. 238, no. 3, pp. 379–410, 2003.
- [13] Y. Feng and Y. Shi, "Characterizing locally indistinguishable orthogonal product states," *IEEE Trans. Inf. Theory*, vol. 55, no. 6, pp. p.2799–2806, 2009.
- [14] J. Niset and N. J. Cerf, "Multipartite nonlocality without entanglement in many dimensions," Phys. Rev. A, vol. 74, p. 052103, 2006.
- [15] Y. Yang, F. Gao, G. Tian, T. Cao, and Q. Wen, "Local distinguishability of orthogonal quantum states in a 2⊗2⊗2 system," Phys. Rev. A, vol. 88, no. 2, p. 024301, 2013.

- [16] S. Halder, "Several nonlocal sets of multipartite pure orthogonal product states," Phys. Rev. A, vol. 98, p. 022303, 2018.
- [17] G. Xu, Q. Wen, F. Gao, S. Qin, and H. Zuo, "Local indistinguishability of multipartite orthogonal product bases," *Quantum Inf. Process.*, vol. 16, no. 11, p. 276, 2017.
- [18] Y.-L. Wang, M.-S. Li, Z.-J. Zheng, and S.-M. Fei, "The local indistinguishability of multipartite product states," Quantum Inf. Process., vol. 16, no. 1, pp. 1–13, 2017.
- [19] Z.-C. Zhang, K.-J. Zhang, F. Gao, Q.-Y. Wen, and C. H. Oh, "Construction of nonlocal multipartite quantum states," *Phys. Rev. A*, vol. 95, p. 052344, 2017.
- [20] S. Ghosh, G. Kar, A. Roy, and D. Sarkar, "Distinguishability of maximally entangled states," Phys. Rev. A, vol. 70, p. 022304, 2004.
- [21] H. Fan, "Distinguishability and indistinguishability by local operations and classical communication," Phys. Rev. Lett., vol. 92, p. 177905, 2004.
- [22] M. Nathanson, "Distinguishing bipartitite orthogonal states using locc: Best and worst cases," J. Math. Phys, vol. 46, no. 6, p. 062103, 2005.
- [23] N. Yu, R. Duan, and M. Ying, "Any $2 \otimes n$ subspace is locally distinguishable," Phys. Rev. A, vol. 84, p. 012304, 2011.
- [24] R. Duan, Y. Feng, Z. Ji, and M. Ying, "Distinguishing arbitrary multipartite basis unambiguously using local operations and classical communication," *Phys. Rev. Lett.*, vol. 98, p. 230502, 2007.
- [25] S. Bandyopadhyay, S. Ghosh, and G. Kar, "Locc distinguishability of unilaterally transformable quantum states," New J. Phys., vol. 13, no. 12, p. 123013, 2011.
- [26] A. Cosentino, "Positive-partial-transpose-indistinguishable states via semidefinite programming," Phys. Rev. A, vol. 87, no. 1, 2013.
- [27] N. Yu, R. Duan, and M. Ying, "Four locally indistinguishable ququad-ququad orthogonal maximally entangled states," *Phys. Rev. Lett.*, vol. 109, no. 2, p. 020506, 2012.
- [28] S. Bandyopadhyay, "Entanglement, mixedness, and perfect local discrimination of orthogonal quantum states," Phys. Rev. A, vol. 85, p. 042319, 2012.
- [29] Y.-L. Wang, M.-S. Li, Z.-J. Zheng, and S.-M. Fei, "Nonlocality of orthogonal product-basis quantum states," Phys. Rev. A, vol. 92, p. 032313, 2015.
- [30] M.-S. Li, Y.-L. Wang, F. Shi, and M.-H. Yung, "Local distinguishability based genuinely quantum nonlocality without entanglement," arXiv:2011.10286, 2020.
- [31] C. H. Bennett, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. M. Terhal, "Unextendible product bases and bound entanglement," *Phys. Rev. Lett.*, vol. 82, no. 26, p. 5385, 1999.
- [32] S. De Rinaldis, "Distinguishability of complete and unextendible product bases," Phys. Rev. A, vol. 70, no. 2, p. 022309, 2004.
- [33] S. Halder, M. Banik, and S. Ghosh, "Family of bound entangled states on the boundary of the Peres set," *Physical Review* A, vol. 99, no. 6, p. 062329, 2019.
- [34] J. Tura, R. Augusiak, P. Hyllus, M. Kuś, J. Samsonowicz, and M. Lewenstein, "Four-qubit entangled symmetric states with positive partial transpositions," *Physical Review A*, vol. 85, no. 6, p. 060302, 2012.
- [35] J. Chen, L. Chen, and B. Zeng, "Unextendible product basis for fermionic systems," Journal of Mathematical Physics, vol. 55, no. 8, 2014.
- [36] R. Augusiak, T. Fritz, M. Kotowski, M. Kotowski, M. Pawlowski, M. Lewenstein, and A. Acin, "Tight bell inequalities with no quantum violation from qubit unextendible product bases," *Physical Review A*, vol. 85, no. 4, p. 042113, 2012.
- [37] R. Augusiak, J. Stasińska, C. Hadley, J. Korbicz, M. Lewenstein, and A. Acin, "Bell inequalities with no quantum violation and unextendable product bases," *Physical review letters*, vol. 107, no. 7, p. 070401, 2011.
- [38] N. Alon and L. Lovász, "Unextendible product bases," Journal of combinatorial theory. Series A, vol. 95, no. 1, pp. 169–179, 2001.
- [39] K. Feng, "Unextendible product bases and 1-factorization of complete graphs," Discrete Appl. Math., vol. 154, pp. 942–949, 2006.
- [40] N. Johnston, "The minimum size of qubit unextendible product bases," arXiv preprint arXiv:1302.1604, 2013.
- [41] J. Chen and N. Johnston, "The minimum size of unextendible product bases in the bipartite case (and some multipartite cases)," Communications in Mathematical Physics, vol. 333, no. 1, pp. 351–365, 2013.
- [42] S. Halder, M. Banik, S. Agrawal, and S. Bandyopadhyay, "Strong quantum nonlocality without entanglement," Phys. Rev. Lett., vol. 122, no. 4, p. 040403, 2019.
- [43] P. Yuan, G. Tian, and X. Sun, "Strong quantum nonlocality without entanglement in multipartite quantum systems," *Phys. Rev. A*, vol. 102, p. 042228, 2020.
- [44] F. Shi, M.-S. Li, M. Hu, L. Chen, M.-H. Yung, Y.-L. Wang, and X. Zhang, "Strong quantum nonlocality from hypercubes," arXiv:2110.08461, 2021.
- [45] ——, "Strongly nonlocal unextendible product bases do exist," arXiv:2101.00735v2, 2021.
- [46] S. Agrawal, S. Halder, and M. Banik, "Genuinely entangled subspace with all-encompassing distillable entanglement across every bipartition," *Phys. Rev. A*, vol. 99, no. 3, 2019.
- [47] F. Shi, M. Hu, L. Chen, and X. Zhang, "Strong quantum nonlocality with entanglement," Phys. Rev. A, vol. 102, p. 042202, 2020.
- [48] Y.-L. Wang, M.-S. Li, and M.-H. Yung, "Graph connectivity based strong quantum nonlocality with genuine entanglement," *Physical Review A*, vol. 104, no. 1, p. 012424, 2021.
- [49] S. M. Cohen, "Understanding entanglement as resource: Locally distinguishing unextendible product bases," Phys. Rev. A, vol. 77, no. 1, p. 012304, 2008.

- [50] S. Ghosh, G. Kar, A. Roy, A. Sen, U. Sen et al., "Distinguishability of bell states," Phys. Rev. Lett., vol. 87, no. 27, p. 277902, 2001.
- [51] S. Bandyopadhyay, S. Halder, and M. Nathanson, "Entanglement as a resource for local state discrimination in multipartite systems," *Phys. Rev. A*, vol. 94, no. 2, p. 022311, 2016.
- [52] Z.-C. Zhang, F. Gao, T.-Q. Cao, S.-J. Qin, and Q.-Y. Wen, "Entanglement as a resource to distinguish orthogonal product states," Sci. Rep., vol. 6, no. 1, pp. 1–7, 2016.
- [53] Ö. Güngör and S. Turgut, "Entanglement-assisted state discrimination and entanglement preservation," Phys. Rev. A, vol. 94, no. 3, p. 032330, 2016.
- [54] Z.-C. Zhang, Y.-Q. Song, T.-T. Song, F. Gao, S.-J. Qin, and Q.-Y. Wen, "Local distinguishability of orthogonal quantum states with multiple copies of 2⊗2 maximally entangled states," *Phys. Rev. A*, vol. 97, no. 2, p. 022334, 2018.
- [55] S. Rout, A. G. Maity, A. Mukherjee, S. Halder, and M. Banik, "Genuinely nonlocal product bases: Classification and entanglement-assisted discrimination," *Phys. Rev. A*, vol. 100, no. 3, p. 032321, 2019.
- [56] Z.-C. Zhang, X. Wu, and X. Zhang, "Locally distinguishing unextendible product bases by using entanglement efficiently," *Phys. Rev. A*, vol. 101, no. 2, p. 022306, 2020.
- [57] F. Shi, X. Zhang, and L. Chen, "Unextendible product bases from tile structures and their local entanglement-assisted distinguishability," *Phys. Rev. A*, vol. 101, no. 6, p. 062329, 2020.
- [58] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, "Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels," vol. 70, no. 13, 1993, p. 1895.
- [59] F. Shi, M.-S. Li, L. Chen, and X. Zhang, "Supplementary material: strongly nonlocal four-partite UPBs," 2021. [Online]. Available: http://staff.ustc.edu.cn/~drzhangx/papers/UPBnonunisupp.pdf
- [60] P. Bej and S. Halder, "Unextendible product bases, bound entangled states, and the range criterion," Physics Letters A, vol. 386, p. 126992, 2021.
- [61] S. Halder and R. Sengupta, "Construction of noisy bound entangled states and the range criterion," *Physics Letters A*, vol. 383, no. 17, pp. 2004–2010, 2019.