NONLINEAR ANDERSON LOCALIZED STATES AT ARBITRARY DISORDER

WENCAI LIU AND W.-M. WANG

ABSTRACT. It is classical, following Furstenberg's theorem on positive Lyapunov exponent for products of random $SL(2,\mathbb{R})$ matrices, that the one dimensional random Schrödinger operator has Anderson localization at arbitrary disorder. This paper proves a nonlinear analogue, thereby establishing a KAM-type persistence result for a *non-integrable* system.

1. INTRODUCTION AND THE MAIN THEOREM

We study the discrete nonlinear random Schrödinger equation (NLRS) in one dimension:

(1)
$$i\frac{\partial u}{\partial t} = -\Delta u + Vu + \delta |u|^{2p} u, \ p \in \mathbb{N},$$

where Δ is the discrete Laplacian:

$$(\Delta u)(x) = u(x+1) + u(x-1),$$

and $V = \{v_x\}$ is a family of independent identically distributed random variables on [0, 1], with distribution density g. Assume that g is bounded, $g \in L^{\infty}$. Let

(2)
$$H = -\Delta + V,$$

be the random Schrödinger operator. It is well-known, as a consequence of Furstenberg's theorem on positive Lyapunov exponent for products of random $SL(2,\mathbb{R})$ matrices [15], that with probability 1, H has Anderson localization, namely, pure point spectrum with exponentially decaying eigenfunctions [19, 22]. (See also [9, 16, 20, 21, 24].) In higher dimensions and for large disorder, i.e., replacing V by λV , $\lambda \gg 1$, Anderson localization has been established using multiscale analysis [13], see also [26], or fractional moment method [1].

Assume that H has Anderson localization. Let $\{\phi_j^V\}_{j\in\mathbb{Z}}$ be the (real) eigenbasis of H. Assume that ℓ_j^V satisfies

$$|\phi_j^V(\ell_j^V)| = \max_x |\phi_j^V(x)|.$$

²⁰¹⁰ Mathematics Subject Classification. 35R60 (primary); 35B15, 82B44 (secondary).

Key words and phrases. Anderson localization, quasi-periodic solution, nonlinear random Schrödinger equation, large deviation theorem, semi-algebraic set.

(If it is not unique, one may choose a maximum arbitrarily.) We call ℓ_j^V , the *localization center*. It suffices to say here that as a consequence of localization, there is an eigenfunction labelling such that when $j_1 < j_2$, $\ell_{j_1}^V \leq \ell_{j_2}^V$ (see sect. 3 and appendix A for details), and that we use this labelling.

So for a given V such that H has Anderson localization, let $j \in \mathbb{Z}$, and denote by ϕ_j^V and μ_j^V the eigenfunctions and corresponding eigenvalues of H. When $\delta = 0$, all solutions to (1) are of the form

$$\sum_{j\in\mathbb{Z}}c_j e^{-i\mu_j^V t}\phi_j^V,$$

with appropriate c_j , which decay to 0, as $j \to \pm \infty$. This paper is concerned with the case $c_j \neq 0$ for finite (but arbitrary) number of j.

Denote by $\mathbb{P}(\cdot)$ the measure of a set. We prove the following nonlinear analogue:

Theorem 1.1. Consider the discrete NLRS in one dimension:

(3)
$$i\frac{\partial u}{\partial t} = -\Delta u + Vu + \delta |u|^{2p}u, p \in \mathbb{N}.$$

Let $a = (a_1, a_2, \dots, a_b) \in [1, 2]^b$. For any $\epsilon > 0$, there exists l_{ϵ} such that the following holds. Fix any $L \ge \ell_{\epsilon}$ and $\beta_k \in \mathbb{Z}$, $k = 1, 2, \dots, b$ satisfying $10L \le |\beta_k| \le L^3$ and $|\beta_k - \beta_{k'}| \ge 10L$ for any distinct $k, k' \in \{1, 2, \dots, b\}$, there exist a probability space X_{ϵ} with $\mathbb{P}(X_{\epsilon}) \ge 1 - \epsilon$ and $\delta_0 > 0$ (depending on g, ϵ and L) such that for any $V \in X_{\epsilon}$ and $0 < \delta \le \delta_0$, there exists a set $A_{\delta} \subset [1, 2]^b$ of measure at least $1 - e^{-|\log \delta|^{1/2}}$, such that for any $a \in A_{\delta}$, any eigenfunction $\phi_{\alpha_k}^V$ with $\ell_{\alpha_k}^V \in B_k = \{l \in \mathbb{Z} : |l - \beta_k| \le L\}$, $k = 1, 2, \dots, b$, the solution to the linear equation:

$$u_0(t,x) = \sum_{k=1}^{b} a_k e^{-i\mu_{\alpha_k}^V t} \phi_{\alpha_k}^V(x),$$

bifurcates to a solution to the nonlinear equation (3):

$$u(t,x) = \sum_{(n,j)\in\mathbb{Z}^b\times\mathbb{Z}} \hat{u}(n,j)e^{in\cdot\omega t}\phi_j^V(x) = \sum_{k=1}^b a_k e^{-i\omega_k t}\phi_{\alpha_k}^V(x) + O(\delta^{1/2}),$$

satisfying $\omega = (\omega_1, \omega_2, \cdots, \omega_b) = (\mu_{\alpha_1}^V, \mu_{\alpha_2}^V, \cdots, \mu_{\alpha_b}^V) + O(\delta)$, and $\hat{u}(n, j)$ decay exponentially as $|(n, j)| \to \infty$.

- **Remark 1.** (1) For any B_k , $k = 1, 2, \dots, b$, there are at least $2(1 \epsilon)L$ normalized eigenfunctions $\phi_{\alpha_k}^V$ such that the localization centers $\ell_{\alpha_k}^V$ lie in B_k .
 - (2) We could replace L^3 with $e^{L^{\kappa}}$, $0 < \kappa < 1$. We could also replace $1 e^{-|\log \delta|^{1/2}}$ with $1 e^{-|\log \delta|^{\kappa}}$, $0 < \kappa < 1$.

1.1. About Theorem 1.1. The linear solution u_0 is localized in space, and quasi-periodic in time, with frequencies the *b* eigenvalues of the linear random Schrödinger operator. Theorem 1.1 shows that under small nonlinear perturbations, for a large set of amplitudes, there is a solution to the nonlinear equation nearby. This nonlinear solution *u* remains localized in space and quasi-periodic in time; moreover the frequencies are harmonics of the modulated *b* Fourier modes of the linear random Schrödinger equation.

Theorem 1.1 is a KAM-type persistence result. Most such results pertain to perturbations of integrable systems. The random Schrödinger equation is, however, *non-integrable*. Nonetheless, Theorem 1.1 shows persistence of time quasi-periodic, localized solutions. Moreover there is an abundance of such solutions. This is the main novelty.

Remark 2. Previously the paper [8] established existence of quasi-periodic solutions at large disorder. It perturbs about the diagonal operator λV , $\lambda \gg 1$, in the canonical \mathbb{Z}^d basis. The method is not applicable here. For d = 1 and large disorder, see also [30]. For other nonlinear random models, see e.g., [14].

Remark 3. The symbol \hat{u} is merely a notation here and does not carry the connotation of being the dual of u.

1.2. Ideas of the proof. One of the main ideas is to fix a "good" random potential, and work in the eigenfunction basis provided by the random Schrödinger operator. We give the requirements to be good in sect. 1.3 below. Fixing a potential circumvents the lack of control of the eigenfunctions as the potentials vary. Note moreover that generally it is not possible to know precisely the eigenfunctions of the random Schrödinger operators, even for large disorder, see [12].

So fix indeed such a good potential V, and let

(4)
$$u_0(t,x) = \sum_{k=1}^b a_k e^{-i\mu_{\alpha_k}^V t} \phi_{\alpha_k}^V(x),$$

be a solution to the linear equation, as in Theorem 1.1. As an ansatz, we seek solutions of the form:

(5)
$$u(t,x) = \sum_{(n,j)\in\mathbb{Z}^b\times\mathbb{Z}} \hat{u}(n,j)e^{in\cdot\omega t}\phi_j^V(x).$$

Note that u(t, x) of the above form are closed under multiplication and complex conjugation. So we may seek solutions to (1) in this form.

Using (5) in (1) leads to the following nonlinear system of equations on $\mathbb{Z}^b \times \mathbb{Z}$:

(6)
$$(n \cdot \omega + \mu_j^V)\hat{u}(n,j) + \delta W_{\hat{u}}(n,j) = 0, \ (n,j) \in \mathbb{Z}^b \times \mathbb{Z},$$

where, to give an idea, when p = 1,

(7)
$$W_{\hat{u}}(n,j) = \sum_{\substack{n_1+n_2-n_3=n\\n_1,n_2,n_3\in\mathbb{Z}^b}} \sum_{j_1,j_2,j_3\in\mathbb{Z}} \hat{u}(n_1,j_1)\hat{u}(n_2,j_2)\overline{\hat{u}(n_3,j_3)} \\ \left(\sum_{x\in\mathbb{Z}} \phi_j^V(x)\phi_{j_1}^V(x)\phi_{j_2}^V(x)\phi_{j_3}^V(x)\right);$$

while for general p,

(8)
$$W_{\hat{u}}(n,j) = \sum_{\substack{n' + \sum_{m=1}^{p} (nm - n'_m) = n \\ n', nm, n'_m \in \mathbb{Z}^b}} \sum_{j', l_m, l'_m \in \mathbb{Z}} \hat{u}(n',j') \prod_{m=1}^{p} \hat{u}(n_m, l_m) \overline{\hat{u}(n'_m, l'_m)} \\ \left(\sum_{x \in \mathbb{Z}} \phi_j^V(x) \phi_{j'}^V(x) \prod_{m=1}^{p} \phi_{l_m}^V(x) \phi_{l'_m}^V(x) \right).$$

1.3. The good potentials. The linear solution u_0 solves (1) to order δ . One may write u_0 in the form (5), with $\hat{u}(-e_k, \alpha_k) = a_k, \ k = 1, 2, \dots, b$, where e_k is the kth basis vector of \mathbb{Z}^b , $\hat{u}(n, j) = 0$ otherwise, and $\omega_k = \mu_{\alpha_k}^V$, $k = 1, 2, \dots, b$. The vector $W_{\hat{u}}$ in (6) depends on $a_k, \ k = 1, 2, \dots, b$. Generally speaking, one would need parameters to solve the nonlinear equation (6) using a Newton scheme, starting from the approximate solution u_0 . Since V is fixed, the a_k 's are the parameters in the problem. There is however, a δ factor in front of $W_{\hat{u}}$.

The small $\mathcal{O}(\delta)$ parameters pose difficulties mainly at small scales: $|(n, j)| \ll \delta^{-1}$, when estimating the inverse of the linearized operators. The key new idea is that we can overcome this difficulty if the diagonals of the linear operator in (6) satisfy a *clustering property*. Roughly speaking, this means that if two diagonal elements are "not close", then they are "far apart". (One may think of the integers, which have this property: if two integers are not equal, then they are at least of distance 1.) This then permits localizing about the diagonals in $\mathcal{O}(\delta)$ intervals, which compensates for the small $\mathcal{O}(\delta)$ parameters. The potentials V that lead to clustering properties, in addition to Anderson localization, are good potentials.

It should be emphasized that the clustering property is only needed at *small* scales, and *not* large ones. This makes the approach robust, potentially applicable to many problems.

Remark 4. See papers [27] and [28], which use deterministic clustering properties. In [28], this was established using number theory.

1.4. Anderson localization and clustering property of the diagonals. We use Anderson localization to establish probabilistic clustering at small scales. So one may set ω to be the frequencies of the u_0 in (4), which are b eigenvalues of the H in (2). The diagonals in (6) then correspond to a family of harmonics, i.e., certain linear combinations of the eigenvalues of H.

The proof of the clustering of these (low lying) harmonics is rather delicate. The Minami estimate [25] on eigenvalue spacing plays a fundamental role. Uniform property of Anderson localization, see [11, 17], is essential. Wegner estimate [29] comes into play as well. This is done in sects. 2 and 3, and the conclusion is summarized in Theorem 3.5, which also provides lower bounds on the diagonals. The clustering property permits the analysis to go beyond small perturbative scales, and is one of the main points of the paper.

1.5. Small scale analysis. The clustering property indicates that at *small* scales the spectrum of the diagonal operator has many gaps. Using perturbation theory, the linearized operators are invertible in the gaps; while away from the gaps, one may work locally in intervals of size $\mathcal{O}(\delta)$. This greatly reduces the number of resonances, and consequently $\mathcal{O}(\delta)$ parameters suffice for the analysis. This is the case for the proof of the large deviation theorem applying Cartan estimates in sect. 4, as well as for the semi-algebraic projection in sect. 6.

1.6. Large scale analysis. Large scale analysis is related to what has been done before in [8], cf. also Chap. 18 [5], which are a priori tailored for parameters of order 1. However, after incorporating the local argument recounted above, it can be adapted and used to prove Theorem 1.1.

1.7. **Organization of the paper.** Sect. 2 establishes the *good* potential space; sect. 3 makes linear estimates for small scales; sect. 4 proves a large deviation theorem, to be used for the nonlinear analysis at large scales; sects. 5 and 6 finally solve (6) and hence (1), using a Lyapunov-Schmidt decomposition and Theorems 3.5 and 4.1.

2. One dimensional random Schrödinger operators in finite volumes

For an operator H on $\ell^2(\mathbb{Z}^d)$ and $\Lambda \subset \mathbb{Z}^d$, let $H_{\Lambda} = R_{\Lambda}HR_{\Lambda}$, where R_{Λ} is the restriction. For $n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d$, let $|n| = \max_{j \in \{1, 2, \cdots, d\}} |n_j|$ denote the ℓ^{∞} norm. In this paper d equals either 1 or b + 1.

In this section, we study the one dimensional random Schrödinger operator $H = -\Delta + V$ restricted to finite volumes. For $\Lambda \subset \mathbb{Z}$, denote by $\tilde{\mu}_j^{\Lambda}$, $j \in \Lambda$, eigenvalues of $H_{\Lambda} = R_{\Lambda} H R_{\Lambda}$, with corresponding normalized eigenvectors $\tilde{\phi}_j^{\Lambda}$.

Remark 5. Note that $\tilde{\mu}_j^{\Lambda}$ and $\tilde{\phi}_j^{\Lambda}$ depend on the realization of the potentials in Λ . It is convenient here to label the eigenvalues and normalized eigenvectors by $j \in \Lambda$, instead of $j \in \{1, 2, \dots, |\Lambda|\}$.

For a ball $B = \{n \in \mathbb{Z} : |n - n_0| \leq l\}$ of size l with center n_0 , denote by rB, the dilation: $rB = \{n \in \mathbb{Z} : |n - n_0| \leq rl\}$. Fix L > 0. Let $\tilde{B}_{(k,L)}$ be a ball of size $L, k = 1, 2, \dots, b$. Assume that

$$L \leq \operatorname{dist}(\tilde{B}_{(k,L)}, 0) \leq L^4,$$

and for any distinct k and k',

(9)
$$\operatorname{dist}(\tilde{B}_{(k,L)}, \tilde{B}_{(k',L)}) \ge 6L$$

Since L will be fixed, we omit the dependence on L of $\tilde{B}_{(k,L)}$ and write simply \tilde{B}_k .

Recall that e_k , $k = 1, 2, \dots, b$, are the canonical basis vectors of \mathbb{Z}^b . Let $C_0 > 0, \gamma_0 > 0, q_0 > 0$ be three fixed constants, which will be determined later. Assume that H_{Λ} has b eigen-pairs $\tilde{\mu}^{\Lambda}_{\tilde{\alpha}_k}$ and $\tilde{\phi}^{\Lambda}_{\tilde{\alpha}_k}$, $k = 1, 2, \dots, b$ such that for any $k = 1, 2, \dots, b$, there exists $\tilde{\ell}^{\Lambda}_{\tilde{\alpha}_k} \in \tilde{B}_k$ such that

(10)
$$|\tilde{\phi}^{\Lambda}_{\tilde{\alpha}_k}(\ell)| \le C_0 (1 + |\tilde{\ell}^{\Lambda}_{\tilde{\alpha}_k}|)^{q_0} e^{-\gamma_0 |\ell - \tilde{\ell}^{\Lambda}_{\tilde{\alpha}_k}|}, \ell \in \Lambda.$$

Let $\tilde{\omega}_k^{\Lambda} = \tilde{\mu}_{\tilde{\alpha}_k}^{\Lambda}$, $k = 1, 2, \dots, b$ and $\tilde{\omega}^{\Lambda} = (\tilde{\omega}_1^{\Lambda}, \tilde{\omega}_2^{\Lambda}, \dots, \tilde{\omega}_b^{\Lambda}) \in \mathbb{R}^b$. When there is no ambiguity, we omit the dependence on Λ . In the following, $\delta > 0$ is sufficiently small.

Remark 6. Condition (10) is motivated by uniform properties of Anderson localization in infinite volume, see Theorem 3.1.

2.1. Estimates on the diagonals. Let $\Lambda_1 = \left[-2\left\lfloor e^{|\log \delta|^{\frac{3}{4}}}\right\rfloor, 2\left\lfloor e^{|\log \delta|^{\frac{3}{4}}}\right\rfloor\right]$, where $\lfloor x \rfloor$ is the integer part of x. (The choice of scales is in view of the later nonlinear analysis.) Denote by S_1 the probability space on which H_{Λ_1} has eigen-pairs $\tilde{\mu}_{\tilde{\alpha}_k}^{\Lambda_1}$ and $\tilde{\phi}_{\tilde{\alpha}_k}^{\Lambda_1}$, $k = 1, 2, \cdots, b$ satisfying (10) (with $\Lambda = \Lambda_1$), and there exists either

(11)
$$(n,j) \in \left[-2\left\lfloor e^{|\log \delta|^{\frac{3}{4}}}\right\rfloor, 2\left\lfloor e^{|\log \delta|^{\frac{3}{4}}}\right\rfloor\right]^{b+1} \setminus \{-e_k, \tilde{\alpha}_k\}_{k=1}^b,$$

such that

(12)
$$|n \cdot \tilde{\omega}^{\Lambda_1} + \tilde{\mu}_j^{\Lambda_1}| \le 4\delta^{\frac{1}{8}}$$

or

(13)
$$(n,j) \in \left[-2\left\lfloor e^{|\log \delta|^{\frac{3}{4}}}\right\rfloor, 2\left\lfloor e^{|\log \delta|^{\frac{3}{4}}}\right\rfloor\right]^{b+1} \setminus \{e_k, \tilde{\alpha}_k\}_{k=1}^b,$$

such that

(14)
$$|-n\cdot\tilde{\omega}^{\Lambda_1}+\tilde{\mu}_j^{\Lambda_1}| \le 4\delta^{\frac{1}{8}}$$

and the normalized eigenvector corresponding to $\tilde{\mu}_j^{\Lambda_1}$ satisfies for some $\tilde{\ell}_j^{\Lambda_1} \in \Lambda_1$,

(15)
$$|\tilde{\phi}_{j}^{\Lambda_{1}}(\ell)| \leq C_{0}(1+|\tilde{\ell}_{j}^{\Lambda_{1}}|)^{q_{0}}e^{-\gamma_{0}|\ell-\tilde{\ell}_{j}^{\Lambda_{1}}|}, \ell \in \Lambda_{1}.$$

Fix a large constant $q_1 > 0$. Denote by S_N the probability space such that for any $j \in [-N, N], j' \in [-N, N]$ and $j \neq j'$,

(16)
$$|\tilde{\mu}_{j}^{\Lambda} - \tilde{\mu}_{j'}^{\Lambda}| \ge \frac{1}{N^{q_1}}, \Lambda = [-N, N].$$

Denote by \mathbb{P} the measure as before, and \mathbb{E} the expectation.

Theorem 2.1. For small δ , we have

$$\mathbb{P}(S_1) \le e^{C|\log \delta|^{\frac{3}{4}}} \delta^{\frac{1}{8}},$$

where C is a large constant independent of δ .

We will use the following three lemmas to prove Theorem 2.1.

Lemma 2.2 (Wegner estimate [29]). Let $\Lambda \subset \mathbb{Z}$. For any $E \in \mathbb{R}$ and $\varepsilon > 0$,

$$\mathbb{E}(\operatorname{dist}(E, \sigma(H_{\Lambda})) \leq \varepsilon) \leq C |\Lambda| \varepsilon.$$

Lemma 2.3 (Minami estimate [25]). Let $\Lambda \subset \mathbb{Z}$ and $J \subset \mathbb{R}$ be an interval. Then we have

$$\mathbb{E}([\operatorname{tr}(\mathbf{1}_J(H_{\Lambda}))] \cdot [\operatorname{tr}(\mathbf{1}_J(H_{\Lambda})) - 1]) \le C |\Lambda|^2 |J|^2,$$

where $\mathbf{1}_J$ is the characteristic function of the interval J. In particular, we have that

 $\mathbb{P}(\text{there exist distinct } j, j' \in \Lambda \text{ such that } |\tilde{\mu}_j^{\Lambda} - \tilde{\mu}_{j'}^{\Lambda}| \le \varepsilon) \le C\varepsilon |\Lambda|^2,$

and hence

$$\mathbb{P}(S_N) \le CN^{-q_1+2}$$

Lemma 2.4. Let $\Lambda \subset \mathbb{Z}$. Assume that eigenvalues $\tilde{\mu}_j$, $j \in \Lambda$, of H_{Λ} are simple and let

(17)
$$d = \min_{j \neq j', j \in \Lambda, j' \in \Lambda} |\tilde{\mu}_j - \tilde{\mu}_{j'}| > 0.$$

Let $(\tilde{\mu}, \tilde{\phi})$ be an eigen-pair of H_{Λ} . Let $l \in \Lambda$, and $(\tilde{\mu}^s, \tilde{\phi}^s)$ (depending continuously on s) be an eigenpair of $H_{\Lambda} + sI_{\{l\}}$ with $|s| \leq \frac{d}{10}$ satisfying $\tilde{\mu}^s|_{s=0} = \tilde{\mu}$ and $\tilde{\phi}^s|_{s=0} = \tilde{\phi}$. Then

$$\frac{d\tilde{\mu}^s}{ds} = |\tilde{\phi}(l)|^2 + |s|^2 O\left(\frac{|\Lambda|}{d^2}\right),$$

and hence

$$\tilde{\mu}^s - \tilde{\mu} = s |\tilde{\phi}(l)|^2 + |s|^3 O\left(\frac{|\Lambda|}{d^2}\right).$$

Proof. Let $\{(\tilde{\mu}_j, \tilde{\phi}_j), j \in \Lambda\}$ be the complete set of eigen-pairs of H_{Λ} . Without loss of generality, assume that $\tilde{\mu} = \tilde{\mu}_1$ and $\tilde{\phi} = \tilde{\phi}_1$.

By a standard perturbation argument and (17), one has that for any $|s| \leq \frac{d}{10}$,

$$|\tilde{\mu}^s - \tilde{\mu}_1| \le \frac{d}{2}$$

and

(18)
$$|\tilde{\mu}^s - \tilde{\mu}_{j'}| \ge \frac{d}{2}, j' \in \Lambda \setminus \{1\}.$$

Let

$$\tilde{\phi}^s = \sum_{j \in \Lambda} c_j^s \tilde{\phi}_j.$$

Then

(19)
$$(H_{\Lambda} + sI_{\{l\}})\tilde{\phi}^s = \tilde{\mu}^s \tilde{\phi}^s = \sum_{j \in \Lambda} \tilde{\mu}^s c_j^s \tilde{\phi}_j$$

and

(20)
$$H_{\Lambda}\tilde{\phi}^s = \sum_{j\in\Lambda} c_j^s \tilde{\mu}_j \tilde{\phi}_j.$$

Since $||(H_{\Lambda} + sI_{\{l\}})\tilde{\phi}^s - H_{\Lambda}\tilde{\phi}^s|| = O(s)$, by (18), (19) and (20), one has that for any $j \in \Lambda \setminus \{1\}$,

$$|c_j^s| \le O\left(\frac{s}{d}\right).$$

This implies that $1 - O\left(\frac{|\Lambda|s^2}{d^2}\right) \le (c_1^s)^2 \le 1$. Therefore, one has that

(21)
$$|\tilde{\phi}^s(l)|^2 = |\tilde{\phi}(l)|^2 + O\left(\frac{|\Lambda|s^2}{d^2}\right).$$

By eigenvalue variations and (21), one has that

$$\begin{aligned} \frac{d\tilde{\mu}^s}{ds} &= \langle \tilde{\phi}^s, \frac{d(H_\Lambda + sI_{\{l\}})}{ds} \tilde{\phi}^s \rangle \\ &= |\tilde{\phi}^s(l)|^2 \\ &= |\tilde{\phi}(l)|^2 + O\left(\frac{|\Lambda|s^2}{d^2}\right). \end{aligned}$$

We conclude that

$$\tilde{\mu}^s - \tilde{\mu} = s |\tilde{\phi}(l)|^2 + |s|^3 O\left(\frac{|\Lambda|}{d^2}\right).$$

For any $n = (n_1, n_2, \cdots, n_b) \in \mathbb{Z}^b$, denote by

supp
$$n = \#\{n_k : n_k \neq 0, k = 1, 2, \cdots, b\},\$$

where # denotes the number of elements in a set.

Proof of Theorem 2.1. Let $N = 2 \left| e^{|\log \delta|^{\frac{3}{4}}} \right|$ and $\tilde{S}_1 = S_N$. By Lemma 2.3 (Minami estimate), it suffices to prove that

$$\mathbb{P}(S_1 \cap \tilde{S}_1) \le e^{C|\log \delta|^{\frac{3}{4}}} \delta^{\frac{1}{8}}.$$

Since the total number of (n, j) in (11) and (13) is bounded by $(2N + 1)^{b+1}$, it suffices to prove that for a fixed (n, j) in the set, with probability at most $e^{C|\log \delta|^{\frac{3}{4}}} \delta^{\frac{1}{8}}$, either (12) or (14) holds. Without loss of generality, we only consider probability space $P_{n,j}$ (P for simplicity) such that (15) holds, and

(22)
$$|n \cdot \tilde{\omega} + \tilde{\mu}_j| \le 4\delta^{\frac{1}{8}}.$$

Our goal is to show that

(23)
$$\mathbb{P}(P_{n,j} \cap \tilde{S}_1) \le e^{C|\log \delta|^{\frac{3}{4}}} \delta^{\frac{1}{8}}.$$

Case 1: n = 0. In this case, Wegner estimate implies (23).

Case 2: supp n = 1

Case 2_1 : supp n = 1 and $n = -e_k, k = 1, 2, \dots, b$.

Without loss of generality, assume that $n = (-1, 0, \dots, 0)$. In this case, $n \cdot \tilde{\omega} + \tilde{\mu}_j = -\tilde{\mu}_{\tilde{\alpha}_1} + \tilde{\mu}_j$ and $j \neq \tilde{\alpha}_1$. This is impossible by (16). **Case** 2₂: supp $n = 1, n = e_k, k = 1, 2, \cdots, b$ or $n = re_k, k = 1, 2, \cdots, b$ with

 $|r| \geq 2.$

Without loss of generality, assume that $n = (n_1, 0, \dots, 0)$ with $n_1 = 1$ or $|n_1| \ge 2.$

Denote by $\tilde{\phi}_{\tilde{\alpha}_1}$ and $\tilde{\phi}_j$ eigenvectors of eigenvalues $\tilde{\mu}_{\tilde{\alpha}_1} = \tilde{\omega}_1$ and $\tilde{\mu}_j$. For any $l \in 2B_1$, denote by P_l the probability space such that

(24)
$$|\tilde{\phi}_{\tilde{\alpha}_1}(l)|^2 \ge \frac{4}{5}|\tilde{\phi}_j(l)|^2 + \frac{1}{L^{10}}.$$

By (10), one has that

$$\sum_{\ell \notin 2\tilde{B}_1} |\tilde{\phi}_{\tilde{\alpha}_1}(\ell)|^2 \le e^{-\frac{\gamma_0}{2}L},$$

and hence

(25)
$$\sum_{\ell \in 2\tilde{B}_1} |\tilde{\phi}_{\tilde{\alpha}_1}(\ell)|^2 \ge 1 - e^{-\frac{\gamma_0}{2}L}.$$

By (24) and (25), one has that

$$P_{n,j} \subset \bigcup_{l \in 2\tilde{B}_1} P_l.$$

Take any $l \in 2\tilde{B}_1$ such that (24) holds. Split [0, 1] into N^{4q_1} intervals of size N^{-4q_1} and take any interval $I = [f_1, f_2]$. Define the probability space $P_{l,I}$ to be $P_{l,I} = \{V \in P_l : V_l \in I\}$. Applying Lemma 2.4, one has that if we fix V_{ℓ} , $\ell \in \Lambda_1 \setminus \{l\}$, then for any $f \in I$,

(27)
$$\frac{d\tilde{\mu}_{\tilde{\alpha}_{1}}^{V_{l}=f}}{df} = |\tilde{\phi}_{\tilde{\alpha}_{1}}^{V_{l}=f_{1}}(l)|^{2} + N^{-5q_{1}}O(1),$$

and

(28)
$$\frac{d\tilde{\mu}_{j}^{V_{l}=f}}{df} = |\tilde{\phi}_{j}^{V_{l}=f_{1}}(l)|^{2} + N^{-5q_{1}}O(1),$$

where $\tilde{\mu}_i^{V_l=f}$ is the eigenvalue with the potential $V_l = f$ and fixed V_ℓ , $\ell \in \Lambda_1 \setminus \{l\}$. Obviously,

(29)
$$\frac{d\tilde{\mu}_j^{V_l=f}}{df} \ge 0$$

By (24), (27), (28) and (29), one has that

(30)
$$\left|\frac{d(n \cdot \tilde{\omega}^{V_l=f} + \tilde{\mu}_j^{V_l=f})}{df}\right| \ge \frac{3}{4} |\tilde{\phi}_{\tilde{\alpha}_1}^{V_l=f_1}(l)|^2 - N^{-5q_1}O(1)$$
$$\ge \frac{3}{4L^{10}} - N^{-5q_1}O(1)$$
$$\ge \frac{1}{2L^{10}}.$$

Since $g \in L^{\infty}[0, 1]$, by (22) and (30), one has that for any $l \in 2\tilde{B}_1$,

(31)
$$\mathbb{P}(P_{n,j} \cap P_l \cap \tilde{S}_1) \le O(1)N^{4q_1}L^{10}\delta^{\frac{1}{8}}.$$

By (26) and (31), one has that

(32)
$$\mathbb{P}(P_{n,j} \cap \tilde{S}_1) \le O(1) N^{4q_1} L^{11} \delta^{\frac{1}{8}}.$$

It implies (23) since δ is sufficiently small (depending on L).

Case 3: supp $n \ge 2$

Let n_i be such that $|n_i| = \max_{i \in \{1,2,\dots,b\}} |n_i|$.

Case $3_1: |n_i| \ge 2$

Without loss of generality, assume that $n_i = n_1$. It is easy to see that for any $\ell \in 2\tilde{B}_1$

(33)
$$|\tilde{\phi}_{\tilde{\alpha}_k}(\ell)|^2 \le e^{-\gamma_0 L}, k = 2, 3, \cdots, b.$$

For any $l \in 2\tilde{B}_1$, denote by P_l^1 the probability space such that

(34)
$$|\tilde{\phi}_{\tilde{\alpha}_1}(\ell)|^2 \ge \frac{4}{5} |\tilde{\phi}_j(\ell)|^2 + L^2 \left(\sum_{k=2}^b \tilde{\phi}_{\tilde{\alpha}_k}(\ell)|^2\right) + \frac{1}{L^{10}}$$

By (25), (33) and (34), one has that

$$P_{n,j} \subset \bigcup_{l \in 2\tilde{B}_1} P_l^1.$$

Replacing (26) with (35), and following the proof of Case 2_2 (using Lemma 2.4), we still have (23).

Case 3_2 : $|n_i| \le 1$, namely $n_k = 0, \pm 1, k = 1, 2, \cdots, b$.

In this case, clearly, there exist at least two non-zero n_k , $k = 1, 2, \dots, b$. Without loss of generality, assume that $n_1 \neq 0$ and $n_2 \neq 0$. In this case, if $\tilde{\ell}_j \geq 2L^4$, by (15), one has that for any $\ell \in \bigcup_{k=1}^b 2\tilde{B}_k$,

$$(36) |\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}$$

If $\tilde{\ell}_j \leq 2L^4$, by (9) and (15), one has that either for any $\ell \in 2\tilde{B}_1$,

(37)
$$|\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}$$

or for any $\ell \in 2\tilde{B}_2$,

(38)
$$|\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}.$$

Without loss of generality assume that (37) holds. Therefore, for any $\ell \in 2\tilde{B}_1$,

(39)
$$|\tilde{\phi}_i(\ell)|^2 \le e^{-\gamma_0 L}, i = \tilde{\alpha}_2, \tilde{\alpha}_3, \cdots, \tilde{\alpha}_b, \text{ and } i = j.$$

For any $l \in 2\tilde{B}_1$, denote by P_l^2 the probability space such that

(40)
$$|\tilde{\phi}_{\tilde{\alpha}_1}(l)|^2 \ge L^2 |\tilde{\phi}_j(l)|^2 + L^2 \left(\sum_{k=2}^b \tilde{\phi}_{\alpha_k}(l)|^2\right) + \frac{1}{L^{10}}$$

By (25), (39) and (40), one has that

(41)
$$P_{n,j} \subset \bigcup_{l \in 2\tilde{B}_1} P_l^2.$$

Replacing (26) with (41), and following the proof of Case 2_2 or Case 3_1 (also using Lemma 2.4), we have (23).

2.2. Spacing of the diagonals. For the purpose of nonlinear analysis, it suffices to work with scales $|\log \delta|^s$, s > 1. So let $\Lambda_2^s = [-2 \lfloor |\log \delta|^s \rfloor, 2 \lfloor |\log \delta|^s \rfloor]$. Denote by S_2^s the probability space such that there exist $\tilde{\omega}^{\Lambda_2^s}$ satisfying (10), and $m \in \mathbb{Z}^b$ with $|m| \leq |2 \log \delta|^s$, either $|m_k| \geq 2$ for some $k \in \{1, 2, \dots, b\}$ or supp $m \geq 3$, and $j, j' \in \Lambda_2^s$ satisfying

(42)
$$|m \cdot \tilde{\omega}^{\Lambda_2^s} + \tilde{\mu}_j^{\Lambda_2^s} - \tilde{\mu}_{j'}^{\Lambda_2^s}| \le 4\delta^{\frac{1}{8}},$$

and the eigenvectors corresponding to $\tilde{\mu}_{j}^{\Lambda_{2}^{s}}$ and $\tilde{\mu}_{j'}^{\Lambda_{2}^{s}}$ satisfy that there exist $\tilde{\ell}_{j}^{\Lambda_{2}^{s}} \in \Lambda_{2}^{s}$ and $\tilde{\ell}_{j'}^{\Lambda_{2}^{s}} \in \Lambda_{2}^{s}$ such that for any $\ell \in \Lambda_{2}^{s}$,

(43)
$$|\tilde{\phi}_{j}^{\Lambda_{2}^{s}}(\ell)| \leq C_{0}(1+|\tilde{\ell}_{j}^{\Lambda_{2}^{s}}|)^{q_{0}}e^{-\gamma_{0}|\ell-\tilde{\ell}_{j}^{\Lambda_{2}^{s}}|}, |\tilde{\phi}_{j'}^{\Lambda_{2}^{s}}(\ell)| \leq C_{0}(1+|\tilde{\ell}_{j'}^{\Lambda_{2}^{s}}|)^{q_{0}}e^{-\gamma_{0}|\ell-\tilde{\ell}_{j'}^{\Lambda_{2}^{s}}|}.$$
We about mention that we allow $\dot{a}=\dot{a}'$

We should mention that we allow j = j'.

Theorem 2.5. For small δ , we have

$$\mathbb{P}(S_2^s) \le (|\log \delta|)^{Cs} \delta^{\frac{1}{8}}.$$

Proof. Let $N = 2 \lfloor |\log \delta|^s \rfloor$ and $\tilde{S}_2^s = S_N$. By Lemma 2.3 (Minami estimate), it suffices to prove that

 $\mathbb{P}(S_2^s \cap \tilde{S}_2^s) \le (|\log \delta|)^{Cs} \delta^{\frac{1}{8}}.$

Denote by $P_{m,j,j'}$ the probability space such that (10) and (42) hold. Let m_i be such that $|m_i| = \max\{|m_k|, k = 1, 2, \dots, b\}$. Without loss of generality, assume that $m_i = m_1$. Let us prove the case $|m_1| \ge 2$ first. Without loss of generality, assume that $m_1 \ge 0$. It is easy to see that for any $\ell \in 2\tilde{B}_1$,

(44)
$$|\tilde{\phi}_{\tilde{\alpha}_k}(\ell)|^2 \le e^{-\gamma_0 L}, \ k = 2, 3, \cdots, b.$$

For any $l \in 2\tilde{B}_1$, denote by P_l^3 the probability space such that

(45)
$$|\tilde{\phi}_{\tilde{\alpha}_1}(l)|^2 \ge \frac{4}{5} |\tilde{\phi}_{j'}(l)|^2 + L^2 (\sum_{k=2}^b \tilde{\phi}_{\tilde{\alpha}_k}(l)|^2) + \frac{1}{L^{10}}$$

By (25), (44) and (45), one has that

(46)
$$P_{m,j,j'} \subset \bigcup_{l \in 2\tilde{B}_1} P_l^3.$$

Now the proof follows from Lemma 2.4, which is similar to the proof of Case 2_2 or Case 3_1 . Here are the details. Take any $l \in 2\tilde{B}_1$ such that (45) holds. Split [0, 1] into N^{4q_1} intervals of size N^{-4q_1} and take any interval $I = [f_1, f_2]$. Denote by the probability space $P_{l,I}^3 = \{V \in P_l : V_l \in I\}$. Applying Lemma 2.4, one has that if we fix $V_{\ell}, \ell \in \Lambda_1 \setminus \{l\}$, then for any $f \in I$,

(47)
$$\frac{d\tilde{\mu}_{\tilde{\alpha}_{1}}^{V_{l}=f}}{df} = |\tilde{\phi}_{\tilde{\alpha}_{1}}^{V_{l}=f_{1}}(l)|^{2} + N^{-5q_{1}}O(1),$$

and

(48)
$$\frac{d\tilde{\mu}_{j'}^{V_l=f}}{df} = |\tilde{\phi}_{j'}^{V_l=f_1}(l)|^2 + N^{-5q_1}O(1).$$

By (47), (48) and (29), one has that

(49)
$$\frac{d(m \cdot \tilde{\omega}^{V_l=f} + \tilde{\mu}_j^{V_l=f} - \tilde{\mu}_{j'}^{V_l=f})}{df} \ge \frac{3}{4} |\tilde{\phi}_{\tilde{\alpha}_1}^{V_l=f_1}(l)|^2 + N^{-5q_1}O(1) \\\ge \frac{3}{4L^{10}} + N^{-5q_1}O(1) \\\ge \frac{1}{2L^{10}}.$$

Now the proof follows that of Case 2_2 or Case 3_1 of Theorem 2.1.

Let us proceed to the case supp $m \geq 3$. Without loss of generality, assume that $m_i \neq 0, i = 1, 2, 3$. In this case, if $\tilde{\ell}_j \geq 2L^4$, by (43), one has that for any $\ell \in \bigcup_{k=1}^b 2\tilde{B}_k$,

(50)
$$|\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}.$$

If $\tilde{\ell}_j \leq 2L^4$, by (9) and (43), one has that either for any $\ell \in 2\tilde{B}_1 \cup 2\tilde{B}_2$,

(51)
$$|\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}$$

or for any $\ell \in 2\tilde{B}_2 \cup 2\tilde{B}_3$,

(52)
$$|\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}$$

or for any $\ell \in 2\tilde{B}_1 \cup 2\tilde{B}_3$,

(53)
$$|\tilde{\phi}_j(\ell)| \le e^{-\gamma_0 L}$$

Clearly, (50)-(53) also hold for j'. Therefore, we have that there exists $i \in \{1, 2, 3\}$, such that for any $m \in \{\tilde{\alpha}_1, \tilde{\alpha}_2, \cdots, \tilde{\alpha}_b\} \setminus \{\tilde{\alpha}_i\}$ and m = j, j',

(54)
$$|\tilde{\phi}_m(\ell)| \le e^{-\gamma_0 L}, \ell \in 2\tilde{B}_i.$$

Now the proof follows that of Case 3_2 of Theorem 2.1.

Denote by \hat{S}_2^s the probability space such that there exist $\tilde{\omega}^{\Lambda_2^s}$ satisfying (10), $m \in \mathbb{Z}^b$ with $\sum_{k=1}^b m_k \neq 0$, $|m_k| \leq 1, k = 1, 2, \cdots, b$, and $j \in \Lambda_2^s, j' \in \Lambda_2^s$ satisfying

(55)
$$|m \cdot \tilde{\omega}^{\Lambda_2^s} + \tilde{\mu}_j^{\Lambda_2^s} - \tilde{\mu}_{j'}^{\Lambda_2^s}| \le 4\delta^{\frac{1}{8}}.$$

Theorem 2.6. For small δ , we have

$$\mathbb{P}(\hat{S}_2^s) \le (|\log \delta|)^{Cs} \delta^{\frac{1}{8}}.$$

Proof. Again, let $N = 2 \lfloor |\log \delta|^s \rfloor$ and $\tilde{S}_2^s = S_N$. By Lemma 2.3 (Minami estimate), it suffices to prove that

$$\mathbb{P}(\hat{S}_2^s \cap \tilde{S}_2^s) \le (|\log \delta|)^{Cs} \delta^{\frac{1}{8}}.$$

Since $\sum_{k=1}^{b} m_k \neq 0$, there exists $l \in [-N, N]$, such that

(56)
$$|(\sum_{k=1}^{b} m_k |\tilde{\phi}_{\tilde{\alpha}_k}(l)|^2) + |\tilde{\phi}_j(l)|^2 - |\tilde{\phi}_{j'}(l)|^2| \ge \frac{1}{N^{10}}.$$

Now we use Lemma 2.4 to conclude as in the proof of Case 2_2 of Theorem 2.1. \Box

Remark 7. We could replace the constant 4 in (12), (14), (42) and (55) with any fixed constant and all theorems in this section still hold.

3. One dimensional random Schrödinger operators

We now proceed to the infinite volume random Schrödinger operator,

$$H = -\Delta + V.$$

Let $\{\varphi_j^V\}_j$ be the eigen-basis and assume that ι_j^V satisfies

$$|\varphi_j^V(\iota_j^V)| = \max_{x \in \mathbb{Z}} |\varphi_j^V(x)|.$$

(If it is not unique, one may choose a maximum arbitrarily.) We have

Theorem 3.1. (See e.g., [16] or [18, sect. 1.6]) There exist some q > 0 and $\gamma_1 > 0$ such that, with probability 1,

(57)
$$|\varphi_j^V(\ell)| \le C_V (1+|\iota_j^V|)^q e^{-\gamma_1 |\ell - \iota_j^V|},$$

where $\mathbb{E}(C_V) < \infty$.

Remark 8. • Recall that ι_i^V is called the localization center.

• γ_1 and q only depend on the distribution g. γ_1 can be arbitrarily close to the Lyapunov exponent.

By Theorem 3.1 and following a similar proof of Theorem 7.1 in [11], one has the following Lemma concerning the localization centers.

Lemma 3.2. For any ϵ , there exist \mathcal{V}_{ϵ} with $\mathbb{P}(\mathcal{V}_{\epsilon}) > 1 - \epsilon$ and a constant l_{ϵ} such that the following holds. For any $V \in \mathcal{V}_{\epsilon}$ and $k \in [-L^4, L^4]$ with $L \ge l_{\epsilon}$,

(58)
$$(1-\epsilon)L \le \#\{j : \iota_j^V \in [k, k+L]\} \le (1+\epsilon)L.$$

See appendix B for a proof.

Basing on (58), one may (re)label the eigenfunctions so that if j > j', then the localization centers of the corresponding eigenfunctions ϕ_j^V and $\phi_{j'}^V$ satisfy $\ell_{j'}^V \ge \ell_j^V$. The construction of such a map is presented in appendix A. Here after relabelling, we use the notations ϕ_j^V instead of φ_j^V and ℓ_j^V instead of ι_j^V . Recall that μ_j^V is the eigenvalue corresponding to eigenfunction ϕ_j^V . When there is no ambiguity, we omit the dependence on V.

Below we summarize properties of the eigenfunction basis in this labelling.

Lemma 3.3. There exist q > 0 and $\gamma > 0$ such that for any ϵ , there exist \mathcal{V}_{ϵ} with $\mathbb{P}(\mathcal{V}_{\epsilon}) > 1 - \epsilon$ and constants C_{ϵ} and ℓ_{ϵ} such that for any $V \in \mathcal{V}_{\epsilon}$, the following statements hold:

• for any $\ell \in \mathbb{Z}$,

(59)
$$|\phi_j^V(\ell)| \le C_\epsilon (1+|\ell_j^V|)^q e^{-\gamma|\ell-\ell_j^V|},$$

• for any $|\ell_i^V| \ge \ell_{\epsilon}$,

$$|\ell_j^V - j| \le \epsilon |j|,$$

• for any $k \in [-L^4, L^4]$,

$$(1 - \epsilon)L \le \#\{j : \ell_j^V \in [k, k + L]\} \le (1 + \epsilon)L.$$

Lemma 3.4. Choose any $V \in \mathcal{V}_{\epsilon} \cap S_{2N}$. Let $\Lambda = [-2N, 2N]$. Consider two distinct eigen-pairs (μ_j, ϕ_j) and $(\mu_{j'}, \phi_{j'}), |j|, |j'| \leq N$, of $H = -\Delta + V$. Then there exist two distinct eigen-pairs $(\tilde{\mu}_{\tilde{j}}^{\Lambda}, \tilde{\phi}_{\tilde{j}}^{\Lambda})$ and $(\tilde{\mu}_{\tilde{j}'}^{\Lambda}, \tilde{\phi}_{\tilde{j}'}^{\Lambda}), \tilde{j} \in \Lambda, \tilde{j}' \in \Lambda$ of H_{Λ} such that

$$|\mu_j - \tilde{\mu}_{\tilde{j}}^{\Lambda}| \le e^{-\frac{\gamma}{2}N}, |\mu_{j'} - \tilde{\mu}_{\tilde{j}'}^{\Lambda}| \le e^{-\frac{\gamma}{2}N},$$

and

(60)

(61)

$$||\phi_j - \tilde{\phi}^{\Lambda}_{\tilde{j}}|| \le e^{-\frac{\gamma}{2}N}, ||\phi_{j'} - \tilde{\phi}^{\Lambda}_{\tilde{j}'}|| \le e^{-\frac{\gamma}{2}N}.$$

Proof. By (60), one has that $|\ell_j| \leq (1+\epsilon)N$ and $|\ell_{j'}| \leq (1+\epsilon)N$. Then

$$\sum_{|\ell| \ge 2N+1} |\phi_j(\ell)|^2 \le e^{-\frac{3\gamma}{2}N}, \sum_{|\ell| \ge 2N+1} |\phi_{j'}(\ell)|^2 \le e^{-\frac{3\gamma}{2}N},$$

and

(62)
$$||H_{\Lambda}\phi_{j} - \mu_{j}\phi_{j}|| \le e^{-\frac{3\gamma}{4}N}, ||H_{\Lambda}\phi_{j'} - \mu_{j}\phi_{j'}|| \le e^{-\frac{3\gamma}{4}N}.$$

Therefore, there exist $j \in \Lambda$ and $j' \in \Lambda$ such that

$$|\mu_j - \tilde{\mu}_{\tilde{j}}^{\Lambda}| \le e^{-\frac{3\gamma}{4}N}, |\mu_{j'} - \tilde{\mu}_{\tilde{j}'}^{\Lambda}| \le e^{-\frac{3\gamma}{4}N}.$$

Since $V \in S_{2N}$, one has that for any distinct j_1 and j_2 in Λ ,

$$|\tilde{\mu}_{j_1}^{\Lambda} - \tilde{\mu}_{j_2}^{\Lambda}| \ge \frac{1}{(2N)^{q_1}},$$

and hence for any $m \neq \tilde{j}$ (or $m \neq \tilde{j}'$),

(63)
$$|\tilde{\mu}_m^{\Lambda} - \mu_j| \ge \frac{1}{2^{q_1+1}N^{q_1}} (\text{or } |\tilde{\mu}_m^{\Lambda} - \mu_{j'}| \ge \frac{1}{2^{q_1+1}N^{q_1}}).$$

Let ϕ_m and $\tilde{\mu}_m$ (as usual, for simplicity we have dropped the dependence on Λ from $\tilde{\phi}_m^{\Lambda}$ and $\tilde{\mu}_m^{\Lambda}$, $m \in \Lambda$), be the eigen-pairs of H_{Λ} . Let

(64)
$$I_{\Lambda}\phi_j = \sum_{m\in\Lambda} c_m \tilde{\phi}_m.$$

From (62) and (64), one has that

(65)
$$H_{\Lambda}\phi_j = \sum_{m\in\Lambda} \tilde{\mu}_m c_m \tilde{\phi}_m,$$

and

(66)
$$||\sum_{m\in\Lambda} \tilde{\mu}_m c_m \tilde{\phi}_m - \sum_{m\in\Lambda} \mu_j c_m \tilde{\phi}_m || = O(1)e^{-\frac{3\gamma}{4}N}.$$

By (63), (65) and (66), one has that for any $m \neq \tilde{j}$,

$$|c_m| \le O(1)N^{q_1} e^{-\frac{3\gamma}{4}N}$$

Therefore, $1 - O(1)N^{3q_1}e^{-\frac{3\gamma}{2}N} \le c_{\tilde{j}}^2 \le 1$. We conclude that

$$||\phi_j - \tilde{\phi}_{\tilde{j}}|| \le ||\phi_j - I_\Lambda \phi_j|| + ||I_\Lambda \phi_j - \tilde{\phi}_{\tilde{j}}|| \le e^{-\frac{\gamma}{2}N}.$$

Similarly,

$$||\phi_{j'} - \tilde{\phi}_{\tilde{j}'}|| \le e^{-\frac{\gamma}{2}N}.$$

Since ϕ_j and $\phi_{j'}$ are ortho-normal eigenfunctions, we have that $\tilde{j} \neq \tilde{j'}$.

We now state the conclusion:

Theorem 3.5. For any $\epsilon > 0$, there exists l_{ϵ} such that the following statements hold. Fix any $L \ge \ell_{\epsilon}$ and $\beta_k \in \mathbb{Z}$, $k = 1, 2, \dots, b$ satisfying $10L \le |\beta_k| \le L^3$ and $|\beta_k - \beta_{k'}| \ge 10L$, for any distinct $k, k' \in \{1, 2, \dots, b\}$, there exists a probability space X_{ϵ} with $\mathbb{P}(X_{\epsilon}) \ge 1 - \epsilon$ and $\delta_0 > 0$ (depending on g, ϵ and L) such that for any $V \in X_{\epsilon}$ and $0 < \delta \le \delta_0$,

(67)
$$|\phi_j(\ell)| \le C_\epsilon (1+|\ell_j|)^q e^{-\gamma|\ell-\ell_j|},$$

(2) for any $|\ell_j| \ge l_{\epsilon}$,

(68)
$$|\ell_j - j| \le \epsilon |j|,$$

(3) for large N (depending on ϵ), $|j|, |j'| \leq N$ and $j \neq j'$,

(69)
$$|\mu_j - \mu_{j'}| \ge \frac{1}{2^{q_1 + 1} N^{q_1}},$$

and

(70)
$$|\mu_j| \ge \frac{1}{2N^{q_1}},$$

(4) for any eigenfunction ϕ_{α_k} with $\ell_{\alpha_k} \in B_k = \{l \in \mathbb{Z} : |l - \beta_k| \leq L\}, k = 1, 2, \cdots, b$, we have that for any $(n, j) \in [-e^{|\log \delta|^{\frac{3}{4}}}, e^{|\log \delta|^{\frac{3}{4}}}]^{b+1} \setminus \{(-e_k, \alpha_k)\}_{k=1}^b$,

(71)
$$|n \cdot \omega^{(0)} + \mu_j| \ge 2\delta^{\frac{1}{8}},$$
where $\omega^{(0)} = (\omega_1^{(0)}, \cdots, \omega_b^{(0)}) = (\mu_{\alpha_1}, \cdots, \mu_{\alpha_b}), \text{ and for any } (n, j) \in [-e^{|\log \delta|^{\frac{3}{4}}}, e^{|\log \delta|^{\frac{3}{4}}}]^{b+1} \setminus \{(e_k, \alpha_k)\}_{k=1}^b,$
(72)
$$|-n \cdot \omega^{(0)} + \mu_j| \ge 2\delta^{\frac{1}{8}},$$

(5) for any $\theta \in \mathbb{R}$, there are at most b vertices $(n, j) \in [-|\log \delta|^s, |\log \delta|^s]^{b+1}$, such that

(73)
$$|(n \cdot \omega^{(0)} + \theta) + \mu_j| \le \delta^{\frac{1}{8}},$$

for any $\theta \in \mathbb{R}$, there are at most b vertices $(n, j) \in [-|\log \delta|^s, |\log \delta|^s]^{b+1}$, such that

(74)
$$|-(n\cdot\omega^{(0)}+\theta)+\mu_j|\leq \delta^{\frac{1}{8}}.$$

Proof. By Lemmas 2.2 (Wegner estimate), 2.3 (Minami estimate), 3.3, 3.4 and Borel-Cantelli type arugments, we have (67)-(70).

We apply the Theorems (with Remark 7) in the previous section with $\tilde{B}_k = B_k$, $k = 1, 2, \dots, b$ and $\delta = 2^{-n}$, $n = 1, 2, \dots$. Then by Borel-Cantelli type arugments, we have that there exists X_{ϵ} with $\mathbb{P}(X_{\epsilon}) \geq 1 - \epsilon$ and $X_{\epsilon} \cap (\tilde{S}_1 \cup S_2^s \cup \hat{S}_2^s) = \emptyset$ for any small δ .

(71) and (72) follow from Lemma 3.4 with $N = \left\lfloor e^{\left|\log \delta\right|^{\frac{3}{4}}} \right\rfloor$.

The proof of (73) and (74) takes more time. Without loss of generality, we only prove (73). Assume that there are $(n^{(m)}, j^{(m)}) \in [-|\log \delta|^s, |\log \delta|^s]^{b+1}, m = 1, 2, \dots, b+1$, satisfying

(75)
$$|(n^{(m)} \cdot \omega^{(0)} + \theta) + \mu_{j^{(m)}}| \le \delta^{\frac{1}{8}}.$$

By Lemma 3.4, one has that there exist $\tilde{\omega}^{\Lambda_2^s}$ satisfying (10) and $(n^{(m)}, \tilde{j}^{(m)}) \in [-|\log \delta|^s, |\log \delta|^s]^b \times [-2|\log \delta|^s, 2|\log \delta|]^s, m = 1, 2, \cdots, b+1$ such that

(76)
$$|(n^{(m)} \cdot \tilde{\omega}^{\Lambda_2^s} + \theta) + \tilde{\mu}_{\tilde{j}^{(m)}}^{\Lambda_2^s}| \le 2\delta^{\frac{1}{8}}$$

We can assume that $n^{(m)}$, $m = 1, 2, \dots, b+1$ are distinct, otherwise Lemma 2.3 (Minami estimate) gives the proof.

Choose any $m_1, m_2 \in \{1, 2, \dots, b+1\}$ such that (76) holds. When $|n_k^{(m_1)} - n_k^{(m_2)}| \ge 2$ for some $k \in \{1, 2, \dots, b\}$ or supp $(n^{(m_1)} - n^{(m_2)}) \ge 3$, the proof follows from Theorem 2.5. When $\sum_{k=1}^{b} (n_k^{(m_1)} - n_k^{(m_2)}) \ne 0$ and $|n_k^{(m_1)} - n_k^{(m_2)}| \le 1$, $k = 1, 2, \dots, b$, the proof follows from Theorem 2.6.

WENCAI LIU AND W.-M. WANG

So the only exceptional case is when for $all m_1, m_2 \in \{1, 2, \dots, b+1\}, n^{(m_1)}$ and $n^{(m_2)}$ satisfy supp $(n^{(m_1)} - n^{(m_2)}) = 2$, $\sum_{k=1}^{b} (n_k^{(m_1)} - n_k^{(m_2)}) = 0$ and $n_k^{(m_1)} - n_k^{(m_2)} = \pm 1, 0, k = 1, 2, \dots, b$. We will show that this is not possible. Shifting $n^{(m)}$ by $n^{(1)}$, one may assume that $n^{(1)} = (0, 0, \dots, 0)$. When b = 2, it is obvious. So let $b \geq 3$. Without loss of generality, assume that $n^{(2)} = (1, -1, 0, \dots, 0)$. Thus either $n_1^{(m)} = 1$ or $n_2^{(m)} = -1$ for all $m \in \{3, 4, \dots, b+1\}$. Without loss of generality, assume that $n^{(3)} = (1, 0, -1, 0, 0, \dots, 0)$. Therefore, for all $m \in \{2, 3, \dots, b+1\}, n_1^{(m)} = 1$. This contradicts with $n^{(m)}, m \in \{1, 2, \dots, b+1\}$ being distinct.

4. LARGE DEVIATION THEOREM

Assume that \tilde{H} is an operator on $\ell^2(\mathbb{Z}^{b+1} \times \{0,1\})$, Töplitz with respect to $n \in \mathbb{Z}^b$. We now write \mathbb{Z}^{b+1} interchangeably with $\mathbb{Z}^b \times \mathbb{Z}$. Assume that there exist functions $h_{r,r'}(n, j, j')$, $r, r' \in \{0, 1\}$, on $\mathbb{Z}^b \times \mathbb{Z} \times \mathbb{Z}$, such that for any $u_r(n, j)$, $r \in \{0, 1\}$ and $(n, j) \in \mathbb{Z}^{b+1}$,

(77)
$$(\tilde{H}u)_r(n,j) := \sum_{(n',j') \in \mathbb{Z}^b \times \mathbb{Z}, \, r' \in \{0,1\}} h_{r,r'}(n-n',j,j')u_{r'}(n',j').$$

Assume that there exist $C_1 > 0$ and $c_1 > 0$ such that

(78)
$$|h_{r,r'}(n,j,j')| \le C_1 e^{-c_1(|n|+|j|+|j'|)}.$$

Let $D(\theta)$ be a family of operators from \mathbb{R} to $Op[\ell^2(\mathbb{Z}^{b+1} \times \{0,1\})]$:

(79)
$$D(\theta) = \begin{bmatrix} D+ & 0\\ 0 & D_{-} \end{bmatrix}$$

where $D_{\pm} = \text{diag}(\pm (n \cdot \omega + \theta) + \mu_j), (n, j) \in \mathbb{Z}^{b+1}$. Define $T = T(\theta)$: $\mathbb{R} \to \text{Op}[\ell^2(\mathbb{Z}^{b+1} \times \{0, 1\})]$ as

(80)
$$T(\theta) = D(\theta) + \delta \tilde{H}.$$

Denote by Q_N an elementary region of size N centered at 0, which is one of the following regions,

$$Q_N = [-N, N]^{b+1}$$

or

$$Q_N = [-N, N]^{b+1} \setminus \{ n \in \mathbb{Z}^{b+1} : n_i \varsigma_i 0, 1 \le i \le b+1 \},\$$

where for $i = 1, 2, \dots, b+1$, $\varsigma_i \in \{<, >, \emptyset\}$ and at least two ς_i are not \emptyset .

Denote by \mathcal{E}_N^0 the set of all elementary regions of size N centered at 0. Let \mathcal{E}_N be the set of all translates of elementary regions with center at 0, namely,

$$\mathcal{E}_N := \{ n + Q_N : n \in \mathbb{Z}^{b+1}, Q_N \in \mathcal{E}_N^0 \}.$$

For simplicity, we call elements in \mathcal{E}_N elementary regions. Let $Q_N(j_0) = \{(n, j) \in \mathbb{Z}^b \times \mathbb{Z} : (n, j - j_0) \in Q_N\}.$

The width of a subset $\Lambda \subset \mathbb{Z}^{b+1}$, is defined as the maximum of $M \in \mathbb{N}$ such that for any $n \in \Lambda$, there exists $\hat{M} \in \mathcal{E}_M$ such that

$$n \in M \subset \Lambda$$

and

dist
$$(n, \Lambda \backslash \hat{M}) \ge M/2$$
.

A generalized elementary region is defined to be a subset $\Lambda \subset \mathbb{Z}^{b+1}$ of the form

$$\Lambda := R \backslash (R+z),$$

where $z \in \mathbb{Z}^{b+1}$ is arbitrary and R is a rectangle,

$$R = \{ (n_1, n_2, \cdots, n_{b+1}) \in \mathbb{Z}^d : |n_1 - n_1'| \le M_1, \cdots, |n_{b+1} - n_{b+1}'| \le M_{b+1} \}.$$

For $\Lambda \subset \mathbb{Z}^{b+1}$, we introduce its diameter,

$$\operatorname{diam}(\Lambda) = \sup_{n,n' \in \Lambda} |n - n'|.$$

Denote by \mathcal{R}_N all generalized elementary regions with diameters less than or equal to N. Denote by \mathcal{R}_N^M all generalized elementary regions in \mathcal{R}_N with width larger than or equal to M.

With a slight abuse of notation, we also use \mathcal{E}_N , \mathcal{E}_N^0 , Q_N , $Q_N(j_0)$, \mathcal{R}_N and \mathcal{R}_N^M to denote $\mathcal{E}_N \times \{0,1\}$, $\mathcal{E}_N^0 \times \{0,1\}$, $Q_N \times \{0,1\}$, $Q_N(j_0) \times \{0,1\}$, $\mathcal{R}_N \times \{0,1\}$ and $\mathcal{R}_N^M \times \{0,1\}$ respectively. Similarly for any $\Lambda \subset \mathbb{Z}^{b+1}$, denote by \mathcal{R}_Λ the restriction to $\Lambda \times \{0,1\}$.

We say that T (given by (80)) satisfies the large deviation theorem (LDT) at scale N with parameter \tilde{c}_1 if there exists a subset $\Theta_N \subset \mathbb{R}$ such that

$$\operatorname{Leb}(\Theta_N) \le e^{-N^{\frac{1}{30}}},$$

and for any $j_0 \in [-2N, 2N]$, $Q_N \in \mathcal{E}_N^0$, and $\theta \notin \Theta_N$,

(81)
$$||(R_{Q_N(j_0)}T(\theta)R_{Q_N(j_0)})^{-1}|| \le e^{N^{\frac{9}{10}}}$$

and for any (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} \ge \frac{N}{10}$,

(82)
$$|(R_{Q_N(j_0)}T(\theta)R_{Q_N(j_0)})^{-1}(n,j;n',j')| \le e^{-\tilde{c}_1 \max\{|n-n'|,|j-j'|\})}.$$

Let K_1 be a large constant depending only on b. Let $K = K_1^{100}, K_2 = K_1^5$.

Theorem 4.1. Assume that ω satisfies

(1) $|\omega_k - \mu_{\alpha_k}| \le C_2 \delta, \ k = 1, 2, \cdots, b;$

(2) for any fixed $N \ge (\log \frac{1}{\delta})^K$, and any \tilde{N} with $(\log \frac{1}{\delta})^K \le \tilde{N} \le N$ and $0 \ne |n| \le 2\tilde{N}$,

(83)
$$|n \cdot \omega| \ge e^{-\tilde{N}^{\frac{1}{K_2}}}$$

and for any $|j| \leq 3\tilde{N}, |j'| \leq 3\tilde{N}, |n| \leq 2\tilde{N}$ with $(n, j - j') \neq 0$,

(84)
$$|n \cdot \omega - \mu_j + \mu_{j'}| \ge e^{-\tilde{N}^{\frac{1}{K_2}}}.$$

Then for small enough δ , the LDT holds at any scale $\tilde{N} \leq N$ with parameter $\frac{1}{2}c_1$.

- **Remark 9.** (1) The proof of Theorem 4.1 uses ideas from the work of Bourgain, Goldstein and Schlag [7]. The implementation follows, however, the more recent paper [23], which streamlined and quantified some of their arguments.
 - (2) From (83) and (84), in order to have LDT at all scales, we only need to remove measure (with respect to ω) less than $e^{-\frac{1}{2}(\log \frac{1}{\delta})^{\frac{K}{K_2}}} \leq e^{-(\log \frac{1}{\delta})^{K_1^{90}}} \ll \delta$.
 - (3) Theorem 4.1 holds for any parameter \tilde{c}_1 with $\tilde{c}_1 < c_1$.

4.1. Preparations.

Lemma 4.2. [5, Prop. 14.1] Let T(x) be a $N \times N$ matrix function of a parameter $x \in [-\tau, \tau]$ satisfying the following conditions:

(i) T(x) is real analytic in $x \in [-\tau, \tau]$ and has a holomorphic extension to

$$\mathcal{D}_{\tau,\tau_1} = \{ z : |\Re z| \le \tau, |\Im z| \le \tau_1 \}$$

satisfying

(85)
$$\sup_{z \in \mathcal{D}_{\tau,\tau_1}} \|T(z)\| \le B_1, B_1 \ge 1.$$

(ii) For all $x \in [-\tau, \tau]$, there is a subset $\Lambda \subset [1, N]$ with

$$|\Lambda| \le M,$$

and

(86)
$$||(R_{[1,N]\setminus\Lambda}T(x)R_{[1,N]\setminus\Lambda})^{-1}|| \le B_2, B_2 \ge 1.$$

(iii)

(87) Leb{
$$x \in [-\tau, \tau]$$
: $||T^{-1}(x)|| \ge B_3$ } $\le 10^{-3} \tau_1 (1+B_1)^{-1} (1+B_2)^{-1}$.

Let

(88)
$$0 < \epsilon \le (1 + B_1 + B_2)^{-10M}.$$

Then

(89) Leb
$$\left\{ x \in \left[-\tau/2, \tau/2 \right] : \|T^{-1}(x)\| \ge \epsilon^{-1} \right\} \le C \tau e^{-c \left(\frac{\log \epsilon^{-1}}{M \log(B_1 + B_2 + B_3)} \right)},$$

where C and c are absolute constants.

To apply Lemma 4.2, we also need to introduce semi-algebraic sets. A set $S \subset \mathbb{R}^d$ is called *semi-algebraic* if it is a finite union of sets defined by a finite number of polynomial equalities and inequalities. More precisely, let $\{P_1, \dots, P_s\} \subset \mathbb{R}[x_1, \dots, x_d]$ be a family of real polynomials whose degrees are bounded by κ . A (closed) semi-algebraic set S is given by an expression

(90)
$$\mathcal{S} = \bigcup_{l} \bigcap_{\ell \in \mathcal{L}_{l}} \left\{ x \in \mathbb{R}^{d} : P_{\ell}(x)\varsigma_{l\ell} 0 \right\},$$

where $\mathcal{L}_l \subset \{1, \dots, s\}$ and $\varsigma_{l\ell} \in \{\geq, \leq, =\}$. Then we say that \mathcal{S} has degree at most $s\kappa$. In fact, the degree of \mathcal{S} which is denoted by deg(\mathcal{S}), means the smallest $s\kappa$ over all representations as in (90).

Following are some basic properties of these sets. They are special cases of that in [2], and restated in [5].

Lemma 4.3. [5, Theorem 9.3] [2, Theorem 1] Let $S \subset [0,1]^d$ be a semi-algebraic set of degree B. Then the number of connected components of S does not exceed $(1+B)^{C(d)}$.

Lemma 4.4. [5, Theorem 9.3] [2, Theorem 1] Let $\mathcal{S} \subset [0,1]^{d_1+d_2}$ be a semialgebraic set of degree B. Let $(x,y) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$. Then the projection $\operatorname{proj}_{x_1}(S)$ is a semi-algebraic set of degree at most $(1+B)^{C(d_1,d_2)}$.

4.2. Large deviation theorem for small scales: $N \leq (\log \frac{1}{\delta})^{10}$.

Proof. In this case, let

$$\hat{\Theta}_N = \{ \theta \in \mathbb{R} : \text{ there exists } (n,j) \in [-N,N]^b \times [-3N,3N] \text{ such that either} \\ |(n \cdot \omega + \theta) + \mu_j| \le 2e^{-N\frac{1}{20}} \text{ or } |(-n \cdot \omega - \theta) + \mu_j| \le 2e^{-N\frac{1}{20}} \}.$$

Clearly, $\operatorname{Leb}(\tilde{\Theta}_N) \leq N^{C(b)} e^{-N^{\frac{1}{20}}}$. When $N \leq (\log \frac{1}{\delta})^{10}$, δ is much smaller than $e^{-N^{\frac{1}{20}}}$. Now (81) and (82) follow from standard perturbation arguments.

4.3. Large deviation theorem for intermediate scales: $(\log \frac{1}{\delta})^{10} \leq N \leq (\log \frac{1}{\delta})^{K}$.

Proof. Let Θ_N be such that at least one of (81) and (82) does not hold for $\theta \in \Theta_N$. Choose any $N \in [(\log \frac{1}{\delta})^{10}, (\log \frac{1}{\delta})^K]$.

Assume that (81) and (82) do not hold for some θ . Then we must have for some $(n, j) \in [-N, N]^b \times [-3N, 3N]$, either $|\theta + n \cdot \omega^{(0)} + \mu_j| \leq C\delta$ or $|\theta + n \cdot \omega^{(0)} - \mu_j| \leq C\delta$. Otherwise, standard perturbation arguments yield that for any j_0 with $|j_0| \leq 2N$,

$$||(R_{Q_N(j_0)}TR_{Q_N(j_0)})^{-1}|| \le \frac{1}{C\delta} \le e^{N^{\frac{9}{10}}}$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} \ge \frac{N}{10}$,

$$|(R_{Q_N(j_0)}TR_{Q_N(j_0)})^{-1}(n,j;n',j')| \le e^{-\tilde{c}_1 \max\{n-n'|,|j-j'|\}|}$$

where \tilde{c}_1 can be any constant smaller than c_1 .

Therefore, we can restrict θ to be in $10^{b+1}N^{b+1}$ intervals of size $C\delta$. Denote all the intervals by $\{I_i\}$ and take one of them, I_0 , into consideration. Without loss of generality, assume that I_0 comes from the + sector, namely

$$I_0 = \{ \theta : |\theta + n_0 \cdot \omega^{(0)} + \mu_{j_0}| \le C\delta \text{ and } (n_0, j_0) \in [-N, N]^b \times [-3N, 3N] \}.$$

For the – sector $I_0 = \{\theta : |\theta + n_0 \cdot \omega^{(0)} - \mu_{j_0}| \leq C\delta \text{ and } (n_0, j_0) \in [-N, N]^b \times [-3N, 3N]\}$, the proof is similar.

Let

$$\mathcal{A}_{1}^{\theta} = \{ (n, j) \in [-N, N]^{b} \times [-3N, 3N] : |\theta + n \cdot \omega^{(0)} + \mu_{j}| \le \delta^{\frac{1}{8}} \},\$$

and

$$\mathcal{A}_{2}^{\theta} = \{ (n, j) \in [-N, N]^{b} \times [-3N, 3N] : |\theta + n \cdot \omega^{(0)} - \mu_{j}| \le \delta^{\frac{1}{8}} \}.$$

By (5) of Theorem 3.5, one has that for any θ ,

(91)
$$\#\mathcal{A}_1^{\theta} \le b, \#\mathcal{A}_2^{\theta} \le b.$$

Since the size of I_0 is $C\delta$, we have that there exist \mathcal{A}_1 and \mathcal{A}_2 independent of $\theta \in I_0$ such that

(92)
$$\#\mathcal{A}_1 \le b, \#\mathcal{A}_2 \le b,$$

and for any $(n, j) \in [-N, N]^b \times [-3N, 3N] \setminus (\mathcal{A}_1 \cup \mathcal{A}_2)$ and $\theta \in I_0$

(93)
$$|\theta + n \cdot \omega^{(0)} \pm \mu_j| \ge \frac{1}{2} \delta^{\frac{1}{8}}.$$

Take any $\tilde{\Lambda} \in R_{N_1}^{\sqrt{N}}$ with $N_1 \in [\sqrt{N}, 6N]$ and $\tilde{\Lambda} \subset [-N, N]^b \times [-3N, 3N]$. By perturbation arguments, we have that for any $\theta \in I_0$,

(94)
$$\| (R_{\tilde{\Lambda} \setminus (\mathcal{A}_1 \cup \mathcal{A}_2)} T(\theta) R_{\tilde{\Lambda} \setminus (\mathcal{A}_1 \cup \mathcal{A}_2)})^{-1} \| \le 3\delta^{-\frac{1}{8}}.$$

We are going to apply Cartan's estimate, Lemma 4.2. For this reason, let $\tau = C\delta$, $\tau_1 = 1$, $\Lambda = \mathcal{A}_1 \cup \mathcal{A}_2$, M = 2b, $B_1 = O(1)(\log \frac{1}{\delta})^K$, $B_2 = 3\delta^{-\frac{1}{8}}$, $B_3 = 1$ and $\epsilon = e^{-N_1^{3/4}}$. We note that since I_0 has size $C\delta$, (87) holds automatically.

Applying Cartan's estimate (Lemma 4.2) in all possible $\tilde{\Lambda} \in R_{N_1}^{\sqrt{N}}$ (in total N^C), there exists a subset $\tilde{\Theta}_{N_1} \subset \mathbb{R}$ such that

$$\operatorname{Leb}(\tilde{\Theta}_{N_1}) \le e^{-\frac{N_1^{\frac{3}{4}}}{|\log \delta|^2}}.$$

and for any $\theta \notin \tilde{\Theta}_{N_1}$ and any $\tilde{\Lambda} \in R_{N_1}^{\sqrt{N}}$ with $\tilde{\Lambda} \subset [-N, N]^b \times [-3N, 3N]$,

(95) $||(R_{\tilde{\Lambda}}TR_{\tilde{\Lambda}})^{-1}|| \le e^{N_1^{\frac{3}{4}}}.$

Let $N_0 = \sqrt{N}$. We call a box $(n_1, j_1) + Q_{N_0} \in \mathcal{E}_{N_0}$, $(n_1, j_1) \in [-N, N]^b \times [-3N, 3N]$ good if

$$||(R_{(n_1,j_1)+Q_{N_0}}TR_{(n_1,j_1)+Q_{N_0}})^{-1}|| \le e^{N_0^{\frac{9}{10}}}$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} \ge \frac{N_0}{10}$,

$$|(R_{(n_1,j_1)+Q_{N_0}}TR_{(n_1,j_1)+Q_{N_0}})^{-1}(n,j;n',j')| \le e^{-\tilde{c}_1 \max\{|n-n'|,|j-j'|\}}$$

Otherwise, we call $(n_1, j_1) + Q_{N_0} \in [-N, N]^b \times [-3N, 3N]$ bad. By (92), (93) and perturbation arguments, we have that there are at most 2*b* disjoint bad boxes of size $N_0 = N^{1/2}$ contained in $[-N, N]^b \times [-3N, 3N]$.

We have sublinear bound and (95). By [23, Theorem 2.1], for any $\theta \notin \bigcup_{\{I_i\}} \bigcup_{N_1 \in [\sqrt{N}, 6N]} \tilde{\Theta}_{N_1}$, (81) and (82) hold for the scale N. Therefore,

$$\Theta_N \subset \bigcup_{\{I_i\}} \bigcup_{N_1 \in [\sqrt{N}, 6N]} \tilde{\Theta}_{N_1}$$

and hence

$$\operatorname{Leb}(\Theta_N) \le e^{-N^{\frac{1}{10}}}.$$

г	_	_	٦
L			т

4.4. Large deviation theorem for large scales: $N \ge (\log \frac{1}{\lambda})^K$.

Proof. Let $N_2 = N_1^{K_1}$ and $N_4 = N_2^{K_1}$. Assume that $N_4 \ge (\log \frac{1}{\delta})^K$ and that the LDT holds at both scales N_1 and N_2 with parameter \tilde{c}_1 .

We will show that there are at most N_1^C bad disjoint boxes of size N_1 contained in $[-N_4, N_4]^b \times [-3N_4, 3N_4]$. Let $(n_1, l_1) \in [-N_4, N_4]^b \times [-3N_4, 3N_4]$ be such that $(n_1, l_1) + Q_{N_1}$ is bad for some $Q_{N_1} \in \mathcal{E}_{N_1}^0$.

We first bound the case when $|l_1| \leq 2N_1$. By the LDT at scale N_1 , there exists a set Θ_{N_1} with $\text{Leb}(\Theta_{N_1}) \leq e^{-N_1^{1/30}}$ such that for any $\theta \notin \Theta_{N_1}$ and any $Q_{N_1} \in \mathcal{E}_{N_1}^0$, Q_{N_1} is good. Since the operator is Töplitz with respect to $n \in \mathbb{Z}^b$, one has that for any (n_1, l_1) with $\theta + n_1 \cdot \omega \notin \Theta_{N_1}$, $(n_1, l_1) + Q_{N_1}$ is good for any $Q_{N_1} \in \mathcal{E}_{N_1}^0$. By standard arguments, we can assume that Θ_{N_1} is a semi-algebraic set of degree at most N_1^C , namely, there exist N_1^C intervals I_i of size $e^{-N_1^{1/30}}$,

such that $\Theta_{N_1} \subset \bigcup_i I_i$. The assumption on ω indicates that, for any nonzero n with $|n| \leq 2N_4$,

(96)
$$|n \cdot \omega| \ge e^{-(2N_4)^{\frac{1}{K_2}}} \ge e^{-N_1^{\frac{1}{30}}}.$$

Therefore, for any $|l_1| \leq 2N_1$, there is at most one bad box (n_1, l_1) such that $n_1 \cdot \omega \in I_i$. This leads to at most N_1^C bad boxes in this case.

When $|l_1| \ge 2N_1$, we will show that there are at most three disjoint bad boxes of size N_1 . First, if a box $(n, j) + Q_{N_1}$ is bad, by (78) and perturbation arguments, we must have that for some $(n_1, l_1) \in (n, j) + Q_{N_1}$, either

(97)
$$|\theta + n_1 \cdot \omega + \mu_{l_1}| \le 2e^{-N_1^{9/1}}$$

or

(98)
$$|\theta + n_1 \cdot \omega - \mu_{l_1}| \le 2e^{-N_1^{9/10}}.$$

Assume that indeed there are three bad boxes. We have that there are two from D_+ , namely (97) (or D_- , namely (98)). Therefore, we have that for two distinct vertices $(n, j) \in [-N_4, N_4]^b \times [-3N_4, 3N_4]$ and $(n', j') \in [-N_4, N_4]^b \times [-3N_4, 3N_4]$,

$$|m \cdot \omega - \mu_j + \mu_{j'}| \le 4e^{-N_1^{9/10}}, m = n - n'.$$

This contradicts the assumption (84).

Let $\tilde{\Theta}_{N_2} \subset \mathbb{R}$ be such that for some $(n, j) \in [-N_4, N_4]^b \times [-3N_4, 3N_4]$ such that either $|\theta + n \cdot \omega + \mu_j| \leq 2e^{-N_2^{9/10}}$ or $|\theta + n \cdot \omega - \mu_j| \leq 2e^{-N_2^{9/10}}$. Since for any $|l_1| \geq 2N_2$, the matrix is essentially diagonal, we have that for any $\theta \notin \tilde{\Theta}_{N_2}$, $(n_1, l_1) \in [-N_4, N_4]^b \times [-3N_4, 3N_4]$ with $|l_1| \geq 2N_2$ and $Q_{N_2} \in \mathcal{E}_{N_2}^0$, $(n_1, l_1) + Q_{N_2}$ is good. Let $\hat{\Theta}_{N_2} = \{\theta$: for some $n \in [-N_4, N_4]^b, \theta + n \cdot \omega \in \Theta_{N_2}\}$. Therefore, for any $\theta \notin \hat{\Theta}_{N_2}$, $(n_1, l_1) \in [-N_4, N_4]^b \times [-2N_2, 2N_2]$ and $Q_{N_2} \in \mathcal{E}_{N_2}^0$, $(n_1, l_1) + Q_{N_2}$ is good. Clearly

$$\operatorname{Leb}(\tilde{\Theta}_{N_2} \cap \hat{\Theta}_{N_2}) \le e^{-N_2^{1/31}},$$

and for any $\theta \notin \tilde{\Theta}_{N_2} \cap \hat{\Theta}_{N_2}$, $(n_1, l_1) \in [-N_4, N_4]^b \times [-3N_4, 3N_4]$ and $Q_{N_2} \in \mathcal{E}_{N_2}^0$, $(n_1, l_1) + Q_{N_2}$ is good.

Applying Lemma 4.2 (see proof of [23, Theorem 2.2] for details), for any $N_3 \in [N_4^{1/2}, N_4]$, there exists a subset $\tilde{\Theta}_{N_3} \subset \mathbb{R}$ such that

$$\operatorname{Leb}(\tilde{\Theta}_{N_3}) \le e^{-N_3^{\frac{1}{4}}},$$

and for any $N \in [N_3^{1/2}, N_3]$, $\tilde{\Lambda} \in R_{6N}^{N_3^{1/2}}$ with $\tilde{\Lambda} \subset [-N_3, N_3]^b \times [-3N_3, 3N_3]$, and for any $\theta \notin \tilde{\Theta}_{N_3}$,

(99)
$$||(R_{\tilde{\Lambda}}TR_{\tilde{\Lambda}})^{-1}|| \le e^{N^{\frac{3}{4}}}.$$

Let $N_0 = N_3^{1/2}$. We call a box $(n_1, l_1) + Q_{N_0} \in \mathcal{E}_{N_0}^0$, $(n_1, l_1) \in [-N_3, N_3]^b \times [-3N_3, 3N_3]$ good if

$$||(R_{(n_1,l_1)+Q_{N_0}}TR_{(n_1,l_1)+Q_{N_0}})^{-1}|| \le e^{N_0^{\frac{9}{10}}}$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} \ge \frac{N_0}{10}$,

$$|(R_{(n_1,l_1)+Q_{N_0}}TR_{(n_1,l_1)+Q_{N_0}})^{-1}(n,j;n',j')| \le e^{-c_2 \max\{|n-n'|,|j-j'|\}},$$

where $c_2 = \tilde{c}_1 - N_4^{-\kappa}$ with a proper $\kappa > 0$. Otherwise, we call $(n_1, l_1) + Q_{N_0} \in [-N_3, N_3]^b \times [-3N_3, 3N_3]$ bad. Since there are at most N_1^C bad boxes of size N_1 , by resolvent identity, we have that there are at most N_1^C disjoint bad boxes of size $N_0 = N_3^{1/2}$ contained in $[-N_3, N_3]^b \times [-3N_3, 3N_3]$.

size $N_0 = N_3^{1/2}$ contained in $[-N_3, N_3]^b \times [-3N_3, 3N_3]$. We have achieved the sublinear bound and (99). By [23, Theorem 2.1], we have that the LDT holds for any scale $N_3 \in [N_4^{1/2}, N_4]$ with parameter $\tilde{c}_1 - N_4^{-\kappa}$, where κ is a proper small positive constant. Now the proof follows from standard inductions.

5. The nonlinear analysis

Fix $V \in X_{\epsilon}$, so that the conclusions of Theorem 3.5 hold and Theorem 4.1 is available. (As before, we omit the superscript V, as it is fixed.) We can now solve the nonlinear matrix equation (6) and therefore the NLRS equation (1).

5.1. Lyapunov-Schmidt decomposition. To simplify notations, we write u for \hat{u} , namely $u(n, j) = \hat{u}(n, j)$. Let v be the complex conjugate of u, more precisely, $v(n, j) = \hat{u}(-n, j)$.

By (8), W_u is a vector on $\ell^2(\mathbb{Z}^{b+1})$, which is now given by

$$W_{u}(n,j) = \sum_{\substack{n' + \sum_{m=1}^{p} (n_{m}+n'_{m})=n \\ n',n_{m},n'_{m} \in \mathbb{Z}^{b}}} \sum_{l_{m},l'_{m},j' \in \mathbb{Z}} u(n',j') \prod_{m=1}^{p} u(n_{m},l_{m})v(n'_{m},l'_{m})} \left(\sum_{x \in \mathbb{Z}} \phi_{j}(x)\phi_{j'}(x) \prod_{m=1}^{p} \phi_{l_{m}}(x)\phi_{l'_{m}}(x) \right).$$
(100)

Let \widetilde{W}_u be a vector on $\ell^2(\mathbb{Z}^{b+1})$, which is given by

$$\widetilde{W}_{u}(n,j) = \sum_{\substack{n' + \sum_{m=1}^{p} (n_{m}+n'_{m})=n \\ n',n_{m},n'_{m} \in \mathbb{Z}^{b}}} \sum_{\substack{j',l_{m},l'_{m} \in \mathbb{Z}}} v(n',j') \prod_{m=1}^{p} v(n_{m},l_{m})u(n'_{m},l'_{m})} \left(\sum_{x \in \mathbb{Z}} \phi_{j}(x)\phi_{j'}(x) \prod_{m=1}^{p} \phi_{l_{m}}(x)\phi_{l'_{m}}(x) \right).$$
(101)

We remark that \widetilde{W}_u and W_u are functions of u and v. We only indicate the dependence on u for simplicity and the fact that v is the conjugate of u.

Writing the equation for v as well, leads to the system of nonlinear equations on $\mathbb{Z}^{b+1} \times \{0, 1\}$:

(102)
$$(D_{+}u)(n,j) + \delta W_{u}(n,j) = 0, (D_{-}u)(n,j) + \delta \widetilde{W}_{u}(n,j) = 0,$$

where D_{\pm} are the diagonal matrices with entries

(103)
$$D_{\pm}(n,j) := D_{\pm}(n,j;n,j) = \pm n \cdot \omega + \mu_j.$$

Define

(104)
$$S = \{(-e_k, \alpha_k) \times \{0\}, (e_k, \alpha_k) \times \{1\}, k = 1, 2, \cdots, b\},\$$

and denote the complement by S^c :

(105)
$$S^c = \mathbb{Z}^{b+1} \times \{0, 1\} \backslash S.$$

Write D for the diagonal matrix composed of the diagonal blocks D_{\pm} and write (102) in the form F(u, v) = 0. Since v is the conjugate of u, we simply write F(u) = 0 for F(u, v) = 0. We make a Lyapunov-Schmidt decomposition of (102) into the P-equations:

(106)
$$F(u)|_{S^c} = 0.$$

and the Q-equations:

(107)
$$F(u)|_S = 0.$$

5.2. The *P*-equations. The *P*-equations are infinite dimensional. They are solved using a Newton scheme, starting from the initial approximation $u^{(0)} = u_0$. Let F' be the linearized operator on $\ell^2(\mathbb{Z}^{b+1}) \times \{0,1\}$,

$$F'(u) = D + \delta \mathcal{W}_u,$$

where

$$\mathcal{W}_u = \begin{pmatrix} \frac{\partial W_u}{\partial u} & \frac{\partial W_u}{\partial v} \\ \frac{\partial W_u}{\partial u} & \frac{\partial W_u}{\partial v} \end{pmatrix}.$$

It is easy to see that $\mathcal{W} = \mathcal{W}_u$ satisfies

(1) \mathcal{W} is Töplitz with respect to $n \in \mathbb{Z}^b$, namely for any $j \in \mathbb{Z}, j' \in \mathbb{Z}, k \in \mathbb{Z}^b, n \in \mathbb{Z}^b, n' \in \mathbb{Z}^b, r \in \{0, 1\}, r' \in \{0, 1\}$

$$\mathcal{W}_{r,r'}(n,j;n',j') = \mathcal{W}_{r,r'}(n+k,j;n'+k,j').$$

(2) Assume that $|u(n,j)| \leq e^{-c(|n|+|j|)}$. Direct computation implies that

$$|\mathcal{W}_{r,r'}(n,j;n',j')| \le Ce^{-c'(|n-n'|+|j|+|j'|)}.$$

The operator F' is to be evaluated near $\omega = \omega^{(0)} = (\mu_{\alpha_1}, \mu_{\alpha_2}, \cdots, \mu_{\alpha_b})$, the linear frequency, and $u = u^{(0)}$ and $v = v^{(0)}$. As earlier, we have that $u^{(0)}(-e_k, \alpha_k) = a_k$, $k = 1, 2, \cdots, b; u^{(0)}(n, j) = 0$, otherwise.

Recall next the formal Newton scheme:

$$\Delta \begin{pmatrix} u \\ v \end{pmatrix} = -[F'_{S^c}(u)]^{-1}F(u)|_{S^c},$$

where the left side denotes the correction to $\binom{u}{v}$, $F'_{S^c}(u)$ is the linearized operator evaluated at (u, v): F'(u), and restricted to S^c :

$$F_{S^c}^\prime(u)(x,y)=F^\prime(u)(x,y),$$

for $x, y \in S^c$; likewise $F(u)|_{S^c}$ is F(u) restricted to S^c :

$$[F(u)|_{S^c}](x) = F(u)(x),$$

for $x \in S^c$.

Since we seek solutions close to $(u^{(0)}, v^{(0)})$, which has compact support in $\mathbb{Z}^{b+1} \times \{0, 1\}$, we adopt a *multiscale* Newton scheme as follows: At iteration step (r + 1), choose an appropriate scale N and estimate $[F'_N]^{-1}$,

At iteration step (r + 1), choose an appropriate scale N and estimate $[F_N]^{-1}$, where F'_N is F' restricted to

$$[-N, N]^{b+1} \times \{0, 1\} \setminus S \subset \mathbb{Z}^{b+1} \times \{0, 1\},\$$

and evaluated at $u^{(r)}$ and $v^{(r)}$: $F'_N = F'_N(u^{(r)})$. Define the (r+1)-th correction to be:

$$\Delta \begin{pmatrix} u^{(r+1)} \\ v^{(r+1)} \end{pmatrix} = -[F'_N(u^{(r)})]^{-1}F(u^{(r)}),$$

and

$$\begin{split} u^{(r+1)} &= u^{(r)} + \Delta u^{(r+1)}, \\ v^{(r+1)} &= v^{(r)} + \Delta v^{(r+1)}, \end{split}$$

for all $r = 0, 1, 2, \cdots$.

At step r, $\mathcal{W}_{u^{(r)}(\omega,a)}$ (depending on $u^{(r)}(\omega,a)$) is a function of ω and a. We write $T_{u^{(r)}}(\theta, \omega, a)$ for the operator $F' = D(\theta) + \delta \mathcal{W}_{u^{(r)}(\omega,a)}$, and $\tilde{T}(\theta, \omega, a)$ the operator $F' = D(\theta) + \delta \mathcal{W}_{u^{(r)}(\omega,a)}$, restricted to $\mathbb{Z}^{b+1} \times \{0,1\} \setminus S$, where

(108)
$$D(\theta) = \begin{bmatrix} \operatorname{diag} (n \cdot \omega + \theta + \mu_j) & 0\\ 0 & \operatorname{diag} (-n \cdot \omega - \theta + \mu_j) \end{bmatrix}, (n, j) \in \mathbb{Z}^{b+1}.$$

For simplicity, write $\tilde{T}_{u^{(r)}}(\omega, a)$ for $\tilde{T}_{u^{(r)}}(0, \omega, a)$ and $T_{u^{(r)}}(\omega, a)$ for $T_{u^{(r)}}(0, \omega, a)$.

The analysis of the linearized operators F'_N uses Theorem 3.5 for small scales; for large scales, it also uses Theorem 4.1 and semi-algebraic projection to convert estimates in θ into that of ω , and finally a.

5.3. The *Q*-equations. The *Q*-equations are 2*b* dimensional, but due to symmetry leading to *b* equations only. They are used to relate ω with *a*. The amplitudes u(n, j) are fixed on *S*, i.e., $u(-e_{\alpha_k}, \alpha_k) = a_k, k = 1, 2, \dots, b$, and the same for the complex conjugate. So we have

(109)
$$\omega_k = \mu_{\alpha_k} + \delta \frac{W_u(-e_k, \alpha_k)}{a_k}, k = 1, 2, \cdots, b.$$

When $u = u^{(0)}$, let us compute the terms in the *Q*-equations (109). For $k \in \{1, 2, \dots, b\}$, we have

$$W_{u^{(0)}}(-e_k,\alpha_k) = \sum_{n'+\sum_{m=1}^p n_m - n'_m = -e_k} u^{(0)}(n',l') \prod_{m=1}^p u^{(0)}(n_m,l_m)v^{(0)}(n'_m,l'_m) \\ \left(\sum_{x \in \mathbb{Z}} \phi_{\alpha_k}(x)\phi_{l'}(x) \prod_{m=1}^p \phi_{l_m}(x)\phi_{l'_m}(x)\right).$$

The sum in (110) runs over $l_m \in \mathbb{Z}$, $l'_m \in \mathbb{Z}$, $n'_m \in \mathbb{Z}^b$, $n_m \in \mathbb{Z}^b$, $n' \in \mathbb{Z}^b$, $l' \in \mathbb{Z}$, $m = 1, 2, \dots, p$.

Since $u^{(0)}$ has support $\{(-e_k, \alpha_k)\}_{k=1}^b$, in order to contribute to (110), one has that

(111)
$$l_m \in \{\alpha_k\}_{k=1}^b, l'_m \in \{\alpha_k\}_{k=1}^b, m = 1, 2, \cdots, b \text{ and } l' \in \{\alpha_k\}_{k=1}^b.$$

Take $\sum_{x \in \mathbb{Z}} \phi_{\alpha_k}(x) \phi_{l'}(x) \prod_{m=1}^p \phi_{l_m}(x) \phi_{l'_m}(x)$ into consideration. Assume $l' = \alpha_k$ and $l_m = l'_m = \alpha_k$, $m = 1, 2, \cdots, b$. It is easy to see that (similar to the proof of (25)),

(112)
$$\sum_{\ell \in \mathbb{Z}, |\ell - \ell_{\alpha_k}| \le \frac{L}{2}} |\phi_{\alpha_k}(\ell)|^2 \ge 1 - e^{-\frac{\gamma}{4}L}.$$

This implies that there exists $\ell \in \mathbb{Z}$ with $|\ell - \ell_{\alpha_k}| \leq \frac{L}{2}$ such that

(113)
$$|\phi_{\alpha_k}(\ell)| \ge \frac{1}{L^3}.$$

Therefore, in this case,

(114)
$$\sum_{x \in \mathbb{Z}} \phi_{\alpha_k}(x) \phi_{l'}(x) \prod_{m=1}^p \phi_{l_m}(x) \phi_{l'_m}(x) = \sum_{x \in \mathbb{Z}} |\phi_{\alpha_k}(x)|^{2p+2} \ge \frac{1}{L^{10p}}.$$

Except for the case $l' = \alpha_k$ and $l_m = l'_m = \alpha_k$, $m = 1, 2, \dots, b$, by (111), we have that

(115)
$$|\sum_{x\in\mathbb{Z}}\phi_{\alpha_k}(x)\phi_{l'}(x)\prod_{m=1}^p\phi_{l_m}(x)\phi_{l'_m}(x)| \le e^{-cL}.$$

Denote by $A_k = \sum_{x \in \mathbb{Z}} |\phi_{\alpha_k}(x)|^{2p+2}$. By (110), (114) and (115) (the leading contribution in the sum of (110) is when $(n', l') = (n'_m, l'_m) = (n_m, l_m) = (-e_k, \alpha_k)$), we have that

$$\omega_k^{(0)} = \mu_{\alpha_k} + \delta(A_k a_k^{2p} + O(1)e^{-cL}),$$

and $\frac{1}{L^{10p}} \leq A_k \leq 1$.

Denote by $\Omega_0 = [\mu_{\alpha_1}, \mu_{\alpha_1} + 2^{2p+1}\delta] \times [\mu_{\alpha_2}, \mu_{\alpha_2} + 2^{2p+1}\delta] \times [\mu_{\alpha_b}, \mu_{\alpha_b} + 2^{2p+1}\delta] \subset \mathbb{R}^b$. Assume that after r steps, we obtain a C^1 function $u^{(r)}(\omega, a)$ on $\Omega_0 \times [1, 2]^b$. Substituting $u^{(r)}(\omega, a)$ and $v^{(r)}(\omega, a)$ into (109), the implicit function theorem yields,

(116)
$$\begin{aligned} \omega &= \omega^{(r+1)}(a), \\ a &= a^{(r+1)}(\omega). \end{aligned}$$

Moreover, for some C^1 functions $f_k, k = 1, 2, \dots, b$,

(117)
$$\omega_k = \mu_{\alpha_k} + \delta(A_k a_k^{2p} + f_k(a_1, a_2, \cdots, a_b)).$$

Denote by Γ_r , the graph of (ω, a) at step r. Denote by P_x the projection onto the x-variable, where $x = a, \omega$ or (ω, a) .

5.4. The induction hypothesis. Let M be a large integer, and denote by B(0, R) the ℓ^{∞} ball on \mathbb{Z}^{b+1} centered at the origin with radius R. Set

$$r_0 = \left\lfloor \frac{|\log \delta|^{\frac{3}{4}}}{\log M} \right\rfloor.$$

The proof of the Theorem is an induction. So we first lay down the induction hypothesis, which we prove in sect. 6. In the following C is a large constant (much larger than K_2 , which appears in Theorem 4.1) and c > 0 is a small constant.

For $r \geq 1$, we assume that the following holds:

- **Hi.** $u^{(r)}(\omega, a)$ is a C^1 map on $\Omega_0 \times [1, 2]^b$, and supp $u^{(r)} \subseteq B(0, M^r)$ (supp $u^{(0)} \subset B(0, M)$).
- **Hii.** $\|\Delta u^{(r)}\| \leq \delta_r$, $\|\partial \Delta u^{(r)}\| \leq \bar{\delta}_r$, where ∂ denotes ∂_x , x stands for ω_i , a_i , $i = 1, 2, \cdots, b$ and $\|\| := \sup_{(\omega, a)} \|\|_{\ell^2(\mathbb{Z}^{b+1})}$.
- **Hiii.** $|u^{(r)}(n,j)| \le Ce^{-c \max\{|n|,|j|\}}.$

Hiv. There exists Λ_r , a set of open sets I in (ω, a) of size M^{-r^C} when $r \ge r_0$ (the total number of open sets is therefore bounded above by M^{r^C}), such that for any $(\omega, a) \in \bigcup_{I \in \Lambda_r} I$ when $r \ge r_0$ and $(\omega, a) \in \Omega_0 \times [1, 2]^b$ when $1 \le r \le r_0 - 1$, (1) $u^{(r)}(\omega, a)$ is a rational function in (ω, a) of degree at most M^{r^3} ; (2)

(118)
$$\|F(u^{(r)})\| \le \kappa_r, \|\partial F(u^{(r)})\| \le \bar{\kappa}_r;$$

(119)
$$\| (R_{[-M^r,M^r]^{b+1}} \tilde{T}_{u^{(r-1)}}(\omega,a) R_{[-M^r,M^r]^{b+1}})^{-1} \| \le \delta^{-\frac{1}{8}} M^{r^C},$$

and for max{ $|n - n'|, |j - j'|$ } > r^C , (120)

(120)
$$|(R_{[-M^r,M^r]^{b+1}}\tilde{T}_{u^{(r-1)}}(\omega,a)R_{[-M^r,M^r]^{b+1}})^{-1}(n,j;n',j')| \le \delta^{-\frac{1}{8}}e^{-c\max\{|n-n'|,|j-j'|\}}.$$

(4) Each
$$I \in \Lambda_r$$
 is contained in an open set $I' \in \Lambda_{r-1}$, $r \ge r_0$, and

(121)
$$\operatorname{Leb}(P_a(\Gamma_{r-1} \cap (\bigcup_{I' \in \Lambda_{r-1}} I' \setminus \bigcup_{I \in \Lambda_r} I))) \le e^{-|\log \delta|^{K_1^{90}}} + M^{-\frac{r_0}{2^b}}, r = r_0;$$

and

(122)
$$\operatorname{Leb}(P_a(\Gamma_{r-1} \cap (\bigcup_{I' \in \Lambda_{r-1}} I' \setminus \bigcup_{I \in \Lambda_r} I))) \le M^{-\frac{r}{2^b}}, r \ge r_0 + 1;$$

- (5) for $(\omega, a) \in \bigcup_{I \in \Lambda_r} I$ with $r \ge r_0$, ω satisfies the conditions (83), and (84) for $n \ne 0$, in the scales \tilde{N} in $[(\log \frac{1}{\delta})^K, M^r];$
- (6) The iteration holds with

(123)
$$\delta_r = \delta^{\frac{1}{2}} M^{-(\frac{4}{3})^r}, \ \bar{\delta}_r = \delta^{\frac{1}{8}} M^{-\frac{1}{2}(\frac{4}{3})^r}; \\ \kappa_r = \delta^{\frac{3}{4}} M^{-(\frac{4}{3})^{r+2}}, \ \bar{\kappa}_r = \delta^{\frac{3}{8}} M^{-\frac{1}{2}(\frac{4}{3})^{r+2}}.$$

Remark 10. As usual in multi-scale arguments, the constant c depends on r. From step r to step r + 1, c becomes slightly smaller. We ignore the dependence since it is essentially irrelevant.

Remark 11. The Lyapunov-Schmidt approach to quasi-periodic solutions was initiated in the paper [10], and greatly generalized by Bourgain starting from the paper [4].

6. Proof of the Theorem

The general scheme of the proof is that, for small scales, we use (71) and (72) of Theorem 3.5 to solve the *P*-equations; while for larger scales, we use (73) and (74) of Theorem 3.5, Theorem 4.1 and semi-algebraic projection.

Let us state the projection lemma.

Lemma 6.1. Let $S \subset [0,1]^{d_1} \times [0,1]^{d_2} := [0,1]^d$, be a semi-algebraic set of degree B and $\operatorname{meas}_d S < \eta, \log B \ll \log 1/\eta$. Denote by $(x,y) \in [0,1]^{d_1} \times [0,1]^{d_2}$ the product variable. Fix $\epsilon > \eta^{1/d}$. Then there is a decomposition

$$\mathcal{S} = \mathcal{S}_1 \bigcup \mathcal{S}_2,$$

with S_1 satisfying

$$\operatorname{Leb}(\operatorname{Proj}_x \mathcal{S}_1) \leq B^C \epsilon,$$

and S_2 the transversality property

$$\operatorname{Leb}(\mathcal{S}_2 \cap L) \leq B^C \epsilon^{-1} \eta^{1/d}$$

for any d_2 -dimensional hyperplane L in $[0,1]^{d_1+d_2}$ such that

$$\max_{1 \le l \le d_1} |\operatorname{Proj}_L(e_l)| \le \frac{1}{100} \epsilon,$$

where e_l are the basis vectors for the x-coordinates.

The above lemma is the basic tool, underlining the semi-algebraic techniques used in the subject. It is stated as (1.5) in [6], cf., Lemma 9.9 [5] and Proposition 5.1 [7], and relies on the Yomdin-Gromov triangulation theorem. For a complete proof of the latter, see [3]. Together with Theorem 4.1, (73) and (74), it enables us to go beyond the perturbative scales in (71) and (72).

Proof of induction hypothesis. Assume that the induction holds for all scales up to r. We will prove that it holds for r + 1. From our construction, it is easy to see that $u^{(r)}(\omega, a)$ is a rational function in (ω, a) of degree at most $M^{(r+1)^3}$.

For $r \leq r_0 - 1$, by (71), (72) and standard perturbation arguments, we have that for any $(\omega, a) \in \Omega_0 \times [1, 2]^b$,

(124)
$$||(R_{[-M^{r+1},M^{r+1}]^{b+1}}\tilde{T}_{u^{(r)}}(\omega,a)R_{[-M^{r+1},M^{r+1}]^{b+1}})^{-1}|| \le 2\delta^{-\frac{1}{8}}M^{(r+1)^{C}},$$

and for any (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} \ge (r + 1)^C$, (125)

$$|(R_{[-M^{r+1},M^{r+1}]^{b+1}}\tilde{T}_{u^{(r)}}(\omega,a)R_{[-M^{r+1},M^{r+1}]^{b+1}})^{-1}(n,j;n',j')| \le 2\delta^{-\frac{1}{8}}e^{-c\max\{|n-n'|,|j-j'|\}}.$$

We are in a position to treat the case $r \ge r_0$. Let $\mathcal{X} \subset \bigcup_{I \in \Lambda_r} I$ be such that $(\omega, a) \in \mathcal{X}$ satisfies (5) of Hiv at step r + 1 and

(126)
$$\| (R_{[-M^r,M^r]^{b+1}} \tilde{T}_{u^{(r)}}(\omega,a) R_{[-M^r,M^r]^{b+1}})^{-1} \| \le \delta^{-\frac{1}{8}} M^{r^C}$$

and for any (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} > r^C$, (127)

$$|(R_{[-M^r,M^r]^{b+1}}\tilde{T}_{u^{(r)}}(\omega,a)R_{[-M^r,M^r]^{b+1}})^{-1}(n,j;n',j')| \le \delta^{-\frac{1}{8}}e^{-c\max\{|n-n'|,|j-j'|\}}.$$

By perturbation arguments, we can essentially assume that \mathcal{X} is the union of a collection of open intervals of size $M^{-(r+1)^C}$. Denoting the collection by Λ_{r+1} , we have constructed Λ_{r+1} . Except for (121) and (122), it is now routine that the rest of Hi-v hold for r + 1, see Chap. 18, IV, (18.36)-(18.41) [5] and Lemmas 5.2 and 5.3 [27]. See appendix C for more details.

We proceed to the proof of the measure estimates (122) and (121). Let $N = M^{r+1}$ and $N_1 = (\log N)^C$ with a large constant C. Let

$$r_1 = \left\lfloor \frac{2\log N_1}{\log \frac{4}{3}} \right\rfloor + 1,$$

so that $\delta_{r_1} < e^{-N_1^2}$. Consider $T_{u^{(r_1)}}$. Pick one interval $I \in \Lambda_{r_1}$ of size $M^{-r_1^C}$ and let $I_1 = P_{\omega}(I \cap \Gamma_{r_1})$. By the *Q*-equation, we have that the size of I_1 is smaller than $C\delta$.

Solving the Q-equation at step $r_1 - 1$, one has that $a = a^{(r_1)}(\omega), \omega \in I_1$. Since $\omega \in \Omega_0$, one has that the first assumption of Theorem 4.1 always holds. By Theorem 4.1, there exists X_{N_1} (depending on ω) such that for any $\theta \notin X_{N_1}$,

(128)
$$||(R_{Q_{N_1}}T_{u^{(r_1)}}(\theta,\omega,a^{(r_1)}(\omega))R_{Q_{N_1}})^{-1}|| \le e^{N_1^{\frac{10}{10}}}$$

and for any $(n, j) \in \mathbb{Z}^{b+1}$ and $(n', j') \in \mathbb{Z}^{b+1}$ with $\max\{|n - n'|, |j - j'|\} \ge \frac{N_1}{10}$, (129) $|(R_{Q_{N_1}}T_{u^{(r_1)}}(\theta, \omega, a^{(r_1)}(\omega))R_{Q_{N_1}})^{-1}(n, j; n', j')| \le e^{-c\max\{|n - n'|, |j - j'|\}}$,

and X_{N_1} satisfies

(130)
$$\operatorname{Leb}(X_{N_1}) \le e^{-N_1^{\frac{1}{30}}}.$$

Let $K_{N_1} = \{\pm n \cdot \omega + \mu_j : |n| \leq N_1, |j| \leq 3N_1\}$ and I_{N_1} (depending on ω) be the $C\delta$ neighbour of K_{N_1} . Assume $\theta \notin I_{N_1}$. Then the diagonal entries $D_+, D_$ are larger than $C\delta$. Perturbation argument leads to, for any $|j_0| \leq 2N_1$ and $Q_{N_1} \in \mathcal{E}_{N_1}^0$,

$$\|(R_{Q_{N_1}(j_0)}T_{u^{(r_1)}}(\theta,\omega,a^{(r_1)}(\omega))R_{Q_{N_1}(j_0)})^{-1}\| \le \frac{1}{\delta},$$

and for any (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} \ge \frac{N_1}{10}$,

$$|(R_{Q_{N_1}(j_0)}T_{u^{(r_1)}}(\theta,\omega,a^{(r_1)}(\omega))Q_{Q_{N_1}(j_0)})^{-1}(n,n')| \le e^{-c\max\{|n-n'|,|j-j'|\}}.$$

Since $N_1 \geq (\log \frac{1}{\delta})^C$ (*C* is large), $\frac{1}{\delta} \leq e^{N_1^{\frac{1}{2}}}$. Therefore, (128) and (129) hold. This implies that $X_{N_1} \subset I_{N_1}$. Since Ω_0 has size $C\delta$, we can assume that X_{N_1} is in a union of a collection of intervals of size δ with total number N_1^C (independent of ω). Pick one interval Θ . Let $\mathcal{X}_{N_1}(\omega, \theta) \subset I_1 \times \Theta$ be such that there exists some $Q_{N_1} \in \mathcal{E}_{N_1}$ such that either (128) or (129) is not true. By (130) and Fubini theorem, one has that

(131)
$$\operatorname{Leb}(\mathcal{X}_{N_1}) \le C\delta e^{-N_1^{\frac{1}{30}}} \le \delta e^{-N_1^{\frac{1}{31}}}.$$

We can assume that $\mathcal{X}_{N_1} \subset I_1 \times \Theta$ is a semi-algebraic set of degree at most $N_1^C M^{Cr_1^3}$. This can be seen as follows. Let $\tilde{X}_{N_1} \subset \Omega_0 \times [1, 2]^b \times \mathbb{R}$ be such that there exists some $Q_{N_1} \in \mathcal{E}_{N_1}$ such that one of the following is not true:

(132)
$$||(R_{Q_{N_1}}T_{u^{(r_1)}}(\theta,\omega,a^{(r_1)}(\omega))R_{Q_{N_1}})^{-1}|| \le e^{N_1^{\frac{9}{10}}},$$

and for any $(n, j) \in \mathbb{Z}^{b+1}$ and $(n', j') \in \mathbb{Z}^{b+1}$ with $\max\{|n - n'|, |j - j'|\} \ge \frac{N_1}{10}$,

(133)
$$|(R_{Q_{N_1}}T_{u^{(r_1)}}(\theta,\omega,a^{(r_1)}(\omega))R_{Q_{N_1}})^{-1}(n,j;n',j')| \le e^{-c\max\{|n-n'|,|j-j'|\}}$$

Therefore, $\mathcal{X}_{N_1} = P_{(\omega,\theta)}(\tilde{X}_{N_1} \cap (\Gamma_{r_1} \times \mathbb{R}))$. Clearly, both \tilde{X}_{N_1} and Γ_{r_1} are semialgebraic sets of degree at most $N_1^C M^{Cr_1^3}$. Lemma 4.4 implies that \mathcal{X}_{N_1} is a semi-algebraic set of degree at most $N_1^C M^{Cr_1^3}$.

Let

(134)
$$\epsilon_l = M^{-\frac{r}{2b-l}}, l = 1, 2, \cdots, b-1, \text{ and } \epsilon_b = 10M^{-r}.$$

Choose any $|j_0| \leq 2N_1$. Recall that $T_{u^{(r_1)}}$ is Töplitz with respect to $n \in \mathbb{Z}^d$. Denote by $\epsilon_{b+1} = e^{-N_1^{\frac{1}{40}}}$. Applying Lemma 6.1 in all possible directions (see (3.26) in [6]) and also on all possible open sets and Θ (the total number is bounded by $N_1^C M^{Cr_1^3}$), there exists a set of ω , $I_1^r \subset I_1$ such that

(135)
$$\operatorname{Leb}(I_1^r) \le \delta M^{Cr_1^3} N_1^C \left(\sum_{k=2}^{b+1} \left(\prod_{l=1}^{k-1} \epsilon_l^{-1} \right) \epsilon_k \right) \le \delta M^{-\frac{r}{2^{b-1}}} N_1^C M^{Cr_1^3},$$

and for any $\omega \in I_1 \setminus I_1^r$, one has that for any $(n_0, j_0) \in [-N, N]^b \times [-2N_1, 2N_1]$ with $\max\{|n_0|, |j_0|\} \geq \frac{M^r}{10}$ and $(\omega, a) \in \Gamma_{r_1}$,

(136)
$$||(R_{(n_0,j_0)+Q_{N_1}}T_{u^{(r_1)}}(\omega,a)R_{(n_0,j_0)+Q_{N_1}})^{-1}|| \le e^{N_1^{\frac{9}{10}}},$$

and for any (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} \ge \frac{N_1}{10}$,

(137)
$$|(R_{(n_0,j_0)+Q_{N_1}}T_{u^{(r_1)}}(\omega,a)R_{(n_0,j_0)+Q_{N_1}})^{-1}(n,j;n',j')| \le e^{-c\max\{|n-n'|,|j-j'|\}}.$$

Let us explain where the factor δ in (135) is from. We apply Lemma 6.1 in $I_1 \times \Theta$, where both I_1 and Θ have sizes $C\delta$. By scaling, we have such a δ factor.

Assume $|j_0| > 2N_1$. In this case, we can assume that $R_{(n_0,j_0)+Q_{N_1}}T_{u^{(r_1)}}(\omega, a)R_{(n_0,j_0)+Q_{N_1}}$ is essentially a diagonal matrix. So, we only need to remove ω such that for some $(n,j) \in [-N,N]^{b+1}, |n \cdot \omega + \mu_j| \leq 2e^{-N_1^{\frac{9}{10}}}$ or $|n \cdot \omega - \mu_j| \leq 2e^{-N_1^{\frac{9}{10}}}$. This can not happen when n = 0 because of (70). Direct computions imply that there exists a set of ω , $\tilde{I}_1^r \subset I_1$ such that

(138)
$$\operatorname{Leb}(\tilde{I}_1^r)) \le N^{C(b)} e^{-N_1^{\frac{9}{10}}} < \delta M^{-\frac{r}{2^{b-1}}},$$

and for any $\omega \in I_1 \setminus \tilde{I}_1^r$, one has that for any $(n_0, j_0) \in [-N, N]^{b+1}$ with $\max\{|n_0|, |j_0|\} \ge \frac{M^r}{10}, |j_0| \ge 2N_1$ and $(\omega, a) \in \Gamma_{r_1}$,

(139)
$$||(R_{(n_0,j_0)+Q_{N_1}}T_{u^{(r_1)}}(\omega,a)R_{(n_0,j_0)+Q_{N_1}})^{-1}|| \le e^{N_1^{\frac{9}{10}}},$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} \ge \frac{N_1}{10}$, (140) $|(R_{(n_0, j_0) + Q_{N_1}} T_{u^{(r_1)}}(\omega, a) R_{(n_0, j_0) + Q_{N_1}})^{-1}(n, j; n', j')| \le e^{-c \max\{|n - n'|, |j - j'|\}}$.

Since the distance between Γ_{r_1} and Γ_r is less than $C\delta_{r_1} \leq Ce^{-N_1^2}$ and $||u^{(r_1)} - u^{(r)}|| \leq C\delta_{r_1} \leq Ce^{-N_1^2}$, by perturbation arguments, for any $(n_0, j_0) \in [-M^{r+1}, M^{r+1}]^{b+1}$ with $\max\{|n_0|, |j_0|\} \geq \frac{M^r}{10}$, $(\omega, a) \in \Gamma_r$ and $\omega \in I_1 \setminus (I_1^r \cup \tilde{I}_1^r)$,

(141)
$$||(R_{(n_0,j_0)+Q_{N_1}}T_{u^{(r)}}(\omega,a)R_{(n_0,j_0)+Q_{N_1}})^{-1}|| \le 2e^{N_1^{\frac{7}{10}}},$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} \ge \frac{N_1}{10}$

$$(142) |(R_{(n_0,j_0)+Q_{N_1}}T_{u^{(r)}}(\omega,a)R_{(n_0,j_0)+Q_{N_1}})^{-1}(n,j;n',j')| \le 2e^{-c\max\{|n-n'|,|j-j'|\}}$$

By (119) and (120) at step r, and using perturbation arguments, one has that for any $(\omega, a) \in \bigcup_{I \in \Lambda_r} I$,

(143)
$$\| (R_{[-M^r,M^r]^{b+1}} \tilde{T}_{u^{(r)}}(\omega,a) R_{[-M^r,M^r]^{b+1}})^{-1} \| \le 2\delta^{-\frac{1}{8}} M^{r^C},$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} > r^C$, (144)

$$|(R_{[-M^r,M^r]^{b+1}}\tilde{T}_{u^{(r)}}(\omega,a)R_{[-M^r,M^r]^{b+1}})^{-1}(n,j;n',j')| \le 2\delta^{-\frac{1}{8}}e^{-c\max\{|n-n'|,|j-j'|\}}.$$
By (141)-(144) and resolvent expansion as in Lemma 5.1 [8] one has that of

By (141)-(144) and resolvent expansion as in Lemma 5.1 [8], one has that on $(\omega, a) \in (\bigcup_{I \in \Lambda_r} I) \cap \Gamma_r, \ \omega \in (I_1 \setminus (I_1^r \cup \tilde{I}_1^r))$

(145)
$$||(R_{[-M^{r+1},M^{r+1}]^{b+1}}\tilde{T}_{u^{(r)}}(\omega,a)R_{[-M^{r+1},M^{r+1}]^{b+1}})^{-1}|| \le \delta^{-\frac{1}{8}}M^{(r+1)^{C}},$$

and for any (n, j) and (n', j') such that $\max\{|n - n'|, |j - j'|\} \ge (r+1)^C$, (146)

$$|(R_{[-M^{r+1},M^{r+1}]^{b+1}}\tilde{T}_{u^{(r)}}(\omega,a)R_{[-M^{r+1},M^{r+1}]^{b+1}})^{-1}(n,j;n',j')| \le \delta^{-\frac{1}{8}}e^{-c\max\{|n-n'|,|j-j'|\}}.$$

In order to have (5) in Hiv, we have to remove ω of measure less than $\delta^2 M^{-\frac{r}{2}}$ from scales M^r to $M^{(r+1)}$ for $r \geq r_0$, and of measure less than $\delta^2 e^{-|\log \delta|^{K_1^{90}}}$ from scales $|\log \delta|^K$ to M^{r_0} . By counting all possible intervals *I*, and by (135), (138), one has that for $r \geq r_0$,

(147)
$$\operatorname{Leb}(P_{\omega}(\Gamma_{r} \cap (\bigcup_{I' \in \Lambda_{r}} I' \setminus \bigcup_{I \in \Lambda_{r+1}} I))) \leq \delta M^{r_{1}^{C}} M^{-\frac{r}{2^{b-1}}} + \delta^{2} M^{-\frac{r}{2}},$$

and for $r = r_0 - 1$,

(148)
$$\operatorname{Leb}(P_{\omega}(\Gamma_{r} \cap (\bigcup_{I' \in \Lambda_{r}} I' \setminus \bigcup_{I \in \Lambda_{r+1}} I))) \leq \delta M^{r_{1}^{C}} M^{-\frac{r}{2^{b-1}}} + \delta^{2} e^{-|\log \delta|^{K_{1}^{90}}}.$$

This implies that for $r \ge r_0$

(149)
$$\operatorname{Leb}(P_a(\Gamma_r \cap (\bigcup_{I' \in \Lambda_r} I' \setminus \bigcup_{I \in \Lambda_{r+1}} I))) \le M^{-\frac{r}{2b}},$$

and for $r = r_0 - 1$

(150)
$$\operatorname{Leb}(P_a(\Gamma_r \cap (\bigcup_{I' \in \Lambda_r} I' \setminus \bigcup_{I \in \Lambda_{r+1}} I))) \le M^{-\frac{r}{2^b}} + e^{-|\log \delta|^{K_1^{90}}}.$$

We have completed the proof.

Proof of main theorem. Theorem 1.1 follows immediately from the (by now verified) hypothesis (Hi-iv). \Box

APPENDIX A. EIGENFUNCTION RELABELLING MAP

For a fixed $V \in \mathcal{V}_{\epsilon}$, basing on (58), one may relabel the eigenfunctions in a more intrinsic way. Write φ_i and ι_i for φ_i^V and ι_i^V , since V is fixed. The goal is that in the new labelling scheme, if j > j', then the localization centers of the corresponding eigenfunctions ϕ_j and $\phi_{j'}$ satisfy $\ell_{j'} \ge \ell_j$. Below we provide such a relabelling map.

For a given eigenfunction φ_i , we first select a vertex among the set of vertices, on which φ_i achieves its maximum. (This selection could be arbitrary, but it is practical to have a rule.) So for $i \in \mathbb{Z}$, let

$$\mathcal{M}_i = \{ x_0 \in \mathbb{Z} : |\varphi_i(x_0)| = \max_{x \in \mathbb{Z}} |\varphi_i(x)| \}.$$

Define $\mathcal{M}_i^+ = \mathcal{M}_i \cap \{\{0\} \cup \mathbb{Z}_+\}$. If $\mathcal{M}_i^+ \neq \emptyset$, define $\iota_i = \min x_0, x_0 \in \mathcal{M}_i^+$; otherwise define $\iota_i = \min -x_0, x_0 \in \mathcal{M}_i$.

Define

$$f_1: \mathbb{Z} \mapsto \mathbb{Z}, f_1(i) = \iota_i.$$

Let \mathcal{L} be the range of f_1 ,

$$\mathcal{L} = \operatorname{Ran}(f_1), \mathcal{L} \subseteq \mathbb{Z}.$$

For a given $l \in \mathcal{L}$, let $I_l = \{i | \iota_i = l\}$. Define

$$f_2: \mathcal{L} \mapsto \bigsqcup_{l \in \mathcal{L}} I_l.$$

From (58): If $|l| < l_{\epsilon}$, $\#\{\bigcup_{|l| \le l_{\epsilon}} I_l\} \le (1+\epsilon)l_{\epsilon}$; and if $|l| \ge l_{\epsilon}$,

$$1-\epsilon)l \le \#\{\bigcup_{|l|\le l_{\epsilon}} I_l\} \le (1+\epsilon)l.$$

The upper bound gives that for all l, I_l is finite. So we may define a map

$$f_3:\bigsqcup_{l\in\mathcal{L}}I_l\mapsto\mathbb{Z},$$

such that if $x \in I_l$ and $y \in I_{l'}$, with l' > l, then $f_3(y) > f_3(x)$.

Finally define the map f to be $f = f_3 \circ f_2 \circ f_1$,

 $f:\mathbb{Z}\mapsto\mathbb{Z}.$

Using the relabelling map f yields our ortho-normal eigen-basis $\{\phi_j\}_{j\in\mathbb{Z}}$.

Appendix B. Proof of Lemma 3.2

Proof. By Theorem 3.1 and Chebyshev's inequality, one has that for any ϵ_1 , there exists \mathcal{V}_{ϵ_1} with $\mathbb{P}(\mathcal{V}_{\epsilon_1}) > 1 - \epsilon_1$ such that for any $V \in \mathcal{V}_{\epsilon_1}$,

(151)
$$|\varphi_j^V(\ell)| \le C_{\epsilon_1} (1+|\iota_j^V|)^q e^{-\gamma_1|\ell-\iota_j^V|}$$

For simplicity, below we drop the superscript V. Clearly, we have

(152)
$$\sum_{\ell \in \mathbb{Z}} |\varphi_j(\ell)|^2 = 1$$

and

(153)
$$\sum_{j\in\mathbb{Z}} |\varphi_j(\ell)|^2 = 1.$$

Let ϵ be an arbitrarily small constant. Assume L is large enough, depending on ϵ and ϵ_1 . If $k \leq \iota_j \leq k + L$ with $|k| \leq L^4$, then

(154)
$$\sum_{\ell \le k - \epsilon L} |\varphi_j(\ell)|^2 + \sum_{\ell \ge k + (1+\epsilon)L} |\varphi_j(\ell)|^2 \le \sum_{m \ge \epsilon L} C_{\epsilon_1} L^{8q} e^{-\gamma_1 m}$$
(155)
$$\le C e^{-\epsilon L},$$

where C depends on ϵ_1 and ϵ . By (152) and (155), one has that for any j with $k \leq \iota_j \leq k + L$,

(156)
$$\sum_{k-\epsilon L \le \ell \le k+(1+\epsilon)L} |\varphi_j(\ell)|^2 \ge 1 - Ce^{-\epsilon L}.$$

By (153) and (156), we have that

(157)
$$(1+\epsilon)L \ge \sum_{\substack{k-\epsilon L \le \ell \le k+(1+\epsilon)L\\ j \in \mathbb{Z}}} |\varphi_j(\ell)|^2$$

(158)
$$\geq \sum_{\substack{k-\epsilon L \leq \ell \leq k+(1+\epsilon)L\\ j \in \mathbb{Z}: k \leq \iota_j \leq k+L}} |\varphi_j(\ell)|^2$$

(159)
$$\geq (1 - Ce^{-\epsilon L}) \# \{j : k \leq \iota_j \leq k + L\}.$$

This implies that for any $k \in [-L^4, L^4]$,

(160)
$$\#\{j: k \le \iota_j \le k+L\} \le (1+\epsilon)L.$$

For any $\ell \in [k + \epsilon L, k + (1 - \epsilon)L]$ with $|k| \leq L^4$, by (151), one has

(161)

$$\sum_{j \in \mathbb{Z}: \iota_j \notin [k, k+L]} |\varphi_j(\ell)|^2 \leq \sum_{m=0}^{\infty} \sum_{j \in \mathbb{Z}: mL \leq |\iota_j| \leq (m+1)L} C(1+|\iota_j|)^{2q} e^{-\gamma_1 |\ell - \iota_j|}$$

$$\leq \sum_{m \leq 10L^4} \sum_{j \in \mathbb{Z}: |\iota_j| \leq 20L^5} CL^{10q} e^{-\gamma_1 \epsilon L} + \sum_{m=10L^4}^{\infty} C(1+mL)^{2q} \#\{j: |\iota_j| \leq (m+1)L\} e^{-\frac{1}{2}\gamma_1 mL}$$

$$\leq Ce^{-\epsilon L} + \sum_{m=1}^{\infty} C(1+mL)^{3q} e^{-\frac{1}{2}\gamma_1 mL}$$

$$\leq Ce^{-\epsilon L},$$

where (161) holds by (160). It implies that

(162)
$$\sum_{\substack{j \in \mathbb{Z}: \iota_j \notin [k,k+L]\\ \ell \in [k+\epsilon L, k+(1-\epsilon)L]}} |\varphi_j(\ell)|^2 \le C e^{-\epsilon L}.$$

By (153) and (162), one has that

$$(1-\epsilon)L = \sum_{\ell \in [k+\epsilon L, k+(1-\epsilon)L], j \in \mathbb{Z}} |\varphi_j(\ell)|^2$$

$$\leq Ce^{-\epsilon L} + \sum_{\substack{j \in \mathbb{Z}: \iota_j \in [k, k+L]\\\ell \in [k+\epsilon L, k+(1-\epsilon)L]}} |\varphi_j(\ell)|^2$$

$$\leq Ce^{-\epsilon L} + \#\{j : k \leq \iota_j \leq k+L\}.$$

This yields that for any k with $|k| \leq L^4$,

(163)
$$\#\{j: k \le \iota_j \le k+L\} \ge (1-\epsilon)L.$$

Now (58) follows from (160) and (163).

APPENDIX C. BOURGAIN'S INDUCTION ESTIMATES

Assume that Hi-v hold at step r, and that Hi-v hold at step r + 1 except for (123). We show that (123) holds at step r + 1. This follows from Chap. 18 [5] with minor modifications.

Let us state the difference between our setting and that of Bourgain's. Bourgain assumed that

(164)
$$\| (R_{[-M^r,M^r]^{b+1}} \tilde{T}_{u^{(r-1)}} R_{[-M^r,M^r]^{b+1}})^{-1} \| \le M^{r^C},$$

and that for (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} > r^C$, (165) $|(R_{[-M^r, M^r]^{b+1}} \tilde{T}_{u^{(r-1)}} R_{[-M^r, M^r]^{b+1}})^{-1} (n, j; n', j')| \le e^{-c \max\{|n - n'|, |j - j'|\}}$. Then he proved that the induction holds if

$$\begin{split} \delta_{r+1} &\geq M^{(r+1)^C} \kappa_r, \\ \bar{\delta}_{r+1} &\geq M^{2(r+1)^C} \bar{\kappa}_r + M^{(r+1)^C} \delta_{r+1}, \\ \kappa_{r+1} &\geq e^{-\frac{c}{3}M^{r+1}} \kappa_r + \delta_{r+1}^2, \\ \bar{\kappa}_{r+1} &\geq M^{2(r+1)^C} \kappa_r + e^{-\frac{c}{3}M^{r+1}} \bar{\kappa}_r + \delta_{r+1} \bar{\delta}_{r+1}. \end{split}$$

We assume that for a proper $0 < \nu < 1$,

(166)
$$\| (R_{[-M^r,M^r]^{b+1}} \tilde{T}_{u^{(r-1)}} R_{[-M^r,M^r]^{b+1}})^{-1} \| \le \delta^{-\nu} M^{r^C},$$

and that for (n, j) and (n', j') satisfying $\max\{|n - n'|, |j - j'|\} > r^C$, (167)

$$|(R_{[-M^r,M^r]^{b+1}}\tilde{T}_{u^{(r-1)}}R_{[-M^r,M^r]^{b+1}})^{-1}(n,j;n',j')| \le \delta^{-\nu}e^{-c\max\{|n-n'|+|j-j'|\}}.$$

Following Bourgain's proof, we obtain the following new relations

$$\begin{split} \delta_{r+1} &\geq \delta^{-\nu} M^{(r+1)^C} \kappa_r, \\ \bar{\delta}_{r+1} &\geq \delta^{-2\nu} M^{2(r+1)^C} \bar{\kappa}_r + \delta^{-\nu} M^{(r+1)^C} \delta_{r+1}, \\ \kappa_{r+1} &\geq \delta^{1-\nu} e^{-\frac{c}{3} M^{r+1}} \kappa_r + \delta_{r+1}^2, \\ \bar{\kappa}_{r+1} &\geq \delta^{-2\nu} M^{2(r+1)^C} \kappa_r + \delta^{1-\nu} e^{-\frac{c}{3} M^{r+1}} \bar{\kappa}_r + \delta_{r+1} \bar{\delta}_{r+1} \end{split}$$

For example, we may take $\nu = \frac{1}{8}$

$$\delta_r = \delta^{\frac{1}{2}} M^{-(\frac{4}{3})^r}, \bar{\delta}_r = \delta^{\frac{1}{8}} M^{-\frac{1}{2}(\frac{4}{3})^r}, \kappa_r = \delta^{\frac{3}{4}} M^{-(\frac{4}{3})^{r+2}}, \bar{\kappa}_r = \delta^{\frac{3}{8}} M^{-\frac{1}{2}(\frac{4}{3})^{r+2}}, \bar{\kappa}_r = \delta^{\frac{3}{8}} M$$

Acknowledgments

W. Liu was supported by NSF DMS-2000345 and DMS-2052572. W.-M. Wang acknowledges support from CY Initiative of Excellence (Grant "Investissements d'Avenir" ANR-16-IDEX-0008).

References

- M. Aizenman and S. Molchanov. Localization at large disorder and at extreme energies: an elementary derivation. *Commun. Math. Phys.*, 157:245– 278, 1993.
- [2] S. Basu. On bounding the Betti numbers and computing the Euler characteristic of semi-algebraic sets. *Discrete Comput. Geom.*, 22(1):1–18, 1999.
- [3] G. Binyamini and D. Novikov. Complex cellular structures. Ann. of Math., 190(2):145–248, 2019.

- [4] J. Bourgain. Quasi-periodic solutions of Hamiltonian perturbations of 2d linear Schrödinger equations. Ann. of Math., 148:363–439, 1998.
- [5] J. Bourgain. Green's function estimates for lattice Schrödinger operators and applications, volume 158 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2005.
- [6] J. Bourgain. Anderson localization for quasi-periodic lattice Schrödinger operators on Z^d, d arbitrary. Geom. Funct. Anal., 17(3):682–706, 2007.
- [7] J. Bourgain, M. Goldstein, and W. Schlag. Anderson localization for Schrödinger equations on Z² with quasi-periodic potential. Acta Math., 188:41–86, 2002.
- [8] J. Bourgain and W.-M. Wang. Quasi-periodic solutions of nonlinear random Schrödinger equations. J. Eur. Math. Soc., 10:1–45, 2008.
- [9] V. Bucaj, D. Damanik, J. Filman, V. Gerbuz, T. VandenBoom, F. Wang, and Z. Zhang. Localization for the one-dimensional Anderson model via positivity and large deviations for the Lyapunov exponent. *Trans. Amer. Math. Soc.*, 372:3619–3667, 2019.
- [10] W. Craig and C. E. Wayne. Newton's method and periodic solutions of nonlinear equations. Commun. Pure Appl. Math., 46:1409–1498, 1993.
- [11] R. del Rio, S. Jitomirskaya, Y. Last, and B. Simon. Operators with singular continuous spectrum. IV. Hausdorff dimensions, rank one perturbations, and localization. J. Anal. Math., 69:153–200, 1996.
- [12] A. Elgart and A. Klein. An eigensystem approach to Anderson localization. J. Funct. Anal., 271:3465–3512, 2016.
- [13] J. Fröhlich and T. Spencer. Absence of diffusion in the Anderson tight binding model. *Commun. Math. Phys.*, 88:151–184, 1983.
- [14] J. Fröhlich, T. Spencer, and C. E. Wayne. Localization in disordered, nonlinear dynamical systems. J. Stat. Phys., 42(3/4):247–274, 1986.
- [15] H. Furstenberg. Non-commuting random products. Trans. Amer. Math. Soc., 108:377–428, 1963.
- [16] L. Ge and X. Zhao. Exponential dynamical localization in expectation for the one dimensional Anderson model. J. Spectr. Theory., 10(3):887–904, 2020.
- [17] F. Germinet and A. Klein. New characterization of the region of complete localization for random Schrödinger operators. J. Stat. Phys., 122:73–94, 2006.
- [18] F. Germinet and F. Klopp. Spectral statistics for random Schrödinger operators in the localized regime. J. Eur. Math. Soc. (JEMS), 16(9):1967–2031, 2014.
- [19] I. Goldshied, S. Molchanov, and L. Pastur. A pure point spectrum of the stochastic one-dimensional Schrödinger equation. *Funct. Anal. Appl.*, 11:1– 10, 1977.

- [20] A. Gorodetski and V. Kleptsyn. Parametric Furstenburg theorem on random products of SL(2, ℝ) matrices. Adv. Math., 378:107522, 2021.
- [21] S. Jitomirskaya and X. Zhu. Large deviation of the Lyapunov exponent and localization for the 1d Anderson model. *Commun. Math. Phys.*, 370:311– 324, 2019.
- [22] H. Kunz and B. Souillard. Sur le spectre des opérateurs aux differences finies aléatoire. Commun. Math. Phys., 78:201–246, 1980.
- [23] W. Liu. Quantitative inductive estimates for Green's functions of non-selfadjoint matrices. to appear in Anal. and PDE; arXiv:2007.00578, 2020.
- [24] D. Macera and S. Sodin. Anderson localization for quasi-one-dimensional random operators. arXiv: 2110.00097, 2021.
- [25] N. Minami. Local fluctuations of a multidimensional Anderson tight binding model. Commun. Math. Phys., 177:709–725, 1996.
- [26] H. von Dreifus and A. Klein. A new proof of localization in the Anderson tight binding model. *Commun. Math. Phys.*, 124:285–299, 1989.
- [27] W.-M. Wang. Energy supercritical nonlinear Schrödinger equations: Quasiperiodic solutions. Duke Math. J., 165(6):1129–1192, 2016.
- [28] W.-M. Wang. Quasi-periodic solutions to a nonlinear Klein-Gordon equation with a decaying nonlinear term. arXiv: 1609.00309, 2021.
- [29] F. Wegner. Bounds on the density of states in disordered systems. Z. Phys. B., 44:9–15, 1981.
- [30] Z. Zhao and J. Geng. Linearly stable quasi-periodic breathers in a class of random Hamiltonian systems. J. Dyn. Diff. Equat., 23:961–997, 2011.

(W. Liu) Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

Email address: liuwencai1226@gmail.com; wencail@tamu.edu

(W.-M. Wang) CNRS AND DÉPARTEMENT DE MATHÉMATIQUE, CERGY PARIS UNIVER-SITÉ, 95302 CERGY-PONTOISE CEDEX, FRANCE Email address: wei-min.wang@math.cnrs.fr