ON INVARIANT OF THE REGULARITY INDEX OF FAT POINTS

Phan Van Thien Department of Mathematics, University of Education, Hue University, Vietnam E-mail: pvthien@hueuni.edu.vn

Abstract

We prove invariant of the regularity index of fat points under changes of the linear subspace containing the support of the fat points (Theorem 3.5). Then we show that Segre's bound is attained by any set of s non-degenerate equimultiple fat points in \mathbb{P}^n , $s \leq n+3$ (Theorem 4.3). We also give an example showing that there always exists a set of n+4 non-degenerate equimultiple fat points in \mathbb{P}^n such that Segre's bound is not attained (Example 4.4).

Key words and phrases. Regularity index; Fat points. 2010 Mathematics Subject Classification. Primary 14C20; Secondary 13D40.

1 Introduction

In this paper, we denote by $\mathbb{P}^n := \mathbb{P}_K^n$ the *n*-dimensional projective space over an algebraically closed K of arbitrary characteristic, and denote by $R := K[X_0, \ldots, X_n]$ the polynomial ring in variables X_0, \ldots, X_n over K. If P_1, \ldots, P_s are distinct points in \mathbb{P}^n ; we denote by \wp_1, \ldots, \wp_r the defining homogeneous prime ideals of P_1, \ldots, P_s in R, respectively.

Let P_1, \ldots, P_s be distinct points in \mathbb{P}^n , and let m_1, \ldots, m_s be positive integers. Then the set of all homogeneous polynomilas that vanish at P_i to order m_i ; for $i = 1, \ldots, s$; is the homogeneous ideal $I := \wp_1^{m_1} \cap \cdots \cap \wp_s^{m_s}$. We call the zero-scheme defined by I to be fat points (or a set of fat points) in \mathbb{P}^n , and we denote it by

$$Z := m_1 P_1 + \dots + m_s P_s$$

The integer m_i is called the multiplicity of P_i ; i = 1, ..., s. If $m_1 = \cdots = m_s = 2$, then fat points $Z = 2P_1 + \cdots + 2P_s$ is said to be double points. If $m_1 = \cdots = m_s = m$, then fat points $Z = mP_1 + \cdots + mP_s$ is said to be equimultiple. We call the points P_i to be the points of support of Z. If the support of Z span \mathbb{P}^n ; then P_1, \ldots, P_s is called non-degenerate in \mathbb{P}^n ; and we also call Z a set of non-degenerate fat points in \mathbb{P}^n .

The homogeneous coordinate ring R/I of Z is a graded ring, $R/I = \bigoplus_{t \ge 0} (R/I)_t$. We call the function

$$H_{R/I}(t) := \dim_K (R/I)_t$$

to be the Hilbert function of Z in \mathbb{P}^n . Since

$$\dim_K I_t = \dim_K R_t - \dim_K (R/I)_t = \binom{t+n}{n} - H_{R/I}(t),$$

we know the dimension of linear space of all homogeneous polynomials of degree t vanishing at P_i to order m_i ; $i = 1, \ldots, s$; if we know the Hilbert function $H_{R/I}(t)$. But it is a fairly difficult problem to determine the Hilbert function of fat points in \mathbb{P}^n .

ON INVARIANT OF THE REGULARITY INDEX OF FAT POINTS

The ring R/I has the multiplicity

$$e(R/I) := \sum_{i=1}^{s} \binom{m_i + n - 1}{n}.$$

The Hilbert function $H_{R/I}(t)$ strictly increases until it reaches the multiplicity e(R/I), at which it stabilizes. We call the least integer t such that $H_{R/I}(t) = e(R/I)$ to be the regularity index of Z in \mathbb{P}^n , and we denote it by $\operatorname{reg}(Z)$. It is well known that $\operatorname{reg}(Z) = \operatorname{reg}(R/I)$, the Castelnuovo-Mumford regularity of the ring R/I.

If $t \geq \operatorname{reg}(Z)$, we have

$$\dim_K I_t = \binom{t+n}{n} - \sum_{i=1}^s \binom{m_i+n-1}{n}.$$

So, it is interesting to determine reg(Z) or, even less, an upper bound for it.

From now on if a is a rational number, we denote by [a] it's integer part.

For generic fat points $Z = m_1 P_1 + \cdots + m_s P_s$ in \mathbb{P}^2 with $m_1 \ge \cdots \ge m_s$, Segre [8] showed that

$$\operatorname{reg}(Z) \le \max\left\{m_1 + m_2 - 1, \left[\frac{m_1 + \dots + m_s}{2}\right]\right\}.$$

It was conjectured by Trung (see [9]) and, independently, by Farabbi and Lorenzini (see [6]) that

$$\operatorname{reg}(R/I) \le \max\{T_j(Z) | j = 1, \dots, n\},\$$

where

$$T_j(Z) := \max\left\{ \left[\frac{\sum_{l=1}^q m_{i_l} + j - 2}{j} \right] \mid P_{i_1}, \dots, P_{i_q} \text{ on a } j \text{-dimensional linear subspace} \right\}$$

The number

$$T(Z) := \max\{T_j(Z) | j = 1, \dots, n\}$$

is called the Segre's bound because it generalizes Segre's upper bound. There were many different results for proving the Segre's bound. In 2016, Ballico et al. [2] successfully proved the Segre's bound for n + 3 non-degenerate fat points in \mathbb{P}^n . Recently, Nagel and Trok (see [7, Theorem 5.3]) have succefully proved the Segre's bound for arbitrary fat points Z in \mathbb{P}^n .

The problem to exactly determine reg(R/I) is more fairly difficult. So far, there are only a few results about computing reg(R/I).

For abitrary fat points $Z = m_1 P_1 + \cdots + m_s P_s$ in \mathbb{P}^n , Davis and Geramita [5, Corollary 2.3] proved that

$$\operatorname{reg}(R/I) = m_1 + \dots + m_s - 1$$

if and only if points P_1, \ldots, P_s lie on a line in \mathbb{P}^n .

The points P_1, \ldots, P_s in \mathbb{P}^n is said to be in linearly general position if no j + 2 of the points P_1, \ldots, P_s are on any *j*-dimensional linear subspace for j < n, then we also call the fat points $Z = m_1 P_1 + \cdots + m_s P_s$ to be in general position in \mathbb{P}^n . A rational normal curve in \mathbb{P}^n to be a curve of degree *n* that may be given parametrically as the image of the map

$$\mathbb{P}^1 \to \mathbb{P}^n.$$

(s,t) $\mapsto (s^n, s^{n-1}t, \dots, t^n)$

For fat points $Z = m_1 P_1 + \cdots + m_s P_s$ in \mathbb{P}^n with $m_1 \ge \cdots \ge m_s$, Catalisano et al. [4] showed formulas to compute $\operatorname{reg}(Z)$ in the following two cases:

If $s \geq 2$ and P_1, \ldots, P_s are on a rational normal curve in \mathbb{P}^n [4, Proposition 7], then

$$\operatorname{reg}(R/I) = \max\left\{m_1 + m_2 - 1, \left[(\sum_{i=1}^s m_i + n - 2)/n\right]\right\}.$$

If
$$n \geq 3, 2 \leq s \leq n+2, 2 \leq m_1 \geq \cdots \geq m_s$$
 and P_1, \ldots, P_s are in general position in \mathbb{P}^n , then

$$\operatorname{reg}(R/I) = m_1 + m_2 - 1.$$

For fat points $Z = m_1 P_1 + \cdots + m_{s+2} P_{s+2}$ in \mathbb{P}^n with P_1, \ldots, P_{s+2} not in a linear (s-1)-space in \mathbb{P}^n , $s \leq n$, Thien [10, Theorem 3.4] showed

$$\operatorname{reg}(R/I) = T(Z).$$

For equimutiple fat points $Z = mP_1 + \cdots + mP_s$ in \mathbb{P}^n with $m \neq 2$; P_1, \ldots, P_s are not on a linear (r-1)-space and $s \leq r+3$, Thien and Sinh [11, Theorem 4.5] showed

$$\operatorname{reg}(R/I) = T(Z).$$

Assume that α is a *r*-dimensional linear subspace of \mathbb{P}^n containing the points P_1, \ldots, P_s . We may consider the *r*-dimensional linear subspace α as a projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and we consider $Z_\alpha := m_1 P_{1\alpha} + \cdots + m_s P_{s\alpha}$ as a zero-scheme in \mathbb{P}^r . What is the relation between $\operatorname{reg}(Z)$ and $\operatorname{reg}(Z_\alpha)$?

In this paper we prove that $\operatorname{reg}(Z) = \operatorname{reg}(Z_{\alpha})$ (Theorem 3.5). Then we calculate regularity index of a set of n+3 non-degenerate double points in \mathbb{P}^n (Proposition 4.2) and show that Segre's bound is attained by any set of s non-degenerate equimultiple fat points in \mathbb{P}^n , $s \leq n+3$ (Theorem 4.3). We also show that there always exists a set of n+4 non-degenerate equimultiple fat points in \mathbb{P}^n such that Segre's bound is not attained (Corollary 4.5).

2 Preliminaries

We use the following lemmas which have been proved yet. The first lemma allows us to compute the regularity index by induction.

Lemma 2.1. [4, Lemma 1] Let P_1, \ldots, P_r, P be distinct points in \mathbb{P}^n and \wp be the defining ideals in R corresponding to the point P. If m_1, \ldots, m_r , a are positive integers, $J := \wp_1^{m_1} \cap \cdots \cap \wp_r^{m_r}$, and $I = J \cap \wp^a$, then

$$\operatorname{reg}(R/I) = \max\left\{a - 1, \operatorname{reg}(R/J), \operatorname{reg}(R/(J + \wp^a))\right\}$$

Note that $R/(J + \wp^a)$ is a zero-dimensional graded ring algebra, the regularity index reg $(R/(J + \wp^a))$ is the least integer such that $[R/(J + \wp^a))]_t = 0$. To estimate reg $(R/(J + \wp^a))$ we shall use the following lemma.

Lemma 2.2. [4, Lemma 3] Let P_1, \ldots, P_r, P be distinct points in \mathbb{P}^n and \wp be the defining ideal in R corresponding to the point P. If m_1, \ldots, m_r, a are positive integers, $J = \wp_1^{m_1} \cap \cdots \cap \wp_r^{m_r}$ and $\wp = (X_1, \ldots, X_n)$, then

$$\operatorname{reg}(R/(J+\wp^a)) \le b$$

if and only if $X_0^{b-i}M \in J + \wp^{i+1}$ for every monomial M of degree i in X_1, \ldots, X_n ; $i = 0, \ldots, a-1$.

Lemma 2.3. [4, Corollary 2] Let $s \ge 2$; P_1, \ldots, P_s be distinct points in \mathbb{P}^n and $m_1 \ge \cdots \ge m_s$ be positive integers. If $I = \wp_1^{m_1} \cap \cdots \cap \wp_s^{m_s}$, then

$$\operatorname{reg}(R/I) \ge m_1 + m_2 - 1.$$

Lemma 2.4. [4, Corollary 8] Let $n \ge 3$, $2 \le s \le n+2$, and let P_1, \ldots, P_s be distinct points in general position in \mathbb{P}^n . If $2 \le m_1 \ge \cdots \ge m_s > 0$ are given integers and $I = \wp_1^{m_1} \cap \cdots \cap \wp_s^{m_s}$, then

$$\operatorname{reg}(R/I) = m_1 + m_2 - 1.$$

In [2], Ballico et al. successfully proved the Segre's for non-degenerate n+3 fat points in \mathbb{P}^n .

Lemma 2.5. [2, Theorem 2.1] Let $Z = m_1P_1 + \cdots + m_{n+3}P_{n+3}$ be a scheme of fat points supported on a non-degenerate set of distinct points in \mathbb{P}^n . Then Z satisfies the Segre's bound, namely

$$\operatorname{reg}(Z) \le \max\{T(Z,L) : L \subseteq \mathbb{P}^n\}$$

where L is a j-dimensional linear subspace of \mathbb{P}^n ; j = 1, ..., n; and

$$T(Z,L) := \max\left\{ \left[\frac{\sum_{l=1}^{q} m_{i_l} + j - 2}{j} \right] | P_{i_1}, \dots, P_{i_q} \in L \right\}.$$

We see that the number $\max\{T(Z,L) : L \subseteq \mathbb{P}^n\}$ is the Segre's bound T(Z) in the Trung's conjecture.

The following lemma was proved in [10], it also has been shown in [7, Lemma 3.1].

Lemma 2.6. [10, Lemma 3.3] Let $X = \{P_1, \ldots, P_s\}$ be a set of distinct points in \mathbb{P}^n and m_1, \ldots, m_s be positive integers. Put $I = \wp_1^{m_1} \cap \cdots \cap \wp_s^{m_s}$. If Y is a subset of X and $J = \bigcap_{P_i \in V} \wp_i^{m_i}$, then

 $\operatorname{reg}(R/J) \le \operatorname{reg}(R/I).$

This implies that if $Z = m_1 P_1 + \cdots + m_s P_s$ is the set of fat points defining by I, and $U = \sum_{P_i \in Y} m_i P_i$ is the set of fat points defining by J, then

$$\operatorname{reg}(U) \le \operatorname{reg}(Z).$$

Lemma 2.7. [10, Theorem 3.4] Let P_1, \ldots, P_{s+2} be distinct points not on a (s-1)-dimensional linear subspace in \mathbb{P}^n ; $s \leq n$; and m_1, \ldots, m_{s+2} be positive integers. Put $I = \wp_1^{m_1} \cap \cdots \cap \wp_{s+2}^{m_{s+2}}$. Then,

$$\operatorname{reg}(R/I) = \max\{T_j | j = 1, \dots, n\},\$$

where

$$T_j := \max\left\{ \left[\frac{\sum_{l=1}^q m_{i_l} + j - 2}{j} \right] \mid P_{i_1}, \dots, P_{i_q} \text{ on a } j \text{-dimensional linear subspace} \right\}.$$

If P_1, \ldots, P_s are on a *r*-dimensional linear subspace $\alpha \subset \mathbb{P}^n$ and there are no j + 2 of the points P_1, \ldots, P_s on any *j*-dimensional linear subspace for j < r; then we call P_1, \ldots, P_s to be in linearly general position on α . The following lemmas were proved in [11].

Lemma 2.8. [11, Theorem 3.1] Let P_1, \ldots, P_s be distinct points in general position on an rdimensional linearly linear subspace α in \mathbb{P}^n , $s \leq r+3$. Let m_1, \ldots, m_s be positive integers and $Z = m_1P_1 + \cdots + m_sP_s$. Then,

$$\operatorname{reg}(Z) = \max\{T_1, T_r\},\$$

where

$$T_{1} = \max\{m_{i} + m_{j} - 1 \mid i \neq j; i, j = 1, \dots, s\},\$$
$$T_{r} = \left[\frac{m_{1} + \dots + m_{s} + r - 2}{r}\right].$$

We see that the Segre's bound is attained for this case. If r = n and s = n + 3, the P_1, \ldots, P_{n+3} are in linearly general position in \mathbb{P}^n . So, there exists a rational normal curve passing through P_1, \ldots, P_{n+3} . Then we get Proposition 7 in [4].

Lemma 2.9. [11, Corollary 3.2] Let P_1, \ldots, P_s be distinct points in \mathbb{P}^n , $s \leq 5$. Let m be a positive integer and $Z = mP_1 + \cdots + mP_s$. Then

$$\operatorname{reg}(Z) = \max\{T_j | j = 1, \dots, n\},\$$

where

$$T_j := \max\left\{ \left[\frac{qm+j-2}{j} \right] \mid P_{i_1}, \dots, P_{i_q} \text{ on a } j \text{-dimensional linear subspace} \right\}.$$

Lemma 2.10. [11, Theorem 4.5] Let $X = \{P_1, \ldots, P_{s+3}\}$ be a set of distinct points not on a (s-1)-dimensional linear subspace in \mathbb{P}^n , $s \leq n$ and m be a positive integer, $m \neq 2$. Let Z = $mP_1 + \cdots + mP_{s+3}$ be a set equimultiple fat points. Then,

$$\operatorname{reg}(Z) = \max\{T_j | j = 1, \dots, n\}$$

where

$$T_{j} = \max\left\{ \left[\frac{mq+j-2}{j} \right] \mid P_{i_{1}}, \dots, P_{i_{q}} \text{ lie on a linear } j\text{-space} \right\}$$

We see that in case s = n and $m \neq 2$, the regularity index of n + 3 non-degenerate equimultiple fat points $Z = m1P_1 + \cdots + mP_{n+3}$ in \mathbb{P}^n reaches the Segre's bound.

Invariant of the regularity index under changes of linear 3 subspace containing the support of fat points

Let $Z = m_1 P_1 + \cdots + m_s P_s$ be a set of fat points in \mathbb{P}^n , and I be the ideal consists all homogeneous polynomials in R vanishing at $P_i \in \mathbb{P}^n$ to order m_i ; $i = 1, \ldots, s$. The regularity index reg(Z) is the least integer such that the Hilbert function

$$H_{R/I}(t) = \dim_K (R/I)_t = \binom{t+n}{n} - \dim_K I_t$$

reaches the multiplicity $e(R/I) = \sum_{i=1}^{s} {\binom{m_i+n-1}{n}}$. Note that the Hilbert function of an ideal is invariant under projective automorphisms. So, the regularity index of a set of fat points is also invariant under projective automorphisms.

Assume that α is a r-dimensional linear subspace of \mathbb{P}^n containing the points P_1, \ldots, P_s . We consider the r-dimensional linear subspace α as a projective space \mathbb{P}^r containing the points $P_{1\alpha} :=$ $P_1, \ldots, P_{s\alpha} := P_s$, and consider $Z_{\alpha} := m_1 P_{1\alpha} + \cdots + m_s P_{s\alpha}$ as a set of fat points in \mathbb{P}^r . We denote by I_{α} the ideal consists all homogeneous polynomials in $R_{\alpha} := K[X_0, \ldots, X_r]$ that vanish at $P_{i\alpha}$ to order m_i ; $i = 1, \ldots, s$. The Hilbert function of Z_{α} in \mathbb{P}^r is

$$H_{Z_{\alpha}}(t) := H_{R_{\alpha}/I_{\alpha}}(t) = \dim_{K}(K[X_{0},\ldots,X_{r}]/I_{\alpha})_{t} = \binom{t+r}{r} - \dim_{K}(I_{\alpha})_{t}.$$

The multiplicity number of Z_{α} in \mathbb{P}^r is

$$e(R_{\alpha}/I_{\alpha}) := \sum_{i=1}^{s} \binom{m_i + r - 1}{r}.$$

The regularity index of Z_{α} in \mathbb{P}^r is the least integer t such that $H_{R/I_{\alpha}}(t) = e(R_{\alpha}/I_{\alpha})$, and we denote it by $\operatorname{reg}(Z_{\alpha})$.

We have $e(R_{\alpha}/I_{\alpha}) \leq e(R/I)$, $\binom{t+r}{r} \leq \binom{t+n}{n}$ and $\dim_{K}(I_{\alpha})_{t} \leq \dim_{K}I_{t}$. Here there are two questions:

- 1) Question 1: $H_{Z_{\alpha}}(t) = H_Z(t)$?
- 2) Question 2: $\operatorname{reg}(Z_{\alpha}) = \operatorname{reg}(Z)$?

For Question 1, we now do not know about the relation between $H_Z(t)$ and $H_{Z_\alpha}(t)$ if $t < \operatorname{reg}(Z)$. But if $t \geq \operatorname{reg}(Z)$, then $H_{Z_{\alpha}}(t) \leq H_{Z}(t)$. Furthermore, if there exists $m_i \geq 2$ and r < n, then $H_{Z_{\alpha}}(t) < H_Z(t)$ for $t \ge \operatorname{reg}(Z)$. (see Corollary 3.6).

For Question 2, due to a result of Benedetti et al. (see [3, Lemma 4.4]), we know that if $Z = m_1 P_1 + \cdots + m_s P_s$ is a set of fat points in \mathbb{P}^n whose support is contained in a (n-1)-dimensional linear subspace $\alpha \cong \mathbb{P}^{n-1}$ and if there is a non-negative integer t such that $\operatorname{reg}(Z_{\alpha}) \leq t$, then

 $\operatorname{reg}(Z) \leq t.$

By inductive argument on the dimension of linear subspace, we get

Lemma 3.1. Let $Z = m_1 P_1 + \cdots + m_s P_s$ be a set of fat points in \mathbb{P}^n whose support is contained in a linear r-space $\alpha \cong \mathbb{P}^r$. We may consider the linear r-space α as a r-dimensional projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$ and consider $Z_\alpha = m_1 P_{1\alpha} + \cdots + m_s P_{s\alpha}$ as a set of fat points in \mathbb{P}^r . If there is a non-negative integer t such that

$$\operatorname{reg}(Z_{\alpha}) \leq t$$

then

$$\operatorname{reg}(Z) \leq t.$$

Now we prove that the converse is also true. The fisrt, we prove the following lemma.

Lemma 3.2. Let α be a (n-1)-dimensional linear subspace in \mathbb{P}^n , and let P_1, \ldots, P_s be distinct points on α . Let m_1, \ldots, m_s be positive integers. We may consider the (n-1)-dimensional linear subspace α as a (n-1)-dimensional projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and consider $Z_\alpha = m_1 P_{1\alpha} + \cdots + m_s P_{s\alpha}$ as a set of fat points in \mathbb{P}^{n-1} . Put $R_\alpha = K[X_0, \ldots, X_{n-1}]$. Let $\varphi_j \subset R$ be the defining prime ideal of $P_j \in \mathbb{P}^n$, and $\varphi_{j\alpha} \subset R_\alpha$ be the defining prime ideal of $P_{j\alpha} \in \mathbb{P}^r$; $j = 1, \ldots, s$. Put $J = \varphi_1^{m_1} \cap \cdots \cap \varphi_{s-1}^{m_{s-1}}$ and $J_\alpha = \varphi_{1\alpha}^{m_1} \cap \cdots \cap \varphi_{(s-1)\alpha}^{m_{s-1}}$. If there is a non-negative integer b such that

$$\operatorname{reg}(R_{\alpha}/(J_{\alpha}+\wp_{s\alpha}^{m_s}))>b,$$

then

$$\operatorname{reg}(R/(J+\wp_s^{m_s})) > b.$$

Proof. Consider distinct points P_1, \ldots, P_s on α , an (n-1)-dimensional linear subspace in \mathbb{P}^n . We may assume that n points $(1, 0, \ldots, 0, 0, 0), \ldots, (0, 0, \ldots, 0, 1, 0)$ span α . Put $P_s = (1, 0, \ldots, 0)$, then $\wp_s = (X_1, \ldots, X_n), P_{s\alpha} = (1, 0, \ldots, 0) \in \mathbb{P}^{n-1}$ and $\wp_{s\alpha} = (X_1, \ldots, X_{n-1})$. If $\operatorname{reg}(R_{\alpha}/(J_{\alpha} + \wp_{s\alpha}^{m_s})) > b$,

then by Lemma 2.2 there exists a monomial $M = X_1^{c_1} \cdots X_{n-1}^{c_{n-1}} \in \wp_{s\alpha}, c_1 + \cdots + c_r = i, i \in \{0, \ldots, m_s - 1\}$, such that

$$X_0^{b-i}M \notin J_\alpha + \wp_{s\alpha}^{i+1}.$$

Now we consider $M = X_1^{c_1} \cdots X_{n-1}^{c_{n-1}}$ in $K[X_0, \dots, X_n]$. We claim that

 $X_0^{b-i}M \notin J + \wp_s^{i+1}.$

In fact, if $X_0^{b-i}M \in J + \wp_s^{i+1}$, then $X_0^{b-i}M \in [J + \wp_s^{i+1}]_b$. Consider the finite dimensional vector space $[J + \wp_s^{i+1}]_b$ over the field K with the monomial $X_0^{b-i}M$ being an element of it's basis, we get

$$X_0^{b-i}M \in J \text{ or } X_0^{b-i}M \in \wp_s^{i+1}$$

In case of $X_0^{b-i}M \in J$, then $X_0^{b-i}M = X_0^{b-i}X_1^{c_1}\cdots X_{n-1}^{c_{n-1}} \in K[X_0,\ldots,X_{n-1}]$ and $X_0^{b-i}M$ vanishing at the points $P_j \in \alpha$ to order m_j ; $j = 1, \ldots, s$. Thus $X_0^{b-i}M \in J_\alpha$, this is a contradiction with $X_0^{b-i}M \notin J_\alpha + \wp_{s\alpha}^{i+1}$. In case of $X_0^{b-i}M \in \wp_s^{i+1}$, then $X_0^{b-i}M = X_0^{b-i}X_1^{c_1}\cdots X_{n-1}^{c_{n-1}} \in K[X_0,\ldots,X_{n-1}]$ and $X_0^{b-i}M$ vanishing at the points $P_s \in \alpha$ to order i+1. Thus $X_0^{b-i}M \in \wp_{s\alpha}^{i+1}$, this is a contradiction with $X_0^{b-i}M \notin J_\alpha + \wp_{s\alpha}^{i+1}$. Thus, we have the claim

$$X_0^{b-i}M \notin J + \wp_s^{i+1}$$

Then, by Lemma 2.2 we get

$$\operatorname{reg}(R/(J+\wp_{s\alpha}^{m_s})) > b.$$

From the above lemma, by inductive argument on the dimension of linear subspace, we get the following lemma.

Lemma 3.3. Let α be an r-dimensional linear subspace in \mathbb{P}^n , $1 \leq r \leq n-1$, and let P_1, \ldots, P_s be distinct points on α . Let m_1, \ldots, m_s be positive integers. We may consider the r-dimensional linear subspace α as a r-dimensional projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and $Z_{\alpha} = m_1 P_{1\alpha} + \cdots + m_s P_{s\alpha}$ as a zero-scheme in \mathbb{P}^r . Put $R_{\alpha} = K[X_0, \ldots, X_r]$. Let $\varphi_j \subset R$ be the defining prime ideal of P_j , and $\varphi_{j\alpha} \subset R_{\alpha}$ be the defining prime ideal of $P_{j\alpha}$; $j = 1, \ldots, s$. Put $J = \varphi_1^{m_1} \cap \cdots \cap \varphi_{s-1}^{m_{s-1}}$ and $J_{\alpha} = \varphi_{1\alpha}^{m_1} \cap \cdots \cap \varphi_{(s-1)\alpha}^{m_{s-1}}$. If there is a non-negative integer b such that

$$\operatorname{reg}(R_{\alpha}/(J_{\alpha}+\wp_s^{m_s}))>b,$$

then

$$\operatorname{reg}(R/(J+\wp_{s\alpha}^{m_s})) > b$$

By using the above lemmas we prove the following result.

Proposition 3.4. Let $Z = m_1P_1 + \cdots + m_sP_s$ be fat points in \mathbb{P}^n whose support is contained in an r-dimensional linear space $\alpha \cong \mathbb{P}^r$. We may consider the r-dimensional linear space α as a projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and $Z_{\alpha} = m_1P_{1\alpha} + \cdots + m_sP_{s\alpha}$ as a zero-scheme in \mathbb{P}^r . If there is a non-negative integer b such that

$$\operatorname{reg}(Z_{\alpha}) > b,$$

then

 $\operatorname{reg}(Z) > b.$

Proof. Without loss of generality we may assume that $m_1 \geq \ldots \geq m_s$. Put $R_{\alpha} = K[X_0, \ldots, X_r]$ and denote by $\wp_{j\alpha}$ the homogeneous prime ideal in R_{α} defining by $P_{j\alpha}$; $j = 1, \ldots, s$. We argue by induction on s. If s = 1, then $Z = m_1 P_1$ in \mathbb{P}^n and $Z_{\alpha} = m_1 P_{1\alpha}$ in \mathbb{P}^r . It is well known that $\operatorname{reg}(m_1 P_1) = m_1 - 1$ and $\operatorname{reg}(m_1 P_{1\alpha}) = m_1 - 1$. Suppose that the lemma is true for s - 1. Put $J = \wp_1^{m_1} \cap \cdots \cap \wp_{s-1}^{m_{s-1}}$, $I = J \cap \wp_s^{m_s}$, $J_{\alpha} = \wp_{1\alpha}^{m_1} \cap \cdots \cap \wp_{(s-1)\alpha}^{m_{s-1}}$, $I_{\alpha} = J_{\alpha} \cap \wp_{s\alpha}^{m_s}$. By Lemma 2.1 we get

$$\operatorname{reg}(R/I) = \max\left\{m_s - 1, \operatorname{reg}(R/J), \operatorname{reg}(R/(J + \wp_s^{m_s}))\right\}$$

and

$$\operatorname{reg}(R_{\alpha}/I_{\alpha}) = \max\left\{m_s - 1, \operatorname{reg}(R_{\alpha}/J_{\alpha}), \operatorname{reg}(R_{\alpha}/(J_{\alpha} + \wp_{s\alpha}^{m_s}))\right\}$$

By Lemma 2.3 we have $\operatorname{reg}(R/J) \ge m_1 + m_2 - 1$ and $\operatorname{reg}(R_\alpha/J_\alpha) \ge m_1 + m_2 - 1$. So, $\operatorname{reg}(R/J) \ge m_s - 1$ and $\operatorname{reg}(R_\alpha/J_\alpha) \ge m_s - 1$. Therefore,

$$\operatorname{reg}(R/I) = \max\left\{\operatorname{reg}(R/J), \operatorname{reg}(R/(J + \wp_s^{m_s}))\right\}$$

and

$$\operatorname{reg}(R_{\alpha}/I_{\alpha}) = \max\left\{\operatorname{reg}(R_{\alpha}/J_{\alpha}), \operatorname{reg}(R_{\alpha}/(J_{\alpha} + \wp_{s\alpha}^{m_s}))\right\}$$

If $\operatorname{reg}(R_{\alpha}/I_{\alpha}) > b$ in \mathbb{P}^r , then we consider two following cases:

Case $\operatorname{reg}(R_{\alpha}/J_{\alpha}) > b$: Then by inductive assumption we have $\operatorname{reg}(R/J) > b$. Therefore,

$$\operatorname{reg}(R/I) > b$$

Case $\operatorname{reg}(R_{\alpha}/(J_{\alpha}+\wp_{s\alpha}^{m_s})) > b$: Then by using Lemma 3.3 we get $\operatorname{reg}(R/(J+\wp_{s\alpha}^{m_s})) > b$. Therefore, $\operatorname{reg}(R/I) > b$.

From Lemma 3.1 and Proposition 3.4 we get the following theorem which answers Question 2.

Theorem 3.5. Let $Z = m_1P_1 + \cdots + m_sP_s$ be fat points in \mathbb{P}^n whose support is contained in a linear r-space $\alpha \cong \mathbb{P}^r$. We may consider the linear r-space α as a r-dimensional projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and consider $Z_{\alpha} = m_1P_{1\alpha} + \cdots + m_sP_{s\alpha}$ as a zero-scheme in \mathbb{P}^r . Denote $\operatorname{reg}(Z)$ the regularity index of Z in \mathbb{P}^n , and denote $\operatorname{reg}(Z_{\alpha})$ the regularity of Z_{α} in \mathbb{P}^n . Then

$$\operatorname{reg}(Z_{\alpha}) = \operatorname{reg}(Z).$$

Hence, the regularity index of a set of fat points is invariant under changes of linear subspace containing the support of fat points.

Corollary 3.6. Let $Z = m_1P_1 + \cdots + m_sP_s$ be fat points in \mathbb{P}^n whose support is contained in a linear r-space $\alpha \cong \mathbb{P}^r$. We may consider the linear r-space α as a r-dimensional projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and consider $Z_\alpha = m_1P_{1\alpha} + \cdots + m_sP_{s\alpha}$ as a zero-scheme in \mathbb{P}^r . If $t \ge \operatorname{reg}(Z)$, then

$$H_{Z_{\alpha}}(t) \leq H_Z(t).$$

Furthermore, if there exists $m_i \geq 2$ and r < n, then

$$H_{Z_{\alpha}}(t) < H_Z(t)$$

for $t \geq \operatorname{reg}(Z)$.

Proof. We know that the Hilbert function $H_Z(t)$ strictly increases until it reaches the multiplicity e(Z), at which it stabilizes, and the regularity index $\operatorname{reg}(Z)$ is the least integer t such that $H_{R/I}(t) = e(Z)$. So, if $t \ge \operatorname{reg}(Z)$ then $H_Z(t) = e(Z) = \sum_{i=1}^{s} {\binom{m_i+n-1}{n}}.$

For fat points Z_{α} in \mathbb{P}^r , we also get: if $t \ge \operatorname{reg}(Z_{\alpha})$, then $H_{Z_{\alpha}}(t) = \sum_{i=1}^{s} \binom{m_i+r-1}{r}$. By Theorem 3.5, $\operatorname{reg}(Z) = \operatorname{reg}(Z_{\alpha})$. Therefore, if $t \ge \operatorname{reg}(Z) = \operatorname{reg}(Z_{\alpha})$ then

$$H_{Z_{\alpha}}(t) = \sum_{i=1}^{s} \binom{m_{i}+r-1}{r} \leq \sum_{i=1}^{s} \binom{m_{i}+n-1}{n} = H_{Z}(t).$$

This result implies that if $m_1 = \cdots = m_s = 1$ then $H_{Z_\alpha}(t) = H_Z(t)$ for $t \ge \operatorname{reg}(Z)$, and if there exists $m_i \ge 2$ and r < n, then $H_{Z_\alpha}(t) < H_Z(t)$ for $t \ge \operatorname{reg}(Z)$.

Corollary 3.7. Let $Z = m_1 P_1 + \cdots + m_s P_s$ be fat points in \mathbb{P}^n with the support in a r-dimensional linear subspace α . If $t \geq \operatorname{reg}(Z)$, then we have the relation between dimension of the t-th graded part I_t of ideal I consisting all homogeneous polynomials in $K[X_0, \ldots, X_n]$ that vanish at Pi to order m_i ; $i = 1, \ldots, s$; and dimension of the t-th graded part $(I_\alpha)_t$ of ideal I_α consisting all homogeneous polynomials in $K[X_0, \ldots, X_n]$ that vanish at Pi to order polynomials in $K[X_0, \ldots, X_r]$ that vanish at Pi to order m_i ; $i = 1, \ldots, s$; as following:

$$(t+r+1)\cdots(t+n)\left[\sum_{i=1}^{s} \binom{m_i+r-1}{r} + \dim_K(I_\alpha)_t\right]$$
$$= (r+1)\cdots(n-1)n\left[\sum_{i=1}^{s} \binom{m_i+n-1}{n} + \dim_K I_t\right].$$

Proof. We may consider α as a *r*-dimensional projective space \mathbb{P}^r containing the points $P_{1\alpha} := P_1, \ldots, P_{s\alpha} := P_s$, and consider $Z_{\alpha} = m_1 P_{1\alpha} + \cdots + m_s P_{s\alpha}$ as a set of fat points in \mathbb{P}^r . Since $I = \bigoplus_{t \ge 0} I_t$ is the ideal consists all homogeneous polynomials in $K[X_0, \ldots, X_n]$ that vanish at Pi to order m_i , $i = 1, \ldots, s$; the Hilbert function of Z is

$$H_Z(t) = \dim_K (K[X_0, \dots, X_n]/I)_t = \dim_K K[X_0, \dots, X_n]_t - \dim_K I_t = \binom{t+n}{n} - \dim_K I_t.$$

Since $I_{\alpha} = \bigoplus_{t \ge 0} (I_{\alpha})_t$ is the ideal consists all homogeneous polynomials in $K[X_0, \ldots, X_r]$ that vanish at $P_{i\alpha}$ to order m_i , $i = 1, \ldots, s$; the Hilbert function of Z_{α} is

$$H_{Z_{\alpha}}(t) = \dim_{K}(K[X_{0},\ldots,X_{r}]/I_{\alpha})_{t} = \dim_{K}K[X_{0},\ldots,X_{r}]_{t} - \dim_{K}(I_{\alpha})_{t} = \binom{t+r}{r} - \dim_{K}(I_{\alpha})_{t}.$$

By Theorem 3.5 we have $\operatorname{reg}(Z) = \operatorname{reg}(Z_{\alpha})$. If $t \ge \operatorname{reg}(Z) = \operatorname{reg}(Z_{\alpha})$, then $H_Z(t)$ reaches the multiplicity $e(Z) = \sum_{i=1}^{s} {m_i+n-1 \choose i}$, and $H_{Z_{\alpha}}(t)$ reaches the multiplicity $e(Z_{\alpha}) = \sum_{i=1}^{s} {m_i+r-1 \choose r}$, and so we have

$$\begin{cases} \binom{t+n}{n} = \sum_{\substack{i=1\\r}}^{s} \binom{m_i+n-1}{n} + \dim_K I_t, \\ \binom{t+r}{r} = \sum_{i=1}^{s} \binom{m_i+r-1}{r} + \dim_K (I_\alpha)_t. \end{cases}$$

Hence if $t \geq \operatorname{reg}(Z)$, then

$$(t+r+1)\cdots(t+n)\left[\sum_{i=1}^{s} \binom{m_i+r-1}{r} + \dim_K(I_{\alpha})_t\right] = (t+r+1)\cdots(t+n)\binom{t+r}{r} = (r+1)\cdots(n-1)n\binom{t+n}{n} = (r+1)\cdots(n-1)n\left[\sum_{i=1}^{s} \binom{m_i+n-1}{n} + \dim_K I_t\right].$$

Let $Z = m_1 P_1 + \cdots + m_s P_s$ be a set of fat points in \mathbb{P}^n , and let α be a r-dimensional linear subspace. We put

$$\alpha \cap Z := \sum_{P_i \in \alpha} m_i P_i$$

and

$$w_{\alpha \cap Z} := \sum_{P_i \in \alpha} m_i.$$

Corollary 3.8. Let Z be a set of fat points in \mathbb{P}^n . Assume that α is a r-dimensional linear subspace such that $\left[\frac{w_{\alpha\cap Z}-2}{r}\right] = T(Z)$. Consider $(\alpha \cap Z)_{\alpha}$ as a set of fat points in \mathbb{P}^r . If the support of $(\alpha \cap Z)_{\alpha}$ lies on a line or on a rational normal curve of \mathbb{P}^r , then $\operatorname{reg}(Z) = T(Z)$.

Proof. By Theorem 3.5 we have

$$\operatorname{reg}(\alpha \cap Z) = \operatorname{reg}((\alpha \cap Z)_{\alpha}).$$

By using Lemma 2.6 and [7, Theorem 5.3] we get

$$\operatorname{reg}(\alpha \cap Z) \le \operatorname{reg}(Z) \le T(Z)$$

Since the support of $(\alpha \cap Z)_{\alpha}$ lies on a line or on a rational normal curve of \mathbb{P}^r , by using [5, Corollary 2.3] and [4, Proposition 7] we get

$$\operatorname{reg}((\alpha \cap Z)_{\alpha}) \ge \left[\frac{w_{\alpha \cap Z} - 2}{r}\right].$$

Note that $\left[\frac{w_{\alpha\cap Z}-2}{r}\right] = T(Z)$ by the hypothesis. Therefore, reg $(Z) = \operatorname{reg}((\alpha \cap Z)_{\alpha}) = T(Z).$

Let $Z = m_1 P_1 + \cdots + m_s P_s$ be a set of fat points in \mathbb{P}^n . If α is a linear subspace of \mathbb{P}^n such that P_1, \ldots, P_s span α , then we say that Z is non-degenerate in α . From Theorem 3.5, we have the following remark.

Remark 3.9. To calculate the regularity index of a set of fat points Z in \mathbb{P}^n , we find a linear subspace $\alpha \cong \mathbb{P}^r$ such that Z is non-degenerate on α . Then we calculate the regularity index of the non-degenerate fat points Z_{α} in \mathbb{P}^r .

4 Segre's bound and the regularity index of non-degenerate equimultiple fat points

Recall that for a set of fat points $Z = m_1 P_1 + \cdots + m_s P_s$ in \mathbb{P}^n , the Segre's bound for the regularity index of Z is the number

$$T(Z) = \max\{T_j(Z) | j = 1, \dots, n\},\$$

where

$$T_j(Z) := \max\left\{ \left[\frac{\sum_{l=1}^q m_{i_l} + j - 2}{j} \right] \mid P_{i_1}, \dots, P_{i_q} \text{ on a } j \text{-dimensional linear subspace} \right\}.$$

Lemma 4.1. Let $Z = 2P_1 + \cdots + 2P_7$ be a set of seven non-degenerate double points in \mathbb{P}^4 . Then

$$\operatorname{reg}(Z) \ge 4.$$

Proof. Let $U = 2Q_1 + \cdots + 2Q_7$ be a set of seven generic double points in \mathbb{P}^4 . Denote by $H_Z(t)$ the Hilbert function of Z and by $H_U(t)$ the Hilbert function of U. By the maximality of the Hilbert function of generic fat points we have

$$H_Z(t) \le H_U(t).$$

Note that Z and U have the same multiplicity

$$e(Z) = e(U) = 7\binom{2+4-1}{4} = 35$$

By the Alexander-Hirschowits Theorem [1] about the Hilbert function of generic double points in \mathbb{P}^4 we have

$$H_U(3) = 34$$

Since $H_Z(3) \leq H_U(3) = 34 < 35 = e(Z)$, by the definition of reg(Z) we get

$$\operatorname{reg}(Z) \ge 4.$$

ON INVARIANT OF THE REGULARITY INDEX OF FAT POINTS

The above lemma and Theorem 3.5 help us to calculate the regularity index of a set of n + 3 non-degenerate double points in \mathbb{P}^n . In this case we see that the regularity index reaches the Segre's bound.

Proposition 4.2. Let $Z = 2P_1 + \cdots + 2P_{n+3}$ be a set of n+3 non-degenerate double points in \mathbb{P}^n . Then

$$\operatorname{reg}(Z) = T(Z)$$

where

$$T_{j}(Z) := \max\left\{ \left[\frac{2q+j-2}{j}\right] \mid P_{i_{1}}, \dots, P_{i_{q}} \text{ on a } j\text{-dimensional linear subspace} \right\}$$

 $T(Z) = \max\{T_i(Z) | j = 1, \dots, n\},\$

Proof. Consider the fat points in the assumption of Lemma 2.5, we see that

$$T(Z,L) = \max\left\{ \left[\frac{\sum_{l=1}^{q} m_{i_l} + j - 2}{j} \right] \mid P_{i_1}, \dots, P_{i_q} \in L \text{ is a } j \text{-dimensional linear subspace} \right\}$$
$$= T_j(Z).$$

Hence

$$\max\{T(Z,L): L \subseteq \mathbb{P}^n\} = T(Z).$$

By using Lemma 2.5 we get $reg(Z) \leq T(Z)$. So, it is sufficient to prove

$$\operatorname{reg}(Z) \ge T(Z).$$

We argue by induction on n. For n = 2, then n + 3 = 5. By Lemma 2.9 we get $\operatorname{reg}(Z) = T(Z)$. For $n \geq 3$, we assume that the proposition is true for any m + 3 non-degenerate double points $2P_1 + \cdots + 2P_{m+3}$ in \mathbb{P}^m , m < s. We now consider n + 3 non-degenerate double points $Z = 2P_1 + \cdots + 2P_{n+3}$ in \mathbb{P}^n .

Since P_1, \ldots, P_{n+3} are not not on any (n-1)-dimensional linear subspace, there are at most j+3 points of $\{P_1, \ldots, P_{n+3}\}$ on a *j*-dimensional linear subspace of \mathbb{P}^n ; $j = 1, \ldots, n-1$. Put

$$p = \min\{j \mid T_j(Z) = T(Z)\}.$$

and denote by α the *p*-dimensional linear subspace containing points P_{i_1}, \ldots, P_{i_r} of $\{P_1, \ldots, P_{n+3}\}$ such that

$$\left[\frac{2r+p-2}{p}\right] = T_p(Z) = T(Z).$$

Then the points P_{i_1}, \ldots, P_{i_r} are not on any (p-1)-dimensional linear subspace. So, $r \ge p+1$. Since P_1, \ldots, P_{n+3} are non-degenerate in \mathbb{P}^n , there are at most p+3 points of $\{P_1, \ldots, P_{n+3}\}$ on a p-dimensional linear subspace. Hence $r \le p+3$.

Consider the set of r double points

$$Y = 2P_{i_1} + \dots + 2P_{i_r}.$$

in \mathbb{P}^n . Put $P_{i,\alpha} = P_j$, $j = 1, \ldots, r$. Consider the set of r non-degenerate double points

$$Y_{\alpha} = 2P_{i_1\alpha} + \dots + 2P_{i_r\alpha}$$

in \mathbb{P}^p . If r = p + 1 or r = p + 2, then by Lemma 2.7 we get

$$\operatorname{reg}(Y_{\alpha}) = T(Y_{\alpha}) = \left[\frac{2r+p-2}{p}\right].$$

So, $\operatorname{reg}(Y_{\alpha}) = T(Z)$. By Theorem 3.5 we get $\operatorname{reg}(Y) = \operatorname{reg}(Y_{\alpha})$. By use Lemma 2.6, we obtain

$$\operatorname{reg}(Z) \ge \operatorname{reg}(Y) = T(Z).$$

If r = p + 3, then we consider two following cases for p:

Case p < n: Since p + 3 points $P_{i_1}, \ldots, P_{i_{p+3}}$ lie on the *p*-dimensional linear subspace α and do not lie on a (p-1)-dimensional linear subspace, by the inductive assumption we have

$$\operatorname{reg}(Y_{\alpha}) = T(Y_{\alpha})$$

By using Theorem 3.5 we get $\operatorname{reg}(Y) = \operatorname{reg}(Y_{\alpha})$. Now use Lemma 2.6 and obtain $\operatorname{reg}(Z) \ge \operatorname{reg}(Y)$. Since $\operatorname{reg}(Y) = \operatorname{reg}(Y_{\alpha}) = T(Y_{\alpha}) = \left[\frac{2r+p-2}{p}\right] = T(Z)$, we get

$$\operatorname{reg}(Z) \ge T_p(Z) = T(Z).$$

Case p = n: Then $T_n(Z) > T_j(Z)$; j = 1, ..., n-1. By defining we have

$$T_n(Z) = \left[\frac{(n+3)2+n-2}{n}\right] = 3 + \left[\frac{4}{n}\right], \text{ and } T_1(Z) \ge 2.2 - 1 = 3$$

Since $T_n(Z) > T_1(Z)$, we have $n \le 4$. But we are considering $n \ge 3$, so n = 3 or n = 4.

For n = 3, then $T_3(Z) = 4 = T(Z)$. Consider the set of 6 non-degenerate double points

$$Z = 2P_1 + \dots + 2P_6$$

in \mathbb{P}^3 . If P_1, \ldots, P_6 are not in general position in \mathbb{P}^3 , then there are 3 points of $\{P_1, \ldots, P_6\}$ on a line or there are 4 points of $\{P_1, \ldots, P_6\}$ on a 2-dimensional linear subspace. So, $T_1(Z) \ge 5$ or $T_2(Z) \ge 4$, this is a contradiction with $T(Z) = T_p(Z) = T_3(Z) = 4 > T_j(Z); j = 1, 2$. So, P_1, \ldots, P_6 are in general position in \mathbb{P}^3 . By using the Lemma 2.8 we get

$$\operatorname{reg}(Z) = T(Z).$$

For n = 4, then $T_4(Z) = 4 = T(Z)$. Consider the set of seven non-degenerate double points

$$Z = 2P_1 + \dots + 2P_7$$

in \mathbb{P}^4 . By Lemma 4.1 we get

$$\operatorname{reg}(Z) \ge 4 = T(Z).$$

The proof of Proposition 4.2 is completed.

The Proposition 4.2 help us to prove the following theorem which shows that the Segre's bound is attained for the regularity index of any s+3 non-degenerate equimultiple fat points in \mathbb{P}^n , $s \leq n+3$.

Theorem 4.3. Let m be a positive integer and $Z = mP_1 + \cdots + mP_s$ be a set of s non-degenerate equimultiple points in \mathbb{P}^n . If $s \leq n+3$, then

$$\operatorname{reg}(Z) = T(Z),$$

where

$$T(Z) = \max\{T_j(Z) \mid j = 1, \dots, n\},\$$
$$T_j(Z) := \max\left\{ \left[\frac{mq + j - 2}{j}\right] \mid P_{i_1}, \dots, P_{i_q} \text{ on a } j \text{-dimensional linear subspace} \right\}$$

Proof. Since P_1, \ldots, P_s are not on any (n-1)-dimensional linear subspace, we have $s \ge n+1$. We consider the three following cases for s.

Case s = n + 1: Then P_1, \ldots, P_{n+1} are in general position in \mathbb{P}^n . By Lemma 2.4 we get

$$\operatorname{reg}(Z) = 2m - 1 = T_1(Z) = T(Z).$$

Case s = n + 2: Then P_1, \ldots, P_{n+2} are not on a (n-1)-dimensional linear subspace in \mathbb{P}^n . By Lemma 2.7 we get

$$\operatorname{reg}(Z) = T(Z).$$

Case s = n + 3: If $m \neq 2$, then by using Lemma 2.10 we get

$$\operatorname{reg}(mP_1 + \dots + mP_{n+3}) = T(Z).$$

If m = 2, then by Proposition 4.2 we get

$$\operatorname{reg}(2P_1 + \dots + 2P_{n+3}) = T(Z).$$

The following simple example shows that Theorem 3.5 is not true for n = 2, s = n + 4.

From now on, if H is a hyperplane in \mathbb{P}^n , we coincide H with the linear form in R defining it. So, if H_j is a hyperplane passing through points P_j and m_j is a positive integer; $j = 1, \ldots, r$; then we get $H_1^{m_1} \cdots H_r^{m_r} \in \wp_1^{m_1} \cap \cdots \cap \wp_r^{m_r}$.

Example 4.4. Let l_1, l_2 be two distinct lines in \mathbb{P}^2 and P_1, \ldots, P_6 be six distinct points such that $P_1, P_2, P_3 \in l_1 \setminus l_2$; $P_4, P_5 \in l_2 \setminus l_1$ and $P_6 \notin l_1 \cup l_2$.



Let m be a positive integer, then $Z = mP_1 + \cdots + mP_6$ is a set of 6 non-degenerate equimultiple fat points in \mathbb{P}^2 . We have

$$T_1(Z) = 3m - 1, \ T_2(Z) = \left[\frac{6m}{2}\right] = 3m = T(Z).$$

Let \wp_j be the homogeneous prime ideal in $\mathbb{R} = K[X_0, X_1, X_2]$ corresponding to the point P_j ; $j = 1, \ldots, 6$. Choose $P_6 = (1, 0, 0)$, then $\wp_6 = (X_1, X_2)$. Put $J = \wp_1^m \cap \cdots \cap \wp_5^m$, $I = J \cap \wp_6^m$. Then R/I is the coordinate ring of Z, so reg $(Z) = \operatorname{reg}(R/I)$, the Castelnuovo-Mumford regularity index of the ring R/I. By Lemma 2.1 we get

$$\operatorname{reg}(R/I) = \max\{m - 1, \operatorname{reg}(R/J), \operatorname{reg}(R/(J + \wp_6^m))\}.$$

Since l_1 is a line passing through three points P_1, P_2, P_3 ; we have $l_1^m \in \wp_1^m \cap \wp_2^m \cap \wp_3^m$. Since l_2 is a line passing through two points P_4, P_5 ; we have $l_2^m \in \wp_4^m \cap \wp_5^m$. Hence, $l_1^m l_2^m \in J$. This implies that for every monomial M of degree i in X_1, X_2 ; $i = 0, \ldots, m-1$; we have

$$l_1^m l_2^m M \in J$$

Since l_j is a line avoiding $P_6 = (1, 0, 0)$, we can write $l_j = X_0 + g_j$ for some $g_j \in \wp_6 = (X_1, X_2)$; j = 1, 2. Thus we have

$$(X_0 + g_1)^m (X_0 + g_2)^m M \in J.$$

Moreover, since $g_1, g_2 \in \wp_6$ and $M \in \wp_6^i$, we get $X_0^{2m}M \in J + \wp_6^{i+1}$ for $i = 0, \ldots, m-1$. This implies that

$$X_0^{3m-1-i}M \in J + \wp_6^{i+1}$$

for $i = 0, \ldots, m - 1$. By Lemma 2.2 we get

$$\operatorname{reg}(R/(J+\wp_6^2)) \le 3m-1.$$

Let $U = mP_1 + \cdots + mP_5$, we have

$$T_1(U) = 3m - 1 \ge T_2(U) = \left[\frac{5m}{2}\right] = 2m + \left[\frac{m}{2}\right].$$

So $T(U) = T_1(U) = 3m - 1$. By Lemma 2.9 we get reg(U) = 3m - 1. Since R/J is the homogeneous coordinate ring of U. Thus

$$\operatorname{reg}(R/J) = \operatorname{reg}(U) = 3m - 1.$$

Therefore,

$$\operatorname{reg}(Z) = \operatorname{reg}(R/I) = \max\{m - 1, 3m - 1, \operatorname{reg}(R/(J + \varphi_6^m))\} = 3m - 1 < T(Z).$$

By using Theorem 3.5 and Example 4.4 we show that Theorem 4.3 is not true for s = n + 4. The Segre's bound is not attained for this case.

Corollary 4.5. Let m, n be positive integers, $n \ge 2$. There always exists a set of n+4 non-degenerate equimultiple fat points $Z = mP_1 + \cdots + mP_{n+4}$ in \mathbb{P}^n such that

$$\operatorname{reg}(Z) = 3m - 1 < T(Z) = 3m,$$

where

$$T(Z) = \max\{T_j(Z) | j = 1, \dots, n\},\$$
$$T_j(Z) := \max\left\{\left[\frac{mq+j-2}{j}\right] | P_{i_1}, \dots, P_{i_q} \text{ on a } j \text{-dimensional linear subspace}\right\}.$$

Proof. We consider the set of fat points Z with the configuration as following.

Let α_1 be the 2-dimensional linear subspace in \mathbb{P}^n defining by n-2 linear forms X_3, \ldots, X_n ; and l_1, l_2 be two distinct lines in $\alpha_1 \cong \mathbb{P}^2$. Let P_1, \ldots, P_3 be three distinct points on $l_1 \setminus l_2$; and P_4, P_5 be two distinct points on $l_2 \setminus l_1$. Let P_6 be a points in α_1 and not on $l_1 \cup l_2$. Consider the set of 6 double points $Z_{\alpha_1} = mP_1 + \cdots + mP_6$ in \mathbb{P}^2 , by Example 4.4 we get

$$\operatorname{reg}(Z_{\alpha_1}) = 3m - 1$$
 and $T(Z_{\alpha_1}) = T_2(Z_{\alpha_1}) = 3m$.

Since P_4, P_5, P_6 do not lie on a line, we can choose P_7, \ldots, P_{n+4} be distinct points in \mathbb{P}^n such that n+1 points $P_4, P_5, P_6, P_7, \ldots, P_{n+4}$ span \mathbb{P}^n . Then P_4, \ldots, P_{n+4} are in linearly general position in \mathbb{P}^n and

$$Z = mP_1 + \dots + mP_{n+4}$$

is the set of n + 4 non-degenerate equimultiple fat points in \mathbb{P}^n . We have

$$T_1(Z) = 3m - 1, T_2(Z) = 3m,$$

$$T_j(Z) = \left[\frac{6m + (j-2)m + j - 1}{j}\right] = m + \left[\frac{4m + j - 2}{j}\right]; j = 3, \dots, n - 1.$$

Thus,

$$T(Z) = T_2(Z) = 3m$$

For j = 2, ..., n-1; the points $P_4, P_5, P_6, ..., P_{j+5}$ span a (j+1)-dimensional linear subspace, say α_j . Put $P_{i\alpha_j} = P_i$; i = 1, ..., j+5; then

$$Z_{\alpha_j} = mP_{1\alpha_j} + \dots + mP_{(j+5)\alpha_j}$$

is the set of j + 5 non-degenerate equimultiple fat points in \mathbb{P}^j . Since $\alpha_{n-1} = \mathbb{P}^n$, we get

$$\operatorname{reg}(Z) = \operatorname{reg}(Z_{\alpha_{n-1}}).$$

We argue by induction on j to prove that

$$\operatorname{reg}(Z_{\alpha_i}) = \operatorname{reg}(Z_{\alpha_1}) = 3m - 1$$

for j = 2, ..., n - 1.

For j = 2, consider the set of 7 non-degenerate equimultiple fat points

$$Z_{\alpha_2} = m_1 P_{1\alpha_2} + \dots + m P_{7\alpha_2}$$

in \mathbb{P}^3 . Put $J_{\alpha_2} = \wp_{1\alpha_2}^m \cap \cdots \cap \wp_{6\alpha_2}^m$ and $I_{\alpha_2} = J_{\alpha_2} \cap \wp_{7\alpha_2}^m$, $R_{\alpha_2} = K[X_0, X_1, X_2, X_3]$. Then $\operatorname{reg}(Z_{\alpha_2}) = \operatorname{reg}(R_{\alpha_2}/I_{\alpha_2})$. By Lemma 2.1 we get

$$\operatorname{reg}(R_{\alpha_2}/I_{\alpha_2}) = \max\{m - 1, \operatorname{reg}(R_{\alpha_2}/J_{\alpha_2}), \operatorname{reg}(R_{\alpha_2}/(J_{\alpha_2} + \wp_{7\alpha_2}^m))\}$$

Since $P_{1\alpha_2} = P_1, \ldots, P_{6\alpha_2} = P_6$ are on $\alpha_1 \cong \mathbb{P}^2$; by using Example 4.4 we get

$$\operatorname{reg}(R_{\alpha_2}/J_{\alpha_2}) = \operatorname{reg}(Z_{\alpha_1}) = 3m - 1.$$

Let $P_{7\alpha_2} = (1, 0, 0, 0)$ in \mathbb{P}^3 . Since $P_{7\alpha_2} \notin \alpha_1$ and $P_{1\alpha_2}, \ldots, P_{6\alpha_2} \in \alpha_1$, we can choose a 2-dimensional linear subspace H in \mathbb{P}^3 passing through $P_{1\alpha_2}, \ldots, P_{6\alpha_2}$ and avoiding $P_{7\alpha_2}$. Then we get

$$H^m \in \wp_{1\alpha_2}^m \cap \dots \cap \wp_{6\alpha_2}^m = J_{\alpha_2}$$

It follows that

$$H^{2m-i}M \in J_{\alpha}$$

for every monomial M of degree i in X_1, X_2, x_3 ; i = 0, ..., m-1. Since $P_{7\alpha_2} \notin H$, we can write $H = X_0 + G$ for some linear form $G \in \wp_{7\alpha_2} = (X_1, X_2, X_3)$. Thus we have

$$(X_0 + G)^{2m-i}M \in J_{\alpha_2}$$

for every monomial M of degree i in X_1, X_2, x_3 ; i = 0, ..., m - 1. Moreover, since $G \in \wp_{7\alpha_2}$ and $M \in \wp_{7\alpha_2}^i$, we get

$$X_0^{2m-i}M \in J_{\alpha_2} + \wp_{7\alpha_2}^{i+1}$$

for every monomial M of degree i in X_1, X_2, x_3 ; $i = 0, \ldots, m-1$. By Lemma 2.2 we get

$$\operatorname{reg}(R_{\alpha_2}/(J_{\alpha_2}+\wp_{7\alpha_2}^m)) \le 2m$$

Therefore,

$$\operatorname{reg}(Z_{\alpha_2}) = 3m - 1.$$

Assume that $\operatorname{reg}(Z_{\alpha_{n-2}}) = 3m - 1$. We need prove that $\operatorname{reg}(Z_{\alpha_{n-1}}) = 3m - 1$. Consider the set of n + 4 non-degenerate equimultiple fat points

$$Z = mP_1 + \dots + mP_{n+4}$$

in $\alpha_{n-1} = \mathbb{P}^n$. Put $J = \wp_1^m \cap \cdots \cap \wp_{n_3}^m$ and $I = J \cap \wp_{n+4}^m$. Then $\operatorname{reg}(Z) = \operatorname{reg}(R/I)$. By Lemma 2.1 we get

$$\operatorname{reg}(R/I) = \max\{m - 1, \operatorname{reg}(R/J), \operatorname{reg}(R/(J + \wp_{n+4}^m))\}$$

By Theorem 3.5 and inductive assumption we get $\operatorname{reg}(R/J) = 3m - 1$. Choose $P_{n+4} = (1, 0, \dots, 0)$. Since α_{n-2} is a (n-1)-dimensional linear subspace passing through P_1, \dots, P_{n+3} , we have

$$\alpha_{n-2}^m N \in J$$

for every monomial N of degree i in X_1, \ldots, X_n ; $i = 0, \ldots, m-1$. By a similar argue as above, since $P_{n+4} \notin \alpha_2$, we can write $\alpha_{n-2} = X_0 + L$ for some linear form $L \in \wp_{n+4}$. Thus, we have

$$(X_0 + L)^{2m-i} N \in J.$$

Moreover, since $L \in \wp_{n+4}$ and $N \in \wp_{n+4}^i$, we get

$$X_0^{2m-i} N \in J + \wp_{n+4}^{i+1}$$

for every monomial N of degree i in X_1, \ldots, X_n ; $i = 0, \ldots, m-1$. By Lemma 2.2 we get

$$\operatorname{reg}(R/(J+\wp_{n+4}^m)) \le 2m-1$$

Therefore,

$$\operatorname{reg}(Z) = 3m - 1$$

References

- J. Alexander et A. Hirschowitz, La méthode d'Horace éclatée: application à l'interpolation en degré quatre, Invent. Math. 107 (1992), 585-602.
- [2] E.Ballico, O. Dumitrescu and E. Postinghel, On Segre's bound for fat points in Pⁿ, J. Pure Appl. Algebra 220 (2016), 2307-2323.
- [3] B. Benedetti, G. Fatabbi and A. Lorenzini, Segre's Bound and the case of n + 2 fat points of P^n , Comm. Algebra **40** (2012), 395-403.
- [4] M.V. Catalisano, N.V. Trung and G. Valla, A sharp bound for the regularity index of fat points in general position, Proc. Amer. Math. Soc. 118 (1993), 717-724.
- [5] E.D. Davis and A.V. Geramita, The Hilbert function of a special class of 1-dimensional Cohen-Macaulay graded algebras, The Curves Seminar at Queen's, Queen's Papers in Pure and Appl. Math. 67 (1984), 1-29.
- [6] G. Fatabbi, A. Lorenzini On a sharp bound for the regularity index of any set of fat points, J. Pure and Appl. Algebra 161 (2001), 91-111.
- [7] U. Nagel and B. Trok, Segre's regularity bound for fat point schemes, Annalli della Scuola Normale Superiore, Vol. XX (2020), 217-237.
- [8] B. Segre, Alcune questioni su insiemi finiti di punti in geometria algebrica, Atti. Convergno. Intern. di Torino 1961, 15-33.
- [9] P.V. Thien, Segre bound for the regularity index of fat points in P³, J. Pure and Appl. Algebra 151 (2000), 197-214.
- [10] P.V. Thien, Regularity index of s + 2 fat points not on a linear (s 1)-space, Comm. Algebra 40 (2012), 3704-3715.
- [11] P.V. Thien and T.N. Sinh, On the regularity index of s fat points not on a linear (r-1)-space, $s \le r+3$, Comm. Algebra, **45** (2017), 4123-4138.