Optimal Reinsurance to Minimize the Probability of Drawdown Under the Mean-Variance Premium Principle:

Asymptotic Analysis

Pablo Azcue*

Xiaoqing Liang[†] Nora Muler[‡]

Virginia R. Young[§]

January 4, 2022

Abstract

In this paper, we consider an optimal reinsurance problem to minimize the probability of drawdown for the scaled Cramér-Lundberg risk model when the reinsurance premium is computed according to the mean-variance premium principle. We extend the work of Liang et al. [16] to the case of minimizing the probability of drawdown. By using the comparison method and the tool of adjustment coefficients, we show that the minimum probability of drawdown for the scaled classical risk model converges to the minimum probability for its diffusion approximation, and the rate of convergence is of order $O(n^{-1/2})$. We further show that using the optimal strategy from the diffusion approximation in the scaled classical risk model is $O(n^{-1/2})$ -optimal.

Keywords: Optimal reinsurance; probability of drawdown; scaled Cramér-Lundberg model; asymptotic analysis; diffusion approximation.

^{*}Departamento de Matematicas, Universidad Torcuato Di Tella. Av. Figueroa Alcorta 7350 (C1428BIJ) Ciudad de Buenos Aires, Argentina, pazcue@utdt.edu.

[†]Corresponding author. Department of Statistics, School of Sciences, Hebei University of Technology, Tianjin 300401, P. R. China, liangxiaoqing115@hotmail.com. X. Liang thanks the National Natural Science Foundation of China (11701139) and the Natural Science Foundation of Hebei Province (A2018202057, A2020202033) and the Research Foundation for Returned Scholars of Hebei Province (C20200102) for financial support.

[‡]Departamento de Matematicas, Universidad Torcuato Di Tella. Av. Figueroa Alcorta 7350 (C1428BIJ) Ciudad de Buenos Aires, Argentina, nmuler@utdt.edu.

[§]Department of Mathematics, University of Michigan, Ann Arbor, Michigan, 48109, vryoung@umich.edu. V. R. Young thanks the Cecil J. and Ethel M. Nesbitt Professorship of Actuarial Mathematics for financial support.

AMS 2020 Subject Classification. 91G05, 93E20, 35B51, 47G20.

JEL Codes. C61, G22, D81.

1 Introduction

Drawdown occurs when the value of a portfolio or company's surplus is lower than a proportion of its historic maximum, which is an important risk metric for fund or corporate managers when evaluating their portfolios or companies, respectively. The frequent occurrence of drawdown implies low financial profit, potential large losses, or even bankruptcy, which may help managers take effective action for the company's operation.

Due to the importance of analyzing the occurrence of drawdown, recently researchers have been considering the problem of minimizing the probability of drawdown. Specifically, a decision maker chooses an optimal strategy to minimize the probability that the fund or surplus decreases to a fixed proportion, say $\alpha \in [0, 1)$, of its historic maximum value. If we set $\alpha = 0$, then minimizing the probability of drawdown degenerates to minimizing the probability of ruin. Angoshtari et al. [1] minimize the probability of drawdown with investment in a Black-Scholes financial market. They find that the strategy for minimizing the probability of drawdown is identical to the one for minimizing the probability of ruin. Han et al. [14] consider the reinsurance problem of minimizing the probability of drawdown under the mean-variance premium principle and under the diffusion approximation of the classical Cramér-Lundberg risk model. They find an explicit form of the optimal reinsurance strategy and show that the optimal reinsurance strategy for minimizing the probability of drawdown coincides with the one for minimizing the probability of ruin, as in Angoshtari et al. [1]. There is also research related to optimization problems under drawdown from the perspective of individual investors; see, for example, Grossman and Zhou [12], Cvitanić and Karatzas [9], Elie and Touzi [10], Chen et al. [6], and Angoshtari et al. [2].

There are many papers that employ the diffusion approximation in the actuarial science literature. By comparison, research on optimization problems with jump-diffusion models or classical Cramér-Lundberg (CL) models are less. The reason is that explicit solutions for the latter models are difficult to derive. Some researchers use probabilistic techniques to verify the convergent relationship between the probability of ruin for the scaled model and that of its corresponding diffusion approximation; see, for example, Iglehart [15], Grandell [11], Asmussen [3], and Bäuerle [5]. More recently, Cohen and Young [7] use the comparison method from differential equations to find upper and lower bounds of the probability of ruin and prove that the probability of ruin for the scaled CL model converges to the probability of ruin for the corresponding diffusion approximation. Liang et al. [16] consider the problem of minimizing the probability of ruin with reinsurance under the mean-variance premium principle. They show that the minimum probability of ruin is the unique viscosity solution of the corresponding Hamilton-Jacobi-Bellman equation with boundary conditions. They prove that, under appropriate scaling of the CL risk model, the probability of ruin converges to its diffusion approximation. Liang and Young [17] extend Cohen and Young [7] to the exponential Parisian ruin model. Cohen and Young [8] prove asymptotic results for the optimal-dividend problem.

In this paper, we analyze the relationship between the probability of drawdown for the scaled classical CL risk model and its corresponding diffusion approximation. We first prove two "smooth" comparison results. Then, we use the comparison results and the analogs of adjustment coefficients to prove that the minimum probability of drawdown for the scaled classical risk model converges to the minimum probability for its diffusion approximation uniformly, with rate of convergence of order $\mathcal{O}(n^{-1/2})$. To justify using the diffusion approximation in this optimization problem, we show that using the optimal retention strategy from the diffusion approximation is $\mathcal{O}(n^{-1/2})$ -optimal in the scaled classical CL risk model.

The rest of the paper is organized as follows. In Section 2.1, we present the classical CL model for the insurer's surplus, we describe the reinsurance market, and we define the minimum probability of drawdown for the classical risk process. Then, in Section 2.2, we provide two comparison theorems that form the backbone of the proofs in Section 4.1. In Section 2.3, we define the scaled model and discuss how to extend the comparison theorems to this model. Section 2.4 provides an explicit expression for the minimum probability of drawdown for the diffusion approximation of the scaled classical risk model, and Section 2.5 gives an outline of the remainder of the paper.

In Sections 3.1 and 3.2, we define analogs of the adjustment coefficient for the scaled classical risk model and for its diffusion approximation, respectively, and in Section 3.3, we prove two lemmas that show how these adjustment coefficients are related. In Section 4.1, we use the comparison lemma from Section 2.2 to modify the minimum probability of drawdown for the diffusion approximation by functions of order $O(n^{-1/2})$ to obtain upper and lower bounds of the corresponding probability of drawdown for the scaled classical risk model. In Section 4.2, we use these bounds to prove that the minimum probability of drawdown for the scaled classical risk model converges to that of its diffusion approximation. We show that the rate of convergence is of order $\mathcal{O}(n^{-1/2})$, uniformly with respect to the surplus. Finally, in Section 4.3, we prove the main result of our paper, namely, that if the insurer uses the optimal strategy from the diffusion approximation, then the resulting probability of drawdown is $\mathcal{O}(n^{-1/2})$ -optimal. This result thereby justifies using the diffusion approximation when minimizing the probability of drawdown.

2 Scaled Cramér-Lundberg model and its diffusion approximation

2.1 Cramér-Lundberg model and the probability of drawdown

In this section, we describe the reinsurance market available to the insurance company, and we formulate the problem of minimizing the probability of drawdown. Assume that all random processes exist on the filtered probability space $(\Omega, \mathcal{F}, \mathbb{F} = \{\mathcal{F}_t\}_{t\geq 0}, \mathbb{P})$.

We model the insurer's claim process $C = \{C_t\}_{t \ge 0}$ according to a compound Poisson process, namely,

$$C_t = \sum_{i=1}^{N_t} Y_i,\tag{2.1}$$

in which the claim severities Y_1, Y_2, \ldots are independent and identically distributed according to a common cumulative distribution function F_Y , with $F_Y(0) = 0$, and in which the claim frequency $N = \{N_t\}_{t\geq 0}$ follows a Poisson process with parameter $\lambda > 0$. Let $S_Y = 1 - F_Y$ denote the survival function of the severity random variable Y, and assume that Y's moment generating function M_Y is finite in a neighborhood of 0. Also, assume that the insurer receives premium payable continuously at a rate $c > \lambda \mathbb{E}Y$, and assume that the Poisson process N is independent of the claim severity process $\{Y_i\}_{i\in\mathbb{N}}$.

Remark 2.1. From Lemma 2.3.1 in Rolski et al. [19], we know that, because $M_Y(t_0) < \infty$ for some $t_0 > 0$, then there exists b > 0 such that

$$S_Y(y) \le b e^{-t_0 y}, \qquad \forall y \ge 0.$$
(2.2)

In other words, if M_Y is finite in a neighborhood of 0, then S_Y has an exponentially decreasing right tail. Conversely, if (2.2) holds, that is, if S_Y has an exponentially decreasing right tail, then $M_Y(t) < \infty$ for all $t < t_0$.

We assume that the insurer can buy per-loss reinsurance, with a continuously payable premium computed according to the so-called *mean-variance* premium principle, which combines the expected-value and variance premium principles, with risk loadings θ and η , respectively. Specifically, if $R_t(\omega, y)$ represents the retained claim at time $t \ge 0$, as a function of the (possible) claim Y = y at that time and state of the world $\omega \in \Omega$, then reinsurance indemnifies the insurer by the amount $y - R_t(\omega, y)$ if there is a claim y at time $t \ge 0$ and $\omega \in \Omega$, and the time-t premium rate is given by

$$(1+\theta)\lambda\mathbb{E}(Y-R_t) + \frac{\eta}{2}\lambda\mathbb{E}((Y-R_t)^2).$$
(2.3)

Assume that

$$c < (1+\theta)\lambda \mathbb{E}Y + \frac{\eta}{2}\lambda \mathbb{E}(Y^2);$$
 (2.4)

in words, the insurer's premium income is not sufficient to buy full reinsurance, and let κ denote the positive difference

$$\kappa = (1+\theta)\lambda\mathbb{E}Y + \frac{\eta}{2}\lambda\mathbb{E}(Y^2) - c.$$
(2.5)

Definition 2.1. A strategy $\mathcal{R} = \{R_t\}_{t\geq 0}$ is an admissible retention strategy if it satisfies the following properties:

- (i) \mathcal{R} is predictable; that is, the function $(\omega, y) \mapsto R_t(\omega, y)$ is $\mathcal{F}_{t^-} \times \mathcal{B}(\mathbb{R}_+)$ -measurable for every $t \geq 0$, in which $\mathcal{B}(\mathbb{R}_+)$ denotes the Borel σ -algebra on \mathbb{R}_+ .
- (ii) $0 \leq R_t(\omega, y) \leq y$, for all $t \geq 0$ and $\omega \in \Omega$.
- (iii) The net premium of the controlled surplus is greater than the expected rate of claim payment, that is,

$$c - (1+\theta)\lambda \mathbb{E}(Y - R_t) - \frac{\eta}{2}\lambda \mathbb{E}((Y - R_t)^2) > \lambda \mathbb{E}R_t,$$
(2.6)

with probability one for all $t \ge 0$. Hald and Schmidli [13] and Liang and Guo [18], among others, refer to inequality (2.6) as the net-profit condition.

- (iv) \mathcal{R} is progressively measurable; that is, the function $(\omega, y, t) \mapsto R_t(\omega, y)$ is $\mathcal{F}_t \times \mathcal{B}(\mathbb{R}_+) \times \mathcal{B}(\mathbb{R}_+)$ measurable.
- When initial surplus equals x, denote the set of admissible strategies by \mathfrak{R}_x . Moreover, a function $R : \mathbb{R}_+ \to \mathbb{R}_+$ is an admissible retention function if

(i) The mapping $y \mapsto R(y)$ is $\mathcal{B}(\mathbb{R}_+)$ -measurable.

(ii)
$$0 \le R(y) \le y$$
, for all $y \ge 0$.

(*iii*)
$$c - (1+\theta)\lambda \mathbb{E}(Y-R) - \frac{\eta}{2}\lambda \mathbb{E}((Y-R)^2) > \lambda \mathbb{E}R$$

In other words, a function R is an admissible retention function if the constant strategy $\mathcal{R} = \{R_t \equiv R\}$ is an admissible retention strategy.

Given a retention strategy $\mathcal{R} \in \mathfrak{R}_x$, the insurer's surplus follows the dynamics

$$dX_t^{\mathcal{R}} = \left(c - (1+\theta)\lambda\mathbb{E}(Y-R_t) - \frac{\eta}{2}\lambda\mathbb{E}((Y-R_t)^2)\right)dt - R_t dN_t$$
$$= \left(-\kappa + \lambda\left((1+\theta)\mathbb{E}R_t + \eta\mathbb{E}(YR_t) - \frac{\eta}{2}\mathbb{E}(R_t^2)\right)\right)dt - R_t dN_t,$$
(2.7)

with $X_{0^-}^{\mathcal{R}} = x \ge 0.^1$ Define the corresponding maximum surplus process $M^{\mathcal{R}} = \{M_t^{\mathcal{R}}\}_{t\ge 0}$ by

$$M_t^{\mathcal{R}} = \max\left\{\sup_{0\le s\le t} X_s^{\mathcal{R}}, M_{0^-}^{\mathcal{R}}\right\},\tag{2.8}$$

with $M_0^{\mathcal{R}} = m \geq x$. We allow the surplus process to have a financial past, as embodied by the term $M_{0^{-}}^{\mathcal{R}}$ in (2.8). Drawdown is the time when the surplus process drops below the proportion $\alpha \in [0, 1)$ of its maximum value, that is, at the hitting time $\tau_{\alpha}^{\mathcal{R}}$ given by

$$\tau_{\alpha}^{\mathcal{R}} = \inf \left\{ t \ge 0 : \ X_t^{\mathcal{R}} < \alpha M_t^{\mathcal{R}} \right\}.$$
(2.9)

If $\alpha = 0$, then the time or event of drawdown is the same as that for ruin under the ruin level 0.

The goal of the insurer is to minimize its probability of drawdown by purchasing per-loss reinsurance. The corresponding minimum probability of drawdown ψ is defined by

$$\psi(x,m) = \inf_{\mathcal{R}\in\mathfrak{R}_x} \mathbb{P}^{x,m} \big(\tau_{\alpha}^{\mathcal{R}} < \infty \big) = \inf_{\mathcal{R}\in\mathfrak{R}_x} \mathbb{E}^{x,m} \big(\mathbf{1}_{\{\tau_{\alpha}^{\mathcal{R}}<\infty\}} \big),$$
(2.10)

in which $\mathbb{P}^{x,m}$ and $\mathbb{E}^{x,m}$ denote the probability and expectation, respectively, conditional on $X_0 = x$ and $M_0 = m$. Note that, if $x < \alpha m$, then $\psi(x,m) = 1$. It remains for us to study the minimum probability of drawdown ψ on the domain

$$\mathcal{D} = \left\{ (x, m) \in \mathbb{R}^2_+ : \alpha m \le x \le m \right\}.$$
(2.11)

In the following proposition, we prove some interesting properties of ψ .

¹We assume that no claim occurs at time 0, which is true with probability 1, so we also have $X_0^{\mathcal{R}} = x \ge 0$.

Proposition 2.1. $\psi(x, \cdot)$ is nondecreasing on \mathcal{D} , and $\psi(\cdot, m)$ is nonincreasing and Lipschitz on \mathcal{D} . Moreover,

$$\lim_{m \to \infty} \psi(m, m) = 0. \tag{2.12}$$

Proof. For a fixed value of x, if $(x \leq) m_1 < m_2$, then the ruin levels are ordered, that is, $\alpha m_1 < \alpha m_2$, which implies that the probability of drawdown under ruin level αm_1 is less than or equal to the probability of drawdown under αm_2 . In other words, $\psi(x, \cdot)$ is nondecreasing on \mathcal{D} .

For a fixed value of m, if $\alpha m \leq x_1 < x_2 \leq m$, then clearly the probability of drawdown when surplus equals x_1 is greater than or equal to the probability of drawdown when surplus equals x_2 . In other words, $\psi(\cdot, m)$ is nonincreasing on \mathcal{D} . It remains to show that $\psi(\cdot, m)$ is Lipschitz on \mathcal{D} .

For a fixed value of m, suppose $\alpha m \leq x_1 < x_2 \leq m$. Choose an admissible retention function \hat{R} , and form the constant admissible retention strategy $\hat{\mathcal{R}} = {\{\hat{R}\}_{t\geq 0} \in \mathfrak{R}_{x_1}}$. Given $\varepsilon > 0$, let $\bar{\mathcal{R}}_1 = {\{R_t^1\}_{t\geq 0} \in \mathfrak{R}_{x_2}}$ be such that

$$\psi(x_2, m; \mathcal{R}_1) \le \psi(x_2, m) + \varepsilon$$

in which $\psi(\cdot, \cdot; \mathcal{R})$ denotes the probability of drawdown when the insurer follows strategy \mathcal{R} .

Now, define $\overline{\mathcal{R}} = \{R_t\}_{t \ge 0} \in \mathfrak{R}_{x_1}$ as follows:

$$R_t = \begin{cases} \hat{R}, & t \le \tau_{x_2} := \inf\{t \ge 0 : X_t^{\bar{\mathcal{R}}} = x_2\}, \\ R_{t-\tau_{x_2}}^1, & t > \tau_{x_2}. \end{cases}$$

If no claims occur, then the process $X^{\bar{\mathcal{R}}}$ with initial surplus x_1 reaches x_2 at time $h = (x_2 - x_1)/p_{\hat{R}}$, in which

$$p_R = c - (1+\theta)\lambda \mathbb{E}(Y-R) - \frac{\eta}{2}\lambda \mathbb{E}((Y-R)^2).$$
(2.13)

Note that condition (iii) in Definition 2.1 implies $p_R > \lambda \mathbb{E}R$ for any admissible retention function R. Thus, if we let τ_1 denote the time of the first claim when initial surplus equals x_1 , we have

$$1 - \psi(x_1, m) \ge 1 - \psi(x_1, m; \mathcal{R})$$

$$\ge (1 - \psi(x_2, m; \overline{\mathcal{R}})) \mathbb{P}(\tau_1 > h)$$

$$= (1 - \psi(x_2, m; \overline{\mathcal{R}})) e^{-\lambda h}$$

$$\ge (1 - \psi(x_2, m) - \varepsilon) e^{-\lambda h}.$$

Because $\varepsilon > 0$ is arbitrary, we deduce

$$1 - \psi(x_1, m) \ge (1 - \psi(x_2, m))e^{-\lambda h}.$$
(2.14)

From inequality (2.14) and $\psi(\cdot, m)$ nonincreasing, we obtain

$$0 \le \psi(x_1, m) - \psi(x_2, m) \le (1 - \psi(x_2, m))(1 - e^{-\lambda h})$$

$$\le 1 - e^{-\lambda h} \le \lambda h = \frac{\lambda}{p_{\hat{R}}} (x_2 - x_1).$$

Because \hat{R} is arbitrary such that $p_{\hat{R}} > \lambda \mathbb{E} \hat{R}$, it follows that

$$\left|\psi(x_1,m) - \psi(x_2,m)\right| \le \frac{\lambda}{\sup_R p_R} |x_2 - x_1|.$$

Thus, ψ is Lipschitz in x.

It remains to prove the limit in (2.12). To that end, for any value $(x,m) \in \mathcal{D}$, let R be an admissible retention function, and let $\mathcal{R} = \{R\}_{t\geq 0} \in \mathfrak{R}$ be the corresponding constant retention strategy. We wish to prove that

$$\psi(x,m;\mathcal{R}) \le e^{-J(x-\alpha m)},\tag{2.15}$$

in which J is the positive solution of the following equation:

$$\lambda \left(\mathbb{E}e^{JR} - 1 - J\mathbb{E}R \right) = J \left[-\kappa + \lambda \left(\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\eta}{2} \mathbb{E}(R^2) \right) \right].$$
(2.16)

As an aside, to see that (2.16) has a unique positive solution, note that condition (iii) of Definition 2.1 implies the coefficient of J on the right side is positive. The two sides equal 0 when J = 0, and the rates of growth with respect to J when J = 0 of the left and right sides equal, respectively, 0 and the positive coefficient in square brackets. Thus, because the left side grows exponentially and the right grows linearly with J, there is a unique positive solution of (2.16).

Next, let $\psi_k(\cdot, \cdot; \mathcal{R})$ denote the probability of drawdown (under the strategy \mathcal{R}) at or before the k^{th} claim. For all $(x,m) \in \mathcal{D}$, $\psi_k(x,m;\mathcal{R})$ increases with respect to k, and $\lim_{k\to\infty} \psi_k(x,m;\mathcal{R}) = \psi(x,m;\mathcal{R})$. Indeed, if we let T_k denote the time of the k^{th} claim, then $\psi_k(x,m;\mathcal{R}) = \mathbb{P}(\tau_{\alpha}^{\mathcal{R}} \leq T_k)$, and $\lim_{k\to\infty} T_k = \infty$, which give us

$$\lim_{k \to \infty} \psi_k(x, m; \mathcal{R}) = \lim_{k \to \infty} \mathbb{P}(\tau_\alpha^{\mathcal{R}} \le T_k) = \mathbb{P}(\bigcup_{k=1}^\infty \{\tau_\alpha^{\mathcal{R}} \le T_k\}) = \mathbb{P}(\tau_\alpha^{\mathcal{R}} < \infty) = \psi(x, m; \mathcal{R}).$$

Thus, to prove (2.15), it is enough to prove

$$\psi_k(x,m;\mathcal{R}) \le e^{-J(x-\alpha m)},\tag{2.17}$$

for all $k \in \mathbb{N}$. By a recursive argument, ψ_k solves

$$\psi_k(x,m;\mathcal{R}) = \int_0^\infty \int_0^\infty \psi_{k-1}(x+p_R t - R(y),m;\mathcal{R})\lambda e^{-\lambda t} dF_Y(y)dt, \qquad (2.18)$$

for $k \in \mathbb{N}$, in which

$$\psi_0(x,m;\mathcal{R}) = \begin{cases} 0, & x \ge \alpha m, \\ 1, & x < \alpha m. \end{cases}$$
(2.19)

For $(x,m) \in \mathcal{D}$, $\psi_0(x,m;\mathcal{R}) = 0 \le e^{-J(x-\alpha m)}$. Assume (2.17) holds for some $k-1 = 0, 1, 2, \ldots$; then,

$$\psi_k(x,m;\mathcal{R}) \leq \int_0^\infty \int_0^\infty e^{-J(x+p_Rt-R(y)-\alpha m)} \lambda e^{-\lambda t} dF_Y(y) dt$$

= $e^{-J(x-\alpha m)} \int_0^\infty \lambda e^{JR(y)} dF_Y(y) \cdot \int_0^\infty e^{-(\lambda+Jp_R)t} dt$
= $e^{-J(x-\alpha m)} (\lambda+Jp_R) \cdot \frac{1}{\lambda+Jp_R} = e^{-J(x-\alpha m)},$

in which the second equality follows from (2.16). Thus, inequality (2.17) follows by recursion, which implies inequality (2.15). From the minimality of ψ and from (2.15), we obtain

$$0 \le \lim_{m \to \infty} \psi(m, m) \le \lim_{m \to \infty} \psi(m, m; \mathcal{R}) \le \lim_{m \to \infty} e^{-J(1-\alpha)m} = 0,$$

which proves the limit in (2.12).

2.2 Comparison theorems

For the classical risk model, we cannot find an explicit expression for the minimum probability of drawdown ψ , so in Section 4, we prove that the minimum probability of drawdown for the diffusion approximation of $X^{\mathcal{R}}$ approximates ψ . To that end, in this section, we prove two comparison theorems which we use in Section 4.

We present the following lemma, which we will use in the proofs of Theorems 2.1 and 2.2 below.

Lemma 2.1. For $(x,m) \in \mathcal{D}$, fix an admissible retention strategy $\mathcal{R} \in \mathfrak{R}_x$. Define $s_b = \inf\{t \ge 0 : X_t^{\mathcal{R}} \ge b\}$ with $X_0^{\mathcal{R}} = x$, and define $s_{\alpha b} = \tau_\alpha \wedge s_b$.² Then, $\mathbb{P}^{x,m}(s_{\alpha b} < \infty) = 1$. ²These stopping times depend on \mathcal{R} via $X^{\mathcal{R}}$, but for simplicity of notation in the remainder of this section, we

²These stopping times depend on \mathcal{R} via $X^{\mathcal{R}}$, but for simplicity of notation in the remainder of this section, we suppress the superscript \mathcal{R} on τ_{α} , s_b , and $s_{\alpha b}$.

Proof. Proposition 2.2 in Azcue and Muler [4] states: with probability one, either ruin occurs in finite time or $X_t^{\mathcal{R}}$ diverges to infinity as t goes to infinity. Because $\alpha M_t \ge 0$ and because $b < \infty$, it follows that $\mathbb{P}^{x,m}(s_{\alpha b} < \infty) = 1$.

In the first theorem, we show that any smooth subsolution of our problem is less than or equal to the value function. Before stating the theorem, we introduce some notation. For an admissible retention function R, define the operator \mathcal{L}^R on $\mathcal{C}^{1,1}(\mathcal{D})$ as follows: for $u \in \mathcal{C}^{1,1}(\mathcal{D})$,

$$\mathcal{L}^{R}u(x,m) = -\kappa u_{x}(x,m) + \lambda \left[\left((1+\theta)\mathbb{E}R + \eta\mathbb{E}(YR) - \frac{\eta}{2}\mathbb{E}(R^{2}) \right) u_{x}(x,m) + \mathbb{E}u(x-R,m) - u(x,m) \right],$$
(2.20)

in which we extend u by defining u(x,m) = 1 for $x < \alpha m$. Note that the HJB equation for ψ is $\inf_R \mathcal{L}^R \psi(x,m) = 0$.

Theorem 2.1 (Subsolution). Suppose $u \in C^{1,1}(D)$ is a bounded function that satisfies the following conditions:

- $(i) \ \lim_{m \to \infty} u(m,m) \leq 0.$
- (ii) u(x,m) is defined for all $x \le m$ and $m \ge 0$, with u(x,m) = 1 for all $x < \alpha m$.
- (iii) $u_m(m,m) \ge 0$ for all $m \ge 0$.
- (iv) $\mathcal{L}^{R}u(x,m) \geq 0$ for all $(x,m) \in \mathcal{D}$ and for all admissible retention functions R.

Then, $u \leq \psi$ on \mathcal{D} .

Proof. Assume that u satisfies the conditions specified in the statement of this theorem, and fix an admissible retention strategy \mathcal{R} .

For a fixed value of $m \ge 0$, let b > m, and define s_b and $s_{\alpha b}$ as in Lemma 2.1. By applying Itô's

formula to u(x, m), we have

$$u(X_{s_{\alpha b}}^{\mathcal{R}}, M_{s_{\alpha b}}^{\mathcal{R}}) = u(x, m) + \int_{0}^{s_{\alpha b}} \left(-\kappa + \lambda \left((1+\theta) \mathbb{E}R_{t} + \eta \mathbb{E}(YR_{t}) - \frac{\eta}{2} \mathbb{E}(R_{t}^{2}) \right) \right) u_{x} \left(X_{t^{-}}^{\mathcal{R}}, M_{t}^{\mathcal{R}} \right) dt + \int_{0}^{s_{\alpha b}} \left(\mathbb{E}u(X_{t^{-}}^{\mathcal{R}} - R_{t}, M_{t}^{\mathcal{R}}) - u(X_{t^{-}}^{\mathcal{R}}, M_{t}^{\mathcal{R}}) \right) dN_{t} + \int_{0}^{s_{\alpha b}} u_{m} \left(X_{t^{-}}^{\mathcal{R}}, M_{t}^{\mathcal{R}} \right) dM_{t}^{\mathcal{R}} = u(x, m) + \int_{0}^{s_{\alpha b}} \mathcal{L}^{R_{t}} u(X_{t^{-}}^{\mathcal{R}}, M_{t}^{\mathcal{R}}) dt + \int_{0}^{s_{\alpha b}} \left(\mathbb{E}u(X_{t^{-}}^{\mathcal{R}} - R_{t}, M_{t}^{\mathcal{R}}) - u(X_{t^{-}}^{\mathcal{R}}, M_{t}^{\mathcal{R}}) \right) d(N_{t} - \lambda t) + \int_{0}^{s_{\alpha b}} u_{m} \left(X_{t^{-}}^{\mathcal{R}}, M_{t}^{\mathcal{R}} \right) dM_{t}^{\mathcal{R}}.$$

$$(2.21)$$

The first integral in (2.21) is non-negative because of condition (iv) of the theorem. The expectation of the second integral equals 0 because u is bounded. The third integral is non-negative almost surely because $dM_t^{\mathcal{R}}$ is non-zero only when $M_t^{\mathcal{R}} = X_t^{\mathcal{R}}$ and $u_m(m,m) \ge 0$ by condition (iii). Here, we also used the fact that $M^{\mathcal{R}}$ is non-decreasing; therefore, the first variation process associated with it is finite almost surely, and we conclude that the cross variation of $M^{\mathcal{R}}$ and $X^{\mathcal{R}}$ is zero almost surely. Thus, by taking expectations in (2.21), we have

$$\mathbb{E}^{x,m}\left[u\left(X_{s_{\alpha b}}^{\mathcal{R}}, M_{s_{\alpha b}}^{\mathcal{R}}\right)\right] \ge u(x,m).$$
(2.22)

Because $s_{\alpha b} < \infty$ with probability 1 and because b > m, it follows from the extension of u to u(x,m) = 1 for all $x < \alpha m$ and from inequality (2.22) that

$$u(x,m) \leq \mathbb{P}^{x,m}(\tau_{\alpha} < s_{b}) \cdot 1 + \mathbb{P}^{x,m}(s_{b} < \tau_{\alpha}) \cdot u(b,b)$$
$$\leq \mathbb{P}^{x,m}(\tau_{\alpha} < \infty) + \mathbb{P}^{x,m}(s_{b} < \tau_{\alpha}) \cdot u(b,b).$$
(2.23)

In (2.23), we purposefully omit $\mathbb{P}^{x,m}(\tau_{\alpha} = s_b)$ because that probability equals 0, which follows from $\mathbb{P}^{x,m}(s_{\alpha b} < \infty) = 1$ and $\alpha m < b$.

By applying the Dominated Convergence Theorem to (2.23) as we take the limit $b \to \infty$ and by using $\lim_{b\to\infty} u(b,b) \leq 0$ from condition (i), we obtain

$$u(x,m) \le \mathbb{P}^{x,m}(\tau_{\alpha} < \infty). \tag{2.24}$$

By taking the infimum over admissible strategies, we obtain $u \leq \psi$ on \mathcal{D} .

In the second theorem, we show that any smooth supersolution of our problem is greater than or equal to the value function. **Theorem 2.2** (Supersolution). Suppose $v \in C^{1,1}(D)$ is a bounded function that satisfies the following conditions:

- $(i) \ \lim_{m \to \infty} v(m,m) \ge 0.$
- (ii) v(x,m) is defined for all $x \le m$ and $m \ge 0$, with v(x,m) = 1 for all $x < \alpha m$.
- (iii) $v_m(m,m) \leq 0$ for all $m \geq 0$.
- (iv) $\mathcal{L}^{\hat{R}}v(x,m) \leq 0$ for all $(x,m) \in \mathcal{D}$ and for some admissible retention function \hat{R} .

Then,
$$\psi \leq v$$
 on \mathcal{D} .

Proof. Assume that v and \hat{R} satisfy the conditions specified in the statement of this theorem, and let $\hat{\mathcal{R}}$ denote the constant retention strategy corresponding to \hat{R} .

For a fixed value of $m \ge 0$, let b > m, and define s_b and $s_{\alpha b}$ as in Lemma 2.1. By applying Itô's formula to v(x, m), we have

$$v\left(X_{s_{\alpha b}}^{\hat{\mathcal{R}}}, M_{s_{\alpha b}}^{\hat{\mathcal{R}}}\right) = v(x, m) + \int_{0}^{s_{\alpha b}} \left(-\kappa + \lambda \left((1+\theta)\mathbb{E}\hat{R}_{t} + \eta\mathbb{E}(Y\hat{R}_{t}) - \frac{\eta}{2}\mathbb{E}(\hat{R}_{t}^{2})\right)\right) v_{x}\left(X_{t^{-}}^{\hat{\mathcal{R}}}, M_{t}^{\hat{\mathcal{R}}}\right) dt
 + \int_{0}^{s_{\alpha b}} \left(\mathbb{E}v\left(X_{t^{-}}^{\hat{\mathcal{R}}} - \hat{R}_{t}, M_{t}^{\hat{\mathcal{R}}}\right) - v\left(X_{t^{-}}^{\hat{\mathcal{R}}}, M_{t}^{\hat{\mathcal{R}}}\right)\right) dN_{t} + \int_{0}^{s_{\alpha b}} v_{m}\left(X_{t^{-}}^{\hat{\mathcal{R}}}, M_{t}^{\hat{\mathcal{R}}}\right) dM_{t}^{\hat{\mathcal{R}}}
 = v(x, m) + \int_{0}^{s_{\alpha b}} \mathcal{L}^{\hat{R}_{t}}v\left(X_{t^{-}}^{\hat{\mathcal{R}}}, M_{t}^{\hat{\mathcal{R}}}\right) dt
 + \int_{0}^{s_{\alpha b}} \left(\mathbb{E}v\left(X_{t^{-}}^{\hat{\mathcal{R}}} - \hat{R}_{t}, M_{t}^{\hat{\mathcal{R}}}\right) - v\left(X_{t^{-}}^{\hat{\mathcal{R}}}, M_{t}^{\hat{\mathcal{R}}}\right)\right) d(N_{t} - \lambda t)
 + \int_{0}^{s_{\alpha b}} v_{m}\left(X_{t^{-}}^{\hat{\mathcal{R}}}, M_{t}^{\hat{\mathcal{R}}}\right) dM_{t}^{\hat{\mathcal{R}}}.$$
(2.25)

The first integral in (2.25) is non-positive because of condition (iv) of the theorem. The expectation of the second integral equals 0 because v is bounded. The third integral is non-positive almost surely because $dM_t^{\hat{\mathcal{R}}}$ is non-zero only when $M_t^{\hat{\mathcal{R}}} = X_t^{\hat{\mathcal{R}}}$ and $v_m(m,m) \leq 0$ by condition (iii). Thus, by taking expectations in (2.25), we have

$$\mathbb{E}^{x,m}\left[v\left(X_{s_{\alpha b}}^{\hat{\mathcal{R}}}, M_{s_{\alpha b}}^{\hat{\mathcal{R}}}\right)\right] \le v(x,m).$$
(2.26)

Because $s_{\alpha b} < \infty$ with probability 1 and because b > m, it follows from the extension of v to v(x,m) = 1 for all $x < \alpha m$ and from inequality (2.26) that

$$v(x,m) \ge \mathbb{P}^{x,m}(\tau_{\alpha} < s_b) \cdot 1 + \mathbb{P}^{x,m}(s_b < \tau_{\alpha}) \cdot v(b,b).$$
(2.27)

Next, we wish to show that

$$\lim_{b \to \infty} \{\tau_{\alpha} < s_b\} = \{\tau_{\alpha} < \infty\}.$$
(2.28)

Because $\{\tau_{\alpha} < s_b\} \subseteq \{\tau_{\alpha} < \infty\}$ for all b > m, it follows that $\lim_{b\to\infty} \{\tau_{\alpha} < s_b\} \subseteq \{\tau_{\alpha} < \infty\}$. To prove the opposite inclusion, suppose $\omega \in \{\tau_{\alpha} < \infty\}$, that is, drawdown occurs in finite time, say, at time t. Thus, $M_t^{\hat{\mathcal{R}}} < \infty$, which implies $\omega \in \{\tau_{\alpha} < s_b\}$ for all $b > M_t^{\hat{\mathcal{R}}}$, and we have shown that $\{\tau_{\alpha} < \infty\} \subseteq \lim_{b\to\infty} \{\tau_{\alpha} < s_b\}$, from which (2.28) follows.

By applying the Dominated Convergence Theorem to (2.27) as we take the limit $b \to \infty$, by using (2.28), and by using $\lim_{b\to\infty} v(b,b) \ge 0$ from condition (i), we obtain

$$v(x,m) \ge \mathbb{P}^{x,m}(\tau_{\alpha} < \infty), \tag{2.29}$$

which implies $v \ge \psi$ on \mathcal{D} .

2.3 Scaled Cramér-Lundberg model

In this section, we scale the Cramér-Lundberg risk model by n > 0. To obtain the scaled model, multiply the Poisson rate λ by n, divide the claim severity by \sqrt{n} , and adjust the premium rate so that net premium income remains constant. Specifically, define $\lambda_n = n\lambda$, so n large is equivalent to λ_n large. Scale the claim severity by defining $Y_n = Y/\sqrt{n}$. Also, define $\theta_n = \theta/\sqrt{n}$ and $c_n = c + (\sqrt{n} - 1)\lambda \mathbb{E}Y$, which implies $c_n - \lambda_n \mathbb{E}_n Y_n = c - \lambda \mathbb{E}Y$, independent of n.³ The parameter η remains unchanged. Finally, define

$$\kappa_n = (1+\theta_n)\lambda_n \mathbb{E}_n Y_n + \frac{\eta}{2}\lambda_n \mathbb{E}_n (Y_n^2) - c_n$$
$$= (1+\theta)\lambda \mathbb{E}Y + \frac{\eta}{2}\lambda \mathbb{E}(Y^2) - c = \kappa,$$

so κ_n is also independent of n.

In the scaled model, define an *n*-admissible retention strategy $\mathcal{R}_n = \{(R_n)_t\}_{t\geq 0}$ is as in Definition 2.1, except with condition (iii) replaced by

$$(iii') \qquad c_n - (1+\theta_n)\lambda_n \mathbb{E}_n(Y_n - (R_n)_t) - \frac{\eta}{2}\lambda_n \mathbb{E}_n((Y_n - (R_n)_t)^2) > \lambda_n \mathbb{E}_n(R_n)_t.$$
(2.30)

³By writing \mathbb{E}_n , we mean expectation with respect to the measure induced by Y_n . Specifically,

$$\mathbb{E}_n(g(Y_n)) = \int_0^\infty g(y)dF_{Y_n}(y) = \int_0^\infty g(y)dF_Y(\sqrt{ny})$$
$$= \int_0^\infty g(t/\sqrt{n})dF_Y(t) = \mathbb{E}(g(Y/\sqrt{n})),$$

in which $\mathbb{E} = \mathbb{E}_1$.

Similarly, define an *n*-admissible retention function R_n . If we are given an *n*-admissible retention strategy $\mathcal{R}_n = \{(R_n)_t\}_{t\geq 0}$, then we can define an admissible retention strategy $\mathcal{R} = \{R_t\}_{t\geq 0}$ by

$$R_t(\omega, y) = \sqrt{n} \left(R_n \right)_t(\omega, y/\sqrt{n} \right).$$
(2.31)

Indeed, by using the assignment in (2.31), condition (iii') in (2.30) implies

$$(c + (\sqrt{n} - 1)\lambda\mathbb{E}Y) - \left(1 + \frac{\theta}{\sqrt{n}}\right)n\lambda\mathbb{E}\left(\frac{Y - R_t}{\sqrt{n}}\right) - \frac{\eta}{2}n\lambda\mathbb{E}\left(\left(\frac{Y - R_t}{\sqrt{n}}\right)^2\right) > n\lambda\mathbb{E}\left(\frac{R_t}{\sqrt{n}}\right)$$
$$\iff c - \lambda(1 + \theta)\mathbb{E}Y + \lambda\theta\mathbb{E}R_t - \frac{\eta}{2}\lambda\mathbb{E}((Y - R_t)^2) > 0$$
$$\iff c - \lambda(1 + \theta)\mathbb{E}(Y - R_t) - \frac{\eta}{2}\lambda\mathbb{E}((Y - R_t)^2) > \lambda\mathbb{E}R_t.$$

Conversely, given an admissible strategy \mathcal{R} , we can define an *n*-admissible strategy \mathcal{R}_n via (2.31). Thus, we deduce there is a one-to-one correspondence between admissible strategies and *n*-admissible strategies. Similarly, via the relationship

$$R(y) = \sqrt{n} R_n(y/\sqrt{n}), \qquad (2.32)$$

we obtain a one-to-one correspondence between admissible retention functions and n-admissible retention functions.

Let $X_n^{\mathcal{R}_n}$ denote the surplus process for the scaled model; thus, $X_n^{\mathcal{R}_n}$ follows the dynamics

$$d(X_n^{\mathcal{R}_n})_t = \left(-\kappa_n + \lambda_n \left((1+\theta_n)\mathbb{E}_n(R_n)_t + \eta\mathbb{E}_n(Y_n(R_n)_t) - \frac{\eta}{2}\mathbb{E}_n((R_n)_t^2)\right)\right)dt - (R_n)_t d(N_n)_t$$
$$= \left(-\kappa + n\lambda \left((1+\theta/\sqrt{n})\mathbb{E}_n(R_n)_t + \eta\mathbb{E}_n(Y(R_n)_t/\sqrt{n}) - \frac{\eta}{2}\mathbb{E}_n((R_n)_t^2)\right)\right)dt - (R_n)_t d(N_n)_t$$
(2.33)

in which N_n denotes the Poisson process with rate $\lambda_n = n\lambda$. Let ψ_n denote the minimum probability of drawdown for the scaled system, and note that Theorems 2.1 and 2.2 apply to ψ_n with \mathcal{L}^R replaced by $\mathcal{L}_n^{R_n}$, in which $\mathcal{L}_n^{R_n}$ is defined as follows: for $u \in \mathcal{C}^{1,1}(\mathcal{D})$ and for an *n*-admissible retention function R_n ,

$$\mathcal{L}_{n}^{R_{n}}u(x,m) = -\kappa_{n}u_{x}(x,m) + \lambda_{n}\left[\left((1+\theta_{n})\mathbb{E}_{n}R_{n}+\eta\mathbb{E}_{n}(Y_{n}R_{n})-\frac{\eta}{2}\mathbb{E}_{n}(R_{n}^{2})\right)u_{x}(x,m) + \mathbb{E}_{n}u(x-R_{n},m)-u(x,m)\right].$$
 (2.34)

In (2.34), we again extend u by defining u(x,m) = 1 for $x < \alpha m$.

2.4 Diffusion approximation

Let $X_{n,D}^{\mathcal{R}}$ denote the diffusion approximation to the *n*-scaled process in (2.33), which we form by

$$(R_n)_t d(N_n)_t \approx n\lambda \mathbb{E}_n(R_n)_t dt - \sqrt{n\lambda \mathbb{E}_n((R_n)_t^2)} dB_t,$$

in which B is a standard Brownian motion on the filtered probability space. Then, $X_{n,D}^{\mathcal{R}_n}$ follows the dynamics

$$d(X_{n,D}^{\mathcal{R}_n})_t = \left(-\kappa + n\lambda\left((\theta/\sqrt{n})\mathbb{E}_n(R_n)_t + \eta\mathbb{E}_n(Y_n(R_n)_t) - \frac{\eta}{2}\mathbb{E}_n((R_n)_t^2)\right)\right)dt + \sqrt{n\lambda\mathbb{E}_n((R_n)_t^2)}dB_t$$
$$= \left(-\kappa + \lambda\left(\theta\mathbb{E}_n(\sqrt{n}(R_n)_t) + \eta\mathbb{E}_n(\sqrt{n}Y_n \cdot \sqrt{n}(R_n)_t) - \frac{\eta}{2}\mathbb{E}_n(n(R_n)_t^2)\right)\right)dt$$
$$+ \sqrt{\lambda\mathbb{E}_n(n(R_n)_t^2)}dB_t$$
$$= \left(-\kappa + \lambda\left(\theta\mathbb{E}R_t + \eta\mathbb{E}(YR_t) - \frac{\eta}{2}\mathbb{E}(R_t^2)\right)\right)dt + \sqrt{\lambda\mathbb{E}(R_t^2)}dB_t,$$
(2.35)

in which we define the strategy $\mathcal{R} = \{R_t\}_{t\geq 0}$ by the assignment in (2.31). Thus, the dynamics of $X_{n,D}^{\mathcal{R}_n}$ are independent of n after this assignment, which implies that the minimum probability of ruin for the diffusion approximation of the *n*-scaled process is independent of n.

Let ψ_D denote the minimum probability of drawdown for the diffusion approximation of the (scaled) Cramér-Lundberg model. Han et al. [14] solved the optimization problem associated with ψ_D . The following theorem is a limiting case (as the riskless rate $r \to 0^+$) of Theorem 3.2, the main result of Han et al. [14].

Theorem 2.3. The minimum probability of drawdown ψ_D on

$$\mathcal{D} = \left\{ (x, m) \in \mathbb{R}^2_+ : \alpha m \le x \le m \right\}$$

under the diffusion approximation in (2.35) equals

$$\psi_D(x,m) = 1 - h_D(m) \left(1 - e^{-\rho_D(x - \alpha m)} \right), \qquad (2.36)$$

in which h_D is defined by

$$h_D(m) = \left(1 - e^{-\rho_D(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}},$$
 (2.37)

and $\rho_D > 0$ uniquely solves

$$c - \lambda \mathbb{E}Y = \lambda \rho \int_0^\infty \left(\frac{\theta + \eta y}{\rho + \eta} \wedge y\right) S_Y(y) dy.$$
(2.38)

The corresponding optimal retention strategy is a constant strategy $\{R_D\}_{t\geq 0}$, in which

$$R_D(y) = \frac{\theta + \eta y}{\rho_D + \eta} \wedge y, \qquad (2.39)$$

with $y \ge 0$ the possible claim size.

Remark 2.2. If we reverse (2.32) as applied to $R = R_D$ in (2.39), then we obtain

$$R_{D,n}(y) := \frac{1}{\sqrt{n}} R_D(\sqrt{n}y) = \frac{1}{\sqrt{n}} \left(\frac{\theta + \eta \sqrt{n}y}{\rho_D + \eta} \wedge \sqrt{n}y \right) = \frac{\theta_n + \eta y}{\rho_D + \eta} \wedge y, \tag{2.40}$$

that is, we get the optimal retention function for the diffusion approximation of the n-scaled model, as expected, from the discussion following (2.35).

Remark 2.3. In Section 3.2, we show that ρ_D is the maximum adjustment coefficient, which parallels a similar result in Liang, Liang, and Young [16].

An important result of this paper is that ψ_D and ψ_n are approximately equal, specifically, to order $\mathcal{O}(n^{-1/2})$. The technique for showing that ψ_D approximates ψ_n is to modify ψ_D by a function of order $\mathcal{O}(n^{-1/2})$ and, then, prove that the modified function is a sub- or supersolution of ψ_n via Theorem 2.1 or 2.2, respectively. In the next section, we detail the steps in the remainder of the paper.

2.5 Outline of paper

To make it easier to follow the material in Sections 3 and 4, we outline our steps in those sections:

- 1. In Section 3, we define analogs of the so-called *adjustment coefficient* from risk theory; for background on the adjustment coefficient see, for example, Section 5.4 in Schmidli [20].
 - (a) In Section 3.1, we define an analog of the maximum adjustment coefficient for the scaled Cramér-Lundberg model, which we denote by ρ_n .
 - (b) In Section 3.2, we show that ρ_D from Section 2.4 is the analog of the maximum adjustment coefficient for the diffusion approximation.
 - (c) In Section 3.3, we prove two lemmas that relate ρ_n and ρ_D . In particular, we show $\lim_{n \to \infty} \rho_n = \rho_D$.
- 2. In Section 4, we justify using the diffusion approximation for the classical risk process when analyzing the minimum probability of drawdown.

- (a) In Section 4.1, we modify ψ_D by functions of order $\mathcal{O}(n^{-1/2})$ to obtain upper and lower bounds of ψ_n ; denote those bounds by u_n and ℓ_n , respectively. We use Theorems 2.1 and 2.2 to prove that, indeed, $\ell_n \leq \psi_n \leq u_n$ on \mathbb{R} . Propositions 4.2 and 4.1 prove these two inequalities, respectively.
- (b) In Section 4.2, we use Propositions 4.1 and 4.2 to prove Theorem 4.1, which states that, as n goes to infinity, ψ_n converges to ψ_D uniformly on D with rate of convergence of order O(n^{-1/2}).
- (c) Finally, in Section 4.3, we prove the main result of our paper, namely, that if the insurer uses the optimal strategy from the diffusion approximation, then the resulting probability of drawdown is $\mathcal{O}(n^{-1/2})$ -optimal. This result thereby justifies using the diffusion approximation when minimizing the probability of drawdown.

3 Analogs of the adjustment coefficient

3.1 Scaled Cramér-Lundberg model

In this section, we define the analog of the adjustment coefficient for the scaled Cramér-Lundberg model for the probability of drawdown. We will use that analog to create an exponential upper bound of ψ_n .

For a given *n*-admissible retention function R_n , formally obtain the adjustment coefficient $\rho_n(R_n) > 0$ by substituting $e^{-\rho x}$ for u in $\mathcal{L}_n^{R_n} u = 0$, including when $x - Y/\sqrt{n} < \alpha m$. When we perform this substitution, we obtain the following equation for $\rho_n(R_n)$:

$$\left[-\kappa_n + \lambda_n \left((1+\theta_n)\mathbb{E}_n R_n + \eta \mathbb{E}_n(Y_n R_n) - \frac{\eta}{2} \mathbb{E}_n(R_n^2)\right)\right] \rho - \lambda_n \left(M_{R_n}(\rho) - 1\right) = 0,$$

in which M_{R_n} denotes the moment generating function of R_n , which is finite in a neighborhood of 0 because M_Y is finite in a neighborhood of 0. After we substitute for the *n*-scaled parameters and set $R(y) = \sqrt{n}R_n(y/\sqrt{n})$ as in (2.32), this equation is equivalent to

$$n\lambda g_n(\rho; R) = \lambda \left(\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\eta}{2} \mathbb{E}(R^2)\right) - \kappa, \qquad (3.1)$$

in which we define g_n by

$$g_n(\rho; R) = \frac{1}{\rho} \left(\mathbb{E}e^{\frac{\rho}{\sqrt{n}}R} - 1 - \frac{\rho}{\sqrt{n}} \mathbb{E}R \right).$$
(3.2)

Condition (iii) in Definition 2.1 for an admissible retention function implies that the right side of (3.1) is positive; in fact, those two statements are equivalent. Also, $g_n(\rho; R)$ increases from 0^+ to infinity; thus, $\rho_n(R_n) > 0$ exists for any *n*-admissible retention function R_n .

Let ρ_n denote the maximum adjustment coefficient for the classical risk model, in which we maximize over *n*-admissible retention functions R_n . By following the argument in Section 4.1 of Liang, Liang, and Young [16], we deduce that ρ_n solves the following maximization problem:

$$\sup_{R_n} \left\{ \left[-\kappa_n + \lambda_n \left((1+\theta_n) \mathbb{E}_n R_n + \eta \mathbb{E}_n (Y_n R_n) - \frac{\eta}{2} \mathbb{E}_n (R_n^2) \right) \right] \rho - \lambda_n \left(M_{R_n}(\rho) - 1 \right) \right\} = 0, \quad (3.3)$$

in which we maximize over *n*-admissible retention functions R_n . By using the assignment in (2.32), this maximization problem is equivalent to

$$-\kappa + \lambda \sup_{R} \left[\left(\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\eta}{2} \mathbb{E}(R^2) \right) - ng_n(\rho; R) \right] = 0, \qquad (3.4)$$

in which we maximize over admissible retention functions R.

In the following proposition, we give an expression for ρ_n and the corresponding optimal *n*-retention function R_n^{ρ} .

Proposition 3.1. The maximum adjustment coefficient $\rho_n > 0$ for the n-scaled risk process in (2.33) uniquely solves

$$c - \lambda \mathbb{E}Y = n\lambda \int_0^\infty \left(e^{\rho R_n^\rho(y)} - 1 \right) S_{Y_n}(y) dy.$$
(3.5)

in which $S_{Y_n}(y) = S_Y(\sqrt{ny})$, for all $y \ge 0$, and the corresponding optimal n-retention function R_n^{ρ} is given by

$$R_{n}^{\rho}(y) = \begin{cases} y, & 0 \le y \le \frac{1}{\rho_{n}} \ln(1+\theta_{n}), \\ R_{c}(y), & y > \frac{1}{\rho_{n}} \ln(1+\theta_{n}). \end{cases}$$
(3.6)

In (3.6), $R_c(y) \in [0, y)$ for $y > \frac{1}{\rho_n} \ln(1 + \theta_n)$ uniquely solves

$$(1+\theta_n) + \eta(y-R) = e^{\rho_n R}.$$
 (3.7)

Proof. Consider the R-dependent terms in (3.4); rewrite them as follows:

$$\int_0^\infty \left[(\sqrt{n} + \theta) R(y) + \eta y R(y) - \frac{\eta}{2} R^2(y) - \frac{n}{\rho} e^{\frac{\rho}{\sqrt{n}} R(y)} \right] dF_Y(y)$$

If we maximize the integrand y-by-y, subject to $0 \le R(y) \le y$, then the integral itself is maximized. To that end, for a fixed value of ρ , define the function j by

$$j(R) = (\sqrt{n} + \theta)R + \eta y R - \frac{\eta}{2} R^2 - \frac{n}{\rho} e^{\frac{\rho}{\sqrt{n}}R}.$$
(3.8)

Then,

$$j'(R) = (\sqrt{n} + \theta) + \eta(y - R) - \sqrt{n} e^{\frac{p}{\sqrt{n}}R}$$

and

$$j''(R) = -\eta - \rho e^{\frac{\rho}{\sqrt{n}R}} < 0.$$

Thus, because j is strictly concave with respect to R, the critical value (truncated on the left by 0 and on the right by y) maximizes j. As R increases from 0 to y, j'(R) decreases from $\theta + \eta y \ge 0$ to

$$\sqrt{n}(1 - e^{\rho y/\sqrt{n}}) + \theta. \tag{3.9}$$

If the expression in (3.9) is negative, then the maximizing R lies in [0, y). If it is non-negative, then the maximizing R equals y. Let $\check{R}(\cdot; \rho)$ denote the maximizing retention function for a given $\rho > 0$; then,

$$\check{R}(y;\rho) = \begin{cases} y, & 0 \le y \le \frac{\sqrt{n}}{\rho} \ln\left(1 + \frac{\theta}{\sqrt{n}}\right), \\
\check{R}_c(y;\rho), & y > \frac{\sqrt{n}}{\rho} \ln\left(1 + \frac{\theta}{\sqrt{n}}\right), \end{cases}$$
(3.10)

in which $\check{R}_c(y;\rho) \in [0,y)$ for $y > \frac{\sqrt{n}}{\rho} \ln\left(1 + \frac{\theta}{\sqrt{n}}\right)$ uniquely solves

$$(\sqrt{n} + \theta) + \eta(y - R) = \sqrt{n} e^{\frac{\rho}{\sqrt{n}}R}.$$
(3.11)

As an aside, Liang, Liang, and Young [16] show that $\lim_{n\to\infty} \mathring{R}(y;\rho_n) = R_D(y)$.

Next, substitute $\check{R}(\cdot; \rho)$ from (3.10) into (3.4), or equivalently,

$$c - \lambda \mathbb{E}Y = \lambda \left(\theta \mathbb{E}(Y - R) + \frac{\eta}{2} \mathbb{E}((Y - R)^2) \right) + \frac{n\lambda}{\rho} \mathbb{E}\left(e^{\frac{\rho}{\sqrt{n}}R} - 1 - \frac{\rho R}{\sqrt{n}} \right),$$

and solve for ρ_n to obtain

$$c - \lambda \mathbb{E}Y = \sqrt{n\lambda} \int_0^\infty \left(\exp\left(\rho \,\frac{\check{R}(t;\rho)}{\sqrt{n}}\right) - 1 \right) S_Y(t) dt.$$
(3.12)

By substituting $t = \sqrt{ny}$ in (3.12), and by reversing the assignment in (2.32), we obtain $R_n^{\rho}(y) = \frac{1}{\sqrt{n}}\check{R}(\sqrt{ny};\rho_n)$ in (3.6) with ρ_n solving (3.5).

It remains to show that (3.5) has a unique positive solution ρ_n . To that end, consider the exponent in the integrand of (3.5), namely, $\rho R_n(y;\rho)$, in which we define $R_n(y;\rho)$ by replacing ρ_n in (3.6) with a generic $\rho > 0$. When $0 < y < \frac{1}{\rho} \ln(1 + \theta_n)$,

$$\frac{\partial}{\partial \rho} \left(\rho R_n^{\rho}(y) \right) = \frac{\partial}{\partial \rho} (\rho y) = y > 0.$$

When $y > \frac{1}{\rho} \ln(1 + \theta_n)$,

$$\frac{\partial}{\partial \rho} \left(\rho R_n^{\rho}(y) \right) = \frac{\partial}{\partial \rho} \left(\rho R_c(y) \right) = R_c(y) + \rho \frac{\partial R_c(y)}{\partial \rho}$$
$$= R_c(y) - \rho \frac{R_c(y) e^{\rho R_c(y)}}{\eta + \rho e^{\rho R_c(y)}} = \frac{\eta R_c(y)}{\eta + \rho e^{\rho R_c(y)}} > 0.$$

Thus, the right side of (3.5) increases with respect to ρ . As $\rho \to 0^+$, the right side of (3.5) approaches 0, which is less than the left side because we assume $c > \lambda \mathbb{E}Y$. As $\rho \to \infty$, $R_n(y; \rho)$ approaches 0 in such a way that

$$\lim_{\rho \to \infty} e^{\rho R_n(y;\rho)} = (1+\theta_n) + \eta y$$

Thus, the right side of (3.5) approaches

$$n\lambda \int_0^\infty \left(\theta_n + \eta y\right) S_{Y_n}(y) dy = \lambda \int_0^\infty \left(\theta + \eta t\right) S_Y(t) dt = \lambda \left(\theta \mathbb{E}Y + \frac{\eta}{2} \mathbb{E}(Y^2)\right),$$

which is greater than $c - \lambda \mathbb{E}Y$, from the assumption in (2.4). It follows that (3.5) has a unique positive solution ρ_n .

3.2 Diffusion approximation

The analog of (3.4) for the maximum adjustment coefficient for the diffusion approximation is

$$-\kappa + \lambda \sup_{R} \left[\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\rho + \eta}{2} \mathbb{E}(R^2) \right] = 0, \qquad (3.13)$$

in which we maximize over admissible retention functions R. In the following proposition, we show that the maximum adjustment coefficient for the diffusion approximation equals ρ_D from Theorem 2.3, with R_D the optimal retention function.

Proposition 3.2. The maximum adjustment coefficient $\rho_D > 0$ for the diffusion approximation in (2.35) uniquely solves (2.38), and the corresponding optimal retention function equals R_D in (2.39).

Proof. By following the argument in the proof of Proposition 3.1, for a given value of $\rho > 0$, the retention function maximizing the expression in (3.13) equals

$$R(y;\rho) = \frac{\theta + \eta y}{\rho + \eta} \wedge y.$$

By substituting this expression into (3.13), we obtain equation (2.38) for ρ_D . The remainder of the proof is similar to, but simpler than, the proof of Proposition 3.1, so we omit the details.

As we expect from the maximization problems in (3.4) and (3.13), the maximum adjustment coefficients ρ_n and ρ_D are related, which we prove in the next section.

3.3 Relationship between ρ_n and ρ_D

In this section, we prove two lemmas that relate ρ_n and ρ_D . We use those lemmas in Section 4.1 to modify ψ_D by function of order $\mathcal{O}(n^{-1/2})$ to bound ψ_n .

In the first lemma, we prove $\rho_n < \rho_D$.

Lemma 3.1. The maximum adjustment coefficient for the n-scaled risk process is less than the maximum adjustment coefficient for the diffusion approximation, that is, $\rho_n < \rho_D$.

Proof. For a given admissible retention function R, let $\rho_n(R)$ and $\rho_D(R)$ denote the solutions of (3.4) and (3.13), respectively, with the sup_R removed. We begin by showing that $\rho_n(R) < \rho_D(R)$. (As an aside, the left side of both (3.4) and (3.13) decrease from a positive number to negative infinity as ρ increases from 0 to infinity. Therefore, positive solutions $\rho_n(R)$ and $\rho_D(R)$ exist.)

 $\rho_D(R) > 0$ solves

$$-\kappa + \lambda \left[\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\rho + \eta}{2} \mathbb{E}(R^2) \right] = 0.$$
(3.14)

Because the left side of this equation decreases with respect to ρ , then $\rho_n(R) < \rho_D(R)$ if and only if

$$-\kappa + \lambda \left[\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\rho_n(R) + \eta}{2} \mathbb{E}(R^2) \right] > 0.$$

By using (3.1) and canceling a factor of $\lambda > 0$, this inequality becomes

$$n g_n(\rho_n(R); R) - \frac{\rho_n(R)}{2} \mathbb{E}(R^2) > 0,$$

or equivalently,

$$\frac{n}{\rho_n(R)} \mathbb{E}\left(e^{\frac{\rho_n(R)\cdot R}{\sqrt{n}}} - 1 - \frac{\rho_n(R)\cdot R}{\sqrt{n}} - \frac{\rho_n^2(R)\cdot R^2}{n}\right) > 0,$$

which is true because $e^x > 1 + x + x^2$ for x > 0.

Thus, we have shown $\rho_n(R) < \rho_D(R)$ for all admissible retention function R. Now, let $R = \check{R}$ from (3.10); then, we have

$$\rho_n = \rho_n(\check{R}) < \rho_D(\check{R}) \le \rho_D,$$

and we have proved this lemma.

In the following lemma, we show that we can modify ρ_D by a constant of order $\mathcal{O}(n^{-1/2})$ to get a lower bound of ρ_n .

Lemma 3.2. Choose C so that

$$C > \frac{\mathbb{E}(R_D^3)}{3 \mathbb{E}(R_D^2)} \rho_D^2. \tag{3.15}$$

Then, there exists N > 0 such that, for all $n \ge N$,

$$0 < \rho_D - \frac{C}{\sqrt{n}} < \rho_n < \rho_D, \tag{3.16}$$

from which it follows that

$$\lim_{n \to \infty} \rho_n = \rho_D, \tag{3.17}$$

with rate of convergence of order $\mathcal{O}(n^{-1/2})$.

Proof. The limit in (3.17) follows directly from the bounds in (3.16). The third inequality in (3.16) follows from Lemma 3.1; thus, it remains to show the first and second inequalities in (3.16). To that end, consider the admissible retention strategy R_D in (2.39). Then, from (3.1) and (3.2), we know $\rho_n(R_D)$ solves

$$-\kappa + \lambda \left[\theta \mathbb{E}R_D + \eta \mathbb{E}(YR_D) - \frac{\eta}{2} \mathbb{E}(R_D^2) - \frac{n}{\rho} \left(\mathbb{E}e^{\frac{\rho}{\sqrt{n}}R_D} - 1 - \frac{\rho}{\sqrt{n}} \mathbb{E}R_D \right) \right] = 0, \qquad (3.18)$$

or equivalently, from the expression for ρ_D in (3.13), $\rho_n(R_D)$ solves

$$\frac{\mathbb{E}(R_D^2)}{2}\rho_D - ng_n(\rho; R_D) = 0$$
(3.19)

The left side of this equation decreases with respect to ρ ; thus, to show that

$$\rho_D - \frac{C}{\sqrt{n}} < \rho_n(R_D)$$

for some value of C, it is enough to show that the left side of (3.19) is positive when we set $\rho = \rho_D - \frac{C}{\sqrt{n}}$. That is, we want to show there exist C and N such that

$$\frac{\mathbb{E}(R_D^2)}{2}\rho_D > ng_n(\rho_D - C/\sqrt{n}; R_D),$$

for all $n \ge N$, with $\rho_D - \frac{C}{\sqrt{n}} > 0$, or equivalently,

$$\frac{\mathbb{E}(R_D^2)}{2}\rho_D > \frac{n}{\rho_D - C/\sqrt{n}} \left(\mathbb{E}e^{\frac{\rho_D - C/\sqrt{n}}{\sqrt{n}}R_D} - 1 - \frac{\rho_D - C/\sqrt{n}}{\sqrt{n}}\mathbb{E}R_D \right).$$
(3.20)

To simplify the right side of inequality (3.20), we use the following identity:

$$e^{x} - 1 - x = \frac{x^{2}}{2} + \frac{x^{3}}{6} + \frac{x^{4}}{6} \int_{0}^{1} (1 - \omega)^{3} e^{\omega x} d\omega$$

Then, inequality (3.20) is equivalent to

$$\frac{\mathbb{E}(R_D^2)}{2} \rho_D \left(\rho_D - \frac{C}{\sqrt{n}} \right) \\
> \mathbb{E} \left(\frac{(\rho_D - C/\sqrt{n})^2 R_D^2}{2} + \frac{(\rho_D - C/\sqrt{n})^3 R_D^3}{6\sqrt{n}} + \frac{(\rho_D - C/\sqrt{n})^4 R_D^4}{6n} \int_0^1 (1-\omega)^3 e^{\omega \frac{\rho_D - C/\sqrt{n}}{\sqrt{n}} R_D} d\omega \right),$$

or

$$\mathbb{E}(R_D^2) \frac{C}{\sqrt{n}} > \mathbb{E}\bigg(\frac{(\rho_D - C/\sqrt{n})^2 R_D^3}{3\sqrt{n}} + \frac{(\rho_D - C/\sqrt{n})^3 R_D^4}{3n} \int_0^1 (1-\omega)^3 e^{\omega \frac{\rho_D - C/\sqrt{n}}{\sqrt{n}} R_D} d\omega\bigg),$$

which holds if the following stronger inequality holds:

$$\mathbb{E}(R_D^2) C > \mathbb{E}\left(\frac{\rho_D^2 R_D^3}{3} + \frac{\rho_D^3 R_D^4}{3\sqrt{n}} e^{\frac{\rho_D}{\sqrt{n}}R_D}\right),$$

or

$$\mathbb{E}(R_D^2) C - \rho_D^2 \frac{\mathbb{E}(R_D^3)}{3} > \frac{\rho_D^3}{3\sqrt{n}} \mathbb{E}\left(R_D^4 e^{\frac{\rho_D}{\sqrt{n}}R_D}\right).$$
(3.21)

If we choose C according to (3.15), then there exists N > 0 such that $\rho_D - C/\sqrt{N} > 0$ and such that inequality (3.21) holds at n = N. Because the right side of (3.21) decreases with n, it follows that inequality (3.21) holds for all $n \ge N$.

We have, thereby, shown that

$$0 < \rho_D - \frac{C}{\sqrt{n}} < \rho_n(R_D), \tag{3.22}$$

for all $n \ge N$, and we know that $\rho_n(R_D) \le \rho_n$ because ρ_n is maximal. Thus, we have proven the first two inequalities in (3.16).

4 Justifying the diffusion approximation

4.1 Bounds for ψ_n

In this section, we modify ψ_D by functions of order $\mathcal{O}(n^{-1/2})$ to obtain upper and lower bounds of ψ_n . In the process of finding an upper bound of ψ_n , we obtain a type of Lundberg bound for ψ_n .

To that end, by analogy with the expression for ψ_D in (2.36), define $\overline{\psi}_n$ for $x \leq m$ and $m \geq 0$ as follows:

$$\overline{\psi}_n(x,m) = \begin{cases} 1 - h_n(m) \left(1 - e^{-\rho_n(x - \alpha m)} \right), & (x,m) \in \mathcal{D}, \\ 1, & x < \alpha m, \end{cases}$$
(4.1)

in which h_n is defined by

$$h_n(m) = \left(1 - e^{-\rho_n(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}},\tag{4.2}$$

for all $m \ge 0$. In the following lemma, we use Theorem 2.2 to prove that $\overline{\psi}_n$ is an upper bound of ψ_n .

Lemma 4.1. For all $(x,m) \in \mathcal{D}$,

$$\psi_n(x,m) \le \overline{\psi}_n(x,m),\tag{4.3}$$

in which ψ_n is the minimum probability of drawdown for the n-scaled model, and $\overline{\psi}_n$ is defined in (4.1).

Proof. We prove this lemma via Theorem 2.2 modified to account for the *n*-scaled model, which essentially replaces $\mathcal{L}^{\hat{R}}$ in condition (iv) with $\mathcal{L}_{n}^{\hat{R}_{n}}$, given in (2.34). First, note that $\overline{\psi}_{n} \in \mathcal{C}^{1,1}(\mathcal{D})$ by its definition. Next, we go through each condition in Theorem 2.2 in turn.

Condition (i):

$$\lim_{m \to \infty} \overline{\psi}_n(m,m) = \lim_{m \to \infty} \left(1 - h_n(m) \left(1 - e^{-\rho_n(1-\alpha)m} \right) \right)$$
$$= 1 - \lim_{m \to \infty} \left(1 - e^{-\rho_n(1-\alpha)m} \right)^{\frac{\alpha}{1-\alpha}+1} = 1 - 1 = 0,$$

so condition (i) is satisfied with equality.

Condition (ii): This condition, namely, that $\overline{\psi}_n(x,m)$ is defined for all $x \leq m$ and $m \geq 0$, with $\overline{\psi}_n(x,m) = 1$ for all $x < \alpha m$, is satisfied by the definition of $\overline{\psi}_n$ in (4.1).

Condition (iii): For m > 0, differentiate $\overline{\psi}_n$ with respect to m and simplify the expression to obtain

$$(\overline{\psi}_n)_m(x,m) = \frac{\alpha \rho_n h_n(m)}{1 - e^{-\rho_n(1-\alpha)m}} \left(e^{-\rho_n(x-\alpha m)} - e^{-\rho_n(1-\alpha)m} \right),$$

which equals 0 when x = m. Therefore, condition (iii) is satisfied with equality.

Condition (iv): Let \hat{R}_n be the *n*-admissible retention function R_n^{ρ} that maximizes the adjustment coefficient. Then, by (3.3), R_n^{ρ} and ρ_n satisfy

$$\lambda_n \left(\mathbb{E}_n e^{\rho_n R_n^{\rho}} - 1 \right) = \left[-\kappa_n + \lambda_n \left((1 + \theta_n) \mathbb{E}_n R_n^{\rho} + \eta \mathbb{E}_n (Y_n R_n^{\rho}) - \frac{\eta}{2} \mathbb{E}_n ((R_n^{\rho})^2) \right) \right] \rho_n$$

Then, for $(x, m) \in \mathcal{D}$, the expression in (2.34) gives us

$$\begin{split} \mathcal{L}_{n}^{R_{n}^{\rho}}\overline{\psi}_{n}(x,m) &= \left[-\kappa_{n} + \lambda_{n} \left((1+\theta_{n})\mathbb{E}_{n}R_{n}^{\rho} + \eta\mathbb{E}_{n}(Y_{n}R_{n}^{\rho}) - \frac{\eta}{2} \mathbb{E}_{n}((R_{n}^{\rho})^{2}) \right) \right] (\overline{\psi}_{n})_{x}(x,m) \\ &+ \lambda_{n} \left(\mathbb{E}_{n}\overline{\psi}_{n}(x-R_{n}^{\rho},m) - \overline{\psi}_{n}(x,m) \right) \\ &= \left[-\kappa_{n} + \lambda_{n} \left((1+\theta_{n})\mathbb{E}_{n}R_{n}^{\rho} + \eta\mathbb{E}_{n}(Y_{n}R_{n}^{\rho}) - \frac{\eta}{2} \mathbb{E}_{n}((R_{n}^{\rho})^{2}) \right) \right] \left(-\rho_{n}h_{n}(m)e^{-\rho_{n}(x-\alpha m)} \right) \\ &+ \lambda_{n} \left[\mathbb{E}_{n} \left(\left(1-h_{n}(m) \left(1-e^{-\rho_{n}(x-R_{n}^{\rho}-\alpha m)} \right) \right) \mathbb{1}_{\{x-R_{n}^{\rho}\geq\alpha m\}} \right) \right] \\ &+ \lambda_{n} \left[\mathbb{E}_{n} (\mathbb{1}_{\{x-R_{n}^{\rho}<\alpha m\}}) - \left(1-h_{n}(m) \left(1-e^{-\rho_{n}(x-\alpha m)} \right) \right) \right] \\ &= -\lambda_{n}h_{n}(m)e^{-\rho_{n}(x-\alpha m)} \left(\mathbb{E}_{n}e^{\rho_{n}R_{n}^{\rho}} - 1 \right) + \lambda_{n}h_{n}(m) \left(1-\mathbb{E}_{n}(\mathbb{1}_{\{x-R_{n}^{\rho}\geq\alpha m\}}) \right) \\ &+ \lambda_{n}h_{n}(m)e^{-\rho_{n}(x-\alpha m)} \left(\mathbb{E}_{n}(e^{\rho_{n}R_{n}^{\rho}}\mathbb{1}_{\{x-R_{n}^{\rho}\geq\alpha m\}}) - 1 \right) \\ &= \lambda_{n}h_{n}(m) \left\{ -\mathbb{E}_{n}(e^{-\rho_{n}(\alpha - R_{n}^{\rho}-\alpha m)}\mathbb{1}_{\{x-R_{n}^{\rho}<\alpha m\}}) + \mathbb{E}_{n}(\mathbb{1}_{\{x-R_{n}^{\rho}<\alpha m\}}) \right\} \\ &= -\lambda_{n}h_{n}(m)\mathbb{E}_{n} \left(\left(e^{\rho_{n}(\alpha m - (x-R_{n}^{\rho}))} - 1 \right) \mathbb{1}_{\{x-R_{n}^{\rho}<\alpha m\}} \right) \\ &\leq 0. \end{split}$$

Thus, we have proved condition (iv). Theorem 2.2, then, implies inequality (4.3) on \mathcal{D} .

We obtain the following proposition from Lemmas 3.2 and 4.1, in which we modify ψ_D in (2.36) by a function of order $\mathcal{O}(n^{-1/2})$ to obtain an upper bound of ψ_n .

Proposition 4.1. Let C and N be as in the statement of Lemma 3.2, and for $n \in \mathbb{N}$, define u_n by

$$u_n(x,m) = \begin{cases} 1 - k_n(m) \left(1 - e^{-(\rho_D - C/\sqrt{n})(x - \alpha m)} \right), & (x,m) \in \mathcal{D}, \\ 1, & x < \alpha m, \end{cases}$$
(4.4)

in which k_n is defined by

$$k_n(m) = \left(1 - e^{-(\rho_D - C/\sqrt{n})(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}},$$
(4.5)

Then, for $n \geq N$,

$$\psi_n \le u_n,\tag{4.6}$$

on \mathcal{D} .

Proof. It is straightforward to show that $\overline{\psi}_n$ in (4.1) decreases with respect to $\rho_n > 0$ on \mathcal{D} . Thus, if we replace ρ_n in $\overline{\psi}_n$'s definition with a (positive) parameter less than ρ_n , then we get a function

that is an upper bound of $\overline{\psi}_n$. That is exactly how we defined u_n in (4.4) because, from Lemma 3.2, we know $0 < \rho_D - C/\sqrt{n} < \rho_n$ for all $n \ge N$; thus, we have, from Lemma 4.1,

$$\psi_n \le \overline{\psi}_n \le u_n$$

on \mathcal{D} .

In the following proposition, we modify ψ_D to obtain a lower bound of ψ_n .

Proposition 4.2. Formally, define the random variable $Z_d = (Y - d) | (Y > d)$ for $d \ge 0$, and suppose ς exists such that $M_Y(\rho_D/\sqrt{\varsigma}) < \infty$, with

$$\sup_{d\geq 0} \mathbb{E}\left(e^{\frac{\rho_D}{\sqrt{\varsigma}}Z_d}\right) < \infty.$$
(4.7)

Choose $\varepsilon > 0$, and define δ by

$$\delta = \sup_{d \ge 0} \left(\rho_D \mathbb{E} Z_d + \varepsilon \right), \tag{4.8}$$

and choose $N > \max(\delta^2, 4\varsigma)$ such that⁴

$$\sup_{d\geq 0} \frac{\rho_D^2}{\sqrt{N}} \mathbb{E}\left(Z_d^2 \, e^{\frac{\rho_D}{\sqrt{N}} Z_d}\right) \leq \varepsilon.$$
(4.9)

For $n \in \mathbb{N}$, define ℓ_n by

$$\ell_n(x,m) = \begin{cases} \left(1 - \frac{\delta}{\sqrt{n}}\right) \psi_D(x,m), & (x,m) \in \mathcal{D}, \\ 1, & x < \alpha m. \end{cases}$$
(4.10)

Then, for all $n \geq N$,

$$\ell_n \le \psi_n,\tag{4.11}$$

on \mathcal{D} .

⁴Condition (4.7) implies that we can find such an N. Indeed, for z large enough, we have

$$z^2 < \frac{e^{\frac{\rho_D z}{\sqrt{\varsigma}}}}{e^{\frac{\rho_D z}{2\sqrt{\varsigma}}}} = e^{\frac{\rho_D z}{2\sqrt{\varsigma}}},$$

which implies there exists M > 0 such that

$$\mathbb{E}\left(Z_d^2 e^{\frac{\rho_D}{2\sqrt{\varsigma}}Z_d}\right) \le \mathbb{E}\left(e^{\frac{\rho_D}{\sqrt{\varsigma}}Z_d}\mathbbm{1}_{\{Z_d > M\}}\right) + \mathbb{E}\left(M^2 e^{\frac{\rho_D}{\sqrt{\varsigma}}M}\mathbbm{1}_{\{Z_d \le M\}}\right)$$

Proof. We prove this lemma via Theorem 2.1 modified to account for the *n*-scaled model, which essentially replaces $\mathcal{L}^{\hat{R}}$ in condition (iv) with $\mathcal{L}_{n}^{\hat{R}_{n}}$, given in (2.34). First, note that $\ell_{n} \in \mathcal{C}^{1,1}(\mathcal{D})$ by its definition. Next, we go through each condition in Theorem 2.1 in turn.

Without loss of generality, assume $n > \delta^2$.

Condition (i):

$$\lim_{m \to \infty} \ell_n(m, m) = \left(1 - \frac{\delta}{\sqrt{n}}\right) \lim_{m \to \infty} \psi_D(m, m) = 0,$$

so condition (i) is satisfied with equality.

Condition (ii): This condition, namely, that $\ell_n(x,m)$ is defined for all $x \leq m$ and $m \geq 0$, with $\ell_n(x,m) = 1$ for all $x < \alpha m$, is satisfied by the definition of ℓ_n in (4.10).

Condition (iii): For m > 0, differentiate ℓ_n with respect to m and simplify the expression to obtain

$$(\ell_n)_m(x,m) = \left(1 - \frac{\delta}{\sqrt{n}}\right) \frac{\alpha \rho_D h_D(m)}{1 - e^{-\rho_D(1-\alpha)m}} \left(e^{-\rho_D(x-\alpha m)} - e^{-\rho_D(1-\alpha)m}\right),$$

which equals 0 when x = m. Therefore, condition (iii) is satisfied with equality.

Condition (iv): Let R_n be any *n*-admissible retention function. We wish to show that $\mathcal{L}_n^{R_n} \ell_n(x, m) \ge 0$ for all $(x, m) \in \mathcal{D}$.

$$\mathcal{L}_{n}^{R_{n}}\ell_{n}(x,m) = \left[-\kappa_{n} + \lambda_{n}\left((1+\theta_{n})\mathbb{E}_{n}R_{n} + \eta\mathbb{E}_{n}(Y_{n}R_{n}) - \frac{\eta}{2}\mathbb{E}_{n}(R_{n}^{2})\right)\right](\ell_{n})_{x}(x,m) + \lambda_{n}\left(\mathbb{E}_{n}\ell_{n}(x-R_{n},m) - \ell_{n}(x,m)\right).$$

$$(4.12)$$

We compute

$$(\ell_n)_x(x,m) = -\left(1 - \frac{\delta}{\sqrt{n}}\right)\rho_D h_D(m) e^{-\rho_D(x - \alpha m)},\tag{4.13}$$

and

$$\mathbb{E}_{n}\ell_{n}(x-R_{n},m) - \ell_{n}(x,m) = \mathbb{E}_{n}\left(\left(1-\frac{\delta}{\sqrt{n}}\right)\left(1-h_{D}(m)\left(1-e^{-\rho_{D}(x-R_{n}-\alpha m)}\right)\right)\mathbb{1}_{\{x-R_{n}\geq\alpha m\}} + \mathbb{1}_{\{x-R_{n}<\alpha m\}} - \left(1-\frac{\delta}{\sqrt{n}}\right)\left(1-h_{D}(m)\left(1-e^{-\rho_{D}(x-\alpha m)}\right)\right)\right).$$
(4.14)

Now, by substituting (4.13) and (4.14) into (4.12) and by rearranging terms, we obtain

$$\begin{split} \mathcal{L}_{n}^{R_{n}}\ell_{n}(x,m) \propto \kappa\rho_{D} - \lambda \left((\sqrt{n} + \theta)\mathbb{E}R + \eta\mathbb{E}(YR) - \frac{\eta}{2}\mathbb{E}(R^{2}) \right)\rho_{D} \\ &+ \frac{n\lambda e^{\rho_{D}(x-\alpha m)}}{h_{D}(m)} \mathbb{E}\left(\left(1 - h_{D}(m)\left(1 - e^{-\rho_{D}(x-R/\sqrt{n}-\alpha m)}\right) \right) \mathbb{1}_{\{x-R/\sqrt{n}\geq\alpha m\}} \right) \\ &+ \frac{n\lambda e^{\rho_{D}(x-\alpha m)}}{h_{D}(m)} \frac{\mathbb{E}\left(\mathbb{1}_{\{x-R/\sqrt{n}<\alpha m\}} \right)}{1 - \delta/\sqrt{n}} - \frac{n\lambda e^{\rho_{D}(x-\alpha m)}}{h_{D}(m)} \mathbb{E}\left(1 - h_{D}(m)\left(1 - e^{-\rho_{D}(x-\alpha m)}\right) \right), \end{split}$$

in which \propto denotes "positively proportional to," and in which R and R_n are related via (2.32). Recall that ρ_D solves

$$\kappa \rho_D = \lambda \sup_R \left\{ \left(\theta \mathbb{E}R + \eta \mathbb{E} (YR) - \frac{\eta}{2} \mathbb{E} (R^2) \right) \rho_D - \frac{1}{2} \mathbb{E} (R^2) \rho_D^2 \right\},\,$$

that is,

$$\kappa \rho_D - \lambda \left\{ \left(\theta \mathbb{E}R + \eta \mathbb{E}(YR) - \frac{\eta}{2} \mathbb{E}(R^2) \right) \rho_D - \frac{1}{2} \mathbb{E}(R^2) \rho_D^2 \right\} \ge 0,$$

for any admissible retention function R. Thus, to prove $\mathcal{L}_n^{R_n}\ell_n(x,m) \ge 0$ for all $(x,m) \in \mathcal{D}$ and for all *n*-admissible retention functions R_n , it is enough to prove

$$\lambda \left\{ \left(\theta \mathbb{E}R + \eta \mathbb{E} \left(YR \right) - \frac{\eta}{2} \mathbb{E} \left(R^2 \right) \right) \rho_D - \frac{1}{2} \mathbb{E} \left(R^2 \right) \rho_D^2 \right\}$$

$$\geq \lambda \left[\left(\left(\sqrt{n} + \theta \right) \mathbb{E}R + \eta \mathbb{E} \left(YR \right) - \frac{\eta}{2} \mathbb{E} \left(R^2 \right) \right) \right] \rho_D$$

$$- \frac{n \lambda e^{\rho_D (x - \alpha m)}}{h_D (m)} \mathbb{E} \left(\left(1 - h_D (m) \left(1 - e^{-\rho_D (x - R/\sqrt{n} - \alpha m)} \right) \right) \mathbb{1}_{\{x - R/\sqrt{n} \ge \alpha m\}} \right)$$

$$- \frac{n \lambda e^{\rho_D (x - \alpha m)}}{h_D (m)} \frac{\mathbb{E} \left(\mathbb{1}_{\{x - R/\sqrt{n} \le \alpha m\}} \right)}{1 - \delta/\sqrt{n}} + \frac{n \lambda e^{\rho_D (x - \alpha m)}}{h_D (m)} \mathbb{E} \left(1 - h_D (m) \left(1 - e^{-\rho_D (x - \alpha m)} \right) \right),$$

$$(4.15)$$

for all admissible retention functions R, or equivalently,

$$\frac{1}{2n} \mathbb{E}(R^2) \rho_D^2$$

$$\leq -1 - \frac{\rho_D \mathbb{E}R}{\sqrt{n}} + e^{\rho_D(x-\alpha m)} \left(\frac{\delta/\sqrt{n}}{h_D(m)(1-\delta/\sqrt{n})} + 1 \right) \mathbb{E}\left(\mathbb{1}_{\{x-R/\sqrt{n}<\alpha m\}}\right) + \mathbb{E}\left(e^{\frac{\rho_D R}{\sqrt{n}}}\mathbb{1}_{\{x-R/\sqrt{n}\geq\alpha m\}}\right)$$

$$= \int_0^\infty \left(e^{\frac{\rho_D R(y)}{\sqrt{n}}} - 1 - \frac{\rho_D R(y)}{\sqrt{n}}\right) dF_Y(y)$$

$$+ \int_0^\infty \left(e^{\rho_D(x-\alpha m)} \left(\frac{\delta/\sqrt{n}}{h_D(m)(1-\delta/\sqrt{n})} + 1\right) - e^{\frac{\rho_D R(y)}{\sqrt{n}}}\right) \mathbb{1}_{\{x-R/\sqrt{n}<\alpha m\}} dF_Y(y). \tag{4.16}$$

From $e^x > 1 - x - x^2/2$ for all x > 0, we deduce

$$\frac{1}{2n}\mathbb{E}(R^2)\rho_D^2 < \int_0^\infty \left(e^{\frac{\rho_D R(y)}{\sqrt{n}}} - 1 - \frac{\rho_D R(y)}{\sqrt{n}}\right) dF_Y(y).$$

Thus, to prove (4.16), it is enough to prove the stronger inequality

$$\int_{0}^{\infty} \left(e^{\frac{\rho_D}{\sqrt{n}} \left(R(y) - \sqrt{n}(x - \alpha m) \right)} - \frac{\delta/\sqrt{n}}{h_D(m)(1 - \delta/\sqrt{n})} - 1 \right) \mathbb{1}_{\{x - R(y)/\sqrt{n} < \alpha m\}} dF_Y(y) \le 0, \quad (4.17)$$

and we wish to find values of δ and $N > \delta^2$ for which inequality (4.17) holds for all n > N and x > 0. Note that the maximal retention function of the integrand in (4.17) is R(y) = y; thus, to prove (4.17), it is sufficient to prove

$$\int_0^\infty \left(e^{\frac{\rho_D}{\sqrt{n}} \left(y - \sqrt{n} (x - \alpha m) \right)} - \frac{\delta/\sqrt{n}}{h_D(m)(1 - \delta/\sqrt{n})} - 1 \right) \mathbb{1}_{\left\{ x - y/\sqrt{n} < \alpha m \right\}} dF_Y(y) \le 0,$$

that is,

$$\int_{\sqrt{n}(x-\alpha m)}^{\infty} \left(e^{\frac{\rho_D}{\sqrt{n}} \left(y - \sqrt{n}(x-\alpha m) \right)} - \frac{\delta/\sqrt{n}}{h_D(m)(1-\delta/\sqrt{n})} - 1 \right) dF_Y(y) \le 0.$$

Let d denote $\sqrt{n}(x - \alpha m)$. If $S_Y(d) = 0$, then the left side is identically 0, so suppose $S_Y(d) > 0$. After replacing $\sqrt{n}(x - \alpha m)$ by d and dividing by $S_Y(d)$, the above inequality becomes

$$\int_{d}^{\infty} \left(e^{\frac{\rho_D}{\sqrt{n}}(y-d)} - \frac{\delta/\sqrt{n}}{h_D(m)(1-\delta/\sqrt{n})} - 1 \right) \frac{dF_Y(y)}{S_Y(d)} \le 0,$$

for $d \ge 0$, or equivalently,

$$\int_{d}^{\infty} \left(e^{\frac{\rho_D}{\sqrt{n}}(y-d)} - 1 \right) \frac{dF_Y(y)}{S_Y(d)} \le \frac{\delta/\sqrt{n}}{h_D(m)(1-\delta/\sqrt{n})} \,. \tag{4.18}$$

Formally, define $Z_d = (Y - d) | (Y > d)$; then, inequality (4.18) becomes

$$\int_0^\infty \left(e^{\frac{\rho_D z}{\sqrt{n}}} - 1 \right) dF_{Z_d}(z) \le \frac{\delta/\sqrt{n}}{h_D(m)(1 - \delta/\sqrt{n})}.$$

Note that $0 < h_D(m) < 1$; thus, if we find δ to satisfy the following even stronger inequality, then the above sequence of inequalities holds:

$$\int_0^\infty \left(e^{\frac{\rho_D z}{\sqrt{n}}} - 1 \right) dF_{Z_d}(z) \le \frac{\delta}{\sqrt{n}}.$$
(4.19)

Rewrite the integrand from the left side of inequality (4.19) as follows:

$$e^{\frac{\rho_D z}{\sqrt{n}}} - 1 = \frac{\rho_D z}{\sqrt{n}} + \frac{\rho_D^2 z^2}{n} \int_0^1 (1-\omega) e^{\frac{\rho_D z}{\sqrt{n}}\omega} d\omega.$$

Thus, inequality (4.19) is equivalent to

$$\int_0^\infty \left(\frac{\rho_D z}{\sqrt{n}} + \frac{\rho_D^2 z^2}{n} \int_0^1 (1-\omega) e^{\frac{\rho_D z}{\sqrt{n}}\omega} d\omega\right) dF_{Z_d}(z) \le \frac{\delta}{\sqrt{n}},$$

or, after multiplying both side by \sqrt{n} and switching the order of integration,

$$\rho_D \mathbb{E} Z_d + \frac{\rho_D^2}{\sqrt{n}} \int_0^1 (1-\omega) \mathbb{E} \left(Z_d^2 e^{\frac{\rho_D \omega}{\sqrt{n}} Z_d} \right) d\omega \le \delta,$$

or more strongly,

$$\rho_D \mathbb{E} Z_d + \frac{\rho_D^2}{\sqrt{n}} \mathbb{E} \left(Z_d^2 \, e^{\frac{\rho_D}{\sqrt{n}} Z_d} \right) \le \delta. \tag{4.20}$$

Note that the left side of (4.20) decreases with increasing n. Define δ and N as in (4.8) and (4.9), respectively; then, inequality (4.20) holds for all $d \ge 0$ and all n > N, which implies that $\mathcal{L}_n^{R_n}\ell_n(x,m) \ge 0$ for all $(x,m) \in \mathcal{D}$ and all n > N. The conclusion in (4.11), then, follows from Theorem 2.1.

In the following two propositions, we show that the condition in (4.7) includes most of the usual claim distributions with light tails.

Proposition 4.3. Suppose Y has bounded support in \mathbb{R}_+ , then (4.7) holds.

Proof. Let b > 0 be such that $F_Y(b) = 1$. Let $a = \rho_D/\sqrt{\varsigma}$, and define the function G by

$$G(d) = \mathbb{E}(e^{aZ_d}),\tag{4.21}$$

which implies

$$G(d) = \frac{\int_{d}^{b} e^{a(y-d)} dF_Y(y)}{S_Y(d)} \mathbb{1}_{\{d < b\}}$$

Because $e^{a(y-d)} \le e^{a(b-d)}$ for all $0 \le y \le b$, we have

$$G(d) \le e^{a(b-d)},$$

so its supremum over \mathbb{R}_+ is finite.

Proposition 4.4. Suppose Y has a probability density function f_Y with unbounded support on \mathbb{R}_+ , and let h_Y denote the hazard rate function of Y, that is,

$$h_Y(y) = \frac{f_Y(y)}{S_Y(y)},$$
(4.22)

for $y \ge 0$. Furthermore, suppose

$$\lim_{y \to \infty} h_Y(y) = \ell. \tag{4.23}$$

If $\ell = 0$, then $M_Y(y) = \infty$ for all y > 0, which contradicts our assumption concerning Y's moment generating function. On the other hand, if $\ell > 0$, then (4.7) holds.

Proof. The first conclusion in the statement of this proposition follows from Theorem 2.5.1 in Rolski et al. [19], so suppose $\ell > 0$. Let ς be large enough so that $a = \rho_D / \sqrt{\varsigma} < \ell/2$ with $M_Y(a) < \infty$. As in the proof of Proposition 4.3, define the function G by (4.21). By applying L'Hôpital's rule, we obtain

$$\lim_{d \to \infty} G(d) = \lim_{d \to \infty} \frac{\int_d^\infty e^{ay} f_Y(y) dy}{e^{ad} S_Y(d)}$$
$$\stackrel{L'H}{=} \lim_{d \to \infty} \frac{-e^{ad} f_Y(d)}{a e^{ad} S_Y(d) - e^{ad} f_Y(d)}$$
$$= \lim_{d \to \infty} \frac{h_Y(d)}{-a + h_Y(d)} = \frac{\ell}{\ell - a} > 0.$$

Because G is continuous on \mathbb{R}_+ , it follows that $\sup_{d>0} G(d) < \infty$.

4.2 Convergence of ψ_n to ψ_D

In this section we prove an important result, namely, that as $n \to \infty$, ψ_n converges to ψ_D uniformly on \mathcal{D} , with rate of convergence of order $\mathcal{O}(n^{-1/2})$. In the following theorem, we combine the results of Propositions 4.1 and 4.2.

Theorem 4.1. If (4.7) holds, then there exist C' > 0 and N' > 0 such that, for all $n \ge N'$ and $(x,m) \in \mathcal{D}$,

$$\left|\psi_n(x,m) - \psi_D(x,m)\right| \le \frac{C'}{\sqrt{n}}.$$
(4.24)

Proof. From Propositions 4.1 and 4.2 it follows that

$$\left(1 - \frac{\delta}{\sqrt{n}}\right)\psi_D(x,m) < \psi_n(x,m) < u_n(x,m).$$

Subtracting $\psi_D(x, m)$ from each side yields

$$-\frac{\delta}{\sqrt{n}}\psi_D(x,m) < \psi_n(x,m) - \psi_D(x,m) < u_n(x,m) - \psi_D(x,m).$$
(4.25)

Clearly, the left side is bounded below by $-\delta/\sqrt{n}$. From (2.36), (2.37), (4.4), and (4.5), we deduce that the right side is positive and equals

$$\left(1 - e^{-\rho_D(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}} \left(1 - e^{-\rho_D(x-\alpha m)}\right) - \left(1 - e^{-(\rho_D - C/\sqrt{n})(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}} \left(1 - e^{-(\rho_D - C/\sqrt{n})(x-\alpha m)}\right)$$

$$= \left(1 - e^{-\rho_D(x-\alpha m)}\right) \left[\left(1 - e^{-\rho_D(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}} - \left(1 - e^{-(\rho_D - C/\sqrt{n})(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}} \right]$$

$$+ \left(1 - e^{-(\rho_D - C/\sqrt{n})(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}} \left[\left(1 - e^{-\rho_D(x-\alpha m)}\right) - \left(1 - e^{-(\rho_D - C/\sqrt{n})(x-\alpha m)}\right) \right].$$

$$(4.26)$$

First, analyze the next-to-the-last line in (4.26): for $n > (C/\rho_D)^2$, we have

$$\begin{aligned} 0 &\leq \left(1 - e^{-\rho_D(x - \alpha m)}\right) \left[\left(1 - e^{-\rho_D(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} - \left(1 - e^{-(\rho_D - C/\sqrt{n})(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} \right] \\ &\leq \left[\left(1 - e^{-\rho_D(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} - \left(1 - e^{-(\rho_D - C/\sqrt{n})(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} \right] \\ &= \left(1 - e^{-\rho_D(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} \left[1 - \left(\frac{1 - e^{-(\rho_D - C/\sqrt{n})(1 - \alpha)m}}{1 - e^{-\rho_D(1 - \alpha)m}}\right)^{\frac{\alpha}{1 - \alpha}} \right] \\ &\leq 1 - \left(\frac{1 - e^{-(\rho_D - C/\sqrt{n})(1 - \alpha)m}}{1 - e^{-\rho_D(1 - \alpha)m}}\right)^{\frac{\alpha}{1 - \alpha}}. \end{aligned}$$

It is straightforward to show that the last line of the above expression decreases with respect to m; therefore, by applying L'Hôpital's rule to take the limit as m goes to 0 of the expression in parentheses, we obtain

$$\begin{split} 0 &\leq \left(1 - e^{-\rho_D(x - \alpha m)}\right) \left[\left(1 - e^{-\rho_D(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} - \left(1 - e^{-(\rho_D - C/\sqrt{n})(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} \right] \\ &\leq 1 - \left(1 - \frac{C}{\rho_D\sqrt{n}}\right)^{\frac{\alpha}{1 - \alpha}}. \end{split}$$

If $b \ge 1$, then $1 - (1 - x)^b$ is concave on [0, 1], so lies below its tangent line at x = 0, which implies $1 - (1 - x)^b \le bx$ for all $0 \le x \le 1$. If 0 < b < 1, then $1 - (1 - x)^b$ is convex on [0, 1], so lies below its secant line between (0, 0) and (1, 1), which implies $1 - (1 - x)^b \le x$ for all $0 \le x \le 1$. Thus, by choosing any $M > (C/\rho_D)^2$, we conclude that $n \ge M$ implies

$$0 \leq \left(1 - e^{-\rho_D(x - \alpha m)}\right) \left[\left(1 - e^{-\rho_D(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} - \left(1 - e^{-(\rho_D - C/\sqrt{n})(1 - \alpha)m}\right)^{\frac{\alpha}{1 - \alpha}} \right]$$

$$\leq \max\left(\frac{\alpha}{1 - \alpha}, 1\right) \cdot \frac{C}{\rho_D\sqrt{n}}.$$
(4.27)

Next, analyze the last line of (4.26): for $n > (C/\rho_D)^2$, calculus shows that

$$\left(1 - e^{-\rho_D(x - \alpha m)}\right) - \left(1 - e^{-(\rho_D - C/\sqrt{n})(x - \alpha m)}\right) \le \left(1 - \frac{C}{\rho_D\sqrt{n}}\right)^{\frac{\rho_D\sqrt{n}}{C}} \left(\frac{\frac{C}{\sqrt{n}}}{\rho_D - \frac{C}{\sqrt{n}}}\right).$$
(4.28)

The first factor on the right side of (4.28) converges to e^{-1} ; thus, there exists $M' > (C/\rho_D)^2$ such that

$$\left(1 - \frac{C}{\rho_D \sqrt{n}}\right)^{\frac{\rho_D \sqrt{n}}{C}} \le 2e^{-1},$$

for all $n \ge M'$, which implies that the last line in (4.26) satisfies

$$0 \le \left(1 - e^{-(\rho_D - C/\sqrt{n})(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}} \left[\left(1 - e^{-\rho_D(x-\alpha m)}\right) - \left(1 - e^{-(\rho_D - C/\sqrt{n})(x-\alpha m)}\right) \right] \le 2e^{-1} \frac{C}{\rho_D - \frac{C}{\sqrt{M'}}} \cdot \frac{1}{\sqrt{n}},$$
(4.29)

for all $n \geq M'$.

By combining (4.27) and (4.29), by setting $N' = \max(N, M, M')$, and by setting

$$C' = \max\left\{\delta, \ \max\left(\frac{\alpha}{1-\alpha}, \ 1\right) \cdot \frac{C}{\rho_D} + 2e^{-1} \frac{C}{\rho_D - \frac{C}{\sqrt{M'}}}\right\},$$

we obtain (4.24).

4.3 $O(n^{-1/2})$ -optimality of retaining $R_{D,n}$ in the *n*-scaled model

We end this paper by showing that, if an insurer follows the optimal retention strategy for the diffusion model but surplus follows the *n*-scaled model, then the resulting probability of drawdown is $\mathcal{O}(n^{-1/2})$ -optimal. To that end, let $\psi_{D,n}$ denote the probability of drawdown when the insurer retains $R_{D,n}$ in (2.40) when surplus follows the *n*-scaled model. Then, we have the following theorem whose proof is similar to the proof of Theorem 2.2, so we omit it.

Theorem 4.2. Suppose $v \in C^{1,1}(\mathcal{D})$ is a bounded function that satisfies the following conditions:

- (i) $\lim_{m \to \infty} v(m,m) \ge 0.$
- (ii) v(x,m) is defined for all $x \le m$ and $m \ge 0$, with v(x,m) = 1 for all $x < \alpha m$.
- (iii) $v_m(m,m) \leq 0$ for all $m \geq 0$.
- (iv) $\mathcal{L}_n^{R_{D,n}}v(x,m) \leq 0$ for all $(x,m) \in \mathcal{D}$.

Then,
$$\psi_{D,n} \leq v \text{ on } \mathcal{D}$$
.

By analogy with the expressions for ψ_D and $\overline{\psi}_n$ in (2.36) and (4.1), respectively, define $\overline{\psi}_{D,n}$ for $x \leq m$ and $m \geq 0$ as follows:

$$\overline{\psi}_{D,n}(x,m) = \begin{cases} 1 - h_{D,n}(m) \left(1 - e^{-\rho_n(R_D)(x - \alpha m)} \right), & (x,m) \in \mathcal{D}, \\ 1, & x < \alpha m, \end{cases}$$
(4.30)

in which $h_{D,n}$ is defined by

$$h_{D,n}(m) = \left(1 - e^{-\rho_n(R_D)(1-\alpha)m}\right)^{\frac{\alpha}{1-\alpha}},$$
(4.31)

for all $m \ge 0$. In (4.30) and (4.31), $\rho_n(R_D) > 0$ uniquely solves (3.4) with the sup_R removed, as defined at the beginning of the proof of Lemma 3.1.

In the following lemma, we use Theorem 4.2 to prove that $\overline{\psi}_{D,n}$ is an upper bound of $\psi_{D,n}$.

Lemma 4.2. For all $(x, m) \in \mathcal{D}$,

$$\psi_{D,n}(x,m) \le \overline{\psi}_{D,n}(x,m). \tag{4.32}$$

Proof. We prove this lemma via Theorem 4.2. First, note that $\overline{\psi}_{D,n} \in \mathcal{C}^{1,1}(\mathcal{D})$ by its definition. Next, we go through each condition in Theorem 4.2 in turn.

Condition (i):

$$\lim_{m \to \infty} \overline{\psi}_{D,n}(m,m) = \lim_{m \to \infty} \left(1 - h_{D,n}(m) \left(1 - e^{-\rho_n(R_D)(1-\alpha)m} \right) \right)$$
$$= 1 - \lim_{m \to \infty} \left(1 - e^{-\rho_n(R_D)(1-\alpha)m} \right)^{\frac{\alpha}{1-\alpha}+1} = 1 - 1 = 0,$$

so condition (i) is satisfied with equality.

Condition (ii): This condition, namely, that $\overline{\psi}_{D,n}(x,m)$ is defined for all $x \leq m$ and $m \geq 0$, with $\overline{\psi}_{D,n}(x,m) = 1$ for all $x < \alpha m$, is satisfied by the definition of $\overline{\psi}_{D,n}$ in (4.30).

Condition (iii): For m > 0, differentiate $\overline{\psi}_{D,n}$ with respect to m and simplify the expression to obtain

$$(\overline{\psi}_{D,n})_m(x,m) = \frac{\alpha \rho_n(R_D) h_{D,n}(m)}{1 - e^{-\rho_n(R_D)(1-\alpha)m}} \left(e^{-\rho_n(R_D)(x-\alpha m)} - e^{-\rho_n(R_D)(1-\alpha)m} \right),$$

which equals 0 when x = m. Therefore, condition (iii) is satisfied with equality.

Condition (iv): From (3.18), we know $\rho_n(R_D)$ satisfies

$$-\kappa + \lambda \left[(\sqrt{n} + \theta) \mathbb{E}R_D + \eta \mathbb{E}(YR_D) - \frac{\eta}{2} \mathbb{E}(R_D^2) \right] = \frac{n\lambda}{\rho_n(R_D)} \left(\mathbb{E}e^{\rho_n(R_D)R_D/\sqrt{n}} - 1 \right).$$

Then, for $(x, m) \in \mathcal{D}$, the expression in (2.34) gives us

$$\begin{aligned} \mathcal{L}_{n}^{R_{D,n}}\overline{\psi}_{D,n}(x,m) &= \left[-\kappa_{n} + \lambda_{n} \left((1+\theta_{n})\mathbb{E}_{n}R_{D,n} + \eta\mathbb{E}_{n}(Y_{n}R_{D,n}) - \frac{\eta}{2}\mathbb{E}_{n}((R_{D,n})^{2}) \right) \right] (\overline{\psi}_{D,n})_{x}(x,m) \\ &+ \lambda_{n} \big(\mathbb{E}_{n}\overline{\psi}_{D,n}(x-R_{D,n},m) - \overline{\psi}_{D,n}(x,m)) \\ &= \left[-\kappa + \lambda \left((\sqrt{n}+\theta)\mathbb{E}R_{D} + \eta\mathbb{E}(YR_{D}) - \frac{\eta}{2}\mathbb{E}(R_{D}^{2}) \right) \right] (\overline{\psi}_{D,n})_{x}(x,m) \\ &+ n\lambda \big(\mathbb{E}\overline{\psi}_{D,n}(x-R_{D}/\sqrt{n},m) - \overline{\psi}_{D,n}(x,m)) \\ &= \frac{n\lambda}{\rho_{n}(R_{D})} \left(\mathbb{E}e^{\rho_{n}(R_{D})R_{D}/\sqrt{n}} - 1 \right) \left(-\rho_{n}(R_{D})h_{D,n}(m)e^{-\rho_{n}(R_{D})(x-\alpha m)} \right) \\ &+ n\lambda \left[\mathbb{E} \left(\left(1 - h_{D,n}(m) \left(1 - e^{-\rho_{n}(R_{D})(x-R_{D}/\sqrt{n}-\alpha m)} \right) \right) \mathbb{1}_{\{x-R_{D}/\sqrt{n} \ge \alpha m\}} \right) \right] \\ &+ n\lambda \left[\mathbb{E} (\mathbb{1}_{\{x-R_{D}/\sqrt{n} < \alpha m\}}) - \left(1 - h_{D,n}(m) \left(1 - e^{-\rho_{n}(R_{D})(x-\alpha m)} \right) \right) \right] \\ &= -n\lambda h_{D,n}(m)\mathbb{E} \left((e^{\rho_{n}(R_{D})(\alpha m - (x-R_{D}/\sqrt{n}))} - 1)\mathbb{1}_{\{x-R_{D}/\sqrt{n} < \alpha m\}} \right) \\ &\leq 0. \end{aligned}$$

Thus, we have proved condition (iv). Theorem 4.2, then, implies inequality (4.32) on \mathcal{D} .

We obtain the following proposition from Lemma 4.2, in which we show that u_n in (4.4) is an upper bound of $\psi_{D,n}$.

Proposition 4.5. Let C and N be as in the statement of Lemma 3.2; then, for $n \ge N$,

$$\psi_{D,n} \le u_n,\tag{4.33}$$

on \mathcal{D} , in which u_n is defined in (4.4).

Proof. It is straightforward to show that $\overline{\psi}_{D,n}$ in (4.30) decreases with respect to $\rho_n(R_D) > 0$ on \mathcal{D} . Thus, if we replace $\rho_n(R_D)$ in $\overline{\psi}_{D,n}$'s definition with a (positive) parameter less than $\rho_n(R_D)$, then we get a function that is an upper bound of $\overline{\psi}_{D,n}$. That is exactly how we defined u_n in (4.4) because, from inequality (3.22) in the proof of Lemma 3.2, we know $0 < \rho_D - C/\sqrt{n} < \rho_n(R_D)$ for all $n \geq N$; thus, we have, from Lemma 4.2,

$$\psi_{D,n} \le \overline{\psi}_{D,n} \le u_n,$$

on \mathcal{D} .

The following theorem is the main result of this paper, and it fully justifies using the optimal retention function for the diffusion approximation in the classical Cramér-Lundberg model when minimizing the probability of drawdown.

Theorem 4.3. Suppose (4.7) holds, and let C' > 0 and N' > 0 be as in Theorem 4.1. Then, for all $n \ge N'$ and $(x, m) \in \mathcal{D}$,

$$|\psi_n(x,m) - \psi_{D,n}(x,m)| \le \frac{2C'}{\sqrt{n}}.$$
 (4.34)

Proof. From the suboptimality of using $R_{D,n}$, we have $\psi_n \leq \psi_{D,n}$ on \mathcal{D} . From Proposition 4.5, we have $\psi_{D,n} \leq u_n$ for $n \geq N$. Thus, from the proof of Theorem 4.1, we deduce

$$-\frac{C'}{\sqrt{n}} \le \psi_n - \psi_D \le \psi_{D,n} - \psi_D \le u_n - \psi_D \le \frac{C'}{\sqrt{n}},$$

on \mathcal{D} for $n \geq N'$; recall $N' \geq N$. Thus,

$$\left|\psi_D - \psi_{D,n}\right| \le \frac{C'}{\sqrt{n}}$$

on \mathcal{D} for $n \geq N'$. This inequality, together with (4.24) and the triangle inequality, proves (4.34).

References

- Angoshtari, Bahman, Erhan Bayraktar, and Virginia R. Young (2016a). Optimal investment to minimize the probability of drawdown. *Stochastics*, 88(6): 946-958.
- [2] Angoshtari, Bahman, Erhan Bayraktar, and Virginia R. Young (2016b). Minimizing the probability of lifetime drawdown under constant consumption. *Insurance: Mathematics and Economics*, 69: 210-223.
- [3] Asmussen, Søren (1984). Approximation for the probability of ruin within finite time. Scandinavian Actuarial Journal, 1984(1): 31-57.
- [4] Azcue, Pablo and Nora Muler (2014). Stochastic Optimization in Insurance: A Dynamic Programming Approach. Springer. New York, New York, USA.
- [5] Bäuerle, Nicole (2004). Approximation of optimal reinsurance and dividend pay-out policies. Mathematical Finance, 14(1): 99-113.
- [6] Chen, Xinfu, David Landriault, Bin Li, and Dongchen Li (2015). On minimizing drawdown risks of lifetime investments. *Insurance: Mathematics and Economics*, 65: 46-54.
- [7] Cohen, Asaf and Virginia R. Young (2020). Rate of convergence of the probability of ruin in the Cramér-Lundberg model to its diffusion approximation. *Insurance: Mathematics and Economics*, 93: 333-340.

- [8] Cohen, Asaf and Virginia R. Young (2021). Optimal dividend problem: asymptotic analysis. SIAM Journal of Financial Mathematics, 12(1): 29-46.
- [9] Cvitanić, Jaksa and Ioannis Karatzas (1995). On portfolio optimization under "drawdown" constraints. IMA Lecture Notes in Mathematical Applications 65: 77-88.
- [10] Elie, Romuald, and Nizar Touzi. (2008). Optimal lifetime consumption and investment under a drawdown constraint. *Finance and Stochastics* 12(3), 299-330.
- [11] Grandell, Jan (1977). A class of approximations of ruin probabilities. Scandinavian Actuarial Journal, 1977(1): 37-52.
- [12] Grossman Sanford J. and Zhou Zhongquan. (1993). Optimal investment strategies for controlling drawdowns. *Mathematical Finance*, 3(3), 241-276.
- [13] Hald, Morten and Hanspeter Schmidli (2004). On the maximization of the adjustment coefficient under proportional reinsurance. ASTIN Bulletin, 34(1): 75-83.
- [14] Han, Xia, Zhibin Liang, and Virginia R. Young (2020). Optimal reinsurance to minimize the probability of drawdown under the mean-variance premium principle. *Scandinavian Actuarial Journal*, 2020(10): 879-903.
- [15] Iglehart, Donald L. (1969). Diffusion approximations in collective risk theory. Journal of Applied Probability, 6(2): 285-292.
- [16] Liang, Xiaoqing, Zhibin Liang, and Virginia R. Young (2020). Optimal reinsurance under the mean-variance premium principle to minimize the probability of ruin. *Insurance: Mathematics* and Economics, 92: 128-146.
- [17] Liang, Xiaoqing and Virginia R. Young (2021). Discounted probability of exponential Parisian ruin: diffusion approximation, to appear, *Journal of Applied Probability*.
- [18] Liang, Zhibin and Junyi Guo (2007). Optimal proportional reinsurance and ruin probability. Stochastic Models, 23(2): 333-350.
- [19] Rolski, Tomasz, Hanspeter Schmidli, Volker Schmidt, and Jozef L. Teugels (1999). Stochastic Process for Insurance and Finance, John Wiley & Sons. Chichester, England.
- [20] Schmidli, Hanspeter (2017). Risk Theory. Springer Actuarial Notes. Springer. Cham, Switzerland.