Three-color Ramsey number of an odd cycle versus bipartite graphs with small bandwidth *

Chunlin You and Qizhong Lin[†]

Abstract

A graph $\mathcal{H} = (W, E_{\mathcal{H}})$ is said to have *bandwidth* at most *b* if there exists a labeling of *W* as w_1, w_2, \ldots, w_n such that $|i - j| \leq b$ for every edge $w_i w_j \in E_{\mathcal{H}}$. We say that \mathcal{H} is a *balanced* (β, Δ) -graph if it is a bipartite graph with bandwidth at most $\beta |W|$ and maximum degree at most Δ , and it also has a proper 2-coloring $\chi : W \to [2]$ such that $||\chi^{-1}(1)| - |\chi^{-1}(2)|| \leq \beta |\chi^{-1}(2)|.$

In this paper, we prove that for every $\gamma > 0$ and every natural number Δ , there exists a constant $\beta > 0$ such that for every balanced (β, Δ) -graph \mathcal{H} on n vertices we have

$$R(\mathcal{H}, \mathcal{H}, C_n) \le (3+\gamma)n$$

for all sufficiently large odd n. The upper bound is sharp for several classes of graphs. Let $\theta_{n,t}$ be the graph consisting of t internally disjoint paths of length n all sharing the same endpoints. As a corollary, for each fixed $t \geq 1$, $R(\theta_{n,t}, \theta_{n,t}, C_{nt+\lambda}) = (3t+o(1))n$, where $\lambda = 0$ if nt is odd and $\lambda = 1$ if nt is even. In particular, we have $R(C_{2n}, C_{2n}, C_{2n+1}) = (6+o(1))n$, which is a special case of a result of Figaj and Luczak (2018).

Keywords: Ramsey number; Small bandwidth; Cycle; Regularity Lemma

Mathematics Subject Classification: 05C55; 05D10

1 Introduction

For graphs G and G_1, G_2, \ldots, G_k we write $G \to (G_1, G_2, \ldots, G_k)$ if for every k-edge-coloring of G, there is a monochromatic copy of G_i for some $i \in [k]$, where $[k] = \{1, 2, \ldots, k\}$. The multicolor Ramsey number $R(G_1, G_2, \ldots, G_k)$ is defined to be the smallest integer N such that $K_N \to (G_1, G_2, \ldots, G_k)$. We denote $R_k(G)$ instead of $R(G_1, G_2, \ldots, G_k)$ when $G_i = G$ for $1 \le i \le k$.

Let P_n (resp. C_n) be the path (resp. cycle) on n vertices. For the Ramsey numbers of $R(C_m, C_n)$, it has been studied and completely determined in Bondy and Erdős [7], Faudree and Schelp [19], and Rosta [42]. Later, Erdős, Faudree, Rousseau and Schelp [16] obtained the exact value of $R(C_\ell, C_m, C_n)$ for sufficiently large n. In 1999, by using the regularity lemma, Luczak [36] proved that $R(C_n, C_n, C_n) = (4 + o(1))n$ for all large odd integer n, and some years later Kohayakawa, Simonovits and Skokan [33] determined the exact value of it for all large odd n. In [36], Luczak introduced a technique that uses the regularity lemma to reduce problems

^{*}Center for Discrete Mathematics, Fuzhou University, Fuzhou, 350108, P. R. China. Email: chun-lin_you@163.com, linqizhong@fzu.edu.cn.

[†]Corresponding author. Supported in part by NSFC (No. 12171088).

about paths and cycles to problems about connected matchings, which are matchings that are contained in a connected component. This technique has become fairly standard. The following papers were based on this technique. For large even cycle, Figaj and Łuczak [21] obtained that $R(C_n, C_n, C_n) = (2 + o(1))n$, and Benevides and Skokan [5] obtained the exact value. Furthermore, Figaj and Łuczak [21, 22] determined the asymptotic values of $R(C_{\ell}, C_m, C_n)$ when ℓ, m , and n are large integers with same order. In particular, Figaj and Łuczak [22] determined

$$R(C_{2n}, C_{2n}, C_{2n+1}) = (6 + o(1))n.$$

Furthermore, Ferguson [23, 24, 25] obtained exact Ramsey numbers for the three-color cases of mixed parity cycles. Recently, Jenssen and Skokan [31] established that

$$R_k(C_n) = 2^{k-1}(n-1) + 1$$

for each fixed $k \ge 2$ and large odd n, which confirms a conjecture of Bondy and Erdős [7]. For more multicolor Ramsey numbers involving large cycles, we refer the reader to [1, 16, 37, 43], etc.

The Ramsey numbers of paths are also fruitful. A well-known result of Gerencsér and Gyárfás [27] states that $R(P_m, P_n) = n + \lfloor \frac{m}{2} \rfloor$ for all $n \ge m \ge 1$. Faudree and Schelp [20] determined $R(P_{n_0}, P_{n_1}, P_{n_2})$ for $n_0 \ge 6(n_1 + n_2)^2$, and they also conjectured that

$$R(P_n, P_n, P_n) = \begin{cases} 2n - 1, & n \text{ is odd,} \\ 2n - 2, & n \text{ is even.} \end{cases}$$
(1)

Gyárfás, Ruszinkó, Sárközy and Szemerédi [29] confirmed this conjecture for large n. For Ramsey numbers of cycles versus paths, Faudree, Lawrence, Parsons and Schelp [18] obtained the Ramsey numbers for all path-cycle pairs. In 2009, Dzido and Fidytek [15] (independently by Bielak [6]) obtained that if $m \ge 3$ is odd, $n \ge m$ and $n > \frac{3t^2 - 14t + 25}{4}$ when t is odd, and $n > \frac{3t^2 - 10t + 20}{8}$ when t is even, then $R(P_t, P_n, C_m) = 2n + 2 \lfloor \frac{t}{2} \rfloor - 3$. In [45], Shao, Xu, Shi and Pan proved that $R(P_4, P_5, C_n) = n + 2$ for $n \ge 23$ and $R(P_4, P_6, C_n) = n + 3$ for $n \ge 18$. Omidi and Raeisi [40] determined $R(P_4, P_n, C_4) = R(P_5, P_n, C_4) = n + 2$ for $n \ge 5$.

We say that a graph $\mathcal{H} = (W, E_{\mathcal{H}})$ has bandwidth at most *b* if there is a labeling of *W* as w_1, w_2, \ldots, w_n such that $|i - j| \leq b$ for every edge $w_i w_j \in E_{\mathcal{H}}$ (see, e.g., [10]). We say $\mathcal{H} = (W, E_{\mathcal{H}})$ is a balanced (β, Δ) -graph if it is a bipartite graph with bandwidth at most $\beta |W|$ and maximum degree at most Δ , and furthermore it has a proper 2-coloring $\chi : W \to [2]$ such that $||\chi^{-1}(1)| - |\chi^{-1}(2)|| \leq \beta |\chi^{-1}(2)|$. In [38], Mota, Sárközy, Schacht and Taraz determined asymptotically the three color Ramsey number $R(\mathcal{H}, \mathcal{H}, \mathcal{H}) \leq (2 + o(1))n$, which implies that

$$R(C_n, C_n, C_n) = (2 + o(1))n$$

for all large even n.

In this paper, we are concerned with the asymptotic behavior of the Ramsey number $R(\mathcal{H}, \mathcal{H}, C_n)$ when n is odd and every balanced (β, Δ) -graph \mathcal{H} on n vertices.

Theorem 1.1 For any sufficiently small γ and every natural number Δ , there exists a constant $\beta > 0$ such that for every balanced (β, Δ) -graph \mathcal{H} on n vertices we have

$$R(\mathcal{H}, \mathcal{H}, C_n) \le (3+\gamma)n$$

for all sufficiently large odd n.

Let $\theta_{n,t}$ be the graph consisting of t internally disjoint paths of length n all sharing the same endpoints. In particular, $\theta_{n,2} = C_{2n}$, i.e. the even cycle on 2n vertices. Observe that $\theta_{n,t}$ is a bipartite graph with (n-1)t+2 vertices, and $\theta_{n,t}$ is a balanced (β, t) -graph with $\beta \leq \frac{2}{n}$.

The following result is clear from Theorem 1.1.

Corollary 1.1 For each fixed $t \ge 1$,

$$R(\theta_{n,t}, \theta_{n,t}, C_{nt+\lambda}) = (3t + o(1))n,$$

where $\lambda = 0$ if nt is odd and $\lambda = 1$ if nt is even. In particular, we have that $R(P_{2n}, P_{2n}, C_{2n+1}) \sim R(C_{2n}, C_{2n}, C_{2n+1}) = (6 + o(1))n$.

Let us point out that the result $R(C_{2n}, C_{2n}, C_{2n+1}) = (6 + o(1))n$ and $R(P_{2n}, P_{2n}, C_{2n+1}) = (6 + o(1))n$ follow from the work of Figaj and Luczak [22] and have been strengthened by Ferguson [23].

2 Preliminaries

In this paper, we shall omit all floors and ceilings since it will not affect our argument. For a graph G, let V(G) and E(G) denote its vertex set and edge set, respectively. Let |E(G)| = e(G) and |V(G)| = v(G). For $v \in V$, we let $N_G(v)$ denote the neighborhood of v in G, and $\deg_G(v) = |N_G(v)|$ which is the degree of a vertex $v \in V(G)$. We denote by $\delta(G)$ and $\Delta(G)$ the minimum and maximum degrees of the vertices of G. For any subset $U \subseteq V$, we use G[U] to denote the subgraph induced by the vertex set U in G. For a vertex $v \in V$ and $U \subset V$, we write $N_G(v, U)$ for the neighbors of v in U in graph G and $\deg(v, U) = |N_G(v, U)|$.

2.1 The regularity method

For disjoint vertex sets $A, B \subseteq V$, let $e_G(A, B)$ denote the number of edges of G with one endpoint in A and the other in B, and the density between A and B is

$$d_G(A,B) = \frac{e_G(A,B)}{|A||B|}.$$

We always omit the subscript when there is no confusion.

Definition 2.1 (ϵ -regular) A pair (A, B) is ϵ -regular if for all $X \subseteq A$ and $Y \subseteq B$ with $|X| > \epsilon |A|$ and $|Y| > \epsilon |B|$ we have $|d(X, Y) - d(A, B)| < \epsilon$.

Definition 2.2 ((ϵ , d)-regular) For $\epsilon > 0, d \leq 1$, a pair (A, B) is said to be (ϵ , d)-regular if it is ϵ -regular and $d(A, B) \geq d$.

Definition 2.3 ((ϵ , d)-super-regular) A pair (A, B) is said to be (ϵ , d)-super-regular if it is ϵ -regular and deg(u, B) > d|B| for all $u \in A$ and deg(v, A) > d|A| for all $v \in B$.

We have a property that any regular pair has a large subgraph which is super-regular, and we include a proof for completeness.

Fact 2.1 For $0 < \epsilon < 1/2$ and $d \le 1$, if (A, B) is (ϵ, d) -regular with |A| = |B| = m, then there exist $A_1 \subseteq A$ and $B_1 \subseteq B$ with $|A_1| = |B_1| \ge (1 - \epsilon)m$ such that (A_1, B_1) is $(2\epsilon, d - 2\epsilon)$ -super-regular.

Proof. Let $X \subseteq A$ consist of vertices with at most $(d - \epsilon)|B|$ neighbors in B. Accounting for $e(X,B) \leq |X| \cdot (d - \epsilon)|B|$, we have $|d(X,B) - d| \geq \epsilon$. Due to the definition of ϵ -regular, it follows that $|X| \leq \epsilon m$. Similarly, assume that $Y \subseteq B$ consists of vertices with at most $(d - \epsilon)|A|$ neighbors in A, then we obtain $|Y| \leq \epsilon m$. Take $A_1 \subseteq A \setminus X$ and $B_1 \subseteq B \setminus Y$ with $|A_1| = |B_1| = (1 - \epsilon)m$. Thus for any vertex $u \in A_1$, $\deg(u, B_1) \geq (d - 2\epsilon)m$. Also, $\deg(w, A_1) \geq (d - 2\epsilon)m$ for any $w \in B_1$.

Now, for any subsets $S \subseteq A_1$ and $T \subseteq B_1$, if $|S| > 2\epsilon |A_1|$ and $|T| > 2\epsilon |B_1|$, then clearly $|S| > \epsilon m$ and $|T| > \epsilon m$. Since (A, B) is (ϵ, d) -regular, it follows that

$$|d(S,T) - d(A_1, B_1)| \le |d(S,T) - d(A,B)| + |d(A_1, B_1) - d(A,B)| < 2\epsilon.$$

We have thus proved the fact.

In this paper, we will use the following three-color version of the regularity lemma. For many applications, we refer the reader to surveys [32, 41] and many recent references [2, 11, 12, 13, 14, 26, 35, 39, 47], etc.

Lemma 2.1 (Szemerédi [46]) For every $\epsilon > 0$ and integer $t_0 \ge 1$, there exists $T_0 = T_0(\epsilon, t_0) \ge t_0$ such that the following holds. For all graphs G_1, G_2 and G_3 with the same vertex set V and $|V| \ge t_0$, there exists a partition $V = \bigcup_{i=0}^t V_i$ satisfying $t_0 \le t \le T_0$ and

- (1) $|V_0| < \epsilon n, |V_1| = |V_2| = \ldots = |V_t|;$
- (2) all but at most $\epsilon\binom{t}{2}$ pairs (V_i, V_j) , $1 \leq i \neq j \leq t$, are ϵ -regular for G_1, G_2 and G_3 .

The next lemma by Benevides and Skokan [5] is a slightly stronger version compared to the original one established by Luczak [36, Claim 3].

Lemma 2.2 (Benevides and Skokan [5]) For every $0 < \beta_0 < 1$, there exists an n_0 such that for every $n > n_0$ the following holds: If (V_1, V_2) is ϵ -regular with $|V_1| = |V_2| = n$ and density at least $\beta_0/4$ for some ϵ satisfying $0 < \epsilon < \beta_0/100$, then for every $\ell, 1 \le \ell \le n - 5\epsilon n/\beta_0$, and for every pair of vertices $v' \in V_1$, $v'' \in V_2$ satisfying $\deg(v', V_2)$, $\deg(v'', V_1) \ge \beta_0 n/5$, G contains a path of length $2\ell + 1$ connecting v' and v''.

Let $\mathcal{H} = (W, E_{\mathcal{H}})$ be a graph, for $S \subseteq W$, denote $N_{\mathcal{H}}(S) = [\bigcup_{v \in S} N_{\mathcal{H}}(v)] \setminus S$. For a graph G = (V, E), a partition $\bigcup_{i=1}^{k} V_i$ of V is said to be (ϵ, d) -regular on a reduced graph H with vertex set contained in [k] if the pair (V_i, V_i) is (ϵ, d) -regular whenever $ij \in E(H)$.

When applying the regularity lemma, we will indeed find a partition of a monochromatic subgraph G of K_N with corresponding reduced graph containing a tree T that contains a "large" matching M, where the bipartite subgraphs of G corresponding to the matching are super-regular pairs. The following definition of ϵ -compatible is due to Mota, Sárközy, Schacht and Taraz [38]. **Definition 2.4** (ϵ -compatible) Let $\mathcal{H} = (W, E_{\mathcal{H}})$ and $T = ([t], E_T)$ be graphs. Let $M = ([t], E_M)$ be a subgraph of T where E_M is a matching. Given a partition $W = \bigcup_{i=1}^t W_i$, let U_i , for $i \in [t]$, be the set of vertices in W_i , with neighbors in some W_j with $ij \in E_T \setminus E_M$. Set $U = \bigcup U_i$ and $U'_i = N_{\mathcal{H}}(U) \cap (W_i \setminus U)$.

We say that $W = \bigcup_{i=1}^{t} W_i$ is (ϵ, T, M) -compatible with a vertex partition $\bigcup_{i=1}^{t} V_i$ of a graph G = (V, E) if the following holds.

- (1) $|W_i| \le |V_i|$ for $i \in [t]$.
- (2) $xy \in E_{\mathcal{H}}$ for $x \in W_i, y \in W_j$ implies $ij \in E_T$ for all distinct $i, j \in [t]$.
- (3) $|U_i| \le \epsilon |V_i|$ for $i \in [t]$.

 $(4) |U'_i|, |U'_j| \le \epsilon \min\{|V_i|, |V_j| : ij \in E_M\}.$

The following corollary of the Blow-up Lemma (see Böttcher, Heinig and Taraz [8, 9]) asserts that in the setup of Definition 2.4 graphs \mathcal{H} of bounded degree can be embedded into G, if Gadmits a partition being sufficiently regular on T and super-regular on M.

Lemma 2.3 (Embedding Lemma [8, 9]) For all $d, \Delta > 0$ there is a constant $\epsilon = \epsilon(d, \Delta) > 0$ such that the following holds. Let G = (V, E) be an N-vertex graph that has a partition $\cup_{i=1}^{t} V_i$ with (ϵ, d) -reduced graph T on [t] which is (ϵ, d) -super-regular on a graph $M \subset T$. Further, let $\mathcal{H} = (W, E_{\mathcal{H}})$ be an n-vertex graph with maximum degree $\Delta(\mathcal{H}) \leq \Delta$ and $n \leq N$ that has a vertex partition $\cup_{i=1}^{t} W_i$ of W which is (ϵ, T, M) -compatible with $\cup_{i=1}^{t} V_i$. Then $\mathcal{H} \subseteq G$.

For a graph $\mathcal{H} = (W, E_{\mathcal{H}})$ with $W = \{w_1, w_2, \ldots, w_n\}$, where w_i is a labeling of the vertices, let $\chi : W \to [2]$ be a 2-coloring. For $W' \subseteq W$, denote $C_i(W') = |\chi^{-1}(i) \cap W'|$ for i = 1, 2. We know that χ is a β -balanced coloring of W if $1 - \beta \leq \frac{C_1(W)}{C_2(W)} \leq 1 + \beta$. A set $I \subseteq W$ is called an interval if there exists p < q such that $I = \{w_p, w_{p+1}, \ldots, w_q\}$. Finally, let $\sigma : [\ell] \to [\ell]$ be a permutation, and for a partition $\{I_1, I_2, \ldots, I_\ell\}$ of W, where I_1, \ldots, I_ℓ are intervals, let $C_i(\sigma, a, b) = \sum_{j=a}^b C_i(I_{\sigma(j)})$ for i = 1, 2.

Lemma 2.4 (Mota et al. [38]) For every $\xi > 0$ and every integer $\ell \ge 1$ there exists n_0 such that if $\mathcal{H} = (W, E_{\mathcal{H}})$ is a graph on $W = \{w_1, w_2, \ldots, w_n\}$ with $n \ge n_0$, then for every β -balanced 2-coloring χ of W with $\beta \le 2/\ell$, and every partition of W into intervals I_1, I_2, \ldots, I_ℓ with $|I_1| \le |I_2| \le \cdots \le |I_\ell| \le |I_1| + 1$ there exists a permutation $\sigma : [\ell] \to [\ell]$ such that for every pair of integers $1 \le a < b \le \ell$ with $b - a \ge 7/\xi$, we have $|C_1(\sigma, a, b) - C_2(\sigma, a, b)| \le \xi C_2(\sigma, a, b)$.

2.2 Structure

Recall that a set M of independent edges in a graph G = (V, E) is called a matching. The size of a connected matching corresponds to the number of edges in the matching. The structure of graphs without large connected matchings play an important role in this paper. As a path on n vertices contains a connected matching on $\lfloor n/2 \rfloor$ edges, extremal results for paths directly give an upper bound for connected matchings.

The following is the well-known extremal result for paths.

Lemma 2.5 (Erdős and Gallai [17]) If G is a graph of order N which contains no P_n , then

$$e(G) \le \frac{n-2}{2}N.$$

The following result describes the structure of a graph without a large matching.

Lemma 2.6 (Knierim and Su [34]) For every connected graph G = (V, E) which contains no matching of size n/2, there is a partition $S_G \cup Q_G \cup I_G$ of the vertex set V such that

- (i) $|Q_G| + 2|S_G| = \min\{v(G), n-1\},\$
- (ii) I_G is an independent set; additionally, if $v(G) \leq n-1$, then $I_G = \emptyset$,
- (iii) every vertex in Q_G has at most one neighbor in I_G ,
- (iv) every vertex in I_G has degree less than n/2.

We will also use the following lemmas. Let us begin with a result due to Łuczak [36] which gives a description of the structure of a graph that contains no large odd cycle as a subgraph.

Lemma 2.7 (Luczak [36]) For every $0 < \delta < 10^{-15}$, $\alpha \ge 2\delta$ and $t \ge exp(\delta^{-16}/\alpha)$ the following holds. Each graph H on t vertices which contains no odd cycles longer than αt contains subgraphs H' and H" such that:

(i) $V(H') \cup V(H'') = V(H)$, $V(H') \cap V(H'') = \emptyset$ and each of the sets V(H') and V(H'') is either empty or contains at least $\alpha \delta t/2$ vertices;

- (ii) H' is bipartite;
- (iii) H'' contains no more than $\alpha t |V(H'')|/2$ edges;
- (iv) all except no more than δt^2 edges of H belong to either H' or H".

The following result due to Gyárfás, Ruszinkó, Sárközy and Szemerédi [29] states that a graph with high density always contains a dense subgraph.

Fact 2.2 (Gyárfás et al. [29]) Let $\epsilon > 0$ be sufficiently small and let H be a graph with v(H) vertices. If $e(H) \ge {\binom{v(H)}{2}} - \epsilon {\binom{t}{2}}$, then H has a subgraph H' with at least $v(H) - \sqrt{\epsilon}t$ vertices and $\delta(H') \ge v(H) - 2\sqrt{\epsilon}t$.

A spanning subgraph of a graph G is a subgraph obtained by edge deletions only, in other words, a subgraph whose vertex set is the entire vertex set of G. We also need the following simple result.

Lemma 2.8 For any $\epsilon > 0$, if H' is a spanning subgraph of H with $e(H') \ge e(H) - \epsilon {t \choose 2}$, then there exists a induced subgraph $H'' \subset H$ with at least $v(H) - \sqrt{\epsilon t}$ vertices and $\deg_{H''}(u) < \deg_{H'}(u) + \sqrt{\epsilon t}$ for any vertex $u \in V(H'')$.

Proof. Let $X = \{u \in V(H) | \deg_H(u) - \deg_{H'}(u) \ge \sqrt{\epsilon}t\}$. Clearly, $e(H) - e(H') \ge \frac{\sqrt{\epsilon}t|X|}{2}$. Thus we have $\frac{\sqrt{\epsilon}t|X|}{2} \le \epsilon \binom{t}{2},$

implying that $|X| < \sqrt{\epsilon t}$. Denote H'' = H - X, i.e. the subgraph obtained from H by deleting all vertices of X and all edges incident to some vertices of X. Thus for any $u \in V(H'')$,

$$\deg_{H''}(u) = \deg_H(u) - \deg_H(u, X) < \left(\deg_{H'}(u) + \sqrt{\epsilon t}\right) - \deg_H(u, X) \le \deg_{H'}(u) + \sqrt{\epsilon t},$$

completing the proof.

2.3 Monochromatic components

In [44], Schelp conjectured that if G is a graph on 3n - 1 vertices with minimum degree at least 3|V(G)|/4, then $G \to (P_{2n}, P_{2n})$ provided n sufficiently large. Gyárfás and Sárközy [30] and independently Benevides, Luczak, Scott, Skokan and White [4] confirmed this conjecture asymptotically. Recently, Balogh, Kostochka, Lavrov and Liu [3] obtain the following result and thus confirm this conjecture thoroughly.

Lemma 2.9 (Balogh et al. [3]) Let G be a graph on 3n - 1 vertices with minimum degree at least (3|V(G)| - 1)/4. If n is sufficiently large, then $G \to (P_{2n}, P_{2n})$.

Erdős and Rado remarked that any 2-colored complete graph contains a monochromatic spanning tree, see [28]. We will apply the following result due to Gyárfás and Sárközy [30] to get a large monochromatic component for every 2-coloring of the edges of graph G with large minimum degree.

Lemma 2.10 (Gyárfás and Sárközy [30]) For any 2-coloring of edges of a graph G with minimum degree $\delta(G) \geq \frac{3|V(G)|}{4}$, there is a monochromatic component of order larger than $\delta(G)$. This estimate is sharp.

3 Proof of Theorem 1.1

Let $N = (3 + \gamma)n$, where $\gamma > 0$ is a sufficiently small real number and n is a sufficiently large odd integer. Consider a 3-edge coloring of K_N on vertex set V, and let G_i (i = 1, 2, 3) be the graph induced by the edges in the *i*th color. We shall show that either G_i contains a copy of \mathcal{H} for some i = 1, 2, or G_3 contains an odd cycle C_n . On the contrary, we suppose that G_i contains no \mathcal{H} for i = 1, 2, and G_3 contains no odd cycle C_n . We aim to find a contradiction.

We write $a \ll b$ if a is much smaller than b. Let $\gamma > 0$ and $\Delta \ge 1$ be given. We apply Lemma 2.3 to G_1 (or G_2) with d = 1/4 and Δ to get ϵ_1 . We set η, ϵ, δ and β such that

$$\eta = \frac{\gamma}{15}, \ \delta = \min\left\{10^{-16}, \ \frac{\eta^2}{100}\right\} \ \epsilon = \min\left\{\frac{\delta}{1296}, \epsilon_1\right\} \ \text{and} \ 0 < \beta \ll \epsilon.$$
(2)

Moreover, set

$$t_0 = \max\left\{\frac{1}{\epsilon}, \ \exp\left(\frac{3(1+\eta)}{\delta^{16}}\right)\right\}.$$
(3)

We apply the regularity lemma (Lemma 2.1) to G_1 , G_2 and G_3 with ϵ, t_0 to obtain a $T_0 = T_0(\epsilon, t_0)$ such that there exists a partition of the vertex set V into t+1 classes $V = V_0 \cup V_1 \cup \cdots \cup V_t$ satisfying $t_0 \leq t \leq T_0$ and (1) $|V_0| < \epsilon n$, $|V_1| = |V_2| = \ldots = |V_t|$; (2) all but at most $\epsilon {t \choose 2}$ pairs $(V_i, V_j), 1 \leq i \neq j \leq t$, are ϵ -regular for G_1, G_2 and G_3 . We construct the reduced graph H with vertex set $\{v_1, v_2, \ldots, v_t\}$ and the edge set formed by pairs $\{v_i, v_j\}$ for which (V_i, V_j) is ϵ -regular with respect to G_1, G_2 and G_3 . In which v_i and v_j are non-adjacent in H if the pairs (V_i, V_j) is not ϵ -regular for some G_i . Thus we obtain a bijection $f: v_i \to V_i$ between the vertices of H and the clusters of the partition.

Clearly, $e(H) \ge (1-\epsilon) {t \choose 2}$ from Lemma 2.1. Accordingly, we assign color $i \ (i \in [3])$ to an edge of H if and only if i is the minimum integer for which

$$d_{G_i}(V_i, V_j) \ge 1/3.$$

Let $H_i(F_i)$ be the spanning subgraph of H(F) induced by all edges that have received color *i*.

Overview of the remaining proof: The remaining part of the proof is straightforward but rather rich in technical details, so we shall briefly outline it first. From the assumption that G_3 contains no odd cycle C_n , we will show that the reduced subgraph H_3 contains no odd cycle of length at least $\frac{(1+0.1\eta)t}{3(1+\eta)}$. Since G_i contains no \mathcal{H} for i = 1 or 2, we shall get that H_i contains no connected matching on more than $(\frac{1}{3}-0.2\eta)t$ vertices for i = 1 or 2. Then, we can easily get that $e(H_3) \ge e(H)/3 \ge (\frac{1}{6} - \epsilon/3)t^2$. Now, since H_3 contains no odd cycle of length at least $\frac{(1+0.1\eta)t}{3(1+\eta)}$, we apply Lemma 2.7 to graph H_3 with $\alpha = \frac{(1+0.1\eta)}{3(1+\eta)}$ to deduce that H_3 contains subgraphs H'_3 which is bipartite and H''_3 such that the desired properties hold. Let A and B be the color classes of the bipartition of H'_3 , and let $X = V(H''_3)$. We will show that $|X| < (1/3 - \eta/5)t$ and $\max\{|A|, |B|\} < (\frac{1}{2} - \sqrt{\delta}/2)t$.

Combining with the above desired properties and Lemma 2.8, we can deduce that the reduced graph H contains a subgraph F on at least $(1 - \frac{3}{2}\sqrt{\delta})t$ vertices such that $\delta(F) > v(F) - \sqrt{\delta t}/2$ and $\deg_{F_3}(u) < (\frac{1}{2} + \sqrt{\delta})t$ for any vertex $u \in V(F)$. Since $F \subset H$ and H_i contains no connected matching on more than $(\frac{1}{3} - 0.2\eta)t$ vertices for i = 1 or 2, we conclude that F_i contains no connected matching on more than $(\frac{1}{3} - 3\sqrt{\delta})v(F)$ vertices for i = 1, 2. Moreover, by Lemma 2.6, the order of the largest component of F_i (i = 1, 2) must be less than $\frac{1}{3}v(F)$. By noting $|X| < (1/3 - \eta/5)t$ and Fact 2.2, we obtain a vertex set $B'' \subseteq B \cap V(F)$ such that each vertex of B'' is adjacent to at least $\max\{v(F)/3, (1 - \sqrt{\delta}/4)|B''|\}$. Now we apply Lemma 2.10 to H[B''] to conclude that H[B''] contains a monochromatic component in color 1 or 2 of order larger than $\frac{1}{3}v(F)$. Then F_i (i = 1, 2) contains a monochromatic component in color 1 or 2 of order at least $\frac{1}{3}v(F)$, which will leads to a contradiction.

Details of the remaining proof: We will have the following claims at first.

Claim 3.1 H_3 contains no odd cycle of length at least $\frac{(1+0.1\eta)t}{3(1+\eta)}$.

Proof of Claim 3.1. On the contrary, let C be an odd cycle of length $s \geq \frac{(1+0.1\eta)t}{3(1+\eta)}$. Without loss of generality, assume that its vertex set is [s]. Thus we have that for $i = 1, 2, \ldots, s$, (V_i, V_{i+1}) is ϵ -regular and $d_{G_3}(V_i, V_{i+1}) \geq 1/3$, where the indices take modula on s. By Fact 2.1, for odd $i = 1, 3, \ldots, s - 2$, we can take $V'_i \subseteq V_i$ with $|V'_i| \geq (1 - \epsilon)|V_i|$ and $|V'_{i+1}| \geq (1 - \epsilon)|V_{i+1}|$ such that (V'_i, V'_{i+1}) is $(2\epsilon, 1/3 - 2\epsilon)$ -super-regular. From the property of ϵ -regularity pairs, we can find an odd cycle $u_1u_2 \ldots u_su_1$ such that $u_i \in V'_i$ for $i = 1, 2, \ldots, s - 1$. We apply the techniques used by Luczak [36] and Lemma 2.2 to show that the G_3 contains a C_n .

Let $m = (1-\epsilon)^2 \frac{N}{t}$, $\epsilon' = 2\epsilon$ and $\beta_0 = 1-6\epsilon$. Thus we have that every pair of vertices $v_i \in V'_i$, $v_{i+1} \in V'_{i+1}$ satisfies $\deg_{G_3}(v_i, V'_{i+1}) \ge (1/3 - 2\epsilon)|V'_{i+1}| > \beta_0 m/5$, and $\deg_{G_3}(v_{i+1}, V'_i) > \beta_0 m/5$. Therefore, by Lemma 2.2, for every ℓ , $1 \le \ell \le m - 5\epsilon m/\beta_0$, and for every pair of vertices $u_i \in V'_i$ and $u_{i+1} \in V'_{i+1}$ and odd $i = 1, 3, \ldots, s-2$, G_3 contains a path of length $2\ell + 1$ connecting u_i

and u_{i+1} . Thus, there are odd cycles of all lengths from s to $(s-1)(m-5\epsilon m/\beta_0)$. Since

$$(s-1)(m-5\epsilon m/\beta_0) \ge \left(\frac{(1+0.1\eta)t}{3(1+\eta)} - 1\right) \left(1 - \frac{5\epsilon}{1-6\epsilon}\right) (1-\epsilon)^2 \frac{N}{t}$$

$$\stackrel{(3)}{\ge} \frac{t}{3(1+\eta)} (1-10\epsilon) \frac{(3+\gamma)n}{t} \stackrel{(2)}{\ge} \frac{1+5\eta}{1+\eta} (1-10\epsilon) n,$$

which is at least n again by noting (2). So G_3 contains an odd cycle C_n as desired.

A connected matching in a graph G is a matching M such that all edges of M are in the same connected component of G. By the assumption of G_i contains no \mathcal{H} for i = 1, 2, we have the following claim.

Claim 3.2 For $i = 1, 2, H_i$ contains no connected matching on more than $(\frac{1}{3} - 0.2\eta)t$ vertices.

Proof of Claim 3.2. On the contrary, without loss of generality, suppose that H_1 contains a connected matching M on at least $(\frac{1}{3} - 0.2\eta)t$ vertices that is contained in a tree $T \subset H_1$. Suppose that the vertex set of T is $\{x_1, \ldots, x_l, x_{l+1}, \ldots, x_{2l}, x_{2l+1}, \ldots, x_{2l+l'}\}$, and the matching M has edge set $E_M = \{x_i x_{l+i} : i = 1, \ldots, l\}$.

We will prove that there exists a copy of \mathcal{H} in G_1 , contradicting the assumption that G_1 contains no \mathcal{H} . Since the proof is similar as in [38, Theorem 1.3], we only give a sketch of the proof as follows.

Firstly, we shall apply Fact 2.1 to get a subgraph G_P of G_1 with classes

$$A_1, \ldots, A_l, A_{l+1}, \ldots, A_{2l}, A_{2l+1}, \ldots, A_{2l+l'}$$

which corresponds to the vertices $x_1, \ldots, x_l, x_{l+1}, \ldots, x_{2l}, x_{2l+1}, \ldots, x_{2l+l'}$, and each of those sets has size at least $(1 - 2\epsilon)N/t$ and the bipartite graphs induced by A_i and A_{l+i} are $(2\epsilon, 1/3 - \epsilon)$ super-regular for $i \in [l]$ and the bipartite graphs induced by all the other pairs are $(2\epsilon, 1/3 - \epsilon)$ regular. It is clear that these dense super-regular pairs covering (1 + o(1))n vertices.

Secondly, we partition the vertices of \mathcal{H} and, since $\mathcal{H} = (W, E_{\mathcal{H}})$ has small bandwidth, we can apply Lemma 2.4 to obtain a partition of W which will be composed of clusters

$$W_1, \ldots, W_l, W_{l+1}, \ldots, W_{2l}, W_{2l+1}, \ldots, W_{2l+l'}.$$

Fix $1 \leq j \leq 2l + l'$, we define U_j as the set of vertices of W_j with neighbors in some W_k with $j \neq k$ and $\{x_j x_k\} \notin M$. Define the set $U'_j = N_H(U) \cap (W_j \setminus U)$, where $U = \bigcup_{I=1}^{2l+l'} U_i$.

Then, we can verify that all of the four conditions of Definition 2.4 hold, i.e.,

(1) $|W_i| \le |A_i|$ for $i \in [2l+l']$.

(2)
$$xy \in E_{\mathcal{H}}$$
 for $x \in W_i, y \in W_j$ implies $x_i x_j \in E_T$ for all distinct $i, j \in [2l + l']$

- (3) $|U_i| \leq \epsilon |A_i|$ for $i \in [2l+l']$.
- (4) $|U'_i|, |U'_j| \le \epsilon \min\{|A_i|, |A_j| : x_i x_j \in E_M\}.$

Thus, the partition $\{W_1, \ldots, W_{2l+l'}\}$ of W is $(2\epsilon, T, M)$ -compatible with $\{A_1, \ldots, A_{2l+l'}\}$, which is a partition of $V(G_P)$.

Finally, we can find a copy of \mathcal{H} in G_P by the Embedding Lemma (Lemma 2.3), completing the proof.

Claim 3.3 $e(H_3) \ge e(H)/3 \ge (\frac{1}{6} - \epsilon/3)t^2$.

Proof of Claim 3.3. On the contrary, suppose that $e(H_3) < e(H)/3$. Thus, without loss of generality, suppose that

$$e(H_1) > e(H)/3 \ge \frac{1}{3}(1-\epsilon)\binom{t}{2} \ge \left(\frac{1}{6} - \epsilon/3\right)t^2.$$
 (4)

However, Claim 3.2 implies that H_1 contains no path with more than $(\frac{1}{3} - 0.2\eta)t + 1$ vertices, it follows by Lemma 2.5 that

$$e(H_1) \le \frac{(\frac{1}{3} - 0.2\eta)t - 1}{2} \cdot t < \left(\frac{1}{6} - 0.1\eta\right)t^2$$

This contradicts (4) by noting (2).

By Claim 3.1, H_3 contains no odd cycle of length at least $\frac{(1+0.1\eta)t}{3(1+\eta)}$. Now we apply Lemma 2.7 to graph H_3 with $\alpha = \frac{(1+0.1\eta)}{3(1+\eta)}$ to deduce that H_3 contains subgraphs H'_3 and H''_3 such that

(i) $V(H'_3) \cup V(H''_3) = V(H_3)$, $V(H'_3) \cap V(H''_3) = \emptyset$ and each of the sets $V(H'_3)$ and $V(H''_3)$ is either empty or contains at least $\alpha \delta t/2$ vertices;

(*ii*) H'_3 is bipartite;

- (*iii*) H_3'' contains no more than $\alpha t |V(H_3'')|/2$ edges;
- (iv) all except no more than δt^2 edges of H_3 belong to either H'_3 or H''_3 .

Let A and B be the sets of the bipartition of H'_3 , and let $X = V(H''_3)$. From Claim 3.3 and the property of H_3 (see (iv)), we have

$$e(H'_3) + e(H''_3) \ge e(H_3) - \delta t^2 \ge \left(\frac{1}{6} - \epsilon/3\right) t^2 - \delta t^2 > \left(\frac{1 - 7\delta}{6}\right) t^2.$$
(5)

Claim 3.4 $|X| < (1/3 - \eta/5)t$.

Proof of Claim 3.4. Let us put $|X| = \lambda t$ and $|A \cup B| = (1 - \lambda)t$. By Lemma 2.7 (*iii*),

$$e(H'_{3}) + e(H''_{3}) \leq e(H'_{3}) + \frac{\alpha t |V(H''_{3})|}{2} \leq \frac{|V(H'_{3})|^{2}}{4} + \frac{\alpha t |V(H''_{3})|}{2} \leq \frac{(1-\lambda)^{2}}{4}t^{2} + \frac{\lambda (1+0.1\eta)}{6(1+\eta)}t^{2},$$
(6)

which together with (5) yield that

$$\frac{(1-\lambda)^2}{4} + \frac{\lambda \left(1-0.8\eta\right)}{6} > \frac{(1-\lambda)^2}{4} + \frac{\lambda \left(1+0.1\eta\right)}{6(1+\eta)} \ge \frac{1}{6} - \frac{7\delta}{6},$$

from which we obtain that $3\lambda^2 - (4 + 1.6\eta)\lambda + (1 + 14\delta) > 0$. Since $\lambda \leq 1$, it follows that

$$\lambda < \frac{1}{6} \left(4 + 1.6\eta - \sqrt{(4 + 1.6\eta)^2 - 12(1 + 14\delta)} \right) < \frac{1}{3} \left(2 + 0.8\eta - \sqrt{1 + 3\eta} \right) < \frac{1}{3} - \frac{\eta}{5}$$

provided η is sufficiently small and by noting (2).

Claim 3.5 max{|A|, |B|} < $(\frac{1}{2} - \sqrt{\delta}/2)t$.

Proof of Claim 3.5. On the contrary, suppose that $|A| \ge (\frac{1}{2} - \sqrt{\delta}/2)t$ without loss of generality. Let H[A] be the subgraph of H induced by A. Note that H[A] only contains edges in color 1 or color 2 as H'_3 is bipartite. Note also that $e(H[A]) \ge \binom{|A|}{2} - \binom{t}{2}$. By Fact 2.2, H[A] contains a subgraph H[A'] such that $|A'| \ge (\frac{1}{2} - \sqrt{\delta}/2)t - \sqrt{\epsilon}t > (\frac{1}{2} - \sqrt{\delta})t$ and

$$\delta(H[A']) \ge |A'| - 2\sqrt{\epsilon}t > |A'| - 6\sqrt{\epsilon}|A'| \stackrel{(2)}{>} (1 - \sqrt{\delta})|A'|,$$

the second inequality due to $t < \frac{|A'|}{(1/2-\sqrt{\delta})}$ and δ is sufficiently small. Now we apply Lemma 2.9 with graph H[A'] and $n = \frac{(1/2-\sqrt{\delta})t+1}{3}$ to conclude that H[A'] contains a monochromatic path $P_{(1-2\sqrt{\delta})t}$ in color 1 or color 2. As $\frac{(1-2\sqrt{\delta})t}{3} > (\frac{1}{3}-0.2\eta)t+1$ from (2), we have that either G_1 or G_2 must contain a copy of \mathcal{H} by Claim 3.2, a contradiction.

For graphs P and Q, we use $P \cup Q$ to denote the graph defined on $V(P) \cup V(Q)$ whose edge set is $E(P) \cup E(Q)$. From Claims 3.4-3.5 noting that $V(H'_3) \cup V(H''_3) = V(H_3) = V(H)$ and $V(H'_3) \cap V(H''_3) = \emptyset$, we have that for any vertex $u \in V(H)$,

$$\deg_{H'_3 \cup H''_3}(u) < \left(\frac{1}{2} - \frac{\sqrt{\delta}}{2}\right)t.$$

$$\tag{7}$$

For an edge-colored graph F, we use F_i to denote the subgraph induced by edges in color i in F.

Claim 3.6 *H* has a subgraph *F* with at least $(1 - \frac{3}{2}\sqrt{\delta})t$ vertices such that $\delta(F) > v(F) - \sqrt{\delta}t/2$ and $\deg_{F_3}(u) < (\frac{1}{2} + \sqrt{\delta})t$ for any vertex $u \in V(F)$.

Proof of Claim 3.6. We know that all but at most δt^2 edges of H_3 are contained in $H'_3 \cup H''_3$. By Lemma 2.8, there exists an induced subgraph H^0_3 of H_3 such that $v(H^0_3) \ge v(H) - \sqrt{2\delta t}$ and

$$\deg_{H_3^0}(u) < \deg_{H_3' \cup H_3''}(u) + \sqrt{2\delta t} \stackrel{(7)}{<} \left(\frac{1}{2} - \sqrt{\delta}/2\right) t + \sqrt{2\delta t} < \left(\frac{1}{2} + \sqrt{\delta}\right) t \tag{8}$$

for any vertex $u \in V(H_3^0)$.

Note that $e(H[V(H_3^0)]) \ge {\binom{v(H_3^0)}{2}} - \epsilon {\binom{t}{2}}$, it follows from Fact 2.2 that $H[V(H_3^0)]$ contains a subgraph F such that

$$v(F) \ge v(H_3^0) - \sqrt{\epsilon}t \ge v(H) - \sqrt{2\delta}t - \sqrt{\epsilon}t \stackrel{(2)}{>} \left(1 - \frac{3}{2}\sqrt{\delta}\right)t \tag{9}$$

and

$$\delta(F) \ge v(H_3^0) - 2\sqrt{\epsilon}t > v(F) - \sqrt{\delta}t/2.$$
(10)

Since $F_3 \subset H_3^0$, by noting (8), we obtain that for any $u \in V(F)$,

$$\deg_{F_3}(u) \le \deg_{H_3^0}(u) < \left(\frac{1}{2} + \sqrt{\delta}\right)t,\tag{11}$$

completing the proof.

According to (10) and (11), we obtain that for any vertex $u \in V(F)$,

$$\deg_{F_1 \cup F_2}(u) = \deg_{F_1}(u) + \deg_{F_2}(u) \ge \delta(F) - d_{F_3}(u) \ge v(F) - \sqrt{\delta}t/2 - \left(\frac{1}{2} + \sqrt{\delta}\right)t$$

$$\stackrel{(9)}{>} \left(1 - \frac{\frac{1}{2} + \frac{3}{2}\sqrt{\delta}}{1 - \frac{3}{2}\sqrt{\delta}}\right)v(F) > \left(\frac{1}{2} - 3\sqrt{\delta}\right)v(F).$$
(12)

Recall that a connected matching in a graph G is a matching M such that all edges of M are in the same connected component of G.

Claim 3.7 For i = 1, 2, F_i contains no connected matching on more than $(\frac{1}{3} - 3\sqrt{\delta})v(F)$ vertices.

Proof of Claim 3.7. Note that since $F \subset H$, by Claim 3.2, F_i contains no connected matching on more than $(\frac{1}{3} - 0.2\eta)t$ vertices for i = 1, 2. Since

$$\left(\frac{1}{3} - 3\sqrt{\delta}\right)v(F) \stackrel{(9)}{>} \left(\frac{1}{3} - 3\sqrt{\delta}\right)\left(1 - \frac{3\sqrt{\delta}}{2}\right)t > \left(\frac{1}{3} - \frac{7\sqrt{\delta}}{2}\right)t \stackrel{(2)}{>} \left(\frac{1}{3} - 0.2\eta\right)t.$$

Thus F_i (i = 1, 2) contains no connected matching on more than $(\frac{1}{3} - 3\sqrt{\delta})v(F)$ vertices. \Box

Claim 3.8 For i = 1, 2, the largest component of F_i has order less than $\frac{1}{3}v(F)$.

Proof of Claim 3.8. Let R_1, \ldots, R_r be the components of F_1 and let B_1, \ldots, B_b be the components of F_2 . Without loss of generality, suppose that $|V(R_i)| \ge |V(R_{i+1})|$ for all $1 \le i \le r-1$, and $|V(B_j) \ge |V(B_{j+1})|$ for all $1 \le j \le b-1$. Let $r', 0 \le r' \le r$, be the maximum integer such that $|V(R_{r'})| \ge (\frac{1}{3} - 3\sqrt{\delta})v(F)$. Similarly, let $b', 0 \le b' \le b$, be the maximum integer such that $|V(R_{b'})| \ge (\frac{1}{3} - 3\sqrt{\delta})v(F)$. We aim to show that r' = b' = 0.

By Claim 3.7, F_1 and hence each R_i contains no connected matching on more than $(\frac{1}{3} - 3\sqrt{\delta})v(F)$ vertices. Thus Lemma 2.6 implies that each R_i has a partition $S_{R_i} \cup Q_{R_i} \cup I_{R_i}$ satisfying

$$|Q_{R_i}| + 2|S_{R_i}| = \min\left\{ |V(R_i)|, \left(\frac{1}{3} - 3\sqrt{\delta}\right)v(F) - 1 \right\}.$$
(13)

Proposition 3.1 For i > r', $I_{R_i} = \emptyset$ and $S_{R_i} = \emptyset$; similarly, for j > b', $I_{B_j} = \emptyset$ and $S_{B_j} = \emptyset$.

Proof. For i > r', $|V(R_i)| \le (\frac{1}{3} - 3\sqrt{\delta})v(F) - 1$, implying that $|Q_{R_i}| + 2|S_{R_i}| = |V(R_i)|$ and $I_{R_i} = \emptyset$ by (13). Thus $|Q_{R_i}| + |S_{R_i}| = |V(R_i)|$, and $S_{R_i} = \emptyset$ follows. The second assertion is similar.

According to Lemma 2.6 and (13), we obtain that for $1 \le i \le r$,

$$|S_{R_i}| < \frac{1}{2} \left(\frac{1}{3} - 3\sqrt{\delta}\right) v(F), \quad |Q_{R_i}| + 2|S_{R_i}| < \left(\frac{1}{3} - 3\sqrt{\delta}\right) v(F), \tag{14}$$

and

$$|I_{R_i}| = |V(R_i) \setminus (Q_{R_i} \cup S_{R_i})| = |V(R_i)| + |S_{R_i}| - (|Q_{R_i}| + 2|S_{R_i}|)$$

> $|V(R_i)| + |S_{R_i}| - \left(\frac{1}{3} - 3\sqrt{\delta}\right)v(F).$ (15)

Note that for any B_j , similar sets S_{B_j} , I_{B_j} and Q_{B_j} can be defined, with analogues of the above bounds. In particular, $|I_{B_j}| \ge |V(B_j)| + |S_{B_j}| - (\frac{1}{3} - 3\sqrt{\delta})v(F)$.

By Lemma 2.6 (*iv*), for $1 \leq i \leq r$, any vertex $u \in I_{R_i}$ satisfies that $\deg_{F_1}(u) \leq \frac{1}{2}(\frac{1}{3} - 3\sqrt{\delta})v(F)$. Similarly, for $1 \leq j \leq b$, any vertex $u \in I_{B_j}$ satisfies that $\deg_{F_2}(u) \leq \frac{1}{2}(\frac{1}{3} - 3\sqrt{\delta})v(F)$.

Note that for $1 \leq i \leq r$, each vertex in Q_{R_i} has at most one neighbor in I_{R_i} in F_1 by Lemma 2.6 (*iii*). Hence, any vertex $u \in Q_{R_i}$ satisfies that $\deg_{F_1}(u) \leq |(Q_{R_i} \cup S_{R_i}) \setminus \{u\}| + 1$, which is less than $(\frac{1}{3} - 3\sqrt{\delta})v(F)$ by noting (14). Similarly, for $1 \leq j \leq b$, any vertex $u \in Q_{B_j}$ satisfies that $\deg_{F_2}(u) < (\frac{1}{3} - 3\sqrt{\delta})v(F)$.

Proposition 3.2 $I_{R_i} \cap (I_{B_j} \cup Q_{B_j}) = \emptyset$.

Proof. Indeed, if there is a vertex $u \in I_{R_i} \cap (I_{B_j} \cup Q_{B_j})$, then from the above observation,

$$\deg_{F_1 \cup F_2}(u) = \deg_{F_1}(u) + \deg_{F_2}(u) < \frac{1}{2} \left(\frac{1}{3} - 3\sqrt{\delta}\right) v(F) + \left(\frac{1}{3} - 3\sqrt{\delta}\right) v(F),$$

which is less than $(\frac{1}{2} - 4\sqrt{\delta})v(F)$, contradicting (12).

Therefore, for $1 \leq i \leq r$, the set $I_{R_i} \subseteq \bigcup_{1 \leq j \leq b} S_{B_j}$ due to Proposition 3.2. Since $S_{B_j} = \emptyset$ for j > b' by Proposition 3.1, we have that $I_{R_i} \subseteq \bigcup_{1 \leq j \leq b'} S_{B_j}$. Similarly, for $1 \leq j \leq b$, $I_{B_j} \subseteq \bigcup_{1 \leq i \leq r'} S_{R_i}$. Consequently, again by Proposition 3.1, $I_{R_i} = \emptyset$ for i > r' and $I_{B_j} = \emptyset$ for j > b', we obtain that

$$\sum_{i=1}^{r'} |I_{R_i}| = \sum_{i=1}^{r} |I_{R_i}| \le \sum_{j=1}^{b'} |S_{B_j}|, \text{ and } \sum_{j=1}^{b'} |I_{B_j}| \le \sum_{i=1}^{r'} |S_{R_i}|,$$
(16)

which together with (15) yield that

$$0 \geq \sum_{j=1}^{b'} |I_{B_j}| - \sum_{i=1}^{r'} |S_{R_i}|$$

$$\geq \sum_{j=1}^{b'} \left(|V(B_j)| + |S_{B_j}| - \left(\frac{1}{3} - 3\sqrt{\delta}\right)t \right) - \sum_{i=1}^{r'} |S_{R_i}|$$

$$\stackrel{(16)}{\geq} \sum_{j=1}^{b'} \left(|V(B_j)| - \left(\frac{1}{3} - 3\sqrt{\delta}\right)t \right) + \sum_{i=1}^{r'} |I_{R_i}| - \sum_{i=1}^{r'} |S_{R_i}|$$

$$\stackrel{(15)}{\geq} \sum_{j=1}^{b'} \left(|V(B_j)| - \left(\frac{1}{3} - 3\sqrt{\delta}\right)v(F) \right) + \sum_{i=1}^{r'} \left(|V(R_i)| - \left(\frac{1}{3} - 3\sqrt{\delta}\right)v(F) \right), \quad (17)$$

which implies that r' = b' = 0 since $|V(R_i)| \ge (\frac{1}{3} - 3\sqrt{\delta})v(F)$ and $|V(B_j)| \ge (\frac{1}{3} - 3\sqrt{\delta})v(F)$ for $1 \le i \le r'$ and $1 \le j \le b'$. This completes Claim 3.8.

By noting Claim 3.4 and $A \cup B \cup X = V(H_3) = V(H)$ and $F \subseteq H$, we obtain that

$$\begin{aligned} |(A \cup B) \cap V(F)| &\ge |V(F)| - |X| > v(F) - \left(\frac{1}{3} - \frac{1}{5}\eta\right)t \stackrel{(9)}{>} v(F) - \frac{\frac{1}{3} - \frac{1}{5}\eta}{1 - \frac{3}{2}\sqrt{\delta}}v(F) \\ &\stackrel{(2)}{>} v(F) - \frac{1}{3}(1 - \sqrt{\delta}/2)v(F) = \left(\frac{2}{3} + \frac{1}{6}\sqrt{\delta}\right)v(F). \end{aligned}$$

Thus, one of $|A \cap V(F)|$ and $|B \cap V(F)|$ must be at least $(\frac{1}{3} + \frac{1}{12}\sqrt{\delta})v(F)$. Without loss of generality, we suppose that $B' \subseteq B \cap V(F)$ satisfies

$$\left|B'\right| = \left(\frac{1}{3} + \frac{1}{12}\sqrt{\delta}\right)v(F).$$
(18)

Note that $e(H[B']) \ge {|B'| \choose 2} - \epsilon {t \choose 2}$. Therefore, by deleting at most $\sqrt{\epsilon t}$ vertices from B' we obtain a vertex set $B'' \subseteq B'$ such that each vertex of B'' is adjacent to at least $|B'| - 2\sqrt{\epsilon t}$ in H[B''] from Fact 2.2. We obtain that

$$\delta(H[B'']) \ge |B'| - 2\sqrt{\epsilon}t \stackrel{(18),(9)}{>} \left(\frac{1}{3} + \frac{1}{12}\sqrt{\delta}\right)v(F) - \frac{2\sqrt{\epsilon}}{1 - \frac{3}{2}\sqrt{\delta}}v(F) \stackrel{(2)}{\ge} \frac{1}{3}v(F).$$
(19)

By noting (18),

$$\delta(H[B'']) > \frac{1}{3}v(F) = \frac{|B'|}{1 + \sqrt{\delta}/4} \ge \frac{|B''|}{1 + \sqrt{\delta}/4} > \left(1 - \frac{\sqrt{\delta}}{4}\right) |B''|.$$

Note that all edges of H[B''] are colored only with color 1 or 2 since $B'' \subseteq B$ and B is one of parts of the bipartite graph H'_3 . Now we apply Lemma 2.10 to H[B''] to conclude that H[B''] contains a monochromatic component in color 1 or 2 of order larger than $\delta(H[B''])$, which is at least $\frac{1}{3}v(F)$ according to (19). Since $H[B''] \subseteq F$, it follows that F contains a monochromatic component in color 1 or 2 of order at least $\frac{1}{3}v(F)$. This contradicts Claim 3.8.

In conclusion, the proof of Theorem 1.1 is complete.

References

- P. Allen, G. Brightwell and J. Skokan, Ramsey-goodness and otherwise, *Combinatorica* 33 (2013), 125–160.
- [2] N. Alon, J. Fox and Y. Zhao, Efficient arithmetic regularity and removal lemmas for induced bipartite patterns, Discrete Anal. 2019, Paper No. 3, 14 pp.
- [3] J. Balogh, A. Kostochka, M. Lavrov and X. Liu, Monochromatic paths and cycles in 2edge-colored graphs with large minimum degree, arXiv:1906.02854.
- [4] F. S. Benevides, T. Luczak, A. Scott, J. Skokan and M. White, Monochromatic cycles in 2-coloured graphs, *Combin. Probab. Comput.* 21 (2012), 57–87.
- [5] F. S. Benevides and J. Skokan, The 3-colored Ramsey number of even cycles, J. Combin. Theory Ser. B 99 (2009), 690–708.
- [6] H. Bielak, Multicolor Ramsey numbers for some paths and cycles, Discuss. Math. Graph Theory 29 (2009), 209–218.
- [7] J. A. Bondy and P. Erdős, Ramsey numbers for cycles in graphs, J. Combin. Theory Ser. B 14 (1973), 46–54.
- [8] J. Böttcher, Embedding large graphs-The Bollobás-Komlós conjecture and beyond, Ph.D. thesis, Technischen Universität München, 2009.
- [9] J. Böttcher, P. Heinig and A. Taraz, Embedding into bipartite graphs, SIAM J. Discrete Math. 24 (2010), 1215–1233.
- [10] J. Böttcher, K. P. Pruessmann, A. Taraz and A. Würfl, Bandwidth, expansion, treewidth, separators and universality for bounded-degree graphs, *European J. Combin.* 31 (2010), 1217–1227.
- [11] X. Chen, Q. Lin and C. You, Ramsey numbers of large books, J. Graph Theory, to appear. DOI: 10.1002/jgt.22815.
- [12] D. Conlon, The Ramsey number of books, Adv. Combin. 3 (2019), 12pp.
- [13] D. Conlon and J. Fox, Bounds for graph regularity and removal lemmas, Geom. Funct. Anal. 22 (2012), 1191–1256.
- [14] D. Conlon, J. Fox and Y. Wigderson, Ramsey number of books and quasirandomness, *Combinatorica*, to appear.
- [15] T. Dzido and R. Fidytek, On some three color Ramsey numbers for paths and cycles, Discrete Math. 309 (2009), 4955–4958.
- [16] P. Erdős, R. J. Faudree, C. C. Rousseau and R. H. Schelp, Generalized Ramsey theory for multiple colors, J. Combin. Theory Ser. B 20 (1976), 250–264.

- [17] P. Erdős and T. Gallai, On maximal paths and circuits of graphs, Acta Math. Hungar. 10 (1959), 337–356.
- [18] R. J. Faudree, S. L. Lawrence, T. D. Parsons and R. H. Schelp, Path-Cycle Ramsey numbers, Discrete Math. 10 (1974), 269–277.
- [19] R. J. Faudree and R. H. Schelp, All Ramsey numbers for cycles in graphs, *Discrete Math.* 8 (1974), 313–329.
- [20] R. J. Faudree and R. H. Schelp, Path Ramsey numbers in multicolorings, J. Combin. Theory Ser. B 19 (1975), 150–160.
- [21] A. Figaj and T. Łuczak, The Ramsey number for a triple of long even cycles, J. Combin. Theory Ser. B 97 (2007), 584–596.
- [22] A. Figaj and T. Łuczak, The Ramsey numbers for a triple of long cycles, *Combinatorica* 38 (2018), 827–845.
- [23] D. G. Ferguson, The Ramsey number of mixed-parity cycles I, arXiv:1508.07154.
- [24] D. G. Ferguson, The Ramsey number of mixed-parity cycles II, arXiv:1508.07171.
- [25] D. G. Ferguson, The Ramsey number of mixed-parity cycles III, arXiv:1508.07176.
- [26] J. Fox, L. M. Lovász and Y. Zhao, On regularity lemmas and their algorithmic applications, Combin. Probab. Comput. 26 (2017), 481–505.
- [27] L. Gerencsér and A. Gyarfás, On Ramsey-type problems, Ann. Univ. Sci. Budapest Eötvös Sect. Math. 10 (1967), 167–170.
- [28] A. Gyárfás, Large monochromatic components in edge colorings of graphs: A survey. In Ramsey Theory: Yesterday, Today and Tomorrow (A. Soifer, ed.), Birkhäuser, pp. 77–96, 2010.
- [29] A. Gyárfás, M. Ruszinkó, N. Sárközy and E. Szemerédi, Three-color Ramsey numbers for Paths, Combinatorica 27 (2007), 35–69.
- [30] A. Gyárfás and G. N. Sárközy, Star versus two stripes Ramsey numbers and a conjecture of Schelp, Combin. Probab. Comput. 21 (2012), 179–186.
- [31] M. Jenssen and J. Skokan, Exact Ramsey numbers of odd cycles via nonlinear optimisation, Adv. Math. 376 (2021), 107444, 46 pp.
- [32] J. Komlós and M. Simonovits, "Szemerédi's regularity lemma and its applications in graph theory," Combinatorics, Paul Erdős is eighty, Vol. 2 (Keszthely, 1993), 295–352, Bolyai Soc. Math. Stud., 2, János Bolyai Math. Soc., Budapest, 1996.
- [33] Y. Kohayakawa, M. Simonovits and J. Skokan, The 3-colored Ramsey number of odd cycles, *Electron. Notes Discrete Math.* 19 (2005), 397–402.
- [34] C. Knierim and P. Su, Improved bounds on the multicolor Ramsey numbers of paths and even cycles, *Electron. J. Combin.* 26 (2019), #P1.26.

- [35] Q. Lin and X. Peng, Large book-cycle Ramsey numbers, SIAM J. Discrete Math. 35 (2021), 532–545.
- [36] T. Łuczak, $R(C_n, C_n, C_n) \leq (4 + o(1))n$, J. Combin. Theory Ser. B 75 (1999), 174–187.
- [37] T. Luczak, M. Simonovits and J. Skokan, On the multi-colored Ramsey numbers of cycles, J. Graph Theory 69 (2012), 169–175.
- [38] G. Mota, G. N. Sárközy, M. Schacht and A. Taraz, Ramsey number for bipartite graphs with small bandwidth, *European J. Combin.* 48 (2015), 165–176.
- [39] V. Nikiforov and C. C. Rousseau, Ramsey goodness and beyond, *Combinatorica* 29 (2009), 227–262.
- [40] G. R. Omidi and G. Raeisi, On multicolor Ramsey number of paths versus cycles, *Electron. J. Combin.* 18 (2011), #P24.
- [41] V. Rödl and M. Schacht, Regularity lemmas for graphs. Fete of combinatorics and computer science, 287–325, Bolyai Soc. Math. Stud., 20, János Bolyai Math. Soc., Budapest, 2010.
- [42] V. Rosta, On a Ramsey-type problem of J. A. Bondy and P. Erdős. I, II, J. Combin. Theory Ser. B 15 (1973), 105–120.
- [43] G. N. Sárkozy, On the multi-colored Ramsey numbers of paths and even cycles, *Electron. J. Combin.* 23 (2016), #P3.
- [44] R. H. Schelp, Some Ramsey-Turán type problems and related questions, Discrete Math. 312 (2012), 2158–2161.
- [45] Z. Shao, X. Xu, X. Shi and L. Pan, Some three-color Ramsey numbers, $R(P_4, P_5, C_k)$ and $R(P_4, P_6, C_k)$, European J. Combin. 30 (2009), 396–403.
- [46] E. Szemerédi, Regular partitions of graphs, Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, 1976), 399–401, Colloq. Internat. CNRS, 260, CNRS, Paris, 1978.
- [47] E. Szemerédi, Arithmetic progressions, different regularity lemmas and removal lemmas, Commun. Math. Stat. 3 (2015), 315–328.