# Effects of concurrency on epidemic spreading in Markovian temporal networks

Ruodan Liu<sup>a</sup>, Masaki Ogura<sup>b</sup>, Elohim Fonseca Dos Reis<sup>a</sup>, Naoki Masuda<sup>a,c,d,\*</sup>

<sup>a</sup>Department of Mathematics, State University of New York at Buffalo, Buffalo, NY 14260-2900, USA. <sup>b</sup>Graduate School of Information Science and Technology, Osaka University, Suita, Osaka 565-0871, Japan <sup>c</sup>Computational and Data-Enabled Sciences and Engineering Program, State University of New York at Buffalo, Buffalo, NY 14260-5030. USA

 $^d$ Faculty of Science and Engineering, Waseda University, 169-8555 Tokyo, Japan

#### Abstract

The concurrency of edges, quantified by the number of edges that share a common node at a given time point, may be an important determinant of epidemic processes in temporal networks. We propose theoretically tractable Markovian temporal network models in which each edge flips between the active and inactive states in continuous time. The different models have different amounts of concurrency while we can tune the models to share the same statistics of edge activation and deactivation (and hence the fraction of time for which each edge is active) and the structure of the aggregate (i.e., static) network. We analytically calculate the amount of concurrency of edges sharing a node for each model. We then numerically study effects of concurrency on epidemic spreading in the stochastic susceptible-infectious-recovered (SIR) dynamics on the proposed temporal network models. We find that the concurrency enhances epidemic spreading near the epidemic threshold while this effect is small in some cases. When the infection rate is substantially larger than the epidemic threshold, the concurrency suppresses epidemic spreading. In sum, our numerical simulations indicate that the impact of concurrency on epidemic spreading is modest. The proposed temporal network models are expected to be useful for investigating effects of concurrency on various collective dynamics on networks including both infectious and other dynamics.

Keywords: Concurrency, temporal network, susceptible-infectious-recovered model, epidemic spreading, Poisson process.

#### 1. Introduction

The structure of contact networks among individuals shapes the dynamics of contagion processes in a population such as the number of individuals infected, their spatial distributions, and speed of spreading [1–4]. For example, a heterogeneous degree distribution, where the degree is the number of neighboring nodes that a node has, and a short average distance between nodes are two factors that usually enhance epidemic spreading on networks. In fact, empirical contact networks often vary over time on a time scale comparable to or faster than that of epidemic dynamics, probably most famously owing to mobility of human or animal individuals. This observation has naturally led to the investigation of how features of such temporal (i.e., time-varying) networks and statistical properties of contact events, such as distributions of inter-contact times, temporal correlation in inter-contact times, and appearance and disappearance of nodes and edges, affect outcomes of contagion processes [5–10].


The concurrency is a feature of temporal contact networks that has been investigated in both mathematical and field epidemiology for over two decades, while many of these studies do not specifically refer to temporal networks [11–19]. Concurrency generally refers to multiple partnerships of an individual that overlap in time. Concurrency has been examined in particular in the context of sexually transmitted infections such as HIV, specifically regarding whether or not the concurrency was high in sub-Saharan Africa and, if so, the high

Email address: naokimas@buffalo.edu (Naoki Masuda)

<sup>\*</sup>Corresponding author

concurrency enhanced HIV spreading there [11–13, 20–23]. Suppose that a partnership between individuals i and j overlaps one between i and j' for some time. Such concurrency may promote epidemic spreading because rapid transmission from j to j' through i and vice versa is possible during the concurrent partnerships (see Fig. 1(a)). In contrast, if a partnership between i and j occurs before that between i and j' without an overlap, i.e., without concurrency, the transmission from j to j' can still occur in various different ways, but the transmission from j' to j does not (see Fig. 1(b)).

Modeling studies are diverse in how to quantify the amount of concurrency. Many studies measure the concurrency through the mean degree, or the average contact rate for nodes [19, 24–26]. However, in these cases, it is hard to say whether an increased epidemic size owes to a high density of edges or a large amount of concurrency [15, 27–29]. In studies of epidemic processes on static networks, it is a stylized fact that epidemic spreading is enhanced if there are many edges with all other things being equal. Another line of theoretical and computational approach to concurrency quantifies concurrency by the heterogeneity in the degree distribution of the network [12, 13, 15]. However, the results that a high concurrency in this sense enhances epidemic spreading is equivalent to an established result in static network epidemiology that heterogeneous degree distributions lead to a small epidemic threshold, thus promoting epidemic spreading [2–4, 30]. In some studies, the authors investigated the effect of concurrency by carefully ensuring that the degree distribution (and hence the average degree) stays the same across the comparisons while they manipulate the amount of concurrency [27–29]. These studies suggest that, despite the different definitions of the concurrency employed in these studies, higher concurrency increases epidemic spreading (but see [27] for different results). However, mathematical models that enable us to analyze the effect of concurrency by fixing the structure of the static network are still scarce.



**Figure 1:** Schematic of part of temporal networks with different amounts of concurrency. We depict two edges sharing a node in each case. (a) Concurrent partnerships. (b) Non-concurrent partnerships. A shaded box represents a duration for which a partnership is present on the edge. The thick lines are examples of time-respecting paths transmitting infection from j to j' (shown in orange) or from j' to j (shown in magenta). This figure is inspired by Fig. 1 in Ref. [15] and Fig. 1 in Ref. [27].

In the present study, we focus on effects of concurrency on epidemic spreading for an arbitrary static network on top of which partnerships, infection events, and recovery events occur. In this manner, we aim to study the effect of concurrency without being affected by the effect of the network structure itself. We consider three temporal network models that have different amounts of concurrency while keeping the probability that each edge is available the same across comparisons. In terms of a concurrency measure, we analytically evaluate the amount of concurrency of edges sharing a node for the three models as well as other properties of the models. Then, we numerically study effects of concurrency on epidemic spreading using the stochastic susceptible-infectious-recovered (SIR) model.

# 2. Temporal network models

We introduce three models of undirected and unweighted temporal networks in continuous time, which we build from independent Markov processes occurring on a given static network. We use these models to compare temporal networks that have the same time-aggregated network but different amounts of concurrency. Let G be an undirected and unweighted static network with node set  $V = \{1, ..., N\}$  and edge set  $E = \{e_i\}_{i=1}^M$ , where N is the number of nodes and M is the number of edges. We assume that the network G has no self-loops. In the temporal network constructed based on undirected and unweighted static network G, by definition, each edge  $e_i \in E$  is either active (i.e., temporarily present) or inactive (i.e., temporarily absent) at any given time  $t \in \mathbb{R}$  and switches between the active and inactive states over time. We denote by G the temporal network and by G the instantaneous network in G observed at time G.

#### 2.1. Model 1

Let  $\tau_1$  and  $\tau_2$  be the duration of the active state of and the inactive state of an edge, respectively. In our model 1, we assume that  $\tau_1$  is independently drawn from probability density  $\psi_1(\tau_1)$  each time the edge switches from the inactive to the active state. Similarly, we draw  $\tau_2$  independently from probability density  $\psi_2(\tau_2)$  each time the edge switches from the active to the inactive state. The duration  $\tau_1$  or  $\tau_2$  for different edges obeys the same distributions but is drawn independently for the different edges. When  $\psi_1(\tau_1)$  and  $\psi_2(\tau_2)$  are both exponential distributions, the stochastic dynamics of the state of each edge obeys independent continuous-time Markov processes with two states, and our model 1 reduces to previously proposed models [31, 32].

#### 2.2. Model 2

In models 2 and 3, we assume that each node independently obeys a continuous-time Markov process with two states, which we refer to as the high-activity and low-activity states. Each node independently switches between the high-activity state, denoted by h, and the low-activity state, denoted by  $\ell$ . Let a be the rate at which the state of a node changes from  $\ell$  to h, and let b be the rate at which the state of a node changes from  $\ell$  to h, and let h be the rate at which the state of a node changes from h to  $\ell$ . In model 2, each edge  $(u,v) \in E$ , where  $u,v \in V$ , is active at any given time if and only if both u and u are in the u state. The dynamics of a single edge in model 2 is schematically shown in Fig. 2. The model resembles the so-called AND model in [33]. An intuitive interpretation of model 2 is that two individuals chat with each other if and only if both of them want to.




Figure 2: Schematic illustration of models 2 and 3. In model 2, edge (1,2) is active if and only if both node 1 and node 2 are in the h state. Otherwise, the edge is inactive. In model 3, edge (1,2) is active if and only if either node 1 or node 2 is in the h state.

# 2.3. Model 3

Model 3 is a variant of model 2. As in model 2, we assume that each node independently switches between the high-activity and low-activity states at rates a and b. In model 3, the edge between two nodes is active if and only if either node is in the b state (see Fig. 2 for a schematic). This model resembles the OR model proposed in [33]. The intuition behind model 3 is that one person can start conversation with another person whenever either person wants to talk regardless of whether the other person wants to.

#### 3. Concurrency for the three models

In this section, we first define a concurrency index for temporal networks in which partnership on each edge appears and disappears over time. Then, we calculate and compare the concurrency index and related measures for models 1, 2, and 3.

#### 3.1. Definition of a concurrency index

Consider a temporal network  $\mathcal{G}$  that is constructed on the underlying static network G and defined for time  $t \in [0,T]$ . We refer to G as the aggregate network. If two edges in  $\mathcal{G}$  sharing a node in G are both active at  $t \in \mathbb{R}$ , we say that the two edges are concurrent at time t; see Fig. 1(a) for an example. Consider a pair of edges sharing a node in the aggregate network, denoted by  $e_i$  and  $e_j$ , where  $e_i, e_j \in E$ . In fact, the likelihood that  $e_i$  and  $e_j$  are concurrent at time t depends on how likely each single edge is active at time t. We define a concurrency index that takes into account of this factor. To this end, we first define the set of time for which an edge  $e \in E$  is active by

$$S(e) = \{ t \in [0, T]; \text{ edge } e \text{ is active at time } t \}.$$
(3.1)

We define the concurrency for the edge pair  $\{e_i, e_j\} \in \mathcal{S}$  by

$$\kappa(e_i, e_j) = \frac{|S(e_i) \cap S(e_j)|}{\min\{|S(e_i)|, |S(e_j)|\}},$$
(3.2)

where S is the set of edge pairs that share a node in G. Note that

$$|\mathcal{S}| = \sum_{i=1}^{N} \frac{\overline{k}_i(\overline{k}_i - 1)}{2},\tag{3.3}$$

where  $\overline{k}_i$  is the degree of the *i*th node in G [12, 13]. The numerator on the right-hand side of Eq. (3.2) is equal to the length of time for which both  $e_i$  and  $e_j$  are active. The denominator is a normalization constant to discount the fact that the numerator would be large if the two edges are active for long time. Because any edge e in the aggregate network G should be active sometime in [0,T], the denominator is always positive; otherwise, we should exclude e from G.

We define the concurrency index for temporal network  $\mathcal{G}$  by

$$\kappa(\mathcal{G}) = \frac{1}{|\mathcal{S}|} \sum_{i,j \text{ such that } 1 \le i < j \le N \text{ and } \{e_i, e_j\} \in \mathcal{S}} \kappa(e_i, e_j).$$
(3.4)

It holds true that  $0 \le \kappa(\mathcal{G}) \le 1$  because  $0 \le \kappa(e_i, e_j) \le 1$ . For empirical or numerical data, [0, T] is the observation time window. For a stochastic temporal network model, we calculate S(e) as the expectation and in the limit of  $T \to \infty$  such that  $\kappa(\mathcal{G})$  is a deterministic quantity.

Differently from  $\kappa_3$ , a concurrency index proposed in seminal studies [12, 13] (also see for [15] a review),  $\kappa(\mathcal{G})$  is not affected by the degree distribution of the aggregate network G. It should be noted that the calculation of  $\kappa(e_i, e_i)$  and hence  $\kappa(\mathcal{G})$  requires the information about the aggregate network, i.e.,  $\mathcal{S}$ .

#### 3.2. Model 1

To calculate the concurrency index for the three models of temporal networks, we first derive the probability of an arbitrary edge being active in the equilibrium for each model. In model 1, consider an arbitrary edge in the static network G. The mean duration for which the edge is active and that for which the edge is inactive are given by

$$\langle \tau_1 \rangle = \int_0^\infty \psi_1(\tau_1) d\tau_1 \tag{3.5}$$

and

$$\langle \tau_2 \rangle = \int_0^\infty \psi_2(\tau_2) d\tau_2, \tag{3.6}$$

respectively. Owing to the renewal reward theorem [34], the probability that the edge is active in the equilibrium, denoted by  $q^*$ , is given by

$$q^* = \frac{\langle \tau_1 \rangle}{\langle \tau_1 \rangle + \langle \tau_2 \rangle}. (3.7)$$

The concurrency index,  $\kappa(\mathcal{G})$ , which we simply refer to as  $\kappa$  in the following text, is equal to the ratio of the time for which the two edges sharing a node are both active to the time for which an edge is active. Because the states of different edges are independent of each other, we obtain

$$\kappa = \frac{q^{*2}}{q^*} = q^* = \frac{\langle \tau_1 \rangle}{\langle \tau_1 \rangle + \langle \tau_2 \rangle}.$$
 (3.8)

The special case of model 1 in which the edge activation and deactivation occur as Poisson processes is equivalent to previously proposed models [31, 32]. In this case, using  $\psi_1(\tau_1) = \lambda_1 e^{-\lambda_1 \tau_1}$  and  $\psi_2(\tau_2) = \lambda_2 e^{-\lambda_2 \tau_2}$ , we obtain

$$\kappa = \frac{\langle \tau_1 \rangle}{\langle \tau_1 \rangle + \langle \tau_2 \rangle} = \frac{\lambda_1^{-1}}{\lambda_1^{-1} + \lambda_2^{-1}} = \frac{\lambda_2}{\lambda_1 + \lambda_2}.$$
 (3.9)

#### 3.3. Model 2

To analyze model 2, let us consider a pair of neighboring nodes  $v_1$  and  $v_2$  in the static network G. Let  $p_h^*$  and  $p_\ell^*$  be the probability that an arbitrary node in G is in state h and  $\ell$  in the equilibrium, respectively. Denote by  $p_{s_1s_2}^*$  the probability that node  $v_i$  is in state  $s_i \in \{h, \ell\}$  in the equilibrium, where  $i \in \{1, 2\}$ . Because the duration of the high-activity state and that of the low-activity state of a node obey the exponential distributions with mean 1/b and 1/a, respectively, we apply the renewal reward theorem [34] to obtain

$$p_h^* = \frac{\frac{1}{b}}{\frac{1}{a} + \frac{1}{b}} = \frac{a}{a+b} \tag{3.10}$$

and

$$p_{\ell}^* = \frac{\frac{1}{a}}{\frac{1}{a} + \frac{1}{b}} = \frac{b}{a+b}.$$
 (3.11)

Because the states of different nodes are independent, we obtain

$$p_{hh}^* = (p_h^*)^2 = \frac{a^2}{(a+b)^2},$$
 (3.12)

$$p_{h\ell}^* = p_{\ell h}^* = p_h^* p_\ell^* = \frac{ab}{(a+b)^2},\tag{3.13}$$

and

$$p_{\ell\ell}^* = (p_\ell^*)^2 = \frac{b^2}{(a+b)^2}.$$
 (3.14)

Therefore, the probability that an edge is active in the equilibrium is given by

$$q^* = p_{hh}^* = \frac{a^2}{(a+b)^2}. (3.15)$$

To derive the concurrency for model 2, we consider a pair of edges sharing a node in G, denoted by  $(v_1, v_2)$  and  $(v_2, v_3)$ . Denote by  $p_{s_1 s_2 s_3}$  the probability that node  $v_i$  is in state  $s_i \in \{h, \ell\}$ , where i = 1, 2, and 3. Similar to the analysis of  $q^*$  for model 2, we obtain the following stationary probabilities:

$$p_{hhh}^* = \frac{a^3}{(a+b)^3},\tag{3.16}$$

$$p_{hh\ell}^* = p_{h\ell h}^* = p_{\ell hh}^* = \frac{a^2 b}{(a+b)^3},$$
 (3.17)

$$p_{h\ell\ell}^* = p_{\ell h\ell}^* = p_{\ell\ell h}^* = \frac{ab^2}{(a+b)^3},$$
 (3.18)

$$p_{\ell\ell\ell}^* = \frac{b^3}{(a+b)^3}. (3.19)$$

Therefore, we obtain

$$\kappa = \frac{p_{hhh}^*}{q^*} = \frac{a}{a+b} = \sqrt{q^*}.$$
 (3.20)

# 3.4. Model 3

Similarly, for model 3, we obtain

$$q^* = p_{hh}^* + p_{h\ell}^* + p_{\ell h}^* = \frac{a(a+2b)}{(a+b)^2}$$
(3.21)

and

$$\kappa = \frac{p_{hhh}^* + p_{hh\ell}^* + p_{hhh}^* + p_{\ell hh}^* + p_{\ell hh}^* + p_{\ell h\ell}^*}{q^*} = \frac{a^2 + 3ab + b^2}{(a+b)(a+2b)} = \frac{q^* + \sqrt{1-q^*}}{1 + \sqrt{1-q^*}}.$$
 (3.22)

# 3.5. Comparison among the three models

A large fluctuation of  $\mathcal{G}(t)$  over time may impact concurrency [15]. In this section, we compare the amount of concurrency between the three models.

**Proposition 1.** Model 2 is more concurrent than model 1 given that  $q^*$  is the same between the two models.

*Proof.* Because  $q^*$  is the same between the two models and  $0 < q^* < 1$ , we obtain  $\sqrt{q^*} > q^*$  using Eqs. (3.8) and (3.20). Therefore, model 2 is more concurrent than model 1.

**Proposition 2.** Model 3 is more concurrent than model 1 given that  $q^*$  is the same between the two models.

*Proof.* Because  $q^*$  is the same between the two models and  $0 < q^* < 1$ , we obtain

$$\frac{q^* + \sqrt{1 - q^*}}{1 + \sqrt{1 - q^*}} - q^* = \frac{(1 - q^*)^{\frac{3}{2}}}{1 + (1 - q^*)^{\frac{1}{2}}} > 0.$$
(3.23)

Therefore, model 3 is more concurrent than model 1.

**Proposition 3.** Given that  $q^*$  is the same between models 2 and 3,

- (i) model 3 is more concurrent than model 2 if  $0 < q^* < \frac{1}{2}$ ,
- (ii) model 3 is less concurrent than model 2 if  $\frac{1}{2} < q^* \le 1$ ,
- (iii) model 3 is equally concurrent to model 2 if  $q^* = \frac{1}{2}$ .

*Proof.* For the sake of the present proof, let  $a_2$  and  $a_3$  be the rate at which the state of a node changes from  $\ell$  to h for models 2 and 3, respectively. Likewise, let  $b_2$  and  $b_3$  be the rate at which the state of a node changes from h to  $\ell$  for models 2 and 3, respectively. By imposing that  $q^*$  is the same between the two models, we obtain

$$q^* = \frac{a_3(a_3 + 2b_3)}{(a_3 + b_3)^2} = \frac{a_2^2}{(a_2 + b_2)^2}.$$
(3.24)

Therefore, the difference in the concurrency between the two models, given by Eqs. (3.20) and (3.22), is given by

$$\frac{a_3^2 + 3a_3b_3 + b_3^2}{(a_3 + b_3)(a_3 + 2b_3)} - \frac{a_2}{a_2 + b_2} = \frac{a_3^2 + 3a_3b_3 + b_3^2}{(a_3 + b_3)(a_3 + 2b_3)} - \frac{\sqrt{a_3(a_3 + 2b_3)}}{a_3 + b_3}$$

$$= \frac{b_3^2 \left[ (b_3 - a_3)^2 - 2a_3^2 \right]}{(a_3 + b_3)(a_3 + 2b_3) \left[ a_3^2 + 3a_3b_3 + b_3^2 + (a_3 + 2b_3)\sqrt{a_3(a_3 + b_3)} \right]}. \quad (3.25)$$

Therefore, the concurrency of model 3 is larger than that of model 2 if and only if  $b_3 > (1+\sqrt{2})a_3$ , which is equivalent to  $q^* = \frac{a_3(a_3+2b_3)}{(a_3+b_3)^2} = \frac{1+2\frac{b_3}{a_3}}{\left(1+\frac{b_3}{a_3}\right)^2} < \frac{1}{2}$ .

Using Eqs. (3.8), (3.20), and (3.22), we compare the amount of concurrency for models 1, 2, and 3 in Fig. 3. We find that, when  $q^* > 1/2$ , the concurrency index for model 2 is only slightly larger than that for model 3.




Figure 3: Concurrency index,  $\kappa$ , as a function of the stationary probability that an edge is active,  $q^*$ , for models 1, 2, and 3.

# 3.6. Fluctuations in the node's degree

In this section, we establish relationships between the concurrency and the statistical fluctuation in the node's degree for each of the three models. Let  $A = (A_{ij})_{N \times N}$  be the adjacency matrix of static network G. To analyze the fluctuation in the node's degree in model 1, we define random variables by

$$x_{ij}(t) = \begin{cases} 0 & \text{if edge } (i,j) \text{ is inactive at time } t, \\ 1 & \text{if edge } (i,j) \text{ is active at time } t. \end{cases}$$
(3.26)

The adjacency matrix of the temporal network at time t,  $\mathcal{G}(t)$ , of model 1 is given by the  $N \times N$  matrix  $B(t) = (B_{ij}(t))$ , where  $B_{ij}(t) = A_{ij}x_{ij}(t)$  (with  $i, j \in \{1, ..., N\}$ ). Let  $k_i(t) \equiv \sum_{j=1}^{N} B_{ij}(t)$  be the degree of the *i*th node in network  $\mathcal{G}(t)$ . The average degree at time t is given by

$$\langle k \rangle(t) \equiv \frac{1}{N} \sum_{i=1}^{N} k_i(t). \tag{3.27}$$

In the following text, we omit t because we discuss the fluctuations in  $k_i(t)$  and  $\langle k \rangle(t)$  in the equilibrium.

**Proposition 4.** For model 1, it holds true that  $k_i \sim B(\overline{k}_i, q^*)$  and  $\langle k \rangle \sim \frac{2}{N} B(M, q^*)$ , where  $B(\cdot, \cdot)$  represents the binomial distribution. (We remind that  $\overline{k}_i$  is the degree of the ith node in G and that M is the number of edges in G.)

*Proof.* For a proposition C, define the indicator function by

$$\mathbb{1}(C) = \begin{cases} 1 & \text{if } C \text{ is true,} \\ 0 & \text{if } C \text{ is false.} \end{cases}$$
(3.28)

For a node i, we obtain

$$k_i = \sum_{j=1}^{N} B_{ij} = \sum_{j=1}^{N} \mathbb{1}(A_{ij} = 1)x_{ij}.$$
(3.29)

Because  $x_{ij}$  are independent and identically distributed Bernoulli random variables and  $k_i$  is the sum of  $\overline{k}_i$  terms,  $k_i$  obeys  $B(\overline{k}_i, q^*)$ .

Because we have assumed that the network is undirected, we obtain

$$\langle k \rangle = \frac{1}{N} \sum_{i=1}^{N} k_{i}$$

$$= \frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} B_{ij}$$

$$= \frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} \mathbb{1}(A_{ij} = 1) x_{ij}.$$
(3.30)

Because there are M terms that comprise the summation,  $\langle k \rangle$  obeys  $\frac{2}{N}B(M,q^*)$ .

We denote the expectation by  $\mathbb{E}$ , and the variance by  $\sigma^2$ ; the standard deviation is equal to  $\sigma$ . Using Proposition 4, we obtain

$$\mathbb{E}[k_i] = \overline{k}_i q^*, \tag{3.31}$$

$$\sigma^{2}[k_{i}] = \overline{k}_{i}q^{*}(1 - q^{*}), \tag{3.32}$$

$$\mathbb{E}[\langle k \rangle] = \frac{2Mq^*}{N},\tag{3.33}$$

$$\sigma^{2}[\langle k \rangle] = \frac{4Mq^{*}(1-q^{*})}{N^{2}}.$$
(3.34)

To analyze models 2 and 3, we define

$$y_i(t) = \begin{cases} 0 & \text{if node } i \text{ is in the } \ell \text{ state at time } t, \\ 1 & \text{if node } i \text{ is in the } h \text{ state at time } t. \end{cases}$$
(3.35)

The adjacency matrix of  $\mathcal{G}(t)$  for model 2 is given by  $B(t) = (B_{ij}(t))$ , where  $B_{ij}(t) = A_{ij}y_i(t)y_j(t)$ . Using this expression, we can prove the following proposition.

Proposition 5. For model 2, we obtain

$$\mathbb{E}[k_i] = \overline{k}_i \left(\frac{a}{a+b}\right)^2,\tag{3.36}$$

$$\sigma^2[k_i] = \frac{\overline{k}_i a^2 b}{(a+b)^3} \left( 1 + \frac{\overline{k}_i a}{a+b} \right), \tag{3.37}$$

$$\mathbb{E}[\langle k \rangle] = \frac{2M}{N} \left(\frac{a}{a+b}\right)^2,\tag{3.38}$$

$$\sigma^2[\langle k \rangle] = \frac{4M}{N^2} \left(\frac{a}{a+b}\right)^2 + \frac{4M(M-1)}{N^2} \left(\frac{a}{a+b}\right)^3 - \frac{8}{N^2} \left(\frac{a}{a+b}\right)^3 \frac{b}{a+b} \sum_{i=1}^{N-1} \sum_{\substack{j=i+1 \ k \neq i}}^{N-1} \sum_{\substack{\ell > k \ k \neq j}}^{N-1} A_{ij} A_{k\ell}$$

$$-\frac{4M^2}{N^2} \left(\frac{a}{a+b}\right)^4. \tag{3.39}$$

We prove Proposition 5 in Appendix A.

The adjacency matrix of  $\mathcal{G}(t)$  for model 3 is given by  $B(t) = (B_{ij}(t))$ , where  $B_{ij} = A_{ij} [1 - (1 - y_i(t)) (1 - y_j(t))]$ . Using this expression, we can prove the following proposition.

**Proposition 6.** For model 3, we obtain

$$\mathbb{E}[k_i] = \frac{\overline{k}_i a(a+2b)}{(a+b)^2},\tag{3.40}$$

$$\sigma^{2}[k_{i}] = \frac{\overline{k}_{i}a(a+2b)}{(a+b)^{2}} + \frac{\overline{k}_{i}(\overline{k}_{i}-1)a(a^{2}+3ab+b^{2})}{(a+b)^{3}} - \left[\frac{\overline{k}_{i}a(a+2b)}{(a+b)^{2}}\right]^{2},$$
(3.41)

$$\mathbb{E}[\langle k \rangle] = \frac{2Ma(a+2b)}{N(a+b)^2},\tag{3.42}$$

$$\sigma^{2}[\langle k \rangle] = \frac{4Ma(a+2b)}{N^{2}(a+b)^{2}} + \frac{4M(M-1)a(a^{2}+3ab+b^{2})}{N^{2}(a+b)^{3}}$$

$$-\frac{8ab^3}{N^2(a+b)^4} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \sum_{\substack{k>i\\k\neq j}}^{N-1} \sum_{\substack{\ell>k\\\ell\neq i,j}}^{N} A_{ij} A_{k\ell} - \frac{4M^2a^2(a+2b)^2}{N^2(a+b)^4}.$$
 (3.43)

We prove Proposition 6 in Appendix B.

Now, let us compare the variance of  $k_i$  and  $\langle k \rangle$  among the three models under the condition that the expectation of  $k_i$  and  $\langle k \rangle$  is the same across the different models. This condition is equivalent to keeping  $q^*$  the same across the models for each edge. The purpose of examining the variance of the node's degree is the following. Similar to Eq. (3.3), the number of concurrent edge pairs at time t is given by  $\sum_{i=1}^{N} k_i(k_i-1)/2$ . Because this expression contains the second moment of the degree, we expect that the variance of the degree may be related to our concurrency measure.

In fact, we obtain the following results for the fluctuation of the degree, which are parallel to those for the concurrency index.

**Proposition 7.** Assume that  $\mathbb{E}[k_i]$  is the same between models 1 and 2 for any  $i \in \{1, ..., N\}$ . For any given  $q^*$ , the variance  $\sigma^2[k_i]$  is larger for model 2 than model 1 if  $\overline{k_i} > 1$ . Likewise,  $\sigma^2[\langle k \rangle]$  is larger for model 2 than model 1 if there exists i such that  $\overline{k_i} > 1$ .

*Proof.* We use subscripts 1 and 2 to represent the variance with respect to the probability distribution for models 1 and 2, respectively. We substitute Eq. (3.15) in Eq. (3.32) and use Eq. (3.37) to obtain

$$\sigma_{2}^{2}[k_{i}] - \sigma_{1}^{2}[k_{i}] = \frac{\overline{k}_{i}a^{2}b}{(a+b)^{3}} \left(1 + \frac{\overline{k}_{i}a}{a+b}\right) - \overline{k}_{i}q^{*}(1-q^{*})$$

$$= \frac{\overline{k}_{i}a^{2}b}{(a+b)^{3}} \left(1 + \frac{\overline{k}_{i}a}{a+b}\right) - \overline{k}_{i} \left(\frac{a}{a+b}\right)^{2} \left[1 - \left(\frac{a}{a+b}\right)^{2}\right]$$

$$= \overline{k}_{i}(\overline{k}_{i}-1) \frac{a^{3}b}{(a+b)^{4}}$$

$$> 0. \tag{3.44}$$

Next, we compare the variance of the average degree. Because M is the number of edges of static network G and there exists i such that  $\overline{k}_i > 1$  in G, we obtain

$$\sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \sum_{\substack{k>i\\k\neq j}}^{N-1} \sum_{\substack{\ell>k\\\ell\neq i,j}}^{N} A_{ij} A_{k\ell} < \frac{M(M-1)}{2}$$
(3.45)

for the following reason. The right-hand side of Eq. (3.45) is the number of pairs of edges. The left-hand side is the number of pairs of edges that do not share a node. These two quantities are equal to each other if and only if there is no pair of edges sharing a node, i.e., when all nodes have the degree at most 1.

By substituting Eq. (3.15) in Eq. (3.34) and using Eqs. (3.39) and (3.45), we obtain

$$\begin{split} &\sigma_{2}^{2}\left[\langle k\rangle\right] - \sigma_{1}^{2}\left[\langle k\rangle\right] \\ &= \frac{4M}{N^{2}}\left(\frac{a}{a+b}\right)^{2} + \frac{4M(M-1)}{N^{2}}\left(\frac{a}{a+b}\right)^{3} - \frac{8}{N^{2}}\left(\frac{a}{a+b}\right)^{3}\frac{b}{a+b}\sum_{i=1}^{N-1}\sum_{j=i+1}^{N}\sum_{\substack{k>i\\k\neq j}}^{N-1}\sum_{\substack{\ell>k\\\ell\neq i,j}}^{N}A_{ij}A_{k\ell} \\ &-\left(\frac{2M}{N}\right)^{2}\left(\frac{a}{a+b}\right)^{4} - \frac{4Mq^{*}(1-q^{*})}{N^{2}} \\ &> \frac{4M}{N^{2}}\left(\frac{a}{a+b}\right)^{2} + \frac{4M(M-1)}{N^{2}}\left(\frac{a}{a+b}\right)^{3} - \frac{4M(M-1)}{N^{2}}\left(\frac{a}{a+b}\right)^{3}\frac{b}{a+b} \\ &-\left(\frac{2M}{N}\right)^{2}\left(\frac{a}{a+b}\right)^{4} - \frac{4M}{N^{2}}\left(\frac{a}{a+b}\right)^{2}\left[1 - \left(\frac{a}{a+b}\right)^{2}\right] \\ = 0. \end{split} \tag{3.46}$$

**Proposition 8.** Assume that  $\mathbb{E}[k_i]$  is the same between models 1 and 3 for any  $i \in \{1, \ldots, N\}$ . For any given  $q^*$ , the variance  $\sigma^2[k_i]$  is larger for model 3 than model 1 if  $\overline{k}_i > 1$ . Likewise,  $\sigma^2[\langle k \rangle]$  is larger for model 3 than model 1 if there exists i such that  $\overline{k}_i > 1$ .

*Proof.* We substitute Eq. (3.21) in Eq. (3.32) and use Eq. (3.41) to obtain

$$\sigma_{3}^{2}[k_{i}] - \sigma_{1}^{2}[k_{i}]$$

$$= \overline{k}_{i} \frac{a(a+2b)}{(a+b)^{2}} + \overline{k}_{i}(\overline{k}_{i}-1) \frac{a(a^{2}+3ab+b^{2})}{(a+b)^{3}} - \left[\overline{k}_{i} \frac{a(a+2b)}{(a+b)^{2}}\right]^{2} - \overline{k}_{i}q^{*}(1-q^{*})$$

$$= \overline{k}_{i}(\overline{k}_{i}-1) \frac{ab^{3}}{(a+b)^{4}}$$

$$> 0. \tag{3.47}$$

By substituting Eq. (3.21) in Eq. (3.34) and using Eqs. (3.43) and (3.45), we obtain

$$\sigma_{3}^{2} [\langle k \rangle] - \sigma_{1}^{2} [\langle k \rangle]$$

$$= \frac{4M}{N^{2}} \cdot \frac{a(a+2b)}{(a+b)^{2}} + \frac{4M(M-1)}{N^{2}} \cdot \frac{a(a^{2}+3ab+b^{2})}{(a+b)^{3}}$$

$$- \frac{8}{N^{2}} \cdot \frac{ab^{3}}{(a+b)^{4}} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \sum_{\substack{k>i\\k\neq j}}^{N-1} \sum_{\substack{\ell>k\\\ell\neq i,j}}^{N} A_{ij} A_{k\ell} - \left(\frac{2M}{N}\right)^{2} \frac{a^{2}(a+2b)^{2}}{(a+b)^{4}} - \frac{4Mq^{*}(1-q^{*})}{N^{2}}$$

$$> \frac{4M}{N^{2}} \cdot \frac{a(a+2b)}{(a+b)^{2}} + \frac{4M(M-1)}{N^{2}} \cdot \frac{a(a^{2}+3ab+b^{2})}{(a+b)^{3}}$$

$$- \frac{4M(M-1)}{N^{2}} \cdot \frac{ab^{3}}{(a+b)^{4}} - \left(\frac{2M}{N}\right)^{2} \frac{a^{2}(a+2b)^{2}}{(a+b)^{4}} - \frac{4M}{N^{2}} \frac{ab^{2}(a+2b)}{(a+b)^{4}}$$

$$= 0.$$

$$(3.48)$$

**Proposition 9.** Assume that  $\mathbb{E}[k_i]$  is the same between models 2 and 3 for any  $i \in \{1, \ldots, N\}$ . For any given  $q^*$ , if  $\overline{k}_i > 1$ , we obtain

(i) 
$$\sigma_3^2[k_i] > \sigma_2^2[k_i]$$
 if  $0 < q^* < \frac{1}{2}$ ,  
(ii)  $\sigma_3^2[k_i] < \sigma_2^2[k_i]$  if  $\frac{1}{2} < q^* \le 1$ ,

11

(iii) 
$$\sigma_3^2[k_i] = \sigma_2^2[k_i]$$
 if  $q^* = \frac{1}{2}$ .

Furthermore,  $\sigma^2[\langle k \rangle]$  satisfies the same relationships if there exists i such that  $\overline{k}_i > 1$ .

We prove Proposition 9 in Appendix C.

# 3.7. Duration for the edge being inactive in model 2

In model 2, an edge is active if and only if both nodes forming the edge are in the h state, and the state of each node (i.e., h or  $\ell$ ) independently obeys a continuous-time Markov process with two states. Therefore, the duration of the edge being active obeys an exponential distribution with rate 2b. In contrast, the duration of the edge being inactive does not obey an exponential distribution, which we characterize as follows.

**Proposition 10.** The probability density function of the duration of the edge being inactive in model 2 is the mixture of two exponential distributions given by

$$f(t) = C_1 \lambda_1 e^{-\lambda_1 t} + C_2 \lambda_2 e^{-\lambda_2 t}, \tag{3.49}$$

where

$$\lambda_1 = \frac{1}{2} \left( 3a + b - \sqrt{a^2 + 6ab + b^2} \right), \tag{3.50}$$

$$\lambda_2 = \frac{1}{2} \left( 3a + b + \sqrt{a^2 + 6ab + b^2} \right), \tag{3.51}$$

$$C_1 = \frac{a\left(1 + \frac{a-b}{\sqrt{a^2 + 6ab + b^2}}\right)}{3a + b - \sqrt{a^2 + 6ab + b^2}},\tag{3.52}$$

and

$$C_2 = \frac{a\left(1 - \frac{a-b}{\sqrt{a^2 + 6ab + b^2}}\right)}{3a + b + \sqrt{a^2 + 6ab + b^2}}.$$
(3.53)

Note that it is straightforward to check  $C_1 + C_2 = 1$ ,  $C_1 > 0$ ,  $C_2 > 0$ , and  $\lambda_1 > 0$ .

*Proof.* Consider a three-state continuous-time Markov process described as follows. Let z (with z=0,1, or 2) denote the number of nodes forming an edge that are in the h state. We refer to the value of z as the state of the three-state Markov chain without arising confusion with the single node's state (i.e., h or  $\ell$ ) or edge's state (i.e., active or inactive). We initialize the stochastic dynamics of nodes by setting z=2, which corresponds to the edge being active. Consider a sequence of the state z that starts from z=2 at time 0 and return to z=2 for the first time. Let  $I_n$  be such a sequence of the z values visiting z=0 in total z=0 in total z=0 the first time, which we denote by

$$I_n = (2, 1, \underbrace{0, 1, ..., 0, 1}_{n \text{ repetitions of 0 and 1}}, 2).$$

The duration of the edge being inactive is the difference between the time of the first passage to z=2 and the time of leaving z=2 last time. We denote this duration by  $T_n$  for the case in which z=0 is visited n times before z=2 is revisited for the first time.

For general n, we obtain

$$T_n = \tau'_{1,1} + \tau''_{0,1} + \tau'_{1,2} + \dots + \tau''_{0,n} + \tau'_{1,n+1}, \tag{3.54}$$

where  $\tau'_{1,i}$  is the *i*th duration of z=1, which obeys the exponential distribution with rate a+b;  $\tau''_{0,i}$  is the *i*th duration of z=0, which obeys the exponential distribution with rate 2a. Variables  $\tau'_{1,1}, \ldots, \tau'_{1,n+1}, \tau''_{0,1}, \ldots, \tau''_{0,n}$  are independent of each other. Therefore, the Laplace transform of the distribution of  $T_n$  is given by

$$\mathcal{L}_n(s) = \mathbb{E}[e^{-sT_n}]$$

$$= \mathbb{E}[e^{-s\tau'_{1,1}}]\mathbb{E}[e^{-s\tau'_{0,1}}]\mathbb{E}[e^{-s\tau'_{1,2}}]\mathbb{E}[e^{-s\tau'_{0,2}}]\cdots\mathbb{E}[e^{-s\tau''_{0,n}}]\mathbb{E}[e^{-s\tau'_{1,n+1}}]$$

$$= \left(\frac{a+b}{s+a+b}\right)^{n+1} \left(\frac{2a}{s+2a}\right)^{n}.$$
(3.55)

The probability that sequence  $I_n$  occurs is given by

$$\tilde{q}(n) = 1 \cdot \left(\frac{b}{a+b}\right)^n \cdot 1^n \cdot \frac{a}{a+b} = \frac{ab^n}{(a+b)^{n+1}}.$$
 (3.56)

Let T be the duration for which the edge is inactive. Using Eqs. (3.55) and (3.56), we obtain the Laplace transform of the distribution of T as follows:

$$\mathcal{L}(s) = \sum_{n=0}^{\infty} \tilde{q}(n) \mathbb{E}[e^{-sT_n}]$$

$$= \sum_{n=0}^{\infty} \frac{ab^n}{(a+b)^{n+1}} \left(\frac{a+b}{s+a+b}\right)^{n+1} \left(\frac{2a}{s+2a}\right)^n$$

$$= \frac{a(s+2a)}{s^2 + (3a+b)s + 2a^2}$$


$$= \frac{a\left(s + \frac{3a+b}{2} + \frac{a-b}{2}\right)}{\left(s + \frac{3a+b}{2}\right)^2 - \left(\sqrt{\frac{a^2 + 6ab + b^2}{4}}\right)^2}.$$
(3.57)


Therefore,

$$\mathcal{L}^{-1}(s) = a \left[ e^{-\frac{3a+b}{2}t} \cosh\left(\sqrt{\frac{a^2+6ab+b^2}{4}}t\right) + \frac{a-b}{\sqrt{a^2+6ab+b^2}} e^{-\frac{3a+b}{2}t} \sinh\left(\sqrt{\frac{a^2+6ab+b^2}{4}}t\right) \right]$$

$$= C_1 \lambda_1 e^{-\lambda_1 t} + C_2 \lambda_2 e^{-\lambda_2 t}. \tag{3.58}$$

We verify Eq. (3.49) with numerical simulations for two parameter sets. The results are shown in Fig. 4. The dashed curves represent the exponential distributions whose mean is the same as that of the corresponding mixture of two exponential distributions. The figure suggests that the actual duration for the edge to be inactive is distributed more heterogeneously than the exponential distribution for both parameter sets.





**Figure 4:** Distributions of the duration of the edge being inactive in model 2. The shaded bars represent numerically obtained distributions calculated on the basis of  $5 \times 10^5$  samples. The solid lines represent the mixture of two exponential distributions, i.e., Eq. (3.49). The dashed lines represent the exponential distribution whose mean is the same as that for Eq. (3.49). (a) a = 2.0, b = 1.0. (b) a = 1.0, b = 2.0.

For model 3, the duration for the edge being inactive, which is equivalent to the time for which the same three-state Markov process stays in state z = 0, obeys an exponential distribution with rate 2a. The duration for the edge being active is the first passage time to z = 0 since the Markov chain has left z = 0. Therefore, the duration for the edge being active for model 3 obeys the mixture of two exponential distributions given by Eq. (3.49), but with a and b being swapped.

#### 4. Impact of concurrency on the dynamics of the SIR model

#### 4.1. Numerical results

# 4.1.1. Methods

To examine the effect of concurrency on epidemic spreading, we run the stochastic SIR model on three static networks. On each static network, we run stochastic dynamics of partnership according to model 1, 2, or 3 and compare the extent of epidemic spreading among the models. Each node takes either the susceptible, infectious, or recovered state, and the node's state may change over time. An infectious node infects each of its susceptible neighbor independently at rate  $\beta$ , which we call the infection rate. An infectious node also recovers at rate  $\mu$ , which we call the recovery rate, independently of the other nodes' states. The infection and recovery events occur as Poisson processes with the respective rates.

We consider three static networks. First, we use the Erdős-Rényi (ER) random graph with N=200 nodes. We independently connect each pair of nodes with probability 0.05. We iterated generating a network from the ER random graph until we obtained a connected network with M=1000 edges. The second network is a network with N=200 nodes that has a heterogeneous degree distribution generated by the Barabási-Albert (BA) model [35]. We assume that each incoming node has five edges to be connected to already existing nodes according to the preferential attachment rule. The initial network is a star graph on 6 nodes. The network is connected and contains M=975 edges. Third, we use the largest connected component of a collaboration network among researchers who had published papers in network science up to 2006 [36]. The network has N=379 nodes and M=914 edges. The edge represents the presence of at least one paper that two authors have coauthored. We use this network as an unweighted network.

We set  $\mu = 1$  without loss of generality; multiplying the same constant to  $\beta$ ,  $\mu$ ,  $a_1$ ,  $b_1$ ,  $a_2$ ,  $b_2$ ,  $a_3$ , and  $b_3$  only rescales the time;  $a_i$  and  $b_i$  represent parameters a and b for model i. We run 5000 simulations for each static network, each partnership model (i.e., model 1, 2, or 3), and each parameter set. In each simulation, just one node selected uniformly at random is initially infectious, and all the other nodes are susceptible. For each static network, the initially infectious node is the same over the different partnership models and the different  $\beta$  values. For each  $\beta$  value, we average the fraction of recovered nodes at the end of the simulation over the 5000 simulations, which we call the final epidemic size. We run the simulation using the Laplace Gillespie algorithm [37, 38], which is an extension of the direct method of Gillespie to the case in which inter-event times are allowed to obey a mixture of exponential distributions like Eq. (3.49).

Apart from the variation in the  $\beta$  value, we consider four sets of parameter values. In the first set of simulations, we set  $a_1 = 1$ ,  $b_1 = 9$ ,  $a_2 = 0.5(\sqrt{10} + 1) \approx 2.08$ ,  $b_2 = 4.5$ ,  $a_3 = 0.5$ , and  $b_3 = 1.5(\sqrt{10} + 3) \approx 9.24$ . Then, we obtain  $q^* = 0.1$  for all the three partnership models. In this manner, we compare the final epidemic size for the three partnership models, which yield different amounts of concurrency, under the condition that each edge is active for the same amount of time on average. It should be noted that a large  $q^*$  value will obviously lead to a larger final epidemic size with the other things being equal, and so it is necessary to compare the models with the  $q^*$  value being fixed. We obtain a second parameter set by making the values of  $a_1, b_1, a_2, b_2, a_3$ , and  $b_3$  for the first parameter set five times smaller, which implies that the nodes and edges flip their states five times more slowly than in the case of the first parameter set. Because multiplying these six parameters by the same constant does not change  $q^*$ , we retain  $q^* = 0.1$  for the second parameter set. We refer to the first and second parameter sets as the fast and slow edge dynamics, respectively. The third parameter set is defined by  $a_1 = 1$ ,  $b_1 = 1$ ,  $a_2 = 0.5(\sqrt{2} + 1) \approx 1.21$ ,  $b_2 = 0.5$ ,  $a_3 = 0.5$ , and  $b_3 = 0.5(\sqrt{2} + 1) \approx 1.21$  such that  $q^* = 0.5$ . We also consider a slower variant of edge dynamics by dividing these  $a_1, b_1, a_2, b_2, a_3$ , and  $c_3$  values by five. In summary, for each of the three networks and each partnership model, we have four cases each of which consists of the combination of  $q^* \in \{0.1, 0.5\}$  and either the fast or slow edge dynamics.

# 4.1.2. Results

We first compare between partnership models 1 and 2. We recall that model 2 has a higher concurrency than model 1 when each edge is active with the same probability in the two models. To exclude the possible effects of the distribution of the duration for which the edge is active and that for which the edge is inactive, we consider model 1 with  $\psi_1(\tau_1) = 2b_2e^{-2b_2\tau_1}$  and  $\psi_2(\tau_2) = C_1\lambda_1e^{-\lambda_1\tau_2} + C_2\lambda_2e^{-\lambda_2\tau_2}$ , where  $\lambda_1, \lambda_2, C_1$ , and  $C_2$  are given by Eqs. (3.50), (3.51), (3.52), and (3.53), respectively, with  $a = a_2$  and  $b = b_2$ . In this manner, models 1 and 2 have the identical distribution of the duration of the edge being active (i.e.,  $\psi_1$ ) and that of the edge being inactive (i.e.,  $\psi_2$ ), whereas they are different in terms of the amount of concurrency.

In Figs. 5(a)-5(d), we show the relationships between the infection rate and final epidemic size for a network generated by the ER random graph. Figures 5(a) and 5(b) correspond to  $q^* = 0.1$ ; Figs. 5(c) and 5(d) correspond to  $q^* = 0.5$ . Figures 5(a) and 5(c) correspond to the slow edge dynamics; Figs. 5(b) and 5(d) correspond to the fast edge dynamics. Figures 5(a)-5(d) indicate that, in each of these combinations of a value of  $q^*$  and speed of the edge dynamics, the final epidemic size is larger for model 2 than model 1 when the infection rate is near the epidemic threshold. This result suggests that concurrency promotes epidemic spreading near the epidemic threshold. In contrast, the concurrency suppresses the epidemic spreading when the infection rate, and hence the final epidemic size, is larger in each of the four cases. We also find that, at small infection rates including near the epidemic threshold, model 2 yields only marginally larger final epidemic sizes than model 1 when  $q^* = 0.5$  (see Figs. 5(c) and 5(d)).

We find that the dependence of the final epidemic size on the infection rate for the BA model network, shown in Figs. 5(e)-5(h), is similar to that for the ER random graph.

We show the results for the collaboration network in Figs. 5(i)–5(l). When  $q^* = 0.1$ , the final epidemic size is larger for model 2 than model 1 across the entire range of the infection rate,  $\beta$ , that we investigated. Because the final epidemic size has approximately plateaued for both models 1 and 2 at the largest  $\beta$  value that we investigated, it is unlikely that model 1 yields a larger final epidemic size than model 2 when  $\beta$  is larger. This result is qualitatively different from that for the ER and BA models and indicates that the effect of concurrency on enhancing epidemic spreading is stronger for the collaboration network than for the ER and BA models. In contrast, when  $q^* = 0.5$ , the final epidemic size is only marginally larger for model 2 than model 1 when  $\beta$  is near the epidemic threshold and larger for model 1 than for model 2 when  $\beta$  is larger. This result is qualitatively the same as that for the ER and BA models.




Figure 5: Comparison of the final epidemic size between models 1 and 2. (a)–(d) ER random graph. (e)–(h) BA model. (i)–(l) Collaboration network. In panels (a), (e), and (i), we set  $q^* = 0.1$  and use the slow edge dynamics. In panels (b), (f), and (j), we set  $q^* = 0.1$  and use the fast edge dynamics. In panels (c), (g), and (k), we set  $q^* = 0.5$  and use the slow edge dynamics. In panels (d), (h), and (l), we set  $q^* = 0.5$  and use the fast edge dynamics. "Collabo." is a shorthand for the collaboration (i.e., co-authorship) network. The inset of (i) shows the magnification of the main panel because the final epidemic size is small in the entire range of the infection rate in this case.

Now we compare models 1 and 3. We use model 1 with  $\psi_1(\tau_1) = C_1\lambda_1e^{-\lambda_1\tau_1} + C_2\lambda_2e^{-\lambda_2\tau_1}$ , where the four parameters are given by Eqs. (3.50), (3.51), (3.52), and (3.53), respectively, with  $a=b_3$  and  $b=a_3$ , and  $\psi_2(\tau_2)=2a_3e^{-2a_3\tau_2}$ . In this manner, models 1 and 3 have the identical  $\psi_1$  and  $\psi_2$ , whereas model 3 has a larger amount of concurrency than model 1 does. We show the final epidemic size for models 1 and 3 for the same three underlying static networks, two values of  $q^*$ , and two speeds of edge dynamics in Fig. 6. The results are largely similar to the comparison between models 1 and 2. A notable difference is that model 3 yields a larger final epidemic size than model 1 across a wider range of the infection rate values when  $q^*=0.1$  and the edge dynamics is slow (see Figs. 6(a) and 6(e)) than in the comparison between models 1 and 2.




Figure 6: Comparison of the final epidemic size between models 1 and 3. (a)–(d) ER random graph. (e)–(h) BA model. (i)–(l) Collaboration network. In panels (a), (e), and (i), we set  $q^* = 0.1$  and use the slow edge dynamics. In panels (b), (f), and (j), we set  $q^* = 0.1$  and use the fast edge dynamics. In panels (c), (g), and (k), we set  $q^* = 0.5$  and use the slow edge dynamics. In panels (d), (h), and (l), we set  $q^* = 0.5$  and use the fast edge dynamics.

#### 4.2. Theoretical results

By assuming that the network at each time is generated as an i.i.d., we analytically derived an expression of the epidemic threshold for models 1, 2, and 3. We assume the susceptible-infectious-susceptible (SIS) model for facilitating theoretical analysis. For simplicity, we assume that each network composed of partnership edges lasts for time h before switching to another partnership network. Such a switching is induced by the flip of the state of an edge. We found that, to the first order of h, the epidemic threshold is the same for the three models. We show the detail in Appendix D.

# 5. Discussion

We investigated three models of edge dynamics. They allow us to generate cases with different amounts of concurrency while keeping the probability that each edge is active (i.e.,  $q^*$ ) and the structure of the aggregate network the same. We theoretically evaluated the concurrency of each model and compared it among the models. Then, we numerically observed using the stochastic SIR model that the final epidemic size is larger in the model with the higher concurrency (i.e., models 2 or 3 compared to model 1) if the infection rate is near or below the epidemic threshold, while the epidemic size is smaller when the concurrency is higher for large infection rates. Furthermore, the concurrency has a larger impact on the final epidemic size near the epidemic threshold when  $q^*$  is smaller (i.e., with  $q^* = 0.1$  than with  $q^* = 0.5$ ).

A previous study numerically showed that concurrency has a greater effect on increasing an epidemic potential in sparser networks [11]. They measured the epidemic potential defined by the proportion of ordered pairs of nodes that are reachable in the sense that it is possible to travel from one node to the other node along a time-respecting path in the given temporal network composed of time-stamped edges.

Our results are consistent with theirs because the effect of concurrency on enhancing epidemic spreading is stronger when  $q^*$  is smaller in all the cases investigated (see Figs. 5 and 6). The larger effect of concurrency on epidemic spreading for sparser networks may be because networks with large  $q^*$ , which leads to a large number of edges, have more time-respecting paths from an infectious node to susceptible nodes transmitting infection even in the absence of concurrency [11].

Another previous study concluded that the impact of concurrency on the epidemic size is fairly limited except in an early stage of epidemic dynamics [27]. Despite the dependence of epidemic spreading on the density of contacts, which we discussed in the last paragraph, our findings are at large consistent with theirs. In other words, we found that the presence of concurrency little affected or even decreased the final epidemic size in the SIR model except when the infection rate is posed near the epidemic threshold for some parameter combinations. Our analytical result showing that the epidemic threshold is independent of the concurrency (i.e., see section 4.2) is also consistent with this result.

In contrast, our previous study analytically showed that the concurrency enhances epidemic spreading in terms of the epidemic threshold value for the SIS model [28, 39]. The diversity of these results may be due to different strategies for modeling concurrency. For example, the authors of Ref. [27] designed their dynamic network models by ensuring that the number of partners that an individual has in a long term is the same across the individuals, rendering the aggregate network the complete graph. A different study that also carefully controlled the amount of interaction for each edge to be the same across comparisons made the same homogeneity assumption [29]. In contrast, Refs. [28, 39] and the present study have assumed that the number of partners that an individual has in a long term may be heterogeneously distributed. Examining similarities and differences between these existing models of concurrency, including the model proposed in the present study, is an outstanding question. This being said, an overall conclusion based on the present numerical results is that the concurrency only modestly affects the final epidemic size, except near the epidemic threshold in some cases. This result indicates that other factors such as static network structure [1–4] and burstiness [5–7, 40] may be a larger contributor to epidemic dynamics than concurrency. Further comparing the impact of concurrency on epidemic spreading and that of other network factors warrants future work.

Concurrency is not only relevant to sexually transmitted infections, as most studies of concurrency have focused on [15, 20–23, 41], or even to epidemic spreading in general. Other network dynamics such as opinion formation, synchronization, and information spreading and cascading on social networks may also be affected by concurrency. Our assumption that the partnership between any given pair of individuals can reoccur after it has disappeared is unrealistic in most cases of sexual partnerships. However, this assumption is considered to be natural for describing other types of dynamic social contacts such as face-to-face encounters and online communications, which underlie social dynamics apart from sexually transmitted infections. Investigating impacts of concurrency on these different social dynamics using the present models also warrants future work.

#### Acknowledgments

M.O. acknowledges support from JSPS KAKENHI (under Grant No. JP21H01352). N.M. acknowledges support from AFOSR European Office (under Grant No. FA9550-19-1-7024), the Nakatani Foundation, the Sumitomo Foundation, and the Japan Science and Technology Agency (JST) Moonshot R&D (under Grant No. JPMJMS2021).

#### Appendix A Proof of Proposition 5

In model 2, random variable  $y_i$  obeys a Bernoulli distribution with expected value  $\mathbb{E}[y_i] = a/(a+b)$  because  $y_i = 1$  with probability a/(a+b) and  $y_i = 0$  with probability b/(a+b). Because  $y_i$  and  $y_j$ , where  $i \neq j$ , are independent of each other, we obtain

$$\mathbb{E}[k_i] = \mathbb{E}\left[y_i \sum_{j=1}^N A_{ij} y_j\right] = \sum_{\substack{j=1\\j \neq i}}^N A_{ij} \mathbb{E}[y_i] \mathbb{E}[y_j] = \overline{k}_i \left(\frac{a}{a+b}\right)^2$$
(A.1)

and

$$\sigma^{2}[k_{i}] = \mathbb{E}[k_{i}^{2}] - (\mathbb{E}[k_{i}])^{2}$$

$$= \mathbb{E}\left[\sum_{j=1}^{N} A_{ij}^{2} y_{i}^{2} y_{j}^{2}\right] + \mathbb{E}\left[\sum_{\substack{j=1\\j\neq i}}^{N} \sum_{\substack{k=1\\k\neq i,j}}^{N} A_{ij} A_{ik} y_{i}^{2} y_{j} y_{k}\right] - (\mathbb{E}[k_{i}])^{2}$$

$$= \mathbb{E}\left[\sum_{j=1}^{N} A_{ij} y_{i} y_{j}\right] + \mathbb{E}\left[\sum_{\substack{j=1\\j\neq i}}^{N} \sum_{\substack{k=1\\k\neq i,j}}^{N} A_{ij} A_{ik} y_{i} y_{j} y_{k}\right] - (\mathbb{E}[k_{i}])^{2}$$

$$= \overline{k}_{i} \left(\frac{a}{a+b}\right)^{2} + \overline{k}_{i} (\overline{k}_{i} - 1) \left(\frac{a}{a+b}\right)^{3} - \left[\overline{k}_{i} \left(\frac{a}{a+b}\right)^{2}\right]^{2}$$

$$= \frac{\overline{k}_{i} a^{2} b}{(a+b)^{3}} \left(1 + \frac{\overline{k}_{i} a}{a+b}\right). \tag{A.2}$$

We also obtain

$$\mathbb{E}[\langle k \rangle] = \mathbb{E}\left[\frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij} y_i y_j\right]$$

$$= \frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij} \mathbb{E}[y_i] \mathbb{E}[y_j]$$

$$= \frac{2M}{N} \left(\frac{a}{a+b}\right)^2 \tag{A.3}$$

and

$$\begin{split} \sigma^{2}[\langle k \rangle] &= \mathbb{E}[\langle k \rangle^{2}] - (\mathbb{E}[\langle k \rangle])^{2} \\ &= \mathbb{E}\left[\left(\frac{2}{N}\sum_{i=1}^{N}\sum_{j=i+1}^{N}A_{ij}y_{i}y_{j}\right)^{2}\right] - \left[\frac{2M}{N}\left(\frac{a}{a+b}\right)^{2}\right]^{2} \\ &= \frac{4}{N^{2}}\mathbb{E}\left[\sum_{i=1}^{N}\sum_{j=i+1}^{N}A_{ij}^{2}y_{i}^{2}y_{j}^{2} + 2\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=j+1}^{N}A_{ij}A_{jk}y_{i}y_{j}^{2}y_{k} + 2\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=j+1}^{N}A_{ij}A_{ik}y_{i}^{2}y_{j}y_{k} \\ &+ 2\sum_{i=1}^{N}\sum_{k=i+1}^{N}\sum_{j=k+1}^{N}A_{ij}A_{kj}y_{i}y_{k}y_{j}^{2} + 2\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k>i}\sum_{\substack{\ell>k\\k\neq j}}^{N}\sum_{\substack{\ell>k\\\ell\neq i,j}}^{N}A_{ij}A_{k\ell}y_{i}y_{j}y_{k}y_{\ell} - \left(\frac{2M}{N}\right)^{2}\left(\frac{a}{a+b}\right)^{4} \\ &= \frac{4}{N^{2}}\left(\mathbb{E}\left[\sum_{i=1}^{N}\sum_{j=i+1}^{N}A_{ij}y_{i}y_{j}\right] + 2\mathbb{E}\left[\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=j+1}^{N}A_{ij}A_{jk}y_{i}y_{j}y_{k}\right] + 2\mathbb{E}\left[\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=j+1}^{N}A_{ij}A_{kj}y_{i}y_{j}y_{k}\right] \\ &+ 2\mathbb{E}\left[\sum_{i=1}^{N}\sum_{k=i+1}^{N}\sum_{j=k+1}^{N}A_{ij}A_{kj}y_{i}y_{k}y_{j}\right] + 2\mathbb{E}\left[\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=j+1}^{N}\sum_{k\neq j}^{N}\sum_{\ell\neq i,j}^{N}A_{ij}A_{k\ell}y_{i}y_{j}y_{k}y_{\ell}\right] - \left(\frac{2M}{N}\right)^{2}\left(\frac{a}{a+b}\right)^{4} \\ &= \frac{4}{N^{2}}\left[M\left(\frac{a}{a+b}\right)^{2} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{i=i+1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{jk} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{ik}\right)^{4} \right] \\ &= \frac{4}{N^{2}}\left[M\left(\frac{a}{a+b}\right)^{2} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{i=i+1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{jk} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{ik}\right)^{4} \right] \\ &= \frac{4}{N^{2}}\left[M\left(\frac{a}{a+b}\right)^{2} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{i=i+1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{jk} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{j=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{ik}\right)^{4} \right] \\ &= \frac{4}{N^{2}}\left[M\left(\frac{a}{a+b}\right)^{2} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{i=i+1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{ik} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{ik}\right)^{4} \right] \\ &= \frac{4}{N^{2}}\left[M\left(\frac{a}{a+b}\right)^{2} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{i=1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}\sum_{k=i+1}^{N}A_{ij}A_{ik}\right)^{2} \right] \\ &= \frac{4}{N^{2}}\left[M\left(\frac{a}{a+b}\right)^{2} + 2\left(\frac{a}{a+b}\right)^{3}\sum_{k=1}^{N}\sum$$

$$+2\left(\frac{a}{a+b}\right)^{3} \sum_{i=1}^{N} \sum_{k=i+1}^{N} \sum_{j=k+1}^{N} A_{ij} A_{kj} + 2\left(\frac{a}{a+b}\right)^{4} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \sum_{k>i}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} \right] - \left(\frac{2M}{N}\right)^{2} \left(\frac{a}{a+b}\right)^{4}$$

$$= \frac{4M}{N^{2}} \left(\frac{a}{a+b}\right)^{2} + \frac{8}{N^{2}} \left(\frac{a}{a+b}\right)^{3} \left(\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{jk} + \sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k>j}^{N} A_{ij} A_{ik} + \sum_{i=1}^{N} \sum_{j=k+1}^{N} A_{ij} A_{kj} \right)$$

$$+ \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \sum_{k>i}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} + \sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k>i}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} + \sum_{i=1}^{N} \sum_{j=k+1}^{N} A_{ij} A_{k\ell} + \sum_{i=1}^{N} \sum_{j=k+1}^{N} A_{ij} A_{kj} + \sum_{i=1}^{N} \sum_{j=k+1}^{N} \sum_{k>i}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} + \sum_{i=1}^{N} \sum_{j=k+1}^{N} \sum_{\ell>k}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} + \sum_{i=1}^{N} \sum_{j=k+1}^{N} \sum_{\ell>k}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} + \sum_{i=1}^{N} \sum_{\ell>k}^{N-1} \sum_{\ell>k}^{N} A_{ij} A_{k\ell} + \sum_{\ell>k}^{N} A_{ij} A_{\ell} + \sum_{$$

# Appendix B Proof of Proposition 6

For model 3, we obtain

$$\mathbb{E}[k_i] = \mathbb{E}\left[\sum_{j=1}^N A_{ij}(y_i + y_j - y_i y_j)\right]$$

$$= \sum_{\substack{j=1\\j \neq i}}^N A_{ij}(\mathbb{E}[y_i] + \mathbb{E}[y_j] - \mathbb{E}[y_i]\mathbb{E}[y_j]) = \frac{\overline{k}_i a(a+2b)}{(a+b)^2}$$
(B.1)

and

$$\sigma^{2}[k_{i}] = \mathbb{E}[k_{i}^{2}] - (\mathbb{E}[k_{i}])^{2}$$

$$= \mathbb{E}\left[\sum_{j=1}^{N} A_{ij}^{2}(y_{i} + y_{j} - y_{i}y_{j})^{2}\right] + \mathbb{E}\left[\sum_{\substack{j=1\\j\neq i}}^{N} \sum_{\substack{k=1\\k\neq i,j}}^{N} A_{ij}A_{ik}(y_{i} + y_{j}y_{k} - y_{i}y_{j}y_{k})\right] - (\mathbb{E}[k_{i}])^{2}$$

$$= \mathbb{E}\left[\sum_{j=1}^{N} A_{ij}(y_{i} + y_{j} - y_{i}y_{j})\right] + \mathbb{E}\left[\sum_{\substack{j=1\\j\neq i}}^{N} \sum_{\substack{k=1\\k\neq i,j}}^{N} A_{ij}A_{ik}(y_{i} + y_{j}y_{k} - y_{i}y_{j}y_{k})\right] - (\mathbb{E}[k_{i}])^{2}$$

$$= \frac{\overline{k}_{i}a(a+2b)}{(a+b)^{2}} + \frac{\overline{k}_{i}(\overline{k}_{i} - 1)a(a^{2} + 3ab + b^{2})}{(a+b)^{3}} - \left[\frac{\overline{k}_{i}a(a+2b)}{(a+b)^{2}}\right]^{2}.$$
(B.2)

We also obtain

$$\mathbb{E}[\langle k \rangle] = \mathbb{E}\left[\frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij} \left(y_i + y_j - y_i y_j\right)\right]$$

$$= \frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij} \left(\mathbb{E}[y_i] + \mathbb{E}[y_j] - \mathbb{E}[y_i]\mathbb{E}[y_j]\right)$$

$$= \frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij} \left[\frac{2a}{a+b} - \left(\frac{a}{a+b}\right)^2\right]$$

$$=\frac{2M}{N} \cdot \frac{a(a+2b)}{(a+b)^2}.\tag{B.3}$$

and

$$\begin{split} \sigma^{2}[\langle k \rangle] &= \mathbb{E}[\langle k \rangle^{2}] - (\mathbb{E}[\langle k \rangle])^{2} \\ &= \mathbb{E}\left[\left(\frac{2}{N} \sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij} \left(y_{i} + y_{j} - y_{i}y_{j}\right)\right)^{2}\right] - \left[\frac{2M}{N} \cdot \frac{a(a+2b)}{(a+b)^{2}}\right]^{2} \\ &= \frac{4}{N^{2}} \mathbb{E}\left[\sum_{i=1}^{N} \sum_{j=i+1}^{N} A_{ij}^{2} \left(y_{i} + y_{j} - y_{i}y_{j}\right)^{2} + 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{jk} \left(y_{i} + y_{j} - y_{i}y_{j}\right) \left(y_{k} + y_{k} - y_{i}y_{k}\right) \\ &+ 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{kk} \left(y_{i} + y_{j} - y_{i}y_{j}\right) \left(y_{k} + y_{k} - y_{k}y_{k}\right) \\ &+ 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{kk} \left(y_{i} + y_{j} - y_{i}y_{j}\right) \left(y_{k} + y_{i} - y_{k}y_{i}\right) \\ &+ 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{kk} \left(y_{i} + y_{j} - y_{i}y_{j}\right) \left(y_{k} + y_{i} - y_{k}y_{i}\right) \\ &+ 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} \left(y_{i} + y_{j} - y_{i}y_{j}\right) + 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{jk} \left(y_{j} + y_{i}y_{k} - y_{i}y_{j}y_{k}\right) \\ &+ 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{kk} \left(y_{i} + y_{j} - y_{i}y_{j}y_{k}\right) + 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} A_{ij} A_{kj} \left(y_{j} + y_{i}y_{k} - y_{i}y_{j}y_{k}\right) \\ &+ 2\sum_{i=1}^{N} \sum_{j=i+1}^{N} \sum_{k=j+1}^{N} \sum_{k=j+1}^{N$$

# Appendix C Proof of Proposition 9

Let  $a_2$  and  $a_3$  be the rate at which the state of a node changes from the  $\ell$  to the h state for models 2 and 3, respectively. Similarly, let  $b_2$  and  $b_3$  be the rate at which the state of a node changes from the h to  $\ell$  state for models 2 and 3, respectively. Because  $q^*$  is the same between models 2 and 3 by assumption, Eq. (3.24) holds true. We apply Eq. (3.24) to Eqs. (3.37) and (3.41) to obtain

$$\sigma_{3}^{2}[k_{i}] - \sigma_{2}^{2}[k_{i}]$$

$$= \overline{k}_{i} \frac{a_{3}(a_{3} + 2b_{3})}{(a_{3} + b_{3})^{2}} + \overline{k}_{i}(\overline{k}_{i} - 1) \frac{a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2})}{(a_{3} + b_{3})^{3}} - \left[\overline{k}_{i} \frac{a_{3}(a_{3} + 2b_{3})}{(a_{3} + b_{3})^{2}}\right]^{2}$$

$$- \overline{k}_{i} \left(\frac{a_{2}}{a_{2} + b_{2}}\right)^{2} - \overline{k}_{i}(\overline{k}_{i} - 1) \left(\frac{a_{2}}{a_{2} + b_{2}}\right)^{3} + \left[\overline{k}_{i} \left(\frac{a_{2}}{a_{2} + b_{2}}\right)^{2}\right]^{2}$$

$$= \overline{k}_{i}(\overline{k}_{i} - 1) \frac{a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2})}{(a_{3} + b_{3})^{3}} - \overline{k}_{i}(\overline{k}_{i} - 1) \left(\frac{a_{2}}{a_{2} + b_{2}}\right)^{3}$$

$$= \overline{k}_{i}(\overline{k}_{i} - 1) \frac{a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2})}{(a_{3} + b_{3})^{3}} - \overline{k}_{i}(\overline{k}_{i} - 1) \frac{\sqrt{a_{3}^{3}(a_{3} + 2b_{3})^{3}}}{(a_{3} + b_{3})^{3}}$$

$$= \frac{\overline{k}_{i}(\overline{k}_{i} - 1)}{(a_{3} + b_{3})^{3}} \left[a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2}) - \sqrt{a_{3}^{3}(a_{3} + 2b_{3})^{3}}\right]. \tag{C.1}$$

Therefore,  $\sigma_3^2[k_i] < \sigma_2^2[k_i]$  if and only if  $b_3 < (1+\sqrt{2})a_3$ , which is equivalent to  $\frac{1}{2} < q^* \le 1$ . By applying Eq. (3.24) to Eqs. (3.39) and (3.43) and using Eq. (3.45), we obtain

$$\begin{split} & \sigma_{2}^{2}[\langle k \rangle] - \sigma_{3}^{2}[\langle k \rangle] \\ & = \frac{4M(M-1)}{N^{2}} \left( \frac{a_{2}}{a_{2} + b_{2}} \right)^{3} - \frac{8}{N^{2}} \left( \frac{a_{2}}{a_{2} + b_{2}} \right)^{3} \frac{b_{2}}{a_{2} + b_{2}} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N-1} \sum_{k>i}^{N-1} \sum_{\substack{l > k \\ k \neq j}}^{N-1} \sum_{\substack{l < k \\ l \neq i,j}}^{N-1} A_{ij} A_{k\ell} \\ & - \frac{4M(M-1)}{N^{2}} \cdot \frac{a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2})}{(a_{3} + b_{3})^{3}} + \frac{8}{N^{2}} \cdot \frac{a_{3}b_{3}^{3}}{(a_{3} + b_{3})^{3}} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N-1} \sum_{k>i}^{N-1} \sum_{\substack{l > k \\ k \neq j}}^{N-1} \sum_{\substack{l < k \\ l \neq i,j}}^{N-1} A_{ij} A_{k\ell} \\ & - \frac{4M(M-1)}{N^{2}} \cdot \frac{a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2})}{(a_{3} + b_{3})^{3}} + \frac{8}{N^{2}} \cdot \frac{a_{3}b_{3}^{3}}{(a_{3} + b_{3})^{3}} \cdot \frac{a_{3} + b_{3} - \sqrt{a_{3}(a_{3} + 2b_{3})}}{a_{3} + b_{3}} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N-1} \sum_{k>i}^{N-1} \sum_{\substack{l > k \\ k \neq j}}^{N-1} \sum_{\substack{l < k \\ l \neq i,j}}^{N-1} A_{ij} A_{k\ell} \\ & = \frac{4M(M-1)}{N^{2}} \cdot \frac{(a_{3} + b_{3})}{(a_{3} + b_{3})} \left[ \sqrt{a_{3}^{3}(a_{3} + 2b_{3})^{3}} - a_{3}(a_{3}^{2} + 3a_{3}b_{3} + b_{3}^{2}) \right]}{(a_{3} + b_{3})^{4}} \\ & - \frac{8}{N^{2}} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \sum_{k > i} \sum_{\substack{l > k \\ k \neq j}}^{N-1} \sum_{\substack{l < k \\ l \neq i,j}}^{N-1} A_{ij} A_{k\ell} \frac{(a_{3} + b_{3})\sqrt{a_{3}^{3}(a_{3} + 2b_{3})^{3}} - a_{3}(a_{3}^{2} + 2b_{3})^{2} - a_{3}b_{3}^{2}}{(a_{3} + b_{3})^{4}} \\ & - \frac{8}{N^{2}} \left[ \frac{M(M-1)}{2} - \sum_{i=1}^{N-1} \sum_{\substack{k > i \\ k \neq j}}^{N-1} \sum_{\substack{l < k \\ l \neq i,j}}^{N-1} A_{ij} A_{k\ell} \frac{(a_{3} + b_{3})\sqrt{a_{3}^{3}(a_{3} + 2b_{3})^{3}} - a_{3}(a_{3}^{2} + 2b_{3})^{2} - a_{3}b_{3}^{2}}{(a_{3} + b_{3})^{2}} \right] \\ & = \frac{8}{N^{2}} \left[ \frac{M(M-1)}{2} - \sum_{i=1}^{N-1} \sum_{\substack{j = i + 1 \\ k \neq j}}^{N-1} \sum_{\substack{l < k \\ l \neq i,j}}^{N-1} A_{ij} A_{k\ell} \frac{(a_{3} + b_{3})\sqrt{a_{3}^{3}(a_{3} + 2b_{3})^{3}} - a_{3}(a_{3}^{2} + 2b_{3})^{2} - a_{3}b_{3}^{2}}{(a_{3} + b_{3})^{2}} \right] \right]$$

Therefore,  $\sigma_3^2[\langle k \rangle] < \sigma_2^2[\langle k \rangle]$  if and only if  $b_3 < (1 + \sqrt{2})a_3$ , which is equivalent to  $\frac{1}{2} < q^* \le 1$ .

# Appendix D Epidemic threshold when the network switches rapidly

In this section, we evaluate the epidemic threshold for models 1, 2, and 3 under the assumption that the time-independent networks at different times are independent of each other in each model.

We denote the identity and the zero matrices by I and O, respectively. A real matrix A (or a vector as its special case) is said to be nonnegative, denoted by  $A \ge 0$ , if all the entries of A are nonnegative. If all the entries of A are positive, then A is said to be positive. We say that  $A \le B$ , where A and B are of the same dimension, if  $B - A \ge 0$ . A square matrix A is said to be Metzler if all its off-diagonal entries are nonnegative [42]. If A is Metzler, it holds true that  $e^{At} \ge 0$  for all  $t \ge 0$  [42]. For a Metzler matrix A, the maximum real part of the eigenvalues of A is denoted by  $\lambda_{\max}(A)$ . For any matrix A, the spectral radius is the largest absolute value of its eigenvalues and denoted by  $\rho(A)$ .

The SIS model is a continuous-time Markov process with  $2^N$  possible states [2, 3, 43] and has a unique absorbing state in which all the N nodes are susceptible. Because this absorbing state is reachable from any other state, the dynamics of the SIS model reaches the disease-free absorbing equilibrium in finite time with probability one.

#### D.1 A lower bound on the decay rate for subgraphs

We refer to the directed edges of a given network  $\overline{\mathcal{G}}$  as  $\overline{\mathcal{E}} = \{e_1, \dots, e_M\}$ , where the  $\ell$ th edge  $(1 \leq \ell \leq M)$  is represented by  $e_{\ell} = (i_{\ell}, j_{\ell})$ , i.e., the edge is directed from node  $v_{i_{\ell}}$  to node  $v_{j_{\ell}}$ . Define the incidence matrix  $\overline{C} \in \mathbb{R}^{N \times M}$  of the network  $\overline{\mathcal{G}}$  by [1, 44]

$$\overline{C}_{i\ell} = \begin{cases}
1, & \text{if } j_{\ell} = i, \\
-1, & \text{if } i_{\ell} = i, \\
0, & \text{otherwise,} 
\end{cases}$$
(D.1)

for all  $i \in \{1, ..., N\}$  and  $\ell \in \{1, ..., M\}$ . We also define the non-backtracking matrix  $\overline{H} \in \mathbb{R}^{M \times M}$  of  $\overline{\mathcal{G}}$  by [45, 46]

$$\overline{H}_{\ell m} = \begin{cases} 1, & \text{if } j_{\ell} = i_m \text{ and } j_m \neq i_{\ell}, \\ 0, & \text{otherwise.} \end{cases}$$
 (D.2)

We prove the following corollary about the SIS spreading processes taking place in a subgraph of  $\mathcal{G}$ .

Corollary 11. Let  $\check{\mathcal{E}}$  be a subset of  $\overline{\mathcal{E}}$ . Consider the SIS model over the network  $\check{\mathcal{G}} = (\mathcal{V}, \check{\mathcal{E}})$  with the infection rate  $\beta > 0$  and recovery rate  $\mu > 0$ . Define matrix  $\check{H} \in \mathbb{R}^{M \times M}$  by

$$\check{H}_{\ell m} = \begin{cases} 1, & \text{if } e_{\ell} \in \check{\mathcal{E}}, \ j_{\ell} = i_{m}, \ \text{and } j_{m} \neq i_{\ell}, \\ 0, & \text{otherwise}, \end{cases}$$
(D.3)

We also define the diagonal matrix

$$\Xi = diag(\xi_1, \dots, \xi_M),$$
(D.4)

where

$$\check{\xi}_{\ell} = \begin{cases}
1, & if \ e_{\ell} \in \check{\mathcal{E}}, \\
0, & otherwise
\end{cases}$$
(D.5)

for each  $\ell \in \{1, ..., M\}$ . Then, we obtain

$$\frac{d}{dt} \begin{bmatrix} \boldsymbol{p} \\ \boldsymbol{q} \end{bmatrix} \le \check{\mathcal{A}} \begin{bmatrix} \boldsymbol{p} \\ \boldsymbol{q} \end{bmatrix}, \tag{D.6}$$

where  $\mathbf{p}(t) = [p_1(t), \dots, p_N(t)]^\top$ ,  $p_i(t) = \mathbb{E}[x_i(t)]$  is the probability that node  $v_i$  is infectious at time t,  $\top$  represents the transposition,  $\mathbf{q}(t) = [q_{i_1j_1}(t), \dots, q_{i_Mj_M}(t)]^\top$ ,  $q_{ij}(t) = \mathbb{E}[x_i(t)(1-x_j(t))]$  is the joint probability

that  $v_i$  is infectious and node  $v_j$  is susceptible at time t, and the  $(N+M)\times(N+M)$  matrix  $\check{\mathcal{A}}$  is defined by

$$\breve{\mathcal{A}} = \begin{bmatrix} -\mu I & \beta \overline{C}_{+} \breve{\Xi} \\ \mu \overline{C}_{-}^{\top} & \beta \breve{H}^{\top} - \beta \breve{\Xi} - 2\mu I \end{bmatrix},$$
(D.7)

where  $\overline{C}_{+} = \max(\overline{C}, 0)$  and  $\overline{C}_{-} = \max(-\overline{C}, 0)$  denote the positive and negative parts of the incidence matrix  $\overline{C}_{+}$  respectively.

Proof. We adapt Eq. (3.20) in Ref. [47] to the case in which the network is defined by  $\check{\mathcal{G}}$ , the infection rate is independent of edges, and the recovery rate is independent of nodes. Specifically, by replacing  $B', D, D'_1, D'_2$  in Eq. (3.20) in Ref. [47] by  $\beta \check{\Xi}, \mu I, \mu I, \mu I$ , respectively, where I is the identity matrix, we obtain

$$\frac{d}{dt} \begin{bmatrix} \mathbf{p} \\ \mathbf{q} \end{bmatrix} = \mathcal{A} \begin{bmatrix} \mathbf{p} \\ \mathbf{q} \end{bmatrix} - \begin{bmatrix} \mathbf{0} \\ \boldsymbol{\epsilon} \end{bmatrix}, \tag{D.8}$$

where  $\epsilon(t)$  is entry-wise nonnegative for every  $t \geq 0$ . Equation (D.8) implies

$$\frac{d}{dt} \begin{bmatrix} \mathbf{p} \\ \mathbf{q} \end{bmatrix} \le \mathcal{A} \begin{bmatrix} \mathbf{p} \\ \mathbf{q} \end{bmatrix}, \tag{D.9}$$

where matrix  $\mathcal{A}$  is given by

$$\mathcal{A} = \begin{bmatrix} -\mu I & \beta \overline{C}_{+} \check{\Xi} \\ \mu \overline{C}_{-}^{\top} & \beta \overline{H}^{\top} \check{\Xi} - \beta \check{\Xi} - 2\mu I \end{bmatrix}. \tag{D.10}$$

Therefore, to complete the proof, it is sufficient to show

$$\overline{H}^{\mathsf{T}} \check{\Xi} = \check{H}^{\mathsf{T}},\tag{D.11}$$

which proves  $A = \check{A}$ .

Let us show Eq. (D.11). For any  $\ell, m \in \{1, ..., M\}$ , one obtains

$$[\overline{H}^{\top} \check{\Xi}]_{\ell m} = \sum_{k=1}^{M} [\overline{H}^{\top}]_{\ell k} [\check{\Xi}]_{k m}$$

$$= [\overline{H}^{\top}]_{\ell m} [\check{\Xi}]_{m m}$$

$$= \overline{H}_{m \ell} \check{\xi}_{m}$$

$$= \begin{cases} 1, & \text{if } e_{m} \in \check{\mathcal{E}}, j_{m} = i_{\ell}, \text{ and } j_{\ell} \neq i_{m}, \\ 0, & \text{otherwise,} \end{cases}$$

$$= [\check{H}^{\top}]_{\ell m}, \qquad (D.12)$$

which proves (D.11).

D.2 A lower bound on the decay rate for temporal networks

Let  $\tilde{\mathcal{G}}_1 = (\mathcal{V}, \tilde{\mathcal{E}}_1), \ldots, \tilde{\mathcal{G}}_L = (\mathcal{V}, \tilde{\mathcal{E}}_L)$  be directed and unweighted networks having the common node set  $\mathcal{V} = \{v_1, \ldots, v_N\}$ . Let  $\{\sigma_k\}_{k=0}^{\infty}$  be independent and identically distributed random variables following a probability distribution on the set  $\{1, \ldots, L\}$ . Variable  $\sigma_k$  indexes the kth network to be used. Let k > 0 be arbitrary. For a real number x, let  $\lfloor x \rfloor$  denote the maximum integer that does not exceed x. Define the stochastic temporal network  $\mathcal{G}$  by

$$\mathcal{G}(t) = \tilde{\mathcal{G}}_{\sigma_{\lfloor t/h \rfloor}} \tag{D.13}$$

for all  $t \geq 0$ . Equation (D.13) implies that  $\mathcal{G}(kh+\tau) = \tilde{\mathcal{G}}_{\sigma_k}$  for all  $k \geq 0$  and  $\tau \in [0,h)$ . In the temporal network models 1, 2, and 3 used in the main text, the states of individual edges or nodes independently repeat flipping in continuous time in a Poissonian manner, which induces network switching. Therefore, the duration for a single time-independent network, h, is not a constant. Furthermore, the time-independent

networks before and after a single flipping of an edge's or node's state are not independent of each other. For these two reasons, the present  $\mathcal{G}$  is not the same as models 1, 2, or 3. However, one can enforce the present  $\mathcal{G}$  and any of models 1, 2, and 3 to have the same probability that each type of time-independent network,  $\tilde{\mathcal{G}}_{\ell}$ , occurs by setting an appropriate probability distribution for  $\{\tilde{\mathcal{G}}_1,\ldots,\tilde{\mathcal{G}}_L\}$  for the present  $\mathcal{G}$ . We assume such a probability distribution in the following text. We consider the stochastic SIS model taking place in  $\mathcal{G}$ .

**Definition 12.** The decay rate of the SIS model over  $\mathcal{G}$  is defined by

$$\gamma = -\limsup_{t \to \infty} \frac{\log \sum_{i=1}^{N} \mathbb{E}[p_i(t)]}{t},$$
(D.14)

where all nodes are assumed to be infected at t = 0.

Definition 12 states that  $\sum_{i=1}^{N} p_i(t)$ , which is equal to the expected number of infected nodes at time t, roughly decays exponentially in time in proportion to  $e^{-\gamma t}$ . Because the number of infected nodes always becomes zero in finite time, the SIS model on networks always has a positive decay rate. In fact, exact computation of the decay rate is computationally demanding because the decay rate is equal to the modulus of the largest real part of the non-zero eigenvalues of a  $2^N \times 2^N$  matrix representing the infinitesimal generator of the Markov chain [43]. Therefore, bounds of the decay rate that only require computation of much smaller matrices are available [43, 47–51]. Here we develop such a bound for temporal network  $\mathcal{G}$ .

Let us label the set of time-aggregated edges,  $\overline{\mathcal{E}} = \bigcup_{\ell=1}^L \tilde{\mathcal{E}}_\ell$ , as  $\overline{\mathcal{E}} = \{e_1, \dots, e_M\}$ . Define the time-aggregated graph  $\overline{\mathcal{G}} = (\mathcal{V}, \overline{\mathcal{E}})$ . Let  $\overline{\mathcal{C}} \in \mathbb{R}^{N \times M}$  denote the incidence matrix of  $\overline{\mathcal{G}}$ . Although the notations  $\overline{\mathcal{G}}$ ,  $\overline{\mathcal{E}}$ , and  $\overline{\mathcal{C}}$  are common to those used in section D.1, this should not arise confusions in the following text. For each  $t \geq 0$ , let  $\mathcal{E}(t)$  denote the set of edges of the network at time t, i.e.,  $\mathcal{G}(t)$ . In other words,  $\mathcal{E}(t) = \tilde{\mathcal{E}}_{\sigma_{\lfloor t/h \rfloor}}$ . We also define the matrix  $H(t) \in \mathbb{R}^{M \times M}$  by

$$H(t)_{\ell m} = \begin{cases} 1 & \text{if } e_{\ell} \in \mathcal{E}(t), j_{\ell} = i_{m}, \text{ and } j_{m} \neq i_{\ell}, \\ 0 & \text{otherwise,} \end{cases}$$
 (D.15)

where  $\ell, m \in \{1, \dots, M\}$ , and the diagonal matrix

$$\Xi(t) = \operatorname{diag}(\xi_1(t), \dots, \xi_M(t)), \tag{D.16}$$

where

$$\xi_{\ell}(t) = \begin{cases} 1 & \text{if } e_{\ell} \in \mathcal{E}(t), \\ 0 & \text{otherwise,} \end{cases}$$
 (D.17)

and  $\ell \in \{1, ..., M\}$ . Note that H(t) is similar to but different from the non-backtracking matrix of network  $\mathcal{G}(t)$  because the definition of H(t) does not require  $e_m \in \mathcal{E}(t)$ .

The following proposition gives an upper bound on the decay rate of the SIS model over the temporal network  $\mathcal{G}$ .

**Proposition 13.** Define an  $(N+M) \times (N+M)$  random matrix  $\mathcal{A}(t)$  by

$$\mathcal{A}(t) = \begin{bmatrix} -\mu I & \beta \overline{C}_{+} \Xi(t) \\ \mu \overline{C}_{-}^{\top} & \beta H(t)^{\top} - \beta \Xi(t) - 2\mu I \end{bmatrix}.$$
 (D.18)

Let

$$\mathcal{F} = \mathbb{E}[e^{h\mathcal{A}(0)}]. \tag{D.19}$$

Then, the decay rate of the SIS model over the temporal network  $\mathcal{G}$  is greater than or equal to  $-h^{-1}\log\rho(\mathcal{F})$ .

*Proof.* From Corollary 11, we obtain

$$\frac{d}{dt} \begin{bmatrix} \boldsymbol{p}(t) \\ \boldsymbol{q}(t) \end{bmatrix} \le \mathcal{A}(t) \begin{bmatrix} \boldsymbol{p}(t) \\ \boldsymbol{q}(t) \end{bmatrix}. \tag{D.20}$$

By combining Eq. (D.20) and the definition of the temporal network given by (D.13), we obtain

$$\frac{d}{d\tau} \begin{bmatrix} \boldsymbol{p}(kh+\tau) \\ \boldsymbol{q}(kh+\tau) \end{bmatrix} \le \mathcal{A}(kh) \begin{bmatrix} \boldsymbol{p}(\tau) \\ \boldsymbol{q}(\tau) \end{bmatrix}$$
(D.21)

for all  $k = 0, 1, \ldots$  and  $\tau \in [0, h)$ , which implies

$$\begin{bmatrix} \boldsymbol{p}((k+1)h) \\ \boldsymbol{q}((k+1)h) \end{bmatrix} \le e^{h\mathcal{A}(kh)} \begin{bmatrix} \boldsymbol{p}(kh) \\ \boldsymbol{q}(kh) \end{bmatrix}.$$
(D.22)

Because the random variables  $\{\sigma_k\}_{k=0}^{\infty}$  are independently and identically distributed, we take the expectation with respect to  $\sigma$  in Eq. (D.22) to obtain

$$\begin{bmatrix}
\mathbb{E}[\boldsymbol{p}((k+1)h)] \\
\mathbb{E}[\boldsymbol{q}((k+1)h)]
\end{bmatrix} \le \mathcal{F}\begin{bmatrix}
\mathbb{E}[\boldsymbol{p}(kh)] \\
\mathbb{E}[\boldsymbol{q}(kh)]
\end{bmatrix},$$
(D.23)

which implies

$$\begin{bmatrix} \mathbb{E}[\boldsymbol{p}(kh)] \\ \mathbb{E}[\boldsymbol{q}(kh)] \end{bmatrix} \le \mathcal{F}^k \begin{bmatrix} \boldsymbol{p}(0) \\ \boldsymbol{q}(0) \end{bmatrix}. \tag{D.24}$$

Now, let  $\mathcal{S} \subset \mathbb{R}^{(N+M)\times (N+M)}$  denote the support of the random matrix  $\mathcal{A}(0)$ . Inequality (D.21) shows that, for each sample path, there exists a matrix  $X_k \in \mathcal{S}$  dependent on k such that

$$\begin{bmatrix} \boldsymbol{p}(kh+\tau) \\ \boldsymbol{q}(kh+\tau) \end{bmatrix} \le e^{\tau X_k} \begin{bmatrix} \boldsymbol{p}(kh) \\ \boldsymbol{q}(kh) \end{bmatrix}$$
(D.25)

for all  $k = 0, 1, \ldots$  and  $\tau \in [0, h)$ . Because S is finite, the maximum  $\Gamma = \max_{0 \le \tau \le h, X \in S} \|e^{\tau X}\|$  exists and is finite. Therefore, Eq. (D.25) implies

$$\left\| \begin{bmatrix} \boldsymbol{p}(kh+\tau) \\ \boldsymbol{q}(kh+\tau) \end{bmatrix} \right\| \le \left\| e^{\tau X_k} \right\| \left\| \begin{bmatrix} \boldsymbol{p}(kh) \\ \boldsymbol{q}(kh) \end{bmatrix} \right\| \le \Gamma \left\| \begin{bmatrix} \boldsymbol{p}(kh) \\ \boldsymbol{q}(kh) \end{bmatrix} \right\|$$
(D.26)

with probability one. By combining Eqs. (D.24) and (D.26), we obtain

$$\|\mathbb{E}[\boldsymbol{p}(kh+\tau)]\| \le \Gamma \|\mathcal{F}^k\| \left\| \begin{bmatrix} \boldsymbol{p}(0) \\ \boldsymbol{q}(0) \end{bmatrix} \right\|. \tag{D.27}$$

Equation (D.27) implies that

$$\|\mathbb{E}[\boldsymbol{p}(kh+\tau)]\| \le c\|\mathcal{F}^k\| \tag{D.28}$$

for a constant c > 0. Because

$$\limsup_{t \to \infty} \frac{\log \sum_{i=1}^{N} \mathbb{E}[p_{i}(t)]}{t} \leq \limsup_{t \to \infty} \frac{\log \|\mathbb{E}[\boldsymbol{p}(t)]\|}{t}$$

$$= \lim_{k \to \infty} \sup_{\ell \geq k} \max_{0 \leq \tau < h} \frac{\log \|\mathbb{E}[\boldsymbol{p}(\ell h + \tau)]\|}{\ell h + \tau}$$

$$\leq \frac{1}{h} \lim_{k \to \infty} \sup_{\ell > k} \max_{0 \leq \tau < h} \frac{\log \|\mathbb{E}[\boldsymbol{p}(\ell h + \tau)]\|}{\ell}, \tag{D.29}$$

inequality (D.28) implies that

$$\limsup_{t \to \infty} \frac{\log \sum_{i=1}^{N} \mathbb{E}[p_i(t)]}{t} \le \frac{1}{h} \lim_{k \to \infty} \sup_{\ell \ge k} \frac{\log \|\mathcal{F}^{\ell}\| + \log c}{\ell}$$
$$= \frac{1}{h} \limsup_{k \to \infty} \frac{\log \|\mathcal{F}^{k}\|}{k}$$

$$= \frac{1}{h} \log \rho(\mathcal{F}). \tag{D.30}$$

In the last equality in Eq. (D.30), we used the Gelfand's formula [52, Corollary 5.6.14].

# D.3 Epidemic threshold for small h

In this section, we consider the case in which the switching interval, h, is sufficiently small. The first-order expansion of matrix  $\mathcal{F}$  with respect to h yields

$$\mathcal{F} = I + \mathbb{E}[h\mathcal{A}(0)] + O(h^2) = I + h\mathbb{E}[\mathcal{A}(0)] + O(h^2). \tag{D.31}$$

Let us assume that matrix  $\mathbb{E}[\mathcal{A}(0)]$  is irreducible. Let  $\lambda_{\text{max}}$  denote the real eigenvalue of  $\mathbb{E}[\mathcal{A}(0)]$  with the largest real part. Then, by the Perron-Frobenius theorem and the differentiability of the eigenvalues of matrices [52, Theorem 6.3.12], we obtain

$$\rho(\mathcal{F}) = 1 + h\lambda_{\text{max}} + O(h^2). \tag{D.32}$$

Therefore,  $h^{-1}\log \rho(\mathcal{F}) = \lambda_{\max} = O(h)$ . The combination of this asymptotic evaluation and Proposition 13 suggests that the epidemic threshold is the value of  $\beta/\mu$  at which  $\lambda_{\max} = 0$ .

We rewrite the epidemic threshold in terms of the spectral radius of the relevant matrices. As in the proof of Corollary 5 in Ref. [47], we rewrite the random matrix  $\mathcal{A}(0)$  as

$$\mathcal{A}(0) = R + \mathcal{P} \tag{D.33}$$

with a constant matrix

$$R = \begin{bmatrix} -\mu I & O\\ \mu \bar{C}_{-}^{\mathsf{T}} & -\beta I - 2\mu I \end{bmatrix}$$
 (D.34)

and a random matrix

$$\mathcal{P} = \beta \begin{bmatrix} O & \overline{C}_{+}\Xi(0) \\ O & I + H(0)^{\top} - \Xi(0) \end{bmatrix}. \tag{D.35}$$

Then, we obtain the decomposition

$$\mathbb{E}[\mathcal{A}(0)] = R + P,\tag{D.36}$$

where

$$P = \mathbb{E}[\mathcal{P}] = \beta \begin{bmatrix} O & \overline{C}_{+}\mathbb{E}[\Xi(0)] \\ O & I + \mathbb{E}[H(0)]^{\top} - \mathbb{E}[\Xi(0)] \end{bmatrix}. \tag{D.37}$$

Matrix R is Metzler, and all the eigenvalues of R have negative real parts. Matrix P is nonnegative because each diagonal of  $\Xi(0)$  is a  $\{0,1\}$ -valued random variable. Therefore, Theorem 2.11 in Ref. [53] implies that  $\lambda_{\max} < 0$  if and only if  $\rho(R^{-1}P) < 1$ . Because

$$R^{-1}P = \begin{bmatrix} O & -\frac{\beta}{\mu}\overline{C}_{+}\mathbb{E}[\Xi(0)] \\ O & -\frac{\beta}{\beta+2\mu}\left(I + \mathbb{E}[H(0)]^{\top} - \mathbb{E}[\Xi(0)] + \overline{C}_{-}^{\top}\overline{C}_{+}\mathbb{E}[\Xi(0)]\right) \end{bmatrix}, \tag{D.38}$$

one obtains

$$\rho(R^{-1}P) = \frac{\beta}{\beta + 2\mu} \rho \left( I + \mathbb{E}[H(0)]^{\top} - \mathbb{E}[\Xi(0)] + \overline{C}_{-}^{\top} \overline{C}_{+} \mathbb{E}[\Xi(0)] \right). \tag{D.39}$$

Equation (D.39) implies that  $\rho(R^{-1}P) < 1$  if and only if

$$\frac{\beta}{\mu} < \frac{2}{\rho \left( I - \mathbb{E}[\Xi(0)] + \mathbb{E}\left[\Xi(0)\right] \overline{C}_{+}^{\top} \overline{C}_{-} + \mathbb{E}\left[H(0)\right] \right) - 1}.$$
 (D.40)

The right-hand side of Eq. (D.40) gives the epidemic threshold, which we can further as follows. Equations (D.15), (D.16), and (D.17) imply that  $H(t) = \Xi(t)\overline{H}$ , where  $\overline{H}$  is the non-backtracking matrix of the time-

aggregated graph. (Note that this notation should not cause confusions with the definition of  $\overline{H}$  given by Eq. (D.2).) Therefore, we rewrite Eq. (D.40) as

$$\frac{\beta}{\mu} < \frac{2}{\rho \left( I - \mathbb{E}[\Xi(0)] + \overline{C}_{+}^{\mathsf{T}} \mathbb{E}[\Xi(0)] \overline{C}_{-} + \mathbb{E}[\Xi(0)] \overline{H} \right) - 1}.$$
 (D.41)

Therefore, the epidemic threshold, i.e., the right-hand side of Eq. (D.41), only depends on the expectation of  $\Xi(0)$ , i.e.,  $\mathbb{E}[\Xi(0)]$ . We keep  $\mathbb{E}[\Xi(0)]$  constant across the different models to inspect the effect of different amounts of concurrency on epidemic spreading under the condition that the aggregate network is the same across the comparisons. Therefore, with this analysis, we do not find a difference in the epidemic threshold across our different models.

#### References

- [1] M. E. J. Newman, (2018). Networks (2nd ed.). Oxford, UK: Oxford University Press.
- [2] R. Pastor-Satorras, C. Castellano, P. V. Mieghem, and A. Vespignani, Epidemic processes in complex networks, Rev. Mod. Phys. (2015) 87: 925–979.
- [3] I. Z. Kiss, J. C. Miller, P. L. Simon, (2017). *Mathematics of Epidemics on Networks*. Cham, Switzerland: Springer.
- [4] A. Barrat, M. Barthélemy, A. Vespignani, (2008). Dynamical Processes on Complex Networks. Cambridge, UK: Cambridge University Press.
- [5] P. Holme and J. Saramäki, Temporal networks, Phys. Rep. (2012) 519: 97–125.
- [6] P. Holme, Modern temporal network theory: A colloquium, Eur. Phys. J. B (2015) 88: 234.
- [7] N. Masuda and R. Lambiotte, (2020). A Guide to Temporal Networks (2nd ed.). Singapore: World Scientific.
- [8] S. Bansal, J. Read, B. Pourbohloul, and L. A. Meyers, The dynamic nature of contact networks in infectious disease epidemiology, J. Biol. Dyn. (2010) 4: 478–489.
- [9] N. Masuda and P. Holme, Predicting and controlling infectious disease epidemics using temporal networks, F1000Prime Rep. (2013) 5: 6.
- [10] J. Moody, The importance of relationship timing for diffusion, Social Forces (2002) 81: 25–56.
- [11] J. Moody and R. A. Benton, Interdependent effects of cohesion and concurrency for epidemic potential, Ann. Epidemiol. (2016) 26: 241–248.
- [12] M. Morris and M. Kretzschmar, Concurrent partnerships and transmission dynamics in networks, Soc. Netw. (1995) 17: 299–318.
- [13] M. Kretzschmar and M. Morris, Measures of concurrency in networks and the spread of infectious disease, Math. Biosci. (1996) 133: 165–195.
- [14] S. M. Goodreau, A decade of modelling research yields considerable evidence for the importance of concurrency: A response to Sawers and Stillwaggon, J. Intl. AIDS Soc. (2011) 14: 12.
- [15] N. Masuda, J. C. Miller, and P. Holme, Concurrency measures in the era of temporal network epidemiology: A review, J. R. Soc. Interface. (2021) 18: 20210019.
- [16] E. Lee, J. Moody, P. J. Mucha, Exploring concurrency and reachability in the presence of high temporal resolution, In: *Temporal Network Theory* (2019). P. Holme and J. Saramäki, editors, Cham, Switzerland: Springer, pp. 129–145.

- [17] M. Morris, H. Epstein, M. Wawer, Timing is everything: International variations in historical sexual partnership concurrency and HIV prevalence, PLoS ONE (2010) 5: e14092.
- [18] M. Morris and M. Kretzschmar, Concurrent partnerships and the spread of HIV, AIDS. (1997) 11: 641–648.
- [19] C. H. Watts and R. M. May, Concurrent partnerships and transmission dynamics in networks, Math. Biosci. (1992) 108: 89–104.
- [20] S. O. Aral, Partner concurrency and the STD/HIV epidemic. Curr. Infect. Dis. Rep. (2010) 12: 134–139.
- [21] M. Kretzschmar, M. Caraël, Is concurrency driving HIV transmission in sub-Saharan African sexual networks? The significance of sexual partnership typology, AIDS Behav. (2012) 16: 1746–1752.
- [22] M. N. Lurie, S. Rosenthal, Concurrent partnerships as a driver of the HIV epidemic in sub-Saharan Africa? The evidence is limited, AIDS Behav. (2010) 14: 17–24.
- [23] B. Foxman, M. Newman, B. Percha, K. K. Holmes, S. O. Aral, Measures of sexual partnerships: lengths, gaps, overlaps, and sexually transmitted infection, Sex. Transm. Dis. (2006) 33: 209–214.
- [24] I. A. Doherty, S. Shiboski, J. M. Ellen, A. A. Adimora, and N. S. Padian, Sexual bridging socially and over time: a simulation model exploring the relative effects of mixing and concurrency on viral sexually transmitted infection transmission, Sex. Transm. Dis. (2006) 33: 368–373.
- [25] K. Gurski and K. Hoffman, Influence of concurrency, partner choice, and viral suppression on racial disparity in the prevalence of HIV infected women, Math. Biosci. (2016) 282: 91–108.
- [26] K. T. D. Eames and M. J. Keeling, Monogamous networks and the spread of sexually transmitted diseases, Math. Biosci. (2004) 189: 115–130.
- [27] J. C. Miller and A. C. Slim, Saturation effects and the concurrency hypothesis: Insights from an analytic model. PLoS ONE (2017) 12: e0187938.
- [28] T. Onaga, J. P. Gleeson, and N. Masuda, Concurrency-induced transitions in epidemic dynamics on temporal networks, Phys. Rev. Lett. (2017) 119: 108301.
- [29] C. Bauch, D. A. Rand, A moment closure model for sexually transmitted disease transmission through a concurrent partnership network, Proc. R. Soc. Lond. B (2000) 267: 2019–2027.
- [30] R. Pastor-Satorras, A. Vespignani, Epidemic spreading in scale-free networks, Phys. Rev. Lett. (2001) 86: 3200–3203.
- [31] X. Zhang, C. Moore, M. E. J. Newman, Random graph models for dynamic networks, Eur. Phys. J. B (2017) 90: 200.
- [32] A. E. Clementi, C. Macci, A. Monti, F. Pasquale, R. Silvestri, Flooding time in edge-Markovian dynamic graphs, In: 27th ACM Symposium on Principles of Distributed Computing (2008) pp. 213–222.
- [33] E. Fonseca dos Reis, A. Li, N. Masuda, Generative models of simultaneously heavy-tailed distributions of inter-event times on nodes and edges, Phys. Rev. E (2020) 102: 052303.
- [34] S. M. Ross, (1995). Stochastic Processes (2nd ed.). Hoboken, NJ: Wiley.
- [35] A.-L. Barabási, R. Albert, Emergence of scaling in random networks, Science (1999) 286: 509–512.
- [36] M. E. J. Newman, Finding community structure in networks using the eigenvectors of matrices, Phys. Rev. E (2006) 74: 036104.
- [37] N. Masuda, L. E. C. Rocha, A Gillespie algorithm for non-markovain stochastic processes, SIAM Rev. (2018) 60: 95–115.

- [38] N. Masuda, C. L. Vestergaard, Gillespie algorithms for stochastic multiagent dynamics in populations and network, (2021) arXiv:2112.05293.
- [39] T. Onaga, J. P. Gleeson, and N. Masuda, The effect of concurrency on epidemic threshold in time-varying networks, In: *Temporal Network Theory* (2019). P. Holme and J. Saramäki, editors, Cham, Switzerland: Springer, pp. 253–267.
- [40] M. Karsai, H. Jo, K. Kaski, (2018). Bursty Human Dynamics. Cham, Switzerland: Springer.
- [41] L. Sawers, Measuring and modelling concurrency, J. Int. AIDS Soc. (2013) 16: 17431.
- [42] L. Farina, S. Rinaldi, (2000). Positive Linear Systems Theory and Applications. New York, NY: John Wiley & Sons, Inc..
- [43] P. Van Mieghem, J. Omic, R. Kooij, Virus spread in networks, IEEE/ACM Transactions on Networking (2009) 17: 1–14.
- [44] R. J. Wilson, (2010) Introduction to Graph Theory (5th ed.). Harlow, UK: Prentice Hall.
- [45] K. Hashimoto, Zeta functions of finite graphs and representations of p-adic groups, Adv. Stud. Pure Math. (1989) 15: 211–280.
- [46] N. Alon, I. Benjamini, E. Lubetzky, S. Sodin, Non-backtracking random walks mix faster, Comm. Contemp. Math. (2007) 9: 585–603.
- [47] N. Masuda, M. Ogura, Analysis of the susceptible-infected-susceptible epidemic dynamics in networks via the non-backtracking matrix, IMA J. Appl. Math. (2020) 85: 214-230
- [48] A. Ganesh, L. Massoulié, D. Towsley, The effect of network topology on the spread of epidemics, In: Proc. IEEE 24th Annual Joint Conference of the IEEE Computer and Communications Societies (INFOCOM'05) (2005) pp. 1455–1466.
- [49] D. Chakrabarti, Y. Wang, C. Wang, J. Leskovec, C. Faloutsos, Epidemic thresholds in real networks, ACM Trans. Info. Syst. Security (2008) 10: 1–26.
- [50] V. M. Preciado, M. Zargham, C. Enyioha, A. Jadbabaie, G. J. Pappas, Optimal resource allocation for network protection against spreading processes, IEEE Trans. Control Netw. Syst. (2014) 1: 99–108.
- [51] M. Ogura, V. M. Preciado, Second-order moment-closure for tighter epidemic thresholds, Syst. Control Lett. (2018) 113: 59–64.
- [52] R. Horn and C. Johnson, (2012). *Matrix Analysis* (2nd ed.). Cambridge, UK: Cambridge University Press.
- [53] T. Damm, D. Hinrichsen, Newton's method for concave operators with resolvent positive derivatives in ordered Banach spaces, Lin. Alg. its Appl. (2003) 363: 43–64.