

On the global behavior of solutions of the Beltrami equations

Ruslan Salimov, Mariia Stefanchuk

Abstract. In this paper, the estimate for growth of homeomorphic solutions of the Beltrami equation at infinity is obtained, provided that the dilatation quotient has a global finite mean oscillation.

MSC 2020. 30C62+31A15

Keywords. Beltrami equations, ring Q -homeomorphisms, modulus, capacity.

1 Introduction

Let D be a domain in the complex plane \mathbb{C} , i.e., a connected and open subset of \mathbb{C} , and let $\mu : D \rightarrow \mathbb{C}$ be a measurable function with $|\mu(z)| < 1$ a.e. (almost everywhere) in D . The *Beltrami equation* is the equation of the form

$$f_{\bar{z}} = \mu(z) f_z \quad (1.1)$$

where $f_{\bar{z}} = \bar{\partial}f = (f_x + if_y)/2$, $f_z = \partial f = (f_x - if_y)/2$, $z = x + iy$, and f_x and f_y are partial derivatives of f in x and y , correspondingly. The function μ is called the *complex coefficient* and

$$K_\mu(z) = \frac{1 + |\mu(z)|}{1 - |\mu(z)|} \quad (1.2)$$

the *dilatation quotient* for the equation (1.1). The Beltrami equation (1.1) is said to be *degenerate* if $\text{ess sup } K_\mu(z) = \infty$. The existence theorem for homeomorphic $W_{\text{loc}}^{1,1}$ solutions was established to many degenerate Beltrami equations, see, e.g., related references in the recent monographs [3], [10], [7]; cf. also [6], [14] – [18].

Recall that the (*conformal*) *modulus* of a family Γ of curves γ in \mathbb{C} is the quantity

$$M(\Gamma) = \inf_{\rho \in \text{adm } \Gamma} \int_{\mathbb{C}} \rho^2(z) \, dx \, dy \quad (1.3)$$

where a Borel function $\rho : \mathbb{C} \rightarrow [0, \infty]$ is *admissible* for Γ , write $\rho \in \text{adm } \Gamma$, if

$$\int_{\gamma} \rho \, ds \geq 1 \quad \forall \gamma \in \Gamma \quad (1.4)$$

where s is a natural parameter of the length on γ .

Throughout this paper,

$$B(z_0, r) = \{z \in \mathbb{C} : |z - z_0| < r\},$$

$$S(z_0, r) = \{z \in \mathbb{C} : |z - z_0| = r\},$$

and

$$\mathbb{A}(z_0, r_1, r_2) = \{z \in \mathbb{C} : r_1 < |z - z_0| < r_2\}.$$

Let $E, F \subset \overline{\mathbb{C}}$ be arbitrary sets. Denote by $\Delta(E, F, D)$ a family of all curves $\gamma : [a, b] \rightarrow \overline{\mathbb{C}}$ joining E and F in D , i.e., $\gamma(a) \in E, \gamma(b) \in F$ and $\gamma(t) \in D$ as $t \in (a, b)$.

Here a *condenser* is a pair $\mathcal{E} = (A, C)$ where $A \subset \mathbb{C}$ is open and C is a non-empty compact set contained in A . \mathcal{E} is a *ringlike condenser* if $B = A \setminus C$ is a ring, i.e., if B is a domain whose complement $\overline{\mathbb{C}} \setminus B$ has exactly two components where $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ is the one-point compactification of \mathbb{C} . \mathcal{E} is a *bounded condenser* if A is bounded. A condenser $\mathcal{E} = (A, C)$ is said to be in a domain G if $A \subset G$.

The following lemma is immediate.

Lemma 1. *If $f : G \rightarrow \mathbb{C}$ is open and $\mathcal{E} = (A, C)$ is a condenser in G , then (fA, fC) is a condenser in fG .*

In the above situation we denote $f\mathcal{E} = (fA, fC)$.

Let $\mathcal{E} = (A, C)$ be a condenser. We set

$$\text{cap } \mathcal{E} = \text{cap } (A, C) = \inf_{u \in W_0(\mathcal{E})} \int_A |\nabla u|^2 \, dx dy$$

and call it the *capacity* of the condenser \mathcal{E} . The set $W_0(\mathcal{E}) = W_0(A, C)$ is the family of nonnegative functions $u : A \rightarrow \mathbb{R}$ such that $u \in C_0(A)$, $u(z) \geq 1$ for $z \in C$, and u is absolutely continuous on lines (ACL). In the above formula

$$|\nabla u| = \sqrt{\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2}.$$

We mention some properties of the capacity of a condenser. It was proven in ([20], Theorem 1) that

$$\text{cap } \mathcal{E} = M(\Delta(\partial A, \partial C; A \setminus C)), \quad (1.5)$$

where $\Delta(\partial A, \partial C; A \setminus C)$ denotes the set of all continuous curves joining the boundaries ∂A and ∂C in $A \setminus C$.

Moreover, the following estimate is known:

$$\text{cap } \mathcal{E} \geq \frac{4\pi}{\log \frac{m(A)}{m(C)}} \quad (1.6)$$

(see, e.g., (8.8) in [11]).

The following notion is motivated by the ring definition of Gehring for quasiconformal mappings, see, e.g., [5], introduced first in the plane, see [17], and extended later on to the space case in [13], see also Chapters 7 and 11 in [10], cf. [1], [2], [4], [12].

Given a domain D in \mathbb{C} , a (Lebesgue) measurable function $Q : D \rightarrow [0, \infty]$, $z_0 \in D$, a homeomorphism $f : D \rightarrow \overline{\mathbb{C}}$ is said to be a *ring Q -homeomorphism at the point z_0* if

$$M(f(\Delta(S_1, S_2, \mathbb{A}(z_0, r_1, r_2)))) \leq \int_{\mathbb{A}(z_0, r_1, r_2)} Q(z) \cdot \eta^2(|z - z_0|) \, dx \, dy \quad (1.7)$$

for every ring $\mathbb{A}(z_0, r_1, r_2)$ and the circles $S_i = S(z_0, r_i)$, $i = 1, 2$, where $0 < r_1 < r_2 < r_0 := \text{dist}(z_0, \partial D)$, and every measurable function $\eta : (r_1, r_2) \rightarrow [0, \infty]$ such that

$$\int_{r_1}^{r_2} \eta(r) \, dr = 1.$$

f is called a *ring Q -homeomorphism in the domain D* if f is a ring Q -homeomorphism at every point $z_0 \in D$.

The following statement was first proved in [9], Theorem 3.1, cf. also Corollary 3.1 in [19].

Proposition 1. *Let f be a homeomorphic $W_{\text{loc}}^{1,1}$ solution of the Beltrami equation (1.1). Then f is a ring Q -homeomorphism at each point $z_0 \in D$ with $Q(z) = K_\mu(z)$.*

2 GFMO functions

Similarly to [8] (cf. also [14], [16]), we say that a function $\varphi: \mathbb{C} \rightarrow \mathbb{R}$ has *global finite mean oscillation at a point* $z_0 \in \mathbb{C}$, abbr. $\varphi \in GFMO(z_0)$, if

$$\limsup_{R \rightarrow \infty} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \overline{\varphi}_R| dx dy < \infty, \quad (2.1)$$

where

$$\overline{\varphi}_R = \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} \varphi(z) dx dy$$

is the mean value of the function $\varphi(z)$ over $B(z_0, R)$, $R > 0$. Here $B(z_0, R) = \{z \in \mathbb{C}: |z - z_0| < R\}$, and condition (2.1) includes the assumption that φ is integrable in $B(z_0, R)$ for $R > 0$.

Proposition 2. *If, for some collection of numbers $\varphi_R \in \mathbb{R}$, $R \in [r_0, +\infty)$, $r_0 > 0$,*

$$\limsup_{R \rightarrow \infty} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \varphi_R| dx dy < \infty,$$

then φ has global finite mean oscillation at z_0 .

Proof. Indeed, by the triangle inequality,

$$\begin{aligned} & \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \overline{\varphi}_R| dx dy \leq \\ & \leq \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \varphi_R| dx dy + |\varphi_R - \overline{\varphi}_R| \leq \\ & \leq \frac{2}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \varphi_R| dx dy. \end{aligned}$$

Corollary 1. *If, for a point $z_0 \in \mathbb{C}$,*

$$\limsup_{R \rightarrow \infty} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \varphi(z_0)| dx dy < \infty,$$

then φ has global finite mean oscillation at z_0 .

Corollary 2. *If, for a point $z_0 \in \mathbb{C}$,*

$$\limsup_{R \rightarrow \infty} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z)| \, dxdy < \infty,$$

then φ has global finite mean oscillation at z_0 .

Lemma 2. *Let $z_0 \in \mathbb{C}$. If a nonnegative function $\varphi: \mathbb{C} \rightarrow \mathbb{R}$ has global finite mean oscillation at z_0 and φ is integrable in $B(z_0, e)$, then, for $R > e^e$,*

$$\int_{\mathbb{A}(z_0, e, R)} \frac{\varphi(z) \, dxdy}{(|z - z_0| \log |z - z_0|)^2} \leq C \cdot \log \log R,$$

where

$$C = \frac{\pi}{6}((24 + \pi^2)e^2\delta_\infty + 2\pi^2\varphi_0),$$

φ_0 is the mean value of φ over the disk $B(z_0, e)$ and

$$\delta_\infty = \delta_\infty(\varphi) = \sup_{R \in (e^e, +\infty)} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \overline{\varphi}_R| \, dxdy$$

is the maximal dispersion of φ .

Proof. Let $R > e^e$, $r_k = e^k$, $\mathbb{A}_k = \{z \in \mathbb{C} : r_k \leq |z - z_0| < r_{k+1}\}$. Clearly,

$$\delta_\infty = \sup_{R \in (e^e, +\infty)} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |\varphi(z) - \overline{\varphi}_R| \, dxdy < \infty,$$

$B_k = B(z_0, r_k)$ and let φ_k be the mean value of $\varphi(z)$ over B_k , $k = 1, 2, \dots$. Take a natural number N such that $R \in [r_N, r_{N+1})$.

Then $\mathbb{A}(z_0, e, R) \subset \Delta(R) = \bigcup_{k=1}^N \mathbb{A}_k$ and

$$I(R) = \int_{\Delta(R)} \varphi(z) \alpha(|z - z_0|) \, dxdy \leq |S_1(R)| + S_2(R),$$

$$\alpha(t) = \frac{1}{(t \log t)^2},$$

$$S_1(R) = \sum_{k=1}^N \int_{\mathbb{A}_k} (\varphi(z) - \varphi_{k+1}) \alpha(|z - z_0|) \, dxdy,$$

and

$$S_2(R) = \sum_{k=1}^N \varphi_{k+1} \int_{\mathbb{A}_k} \alpha(|z - z_0|) dx dy.$$

Since $\mathbb{A}_k \subset B_{k+1}$, $\frac{1}{|z - z_0|^2} \leq \frac{\pi e^2}{m(B_{k+1})}$ for $z \in \mathbb{A}_k$ and $\log |z - z_0| > k$ in \mathbb{A}_k , then

$$\begin{aligned} |S_1(R)| &\leq \pi e^2 \sum_{k=1}^N \frac{1}{k^2} \cdot \frac{1}{m(B_{k+1})} \int_{B_{k+1}} |\varphi(z) - \varphi_{k+1}| dx dy \leq \\ &\leq \pi e^2 \delta_\infty \sum_{k=1}^N \frac{1}{k^2} \leq \pi e^2 \delta_\infty \sum_{k=1}^\infty \frac{1}{k^2} = \frac{\pi^3 e^2 \delta_\infty}{6}. \end{aligned}$$

Now,

$$\int_{\mathbb{A}_k} \alpha(|z - z_0|) dx dy \leq \frac{1}{k^2} \int_{\mathbb{A}_k} \frac{dx dy}{|z - z_0|^2} = \frac{2\pi}{k^2}.$$

Moreover,

$$\begin{aligned} |\varphi_{k-1} - \varphi_k| &= \left| \frac{1}{m(B_{k-1})} \int_{B_{k-1}} \varphi(z) dx dy - \frac{1}{m(B_{k-1})} \int_{B_{k-1}} \varphi_k dx dy \right| \leq \\ &\leq \frac{1}{m(B_{k-1})} \int_{B_{k-1}} |\varphi(z) - \varphi_k| dx dy \leq \frac{e^2}{m(B_k)} \int_{B_k} |\varphi(z) - \varphi_k| dx dy \leq e^2 \delta_\infty, \end{aligned}$$

and by the triangle inequality, for $k \geq 1$

$$\begin{aligned} \varphi_{k+1} = |\varphi_{k+1}| &= \left| \varphi_1 + \sum_{l=2}^{k+1} (\varphi_l - \varphi_{l-1}) \right| \leq \\ &\leq |\varphi_1| + \sum_{l=2}^{k+1} |\varphi_l - \varphi_{l-1}| \leq |\varphi_1| + e^2 \delta_\infty k. \end{aligned}$$

Hence,

$$\begin{aligned} S_2(R) = |S_2(R)| &\leq 2\pi \sum_{k=1}^N \frac{\varphi_{k+1}}{k^2} \leq 2\pi \sum_{k=1}^N \frac{|\varphi_1| + e^2 \delta_\infty k}{k^2} \leq \\ &\leq 2\pi |\varphi_1| \sum_{k=1}^\infty \frac{1}{k^2} + 2\pi e^2 \delta_\infty \sum_{k=1}^N \frac{1}{k} = \end{aligned}$$

$$= \frac{\pi^3 \varphi_1}{3} + 2\pi e^2 \delta_\infty \sum_{k=1}^N \frac{1}{k}.$$

But

$$\sum_{k=2}^N \frac{1}{k} < \int_1^N \frac{dt}{t} = \log N$$

and, for $R > r_N$,

$$N = \log r_N < \log R.$$

Consequently,

$$\sum_{k=1}^N \frac{1}{k} < 1 + \log \log R$$

and thus, for $R \in (e^e, +\infty)$

$$\begin{aligned} I(R) &\leq \frac{\pi^3 e^2 \delta_\infty}{6} + \frac{\pi^3 \varphi_1}{3} + 2\pi e^2 \delta_0 (1 + \log \log R) = \\ &= \left(\frac{\pi^3 e^2 \delta_\infty + 12\pi e^2 \delta_\infty + 2\pi^3 \varphi_1}{6 \log \log R} + 2\pi e^2 \delta_\infty \right) \log \log R \leq \\ &\leq \frac{\pi}{6} ((24 + \pi^2) e^2 \delta_\infty + 2\pi^2 \varphi_1) \log \log R. \end{aligned}$$

Finally,

$$\begin{aligned} \int_{\mathbb{A}(z_0, e, R)} \frac{\varphi(z) dx dy}{(|z - z_0| \log |z - z_0|)^2} &\leq I(R) \leq \\ &\leq \frac{\pi}{6} ((24 + \pi^2) e^2 \delta_\infty + 2\pi^2 \varphi_1) \log \log R. \end{aligned}$$

3 The behavior of homeomorphic solutions of the Beltrami equations at infinity

Set

$$\begin{aligned} l_f(z_0, e) &= \min_{|z - z_0| = e} |f(z) - f(z_0)|, \\ \delta_\infty &= \delta_\infty(K_\mu, z_0) = \\ &= \sup_{R \in (e^e, +\infty)} \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} |K_\mu(z) - K_{\mu, z_0}(R)| dx dy, \\ K_{\mu, z_0}(R) &= \frac{1}{m(B(z_0, R))} \int_{B(z_0, R)} K_\mu(z) dx dy, \quad k_0 = K_{\mu, z_0}(e). \end{aligned}$$

Theorem 1. Let $\mu: \mathbb{C} \rightarrow \mathbb{C}$ be a measurable function with $|\mu(z)| < 1$ a.e. and $f: \mathbb{C} \rightarrow \mathbb{C}$ be a homeomorphic $W_{\text{loc}}^{1,1}$ solution of the Beltrami equation (1.1). If $K_\mu \in GFM O(z_0)$, $z_0 \in \mathbb{C}$, then

$$\liminf_{R \rightarrow \infty} \frac{\max_{|z-z_0|=R} |f(z) - f(z_0)|}{(\log R)^{\frac{2\pi}{C}}} \geq l_f(z_0, e), \quad (3.1)$$

where $C = \frac{\pi}{6}((24 + \pi^2)e^2\delta_\infty + 2\pi^2k_0)$.

Proof. Consider the ring $\mathbb{A}(R) = \mathbb{A}(z_0, e, R)$, with $R > e^e$. Set $\mathcal{E} = (B(z_0, R), \overline{B(z_0, e)})$. Then, by lemma 1 $f\mathcal{E} = (fB(z_0, R), \overline{fB(z_0, e)})$ is a condenser in \mathbb{C} , according (1.5),

$$\text{cap}(fB(z_0, R), \overline{fB(z_0, e)}) = M(\Delta(\partial fB(z_0, e), \partial fB(z_0, R); f\mathbb{A}(R)))$$

and, in view of the homeomorphism of f ,

$$\Delta(\partial fB(z_0, e), \partial fB(z_0, R); f\mathbb{A}(R)) = f\Delta(\partial B(z_0, e), \partial B(z_0, R); \mathbb{A}(R)).$$

By Proposition 1 f is a ring Q -homeomorphism with $Q = K_\mu(z)$

$$\text{cap}(fB(z_0, R), \overline{fB(z_0, e)}) \leq \int_{\mathbb{A}(R)} K_\mu(z) \eta^2(|z - z_0|) dx dy \quad (3.2)$$

for every measurable function $\eta: (e, R) \rightarrow [0, +\infty]$ such that

$$\int_e^R \eta(t) dt = 1.$$

Choosing in (3.2) $\eta(t) = \frac{1}{t \log t \cdot \log \log R}$, we obtain

$$\text{cap}(fB(z_0, R), \overline{fB(z_0, e)}) \leq \frac{1}{(\log \log R)^2} \cdot \int_{\mathbb{A}(R)} \frac{K_\mu(z) dx dy}{(|z - z_0| \log |z - z_0|)^2}.$$

Since $K_\mu \in GFM O(z_0)$, then by lemma 2

$$\text{cap}(fB(z_0, R), \overline{fB(z_0, e)}) \leq \frac{C}{\log \log R}, \quad (3.3)$$

where $C = \frac{\pi}{6}((24 + \pi^2)e^2\delta_\infty + 2\pi^2k_0)$. On the other hand, by (1.6), we have

$$\text{cap}(fB(z_0, R), \overline{fB(z_0, e)}) \geq \frac{4\pi}{\log \frac{m(fB(z_0, R))}{m(fB(z_0, e))}}. \quad (3.4)$$

Combining (3.3) and (3.4), we obtain

$$\frac{4\pi}{\log \frac{m(fB(z_0, R))}{m(fB(z_0, e))}} \leq \frac{C}{\log \log R}.$$

This gives

$$m(\overline{fB(z_0, e)}) \leq \frac{m(fB(z_0, R))}{(\log R)^{\frac{4\pi}{C}}}.$$

Using the inequalities

$$\begin{aligned} \pi \left(\min_{|z-z_0|=e} |f(z) - f(z_0)| \right)^2 &\leq m(\overline{fB(z_0, e)}) \leq \\ &\leq m(fB(z_0, R)) \leq \pi \left(\max_{|z-z_0|=R} |f(z) - f(z_0)| \right)^2, \end{aligned}$$

we obtain

$$\min_{|z-z_0|=e} |f(z) - f(z_0)| \leq \frac{\max_{|z-z_0|=R} |f(z) - f(z_0)|}{(\log R)^{\frac{2\pi}{C}}}. \quad (3.5)$$

Set

$$l_f(z_0, e) = \min_{|z-z_0|=e} |f(z) - f(z_0)|.$$

Passing to the lower limit as $R \rightarrow \infty$ in (3.5), we obtain relation (3.1).

References

- [1] C. Andreian Cazacu, On the length-area dilatation, Complex Var. Theory Appl. (7 – 11), **50** (2005), 765 – 776.
- [2] C. Andreian Cazacu, V. Stanciu, BMO-mappings in the plane, Topics in analysis and its applications, NATO Sci. Ser. II Math. Phys. Chem., **147** (2004), 11 – 30.
- [3] K. Astala, T. Iwaniec, G. Martin, Elliptic Partial Differential Equations and Quasiconformal Mappings in the Plane, Princeton University Press (2009).
- [4] M. Cristea, Local homeomorphisms having local ACLⁿ inverses, Complex Var. Elliptic Equ. (1), **53** (2008), 77 – 99.
- [5] F.W. Gehring, Rings and quasiconformal mappings in space, Trans. Amer. Math. Soc., **103** (1962), 353 – 393.

- [6] V. Gutlyanskii, V. Ryazanov, U. Srebro, E. Yakubov, On recent advances in the Beltrami equations, *Ukr. Mat. Visn.* (4), **7** (2010), 467 – 515 (translated into English by Springer starting from 2010).
- [7] V. Gutlyanskii, V. Ryazanov, U. Srebro and E. Yakubov, The Beltrami Equation: A Geometric Approach, *Developments in Mathematics*, Vol. 26, Springer, New York etc., 2012.
- [8] A. Ignat’ev, V. Ryazanov, Finite mean oscillation in the mapping theory, *Ukr. Mat. Visn.* (3), **2** (2005), 403 – 424 (translated by AMS till 2009 as *Ukrainian Math. Bull.*).
- [9] T. Lomako, R. Salimov., E. Sevostyanov, On equicontinuity of solutions to the Beltrami equations, *Ann. Univ. Bucharest, Ser. Math.* (2), **59** (2010), 263 – 274.
- [10] O. Martio, V. Ryazanov, U. Srebro, E. Yakubov, *Moduli in Modern Mapping Theory*, Springer Monographs in Mathematics, Springer, New York (2009).
- [11] V. Maz’ya, *Lectures on isoperimetric and isocapacitary inequalities in the theory of Sobolev spaces. Heat kernels and analysis on manifolds, graphs, and metric spaces* (Paris, 2002), 307 – 340, *Contemp. Math.*, 338, Amer. Math. Soc., Providence, RI, 2003.
- [12] V. Ryazanov, E. Sevostyanov, Equicontinuity of mappings quasiconformal in the mean, *Ann. Acad. Sci. Fenn. Math.*, **36** (2011), 231 – 244.
- [13] V. Ryazanov, E. Sevostyanov, Toward the theory of ring Q -homeomorphisms, *Israel J. Math.*, **168** (2008), 101 – 118.
- [14] Ryazanov V., Srebro U., Yakubov E., Beltrami equation and FMO functions. *Contemp. Math.* **382** (2005), 357 – 364.
- [15] V. Ryazanov, U. Srebro, E. Yakubov, On ring solutions of Beltrami equation, *J. d’Analyse Math.*, **96** (2005), 117 – 150.
- [16] Ryazanov V., Srebro U., Yakubov E., Finite mean oscillation and the Beltrami equation, *Israel J. Math.*, **153** (2006), 247 – 266.
- [17] Ryazanov V., Srebro U., Yakubov E., The Beltrami equation and ring homeomorphisms, *Ukrain. Math. Bull* (1), **4** (2007), 79 – 115 (in Russian).

- [18] V. Ryazanov, U. Srebro, E. Yakubov, On integral conditions in the mapping theory, *Ukr. Mat. Visn.*, **7** (2010), 73 – 87 (translated into English by Springer starting from 2010).
- [19] R. Salimov, On regular homeomorphisms in the plane, *Ann. Acad. Sci. Fenn. Math.*, **35** (2010), 285 – 289.
- [20] V. A. Shlyk, The equality between p -capacity and p -modulus, *Sib. Math. J.*, 34:6 (1993), 1196 – 1200.

Ruslan Salimov,
Institute of Mathematics of the National Academy of Sciences of
Ukraine,
01601, Kyiv, Tereshchenkivska st. 3;
ruslan.salimov1@gmail.com

Mariia Stefanchuk,
Institute of Mathematics of the National Academy of Sciences of
Ukraine,
01601, Kyiv, Tereshchenkivska st. 3;
stefanmv43@gmail.com