Global Gradient Estimates for Dirichlet Problems of Elliptic Operators with a BMO Anti-Symmetric Part

Sibei Yang, Dachun Yang and Wen Yuan*

Abstract. Let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain. In this article, the authors investigate (weighted) global gradient estimates for Dirichlet boundary value problems of second order elliptic equations of divergence form with an elliptic symmetric part and a BMO anti-symmetric part in Ω . More precisely, for any given $p \in (2, \infty)$, the authors prove that a weak reverse Hölder inequality with exponent p implies the global $W^{1,p}$ estimate and the global weighted $W^{1,q}$ estimate, with $q \in [2, p]$ and some Muckenhoupt weights, of solutions to Dirichlet boundary value problems. As applications, the authors establish some global gradient estimates for solutions to Dirichlet boundary value problems of second order elliptic equations of divergence form with small BMO symmetric part and small BMO anti-symmetric part, respectively, on bounded Lipschitz domains, quasi-convex domains, Reifenberg flat domains, C^1 domains, or (semi-)convex domains, in weighted Lebesgue spaces. Furthermore, as further applications, the authors obtain the global gradient estimate, respectively, in (weighted) Lorentz spaces, (Lorentz–)Morrey spaces, (Musielak–)Orlicz spaces, and variable Lebesgue spaces. Even on global gradient estimates in Lebesgue spaces, the results obtained in this article improve the known results via weakening the assumption on the coefficient matrix.

1 Introduction

It is well known that the research of global regularity estimates in various function spaces for (non-)linear elliptic equations (or systems) in non-smooth domains is one of the most interesting and important topics in partial differential equations (see, for instance, [1, 5, 12, 16, 21, 23, 25, 28, 35, 55] for the linear case, and [2, 8, 9, 17, 18, 44, 45, 63, 66] for the non-linear case). Moreover, the global regularity estimates for solutions to elliptic boundary problems depend not only on the structure of equations, the integrability of the right-hand side datum, and the properties of coefficients appearing in equations, but also on the smooth property or the geometric property of the boundary of domains (see, for instance, [1, 5, 16, 24, 25, 28, 36, 41, 44, 62]).

Motivated by [28, 42, 51, 55, 62], in this article, we study global gradient estimates in various function spaces for Dirichlet boundary value problems of second order elliptic equations of divergence form with an elliptic symmetric part and a BMO anti-symmetric part on non-smooth

²⁰²⁰ Mathematics Subject Classification. Primary 35J25; Secondary 35J15, 42B35, 42B37.

Key words and phrases. elliptic operator, Dirichlet problem, NTA domain, quasi-convex domain, weak reverse Hölder inequality, gradient estimate, Muckenhoupt weight.

This work is supported by the National Natural Science Foundation of China (Grant Nos. 11871254, 12071431, 11971058, 12071197 and 11871100) and the National Key Research and Development Program of China (Grant No. 2020YFA0712900).

^{*}Corresponding author, E-mail: wenyuan@bnu.edu.cn/November 17, 2021/Final version.

domains of \mathbb{R}^n . More precisely, let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded non-tangentially accessible domain (for short, NTA domain). For any given $p \in (2, \infty)$, using a real-variable argument, we show that a weak reverse Hölder inequality with exponent p implies the global $W^{1,p}$ estimate and the global weighted $W^{1,q}$ estimate, with $q \in [2, p]$ and some Muckenhoupt weights, of solutions to Dirichlet boundary value problems. As applications, we obtain some global gradient estimates for solutions to Dirichlet boundary value problems of second order elliptic equations of divergence form with small BMO symmetric part and small BMO anti-symmetric part, respectively, on bounded Lipschitz domains, quasi-convex domains, Reifenberg flat domains, C^1 domains, or (semi-)convex domains, in the scale of weighted Lebesgue spaces. Applying these weighted global estimates and some technique from harmonic analysis, such as properties of Muckenhoupt weights, the interpolation theorem of operators, and the extrapolation theorem, we further establish the global gradient estimate, respectively, in (weighted) Lorentz spaces, (Lorentz–)Morrey spaces, (Musielak–)Orlicz spaces, and variable Lebesgue spaces. Even on global gradient estimates in Lebesgue spaces, the results obtained in this article improve the corresponding results in [1, 16, 36, 55] via weakening the assumption on the coefficient matrix.

To describe the main results of this article, we first recall the notions of the Muckenhoupt weight class and the reverse Hölder class (see, for instance, [4, 32, 56]).

Definition 1.1. Let $q \in [1, \infty)$. A non-negative and locally integrable function ω on \mathbb{R}^n is said to belong to the *Muckenhoupt weight class* $A_q(\mathbb{R}^n)$, denoted by $\omega \in A_q(\mathbb{R}^n)$, if, when $q \in (1, \infty)$,

$$[\omega]_{A_q(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \left[\frac{1}{|B|} \int_B \omega(x) \, dx \right] \left\{ \frac{1}{|B|} \int_B [\omega(x)]^{-\frac{1}{q-1}} \, dx \right\}^{q-1} < \infty,$$

or

$$[\omega]_{A_1(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \left[\frac{1}{|B|} \int_B \omega(x) \, dx \right] \left\{ \operatorname{ess\,sup}_{y \in B} [\omega(y)]^{-1} \right\} < \infty,$$

where the suprema are taken over all balls *B* of \mathbb{R}^n .

Let $r \in (1, \infty]$. A non-negative and locally integrable function ω on \mathbb{R}^n is said to belong to the *reverse Hölder class* $RH_r(\mathbb{R}^n)$, denoted by $\omega \in RH_r(\mathbb{R}^n)$, if, when $r \in (1, \infty)$,

$$[\omega]_{RH_r(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \left\{ \frac{1}{|B|} \int_B [\omega(x)]^r \, dx \right\}^{\frac{1}{r}} \left[\frac{1}{|B|} \int_B \omega(x) \, dx \right]^{-1} < \infty,$$

or

$$[\omega]_{RH_{\infty}(\mathbb{R}^{n})} := \sup_{B \subset \mathbb{R}^{n}} \left[\operatorname{ess\,sup}_{y \in B} \omega(y) \right] \left[\frac{1}{|B|} \int_{B} \omega(x) \, dx \right]^{-1} < \infty,$$

where the suprema are taken over all balls *B* of \mathbb{R}^n .

Furthermore, we recall the definition of the so-called NTA domain introduced by Jerison and Kenig [34] (see also [40, 57]) as follows. We begin with recalling several notions. For any given $x \in \mathbb{R}^n$ and measurable subset $E \subset \mathbb{R}^n$, let dist $(x, E) := \inf\{|x - y| : y \in E\}$. Meanwhile, for any measurable subsets $E, F \subset \mathbb{R}^n$, let dist $(E, F) := \inf\{|x - y| : x \in E, y \in F\}$ and diam $(E) := \sup\{|x - y| : x, y \in E\}$. Moreover, for any given $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $B(x, r) := \{y \in \mathbb{R}^n : |y - x| < r\}$.

Definition 1.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a *domain* which means that Ω is a connected open set, and $\Omega^{\complement} := \mathbb{R}^n \setminus \Omega$. Denote by $\partial \Omega$ the *boundary* of Ω .

- (i) Then the domain Ω is said to satisfy the *interior* [resp., the *exterior*] *corkscrew condition* if there exist positive constants $R \in (0, \infty)$ and $\sigma \in (0, 1)$ such that, for any $x \in \partial \Omega$ and $r \in (0, R)$, there exists a point $x_0 \in \Omega$ [resp., $x_0 \in \Omega^{\mathbb{C}}$], depending on x, such that $B(x_0, \sigma r) \subset \Omega \cap B(x, r)$ [resp., $B(x_0, \sigma r) \subset \Omega^{\mathbb{C}} \cap B(x, r)$].
- (ii) The domain Ω is said to satisfy the *Harnack chain condition* if there exist constants $m_1 \in (1, \infty)$ and $m_2 \in (0, \infty)$ such that, for any $x_1, x_2 \in \Omega$ satisfying

$$M := \frac{|x_1 - x_2|}{\min\{\operatorname{dist}(x_1, \partial\Omega), \operatorname{dist}(x_2, \partial\Omega)\}} > 1,$$

there exists a chain $\{B_i\}_{i=1}^N$ of open Harnack balls, $B_i \subset \Omega$ for any $i \in \{1, ..., N\}$, that connects x_1 to x_2 ; namely, $x_1 \in B_1$, $x_2 \in B_N$, $B_i \cap B_{i+1} \neq \emptyset$ for any $i \in \{1, ..., N-1\}$, and, for any $i \in \{1, ..., N\}$,

$$m_1^{-1}$$
diam $(B_i) \leq$ dist $(B_i, \partial \Omega) \leq m_1$ diam (B_i) ,

where the integer N satisfies $N \le m_2 \log_2 M$.

(iii) The domain Ω is called a *non-tangentially accessible domain* (for short, NTA *domain*) if Ω satisfies the interior and the exterior corkscrew conditions, and the Harnack chain condition.

We point out that NTA domains include Lipschitz domains, Zygmund domains, quasi-spheres, and some Reifenberg flat domains as special examples (see, for instance, [34, 40, 57]).

Let $n \ge 2$ and Ω be a bounded NTA domain in \mathbb{R}^n . Assume that $p \in [1, \infty)$ and $\omega \in A_q(\mathbb{R}^n)$ with some $q \in [1, \infty)$. Recall that the *weighted Lebesgue space* $L^p_{\omega}(\Omega)$ is defined by setting

(1.1)
$$L^{p}_{\omega}(\Omega) := \left\{ f \text{ is measurable on } \Omega : \|f\|_{L^{p}_{\omega}(\Omega)} := \left[\int_{\Omega} |f(x)|^{p} \omega(x) \, dx \right]^{\frac{1}{p}} < \infty \right\}.$$

Moreover, let

(1.2)
$$L^{p}_{\omega}(\Omega; \mathbb{R}^{n}) := \left\{ \mathbf{f} := (f_{1}, \dots, f_{n}) : \text{ for any } i \in \{1, \dots, n\}, f_{i} \in L^{p}_{\omega}(\Omega) \right\}$$

and

$$\|\mathbf{f}\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)} := \sum_{i=1}^n \|f_i\|_{L^p_{\omega}(\Omega)}.$$

Denote by $W^{1,p}_{\omega}(\Omega)$ the weighted Sobolev space on Ω equipped with the norm

$$\|f\|_{W^{1,p}_{\omega}(\Omega)} := \|f\|_{L^p_{\omega}(\Omega)} + \|\nabla f\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)},$$

where ∇f denotes the *distributional gradient* of f. Furthermore, $W_{0,\omega}^{1,p}(\Omega)$ is defined to be the *closure* of $C_c^{\infty}(\Omega)$ in $W_{\omega}^{1,p}(\Omega)$, where $C_c^{\infty}(\Omega)$ denotes the set of all *infinitely differentiable functions* on Ω with compact support contained in Ω . In particular, when $\omega \equiv 1$, the weighted spaces $L_{\omega}^p(\Omega)$, $W_{\omega}^{1,p}(\Omega)$, and $W_{0,\omega}^{1,p}(\Omega)$ are denoted simply, respectively, by $L^p(\Omega)$, $W^{1,p}(\Omega)$, and $W_0^{1,p}(\Omega)$, which are just, respectively, the classical Lebesgue space and the classical Sobolev spaces.

Let Ω be a domain of \mathbb{R}^n . Denote by $L^1_{loc}(\Omega)$ the set of all locally integrable functions on Ω .

Definition 1.3. Let $\Omega \subset \mathbb{R}^n$ be a domain and $f \in L^1_{loc}(\Omega)$. Then f is said to belong to the *space* BMO(Ω) if

$$||f||_{\mathrm{BMO}(\Omega)} := \sup_{B \subset \Omega} \frac{1}{|B|} \int_{B} |f(x) - f_{B}| \, dx < \infty,$$

where the supremum is taken over all balls $B \subset \Omega$ and $f_B := \frac{1}{|B|} \int_B f(y) dy$.

For any given $x \in \Omega$, let $a(x) := \{a_{ij}(x)\}_{i,j=1}^n$ denote an $n \times n$ matrix with real-valued, bounded and measurable entries. Then *a* is said to satisfy the *uniform ellipticity condition* if there exists a positive constant $\mu_0 \in (0, 1]$ such that, for any $x \in \Omega$ and $\xi := (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$,

(1.3)
$$\mu_0 |\xi|^2 \le \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \le \mu_0^{-1} |\xi|^2.$$

Recall that the matrix $b := \{b_{ij}\}_{i,j=1}^n$ is said to be *anti-symmetric* if $b_{ij} = -b_{ji}$ for any $i, j \in \{1, ..., n\}$. Throughout this article, we *always assume* that the matrix A satisfies the following assumption.

Assumption 1.4. Let $\Omega \subset \mathbb{R}^n$ be a domain. Assume that, for any given $x \in \Omega$, A(x) is an $n \times n$ matrix satisfying that A(x) = a(x) + b(x), where the matrix $a(x) := \{a_{ij}(x)\}_{i,j=1}^n$ is real-valued, symmetric, and measurable, and satisfies the uniform ellipticity condition (1.3), and the matrix $b(x) := \{b_{ij}(x)\}_{i,j=1}^n$ is real-valued, anti-symmetric, and measurable, and satisfies $b_{ij} \in BMO(\Omega)$ for any $i, j \in \{1, ..., n\}$.

Remark 1.5. Let $\Omega \subset \mathbb{R}^n$ be a domain and A := a + b satisfy Assumption 1.4.

- (i) By the assumption that *a* satisfies (1.3), we conclude that $a \in L^{\infty}(\Omega; \mathbb{R}^{n^2})$, which, together with the facts that $L^{\infty}(\Omega) \subset BMO(\Omega)$ and $b \in BMO(\Omega; \mathbb{R}^{n^2})$, further implies that $A \in BMO(\Omega; \mathbb{R}^{n^2})$.
- (ii) Via replacing Ω by \mathbb{R}^n in Definition 1.3, we obtain the definition of the space BMO(\mathbb{R}^n). Jones [38] proved that any given function $f \in BMO(\Omega)$ admits an extension to some $\tilde{f} \in BMO(\mathbb{R}^n)$ if and only if the domain Ω is a *uniform domain* (namely, the domain satisfying the interior corkscrew condition and the Harnack chain condition). Thus, if Ω is an NTA domain, then, for any given $f \in BMO(\Omega)$, there exists an $\tilde{f} \in BMO(\mathbb{R}^n)$ such that

$$\widetilde{f}|_{\Omega} = f$$
 and $\|\widetilde{f}\|_{BMO(\mathbb{R}^n)} \le C \|f\|_{BMO(\Omega)}$

where *C* is a positive constant depending only on Ω and *n*.

(iii) By the assumptions that *a* satisfies (1.3) and *b* is anti-symmetric, we conclude that, for any $x \in \Omega$ and $\xi \in \mathbb{R}^n$,

$$(A(x)\xi) \cdot \xi = (a(x)\xi) \cdot \xi \ge \mu_0 |\xi|^2$$

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and the matrix A satisfy Assumption 1.4. Assume that $p \in (1, \infty)$, $\omega \in A_q(\mathbb{R}^n)$ with some $q \in [1, \infty)$, and $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$. Then a function u is called a *weak solution* of the following *weighted Dirichlet boundary value problem*

(1.4)
$$\begin{cases} -\operatorname{div}(A\nabla u) = \operatorname{div}(\mathbf{f}) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases}$$
 $(D)_{p,\omega}$

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

if
$$u \in W^{1,p}_{0,\omega}(\Omega)$$
 and, for any $\varphi \in C^{\infty}_{c}(\Omega)$,

(1.5)
$$\int_{\Omega} A(x)\nabla u(x) \cdot \nabla \varphi(x) \, dx = -\int_{\Omega} \mathbf{f}(x) \cdot \nabla \varphi(x) \, dx.$$

In particular, when $\omega \equiv 1$, the weighted Dirichlet problem $(D)_{p,\omega}$ is just the Dirichlet problem $(D)_{p,\omega}$. The weighted Dirichlet problem $(D)_{p,\omega}$ is said to be *uniquely solvable* if, for any given $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$, there exists a *unique* $u \in W^{1,p}_{0,\omega}(\Omega)$ such that (1.5) holds true for any $\varphi \in C^{\infty}_{c}(\Omega)$.

Let $L := -\text{div}(A\nabla)$ with the matrix A satisfying Assumption 1.4. The elliptic operator L naturally arises in the study of the elliptic equation of the form

$$(1.6) \qquad -\Delta u + \mathbf{c} \cdot \nabla u = f$$

(see, for instance, [43, 51]) and the parabolic equation

$$\frac{\partial u}{\partial t} - \Delta u + \mathbf{c} \cdot \nabla u = f,$$

where the drift term **c** satisfies $\operatorname{div}(\mathbf{c}) = 0$. By $\operatorname{div}(\mathbf{c}) = 0$, we know that $\mathbf{c} = \operatorname{div}(b)$ for some anti-symmetric tensor $b := \{b_{ij}\}_{i,j=1}^{n}$. Therefore, the equation (1.6) becomes

$$-\operatorname{div}(I-b)\nabla u = f,$$

where *I* denotes the unit matrix on \mathbb{R}^n . In particular, Seregin et al. [51] discovered that the wellknown Moser iteration works for such an elliptic operator *L*. Via the Moser iteration, Seregin et al. [51] proved the Liouville theorem and the Harnack inequality for solutions to the equation (1.6) or its parabolic case. Moreover, Li and Pipher [42] studied the boundary behavior of solutions of the equation Lu = 0 in NTA domains. Furthermore, Dong and Phan [26] investigated the mixednorm Sobolev estimate for solutions to non-stationary Stokes systems with coefficients having unbounded anti-symmetric part in cylindrical domains.

Remark 1.6. Let $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain and the matrix A := a+b satisfy Assumption 1.4. For any $u, v \in W_0^{1,2}(\Omega)$, let

$$B[u,v] := \int_{\Omega} A(x) \nabla u(x) \cdot \nabla v(x) \, dx.$$

From Remark 1.5(iii), it follows that, for any $u \in W_0^{1,2}(\Omega)$,

$$B[u,u] \ge \mu_0 \|\nabla u\|_{L^2(\Omega;\mathbb{R}^n)}^2.$$

Moreover, it was showed in [42, (2.11)], via using the div-curl lemma, that, for any $u, v \in W_0^{1,2}(\Omega)$,

$$|B[u,v]| \le C \|\nabla u\|_{L^2(\Omega;\mathbb{R}^n)} \|\nabla v\|_{L^2(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant depending only on *A* and Ω . Thus, by the Lax–Milgram theorem (see, for instance, [31, Theorem 5.8]), we conclude that the Dirichlet problem $(D)_2$ is uniquely solvable and, for any given $\mathbf{f} \in L^2(\Omega; \mathbb{R}^n)$, the weak solution $u \in W_0^{1,2}(\Omega)$ of $(D)_2$ satisfies that

$$\|\nabla u\|_{L^2(\Omega;\mathbb{R}^n)} \le \mu_0^{-1} \|\mathbf{f}\|_{L^2(\Omega;\mathbb{R}^n)},$$

where μ_0 is as in (1.3).

Moreover, via an example given by Meyers [46, Section 5] (see also [16, p. 1285]), we know that, for any $p \in (1, \infty)$ with $p \neq 2$, the Dirichlet problem $(D)_p$ may not be uniquely solvable, even when the domain Ω is smooth.

Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and the matrix A satisfy Assumption 1.4. Assume further that A satisfies the (δ, R) -BMO condition (see Definition 2.1 below) or A belongs to the space VMO(Ω) (see, for instance, [50]). In this article, our aim is to establish the weighted Calderón–Zygmund type estimates

(1.7)
$$\|\nabla u\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)}$$

for the Dirichlet problem (1.4), with $p \in (1, \infty)$ and $\omega \in A_q(\mathbb{R}^n)$ for some $q \in [1, \infty)$, and then give their applications, where *C* is a positive constant independent of *u* and **f**.

Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded domain, and the matrix A := a + b satisfy Assumption 1.4. For the Dirichlet problem $(D)_p$, the estimate (1.7) with $p \in (1, \infty)$ and $\omega \equiv 1$ was established by Di Fazio [21], under the assumptions that $a \in VMO(\mathbb{R}^n)$, $b \equiv 0$, and $\partial \Omega \in C^{1,1}$, which was weakened to $\partial \Omega \in C^1$ by Auscher and Qafsaoui [5]. Moreover, for the Dirichlet problem $(D)_p$, the estimate (1.7) with $p \in (1, \infty)$ and $\omega \equiv 1$ was obtained by Byun and Wang in [12, 16], under the assumptions that a satisfies the (δ, R) -BMO condition for sufficiently small $\delta \in (0, \infty), b \equiv 0$, and Ω is a bounded Lipschitz domain with a small Lipschitz constant or a bounded Reifenberg flat domain (see, for instance, [49, 57]). Furthermore, for the Dirichlet problem $(D)_p$ with a having partial small BMO coefficients and $b \equiv 0$, the estimate (1.7) with $p \in (1, \infty)$ and $\omega \equiv 1$ was studied, respectively, by Dong and Kim [23], and Krylov [41], under the assumption that Ω is a bounded Lipschitz domain with small Lipschitz constant. Moreover, if Ω is a bounded quasiconvex domain, a satisfies the (δ, R) -BMO condition for sufficiently small $\delta \in (0, \infty)$, and $b \equiv 0$, the estimate (1.7) with $p \in (1, \infty)$ and $\omega \equiv 1$ was established by Jia et al. [36] for the Dirichlet problem $(D)_p$. For the Dirichlet problem $(D)_p$ in a general Lipschitz domain Ω , it was proved by Shen [55] that, if $a \in VMO(\mathbb{R}^n)$ and $b \equiv 0$, then (1.7) with $\omega \equiv 1$ holds true for any given $p \in (\frac{3}{2} - \varepsilon, 3 + \varepsilon)$ when $n \ge 3$, or $p \in (\frac{4}{3} - \varepsilon, 4 + \varepsilon)$ when n = 2, where $\varepsilon \in (0, \infty)$ is a positive constant depending on Ω . It is worth pointing out that, when A := I (the identity matrix) in (1.7), the range of p obtained in [55] is even sharp for general Lipschitz domains (see, for instance, [35]).

For the weighted Dirichlet problem $(D)_{p,\omega}$ with *a* having partial small BMO coefficients and $b \equiv 0, (1.7)$ with $p \in (2, \infty)$ and $\omega \in A_{p/2}(\mathbb{R}^n)$ was obtained by Byun and Palagachev [14] under the assumption that Ω is a bounded Reifenberg flat domain. Furthermore, for the problem $(D)_{p,\omega}$ with *a* having partial small BMO coefficients and $b \equiv 0$, the estimate (1.7) with $p \in (1, \infty)$ and $\omega \in A_p(\mathbb{R}^n)$ was established by Dong and Kim [25] under the assumption that Ω is a bounded Reifenberg flat domain. For the problem $(D)_{p,\omega}$ with *a* having small BMO coefficients and $b \equiv 0$, (1.7) with $p \in (1, \infty)$ and $\omega \in A_p(\mathbb{R}^n)$ was obtained by Adimurthi et al. [1] under the assumption that Ω is a bounded Lipschitz domain with small Lipschitz constant.

Now, we state the main results of this article as follows.

Theorem 1.7. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, the matrix A satisfy Assumption 1.4, and $p \in (2, \infty)$. Assume that there exist positive constants $C_0 \in (0, \infty)$ and $r_0 \in (0, \text{diam}(\Omega))$ such that, for any ball $B(x_0, r)$ having the property that $r \in (0, r_0/4)$ and either $x_0 \in \partial \Omega$ or

 $B(x_0, 2r) \subset \Omega$, the weak reverse Hölder inequality

(1.8)
$$\left[\frac{1}{|B_{\Omega}(x_0,r)|}\int_{B_{\Omega}(x_0,r)}|\nabla v(x)|^p dx\right]^{\frac{1}{p}} \le C_0 \left[\frac{1}{|B_{\Omega}(x_0,2r)|}\int_{B_{\Omega}(x_0,2r)}|\nabla v(x)|^2 dx\right]^{\frac{1}{p}}$$

holds true for any function $v \in W^{1,2}(B_{\Omega}(x_0, 2r))$ satisfying $\operatorname{div}(A\nabla v) = 0$ in $B_{\Omega}(x_0, 2r)$, and v = 0on $B(x_0, 2r) \cap \partial\Omega$ when $x_0 \in \partial\Omega$, where $B_{\Omega}(x_0, r) := B(x_0, r) \cap \Omega$.

(i) Then the weak solution $u \in W_0^{1,2}(\Omega)$ of the Dirichlet problem $(D)_p$ with $\mathbf{f} \in L^p(\Omega; \mathbb{R}^n)$ exists and, moreover, $u \in W_0^{1,p}(\Omega)$ and there exists a positive constant *C*, depending only on *n*, *p*, and Ω , such that

(1.9)
$$\|\nabla u\|_{L^p(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p(\Omega;\mathbb{R}^n)}.$$

(ii) Let $q \in [2, p]$, $q_0 \in [1, \frac{q}{p'}]$, $r_0 \in [(\frac{p}{q})', \infty]$, and $\omega \in A_{q_0}(\mathbb{R}^n) \cap RH_{r_0}(\mathbb{R}^n)$. Then a weak solution u of the weighted Dirichlet problem $(D)_{q,\omega}$ with $\mathbf{f} \in L^q_{\omega}(\Omega; \mathbb{R}^n)$ exists and, moreover, $u \in W^{1,q}_{0,\omega}(\Omega)$ and there exists a positive constant C, depending only on n, p, q, $[\omega]_{A_{q_0}(\mathbb{R}^n)}$, $[\omega]_{RH_{r_0}(\mathbb{R}^n)}$, and Ω , such that

(1.10)
$$\|\nabla u\|_{L^q_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^q_{\omega}(\Omega;\mathbb{R}^n)}.$$

Here and thereafter, for any $s \in [1, \infty]$, s' denotes its conjugate exponent, namely, 1/s + 1/s' = 1.

We prove Theorem 1.7 via using a (weighted) real-variable argument (see Theorem 3.1 below), which was essentially established in [54, Theorem 3.4] (see also [28, 29, 53, 55, 62]) and inspired by [18, 58]. It is worth pointing out that a similar real-variable argument with the different motivation was used in [3, 4]. Moreover, a different weighted real-variable argument was obtained by Shen [52, Theorem 2.1]. Furthermore, the linear structure of second order elliptic operators of divergence form and the properties of Muckenhoupt weights are subtly used in the proof of Theorem 1.7.

Let the (δ, R) -BMO condition and the space VMO(Ω) be as in Definition 2.1 below. As an application of Theorem 1.7, we obtain the (weighted) global gradient estimates for solutions to Dirichlet boundary problems on bounded Lipschitz domains as follows.

Theorem 1.8. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, and the matrix A satisfy Assumption 1.4.

(i) Then there exist positive constants ε₀, δ₀ ∈ (0,∞), depending only on n and the Lipschitz constant of Ω, such that, for any given p ∈ ((3 + ε₀)', 3 + ε₀) when n ≥ 3, or p ∈ ((4 + ε₀)', 4 + ε₀) when n = 2, if A satisfies the (δ, R)-BMO condition for some δ ∈ (0, δ₀) and R ∈ (0,∞), or A ∈ VMO(Ω), then the Dirichlet problem (D)_p with **f** ∈ L^p(Ω; ℝⁿ) is uniquely solvable and there exists a positive constant C, depending only on n, p, and the Lipschitz constant of Ω, such that, for any weak solution u, u ∈ W₀^{1,p}(Ω) and

(1.11)
$$\|\nabla u\|_{L^p(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p(\Omega;\mathbb{R}^n)}.$$

(ii) Let ε_0 be as in (i) and $p_0 := 3 + \varepsilon_0$ when $n \ge 3$, and $p_0 := 4 + \varepsilon_0$ when n = 2. Then, for any given $p \in (p'_0, p_0)$ and any $\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)$, there exists a positive constant $\delta_0 \in (0, \infty)$, depending only on n, p, the Lipschitz constant of Ω , $[\omega]_{A_{\frac{p}{p'_0}}(\mathbb{R}^n)}$, and $[\omega]_{RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)}$, such that, if A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in$ $(0, \infty)$, or $A \in VMO(\Omega)$, then the weighted Dirichlet problem $(D)_{p,\omega}$ with $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$ is uniquely solvable and there exists a positive constant C, depending only on n, p, $[\omega]_{A_{\frac{p}{p'_0}}(\mathbb{R}^n)}$, $[\omega]_{RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)}$, and the Lipschitz constant of Ω , such that, for any weak solution u, $u \in$ $W^{1,p}_{0,\omega}(\Omega)$ and

(1.12)
$$\|\nabla u\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)}$$

Let A := a + b satisfy Assumption 1.4 and p_0 be as in Theorem 1.8(ii). The key to proving Theorem 1.8 is to show that the weak reverse Hölder inequality (1.8) is valid for any $p \in (2, p_0)$. To do this, we flexibly apply the real-variable argument established in Theorem 3.1 below, the method of perturbation, the assumptions that $b \in BMO(\Omega; \mathbb{R}^{n^2})$ and b is anti-symmetric, and the properties of Muckenhoupt weights.

Remark 1.9. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, and A := a+b satisfy Assumption 1.4. If $a \in \text{VMO}(\Omega)$ and $b \equiv 0$, then Theorem 1.8 in this case was essentially established by Shen [55, Theorem C]. Thus, Theorem 1.8 improves [55, Theorem C] via weakening the condition for the matrix A.

Let the quasi-convex domain be as in Definition 2.3 below. We further obtain the following (weighted) global gradient estimates for the Dirichlet problems on quasi-convex domains by using Theorem 1.7.

Theorem 1.10. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and $p \in (1, \infty)$. Assume further that the matrix A satisfies Assumption 1.4, and Ω is a (δ, σ, R) quasi-convex domain with some $\delta, \sigma \in (0, 1)$ and $R \in (0, \infty)$.

(i) Then there exists a positive constant δ₀ ∈ (0, 1), depending only on n, p, and Ω, such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R)-BMO condition for some δ ∈ (0, δ₀), σ ∈ (0, 1), and R ∈ (0,∞), or A ∈ VMO(Ω), then the Dirichlet problem (D)_p with **f** ∈ L^p(Ω; ℝⁿ) is uniquely solvable and, for any weak solution u of the problem (D)_p, u ∈ W₀^{1,p}(Ω) and

 $\|\nabla u\|_{L^p(\Omega;\mathbb{R}^n)} \leq C \|\mathbf{f}\|_{L^p(\Omega;\mathbb{R}^n)},$

where C is a positive constant depending only on n, p, and Ω .

(ii) Let $\omega \in A_p(\mathbb{R}^n)$. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, Ω , and $[\omega]_{A_p(\mathbb{R}^n)}$, such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then the weighted Dirichlet problem $(D)_{p,\omega}$ with $\mathbf{f} \in L^{\omega}_{\omega}(\Omega; \mathbb{R}^n)$ is uniquely solvable and there exists a positive constant C, depending only on n, p, $[\omega]_{A_p(\mathbb{R}^n)}$, and Ω , such that, for any weak solution $u, u \in W^{1,p}_{0,\omega}(\Omega)$ and

(1.13)
$$\|\nabla u\|_{L^{p}_{\omega}(\Omega;\mathbb{R}^{n})} \leq C \|\mathbf{f}\|_{L^{p}_{\omega}(\Omega;\mathbb{R}^{n})}.$$

Assume that $p \in [2, \infty)$. To prove Theorem 1.10 via using Theorem 1.7, we need to prove that there exists a $\delta_0 \in (0, 1)$, depending on p and Ω , such that, if Ω is a bounded (δ, σ, R) quasi-convex domain with some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then the weak reverse Hölder inequality (1.8) is valid for the exponent p. To this end, we adequately use the realvariable argument obtained in Theorem 3.1 below, the method of perturbation, and the geometric properties of quasi-convex domains (see Lemma 5.3 below).

Let the (semi-)convex domain be as in Remark 2.5(i) below. As a corollary of Theorem 1.10, we have the following conclusion.

Corollary 1.11. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded C^1 or (semi-)convex domain, $p \in (1, \infty)$, and $\omega \in A_p(\mathbb{R}^n)$. Assume that the matrix A satisfies Assumption 1.4. Then there exists a positive constant $\delta_0 \in (0, \infty)$, depending only on n, p, Ω , and $[\omega]_{A_p(\mathbb{R}^n)}$, such that, if A satisfies the (δ, \mathbb{R}) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then the weighted Dirichlet problem $(D)_{p,\omega}$ with $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$ is uniquely solvable and there exists a positive constant C, depending only on n, p, $[\omega]_{A_p(\mathbb{R}^n)}$, and Ω , such that, for any weak solution u, $u \in W^{1,p}_{0,\omega}(\Omega)$ and

$$\|\nabla u\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)}.$$

Moreover, let the Reifenberg flat domain be as in Remark 2.5(ii) below. By Theorem 1.10 and Remark 2.5(ii), we have the following corollary.

Corollary 1.12. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded (δ, R) -Reifenberg flat domain with some $\delta \in (0, 1)$ and $R \in (0, \infty)$, $p \in (1, \infty)$, and $\omega \in A_p(\mathbb{R}^n)$. Assume that the matrix A satisfies Assumption 1.4. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, Ω , and $[\omega]_{A_p(\mathbb{R}^n)}$, such that, if Ω is a bounded (δ, R) -Reifenberg flat domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then the weighted Dirichlet problem $(D)_{p,\omega}$ with $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$ is uniquely solvable and there exists a positive constant C, depending only on $n, p, [\omega]_{A_p(\mathbb{R}^n)}$, and Ω , such that, for any weak solution $u, u \in W^{1,p}_0(\Omega)$ and

$$\|\nabla u\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)}.$$

Remark 1.13. Let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded (δ, σ, R) quasi-convex domain with some $\delta, \sigma \in (0, 1)$ and $R \in (0, \infty)$. Assume that the matrix A := a + b satisfies Assumption 1.4.

- (i) If *a* satisfies the (δ, R) -BMO condition for some sufficiently small $\delta \in (0, \infty)$ and $b \equiv 0$, then Theorem 1.10(i) in this case was established by Jia et al. in [36, Theorem 1.1]. Thus, Theorem 1.10(i) improves [36, Theorem 1.1] via weakening the condition for the matrix *A*. Moreover, even when $b \equiv 0$, the conclusion of Theorem 1.10(ii) in this case is also new.
- (ii) When *a* satisfies the (δ, R) -BMO condition for some sufficiently small $\delta \in (0, \infty)$, $b \equiv 0$, and $\omega \equiv 1$, Corollary 1.12 in this case was obtained by Byun and Wang in [16, Theorem 1.5].

Therefore, even when $\omega \equiv 1$, Corollary 1.12 in this case also improves [16, Theorem 1.5] via weakening the assumption on A.

Furthermore, we point out that the approach used in this article to establish the global gradient estimates is different from that used in [12, 16, 36]. Indeed, the global estimates were obtained in [12, 16, 36] by using an approximation argument, the modified Vitali covering lemma, and a compactness method. However, in this article, we establish the (weighted) global estimates via using a (weighted) real-variable argument (see Theorem 3.1 below), the method of perturbation, and the geometric properties of quasi-convex domains.

(iii) We point out that, even when Ω is a bounded convex domain and $\omega \equiv 1$, the conclusion of Corollary 1.11 in this case is also new.

Applying the weighted global regularity estimates obtained in Theorems 1.8(ii) and 1.10(ii), and some tools from harmonic analysis, such as the properties of Muckenhoupt weights, the interpolation theorem of operators, and the Rubio de Francia extrapolation theorem, we obtain the global gradient estimates for the Dirichlet problem (1.4), respectively, in (weighted) Lorentz spaces, (Lorentz–)Morrey spaces, (Musielak–)Orlicz spaces, and variable Lebesgue spaces, which have independent interests and are presented in Section 6 below. It is worth pointing out that the approach used in this article to establish the global estimates in both Orlicz spaces and variable Lebesgue spaces is quite different from that used in [13, 17]. In [13, 17], the global estimates in variable Lebesgue spaces or in Orlicz spaces were established via the so-called "maximum function free technique". However, in this article, we obtain the global gradient estimates in both Orlicz spaces and variable Lebesgue spaces destimates in both Orlicz spaces and variable Lebesgue spaces in the approach used in 1.10(ii), and the Rubio de Francia extrapolation theorem. Furthermore, we point out that the extrapolation theorem used in this article is also valid for the boundary value problem studied in [13, 17] and independent of the boundary value condition and the considered equation.

Moreover, the global estimates in Orlicz spaces were obtained in [37] through an approximation argument, the modified Vitali covering lemma, and a compactness method. However, in this article, the global gradient estimates in Orlicz spaces are obtained as corollaries of both the weighted norm inequality in Theorem 1.10(ii) and the extrapolation theorem.

The remainder of this article is organized as follows.

In Section 2, we present several notions on the (δ , *R*)-BMO condition, the space VMO(Ω), and several domains, and clarify their relations. We then prove Theorems 1.7, and 1.8, and 1.10, and Corollary 1.11, respectively, in Sections 3, and 4, and 5. Several applications of Theorems 1.8 and 1.10 are given in Section 6.

Finally, we make some conventions on notation. Throughout this article, we always denote by *C* a *positive constant* which is independent of the main parameters, but it may vary from line to line. We also use $C_{(\gamma,\beta,...)}$ or $c_{(\gamma,\beta,...)}$ to denote a *positive constant* depending on the indicated parameters γ, β, \ldots . The symbol $f \leq g$ means that $f \leq Cg$. If $f \leq g$ and $g \leq f$, then we write $f \sim g$. If $f \leq Cg$ and g = h or $g \leq h$, we then write $f \leq g \sim h$ or $f \leq g \leq h$, rather than $f \leq g = h$ or $f \leq g \leq h$. For each ball $B := B(x_B, r_B)$ of \mathbb{R}^n , with some $x_B \in \mathbb{R}^n$ and $r_B \in (0, \infty)$, and $\alpha \in (0, \infty)$, let $\alpha B := B(x_B, \alpha r_B)$; furthermore, denote the set $B(x, r) \cap \Omega$ by $B_{\Omega}(x, r)$, and the set $(\alpha B) \cap \Omega$ by αB_{Ω} . For any subset *E* of \mathbb{R}^n , we denote the set $\mathbb{R}^n \setminus E$ by $E^{\mathbb{C}}$, and its *characteristic function* by $\mathbf{1}_E$. For any $\omega \in A_p(\mathbb{R}^n)$ with $p \in [1, \infty)$, and any measurable set $E \subset \mathbb{R}^n$, let $\omega(E) := \int_E \omega(x) dx$. For any given $q \in [1, \infty]$, we denote by q' its *conjugate exponent*, namely, 1/q + 1/q' = 1. Finally, for any measurable set $E \subset \mathbb{R}^n$, $\omega \in A_q(\mathbb{R}^n)$ with some $q \in [1, \infty)$, and $f \in L^1(E)$, we denote the integral $\int_E |f(x)| \omega(x) dx$ simply by $\int_E |f| \omega dx$ and, when $|E| < \infty$, we use the notation

$$\int_E f dx := \frac{1}{|E|} \int_E f(x) dx.$$

2 Several notions

In this section, we present the definitions of the (δ, R) -BMO condition, the space VMO(Ω), and several domains including quasi-convex domains, (semi-)convex domains, and Reifenberg flat domains. Furthermore, we also clarify the relations between NTA domains, Lipschitz domains, quasi-convex domains, Reifenberg flat domains, and semi-convex domains.

First, we recall the notions of the (δ , R)-BMO condition and the space VMO(Ω) as follows (see, for instance, [15, 16, 50]).

Definition 2.1. Let $\Omega \subset \mathbb{R}^n$ be a domain and $R, \delta \in (0, \infty)$.

(i) A function $f \in L^1_{loc}(\Omega)$ is said to satisfy the (δ, R) -BMO condition if

$$||f||_{*,R} := \sup_{B(x,r) \subset \Omega, r \in (0,R)} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y) - f_{B(x,r)}| \, dy \le \delta,$$

where the supremum is taken over all balls $B(x, r) \subset \Omega$ with $r \in (0, R)$. Furthermore, f is said to belong to the *space* VMO(Ω) if f satisfies the (δ, R) -BMO condition for some $\delta, R \in (0, \infty)$, and

$$\lim_{r \to 0^+} \sup_{B(x,r) \subset \Omega} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y) - f_{B(x,r)}| \, dy = 0,$$

where $r \to 0^+$ means $r \in (0, \infty)$ and $r \to 0$.

(ii) A matrix $A := \{a_{ij}\}_{i,j=1}^{n}$ is said to satisfy the (δ, R) -BMO *condition* [resp., $A \in \text{VMO}(\Omega)$] if, for any $i, j \in \{1, ..., n\}, a_{ij}$ satisfies the (δ, R) -BMO condition [resp., $a_{ij} \in \text{VMO}(\Omega)$].

Remark 2.2. Let $\Omega \subset \mathbb{R}^n$ be a domain and $\delta \in (0, \infty)$. If $f \in BMO(\Omega)$ and $||f||_{BMO(\Omega)} \leq \delta$, then f satisfies the (δ, R) -BMO condition for any $R \in (0, \infty)$. Moreover, if $f \in VMO(\Omega)$, then f satisfies the (γ, R) -BMO condition for any $\gamma \in (0, \infty)$ and some $R \in (0, \infty)$.

Now, we recall the notion of quasi-convex domains as follows. Let $E_1, E_2 \subset \mathbb{R}^n$ be non-empty measurable subsets. Then the *Hausdorff distance* between E_1 and E_2 is defined by setting

$$d_H(E_1, E_2) := \max \left\{ \sup_{x \in E_1} \inf_{y \in E_2} |x - y|, \sup_{y \in E_2} \inf_{x \in E_1} |x - y| \right\}$$

Definition 2.3. Let $\Omega \subset \mathbb{R}^n$ be a domain, δ , $\sigma \in (0, 1)$, and $R \in (0, \infty)$. Then Ω is called a (δ , σ , R) *quasi-convex domain* if, for any $x \in \partial \Omega$ and $r \in (0, R]$,

- (a) there exists an $x_0 \in \Omega$, depending on *x*, such that $B(x_0, \sigma r) \subset \Omega \cap B(x, r)$;
- (b) there exists a convex domain V := V(x, r), depending on x and r, such that $B(x, r) \cap \Omega \subset V$ and $d_H(\partial(B(x, r) \cap \Omega), \partial V) \leq \delta r$.
- **Remark 2.4.** (i) The notion of quasi-convex domains was introduced by Jia et al. [36] to study the global regularity of second order elliptic equations. Roughly speaking, a quasi-convex domain is a domain satisfying that the local boundary is close to be convex at small scales. It is easy to see that, if Ω is a convex domain, then Ω is a (δ , σ , R) quasi-convex domain for any $\delta \in (0, 1)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$.
 - (ii) We may *always* assume that the convex domain *V* in Definition 2.3(b) is the convex hull of $B(x, r) \cap \Omega$, which is the smallest convex domain containing $B(x, r) \cap \Omega$ (see [36, Theorem 3.1] and [67, Remark 1.2] for more details).
- (iii) It was showed in [36, Theorem 3.10] that Definition 2.3(b) can be replaced by the following condition:
 - (c) For any $x \in \partial \Omega$ and $r \in (0, R]$, there exist an (n 1)-dimensional plane L(x, r) containing x, a choice of the unit normal vector to L(x, r), denoted by $v_{x,r}$, and the half space

 $H(x, r) := \{y + t v_{x,t} : y \in L(x, t), t \in [-\delta r, \infty)\}$

such that

$$\Omega \cap B(x,r) \subset H(x,r) \cap B(x,r).$$

More precisely, it was proved in [36, Theorem 3.10] that, if Ω is a domain satisfying the assumptions (a) and (c) with some δ , $\sigma \in (0, 1)$ and $R \in (0, \infty)$, then Ω is a (δ_1, σ, R) quasi-convex domain with $\delta_1 := 8\delta/\sigma$.

In the following remark, we recall the notions of semi-convex domains and Reifenberg flat domains, and then clarify the relations between NTA domains, Lipschitz domains, quasi-convex domains, Reifenberg flat domains, and semi-convex domains.

Remark 2.5. (i) A set $E \subset \mathbb{R}^n$ is said to satisfy an *exterior ball condition* at $x \in \partial E$ if there exist a $\mathbf{v} \in S^{n-1}$ and an $r \in (0, \infty)$ such that

(2.1)
$$B(x + r\mathbf{v}, r) \subset (\mathbb{R}^n \setminus E),$$

where S^{n-1} denotes the *unit sphere* of \mathbb{R}^n . For such an $x \in \partial E$, let

 $r(x) := \sup \left\{ r \in (0, \infty) : (2.1) \text{ holds true for some } v \in S^{n-1} \right\}.$

A set *E* is said to satisfy a *uniform exterior ball condition* (for short, UEBC) with radius $r \in (0, \infty]$ if

(2.2)
$$\inf_{x \in \partial E} r(x) \ge r,$$

and the value r in (2.2) is referred to the UEBC *constant*. A set E is said to satisfy a UEBC if there exists an $r \in (0, \infty]$ such that E satisfies the uniform exterior ball condition with radius r. Moreover, the largest positive constant r as above is called the *uniform ball constant* of E.

It is known that, for any open set $\Omega \subset \mathbb{R}^n$ with compact boundary, Ω is a *Lipschitz domain* satisfying a UEBC if and only if Ω is a semi-convex domain in \mathbb{R}^n (see, for instance, [47, Theorem 2.5] or [48, Theorem 3.9]). Moreover, more equivalent characterizations of semi-convex domains are given in [47, 48].

It is worth pointing out that, if $\Omega \subset \mathbb{R}^n$ is convex, then Ω satisfies a UEBC with the uniform ball constant ∞ (see, for instance, [27]). Thus, convex domains of \mathbb{R}^n are semi-convex domains (see, for instance, [47, 48, 60, 61]). Moreover, (semi-)convex domains are special cases of Lipschitz domains. More precisely,

class of convex domains \subsetneq class of semi-convex domains \subsetneq class of Lipschitz domains.

(ii) Let $n \ge 2$, $\delta \in (0, 1)$, and $R \in (0, \infty)$. A domain $\Omega \subset \mathbb{R}^n$ is called a (δ, R) -*Reifenberg flat domain* if, for any $x_0 \in \partial \Omega$ and $r \in (0, R]$, there exists a system of coordinates, $\{y_1, \ldots, y_n\}$, which may depend on x_0 and r, such that, in this coordinate system, $x_0 = \mathbf{0}$ and

$$B(\mathbf{0},r) \cap \{y \in \mathbb{R}^n : y_n > \delta r\} \subset B(\mathbf{0},r) \cap \Omega \subset B(\mathbf{0},r) \cap \{y \in \mathbb{R}^n : y_n > -\delta r\},\$$

where **0** denotes the *origin* of \mathbb{R}^n . The Reifenberg flat domain was introduced by Reifenberg in [49], which naturally appears in the theory of minimal surfaces and free boundary problems. A typical example of Reifenberg flat domains is the well-known Van Koch snowflake (see, for instance, [57]). In recent years, boundary value problems of elliptic or parabolic equations on Reifenberg flat domains have been widely concerned and studied (see, for instance, [8, 10, 11, 13, 16, 17, 25, 44, 64, 65]).

Moreover, for any given $\delta \in (0, 1)$ and $R \in (0, \infty)$, a (δ, R) -Reifenberg flat domain is a (δ, σ, R) quasi-convex domain with $\sigma := \frac{1-\delta}{2}$ (see, for instance, [36]). However, a quasi-convex domain may not be a Reifenberg flat domain. Indeed, let

$$\Omega := \left\{ (x_1, x_2) \in \mathbb{R}^2 : 1 > x_2 > |x_1| \right\}.$$

Then Ω is convex and hence a quasi-convex domain, but Ω is not a Reifenberg flat domain. Thus,

class of Reifenberg flat domains \subsetneq class of quasi-convex domains.

Furthermore, it was showed by Kenig and Toro [40] that, if δ is sufficiently small, then (δ, R) -Reifenberg flat domains are also NTA domains.

(iii) By the facts that Lipschitz domains with small Lipschitz constants are Reifenberg flat domains and that C^1 domains are Lipschitz domains with small Lipschitz constants (see, for instance, [57]), we conclude that C^1 domains are (δ, σ, R) quasi-convex domains with any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$, where $\delta_0 \in (0, 1)$ is a positive constant depending only on Ω .

Let $\Omega \subset \mathbb{R}^n$ be a bounded semi-convex domain. Then Ω is a (δ, σ, R) quasi-convex domain for any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$, where $\delta_0 \in (0, 1)$ is a positive constant depending only on Ω (see Lemma 5.4 below).

(iv) On NTA domains, quasi-convex domains, Reifenberg flat domains, Lipschitz domains, C^1 domains, and (semi-)convex domains, we have the following relations.

 $(iv)_1$

class of C^1 domains \subsetneq class of Lipschitz domains with small Lipschitz constants \subsetneq class of Lipschitz domains \subsetneq class of NTA domains;

class of (semi-)convex domains \subsetneq class of Lipschitz domains.

 $(iv)_2$

class of C^1 domains \subsetneq class of Lipschitz domains with small Lipschitz constants

- ⊊ class of Reifenberg flat domains
- \subseteq class of quasi-convex domains;

class of (semi-)convex domains \subsetneq class of quasi-convex domains.

(iv)₃ Reifenberg flat domains or quasi-convex domains may not be Lipschitz domains, and generally Lipschitz domains may also not be Reifenberg flat domains (see, for instance, [57]). Moreover, (semi-)convex domains may not be Lipschitz domains with small Lipschitz constants, or Reifenberg flat domains.

3 Proof of Theorem 1.7

In this section, we prove Theorem 1.7 via using a real-variable argument for (weighted) $L^p(\Omega)$ estimates, which is inspired by the work of Caffarelli and Peral [18] (see also [58]). When Ω is a bounded Lipschitz domain in \mathbb{R}^n , the conclusion of Theorem 3.1 was essentially established in [54, Theorem 3.4] (see also [28, Theorems 2.1 and 2.2], [53, Theorem 4.2.6], [55, Theorem 3.3], and [62, Theorem 3.1]). It is worth pointing out that the proofs of [54, Theorem 3.4], [53, Theorem 4.2.6], and [62, Theorem 3.1] are also valid in the case of bounded NTA domains. Thus, we omit the proof of Theorem 3.1 here. Furthermore, we also mention that a similar argument with a different motivation was established in [3, 4]. Moreover, a different weighted real-variable argument was obtained by Shen [52, Theorem 2.1].

Theorem 3.1. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $p_1, p_2 \in [1, \infty)$ satisfy $p_2 > p_1$, $F \in L^{p_1}(\Omega)$, and $f \in L^q(\Omega)$ with some $q \in (p_1, p_2)$. Assume that, for any ball $B := B(x_B, r_B) \subset \mathbb{R}^n$ having the property that $|B| \le \beta_1 |\Omega|$ and either $2B \subset \Omega$ or $x_B \in \partial\Omega$, there exist two measurable functions F_B and R_B on 2B such that $|F| \le |F_B| + |R_B|$ on $2B \cap \Omega$,

(3.1)
$$\left(\int_{2B_{\Omega}} |R_B|^{p_2} dx\right)^{\frac{1}{p_2}} \le C_1 \left[\left(\int_{\beta_2 B_{\Omega}} |F|^{p_1} dx\right)^{\frac{1}{p_1}} + \sup_{\widetilde{B} \supset B} \left(\int_{\widetilde{B}_{\Omega}} |f|^{p_1} dx\right)^{\frac{1}{p_1}} \right],$$

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

and

(3.2)
$$\left(\int_{2B_{\Omega}} |F_B|^{p_1} dx\right)^{\frac{1}{p_1}} \le \varepsilon \left(\int_{\beta_2 B_{\Omega}} |F|^{p_1} dx\right)^{\frac{1}{p_1}} + C_2 \sup_{\widetilde{B} \supset B} \left(\int_{\widetilde{B}_{\Omega}} |f|^{p_1} dx\right)^{\frac{1}{p_1}}$$

where C_1 , C_2 , ε , and $\beta_1 < 1 < \beta_2$ are positive constants independent of F, f, R_B , F_B , and B, and the suprema are taken over all balls $\widetilde{B} \supset B$.

Then, for any $\omega \in A_{q/p_1}(\mathbb{R}^n) \cap RH_s(\mathbb{R}^n)$ with $s \in ((\frac{p_2}{q})', \infty]$, there exists a positive constant ε_0 , depending only on C_1 , C_2 , n, p_1 , p_2 , q, β_1 , β_2 , $[\omega]_{A_{q/p_1}(\mathbb{R}^n)}$, and $[\omega]_{RH_s(\mathbb{R}^n)}$, such that, if $\varepsilon \in [0, \varepsilon_0)$, then

$$\left[\frac{1}{\omega(\Omega)}\int_{\Omega}|F|^{q}\omega\,dx\right]^{\frac{1}{q}} \leq C\left\{\left(\frac{1}{|\Omega|}\int_{\Omega}|F|^{p_{1}}\,dx\right)^{\frac{1}{p_{1}}} + \left[\frac{1}{\omega(\Omega)}\int_{\Omega}|f|^{q}\omega\,dx\right]^{\frac{1}{q}}\right\},$$

where *C* is a positive constant depending only on *C*₁, *C*₂, *n*, *p*₁, *p*₂, *q*, β_1 , β_2 , $[\omega]_{A_{q/p_1}(\mathbb{R}^n)}$, and $[\omega]_{RH_s(\mathbb{R}^n)}$.

To show Theorem 1.7 via using Theorem 3.1, we need the following auxiliary conclusion.

Lemma 3.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $p \in (1, \infty)$, and $p' \in (1, \infty)$ be given by 1/p + 1/p' = 1. Assume that, if the matrix A satisfies Assumption 1.4 and $\mathbf{f} \in L^p(\Omega; \mathbb{R}^n)$, then the weak solution $u \in W_0^{1,p}(\Omega)$ of the Dirichlet problem

(3.3)
$$\begin{cases} -\operatorname{div}(A\nabla u) = \operatorname{div}(\mathbf{f}) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

satisfies the estimate

$$(3.4) \|\nabla u\|_{L^p(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant independent of *u* and **f**. Let $\mathbf{g} \in L^{p'}(\Omega; \mathbb{R}^n)$ and $v \in W_0^{1,p'}(\Omega)$ be the weak solution of the Dirichlet problem (3.3) with **f** replaced by **g**. Then

$$\|\nabla v\|_{L^{p'}(\Omega;\mathbb{R}^n)} \leq C \|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^n)},$$

where C is a positive constant independent of v and g.

Proof. Let $\mathbf{f} \in L^p(\Omega; \mathbb{R}^n)$ and $w \in W_0^{1,p}(\Omega)$ be the weak solution of the Dirichlet problem

(3.5)
$$\begin{cases} -\operatorname{div}(A^*\nabla w) = \operatorname{div}(\mathbf{f}) & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega \end{cases}$$

where A^* denotes the transpose of A. By the assumption that A satisfies Assumption 1.4, we find that A^* also satisfies Assumption 1.4, which, combined with the assumption (3.4), further implies that (3.4) also holds true for the weak solution w of (3.5). Moreover, from the fact that v is the weak solution of (3.3) with **f** replaced by **g**, it follows that

$$\int_{\Omega} \mathbf{g} \cdot \nabla w \, dx = -\int_{\Omega} A \nabla v \cdot \nabla w \, dx = -\int_{\Omega} A^* \nabla w \cdot \nabla v \, dx = \int_{\Omega} \mathbf{f} \cdot \nabla v \, dx,$$

which, together with (3.4) and the Hölder inequality, further implies that

$$\begin{split} \|\nabla v\|_{L^{p'}(\Omega;\mathbb{R}^{n})} &= \sup_{\|\mathbf{f}\|_{L^{p}(\Omega;\mathbb{R}^{n})} \leq 1} \left| \int_{\Omega} \mathbf{f} \cdot \nabla v \, dx \right| = \sup_{\|\mathbf{f}\|_{L^{p}(\Omega;\mathbb{R}^{n})} \leq 1} \left| \int_{\Omega} \mathbf{g} \cdot \nabla w \, dx \right| \\ &\leq \sup_{\|\mathbf{f}\|_{L^{p}(\Omega;\mathbb{R}^{n})} \leq 1} \|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \|\nabla w\|_{L^{p}(\Omega;\mathbb{R}^{n})} \\ &\lesssim \sup_{\|\mathbf{f}\|_{L^{p}(\Omega;\mathbb{R}^{n})} \leq 1} \|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \|\mathbf{f}\|_{L^{p}(\Omega;\mathbb{R}^{n})} \lesssim \|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})}. \end{split}$$

This finishes the proof of Lemma 3.2.

Moreover, to prove Theorem 1.7, we need the following properties of $A_p(\mathbb{R}^n)$ weights, which are well known (see, for instance, [32, Chapter 7] and [4, Proposition 2.1 and Lemma 4.4]).

Lemma 3.3. Let $q \in (1, \infty)$, $\omega \in A_q(\mathbb{R}^n)$, and $\Omega \subset \mathbb{R}^n$ be a bounded domain.

- (i) There exists a $q_1 \in (1, q)$, depending only on n, q, and $[\omega]_{A_a(\mathbb{R}^n)}$, such that $\omega \in A_{q_1}(\mathbb{R}^n)$.
- (ii) There exists a $\gamma \in (0, \infty)$, depending only on n, q, and $[\omega]_{A_q(\mathbb{R}^n)}$, such that $\omega \in RH_{1+\gamma}(\mathbb{R}^n)$.
- (iii) If q' denotes the conjugate number of q, namely, 1/q + 1/q' = 1, then $\omega^{-q'/q} \in A_{q'}(\mathbb{R}^n)$ and $[\omega^{-q'/q}]_{A_{q'}(\mathbb{R}^n)} = [\omega]_{A_{q'}(\mathbb{R}^n)}^{q'/q}$.
- (iv) If $\omega \in RH_p(\mathbb{R}^n)$ with $p \in (1, \infty)$, then there exists a $p_1 \in (p, \infty]$, depending only on n, p, and $[\omega]_{RH_p(\mathbb{R}^n)}$, such that $\omega \in RH_{p_1}(\mathbb{R}^n)$.
- (v) Let $p_0, q_0 \in (1, \infty), p \in (p_0, q_0)$, and $v \in L^1_{loc}(\mathbb{R}^n)$. Then $v \in A_{\frac{p}{p_0}}(\mathbb{R}^n) \cap RH_{(\frac{q_0}{p})'}(\mathbb{R}^n)$ if and only if $v^{1-p'} \in A_{\frac{p'}{(q_0)'}}(\mathbb{R}^n) \cap RH_{(\frac{(p_0)'}{p'})'}(\mathbb{R}^n)$.
- (vi) Let $\omega \in A_p(\mathbb{R}^n)$ with some $p \in [1, \infty)$ and $\gamma \in (0, 1)$. Then $\omega^{\gamma} \in RH_{\gamma^{-1}}(\mathbb{R}^n)$ and there exists a positive constant C, depending only on $[\omega]_{A_p(\mathbb{R}^n)}$ and γ , such that $[\omega^{\gamma}]_{RH_{\gamma^{-1}}(\mathbb{R}^n)} \leq C$.
- (vii) Let $q_2 := q(1 + \frac{1}{\gamma})$ with γ as in (ii) and let q_1 be as in (i). Then $L^{q_2}(\Omega) \subset L^q_{\omega}(\Omega) \subset L^{\frac{q}{q_1}}(\Omega)$.

Furthermore, we also need the following Lemma 3.4, whose proof is similar to that of [7, Lemma 4.38], and we omit the details here.

Lemma 3.4. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $0 < p_0 < q \le \infty$, and $r_0 \in (0, \operatorname{diam}(\Omega))$. Assume that $x \in \overline{\Omega}$ and the weak reverse Hölder inequality

$$\left[\int_{B_{\Omega}(x,r)} |g|^{q} dx\right]^{\frac{1}{q}} \leq C_{3} \left[\int_{B_{\Omega}(x,2r)} |g|^{p_{0}} dx\right]^{\frac{1}{p_{0}}}$$

holds true for a given measurable function g on Ω and any $r \in (0, r_0)$, where C_3 is a positive constant, independent of x and r, which may depend on g. Then, for any given $p \in (0, \infty]$, there exists a positive constant C, depending only on p, p_0 , q, and C_3 , such that

$$\left[\int_{B_{\Omega}(x,r)} |g|^q \, dx\right]^{\frac{1}{q}} \le C \left[\int_{B_{\Omega}(x,2r)} |g|^p \, dx\right]^{\frac{1}{p}}$$

holds true for any $r \in (0, r_0)$ *.*

Now, we prove Theorem 1.7 by using Theorem 3.1 and Lemmas 3.2, 3.3, and 3.4.

Proof of Theorem 1.7. We first show (i). Let $B := B(x_B, r_B) \subset \mathbb{R}^n$ be a ball satisfying $r_B \in (0, r_0/4)$ and either $2B \subset \Omega$ or $x_B \in \partial \Omega$. Take $\phi \in C_c^{\infty}(\mathbb{R}^n)$ such that $\phi \equiv 1$ on 2B, $0 \le \phi \le 1$, and $\operatorname{supp}(\phi) \subset 4B$. Let $w, v \in W_0^{1,2}(\Omega)$ be respectively the weak solutions of the Dirichlet problems

(3.6)
$$\begin{cases} -\operatorname{div}(A\nabla w) = \operatorname{div}(\phi \mathbf{f}) & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega \end{cases}$$

and

(3.7)
$$\begin{cases} -\operatorname{div}(A\nabla v) = \operatorname{div}((1-\phi)\mathbf{f}) & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Then u = w + v and $\nabla u = \nabla w + \nabla v$. Let $F := |\nabla u|$, $f := |\mathbf{f}|$, $F_B := |\nabla w|$, and $R_B := |\nabla v|$. It is easy to see that $0 \le F \le F_B + R_B$. By $\mathbf{f} \in L^2(\Omega; \mathbb{R}^n)$, (3.6), and the fact that (1.9) holds true for p = 2 (see Remark 1.6), we conclude that

(3.8)
$$\int_{2B_{\Omega}} F_B^2 dx = \int_{2B_{\Omega}} |\nabla w|^2 dx \leq \frac{1}{|2B_{\Omega}|} \int_{\Omega} |\mathbf{f}\phi|^2 dx \leq \int_{4B_{\Omega}} |\mathbf{f}|^2 dx \sim \int_{4B_{\Omega}} f^2 dx.$$

Moreover, from (3.7) and the assumption (1.8) of this theorem, it follows that (1.8) holds true for the above v, which, together with the self-improvement property of the weak reverse Hölder inequality (see, for instance, [30, pp. 122-123]), further implies that there exists an $\varepsilon_0 \in (0, \infty)$ such that the inequality (1.8) holds true with p replaced by $p + \varepsilon_0$. By this and Lemma 3.4, we conclude that, for any $q \in (0, 2]$, the weak reverse Hölder inequality

(3.9)
$$\left(\int_{B_{\Omega}} |\nabla v|^{p+\varepsilon_0} dx\right)^{1/(p+\varepsilon_0)} \lesssim \left(\int_{2B_{\Omega}} |\nabla v|^q dx\right)^{1/q}$$

holds true, which, combined with (3.8), further implies that

$$(3.10) \qquad \left(\int_{B_{\Omega}} R_{B}^{p+\varepsilon_{0}} dx\right)^{1/(p+\varepsilon_{0})} \lesssim \left(\int_{2B_{\Omega}} |\nabla v|^{2} dx\right)^{1/2} \lesssim \left(\int_{4B_{\Omega}} |\nabla u|^{2} dx\right)^{1/2} + \left(\int_{4B_{\Omega}} f^{2} dx\right)^{1/2} \\ \sim \left(\int_{4B_{\Omega}} F^{2} dx\right)^{1/2} + \left(\int_{4B_{\Omega}} f^{2} dx\right)^{1/2}.$$

From (3.8) and (3.10), we deduce that (3.1) and (3.2) hold true with $p_2 := p + \varepsilon_0$ and $p_1 := 2$. Thus, by Theorem 3.1 with $\omega \equiv 1$ and q := p, and the Hölder inequality, we conclude that

$$\left(\int_{\Omega} |\nabla u|^p \, dx \right)^{1/p} \lesssim \left(\int_{\Omega} |\nabla u|^2 \, dx \right)^{1/2} + \left(\int_{\Omega} |\mathbf{f}|^p \, dx \right)^{1/p}$$
$$\lesssim \left(\int_{\Omega} |\mathbf{f}|^2 \, dx \right)^{1/2} + \left(\int_{\Omega} |\mathbf{f}|^p \, dx \right)^{1/p} \lesssim \left(\int_{\Omega} |\mathbf{f}|^p \, dx \right)^{1/p},$$

which implies that (1.9) holds true. Thus, (i) holds true.

Next, we prove (ii). Let $q \in [2, p]$, $q_0 \in [1, \frac{q}{p'}]$, $r_0 \in [(\frac{p}{q})', \infty]$, and $\omega \in A_{q_0}(\mathbb{R}^n) \cap RH_{r_0}(\mathbb{R}^n)$. Assume that $\varepsilon_0 \in (0, \infty)$ is as in (3.9). Then $\omega \in A_{\frac{q}{(p+\varepsilon_0)'}}(\mathbb{R}^n) \cap RH_s(\mathbb{R}^n)$ with $s \in ((\frac{p+\varepsilon_0}{q})', \infty]$. Let u be the weak solution of the Dirichlet problem $(D)_{q,\omega}$ with $\mathbf{f} \in L^q_{\omega}(\Omega; \mathbb{R}^n)$. Then, from Lemma 3.3(vii), it follows that

$$(3.11) L^q_{\omega}(\Omega) \subset L^{q/q_0}(\Omega).$$

By the fact that $q_0 \leq q/p' < q/(p + \varepsilon_0)'$, we find that $(p + \varepsilon_0)' < q/q_0$, which, together with the Hölder inequality and the assumption that Ω is bounded, implies that $L^{q/q_0}(\Omega) \subset L^{(p+\varepsilon_0)'}(\Omega)$. From this and (3.11), we deduce that $\mathbf{f} \in L^{q/q_0}(\Omega; \mathbb{R}^n) \subset L^{(p+\varepsilon_0)'}(\Omega; \mathbb{R}^n)$.

Let $B := B(x_B, r_B) \subset \mathbb{R}^n$ be a ball satisfying $|B| \leq \beta_1 |\Omega|$ and either $2B \subset \Omega$ or $x_B \in \partial\Omega$, where $\beta_1 \in (0, 1)$ is as in Theorem 3.1. Take $\phi \in C_c^{\infty}(\mathbb{R}^n)$ such that $\phi \equiv 1$ on 2B, $0 \leq \phi \leq 1$, and $\sup p(\phi) \subset 4B$. Let w and v be, respectively, as in (3.6) and (3.7). Then u = w + v and $\nabla u = \nabla w + \nabla v$. Recall that $F := |\nabla u|$, $f := |\mathbf{f}|$, $F_B := |\nabla w|$, and $R_B := |\nabla v|$. By the proof of (i), we know that (1.9) holds true with p replaced by $p + \varepsilon_0$, which, combined with Lemma 3.2 and (3.6), further implies that

$$\left\|\nabla w\right\|_{L^{(p+\varepsilon_0)'}(\Omega;\mathbb{R}^n)} \lesssim \left\|\phi \mathbf{f}\right\|_{L^{(p+\varepsilon_0)'}(\Omega;\mathbb{R}^n)} \lesssim \left\|\mathbf{f}\right\|_{L^{(p+\varepsilon_0)'}(4B_\Omega;\mathbb{R}^n)}.$$

From this, it follows that

(3.12)
$$\int_{2B_{\Omega}} F_B^{(p+\varepsilon_0)'} dx \lesssim \int_{4B_{\Omega}} f^{(p+\varepsilon_0)'} dx.$$

Furthermore, by the assumption (1.8) of this theorem and (3.7), we conclude that (1.8) holds true for the above *v*. Thus, (3.10) holds true for the above *v*, which, together with (3.12), further implies that

$$(3.13) \qquad \left[\int_{B_{\Omega}} R_{B}^{p+\varepsilon_{0}} dx \right]^{1/(p+\varepsilon_{0})} = \left[\int_{B_{\Omega}} |\nabla v|^{p+\varepsilon_{0}} dx \right]^{1/(p+\varepsilon_{0})} \lesssim \left[\int_{2B_{\Omega}} |\nabla v|^{(p+\varepsilon_{0})'} dx \right]^{1/(p+\varepsilon_{0})'} \\ \lesssim \left[\int_{2B_{\Omega}} |\nabla u|^{(p+\varepsilon_{0})'} dx \right]^{1/(p+\varepsilon_{0})'} + \left[\int_{4B_{\Omega}} |\mathbf{f}|^{(p+\varepsilon_{0})'} dx \right]^{1/(p+\varepsilon_{0})'} \\ \lesssim \left[\int_{4B_{\Omega}} F^{(p+\varepsilon_{0})'} dx \right]^{1/(p+\varepsilon_{0})'} + \left[\int_{4B_{\Omega}} f^{(p+\varepsilon_{0})'} dx \right]^{1/(p+\varepsilon_{0})'}.$$

From (3.12) and (3.13), we deduce that (3.1) and (3.2) hold true with $p_2 := p + \varepsilon_0$ and $p_1 := (p + \varepsilon_0)'$, which, combined with q , Theorem 3.1, and (3.11), further implies that

$$\begin{bmatrix} \frac{1}{\omega(\Omega)} \int_{\Omega} |\nabla u|^{q} \,\omega \,dx \end{bmatrix}^{1/q} = \begin{bmatrix} \frac{1}{\omega(\Omega)} \int_{\Omega} F^{q} \,\omega \,dx \end{bmatrix}^{1/q}$$

$$\lesssim \begin{bmatrix} \frac{1}{|\Omega|} \int_{\Omega} |F|^{(p+\varepsilon_{0})'} \,dx \end{bmatrix}^{1/(p+\varepsilon_{0})'} + \begin{bmatrix} \frac{1}{\omega(\Omega)} \int_{\Omega} |\mathbf{f}|^{q} \,\omega \,dx \end{bmatrix}^{1/q}$$

$$\sim \begin{bmatrix} \frac{1}{|\Omega|} \int_{\Omega} |\nabla u|^{(p+\varepsilon_{0})'} \,dx \end{bmatrix}^{1/(p+\varepsilon_{0})'} + \begin{bmatrix} \frac{1}{\omega(\Omega)} \int_{\Omega} |\mathbf{f}|^{q} \,\omega \,dx \end{bmatrix}^{1/q}$$

$$\lesssim \left[\frac{1}{|\Omega|} \int_{\Omega} |\mathbf{f}|^{(p+\varepsilon_0)'} dx\right]^{1/(p+\varepsilon_0)'} + \left[\frac{1}{\omega(\Omega)} \int_{\Omega} |\mathbf{f}|^q \omega dx\right]^{1/q} \\ \lesssim \left[\frac{1}{\omega(\Omega)} \int_{\Omega} |\mathbf{f}|^q \omega dx\right]^{1/q}.$$

Therefore, (1.10) holds true, which shows (ii). This finishes the proof of Theorem 1.7.

4 **Proof of Theorem 1.8**

In this section, we prove Theorem 1.8 by using Theorem 1.7 and the method of perturbation. We begin with the following reverse Hölder inequality established in [42, Lemma 3.2 and Corollary 4.1].

Lemma 4.1. Let $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $B(x_0, r)$ a ball such that $r \in (0, r_0/4)$ and either $x_0 \in \partial \Omega$ or $B(x_0, 2r) \subset \Omega$, where $r_0 \in (0, \text{diam}(\Omega))$ is a constant. Assume that the matrix A := a + b satisfies Assumption 1.4 and $u \in W^{1,2}(B_\Omega(x_0, 2r))$ is the weak solution of the following Dirichlet problem

$$\begin{cases} -\operatorname{div}(A\nabla u) = 0 & \text{ in } B_{\Omega}(x_0, 2r), \\ u = 0 & \text{ on } B(x_0, 2r) \cap \partial \Omega \end{cases}$$

Then there exists a positive constant $p \in (2, \infty)$, depending on Ω , n, and μ_0 in (1.3), such that

$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla u|^p \, dx\right]^{1/p} \leq C \left[1 + ||b||_{\text{BMO}(\Omega)}\right] \left[\int_{B_{\Omega}(x_0,2r)} |\nabla u|^2 \, dx\right]^{1/2},$$

where *C* is a positive constant depending only on *n*, Ω , and *p*.

To show Theorem 1.8, we need the following perturbation argument which is motivated by [18].

Lemma 4.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and $r_0 \in (0, \text{diam}(\Omega))$ a constant. Assume that the matrix A satisfies Assumption 1.4. Let $u \in W^{1,2}(B_{\Omega}(x_0, 4r))$ be a solution of the equation $\operatorname{div}(A\nabla u) = 0$ in $B_{\Omega}(x_0, 4r)$ with u = 0 on $B(x_0, 4r) \cap \partial\Omega$, where $x_0 \in \overline{\Omega}$ and $r \in (0, r_0/4)$. Then there exist a function $\theta := \theta(r)$, $p \in (2, \infty)$, and a function $v \in W^{1,p}(B_{\Omega}(x_0, r))$ such that

(4.1)
$$\left[\oint_{B_{\Omega}(x_0,r)} |\nabla v|^p \, dx \right]^{1/p} \le C \left[\oint_{B_{\Omega}(x_0,4r)} |\nabla u|^2 \, dx \right]^{1/2}$$

and

(4.2)
$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla(u-v)|^2 \, dx \right]^{1/2} \le \theta(r) \left[\int_{B_{\Omega}(x_0,4r)} |\nabla u|^2 \, dx \right]^{1/2},$$

where C is a positive constant independent of u, v, x_0 , and r.

Proof. By the assumption that *A* satisfies Assumption 1.4, we find that A = a + b, where $a := \{a_{ij}\}_{i,j=1}^{n}$ is real-valued, symmetric, and measurable, and satisfies (1.3), and $b := \{b_{ij}\}_{i,j=1}^{n}$ is real-valued, anti-symmetric, and measurable, and satisfies $b_{ij} \in BMO(\Omega)$ for any $i, j \in \{1, ..., n\}$.

Let $a_0 := \{c_{ij}\}_{i,j=1}^n$, where, for any $i, j \in \{1, ..., n\}$,

$$c_{ij} := \int_{B_{\Omega}(x_0,2r)} a_{ij} \, dx.$$

Assume that $v \in W^{1,2}(B_{\Omega}(x_0, 2r))$ is the solution of the following boundary value problem

(4.3)
$$\begin{cases} \operatorname{div}(a_0 \nabla v) = 0 & \text{ in } B_{\Omega}(x_0, 2r), \\ v = u & \text{ on } \partial B_{\Omega}(x_0, 2r). \end{cases}$$

Then, from (4.3) and the fact that $u - v \in W_0^{1,2}(B_\Omega(x_0, 2r))$, we deduce that

(4.4)
$$\int_{B_{\Omega}(x_0,2r)} a_0 \nabla(u-v) \cdot \nabla(u-v) \, dx = \int_{B_{\Omega}(x_0,2r)} (a_0-A) \nabla u \cdot \nabla(u-v) \, dx.$$

Denote by \widetilde{u} the $W^{1,2}$ -extension of u to \mathbb{R}^n ; namely, $\widetilde{u} \in W^{1,2}(\mathbb{R}^n)$ and $\widetilde{u}|_{B_{\Omega}(x_0,2r)} = u$. Furthermore, denote by $\widetilde{u-v}$ the zero extension of $(u-v)|_{B_{\Omega}(x_0,2r)}$ to \mathbb{R}^n . Obviously, $\widetilde{u-v} \in W^{1,2}(\mathbb{R}^n)$ and supp $(\widetilde{u-v}) \subset B_{\Omega}(x_0,2r)$, which, combined with the divergence theorem and the assumption that

$$b_{B_{\Omega}(x_0,2r)} := \left\{ \int_{B_{\Omega}(x_0,2r)} b_{ij} \, dx \right\}_{i,j=1}^n$$

is anti-symmetric, further implies that

$$\begin{split} \int_{B_{\Omega}(x_0,2r)} b_{B_{\Omega}(x_0,2r)} \nabla u \cdot \nabla (u-v) \, dx &= \int_{B(x_0,2r)} b_{B_{\Omega}(x_0,2r)} \nabla \widetilde{u} \cdot \nabla (\widetilde{u-v}) \, dx \\ &= \int_{B(x_0,2r)} \operatorname{div} \left(b_{B_{\Omega}(x_0,2r)} \nabla \widetilde{u} \right) (\widetilde{u-v}) \, dx \\ &- \int_{\partial B(x_0,2r)} b_{B_{\Omega}(x_0,2r)} \nabla \widetilde{u} \cdot v(\widetilde{u-v}) \, d\sigma(x) \\ &= 0, \end{split}$$

where v denotes the *outward unit normal* to $\partial B(x_0, 2r)$. By this, (4.4), and the definitions of a_0 and $b_{B_{\Omega}(x_0,2r)}$, we find that

(4.5)
$$\int_{B_{\Omega}(x_0,2r)} a_0 \nabla(u-v) \cdot \nabla(u-v) \, dx = \int_{B_{\Omega}(x_0,2r)} \left(a_0 - A + b_{B_{\Omega}(x_0,2r)}\right) \nabla u \cdot \nabla(u-v) \, dx$$
$$= \int_{B_{\Omega}(x_0,2r)} \left(A_{B_{\Omega}(x_0,2r)} - A\right) \nabla u \cdot \nabla(u-v) \, dx,$$

where

$$A_{B_{\Omega}(x_0,2r)} := \left\{ \int_{B_{\Omega}(x_0,2r)} (a_{ij} + b_{ij}) \, dx \right\}_{i,j=1}^n.$$

Therefore, from (4.5), (1.3), and the Hölder inequality, it follows that, for any given $\varepsilon \in (0, \infty)$, there exists a positive constant $C_{(\varepsilon)}$, depending on ε , such that

(4.6)
$$\mu_0 \int_{B_{\Omega}(x_0,2r)} |\nabla(u-v)|^2 dx$$

$$\leq \int_{B_{\Omega}(x_0,2r)} |A - A_{B_{\Omega}(x_0,2r)}| |\nabla u| |\nabla(u-v)| dx$$

$$\leq \varepsilon \int_{B_{\Omega}(x_0,2r)} |\nabla(u-v)|^2 dx + C_{(\varepsilon)} \int_{B_{\Omega}(x_0,2r)} |A - A_{B_{\Omega}(x_0,2r)}|^2 |\nabla u|^2 dx,$$

where $\mu_0 \in (0, 1)$ is as in (1.3). Take $\varepsilon := \mu_0/2$ in (4.6). Then, by (4.6), we conclude that

(4.7)
$$\left[\int_{B_{\Omega}(x_0,2r)} |\nabla(u-v)|^2 \, dx\right]^{1/2} \le C_4 \left[\int_{B_{\Omega}(x_0,2r)} \left|A - A_{B_{\Omega}(x_0,2r)}\right|^2 |\nabla u|^2 \, dx\right]^{1/2}.$$

Moreover, from Lemma 4.1, we deduce that there exist positive constants $\tilde{p} \in (1, \infty)$ and $C_5 \in (0, \infty)$, independent of u, x_0 , and r, such that

(4.8)
$$\left[\int_{B_{\Omega}(x_0,2r)} |\nabla u|^{2\widetilde{p}} dx \right]^{1/(2\widetilde{p})} \le C_5 \left[1 + ||b||_{\text{BMO}(\Omega)} \right] \left[\int_{B_{\Omega}(x_0,4r)} |\nabla u|^2 dx \right]^{1/2}.$$

Furthermore, by [38, Theorem 1], we know that there exist an $\widetilde{A} \in BMO(\mathbb{R}^n; \mathbb{R}^{n^2})$ and a positive constant *C*, independent of x_0 and *r*, such that

(4.9)
$$\widetilde{A}|_{B_{\Omega}(x_0,2r)} = A \quad \text{and} \quad \|\widetilde{A}\|_{BMO(\mathbb{R}^n;\mathbb{R}^{n^2})} \le C\|A\|_{BMO(B_{\Omega}(x_0,2r);\mathbb{R}^{n^2})}.$$

From the Hölder inequality and the well-known John–Nirenberg inequality on $BMO(\mathbb{R}^n)$ (see, for instance, [33, 56]), it follows that

$$\begin{split} \left[\int_{B_{\Omega}(x_{0},2r)} \left| A - A_{B_{\Omega}(x_{0},2r)} \right|^{2\widetilde{p}'} dx \right]^{1/(2\widetilde{p}')} \\ &\lesssim \left[\int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right|^{2\widetilde{p}'} dx \right]^{1/(2\widetilde{p}')} + \left[\int_{B(x_{0},2r)} \left| \widetilde{A}_{B(x_{0},2r)} - A_{B_{\Omega}(x_{0},2r)} \right|^{2\widetilde{p}'} dx \right]^{1/(2\widetilde{p}')} \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \left| \widetilde{A}_{B(x_{0},2r)} - A_{B_{\Omega}(x_{0},2r)} \right| \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B_{\Omega}(x_{0},2r)} \left| A - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} - \widetilde{A}_{B(x_{0},2r)} \right| dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} + \int_{B(x_{0},2r)} \left| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}})} dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}}) + \int_{B(x_{0},2r)} \left| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^{2}}) dx \\ &\lesssim \left\| \widetilde{A} \right\|_{BMO(\mathbb{R}^{n};\mathbb{R}^{n^$$

where $\tilde{p}' \in (1, \infty)$ is given by $1/\tilde{p} + 1/\tilde{p}' = 1$, which, together with (4.9), further implies that there exists a positive constant C_6 , independent of x_0 , r, and A, such that

$$\left[\int_{B_{\Omega}(x_0,2r)} \left|A - A_{B_{\Omega}(x_0,2r)}\right|^{2\widetilde{p}'} dx\right]^{1/(2\widetilde{p}')} \le C_6 ||A||_{\mathrm{BMO}(B_{\Omega}(x_0,2r);\mathbb{R}^{n^2})}.$$

By this, (4.7) and (4.8), we conclude that

$$(4.10) \qquad \left[\int_{B_{\Omega}(x_{0},2r)} |\nabla(u-v)|^{2} dx \right]^{1/2} \\ \leq C_{4} \left[\int_{B_{\Omega}(x_{0},2r)} |A - A_{B_{\Omega}(x_{0},2r)}|^{2\widetilde{p}'} dx \right]^{1/(2\widetilde{p}')} \left[\int_{B_{\Omega}(x_{0},2r)} |\nabla u|^{2\widetilde{p}} dx \right]^{1/(2\widetilde{p})} \\ \leq C_{4}C_{5}C_{6} \left[1 + ||b||_{BMO(\Omega)} \right] ||A||_{BMO(B_{\Omega}(x_{0},2r);\mathbb{R}^{n^{2}})} \left[\int_{B_{\Omega}(x_{0},4r)} |\nabla u|^{2} dx \right]^{1/2} \\ \leq \theta(r) \left[\int_{B_{\Omega}(x_{0},4r)} |\nabla u|^{2} dx \right]^{1/2},$$

where

(4.11)
$$\theta(r) := C_4 C_5 C_6 \left[1 + ||b||_{\text{BMO}(\Omega)} \right] \sup_{B(x,2t) \subset \Omega, \ t \in (0,r]} \int_{B(x,2t)} \left| A - \int_{B(x,2t)} A \ dz \right| \ dy.$$

Thus, (4.2) holds true. Furthermore, from the known regularity theory of second order elliptic equations (see, for instance, [39, Chapter 1] and [30, Chapter V]), we deduce that there exists a $p \in (2, \infty)$ such that

$$\left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^p \, dx\right]^{1/p} \lesssim \left[\int_{B_{\Omega}(x_0,4r)} |\nabla v|^2 \, dx\right]^{1/2},$$

which, combined with (4.10), (4.11), and the fact that $A \in BMO(\Omega; \mathbb{R}^{n^2})$, further implies that (4.1) holds true. This finishes the proof of Lemma 4.2.

Furthermore, to prove Theorem 1.8, we need the following conclusion for the constant coefficient boundary value problems, which was established in [55, Lemma 4.1] (see also [28, Lemma 4.1]).

Lemma 4.3. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, and $a_0 := \{a_{ij}\}_{i,j=1}^n$ a symmetric constant coefficient matrix that satisfies (1.3). Assume that $v \in W^{1,2}(B_\Omega(x_0, 2r))$ is a weak solution of the equation $\operatorname{div}(a_0 \nabla v) = 0$ in $B_\Omega(x_0, 2r)$ with v = 0 on $B(x_0, 2r) \cap \partial \Omega$, where $B(x_0, r)$ is a ball such that $r \in (0, r_0/4)$ and either $x_0 \in \partial \Omega$ or $B(x_0, 2r) \subset \Omega$, and $r_0 \in (0, \operatorname{diam}(\Omega))$ is a constant. Then the weak reverse Hölder inequality

$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla v|^p \, dx\right]^{\frac{1}{p}} \le C \left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^2 \, dx\right]^{\frac{1}{2}}$$

holds true for $p := 3 + \varepsilon$ when $n \ge 3$, or $p := 4 + \varepsilon$ when n = 2, where C and ε are positive constants depending only on n, the Lipschitz constant of Ω , and μ_0 in (1.3).

Now, we prove Theorem 1.8 via using Theorem 3.1 and Lemmas 4.2 and 4.3.

Proof of Theorem 1.8. We first show (i). Let *A* satisfy Assumption 1.4 and *u* be the weak solution of the following Dirichlet problem

(4.12)
$$\begin{cases} -\operatorname{div}(A\nabla u) = \operatorname{div}(\mathbf{f}) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Based on Lemma 3.2, to prove (1.11), it suffices to show that there exists a positive constant $\varepsilon_0 \in (0, \infty)$, depending only on *n* and the Lipschitz constant of Ω , such that, for any given $p \in [2, 3 + \varepsilon_0)$ when $n \ge 3$, or $p \in [2, 4 + \varepsilon_0)$ when n = 2, there exists a $\delta_0 \in (0, \infty)$, depending only on *n*, *p*, and the Lipschitz constant of Ω , such that, if *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then (1.11) holds true.

By the definition of VMO(Ω), we conclude that, if $A \in \text{VMO}(\Omega)$, then there exists an $r_1 \in (0, \infty)$ such that, for any $r \in (0, r_1)$, $\theta(r) < \varepsilon_0/2$, where $\theta(r)$ and ε_0 are, respectively, as in (4.11) and Theorem 3.1. Let $B(x_0, r) \subset \mathbb{R}^n$ be such that $r \in (0, \min\{r_0, r_1\}/4)$ and either $x_0 \in \partial\Omega$ or $B(x_0, 2r) \subset \Omega$, where $r_0 \in (0, \operatorname{diam}(\Omega))$ is as in Lemma 4.3. Assume that $v \in W^{1,2}(B_\Omega(x_0, 2r))$ is a weak solution of the equation $\operatorname{div}(A\nabla v) = 0$ in $B_\Omega(x_0, 2r)$ with v = 0 on $B(x_0, 2r) \cap \partial\Omega$. Let $w \in W^{1,2}(B_\Omega(x_0, 2r))$ be a weak solution of the equation $\operatorname{div}(a\nabla v) = 0$ in $B_\Omega(x_0, 2r)$ with v = 0 on $B(x_0, 2r) \cap \partial\Omega$. Let $w \in W^{1,2}(B_\Omega(x_0, 2r))$ be a weak solution of the equation $\operatorname{div}(a_0\nabla w) = 0$ in $B_\Omega(x_0, 2r)$ with w = v on $\partial B_\Omega(x_0, 2r)$, where $a_0 := \{c_{ij}\}_{i,j=1}^n$ with $c_{ij} := \int_{B_\Omega(x_0, 2r)} a_{ij} dx$ for any $i, j \in \{1, \ldots, n\}$. Then, from Lemmas 4.2 and 4.3, it follows that

(4.13)
$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla w|^p \, dx \right]^{\frac{1}{p}} \lesssim \left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^2 \, dx \right]^{\frac{1}{2}}$$

and

(4.14)
$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla(v-w)|^2 dx\right]^{\frac{1}{2}} \le \theta(r) \left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^2 dx\right]^{\frac{1}{2}},$$

where $\theta(r)$ is as in (4.11). By the well-known John–Nirenberg inequality on BMO(Ω) (see, for instance, [33, 56]), we conclude that there exists a $\delta_0 \in (0, \infty)$ sufficiently small such that, if *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, then, for any $r \in (0, r_0)$, $\theta(r) < \varepsilon_0/2$, where ε_0 is as in Theorem 3.1. From this, (4.13), (4.14), $r \in (0, \min\{r_0, r_1\}/4)$, and Theorem 3.1 with $\omega \equiv 1$, we deduce that, if *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then $|\nabla v| \in L^p(B_\Omega(x_0, 2r))$ and

(4.15)
$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla v|^p \, dx\right]^{\frac{1}{p}} \lesssim \left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^2 \, dx\right]^{\frac{1}{2}},$$

which, together with Theorem 1.7 and the fact that the Dirichlet problem $(D)_2$ is uniquely solvable, further implies that (1.11) holds true in the case $p \in [2, 3 + \varepsilon_0)$ when $n \ge 3$, or $p \in [2, 4 + \varepsilon_0)$ when n = 2. This finishes the proof of (i).

Now, we prove (ii). Let ε_0 be as in (i) and u the weak solution of the Dirichlet problem (4.12) with $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$. We first assume that $p \in [2, p_0)$ and $\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)$, where $p_0 := 3 + \varepsilon_0$ when $n \ge 3$, or $p_0 := 4 + \varepsilon_0$ when n = 2. Then, by (4.15) and Theorem 1.7, we conclude that there exists a $\delta_0 \in (0, \infty)$, depending only on $n, p, [\omega]_{A_{\frac{p}{p'}}(\mathbb{R}^n)}, [\omega]_{RH_{(\frac{p_0}{p'})'}(\mathbb{R}^n)}$, and the

Lipschitz constant of Ω , such that, if *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then (1.12) holds true.

Next, we assume that $p \in (p'_0, 2)$ and $\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p'})'}(\mathbb{R}^n)$. Then, from Lemma 3.3(v), we deduce that $p' \in (2, p_0)$ and $\omega^{1-p'} \in A_{\frac{p'}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p'})'}(\mathbb{R}^n)$. Let $\mathbf{g} \in L^{p'}_{\omega^{1-p'}}(\Omega; \mathbb{R}^n)$ and v be the weak solution of the following Dirichlet problem

$$\begin{cases} -\operatorname{div}(A^*\nabla v) = \operatorname{div}(\mathbf{g}) & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega \end{cases}$$

where A^* denotes the transpose of A. By the assumption that A satisfies Assumption 1.4, we find that A^* also satisfies Assumption 1.4. Thus, we have

(4.16)
$$\|\nabla v\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^n)} \lesssim \|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^n)}.$$

Moreover,

$$\int_{\Omega} \mathbf{g} \cdot \nabla u \, dx = -\int_{\Omega} A^* \nabla v \cdot \nabla u \, dx = -\int_{\Omega} A \nabla u \cdot \nabla v \, dx = \int_{\Omega} \mathbf{f} \cdot \nabla v \, dx,$$

which, combined with (4.16) and the Hölder inequality, further implies that

$$(4.17) \|\nabla u\|_{L^{p}_{\omega}(\Omega;\mathbb{R}^{n})} = \sup_{\|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^{n})} \leq 1} \left| \int_{\Omega} \mathbf{g} \cdot \nabla u \, dx \right| = \sup_{\|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^{n})} \leq 1} \left| \int_{\Omega} \mathbf{f} \cdot \nabla v \, dx \right|$$
$$\leq \sup_{\|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^{n})} \leq 1} \|\mathbf{f}\|_{L^{p}_{\omega}(\Omega;\mathbb{R}^{n})} \|\nabla v\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^{n})}$$
$$\lesssim \sup_{\|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^{n})} \leq 1} \|\mathbf{f}\|_{L^{p}_{\omega}(\Omega;\mathbb{R}^{n})} \|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^{n})} \leq \|\mathbf{f}\|_{L^{p}_{\omega}(\Omega;\mathbb{R}^{n})}.$$

From this, it follows that (1.12) holds true when $p \in (p'_0, 2)$. This finishes the proof of (ii) and hence of Theorem 1.8.

5 Proof of Theorem 1.10

In this section, we give the proofs of Theorem 1.10 and Corollary 1.11 by using Theorem 1.7 and some properties of quasi-convex domains. We begin with the following Lemma 5.1.

Lemma 5.1. Let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain. Assume that $p \in (2, \infty)$, Ω is a (δ, σ, R) quasi-convex domain for some δ , $\sigma \in (0, 1)$ and $R \in (0, \infty)$, $a_0 := \{a_{ij}\}_{i,j=1}^n$ a symmetric constant coefficient matrix that satisfies (1.3). Let $v \in W^{1,2}(B_\Omega(x_0, 2r))$ be a weak solution of the equation div $(a_0 \nabla v) = 0$ in $B_\Omega(x_0, 2r)$ with v = 0 on $B(x_0, 2r) \cap \partial\Omega$, where $B(x_0, r)$ is a ball such that $r \in (0, r_0/4)$ and either $x_0 \in \partial\Omega$ or $B(x_0, 2r) \subset \Omega$, and $r_0 \in (0, R)$ is a constant. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, and Ω , such that, if Ω is a (δ, σ, R)

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

quasi-convex domain for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then the weak reverse Hölder inequality

(5.1)
$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla v|^p \, dx \right]^{\frac{1}{p}} \le C \left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^2 \, dx \right]^{\frac{1}{2}}$$

holds true, where C is a positive constant depending only on n, δ , σ , R, and diam (Ω).

To show Lemma 5.1, we need the following Lemma 5.2, which is a special case of Theorem 3.1.

Lemma 5.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $p_1, p_2 \in [1, \infty)$ with $p_2 > p_1$, $F \in L^{p_1}(\Omega)$, and $q \in (p_1, p_2)$. Suppose that, for any ball $B := B(x_B, r_B) \subset \mathbb{R}^n$ having the property that $|B| \le \beta_1 |\Omega|$ and either $2B \subset \Omega$ or $x_B \in \partial \Omega$, there exist two measurable functions F_B and R_B on 2B such that $|F| \le |F_B| + |R_B|$ on $2B \cap \Omega$,

$$\left(\int_{2B_{\Omega}} |R_B|^{p_2} dx\right)^{\frac{1}{p_2}} \le C_7 \left(\int_{\beta_2 B_{\Omega}} |F|^{p_1} dx\right)^{\frac{1}{p_1}}$$

and

$$\left(\int_{2B_{\Omega}}|F_B|^{p_1}\,dx\right)^{\frac{1}{p_1}}\leq \varepsilon\left(\int_{\beta_2B_{\Omega}}|F|^{p_1}\,dx\right)^{\frac{1}{p_1}},$$

where C_7 , ε , and $\beta_1 < 1 < \beta_2$ are positive constants independent of F, R_B , F_B , and B. Then there exists a positive constant ε_0 , depending only on C_7 , n, p_1 , p_2 , q, β_1 , and β_2 , such that, if $\varepsilon \in [0, \varepsilon_0)$, then

$$\left(\int_{\Omega} |F|^q \, dx\right)^{\frac{1}{q}} \leq C \left(\int_{\Omega} |F|^{p_1} \, dx\right)^{\frac{1}{p_1}},$$

where C is a positive constant depending only on C_7 , ε_0 , n, p_1 , p_2 , q, β_1 , and β_2 .

Moreover, we also need the following Lemma 5.3, which was established in [67, Lemmas 3.4 and 3.5].

Lemma 5.3. Let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded (δ, σ, R) quasi-convex domain for some $\delta, \sigma \in (0, 1)$ and $R \in (0, \infty)$. Assume that $x_0 \in \partial \Omega$, $r \in (0, R/4)$, and V_{4r} is the convex hull of $B_{\Omega}(x_0, 4r)$. For any $t \in (0, 1)$, let

(5.2)
$$\Omega_{tr} := \{ x \in \Omega : \operatorname{dist} (x, \partial \Omega)$$

and

(5.3)
$$W_{r,t} := \{ x \in V_{4r} : \operatorname{dist}(x, \partial V_{4r} \cap B(x_0, 3r)) \le (t + \delta)r \}.$$

(i) Then, for any $t \in (0, 1)$, $\Omega_{tr} \cap B(x_0, r) \subset W_{r,t}$. Moreover, there exists a positive constant C, depending only on n and σ , such that $|W_{r,t}| \leq C(t + \delta)r^n$.

(ii) Let $u \in W^{1,2}(B(x_0, 4r))$ satisfy u = 0 on $B(x_0, 4r) \setminus V_{4r}$. Then, for any $t \in (0, 1 - \delta)$, there exists a positive constant *C*, depending only on *n* and σ , such that

$$\int_{B(x_0,r)\cap\Omega_{tr}} u^2 \, dx \le C(t+\delta)^2 r^2 \int_{W_{r,t}} |\nabla u|^2 \, dx.$$

Now, we prove Lemma 5.1 via using Lemmas 4.1, 5.2, and 5.3.

Proof of Lemma 5.1. We borrow some ideas from [67]. Observing that the matrix a_0 is symmetric and elliptic, and has constant coefficients, without loss of generality, by a change of the coordinate system, we may assume that $a_0 = I$ (the unit matrix), namely, $\Delta v = 0$ in $B(x_0, 2r) \cap \Omega$ and v = 0 on $B(x_0, 2r) \cap \partial \Omega$. If $B(x_0, 2r) \subset \Omega$, from the interior gradient estimates of harmonic functions (see, for instance, [31, Theorem 2.10]), it follows that (5.1) holds true for any given $p \in (2, \infty)$.

Next, we assume that $x_0 \in \partial\Omega$. By the assumption that v = 0 on $B(x_0, 2r) \cap \partial\Omega$, we know that v can be extended to a function $\tilde{v} \in W^{1,2}(B(x_0, 2r))$ by zero-extension. Let $s \in (0, r/16)$. Since Ω is a bounded (δ, σ, R) quasi-convex domain, from Definition 2.3 and Remark 2.4(ii), it follows that the convex hull of $B_{\Omega}(x_0, s)$, denoted by V_s , satisfies that

$$V_s \cap \Omega = B(x_0, s) \cap \Omega$$
 and $d_H(\partial V_s, \partial (B(x_0, s) \cap \Omega)) \le \delta s$.

Let $w \in W^{1,2}(V_s)$ be the weak solution of the following Dirichlet problem

$$\begin{cases} -\Delta w = 0 & \text{ in } V_s, \\ w = \widetilde{v} & \text{ on } \partial V_s. \end{cases}$$

By the assumptions that $\tilde{v} = 0$ on $\partial V_s \cap B(x_0, s)$ and that V_s is convex, and the well known result for the boundary gradient estimates of harmonic functions in convex domains (see, for instance, [6, Theorem 1.1]), we find that

(5.4)
$$\|\nabla w\|_{L^{\infty}(B_{\Omega}(x_0,s))} \lesssim \left[\int_{B_{\Omega}(x_0,2s)} |\nabla w|^2 dx \right]^{1/2}.$$

We now clam that there exists a positive constant ϵ such that

(5.5)
$$\left[\int_{B_{\Omega}(x_0,s)} |\nabla(v-w)|^2 dx\right]^{1/2} \le C\delta^{\epsilon} \left[\int_{B_{\Omega}(x_0,2s)} |\nabla v|^2 dx\right]^{1/2},$$

where *C* is a positive constant independent of x_0 , s, δ , and v. If (5.5) holds true, then, from (5.5), (5.4), and Lemma 5.2, we deduce that there exists a $\delta_0 \in (0, 1)$ such that, if Ω is a (δ, σ, R) quasi-convex domain for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then (5.1) holds true.

Next, we prove (5.5). For any given $t \in (0, 1)$, let Ω_{ts} be as in (5.2). Meanwhile, take $\theta_{\delta s} \in C_c^{\infty}(\mathbb{R}^n)$ satisfying that $0 \le \theta_{\delta s} \le 1$, $\theta_{\delta s} \equiv 1$ on $\Omega \setminus \Omega_{2\delta s}$, $\theta_{\delta s} = 0$ in $\Omega_{\delta s}$, and $|\nabla \theta_{\delta s}| \le (\delta s)^{-1}$. Then

(5.6)
$$\int_{V_s} |\nabla(w - \widetilde{v})|^2 dx = \int_{V_s} \nabla w \cdot \nabla(w - \widetilde{v}) dx - \int_{V_s} \nabla \widetilde{v} \cdot \nabla(w - \widetilde{v}) dx$$
$$= -\int_{V_s} \nabla \widetilde{v} \cdot \nabla(w - \widetilde{v}) dx = -\int_{B_{\Omega}(x_0, s)} \nabla v \cdot \nabla(w - v) dx$$

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

$$= -\int_{B_{\Omega}(x_0,s)} \nabla(\theta_{\delta s} v) \cdot \nabla(w - v) \, dx$$
$$- \int_{B_{\Omega}(x_0,s)} \nabla((1 - \theta_{\delta s}) v) \cdot \nabla(w - v) \, dx$$

By the assumptions that $\Delta v = 0$ in $B_{\Omega}(x_0, s)$, and that v = 0 on $B(x_0, 2s) \cap \partial \Omega$, and Lemma 5.3, we conclude that

$$\left[\int_{B_{\Omega}(x_0,s)\cap\Omega_{2\delta s}}|v|^2\,dx\right]^{1/2}\lesssim\delta s\left(\int_{W_{s,2\delta}}|\nabla v|^2\,dx\right)^{1/2}\lesssim\delta s(\delta s^n)^\epsilon\left[\int_{B_{\Omega}(x_0,2s)}|\nabla v|^p\,dx\right]^{1/p},$$

where $W_{s,2\delta}$ is as in (5.3), $p \in (2, \infty)$ as in Lemma 4.1, and $\epsilon := \frac{1}{2} - \frac{1}{p}$, which, combined with $|\nabla \theta_{\delta s}| \leq (\delta s)^{-1}$ and the Hölder inequality, further implies that

$$(5.7) \qquad \left| \int_{B_{\Omega}(x_{0},s)} \nabla(\theta_{\delta s} v) \cdot \nabla(w-v) \, dx \right| \\ \leq \left| \int_{B_{\Omega}(x_{0},s)} v \nabla \theta_{\delta s} \cdot \nabla(w-v) \, dx \right| + \left| \int_{B_{\Omega}(x_{0},s)} \theta_{\delta s} \nabla v \cdot \nabla(w-v) \, dx \right| \\ \leq \left[(\delta s)^{-1} ||v||_{L^{2}(B_{\Omega}(x_{0},s) \cap \Omega_{2\delta s})} + ||\nabla v||_{L^{2}(B_{\Omega}(x_{0},s) \cap \Omega_{2\delta s};\mathbb{R}^{n})} \right] ||\nabla(w-v)||_{L^{2}(B_{\Omega}(x_{0},s);\mathbb{R}^{n})} \\ \leq (\delta s^{n})^{\epsilon} ||\nabla v||_{L^{p}(B_{\Omega}(x_{0},2s);\mathbb{R}^{n})} ||\nabla(w-v)||_{L^{2}(B_{\Omega}(x_{0},s);\mathbb{R}^{n})}.$$

Furthermore, similarly to (5.7), we have

(5.8)
$$\left| \int_{B_{\Omega}(x_0,s)} \nabla((1-\theta_{\delta s})v) \cdot \nabla(w-v) \, dx \right| \\ \lesssim (\delta s^n)^{\epsilon} ||\nabla v||_{L^p(B_{\Omega}(x_0,2s);\mathbb{R}^n)} ||\nabla(w-v)||_{L^2(B_{\Omega}(x_0,s);\mathbb{R}^n)}.$$

Thus, from (5.6), (5.7), and (5.8), it follows that

$$\|\nabla(w-\widetilde{v})\|_{L^2(V_s;\mathbb{R}^n)}^2 \lesssim (\delta s^n)^{\epsilon} \|\nabla v\|_{L^p(B_{\Omega}(x_0,2s);\mathbb{R}^n)} \|\nabla(w-v)\|_{L^2(B_{\Omega}(x_0,s);\mathbb{R}^n)},$$

which further implies that

(5.9)
$$\|\nabla(w-v)\|_{L^2(B_{\Omega}(x_0,s);\mathbb{R}^n)} \leq (\delta s^n)^{\epsilon} \|\nabla v\|_{L^p(B_{\Omega}(x_0,2s);\mathbb{R}^n)}.$$

Then, by (5.9), Lemma 4.1, and $\epsilon := \frac{1}{2} - \frac{1}{p}$, we find that

$$\left[\int_{B_{\Omega}(x_0,s)} |\nabla(v-w)|^2 \, dx\right]^{1/2} \lesssim \delta^{\epsilon} \left[\int_{B_{\Omega}(x_0,2s)} |\nabla v|^p \, dx\right]^{1/p} \lesssim \delta^{\epsilon} \left[\int_{B_{\Omega}(x_0,4s)} |\nabla v|^2 \, dx\right]^{1/2}.$$

This finishes the proof of (5.5) and hence of Lemma 5.1.

Now, we prove Theorem 1.10 via using Lemma 5.1 and Theorem 1.7.

Proof of Theorem 1.10. We first show (i). Via replacing Lemma 4.3 by Lemma 5.1, and repeating the proof of Theorem 1.8(i), we can prove (i). We omit the details here.

Next, we show (ii). Let $p \in (1, \infty)$ and $\omega \in A_p(\mathbb{R}^n)$. Assume that u is the weak solution of the Dirichlet problem (4.12) with $\mathbf{f} \in L^p_{\omega}(\Omega; \mathbb{R}^n)$.

We first assume that $p \in [2, \infty)$. From $\omega \in A_p(\mathbb{R}^n)$ and (i) and (ii) of Lemma 3.3, it follows that there exists a sufficiently large $p_0 \in (p, \infty)$ such that

(5.10)
$$\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p})'}(\mathbb{R}^n).$$

Let *v* be as in (4.14). Using Lemmas 5.1 and 3.4, and repeating the proof of (4.15), we find that there exists a positive constant $\delta_0 \in (0, 1)$, depending only on *n*, *p*, and Ω , such that, if Ω is a (δ, σ, R) quasi-convex domain and *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, the inequality (4.15) holds true with *p* replaced by p_0 , namely,

(5.11)
$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla v|^{p_0} dx \right]^{\frac{1}{p_0}} \lesssim \left[\int_{B_{\Omega}(x_0,2r)} |\nabla v|^2 dx \right]^{\frac{1}{2}}.$$

Then, by (5.10), (5.11), and Theorem 1.7(ii), we conclude that there exists a positive constant $\delta_0 \in (0, 1)$, depending only on *n*, *p*, and Ω , such that, if Ω is a (δ, σ, R) quasi-convex domain and *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then the weighted Dirichlet problem $(D)_{p,\omega}$ is uniquely solvable and (1.13) holds true in this case.

Now, let $p \in (1, 2)$. From this, $\omega \in A_p(\mathbb{R}^n)$, and Lemma 3.3(iii), we deduce that $p' \in (2, \infty)$ and $\omega^{1-p'} \in A_{p'}(\mathbb{R}^n)$. Let $\mathbf{g} \in L^{p'}_{\omega^{1-p'}}(\Omega; \mathbb{R}^n)$ and w be the weak solution of the following Dirichlet problem

$$\begin{cases} -\operatorname{div}(A^*\nabla w) = \operatorname{div}(\mathbf{g}) & \text{ in } \Omega, \\ w = 0 & \text{ on } \partial\Omega \end{cases}$$

where A^* denotes the transpose of A. By the assumption that A satisfies Assumption 1.4, we find that A^* also satisfies Assumption 1.4. Thus, we know that there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, and Ω , such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then

(5.12)
$$\|\nabla w\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^n)} \lesssim \|\mathbf{g}\|_{L^{p'}_{\omega^{1-p'}}(\Omega;\mathbb{R}^n)}.$$

Using (5.12) and repeating the proof of (4.17), we conclude that, when $p \in (1, 2)$ and $\omega \in A_p(\mathbb{R}^n)$, the weighted Dirichlet problem $(D)_{p,\omega}$ is uniquely solvable and (1.13) holds true. This finishes the proof of (ii) and hence of Theorem 1.10.

To prove Corollary 1.11 by using Theorem 1.10, we need the following Lemma 5.4.

Lemma 5.4. Let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded semi-convex domain. Then there exists a $\delta_0 \in (0, 1)$ such that Ω is a (δ, σ, R) quasi-convex domain for any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$.

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

Proof. Assume that $x \in \partial \Omega$ and Ω has the UEBC constant $R_0 \in (0, \infty)$. Then there exists a $\mathbf{v}_x \in S^{n-1}$, depending on *x*, such that $B(x + R_0 \mathbf{v}_x, R_0) \subset \Omega^{\mathbb{C}}$, which further implies that, for any $r \in (0, R_0)$,

(5.13)
$$B(x + r\mathbf{v}_x, r) \subset B(x + R_0\mathbf{v}_x, R_0) \subset \Omega^{\cup}.$$

Denote by L_x the (n-1)-dimensional plane such that L_x contains x and has a unit normal direction \mathbf{v}_x . For any given $\delta \in (0, 1)$ and any $r \in (0, 2R_0\delta)$, let

$$H(x,\delta,r) := \{ y + t(-\mathbf{v}_x) : y \in L_x, t \in [-\delta r,\infty) \}.$$

Then, by (5.13) and a simple geometric observation, we conclude that, for any given $\delta \in (0, 1)$ and any $r \in (0, 2R_0\delta)$,

(5.14)
$$\Omega \cap B(x,r) \subset H(x,\delta,r) \cap B(x,r).$$

Moreover, from Remark 2.5(i), it follows that Ω is a Lipschitz domain, which, together with the fact that Lipschitz domains are NTA domains, implies that there exist a $\sigma \in (0, 1)$ and an $R_1 \in (0, \infty)$ such that, for any $x \in \partial \Omega$ and $r \in (0, R_1)$, there exists an $x_0 \in \Omega$, depending on x, such that $B(x_0, \sigma r) \subset \Omega \cap B(x, r)$. By this, (5.14), and Remark 2.4(iii), we conclude that there exists a $\delta_0 \in (0, 1)$ such that Ω is a (δ, σ, R) quasi-convex domain for any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$. This finishes the proof of Lemma 5.4.

Next, we show Corollary 1.11 via using Theorem 1.10(ii) and Lemma 5.4.

Proof of Corollary 1.11. By Remark 2.5(iii), we know that, for any given bounded C^1 domain Ω , Ω is a (δ, σ, R) quasi-convex domain for any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$, where $\delta_0 \in (0, 1)$ is a constant depending on Ω . Furthermore, from Lemma 5.4, it follows that, for any bounded semi-convex domain Ω , Ω is a (δ, σ, R) quasi-convex domain for any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$, where $\delta_0 \in (0, 1)$ is a constant depending on Ω . Thus, for any given bounded C^1 domain or semi-convex domain Ω , Ω is a (δ, σ, R) quasi-convex domain for any $\delta \in (0, \delta_0)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$, where $\delta_0 \in (0, 1)$ is a constant depending on Ω . By this and Theorem 1.10(ii), we conclude that the conclusion of this corollary holds true, which completes the proof of Corollary 1.11.

6 Several applications of Theorems 1.8 and 1.10

In this section, we give several applications of the weighted global estimates obtained in Theorems 1.8 and 1.10. More precisely, using Theorems 1.8(ii) and 1.10(ii), we obtain the global gradient estimates, respectively, in (weighted) Lorentz spaces, (Lorentz–)Morrey spaces, (Musielak–)Orlicz spaces (also called generalized Orlicz spaces), and variable Lebesgue spaces. We begin with recalling the following notion of the weighted Lorentz space $L^{q,r}_{\omega}(\Omega)$ on the domain Ω .

Definition 6.1. Let $n \ge 2$ and Ω be a bounded NTA domain in \mathbb{R}^n . Assume that $q \in [1, \infty)$, $r \in (0, \infty]$, and $\omega \in A_p(\mathbb{R}^n)$ with some $p \in [1, \infty)$. The weighted Lorentz space $L^{q,r}_{\omega}(\Omega)$ is defined by setting

$$L^{q,r}_{\omega}(\Omega) := \left\{ f \text{ is measurable on } \Omega : \|f\|_{L^{q,r}_{\omega}(\Omega)} < \infty \right\},$$

where, when $r \in (0, \infty)$,

$$\|f\|_{L^{q,r}_{\omega}(\Omega)} := \left\{ q \int_0^\infty \left[t^q \omega \left(\{ x \in \Omega : |f(x)| > t \} \right) \right]^{r/q} \frac{dt}{t} \right\}^{1/q}$$

and

$$\|f\|_{L^{q,\infty}_{\omega}(\Omega)} := \sup_{t \in (0,\infty)} t[\omega \left(\{x \in \Omega : |f(x)| > t\}\right)]^{1/q}$$

Moreover, the space $L^{q,r}_{\omega}(\Omega; \mathbb{R}^n)$ is defined via replacing $L^p_{\omega}(\Omega)$ in (1.1) by the above $L^{q,r}_{\omega}(\Omega)$ in the definition of $L^p_{\omega}(\Omega; \mathbb{R}^n)$ in (1.2).

It is easy to see that, when $q, r \in [1, \infty)$ and q = r, $L^{q,r}_{\omega}(\Omega) = L^{q}_{\omega}(\Omega)$ and $L^{q,r}_{\omega}(\Omega; \mathbb{R}^{n}) = L^{q}_{\omega}(\Omega; \mathbb{R}^{n})$.

As applications of both Theorems 1.8(ii) and 1.10(ii) and the interpolation theorem of operators in the scale of (weighted) Lorentz spaces, we have the following global gradient estimates for the Dirichlet problem $(D)_p$ in (weighted) Lorentz spaces.

Theorem 6.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, and the matrix A satisfy Assumption 1.4. Assume that $r \in (0, \infty]$, ε_0 is as in Theorem 1.8(i), $p_0 := 3 + \varepsilon_0$ when $n \ge 3$, or $p_0 := 4 + \varepsilon_0$ when n = 2. Then, for any given $p \in (p'_0, p_0)$ and any $\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)$, there exists a positive constant $\delta_0 \in (0, \infty)$, depending only on n, p, the Lipschitz constant of Ω , $[\omega]_{A_{\frac{p}{p'_0}}(\mathbb{R}^n)}$, and $[\omega]_{RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)}$, such that, if A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$ or $A \in VMO(\Omega)$ then for any weak solution $u \in W^{1,2}(\Omega)$ of the problem (D)

and $R \in (0, \infty)$, or $A \in VMO(\Omega)$, then, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the problem $(D)_2$ with $\mathbf{f} \in L^{p,r}_{\omega}(\Omega; \mathbb{R}^n)$, $\nabla u \in L^{p,r}_{\omega}(\Omega; \mathbb{R}^n)$ and

(6.1)
$$\|\nabla u\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant depending only on *n*, *p*, *r*, $[\omega]_{A_{\frac{p}{p'_0}}(\mathbb{R}^n)}$, $[\omega]_{RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)}$, and the Lips-

chitz constant of Ω .

Proof. Let p_0 be as in Theorem 6.2, $p \in (p'_0, p_0), r \in (0, \infty]$, and $\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)$. By (i) and (iv) of Lemma 3.2, we know that there exists an $\epsilon \in (0, \min\{p - p'_0, p_0 - p\})$ such that $\omega \in A_{\frac{p+\epsilon}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p+\epsilon})'}(\mathbb{R}^n)$ and $\omega \in A_{\frac{p-\epsilon}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p-\epsilon})'}(\mathbb{R}^n)$.

For any $\mathbf{f} \in L^2(\Omega; \mathbb{R}^n)$, let $T : \mathbf{f} \mapsto \nabla u_{\mathbf{f}}$, where $u_{\mathbf{f}}$ is the weak solution of the following Dirichlet problem

(6.2)
$$\begin{cases} -\operatorname{div}(A\nabla u_{\mathbf{f}}) = \operatorname{div}(\mathbf{f}) & \text{in } \Omega, \\ u_{\mathbf{f}} = 0 & \text{on } \partial\Omega. \end{cases}$$

From (1.12), it follows that *T* is a well-defined linear operator on both the spaces $L^{p+\epsilon}_{\omega}(\Omega; \mathbb{R}^n)$ and $L^{p-\epsilon}_{\omega}(\Omega; \mathbb{R}^n)$. Let $\theta_0 := \frac{p+\epsilon}{2p}$. Then $\theta_0 \in (0, 1)$ and $\frac{1}{p} = \frac{1-\theta_0}{p-\epsilon} + \frac{\theta_0}{p+\epsilon}$, which, together with (1.12) and the interpolation theorem of operators on Lorentz spaces (see, for instance, [32, Theorem 1.4.19]), further implies that, for any $\mathbf{f} \in L^{p,r}_{\omega}(\Omega; \mathbb{R}^n)$,

$$\|\nabla u_{\mathbf{f}}\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^{n})} = \|T(\mathbf{f})\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^{n})} \lesssim \|\mathbf{f}\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^{n})},$$

where $u_{\rm f}$ is as in (6.2). This finishes the proof of (6.1) and hence of Theorem 6.2.

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

Theorem 6.3. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $p \in (1, \infty)$, $r \in (0, \infty]$, and $\omega \in A_p(\mathbb{R}^n)$. Assume that the matrix A satisfies Assumption 1.4 and Ω is a (δ, σ, R) quasi-convex domain with some δ , $\sigma \in (0, 1)$ and $R \in (0, \infty)$. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, Ω , and $[\omega]_{A_p(\mathbb{R}^n)}$, such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in VMO(\Omega)$, then, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the problem $(D)_2$ with $\mathbf{f} \in L^{p,r}_{\omega}(\Omega; \mathbb{R}^n)$, $\nabla u \in L^{p,r}_{\omega}(\Omega; \mathbb{R}^n)$ and

(6.3)
$$\|\nabla u\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^{p,r}_{\omega}(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant depending only on *n*, *p*, *r*, $[\omega]_{A_p(\mathbb{R}^n)}$, and Ω .

The proof of Theorem 6.3 is similar to that of Theorem 6.2. We omit the details here. Next, we recall the definition of the (Lorentz–)Morrey space on the domain Ω as follows.

Definition 6.4. Assume that $n \ge 2$ and Ω is a bounded NTA domain in \mathbb{R}^n . Let $p \in (1, \infty)$, $r \in (0, \infty]$, and $\theta \in [0, n]$. The Lorentz–Morrey space $L^{p,r;\theta}(\Omega)$ is defined by setting

$$L^{p,r;\theta}(\Omega) := \left\{ f \text{ is measurable on } \Omega : \|f\|_{L^{p,r;\theta}(\Omega)} < \infty \right\},\$$

where

$$||f||_{L^{p,r;\theta}(\Omega)} := \sup_{s \in (0,\text{diam}(\Omega)]} \sup_{x \in \Omega} \left\{ s^{\frac{\theta-n}{p}} ||f||_{L^{p,r}(B(x,s) \cap \Omega)} \right\}.$$

Moreover, the space $L^{p,r;\theta}(\Omega; \mathbb{R}^n)$ is defined via replacing $L^p_{\omega}(\Omega)$ in (1.1) by the above $L^{p,r;\theta}(\Omega)$ in the definition of $L^p_{\omega}(\Omega; \mathbb{R}^n)$ in (1.2).

It is worth pointing out that, when $\theta = n$, the Lorentz–Morrey space $L^{p,r;\theta}(\Omega)$ is just the Lorentz space; in this case, we denote the spaces $L^{p,r;\theta}(\Omega)$ and $L^{p,r;\theta}(\Omega; \mathbb{R}^n)$ simply, respectively, by $L^{p,r}(\Omega)$ and $L^{p,r}(\Omega; \mathbb{R}^n)$. Moreover, when p = r, the space $L^{p,r;\theta}(\Omega)$ is just the *Morrey space*; in this case, we denote the spaces $L^{p,r;\theta}(\Omega)$ and $L^{p,r;\theta}(\Omega; \mathbb{R}^n)$ simply by $\mathcal{M}^{\theta}_{n}(\Omega)$ and $\mathcal{M}^{\theta}_{n}(\Omega; \mathbb{R}^n)$, respectively.

Applying Theorems 6.2 and 6.3, we further obtain the global gradient estimates for the Dirichlet problem $(D)_p$ in Lorentz–Morrey spaces as follows.

Theorem 6.5. Let A and Ω be as in Theorem 6.2, ε_0 as in Theorem 1.8(i), $p_0 := 3 + \varepsilon_0$ when $n \ge 3$, or $p_0 := 4 + \varepsilon_0$ when n = 2, $p \in (p'_0, p_0)$, $r \in (0, \infty]$, and $\theta \in (pn/p_0, n]$. Then there exists a positive constant $\delta_0 \in (0, \infty)$, depending on n, p, r, θ , and Ω , such that, if A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then there exists a positive constant C, depending on n, p, r, θ , and the Lipschitz constant of Ω , such that, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the problem $(D)_2$ with $\mathbf{f} \in L^{p,r;\theta}(\Omega; \mathbb{R}^n)$, $\nabla u \in L^{p,r;\theta}(\Omega; \mathbb{R}^n)$ and

(6.4)
$$\|\nabla u\|_{L^{p,r;\theta}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^{p,r;\theta}(\Omega;\mathbb{R}^n)}.$$

Theorem 6.6. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $p \in (1, \infty)$, $r \in (0, \infty]$, and $\theta \in (0, n]$. Assume that the matrix A satisfies Assumption 1.4 and Ω is a (δ, σ, R) quasi-convex domain with some δ , $\sigma \in (0, 1)$ and $R \in (0, \infty)$. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, r, θ , and Ω , such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the problem $(D)_2$ with $\mathbf{f} \in L^{p,r;\theta}(\Omega; \mathbb{R}^n)$, $\nabla u \in L^{p,r;\theta}(\Omega; \mathbb{R}^n)$ and

(6.5)
$$\|\nabla u\|_{L^{p,r;\theta}(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^{p,r;\theta}(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant depending only on *n*, *p*, *r*, θ , and Ω .

As corollaries of Theorems 6.5 and 6.6, we have the following global gradient estimates in Morrey spaces.

Corollary 6.7. Let $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain and A satisfy Assumption 1.4.

(i) Assume further that Ω is a bounded Lipschitz domain, ε₀ is as in Theorem 1.8(i), p₀ := 3+ε₀ when n ≥ 3, or p₀ := 4 + ε₀ when n = 2, p ∈ (p'₀, p₀), and θ ∈ (pn/p₀, n]. Then there exists a positive constant δ₀ ∈ (0,∞), depending on n, p, θ, and the Lipschitz constant of Ω, such that, if A satisfies the (δ, R)-BMO condition for some δ ∈ (0,δ₀) and R ∈ (0,∞), or A ∈ VMO(Ω), then there exists a positive constant C, depending only on n, p, θ, and the Lipschitz constant of Ω, such that, for any weak solution u ∈ W^{1,2}₀(Ω) of the problem (D)₂ with **f** ∈ M^θ_p(Ω; ℝⁿ), ∇u ∈ M^θ_p(Ω; ℝⁿ) and

 $\|\nabla u\|_{\mathcal{M}^{\theta}_{n}(\Omega;\mathbb{R}^{n})} \leq C \|\mathbf{f}\|_{\mathcal{M}^{\theta}_{n}(\Omega;\mathbb{R}^{n})}.$

(ii) Let $p \in (1, \infty)$ and $\theta \in (0, n]$. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, p, θ , and Ω , such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in \text{VMO}(\Omega)$, then, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the problem $(D)_2$ with $\mathbf{f} \in \mathcal{M}_p^{\theta}(\Omega; \mathbb{R}^n)$, $\nabla u \in \mathcal{M}_p^{\theta}(\Omega; \mathbb{R}^n)$ and

 $\|\nabla u\|_{\mathcal{M}^{\theta}_{n}(\Omega;\mathbb{R}^{n})} \leq C \|\mathbf{f}\|_{\mathcal{M}^{\theta}_{n}(\Omega;\mathbb{R}^{n})},$

where *C* is a positive constant depending only on *n*, *p*, θ , and Ω .

Remark 6.8. Let $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain and A := a + b satisfy Assumption 1.4. For the Dirichlet problem (1.4), the estimates (6.3) and (6.5) were established in [1, Corollary 2.2 and Theorem 2.3] under the assumptions that *a* satisfies the (δ, R) -BMO condition for some small $\delta \in (0, \infty)$ and some $R \in (0, \infty)$, $b \equiv 0$, and Ω is a bounded Lipschitz domain with small Lipschitz constants. Thus, the estimates (6.3) and (6.5) improve [1, Corollary 2.2 and Theorem 2.3] via weakening the assumptions on the matrix *A* and the domain Ω .

Moreover, some estimates similar to (6.3) and (6.5) for the Dirichlet problem of some nonlinear elliptic or parabolic equations on Reifenberg flat domains were obtained in [2, 10, 11, 44, 45].

To show Theorem 6.5 via using Theorem 6.2, we need the following lemma, which is well known (see, for instance, [32, Section 7.1.2] and [44, Lemma 3.4]).

Lemma 6.9. (i) Let $s \in [1, \infty)$, $\omega \in A_s(\mathbb{R}^n)$, $z \in \mathbb{R}^n$, and $k \in (0, \infty)$ be a constant. Assume that $\tau^z(\omega)(\cdot) := \omega(\cdot - z)$ and $\omega_k := \min\{\omega, k\}$. Then $\tau^z(\omega) \in A_s(\mathbb{R}^n)$ and $[\tau^z(\omega)]_{A_s(\mathbb{R}^n)} = [\omega]_{A_s(\mathbb{R}^n)}$, and $\omega_k \in A_s(\mathbb{R}^n)$ and $[\omega_k]_{A_s(\mathbb{R}^n)} \leq c_{(s)}[\omega]_{A_s(\mathbb{R}^n)}$, where $c_{(s)} := 1$ when $s \in [1, 2]$, and $c_{(s)} := 2^{s-1}$ when $s \in (2, \infty)$.

(ii) For any $x \in \mathbb{R}^n$, let $\omega_{\gamma}(x) := |x|^{\gamma}$, where $\gamma \in \mathbb{R}$ is a constant. Then, for any given $s \in (1, \infty)$, $\omega_{\gamma} \in A_s(\mathbb{R}^n)$ if and only if $\gamma \in (-n, n[s-1])$. Moreover, $[\omega_{\gamma}]_{A_s(\mathbb{R}^n)} \leq C_{(n, s, \gamma)}$, where $C_{(n, s, \gamma)}$ is a positive constant depending only on n, s, and γ .

Now, we show Theorem 6.5 by using Theorem 6.2 and Lemma 6.9.

Proof of Theorem 6.5. We prove this theorem via borrowing some ideas from [44, 45]. Let $p \in (p'_0, p_0), r \in (0, \infty]$, and $\theta \in (pn/p_0, n]$, where p_0 is as in Theorem 6.5. Assume that u is the weak solution of the Dirichlet problem (1.4) with $\mathbf{f} \in L^{q,r;\theta}(\Omega; \mathbb{R}^n)$. For any $x, z \in \Omega, \rho \in (0, \text{diam}(\Omega)]$, and $\epsilon \in (0, \theta - \frac{pn}{p_0})$, let

$$\omega_{z}(x) := \min\left\{ |x-z|^{-n+\theta-\epsilon}, \rho^{-n+\theta-\epsilon} \right\}.$$

Then, by Lemma 6.9, we conclude that, for any given $z \in \Omega$, $\omega_z \in A_s(\mathbb{R}^n)$ for any given $s \in (1, \infty)$, and there exists a positive constant $C_{(n, s, \theta)}$, depending only on n, s, and θ , such that $[\omega_z]_{A_s(\mathbb{R}^n)} \leq C_{(n, s, \theta)}$. Moreover, from the assumptions $\theta > np/p_0$ and $\epsilon \in (0, \theta - \frac{pn}{p_0})$, it follows that $\theta - n - \epsilon > -n/(\frac{p_0}{p})'$. By this, and Lemmas 3.3(vi) and 6.9, we conclude that, for any given $z \in \Omega$, $\omega_z \in RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)$ and $[\omega_z]_{RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)} \leq 1$, which, combined with Theorem 6.2 and the assumption that, for any $x \in B(z, \rho)$, $\omega_z(x) = \rho^{-n+\theta-\epsilon}$, further implies that, for any $z \in \Omega$ and $\rho \in (0, \text{diam}(\Omega)]$,

(6.6)
$$\|\nabla u\|_{L^{p,r}(B(z,\rho)\cap\Omega;\mathbb{R}^n)} = \rho^{\frac{n-\theta+\epsilon}{p}} \|\nabla u\|_{L^{p,r}_{\omega_z}(B(z,\rho)\cap\Omega;\mathbb{R}^n)} \lesssim \rho^{\frac{n-\theta+\epsilon}{p}} \|\mathbf{f}\|_{L^{p,r}_{\omega_z}(\Omega;\mathbb{R}^n)}$$

Moreover, similarly to the proofs of [44, (5.12) and (5.14)], we know that, for any $z \in \Omega$ and $\rho \in (0, \operatorname{diam}(\Omega)]$,

$$\|\mathbf{f}\|_{L^{p,r}_{\omega_{z}}(\Omega;\mathbb{R}^{n})} \lesssim \|\mathbf{f}\|_{L^{p,r;\theta}(\Omega;\mathbb{R}^{n})} \rho^{-\frac{r}{p}},$$

which, together with (6.6), implies that, for any $z \in \Omega$ and $\rho \in (0, \operatorname{diam}(\Omega)]$,

$$\|\nabla u\|_{L^{p,r}(B(z,\rho)\cap\Omega;\mathbb{R}^n)} \lesssim \rho^{\frac{n-\theta}{p}} \|\mathbf{f}\|_{L^{p,r;\theta}(\Omega;\mathbb{R}^n)}.$$

From this and the definition of $L^{p,r;\theta}(\Omega; \mathbb{R}^n)$, we deduce that (6.4) holds true, which completes the proof of Theorem 6.5.

Proof of Theorem 6.6. The proof of this theorem is similar to that of Theorem 6.5. We omit the details here. \Box

In what follows, a function $f : [0, \infty) \to [0, \infty]$ is said to be *almost increasing* (resp., *almost decreasing*) if there exists a positive constant $L \in [1, \infty)$ such that, for any $s, t \in [0, \infty)$ satisfying $s \le t, f(s) \le Lf(t)$ [resp., $f(s) \ge Lf(t)$]; in particular, if L := 1, then f is said to be increasing (resp., decreasing). Now, we recall the definitions of weak Φ -functions and Musielak–Orlicz spaces (also called generalized Orlicz spaces) as follows (see, for instance, [20, 59]). Recall that the symbol $t \to 0^+$ means $t \in (0, \infty)$ and $t \to 0$.

Definition 6.10. Let $\varphi : [0, \infty) \to [0, \infty]$ be an increasing function satisfying that

$$\varphi(0) = \lim_{t \to 0^+} \varphi(t) = 0$$
 and $\lim_{t \to \infty} \varphi(t) = \infty$.

- (i) Then φ is called a *weak* Φ -function, denoted by $\varphi \in \Phi_w$, if $t \to \frac{\varphi(t)}{t}$ is almost increasing on $(0, \infty)$.
- (ii) The *left-continuous generalized inverse* of φ, denoted by φ⁻¹, is defined by setting, for any s ∈ [0,∞],

$$\varphi^{-1}(s) := \inf \{t \in [0, \infty) : \varphi(t) \ge s\}.$$

(iii) The *conjugate* Φ -function of φ , denoted by φ^* , is defined by setting, for any $t \in [0, \infty)$,

$$\varphi^*(t) := \sup_{s \in [0,\infty)} \{st - \varphi(s)\}.$$

- (iv) Let $E \subset \mathbb{R}^n$ be a measurable set. A function $\varphi : E \times [0, \infty) \to [0, \infty]$ is called a *Musielak–Orlicz function* (or a *generalized* Φ *-function*) on E if it satisfies
 - (iv)₁ for any $t \in [0, \infty)$, $\varphi(\cdot, t)$ is measurable;
 - (iv)₂ for almost every $x \in E$, $\varphi(x, \cdot) \in \Phi_w$.

Then the set $\Phi_w(E)$ is defined to be the collection of all Musielak–Orlicz functions on *E*.

Definition 6.11. Let $E \subset \mathbb{R}^n$ be a measurable set and $\varphi \in \Phi_w(E)$. For any given $f \in L^1_{loc}(E)$, the *Musielak–Orlicz modular* of f is defined by setting

$$\rho_{\varphi}(f) := \int_{E} \varphi(x, |f(x)|) \, dx.$$

Then the Musielak–Orlicz space (also called generalized Orlicz space) $L^{\varphi}(E)$ is defined by setting

$$L^{\varphi}(E) := \{ u \text{ is measurable on } E :$$

there exists a $\lambda \in (0, \infty)$ such that $\rho_{\varphi}(\lambda f) < \infty$

equipped with the Luxembourg (also called the Luxembourg-Nakano) norm

$$\|u\|_{L^{\varphi}(E)}:=\inf\left\{\lambda\in(0,\infty):\;\rho_{\varphi}\left(\frac{u}{\lambda}\right)\leq 1\right\}.$$

To obtain the global gradient estimates for the Dirichlet problem in the scale of Musielak– Orlicz spaces, we need several additional assumptions for the Musielak–Orlicz function φ . Let $E \subset \mathbb{R}^n$ be a measurable set, $\varphi \in \Phi_w(E)$, and $p \in (0, \infty)$.

Assumption (A0). There exist positive constants $\beta \in (0, 1)$ and $\gamma \in (0, \infty)$ such that, for any $x \in E$, $\varphi(x, \beta\gamma) \le 1 \le \varphi(x, \gamma)$.

Assumption (A1). There exists a $\beta \in (0, 1)$ such that, for any $x, y \in E$ satisfying $|x - y| \le 1$, and any $t \in [1, |x - y|^{-n}], \beta \varphi^{-1}(x, t) \le \varphi^{-1}(y, t)$.

Assumption (A2). There exist β , $\sigma \in (0, \infty)$ and $h \in L^1(E) \cap L^{\infty}(E)$ such that, for any $t \in [0, \sigma]$ and $x, y \in E$,

$$\varphi(x,\beta t) \le \varphi(y,t) + h(x) + h(y).$$

Assumption $(aInc)_p$. The function $s \to \frac{\varphi(x,s)}{s^p}$ is almost increasing uniformly in $x \in E$. Assumption $(aDec)_p$. The function $s \to \frac{\varphi(x,s)}{s^p}$ is almost decreasing uniformly in $x \in E$.

Using the weighted global gradient estimates obtained in Theorems 1.8(ii) and 1.10(ii), and the limited range extrapolation theorem established in [20, Theorem 4.18 and Corollary 4.21] in the scale of Musielak–Orlicz spaces, we obtain the following global gradient estimates in Musielak–Orlicz spaces for the Dirichlet problem $(D)_p$ on bounded Lipschitz domains and (δ, σ, R) quasi-convex domains.

Theorem 6.12. Let A and Ω be as in Theorem 6.2, ε_0 as in Theorem 1.8(i), $p_0 := 3 + \varepsilon_0$ when $n \ge 3$, or $p_0 := 4 + \varepsilon_0$ when n = 2, and p_1 , $p_2 \in (p'_0, p_0)$ with $p_1 \le p_2$. Assume that $\varphi \in \Phi_w(\Omega)$ satisfies Assumptions (A0) – (A2), $(\operatorname{aInc})_{p_1}$, and $(\operatorname{aDec})_{p_2}$. Then there exists a positive constant $\delta_0 \in (0, \infty)$, depending only on n, φ , and the Lipschitz constant of Ω , such that, if A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in \operatorname{VMO}(\Omega)$, then, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the Dirichlet problem (D)₂ with $\mathbf{f} \in L^{\varphi}(\Omega; \mathbb{R}^n)$, $\nabla u \in L^{\varphi}(\Omega; \mathbb{R}^n)$ and

$$\|\nabla u\|_{L^{\varphi}(\Omega;\mathbb{R}^n)} \leq C \|\mathbf{f}\|_{L^{\varphi}(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant depending only on *n*, φ , diam (Ω), and the Lipschitz constant of Ω .

Theorem 6.13. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and p_1 , $p_2 \in (1, \infty)$ with $p_1 \le p_2$. Assume that the matrix A satisfies Assumption 1.4 and $\varphi \in \Phi_w(\Omega)$ satisfies Assumptions (A0)–(A2), $(aInc)_{p_1}$, and $(aDec)_{p_2}$. Then there exists a positive constant $\delta_0 \in (0, 1)$, depending only on n, φ , and Ω , such that, if Ω is a (δ, σ, R) quasi-convex domain and A satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, or $A \in VMO(\Omega)$, then, for any weak solution $u \in W_0^{1,2}(\Omega)$ of the problem $(D)_2$ with $\mathbf{f} \in L^{\varphi}(\Omega; \mathbb{R}^n)$, $\nabla u \in L^{\varphi}(\Omega; \mathbb{R}^n)$ and

(6.7)
$$\|\nabla u\|_{L^{\varphi}(\Omega;\mathbb{R}^n)} \leq C \|\mathbf{f}\|_{L^{\varphi}(\Omega;\mathbb{R}^n)},$$

where *C* is a positive constant depending only on *n*, φ , and Ω .

To prove Theorem 6.12 via using the Rubio de Francia extrapolation theorem in the scale of Musielak–Orlicz spaces, we need the following Lemma 6.14, which is just [20, Corollary 4.21].

Lemma 6.14. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, f and h be two given non-negative measurable functions on Ω , and $1 < p_1 < p < p_2 < \infty$. Assume that, for any given $\omega \in A_{p/p_1}(\mathbb{R}^n) \cap RH_{(p_2/p)'}(\mathbb{R}^n)$,

$$\|f\|_{L^p_{\omega}(\Omega)} \le C \|h\|_{L^p_{\omega}(\Omega)},$$

where *C* is a positive constant depending only on *n*, *p*, Ω , $[\omega]_{A_{p/p_1}(\mathbb{R}^n)}$, and $[\omega]_{RH_{(p_2/p)'}(\mathbb{R}^n)}$. If $\varphi \in \Phi_w(\Omega)$ satisfies Assumptions (A0)–(A2), $(\operatorname{aInc})_{q_1}$, and $(\operatorname{aDec})_{q_2}$ for some $p_1 < q_1 \leq q_2 < p_2$, then there exists a positive constant *C*, depending only on *n*, Ω , and φ , such that $||f||_{L^{\varphi}(\Omega)} \leq C||h||_{L^{\varphi}(\Omega)}$.

Now, we show Theorem 6.12 via using Theorem 1.8 and Lemma 6.14.

Proof of Theorem 6.12. Let *u* be the weak solution of the Dirichlet problem

$$\begin{cases} -\operatorname{div}(A\nabla u) = \operatorname{div}(\mathbf{f}) & \text{ in } \Omega, \\ u = 0 & \text{ on } \partial\Omega. \end{cases}$$

Assume that $p \in (p'_0, p_0)$, where p_0 is as in Theorem 6.2. Then, by Theorem 1.8, we conclude that, for any given $\omega \in A_{\frac{p}{p'_0}}(\mathbb{R}^n) \cap RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)$, there exists a positive constant $\delta_0 \in (0, \infty)$, depending only on *n*, *p*, the Lipschitz constant of Ω , $[\omega]_{A_{\frac{p}{p'_0}}(\mathbb{R}^n)}$, and $[\omega]_{RH_{(\frac{p_0}{p})'}(\mathbb{R}^n)}$, such that, if *A* satisfies the (δ, R) -BMO condition for some $\delta \in (0, \delta_0)$ and $R \in (0, \infty)$, or $A \in VMO(\Omega)$, then

$$\|\nabla u\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)} \lesssim \|\mathbf{f}\|_{L^p_{\omega}(\Omega;\mathbb{R}^n)}$$

From this and Lemma 6.14 with $f := |\nabla u|$, $h := |\mathbf{f}|$, $p_1 := p'_0$, and $p_2 := p_0$, it follows that

$$\|\nabla u\|_{L^{\varphi}(\Omega;\mathbb{R}^n)} \sim \|f\|_{L^{\varphi}(\Omega)} \lesssim \|h\|_{L^{\varphi}(\Omega)} \sim \|\mathbf{f}\|_{L^{\varphi}(\Omega;\mathbb{R}^n)},$$

which completes the proof of Theorem 6.12.

Proof of Theorem 6.13. The proof of this theorem is similar to that of Theorem 6.12. We omit the details here. \Box

To give more corollaries of Theorems 6.12 and 6.13, we recall some necessary notions for variable exponent functions $p(\cdot)$ as follows (see, for instance, [19, 22]). Let $\mathcal{P}(\mathbb{R}^n)$ be the set of all measurable functions $p : \mathbb{R}^n \to [1, \infty)$. For any $p \in \mathcal{P}(\mathbb{R}^n)$, let

(6.8)
$$p_+ := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} p(x) \text{ and } p_- := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} p(x).$$

Recall that a function $p : \mathbb{R}^n \to \mathbb{R}$ is said to satisfy the *local log-Hölder continuity condition* if there exists a positive constant C_{loc} such that, for any $x, y \in \mathbb{R}^n$ with $|x - y| \le 1/2$,

$$|p(x) - p(y)| \le \frac{C_{\text{loc}}}{-\log(|x - y|)};$$

a function $p : \mathbb{R}^n \to \mathbb{R}$ is said to satisfy the *log-Hölder decay condition* (at infinity) if there exist positive constants $C_{\infty} \in (0, \infty)$ and $p_{\infty} \in [1, \infty)$ such that, for any $x \in \mathbb{R}^n$,

$$|p(x) - p_{\infty}| \le \frac{C_{\infty}}{\log(e + |x|)}.$$

If a function *p* satisfies both the local log-Hölder continuity condition and the log-Hölder decay condition, then the function *p* is said to satisfy the *log-Hölder continuity condition*.

Moreover, recall that, for any $\alpha \in (0, 1]$, the *Hölder space* $C^{0,\alpha}(\Omega)$ is defined by setting

$$C^{0,\alpha}(\Omega) := \left\{ g \text{ is continuous on } \Omega : [g]_{C^{0,\alpha}(\Omega)} := \sup_{x, y \in \Omega, x \neq y} \frac{|g(x) - g(y)|}{|x - y|^{\alpha}} < \infty \right\}.$$

Then we have the following two corollaries of Theorems 6.12 and 6.13.

Corollary 6.15. Assume that $p \in \mathcal{P}(\mathbb{R}^n)$ satisfies the log-Hölder continuity condition, ε_0 is as in Theorem 1.8(i), $p_0 := 3 + \varepsilon_0$ when $n \ge 3$, or $p_0 := 4 + \varepsilon_0$ when n = 2, and $p'_0 < p_- \le p_+ < p_0$, where p_- and p_+ are as in (6.8). Then the conclusion of Theorem 6.12 holds true if φ satisfies one of the following cases:

GLOBAL GRADIENT ESTIMATES FOR DIRICHLET PROBLEMS

- (i) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := \phi(t)$, where $\phi \in \Phi_w$ satisfies Assumptions $(aInc)_{p_1}$ and $(aDec)_{p_2}$ with $p'_0 < p_1 \le p_2 < p_0$.
- (ii) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := a(x)t^{p(x)}$, where $C^{-1} \le a \le C$ with C being a positive constant.
- (iii) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^{p(x)} \log(e + t)$.
- (iv) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^p + a(x)t^q$, where $p'_0 satisfy <math>\frac{q}{p} < 1 + \frac{1}{n}$, and $0 \le a \in L^{\infty}(\Omega) \cap C^{0, \frac{n}{p}(q-p)}(\Omega)$.
- (v) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^p + a(x)t^p \log(e + t)$, where $p'_0 , and <math>0 \le a \in L^{\infty}(\Omega)$ satisfies the local log-Hölder continuity condition.

Corollary 6.16. Assume that $p \in \mathcal{P}(\mathbb{R}^n)$ satisfies the log-Hölder continuity condition and $1 < p_- \le p_+ < \infty$, where p_- and p_+ are as in (6.8). Then the conclusion of Theorem 6.13 holds true if φ satisfies one of the following cases:

- (i) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := \phi(t)$, where $\phi \in \Phi_w$ satisfies Assumptions $(aInc)_{p_1}$ and $(aDec)_{p_2}$ with $1 < p_1 \le p_2 < \infty$.
- (ii) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := a(x)t^{p(x)}$, where $C^{-1} \le a \le C$ with C being a positive constant.
- (iii) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^{p(x)} \log(e + t)$.
- (iv) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^p + a(x)t^q$, where $1 satisfy <math>\frac{q}{p} < 1 + \frac{1}{n}$ and $0 \le a \in L^{\infty}(\Omega) \cap C^{0, \frac{n}{p}(q-p)}(\Omega)$.
- (v) for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^p + a(x)t^p \log(e + t)$, where $p \in (1, \infty)$ and $0 \le a \in L^{\infty}(\Omega)$ satisfies the local log-Hölder continuity condition.

Remark 6.17. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and A satisfy Assumption 1.4.

- (i) Assume that, for any x ∈ Ω and t ∈ [0,∞), φ(x,t) := φ(t), where φ is as in Corollary 6.16(i). For the Dirichlet problem (1.4), the estimate (6.7) in this case was obtained in [37, Theorem 3.1] under the assumptions that a satisfies the (δ, R)-BMO condition for some small δ ∈ (0,∞) and some R ∈ (0,∞), b ≡ 0, and Ω is a bounded Reifenberg flat domain. Thus, Corollary 6.16(i) improves [37, Theorem 3.1] via weakening the assumptions on the matrix A and the domain Ω.
- (ii) Assume that, for any $x \in \Omega$ and $t \in [0, \infty)$, $\varphi(x, t) := t^{p(x)}$, where $p(\cdot)$ is as in Corollary 6.16. For the Dirichlet problem (1.4), the estimate (6.7) in this case was established in [13, Theorem 2.5] under the assumptions that *a* has partial small BMO coefficients, $b \equiv 0$, and Ω is a bounded Reifenberg flat domain.

Moreover, a variable exponent type estimate similar to (6.7) for the Dirichlet problem of some *p*-Laplace type elliptic equations on Reifenberg flat domains was also obtained in [8, Theorem 1.4].

References

- [1] K. Adimurthi, T. Mengesha and N. C. Phuc, Gradient weighted norm inequalities for linear elliptic equations with discontinuous coefficients, Appl. Math. Optim. 83 (2021), 327-371.
- [2] K. Adimurthi and N. C. Phuc, Global Lorentz and Lorentz–Morrey estimates below the natural exponent for quasilinear equations, Calc. Var. Partial Differential Equations 54 (2015), 3107-3139.
- [3] P. Auscher, On necessary and sufficient conditions for L^p-estimates of Riesz transforms associated to elliptic operators on ℝⁿ and related estimates, Mem. Amer. Math. Soc. 186 (2007), no. 871, xviii+75 pp.
- [4] P. Auscher and J. M. Martell, Weighted norm inequalities, off-diagonal estimates and elliptic operators. I. General operator theory and weights, Adv. Math. 212 (2007), 225-276.
- [5] P. Auscher and M. Qafsaoui, Observations on $W^{1,p}$ estimates for divergence elliptic equations with VMO coefficients, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) 5 (2002), 487-509.
- [6] A. Banerjee and J. L. Lewis, Gradient bounds for *p*-harmonic systems with vanishing Neumann (Dirichlet) data in a convex domain, Nonlinear Anal. 100 (2014), 78-85.
- [7] A. Barton and S. Mayboroda, Layer potentials and boundary-value problems for second order elliptic operators with data in Besov spaces, Mem. Amer. Math. Soc. 243 (2016), no. 1149, vi+110 pp.
- [8] T. A. Bui, Global $W^{1,p(\cdot)}$ estimate for renormalized solutions of quasilinear equations with measure data on Reifenberg domains, Adv. Nonlinear Anal. 7 (2018), 517-533.
- [9] T. A. Bui and X. T. Duong, Weighted variable exponent Sobolev estimates for elliptic equations with non-standard growth and measure data, NoDEA Nonlinear Differential Equations Appl. 25 (2018), no. 4, Paper No. 28, 37 pp.
- [10] T. A. Bui and X. T. Duong, Weighted Lorentz estimates for parabolic equations with nonstandard growth on rough domains, Calc. Var. Partial Differential Equations 56 (2017), no. 6, Art. 177, 27 pp.
- [11] T. A. Bui and X. T. Duong, Global Lorentz estimates for nonlinear parabolic equations on nonsmooth domains, Calc. Var. Partial Differential Equations 56 (2017), no. 2, Art. 47, 24 pp.
- [12] S.-S. Byun, Elliptic equations with BMO coefficients in Lipschitz domains, Trans. Amer. Math. Soc. 357 (2005), 1025-1046.
- [13] S.-S. Byun, J. Ok and L. Wang, $W^{1,p(\cdot)}$ -regularity for elliptic equations with measurable coefficients in nonsmooth domains, Comm. Math. Phys. 329 (2014), 937-958.
- [14] S.-S. Byun and D. K. Palagachev, Weighted L^p-estimates for elliptic equations with measurable coefficients in nonsmooth domains, Potential Anal. 41 (2014), 51-79.
- [15] S.-S. Byun and L. Wang, The conormal derivative problem for elliptic equations with BMO coefficients on Reifenberg flat domains, Proc. London Math. Soc. (3) 90 (2005), 245-272.
- [16] S.-S. Byun and L. Wang, Elliptic equations with BMO coefficients in Reifenberg domains, Comm. Pure Appl. Math. 57 (2004), 1283-1310.
- [17] S.-S. Byun, F. Yao and S. Zhou, Gradient estimates in Orlicz space for nonlinear elliptic equations, J. Funct. Anal. 255 (2008), 1851-1873.
- [18] L. Caffarelli and I. Peral, On W^{1,p} estimates for elliptic equations in divergence form, Comm.
 Pure Appl. Math. 51 (1998), 1-21.

- [19] D. Cruz-Uribe and A. Fiorenza, Variable Lebesgue Spaces, Foundations and Harmonic Analysis, Applied and Numerical Harmonic Analysis, Birkhäuser/Springer, Heidelberg, 2013.
- [20] D. Cruz-Uribe and P. Hästö, Extrapolation and interpolation in generalized Orlicz spaces, Trans. Amer. Math. Soc. 370 (2018), 4323-4349.
- [21] G. Di Fazio, L^p estimates for divergence form elliptic equations with discontinuous coefficients, Boll. Un. Mat. Ital. A (7) 10 (1996), 409-420.
- [22] L. Diening, P. Harjulehto, P. Hästö and M. Růžička, Lebesgue and Sobolev Spaces with Variable Exponents, Lecture Notes in Mathematics 2017, Springer, Heidelberg, 2011.
- [23] H. Dong and D. Kim, Elliptic equations in divergence form with partially BMO coefficients, Arch. Ration. Mech. Anal. 196 (2010), 25-70.
- [24] H. Dong and D. Kim, The conormal derivative problem for higher order elliptic systems with irregular coefficients, in: Recent Advances in Harmonic Analysis and Partial Differential Equations, 69-97, Contemp. Math. 581, Amer. Math. Soc., Providence, RI, 2012.
- [25] H. Dong and D. Kim, On L^p -estimates for elliptic and parabolic equations with A_p weights, Trans. Amer. Math. Soc. 370 (2018), 5081-5130.
- [26] H. Dong and T. Phan, Mixed-norm L^p-estimates for non-stationary Stokes systems with singular VMO coefficients and applications, J. Differential Equations 276 (2021), 342-367.
- [27] H. Eggleston, Convexity, Cambridge Tracts in Mathematics and Mathematical Physics 47, Cambridge University Press, New York, 1958.
- [28] J. Geng, W^{1,p} estimates for elliptic problems with Neumann boundary conditions in Lipschitz domains, Adv. Math. 229 (2012), 2427-2448.
- [29] J. Geng, Homogenization of elliptic problems with Neumann boundary conditions in nonsmooth domains, Acta Math. Sin. (Engl. Ser.) 34 (2018), 612-628.
- [30] M. Giaquinta, Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems, Annals of Mathematics Studies 105, Princeton University Press, Princeton, NJ, 1983.
- [31] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Reprint of the 1998 edition, Springer-Verlag, Berlin, 2001.
- [32] L. Grafakos, Classical Fourier Analysis, third edition, Graduate Texts in Mathematics 249, Springer, New York, 2014.
- [33] L. Grafakos, Modern Fourier Analysis, third edition, Graduate Texts in Mathematics 250, Springer, New York, 2014.
- [34] D. Jerison and C. E. Kenig, Boundary behavior of harmonic functions in nontangentially accessible domains, Adv. Math. 46 (1982), 80-147.
- [35] D. Jerison and C. E. Kenig, The inhomogeneous Dirichlet problem in Lipschitz domains, J. Funct. Anal. 130 (1995), 161-219.
- [36] H. Jia, D. Li and L. Wang, Global regularity for divergence form elliptic equations on quasiconvex domains, J. Differential Equations 249 (2010), 3132-3147.
- [37] H. Jia, D. Li and L. Wang, Regularity theory in Orlicz spaces for elliptic equations in Reifenberg domains, J. Math. Anal. Appl. 334 (2007), 804-817.
- [38] P. W. Jones, Extension theorems for BMO, Indiana Univ. Math. J. 29 (1980), 41-66.
- [39] C. E. Kenig, Harmonic Analysis Techniques for Second Order Elliptic Boundary Value Problems, CBMS Regional Conference Series in Mathematics 83, American Mathematical Society, Providence, RI, 1994.

- [40] C. E. Kenig and T. Toro, Harmonic measure on locally flat domains, Duke Math. J. 87 (1997), 509-551.
- [41] N. V. Krylov, Parabolic and elliptic equations with VMO coefficients, Comm. Partial Differential Equations 32 (2007), 453-475.
- [42] L. Li and J. Pipher, Boundary behavior of solutions of elliptic operators in divergence form with a BMO anti-symmetric part, Comm. Partial Differential Equations 44 (2019), 156-204.
- [43] V. G. Maz'ya and I. E. Verbitsky, Form boundedness of the general second-order differential operator, Comm. Pure Appl. Math. 59 (2006), 1286-1329.
- [44] T. Mengesha and N. C. Phuc, Global estimates for quasilinear elliptic equations on Reifenberg flat domains, Arch. Ration. Mech. Anal. 203 (2012), 189-216.
- [45] T. Mengesha and N. C. Phuc, Weighted and regularity estimates for nonlinear equations on Reifenberg flat domains, J. Differential Equations 250 (2011), 2485-2507.
- [46] N. G. Meyers, An L^p-estimate for the gradient of solutions of second order elliptic divergence equations, Ann. Scuola Norm. Sup. Pisa (3) 17 (1963), 189-206.
- [47] D. Mitrea, I. Mitrea, M. Mitrea and L. Yan, Coercive energy estimates for differential forms in semi-convex domains, Commun. Pure Appl. Anal. 9 (2010), 987-1010.
- [48] D. Mitrea, M. Mitrea and L. Yan, Boundary value problems for the Laplacian in convex and semiconvex domains, J. Funct. Anal. 258 (2010), 2507-2585.
- [49] E. R. Reifenberg, Solution of the Plateau Problem for *m*-dimensional surfaces of varying topological type, Acta Math. 104 (1960), 1-92.
- [50] D. Sarason, Functions of vanishing mean oscillation, Trans. Amer. Math. Soc. 207 (1975), 391-405.
- [51] G. Seregin, L. Silvestre, V. Šverák and A. Zlatoš, On divergence-free drifts, J. Differential Equations 252 (2012), 505-540.
- [52] Z. Shen, Weighted L^2 estimates for elliptic homogenization in Lipschitz domains, arXiv: 2004.03087.
- [53] Z. Shen, Periodic Homogenization of Elliptic Systems, Operator Theory: Advances and Applications 269, Advances in Partial Differential Equations (Basel), Birkhäuser/Springer, Cham, 2018.
- [54] Z. Shen, The L^p boundary value problems on Lipschitz domains, Adv. Math. 216 (2007), 212-254.
- [55] Z. Shen, Bounds of Riesz transforms on L^p spaces for second order elliptic operators, Ann. Inst. Fourier (Grenoble) 55 (2005), 173-197.
- [56] E. M. Stein, Harmonic Analysis: Real-variable Methods, Orthogonality, and Oscillatory Integrals, Princeton University Press, Princeton, NJ, 1993.
- [57] T. Toro, Doubling and flatness: geometry of measures, Notices Amer. Math. Soc. 44 (1997), 1087-1094.
- [58] L. Wang, A geometric approach to the Calderón–Zygmund estimates, Acta Math. Sin. (Engl. Ser.) 19 (2003), 381-396.
- [59] D. Yang, Y. Liang and L. D. Ky, Real-Variable Theory of Musielak–Orlicz Hardy Spaces, Lecture Notes in Mathematics 2182, Springer, Cham, 2017.
- [60] S. Yang, The Neumann problem of Laplace's equation in semiconvex domains, Nonlinear Anal. 133 (2016), 275-291.

- [61] S. Yang, D. Yang and W. Yuan, The *L^p* Robin problem for Laplace equations in Lipschitz and (semi-)convex domains, J. Differential Equations 264 (2018), 1348-1376.
- [62] S. Yang, D.-C. Chang, D. Yang and W. Yuan, Weighted gradient estimates for elliptic problems with Neumann boundary conditions in Lipschitz and (semi-)convex domains, J. Differential Equations 268 (2020), 2510-2550.
- [63] F. Yao, C. Zhang and S. Zhou, Gradient estimates for a class of quasilinear elliptic equations with measure data, Sci. China Math. 62 (2019), 1719-1730.
- [64] C. Zhang, Gradient estimates for *p*-Laplacian equation in composite Reifenberg domains, Nonlinear Anal. 133 (2016), 134-143.
- [65] C. Zhang, Global weighted estimates for the nonlinear parabolic equations with non-standard growth, Calc. Var. Partial Differential Equations 55 (2016), no. 5, Art. 109, 27 pp.
- [66] C. Zhang and S. Zhou, Global weighted estimates for quasilinear elliptic equations with non-standard growth, J. Funct. Anal. 267 (2014), 605-642.
- [67] J. Zhuge, Weak maximum principle for biharmonic equations in quasiconvex Lipschitz domains, J. Funct. Anal. 279 (2020), no. 12, 108786, 36 pp.

Sibei Yang

School of Mathematics and Statistics, Gansu Key Laboratory of Applied Mathematics and Complex Systems, Lanzhou University, Lanzhou 730000, People's Republic of China

E-mail: yangsb@lzu.edu.cn

Dachun Yang and Wen Yuan (Corresponding author)

Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, People's Republic of China

E-mails: dcyang@bnu.edu.cn(D. Yang) wenyuan@bnu.edu.cn(W. Yuan)