

# On the freeness of certain determinantal hypersurface arrangements in $\mathbb{P}^{14}$

Paulina Wiśniewska and Marek Janasz

January 5, 2022

## Abstract

In the present note we study determinantal arrangements constructed with use of the 3-minors of a  $3 \times 5$  generic matrix of indeterminates. In particular, we show that certain naturally constructed hypersurface arrangements in  $\mathbb{P}_{\mathbb{C}}^{14}$  are free.

**Keywords** hypersurface arrangements, freeness, determinantal arrangements

**Mathematics Subject Classification (2020)** 14N20, 14C20

## 1 Introduction

The main aim of the present note is to find new examples of free hypersurfaces arrangements constructed as the so-called determinantal arrangements. Such determinantal arrangements possess many interesting homological property and some of them will be outlined. On the other side, these arrangements are computationally involving and probably this is the main reason why these object are not well-studied. In the note we focus on determinantal arrangements constructed via the 3 minors of a  $3 \times 5$  generic matrix. Before we present our main results, let us summarize briefly the basic concepts (see [4, 5] for more details). Let  $\mathcal{C} \subset \mathbb{P}^n$  be an arrangement of reduced and irreducible hypersurfaces and let  $\mathcal{C} = V(F)$ , where  $F = f_1 \cdot \dots \cdot f_d$  with  $\text{GCD}(f_i, f_j) = 1$ . Denote by  $\text{Der}(S) = S \cdot \partial_{x_0} \oplus \dots \oplus S \cdot \partial_{x_n}$  the ring of polynomial derivations, where  $S = \mathbb{K}[x_0, \dots, x_n]$  and  $\text{char. } \mathbb{K} = 0$ . If we take  $\theta \in \text{Der}(S)$ , then

$$\theta(f_1 \cdot \dots \cdot f_d) = f_1 \cdot \theta(f_2 \cdot \dots \cdot f_d) + f_2 \cdot \dots \cdot f_d \cdot \theta(f_1).$$

Now we can define the ring of polynomial derivations tangent to  $\mathcal{C}$  as

$$D(\mathcal{C}) = \{\theta \in \text{Der}(S) : \theta(F) \in F \cdot S\}.$$

An inductive application of the Leibniz formula leads us to the following characterization of  $D(\mathcal{C})$ , namely

$$D(\mathcal{C}) = \{\theta \in \text{Der}(S) : \theta(f_i) \in f_i \cdot S \text{ for } i \in \{1, \dots, d\}\}.$$

We have the following (automatic) decomposition

$$D(\mathcal{C}) \simeq E \oplus D_0(\mathcal{C}),$$

where  $E$  is the Euler derivation and  $D_0(\mathcal{C}) = \text{syz}(J_F)$  is the module of syzygies on the Jacobian ideal  $J_F = \langle \partial_{x_0} F, \dots, \partial_{x_n} F \rangle$  of the defining polynomial  $F$ . The freeness of  $\mathcal{C}$  boils

down to a question whether  $\text{pdim}(S/J_F) = 2$ , which is equivalent to  $J_F$  being Cohen-Macaulay. One can show that  $\mathcal{C} \subset \mathbb{P}^n$  given by  $F = 0$  is free if the following condition holds: the minimal resolution of the Milnor algebra  $M(F) = S/J_F$  has the following short form

$$0 \rightarrow \bigoplus_{i=1}^n S(-d_i - (d-1)) \rightarrow S^{n+1}(-d+1) \rightarrow S,$$

and the multiset of integers  $(d_1, \dots, d_n)$ , with  $d_1 \leq \dots \leq d_n$ , is called the set of exponents for  $D_0(\mathcal{C})$ .

The literature devoted to determinantal arrangements is not robust. In this context it is worth recalling a general result by Yim [6, Theorem 3.3], where he is focusing on determinantal arrangements in  $\mathbb{P}_{\mathbb{C}}^{2n-1}$  defined by the products of the 2 minors. For  $i < j$  we denote the 2 minor of the matrix

$$N = \begin{pmatrix} x_1 & x_2 & x_3 & \dots & x_n \\ y_1 & y_2 & y_3 & \dots & y_n \end{pmatrix}$$

by  $\Delta_{ij} = x_i y_j - x_j y_i$ . Consider arrangement  $\mathcal{A}$  defined by the polynomial  $F = \prod_{1 \leq i < j \leq n} \Delta_{ij}$  with  $n \geq 3$ . Then the arrangement  $\mathcal{A}$  is free and a basis of  $D(\mathcal{A})$  is explicitly described.

Our research is motivated by the following question [6, Question 3.4].

**Question 1.1.** Let  $M$  be the  $m \times n$  matrix of indeterminates, and let  $F$  be the product of all maximal minors of  $M$ . Is the arrangement defined by  $F$  free for any  $n > m > 2$ ?

**Remark 1.2.** First of all, if  $\mathcal{C} : F = 0$  is the hypersurfaces defined by the determinant of a generic  $3 \times 3$  matrix of indeterminates, then  $\mathcal{C}$  is far away from being free. Buchweitz and Mond in [1] showed that the arrangement defined by the product of the maximal minors of a generic  $n \times (n+1)$  matrix of indeterminates is free (and it means that we have the freeness property when  $m = 3$  and  $n = 4$ ), so the first non-trivial and unsolved case (to the best of our knowledge) is when  $m = 3$  and  $n = 5$ .

Let us consider the  $3 \times 5$  matrix of indeterminates

$$M = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 & x_5 \\ y_1 & y_2 & y_3 & y_4 & y_5 \\ z_1 & z_2 & z_3 & z_4 & z_5 \end{pmatrix}.$$

Now for a triple  $\{i, j, k\}$  with  $i < j < k$  we construct the 3-minor of  $M$  by taking  $i$ -th,  $j$ -th, and  $k$ -th column. Using the 3-minors we can get 10 hypersurfaces  $H_l \subset \mathbb{P}^{14}$  which are given by the following defining polynomials:

$$\begin{aligned} f_1 &= -x_3 y_2 z_1 + x_2 y_3 z_1 + x_3 y_1 z_2 - x_1 y_3 z_2 - x_2 y_1 z_3 + x_1 y_2 z_3, \\ f_2 &= -x_4 y_2 z_1 + x_2 y_4 z_1 + x_4 y_1 z_2 - x_1 y_4 z_2 - x_2 y_1 z_4 + x_1 y_2 z_4, \\ f_3 &= -x_4 y_3 z_1 + x_3 y_4 z_1 + x_4 y_1 z_3 - x_1 y_4 z_3 - x_3 y_1 z_4 + x_1 y_3 z_4, \\ f_4 &= -x_4 y_3 z_2 + x_3 y_4 z_2 + x_4 y_2 z_3 - x_2 y_4 z_3 - x_3 y_2 z_4 + x_2 y_3 z_4, \\ f_5 &= -x_5 y_2 z_1 + x_2 y_5 z_1 + x_5 y_1 z_2 - x_1 y_5 z_2 - x_2 y_1 z_5 + x_1 y_2 z_5, \\ f_6 &= -x_5 y_3 z_1 + x_3 y_5 z_1 + x_5 y_1 z_3 - x_1 y_5 z_3 - x_3 y_1 z_5 + x_1 y_3 z_5, \\ f_7 &= -x_5 y_3 z_2 + x_3 y_5 z_2 + x_5 y_2 z_3 - x_2 y_5 z_3 - x_3 y_2 z_5 + x_2 y_3 z_5, \\ f_8 &= -x_5 y_4 z_1 + x_4 y_5 z_1 + x_5 y_1 z_4 - x_1 y_5 z_4 - x_4 y_1 z_5 + x_1 y_4 z_5, \end{aligned}$$

$$\begin{aligned} f_9 &= -x_5y_4z_2 + x_4y_5z_2 + x_5y_2z_4 - x_2y_5z_4 - x_4y_2z_5 + x_2y_4z_5, \\ f_{10} &= -x_5y_4z_3 + x_4y_5z_3 + x_5y_3z_4 - x_3y_5z_4 - x_4y_3z_5 + x_3y_4z_5. \end{aligned}$$

Using these 3-minors we would like to explore new examples of free divisors constructed as determinantal arrangements of hypersurfaces.

In order to show the freeness of such arrangements, we are going to use the following criterion due to Saito – see for instance [4, Theorem 8.1]. Let  $\mathcal{C} \subset \mathbb{P}^n$  be a reduced effective divisor defined by a homogeneous equation  $f = 0$ . Now we define the graded module of all Jacobian syzygies as

$$\text{AR}(f) := \left\{ r = (a_0, \dots, a_n) \in S^{n+1} : a_0 \cdot \partial_{x_0}(f) + \dots + a_n \cdot \partial_{x_n}(f) = 0 \right\}.$$

To each Jacobian relation  $r \in \text{AR}(f)$  one can associate a derivation

$$D(r) = a_0 \cdot \partial_{x_0} + \dots + a_n \cdot \partial_{x_n}$$

that kills  $f$ , i.e.,  $D(r)(f) = 0$ . One can additionally show that in fact  $\text{AR}(f)$  is isomorphic, as a graded  $S$ -module, with  $D_0(\mathcal{C})$ .

**Theorem 1.3.** *The homogeneous Jacobian syzygies  $r_i \in \text{AR}(f)$  for  $i \in \{1, \dots, n\}$  form a basis of this  $S$ -module if and only if*

$$\phi(f) = c \cdot f,$$

where  $\phi(f)$  is the determinant of the  $(n+1) \times (n+1)$  matrix  $\Phi(f) = (r_{ij})_{0 \leq i, j \leq n}$ ,  $r_0 := (x_0, \dots, x_n)$ , and  $c$  is a non-zero constant.

Saito's criterion is a very powerful tool under the assumption that we have a set of potential candidates that might form a basis of  $\text{AR}(f)$ , so our work boils down to finding appropriate sets of Jacobian relations that will lead us to a basis of  $\text{AR}(f)$  for a given arrangement  $\mathcal{C} : f = 0$ .

Here is our first result of the note.

**Theorem 1.4.** *Let us consider the following hypersurfaces arrangements*

$$\mathcal{C}_j : F_j = f_1 f_2 f_3 f_4 f_j \quad \text{for } j \in \{5, \dots, 10\}.$$

Then  $C_j$  is free with the exponents  $(\underbrace{1, \dots, 1}_{14 \text{ times}})$ .

**Corollary 1.5.** *In the setting of the above theorem, one has*

$$\text{reg}(S/J_{F_j}) = 13$$

for each  $j \in \{5, \dots, 10\}$ , so we reach an upper bound for the regularity according to the content of [2, Proposition 2.6].

**Remark 1.6.** Of course not every combination of 5 defining equations  $f_i, f_j, f_k, f_l, f_m$  leads to an example of a free determinantal arrangement. Consider  $\mathcal{A} : V(f_1 f_2 f_3 f_5 f_{10}) = 0$ , then the minimal free resolution of the Milnor algebra  $M(F) = S/J_F$  with  $F = f_1 f_2 f_3 f_5 f_{10}$  has the following form:

$$0 \rightarrow S(-19)^3 \rightarrow S^4(-18) \oplus S^{13}(-15) \rightarrow S^{15}(-14) \rightarrow S,$$

so the projective dimension is equal to 3.

Moreover, not every choice of 5 *consecutive* hyperplanes leads to a free arrangement. Consider  $\mathcal{B} : V(f_6 f_7 f_8 f_9 f_{10}) = 0$ , then the minimal free resolution of the Milnor algebra has the following form

$$0 \rightarrow S(-16)^3 \rightarrow S^1(-18) \oplus S^{16}(-15) \rightarrow S^{15}(-14) \rightarrow S,$$

so  $\mathcal{B}$  is not free.

The ultimate goal of the present paper is the understand whether we can expect a positive answer on a (sub)question devoted to the freeness of the full determinantal arrangement in  $\mathbb{P}^{14}$ .

**Question 1.7.** Let us consider the following hypersurfaces arrangements  $\mathcal{H} : V(F) = 0$  defined by  $F = f_1 f_2 f_3 f_4 f_5 f_6 f_7 f_8 f_9 f_{10}$ . Is it true that  $\mathcal{H}$  is free?

Towards approaching the above question, we study mid-step defined arrangements, namely those having the defining equation  $Q_k = f_1 f_2 f_3 f_4 f_5 f_k$  with  $k \in \{6, 7, 8, 9, 10\}$ . In particular, we can show the following results.

**Theorem 1.8.** *Let us consider the hypersurfaces arrangement*

$$\mathcal{H}_k : V(Q_k) = 0$$

*given by  $Q_k = f_1 f_2 f_3 f_4 f_5 f_k$  with  $k \in \{6, 7, 8, 9\}$ . Then  $\mathcal{H}_k$  is free with the exponents  $(\underbrace{1, \dots, 1}_{13 \text{ times}}, 4)$ .*

**Corollary 1.9.** *In the setting of the above theorem, one has*

$$\text{reg}(S/J_{Q_k}) = 19$$

*for each  $k \in \{6, 7, 8, 9\}$ , so we reach an upper bound for the regularity according to the content of [2, Proposition 2.6].*

**Remark 1.10.** If we consider the arrangement  $\mathcal{H}_{10}$  defined by  $Q_{10}$ , then it is not free since the minimal free resolution of the Milnor algebra has the following form:

$$0 \rightarrow S(-22)^3 \rightarrow S^5(-21) \oplus S^{12}(-18) \rightarrow S^{15}(-17) \rightarrow S,$$

which is quite surprising.

Our very ample numerical experiments suggest that the full determinantal arrangement  $\mathcal{H} : f_1 \cdots f_{10} = 0$  should be free with the exponents  $(\underbrace{1, \dots, 1}_{9 \text{ times}}, \underbrace{4, \dots, 4}_{5 \text{ times}})$ . In order to verify our claim we also checked other larger arrangements of hyperplanes, for instance we can verify that  $\mathcal{C} : f_1 f_2 f_3 f_4 f_7 f_8 f_9 = 0$  is free with the exponents  $(\underbrace{1, \dots, 1}_{12 \text{ times}}, 4, 4)$ . However, the derivations of degree 4 seem to us that they do not have a natural geometric or algebraic explanation so it is very hard to find the basis of derivations for  $\mathcal{H}$ . We hope to solve this problem in the nearest future.

## 2 Proofs of the results

We start we our proof of Theorem 1.7.

*Proof.* We are going to apply directly Saito's criterion. In order to do so, we need to find a basis of the  $S$ -modules  $\text{AR}(F_j)$  for each  $j \in \{5, \dots, 10\}$ . This means that in each case we need to find 14 derivations for each  $\text{AR}(F_j)$ . Since for each choice of  $F_j$  the procedure goes along the same lines, let us focus on the first case  $F_5 = f_1 f_2 f_3 f_4 f_5$ .

We start with a group of (obvious to see) derivations, namely

$$\begin{aligned}\theta_1 &= z_1 \cdot \partial_{x_1} + z_2 \cdot \partial_{x_2} + z_3 \cdot \partial_{x_3} + z_4 \cdot \partial_{x_4} + z_5 \cdot \partial_{x_5}, \\ \theta_2 &= z_1 \cdot \partial_{y_1} + z_2 \cdot \partial_{y_2} + z_3 \cdot \partial_{y_3} + z_4 \cdot \partial_{y_4} + z_5 \cdot \partial_{y_5}, \\ \theta_3 &= y_1 \cdot \partial_{x_1} + y_2 \cdot \partial_{x_2} + y_3 \cdot \partial_{x_3} + y_4 \cdot \partial_{x_4} + y_5 \cdot \partial_{x_5}, \\ \theta_4 &= y_1 \cdot \partial_{z_1} + y_2 \cdot \partial_{z_2} + y_3 \cdot \partial_{z_3} + y_4 \cdot \partial_{z_4} + y_5 \cdot \partial_{z_5}, \\ \theta_5 &= x_1 \cdot \partial_{y_1} + x_2 \cdot \partial_{y_2} + x_3 \cdot \partial_{y_3} + x_4 \cdot \partial_{y_4} + x_5 \cdot \partial_{y_5}, \\ \theta_6 &= x_1 \cdot \partial_{z_1} + x_2 \cdot \partial_{z_2} + x_3 \cdot \partial_{z_3} + x_4 \cdot \partial_{z_4} + x_5 \cdot \partial_{z_5}, \\ \theta_7 &= x_2 \cdot \partial_{x_5} + y_2 \cdot \partial_{y_5} + z_2 \cdot \partial_{z_5}, \\ \theta_8 &= x_1 \cdot \partial_{x_5} + y_1 \cdot \partial_{y_5} + z_1 \cdot \partial_{z_5}, \\ \theta_9 &= y_1 \cdot \partial_{y_1} + y_2 \cdot \partial_{y_2} + y_3 \cdot \partial_{y_3} + y_4 \cdot \partial_{y_4} + y_5 \cdot \partial_{y_5} - z_1 \partial_{z_1} - z_2 \partial_{z_2} - z_3 \partial_{z_3} - z_4 \partial_{z_4} - z_5 \partial_{z_5}.\end{aligned}$$

We have additionally 5 non-obvious-to-see relations among the partials derivatives (we have found them with use of **Singular** [3]), namely:

$$\begin{aligned}\theta_{10} &= 5x_5 \cdot \partial_{x_5} + 5y_5 \cdot \partial_{y_5} - z_1 \cdot \partial_{z_1} - z_2 \cdot \partial_{z_2} - z_3 \cdot \partial_{z_3} - z_4 \cdot \partial_{z_4} + 4z_5 \cdot \partial_{z_5}, \\ \theta_{11} &= 5x_4 \cdot \partial_{x_4} + 5y_4 \cdot \partial_{y_4} - 3z_1 \cdot \partial_{z_1} - 3z_2 \cdot \partial_{z_2} - 3z_3 \cdot \partial_{z_3} + 2z_4 \cdot \partial_{z_4} - 3z_5 \cdot \partial_{z_5}, \\ \theta_{12} &= 5x_3 \cdot \partial_{x_3} - 3y_1 \cdot \partial_{y_1} - 3y_2 \cdot \partial_{y_2} + 2y_3 \cdot \partial_{y_3} - 3y_4 \cdot \partial_{y_4} - 3y_5 \cdot \partial_{y_5} + 5z_3 \cdot \partial_{z_3}, \\ \theta_{13} &= 5x_1 \cdot \partial_{x_1} + 5y_1 \cdot \partial_{y_1} + z_1 \cdot \partial_{z_1} - 4z_2 \cdot \partial_{z_2} - 4z_3 \cdot \partial_{z_3} - 4z_4 \cdot \partial_{z_4} - 4z_5 \cdot \partial_{z_5},\end{aligned}$$

and

$$\begin{aligned}\theta_{14} &= 5x_2 \cdot \partial_{x_2} - 3y_1 \cdot \partial_{y_1} + 2y_2 \cdot \partial_{y_2} - 3y_3 \cdot \partial_{y_3} - 3y_4 \cdot \partial_{y_4} - 3y_5 \cdot \partial_{y_5} - z_1 \cdot \partial_{z_1} + 4z_2 \cdot \partial_{z_2} \\ &\quad - z_3 \cdot \partial_{z_3} - z_4 \cdot \partial_{z_4} - z_5 \cdot \partial_{z_5}.\end{aligned}$$

Now we are going to construct Saito's matrix. In order to do so, let us write the coefficients of all  $\theta_i$ 's as the columns, and for the Euler derivation  $E = \sum_{i=1}^5 x_i \cdot \partial_{x_i} + \sum_{j=1}^5 y_j \cdot \partial_{y_j} + \sum_{k=1}^5 z_k \cdot \partial_{z_k}$  we write  $r_0 = (x_1, \dots, x_5, y_1, \dots, y_5, z_1, \dots, z_5)^t$ .

Then we get the following matrix

$$A = \begin{pmatrix} x_1 & z_1 & 0 & y_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 5x_1 & 0 & 0 & 0 \\ x_2 & z_2 & 0 & y_2 & 0 & 0 & 0 & 0 & 0 & 5x_2 & 0 & 0 & 0 & 0 & 0 \\ x_3 & z_3 & 0 & y_3 & 0 & 0 & 0 & 0 & 5x_3 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_4 & z_4 & 0 & y_4 & 0 & 0 & 0 & 5x_4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_5 & z_5 & 0 & y_5 & 0 & 0 & 5x_5 & 0 & 0 & 0 & x_2 & 0 & x_1 & 0 & 0 \\ y_1 & 0 & z_1 & 0 & y_1 & 0 & 0 & 0 & -3y_1 & -3y_1 & 0 & 5y_1 & 0 & x_1 & 0 \\ y_2 & 0 & z_2 & 0 & y_2 & 0 & 0 & 0 & -3y_2 & 2y_2 & 0 & 0 & 0 & x_2 & 0 \\ y_3 & 0 & z_3 & 0 & y_3 & 0 & 0 & 0 & 2y_3 & -3y_3 & 0 & 0 & 0 & x_3 & 0 \\ y_4 & 0 & z_4 & 0 & y_4 & 0 & 0 & 5y_4 & -3y_4 & -3y_4 & 0 & 0 & 0 & x_4 & 0 \\ y_5 & 0 & z_5 & 0 & y_5 & 0 & 5y_5 & 0 & -3y_5 & -3y_5 & y_2 & 0 & y_1 & x_5 & 0 \\ z_1 & 0 & 0 & 0 & -z_1 & y_1 & -z_1 & -3z_1 & 0 & -z_1 & 0 & z_1 & 0 & 0 & x_1 \\ z_2 & 0 & 0 & 0 & -z_2 & y_2 & -z_2 & -3z_2 & 0 & 4z_2 & 0 & -4z_2 & 0 & 0 & x_2 \\ z_3 & 0 & 0 & 0 & -z_3 & y_3 & -z_3 & -3z_3 & 5z_3 & -z_3 & 0 & -4z_3 & 0 & 0 & x_3 \\ z_4 & 0 & 0 & 0 & -z_4 & y_4 & -z_4 & 2z_4 & 0 & -z_4 & 0 & -4z_4 & 0 & 0 & x_4 \\ z_5 & 0 & 0 & 0 & -z_5 & y_5 & 4z_5 & -3z_5 & 0 & -z_5 & z_2 & -4z_5 & z_1 & 0 & x_5 \end{pmatrix}.$$

After some cumbersome computations we obtain

$$\text{Det}(A) = 9375 \cdot F_5,$$

which completes the proof.  $\square$

Now we are going to sketch the proof of Theorem 1.8.

*Proof.* Once again, we are going to apply Saito's criterion. We focus on the case  $k = 7$  since other cases can be show in analogical way. The proof is heavily based on **Singular** computations and experiments. We can find polynomial derivations that preserves  $\mathcal{H}$ , namely

$$\begin{aligned}\theta_1 &= z_1 \cdot \partial_{x_1} + z_2 \cdot \partial_{x_2} + z_3 \cdot \partial_{x_3} + z_4 \cdot \partial_{x_4} + z_5 \cdot \partial_{x_5}, \\ \theta_2 &= z_1 \cdot \partial_{y_1} + z_2 \cdot \partial_{y_2} + z_3 \cdot \partial_{y_3} + z_4 \cdot \partial_{y_4} + z_5 \cdot \partial_{y_5}, \\ \theta_3 &= y_1 \cdot \partial_{x_1} + y_2 \cdot \partial_{x_2} + y_3 \cdot \partial_{x_3} + y_4 \cdot \partial_{x_4} + y_5 \cdot \partial_{x_5}, \\ \theta_4 &= y_1 \cdot \partial_{z_1} + y_2 \cdot \partial_{z_2} + y_3 \cdot \partial_{z_3} + y_4 \cdot \partial_{z_4} + y_5 \cdot \partial_{z_5}, \\ \theta_5 &= 3x_5 \cdot \partial_{x_5} + 3y_5 \cdot \partial_{y_5} - z_1 \cdot \partial_{z_1} - z_2 \cdot \partial_{z_2} - z_3 \cdot \partial_{z_3} - z_4 \cdot \partial_{z_4} + 2z_5 \cdot \partial_{z_5}, \\ \theta_6 &= 2x_4 \cdot \partial_{x_4} + 2y_4 \cdot \partial_{y_4} - z_1 \cdot \partial_{z_1} - z_2 \cdot \partial_{z_2} - z_3 \cdot \partial_{z_3} + z_4 \cdot \partial_{z_4} - z_5 \cdot \partial_{z_5}, \\ \theta_7 &= 3x_3 \cdot \partial_{x_3} + 3y_3 \cdot \partial_{y_3} - 2z_1 \cdot \partial_{z_1} - 2z_2 \cdot \partial_{z_2} + z_3 \cdot \partial_{z_3} - 2z_4 \cdot \partial_{z_4} - 2z_5 \cdot \partial_{z_5}, \\ \theta_8 &= 6x_2 \cdot \partial_{x_2} + 6y_2 \cdot \partial_{y_2} - 5z_1 \cdot \partial_{z_1} + z_2 \cdot \partial_{z_2} - 5z_3 \cdot \partial_{z_3} - 5z_4 \cdot \partial_{z_4} - 5z_5 \cdot \partial_{z_5}, \\ \theta_9 &= x_2 \cdot \partial_{x_5} + y_2 \cdot \partial_{y_5} + z_2 \cdot \partial_{z_5}, \\ \theta_{10} &= 3x_1 \cdot \partial_{x_1} + 3y_1 \cdot \partial_{y_1} + z_1 \cdot \partial_{z_1} - 2z_2 \cdot \partial_{z_2} - 2z_3 \cdot \partial_{z_3} - 2z_4 \cdot \partial_{z_4} - 2z_5 \cdot \partial_{z_5}, \\ \theta_{11} &= x_1 \cdot \partial_{y_1} + x_2 \cdot \partial_{y_2} + x_3 \cdot \partial_{y_3} + x_4 \cdot \partial_{y_4} + x_5 \cdot \partial_{y_5}, \\ \theta_{12} &= x_1 \cdot \partial_{z_1} + x_2 \cdot \partial_{z_2} + x_3 \cdot \partial_{z_3} + x_4 \cdot \partial_{z_4} + x_5 \cdot \partial_{z_5},\end{aligned}$$

$$\begin{aligned}\theta_{13} &= y_1 \cdot \partial_{y_1} + y_2 \cdot \partial_{y_2} + y_3 \cdot \partial_{y_3} + y_4 \cdot \partial_{y_4} + y_5 \cdot \partial_{y_5} - z_1 \cdot \partial_{z_1} - z_2 \cdot \partial_{z_2} - z_3 \cdot \partial_{z_3} - z_4 \cdot \partial_{z_4} \\ &\quad - z_5 \cdot \partial_{z_5},\end{aligned}$$

and

$$\begin{aligned}\theta_{14} &= 3x_1x_3y_2z_2 \cdot \partial_{x_2} + 180x_1x_2y_3z_3 \cdot \partial_{x_3} + (192x_1x_2y_4z_3 - 9x_1x_3y_4z_2 + 12x_1x_3y_2z_4 - \\ &\quad 12x_1x_2y_3z_4) \cdot \partial_{x_4} + (15x_1x_3y_5z_2 - 12x_1x_3y_2z_5) \cdot \partial_{x_5} + (3x_3y_1y_2z_2 + 60x_2y_1y_2z_3 - \\ &\quad 60x_1y_2^2z_3) \cdot \partial_{y_2} + (3x_3y_1y_3z_2 - 3x_1y_3^2z_2 - 120x_3y_1y_2z_3 + 180x_2y_1y_3z_3 + \\ &\quad 120x_1y_2y_3z_3) \cdot \partial_{y_3} + (12x_4y_1y_3z_2 - 9x_3y_1y_4z_2 - 12x_1y_3y_4z_2 - 132x_4y_1y_2z_3 + \\ &\quad 192x_2y_1y_4z_3 + 132x_1y_2y_4z_3 + 12x_3y_1y_2z_4 - 12x_2y_1y_3z_4) \cdot \partial_{y_4} + (15x_3y_1y_5z_2 - \\ &\quad 12x_5y_1y_3z_2 + 12x_1y_3y_5z_2 + 60x_2y_1y_5z_3 - 60x_1y_2y_5z_3 - 12x_3y_1y_2z_5 + 12x_2y_1y_3z_5 - \\ &\quad 12x_1y_2y_3z_5) \cdot \partial_{y_5} + (4x_1y_3z_2^2 - x_3y_1z_2^2 + 4x_3y_2z_1z_2 - 4x_2y_3z_1z_2 + 176x_2y_2z_1z_3 - \\ &\quad 204x_2y_1z_2z_3 + 28x_1y_2z_2z_3) \cdot \partial_{z_2} + (204x_2y_3z_1z_3 - 28x_3y_2z_1z_3 - 24x_2y_1z_3^2 + \\ &\quad 28x_1y_2z_3^2 + 181x_1y_3z_2z_3 - 181x_3y_1z_2z_3) \cdot \partial_{z_3} + (8x_4y_3z_1z_2 - 8x_3y_4z_1z_2 - \\ &\quad 40x_4y_2z_1z_3 + 216x_2y_4z_1z_3 - 180x_4y_1z_2z_3 + 180x_1y_4z_2z_3 + 12x_3y_2z_1z_4 - \\ &\quad 12x_2y_3z_1z_4 - x_3y_1z_2z_4 - 8x_1y_3z_2z_4 - 24x_2y_1z_3z_4 + 40x_1y_2z_3z_4) \cdot \partial_{z_4} + (16x_3y_5z_1z_2 - \\ &\quad 16x_5y_3z_1z_2 - 16x_5y_2z_1z_3 + 192x_2y_5z_1z_3 - 72x_5y_1z_2z_3 + 12x_1y_5z_2z_3 - 12x_3y_2z_1z_5 + \\ &\quad 12x_2y_3z_1z_5 - x_3y_1z_2z_5 + 4x_1y_3z_2z_5 - 132x_2y_1z_3z_5 + 16x_1y_2z_3z_5) \cdot \partial_{z_5}.\end{aligned}$$

We claim that the set  $\{E, \theta_1, \theta_2, \dots, \theta_{14}\}$  gives us a basis for  $D(\mathcal{H})$ . It is enough to observe that the determinant of Saito's matrix  $A$  is equal to

$$\text{Det}(A) = 23328 \cdot Q_7,$$

which completes the proof.  $\square$

## Acknowledgments

We would like to warmly thank Piotr Pokora for useful comments and suggestions. The first author was partially supported by the National Science Center (Poland) Sonata Grant Nr **2018/31/D/ST1/00177**.

## References

- [1] R.-O. Buchweitz and D. Mond, Singularities and Computer Algebra, London Math. Soc. Lecture Note Ser., vol. 324, Cambridge Univ. Press, Cambridge (2006), pp. 41-77.
- [2] L. Busé, A. Dimca, H. Schenck, and G. Sticlaru, The Hessian polynomial and the Jacobian ideal of a reduced surface in  $\mathbb{P}^n$ . *Adv. Math.* **392**: Article ID 108035, 22 p (2021).
- [3] W. Decker, G.-M. Greuel, G. Pfister, and H. Schönemann, SINGULAR 4-1-1 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de>, 2018.
- [4] A. Dimca, *Hyperplane arrangements. An introduction*. Universitext Cham: Springer. xii, 200 p. (2017).
- [5] H. Schenck and Ștefan O. Tohăneanu, Freeness of Conic-Line Arrangements in  $\mathbb{P}^2$ . *Comment. Math. Helv.* **84**: 235 – 258 (2009).
- [6] A. Yim, Homological properties of determinantal arrangements. *J. Algebra* **471**: 220–239 (2017).

Paulina Wiśniewska, Department of Mathematics and Doctoral School, Pedagogical University of Krakow, ul. Podchorążych 2, PL-30-084 Kraków, Poland.

*E-mail address:* `wisniewska.paulina.m@gmail.com`

Marek Janasz, Department of Mathematics, Pedagogical University of Krakow, ul. Podchorążych 2, PL-30-084 Kraków, Poland.

*E-mail address:* `marek.janasz@up.krakow.pl`