

ON THE LOG ABUNDANCE FOR COMPACT KÄHLER 3-FOLDS

OMPROKASH DAS AND WENHAO OU

ABSTRACT. In this article we show that if (X, Δ) is a log canonical compact Kähler 3-fold such that $K_X + \Delta$ is nef and the numerical dimension $\nu(K_X + \Delta) \neq 2$, then $K_X + \Delta$ is semi-ample.

CONTENTS

Part 1. Introduction	2
1. Introduction	2
2. Preliminaries	4
3. Some MMP Results	9
3.1. Termination of dlt flips	10
4. Length of Extremal rays and MMP with scaling	18
4.1. Length of extremal rays	18
4.2. Running MMP with scaling	21
5. Relative MMP for dlt pairs and existence of dlt model	23
6. LMMP and Abundance for Kähler surfaces	25
7. Abundance for 3-fold pairs (X, Δ) with $\kappa(X, K_X + \Delta) \geq 1$	26
7.1. Iitaka Fibration	27
8. $\mathbb{Q}$ -linear system	29
Part 2. Construction of LMMP for pseudo-effective pair	34
9. Non-vanishing theorem	34
9.1. Case of Mori fiber space	35
9.2. Proof of the non-vanishing theorem	47
10. Cone and Contraction Theorems for pseudo-effective pairs	48
10.1. Negative part of $K_X + \Delta$	49
10.2. Bend-and-break theorem for pseudo-effective $K_X + \Delta$	51
10.3. Cone theorem for pseudo-effective DLT pairs	54
10.4. Contraction theorems	55

Omprokash Das is supported by the Start-Up Research Grant(SRG), Grant No. # SRG/2020/000348 of the Science and Engineering Research Board (SERB), Govt. Of India.

Part 3. Log abundance theorem for log canonical pairs	58
11. Reduction step	58
12. Construction of bimeromorphic models	60
13. $\nu = 1$ implies $\kappa > 0$	65
14. Proof of log abundance for log canonical pairs	66
Appendix A. Ramifications for Curves	69
References	71

Part 1. Introduction

1. INTRODUCTION

The Minimal Model Program is one of the fundamental tools for birational classification of algebraic varieties. For a smooth projective variety X , it predicts that after finitely many elementary birational contractions we can find a variety X' with mild singularities such that it is birational to X and satisfies exactly one of the following two properties:

- (1) $K_{X'}$ is nef,
- (2) there is a fibration $g : X' \rightarrow Z$ such that $\dim Z < \dim X'$ and the general fibers of g are (log) Fano varieties.

Observe that $K_{X'}$ being nef is a numerical property, so it doesn't reveal much geometric information about X' . The Abundance Conjecture says that when $K_{X'}$ is nef, it is in fact semi-ample, i.e. there is a $m \in \mathbb{N}$ such that line bundle $\mathcal{O}_{X'}(mK_{X'})$ is generated by its global sections. This leads to a fibration of X' whose general fibers are (log) Calabi-Yau varieties. So one can hope to classify X' by studying these fibers.

For complex projective varieties, the abundance conjecture was proved in dimension 3 by Miyaoka and Kawamata [Miy87, Miy88b, Miy88a], [Kaw92]. The more general case of the log abundance was settled in [KMM94] for 3-dimensional complex projective varieties.

Between 2015 and 2016, Campana, Peternell and Höring established the fundamental results of the minimal model program for Kähler 3-folds in [HP15, HP16, CHP16]. More specifically, in [CHP16] they showed that if X is a compact Kähler 3-fold with terminal singularities and K_X is nef, then K_X is semi-ample. Building upon their results, in this article we prove the following case of the log abundance conjecture.

Theorem 1.1. *Let (X, Δ) be a compact Kähler 3-fold log canonical pair. Assume that $K_X + \Delta$ is nef. Then the following hold:*

- (1) (Non-Vanishing) *There is a sufficiently divisible positive integer $m \in \mathbb{N}$ such that $H^0(X, \mathcal{O}_X(m(K_X + \Delta))) \neq \{0\}$.*

- (2) If the Kodaira dimension $\kappa(X, K_X + \Delta) > 0$, then $K_X + \Delta$ is semi-ample.
- (3) If the numerical dimension $\nu(X, K_X + \Delta) = 0, 1$ or 3 , then $K_X + \Delta$ is semi-ample.

Our strategy is similar to the proof of the log abundance theorem for projective 3-folds as presented in [Kol92] and [KMM94]. Note that the proofs in [Kol92, KMM94] make rigorous use of the log minimal model program for various reduction steps. Unfortunately, the log MMP for compact Kähler 3-folds is not established in full generality in [HP15, HP16, CHP16]. More specifically, if (X, Δ) is a $\mathbb{Q}$ -factorial compact Kähler 3-dimensional klt pair such that X is uniruled and $K_X + \Delta$ is pseudo-effective, then the cone theorem and the existence of log minimal model for (X, Δ) is not established in the papers of Campana, Peternell and Höring. This is one of first obstacles we overcome in this article. Of course our proof of these results are inspired from and quite similar to those in [HP15, HP16, CHP16]. There are numerous other technical difficulties we encounter while trying to follow the strategy as in [Kol92] and [KMM94], many of these have to do with the fact that we work with varieties which are not necessarily projective.

Remark 1.2. Note that in our main Theorem 1.1 above we excluded the case $\nu(K_X + \Delta) = 2$. For projective varieties this case is established in [Kaw92], [Kol92] and [KMM94, KMM04]. When X is a compact Kähler 3-fold with $\mathbb{Q}$ -factorial terminal singularities and K_X is nef and $\nu(K_X) = 2$, it is claimed in [CHP16, Theorem 8.2, page 1013] that K_X is semi-ample. However, we were not able to follow the proof in [CHP16, Theorem 8.2]. The issue arises in the ‘Step 1’ of the proof of [CHP16, Theorem 8.2] on page 1016, where a result of Kawamata, namely [Kaw88, Theorem 9.6] is incorrectly applied.

We note that the authors of [CHP16] recently informed us that there is a more direct proof of [CHP16, Theorem 8.2] for the following cases (see [CHP21]): (i) the algebraic dimension $a(X) \geq 1$, and (ii) X is covered by a family of curves. So the only remaining case is when $a(X) = 0$ and there are no subvarieties through very general points of X . In this case X is conjectured to be bimeromorphic to a quotient of a torus.

It still appears possible that the original proof of Kawamata in [Kaw92] could lead to a proof in the Kähler case. We remark that Kawamata’s original proof (and also the one in [Kol92, Chapter 14]) uses orbifold Chern classes on klt 3-folds and then proves an analog of the Bogomolov-Miyaoka-Yau (BMY) inequality for these Chern classes, see [Kaw92, Lemma 2.7] and [Kol92, Theorem 10.13]. The proof of this new BMY inequality uses ample divisors in an essential way, namely, cutting down the dimension of the ambient variety

to reduce all the computations to a surface. Obviously such a trick does not work for compact Kähler spaces. At the moment we do not know whether this inequality can be proved more directly by some analytic methods on a compact Kähler 3-fold with klt singularities.

Acknowledgement We would like to thank Burt Totaro, Christopher Hacon, Andreas H\"oring and Fr\'ed\'eric Campana for many fruitful conversations.

2. PRELIMINARIES

An *analytic variety* or simply a *variety* is a reduced and irreducible complex space. A *morphism* $f : X \rightarrow Y$ between two complex spaces is a *holomorphic map* between complex spaces. An open set $U \subseteq X$ of a complex space X is called *Zariski open* if the complement $X \setminus U$ is a closed *analytic* subset of X , i.e. there is a sheaf of ideals $\mathcal{I} \subseteq \mathcal{O}_X$ such that $X \setminus U = \text{Supp}(\mathcal{O}_X/\mathcal{I})$. Let $Z \subseteq X$ be a closed analytic subset of X . We say that an open set $U \subseteq X$ contains the *general points of* Z if $\emptyset \neq U \cap Z$ contains a Zariski open subset of Z . Let $f : X \rightarrow Y$ be a morphism between two complex spaces and $\mathcal{P}$ is a property. We say that the *general fibers of* f satisfy $\mathcal{P}$ if the set $Z := \{y \in Y : X_y = f^{-1}(y) \text{ does not satisfy } \mathcal{P}\} \subseteq Y$ is contained in a proper closed analytic subset of Y or equivalently, there is a Zariski open dense subset $U \subseteq Y$ such that all the fibers of f over U satisfy $\mathcal{P}$. Similarly, we say the *very general fibers of* f satisfy $\mathcal{P}$, if the set Z is contained in an at most countable union of proper closed analytic subsets of Y .

A proper morphism $f : X \rightarrow Y$ between two complex spaces is called *generically finite* if there exists a Zariski open dense subset $V \subseteq Y$ such that $f|_{f^{-1}(V)} : f^{-1}(V) \rightarrow V$ is a finite morphism. Let $f : X \rightarrow Y$ be a proper morphism from a normal variety X and D is a prime Weil divisor on X . We say that D is *horizontal* over Y if $f(D) = Y$, otherwise D is called *vertical* over Y . For any $\mathbb{R}$ -Cartier divisor Δ on X , we can uniquely decompose Δ into horizontal and vertical parts over Y as: $\Delta = \Delta_{\text{hor}} + \Delta_{\text{ver}}$, where Δ_{hor} corresponds to the horizontal part of Δ over Y and Δ_{ver} corresponds to the vertical part.

Let $f : X \rightarrow Y$ be a morphism of normal varieties and D is a $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor on X . We will write $D \sim_{\mathbb{Q},f} 0$ or $D \sim_{\mathbb{Q},Y} 0$ to mean that there is a $\mathbb{Q}$ -Cartier divisor L on Y such that $D \sim_{\mathbb{Q}} f^*L$.

Canonical Divisor. Let X be a normal analytic variety. Then the canonical sheaf ω_X is defined as $\omega_X := (\wedge^{\dim X} \Omega_X^1)^{**}$. Note that, in general ω_X does

not correspond to an Weil divisor on X . However, for simplicity and to keep notations as similar as possible to the projective MMP, we will abuse notation in this article and will denote the canonical sheaf by the standard divisorial notation K_X . We will often write $\mathcal{O}_X(K_X)$ to mean ω_X . We note that this notational abuse does not create any complication in this article, except one case, see the proof of Claim 9.9. This is the only place where we had to recognize that K_X is truly not represented by an Weil divisor and thus we can not push it forward by a morphism as a cycle.

An $\mathbb{R}$ -divisor D on X is called a *boundary* (resp. *pure boundary*, resp. *sub-boundary*, resp. *pure sub-boundary*) if the coefficients of D belong to the interval $[0, 1]$ (resp. $[0, 1)$, resp. $(-\infty, 1]$, resp. $(-\infty, 1)$).

Let X be a normal variety and Δ a $\mathbb{Q}$ -divisor on X such that $K_X + \Delta$ is $\mathbb{Q}$ -Cartier. We say that the pair (X, Δ) has *terminal* (resp. *canonical*, resp. *klt*, resp. *plt*, resp. *lc*) singularities if Δ is *effective* and the pair satisfies the definition as in [Kol13, Definition 2.8]. Moreover, if Δ is not necessarily effective, then we will refer the corresponding singularities as *sub-terminal* (resp. *sub-canonical*, resp. *sub-klt*, resp. *sub-plt*, resp. *sub-lc*). We say that (X, Δ) has *dlt* singularities if Δ is effective and the pair satisfy the definition as in [Kol13, Def. 2.8].

In the following we collect some important technical definitions. For a more detailed discussion, we encourage the reader to look at [HP16, HP15, CHP16] and the references therein.

- Definition 2.1.** (i) An analytic variety X is called *Kähler* if there exists a closed positive $(1, 1)$ -form $\omega \in \mathcal{A}_{\mathbb{R}}^{1,1}(X)$ such that the following holds: for every point $x \in X$ there exists an open neighborhood $x \in U$ and a closed embedding $\iota_U : U \rightarrow V$ into an open set $V \subseteq \mathbb{C}^N$, and a strictly plurisubharmonic C^∞ function $f : V \rightarrow \mathbb{R}$ such that $\omega|_{U \cap X_{\text{sm}}} = (i\partial\bar{\partial}f)|_{U \cap X_{\text{sm}}}$. Here X_{sm} is the smooth locus of X .
- (ii) A compact analytic variety X is said to belong to *Fujiki's class $\mathcal{C}$* if one of the following equivalent conditions are satisfied:
- (a) X is a meromorphic image of a compact Kähler variety Y , i.e., there exists a dominant meromorphic map $f : Y \dashrightarrow X$ from a compact Kähler variety Y (see [Fuj79, 4.3, page 34]).
 - (b) X is a holomorphic image of compact Kähler manifold, i.e., there is a surjective morphism $f : Y \rightarrow X$ from a compact Kähler manifold Y (see [Fuj79, Lemma 4.6]).
 - (c) X is bimeromorphic to a compact Kähler manifold (see [Var89, Theorem 3.2, page 51]).

- (iii) On a normal compact analytic variety X we replace the use of Néron-Severi group $\text{NS}(X)_{\mathbb{R}}$ by $H_{\text{BC}}^{1,1}(X)$, the Bott-Chern cohomology of real closed $(1,1)$ -forms with local potentials or equivalently, the closed bi-degree $(1,1)$ -currents with local potentials. See [HP16, Definition 3.1 and 3.6] for more details. More specifically, we define

$$N^1(X) := H_{\text{BC}}^{1,1}(X).$$

- (iv) If X is in Fujiki's class $\mathcal{C}$ and has *rational singularities*, then from [HP16, Eqn. (3)] we know that $N^1(X) = H_{\text{BC}}^{1,1}(X) \subseteq H^2(X, \mathbb{R})$. In particular, the intersection product can be defined in $N^1(X)$ via the cup product of $H^2(X, \mathbb{R})$.
- (v) Let X be a normal compact analytic variety contained in Fujiki's class $\mathcal{C}$. We define $N_1(X)$ to be the vector space of real closed currents of bi-dimension $(1,1)$ modulo the following equivalence relation: $T_1 \equiv T_2$ if and only is

$$T_1(\eta) = T_2(\eta)$$

for all real closed $(1,1)$ -form with local potentials.

- (vi) We define $\overline{\text{NA}}(X) \subseteq N_1(X)$ to be the closed cone generated by the classes of positive closed currents. The Mori cone $\overline{\text{NE}}(X) \subseteq \overline{\text{NA}}(X)$ is defined as the closure of the cone of currents of integration T_C , where $C \subseteq X$ is an irreducible curve.
- (vii) Let X be a normal compact analytic variety and $u \in H_{\text{BC}}^{1,1}(X)$. Then u is called *pseudo-effective* if it can be represented by a bi-degree $(1,1)$ -current $T \in \mathcal{D}^{1,1}(X)$ which is locally of the form $\partial\bar{\partial}f$ for some psh function f . It is called *nef* if it can be represented by a form α with local potentials such that for some Kähler form ω on X and for every $\epsilon > 0$, there exists a $\mathcal{C}^\infty$ function $f_\epsilon \in \mathcal{A}^0(X)$ such that $\alpha + i\partial\bar{\partial}f_\epsilon \geq -\epsilon\omega$.
- (viii) The *nef cone* $\text{Nef}(X) \subseteq N^1(X)$ is the cone generated by nef cohomology classes. Let $\mathcal{K}$ be the open cone in $N^1(X)$ generated by the classes of Kähler forms. By [Dem92, Proposition 6.1.(iii)] (also see [HP16, Remark 3.12]) it follows that the nef cone $\text{Nef}(X)$ is the closure of $\mathcal{K}$, i.e. $\text{Nef}(X) = \overline{\mathcal{K}}$.
- (ix) We say that a normal variety X is $\mathbb{Q}$ -factorial if for every prime Weil divisor $D \subseteq X$, there is a positive integer $k > 0$ such that kD is a Cartier divisor, and for the canonical sheaf $\omega_X := \wedge^{\dim X} \Omega_X^1$, there is a positive integer $m > 0$ such that $(\omega_X^{\otimes m})^{**}$ is a line bundle. It is well known that if X is a $\mathbb{Q}$ -factorial 3-fold and $X \dashrightarrow X'$ is a flip or divisorial contraction, then X' is also $\mathbb{Q}$ -factorial.
- (x) A $\mathbb{Q}$ -Cartier divisor D is called *$\mathbb{Q}$ -effective* if mD is a Cartier divisor and the linear system $|mD|$ is not empty for some $m \in \mathbb{N}$.

- (xi) Let $f : X \rightarrow Y$ be a projective fibration between two normal Kähler varieties. Then f is called a K_X -Mori fiber space if the following conditions are satisfied:
- (a) X and Y are $\mathbb{Q}$ -factorial varieties and X has terminal singularities,
 - (b) $-K_X$ is f -ample, and
 - (c) the relative Picard number $\rho(X/Y) = 1$.

The following are some easy but useful results which will be used throughout the paper.

Lemma 2.2 (Negativity Lemma). [Wan19, Lemma 1.3] *Let $f : X \rightarrow Y$ be a proper bimeromorphic morphism between two normal analytic varieties. Let B be a $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor on X such that $-B$ is f -nef. Then B is effective if and only if f_*B is effective.*

Lemma 2.3. *Let $f : X \rightarrow Y$ be a finite surjective morphism between two normal varieties. Let $D \geq 0$ be an effective integral Weil divisor on Y . Then f^*D is a well defined effective integral Weil divisor on X such that $\text{Supp}(f^*D) = f^{-1}(\text{Supp } D)$.*

Proof. First we explain how to take pullback of a Weil divisor under a finite map. Since f is a finite surjective proper morphism and X and Y are both normal, there exists a dense Zariski open set V contained in the smooth locus of Y such that $f^{-1}(V)$ is also contained in the smooth locus of X , and that $\text{codim}_Y(Y \setminus V) \geq 2$ and $\text{codim}_X(X \setminus f^{-1}(V)) \geq 2$. Then by restricting D to V we do not lose any irreducible component of D . Since V is smooth, $D|_V$ is a Cartier divisor on V . Since $\text{codim}_X(X \setminus f^{-1}(V)) \geq 2$, $f^*(D|_V)$ extends to a unique divisor on X which we define to be the f^*D . By construction, we have $\text{Supp}(f^*D) = f^{-1}(\text{Supp } D)$. $\square$

Lemma 2.4. *Let $f : X \rightarrow Y$ be a finite surjective morphism between two normal varieties. Let $B = \sum b_i B_i$ be a $\mathbb{Q}$ -divisor on Y such that $\sum B_i$ contains the branch locus of f (here we allow $b_i = 0$). Let $(f^* \sum B_i)_{\text{red}} = \sum D_{ij}$, where $f(D_{ij}) = B_i$ for all j and i . Let e_{ij} be the branching order of f around general points of D_{ij} . Then we have the following ramification formula*

$$(2.1) \quad f^*(K_Y + B) = K_X + \sum_{i,j} (1 - (1 - b_i)e_{ij}) D_{ij}.$$

Moreover, if we set $B' := \sum_{i,j} (1 - (1 - b_i)e_{ij}) D_{ij}$, then (X, B') is sub-klt (resp. sub-lc, resp. sub-lc) if and only if (Y, B) is sub-klt (resp. sub-lc, resp. sub-lc).

Proof. Since in the first part we are only interested about divisors, and X and Y are normal, by removing some appropriate closed subsets of codimension at least 2 from X and Y , we may assume that X and Y are both smooth. Then we have

$$(2.2) \quad K_X = f^*K_Y + R,$$

where R is the ramification divisor.

Now observe that we have $f^*B = \sum_{i,j} b_i e_{ij} D_{ij}$, and $R = \sum_{i,j} (e_{ij} - 1) D_{ij}$. Then (2.1) follows from (2.2).

The second part follows from [KM98, Proposition 5.20]. □

Lemma 2.5. *Let $f : X \rightarrow Y$ be a proper surjective morphism between two normal varieties and $\mathcal{L}$ is a line bundle on Y . Then $\mathcal{L}$ is semi-ample if and only if $f^*\mathcal{L}$ is semi-ample.*

Proof. Clearly if $\mathcal{L}$ is semi-ample, then $f^*\mathcal{L}$ is semi-ample. So assume that $f^*\mathcal{L}$ is semi-ample. Replacing $f^*\mathcal{L}$ by a suitable multiple we may assume that $f^*\mathcal{L}$ is generated by global sections. By Stein factorization it is enough to consider two cases: (i) f has connected fibers, i.e. $f_*\mathcal{O}_X = \mathcal{O}_Y$, and (ii) f is a finite morphism.

In the case (i) we have $H^0(Y, \mathcal{L}) \cong H^0(X, f^*\mathcal{L}) \neq \{0\}$. Choose $y \in Y$ and $x \in f^{-1}(y)$. Since $f^*\mathcal{L}$ is globally generated, there is a global section $s \in H^0(X, f^*\mathcal{L})$ such that s does not vanish at x . Let $t \in H^0(Y, \mathcal{L})$ be the unique global section of $\mathcal{L}$ such that $f^*t = s$. Then t does not vanish at y , and hence $\mathcal{L}$ is globally generated.

In case (ii) f is a finite morphism. Replacing f by its Galois closure we may assume that $f : X \rightarrow Y$ is Galois with Galois group G . Choose $y \in Y$ and let $s \in H^0(X, f^*\mathcal{L})$ be a section such that s does not vanish at any point of $f^{-1}(y)$. Let $s_1, s_2, \dots, s_d$ be the set of all Galois conjugates of s . Then s_i does not vanish at any point of $f^{-1}(y)$ for any $i \in \{1, \dots, d\}$. Now observe that $\otimes_{i=1}^d s_i$ is a G -invariant global section of $f^*\mathcal{L}^d$. Therefore there exists a section $t \in H^0(Y, \mathcal{L}^d)$ such that $f^*t = \otimes_{i=1}^d s_i$. Then clearly t does not vanish at $y \in Y$, and hence $\mathcal{L}^d$ is globally generated. □

Lemma 2.6. *Let X be a variety with $\mathbb{Q}$ -factorial terminal singularities such that (X, Δ) has canonical singularities for some $\Delta \geq 0$. Let $f : X \rightarrow Z$ be a projective morphism. Suppose that we run a $(K_X + \Delta)$ -MMP over Z which*

ends with a minimal model $g : (X', \Delta') \rightarrow Z$ over Z , i.e., $K_{X'} + \Delta'$ is g -nef. Further, assume that this MMP does not contract any component of Δ . Then X' has terminal singularities.

Proof. By induction it is enough to show that every $(K_X + \Delta)$ -flip or divisorial contraction over Z preserves the singularity type of X . Let $f : X \rightarrow Y$ be a $(K_X + \Delta)$ -flipping contraction over Z and $f' : X' \rightarrow Y$ its flip over Z . Set $\Delta' := \phi_* \Delta$, where $\phi : X \dashrightarrow X'$ is the induced bimeromorphic map. Then (X', Δ') has canonical singularities. We will show that X' has terminal singularities. To the contrary assume that there exists an exceptional divisor E over X such that $a(E, X') \leq 0$. Then $a(E, X', \Delta') = 0$, since (X', Δ') is canonical. Note that E is also an exceptional divisor over X , since $\phi : X \dashrightarrow X'$ is an isomorphism in codimension 1. We consider two cases below.

Case I: $\text{Center}_X(E) \subseteq \text{Ex}(f)$ or $\text{Center}_{X'}(E) \subseteq \text{Ex}(f')$. In this case from [KM98, Lemma 3.38] it follows that $a(E, X, \Delta) < a(E, X', \Delta') = 0$. This is a contradiction, since (X, Δ) is canonical.

Case II: ϕ is an isomorphism around some neighborhoods of general points of $\text{Center}_X(E)$ and $\text{Center}_{X'}(E)$. In this case $a(E, X) = a(E, X') \leq 0$, this is a contradiction since X has terminal singularities.

Now we will consider the case of divisorial contraction. Assume that $\psi : X \rightarrow Y$ is a $(K_X + \Delta)$ -divisorial contraction over Z . Set $\Delta_Y := \psi_* \Delta$. Since (X, Δ) is canonical and ψ does not contract any component of Δ , by [KM98, Corollary 4.3(2)] (Y, Δ_Y) has canonical singularities. The rest of the argument works similarly as above. □

3. SOME MMP RESULTS

In this section we will collect some results of MMP which are known for projective varieties but not for Kähler spaces. One such result is the termination of flips for dlt pairs; we will show that Kawamata's proof in [Kaw92] works of Kähler spaces after some modifications.

In [Sho92] Shokurov stated and proved the existence of flips for 3-folds klt pairs where the underlying variety X is allowed to be analytic. Based on this result one can easily deduce the following version of the existence of dlt flips; alternatively, see the arguments in [CHP16, Theorem 4.3].

Theorem 3.1 (Existence of flips). [Sho92][CHP16, Theorem 4.3] *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Let $\varphi : X \rightarrow Y$ be a $(K_X + \Delta)$ -flipping contraction. Then the flip $\varphi^+ : X^+ \rightarrow Y$ exists. Moreover, X^+*

is a normal $\mathbb{Q}$ -factorial compact Kähler 3-fold and (X^+, Δ^+) is dlt, where $\Delta^+ = (\varphi^+)_*^{-1}(\varphi_*\Delta)$.

Special termination follows similarly from [Sho92], more specifically, the proof as explained in [Fuj07] works since the MMP for dlt Kähler surfaces is known (see Section 6).

Theorem 3.2 (Special Termination). [CHP16, Theorem 4.20][Fuj07, Sho92, Kol92] *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Set $(X_0, \Delta_0) = (X, \Delta)$ and let*

$$(X_0, \Delta_0) \dashrightarrow (X_1, \Delta_1) \dashrightarrow \cdots (X_i, \Delta_i) \dashrightarrow \cdots$$

be a sequence of $(K_X + \Delta)$ -flips. Then for all $i \gg 0$, the flipping locus (and the flipped locus) is disjoint from $\lfloor \Delta_i \rfloor$. In particular, if for every i the flipping locus is contained in the support of $\lfloor \Delta_i \rfloor$, then the sequence of $(K_{X_i} + \Delta_i)$ -flips terminates.

3.1. Termination of dlt flips. In this subsection we will show that Kawamata's proof [Kaw92] of termination of dlt flips (for algebraic varieties) works for compact Kähler 3-folds after appropriate modifications. First note that Kawamata's notion of 'weakly log terminal' singularity is what we call *divisorially log terminal* or *dlt* singularity in this article.

Theorem 3.3. *Let (X, B) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Let*

$$(3.1) \quad \begin{array}{ccccccc} (X_0, B_0) & \dashrightarrow & (X_1, B_1) & \dashrightarrow & (X_2, B_2) & \dashrightarrow & \cdots \\ & \searrow \varphi_0 & & \swarrow \varphi_0^+ & & \searrow \varphi_1 & \\ & Y_0 & & & & Y_1 & \\ & & & \swarrow \varphi_1^+ & & \searrow \varphi_2 & \\ & & & & & Y_2 & \end{array}$$

be a sequence of $(K_X + B)$ -flips, where $(X_0, B_0) := (X, B)$. Then this is a finite sequence, i.e., every sequence of $(K_X + B)$ -flips terminates.

Remark 3.4. Note that in Kawamata's proof in [Kaw92] there are some terminologies, arguments and tools which are either not available on analytic varieties or hard to work with. We list them out below.

- (1) By special termination (see Theorem 3.2) after a finitely many steps we may assume that $\lfloor B_i \rfloor \cap \text{Ex}(\varphi_i) = \emptyset$ for $i \gg 0$. In Kawamata's proof he replaces X by $X \setminus \lfloor B_i \rfloor$ to assume that (X, B) is klt for all $i \gg 0$. However, we can not do the same since the MMP is not known for open Kähler spaces. Instead in our case we will replace B by $B_i - \lfloor B_i \rfloor$ so that (X, B_i) is klt for $i \gg 0$; note that we can do this since X is $\mathbb{Q}$ -factorial.
- (2) For a klt pair (X, B) , the number of exceptional divisors E over X with discrepancy $a(E, X, B) \leq 0$ is an invariant of the pair (X, B) .

Kawamata denotes this invariant by $e(X, B)$. In page 655 of [Kaw92] he redefines the invariant $e(X, B)$ using *exceptional discrete valuations* of the function field $\mathbb{C}(X)$. However, for an analytic variety X , the field of meromorphic functions $\mathbb{C}(X)$ has transcendental degree over $\mathbb{C} \leq \dim X$ in general, consequently, we can not characterize $e(X, B)$ using valuations in the analytic case. However, this does not create any problem for us, since Kawamata only used valuative characterization of $e(X, B)$ for convenience.

- (3) We note that there are other more subtle technical difficulties which appear when we try to use Kawamata's arguments in the analytic case. They are discussed in the proof of Theorem 3.3 below.

Now we will start proving Theorem 3.3 step by step following Kawamata's idea as in [Kaw92]. We will start with the special case of $e(X, B) = 0$ as in [Kaw92, Lemma 4].

Theorem 3.5. [Kaw92, Lemma 4] *Theorem 3.3 holds if (X, B) is klt and $e(X, B) = e(X_0, B_0) = 0$.*

Proof. Write $B = \sum_i b_i S_i$, $0 \leq b_i < 1$ for all i and let S_i^n be the pushforward of S_i on X_n . Note that we may assume that either $B = 0$ or there is at least one component, say S_1 of B such that $S_1^n \cap \text{Ex}(\varphi_n) \neq \emptyset$ for infinitely many n , since otherwise we can remove all the components S_i of B (as X is $\mathbb{Q}$ -factorial) and replace the pairs (X_n, B_n) by $(X_n, 0)$ for all $n \gg 0$. The rest of the proof works as in the proof of [Kaw92, Lemma 4]. □

Now we will show the existence of a terminal model for klt pairs as in [Kaw92, Theorem 5]. Note that Kawamata's original proof works in our case with minor changes, however, since this result is of independent interest to us and will be repeatedly used in other sections, we will present a detailed proof here.

Theorem 3.6 (Terminalization). [Kaw92, Theorem 5] *Let (X, B) be a compact Kähler 3-fold klt pair. Then there exists a $\mathbb{Q}$ -factorial terminal pair (Y, B_Y) and a projective bimeromorphic morphism $f : Y \rightarrow X$ such that*

- (1) $K_Y + B_Y = f^*(K_X + B)$, and
- (2) *the number of exceptional divisors of f is equal to $e(X, B)$.*

Proof. Let $g_0 : Y_0 \rightarrow X$ be a log resolution of (X, B) such that $K_{Y_0} + B_0 = g_0^*(K_X + B) + F$, where B_0 and F are both effective $\mathbb{Q}$ -divisors without common components, and $\text{Supp } B_0$ is smooth (see [KM98, Proposition 2.36(1)]). Then $g_{0*} B_0 = B$ and (Y_0, B_0) is terminal. Now we will run $(K_{Y_0} + B_0)$ -MMP over X . Note that since g_0 is a projective morphism, X is compact and (Y_0, B_0) is

klt, the existence of $(K_{Y_0} + B_0)$ -negative extremal rays over X and the corresponding contractions (over X) are guaranteed by the Cone and Contraction theorem of Nakayama [Nak87, Theorem 4.12]. Existence of flips follows from Theorem 3.1, and termination from Theorem 3.5, since (Y_0, B_0) has terminal singularities. So we only need to show that the terminal singularities are preserved at every step of this MMP. By [KM98, Corollary 3.43] it is enough to show that if (Y_i, B_i) is terminal and $\varphi_i : (Y_i, B_i) \rightarrow (Y_{i+1}, B_{i+1})$ is a divisorial contraction associated to a $(K_{Y_0} + B_0)$ -MMP over X , then $\text{Ex}(\varphi_i) \not\subseteq \text{Supp}(B_i)$. To the contrary assume that E is an irreducible component of B_i which is contracted by φ_i . Then $a(E, X, B) < 0$. Let $g_i : Y_i \rightarrow X$ be the induced projective bimeromorphic morphism. Then we have $K_{Y_i} + B_i = g_i^*(K_X + B) + F_i$, where $F_i \geq 0$ is the pushforward of F on Y_i . Let $C \subseteq Y_i$ be a curve contained in a fiber of φ . Then $-(K_{Y_i} + B_i) \cdot C > 0$ and $g_i(C) = \text{pt.}$ In particular, $F_i \cdot C < 0$ and thus E is a component of F_i , and hence $a(E, X, B) > 0$, which is a contradiction.

Finally assume that this MMP ends with (Y_N, B_N) , i.e., $K_{Y_N} + B_N$ is g_N -nef and $K_{Y_N} + B_N = g_N^*(K_X + B) + F_N$. Then by the Negativity Lemma 2.2 $F_N \leq 0$, and thus $F_N = 0$, since it is effective. In particular, we have $K_{Y_N} + B_N = g_N^*(K_X + B)$. Set $(Y, B_Y) := (Y_N, B_N)$ and $f := g_N$ and we are done. □

Now we will prove Theorem 3.3 as in [Kaw92, Theorem 1].

Proof of Theorem 3.3. First we reduce the problem to the klt case. By special termination (Theorem 3.2), after finitely many steps the flipping locus does not intersect the support of $\lfloor B_i \rfloor$. Thus reindexing X_i 's we may assume that the flipping locus does not intersect the support of $\lfloor \Delta_i \rfloor$ for all $i \geq 0$. In particular, any $(K_X + B)$ -flip is also a $(K_X + \{B\})$ -flip, where $\{B\}$ is the fractional part of B . Therefore replacing B by $\{B\}$ we may assume that (X, B) is klt and the given sequence is a sequence of klt flips. Now we follow the steps 0, 1, 2 and 3 as in Kawamata's proof of [Kaw92, Theorem 1].

Step 0, 1 and 2 work exactly as in the proof of [Kaw92, Theorem 1] without any change. Note that in the Step 2, for the bend-break type result we can use Theorem 4.2.

Step 3: For a fixed $e \in \mathbb{Z}^+$, [Kaw92, Lemma 7] for $e(X, B) = e$ and Theorem 3.3 for $e(X, B) < e$ together imply Theorem 3.3 for all $\mathbb{Q}$ -factorial klt pairs (X, B) with $e(X, B) = e$.

This step works as in the proof in [Kaw92] for the most part, but there is a non-trivial component to it. More specifically, at a critical point in his proof Kawamata considers some divisors $F_1, F_2, \dots, F_e$ over X such that $a(F_j, X_n, B_n) \leq 0$ for all $j \in \{1, 2, \dots, e\}$ and $n \geq 0$. Then he claims that if $\text{Center}_{X_n}(F_j) \cap \text{Ex}(\varphi_n) = \emptyset$ for all $j \in \{1, 2, \dots, e\}$ and for all $n \gg 0$, then the termination of flips holds. He proves this claim by simply replacing X_n by the open set $U_n = X_n \setminus \text{Center}_{X_n}(F_j)$, which leads to $e(U_n, B_n|_{U_n}) = e - 1 < e$. Then he invokes induction hypothesis to conclude the termination. However, in our case we cannot argue like this, since the MMP is not known to work on open Kähler varieties. Instead, we extract the divisor F_j on a bimeromorphic model of X_n and then work on this model (see Claim 3.7 below). We will discuss our arguments below.

Let (X, B) be a $\mathbb{Q}$ -factorial klt pair with $e(X, B) = e$. If $e(X_n, B_n) < e$ for some $n \geq 0$, then the sequence of the flips terminates by induction hypothesis, so we may assume that $e(X_n, B_n) = e$ for all $n \geq 0$. Now we will define an invariant $\text{discrep}(X, B)^+$ for compact Kähler pairs (X, B) as follows:

$$\text{discrep}(X, B)^+ := \inf\{a(E, X, B) > 0 : E \text{ is an exceptional divisor over } X\}.$$

Since in our case $(X, B) = (X_0, B_0)$ is klt, by passing to a log resolution of (X, B) and then using [KM98, Corollary 2.31(3)] and its proof, it follows that $\text{discrep}(X, B)^+ = c > 0$ for some positive rational number $c > 0$. Then from [KM98, Lemma 3.38] it follows that $\text{discrep}(X_n, B_n)^+ \geq c$ for all $n \geq 0$. Note that under our induction hypothesis, [Kaw92, Lemma 7] holds, and hence there exists a positive integer $q > 0$ such that qD is Cartier and $\mathcal{O}_{X_n}(qK_{X_n})$ is a line bundle, for every Weil divisor D on X_n and for all $n \geq 0$. Now replacing q by a suitable multiple so that qB_0 is Cartier, we may assume that $q(K_{X_n} + B_n)$ is Cartier for all $n \geq 0$. In particular, it follows that for every divisor E over X_n and for all $n \geq 0$, the discrepancy $a(E, X_n, B_n)$ is contained the discrete set $\frac{1}{q}\mathbb{Z}$.


Let $F_1, F_2, \dots, F_e$ be the divisors over X such that $a(F_j, X, B) \leq 0$. Now we claim that $a(F_j, X_n, B_n) \leq 0$ for all $1 \leq j \leq e$ and for every $n \geq 0$. Indeed, if not, then since $e(X_n, B_n) = e$ for all $n \geq 0$, there is an integer $n_0 > 0$ and a divisor E over X_{n_0} distinct from F_j for all $1 \leq j \leq e$ such that $a(E, X_{n_0}, B_{n_0}) \leq 0$. Then from [KM98, Lemma 3.38] it follows that $a(E, X_{n_0-1}, B_{n_0-1}) \leq a(E, X_{n_0}, B_{n_0}) \leq 0$. This implies that $e(X, B) \geq e + 1$, a contradiction.

Now if there is a $j \in \{1, \dots, e\}$ and $n \geq 0$ such that the center $C_{j,n}$ of F_j on X_n is contained $\text{Ex}(\varphi_n)$, then $\sum_{j=1}^e a(F_j, X_n, B_n)$ strictly increases by [KM98, Lemma 3.38]. But since the sum $\sum_{j=1}^e a(F_j, X_n, B_n) \leq 0$ for all $n \geq 0$ and is also contained in the discrete set $\frac{1}{q}\mathbb{Z}$, this is a contradiction. So after reindexing

if necessary, we may assume that C_{jn} is not contained in $\text{Ex}(\varphi_n)$ for any $n \geq 0$. In particular, $|a(F_j, X_n, B_n)| = c_j$ is constant for all $n \geq 0$ (as it is an increasing sequence in $\frac{1}{q}\mathbb{Z}$ bound above by 0). Suppose that $c_1 \geq c_2 \geq \dots \geq c_e$. Next we make the following claim.

Claim 3.7. If $C_{1n} \cap \text{Ex}(\varphi_n) = \emptyset$ for all $n \gg 0$, then Theorem 3.3 holds.

Proof of Claim 3.7. Let $f_n : W_n \rightarrow X_n$ be the extraction of the divisor F_1 with $\text{Center}_X(F_1) = C_{1n}$ by running an appropriate MMP over X ; note that this MMP terminates by our induction hypothesis in this step that Theorem 3.3 holds for $e(X, B) < e$. Then $\rho(W_n/X_n) = 1$, since X_n is $\mathbb{Q}$ -factorial, and hence $\rho(W_n/Y_n) = 2$. Let $K_{W_n} + B_{W_n} = f_n^*(K_{X_n} + B_n)$. Since $f_n(F_1) \cap \text{Ex}(\varphi_n) = \emptyset$, the strict transform of the curves in $\text{Ex}(\varphi_n)$ generate a $(K_{W_n} + B_{W_n})$ -negative extremal ray R_n of the relative Mori cone $\overline{\text{NE}}(W_n/Y_n)$. Let $g_n : W_n \rightarrow Z_n$ be the flipping contraction over Y_n corresponding to extremal ray R_n (see [Nak87, Theorem 4.12(2)]) and $g_n^+ : W_{n+1} \rightarrow Z_n$ is the associated flip over Y_n (see Theorem 3.1), and $h_n : Z_n \rightarrow Y_n$ is the induced morphism. Let $B_{W_{n+1}}$ be the pushforward of B_{W_n} on W_{n+1} . Then $K_{W_{n+1}} + B_{W_{n+1}}$ is g_n^+ -ample and $\rho(W_{n+1}/Z_n) = 1$. Now observe that, since $f_n(F_j) \cap \text{Ex}(\varphi_n) = \emptyset$, from our construction above it follows that $F_j \cong g_n(F_j) \cong g_n^{+^{-1}}(g_n(F_j)) =: F_j^{n+1}$. Moreover, since $\text{Ex}(h_n \circ g_n) = \text{Ex}(\varphi_n \circ f_n)$, it follows that $K_{W_{n+1}} + B_{W_{n+1}}$ is $(h_n \circ g_n^+)$ -nef, and for a curve $C \subseteq W_{n+1}$, $(K_{W_{n+1}} + B_{W_{n+1}}) \cdot C = 0$ if and only if $C \subseteq F_j^{n+1}$ such that $(h_n \circ g_n^+)(C) = \text{pt.}$ Then from the relative Base-point free theorem for projective morphism as in [Nak87, Theorem 4.10], it follows that $K_{W_{n+1}} + B_{W_{n+1}}$ is semi-ample over Y_n . Then there exist projective bimeromorphic morphisms $f_{n+1} : W_{n+1} \rightarrow V_{n+1}$ and $\varphi_{V_{n+1}} : V_{n+1} \rightarrow Y_n$ such that $K_{V_{n+1}} + B_{V_{n+1}} := f_{n+1*}(K_{W_{n+1}} + B_{W_{n+1}})$ is $\varphi_{V_{n+1}}$ -ample and $K_{W_{n+1}} + B_{W_{n+1}} = f_{n+1}^*(K_{V_{n+1}} + B_{V_{n+1}})$. Note that f_{n+1} contracts the divisor F_j^{n+1} .



From the diagram above it follows that $(\pi_{n+1} \circ \phi_n)_*(K_{X_n} + B_n) = \pi_{n+1*}(K_{X_{n+1}} + B_{n+1}) = K_{V_{n+1}} + B_{V_{n+1}}$ and $\text{codim}_{V_{n+1}} \text{Ex}(\varphi_{V_{n+1}}) \geq 2$. Thus $\varphi_{V_{n+1}} : V_{n+1} \rightarrow Y_n$ is a flip of $\varphi_n : X_n \rightarrow Y_n$, and since flip of a flipping contraction is unique, it follows that $V_{n+1} \cong X_{n+1}$ and $\varphi_{V_{n+1}} = \varphi_n^+$.

The above construction gives us a sequence of klt-flips $\psi : (W_n, B_{W_n}) \dashrightarrow (W_{n+1}, B_{W_{n+1}})$ such that $e(W_n, B_{W_n}) = e - 1$ for all $n \geq 0$. Thus by induction this is a finite sequence, i.e. there exists an integer $n_0 \geq 0$ such that $\psi_n : W_n \dashrightarrow W_{n+1}$ is an isomorphism for all $n \geq n_0$. We claim that $\phi_n : X_n \dashrightarrow X_{n+1}$ is an isomorphism for all $n \geq n_0$. Indeed, let Γ_n be the normalization of the graph of ϕ_n for $n \geq n_0$, and $p_n : \Gamma_n \rightarrow X_n$ and $q_n : \Gamma_n \rightarrow X_{n+1}$ are the induced bimeromorphic morphisms. Then by the universal property of graph it follows that there is a proper bimeromorphic morphism $\theta_n : W_n \rightarrow \Gamma_n$ for $n \geq n_0$ such that $f_n = p_n \circ \theta_n$ and $f_{n+1} = q_n \circ \theta_n$ (note that here we are identifying W_n and W_{n+1} via the isomorphism ψ_n). Now since $\rho(W_n/X_n) = 1$, either θ_n is finite or p_n is finite. If θ_n is finite and hence isomorphism, then identifying W_n with Γ_n and f_n with p_n for $n \geq n_0$ we see that $p_n(F_j) = \text{Ex}(\varphi_n)$, this is a contradiction, since $p_n(F_j) \cap \text{Ex}(\varphi_n) = f_n(F_j) \cap \text{Ex}(\varphi_n) = \emptyset$. On the other hand, if p_n is finite, then it is in fact an isomorphism, and thus ϕ_n is a morphism. Then since X_{n+1} is $\mathbb{Q}$ -factorial and ϕ_n is an isomorphism is codimension 1, it follows that ϕ_n is an isomorphism for all $n \geq n_0$. This completes the proof of the claim. □

Thus we may assume that $C_{1n} \cap \text{Ex}(\varphi_n) \neq \emptyset$ for all $n \geq 0$. Then the rest of the proof works exactly as in the proof of Step 3 of [Kaw92, Theorem 1]. □

Using the results above we can run a relative MMP for projective morphism between Kähler varieties of dimension ≤ 3 .

Proposition 3.8 (MMP for Projective Morphisms). *Let (X, Δ) be a $\mathbb{Q}$ -factorial dlt pair and $f : X \rightarrow Y$ a projective surjective morphism between two normal compact Kähler varieties. If $\dim X \leq 3$, then we can run a relative $(K_X + \Delta)$ -MMP over Y which terminates with either a minimal model or a Mori fiber space, according to whether $K_X + \Delta$ is pseudo-effective over Y or not. Moreover, if $X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_n \cdots$ are the steps of a $(K_X + \Delta)$ -MMP over Y , then every X_i is a Kähler variety for $i \geq 0$; additionally, if $\psi : X_n \rightarrow Y'$ is a Mori fiber space over Y , then Y' is also Kähler.*

Proof. Since f is projective, the (relative) cone and contraction theorems are known due to [Nak87, Theorem 4.12]. Since $\dim X \leq 3$, the existence of flips (over Y) follows from [Sho92]. The termination of flips (over Y) follow

from Theorem 3.3. The proof of the fact that a $(K_X + \Delta)$ -MMP over Y terminates either with a minimal model or a Mori fiber space according to whether $K_X + \Delta$ is pseudo-effective over Y or not, works exactly as in the algebraic case, since $f : X \rightarrow Y$ is a projective morphism. If $g_i : X_i \rightarrow Z_i$ is a contraction of a $(K_{X_i} + \Delta_i)$ -negative extremal ray of $\overline{\text{NE}}(X_i/Y)$, then from the relative base-point free theorem [Nak87, Theorem 4.10] it follows that the induced morphism $h_i : Z_i \rightarrow Y$ is projective. Then from [Var89, Proposition 1.3.1, page 24] it follows that Z_i is Kähler. If g_i is a flipping contraction and $g_i^+ : X_{i+1} \rightarrow Z_i$ is the flip, then again X_{i+1} is Kähler by the same argument. In the Mori fiber space case again by a similar argument it follows that Y' is Kähler. $\square$

Corollary 3.9. *Let X be a $\mathbb{Q}$ -factorial compact Kähler 3-fold with klt singularities and $\mu : X' \rightarrow X$ a proper bimeromorphic morphism. Then every fiber of μ is rationally chain connected.*

Proof. Let $\nu : \tilde{X} \rightarrow X$ be a resolution of singularities of X dominating X' . Then it suffices to show that every fiber of ν is rationally chain connected. Thus replacing X' by $\tilde{X}$ and μ by ν we may assume that X' is smooth. We run a $(K_{X'} + \text{Ex}(\mu))$ -MMP over X via Proposition 3.8:

$$(3.2) \quad X' = X'_0 \dashrightarrow X'_1 \dashrightarrow \cdots \dashrightarrow X'_n.$$

Since X is $\mathbb{Q}$ -factorial and has klt singularities, we have $X'_n = X$. Let $\phi_i : X' \dashrightarrow X'_i$ and E_i be the push-forward of $\text{Ex}(\mu)$ by ϕ_i .

If $f_i : X'_i \rightarrow X'_{i+1}$ is a $(K_{X'_i} + E_i)$ -divisorial contraction of a divisor E in the above MMP (3.2), then by construction E is contained in the support of E_i . In this case f_i is also a $(K_{X'_i} + E)$ -divisorial contraction. Now by adjunction $K_E + B_E = (K_{X'_i} + E)|_E$ and (E, B_E) is a klt surface. Note that $-(K_E + B_E)$ is relatively ample with respect to the morphism $f_i|_E : E \rightarrow f_i(E)$. It is well known that E is relatively rationally chain connected, for example, see [HM07, Corollary 1.3].

If instead we have $f_i : X'_i \dashrightarrow X'_{i+1}$ a $(K_{X'_i} + E_i)$ -flip, then the flipping locus is contained in a component E of $\text{Ex}(\mu)$. Let $F = E + (1 - \epsilon)(E_i - E)$ for some $0 < \epsilon \ll 1$. Then f_i is also a $(K_{X'_i} + F)$ -flip. Now by adjunction $(K_{X'_i} + F)|_E = K_E + B_E$ such that (E, B_E) has klt singularities, and in particular, the surface E is $\mathbb{Q}$ -factorial. Note that if $g : X'_i \rightarrow Z$ is the $(K_{X'_i} + F)$ -flipping contraction associated to the flip f_i , then $-(K_E + B_E)$ is relatively ample with respect to the morphism $g|_E : E \rightarrow g(E)$. Let C be any flipping curve, then $(K_E + B_E) \cdot C < 0$ and $C^2 < 0$ (since it is an exceptional curve contained in the surface E). It follows that $(K_E + C) \cdot C < 0$ and hence that C is rational. Finally, we argue

that every flipped curve C^+ is rational. Note that C^+ is exceptional over X and hence must be contained in a component say E^+ of $E_{i+1} := f_{i,*}(E_i)$ (since E_{i+1} is the exceptional locus of $X'_{i+1} \rightarrow X$). Let $g^+ : X'_{i+1} \rightarrow Z$ be the flipped contraction and $E_Z^+ = g_*^+(E^+)$. Let $E^- = (f_i^{-1})_* E^+$ be its strict transform on X_i and $K_{E^-} + B_{E^-} = (K_{X'_i} + E_i)|_{E^-}$. Since (E^-, B_{E^-}) is dlt and $-(K_{E^-} + B_{E^-})$ is $g|_{E^-}$ ample, it follows that $(E_Z, (g|_{E^-})_* B_{E^-})$ is dlt, where $E_Z := g_* E^-$, and in particular E_Z has rational $\mathbb{Q}$ -factorial singularities. Thus C^+ is a rational curve.

From what we have seen above, it follows easily that the fibers of $X'_i \rightarrow X$ are rationally chain connected if and only if the fibers of $X'_{i+1} \rightarrow X$ are rationally chain connected. Since $X'_n = X$, this concludes the proof. $\square$

The following result shows that if X is a uniruled compact Kähler non-algebraic 3-fold with klt singularities, then the base of the MRCC fibration of X is 2-dimensional.

Lemma 3.10. *Let X be a uniruled normal $\mathbb{Q}$ -factorial compact Kähler 3-fold with klt singularities and $\phi : X \dashrightarrow X'$ a bimeromorphic contraction to a $\mathbb{Q}$ -factorial compact Kähler 3-fold X' with klt singularities such that ϕ does not extract any divisor, i.e. $\phi^{-1} : X' \dashrightarrow X$ does not contract any divisor. If the base of the $\text{MRC}(C)$ fibration of X has dimension less than 2, then X' is a projective variety. In particular, if X is non-algebraic, then the base of the $\text{MRC}(C)$ fibration of X has dimension 2.*

Proof. Let $\pi : X \dashrightarrow Z$ be the $\text{MRC}(C)$ fibration (see [CH20, Remark 6.10]). By assumption $\dim Z = 0$ or 1 . First note that, since X has rational singularities, by [Nam02, Corollary 1.7], X is projective if and only if any resolution $\tilde{X}$ of the singularities of X is projective. Note also that by Corollary 3.9, the fibers of $\nu : \tilde{X} \rightarrow X$ are rationally chain connected, thus it follows that $\tilde{X} \dashrightarrow Z$ is also a $\text{MRC}(C)$ fibration. Note that as $\tilde{X}$ is smooth, then its MRC and MRCC fibrations coincide. Thus replacing X by a resolution of singularities we may assume that X is a compact Kähler manifold and $\phi : X \rightarrow X'$ is a morphism. Possibly replacing X further by a higher resolution we may assume that π and ϕ are both morphisms. Since the general fibers of π are rationally connected, by [Deb01, Corollary 4.18] $H^0(F, \Omega_F^i) = 0$ for all $i \geq 1$ where F is a general fiber. We claim that $H^0(X, \Omega_X^2) = 0$. If $\dim Z = 0$, then this is clear, so assume that $\dim Z = 1$. Then observe that the following exact sequence

$$\pi^* \Omega_Z \longrightarrow \Omega_X \longrightarrow \Omega_{X/Z} \longrightarrow 0$$

is left exact over Zariski open dense subset of Z . This follows from the generic smoothness of π and the fact that the MRC fibration is an almost holomorphic map. Restricting this sequence to a general fiber F of π we get the following

short exact sequence

$$(3.3) \quad 0 \longrightarrow \mathcal{O}_F \longrightarrow \Omega_X|_F \longrightarrow \Omega_F \longrightarrow 0.$$

Thus we have a short exact sequence

$$(3.4) \quad 0 \longrightarrow \Omega_F \longrightarrow \Omega_X^2|_F \longrightarrow \Omega_F^2 \longrightarrow 0.$$

Since, as observed above, $H^0(\Omega_F^i) = 0$ for $i = 1, 2$, we have $H^0(F, \Omega_X^2|_F) = 0$. In particular, $s|_F = 0$ for any section $s \in H^0(X, \Omega_X^2)$. Since Ω_X^2 is torsion free, it follows that $H^0(X, \Omega_X^2) = 0$. Then $H^2(X, \mathcal{O}_X) = \overline{H^0(X, \Omega_X^2)} = 0$ and by Kodaira's projectivity criterion we have that X is projective.

Finally, since X' has $\mathbb{Q}$ -factorial klt singularities, running a $(K_X + \text{Ex}(\phi))$ -MMP over X' will recover X' , and hence X' is also projective. $\square$

Lemma 3.11. *Any compact analytic variety (possibly singular) of dimension 1 is a projective algebraic curve.*

Proof. Let C be a compact analytic curve and $\pi : \bar{C} \rightarrow C$ the normalization morphism. Then $\bar{C}$ is a smooth projective algebraic curve. Let H be a very ample divisor on $\bar{C}$. Let $s \in H^0(\bar{C}, \mathcal{O}_{\bar{C}})$ be a section such that the support of the corresponding divisor $\bar{D} := \text{div}(s)_0$ of zeros of s is not contained in $\pi^{-1}(C_{\text{sing}})$. Then there is an effective divisor $D \geq 0$ on C supported on C_{sm} such that $\bar{D} = \pi^*D$. Thus $\mathcal{O}_C(D)$ is a line bundle on C with a section $t \in H^0(C, \mathcal{O}_C(D))$ such that $\text{div}(t)_0 = D$. Then by Grauert's criterion [BHPVdV04, Theorem I.19.3], C is projective. $\square$

Remark 3.12. The above lemma holds more generally for any *reduced* compact complex space of dimension 1 with finitely many irreducible components. The proof follows similarly by working on each component separately.

4. LENGTH OF EXTREMAL RAYS AND MMP WITH SCALING

4.1. Length of extremal rays. In this section following Kawamata's article [Kaw91] we will show that for a compact Kähler 3-fold klt pair (X, Δ) , the length of the rational curves generating the $(K_X + \Delta)$ -negative extremal rays is bounded above by a universal constant.

Lemma 4.1. [Kaw91, Lemma] *Let $f : X \rightarrow Y$ be projective bimeromorphic morphism of normal analytic varieties, H an f -ample Cartier divisor on X and E an irreducible component of $\text{Ex}(f)$ of maximal dimension. Let $e = \dim E$ and $\nu : \bar{E} \rightarrow E$ be the normalization morphism. Further suppose that there is an effective $\mathbb{Q}$ -divisor $\Delta \geq 0$ such that (X, Δ) is klt, and $f(E)$ is a point in Y . Then*

$$(H^{e-1} \cdot (K_X + \Delta) \cdot E) > ((\nu^*H)^{e-1} \cdot K_{\bar{E}}).$$

Proof. This proof is essentially same as Kawamata's original proof in [Kaw91] with minor modifications, which we will discuss now. First observe that the question is local on the base Y and $f(E)$ is a point in Y . Moreover, we have a f -ample Cartier divisor H on X . So we can choose a small Stein open subset $U \subseteq Y$ containing $f(E)$ and positive integers $m, n > 0$ such that mH gives a closed embedding $f^{-1} \hookrightarrow \mathbb{P}^n \times U$. Then replacing Y by U and X by $f^{-1}U$ we may assume that $\iota : X \hookrightarrow \mathbb{P}^n \times Y$ is a closed embedding such that $\text{pr}_2 \circ \iota = f$. Then $\text{pr}_1^* \mathcal{O}(1)$ is a globally generated line bundle on $\mathbb{P}^n \times Y$. We can use the global sections of $(\text{pr}_1 \circ \iota)^* \mathcal{O}(1)$ to cut down the dimension of X as in the proof of [Kaw91, Lemma]. We will need a Bertini-type theorem here to guaranteed that the general hyperplane sections of X are irreducible and normal. This is achieved by [Man82, Corollary II.5]. We will also need a relative Kawamata-Viehweg vanishing theorem in this proof. Such a statement is proved in [Nak87, Theorem 3.7]. With these tools Kawamata's arguments in [Kaw91, Lemma] work in our settings. Interested reader may also consult [Deb01, Lemma 7.47] for a more detailed account of this proof in the algebraic case. $\square$

Theorem 4.2. *Let $f : X \rightarrow Y$ be a projective morphism between two analytic varieties, and $\Delta \geq 0$ a $\mathbb{Q}$ -divisor on X such that (X, Δ) has klt singularities. Let E be a irreducible component of $\text{Ex}(f)$ of maximal dimension, and $n = \dim E - \dim f(E)$. Assume further that $-(K_X + \Delta)$ is f -ample. Then a dense Zariski open subset of E is covered by a family of rational curves $\{\Gamma_\lambda\}_{\lambda \in \Lambda}$ such that $f(\Gamma_\lambda)$ are points in Y and that $-(K_X + \Delta) \cdot \Gamma_\lambda \leq 2n$.*

Proof. First replacing f by its Stein factorization we may assume that f has connected fibers and Y is normal. There are two cases now depending on whether f is bimeromorphic or not. First we consider the case when f is not bimeromorphic, in this case $E = X$. Let F be a general fiber of f . Then F is a normal projective algebraic variety, since f is projective, and $(K_X + \Delta)|_F = K_F + \Delta_F$ for some $\mathbb{Q}$ -divisor $\Delta_F \geq 0$ on F . Let C be a general curve obtained by intersecting $\dim F - 1$ general hyperplanes in F . Then C is a smooth curve contained in the smooth locus F and not contained in $\text{Supp } \Delta$. Since $-(K_F + \Delta_F)$ is ample, $(K_F + \Delta_F) \cdot C < 0$. This implies that $K_F \cdot C \leq (K_F + \Delta_F) \cdot C < 0$, since $\Delta_F \cdot C \geq 0$. Then by [Kol96, Theorem 5.8] through every point $x \in C$ there is a rational curve Γ_x in F such that

$$(4.1) \quad -(K_F + \Delta_F) \cdot \Gamma_x \leq 2 \dim F \frac{-(K_F + \Delta_F) \cdot C}{-K_F \cdot C}.$$

Since F is a general fiber of f , it follows that a dense Zariski open subset of X is covered by a family of rational curves $\{\Gamma_\lambda\}_{\lambda \in \Lambda}$ such that $-(K_X + \Delta) \cdot \Gamma_\lambda \leq 2n$,

where $n = \dim E = \dim X$ in this case.

Now we consider the case when f is bimeromorphic. Since the question is local on the base, replacing Y by a small analytic open set we may assume that Y is embedded (as an analytic subset) into an open subset of $\mathbb{C}^N$. Let $d = \dim f(E)$. Let Y_0 be the intersection of Y by d very general hyperplanes of $\mathbb{C}^N$. Then Y_0 is normal by [Man82, Corollary II.5]. Moreover, $Y_0 \cap f(E)$ is a set of finite points. Shrinking Y further if necessary we may assume that $Y_0 \cap f(E)$ is a single point. Let $X_0 = f^{-1}Y_0$, $\Delta_0 = \Delta \cap X_0$, $E_0 = E \cap X_0$ and $f_0 = f|_{X_0}$. Then again by [Man82, Theorem II.5] and [KM98, Lemma 5.17] it follows that X_0 is normal and (X_0, Δ_0) is klt. Moreover, we also have f_0 is projective and $-(K_{X_0} + \Delta_0)$ is f_0 -ample. Let $\nu : \bar{E}_0 \rightarrow E_0$ be the normalization morphism, and H a f_0 -very ample Cartier divisor on X_0 . Note that E_0 is a projective algebraic variety, since f_0 is a projective morphism and $f_0(E_0)$ is a point in Y_0 . Let C be the intersection of $n - 1$ general members of the linear system $|\nu^*H|$ (note that $\dim E_0 = n$). Then by Bertini's theorem, C is a smooth curve contained in the smooth locus of $\bar{E}_0$. Then by Lemma 4.1 applied to f_0 , we get

$$(4.2) \quad (H^{n-1} \cdot (K_{X_0} + \Delta_0) \cdot E_0) > ((\nu^*H)^{n-1} \cdot K_{\bar{E}_0}).$$

From the left hand side of (4.2) we get $H^{n-1} \cdot (K_{X_0} + \Delta_0) \cdot E_0 = \nu^*(K_{X_0} + \Delta_0) \cdot C < 0$, since $-(K_{X_0} + \Delta_0)$ is ample on E_0 . Thus from the right hand side of (4.2) it follows that $K_{\bar{E}_0} \cdot C < 0$. Then by [Kol96, Theorem 5.8] through every point of $x \in C$ there is a rational curve $\bar{\Gamma}_x$ in $\bar{E}_0$ such that

$$-\nu^*(K_X + \Delta) \cdot \bar{\Gamma}_x \leq 2 \dim \bar{E}_0 \frac{-\nu^*(K_X + \Delta) \cdot C}{-K_{\bar{E}_0} \cdot C}.$$

Taking the images of $\bar{\Gamma}_x$'s in E_0 we see that E_0 is covered by a family of rational curves $\{\Gamma_\lambda\}_{\lambda \in \Lambda}$ such that $-(K_X + \Delta) \cdot \Gamma_\lambda < 2n$, where $n = \dim E - \dim f(E)$. $\square$

In this subsection we discuss how to run a MMP with scaling of an effective divisor. We need the following proposition first.

Proposition 4.3. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold klt pair. Let D be a nef $\mathbb{Q}$ -Cartier divisor and $K_X + \Delta$ is not nef. Further assume that the cone theorem and contraction theorem hold for the $(K + \Delta)$ -MMP. Set*

$$\lambda := \sup\{t \geq 0 : D + t(K_X + \Delta) \text{ is nef}\}.$$

Then λ is a rational number and there is a $(K_X + \Delta)$ -negative extremal ray R such that

$$(D + \lambda(K_X + \Delta)) \cdot R = 0.$$

Proof. Let k be a natural number such that $k(K_X + \Delta)$ and kD are both Cartier. Then by Theorem 4.2, for each $(K_X + \Delta)$ -negative extremal ray R_i for $i \in I$, there is a rational curve C_i such that $R_i = \mathbb{R}_{\geq 0}[C_i]$ and $-(K_X + \Delta) \cdot C_i \leq 2n$, where n is the dimension of X . In particular, we have

$$(4.3) \quad \mu_i = \frac{D \cdot C_i}{-(K_X + \Delta) \cdot C_i} = \frac{kD \cdot C_i}{-k(K_X + \Delta) \cdot C_i} \geq \frac{(kD) \cdot C_i}{2nk} \text{ for all } i \in I.$$

Note that $-k(K_X + \Delta) \cdot C_i$ is an integer contained in the set $\{1, 2, \dots, 2nk\}$ for all i , and the $(kD) \cdot C_i$'s are also integers for all i . Therefore the μ_i 's are contained in the discrete set $\mathcal{S} = \{\frac{m}{l} : m, l \in \mathbb{Z}, m \geq 0, 1 \leq l \leq 2nk\}$. In particular, $\mu := \inf\{\mu_i : i \in I\}$ is contained in $\{\mu_i : i \in I\}$, i.e., μ is rational, and there is an extremal ray $R = R_{i_0}$ for some $i_0 \in I$ such that

$$(D + \mu(K_X + \Delta)) \cdot R = 0.$$

From the cone theorem it follows that $D + \mu(K_X + \Delta)$ is nef and $D + s(K_X + \Delta)$ is not nef for any $s > \mu$, since μ is the minimum of all μ_i . Therefore μ is the nef threshold, and we set $\lambda = \mu$. □

Corollary 4.4. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold klt pair. Suppose that $K_X + \Delta$ is not nef and there is an effective $\mathbb{Q}$ -divisor H on X such that $K_X + \Delta + H$ is nef. Then there is a $(K_X + \Delta)$ -negative extremal ray R and a rational number $0 < \lambda \leq 1$ such that $K_X + \Delta + \lambda H$ is nef and $(K_X + \Delta + \lambda H) \cdot R = 0$.*

Proof. It follows directly from Proposition 4.3 by setting $D = K_X + \Delta + H$. □

4.2. Running MMP with scaling. Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold klt pair. Suppose that $K_X + \Delta$ is not nef and there is an effective $\mathbb{Q}$ -divisor H on X such that $K_X + \Delta + H$ is nef. We will run a $(K_X + \Delta)$ -MMP with the scaling of H . Set $(X_0, \Delta_0) = (X, \Delta)$, $H_0 = H$, and let $\lambda_0 = \inf\{t \geq 0 : K_X + \Delta + tH \text{ is nef}\}$. If $\lambda_0 = 0$, then $K_X + \Delta$ is nef and we stop. Otherwise, by Corollary 4.4 there exists a $(K_{X_0} + \Delta_0)$ -negative extremal ray, say R_0 , such that $(K_{X_0} + \Delta_0 + \lambda_0 H) \cdot R_0 = 0$. Let $\varphi_0 : X_0 \rightarrow Y$ be the associated contraction. Then $K_{X_0} + \Delta_0 + \lambda_0 H = \varphi_0^* L$, for some $\mathbb{Q}$ -Cartier divisor L on Y . There are two cases now

- (1) $\varphi : X_0 \rightarrow Y$ is a divisorial contraction, or
- (2) φ is a flipping contraction.

If φ is a divisorial contraction, then Y is a normal $\mathbb{Q}$ -factorial compact Kähler 3-fold. In this case we set $X_1 = Y$, $\Delta_1 = \varphi_* \Delta_0$ and $H_1 = \varphi_* H_0$. Then we have (X_1, Δ_1) a klt pair such that $K_{X_1} + \Delta_1 + \lambda_0 H_1$ is nef. We then set

$\lambda_1 = \inf\{t \geq 0 : K_{X_1} + \Delta_1 + tH_1 \text{ is nef}\}$ and continue the program.

If φ is a flipping contraction, then Y is not $\mathbb{Q}$ -factorial, so we take a flip $\varphi^+ : (X_0^+, \Delta_0^+) \rightarrow Y$ of φ , where $\Delta_0^+ = \phi_*\Delta$ and $H^+ = \phi_*H$, and $\phi : X \rightarrow X^+$ is the induced birational map. Then X^+ is a normal $\mathbb{Q}$ -factorial compact Kähler 3-fold. Set $X_1 = X^+$, $\Delta_1 = \Delta^+$ and $H_1 = H^+$. Then we have (X_1, Δ_1) a klt pair such that $K_{X_1} + \Delta_1 + \lambda_0 H_1$ is nef. We then set $\lambda_1 = \inf\{t \geq 0 : K_{X_1} + \Delta_1 + tH_1 \text{ is nef}\}$ and continue the program.

Note that in way we get a monotonically decreasing sequence $\lambda_0 \geq \lambda_1 \geq \dots$. This program will stop if either $\lambda_i = 0$ for some i , in which case $K_{X_i} + \Delta_i$ is nef, or there is a contraction $\varphi : X_i \rightarrow Y_i$ such that $\dim Y_i < \dim X_i$ and $-(K_{X_i} + \Delta_i)$ is φ_i -ample; this is called a $(K_{X_i} + \Delta_i)$ -Mori fiber space.

Theorem 4.5 (Special MMP). *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold klt pair. Suppose that $K_X + \Delta$ is not nef and there is an effective $\mathbb{Q}$ -divisor H on X such that $K_X + \Delta + H$ is nef. Then we can run a $(K_X + \Delta)$ -MMP with the scaling of H which satisfies exactly one of following two properties:*

$$(X, \Delta) = (X_0, \Delta_0) \dashrightarrow (X_1, \Delta_1) \dashrightarrow \dots (X_i, \Delta_i) \dashrightarrow \dots$$

- (1) *Every step of this MMP is $(K_X + \Delta + H)$ -trivial, i.e., $(K_{X_i} + \Delta_i + H_i) \cdot R_i = 0$ for all $i \geq 0$, where R_i is the corresponding extremal ray. In this case the program terminates with either a $(K_X + \Delta)$ -minimal model or $(K_X + \Delta)$ -Mori fiber space,*

or

- (2) *there is a smallest $i \geq 0$ such that $K_{X_i} + \Delta_i + (1 - \varepsilon)H_i$ is nef for all $0 < \varepsilon \ll 1$. In this case we stop the program at this stage.*

Proof. Using the same notations as in the beginning of Subsection 4.2 we see that the property (1) holds if $\lambda_i = 1$ for all $i \leq k - 1$ and $\lambda_k = 0$ for some $k \geq 1$, such that $K_{X_k} + \Delta_k$ is either nef or there is a $(K_{X_k} + \Delta_k)$ -Mori fiber space $g : X_k \rightarrow Z$. Otherwise, assume that $0 < \lambda_i < 1$ with the smallest index $i \geq 0$. Then $\lambda_{i-1} = 1$ and $K_{X_i} + \Delta_i + (1 - \varepsilon)H_i$ is nef for all $0 < \varepsilon \ll 1$ (we choose $\lambda_{-1} = 1$). $\square$

We will also need the following version of relative directed MMP.

Lemma 4.6. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold lc pair such that X has klt singularities. Let $f : X \rightarrow Y$ be a projective contraction of relative dimension 1 such that the general fibers of f are isomorphic to $\mathbb{P}^1$. Assume that $K_X + \Delta$ is nef over Y and $(K_X + \Delta) \cdot F = 0$ for general fibers F of f . We run a relative K_X -MMP over Y with the scaling of Δ*

$$(X, \Delta) =: (X_0, \Delta_0) \dashrightarrow (X_1, \Delta_1) \dashrightarrow \dots \dashrightarrow (X_n, \Delta_n) =: X' \rightarrow Y'$$

and end with a Mori fiber space $f' : X' \rightarrow Y'$ over Y .

Then every step of this MMP is $(K_X + \Delta)$ -trivial.

Proof. From the construction of MMP with scaling we understand that it is enough to show that the nef thresholds

$$\lambda_i := \inf\{t \geq 0 : K_{X_i} + t\Delta \text{ is nef}/Y\} = 1,$$

for all $0 \leq i \leq n$. Note that $\{\lambda_i\}$ form a monotonically decreasing sequence and $\lambda_i \leq 1$ for all $0 \leq i \leq n$. Therefore it is enough to show that $\lambda_n = 1$. To that end, first observe that the composite bimeromorphic map $\phi : X \dashrightarrow X'$ is an isomorphism over a dense Zariski open subset of Y . In particular, if $F(\cong \mathbb{P}^1)$ is a general fiber of $X \rightarrow Y$, then $(K_{X'} + \Delta') \cdot F = (K_X + \Delta) \cdot F = 0$, i.e. $\Delta' \cdot F = 2$. Note that the general fibers of $X \rightarrow Y$ are also general fibers of $f' : X' \rightarrow Y'$, and according to the construction of MMP with scaling, f' is a $(K_{X'} + \lambda_n \Delta')$ -trivial contraction, i.e. $(K_{X'} + \lambda_n \Delta') \cdot F = 0$ for a general fiber F of f' . Thus $\Delta' \cdot F = \frac{2}{\lambda_n}$ and therefore $\lambda_n = 1$. $\square$

5. RELATIVE MMP FOR DLT PAIRS AND EXISTENCE OF DLT MODEL

In this section we will prove a relative cone theorem for dlt pairs and then establish the existence of dlt models for log canonical pairs.

Theorem 5.1. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Let $f : X \rightarrow Z$ be a projective bimeromorphic morphism to a normal compact analytic 3-fold Z . Suppose that $\Delta = \Delta' + \sum_{i=1}^k E_i$ and $K_X + \Delta \equiv_f \sum_{i=1}^k a_i E_i$, where $a_i > 0$ for all i and E_i 's are f -exceptional divisors such that $\text{Ex}(f) = \cup_{i=1}^k E_i$. Then there are at most countably many rational curves $\{C_j\}_{j \in J}$ such that $-(K_X + \Delta) \cdot C_j \leq 3$ and $f(C_j) = \text{pt}$ for all $j \in J$ and the following holds:*

$$(5.1) \quad \overline{\text{NE}}(X/Z) = \overline{\text{NE}}(X/Z)_{(K_X + \Delta) \geq 0} + \sum_{j \in J} \mathbb{R}^{\geq 0}[C_j].$$

Proof. First note that there are at most countably many family of curves in X . In particular, there are at most countably many curve classes $[C] \in \overline{\text{NE}}(X/Z)$ such that $0 < -(K_X + \Delta) \cdot C \leq 3$. Choose an irreducible representative Γ_i from each such class $[C]$ and set

$$(5.2) \quad V := \overline{\text{NE}}(X/Z)_{(K_X + \Delta) \geq 0} + \sum_{0 < -(K_X + \Delta) \cdot \Gamma_i \leq 3} \mathbb{R}^{\geq 0}[\Gamma_i].$$

Next we claim that it is enough to show that $\overline{\text{NE}}(X/Z) = \overline{V}$. Indeed, the claim follows from [HP16, Lemma 6.1] and its proof by replacing K_X by $K_X + \Delta$.

Now clearly $\overline{V} \subseteq \overline{\text{NE}}(X/Z)$. So let $[C]$ be a class of an irreducible curve $C \subseteq X$ such that $f(C) = \text{pt}$. It is enough to show that $[C] \in V$. If

$-(K_X + \Delta) \cdot C \leq 3$, then $[C] \in V$ by definition of V . So assume that $-(K_X + \Delta) \cdot C > 3$, i.e. $(K_X + \Delta) \cdot C < 0$. Then from our hypothesis it follows that C is contained in the support of one of the E_i 's, say $F = E_{i_0}$ for some $i_0 \in \{1, \dots, k\}$, and $F \cdot C < 0$. Now by adjunction $(K_X + \Delta)|_F = K_F + \Delta_F$ and (F, Δ_F) is a dlt surface such that $(K_F + \Delta_F) \cdot C = (K_X + \Delta) \cdot C < 0$. Thus C generates a $(K_F + \Delta)$ -negative extremal ray of $\overline{\text{NE}}(F)$. Now since f is a projective morphism and $f(F)$ is either a point or a projective curve (see Lemma 3.11), it follows that $f^{-1}(f(F))$ is a projective scheme, and hence F is a projective surface. Then by the cone and contraction theorem for log canonical projective surfaces as in the Theorem 3.2 and Proposition 3.8 of [Fuj12] it follows that there is a rational curve $\Gamma \subseteq F$ such that $[\Gamma] = [C]$ in $\overline{\text{NE}}(F)$ and $-(K_F + \Delta_F) \cdot \Gamma \leq -3$. But this implies that $-(K_X + \Delta) \cdot C = -(K_X + \Delta) \cdot \Gamma \leq 3$, this is a contradiction.

Finally if $\mathbb{R}^{\geq 0}[\Gamma_j]$ is a $(K_X + \Delta)$ -negative extremal ray of $\overline{\text{NE}}(X/Z)$, then as in the argument in the previous paragraph we can show that $\Gamma_j \subseteq E_{i_0}$ for some $i_0 \in \{1, \dots, k\}$. Then again by [Fuj12, Thm. 3.2 and Pro. 3.8] there is a rational curve $C_j \subseteq X$ such that $[C_j] = [\Gamma_j]$. This completes the proof. $\square$

Theorem 5.2 (DLT Modification). *Let (X, Δ) be a compact Kähler 3-fold log canonical pair. Then there exists a projective bimeromorphic morphism $f : (X', \Delta') \rightarrow (X, \Delta)$ from a $\mathbb{Q}$ -factorial dlt pair (X', Δ') such that $K_{X'} + \Delta' = f^*(K_X + \Delta)$.*

Proof. Let $g : Y \rightarrow X$ be the log resolution of (X, Δ) . Define $\Delta_Y := f_*^{-1}\Delta + E$, where $E = \text{Ex}(f)$ is the reduced f -exceptional divisor. Then (Y, Δ_Y) has dlt singularities. We will run a $(K_Y + \Delta_Y)$ -MMP over X . Let $C \subseteq Y$ be a curve generating a $(K_Y + \Delta)$ -negative extremal ray of $\overline{\text{NE}}(Y/X)$ as in Theorem 5.1. Then for some $0 < \varepsilon \ll 1$ we still have $(K_X + (1 - \varepsilon)\Delta_Y) \cdot C < 0$ and $(K_Y, (1 - \varepsilon)\Delta_Y)$ is klt. Therefore by [Nak87, Theorem 4.12(2)] there exist projective bimeromorphic morphisms $\varphi : Y \rightarrow Z$ and $h : Z \rightarrow X$ such that φ contracts the extremal ray $R = \mathbb{R}^{\geq 0}[C]$ and $g = h \circ \varphi$. Now since X is Kähler and $h : Z \rightarrow X$ is a projective morphism, from [Var89, Pro. 1.3.1, page 24] it follows that Z is Kähler. Now if φ is a flipping contraction, then the flip, say $\varphi^+ : Y^+ \rightarrow Z$ exists by Theorem 3.1. The fact that Y^+ is Kähler again follows from [Var89, Pro. 1.3.1, page 24], since Z is Kähler and φ^+ is projective. The termination of flips follows from Theorem 3.3. Finally, since f is bimeromorphic, $K_Y + \Delta_Y$ is f -big, and consequently this MMP terminates with a minimal model over X . Then applying the negativity lemma we get our required dlt model. $\square$

Corollary 5.3. *Let (X, Δ) be a compact Kähler 3-fold lc pair. Then there exists a bimeromorphic model $f : (X', \Delta') \rightarrow (X, \Delta)$ such that*

- (1) X' has $\mathbb{Q}$ -factorial terminal singularities,
- (2) (X', Δ') is a dlt pair, and
- (3) $K_{X'} + \Delta' = f^*(K_X + \Delta)$.

Proof. By Theorem 5.2, there is a dlt model $g : (\hat{X}, \hat{\Delta}) \rightarrow (X, \Delta)$ such that $K_{\hat{X}} + \hat{\Delta} = g^*(K_X + \Delta)$. Hence, by replacing (X, Δ) by $(\hat{X}, \hat{\Delta})$, we may assume that (X, Δ) is dlt. Let $U \subseteq X$ be the largest open set such that $(U, \Delta|_U)$ is a SNC pair. Then $\text{codim}_X(X \setminus U) \geq 2$. Let $f : (X', \Theta') \rightarrow (X, 0)$ be a terminal model of $(X, 0)$ as in Theorem 3.6 such that $K_{X'} + \Theta' = f^*K_X$. Then f is an isomorphism over the smooth locus of X ; in particular f is an isomorphism over U . Let $Z = X \setminus U$. Define $\Delta' := \Theta' + f^*\Delta$ on X' so that

$$K_{X'} + \Delta' = f^*(K_X + \Delta),$$

and (X', Δ') is lc.

It remains to show that (X', Δ') is a dlt pair. Let $U' = f^{-1}(U)$ and $Z' = X' \setminus U'$. Then $(U', \Delta'|_{U'})$ is a SNC pair. If E is an exceptional divisor with center in Z' , then its center in X is contained in Z . Hence $a(E; X', \Delta') = a(E; X, \Delta) > -1$. This completes the proof. $\square$

6. LMMP AND ABUNDANCE FOR KÄHLER SURFACES

In this section, we will recall the log minimal model programs for dlt pair (X, Δ) , where X is a compact Kähler surface. Since (X, Δ) is dlt, X has $\mathbb{Q}$ -factorial rational singularity. Now if X is uniruled, then from the classification of compact complex surfaces it follows that X is algebraic. So we will only focus on the case when X is not uniruled. Let $\varphi : \tilde{X} \rightarrow X$ be the minimal resolution of X . Then there is an effective $\mathbb{Q}$ -divisor $E \geq 0$ such that

$$K_{\tilde{X}} \sim_{\mathbb{Q}} \varphi^*K_X - E.$$

Since $\tilde{X}$ is not uniruled, by abundance for smooth compact Kähler surfaces it follows that $K_{\tilde{X}}$ is $\mathbb{Q}$ -effective. As a consequence, K_X is also $\mathbb{Q}$ -effective. We can write

$$K_X \sim_{\mathbb{Q}} \sum_{\text{finite}} \lambda_i C_i,$$

where C_i are irreducible curves. By [Bou04, Proposition 3.4], $K_X + \Delta$ is nef if and only if it has positive intersection with any curves in X . In this case, we can obtain the following cone theorem.

Theorem 6.1. *Let (X, Δ) be a dlt pair such that X is a non-uniruled compact Kähler surface. If $K_X + \Delta$ is not nef, then there are finitely many irreducible*

curves Γ_i such that

$$-(K_X + \Delta) \cdot \Gamma_i > 0$$

and that

$$\overline{\text{NA}}(X) = \overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_i \mathbb{R}^+[\Gamma_i].$$

In order to establish the MMP, we still need the following contraction theorem.

Theorem 6.2. *With the notation and hypothesis as in Theorem 6.1, for each irreducible curve Γ with $-(K_X + \Delta) \cdot \Gamma > 0$, there is a bimeromorphic morphism $f : X \rightarrow Y$ which contracts precisely the curve Γ and Y is a compact Kähler surface.*

Proof. Since K_X is $\mathbb{Q}$ -effective as discussed above, it follows that $\Gamma \cdot \Gamma < 0$. In particular, $\mathcal{O}_\Gamma(-m\Gamma)$ is an ample line bundle on Γ , where m is the Cartier index of Γ on X . Then the existence of the contraction f follows from [HP16, Proposition 7.4]. A similar argument as in proof of [CHP16, Corollary 3.1] shows that Y is Kähler. $\square$

Theorem 6.3 (Abundance). *Let X be a normal compact Kähler space of dimension 2 and (X, Δ) a lc pair. If $K_X + \Delta$ is nef, then it is semi-ample.*

Proof. Replacing X by its minimal resolution we may assume that X is smooth. If $\nu(K_X + \Delta) = 0$ or 1, then from [CHP16, Pro. 5.1 and Lemma 5.1] it follows that $K_X + \Delta$ is semi-ample.

If $\nu(K_X + \Delta) = 2$, then by [Dem96, Corollary 6.8] $K_X + \Delta$ is big; in particular X is a Moishezon manifold. Then by [Nam02, Theorem 1.6] X is projective, and the semi-ameness of $K_X + \Delta$ follows from [Fuj12, Theorem 4.1] $\square$

7. ABUNDANCE FOR 3-FOLD PAIRS (X, Δ) WITH $\kappa(X, K_X + \Delta) \geq 1$

In this section we will prove the log abundance for a 3-fold log canonical pairs (X, Δ) such that $\kappa(X, K_X + \Delta) \geq 1$. The proof is based on the proof of [DW19, Theorem 3.1] which is originally inspired from the proof of [Kaw85, Theorem 7.8].

Lemma 7.1. *Let $f : X \rightarrow Y$ be a proper surjective morphism from a normal compact Kähler variety X to a normal projective variety Y of $\dim Y > 0$. Suppose that $\mathcal{L}$ is a line bundle on X such that $H^0(X, \mathcal{L}) \neq 0$ and $\mathcal{L}|_F \sim \mathcal{O}_F$ for general fibers F of f . Let D be a non-zero effective Cartier divisor on X*

given by the zeros of a non-zero global section of $\mathcal{L}$. Then there exists an effective ample Cartier divisor H on Y such that $D \leq f^*H$.

Proof. Since $\mathcal{L}|_F \cong \mathcal{O}_F$ for general fibers F of f , it follows that $D|_F \sim 0$. We claim that $\text{Supp}(D) \cap F = \emptyset$ for general fibers F of f . Indeed, since D is effective, for a general fiber F , $D|_F$ is an effective divisor on F . Since $D|_F \sim 0$ and F is a normal compact Kähler variety, it follows that $D|_F = 0$, i.e. $\text{Supp } D \cap F = \emptyset$ for general fibers F of f . In particular, $f(D)$ does not dominate Y . Let H be an effective ample Cartier divisor Y such that $\text{Supp } H$ contains $f(D)$. Then $\text{Supp } D$ is contained in the $\text{Supp } f^*H$. Thus replacing H by a sufficiently high multiple we may assume that $D \leq f^*H$. $\square$

7.1. Iitaka Fibration. Let (X, Δ) be a terminal pair, where X is compact Kähler 3-fold. Assume that with $\kappa(X, K_X + \Delta) \geq 0$. Then the Iitaka fibration as in [Laz04, Theorem 2.1.33] gives a diagram

$$\begin{array}{ccc} & Y & \\ \mu \swarrow & & \searrow f \\ X & \text{-----} & Z \end{array}$$

satisfying the following conditions:

- (1) Y is a smooth compact Kähler 3-fold, and Z is a smooth projective variety with $\dim Z = \kappa(X, K_X + \Delta)$.
- (2) $\mu : Y \rightarrow X$ is a log resolution of (X, Δ) and f is a surjective morphism with connected fibers.
- (3) $\kappa(W, (K_Y + \Delta_Y)|_W) = 0$, where W is a general fiber of f , and $\Delta_Y = \mu_*^{-1}\Delta$ (see [LP18, Lemma 2.3]).

Theorem 7.2. *Let (X, Δ) be a compact Kähler 3-fold dlt pair. If $\kappa(X, K_X + \Delta) \geq 1$ and $K_X + \Delta$ is nef, then it is semi-ample.*

Proof. If $\kappa(X, K_X + \Delta) = 3$, then X is a Moishezon space. Since dlt singularities are rational, it follows from [Nam02, Theorem 1.6] that X is projective algebraic varieties. Then semi-ameness follows from [KMM94, Theorem 1.1].

In the following discussion we will consider the remaining cases where $\kappa(X, K_X + \Delta) = 1$ or 2. By passing to a terminal model by Theorem 3.6 we may assume that (X, Δ) has $\mathbb{Q}$ -factorial terminal singularities.

Let the following diagram be the Iitaka fibration of $K_X + \Delta$ as defined above

$$\begin{array}{ccc} & Y & \\ \mu \swarrow & & \searrow f \\ X & \text{-----} & Z. \end{array}$$

Let $\{E_i\}$ be the exceptional divisors of μ , and set $\Delta_Y = \mu_*^{-1}\Delta$. Then we have

$$(7.1) \quad K_Y + \Delta_Y = \mu^*(K_X + \Delta) + \sum r_i E_i, \quad r_i > 0 \text{ for all } i.$$

Let W be a general fiber of $f : Y \rightarrow Z$. Since Y is smooth, by generic smoothness it follows that W is smooth and (W, Δ_W) is a SNC pair, where $\Delta_W = \Delta_Y|_W$. In particular, $(W, K_W + \Delta_W)$ has dlt singularities.

Claim 7.3. $\mu^*(K_X + \Delta)|_W \sim_{\mathbb{Q}} 0$.

Assuming the claim for the time being first we will complete the proof here. Let $m > 0$ be an integer such that $\mu^*(m(K_X + \Delta))$ is a Cartier divisor. Let D be an effective Cartier divisor on Y given by the zeros of a non-zero global section of the line bundle $\mathcal{O}_Y(m(\mu^*(K_X + \Delta)))$. Then by Lemma 7.1, there exists an ample Cartier divisor H on Z such that $D \leq f^*H$. Therefore $\nu(X, K_X + \Delta) = \nu(Y, \mu^*(m(K_X + \Delta))) = \nu(Y, D) \leq \nu(Y, f^*H_2) = \kappa(Y, f^*H_2) = \kappa(Z, H_2) = \dim Z = \kappa(X, K_X + \Delta)$. Thus $\kappa(X, K_X + \Delta) = \nu(X, K_X + \Delta)$ by [Dem96, Proposition 6.10]. Then by [Fuj11, Theorem 4.8] $K_X + \Delta$ is semi-ample. $\square$

Proof of Claim 7.3. We split the proof into two cases depending on the dimension of W .

Case I: $\dim W = 1$. In this case W is a smooth proper curve, hence it is a smooth projective algebraic curve. Since $\kappa(W, K_W + \Delta) = 0$, it follows that $K_W + \Delta_W \sim_{\mathbb{Q}} 0$. Restricting both side of (7.1) to W we get

$$\mu^*(K_X + \Delta)|_W \sim_{\mathbb{Q}} - \sum r_i E_i|_W.$$

Since $K_X + \Delta$ is nef and $r_i > 0$ for all i , we conclude that $E_i|_W = 0$ for all i , i.e., $\mu^*(K_X + \Delta)|_W \sim_{\mathbb{Q}} 0$.

Case II: $\dim W = 2$. In this case W is a smooth Kähler surface and (W, Δ_W) is a dlt pair. Now by Theorem 6.1, 6.2 and 6.3 we run the $(K_W + \Delta_W)$ -MMP which terminates with a dlt pair $(W', \Delta_{W'})$ such that $K_{W'} + \Delta_{W'}$ is semi-ample. Let $\varphi : W \rightarrow W'$ be the induced bimeromorphic morphism such

that $K_{W'} + \Delta_{W'} = \varphi_*(K_W + \Delta_W)$. Then we have

$$(7.2) \quad K_W + \Delta_W = \varphi^*(K_{W'} + \Delta_{W'}) + \sum s_j F_j, \text{ where } s_j > 0 \text{ for all } j,$$

where F_j 's are the φ -exceptional divisors.

Since $\kappa(W', K_{W'} + \Delta_{W'}) = \kappa(W, K_W + \Delta_W) = 0$ and $K_{W'} + \Delta_{W'}$ is semi-ample, $K_{W'} + \Delta_{W'} \sim_{\mathbb{Q}} 0$. Therefore we have

$$(7.3) \quad K_W + \Delta_W \sim_{\mathbb{Q}} \sum s_j F_j, \text{ where } s_j > 0 \text{ for all } j.$$

From adjunction on W we also get that

$$(7.4) \quad K_W + \Delta_W \sim_{\mathbb{Q}} \mu^*(K_X + \Delta)|_W + \sum r_i E_i|_W$$

Thus we have

$$(7.5) \quad \mu^*(K_X + \Delta)|_W \sim_{\mathbb{Q}} \sum s_j F_j - \sum r_i E_i|_W = G^+ - G^-,$$

such that $G^+ \geq 0$ and $G^- \geq 0$ are two effective $\mathbb{Q}$ -divisors on W with no common irreducible components.

We will show that $G^+ = G^- = 0$. On the contrary first assume that $G^+ \neq 0$. It is clear that G^+ is φ -exceptional. Now by [BHPVdV04, Theorem III.2.1] the intersection matrix of the exceptional divisors of $\varphi : W \rightarrow W'$ is a negative definite matrix. Thus in particular $(G^+)^2 < 0$.

On the other hand, since $\mu^*(K_X + \Delta)|_W$ is nef, from relation (7.5) we get $G^+ \cdot (G^+ - G^-) \geq 0$, which is a contradiction. Therefore $G^+ = 0$ and $\mu^*(K_X + \Delta)|_W \sim_{\mathbb{Q}} -G^-$. Again since $\mu^*(K_X + \Delta)|_W$ is nef, $G^- = 0$. Therefore we have

$$(7.6) \quad \mu^*(K_X + \Delta)|_W \sim_{\mathbb{Q}} 0.$$

□

Corollary 7.4. *Let (X, Δ) be a compact Kähler 3-fold lc pair. If $\kappa(X, K_X + \Delta) \geq 1$ and $K_X + \Delta$ is nef, then it is semi-ample.*

Proof. It result follows immediately from Theorem 7.2 by passing to a dlt model using Theorem 5.2. □

8. $\mathbb{Q}$ -LINEAR SYSTEM

Let X be a normal compact analytic variety and L a $\mathbb{Q}$ -Cartier divisor on X . Then the $\mathbb{Q}$ -linear system $|L|_{\mathbb{Q}}$ is defined as the set of all effective $\mathbb{Q}$ -Cartier divisors $L' \geq 0$ such that $m(L - L')$ is an integral Weil divisor for some positive integer $m \in \mathbb{N}$ and that $m(L - L') \sim 0$. The *stable of base locus* of L is defined as

$$\text{SBs } |L| := \bigcap_{m \in \mathbb{N}} \text{Bs } |mkL|,$$

where k is the small positive integer such that kL is Cartier.

Note that since X is compact, it follows that $\text{SBs } |L|$ is an analytic subset of X , in fact $\text{SBs } |L| = \text{Bs } |mkL|$ for some $m \in \mathbb{N}$ (see [Laz04, Pro. 2.1.21]). The *base locus* of the $\mathbb{Q}$ -linear system $|L|_{\mathbb{Q}}$ is defined as

$$\text{Bs } |L|_{\mathbb{Q}} = \bigcap_{L' \in |L|_{\mathbb{Q}}} \text{Supp } L'.$$

Note that $\text{SBs } |L| = \text{Bs } |L|_{\mathbb{Q}}$.

For a $\mathbb{Q}$ -Cartier divisor D on X , we define

$$D + |L|_{\mathbb{Q}} = \{D + P \mid P \in |L|_{\mathbb{Q}}\}.$$

Assume that $|L|_{\mathbb{Q}}$ is not empty. Then for any prime Weil divisor E on X , we define the *multiplicity* of $D + |L|_{\mathbb{Q}}$ along E as

$$\text{mult}_E(D + |L|_{\mathbb{Q}}) = \inf\{\text{mult}_E(D + P) \mid P \in |L|_{\mathbb{Q}}\}.$$

We call $D + |L|_{\mathbb{Q}}$ a *boundary* (resp. *pure boundary*, resp. *sub-boundary*, and resp. *pure sub-boundary*) if its multiplicity along every prime divisor lies in the interval $[0, 1]$ (resp. $[0, 1)$, resp. $(-\infty, 1]$, and resp. $(-\infty, 1)$).

We also consider the pair $(X, D + |L|_{\mathbb{Q}})$. For a proper *generically finite* morphism $f : X' \rightarrow X$ from a normal variety X' , we define the log pullback of $K_X + D + |L|_{\mathbb{Q}}$ as $K_{X'} + D' + |L'|_{\mathbb{Q}}$, where

$$L' = f^*L \quad \text{and} \quad K_{X'} + D' = f^*(K_X + D).$$

We then define the discrepancy of a divisorial valuation ν on X for the pair $(X, D + |L|_{\mathbb{Q}})$ as follows: choose a proper bimeromorphic morphism $f : X' \rightarrow X$ from a smooth variety X' such that ν corresponds to a prime divisor E on X' . Then the *discrepancy* of ν with respect to $(X, D + |L|_{\mathbb{Q}})$ is defined as

$$a(\nu; X, D + |L|_{\mathbb{Q}}) = a(E; X, D + |L|_{\mathbb{Q}}) := -\text{mult}_E(D' + |L'|_{\mathbb{Q}}).$$

The pair $(X, D + |L|_{\mathbb{Q}})$ is called *sub-klt* if the discrepancies of all divisorial valuations are strictly greater than -1 . It is called *klt* if it is sub-klt and D is effective. Other singularities are defined similarly.

Now suppose that L is a Cartier divisor. Then we can decompose it as:

$$|L| = |M| + F,$$

where $|M|$ is the movable part, i.e. $\text{Bs } |M|$ does not have any codimension 1 component, and $F \geq 0$ is the fixed part. Note that the divisor M may not be Cartier here, so in general $|M|$ is a linear system of Weil divisors.

If D is another $\mathbb{Q}$ -Cartier divisor, then we define the *fixed part* of $D + |L|$ as $D + F$, and the *movable part* as $|M|$.

In the remainder of this section we establish some properties of $\mathbb{Q}$ -linear systems of the form $D + |L|_{\mathbb{Q}}$ and the corresponding pairs $(X, D + |L|_{\mathbb{Q}})$.

Lemma 8.1. *Let X be normal $\mathbb{Q}$ -factorial compact analytic variety, $|L|_{\mathbb{Q}}$ a $\mathbb{Q}$ -linear system on X , and D is a $\mathbb{Q}$ -divisor on X . If $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary, then there exist an element $N \in |L|_{\mathbb{Q}}$ such that $D + N$ is a pure sub-boundary.*

Proof. First replacing L by an effective $\mathbb{Q}$ -Cartier divisor in $|L|_{\mathbb{Q}}$ we may assume that L is effective. Let k be the smallest positive integer such that kL is Cartier. Then for any $m \geq 1$ we decompose $|mkL| = |M_m| + F_m$ into movable and fixed parts. Note that these fixed parts give a monotonically decreasing sequence of divisors $\{\frac{1}{mk}F_m\}$ on X , i.e., $\frac{1}{mk}F_m \geq \frac{1}{(m+1)k}F_{m+1}$ for all $m \geq 1$. Now since $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary, there is a $m_0 \gg 0$ such that the coefficients of the divisor $D + \frac{1}{m_0k}F_{m_0}$ along each of its irreducible component is strictly less than 1. Now we focus on the movable part $\frac{1}{m_0k}|M_{m_0}|$. We make the following claim.

Claim 8.2. There is an effective reduced divisor $N' \geq 0$ in the linear system of $|M_{m_0}|$, i.e., the coefficients of N' are all 1, such that the support of N' does not contain any irreducible components of $D + \frac{1}{m_0k}F_{m_0}$.

Assuming the claim for the time being we will complete the proof first. Set $N := \frac{1}{m_0k}N' + \frac{1}{m_0k}F_{m_0} \in |L|_{\mathbb{Q}}$; then clearly $D + N$ is a pure sub-boundary.

Proof of Claim 8.2. First note that, since X is compact, the linear system $|mkL|$ is finite dimensional for all $m \geq 1$; more specifically, $|mkL|$ is given by the finite dimensional $\mathbb{C}$ -vector space $H^0(X, \mathcal{O}_X(kL))$ for all $m \geq 1$. In particular, $|M_m|$ is a finite dimensional linear system of Weil divisors on X , for all $m \geq 1$. Let X_{sing} be the singular locus of X ; note that $\text{codim}_X X_{\text{sing}} \geq 2$, since X is normal. Now consider the closed subset $Y = \text{Supp}(D + \frac{1}{m_0k}F_{m_0}) \cup \text{Bs } |M_{m_0}| \cup X_{\text{sing}} \subseteq X$ and set $U = X \setminus Y$. Then U is a smooth complex space. Notice that the restricted linear system $|M_{m_0}|_U$ is a finite dimensional linear sub-system of $|M_{m_0}|$ of Cartier divisors (since U is smooth). Furthermore, $|M_{m_0}|_U$ is base-point free on U . Then by a Bertini type theorem for complex spaces [Man82, Theorem II.5], a general enough member N'_U of $|M_{m_0}|_U$ is a reduced divisor on U . Note that a very general member of $|M_{m_0}|$ does not contain any codimension 1 component of $X \setminus U$, and thus N'_U extends to a reduced divisor N' on X such that $N' \in |M_{m_0}|$. $\square$

$\square$

Lemma 8.3. *Let X be a normal $\mathbb{Q}$ -factorial compact analytic variety, $|L|_{\mathbb{Q}}$ a $\mathbb{Q}$ -linear system on X , and D is a $\mathbb{Q}$ -divisor on X . Then the following assertions are all equivalent:*

- (1) $(X, D + |L|_{\mathbb{Q}})$ is sub-*klt*;
- (2) For any projective bimeromorphic morphism $f : X' \rightarrow X$ with X' normal, $D' + |L'|_{\mathbb{Q}}$ is a pure sub-boundary, where $D' + |L'|_{\mathbb{Q}}$ is the log pullback of $D + |L|_{\mathbb{Q}}$, i.e., $L' = f^*L$ and $K_{X'} + D' = f^*(K_X + D)$.

Proof. It follows easily from the definition above. □

Lemma 8.4. *Let X be a normal $\mathbb{Q}$ -factorial compact analytic variety, $|L|_{\mathbb{Q}}$ a $\mathbb{Q}$ -linear system on X , and D is a $\mathbb{Q}$ -divisor on X . If $(X, D + |L|_{\mathbb{Q}})$ is sub-*klt*, then there is an element $N \in |L|_{\mathbb{Q}}$ such that $(X, D + N)$ is sub-*klt*.*

Proof. Let $f : X' \rightarrow X$ be a resolution of singularities of X such that $\text{Ex}(f)$, f^*D , f^*L and $f^{-1}\text{Bs}(|L|_{\mathbb{Q}})$ are all simple normal crossing divisors and they intersect each other transversally. Let $D' + |L'|_{\mathbb{Q}}$ be the log pullback of $D + |L|_{\mathbb{Q}}$, i.e.,

$$K_{X'} + D' = f^*(K_X + D) \quad \text{and} \quad L' = f^*L.$$

Then by Lemma 8.3, $D' + |L'|_{\mathbb{Q}}$ is a pure sub-boundary. We need to show that there is some $N' \in |L'|_{\mathbb{Q}}$ such that $D' + N'$ is a pure sub-boundary and it also has simple normal crossing support.

By a similar argument as in the proof of Lemma 8.1 we see that there is a sufficiently large and divisible positive integer m such that $D' + \frac{1}{m}F'$ is a pure sub-boundary, where $|mL| = |M'| + F'$ is the decomposition of mL into movable and fixed parts. Indeed, since m is sufficiently large and divisible and X is compact, it follows that $\text{Bs } |mL| = \text{Bs } |L|_{\mathbb{Q}}$. Then from the construction of D' and L' above it follows that $D' + \frac{1}{m}F'$ is a simple normal crossing divisor. Note that since $(X, D + |L|_{\mathbb{Q}})$ is sub-*klt* and m is sufficiently large, it follows from the arguments in the proof of Lemma 8.1 that $D' + \frac{1}{m}F'$ is a pure sub-boundary. Now M' is a Cartier divisor on the compact complex manifold X' such that $\text{Bs } |M'|$ is a finite set. Thus by [Fuj83, Corollary 1.14], $\text{Bs } |M'| = \emptyset$, i.e., $|M'|$ is base point free. Then by a Bertini type theorem [Man82, Corollary II.7], there is a smooth divisor $N'' \in |M'|$ such that the support of N'' does not contain any component of $D + \frac{1}{m}F'$ and $D + \frac{1}{m}N'' + \frac{1}{m}F'$ has simple normal crossing support. Set $N' := \frac{1}{m}(N'' + F') \in |L'|_{\mathbb{Q}}$; then clearly $D + N'$ is a pure sub-boundary. □

In the following two lemmas we compare $D + |L|_{\mathbb{Q}}$ with its log pullback by a generically finite morphism.

Lemma 8.5. *Let X be a normal $\mathbb{Q}$ -factorial compact analytic variety, $|L|_{\mathbb{Q}}$ a $\mathbb{Q}$ -linear system on X , and D is a $\mathbb{Q}$ -divisor on X . Let $p : X' \rightarrow X$ be a proper finite morphism with X' normal. Let $D' + |L'|_{\mathbb{Q}}$ be the log pullback of $D + |L|_{\mathbb{Q}}$*

under p . Then $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary if and only if $D' + |L'|_{\mathbb{Q}}$ is a pure sub-boundary.

Proof. Assume that $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary. Then by Lemma 8.1, there is some $N \in |L|_{\mathbb{Q}}$ such that $D + N$ is a pure sub-boundary. We claim that $D' + N'$ is a pure sub-boundary, where $N' = p^*N$. Let E' be a prime divisor on X' and set $E := p(E')$. We will compute the multiplicity of $D' + N'$ along E' . Assume that p^*E has multiplicity m along E' . Let d and n be the multiplicities of D and N respectively along E . Then we have

$$\text{mult}_E(D + N) - 1 = d + n - 1 < 0.$$

From the ramification formula we have

$$\text{mult}_{E'} N' = nm, \quad \text{mult}_{E'} D' = dm - (m - 1).$$

Thus

$$\begin{aligned} \text{mult}_{E'}(D' + N') - 1 &= dm - (m - 1) + nm - 1 \\ &= m(d + n - 1) \\ &= m(\text{mult}_E(D + N) - 1) \\ &< 0. \end{aligned}$$

This shows that $D' + N'$ is a pure sub-boundary.

Now we assume that $D' + |L'|_{\mathbb{Q}}$ is a pure sub-boundary. Using the result we just proved above we may replace X' by the Galois closure of the morphism p , and thus assume that $p : X' \rightarrow X$ is a Galois morphism with Galois group G . Now since $D' + |L'|$ is a pure sub-boundary, by Lemma 8.1 there is an element $B' \in |L'|_{\mathbb{Q}}$ such that $D' + B'$ is a pure sub-boundary. Now set $N' := \frac{1}{\#G} \sum_{g \in G} g(B')$. Then $N' \in |L'|_{\mathbb{Q}}$ and $D' + N'$ is also a pure sub-boundary. Moreover, since N' is G -invariant, there is an element $N \in |L|_{\mathbb{Q}}$ such that $N' = p^*N$. We claim that $D + N$ is a pure sub-boundary. Let E be a prime Weil divisor on X and let E' be an irreducible component of p^*E . Then as in the computation above we have

$$\text{mult}_E(D + N) - 1 = \frac{1}{m}(\text{mult}_{E'}(D' + N') - 1) < 0.$$

Therefore $D + N$ is a pure sub-boundary, and so is $D + |L|_{\mathbb{Q}}$. $\square$

Lemma 8.6. *Let X be a normal $\mathbb{Q}$ -factorial compact analytic variety, $|L|_{\mathbb{Q}}$ a $\mathbb{Q}$ -linear system on X , and D is a $\mathbb{Q}$ -divisor on X . Let $p : X' \rightarrow X$ be a generically finite proper morphism from a normal variety X' . Let $D' + |L'|_{\mathbb{Q}}$ be the log pullback of $D + |L|_{\mathbb{Q}}$ under p . Then $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary if $D' + |L'|_{\mathbb{Q}}$ is a pure sub-boundary.*

Proof. By Stein factorization, we only need to prove the lemma in the following two cases: either p is a proper bimeromorphic morphism, or p is a finite proper morphism. The case when p is bimeromorphic is trivial. The case when π is finite follows from Lemma 8.5. $\square$

Part 2. Construction of LMMP for pseudo-effective pair

9. NON-VANISHING THEOREM

The objective of this section is to prove the following non-vanishing theorem.

Theorem 9.1. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Assume that X is non-algebraic and that the base of the MRC fibration for X has dimension 2. Then the following assertions are equivalent:*

- (1) $K_X + \Delta$ is $\mathbb{Q}$ -effective;
- (2) $K_X + \Delta$ is pseudo-effective;
- (3) $(K_X + \Delta) \cdot F \geq 0$ for general fibers F of the MRC fibration of X .

Remark 9.2. Note that if $(K_X + \Delta) \cdot F > 0$ for a general fiber F of the MRC fibration, then Δ has at least one component which dominates the base of the MRC fibration. Reducing the coefficients of these components appropriately we may assume that $(K_X + \Delta) \cdot F = 0$. Thus in order to prove the above theorem it is enough to show that, if $(K_X + \Delta) \cdot F = 0$, then $K_X + \Delta$ is $\mathbb{Q}$ -effective.

Using Lemma 3.10 we will reduce the general non-vanishing Theorem 9.3 to Theorem 9.1.

Theorem 9.3. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. If $K_X + \Delta$ is pseudo-effective, then it is $\mathbb{Q}$ -effective.*

Proof. If X is projective, then the theorem follows from [KMM94, Theorem 1.1] after running a $(K_X + \Delta)$ -MMP. Note that it is not possible for X to be non-projective but still algebraic in this case due to [Nam02, Corollary 1.7].

If X is not uniruled, then first using Corollary 5.3 we may assume that X has terminal singularity. Then the non-vanishing follows from [DP03, Theorem 0.3] after running a K_X -MMP as in [HP16, Theorem 1.1]. It remains to treat the case when X is uniruled and non-algebraic. In this case the base of the MRC fibration of X has dimension 2, by Lemma 3.10, and thus our result follows from Theorem 9.1. $\square$

9.1. Case of Mori fiber space. The aim of this subsection is to prove the Theorem 9.5 below, which is a generalization of [KMM94, Corollary 3.8]. First we state a combinatorial result from [KMM94] which will be very useful to us in this section.

Let $I_1, I_2, \dots, I_p$ be a partition of the set $\{1, 2, \dots, k\}$. Let $a_1, a_2, \dots, a_k$ be a sequence of rational numbers such that $0 \leq a_i \leq 1$ and $\sum_{i=1}^k a_i = 2$. Set $\alpha_q = \sum_{i \in I_q} a_i$.

Lemma 9.4. [KMM94, Lemma 3.2] *With the notations and hypothesis as above assume that α_q satisfies $0 \leq \alpha_q \leq 1$ for all $q \in \{1, \dots, p\}$. Then the system of equations*

$$\sum_{j=1}^k b_{ij} = a_i \quad \text{for all } i = 1, 2, \dots, k$$

has a non-negative rational solution $b_{ij} \geq 0$ such that $b_{ij} = b_{ji}$ for all i, j , and $b_{ij} = 0$ if i and j belong to the same indexing set I_q .

Theorem 9.5. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Assume that X is non-algebraic and that the base of the MRC fibration of X has dimension 2. Let $f : X \rightarrow Y$ be a Mori fibration of relative dimension 1 such that $K_X + \Delta \sim_{\mathbb{Q}, f} 0$. Then there is an effective $\mathbb{Q}$ -divisor $B \geq 0$ on Y such that $K_X + \Delta \sim_{\mathbb{Q}} f^*(K_Y + B)$. Moreover, if (X, Δ) is klt, then (Y, B) is klt.*

We first deal with the case where every component of Δ is bimeromorphic to the base Y .

The following elementary lemmas will be useful in our proof.

Lemma 9.6. *Let $f : Y \rightarrow X$ be a proper bimeromorphic morphism between compact analytic varieties and X is normal. Let ω_Y be the canonical sheaf of Y such that $\omega_Y^{[m]}$ is a line bundle for some $m \in \mathbb{N}$. Let $g : Z \rightarrow Y$ be a resolution of singularities of Y and $\omega_Z^m \cong g^*\omega_Y^{[m]}(E)$ for some Weil divisor E on Z . Then E is $(f \circ g)$ -exceptional.*

Proof. Let $\nu : \bar{Y} \rightarrow Y$ be the normalization of Y . Then $\omega_{\bar{Y}}^{[m]} \cong \nu^*\omega_Y^{[m]}(B)$ for some effective Weil divisor B on $\bar{Y}$. The divisor B is determined by the conductor ideal of the normalization. Since $f : Y \rightarrow X$ is bimeromorphic and X is normal, $\text{Supp}(\nu_*B) \subseteq \text{Ex}(f)$. Now since the resolution $g : Z \rightarrow Y$ factors through the normalization $\bar{Y}$ and $\omega_Z^m \cong g^*\omega_Y^{[m]}(E)$, it follows that the components of E are either exceptional for the induced morphism $Z \rightarrow \bar{Y}$ or

they are strict transform of the components of the conductor divisor B . In particular, we have $\text{Supp } E \subseteq \text{Ex}(f \circ g)$. $\square$

In the next two lemmas we will deal with case when the horizontal components of Δ are bimeromorphic to the base of the MRC fibration.

Lemma 9.7. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Let $f : X \rightarrow Y$ be a contraction of a $(K_X + \Delta)$ -negative extremal ray of $\overline{\text{NA}}(X)$. Let D be a $\mathbb{Q}$ -divisor on X such that D is vertical over Y , i.e. $f(\text{Supp } D)$ does not dominate Y . Then there is a $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor B on Y such that $D = f^*B$.*

Proof. Let m be a positive integer such that mD is a Cartier divisor. Write $D = D^+ - D^-$, where D^+ and D^- effective $\mathbb{Q}$ -divisors without common components. Note that by hypothesis D^+ and D^- are both vertical over Y . Thus $mD^+ \cdot C = 0$ for curves $C \subseteq X$ contained in the fibers of f . Then by [CHP16, Proposition 3.1(6)] there exists a line bundle $\mathcal{L}$ on Y such that $\mathcal{O}_X(mD^+) \cong f^*\mathcal{L}$. Now since $f_*\mathcal{O}_X = \mathcal{O}_Y$, from the projection formula it follows that $\{0\} \neq H^0(X, \mathcal{O}_X(mD^+)) = H^0(X, f^*\mathcal{L}) = H^0(Y, \mathcal{L})$. Therefore there exists a non-zero section $s \in H^0(Y, \mathcal{L})$ such that the divisor of zeros of f^*s is the divisor mD^+ , i.e. $mD^+ = \text{div}(f^*s)_0$. Set $L := \text{div}(s)_0$. Then L is an effective Cartier divisor on Y such that $mD^+ = f^*L$. Similarly, there exists an effective Cartier divisor M on Y such that $mD^- = f^*M$. Define $B := \frac{1}{m}(L - M)$. Thus we have $D = (D^+ - D^-) = f^*B$. $\square$

Lemma 9.8. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold pair with canonical singularities such that X has terminal singularities. Let $f : X \rightarrow Y$ be a K_X -Mori fibration of relative dimension 1 such that $K_X + \Delta \equiv_f 0$. We assume that every component of Δ is bimeromorphic to Y via f . Then there is an effective $\mathbb{Q}$ -divisor $L \geq 0$ on Y such that $K_X + \Delta \sim_{\mathbb{Q}} f^*(K_Y + L)$ and that the base locus of the $\mathbb{Q}$ -linear system $|L|_{\mathbb{Q}}$ has no divisorial component.*

Proof. Write $\Delta = \sum_{i=1}^k a_i \Delta_i$ such that Δ_i 's are the irreducible components of Δ and $a_i \in (0, 1]$ for all i . Since f is a K_X -Mori fiber space of relative dimension 1, the general fibers of f are isomorphic to $\mathbb{P}^1$; in particular, from $K_X + \Delta \equiv_f 0$ we obtain that $\sum_{i=1}^k a_i = 2$. Then by Lemma 9.4 there exist

nonnegative rational numbers $b_{ij} \geq 0$ with $b_{ij} = b_{ji}$ for $1 \leq i, j \leq k$ such that

$$a_i = \sum_{j; j \neq i} b_{ij}.$$

Then we have

$$(9.1) \quad \sum_{i,j; i \neq j} \frac{1}{2} b_{ij} (K_{X/Y} + \Delta_i + \Delta_j) = K_{X/Y} + \sum_{i=1}^k a_i \Delta_i.$$

Set $B_{ij} := f_*(\Delta_i \cdot \Delta_j)$ for any $i \neq j$, and let $L = \frac{1}{2} \sum_{i \neq j} b_{ij} B_{ij} \geq 0$. We make the following claim.

Claim 9.9. The effective $\mathbb{Q}$ -divisor $L \geq 0$ on Y satisfies $(K_{X/Y} + \Delta) \sim_{\mathbb{Q}} f^* L$.

Proof of Claim 9.9. Let F be a general fiber of f ; then $F \cong \mathbb{P}^1$. Since Δ_i 's are bimeromorphic to Y via f , we have $(K_{X/Y} + \Delta_i + \Delta_j) \cdot F = 0$. Now since f is a contraction of a K_X -negative extremal ray, by [CHP16, Proposition 3.1(6)] there exists a positive integer $m > 0$ and a line bundle $\mathcal{L}_{ij}$ on Y such that

$$(9.2) \quad \mathcal{O}_X(m(K_{X/Y} + \Delta_i + \Delta_j)) \cong f^* \mathcal{L}_{ij}$$

is an isomorphism of line bundles.

Now since X has terminal singularities, X is smooth outside a set of finitely many points. Thus by adjunction we have $(K_X + \Delta_i + \Delta_j)|_{\Delta_i} = K_{\Delta_i} + \Delta_j|_{\Delta_i}$. Recall that $g = f|_{\Delta_i} : \Delta_i \rightarrow Y$ is bimeromorphic according to our hypothesis. Now observe that if K_{Δ_i} and K_Y were given by Weil divisors, then we would have $g_* K_{\Delta_i} = K_Y$. This would have made our proof significantly simpler by simply restricting both sides of (9.2) onto Δ_i , and then pushing forward by $f|_{\Delta_i}$ would show that $\mathcal{L}_{ij}$ is given by a Cartier divisor on Y . However, the canonical bundle of an analytic variety only exists as a sheaf and it may not correspond to any Weil divisor in general.

To remedy this problem we pullback everything to a resolution of Δ_i . Let $h : Z \rightarrow \Delta_i$ be a resolution of singularities of Δ_i . Then by Lemma 9.6 there exists a $(g \circ h)$ -exceptional Weil divisor E on Z such that $h^* \omega_{\Delta_i}^{[m]} \cong \omega_Z^m(E)$. Moreover, since Y is $\mathbb{Q}$ -factorial, by our definition of $\mathbb{Q}$ -factoriality, $\omega_Y^{[m]}$ is a line bundle for $m \in \mathbb{N}$ sufficiently divisible. Thus there exists a $(g \circ h)$ -exceptional divisor F on Z such that $\omega_Z^m \cong (g \circ h)^* \omega_Y^{[m]}(F)$. Now restricting both sides of (9.2) to Δ_i and then pulling back to Z we get

$$(9.3) \quad h^*(\mathcal{O}_X(m(K_{X/Y} + \Delta_i + \Delta_j))|_{\Delta_i}) \cong h^*((f^* \mathcal{L}_{ij})|_{\Delta_i}) \cong (g \circ h)^* \mathcal{L}_{ij}.$$

Now

$$\begin{aligned}
h^*(\mathcal{O}_X(m(K_{X/Y} + \Delta_i + \Delta_j))|_{\Delta_i}) &\cong \omega_Z^m(E) \otimes (g \circ h)^* \omega_Y^{[-m]}(mh^*(\Delta_j|_{\Delta_i})) \\
&\cong (g \circ h)^* \omega_Y^{[m]}(E + F) \otimes (g \circ h)^* \omega_Y^{[-m]}(mh^*(\Delta_j|_{\Delta_i})) \\
&\cong \mathcal{O}_Z(E + F + mh^*(\Delta_j|_{\Delta_i})).
\end{aligned}$$

Therefore from (9.3) it follows that the line bundle $(g \circ h)^* \mathcal{L}_{ij}$ is given by a Cartier divisor on Z . Since $g \circ h$ is bimeromorphic, it follows that $\mathcal{L}_{ij}$ corresponds to a Cartier divisor, say M_{ij} on Y . Let $L_{ij} := \frac{1}{m} M_{ij}$; then from (9.3) we have

$$(9.4) \quad \frac{1}{m}(E + F) + h^*(\Delta_j|_{\Delta_i}) \sim_{\mathbb{Q}} (g \circ h)^* L_{ij}.$$

Now according to Lemma 9.6, E , F and the conductor divisor of the normalization of Δ_i are all exceptional over Y . Thus pushing forward both sides of (9.4) we get $L_{ij} \sim_{\mathbb{Q}} (g \circ h)_*(h^*(\Delta_j)|_{\Delta_i}) = f_*(\Delta_i \cdot \Delta_j) = B_{ij}$. Then from (9.1) we have $K_{X/Y} + \Delta \sim_{\mathbb{Q}} f^* L$. $\square$

In the following we will show that L satisfies the other two properties.

Let $E_1, \dots, E_s$ be the distinct irreducible components of L . We will show that the base locus of $|L|_{\mathbb{Q}}$ does not have a divisorial component. To that end it is enough to show that for each $p \in \{1, \dots, s\}$, there exists an effective $\mathbb{Q}$ -divisor $L' \geq 0$ on Y such that $L' \sim_{\mathbb{Q}} L$ and E_p is not contained in the support of L' . We fix some $p \in \{1, 2, \dots, s\}$, and let $C_1, \dots, C_q$ be the curves in X contained in the support of the 1-cycles $\Delta_i \cdot \Delta_j$ for all $i \neq j \in \{1, \dots, k\}$ and dominate E_p , i.e., $f(C_i) = E_p$ for all $i = 1, \dots, q$. Now for each $r \in \{1, \dots, q\}$ we define

$$I_r := \{i \mid C_r \subseteq \text{Supp } \Delta_i\}.$$

Then $I_1, \dots, I_q$ forms a partition of the set $\{1, \dots, k\}$ (recall that $\Delta = \sum_i^k a_i \Delta_i$). Now since X has terminal singularities, X is smooth in a neighborhood of the general points of each C_r . Moreover, since (X, Δ) is canonical, by passing to a log resolution of (X, Δ) which extracts an exceptional divisor dominate C_r , it is easy to see that $\sum_{i \in I_r} a_i \leq 1$. Then by Lemma 9.4 there exist non-negative rational numbers $c_{ij} \geq 0$ such that $c_{ij} = c_{ji}$ for all i, j , and that $c_{ij} = 0$ whenever i, j belong to the same indexing set I_r . Following a similar construction as L given above we define $L' := \frac{1}{2} \sum_{i \neq j} c_{ij} f_*(\Delta_i \cdot \Delta_j)$. Then it is clear that E_p is not contained in the support of L' . Finally, from the construction of L and L' as above it also follows that $0 \leq L' \sim_{\mathbb{Q}} L$. $\square$

Lemma 9.10. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold sub-lc pair such that X has terminal singularities. Let $f : X \rightarrow Y$ be a K_X -Mori fiber*

space of relative dimension 1 such that $K_X + \Delta \equiv_f 0$. We can decompose $\Delta = \Delta_{\text{hor}} + \Delta_{\text{ver}}$ into horizontal and vertical parts over Y , and assume that Δ_{hor} is an effective $\mathbb{Q}$ -divisor. Assume that (X, Δ_{hor}) is a canonical pair and that all components of Δ_{hor} are bimeromorphic to Y via f . Then there exist $\mathbb{Q}$ -divisors D and L on Y such that

- (1) $f^*D = \Delta_{\text{ver}}$,
- (2) L is effective and $\text{Bs } |L|_{\mathbb{Q}}$ has no divisorial component,
- (3) $K_X + \Delta \sim_{\mathbb{Q}} f^*(K_Y + D + L)$, and
- (4) $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary if (X, Δ) is sub-klt.

Proof. Let F be a general fiber of f . Then $\Delta_{\text{ver}} \cdot F = 0$. Since f is the contraction of a K_X -negative extremal ray, by Lemma 9.7 there exists a $\mathbb{Q}$ -Cartier divisor D on Y such that $\Delta_{\text{ver}} = f^*D$. By Lemma 9.8, there is an effective $\mathbb{Q}$ -divisor $L \geq 0$ on Y such that $K_X + \Delta_{\text{hor}} \sim_{\mathbb{Q}} f^*(K_Y + L)$ and $\text{Bs } |L|_{\mathbb{Q}}$ has no divisorial component. Now if (X, Δ) is sub-klt, then Δ_{ver} is a pure sub-boundary, and thus so is D , since $\Delta_{\text{ver}} = f^*D$. Finally, since $\text{Bs } |L|_{\mathbb{Q}}$ does not have a divisorial component, it follows that $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary. $\square$

We will now deal with the case where the components of Δ_{hor} are not all bimeromorphic to the base Y . We will use the following covering trick.

Lemma 9.11 (Covering Trick). *Let $f : X \rightarrow Y$ be a proper fibration of relative dimension 1 between normal compact analytic varieties. Let S be a prime Weil divisor on X such that it is horizontal over Y . Then there is a commutative diagram*

$$\begin{array}{ccc} X' & \xrightarrow{p} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{\pi} & Y \end{array}$$

such that

- (1) Y' is normal and π is a proper finite morphism,
- (2) X' is the normalization of the main component of $X \times_Y Y'$ which dominates Y' ,
- (3) p is a proper finite morphism such that it is étale over a dense Zariski open subset of X ,
- (4) $S' = p^*S$ is an integral Weil divisor, and every component of S' is bimeromorphic to Y' via f' .

Proof. Note that the induced morphism $f|_S : S \rightarrow Y$ is generically finite. Let $S \rightarrow Y'' \rightarrow Y$ be the Stein factorization of $f|_S : S \rightarrow Y$, and $\pi : Y' \rightarrow Y''$

the normalization of Y'' . Then $S \rightarrow Y''$ is a bimeromorphic morphism. Let X' be the normalization of the main component of $X \times_Y Y'$ dominating Y' , and $f' : X' \rightarrow Y'$ and $p : X' \rightarrow X$ are the projections. Note that since $Y' \rightarrow Y$ is etale over a dense Zariski open subset of Y and etale morphisms are stable under base change, it follows that p is etale over a dense Zariski open subset of X .

Now let $S' := p^*S$ be defined as in Lemma 2.3, and S'' is the main component of $S \times_{Y''} Y'$ dominating Y' . Then $S'' \rightarrow Y'$ is bimeromorphic. The morphisms $S'' \rightarrow S \rightarrow X \rightarrow Y$ and $S'' \rightarrow Y' \rightarrow Y$ induce a morphism $S'' \rightarrow X'$. Note that the image of S'' on X' is an irreducible component of S' ; we denote this image again by S'' in X' . Then we can decompose S' as $S' = kS'' + T'$, where $k \in \mathbb{Z}^+$ and S'' is not contained in the support of T' . Thus S' has a component S'' which is bimeromorphic to Y' . We can continue this process until all the components of S' becomes bimeromorphic to Y' . It is easy to see that this process stops after a finitely many steps, since the degree of T' with respect to the general fiber of $f' : X' \rightarrow Y'$ is strictly smaller than the degree of S with respect to the general fibers of $f : X \rightarrow Y$. $\square$

Lemma 9.12. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold sub-lc pair such that X has klt singularities. Assume that X is non-algebraic and that the base of the MRC fibration of X has dimension 2. Let $f : X \rightarrow Y$ be a K_X -Mori fiber space of relative dimension 1 such that $K_X + \Delta \sim_{\mathbb{Q},f} 0$. We decompose $\Delta = \Delta_{\text{hor}} + \Delta_{\text{ver}}$, and assume that Δ_{hor} is effective. Then there exist two divisors L and D on Y such that*

- (1) $f^*D = \Delta_{\text{ver}}$,
- (2) $|L|_{\mathbb{Q}}$ is non-empty,
- (3) $K_X + \Delta \sim_{\mathbb{Q}} f^*(K_Y + D + L)$, and
- (4) $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary if (X, Δ) is sub-klt.

Proof. Since f is the contraction of a K_X -negative extremal ray and X has $\mathbb{Q}$ -factorial klt singularities, by [CHP16, Proposition 3.5(6)] there exists a line bundle $\mathcal{L}$ on Y such that

$$(9.5) \quad \mathcal{O}_X(m(K_X + \Delta_{\text{hor}})) \cong f^*(\mathcal{O}_X(mK_Y) \otimes \mathcal{L})$$

is an isomorphism of lines bundles for some $m \in \mathbb{N}$.

Next by Lemma 9.7 there exists a $\mathbb{Q}$ -Cartier divisor D on Y such that $\Delta_{\text{ver}} = f^*D$. We will now show that $\mathcal{L} \cong \mathcal{O}_X(mL)$ for some $\mathbb{Q}$ -Cartier divisor L such that the $\mathbb{Q}$ -linear system $|L|_{\mathbb{Q}}$ is non-empty. To that end, first applying Lemma 9.11 to the components of Δ_{hor} several times and then taking log resolution at the end we obtain a commutative diagram satisfying the following properties:

$$(9.6) \quad \begin{array}{ccc} (Z, \Theta) & \xrightarrow{p} & (X, \Delta) \\ g \downarrow & & \downarrow f \\ Y' & \xrightarrow{\pi} & Y \end{array}$$

- (i) π and p are generically finite proper morphisms, and g is a projective contraction,
- (ii) Y' is smooth and Z is a desingularization of the normalization of the main component of $X \times_Y Y'$ dominating Y' ,
- (iii) $K_Z + \Theta \sim_{\mathbb{Q}} p^*(K_X + \Delta)$,
- (iv) (Z, Θ) is a log smooth sub-lc pair such that $\text{Supp } \Theta$ is a smooth closed subset of Z ,
- (v) if we decompose $\Theta = \Theta_{\text{hor}} + \Theta_{\text{ver}}$ into horizontal and vertical parts over Y' , then Θ_{hor} is effective,
- (vi) every component of Θ_{hor} is bimeromorphic to Y' via g , and
- (vii) (Z, Θ_{hor}) has canonical singularities.

Note that (i) and (vi) follows from the repeated application of Lemma 9.11; (iii) follows from (ii) and Lemma 2.4; (iv) holds by [KM98, Corollary 2.32(2)]. Now we explain (v) and (vii). Since the desingularization Z is constructed via successive blow ups of a normal variety along smooth centers of codimension at least 2, the exceptional divisors of p are all vertical over Y' . Moreover, from Lemma 9.11 it follows that p is etale over a dense Zariski open subset of X , in particular, the ramification divisors of p are vertical over Y' . Therefore we have $\Theta_{\text{hor}} = p_*^{-1} \Delta_{\text{hor}}$, and hence Θ_{hor} is an effective boundary divisor; this is (v). Then (vii) follows from [KM98, Corollary 2.31(3)] using the fact that Θ_{hor} is a boundary divisor and $\text{Supp } \Theta_{\text{hor}}$ is smooth.

Now we run a $(K_Z + \Theta_{\text{hor}})$ -MMP over Y' as in Proposition 3.8, since $g : X' \rightarrow Y'$ is projective. Note that from our construction above it follows that $K_Z + \Theta_{\text{hor}} + \Theta_{\text{ver}} \sim_{\mathbb{Q}, g} 0$. Thus this MMP yields a pair $(Z', \Theta'_{\text{hor}})$ with $g' : Z' \rightarrow Y'$ such that $K_{Z'} + \Theta'_{\text{hor}}$ is g' -nef. We note that none of the components of Θ_{hor} is contracted by this MMP. Consequently, $(Z', \Theta'_{\text{hor}})$ has canonical singularities, and Z' has terminal singularities by Lemma 2.6. Let Θ'_{ver} be the bimeromorphic transformation of Θ_{ver} by the induced bimeromorphic map $Z \dashrightarrow Z'$. Then we have $K_{Z'} + \Theta'_{\text{hor}} + \Theta'_{\text{ver}} \sim_{\mathbb{Q}, g'} 0$, and $(Z', \Theta'_{\text{hor}} + \Theta'_{\text{ver}})$ has sub-lc singularities, since every step of this MMP is $(K_Z + \Theta)$ -trivial.

Now since $g' : Z' \rightarrow Y'$ is a projective morphism, by Proposition 3.8 we can run a $K_{Z'}$ -MMP over Y' with the scaling of Θ'_{hor} . Note that the general fibers of g' are isomorphic to $\mathbb{P}^1$ and $(K_{Z'} + \Theta'_{\text{hor}}) \cdot F = 0$ for a general fiber F of g' . Therefore this MMP terminates with a Mori fiber space, say $f' : X' \rightarrow Y''$ such that X' has $\mathbb{Q}$ -factorial terminal singularities. Let Δ'_{hor} and Δ'_{ver} be the bimeromorphic transform of Θ'_{hor} and Θ_{ver} , respectively, and $\Delta' := \Delta'_{\text{hor}} + \Delta'_{\text{ver}}$. From Lemma 4.6 it also follows that every step of this MMP is $(K_{Z'} + \Theta'_{\text{hor}})$ -trivial. In particular, $(K_{X'} + \Delta'_{\text{hor}}) \sim_{\mathbb{Q}, \tau \circ f'} 0$ and $(X', \Delta'_{\text{hor}})$ has canonical singularities, where $\tau : Y'' \rightarrow Y'$ is the induced bimeromorphic morphism. Moreover, we also have $(K_{X'} + \Delta') \sim_{\mathbb{Q}, \tau \circ f'} 0$ and (X', Δ') is sub-lc.

$$(9.7) \quad \begin{array}{ccccccc} (X', \Delta') & \leftarrow & \cdots & (Z', \Theta') & \leftarrow & \cdots & (Z, \Theta) \xrightarrow{p} (X, \Delta) \\ \downarrow f' & & & \searrow g' & & \downarrow g & \downarrow f \\ Y'' & \xrightarrow{\tau} & & Y' & \xrightarrow{\pi} & & Y \end{array}$$

Now applying Lemma 9.10 to $f' : (X', \Delta') \rightarrow Y''$ we see that there exist $\mathbb{Q}$ -Cartier divisors L'' and D'' on Y'' such that

- a) $f'^* D'' = \Delta'_{\text{ver}}$,
- b) L'' is effective, and $\text{Bs } |L''|_{\mathbb{Q}}$ does not have any divisorial component,
- c) $K_{X'} + \Delta'_{\text{hor}} \sim_{\mathbb{Q}} f'^*(K_{Y''} + L'')$, and
- d) $D'' + |L''|_{\mathbb{Q}}$ is a pure sub-boundary if (X', Δ') is sub-klt.

From our hypothesis it follows that Y' is a smooth compact Kähler non-uniruled surface, and thus both of the line bundles $\omega_{Y'}^m$ and $\omega_{Y''}^{[m]}$ correspond to effective Cartier divisors for sufficiently divisible $m \in \mathbb{N}$. Combining all of these with the Part (c) above it follows that $K_{Y''} + L'' \sim_{\mathbb{Q}} \tau^*(K_{Y'} + L')$, where $L' = \tau_* L''$. Also, since $K_{X'} + \Delta' \sim_{\mathbb{Q}, \tau \circ f'} 0$ and $K_{X'} + \Delta'_{\text{hor}} \sim_{\mathbb{Q}, \tau \circ f'} 0$, we have $\Delta'_{\text{ver}} \sim_{\mathbb{Q}, \tau \circ f'} 0$; in particular, from Part (a) above it follows that $D'' = \tau^* D'$, where $D' = \tau_* D''$. Thus we have

$$\Delta'_{\text{ver}} = (\tau \circ f')^* D', \quad K_{X'} + \Delta'_{\text{hor}} \sim_{\mathbb{Q}} (\tau \circ f')^*(K_{Y'} + L')$$

and

$$K_{X'} + \Delta' \sim_{\mathbb{Q}} (\tau \circ f')^*(K_{Y'} + L' + D').$$

Finally, from Part (d) it follows that $D' + |L'|_{\mathbb{Q}}$ is a pure sub-boundary if (X', Δ') is sub-klt. Indeed, if F' is a prime Weil divisor on Y' and $F'' := \tau_*^{-1} F'$, then there exists an effective $\mathbb{Q}$ -Cartier divisor $M'' \in |L''|_{\mathbb{Q}}$ such that $\text{mult}_{F''}(D'' + M'') < 1$, since $D'' + |L''|_{\mathbb{Q}}$ is a pure sub-boundary. Thus by

pushing forward (as τ is bimeromorphic) we get that $\text{mult}_{F'}(D' + M') < 1$, where $M' := \tau_* M'' \in |L'|$; hence

$$(9.8) \quad D' + |L'|_{\mathbb{Q}} \text{ is a pure sub-boundary.}$$

From (9.5) we also have $\mathcal{O}_X(m(K_Z + \Theta)) \cong (\pi \circ g)^*(\mathcal{O}_Y(m(K_Y + D))) \otimes \mathcal{L}$ for sufficiently divisible $m \in \mathbb{N}$. Let W be the normalization of the graph of the bimeromorphic contraction $\phi : Z \dashrightarrow X'$ over Y' , and $q : W \rightarrow Z$ and $r : W \rightarrow X'$ are the corresponding projections. Then from our construction above and the negativity lemma it follows that $q^*(K_Z + \Theta) = r^*(K_{X'} + \Delta')$. In particular, we have

$$(\tau \circ f' \circ r)^* \mathcal{O}_{Y'}(m(K_{Y'} + D' + L')) \cong (\pi \circ g \circ q)^*(\mathcal{O}_Y(m(K_Y + D))) \otimes \mathcal{L}$$

for sufficiently divisible $m \in \mathbb{N}$.

Pushing forward both side by $\tau \circ f' \circ r = g \circ p$ and then applying the projection formula we get

$$(9.9) \quad \mathcal{O}_{Y'}(m(K_{Y'} + L' + D')) \cong \pi^*(\mathcal{O}_Y(m(K_Y + D))) \otimes \mathcal{L}.$$

Now since Y and Y' are both non-uniruled compact Kähler surface with klt singularities, it follows that $\mathcal{O}_Y(mK_Y)$ and $\mathcal{O}_{Y'}(mK_{Y'})$ are both given by Cartier divisors for sufficiently divisible $m \in \mathbb{N}$. Thus from (9.9) we see that $\mathcal{L}$ correspond to a $\mathbb{Q}$ -Cartier divisor L on Y such that $\mathcal{L} \cong \mathcal{O}_Y(mL)$ for some sufficiently divisible m . Therefore from (9.9) we have

$$(9.10) \quad K_{Y'} + D' + L' \sim_{\mathbb{Q}} \pi^*(K_Y + L + D).$$

Now our main goal is to show that $|L|_{\mathbb{Q}}$ is non-empty. To that end, first note that since $\pi : Y' \rightarrow Y$ is a generically finite morphism, we can write

$$(9.11) \quad K_{Y'/Y} = R_{\text{nex}} + R_{\text{ex}},$$

where R_{ex} is exceptional over Y , R_{nex} is not exceptional over Y .

Note that R_{nex} is effective. Then from (9.10) we can write

$$(9.12) \quad \pi^* L \sim_{\mathbb{Q}} L' + E + F,$$

where $E = (D' - \pi^* D)_{\text{ex}} + R_{\text{ex}}$ and $F = (D' - \pi^* D)_{\text{nex}} + R_{\text{nex}}$.

Claim 9.13. Then divisor $F = (D' - \pi^* D)_{\text{nex}} + R_{\text{nex}}$ is effective.

Assuming the claim for the time being we will first show that $|L|_{\mathbb{Q}}$ is non-empty. Indeed, recall that L'' is an effective divisor such that the base-locus of $|L''|_{\mathbb{Q}}$ contains only finitely many points, and thus by [Fuj83, Corollary 1.14] L'' is semi-ample. Then from the projection formula it follows that $L' = \tau_* L''$ is algebraically nef on Y' , i.e. $L' \cdot C \geq 0$ for all curves $C \subseteq Y'$. Now if F is effective, then it is algebraically π -nef, i.e., $F \cdot C \geq 0$ for every π -exceptional curve $C \subseteq Y'$, since none of the components of F are π -exceptional. Thus by

the negativity lemma it follows from (9.12) that E is effective. In particular, it again follows from (9.12) that $|L|_{\mathbb{Q}}$ is non-empty.

Proof of Claim 9.13. Recall the resolution $q : W \rightarrow Z$ and $r : W \rightarrow X'$ of the map $\phi : Z \dashrightarrow X'$. We can write

$$(9.13) \quad r^*(K_{X'} + \Delta'_{\text{hor}}) = q^*(K_Z + \Theta_{\text{hor}}) + q^*(\Theta_{\text{ver}} - g^*D').$$

Recall that $\phi : Z \dashrightarrow X'$ is a composition of a $(K_Z + \Theta_{\text{hor}})$ -MMP followed by a $K_{Z'}$ -MMP. Since every step of the $K_{Z'}$ was $(K_Z + \Theta_{\text{hor}})$ -trivial, it follows that the difference $q^*(K_Z + \Theta_{\text{hor}}) - r^*(K_{X'} + \Delta'_{\text{hor}})$ is an effective $\mathbb{Q}$ -divisor. Therefore from (9.13) we have

$$(9.14) \quad \Theta_{\text{ver}} - g^*D' \leq 0.$$

Let D'_1 be a prime Weil divisor on Y' which is not π -exceptional. We will show that $\text{mult}_{D'_1}(F) \geq 0$. To that end set $D_1 := \pi(D'_1)$ and let Δ_1 be a Weil divisor on X such that $\text{Supp}(\Delta_1) = \text{Supp}(f^*D_1)$. Note that Δ_1 is irreducible, since f has equidimensional fibers. Now let Θ_1 be a prime Weil divisor on Z such that it dominates Δ_1 . Let m, n, m' and n' be the coefficients of D'_1 in π^*D_1 , Δ_1 in f^*D_1 , Θ_1 in $p^*\Delta_1$ and Θ_1 in $g^*D'_1$, respectively, i.e., $\text{mult}_{D'_1}(\pi^*D_1) = m$, $\text{mult}_{\Delta_1}(f^*D_1) = n$, $\text{mult}_{\Theta_1}(p^*\Delta_1) = m'$ and $\text{mult}_{\Theta_1}(g^*D'_1) = n'$.

Now we separate our discussion into two cases depending on whether $\text{mult}_{D_1}(D)$ is 0 or not.

Case I: $\text{mult}_{D_1}(D) \neq 0$, where $f^*D = \Delta_{\text{ver}}$. In this case Δ_1 is a component of Δ_{ver} . Let $\text{mult}_{\Delta_1}(\Delta_{\text{ver}}) = \lambda$. Then $\text{mult}_{D_1}(D) = \frac{\lambda}{n}$ and $\text{mult}_{D'_1}(\pi^*D) = \frac{m\lambda}{n}$. Recall that by our construction the generically finite morphism $p : Z \rightarrow X$ is etale over a dense Zariski open subset of X . In particular, the ramification divisor of p is vertical over Y' , and thus $\Theta_{\text{hor}} = p_*^{-1}\Delta_{\text{hor}}$ and $K_Z + \Theta_{\text{ver}} \sim_{\mathbb{Q}} p^*(K_X + \Delta_{\text{ver}})$. Therefore $\text{mult}_{\Theta_1}(\Theta_{\text{ver}}) = m'\lambda - (m' - 1)$. Moreover, from (9.14) it follows that $\text{mult}_{D'_1}(D') \geq \frac{1}{n'} \text{mult}_{\Theta_1}(\Theta_{\text{ver}}) = \frac{m'\lambda - (m' - 1)}{n'}$. Also note

that $\text{mult}_{D'_1}(R_{\text{nex}}) = m - 1$. Therefore we have

$$\begin{aligned}
\text{mult}_{D'_1}((D' - \pi^*D)_{\text{nex}} + R_{\text{nex}}) &\geq \frac{m'\lambda - (m' - 1)}{n'} - \frac{m\lambda}{n} + (m - 1) \\
&= \frac{m'\lambda - (m' - 1)}{n'} - \frac{m'\lambda}{n'} + (m - 1) \quad (\text{since } mn' = m'n) \\
&= (m - 1) - \frac{m' - 1}{n'} \\
&= \frac{1}{n'}(mn' - n' - m' + 1) \\
&= \frac{1}{n'}(dm'n' - n' - m' + 1) \quad (\text{since } m' = \frac{m}{d} \text{ by Lemma A.3, where } d = \gcd(m, n)) \\
&\geq \frac{1}{n'}(m' - 1)(n' - 1) \geq 0.
\end{aligned}$$

Case II: $\text{mult}_{D_1}(D) = 0$. In this case Δ_1 is not contained in the support of Δ_{ver} , however $\text{mult}_{\Theta_1}(\Theta_{\text{ver}})$ could still be non-zero if Θ_1 is a ramification divisor of p . First assume that $\text{mult}_{\Theta_1}(\Theta_{\text{ver}}) = 0$. Then from (9.14) it follows that $\text{mult}_{D'_1}(D') \geq 0$. Thus we have $\text{mult}_{D'_1}((D' - \pi^*D)_{\text{nex}} + R_{\text{nex}}) \geq 0$, since $\text{mult}_{D'_1}(\pi^*D) = 0$ and $R_{\text{nex}} \geq 0$ is an effective divisor.

Now we consider the possibility that $\text{mult}_{\Theta_1}(\Theta_{\text{ver}}) \neq 0$. In this case we have $\text{mult}_{D'_1}(\pi^*D) = 0$, $\text{mult}_{D'_1}(R_{\text{nex}}) = m - 1$, $\text{mult}_{\Delta_1}(\Delta_{\text{ver}}) = 0$ and $\text{mult}_{\Theta_1}(\Theta_{\text{ver}}) = -(m' - 1)$. From (9.14) we have $\text{mult}_{D'_1}(D') \geq -\frac{m'-1}{n'}$. Therefore we have

$$\begin{aligned}
\text{mult}_{D'_1}((D' - \pi^*D)_{\text{nex}} + R_{\text{nex}}) &\geq -\frac{m' - 1}{n'} + (m - 1) \\
&= \frac{1}{n'}(mn' - n' - m' + 1) \\
&= \frac{1}{n'}(dm'n' - n' - m' + 1) \quad (\text{since } m' = \frac{m}{d} \text{ by Lemma A.3, where } d = \gcd(m, n)) \\
&\geq \frac{1}{n'}(m' - 1)(n' - 1) \geq 0.
\end{aligned}$$

□

It remains to show that $D + |L|_{\mathbb{Q}}$ is a pure sub-boundary if (X, Δ) is sub-plt. By Lemma 8.6 we only need to show that

$$(\pi^*D - K_{Y'/Y}) + |\pi^*L|_{\mathbb{Q}}$$

is a pure sub-boundary. From (9.11) and (9.12) we see that

$$\begin{aligned}
(\pi^*D - K_{Y'/Y}) &\sim_{\mathbb{Q}} \pi^*D - (R_{\text{ex}} + R_{\text{nex}}) = D' - (E + F) \quad \text{and} \\
\pi^*L &\sim_{\mathbb{Q}} L' + E + F,
\end{aligned}$$

where $E = (D' - \pi^* D)_{\text{ex}} + R_{\text{ex}}$ and $F = (D' - \pi^* D)_{\text{nex}} + R_{\text{nex}}$ are both effective divisors as proved above.

Therefore it is enough to show that

$$(9.15) \quad D' - (E + F) + |L' + E + F|$$

is a pure sub-boundary.

It is easy to see that (9.15) is a pure sub-boundary if $D' + |L'|$ is a pure sub-boundary, as $E + F$ is effective. But the fact that $D' + |L'|$ is a pure sub-boundary is already proved in (9.8), so we are done. $\square$

We can now prove Theorem 9.5.

Proof of Theorem 9.5. Let D and L be the two divisors in Y given by Lemma 9.12. Since Δ is effective, D is also effective. Hence we can take B as any element in $D + |L|_{\mathbb{Q}}$. It remains to show that (Y, B) can be chosen to be klt if (X, Δ) is klt. By Lemma 8.4 it is enough to show that $(Y, D + |L|_{\mathbb{Q}})$ is sub-klt. We will check this using Lemma 8.3 and 8.6. Let $\bar{\pi} : \bar{Y} \rightarrow Y$ be any proper bimeromorphic morphism with $\bar{Y}$ smooth. Let $\bar{X}$ be the resolution of singularities of the unique component of $X \times_Y \bar{Y}$ dominating $\bar{Y}$ so that $\bar{p} : \bar{X} \rightarrow X$ is a log resolution of the pair (X, Δ) . Let $\bar{f} : \bar{X} \rightarrow \bar{Y}$ be the induced morphism and $K_{\bar{X}} + \bar{\Delta} = \bar{p}^*(K_X + \Delta)$. Then clearly $\bar{\Delta}_{\text{hor}} = \bar{p}_*^{-1} \Delta_{\text{hor}}$ and $K_{\bar{X}} + \bar{\Delta}_{\text{ver}} = \bar{p}^*(K_X + \Delta_{\text{ver}})$. Then as in the proof of Lemma 9.12 we can construct a diagram like (9.7) such that $\bar{f} : \bar{X} \rightarrow \bar{Y}$ sits in between the morphism $g : Z \rightarrow Y'$ and $f : X \rightarrow Y$ and Z is obtained from $\bar{X}$ via base-change by the components of $\bar{\Delta}_{\text{hor}}$ followed a resolution of singularities.

$$(9.16) \quad \begin{array}{ccccccc} (X', \Delta') & \leftarrow \cdots \leftarrow & (Z', \Theta') & \leftarrow \cdots \leftarrow & (Z, \Theta) & \xrightarrow{\quad p \quad} & (\bar{X}, \bar{\Delta}) \xrightarrow{\quad \bar{p} \quad} (X, \Delta) \\ \downarrow f' & & \searrow g' & \downarrow g & & \downarrow \bar{f} & \downarrow f \\ Y'' & \xrightarrow{\quad \tau \quad} & Y' & \xrightarrow{\quad \alpha \quad} & \bar{Y}_{\pi} & \xrightarrow{\quad \bar{\pi} \quad} & Y \end{array}$$

Let $\alpha : Y' \rightarrow \bar{Y}$ be generically finite morphism constructed in the process above and $\pi = \bar{\pi} \circ \alpha : Y' \rightarrow Y$ is the composite morphism. Then as in the proof of Lemma 9.12 we know that

$$(9.17) \quad (\pi^* D - K_{Y'/Y}) + |\pi^* L| \quad \text{is a pure sub-boundary.}$$

Recall that we want to show that $(Y, D + |L|)$ sub-plt, and by Lemma 8.3 it is equivalent to show that the log pullback of $D + |L|$ onto $\bar{Y}$ defined as

$$(9.18) \quad (\bar{\pi}^*D - K_{\bar{Y}/Y}) + |\bar{\pi}^*L|$$

is a pure sub-boundary.

Now since $\alpha : Y' \rightarrow \bar{Y}$ is a generically proper morphism, by Lemma 8.6 it is enough to show that the log pullback of (9.18) onto Y' defined as

$$(\alpha^*(\bar{\pi}^*D - K_{\bar{Y}/Y}) - K_{Y'/\bar{Y}}) + |\alpha^*\bar{\pi}^*L|$$

is a pure sub-boundary.

Note that the above expression is same as $(\pi^*D - K_{Y'/Y}) + |\pi^*L|$ and this is a pure sub-boundary by (9.17), so we are done. $\square$

9.2. Proof of the non-vanishing theorem. We will first prove the following lemma.

Lemma 9.14. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Assume that X is non-algebraic and $f : X \rightarrow Y$ is a projective MRC fibration of X to a compact Kähler surface Y . Assume that every component of Δ is horizontal over Y and $(K_X + \Delta) \cdot F \geq 0$ for a general fiber F of f . Then $K_X + \Delta$ is $\mathbb{Q}$ -effective.*

Proof. It is enough to prove the case when $K_X + \Delta$ has intersection number 0 with general fibers of f . Since f is projective, by Proposition 3.8 we can run a relative $(K_X + \Delta)$ -MMP over Y . Such a MMP terminates by Theorem 3.3. We then obtain a dlt pair (Z, D) such that $K_Z + D$ is relatively nef over Y . It is enough to prove that $K_Z + D$ is $\mathbb{Q}$ -effective.

Let $g : Z \rightarrow Y$ be the induced projective contraction. We can run a relative K_Z -MMP over Y such that every step is $(K_Z + D)$ -trivial. Again, by Theorem 3.3 this MMP terminates and we can obtain a dlt pair (X', Δ') with a induced projective morphism $h : X' \rightarrow Y$ such that

- (1) either there is a $K_{X'}$ -Mori fiber space $f' : X' \rightarrow Y'$ such that $(K_{X'} + \Delta') \sim_{\mathbb{Q}, f'} 0$,
- (2) or every $K_{X'}$ -negative extremal ray of $\overline{\text{NE}}(X'/Y)$ is $(K_{X'} + \Delta')$ -positive.

We claim that the case (2) above is not possible; indeed observe that there is a dense Zariski open subset $V \subseteq Y$ such that $g^{-1}V \cong h^{-1}V$. In particular, $(K_{X'} + \Delta') \cdot F = (K_Z + D) \cdot F = 0$ for general fibers F of h . However, $K_{X'} \cdot F = -2$, and thus by case (2) above we must have $(K_{X'} + \Delta') \cdot F > 0$, this is a contradiction.

Therefore we are in the first case. By Theorem 9.5 there is an effective divisor B' in Y' such that

$$K_{X'} + \Delta' \sim_{\mathbb{Q}} f'^*(K_{Y'} + B').$$

Since f' is the MRC fibration of X' and X' is non-algebraic, it follows that Y' is non-uniruled (see [HP15, Remark 3.2]). Hence $K_{Y'}$ is pseudo-effective and by abundance for Kähler surfaces (see Theorem 6.3) we obtain that $K_{Y'} + B'$ is $\mathbb{Q}$ -effective. This completes the proof of the lemma. $\square$

Now we can deduce Theorem 9.1.

Proof of Theorem 9.1. It is enough to prove that (3) implies (1). By passing to a dlt model as in Corollary 5.3 we may assume that (X, Δ) has $\mathbb{Q}$ -factorial dlt singularities such that X has terminal singularities. By [HP15, Theorem 1.1] there is a K_X -MMP which terminates with a Mori fiber space $f : X' \rightarrow Y$ such that Y is a $\mathbb{Q}$ -factorial compact Kähler surface with klt singularities. Let $p : W \rightarrow X$ and $q : W \rightarrow X'$ be the resolution of the graph of the bimeromorphic map $\phi : X \dashrightarrow X'$ such that p is a log resolution of the pair (X, Δ) and q is a projective morphism. Then we can write

$$K_W + \Delta_W = p^*(K_X + \Delta) + E,$$

where $E \geq 0$ and $\Delta_W \geq 0$ are two effective $\mathbb{Q}$ -divisors without common components and $p_*\Delta_W = \Delta$ and $p_*E = 0$. Thus it is enough to show that $K_W + \Delta_W$ is $\mathbb{Q}$ -effective. We note that (W, Δ_W) is dlt and $f \circ q : W \rightarrow Y$ is the MRC fibration of W , since $f : X' \rightarrow Y$ is the MRC fibration of X' and the fibers of $W \rightarrow X'$ are rationally chain connected by Proposition 3.9. Moreover, observe that there is a dense Zariski open subset $V \subseteq Y$ such that ϕ is an isomorphism over $f^{-1}V$. In particular, for general fibers F of f (also of $f \circ q$) we have $(K_W + \Delta_W) \cdot F = (K_{X'} + \Delta') \cdot F = (K_X + \Delta) \cdot F \geq 0$. Thus by Lemma 9.14, $K_W + \Delta_W$ is $\mathbb{Q}$ -effective, this completes the proof. $\square$

10. CONE AND CONTRACTION THEOREMS FOR PSEUDO-EFFECTIVE PAIRS

In this section we will establish the cone and contraction theorem for a dlt pair (X, Δ) such that X is a $\mathbb{Q}$ -factorial uniruled non-algebraic compact Kähler threefold such that $K_X + \Delta$ is pseudo-effective. The proof is based on the same philosophy as in [HP16, CHP16]. We work under the following hypothesis throughout this section.

Assumption 10.1. Let X be a non-algebraic compact Kähler threefold with $\mathbb{Q}$ -factorial singularities. Assume that X is uniruled. Then the base of the MRC fibration of X is a non-uniruled surface (see [HP15, Remark 3.2]). Assume that there is a boundary Δ such that (X, Δ) is dlt, and that $K_X + \Delta$ is pseudo-effective. Then by Theorem 9.1, $K_X + \Delta$ is $\mathbb{Q}$ -effective and we can write

$$(10.1) \quad K_X + \Delta \sim_{\mathbb{Q}} \sum_{i=1}^N \lambda_i S_i,$$

where $\lambda_i \geq 0$ are rational numbers and S_i are prime Weil divisors on X .

10.1. Negative part of $K_X + \Delta$. The goal of this subsection is to prove the following lemma, which is a key ingredient for various results in this section.

Lemma 10.2. *With notations and hypothesis as in Assumption 10.1, let $S \subseteq X$ be a surface such that $(K_X + \Delta)|_S$ is not pseudo-effective. Then $S = S_j$ is one of the component of the decomposition 10.1. Moreover, S is Moishezon and its minimal resolution $\hat{S}$ is projective and uniruled.*

We follow the same strategy as in [HP16, Lemma 4.1]. We will consider two cases according to whether S is horizontal over the base of the MRC fibration of X or not.

Lemma 10.3. *With notations and hypothesis as in Assumption 10.1, let $S \subseteq X$ be a surface such that S is horizontal over the base of the MRC fibration of X . Then $(K_X + \Delta)|_S$ is pseudo-effective.*

Proof. Let λ be the positive rational number such that $(K_X + \lambda\Delta) \cdot F = 0$, where F is the general fiber of the base of the MRC fibration $f : X \dashrightarrow Z$. Then $K_X + \lambda\Delta$ is $\mathbb{Q}$ -effective by Theorem 9.1. Moreover, since S is horizontal over Z , $S \cdot F > 0$ for general fibers F of f . In particular, S is not a component of any element of $|K_X + \lambda\Delta|_{\mathbb{Q}}$. Hence $(K_X + \lambda\Delta)|_S$ is pseudo-effective. Thus if S is not a component of Δ , then it follows that $(K_X + \Delta)|_S$ is pseudo-effective. So now we assume that S is a component of Δ . Since Δ is a boundary, there is some rational number $\mu \geq 1 - \lambda > 0$ such that the multiplicity of $(\lambda + \mu)\Delta$ along S is 1. Let $\pi : \hat{S} \rightarrow X$ be the minimal resolution of S . Then there is an effective divisor E on $\hat{S}$ such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*((K_X + (\lambda + \mu)\Delta)|_S).$$

We note that S is not uniruled, since $f|_S : S \dashrightarrow Z$ is a generically finite map and Z is not uniruled. Hence $K_{\hat{S}}$ is pseudo-effective and consequently, $(K_X + (\lambda + \mu)\Delta)|_S$ is pseudo-effective. Since $\mu \geq 1 - \lambda$, we obtain that

$\gamma(K_X + \Delta)$ is a convex combination of $K_X + \lambda\Delta$ and $K_X + (\lambda + \mu)\Delta$ for some rational number $\gamma > 0$. This proves that $(K_X + \Delta)|_S$ is pseudo-effective. $\square$

Lemma 10.4. *Let X be a non-algebraic compact Kähler 3-fold with $\mathbb{Q}$ -factorial klt singularities. Let $S \subseteq X$ be a surface such that S is vertical over the base of the MRC fibration of X . Then S is Moishezon and its minimal resolution $\hat{S}$ is projective and uniruled.*

Proof. First note that according to [HP15, Remark 3.2] the base of the MRC fibration of X is a surface. Now replacing X by a terminal model as in Theorem 3.6 we may assume that X has $\mathbb{Q}$ -factorial terminal singularity. Then by [HP15, Theorem 1.1] X is bimeromorphic to a Mori fiber space $X' \rightarrow Y$ over some compact Kähler surface Y . Let $\tilde{X}$ be a common resolution of X and X' such that $\tilde{X} \rightarrow X'$ is projective. Let $\tilde{S}$ be the strict transform of S in $\tilde{X}$. Then $\tilde{S}$ is vertical over Y for the induced projective fibration $g : \tilde{X} \rightarrow Y$. Now we run a $K_{\tilde{X}}$ -MMP over Y , and obtain a Mori fiber space $\pi : \bar{X} \rightarrow T$ over Y . If $\tilde{S}$ is contracted by this MMP, then it is uniruled by Proposition 3.9. Otherwise, let $\bar{S}$ be the bimeromorphic transform of $\tilde{S}$ onto $\bar{X}$ such that $\bar{S}$ is vertical over T . In this case, $\bar{S}$ is again uniruled by Proposition 3.9, since the Mori fiber space $\pi : \bar{X} \rightarrow T$ is the MRC fibration of $\bar{X}$. We can then conclude that S is uniruled, and so is the minimal resolution $\hat{S}$. This implies that $\hat{S}$ is projective by classification, and thus S is Moishezon. $\square$

Now we are ready to conclude Lemma 10.2.

Proof of Lemma 10.2. By Lemma 10.3, the surface S is vertical over the base of the MRC fibration. The lemma then follows from Lemma 10.4. $\square$

Corollary 10.5. *With notations and hypothesis as in Assumption 10.1, $K_X + \Delta$ is nef if and only if it has non-negative intersection with every curve in X .*

Proof. If $K_X + \Delta$ is nef, then it is clear that $(K_X + \Delta) \cdot C \geq 0$ for all curves $C \subseteq X$. So now assume that $K_X + \Delta$ has non-negative intersection with every curve in X . Then by [Pau98, Theorem 3, Page 413] it is enough to prove that $(K_X + \Delta)|_S$ is pseudo-effective for every surface $S \subseteq X$. Assume by contradiction that this is not the case for some surface S . Then by Lemma 10.2, the minimal resolution $\hat{S}$ of S is projective. By assumption, $\pi^*((K_X + \Delta)|_S)$ is not pseudo-effective, where $\pi : \hat{S} \rightarrow X$ is the induced morphism. By [BDPP13, Theorem 0.2], $\hat{S}$ is covered by a family of curves which have negative intersection with $\pi^*((K_X + \Delta)|_S)$. Since π is bimeromorphic onto its image, the image of these curves determine a covering family of curves on $S \subseteq X$ such that they have negative intersection with $K_X + \Delta$; this is a contradiction. $\square$

10.2. Bend-and-break theorem for pseudo-effective $K_X + \Delta$.

In this subsection, we will show the following bend-and-break theorem.

Theorem 10.6. *With notations and hypothesis as in Assumption 10.1, $K_X + \Delta$ has the bend-and-break property, i.e. there is a rational number $d > 0$ such that if $C \subseteq X$ is a curve with $-(K_X + \Delta) \cdot C > d$, then*

$$[C] = [C_1] + [C_2],$$

where C_1, C_2 are two integral nonzero effective 1-cycles.

We will first prove this theorem under an additional assumption that X has terminal singularities. The proof of the general case then follows from [CHP16, Lemma 4.7].

Lemma 10.7. *With notations and hypothesis as in Assumption 10.1, we further assume that X has terminal singularities. Let $C \subseteq X$ be a curve such that $(K_X + \Delta) \cdot C < 0$ and $\dim_C \text{Chow}(X) > 0$. Then there is a unique surface S_j in the decomposition (10.1) such that C and all of its deformations are contained in S_j . Moreover, we have $K_{S_j} \cdot C < K_X \cdot C$.*

Proof. As in the first part of the proof of [HP16, Lemma 5.4] we see that there is some S_j in the decomposition (10.1) which contains a family of deformations of C . Since C deforms, $S_k \cdot C \geq 0$ for any $k \neq j$. Then from the decomposition (10.1) it follows that $S_j \cdot C < 0$. In particular, S_j is the unique surface which contains all of the deformations of C . Moreover, by adjunction we have

$$K_X \cdot C = K_X|_{S_j} \cdot C > (K_X + S_j)|_{S_j} \cdot C \geq K_{S_j} \cdot C.$$

□

Lemma 10.8. *With notations and hypothesis as in Assumption 10.1, we further assume that X has terminal singularities. Let $\mathcal{A}$ be the collection of curves $C \subseteq X$ which satisfy all of the following conditions.*

- (1) $(K_X + \Delta) \cdot C < 0$,
- (2) $\Delta \cdot C < 0$
- (3) C is contained in a component of Δ which is horizontal over the base of the MRC fibration $f : X \dashrightarrow Z$.

Then $\mathcal{A}$ is a finite set.

Proof. Let S be a component of Δ which is horizontal over Z and $C \subseteq S$ a curve from the set $\mathcal{A}$. Then there is a rational number $\mu \geq 0$ such that the coefficient of S in $(1 + \mu)\Delta$ is 1. Moreover, we have

$$K_X + (1 + \mu)\Delta \cdot C < 0.$$

Let $\hat{S} \rightarrow S$ be the minimal resolution of S and $\pi : \hat{S} \rightarrow X$ is the induced morphism. Observe that in order to prove the lemma it is enough to show that there are only finitely many curves in $\hat{S}$ which have negative intersection with $\pi^*(K_X + (1 + \mu)\Delta)$.

Now by adjunction there is an effective $\mathbb{Q}$ -divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + (1 + \mu)\Delta).$$

Note that S is not uniruled, since $f|_S : S \dashrightarrow Z$ is generically finite and Z is not uniruled. Thus $K_{\hat{S}}$ is pseudo-effective and by Theorem 6.3 it follows that there is some effective $\mathbb{Q}$ -divisor $0 \leq D \sim_{\mathbb{Q}} K_{\hat{S}}$. Then any curve in $\hat{S}$ which has negative intersection with $\pi^*(K_X + (1 + \mu)\Delta)$ must be a component of $D + E$. Hence $\mathcal{A}$ is a finite set. $\square$

For every component Δ_i of Δ which is vertical over the base of the MRC fibration $f : X \dashrightarrow Z$, there is a rational number $\mu_i \geq 0$ such that the multiplicity of $(1 + \mu_i)\Delta$ along Δ_i is 1. Let $\hat{\Delta}_i \rightarrow \Delta_i$ be the minimal resolution of Δ_i and $\pi : \hat{\Delta}_i \rightarrow X$ is the induced morphism. Now by adjunction there is an effective $\mathbb{Q}$ -divisor E_i such that

$$K_{\hat{\Delta}_i} + E_i \sim_{\mathbb{Q}} \pi_i^*(K_X + (1 + \mu_i)\Delta).$$

We let $\mathcal{B}$ be the collection of all curves in X which are contained in $\pi_i(\text{Supp } E_i)$ for some i . Then $\mathcal{B}$ is a finite set.

Finally, we let $\mathcal{C}$ be the collection of curves which lies either in the singular locus of the support of Δ or in the singular locus of $\sum S_i$, where S_i are the components of the decomposition (10.1). Then again $\mathcal{C}$ is a finite set.

Lemma 10.9. *With notations and hypothesis as in Assumption 10.1 we further assume that X has terminal singularities. Define a rational number $b > 0$ as follows:*

$$b = \max\{2, -(K_X + \Delta) \cdot Z \mid Z \subseteq X \text{ is a curve and } Z \in \mathcal{A} \cup \mathcal{B} \cup \mathcal{C}\},$$

where $\mathcal{A}$ is a finite set defined in Lemma 10.8. Then for any curve $C \subseteq X$, the inequality

$$-(K_X + \Delta) \cdot C > b$$

implies that $\dim_C \text{Chow}(X) > 0$.

Proof. Let $C \subseteq X$ be a curve such that $-(K_X + \Delta) \cdot C > b$. Then from the definition of b it follows that $C \notin \mathcal{A} \cup \mathcal{B} \cup \mathcal{C}$. Now first we assume that $\Delta \cdot C < 0$. Then there is some component S of Δ which contains C . Since $C \notin \mathcal{A}$, S is not horizontal over the base of the MRC fibration $f : X \dashrightarrow Z$. We also note that S is a unique component of Δ containing C , since otherwise

$C \in \mathcal{C}$. Since (X, Δ) is dlt, there is a rational number $\mu \geq 0$ such that the multiplicity of $(1 + \mu)\Delta$ along S is 1. Then

$$-(K_X + (1 + \mu)\Delta) \cdot C > b.$$

Let $\hat{S} \rightarrow S$ be the minimal resolution of S and $\pi : \hat{S} \rightarrow X$ is the induced morphism. By adjunction there is an effective $\mathbb{Q}$ -divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + (1 + \mu)\Delta).$$

Since $C \notin \mathcal{B}$, C is not contained in $\pi(\text{Supp } E)$. Hence the strict transform of C in $\hat{S}$ is a curve $\hat{C}$ and we have

$$K_{\hat{S}} \cdot \hat{C} \leq (K_{\hat{S}} + E) \cdot \hat{C} = (K_X + (1 + \mu)\Delta) \cdot C < -b.$$

Since $\hat{S}$ is a smooth projective surface and $b \geq 2$, from [Kol96, Theorem II.1.15] we have

$$\dim_{\hat{C}} \text{Chow}(\hat{S}) > 0.$$

Pushing forward a deformation family of $\hat{C}$ onto S gives deformation family of C on S , in particular, $\dim_C \text{Chow}(X) > 0$.

Now we assume that $\Delta \cdot C \geq 0$. Then we have

$$K_X \cdot C \leq (K_X + \Delta) \cdot C < -b \leq -2.$$

Then by [HP16, Theorem 4.5] C is not very rigid and some multiple mC deforms in X . Hence there is some component $S = S_j$ in the decomposition (10.1) such that $S \cdot C < 0$ and that S contains a family of deformations of mC . Moreover, since $C \notin \mathcal{C}$, the component $S = S_j$ is unique.

Let $\hat{S} \rightarrow S$ be the minimal resolution of S and $\pi : \hat{S} \rightarrow X$ is the induced morphism. By adjunction there is an effective $\mathbb{Q}$ -divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + S).$$

Since $C \notin \mathcal{C}$, it is not contained in $\pi(\text{Supp } E)$. Hence the strict transform of C in $\hat{S}$ is a curve $\hat{C}$ and we have

$$K_{\hat{S}} \cdot \hat{C} \leq (K_{\hat{S}} + E) \cdot \hat{C} = (K_X + S) \cdot C < K_X \cdot C < -b \leq -2.$$

As in the previous case, this implies that $\dim_C \text{Chow}(X) > 0$. $\square$

Lemma 10.10. *With notations and hypothesis as in Assumption 10.1 we further assume that X has terminal singularities. Define*

$$d := \max\{3, b\},$$

where b is the positive rational number defined in Lemma 10.9. Then if $C \subseteq X$ is a curve, the inequality $-(K_X + \Delta) \cdot C > d$ implies that

$$[C] = [C_1] + [C_2],$$

where C_1, C_2 are two non-zero integral effective 1-cycles. In particular, $K_X + \Delta$ has the bend-and-break property.

Proof. As in the proof of Lemma 10.9, we need to consider two cases: either $\Delta \cdot C < 0$ or $\Delta \cdot C \geq 0$. As in the proof of Lemma 10.9, in both cases we end up with a unique uniruled surface S such that C is not contained in the singular locus of S . Let $\hat{S} \rightarrow S$ be the minimal resolution of S and $\pi : \hat{S} \rightarrow X$ is the induced morphism. Let $\hat{C}$ be the strict transform of C in $\hat{S}$. Then as in the proof of Lemma 10.9 we have $K_{\hat{S}} \cdot \hat{C} < -d \leq -3$ in both cases. The rest of proof follows from a similar argument as in the proof of [HP16, Lemma 5.7]. $\square$

We can now deduce Theorem 10.6.

Proof of Theorem 10.6. By Corollary 5.3, there is a bimeromorphic morphism $f : (X', \Delta') \rightarrow (X, \Delta)$ such that

- (1) X' has $\mathbb{Q}$ -factorial terminal singularities, and
- (2) (X', Δ') is a dlt pair such that

$$K_{X'} + \Delta' \sim_{\mathbb{Q}} f^*(K_X + \Delta)$$

By Lemma 10.10, $K_{X'} + \Delta'$ has the bend-and-break property. Hence by [CHP16, Lemma 4.2], $K_X + \Delta$ also has the bend-and-break property. $\square$

10.3. Cone theorem for pseudo-effective DLT pairs. We recall that the cone $\overline{\text{NA}}(X)$ for a normal compact Kähler variety is the closed cone generated by positive closed current of bidimension $(1, 1)$. It replaces the Mori cone $\overline{\text{NE}}(X)$ in the non-algebraic setting, and is dual to the closure of the cone generated by the classes of Kähler forms (see [HP16, Section 1] for details).

Theorem 10.11. *With notations and hypothesis as in Assumption 10.1, there is a rational number $d > 0$ and an at most countable set of curves $\{\Gamma_i\}_{i \in I}$ such that*

$$0 < (K_X + \Delta) \cdot \Gamma_i \leq d$$

and that

$$\overline{\text{NA}}(X) = \overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

Proof. The proof works with almost verbatim argument as in the proof of [CHP16, Theorem 4.1, page 985]. We note that in [CHP16, Theorem 4.1] K_X is assumed to be pseudo-effective, which we don't have, but in the proof of [CHP16, Theorem 4.1, page 985] only $K_X + \Delta$ being pseudo-effective is used, which we have in our case. Moreover, we replace the use of [CHP16, Lem. 4.1, Pro. 4.1] by Lemma 10.2 and Theorem 10.6, respectively. $\square$

Together with [CHP16, Theorem 1.3], we obtain the following cone theorem for dlt pair (X, Δ) with pseudo-effective $K_X + \Delta$.

Theorem 10.12. *Let (X, Δ) be a compact Kähler $\mathbb{Q}$ -factorial 3-fold dlt pair. Assume that $K_X + \Delta$ is pseudo-effective. Then there is a rational number $d > 0$ and a countable set of curves $\{\Gamma_i\}_{i \in I}$ such that*

$$0 < (K_X + \Delta) \cdot \Gamma_i \leq d$$

and

$$\overline{\text{NA}}(X) = \overline{\text{NA}}(X)_{(K_X + \Delta) \geq 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

10.4. Contraction theorems.

We will study the contraction of $(K_X + \Delta)$ -negative extremal rays of $\overline{\text{NA}}(X)$ in this subsection.

Definition 10.13. [HP16, Def. 7.1, Rmk. 7.2] Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair such that $K_X + \Delta$ is pseudo-effective. Let R be a $(K_X + \Delta)$ -negative extremal ray of $\overline{\text{NA}}(X)$. We say that R is small if every curve $C \subseteq X$ with $[C] \in R$ is very rigid in the sense of [HP16, Definition 4.3]. Otherwise we say R is of divisorial-type (See [HP16, Remark 7.2]).

10.4.1. Contraction of small rays.

Theorem 10.14. *With notations and hypothesis as in Assumption 10.1, let R is a $(K_X + \Delta)$ -negative extremal ray in $\overline{\text{NA}}(X)$. If R is small, then the contraction $c_R : X \rightarrow Y$ of R exists.*

Proof. First of all, since R is a $(K_X + \Delta)$ -negative extremal ray, there is a supporting nef and big $(1, 1)$ class α on X such that $R = \{\gamma \in \overline{\text{NA}}(X) : \alpha \cdot \gamma = 0\}$; note that the existence of α follows similarly as in the proof of [HP16, Proposition 7.3] since it is a formal proof. Also, the bigness of α follows from [CHP16, Eqn. (10), page 987]. Next we observe that the contraction of R follows exactly as in the proof of [CHP16, Theorem 4.2] provided we can show that $\alpha^2 \cdot S > 0$ holds for every irreducible surface $S \subseteq X$ ([CHP16, Pro. 4.4]). Finally, this result holds by the following proposition. $\square$

Proposition 10.15. *With notations and hypothesis as in Theorem 10.14, if α is a nef supporting class of R , then $\alpha^2 \cdot S > 0$ for any surface $S \subseteq X$.*

Proof. We follow a similar idea as in the proof of [CHP16, Proposition 4.4]. By scaling α if necessary we may assume that $\omega = \alpha - (K_X + \Delta)$ is a Kähler

class. By contradiction assume that $\alpha^2 \cdot S = 0$ for some surface $S \subseteq X$. Now if $\alpha|_S = 0$, then $-(K_X + \Delta)|_S$ is ample. In particular, S is projective and covered by uncountably many curves. Since $\alpha|_S = 0$, the classes of these curves in X are contained in R . This contradicts that R is small.

Next assume that $\alpha|_S \neq 0$ and $\alpha^2 \cdot S = 0$. Then we have

$$(10.2) \quad (K_X + \Delta) \cdot \alpha \cdot S = (\alpha - \omega) \cdot \alpha \cdot S = -\omega \cdot \alpha \cdot S < 0,$$

as $\alpha|_S$ is nef and non-zero. Therefore $(K_X + \Delta)|_S$ is not pseudo-effective. Now let $\hat{S} \rightarrow S$ be the minimal resolution of S and $\pi : \hat{S} \rightarrow X$ is the induced morphism. Then as in the proof of [CHP16, Proposition 4.4] we see that in order to conclude, it is enough to find an effective $\mathbb{Q}$ -divisor E on $\hat{S}$ such that

$$(K_{\hat{S}} + E) \cdot \pi^*(\alpha) < 0.$$

To this end we will consider two cases related to (10.2).

Case I: Assume that $\Delta \cdot \alpha \cdot S < 0$. Since α is nef, this implies that S is a component of Δ . Since Δ is a boundary divisor, there is a rational number $\mu \geq 0$ such that the multiplicity of $(1 + \mu)\Delta$ along S is 1. Then

$$(K_X + (1 + \mu)\Delta) \cdot \alpha \cdot S \leq (K_X + \Delta) \cdot \alpha \cdot S < 0.$$

By adjunction there is an effective $\mathbb{Q}$ -divisor E such that

$$(K_{\hat{S}} + E) \cdot \pi^*\alpha = \pi^*(K_X + (1 + \mu)\Delta) \cdot \pi^*\alpha = (K_X + (1 + \mu)\Delta) \cdot \alpha \cdot S < 0.$$

Case II: Assume that $\Delta \cdot \alpha \cdot S \geq 0$. Then we have

$$K_X \cdot \alpha \cdot S \leq (K_X + \Delta) \cdot \alpha \cdot S < 0.$$

Since $(K_X + \Delta) \cdot \alpha \cdot S < 0$, the surface $S = S_j$ must be a component in the decomposition (10.1), and we must have

$$S \cdot \alpha \cdot S < 0.$$

Therefore $(K_X + S) \cdot \alpha \cdot S < 0$. Hence by adjunction, there is an effective $\mathbb{Q}$ -divisor E on $\hat{S}$ such that

$$(K_{\hat{S}} + E) \cdot \pi^*\alpha = \pi^*(K_X + S) \cdot \pi^*\alpha = (K_X + S) \cdot \alpha \cdot S < 0.$$

□

10.4.2. *Contraction of divisorial rays.*

With notations and hypothesis as in Assumption 10.1, let R is a $(K_X + \Delta)$ -negative extremal ray of $\overline{NA}(X)$ of divisorial type. Then there is a unique surface $S \subseteq X$ such that S contains all curves $C \subseteq X$ such that $[C] \in R$ and also that S is covered by these curves. Let $\nu : \tilde{S} \rightarrow S$ be the normalization of S . Let α be a nef supporting class of R . Then the nef reduction of $\nu^*(\alpha|_S)$ gives a fibration

$$\tilde{f} : \tilde{S} \rightarrow \tilde{T}.$$

We define

$$n(\alpha) = \dim \tilde{T}.$$

Since $\alpha|_S$ intersects a covering of curves in S trivially, it follows that $n(\alpha) \in \{0, 1\}$.

Theorem 10.16. *With notations and hypothesis as in Assumption 10.1, let R is a $(K_X + \Delta)$ -negative extremal ray of $\overline{NA}(X)$ of divisorial type. Let α be a nef supporting class of R . If $n(\alpha) = 0$, then the contraction $c_R : X \rightarrow Y$ of R exists.*

Proof. The same proof as in [HP16, Corollary 7.7] works here using the Cone Theorem 10.12. $\square$

In case of $n(\alpha) = 1$, we have the following result.

Proposition 10.17. *With notations and hypothesis as in Assumption 10.1, let R is a $(K_X + \Delta)$ -negative extremal ray of $\overline{NA}(X)$ of divisorial type. Let $S \subseteq X$ be the unique surface which contains all curves $C \subseteq X$ such that $[C] \in R$ and also that S is covered by these curves. Let α be a nef supporting class of R and assume that $n(\alpha) = 1$. Then the contraction $c_R : X \rightarrow Y$ of R exists if one of the following conditions holds:*

- (1) S has slc singularities, or
- (2) X is terminal and $K_X \cdot R < 0$.

Proof. If the first condition holds, then the same proof as in [CHP16, Proposition 4.5] works without any change here. In the second case R is a K_X -negative extremal ray with X having terminal singularities, so it follows similarly as in the proof of [HP16, Lem. 7.8, Cor. 7.9]. $\square$

Part 3. Log abundance theorem for log canonical pairs

11. REDUCTION STEP

In this section we will prove reduction theorem which will split the abundance problem for klt pairs into two cases.

Definition 11.1. Let $\phi : X \dashrightarrow X'$ be a bimeromorphic map between two normal $\mathbb{Q}$ -factorial varieties and D is an $\mathbb{R}$ -Cartier divisor on X . Then ϕ is called a D -crepant if there is a normal variety W and morphisms $p : W \rightarrow X$ and $q : W \rightarrow X'$ resolving ϕ , i.e., $\phi \circ p = q$, such that $p^*D = q^*(\phi_*D)$ holds.

Theorem 11.2. *Let (X, Δ) be a klt (resp. lc) pair such that X is a $\mathbb{Q}$ -factorial compact Kähler 3-fold with terminal singularities. Assume that $K_X + \Delta$ is nef. Then there is a $(K_X + \Delta)$ -crepant bimeromorphic map $\phi : (X, \Delta) \dashrightarrow (X', \Delta')$ such that exactly one of the following two holds:*

- (1) *There is a $K_{X'}$ -Mori fiber space $X' \rightarrow Y$ such that $K_{X'} + \Delta'$ is numerically trivial over Y .*
- (2) *$K_{X'} + (1 - \epsilon)\Delta'$ is nef for all $0 \leq \epsilon \ll 1$.*

Moreover, X' has $\mathbb{Q}$ -factorial terminal singularities and (X', Δ') is klt (resp. lc).

We need the following lemma to prove this theorem.

Lemma 11.3. *Let (X, Δ) be a klt (resp. lc) pair such that X is a $\mathbb{Q}$ -factorial terminal compact Kähler 3-fold. Assume that X is uniruled and non-algebraic. Further assume that $K_X + \Delta$ is nef and it intersects the general fibers of the MRC fibration of X trivially. Then there is a $(K_X + \Delta)$ -crepant bimeromorphic model $\phi : (X, \Delta) \dashrightarrow (X', \Delta')$ such that*

- (1) $\Delta' := \phi_*\Delta$,
- (2) X' has $\mathbb{Q}$ -factorial terminal singularities,
- (3) (X', Δ') is klt (resp. lc),
- (4) there is a $K_{X'}$ -Mori fiber space $X' \rightarrow Y'$, and
- (5) $(K_{X'} + \Delta') \sim_{\mathbb{Q}, Y'} 0$.

Proof. First of all, since X is uniruled and non-algebraic, from Lemma 3.10 it follows that the base of the MRC fibration of X has dimension 2. We recall that a Kähler class ω on X is called normalized if $K_X + \omega$ intersects the general fibers of the MRC fibration of X trivially (see [HP15, Definition 1.2]). Set $D := K_X + \Delta$; then we claim that one of the following holds:

- (1) either there is a $(K_X + \omega)$ -negative extremal ray R such that $D \cdot R = 0$,
- (2) or there exist a smallest non-negative real number $\lambda \geq 0$ such that $K_X + \omega + \lambda D$ is nef.

Indeed, if (1) does not hold, then either $K_X + \omega$ is nef or every $(K_X + \omega)$ -negative extremal ray R satisfies $D \cdot R > 0$. In the first case we set $\lambda = 0$, and in the second case the existence of smallest λ can be proved as in Proposition 4.3 using the boundedness of the length of extremal rays from [HP15, Theorem 3.6].

Now if we are in case (1), then the K_X -negative extremal ray R can be contracted as in [HP15, Lem. 3.21, Thm. 3.24] by a projective morphism; Note that D is crepant with respect to this morphism. By induction and [HP15], we can run a K_X -MMP, trivial with respect to D , and obtain a D -crepant bimeromorphic map $\psi : (X, \Delta) \dashrightarrow (\hat{X}, \hat{\Delta})$ such that for any normalized Kähler class $\hat{\omega}$ on $\hat{X}$ there exist a smallest non-negative real number λ such that $K_{\hat{X}} + \hat{\omega} + \lambda \hat{D}$ is nef, where $\hat{D} = K_{\hat{X}} + \hat{\Delta} = \phi_* D = \phi_*(K_X + \Delta)$ is nef. Moreover, $\hat{X}$ has $\mathbb{Q}$ -factorial terminal singularities and $\hat{D}$ intersects the general fibers of the MRC fibration of $\hat{X}$ trivially. Therefore $\hat{\omega} + (K_{\hat{X}} + \hat{\omega} + 2\lambda \hat{D})$ is then a normalized Kähler class on $\hat{X}$. Then

$$K_{\hat{X}} + (\hat{\omega} + K_{\hat{X}} + \hat{\omega} + 2\lambda \hat{D}) = 2(K_{\hat{X}} + \hat{\omega} + \lambda \hat{D}) \quad \text{is nef.}$$

If we are in case (2) above, then $K_X + \omega + \lambda D$ is nef for the smallest $\lambda \in \mathbb{R}^{\geq 0}$ and $\omega + \lambda D$ is a normalized Kähler class, since $D = K_X + \Delta$ intersects the general fibers of the MRC fibration of X trivially. Renaming X, ω and D by $\hat{X}, \hat{\omega}$ and $\hat{D} = K_{\hat{X}} + \hat{\Delta}$ we see that, in either case above, by [HP15, Theorem 1.4] there is a projective Fano fibration $\hat{f} : \hat{X} \rightarrow \hat{Y}$ onto a surface $\hat{Y}$.

We remark that $\hat{f}$ is indeed the MRC fibration, as the base of the MRC fibration of X , and hence that of $\hat{X}$, has dimension 2. Note that $\hat{D}$ is nef on $\hat{X}$ and $\hat{D} \cdot F = (K_{\hat{X}} + \hat{\Delta}) \cdot F = 0$ for general fibers of $F(\cong \mathbb{P}^1)$ of $\hat{f}$. Now since $\hat{f}$ is a projective morphism, by Proposition 3.8 and Section 4.2 we run a relative $K_{\hat{X}}$ -MMP over $\hat{Y}$ with the scaling of $\hat{\Delta}$. This MMP terminates with a Mori fiber space, say $f' : X' \rightarrow Y'$ over Y such that X' has $\mathbb{Q}$ -factorial terminal singularities. By Lemma 4.6 every step of this MMP is $\hat{D}$ -trivial. In particular, the induced bimeromorphic map $\phi : \hat{X} \dashrightarrow X'$ is $\hat{D}$ -crepant. Let $\Delta' := \phi_* \hat{\Delta}$; then $(K_{X'} + \Delta') \sim_{\mathbb{Q}, f'} 0$. The composite of ϕ and ψ gives the required map. $\square$

Now we conclude Theorem 11.2.

Proof of Theorem 11.2. If X is projective, then this is just [KMM94, Lemma 5.3 and Lemma 7.1]. Assume that X is non-algebraic and uniruled. If (X, Δ) is lc, by taking a dlt model as in Corollary 5.3, we may assume that (X, Δ) is a $\mathbb{Q}$ -factorial dlt pair such that X has terminal singularities. Let λ be the positive rational number such that $K_X + \lambda \Delta$ has zero intersection with general

fibers of the MRC fibration of X . If $\lambda = 1$, then we can apply Lemma 11.3. So assume from now on that $\lambda < 1$. Then $(X, \lambda\Delta)$ is klt. By Theorem 9.1, $K_X + \lambda\Delta$ is $\mathbb{Q}$ -effective. If $K_X + \lambda\Delta$ is nef, then we can take $(X', \Delta') = (X, \Delta)$ and we are in the second situation. Assume that $K_X + \lambda\Delta$ is not nef. We will show that there exist a terminating $(K_X + \lambda\Delta)$ -MMP, which is trivial with respect to $K_X + \Delta$.

We first assume that there is a $(K_X + \lambda\Delta)$ -negative extremal ray R of $\overline{\text{NA}}(X)$ which is $(K_X + \Delta)$ -trivial. Then R is also K_X -negative. Hence the contraction of R exists by Theorem 10.16 and Proposition 10.17, if R is divisorial, and by Theorem 10.14, if R is small. Moreover, if R is small, then the corresponding flip exists by [CHP16, Theorem 4.19]. Let $h_1 : (X, \Delta) \dashrightarrow (X_1, \Delta_1)$ be the divisorial contraction or the flip corresponding to the contraction of R . Then h_1 is $(K_X + \Delta)$ -crepant and X_1 has terminal singularities. By induction, we can run a $(K_X + \lambda\Delta)$ -MMP, trivial with respect to $K_X + \Delta$. By Theorem 3.3 this MMP terminates. Therefore we obtain a $(K_X + \Delta)$ -crepant bimeromorphic map $\phi : (X, \Delta) \dashrightarrow (X', \Delta')$ such that $K_{X'} + \Delta'$ is nef and X' has $\mathbb{Q}$ -factorial terminal singularities, and

- (i) either $K_{X'} + \lambda\Delta'$ is nef,
- (ii) or every $(K_{X'} + \lambda\Delta')$ -negative extremal ray is $(K_{X'} + \Delta')$ -positive.

In the first case, $K_{X'} + (1 - \epsilon)\Delta'$ is nef for all $0 \leq \epsilon \ll 1$. In the second case, as in the proof of Proposition 4.3 using the boundedness of the lengths of extremal rays of Theorem 10.11 we see that there is a smallest positive rational number $\mu \in \mathbb{Q}^+$ such that $(K_{X'} + \Delta') + \mu(K_{X'} + \lambda\Delta')$ is nef. This shows that $K_{X'} + (1 - \epsilon)\Delta'$ is nef for all $0 \leq \epsilon \ll 1$. Since this MMP is $(K_X + \Delta)$ -trivial, we obtain that (X', Δ') is also klt (respectively lc).

Finally we assume that X is not uniruled. Then we can run a K_X -MMP, trivial with respect to $K_X + \Delta$, by [HP16, Theorem 1.1 and Theorem 1.2] and the rest of the arguments work as above. $\square$

If we are in the first case of Theorem 11.2, then we can apply log abundance for surfaces to the base Y to conclude the log abundance. If we are in the second case, we will need the following theorem.

12. CONSTRUCTION OF BIMEROMORPHIC MODELS

In this section we will construct several bimeromorphic models which will help us in the next section to show that $\nu(X, K_X + \Delta) = 1$ implies $k(X, K_X + \Delta) > 0$.

Lemma 12.1. *Let (X, Δ) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold klt pair. Assume that $K_X + (1 - \epsilon)\Delta$ is nef for all $0 \leq \epsilon \ll 1$. Then there is a boundary Δ' in X such that*

- (1) (X, Δ') is klt,
- (2) $K_X + (1 - \epsilon)\Delta'$ is nef for all $0 \leq \epsilon \ll 1$,
- (3) there is some $D \in |K_X + \Delta'|_{\mathbb{Q}}$ such that $\text{Supp } D = \text{Supp } \Delta'$, and
- (4) $\nu(K_X + \Delta) = \nu(K_X + \Delta')$ and $\kappa(K_X + \Delta) = \kappa(K_X + \Delta')$.

Proof. Since $K_X + \lambda\Delta$ is nef for some $\lambda < 1$, by Theorem 9.3 there is an effective $\mathbb{Q}$ -divisor $P \sim_{\mathbb{Q}} (K_X + \lambda\Delta)$. Set $Q := P + (1 - \lambda)\Delta$; then $Q \sim_{\mathbb{Q}} K_X + \Delta$ is nef and its support contains that of Δ . Choose a rational number $0 < \gamma \ll 1$ such that $(X, \Delta + \gamma Q)$ is klt. Set $\Delta' = \Delta + \gamma Q$; then

$$K_X + (1 - \epsilon)\Delta' = K_X + (1 - \epsilon)\Delta + (1 - \epsilon)\gamma Q$$

is nef for all $0 \leq \epsilon \ll 1$. Moreover, we note that

$$K_X + \Delta' \sim_{\mathbb{Q}} (1 + \gamma)(K_X + \Delta).$$

This proves property (4). Let $D = \frac{1}{1+\gamma}Q$, then $D \in |K_X + \Delta'|_{\mathbb{Q}}$ and its support is the same as that of Δ' . $\square$

Lemma 12.2. *Let (X, B) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold dlt pair. Assume that B is reduced, and that there is an element $D \in |K_X + B|_{\mathbb{Q}}$ such that $\text{Supp } D = B$. Then there is a $(K_X + B)$ -MMP which terminates to a dlt pair (X', B') such that $K_{X'} + B'$ is nef.*

Proof. If X is projective, then we can apply the classical algebraic MMP. If K_X is pseudo-effective, then we can apply [CHP16, Theorem 4.6]. It remains to deal with the case when X is uniruled and non-algebraic.

Assume that $(K_X + B)$ is not nef. We will show that a $(K_X + B)$ -MMP exists. By Theorem 10.12, there is a $(K_X + B)$ -negative extremal ray R . If R is small, then there exist a small contraction $c_R : X \rightarrow Y$ by Theorem 10.14. The flip of c_R exists by Theorem 3.1.

Assume that R is divisorial. Let S be the unique surface containing and covered by all the curves $C \subseteq X$ such that $[C] \in R$. Then S is a component of B , since $B = \text{Supp } D$ and $D \cdot R < 0$. Thus S is normal (being a dlt center) and by adjunction $(S, 0)$ has klt singularities. Hence the contraction exists by Theorem 10.16 and Proposition 10.17. Termination of the $(K_X + B)$ -MMP follows from Theorem 3.3. $\square$

Proposition 12.3. *Let (X, Δ) be a klt pair, where X is a $\mathbb{Q}$ -factorial compact Kähler 3-fold with terminal singularities. Suppose that there is a nef $\mathbb{Q}$ -divisor $D \in |K_X + \Delta|_{\mathbb{Q}}$ such that $\text{Supp } \Delta = \text{Supp } D$. Then there is a $\mathbb{Q}$ -factorial pair $(\hat{X}, \hat{B})$ such that*

- (1) $(\hat{X}, \hat{B})$ is dlt,
- (2) $\hat{B}$ is reduced,

- (3) $\hat{X} \setminus \hat{B}$ has terminal singularities,
- (4) $K_{\hat{X}} + \hat{B}$ is nef,
- (5) there is some $\hat{D} \in |K_{\hat{X}} + \hat{B}|_{\mathbb{Q}}$ such that $\text{Supp } \hat{D} = \hat{B}$, and
- (6) $\nu(K_X + \Delta) = \nu(K_{\hat{X}} + \hat{B})$ and $\kappa(K_X + \Delta) = \kappa(K_{\hat{X}} + \hat{B})$.

Proof. Since X has isolated (terminal) singularities, there is a partial log resolution $r : \tilde{X} \rightarrow X$ of (X, Δ) such that

- r is an isomorphism over $X \setminus \text{Supp } \Delta$,
- $\tilde{X}$ is smooth in a neighborhood of $r^*\Delta$ and $r^*\Delta$ has simple normal crossing support.

Since (X, Δ) is klt, we can write

$$K_{\tilde{X}} + B' = r^*(K_X + \Delta) + E \sim_{\mathbb{Q}} r^*D + E,$$

where $B' = r_*^{-1}\Delta + \text{Ex}(r)$, and $E \geq 0$ is an effective $\mathbb{Q}$ -divisor supported on the exceptional locus of r .

Set $\tilde{B} := \lceil B' \rceil = \lceil r_*^{-1}\Delta \rceil + \text{Ex}(r)$ and $\tilde{D} := r^*D + E + (\lceil r_*^{-1}\Delta \rceil - r_*^{-1}\Delta) \geq 0$. Then $\tilde{D} \in |K_{\tilde{X}} + \tilde{B}|_{\mathbb{Q}}$, $\text{Supp } \tilde{D} = \text{Supp}(r^*D + \text{Ex}(r)) = \tilde{B}$, since $\text{Supp } D = \text{Supp } \Delta$. Moreover, from the construction it also follows that $\tilde{X} \setminus \tilde{B}$ has terminal singularity and $(\tilde{X}, \tilde{B})$ is dlt. We claim that $\tilde{X}$ is $\mathbb{Q}$ -factorial. Let F be a prime Weil divisor on $\tilde{X}$. If F is r -exceptional, then by construction, $\tilde{X}$ is smooth in a neighborhood of F and thus F is Cartier. If F is not r -exceptional, then r_*F is a Weil divisor on X and hence it is $\mathbb{Q}$ -Cartier, since X is $\mathbb{Q}$ -factorial. Then $r^*r_*F = F + G$, where G is a r -exceptional divisor. Since we have just showed that G is $\mathbb{Q}$ -Cartier, it follows that F is $\mathbb{Q}$ -Cartier. Finally, it is easy to see that $\omega_{\tilde{X}}^{[k]}$ is a line bundle for some $k \geq 1$, since the same is true for X and the exceptional divisor of r is $\mathbb{Q}$ -Cartier.

Now by Lemma 12.2 we can run a terminating $(K_{\tilde{X}} + \tilde{B})$ -MMP to obtain a $\mathbb{Q}$ -factorial dlt pair $(\hat{X}, \hat{B})$ such that $K_{\hat{X}} + \hat{B}$ is nef. Let $\hat{D}$ be the birational transform of $\tilde{D}$ in $\hat{X}$. Then $\hat{D} \in |K_{\hat{X}} + \hat{B}|_{\mathbb{Q}}$ and $\text{Supp } \hat{D} = \hat{B}$. Moreover, since $0 \leq \tilde{D} \sim_{\mathbb{Q}} (K_{\tilde{X}} + \tilde{B})$ and $\text{Supp } \tilde{D} = \text{Supp } \tilde{B}$, all the curves and divisors contracted by the MMP is contained in the $\text{Supp } \tilde{D} = \tilde{B}$. In particular, we have

$$\tilde{X} \setminus \text{Supp } \tilde{B} \cong \hat{X} \setminus \text{Supp } \hat{B},$$

and thus $\hat{X} \setminus \hat{B}$ has terminal singularities.

Now we will show (6). Note that since MMP preserves Kodaira dimension, we have

$$\kappa(K_{\hat{X}} + \hat{B}) = \kappa(K_{\tilde{X}} + \tilde{B}) = \kappa(\tilde{D}).$$

On the other hand, since $\text{Supp } \Delta = \text{Supp } D$, there is a positive number m such that

$$\tilde{D} = r^*D + E + (\lceil r_*^{-1}\Delta \rceil - r_*^{-1}\Delta) \leq r^*D + E + mr^*D.$$

Now since E is a r -exceptional effective divisor, we have

$$\kappa(D) = \kappa(r^*D) \leq \kappa(\tilde{D}) \leq \kappa((m+1)r^*D) = \kappa(D).$$

Therefore $\kappa(K_X + \Delta) = \kappa(D) = \kappa(\tilde{D}) = \kappa(K_{\hat{X}} + \hat{B})$.

For the equality of numerical dimensions, we take a desingularization W of the graph of $\phi : \tilde{X} \dashrightarrow \hat{X}$, with induced morphisms $p : W \rightarrow \tilde{X}$ and $q : W \rightarrow \hat{X}$. Since every step of the MMP $\phi : \tilde{X} \dashrightarrow \hat{X}$ is $\tilde{D}$ -negative and $\hat{D} = \phi_*\tilde{D}$, we obtain that $p^*\tilde{D}$ and $q^*\hat{D}$ have same support. In particular, $p^*(r^*D)$ and $q^*\hat{D}$ have same support. Then the effective divisors

$$(r \circ p)^*D \sim_{\mathbb{Q}} p^*(K_X + \Delta) \text{ and } q^*\hat{D} \sim_{\mathbb{Q}} q^*(K_{\hat{X}} + \hat{B})$$

are both nef and have same support.

Therefore by [Kol92, Lemma 11.3.3] (using a Kähler class in place of an ample divisor), we obtain that $\nu(K_X + \Delta) = \nu(K_{\hat{X}} + \hat{B})$. $\square$

The following lemma is an analogue of [CHP16, Lemma 6.5].

Lemma 12.4. *Let (X, B) be a reduced dlt pair such that X is a $\mathbb{Q}$ -factorial compact Kähler threefold. Assume that*

- *there is a nef divisor $D \in |K_X + B|_{\mathbb{Q}}$ such that $\text{Supp } D = B$,*
- *$X \setminus B$ has terminal singularities.*

Choose a rational number $\lambda \in (0, 1)$ such that $D - (1 - \lambda)B$ is effective with support equal to B . Let S be a reduced Weil divisor contained in the support of B and set $Z := B - S$. Assume that $(K_X + B)|_T \equiv 0$ for every component T of Z and that $S \cap Z \neq \emptyset$. Set $X_0 := X$, $S_0 := S$, $Z_0 := Z$ and $B_0 := B$. Then there is a finite sequence of $(K_X + \lambda S + Z)$ -flips and divisorial contractions

$$\{f_i : X_i \dashrightarrow X_{i+1}\}_{i=0, \dots, n}$$

such that for each $i \geq 0$ we have

- (1) D_{i+1} , S_{i+1} , Z_{i+1} and B_{i+1} are bimeromorphic transform of D , S , Z and B onto X_{i+1} ,
- (2) f_i is $(K_{X_i} + B_i)$ -crepant, i.e., for any resolution $p_i : W \rightarrow X_i$, $q_i : W \rightarrow X_{i+1}$ of the graph of f_i , $p_i^*(K_{X_i} + B_i) = q_i^*(K_{X_{i+1}} + B_{i+1})$,
- (3) $K_{X_{i+1}} + B_{i+1}$ has the same numerical dimension and Kodaira dimension as $K_X + B$,
- (4) $D_{i+1} \in |K_{X_{i+1}} + B_{i+1}|_{\mathbb{Q}}$ and $\text{Supp } D_{i+1} = B_{i+1}$,
- (5) $D_{i+1} - (1 - \lambda)B_{i+1}$ is effective with support equal to B_{i+1} ,

- (6) $(X_{i+1}, \lambda S_{i+1} + Z_{i+1})$ is $\mathbb{Q}$ -factorial dlt and $(K_{X_{i+1}}, B_{i+1})$ is lc,
- (7) $(K_{X_{i+1}} + B_{i+1})|_{T_{i+1}} \equiv 0$ for every component T_{i+1} of Z_{i+1} ,
- (8) $X_{i+1} \setminus B_{i+1} \cong X \setminus B$, and
- (9) $S_{n+1} \neq 0$ and $Z_{n+1} \cap S_{n+1} = \emptyset$.

Proof. Proceeding by induction assume that we have already constructed $f_{i-1} : X_{i-1} \rightarrow X_i$ satisfying the properties (3) – (8). If $S_i \cap Z_i = \emptyset$ then we stop and set $i = n + 1$. So assume that $S_i \cap Z_i \neq \emptyset$. Then there is some component T_i of Z_i such that $T_i \cap S_i \neq \emptyset$. Then we have that $(K_{X_i} + \lambda S_i + Z_i)|_{T_i} \equiv -(1 - \lambda)S_i|_{T_i}$ is not pseudo-effective. We claim that T_i is a Moishezon surface. Indeed, if X is not uniruled, then this follows from [CHP16, Lemma 4.1], and otherwise from Lemma 10.2. Therefore there is a curve $\Gamma_i \subseteq T_i$ such that $(K_{X_i} + \lambda B_i + Z_i) \cdot \Gamma_i < 0$. Then from (7) (with $i+1$ replaced by i) it follows that $(K_{X_i} + B_i) \cdot \Gamma_i = 0$. Now by the cone theorem with respect to $K_{X_i} + \lambda S_i + Z_i$ (see Theorem 10.12), we can decompose the class $[\Gamma_i] \in \overline{NA}(X_i)$ as

$$[\Gamma_i] = M_i + \sum_{j=1}^{m_i} [C_{ij}],$$

where M_i is $(K_{X_i} + \lambda S_i + Z_i)$ -nonnegative and C_{ij} generate $(K_{X_i} + \lambda S_i + Z_i)$ -negative extremal rays. Since $K_{X_i} + B_i$ is nef and $(K_{X_i} + B_i) \cdot \Gamma_i = 0$, we obtain that $(K_{X_i} + B_i) \cdot C_{ij} = 0$ for all j . Therefore there is a $(K_{X_i} + \lambda S_i + Z_i)$ -negative extremal ray R_i which is $(K_{X_i} + B_i)$ -trivial.

If R_i is small, then the contraction of R_i exists by Theorem 10.14, and the corresponding flip exists by Theorem 3.1. If R_i is divisorial, then there is a unique surface V_i which contains and covered by all the curves in R_i . This surface V_i must be a component of B_i as $K_{X_i} + \lambda B_i + Z_i \sim_{\mathbb{Q}} D_i - (1 - \lambda)B_i \geq 0$, and the right hand side is supported on B_i . Now since $(X_i, \lambda_i S_i + Z_i)$ is a $\mathbb{Q}$ -factorial dlt pair, X_i has $\mathbb{Q}$ -factorial klt singularity. Since (X_i, B_i) is lc, by adjunction $(V_i, 0)$ is slc (see [Fuj00, Remark 1.2(5)]). Hence the contraction of V_i exists by Theorem 10.16 and Proposition 10.17. Let $f_i : X_i \dashrightarrow X_{i+1}$ be the corresponding divisorial contraction or flip. Since R_i is $(K_{X_i} + B_i)$ -trivial, f_i is $(K_{X_i} + B_i)$ -crepant. Then by [CHP16, Lemma 3.2] we also get the properties (3)–(7) at the $(i + 1)$ -th level. Since the exceptional locus of f_i is contained in B_i , we also have $X_{i+1} \setminus B_{i+1} \cong X_i \setminus B_i \cong X \setminus B$; this is (8).

Next we note that, since $(X, \lambda S + Z)$ is a dlt pair, this MMP terminates by Theorem 3.3. At the end of this MMP by our hypothesis we have $Z_{n+1} \cap S_{n+1} = \emptyset$. So it remains to show that $S_{n+1} \neq 0$. To the contrary assume that $S_{n+1} = 0$; then all the components of S are contracted by the steps of this MMP. Let $f_i : X_i \rightarrow X_{i+1}$ be one of these steps which contracts the last component of S , i.e. $S_i \subseteq X_i$ is irreducible and contracted by f_i . We know that the

corresponding extremal ray R_i is $(K_{X_i} + B_i)$ -trivial and $(K_{X_i} + \lambda S_i + Z_i)$ -negative. Thus $S_i \cdot R_i > 0$, this is a contradiction. $\square$

13. $\nu = 1$ IMPLIES $\kappa > 0$

In this section we will prove following Theorem.

Theorem 13.1. *Let (X, Δ) be a klt pair such that X is a $\mathbb{Q}$ -factorial compact Kähler 3-fold with terminal singularities. Assume that $K_X + (1 - \epsilon)\Delta$ is nef for all $0 \leq \epsilon \ll 1$. If $\nu(K_X + \Delta) = 1$, then $\kappa(K_X + \Delta) = 1$.*

We need the following reduction result first.

Lemma 13.2. *Let (X, B) be a reduced dlt pair such that X is a $\mathbb{Q}$ -factorial compact Kähler 3-fold with klt singularities. Assume that there is a nef $\mathbb{Q}$ -divisor $D \in |K_X + B|_{\mathbb{Q}}$ such that $\text{Supp } D = B$ and that $\nu(K_X + B) = 1$. Moreover, assume that $X \setminus B$ has terminal singularities. Then there is a $(K_X + B)$ -crepant bimeromorphic map $\phi : (X', B') \dashrightarrow (X, B)$ such that*

- (1) (X', B') is a $\mathbb{Q}$ -factorial reduced lc pair, where $B' := \phi_* B$,
- (2) $(X', 0)$ is klt and $X' \setminus B'$ has terminal singularities,
- (3) there is a nef $\mathbb{Q}$ -divisor $D' \in |K_{X'} + B'|_{\mathbb{Q}}$ such that $\text{Supp } D' = B'$,
- (4) $\nu(K_{X'} + B') = 1$ and $\kappa(K_X + B) = \kappa(K_{X'} + B')$, and
- (5) there is a connected component S' of B' which is irreducible.

Proof. Since $\nu(K_X + B) = 1$, we have $(K_X + B) \cdot D = (K_X + B)^2 \equiv 0$. Since $K_X + B$ is nef, this implies that $(K_X + B)|_T \equiv 0$ for every component T of B . Let S be an irreducible component of B and $Z := B - S$. Then applying Lemma 12.4 gives us the required bimeromorphic map $\phi : (X, B) \dashrightarrow (X', B')$. $\square$

Proof of Theorem 13.1. We will follow the same strategy as in [CHP16, §8.A] which is same as in [Kol92, Chapter 13]. First replacing $(X, 0)$ by a terminal model as in Theorem 3.6 we may assume that X has terminal singularities. Then combining 12.1, Proposition 12.3 and Lemma 13.2 we reduce our problem to proving the following statement:

Let (X, B) be a $\mathbb{Q}$ -factorial compact Kähler 3-fold reduced lc pair and $D \in |K_X + B|$ is a $\mathbb{Q}$ -divisor satisfying the following properties:

- (1) $\text{Supp } D = B$,
- (2) $(X, 0)$ is klt and $X \setminus B$ has terminal singularities,
- (3) $K_X + B$ is nef and $\nu(K_X + B) = 1$, and
- (4) there is a connected component S of B such that S is irreducible.

Then $\kappa(K_X + B) \geq 1$.

The rest of the proof works exactly the same way as in the proof of [CHP16, Theorem 8.1]. $\square$

14. PROOF OF LOG ABUNDANCE FOR LOG CANONICAL PAIRS

We first prove the klt log abundance for $\nu(X, K_X + \Delta) = 1$.

Theorem 14.1. *Let (X, Δ) be a klt pair and $K_X + \Delta$ is nef. If the numerical dimension $\nu(X, K_X + \Delta) = 1$, then $K_X + \Delta$ is semi-ample.*

Proof. First replacing (X, Δ) by its terminal model as in Theorem 3.6 we may assume that X has $\mathbb{Q}$ -factorial terminal singularities. Now by Theorem 11.2 there is a $(K_X + \Delta)$ -crepant bimeromorphic model (X', Δ') such that

- (1) either there is a Mori fibration $f : X' \rightarrow Y$ such that $(K_{X'} + \Delta') \sim_{\mathbb{Q}, f} 0$,
or
- (2) $K_{X'} + (1 - \epsilon)\Delta'$ is nef for all $0 \leq \epsilon \ll 1$.

Moreover, X' has $\mathbb{Q}$ -factorial terminal singularities and (X', Δ') is a klt pair. Since the model is $(K_X + \Delta)$ -crepant, we only need to prove that $K_{X'} + \Delta'$ is semi-ample. If we are in the first case, then by Theorem 9.5, there is an effective $\mathbb{Q}$ -divisor $B \geq 0$ on Y such that (Y, B) is klt and

$$K_{X'} + \Delta' \sim_{\mathbb{Q}} f^*(K_Y + B).$$

Now by the log abundance for Kähler surfaces as in Theorem 6.3 it follows that $K_Y + B$ is semi-ample. Hence is $K_{X'} + \Delta'$ semi-ample.

If we are in the second case, then by Theorem 13.1, $\kappa(X', K_{X'} + \Delta') = 1$, and hence $K_{X'} + \Delta'$ is semi-ample by Corollary 7.4. $\square$

In the remainder of this section we will establish the log abundance theorem for log canonical pairs. First we need the following reduction lemma.

Lemma 14.2. *Let (X, Δ) be a log canonical pair and $K_X + \Delta$ is nef, where X is a $\mathbb{Q}$ -factorial compact Kähler 3-fold with terminal singularities. Then there exists a $\mathbb{Q}$ -factorial log canonical pair $(X', \Delta' := S' + B')$, where $S' = \lfloor \Delta' \rfloor$ and X' is a compact Kähler 3-fold with terminal singularities, such that $K_{X'} + \Delta'$ is semi-ample if and only if $K_X + \Delta$ is semi-ample and that one of the following conditions is satisfied:*

- (1) $K_{X'} + (1 - \epsilon)\Delta'$ is nef for all $0 \leq \epsilon \ll 1$
- (2) There is a $K_{X'}$ -Mori fiber space $\pi : X' \rightarrow Y'$ over a normal compact Kähler surface Y' such that $(K_{X'} + \Delta') \sim_{\mathbb{Q}, \pi} 0$ and some component of S' dominates Y' .

- (3) *There is a $K_{X'}$ -Mori fiber space $\pi : X' \rightarrow Y'$ over a normal compact Kähler surface Y' such that $(K_{X'} + \Delta') \sim_{\mathbb{Q}, \pi} 0$, no component of S' dominates Y' and $K_{X'} + B' + \lambda S'$ klt for all $0 \leq \lambda < 1$.*

Moreover, we have $\kappa(X, K_X + \Delta) = \kappa(X', K_{X'} + \Delta')$ and $\nu(X, K_X + \Delta) = \nu(X', K_{X'} + \Delta')$.

Proof. Using Theorem 11.2 we may assume that one of the following holds:

- (i) either there is a K_X -Mori fibration $f : X \rightarrow Y$ such that $(K_X + \Delta) \sim_{\mathbb{Q}, f} 0$, or
- (ii) $K_X + (1 - \epsilon)\Delta$ is nef for all $0 \leq \epsilon \ll 1$.

If we are in the case (ii), then set $(X', \Delta') := (X, \Delta)$ and we are done. So now assume that we are in case (i). If there is a component of $\lfloor \Delta \rfloor$ which dominates Y , then we are again done by setting $(X', \Delta') := (X, \Delta)$. So assume that $\lfloor \Delta \rfloor \neq 0$ and none of the components of $\lfloor \Delta \rfloor$ dominate Y . Let (X_1, Δ_1) be a dlt model of (X, Δ) as in Corollary 5.3. Write $\Delta_1 = S_1 + B_1$ with $S_1 = \lfloor \Delta_1 \rfloor$. Then $K_{X_1} + \Delta_1$ is nef, $(K_{X_1} + \Delta_1) \sim_{\mathbb{Q}, Y} 0$ and $K_X + \Delta$ is semi-ample if and only if $K_{X_1} + \Delta_1$ is semi-ample.

We run a $(K_{X_1} + B_1)$ -MMP over Y as in Proposition 3.8 and obtain a pair $(X_2, \Delta_2 = S_2 + B_2)$ such that $(K_{X_2} + B_2)$ is nef over Y . Note that this MMP is $(K_{X_1} + \Delta_1)$ -trivial. Thus (X_2, Δ_2) is lc and $(K_{X_2} + \Delta_2) \sim_{\mathbb{Q}, Y} 0$.

We run a K_{X_2} -MMP over Y with the scaling of B_2 and obtain a Mori fiber space $f' : X' \rightarrow Y'$. Then by Lemma 4.6 every step of this MMP is $(K_{X_2} + B_2)$ -trivial. Hence (X', B') is klt, where B' is the bimeromorphic transform of B_2 . We also remark that $(K_{X'} + B') \sim_{\mathbb{Q}, Y'} 0$ since $X' \rightarrow Y'$ is a Mori fiber space. Let S' be the bimeromorphic transform of S_2 and $\Delta' := B' + S'$. Since this MMP is also $(K_{X_2} + \Delta_2)$ -trivial, we see that (X', Δ') is lc, $K_{X'} + \Delta'$ is nef and $(K_{X'} + \Delta') \sim_{\mathbb{Q}, Y'} 0$. Moreover, $K_{X'} + \Delta'$ is semi-ample if and only if $K_X + \Delta$ is.

Now we claim that $(X', B' + \lambda S')$ is klt for all $0 \leq \lambda < 1$. To the contrary assume that there is a $\lambda_0 \in (0, 1)$ such that $(X', B' + \lambda_0 S')$ is not klt. Then there is an exceptional divisor E over X' such that $a(E, X', B' + \lambda_0 S') \leq -1$. Note that since (X', B') has $\mathbb{Q}$ -factorial klt singularities, it follows that $\text{Center}_{X'}(E) \subseteq \text{Supp } S'$. Therefore $a(E, X', B' + S') < -1$, this is a contradiction, since $(X', B' + S')$ has lc singularities. This completes the proof. $\square$

Finally we will prove our main theorem.

Proof of Theorem 1.1. First replacing (X, Δ) by a dlt model as in Corollary 5.3, we may assume that (X, Δ) is a dlt pair such that X has $\mathbb{Q}$ -factorial terminal singularities. Now recall that the non-vanishing is proved in Theorem

9.1. When $\kappa(X, K_X + \Delta) > 0$, the semi-ampleness of $K_X + \Delta$ is proved in Corollary 7.4. If $\nu(X, K_X + \Delta) = 0$, i.e. $K_X + \Delta \equiv 0$, then from the non-vanishing theorem it follows that $K_X + \Delta \sim_{\mathbb{Q}} 0$. If $\nu(X, K_X + \Delta) = 3$, then $K_X + \Delta$ is big; in particular, X is a compact Kähler Moishezon space with rational singularities. Hence X is a projective variety in this case by [Nam02, Theorem 1.2], and the semi-ampleness follows from [KMM94, Theorem 1.1]. So the only remaining case is $\nu(X, K_X + \Delta) = 1$. Using Lemma 14.2 we may assume that (X, Δ) satisfies one of the conditions in this lemma. Note that, this in particular implies that (X, Δ) is a lc pair such that X has $\mathbb{Q}$ -factorial terminal singularities.

Now suppose that we are in the first case of Lemma 14.2. If $\Delta = 0$, then the theorem follows from [CHP16, Theorem 8.1], as X has terminal singularities. We may then assume that $\Delta \neq 0$. Then $(X, (1 - \epsilon)\Delta)$ is klt and $K_X + (1 - \epsilon)\Delta$ is nef for all $0 < \epsilon \ll 1$. By Corollary 7.4 it is enough to show that $\kappa(X, (1 - \epsilon)\Delta) > 0$ for some $0 < \epsilon < 1$, because then we will have $\kappa(X, K_X + \Delta) > 0$. To this end, we will first show that $\nu(X, K_X + (1 - \epsilon_0)\Delta) = 1$ for some $0 < \epsilon_0 \ll 1$. First observe that if $K_X + (1 - \epsilon)\Delta \equiv 0$ for all $0 < \epsilon \ll 1$, then taking limit as $\epsilon \rightarrow 0$ we see that $K_X + \Delta \equiv 0$, this is a contradiction. Thus there is a $0 < \epsilon_0 \ll 1$ such that $\nu(X, K_X + (1 - \epsilon_0)\Delta) \geq 1$. Next we will show that $\nu(X, K_X + (1 - \epsilon_0)\Delta) < 2$. If $\nu(X, K_X + (1 - \epsilon_0)\Delta) = 3$, then $K_X + (1 - \epsilon_0)\Delta$ is big and then so is $K_X + \Delta$; this is a contradiction, since $\nu(X, K_X + \Delta) = 1$. So by contradiction assume now that $\nu(X, K_X + (1 - \epsilon_0)\Delta) = 2$. Since $\nu(X, K_X + \Delta) = 1$, we have $(K_X + \Delta)^2 \cdot \omega = 0$ for all Kähler class ω on X . In particular, this shows that $\Delta^2 \cdot \omega \leq 0$. But since $\nu(X, K_X + (1 - \epsilon_0)\Delta) = 2$, there is a Kähler class ω_0 such that $(K_X + (1 - \epsilon_0)\Delta)^2 \cdot \omega_0 > 0$. This implies that $(K_X + \Delta)^2 \cdot \omega_0 > 0$, i.e. $\nu(X, K_X + \Delta) \geq 2$, this is a contradiction. Therefore $\nu(X, K_X + (1 - \epsilon_0)\Delta) = 1$ for some $0 < \epsilon_0 \ll 1$. Then by Theorem 14.1, $K_X + (1 - \epsilon_0)\Delta$ is semi-ample, and hence $\kappa(X, K_X + (1 - \epsilon_0)\Delta) = 1$.

Assume now that we are in the second case of Lemma 14.2 and $\pi : X \rightarrow Y$ is the K_X -Mori fiber space as in there, and $\Delta = B + S$, where $S = \lfloor \Delta \rfloor$. Suppose that $\mathcal{L}$ is a line bundles on Y such that $\mathcal{O}_X(m(K_X + \Delta)) \cong \pi^* \mathcal{L}$ for some $m \in \mathbb{N}$ sufficiently divisible. Then it is enough to prove that $\mathcal{L}$ is semi-ample. Let T be a component of S which dominates Y . Then (T^n, Δ_{T^n}) is lc, where $\tau : T^n \rightarrow T$ is the normalization and $K_{T^n} + \Delta_{T^n} \sim_{\mathbb{Q}} (K_X + \Delta)|_{T^n}$ is defined by adjunction. Then by Theorem 6.3, $K_{T^n} + \Delta_{T^n}$ is semi-ample. Since $(K_{T^n} + \Delta_{T^n}) \sim_{\mathbb{Q}} (\pi|_T \circ \tau)^* \mathcal{L}$, from Lemma 2.5 it follows that $\mathcal{L}$ is semi-ample, and hence $K_X + \Delta$ is semi-ample.

Now assume that we are in the third case of Lemma 14.2 and $\pi : X \rightarrow Y$ is the K_X -Mori fiber space as in there, and $\Delta = B + S$, where $S = \lfloor \Delta \rfloor$. If $S = 0$,

then (X, Δ) is klt and we can apply Theorem 14.1. So from now on we assume that $S \neq 0$. We will show that $K_X + \Delta$ has positive Kodaira dimension. Since S is vertical over Y , by Lemma 9.7 there is an effective non-zero $\mathbb{Q}$ -Cartier divisor $D \geq 0$ on Y such that $f^*D = S$. Choose $0 < \epsilon < 1$ so that $(X, B + (1 - \epsilon)S)$ is klt as in the hypothesis. Note that $(K_X + B + (1 - \epsilon)S) \sim_{\mathbb{Q}, \pi} 0$. Then from Theorem 9.5 and its proof it follows that there is an effective $\mathbb{Q}$ -divisor $\Theta \geq 0$ on Y such that $K_X + B + (1 - \epsilon)S \sim_{\mathbb{Q}} f^*(K_Y + \Theta + (1 - \epsilon)D)$ and that $(Y, \Theta + (1 - \epsilon)D)$ is klt. We also have

$$(14.1) \quad (K_X + \Delta) \sim_{\mathbb{Q}} \pi^*(K_Y + \Theta + D),$$

and thus $K_Y + \Theta + D$ is nef on Y . Note that K_Y is pseudo-effective, since Y is not uniruled. Now we run a $(K_Y + \Theta)$ -MMP with the scaling of D . This MMP ends with a minimal model (Y', Θ') such that $K_{Y'} + \Theta'$ is nef. Let D' be the bimeromorphic transformation of D . Then $D' \neq 0$, $(Y', \Theta' + \lambda D')$ is klt and $K_{Y'} + \Theta' + \lambda D'$ is nef for some rational number $0 < \lambda < 1$. Now by Theorem 6.3, $K_{Y'} + \Theta' + \lambda D'$ is semi-ample. Moreover, notice that $\kappa(Y', K_{Y'} + \Theta' + \lambda D') > 0$, since otherwise $(K_{Y'} + \Theta') \sim_{\mathbb{Q}} -\lambda D'$, a contradiction to the fact that $K_{Y'} + \Theta'$ is nef. This implies that $K_{Y'} + \Theta' + D'$ has positive Kodaira dimension, and thus from (14.1) it follows that $\kappa(X, K_X + \Delta) > 0$. Then by Corollary 7.4, $K_X + \Delta$ is semi-ample. $\square$

APPENDIX A. RAMIFICATIONS FOR CURVES

Lemma A.1. *Let k be an algebraically closed field of characteristic 0. We consider the following commutative diagram*

$$\begin{array}{ccc} X' & \xrightarrow{p} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{\pi} & Y \end{array}$$

where $X = \operatorname{Spec} k[a]$, $Y = \operatorname{Spec} k[b]$, $Y' = \operatorname{Spec} k[c]$ and X' is the normalization of the fiber product $X \times_Y Y'$. Assume that f is given by $b \mapsto a^n$ and π by $b \mapsto c^m$. Let $y \in Y$ be the point defined by $b = 0$, $x \in X$ defined by $a = 0$. Let $x' \in X'$ be a point lying over x . Then the multiplicity of p^*x at x' is equal to $\frac{m}{\gcd(m, n)}$.

Proof. Note that $X' = \operatorname{Spec} A$, where A is the normalization of the ring $k[a] \otimes_{k[b]} k[c] \cong k[r, s]/(r^m - s^n)$ such that the projections p and f' are induced by $a \mapsto s$ and $c \mapsto r$, respectively. Let $d = \gcd(m, n)$ and γ be a primitive d -th root of unity. We write $m = m'd$ and $n = n'd$. Then A is also

the normalization of the product ring

$$\prod_{i=0}^{d-1} k[r, s]/(r^{m'} - \gamma^i s^{n'}).$$

Hence X' is the disjoint union of $X'_i = \text{Spec } B_i$, $i = 0, \dots, d-1$, where B_i is the normalization of the ring $k[r, s]/(r^{m'} - \gamma^i s^{n'})$.

Now since $x' \in p^{-1}(x)$, there is an $i \in \{0, 1, \dots, d-1\}$ such that x' is contained in X'_i . Pick $\delta \in k$ so that $\delta^{m'} = \gamma^i$. Then we can write $B_i = k[t]$ and the normalization map

$$k[r, s]/(r^{m'} - \gamma^i s^{n'}) \rightarrow B_i$$

is defined by $r \mapsto \delta t^{n'}$ and $s \mapsto t^{m'}$. Moreover, the point x' is defined by $t = 0$. Then the restriction of p on X'_i is defined by $a \mapsto t^{m'}$. This proves the lemma. $\square$

Definition A.2. Let $f : X \rightarrow Y$ be a proper surjective morphism between normal varieties and $D \subseteq Y$ a prime Weil divisor on Y . U be the smooth locus of X and $D_U := D|_U$. Then the *generic pullback* of under f is defined as

$$f^*D := \overline{f^*D_U} \subseteq X.$$

Note that f^*D could be a reducible divisor with multiplicities along its components. For an integral Weil divisor $B = \sum n_i D_i$, f^*B is defined by extending it linearly.

Lemma A.3. *We consider the following commutative diagram*

$$\begin{array}{ccc} X' & \xrightarrow{p} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{\pi} & Y \end{array}$$

where X , Y and Y' are normal varieties and X' is the normalization of the main component of $X \times_Y Y'$ dominating Y' . Assume that π is generically finite and f is a proper morphism with equidimensional fibers. Let D and E be two prime Weil divisor on Y and X , respectively, such that $E = \text{Supp}(f^*D)$. Suppose that the multiplicity of f^*D along E is n . Let D' be a prime Weil divisor on Y' which dominates D such that the multiplicity of π^*D along D' is m . Let E' be a prime Weil divisor on X' which dominates both D' and E . Then the multiplicity of p^*E along E' is $\frac{m}{\gcd(m, n)}$.

Proof. The problem is local around general points of D . By taking local hyperplane sections in Y , we may assume that Y and Y' are curves. Then by taking local hyperplane sections in X we may also assume that X is a curve. By choosing local analytic coordinates for X , Y and Y' , we may assume that the maps f and π are algebraic. We can then apply Lemma A.1 to conclude. $\square$

REFERENCES

- [BDPP13] S. Boucksom, J.-P. Demailly, M. Păun and T. Peternell, *The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension*, J. Algebraic Geom. **22**(2), 201–248 (2013).
- [BHPVdV04] W. P. Barth, K. Hulek, C. A. M. Peters and A. Van de Ven, *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, Springer-Verlag, Berlin, second edition, 2004.
- [Bou04] S. Boucksom, *Divisorial Zariski decompositions on compact complex manifolds*, Ann. Sci. École Norm. Sup. (4) **37**(1), 45–76 (2004).
- [CH20] J. Cao and A. Höring, *Rational curves on compact Kähler manifolds*, J. Differential Geom. **114**(1), 1–39 (2020).
- [CHP16] F. Campana, A. Höring and T. Peternell, *Abundance for Kähler threefolds*, Ann. Sci. Éc. Norm. Supér. (4) **49**(4), 971–1025 (2016).
- [CHP21] F. Campana, A. Höring and T. Peternell, *Addendum to the paper: Abundance for Kähler threefolds*, Available at <https://math.unice.fr/~hoering/articles/addendum-abundance.pdf> (2021).
- [Deb01] O. Debarre, *Higher-dimensional algebraic geometry*, Universitext, Springer-Verlag, New York, 2001.
- [Dem92] J.-P. Demailly, *Regularization of closed positive currents and intersection theory*, J. Algebraic Geom. **1**(3), 361–409 (1992).
- [Dem96] J.-P. Demailly, *L^2 vanishing theorems for positive line bundles and adjunction theory*, in *Transcendental methods in algebraic geometry (Cetraro, 1994)*, volume 1646 of *Lecture Notes in Math.*, pages 1–97, Springer, Berlin, 1996.
- [DP03] J.-P. Demailly and T. Peternell, *A Kawamata-Viehweg vanishing theorem on compact Kähler manifolds*, J. Differential Geom. **63**(2), 231–277 (2003).
- [DW19] O. Das and J. Waldron, *On the abundance problem for 3-folds in characteristic $p > 5$* , Math. Z. **292**(3–4), 937–946 (2019).
- [Fuj83] T. Fujita, *Semipositive line bundles*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **30**(2), 353–378 (1983).
- [Fuj00] O. Fujino, *Abundance theorem for semi log canonical threefolds*, Duke Math. J. **102**(3), 513–532 (2000).
- [Fuj07] O. Fujino, *Special termination and reduction to pl flips*, in *Flips for 3-folds and 4-folds*, volume 35 of *Oxford Lecture Ser. Math. Appl.*, pages 63–75, Oxford Univ. Press, Oxford, 2007.
- [Fuj11] O. Fujino, *On Kawamata’s theorem*, in *Classification of algebraic varieties*, EMS Ser. Congr. Rep., pages 305–315, Eur. Math. Soc., Zürich, 2011.
- [Fuj12] O. Fujino, *Basepoint-free theorems: saturation, b -divisors, and canonical bundle formula*, Algebra Number Theory **6**(4), 797–823 (2012).
- [Fuj79] A. Fujiki, *Closedness of the Douady spaces of compact Kähler spaces*, Publ. Res. Inst. Math. Sci. **14**(1), 1–52 (1978/79).
- [HM07] C. D. Hacon and J. McKernan, *On Shokurov’s rational connectedness conjecture*, Duke Math. J. **138**(1), 119–136 (2007).
- [HP15] A. Höring and T. Peternell, *Mori fibre spaces for Kähler threefolds*, J. Math. Sci. Univ. Tokyo **22**(1), 219–246 (2015).

- [HP16] A. Höring and T. Peternell, *Minimal models for Kähler threefolds*, Invent. Math. **203**(1), 217–264 (2016).
- [Kaw85] Y. Kawamata, *Pluricanonical systems on minimal algebraic varieties.*, Inventiones mathematicae **79**, 567–588 (1985).
- [Kaw88] Y. Kawamata, *Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces*, Ann. of Math. (2) **127**(1), 93–163 (1988).
- [Kaw91] Y. Kawamata, *On the length of an extremal rational curve*, Invent. Math. **105**(3), 609–611 (1991).
- [Kaw92] Y. Kawamata, *Termination of log flips for algebraic 3-folds*, Internat. J. Math. **3**(5), 653–659 (1992).
- [KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KMM94] S. Keel, K. Matsuki and J. McKernan, *Log abundance theorem for threefolds*, Duke Math. J. **75**(1), 99–119 (1994).
- [KMM04] S. Keel, K. Matsuki and J. McKernan, *Corrections to: “Log abundance theorem for threefolds”* [Duke Math. J. **75** (1994), no. 1, 99–119; MR1284817], Duke Math. J. **122**(3), 625–630 (2004).
- [Kol92] J. Kollár, *Flips and abundance for algebraic threefolds*, Société Mathématique de France, Paris, 1992, Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991, Astérisque No. 211 (1992).
- [Kol96] J. Kollár, *Singularities of Pairs*, in *eprint arXiv:alg-geom/9601026*, page 1026, January 1996.
- [Kol13] J. Kollár, *Singularities of the minimal model program*, volume 200 of *Cambridge Tracts in Mathematics*, Cambridge University Press, Cambridge, 2013, With a collaboration of Sándor Kovács.
- [Laz04] R. Lazarsfeld, *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series.
- [LP18] V. Lazić and T. Peternell, *Abundance for varieties with many differential forms*, Épijournal de Géométrie Algébrique **Volume 2** (February 2018).
- [Man82] M. Manaresi, *Sard and Bertini type theorems for complex spaces*, Ann. Mat. Pura Appl. (4) **131**, 265–279 (1982).
- [Miy87] Y. Miyaoka, *The Chern classes and Kodaira dimension of a minimal variety*, in *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 449–476, North-Holland, Amsterdam, 1987.
- [Miy88a] Y. Miyaoka, *Abundance conjecture for 3-folds: case $\nu = 1$* , Compositio Math. **68**(2), 203–220 (1988).
- [Miy88b] Y. Miyaoka, *On the Kodaira dimension of minimal threefolds*, Math. Ann. **281**(2), 325–332 (1988).
- [Nak87] N. Nakayama, *The lower semicontinuity of the plurigeners of complex varieties*, in *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 551–590, North-Holland, Amsterdam, 1987.

- [Nam02] Y. Namikawa, *Projectivity criterion of Moishezon spaces and density of projective symplectic varieties*, Internat. J. Math. **13**(2), 125–135 (2002).
- [Pau98] M. Paun, *Sur l'effectivité numérique des images inverses de fibrés en droites*, Math. Ann. **310**(3), 411–421 (1998).
- [Sho92] V. V. Shokurov, *3-fold log flips. Appendix by Yujiro Kawamata: The minimal discrepancy coefficients of terminal singularities in dimension three*, Russ. Acad. Sci., Izv., Math. **40**(1), 95–202 (1992).
- [Var89] J. Varouchas, *Kähler spaces and proper open morphisms*, Math. Ann. **283**(1), 13–52 (1989).
- [Wan19] J. Wang, *On the Iitaka Conjecture $C_{n,m}$ for Kähler Fibre Spaces*, arXiv e-prints, arXiv:1907.06705 (July 2019), 1907.06705.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, NAVY NAGAR, COLABA, MUMBAI 400005, INDIA

Email address: `omprokash@gmail.com`

Email address: `omdas@math.tifr.res.in`

INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

Email address: `wenhaoou@amss.ac.cn`