

Information geometry of quantum critical submanifolds: relevant, marginal and irrelevant operators

Bruno Mera*

*Instituto de Telecomunicações, 1049-001 Lisboa, Portugal
Departamento de Matemática, Instituto Superior Técnico,
Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal and
Departamento de Física, Instituto Superior Técnico,
Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal*

Nikola Paunković†

*Instituto de Telecomunicações, 1049-001 Lisboa, Portugal and
Departamento de Matemática, Instituto Superior Técnico,
Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal*

Syed Tahir Amin‡

*Instituto de Telecomunicações, Av. Rovisco Pais 1, 1049-001 Lisbon, Portugal
Departamento de Física, Instituto Superior Técnico,
Universidade de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal
CeFEMA, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal
Department of Physics and I3N, University of Aveiro, 3810-193 Aveiro, Portugal and
Center of Physics of University of Minho and University of Porto, P-4169-007 Oporto, Portugal*

Vítor R. Vieira§

*Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa,
Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal and
CeFEMA, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal
(Dated: January 6, 2022)*

We analyze the thermodynamical limit of the quantum metric along critical submanifolds of theory space. We relate its singular behaviour to normal directions, which are naturally associated to relevant operators in the renormalization group sense. We exemplify our theory through the paradigmatic examples of the XY and Haldane models, where the normal directions to the critical submanifolds are seen to be precisely those along which the metric has singular behaviour, while for the tangent ones it vanishes—these directions lie in the kernel of the metric.

I. INTRODUCTION

Recently, information geometry has seen a lot of application in the study of quantum systems and their phase transitions [1–5]. Of particular interest are the exotic topological phases of matter that go beyond the standard Landau-Ginzburg classification paradigm and for which there is no well-defined local order parameter, used to infer phase transitions. The quantum metric [6]—a Riemannian metric over the space of quantum states based on state distinguishability—was shown to be a quantity able to probe zero temperature quantum phase transitions among these phases as one varies some parameter in the system [7–9]. In the context of Bloch bands, the quantum metric in momentum space, in the particular setting of flat bands, has received a lot of attention recently, as it gives geometrical contributions to the characterization of a range of different phenomena, such as

exotic superconductivity and superfluidity [10–13], the stability of fractional Chern insulating phases [14–19] and light-matter coupling [20]. The quantum metric is also central in determining maximally-localized Wannier functions [21, 22], and it can be used as a practical indicator for exotic momentum-space monopoles [23, 24]. The integral of the momentum-space quantum metric is a measure of electron localization and it can be extracted through spectroscopy measurements [25]. In two-spatial dimensions, one can define an associated complex structure, which is a measure of anisotropy in localization [26].

More recently, relations between the quantum metric and the Berry curvature, which determines topological invariants of the system and gives rise to Berry-phase effects, have been established and understood based on the Kähler structure of the space of quantum states [27, 28]. These relations have come to play an important role in recent studies concerning the so-called ideal Chern bands, which are presumed to be ideal candidates for hosting fractional Chern insulating phases, the associated band structure engineering and transport [19, 29–34]. More general relations between the quantum metric and topology of quantum states were derived in the context of

* bruno.mera@tecnico.ulisboa.pt

† npaunkov@math.tecnico.ulisboa.pt

‡ tahiramin811@gmail.com

§ vitor.rocha.vieira@tecnico.ulisboa.pt

Dirac Hamiltonians [35].

The quantum metric over the space of parameters describing the system, being the “infinitesimal” distance between two ground states, allows us to probe phase transitions. As long as the quantum metric is regular, the ground state of the system does not change substantially, keeping the system in the same phase. On the other hand, if the quantum metric becomes singular, it means that the ground state has undergone a dramatic change and, thus, the system has experienced a phase transition. However, this singular behaviour has its subtleties regarding the scaling behaviour when performing the thermodynamical infinite volume limit. In the following, we will show how the quantum metric behaves along a submanifold of parameter space composed of critical points—described, in the long wavelength and infinite volume limit, by conformal field theories. We will show that there exist two types of directions of the parameter change: those which take the system away from, and those which move it within the critical submanifold. By performing a scaling analysis, we show that the behaviours of these directions can be interpreted in light of the theory of the renormalization group (RG), namely, the former directions correspond to relevant, while the latter correspond to irrelevant and marginal operators in the quantum theory. We also illustrate our results on the examples of the XY and the Haldane models. Finally, we present conclusions and future lines of research.

II. QUANTUM METRIC ALONG A CRITICAL SUBMANIFOLD

Suppose we are given a many-body Hamiltonian depending smoothly on some parameters, collectively denoted by x , living in a smooth manifold \mathcal{T} of dimension n . We refer to \mathcal{T} as *theory space*. These parameters will typically correspond to couplings and external fields determining the system’s Hamiltonian $H(x)$, with $x \in \mathcal{T}$. We will be interested in the ground state, i.e., zero temperature properties of the system. In particular, we consider the pullback of the *Fubini-Study metric* [which is the natural unitarily invariant metric in the space of pure quantum states], also known as the fidelity susceptibility or the information metric, with respect to the family of ground states $\rho(x) = |\psi(x)\rangle\langle\psi(x)|$ of $H(x)$, with $x \in \mathcal{T}$. Note that there may be parameters for which the ground state is not unique, in which case we will adopt a regularization procedure where we consider the pullback of the *Bures metric*—a Riemannian metric on the space of density operators—under the map $x \mapsto e^{-H(x)/T} / \text{Tr}(e^{-H(x)/T})$ and eventually take the $T \rightarrow 0$ limit. The Bures metric has two independent contributions coming from the “classical” and “quantum” parts of the density matrix $\rho = \sum_j p_j |\psi_j\rangle\langle\psi_j|$, corresponding to the variations on the probabilities and the

eigenstates of ρ , respectively, see Ref. [3]:

$$ds^2 = \frac{1}{4} \sum_j \frac{dp_j^2}{p_j} + \frac{1}{2} \sum_{j \neq i} |\langle\psi_j|d|\psi_i\rangle|^2 \frac{(p_j - p_i)^2}{p_i + p_j}. \quad (1)$$

The first term is the classical Fisher metric corresponding to the statistical model determined by the probability distribution (p_j) . The second term, measuring the variations in the eigenstates of ρ , has the nice property that for pure states $\rho = |\psi\rangle\langle\psi|$ it reduces to the Fubini-Study metric

$$g_{FS} = \langle d\psi|(1 - |\psi\rangle\langle\psi|)d\psi\rangle. \quad (2)$$

Under the map $x \mapsto |\psi_i(x)\rangle\langle\psi_i(x)|$, where $|\psi_i(x)\rangle$ is an eigenstate of $H(x)$ [assuming such state is globally defined, up to a phase], the pullback of the Fubini-Study metric g_{FS} describes a metric in the space of theories \mathcal{T} —the *quantum metric*, which has the nice formula in terms of the remaining eigenstates

$$g = \sum_{j \neq i} \langle d\psi_i(x)|\psi_j(x)\rangle\langle\psi_j(x)|d\psi_i(x)\rangle. \quad (3)$$

We will now take $|\psi_i(x)\rangle = |\psi(x)\rangle$ to be the ground state of $H(x)$. It is convenient to interpret the quantum metric g in terms of operators acting on the Hilbert space. At a given point $x \in \mathcal{T}$, the metric assigns a bilinear nonnegative pairing $g(u, v)$ of tangent vectors $u, v \in T_x\mathcal{T}$. Physically, an element $v \in T_x\mathcal{T}$ generates a first order variation in the parameters of the theory at $x \in \mathcal{T}$. As such, it has a corresponding operator, \mathcal{O}_v , acting on the Hilbert space, which, using local coordinates x^i , $i = 1, \dots, n$, where $v = v^i \frac{\partial}{\partial x^i}$, is given by

$$\mathcal{O}_v = v^i \frac{\partial H}{\partial x^i}, \quad (4)$$

where we assume the Einstein summation convention. It is clear that the operator \mathcal{O}_v is associated to perturbation theory in the sense that $H(x + \varepsilon v) = H(x) + \varepsilon \mathcal{O}_v + \mathcal{O}(\varepsilon^2)$. At each parameter point x from our space of theories \mathcal{T} , once we fix local coordinates x^i , $i = 1, \dots, n$, we have a basis for an associated linear space of operators given by $\mathcal{O}_i(x) = \mathcal{O}_{\partial/\partial x^i}$, $i = 1, \dots, n$. The pullback of the Fubini-Study metric under the map $x \mapsto |\psi(x)\rangle\langle\psi(x)|$, describes the quantum metric in the space of theories \mathcal{T} . Note, however, that it may happen that the metric is degenerate, meaning it is not necessarily invertible at every point x where it is defined—simply because the Jacobian of the transformation can be singular. Alternatively, we can view the degeneracy of the metric in light of the tangent vector to operator correspondence $v \mapsto \mathcal{O}_v$. Assuming the eigenstates are non-degenerate in energy, let $|\psi_i(x)\rangle$ be the eigenstates for $H(x)$ with corresponding energies $E_i(x)$. The eigenstates of $H(x + \varepsilon v)$ can be obtained from first order perturbation theory around $H(x)$ as follows

$$|\psi_i(x + \varepsilon v)\rangle = |\psi_i(x)\rangle + \varepsilon \sum_{j \neq i} \frac{\langle\psi_j(x)|\mathcal{O}_v|\psi_i(x)\rangle}{E_i(x) - E_j(x)} |\psi_j(x)\rangle, \quad (5)$$

where we assume a parallel transport gauge $\langle \psi_i(x) | v^k \frac{\partial}{\partial x^k} | \psi_i(x) \rangle = 0$. The order ε term can vanish if $v \in T_x \mathcal{T}$ is in the kernel of the assignment $v \mapsto \mathcal{O}_v$. More generally, it can vanish if and only if \mathcal{O}_v preserves the eigenspace $L_x^{(i)} = \text{span}_{\mathbb{C}}\{|\psi_i(x)\rangle\}$, the fiber at x of the i th eigenbundle of H . Observe that from Eq. (3), it follows that this precisely yields the vanishing of the pullback of the Fubini-Study metric at $x \in \mathcal{T}$. In short, for the ground state $\rho(x) = |\psi(x)\rangle\langle\psi(x)|$, the above statement can be expressed as

$$[\mathcal{O}_v, \rho(x)] = 0 \iff g(v, \cdot) = 0. \quad (6)$$

Suppose now we have a set $C \subset \mathcal{T}$ of *quantum critical points* where the theory is gapless and our geometric quantity of interest g is presumably singular in the thermodynamical limit. The thermodynamical limit assumes, in particular, that the volume of the system L^d , where L is the linear length of the system and d is the spatial dimension, goes to infinity. For considerations involving the thermodynamical limit, it will be convenient to consider the rescaled quantum metric g/L^d . We also make the assumption that the quantum critical theories, as described by points $x \in C \subset \mathcal{T}$, are conformally invariant so that they are described, in the continuum limit and at large distances, by a conformal field theory (CFT $_d$). For simplicity, we assume C is a submanifold of \mathcal{T} . Of particular interest to us will be the tangent bundle of C in \mathcal{T} , i.e., the vector bundle over C that at each point $x \in C$ associates the tangent space $T_x C \subset T_x \mathcal{T}$. The point that we want to make, which will be clear from the discussion below, is that the rescaled metric g/L^d is actually, in the thermodynamical limit, only divergent along directions which are complementary to TC , which we refer to as directions of *increased ground state distinguishability* and are associated to *relevant operators* in the framework of the RG. To be precise, the thermodynamical limit of g/L^d , which we denote by g_* , vanishes exactly along tangent vectors to C , i.e., $g_*(v, \cdot) = 0$ for any vector $v \in T_x C \subset T_x \mathcal{T}$. As a consequence, the directions where g_* is non-vanishing are normal to TC —i.e., each can be seen, effectively, as orthogonal to TC with respect to some Riemannian metric in \mathcal{T} . The rigorous statement is that, for each $x \in C$, there is an *exact sequence of vector spaces*

$$0 \longrightarrow T_x C \longrightarrow T_x \mathcal{T} \longrightarrow N_x C \longrightarrow 0, \quad (7)$$

defining the vector space of normal directions at x , $N_x C = T_x \mathcal{T} / T_x C$, giving rise, globally, to the normal bundle $NC = T\mathcal{T}|_C / TC \rightarrow C$, see Fig. 1 [by an exact sequence of vector spaces, we mean a sequence of vector spaces and linear maps where the kernel of the next is equal to the image of the current]. Observe that for $x \in C$ and any $u \in T_x C$, if $g_*(u, u) \neq 0$ we have that $g_*(u + v, u) = g_*(u, u)$ for any $v \in T_x C$, since v is annihilated by g_* . It follows that u defines a non-trivial element of N_ϕ . In practice, one chooses an auxiliary Riemannian metric h defined over \mathcal{T} (actually it is

enough to have it defined over $T\mathcal{T}|_C$), and identifies $N_x C \cong (T_x C)^\perp$ using the notion of orthogonal complement provided by h . Importantly, the isomorphism class of this vector bundle is independent of this choice. Finally, we remark that, in light of Eq. (6), we see that the operators associated with $T_x C$ commute with the ground state at x modulo subextensive terms, i.e., terms that scale as L^r , with $r < d$.




FIG. 1. An illustration of the space of theories \mathcal{T} (here a compact surface of genus 1), together with a quantum critical submanifold $C \subset \mathcal{T}$ (here a circle). The red arrow is meant to illustrate a tangent vector to C at $x \in C$, which maps to a marginal or irrelevant operator. The blue arrow illustrates a normal vector to C at $x \in C$, mapping to a relevant operator.

III. RELATION TO RG

We consider a critical point $x_0 \in C \subset \mathcal{T}$ and assume that the \mathcal{O}_i 's are local operators, meaning that they can be written as

$$\mathcal{O}_i = \sum_{\mathbf{r}} \mathcal{O}_i(\mathbf{r}), \quad (8)$$

for $\mathcal{O}_i(\mathbf{r})$ having support in some bounded neighbourhood of \mathbf{r} . The sum over lattice sites \mathbf{r} makes sense before taking the continuum limit. It is convenient to introduce the imaginary time evolved operators, namely,

$$\mathcal{O}_i(\tau, \mathbf{r}) = e^{\tau H(x_0)} \mathcal{O}_i(\mathbf{r}) e^{-\tau H(x_0)}, \quad (9)$$

where τ denotes the imaginary time. At the critical point $x_0 \in C$, we may assume that the \mathcal{O}_i 's have well-defined scaling dimensions Δ_i such that

$$\mathcal{O}_i(\tau, \mathbf{r}) \longrightarrow \mathcal{O}'_i(\tau, \mathbf{r}) = \mathcal{O}_i(\zeta^z \tau, \zeta \mathbf{r}) \zeta^{\Delta_i}, \quad (10)$$

under a transformation $\mathbf{r} \rightarrow \zeta \mathbf{r}$ and $\tau \rightarrow \zeta^z \tau$, where z is the dynamical critical exponent. If it is not the case, assuming $\dim \mathcal{T}$ is big enough, we may use a linear coordinate change, so that the above equation holds. These scaling dimensions arise from the linearization of the renormalization group flow near the critical point $x_0 \in C$, see Ref. [36] for details.

Due to the expression

$$g_{ij}(x) = \int_0^\infty d\tau \tau e^{-\varepsilon\tau} \left[\frac{1}{2} \langle \{\mathcal{O}_i(\tau), \mathcal{O}_j(0)\} \rangle - \langle \mathcal{O}_i(\tau) \rangle \langle \mathcal{O}_j(0) \rangle \right], \quad (11)$$

with $\varepsilon \rightarrow 0^+$ and $\{\cdot, \cdot\}$ being the anticommutator, one discovers that $g_{ij}(x_0)$ has the finite size scaling law originally found by Venuti, Zanardi and coworkers [2, 4] and later found in the work of Miyaji, Numasawa *et. al* [37], in the context of gauge-gravity duality,

$$g_{ij}(x_0) \sim L^{2d+2z-\Delta_i-\Delta_j}, \quad (12)$$

where L is the linear size of the system and d its spatial dimension. In particular, we have the following classification of operators according to how their couplings are renormalized under an RG transformation, as we flow onto the infrared energy scales,

- \mathcal{O}_i is *relevant* if $\Delta_i < d + z$.
- \mathcal{O}_i is *marginal* if $\Delta_i = d + z$.
- \mathcal{O}_i is *irrelevant* if $\Delta_i > d + z$.

We then see that for tangent vectors whose associated operators are *relevant* operators, the metric should blow up in the $L \rightarrow \infty$ thermodynamical limit. For tangent vectors associated with *irrelevant* operators the metric should vanish in the thermodynamical limit. Finally, for tangent vectors associated with *marginal* operators, the metric should scale as L^0 , meaning it should be finite, with possible logarithmic corrections. Accordingly, if we are in a point x taken from a small neighbourhood of $x_0 \in C$, the relevant operators moves us outside C , the irrelevant operators move us back to x_0 , while the marginal operators move us within C , since the corresponding theory is also *scale invariant*. Thus, we expect that the tangent space $T_{x_0}C$ can be identified with the subspace of all irrelevant and marginal operators in the image of $T_{x_0}\mathcal{T} \ni v \mapsto \mathcal{O}_v$. The marginal operators in the image of $T_{x_0}\mathcal{T} \ni v \mapsto \mathcal{O}_v$ then form a complementary subspace $N_{x_0}C$, such that $T_{x_0}\mathcal{T} = T_{x_0}C \oplus N_{x_0}C$.

Consider the rescaled metric $g_{ij}(x_0)/L^d$. The scaling behaviour of its diagonal components is given by

$$\frac{g_{ii}}{L^d} \sim \frac{1}{L^d} L^{2z+2d-2\Delta_i}, \quad (13)$$

and hence it vanishes for $2z + d - 2\Delta_i < 0$ and it is finite or blows up for $2z + d - 2\Delta_i \geq 0$ in the thermodynamic limit. In particular, for irrelevant and marginal operators, it goes to zero. Surprisingly, it also vanishes for those relevant operators that are not “sufficiently relevant”, namely those that satisfy

$$z + d > \Delta_i > z + \frac{d}{2}. \quad (14)$$

Its thermodynamical limit g_* vanishes along $T_{x_0}C$, so we can identify $T_{x_0}C$ as a subspace of $T_{x_0}\mathcal{T}$ where the rescaled metric has nontrivial kernel. In the following, we consider two particular examples for which the tangent bundle to C in \mathcal{T} is exactly the kernel of g_* —the XY and the Haldane models.

IV. XY MODEL

In the XY anisotropic spin-half chain with $N = 2M + 1$ sites on a circle in the presence of an external magnetic field, we have the family of Hamiltonians

$$H(\gamma, \lambda) = - \sum_{i=-M}^M \left(\frac{1+\gamma}{2} \sigma_i^x \sigma_{i+1}^x + \frac{1-\gamma}{2} \sigma_i^y \sigma_{i+1}^y + \frac{\lambda}{2} \sigma_i^z \right), \quad (15)$$

parametrized by $(\gamma, \lambda) \in \mathbb{R}^2$, where γ is the anisotropy and λ the magnetic field. The critical region consists of three critical submanifolds given, respectively, by the segment defined by $\gamma = 0$ and $|\lambda| \leq 1$, and the two lines given by $|\lambda| = 1$ (see for example Ref. [1]). The segment and the lines intersect transversally. The whole set, i.e., the union of the critical submanifolds, is not a submanifold itself, as the intersection points do not have neighbourhoods homeomorphic to \mathbb{R} .

For this model, we see that g_* restricted to the critical lines vanishes precisely on tangent vectors to the critical lines, except on the intersections $(\lambda, \gamma) = (\pm 1, 0)$ where it blows up in all directions. On the complementary subspaces to the tangent spaces it always blows up. In this case there are no operators which are not sufficiently relevant. We remark that at the two critical points $(\gamma, \lambda) = (0, \pm 1)$, corresponding to the intersection between the segment and the two lines, the metric blows up along any direction and, hence, they behave as if they were two isolated critical points [because the normal spaces in this case coincide with the whole of $T_{(0, \pm 1)}\mathcal{T}$]. Note, however, that they are not isolated critical points in the strict sense, as any neighbourhood of them finds other critical points. We refer the reader to Appendix A for details.

V. HALDANE MODEL

We consider the Haldane model Hamiltonian [38]

$$\mathcal{H} = t \sum_{\langle l, m \rangle} c_l^\dagger c_m + t' \sum_{\langle\langle l, m \rangle\rangle} e^{-i\nu_{lm}\phi} c_l^\dagger c_m + M \sum_l \epsilon_l c_l^\dagger c_l, \quad (16)$$

where $\epsilon_l = 1$ on site A and -1 on site B; $\nu_{lm} = \pm 1$ depending on the direction of next nearest neighbor hopping. This model supports quantized conductance without applying an external magnetic field. We also fix $t = 1$ and $t' = 1/3$. In this case, (ϕ, M) are the coordinates describing the parameter manifold in which different topological phases exist. We remark that, in this case, the

manifold has the topology of an infinite cylinder $S^1 \times \mathbb{R}$, since we have to identify ϕ and $\phi + 2\pi$. The critical region for this model is well-known and in the (ϕ, M) plane they have the shape of a *figure eight*, see [38]. Once we identify the points $\phi \sim \phi + 2\pi$, the resulting critical region consists of two circles which intersect at two points described by $(M, \phi) = (0, 0)$ and $(\pi, 0)$, both intersecting transversally.

On Fig. 2 we present fidelity susceptibilities (rescaled quantum metric diagonal components) $\chi_{MM} = g_{MM}/L^2$ (the upper plot) and $\chi_{\phi\phi} = g_{\phi\phi}/L^2$ (the lower plot) calculated at temperature $T = 2 \times 10^{-5}$ and $N_x = N_y = 30$ sites. We see that both plots clearly show the critical lines, apart from parts of the second plot when the change of the parameter ϕ is precisely tangent to the critical manifold, showing that the tangent directions have decreased distinguishability. Just like the case of the XY model, the points $(M, \phi) = (0, 0)$ and $(\pi, 0)$ behave as isolated critical points and the metric blows up in all direction in the thermodynamical limit.

VI. CONCLUSIONS

In this paper, we analyzed the thermodynamical limit of the quantum metric along critical submanifolds of theory space. We related its singular behaviour to normal directions, which are naturally associated to relevant operators. In the paradigmatic examples of the XY and Haldane models, we have seen (exactly for the first model and numerically for the second) that the normal directions to the critical submanifolds are precisely those where the metric has singular behaviour in the thermodynamical limit, while the tangent ones vanish. In both of these models, the critical regions consist of critical submanifolds, which intersect transversally, and the intersection points behave as isolated critical points with enhanced distinguishability. Further analysis of this phenomenon can serve as a future line of research. As we have seen, our theory predicts that there can also be directions associated to relevant operators that lie in the kernel of g_* , a remark which also deserves further investigation. It would be interesting to understand how the above results generalize to the finite temperature case, where Eq. (11) relating the metric and two-point correlation functions gets modified for the case of the Bures metric. For the case of the interferometric metric considered in Ref. [39], the same functional form is retained, and it would be interesting to understand, within the present context, what is the physical relevance of considering either metric.

Acknowledgements.—BM and NP's work was partially supported by SQIG – Security and Quantum Information Group of Instituto de Telecomunicações, by

Programme (COMPETE 2020) of the Portugal 2020 framework [Project Q.DOT with Nr. 039728 (POCI-01-0247-FEDER-039728)] and the Fundação para a




FIG. 2. Fidelity susceptibilities $\chi = \chi_{MM}(\phi, M)$ (the upper plot) and $\chi = \chi_{\phi\phi}(\phi, M)$ (the lower plot) calculated at for temperature $T = 2 \times 10^{-5}$ and $N_x = N_y = 30$ sites.

Ciência e a Tecnologia (FCT) through national funds, by FEDER, COMPETE 2020, and by Regional Operational Program of Lisbon, under UIDB/50008/2020 (actions QuRUNNER, QUESTS), Project QuantumMining POCI-01-0145-FEDER-031826 and Project PREDICT PTDC/CCI-CIF/29877/2017.

NP acknowledges FCT Projects CERN/FIS-PAR/0023/2019, QuantumPrime PTDC/EEI-TEL/8017/2020, as well as the FCT Estímulo ao Emprego Científico grant no. CEECIND/04594/2017/CP1393/CT000.

This work was also supported through grant No. UID/CTM/04540/2019.

Appendix A: Thermodynamic limit of the quantum metric

In this Appendix, we analyze the thermodynamical limit of the quantum metric in the case of the anisotropic XY spin chain in external magnetic field. A straightforward calculation shows that the diagonal components of the quantum metric are given by

$$g_{\lambda\lambda} = \frac{1}{4} \sum_k \frac{\gamma^2 \sin^2(k)}{\left[(\lambda - \cos(k))^2 + \gamma^2 \sin^2 k \right]^2},$$

$$g_{\gamma\gamma} = \frac{1}{4} \sum_k \frac{(\lambda - \cos(k))^2 \sin^2(k)}{\left[(\lambda - \cos(k))^2 + \gamma^2 \sin^2 k \right]^2}.$$

Below we analyse the large N behaviour of the above expressions in the critical region.

1. Segment $\gamma = 0$ and $|\lambda| \leq 1$

For the critical submanifold defined by $\gamma = 0$ and $|\lambda| \leq 1$, with tangent vector $\frac{\partial}{\partial \lambda}$, we immediately see that $g_{\lambda\lambda} = 0$ independently of N . For $g_{\gamma\gamma}$ the metric becomes

$$g_{\gamma\gamma}(\lambda, 0) = \frac{1}{4} \sum_k \frac{\sin^2(k)}{(\lambda - \cos(k))^2}.$$

For $g_{\gamma\gamma}$ the metric is finite for $|\lambda| > 1$, but for $|\lambda| \leq 1$ we have to be careful because the equation

$$\cos(k) = \lambda,$$

will have exactly one solution in the thermodynamical limit. In that case, let k_* be the solution. If k is an allowed momentum in a small neighbourhood of k_* , i.e., $k = k_* + \delta q$, then we may write, $\cos(k) \approx \lambda - \sin(k_*)\delta k = \lambda - \sqrt{1 - \lambda^2}\delta k$ and $\sin(k) \approx \sin(k_*) + \lambda\delta k = \sqrt{1 - \lambda^2} + \lambda\delta k$ (where the sine is positive, since $k \in [0, \pi]$). Then,

$$\frac{\sin^2(k)}{(\lambda - \cos(k))^2} \sim \frac{(1 - \lambda^2)}{(1 - \lambda^2)\delta k^2} \sim N^2,$$

so we see that

$$g_{\gamma\gamma}(\lambda, 0) \sim N^2, \text{ as } N \rightarrow \infty, \text{ for } |\lambda| < 1.$$

For $|\lambda| = 1$ then $\cos(k) \approx \pm 1 + O(\delta k^2)$ and $\sin(k) \approx -\delta k$ (with $\delta k < 0$ since the sine is positive). Hence,

$$\frac{\sin^2(k)}{(\lambda - \cos(k))^2} \sim \frac{\delta k^2}{\delta k^4} \sim \frac{1}{\delta k^2} \sim N^2,$$

as before.

2. Line $\lambda = 1$

For the critical submanifold defined by $\lambda = 1$, with tangent vector $\frac{\partial}{\partial \gamma}$, we have

$$g_{\lambda\lambda}(1, \gamma) = \frac{1}{4} \sum_k \frac{\gamma^2 \sin^2(k)}{\left[(1 - \cos(k))^2 + \gamma^2 \sin^2 k \right]^2},$$

we have to look at momenta in the neighbourhood of $k = 0$, i.e., $k = \delta k$ (with $\delta k > 0$), and then

$$\sin(k) \approx \delta k \text{ and } \cos(k) \approx 1 - \frac{1}{2}\delta k^2,$$

so

$$\begin{aligned} \frac{\gamma^2 \sin^2(k)}{\left[(1 - \cos(k))^2 + \gamma^2 \sin^2 k\right]^2} &\approx \frac{\gamma^2 \delta k^2}{\left[\frac{1}{4} \delta k^4 + \gamma^2 \delta k^2\right]^2} \\ &\approx \frac{1}{\gamma^2 \delta k^2} \sim \frac{N^2}{\gamma^2}, \end{aligned}$$

thus

$$g_{\lambda\lambda}(1, \gamma) \sim \frac{N^2}{\gamma^2}.$$

This means that as $N \rightarrow \infty$ it blows up. Now,

$$g_{\gamma\gamma}(1, \gamma) = \frac{1}{4} \sum_k \frac{(1 - \cos(k))^2 \sin^2(k)}{\left[(1 - \cos(k))^2 + \gamma^2 \sin^2 k\right]^2}.$$

Looking at the momenta in the neighbourhood of $k = 0$ as before, we see that

$$\frac{(1 - \cos(k))^2 \sin^2(k)}{\left[(1 - \cos(k))^2 + \gamma^2 \sin^2 k\right]^2} \approx \frac{1}{4} \frac{\delta k^6}{\left(\frac{1}{4} \delta k^4 + \gamma^2 \delta k^2\right)^2} \approx \frac{1}{4\gamma^4} \delta k^2,$$

so

$$g_{\gamma\gamma}(1, \gamma) \text{ is finite for } \gamma \neq 0 \text{ and } g_{\gamma\gamma}(1, \gamma) \sim \frac{1}{N^2 \gamma^4} \text{ as } \gamma \rightarrow 0.$$

3. Line $\lambda = -1$

For the critical submanifold defined by $\lambda = -1$, with tangent vector $\frac{\partial}{\partial \gamma}$, we have

$$g_{\lambda\lambda}(-1, \gamma) = \frac{1}{4} \sum_k \frac{\gamma^2 \sin^2(k)}{\left[(1 + \cos(k))^2 + \gamma^2 \sin^2 k\right]^2},$$

we have to look at momenta in the neighbourhood of $k = \pi$, i.e., $k = \pi + \delta k$ (with $\delta k < 0$), and then

$$\sin(k) \approx -\delta k \text{ and } \cos(k) \approx -1 + \frac{1}{2} \delta k^2,$$

so

$$\frac{\gamma^2 \sin^2(k)}{\left[(1 + \cos(k))^2 + \gamma^2 \sin^2 k\right]^2} \approx \frac{\gamma^2 \delta k^2}{\left[\frac{1}{4} \delta k^4 + \gamma^2 \delta k^2\right]^2} \approx \frac{1}{\gamma^2 \delta k^2} \sim \frac{N^2}{\gamma^2},$$

thus

$$g_{\lambda\lambda}(1, \gamma) \sim \frac{N^2}{\gamma^2}.$$

This means that as $N \rightarrow \infty$ it blows up. Now,

$$g_{\gamma\gamma}(-1, \gamma) = \frac{1}{4} \sum_k \frac{(1 + \cos(k))^2 \sin^2(k)}{\left[(1 + \cos(k))^2 + \gamma^2 \sin^2 k\right]^2}.$$

Looking at the momenta in the neighbourhood of $k = \pi$ as before, we see that

$$\frac{(1 + \cos(k))^2 \sin^2(k)}{\left[(1 + \cos(k))^2 + \gamma^2 \sin^2 k\right]^2} \approx \frac{1}{4} \frac{\delta k^6}{\left(\frac{1}{4}\delta k^4 + \gamma^2 \delta k^2\right)^2} \approx \frac{1}{4\gamma^4} \delta k^2,$$

so

$$g_{\gamma\gamma}(-1, \gamma) \text{ is finite for } \gamma \neq 0 \text{ and } g_{\gamma\gamma}(-1, \gamma) \sim \frac{1}{N^2\gamma^4} \text{ as } \gamma \rightarrow 0.$$

4. Summary

- Critical line $\gamma = 0$, tangent vector $\partial/\partial\lambda$:

$$g_{\lambda\lambda}(\lambda, 0) = 0 \text{ and } g_{\gamma\gamma}(\lambda, 0) = \begin{cases} \sim N^2, \text{ as } N \rightarrow \infty, \text{ for } |\lambda| \leq 1, \\ \text{finite, otherwise.} \end{cases}.$$

- Critical line $\lambda = 1$, tangent vector $\partial/\partial\gamma$:

$$g_{\lambda\lambda}(1, \gamma) \sim \frac{N^2}{\gamma^2} \text{ and } g_{\gamma\gamma}(1, \gamma) = \begin{cases} \text{is finite for } \gamma \neq 0, \\ \sim \frac{1}{N^2\gamma^4} \text{ as } \gamma \rightarrow 0. \end{cases}.$$

- Critical line $\lambda = -1$, tangent vector $\partial/\partial\gamma$:

$$g_{\lambda\lambda}(-1, \gamma) \sim \frac{N^2}{\gamma^2} \text{ and } g_{\gamma\gamma}(-1, \gamma) = \begin{cases} \text{is finite for } \gamma \neq 0, \\ \sim \frac{1}{N^2\gamma^4} \text{ as } \gamma \rightarrow 0. \end{cases}.$$

In this case we see that $g_* = \lim_{N \rightarrow \infty} \frac{g}{N}$ restricted to the critical lines vanishes precisely on tangent vectors to the critical lines, except on the intersections $(\lambda, \gamma) = (\pm 1, 0)$ where it blows up in all directions. On the complementary subspaces to the tangent spaces it always blows up. In this case there are no operators which are not sufficiently relevant.

-
- [1] Paolo Zanardi and Nikola Paunković. Ground state overlap and quantum phase transitions. *Phys. Rev. E*, 74:031123, Sep 2006.
 - [2] Lorenzo Campos Venuti and Paolo Zanardi. Quantum critical scaling of the geometric tensors. *Physical review letters*, 99(9):095701, 2007.
 - [3] Paolo Zanardi, Lorenzo Campos Venuti, and Paolo Giorda. Bures metric over thermal state manifolds and quantum criticality. *Phys. Rev. A*, 76:062318, Dec 2007.
 - [4] Ali T Rezakhani, Damian F Abasto, Daniel A Lidar, and Paolo Zanardi. Intrinsic geometry of quantum adiabatic evolution and quantum phase transitions. *Physical Review A*, 82(1):012321, 2010.
 - [5] Nikola Paunković and Vitor Rocha Vieira. Macroscopic distinguishability between quantum states defining different phases of matter: Fidelity and the Uhlmann geometric phase. *Physical Review E*, 77(1):011129, 2008.
 - [6] JP Provost and G Vallee. Riemannian structure on manifolds of quantum states. *Communications in Mathematical Physics*, 76(3):289–301, 1980.
 - [7] Bruno Mera, Chrysoula Vlachou, Nikola Paunković, and Vítor R. Vieira. Uhlmann connection in fermionic systems undergoing phase transitions. *Phys. Rev. Lett.*, 119:015702, Jul 2017.
 - [8] Bruno Mera, Chrysoula Vlachou, Nikola Paunković, Vítor R. Vieira, and Oscar Viyuela. Dynamical phase transitions at finite temperature from fidelity and interferometric Loschmidt echo induced metrics. *Phys. Rev. B*, 97:094110, Mar 2018.
 - [9] Syed Tahir Amin, Bruno Mera, Chrysoula Vlachou, Nikola Paunković, and Vítor R. Vieira. Fidelity and Uhlmann connection analysis of topological phase transitions in two dimensions. *Phys. Rev. B*, 98:245141, Dec 2018.
 - [10] Sebastiano Peotta and Päivi Törmä. Superfluidity in topologically nontrivial flat bands. *Nature communications*, 6(1):1–9, 2015.
 - [11] Aleksu Julku, Sebastiano Peotta, Tuomas I Vanhala, Dong-Hee Kim, and Päivi Törmä. Geometric origin of superfluidity in the Lieb-lattice flat band. *Phys. Rev.*

- Lett.*, 117(4):045303, 2016.
- [12] Long Liang, Tuomas I Vanhala, Sebastiano Peotta, Topi Siro, Ari Harju, and Päivi Törmä. Band geometry, Berry curvature, and superfluid weight. *Phys. Rev. B*, 95(2):024515, 2017.
- [13] M Iskin. Quantum-metric contribution to the pair mass in spin-orbit-coupled Fermi superfluids. *Phys. Rev. A*, 97(3):033625, 2018.
- [14] Siddharth A Parameswaran, Rahul Roy, and Shivaji L Sondhi. Fractional quantum Hall physics in topological flat bands. *Comptes Rendus Physique*, 14(9-10):816–839, 2013.
- [15] Rahul Roy. Band geometry of fractional topological insulators. *Phys. Rev. B*, 90(16):165139, 2014.
- [16] Martin Claassen, Ching Hua Lee, Ronny Thomale, Xiao-Liang Qi, and Thomas P Devereaux. Position-momentum duality and fractional quantum Hall effect in Chern insulators. *Phys. Rev. Lett.*, 114(23):236802, 2015.
- [17] Thomas S Jackson, Gunnar Möller, and Rahul Roy. Geometric stability of topological lattice phases. *Nature communications*, 6(1):1–11, 2015.
- [18] Ching Hua Lee, Martin Claassen, and Ronny Thomale. Band structure engineering of ideal fractional Chern insulators. *Phys. Rev. B*, 96:165150, Oct 2017.
- [19] Jie Wang, Jennifer Cano, Andrew J Millis, Zhao Liu, and Bo Yang. Exact Landau Level Description of Geometry and Interaction in a Flatband. *arXiv preprint arXiv:2105.07491*, 2021.
- [20] Gabriel E Topp, Christian J Eckhardt, Dante M Kennes, Michael A Sentef, and Päivi Törmä. Light-matter coupling and quantum geometry in moiré materials. *arXiv preprint arXiv:2103.04967*, 2021.
- [21] Nicola Marzari and David Vanderbilt. Maximally localized generalized Wannier functions for composite energy bands. *Phys. Rev. B*, 56(20):12847, 1997.
- [22] Tomoki Ozawa and Nathan Goldman. Extracting the quantum metric tensor through periodic driving. *Phys. Rev. B*, 97(20):201117, 2018.
- [23] Giandomenico Palumbo and Nathan Goldman. Tensor Berry connections and their topological invariants. *Phys. Rev. B*, 99(4):045154, 2019.
- [24] Grazia Salerno, Nathan Goldman, and Giandomenico Palumbo. Floquet-engineering of nodal rings and nodal spheres and their characterization using the quantum metric. *Phys. Rev. Res.*, 2(1):013224, 2020.
- [25] Tomoki Ozawa and Nathan Goldman. Probing localization and quantum geometry by spectroscopy. *Phys. Rev. Res.*, 1(3):032019, 2019.
- [26] Bruno Mera. Localization anisotropy and complex geometry in two-dimensional insulators. *Phys. Rev. B*, 101:115128, Mar 2020.
- [27] Tomoki Ozawa and Bruno Mera. Relations between topology and the quantum metric for Chern insulators. *Phys. Rev. B*, 104:045103, Jul 2021.
- [28] Bruno Mera and Tomoki Ozawa. Kähler geometry and Chern insulators: Relations between topology and the quantum metric. *Phys. Rev. B*, 104:045104, Jul 2021.
- [29] Bruno Mera and Tomoki Ozawa. Engineering geometrically flat Chern bands with Fubini-Study Kähler structure. *Phys. Rev. B*, 104:115160, Sep 2021.
- [30] Jie Wang and Zhao Liu. Hierarchy of ideal flatbands in chiral twisted multilayer graphene models. *arXiv preprint arXiv:2109.10325*, 2021.
- [31] Patrick J. Ledwith, Ashvin Vishwanath, and Eslam Khalaf. A family of ideal Chern flat bands with arbitrary Chern number in chiral twisted graphene multilayers, 2021.
- [32] P. Törmä, S. Peotta, and B. A. Bernevig. Superfluidity and quantum geometry in twisted multilayer systems, 2021.
- [33] Daniel Parker, Patrick Ledwith, Eslam Khalaf, Tomohiro Soejima, Johannes Hauschild, Yonglong Xie, Andrew Pierce, Michael P. Zaletel, Amir Yacoby, and Ashvin Vishwanath. Field-tuned and zero-field fractional Chern insulators in magic angle graphene. *arXiv preprint arXiv:2112.13837*, 2021.
- [34] Christian Northe, Giandomenico Palumbo, Jonathan Sturm, Christian Tutschku, and Ewelina M. Hankiewicz. Interplay of band geometry and topology in ideal Chern insulators in presence of external electromagnetic fields. *arXiv preprint arXiv:2112.15559*, 2021.
- [35] Bruno Mera, Anwei Zhang, and Nathan Goldman. Relating the topology of Dirac Hamiltonians to quantum geometry: When the quantum metric dictates Chern numbers and winding numbers. *arXiv preprint arXiv:2106.00800*, 2021.
- [36] David Tong. Lectures on statistical field theory, 2017.
- [37] Masamichi Miyaji, Tokiro Numasawa, Noburo Shiba, Tadashi Takayanagi, and Kento Watanabe. Distance between quantum states and gauge-gravity duality. *Phys. Rev. Lett.*, 115:261602, Dec 2015.
- [38] F Duncan M Haldane. Model for a quantum Hall effect without Landau levels: Condensed-matter realization of the "parity anomaly". *Physical Review Letters*, 61(18):2015, 1988.
- [39] Hector Silva, Bruno Mera, and Nikola Paunković. Interferometric geometry from symmetry-broken Uhlmann gauge group with applications to topological phase transitions. *Physical Review B*, 103(8):085127, 2021.