

An approximate Bayes factor based high dimensional MANOVA using Random Projections

Roger S. Zoh

January 6, 2022

Abstract

High-dimensional mean vector testing problem for two or more groups remain a very active research area. In these setting, traditional tests are not applicable because they involve the inversion of rank deficient group covariance matrix. In current approaches, this problem is addressed by simply looking at a test assuming a sparse or diagonal covariance matrix potentially ignoring complex dependency between features. In this paper, we develop a Bayes factor (BF) based testing procedure for comparing two or more population means in (very) high dimensional settings. Two versions of the Bayes factor based test statistics are considered which are based on a Random projection (RP) approach. RPs are appealing since they make not assumption about the form of the dependency across features in the data. The final test statistic is based on an ensemble of Bayes factors corresponding to multiple replications of randomly projected data. Both proposed test statistics are compared through a battery of simulation settings. Finally they are applied to the analysis of a publicly available genomic single cell RNA-seq (scRNA-seq) dataset.

1 Introduction

The problem of comparing multiple group means continue to receive considerable attentions in literature, especially in the ‘large-p-small-n’ setting where $p \gg n$. For the two (or sample) samples testing problem, the approaches proposed in literature all center around a version of the Hotelling’s T^2 statistic. Namely the statistic used is

$$T^2 = C_n(\bar{\mathbf{X}} - \bar{\mathbf{Y}})^T \mathbf{S}^{-1}(\bar{\mathbf{X}} - \bar{\mathbf{Y}}), \quad (1)$$

where C_n is free a data free quantity; \mathbf{S} is the (pooled) sample covariance; the sample mean vectors are $\bar{\mathbf{Y}}$ and $\bar{\mathbf{X}}$. Unfortunately, in its original (1), the T^2 statistic can be quickly become ill formed since it involves the inversion of a sample covariance matrix that is not positive definite when the dimension of the vector exceeds or supersedes the combined sample sizes. Various approaches exists in literature to help circumvent these limitations. Solutions to that problem have centered around the following approaches. One approach ignores dependency between the features or groups of features. This has the direct effect of removing the issue of inverting ill-formed covariance matrices. References of these approaches can be found in (Bai and Saranadasa, 1996, Chen and Qin, 2010, Ahmad, 2014, Feng et al., 2017) among others. A second approach can be viewed as a regularization scheme with a goal of making the sample covariance invertible. Two regularization schemes have emerged (Hu et al., 2020). One regularization scheme uses a ridge type estimator for the sample covariance matrix (see Chen et al., 2011, Li et al., 2020). Another regularization approach, which is in principle closer to a data perturbation approach than a regularization approach as we commonly know it, is based on a random projection approach. This approach works by projecting the originally high dimensional data to a low-dimensional embedding and perform the test in this lower dimension data, completely eliminating the need to inverse a rank degenerate sample covariance matrix. Reference of paper using this approach in the two group testing setting are Lopes et al. (2011); Thulin (2014); Srivastava et al. (2016) in the frequentist setting and Zoh et al. (2018) in the Bayesian setting. Recently, there is a growing effort towards combining these two approach in the two-group mean testing problem (Hu et al., 2020).

Two-sample mean testing problem in high dimension setting is a special case of the more general MANOVA (Multivariate Analysis of Variance) problem. However, extending two group mean testing procedures the case of multiple group testing is no trivial task (Cai and Xia, 2014). Suppose K populations of dimension p , mean vector specified respectively as $\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_K$, and common covariance matrix $\boldsymbol{\Sigma}$. In the MANOVA testing problem is formulated as:

$$H_0 : \boldsymbol{\mu}_i = \boldsymbol{\mu}_j \quad \forall (i, j) \in \mathcal{P} \quad \text{versus} \quad H_1 : \exists (i, j) \in \mathcal{P} \text{ s.t } \boldsymbol{\mu}_i \neq \boldsymbol{\mu}_j \in \mathcal{P} \quad (2)$$

where $\mathcal{P} = \{(i, j) : 1 \leq i < j \leq p\}$. Early references on the more general (more than two groups) MANOVA approach when p exceeds the sample sizes in literature can be found in the work of Dempster (1958); Dempster (1960). In general, the approaches presented in literature center around two major assumptions. One approach derives the test under the assumption of common variances across groups (Fujikoshi et al. (2004)). Another approach removes the assumption of common co-

variances (Srivastava, 2007).

To our knowledge, random projections have not been considered for testing multiple group mean vector in high-dimension. Recently, random matrix approaches in general and random projections (RP) in particular have emerged as effective (linear) data reduction techniques which have been used in many fields. See Wan et al., 2020; López-Sánchez et al., 2021 to list just a few. Additionally, RP have already been proven very successful in the two-sample group tests Lopes et al. (2011); Srivastava et al. (2016); Zoh et al. (2018). However, to the best of our knowledge, they have not been used or evaluated in the multiple (MANOVA) groups mean testing problem. The goal of this paper is to investigate the performance of RPs in the Bayes factor based test in the MANOVA setting. The paper is structure as follows. In section 2, we derive the Bayes factor based tests; section 3 provides some theoretical results of our test along with some simulation results. In section 4, we apply the proposed method to the analysis of an actual data set from single cell sequencing (scRNA-seq). We end with some concluding remarks in section 5.

2 Bayes factor based tests

Here, we describe our approach. Suppose the following data generating model. $\mathbf{X}_{ig} = \boldsymbol{\mu}_i + \boldsymbol{\epsilon}_{ig}$, $g = 1, \dots, G$. Note here that we assume that $G \geq 2$ and $\boldsymbol{\epsilon}_{ig} \stackrel{iid}{\sim} \text{MVN}_p(\mathbf{0}, \boldsymbol{\Sigma})$; MVN_p denotes a multivariate-normal distribution with dimension p . Suppose the data matrices are observed (independently) for each of $G > 1$ groups as $\mathbf{X}_1 \in \mathbb{R}^{n_1 \times p}, \dots, \mathbf{X}_G \in \mathbb{R}^{n_G \times p}$. Let's denote $\mathcal{P} = \{(i, j) : 1 \leq i < j \leq G\}$ the collection all unique group indices. Let's $\boldsymbol{\delta}_{ij} = \boldsymbol{\mu}_i - \boldsymbol{\mu}_j$, the compound hypothesis in (2) is expressed as

$$\begin{aligned}
 H_0 : \boldsymbol{\delta}_{ij} &= \mathbf{0} \quad \forall (i, j) \in \mathcal{P} \\
 &\text{vs} \\
 H_1 : \exists (i, j) \in \mathcal{P} \text{ s.t. } \boldsymbol{\delta}_{ij} &\sim \text{MVN}_p(\mathbf{0}, \boldsymbol{\Sigma}/\tau_{0,ij}) \text{ and } \boldsymbol{\Sigma} \propto |\boldsymbol{\Sigma}|^{-(p+1)/2}
 \end{aligned} \tag{3}$$

which is equivalent to performing $G(G - 1)/2$ pairwise comparisons similar to the approach taken in Tony Cai et al. (2014) and Ahmad (2014). Although the tests proposed in literature almost all adopt a component wise (vector entry) test, here we approach this testing problem while preserving (or minimally disturbing) the dependencies between the vector coordinates. To that end our test uses the Bayes Factor approach and we discuss in detail the two tests we consider.

2.1 Bayes factor Based on Pooled covariance matrix (BF^P)

For the case when $G = 2$ in high-dimensional settings $p \gg n_1 + n_2$, Zoh et al. (2018) proposed a Bayes Factor based test using the idea of Random projection (RP). Namely, in a two-group setting, the BF based test using RP matrix Φ is defined as

$$BF_{10}(\Phi) = (1 + \eta_{12})^{-m/2} \left\{ \frac{1 + \frac{mf}{(1+\eta_{12})(N-m-G+1)}}{1 + \frac{pf}{(N-m-G+1)}} \right\}^{-(N-1)/2}, \quad (4)$$

where τ_0 is the scale parameter for the prior under the alternative; $\eta_{12} = n_{0,12}/\tau_0$ and $f_{1,2} = \frac{N-m-G+1}{(N-G)m} n_{0,12} (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2)^T \Phi (\Phi^T \mathbf{S}_p \Phi^T)^{-1} \Phi^T (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2)$; $N = n_1 + n_2$; $1/n_{0,12} = 1/n_1 + 1/n_2$; $\bar{\mathbf{X}}_g$ is group g sample mean; \mathbf{S}_p is the $p \times p$ pooled sample covariance matrix; $\Phi \in \mathbb{R}^{p \times m}$ is the projection matrix and m is the lower dimensional projection space chosen so that $m < n_1 + n_2 - 2 \ll p$. Zoh et al. (2018) also proposed an automatic approach to select m , τ_0 . Note here that m depends on n_1 and n_2 but is totally independent of p . Similarly, to test the (complex) hypothesis in (3), we can use the following test statistic:

$$BF_{10}^p = (1 + \eta_{ij})^{-p/2} \left\{ \frac{1 + \frac{pf^{max}}{(1+\eta_{ij})(N-p-(G-1))}}{1 + \frac{pf^{max}}{(N-p-(G-1))}} \right\}^{-(N-1)/2}, \quad (5)$$

where $f^{max} = \underset{(i,j)}{\operatorname{argmax}} f_{ij}$ is the maximum over all the pairwise f_{ij} (data dependent) statistics; $N = \sum_{g=1}^G n_g$. Equivalently, $\eta_{ij} = n_{0,ij}/\tau_{0,ij}$ and is based on the pair that yields the highest \mathbf{F} (f_{ij}) statistic. Note that \mathbf{S}_p is the pooled (all the groups) sample covariances. It is important to note that under the data model, we have that $\forall (i, j) \in \mathcal{P}, f_{ij} \stackrel{id}{\sim} \mathbf{F}_{m, N-m-(G-1)}$ (identically but not independently distributed) when the NULL hypothesis is true. The lack of independence renders the derivation of the NULL distribution (or quantiles of the NULL distribution) of f^{max} difficult. We defer the discussion about the choice of m and $\tau_{0,ij}$ to later.

2.2 Bayes factor Based on paired covariance matrix (BF^I)

The Bayes factor proposed in Equation 5 relies on a pooled single covariance matrix \mathbf{S}_p implying similar covariance matrices across all groups. The assumption of common covariance matrix across groups can reveal very powerful as it allows to borrow information across for amore precise estimate of the common covariance matrix Σ .

However, it can also be detrimental if grossly wrong. We relax that assumption by instead using a pooled pairwise covariance matrices, which is a less stringent assumption than an assumption of covariance matrices. We then get

$$BF_{10}^I = (1 + \eta_{ij})^{-m/2} \left\{ \frac{1 + \frac{mf^{max}}{(1+\eta_{ij})(N_{ij}-m-1)}}{1 + \frac{mf^{max}}{(N_{ij}-m-1)}} \right\}^{-(N_{ij}-1)/2}, \quad (6)$$

where $f^{max} = \underset{(i,j)}{\operatorname{argmax}} f_{ij}$; When $m < \underset{(i,j)}{\operatorname{argmin}} N_{ij} - 2 \ll p$, again $f_{i,j} = \frac{(N_{ij}-m-1)}{(N_{ij}-2)m} n_{0,ij} (\overline{\mathbf{X}}_i - \overline{\mathbf{X}}_j)^T \Phi (\Phi^T \mathbf{S}_{p,ij} \Phi)^{-1} \Phi^T (\overline{\mathbf{X}}_i - \overline{\mathbf{X}}_j)$; $N_{ij} = n_i + n_j$; $\mathbf{S}_{p,ij}$ is the sample covariance for both group i and j . Note that $\forall (i, j) \in \mathcal{P}$, $f_{ij} \stackrel{id}{\sim} \mathbf{F}_{m, n_i + n_j - m - 1}$ (identically but not independently distributed). Again, similarly to the observation made above, here also the lack of independence renders the derivation of the NULL distribution (or quantiles of the NULL distribution) of f^{max} difficult.

2.3 Ensemble test

Based on the BF statistics in (5) and (6), we will opt in favor of the alternative if the BF (maximum BF) exceeds a given evidence threshold γ (see Kass and Raftery, 1995). We provide a way to objectively choose γ later. A test based on a single RP Φ can hugely depend on the choice of that single RP matrix. Instead, we based our final decision on multiple RPs and the final decision is reached based on an ensemble test. Hence, for N chosen sufficiently large, our final test statistic is obtained as

$$\tilde{\psi}_N = \frac{1}{N} \sum_{i=1}^N \mathbf{1}\{BF_{10}^*(\Phi_i) \geq \gamma\}; \quad (7)$$

where $\mathbf{1}\{A\}$ is the indicator function which equals 1 if A is true and zero otherwise; BF_{10}^* is the either the BF BF_{10}^P or BF_{10}^I . Note here that γ depends on n . The NULL distribution of the test statistics (7) is difficult to derive analytically but under our assumed data generating model, it is (cheaply) approximated. Additionally, the NULL distribution of the test statistics is invariant under an arbitrary common mean vector and common (unknown) covariance matrix Σ . Provided the NULL distribution our test statistic $\tilde{\psi}^0$, the decision is made as follows:

$$\begin{cases} \text{Reject } H_0 & \text{if } \tilde{\psi}(N) > \tilde{\psi}_\alpha^0, \\ \text{Accept } H_0 & \text{Otherwise,} \end{cases} \quad (8)$$

Here $\tilde{\psi}_\alpha^0$ denotes the upper α percentile of the NULL distribution. We show that the test is unbiased and its power converges to 1 with increasing sample sizes. We formalize these results in the theory section below (Section 3).

2.4 Choices of m , $\tau_{0,ij}$ and γ

We obtain values for m , $\tau_{0,ij}$, and γ using the idea of restricted most powerful Bayesian test (RMPBT) of Goddard and Johnson, 2016; Goddard, 2015. To find the RMPBT, we would like to choose the parameters for the alternative (parameters of the distribution under the alternative) so to maximize the probability of rejecting the NULL under all possible data generating distribution. Namely, we select $\tau_{0,ij}$ so that

$$Pr\{BF_{10}^{max}(\tau_0) \geq \gamma\} \geq Pr\{BF_{20}^{max}(\tau_1) \geq \gamma\},$$

for two different choices of the prior parameters under the alternative. This is equivalent to choosing $\tau_{0,ij}$ so that $P\{f^{max} > f_0^{max}(\tau_{0,ij})\}$ is maximized. It is easy to see that we maximize that probability if $f_0^{max}(\tau_{0,ij})$ is minimized over all possible values of $\tau_{0,ij}$. Note that $f_0^{max}(\tau_{0,ij}) = \frac{1+\eta_{ij}}{\eta_{ij}} \left\{ \frac{(N-m-G+1)C_{ij}}{m(1-C_{ij})} \right\}$, where $C_{ij} = \frac{1+\eta_{ij}}{\eta_{ij}} \{1 - \{\gamma(1 + \eta_{ij})^{m/2}\}^{-2/(N-1)}\}$ (for Bayes Factor in 5) and $C_{ij} = \frac{1+\eta_{ij}}{\eta_{ij}} \{1 - \{\gamma(1 + \eta_{ij})^{m/2}\}^{-2/(n_i+n_j)}\}$ and $N_{ij} = n_i + n_j$ (for Bayes Factor in 6); and $\eta_{ij} = n_{0,ij}/\tau_{0,ij}$. Note that $f_0^{max}(\tau_{0,ij})$ also depends on m , the dimension of the projection space. Noting that for each BF based test, f_{ij} for all $1 \leq i < j \leq G$ have a marginal F distribution but a complex joint distribution. Then we have that for a chosen value of α if $f_0^{max}(\tau_{0,ij})$ is the upper α percentile of the distribution of f^{max} , then $F_{m,N-m-(G-1)}(\alpha) \leq f_0^{max}(\tau_{0,ij})$, for the BF in (5). We can use that fact to obtain an approximate value of m in both cases as:

$$\arg \min_m F_{m,N-m-(G-1)}(\alpha) \text{ or } \min_m \left\{ m : \forall (i, j) \in \mathcal{P}, \arg \min_m F_{m,n_i+n_j-m-1}(\alpha) \right\}, \quad (9)$$

for the BF in (5) and (6) respectively; $F_{a,b}(\theta)$ is the upper $\theta \in (0, 0.5)$ percentile of a F distribution with a and b degrees of freedom. We discuss the choice of α shortly. Next, given n_1, n_2, \dots, n_G and m , the null distribution of f^{max} can simply be approximated under the assumption of common covariance matrix using a monte carlo approach. Next we obtain, $\tau_{0,ij}$ as

$$\tau_{0,ij}(n) = \frac{n_{0,ij}}{f^{max,0}(\alpha) - 1}, \quad (10)$$

for a significance level α ; $1/n_{0,ij} = 1/n_i + 1/n_j$. Recall $f^{max,0}(\alpha)$ denotes the upper α percentile of a null distribution of f^{max} . The choice the $f^{max,0}(\alpha)$ allows us to easily compare our proposed test with a non-Bayesian competitor using a chosen significance level α . Additionally, we can choose the threshold as follows $\gamma_{ij,\alpha} = \{1 + \eta_{ij}\}^{-m/2} \left\{1 - \frac{\eta_{ij}}{1+\eta_{ij}} C_{ij}\right\}^{-(N-1)/2}$ (for Bayes Factor in 5) and $\gamma_{ij,\alpha} = \{1 + \eta_{ij}\}^{-m/2} \left\{1 - \frac{\eta_{ij}}{1+\eta_{ij}} C_{ij}\right\}^{-(n_i+n_j-1)/2}$ (for Bayes Factor in 6). Finally, the RP matrices are chosen orthogonal matrices so that for a given RP matrix $\Phi \in \mathbb{R}^{p \times m}$, $\Phi^T \Phi = \mathbf{I}_m$. We use the sparse and dense version of the RP matrices proposed by Srivastava et al. (2016). Additionally, the normalization step can be completely skipped based on the results of the QR factorization of a matrix (see Zoh et al., 2018 for more details).

3 Theoretical justifications and Simulation Results

3.1 Theoretical Justifications

We assume the following conditions.

Assumption 1 $n_g / \sum_{g=1}^G n_g \rightarrow \theta \in (0, 1)$

Assumption 2 $G / \min(n_1, \dots, n_G) \rightarrow 0$ as

Assumption 3 $\max\{n_1, n_2, \dots, n_G\} / p \rightarrow \infty$ as

Assumption 4 We denote a m_n , $\tau_{ij,n}$, and $\gamma_{ij,n}$ the values of m , $\tau_{0,ij}$ and γ we derived in session 2.4. Further, we assume that $m_n/n \rightarrow \theta \in (0, 1)$ so that $m_n n_{0,n} / \tau_{0,ij,n} \rightarrow \infty$ as $n_{\min} \rightarrow \infty$. These conditions are satisfied by our construction (see Appendix C of Zoh et al., 2018).

3.1.1 Consistency of BF^P

Theorem 1. Suppose that $X_{ig} = \boldsymbol{\mu}_i + \boldsymbol{\epsilon}_{ig}$, where $\boldsymbol{\epsilon}_{ig} \sim MVN_p(\mathbf{0}, \boldsymbol{\Sigma})$, for $g = 1, \dots, G$ independent groups and $p \gg \max\{n_1, \dots, n_G\}$, where n_1, \dots, n_G are the respective sample sizes. Suppose $1 \leq m \leq \min\{n_1, \dots, n_G\}$ and $n = \sum_{g=1}^G n_g$. Let's $BF_{max}^P = \max\{BF_{12}^P, \dots, BF_{(G-1)G}^P\}$, where BF_{ij}^P is the BF based on the group i and j data. As $\min\{n_1, \dots, n_G\} \rightarrow \infty$,

1. If m is fixed, then $\log(BF_{max}^P) \rightarrow -\infty$ under H_0 and $\log(BF_{max}^P) \rightarrow \infty$ under H_1 .

2. If $m(n)$ so that $m(n)/n \rightarrow \theta \in (0, 1)$, then $\log(BF_{max}^P) = \mathcal{O}(n)$ under H_0 and $\log(BF_{max}^P) \rightarrow \infty$ under as sequence of alternative H_{1n} .

Proof: Part(1) For $1 < m < \min\{n_1, \dots, n_G\}$ and $n = \sum_g^G n_g$, we integrate out the parameters with respect to the conjugate priors to obtain the Bayes factor in favor of the alternative as

$$BF_{ij}^P(\Phi) = (1 + \eta)^{-m/2} \left\{ 1 - \frac{mf_{ij}^* \eta / (1 + \eta)}{mf_{ij}^* + n - m - (G - 1)} \right\}^{-(n-1)/2},$$

where

$$f_{ij}^* = \frac{n - m - (G - 1)}{(n - G)m} n_{0,ij} (\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_j)^T \Phi (\Phi \mathbf{S} \Phi^T)^{-1} \Phi^T (\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_j).$$

where

$$\mathbf{S} = \frac{1}{n - G} \sum_{g=1}^G (n_g - 1) \mathbf{S}_g \text{ and } \mathbf{S}_g = \frac{1}{n_g - 1} \sum_{i=1}^{n_g} (\mathbf{X}_{ig} - \bar{\mathbf{X}}_g)(\mathbf{X}_{ig} - \bar{\mathbf{X}}_g)^T.$$

Recall that $1/n_{0,ij} = 1/n_i + 1/n_j$, $\eta_{ij} = n_{0,ij}/\tau_{0,ij}$, and $n_{\min} = \min\{n_1, \dots, n_G\}$. Since $\tau_{0,ij}$ is fixed, $\eta_{ij} \rightarrow \infty$ as $n_{\min} \rightarrow \infty$. For a randomly chosen Φ , under H_0 , $f_{ij}^* \sim F_{m, n-m-(G-1)}$ with m and $n - m - (G - 1)$ degrees of freedom. Thus, $f_{ij}^* = O_p(1)$ and $f_{ij}^{max} = \max_{ij} \{f_{ij}^*\}$ for all $1 \leq i, j \leq G$. Also, from well-known properties of the F distribution, we have that

$$U_{ij} = \frac{mf_{ij}^*/(n - m - (G - 1))}{\{mf_{ij}^*/(n - m - (G - 1))\}} = \frac{mf_{ij}^*}{(mf_{ij}^* + n - m - (G - 1))} \sim \text{Beta}\{m/2, (n - m - (G - 1))/2\},$$

for each ij with $1 \leq i, j \leq G$, where $\text{Beta}(a, b)$ denotes a Beta distribution. Therefore, $\{\eta_{ij}/(1 + \eta_{ij})\} U_{ij} = O_p(1)$. Hence, $\log\{1 - \eta_{ij} U_{ij}/(1 + \eta_{ij})\} = O_p(1)$ as $n_{\min} \rightarrow \infty$. We then get

$$-\frac{m}{2n} \log(1 + \eta_{ij}) - \frac{(n - 1)}{2n} \log\{1 - \eta_{ij} U_{ij}/(1 + \eta_{ij})\} \xrightarrow{p} -\infty,$$

since $\log(1 + \eta_{ij}) \rightarrow \infty$ as $n_{\min} \rightarrow \infty$ and $\lim_{n_{\min} \rightarrow \infty} m/n = \theta \in (0, 1)$. We conclude that $\log\{BF_{ij}^P(\Phi)\} \xrightarrow{p} -\infty$ under the null hypothesis for all $1 \leq i < j \leq G$. Hence $\log\{BF_{10}^P(\Phi)\} \xrightarrow{p} -\infty$.

Under the alternative, $\boldsymbol{\mu}_i \neq \boldsymbol{\mu}_j$ and $\boldsymbol{\delta}_{ij} \sim \mathbf{N}_p(\mathbf{0}, \boldsymbol{\Sigma}/\tau_0)$. Then, $f_{ij}^* \mid \lambda_{ij} \sim F_{m, n-m-(G-1)}(\lambda_{ij})$ with non-centrality $\lambda_{ij} = n_{0,ij} \boldsymbol{\delta}_{ij}^T \boldsymbol{\Phi} (\boldsymbol{\Phi}^T \boldsymbol{\Sigma} \boldsymbol{\Phi})^{-1} \boldsymbol{\Phi}^T \boldsymbol{\delta}_{ij}$. Since $\boldsymbol{\delta}_{ij} \sim \mathbf{N}_p(\mathbf{0}, \boldsymbol{\Sigma}/\tau_0)$, $\lambda \sim n_{0,ij} \chi_m^2/\tau_0$, where χ_m^2 denotes a χ^2 distribution with m degrees of freedom. The non-centrality parameter depends on n through $n_{0,ij}$. We can show that the unconditional distribution of $f_{ij}^*/(1+\eta_{ij}) \sim F_{m, n-m-(G-1)}$ (see Johnson, 2005, page 704). If we denote $f_{ij}^0 = f_{ij}^*/(1+\eta_{ij})$, we have $f_{ij}^0 = O_p(1)$, and $mf_{ij}^0/n = O_p(1)$, as $n_{\min} \rightarrow \infty$. We have that

$$\frac{\eta_{ij} U_{ij}}{(1+\eta_{ij})} = \frac{mf_{ij}^0 \eta_{ij}}{mf_{ij}^0(1+\eta_{ij}) + n - m - (G-1)}$$

From the above equation, we get

$$-\log \left\{ 1 - \frac{\eta_{ij} U_{ij}}{(1+\eta_{ij})} \right\} = \log \left\{ \frac{mf_{ij}^0(1+\eta_{ij})/(n-m-(G-1)) + 1}{\{mf_{ij}^0/(n-m-(G-1))\} + 1} \right\}.$$

Since $f_{ij}^0 = O_p(1)$, and $m/(n-m-1)$ converges, we have

$$-\log \left\{ 1 - \frac{\eta_{ij} U_{ij}}{(1+\eta_{ij})} \right\} \xrightarrow{p} \infty.$$

Since this is true for all (i, j) with $\boldsymbol{\mu}_i \neq \boldsymbol{\mu}_j$ with $1 \leq i < j \leq G$, we conclude that $\log \{BF_{10}^P(\boldsymbol{\Phi})\} \xrightarrow{p} \infty$, under the alternative hypothesis.

Part(2) We now assume that $\eta_{ij} \rightarrow 0$ and $m_n \eta_{ij} \rightarrow \infty$. We have

$$\log \{BF_{10}^P(\boldsymbol{\Phi})\} = \frac{n}{2} \left(1 - \frac{m_n}{n} \right) \log(1+\eta_{ij}) - \frac{n}{2} \log \{1 + \eta_{ij}(1 - U_{ij})\} + \frac{1}{2} \log \left\{ 1 - \frac{\eta_{ij} U_{ij}}{1 + \eta_{ij}} \right\},$$

where $U_{ij} \sim \text{Beta} \{m/2, (n - m_n - (G-1))/2\}$ under H_0 for each $1 \leq i < j \leq G$. For large n , none of the terms with n dominates and their difference converges. The distribution $\log \{BF_{10}(\mathbf{X}^*, \mathbf{Y}^*)\}$ then depends on that of U , which is bounded in probability. Therefore, under H_0 , $\log \{BF_{10,ij}^P(\boldsymbol{\Phi})\} = O_p(1)$ and we conclude $\log \{BF_{10}^P(\boldsymbol{\Phi})\} = O_p(1)$ for each $1 \leq i < j \leq G$.

Under H_1^n , again we have

$$\log \{BF_{10,ij}^P(\boldsymbol{\Phi})\} = -\frac{m}{2} \log(1+\eta) - \frac{(n-1)}{2} \log \left\{ 1 - \frac{\eta U}{1+\eta} \right\},$$

where $U_{ij} \xrightarrow{p} 1$ with $f_{ij}^* \xrightarrow{p} \infty$. Since $\log(1+\eta_{ij})\{(n-1)/2 - m_n/2\} \rightarrow \infty$, we conclude that $\log \{BF_{10,ij}^P(\boldsymbol{\Phi})\} \xrightarrow{p} \infty$ for (i, j) where $\boldsymbol{\mu}_i \neq \boldsymbol{\mu}_j$. Hence $\log \{BF_{10}^P(\boldsymbol{\Phi})\} \xrightarrow{p} \infty$

□

3.1.2 Consistency of BF^I

Theorem 2. Suppose that $X_{ig} = \boldsymbol{\mu}_i + \boldsymbol{\epsilon}_{ig}$, where $\boldsymbol{\epsilon}_{ig} \sim MVN_p(\mathbf{0}, \boldsymbol{\Sigma})$, for $g = 1, \dots, G$ independent groups and $p \gg \max\{n_1, \dots, n_G\}$, where n_1, \dots, n_G are the respective sample sizes. Suppose $1 \leq m \leq \min\{n_1, \dots, n_G\}$ and $n_{ij} = n_i + n_j$. Let's $BF_{max}^P = \max\{BF_{12}^P, \dots, BF_{(G-1)G}^P\}$, where BF_{ij}^P is the BF based on the group i and j data. As $\min\{n_1, \dots, n_G\} \rightarrow \infty$,

1. If m is fixed, then $\log(BF_{max}^I) \rightarrow -\infty$ under H_0 and $\log(BF_{max}^I) \rightarrow \infty$ under H_1
2. If $m(n)$ so that $m(n)/n \rightarrow \theta \in (0, 1)$, then $\log(BF_{max}^I) = O_p(n)$ under H_0 and $\log(BF_{max}^I) = \infty$ under a sequence of alternative H_{1n} .

Proof: Part(1) For $1 < m < \min\{n_1, \dots, n_G\}$ and $n = n_i + n_j$, we integrate out the parameters with respect to the conjugate priors to obtain the Bayes factor in favor of the alternative as

$$BF_{10,ij}^I(\boldsymbol{\Phi}) = (1 + \eta_{ij})^{-m/2} \left\{ 1 - \frac{mf_{ij}^* \eta_{ij} / (1 + \eta_{ij})}{mf_{ij}^* + n_{ij} - m - 1} \right\}^{-(n_{ij}-1)/2},$$

where

$$f_{ij}^* = \frac{n_{ij} - m - 1}{(n_{ij} - 2)m} n_{0,ij} (\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_j)^\top \boldsymbol{\Phi} (\boldsymbol{\Phi} \mathbf{S}_{ij} \boldsymbol{\Phi}^\top)^{-1} \boldsymbol{\Phi}^\top (\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_j).$$

where

$$\mathbf{S}_{ij} = \frac{1}{n_{ij} - 2} \sum_{g=(i,j)} (n_g - 1) \mathbf{S}_g \text{ and } \mathbf{S}_g = \frac{1}{n_g - 1} \sum_{l=1}^{n_g} (\mathbf{X}_{lg} - \bar{\mathbf{X}}_g)(\mathbf{X}_{lg} - \bar{\mathbf{X}}_g)^\top.$$

Recall that $1/n_{0,ij} = 1/n_i + 1/n_j$, $\eta_{ij} = n_{0,ij}/\tau_{0,ij}$, and $n_{\min} = \min\{n_i, n_j\}$. Since $\tau_{0,ij}$ is fixed, $\eta_{ij} \rightarrow \infty$ as $n_{\min} \rightarrow \infty$. For a randomly chosen $\boldsymbol{\Phi}$, under H_0 , $f_{ij}^* \sim F_{m, n_{ij}-m-1}$ with m and $n_{ij} - m - 1$ degrees of freedom. Thus, $f_{ij}^* = O_p(1)$ and $f_{ij}^{max} = \max_{ij}\{f_{ij}^*\}$ for all $1 \leq i, j \leq G$. Also, from well-known properties of the F distribution, we have that

$$U_{ij} = \frac{mf_{ij}^*/(n_{ij} - m - 1)}{\{mf_{ij}^*/(n_{ij} - m - 1)\}} = \frac{mf_{ij}^*}{(mf_{ij}^* + n_{ij} - m - 1)} \sim \text{Beta}\{m/2, (n_{ij} - m - 1)/2\},$$

for each ij with $1 \leq i, j \leq G$, where $\text{Beta}(a, b)$ denotes a Beta distribution. Therefore, $\{\eta_{ij}/(1+\eta_{ij})\}U_{ij} = O_p(1)$. Hence, $\log\{1 - \eta_{ij}U_{ij}/(1 + \eta_{ij})\} = O_p(1)$ as $n_{\min} \rightarrow \infty$. We then get

$$-\frac{m}{2n_{ij}}\log(1 + \eta_{ij}) - \frac{(n_{ij} - 1)}{2n_{ij}}\log\{1 - \eta_{ij}U_{ij}/(1 + \eta_{ij})\} \xrightarrow{p} -\infty,$$

since $\log(1 + \eta_{ij}) \rightarrow \infty$ as $n_{\min} \rightarrow \infty$ and $\lim_{n_{\min} \rightarrow \infty} m/n_{ij} = \theta \in (0, 1)$. We conclude that $\log\{BF_{ij}^I(\Phi)\} \xrightarrow{p} -\infty$ under the null hypothesis for all $1 \leq i < j \leq G$. Hence $\log\{BF_{10}^I(\Phi)\} \xrightarrow{p} -\infty$.

Under the alternative, $\boldsymbol{\mu}_i \neq \boldsymbol{\mu}_j$ and $\boldsymbol{\delta}_{ij} \sim \mathbf{N}_p(\mathbf{0}, \Sigma/\tau_0)$. Then, $f_{ij}^* \mid \lambda_{ij} \sim F_{m, n_{ij}-m-1}(\lambda_{ij})$ with non-centrality $\lambda_{ij} = n_{0,ij}\boldsymbol{\delta}_{ij}^T \Phi (\Phi^T \Sigma \Phi)^{-1} \Phi^T \boldsymbol{\delta}_{ij}$. Since $\boldsymbol{\delta}_{ij} \sim \mathbf{N}_p(\mathbf{0}, \Sigma/\tau_0)$, $\lambda \sim n_{0,ij}\chi_m^2/\tau_0$, where χ_m^2 denotes a χ^2 distribution with m degrees of freedom. The non-centrality parameter depends on n_{ij} through $n_{0,ij}$. We can show that the unconditional distribution of $f_{ij}^*/(1 + \eta_{ij}) \sim F_{m, n_{ij}-m-1}$ (see Johnson, 2005, page 704). If we denote $f_{ij}^0 = f_{ij}^*/(1 + \eta_{ij})$, we have $f_{ij}^0 = O_p(1)$, and $m f_{ij}^0/n_{ij} = O_p(1)$, as $n_{\min} \rightarrow \infty$. We have that

$$\frac{\eta_{ij}U_{ij}}{(1 + \eta_{ij})} = \frac{m f_{ij}^0 \eta_{ij}}{m f_{ij}^0 (1 + \eta_{ij}) + n_{ij} - m - 1}$$

From the above equation, we get

$$-\log\left\{1 - \frac{\eta_{ij}U_{ij}}{(1 + \eta_{ij})}\right\} = \log\left\{\frac{m f_{ij}^0 (1 + \eta_{ij}) / (n_{ij} - m - 1) + 1}{\{m f_{ij}^0 / (n - m - 1)\} + 1}\right\}.$$

Since $f_{ij}^0 = O_p(1)$, and $m/(n_{ij} - m - 1)$ converges, we have

$$-\log\left\{1 - \frac{\eta_{ij}U_{ij}}{(1 + \eta_{ij})}\right\} \xrightarrow{p} \infty.$$

Since this is true for all (i, j) with $\boldsymbol{\mu}_i \neq \boldsymbol{\mu}_j$ with $1 \leq i < j \leq G$, we conclude that $\log\{BF_{10}^I(\Phi)\} \xrightarrow{p} \infty$, under the alternative hypothesis.

Part(2) We now assume that $\eta_{ij} \rightarrow 0$ and $m_n \eta_{ij} \rightarrow \infty$. We have

$$\log\{BF_{10}^I(\Phi)\} = \frac{n_{ij}}{2} \left(1 - \frac{m_n}{n_{ij}}\right) \log(1 + \eta_{ij}) - \frac{n_{ij}}{2} \log\{1 + \eta_{ij}(1 - U_{ij})\} + \frac{1}{2} \log\left\{1 - \frac{\eta_{ij}U_{ij}}{1 + \eta_{ij}}\right\},$$

where $U_{ij} \sim \text{Beta}\{m/2, (n_{ij} - m_n - (G - 1))/2\}$ under H_0 for each $1 \leq i < j \leq G$. For large n_{ij} , none of the terms with n_{ij} dominates and their difference

converges. The distribution of $\log\{BF_{10}^I(\Phi)\}$ then depends on that of U , which is bounded in probability. Therefore, under H_0 , $\log\{BF_{10,ij}^I(\Phi)\} = \mathcal{O}_p(1)$ and we conclude $\log\{BF_{10}^P(\Phi)\} = \mathcal{O}_p(1)$ for all $1 \leq i < j \leq G$.

Under H_1^n , again we have

$$\log\{BF_{10,ij}^I(\Phi)\} = -\frac{m}{2}\log(1 + \eta) - \frac{(n_{ij} - 1)}{2}\log\left\{1 - \frac{\eta U}{1 + \eta}\right\},$$

where $U_{ij} \xrightarrow{p} 1$ with $f_{ij}^* \xrightarrow{p} \infty$. Since $\log(1 + \eta_{ij})\{(n_{ij} - 1)/2 - m_n/2\} \rightarrow \infty$, we conclude that $\log\{BF_{10,ij}^I(\Phi)\} \xrightarrow{p} \infty$ for (i, j) where $\mu_i \neq \mu_j$. Hence $\log\{BF_{10}^I(\Phi)\} \xrightarrow{p} \infty$

□

3.1.3 Power the test

Theorem 3. *Suppose the assumptions of Theorems 1 and 2 hold. Given a collection Φ_1, \dots, Φ_N of random projections matrices, where $\Phi_i^T \Phi_i = \mathbf{I}$ for all $i = 1, \dots, N$, then $\lim_{n_{\min} \rightarrow \infty} Pr\{\tilde{\psi}(N) > \tilde{\psi}_\alpha^0\} = 1$ under the sequence H_1^n of alternatives.*

Proof: The proof is similar to that of Zoh et al., 2018. The power of our test is $Pr\{\tilde{\psi}(N) > \tilde{\psi}_\alpha^0 \mid H_1^n\}$. Henceforth, we make it explicit that $\tilde{\psi}_\alpha^0$ depends on (n_1, \dots, n_G) and write $\tilde{\psi}_\alpha^0(n_1, \dots, n_G)$ instead.

For given n_1, \dots, n_G and α , we choose $\tilde{\psi}_\alpha^0(n_1, n_2)$ so that $Pr\{\tilde{\psi}(N) > \tilde{\psi}_\alpha^0(n_1, n_2) \mid H_0\} = \alpha$. Since $0 < \tilde{\psi}_\alpha^0(n_1, n_2) < 1$, for $0 < \alpha < 1$, we have that $Pr\{\sum_{i=1}^N \tilde{\psi}(\Phi_i) \geq 0 \mid H_1^n\} \geq Pr\{\sum_{i=1}^N \tilde{\psi}(\Phi_i) > N\phi_\alpha^0(n_1, n_2) \mid H_1^n\} \geq Pr\{\sum_{i=1}^N \tilde{\psi}(\Phi_i) \geq N \mid H_1^n\}$.

We have that $Pr\{\tilde{\psi}(\Phi_i) = 1 \mid H_1^n\} \rightarrow 1$ as $n_{\min} \rightarrow \infty$, under the alternative for $i = 1, \dots, N$. So, $Pr\{\sum_{i=1}^N \phi(\Phi_i) \geq 0 \mid H_1^n\} = 1 - \prod_{i=1}^N Pr\{\phi(\Phi_i) = 0 \mid H_1^n\} \rightarrow 1$. Additionally, $Pr\{\sum_{i=1}^N \tilde{\psi}(\Phi_i) \geq N \mid H_1^n\} = Pr\{\sum_{i=1}^N \tilde{\psi}(\Phi_i) = N \mid H_1^n\} = \prod_{i=1}^N Pr\{\tilde{\psi}(\Phi_i) = 1 \mid H_1^n\} \rightarrow 1$ for fixed N as $n_{\min} \rightarrow \infty$. We conclude that $Pr\{\tilde{\psi}(N) \geq \phi_\alpha^0 \mid H_1^n\} \rightarrow 1$ as $n_{\min} \rightarrow \infty$. □

3.2 Simulation

3.2.1 Simulation Study design

We designed a simulation study aimed at investigating the power of the tests proposed in section 2 with respect to the proportion of true elements of μ that are actually zero (for none all zero vectors) for various choices of covariance matrices and different

proportions of non zeros vectors. Broadly, we consider two settings for our simulation. In each case, we have two conditions for each choice of the covariance matrix. In the first condition, we assume $p = 200$, $G = 3, 5$ and $n_g = 50, \forall 1 \leq g \leq G$. Using the approach described above (Section 2), we find $m = 43$. In the second condition, $p = 1000$, $G = 3, 5$ and $n_g = 70, \forall 1 \leq g \leq G$ and we get $m = 62$. We denote by p_0 the proportion of entries of the vector $\boldsymbol{\delta}$ that are exactly zero. We choose $p_0 = 0.5, .75, .80, 0.95, 0.99, 1.00$ (NULL hypothesis). In each setting, the values of τ_α and γ_α are chosen according to our discussion in Section 2.4. We consider two types of random projections matrices, Φ_1 (full matrix) and Φ_2 (sparse matrix) as described in Srivastava et al., 2016; Zoh et al., 2018. Finally, we assume $\alpha = 0.05$.

In **case 1**, only the last group G has a non-zero mean vector $\boldsymbol{\mu}_G$. But all the others groups have vector mean zero. In **case 2**, only the last group G has a zero vector mean $\boldsymbol{\mu}_G$ and a fixed proportions of the entries of the other group mean vectors are selected to be non-zeros. We consider the following choices of covariance matrix $\boldsymbol{\Sigma} = (\sigma_{ij})$.

1. $\boldsymbol{\Sigma}_1 = \mathbf{I}_{p \times p}$ is the identity matrix.
2. $\boldsymbol{\Sigma}_2$ is block diagonal matrix, with block $\mathbf{B} = 0.85\mathbf{I}_{25 \times 25} + 0.15\mathbf{J}_{25 \times 25}$. \mathbf{J} denotes a matrix with 1 in all of its entries.
3. $\boldsymbol{\Sigma}_3$ is a diagonal matrix where the 20% of the entries of the diagonal elements are $\sigma_j^2 = 0.2p/j$ for $j = 1, \dots, 0.2p$ and the remaining $\sigma_j^2 = 1$ for $j > 0.2p$.
4. $\boldsymbol{\Sigma}$ is an AR(1) covariance matrix with $\sigma_{ij} = \sigma^2 \rho^{|i-j|} \mathbf{1}(|i-j| < 2)$. We chose $\sigma^2 = 1$ and $\rho = 0.4$.
5. $\boldsymbol{\Sigma}_5$ is an AR(1) covariance matrix with $\sigma_{ij} = \sigma^2 \rho^{|i-j|}$. We chose $\sigma^2 = 1$ and $\rho = 0.6$.
6. $\boldsymbol{\Sigma}_6 = \mathbf{D}^{1/2} \{ \mathbf{I}_K \otimes (.2\mathbf{I}_2 + \mathbf{J}_2 0.8) \} \mathbf{D}^{1/2}$, where $\text{diag}(\mathbf{D}) = (d_1, \dots, d_p)^T$ and $d_1, \dots, d_p \sim \text{Uniform}(1, 3)$ and $K = p/2$; \mathbf{I} is the identity matrix and \mathbf{J} a matrix of all ones.

For each case, we also consider two possible alternatives. We simulate the non-all zeros mean vectors as follow

Alt.2: $\boldsymbol{\mu}_g \sim \mathbf{N}_p(\mathbf{1}, \mathbf{I})$, set p_0 randomly selected elements to zero, and re-scale $\boldsymbol{\mu}_g$ so that $\boldsymbol{\mu}_g^T \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_g = 2$

Alt.1: $\boldsymbol{\mu}_g \sim N_p(\mathbf{1}, \mathbf{I})$ and set p_0 of its elements to zero and re-scale $\boldsymbol{\mu}_g$ so that

$$\frac{\|\boldsymbol{\mu}_g\|^2}{\sqrt{\text{trace}(\boldsymbol{\Sigma}^2)}} = 0.1.$$

The two alternatives described above were also considered by Srivastava et al. (2016); Zoh et al. (2018).

3.2.2 Simulation Results

We first look at the performance of the both tests (BF^P and BF^I) in term of their empirical power for simulation **case 1** under alternative 1 (See Tables 1, 2). Overall both tests tended to have an empirical type 1 error estimate around 5%; although in some case the estimated type error seems slight inflated for the case of complex covariance matrices. We note a significant difference between both tests in term of the estimated empirical power. For the same setting, now looking at **case 2**, we observations made in **case1** still hold (see Tables S.1 and S.2 from the Supplemental Material), except we observe a higher estimated power for the test on BF^I . Recall that in case 2, only the last group non-zero mean vector. The test based on the paired groups(BF^I) performs much better when compared to the test based on the pooled covariance for data simulated under the alternative 1. Note that the data were simulated under the common covariance model. However, for data simulated under Alternative 2, again assuming common group covariance matrices, we see that both the tests based on pooled covariance (BF^P) and the test based on the pairwise groups (BF_I) performs very similarly. Although the test based on BF^P tended to have slightly higher power and estimated Type 1 error near 0.05 (see Tables 3 4).

We also look at the performance of both tests for the case of 5 groups (see Tables S.3-S.6 of the Supplemental Material). The observation made for the case of 3 groups also hold for the case of 5.

In the second part of the simulation, we simulate data assuming different covariance matrices between the active group (non-zero) mean vector and the non-active group (all zeros) mean vector. Namely, in **case 1** all groups are assumed to have an identity covariance and the last group G is assumed to have one of the covariance matrix $\boldsymbol{\Sigma}_k$, for each $k = 1, \dots, 6$ (see Table 5). In **case 2**, however, the active group (Group G) has an identity covariance matrix and the non-active groups have the same covariance matrix which is one of the $\boldsymbol{\Sigma}_k$, for each $k = 1, \dots, 6$. Under these settings, we see that the test statistic based on BF^P (pooled covariance) is poorly calibrated when the covariance matrices are (very) different. So we do not discuss its estimated power here. We instead focus on the test based on BF^I (Table 6). The test based on on the BF^I seems to hold the nominal Type I error for the case small

Table 1: Empirical estimates of the power for the test based on BF^I when data are simulated as in case (1) and based on the Alternative 1. Note here cp refers to the true covariance matrix ($G = 3$).

		Φ_1						Φ_2					
cp		1	0.99	0.95	0.8	0.75	0.5	1	0.99	0.95	0.8	0.75	0.5
$n = 50, p = 200$	1	0.033	0.833	0.711	0.675	0.633	0.658	0.033	0.833	0.711	0.675	0.633	0.658
	2	0.034	0.796	0.654	0.666	0.616	0.668	0.034	0.796	0.654	0.666	0.616	0.668
	3	0.038	0.780	0.643	0.610	0.586	0.589	0.038	0.780	0.643	0.610	0.586	0.589
	4	0.049	0.494	0.408	0.396	0.402	0.451	0.049	0.494	0.408	0.396	0.402	0.451
	5	0.064	0.437	0.347	0.343	0.329	0.375	0.064	0.437	0.347	0.343	0.329	0.375
	6	0.073	0.283	0.235	0.224	0.217	0.237	0.073	0.283	0.235	0.224	0.217	0.237
$n = 70, p = 1000$	1	0.021	0.340	0.321	0.305	0.310	0.288	0.021	0.340	0.321	0.305	0.310	0.288
	2	0.031	0.322	0.278	0.286	0.293	0.331	0.031	0.322	0.278	0.286	0.293	0.331
	3	0.062	0.292	0.262	0.237	0.237	0.233	0.062	0.292	0.262	0.237	0.237	0.233
	4	0.036	0.179	0.174	0.157	0.170	0.203	0.036	0.179	0.174	0.157	0.170	0.203
	5	0.065	0.179	0.180	0.151	0.160	0.192	0.065	0.179	0.180	0.151	0.160	0.192
	6	0.047	0.143	0.126	0.104	0.119	0.134	0.047	0.143	0.126	0.104	0.119	0.134

sample size, even though the estimated Type error 1 seems inflated for the covariance matrix very different from the identity matrix.

4 Application

The data set used in our application originated from a head and neck squamous cell carcinoma (HNSCC) study where the profile of 5902 single cells from 18 patients with oral cavity tumors by single cell RNA-seq are obtained. See (Puram et al., 2017) for the detailed description of the study designed. The data set used for our analysis can be downloaded from the Gene expression Omnibus (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE102707>). Subsequently, each of the 5902 cells are identified and labelled. We would like to know if there is evidence that cell types have different gene expression profile while accounting for the potential dependency between the genes. We frame this problem as a high-dimensional mean vector testing. Under the NULL hypothesis, all cell types have equal or similar mean gene expression profile. Before we can apply our proposed test, we perform a feature reduction step to root out lowly expressed genes. Similar to the approach considered by Puram et al. (2017), we choose genes with $E_a(i) > 6$, where $E_a(i) = \log_2(\text{average}(\text{TPM}(i)_{1..k}) + 1)$. This results in $p = 2641$ genes selected. We consider the less popular (Tumor free) cell types (3 types): B-cells (n=138), Macrophage (n=98), and Mast(n=120). To perform our test, we select the

Table 2: Empirical estimates of the power for the test based on BF^P when data are simulated as in case (1) and based on the Alternative 1. Note here cp refers to the true covariance matrix ($G = 3$).

		Φ_1						Φ_2					
cp		1	0.99	0.95	0.8	0.75	0.5	1	0.99	0.95	0.8	0.75	0.5
$n = 50, p = 200$	1	0.034	0.050	0.042	0.033	0.051	0.045	0.034	0.050	0.042	0.033	0.051	0.045
	2	0.031	0.059	0.049	0.051	0.058	0.043	0.031	0.059	0.049	0.051	0.058	0.043
	3	0.038	0.039	0.037	0.035	0.042	0.044	0.038	0.039	0.037	0.035	0.042	0.044
	4	0.049	0.064	0.053	0.067	0.059	0.055	0.049	0.064	0.053	0.067	0.059	0.055
	5	0.063	0.076	0.068	0.074	0.076	0.067	0.063	0.076	0.068	0.074	0.076	0.067
	6	0.056	0.066	0.065	0.064	0.072	0.065	0.056	0.066	0.065	0.064	0.072	0.065
$n = 70, p = 1000$	1	0.024	0.025	0.027	0.026	0.033	0.026	0.024	0.025	0.027	0.026	0.033	0.026
	2	0.051	0.040	0.043	0.036	0.044	0.039	0.051	0.040	0.043	0.036	0.044	0.039
	3	0.065	0.071	0.068	0.050	0.056	0.061	0.065	0.071	0.068	0.050	0.056	0.061
	4	0.043	0.031	0.043	0.030	0.042	0.043	0.043	0.031	0.043	0.030	0.042	0.043
	5	0.065	0.069	0.063	0.066	0.070	0.071	0.065	0.069	0.063	0.066	0.070	0.071
	6	0.049	0.036	0.058	0.053	0.042	0.066	0.049	0.036	0.058	0.053	0.042	0.066

dimension of the projections space, according to our discussion above, to be $m = 88$, and the evidence threshold is set at $\gamma = 6.340$, and $\alpha = 0.05$. Since our test is sensible to significant departure of assumption of common covariance, we perform a test comparing covariance matrices (Ahmad, 2017; Srivastava and Yanagihara, 2010). Interesting enough, the test proposed by Ahmad 2017 reject the NULL hypothesis ($p - value < 0.001$). But the covariance test proposed by Srivastava and Yanagihara (2010) does not reject the NULL ($p - value = 0.1878$). Finally, applying our test to the data set, we reject the NULL hypothesis with a $p - value < 0.001$ based on the NULL hypothesis of the test statistic in (7) approximated assuming zero mean vectors for each group and a common identity covariance matrix. In all, the test based on Bayes factor assuming common over all covariance matrix (BF^P) and the test based on pairwise Bayes factor (BF^I) both yield the same result for the projection matrices Φ_1 and Φ_2 . Our testing procure also provides an automatic way to extract information about all pairwise comparisons since the value of each f_{ij} statistic is retained. A useful summary statistic we can look at is the proportions of f_{ij} statistic that exceeded the threshold of significance across all random projections (see table 7). We note that the test assuming common covariance across all group BF^P tended to have a larger proportion of significant test for all pairwise comparisons when compared to its BF^I counterpart.

Table 3: Empirical estimates of the power for the test based on BF^I when data are simulated as in case (1) and based on the Alternative 2. Note here cp refers to the true covariance matrix ($G = 3$).

		Φ_1						Φ_2					
cp		1	0.99	0.95	0.8	0.75	0.5	1	0.99	0.95	0.8	0.75	0.5
$n = 50, p = 200$	1	0.033	0.570	0.450	0.429	0.408	0.408	0.033	0.570	0.450	0.429	0.408	0.408
	2	0.034	0.790	0.637	0.598	0.537	0.508	0.034	0.790	0.637	0.598	0.537	0.508
	3	0.038	0.989	0.997	1	0.996	0.998	0.038	0.989	0.997	1	0.996	0.998
	4	0.049	0.716	0.603	0.563	0.518	0.518	0.049	0.716	0.603	0.563	0.518	0.518
	5	0.064	0.944	0.843	0.802	0.755	0.706	0.064	0.944	0.843	0.802	0.755	0.706
	6	0.073	0.869	0.770	0.717	0.694	0.680	0.073	0.869	0.770	0.717	0.694	0.680
$n = 70, p = 1000$	1	0.021	0.704	0.655	0.628	0.635	0.606	0.021	0.704	0.655	0.628	0.635	0.606
	2	0.031	0.864	0.830	0.789	0.759	0.716	0.031	0.864	0.830	0.789	0.759	0.716
	3	0.062	1	1	1	1	1	0.062	1	1	1	1	1
	4	0.036	0.818	0.759	0.739	0.730	0.696	0.036	0.818	0.759	0.739	0.730	0.696
	5	0.065	0.943	0.905	0.872	0.862	0.827	0.065	0.943	0.905	0.872	0.862	0.827
	6	0.047	0.899	0.859	0.823	0.818	0.794	0.047	0.899	0.859	0.823	0.818	0.794

5 Conclusion

When the dimension of the variables space exceeds or supersedes the sample sizes in each group, classical test statistic can no longer be directly employed and some regularization steps are needed. Here we consider the problem of multiple group mean vector testing using random projections(RPs). We formulate two test based on Bayes factors which differ by the assumption placed on the covariance matrix. One test statistics assume one overall covariance matrix which results in a Bayes factor we denote BF^P . The second test only assume pairwise common covariance matrix and is denoted as BF^I . When the assumption of homogeneous covariance matrix is reasonable, both test statistics perform similarly and very well. However, for moderate departure from the assumption of common covariance, the test based on BF^I seems robust in the simulation setting we considered. Although, we should note that the test based on BF^I seems to have an inflated estimated type error in some settings. A natural extension of this work is in the case of potentially very different covariance matrices. Additionally, the test statistic was derived assuming a normal distribution, it will be interesting to relax that potentially constraining assumption. Our approach was implemented in the statistical software **Julia**(Bezanson et al. (2012)) and will be made available for used on the author Github page.

Table 4: Empirical estimates of the power for the test based on BF^P when data are simulated as in case 1 and based on the Alternative 2. Note here cp refers to the true covariance matrix ($G = 3$).

		Φ_1						Φ_2						
cp		1	0.99	0.95	0.8	0.75	0.5	1	0.99	0.95	0.8	0.75	0.5	
$n = 50, p = 200$	1	0.034	0.564	0.450	0.438	0.437	0.427	0.034	0.564	0.450	0.438	0.437	0.427	
	2	0.031	0.781	0.674	0.590	0.596	0.530	0.031	0.781	0.674	0.590	0.596	0.530	
	3	0.038	0.986	0.996		1	0.997	0.999	0.038	0.986	0.996	1	0.997	0.999
	4	0.049	0.715	0.612	0.559	0.577	0.532	0.049	0.715	0.612	0.559	0.577	0.532	
	5	0.063	0.944	0.868	0.803	0.788	0.733	0.063	0.944	0.868	0.803	0.788	0.733	
	6	0.056	0.872	0.783	0.731	0.739	0.708	0.056	0.872	0.783	0.731	0.739	0.708	
$n = 70, p = 1000$	1	0.024	0.738	0.662	0.664	0.650	0.659	0.024	0.738	0.662	0.664	0.650	0.659	
	2	0.051	0.894	0.817	0.811	0.812	0.775	0.051	0.894	0.817	0.811	0.812	0.775	
	3	0.065	1	1	1	1	1	0.065	1	1	1	1	1	
	4	0.043	0.839	0.772	0.764	0.776	0.748	0.043	0.839	0.772	0.764	0.776	0.748	
	5	0.065	0.947	0.916	0.889	0.910	0.859	0.065	0.947	0.916	0.889	0.910	0.859	
	6	0.049	0.910	0.864	0.833	0.854	0.839	0.049	0.910	0.864	0.833	0.854	0.839	

Acknowledgments

Zoh's research was supported by National Cancer Institute Supplemental under award U01-CA057030-29S1

References

- M. R. Ahmad, A u-statistic approach for a high-dimensional two-sample mean testing problem under non-normality and behrens-fisher setting, *Annals of the Institute of Statistical Mathematics* 66 (2014) 33–61.
- M. R. Ahmad, Location-invariant tests of homogeneity of large-dimensional covariance matrices, *Journal of Statistical Theory and Practice* 11 (2017) 731–745.
- Z. D. Bai, H. Saranadasa, Effect of high dimension: by an example of a two sample problem, *Statistica Sinica* 6 (1996) 311–329.
- J. Bezanson, S. Karpinski, V. B. Shah, A. Edelman, Julia: A fast dynamic language for technical computing, arXiv preprint arXiv:1209.5145 (2012).
- T. T. Cai, Y. Xia, High-dimensional sparse manova, *Journal of Multivariate Analysis* 131 (2014) 174–196.

Table 5: Empirical estimates of the power for the test based on BF^I when data are simulated as in case 2 and based on the Alternative 2. Note here cp refers to the true covariance matrix ($G = 3$).

		Φ_1						Φ_2					
cp		1	0.99	0.95	0.8	0.75	0.5	1	0.99	0.95	0.8	0.75	0.5
$n = 50, p = 200$	1	0.030	0.551	0.435	0.410	0.392	0.392	0.030	0.551	0.435	0.410	0.392	0.392
	2	0.033	0.745	0.608	0.562	0.504	0.505	0.033	0.745	0.608	0.562	0.504	0.505
	3	0.039	0.997	1	1	0.999	1	0.039	0.997	1	1	0.999	1
	4	0.041	0.671	0.544	0.538	0.476	0.485	0.041	0.671	0.544	0.538	0.476	0.485
	5	0.053	0.845	0.738	0.700	0.651	0.631	0.053	0.845	0.738	0.700	0.651	0.631
	6	0.086	0.980	0.942	0.925	0.915	0.914	0.086	0.980	0.942	0.925	0.915	0.914
$n = 70, p = 1000$	1	0.024	0.621	0.554	0.539	0.526	0.509	0.024	0.621	0.554	0.539	0.526	0.509
	2	0.015	0.811	0.763	0.714	0.708	0.667	0.015	0.811	0.763	0.714	0.708	0.667
	3	0.059	1	1	1	1	1	0.059	1	1	1	1	1
	4	0.011	0.738	0.692	0.659	0.659	0.617	0.011	0.738	0.692	0.659	0.659	0.617
	5	0.034	0.916	0.864	0.838	0.845	0.810	0.034	0.916	0.864	0.838	0.845	0.810
	6	0.087	1	0.999	0.998	0.999	0.997	0.087	1	0.999	0.998	0.999	0.997

- L. S. Chen, D. Paul, R. L. Prentice, P. Wang, A regularized hotelling's t^2 test for pathway analysis in proteomic studies, *Journal of the American Statistical Association* 106 (2011) 1345–1360.
- S. X. Chen, Y.-L. Qin, A two-sample test for high-dimensional data with applications to gene-set testing, *The Annals of Statistics* 38 (2010) 808–835.
- A. P. Dempster, A high dimensional two sample significance test, *The Annals of Mathematical Statistics* (1958) 995–1010.
- A. P. Dempster, A significance test for the separation of two highly multivariate small samples, *Biometrics* 16 (1960) 41–50.
- L. Feng, C. Zou, Z. Wang, L. Zhu, Composite t^2 test for high-dimensional data, *Statistica Sinica* (2017) 1419–1436.
- Y. Fujikoshi, T. Himeno, H. Wakaki, Asymptotic results of a high dimensional manova test and power comparison when the dimension is large compared to the sample size, *Journal of the Japan Statistical Society* 34 (2004) 19–26.
- S. Goddard, Restricted most powerful Bayesian tests, Ph.D Thesis, Texas A&M University (2015).

Table 6: Empirical estimates of the power for the test based on BF^I when data are simulated as in case 1 and assuming the Alternative 2. Note here cp refers to the true covariance matrix ($G = 3$). Here the data is simulated assuming different covariance matrices.

		Φ_1						Φ_2					
cp		1	0.99	0.95	0.8	0.75	0.5	1	0.99	0.95	0.8	0.75	0.5
$n = 50, p = 200$	1	0.030	0.921	0.866	0.782	0.781	0.712	0.030	0.921	0.866	0.782	0.781	0.712
	2	0.033	0.993	0.966	0.922	0.931	0.881	0.033	0.993	0.966	0.922	0.931	0.881
	3	0.039	1	1	1	1	1	0.039	1	1	1	1	1
	4	0.041	0.979	0.952	0.885	0.910	0.821	0.041	0.979	0.952	0.885	0.910	0.821
	5	0.053	1	0.995	0.992	0.986	0.970	0.053	1	0.995	0.992	0.986	0.970
	6	0.086	1	0.997	0.994	0.994	0.983	0.086	1	0.997	0.994	0.994	0.983
$n = 70, p = 1000$	1	0.024	0.989	0.981	0.956	0.960	0.906	0.024	0.989	0.981	0.956	0.960	0.906
	2	0.015	0.999	0.999	0.995	0.994	0.973	0.015	0.999	0.999	0.995	0.994	0.973
	3	0.059	1	1	1	1	1	0.059	1	1	1	1	1
	4	0.011	0.996	0.993	0.989	0.980	0.966	0.011	0.996	0.993	0.989	0.980	0.966
	5	0.034	1	1	1	1	1	0.034	1	1	1	1	1
	6	0.087	1	1	1	1	1	0.087	1	1	1	1	1

Table 7: Proportion of pairwise tests that were declared significant across 1000 random projections. The results are reported for the both tests (BF^I, BF^P).

	Macrophage	B Cell
Mast	(0.763, 0.943)	(0.689, 0.819)
Macrophage	-	(0.987, 1.0)

S. D. Goddard, V. E. Johnson, Restricted most powerful Bayesian tests for linear models, *Scandinavian Journal of Statistics* 43 (2016) 1162–1177.

Z. Hu, T. Tong, M. G. Genton, A pairwise hotelling method for testing high-dimensional mean vectors, *arXiv preprint arXiv:2003.04636* (2020).

V. E. Johnson, Bayes factors based on test statistics, *Journal of the Royal Statistical Society: Series B* 67 (2005) 689–701.

R. E. Kass, A. E. Raftery, Bayes factors, *Journal of the American Statistical Association* 90 (1995) 773–795.

H. Li, A. Aue, D. Paul, J. Peng, P. Wang, An adaptable generalization of hotelling’s t^2 test in high dimension, *The Annals of Statistics* 48 (2020) 1815–1847.

- M. Lopes, L. Jacob, M. J. Wainwright, A more powerful two-sample test in high dimensions using random projection, in: J. Shawe-Taylor, R. S. Zemel, P. L. Bartlett, F. Pereira, K. Q. Weinberger (Eds.), *Advances in Neural Information Processing Systems 24*, Curran Associates, Inc., 2011, pp. 1206–1214.
- D. López-Sánchez, C. de Bodt, J. A. Lee, A. G. Arrieta, J. M. Corchado, Tuning database-friendly random projection matrices for improved distance preservation on specific data, *Applied Intelligence* (2021) 1–13.
- S. V. Puram, I. Tirosh, A. S. Parikh, A. P. Patel, K. Yizhak, S. Gillespie, C. Rodman, C. L. Luo, E. A. Mroz, K. S. Emerick, et al., Single-cell transcriptomic analysis of primary and metastatic tumor ecosystems in head and neck cancer, *Cell* 171 (2017) 1611–1624.
- M. S. Srivastava, Multivariate theory for analyzing high dimensional data, *Journal of the Japan Statistical Society* 37 (2007) 53–86.
- M. S. Srivastava, H. Yanagihara, Testing the equality of several covariance matrices with fewer observations than the dimension, *Journal of Multivariate Analysis* 101 (2010) 1319–1329.
- R. Srivastava, P. Li, D. Ruppert, RAPTT: An exact two-sample test in high dimensions using random projections, *Journal of Computational and Graphical Statistics* 25 (2016) 954–970.
- M. Thulin, A high-dimensional two-sample test for the mean using random subspaces, *Computational Statistics & Data Analysis* 74 (2014) 26–38.
- T. Tony Cai, W. Liu, Y. Xia, Two-sample test of high dimensional means under dependence, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 76 (2014) 349–372.
- S. Wan, J. Kim, K. J. Won, Sharp: hyperfast and accurate processing of single-cell rna-seq data via ensemble random projection, *Genome research* 30 (2020) 205–213.
- R. S. Zoh, A. Sarkar, R. J. Carroll, B. K. Mallick, A powerful bayesian test for equality of means in high dimensions, *Journal of the American Statistical Association* 113 (2018) 1733–1741.