

ON QUASI-TAME LOOIJENGA PAIRS

ANDREA BRINI AND YANNIK SCHÜLER

ABSTRACT. We prove a conjecture of Bousseau, van Garrel and the first-named author relating, under suitable positivity conditions, the higher genus maximal contact log Gromov–Witten invariants of Looijenga pairs to other curve counting invariants of Gromov–Witten/Gopakumar–Vafa type. The proof consists of a closed-form q -hypergeometric resummation of the quantum tropical vertex calculation of the log invariants in presence of infinite scattering. The resulting identity of q -series appears to be new and of independent combinatorial interest.

1. INTRODUCTION

1.1. Quasi-tame Looijenga pairs. A Looijenga pair $Y(D) := (Y, D)$ is the datum of a smooth rational complex projective surface Y and an anticanonical singular curve $D \in |-K_Y|$. A Looijenga pair $Y(D)$ is called *nef* if the singular curve D is a simple normal crossings divisor $D = \cup_{i=1}^l D_i$ with each D_i smooth, irreducible, and nef¹ for all $i = 1, \dots, l$. A *tame* Looijenga pair is a nef pair with either $l > 2$, or $D_i^2 > 0$ for all i . Writing $E_{Y(D)} := \text{Tot}(\oplus_i \mathcal{O}_Y(-D_i))$, we will say that a nef pair $Y(D)$ is *quasi-tame* if there exists a tame pair $Y'(D')$ such that $E_{Y(D)}$ is deformation-equivalent to $E_{Y'(D')}$. By definition, there is an obvious sequence of nested inclusions

nef Looijenga pairs \supset quasi-tame Looijenga pairs \supset tame Looijenga pairs .

Looijenga pairs have been the focus of much attention lately due to their intertwined role in mirror symmetry for surfaces [2–4, 16, 19, 32, 36, 37] and the study of cluster varieties [15, 33, 39]. In a recent series of papers [7–9], the log Gromov–Witten theory of quasi-tame pairs was further conjectured to be at the centre of a web of correspondences relating it to several superficially distant enumerative theories. We recall the relevant context and fix notation below.

1.2. Enumerative theories. The authors of [7] consider four different geometries, and associated enumerative invariants, attached to the datum of a quasi-tame Looijenga pair $Y(D)$:

- (i) the log Calabi–Yau surface obtained by viewing $Y(D)$ as a log-scheme for the divisorial log structure induced by D . For a given genus g and effective curve class $d \in H_2(Y, \mathbb{Z})$, the corresponding set of invariants are the log Gromov–Witten invariants [1, 10, 18] of $Y(D)$ with maximal tangency at each component D_i , $l - 1$ point insertions on the surface, and one insertion of the top Chern class of the Hodge bundle:

$$N_{g,d}^{\log}(Y(D)) := \int_{[\mathcal{M}_{g,l-1}^{\log}(Y(D),d)]^{\text{vir}}} \prod_{i=1}^{l-1} \text{ev}_i^*[\text{pt}_Y] (-1)^g \lambda_g, \quad (1)$$

or equivalently, their all-genus generating function

$$\mathbb{N}_d^{\log}(Y(D))(\hbar) := \left(2 \sin\left(\frac{\hbar}{2}\right)\right)^{2-l} \sum_{g \geq 0} N_{g,d}^{\log}(Y(D)) \hbar^{2g-2+l}; \quad (2)$$

¹Since we require D to be singular, an l -component nef Looijenga pair must have $l > 1$.

- (ii) the quasi-projective Calabi–Yau variety $E_{Y(D)}$, and its genus zero local Gromov–Witten invariants [11, 12]

$$N_d(E_{Y(D)}) := \int_{[\mathcal{M}_{0,l-1}(E_{Y(D)},d)]^{\text{vir}}} \prod_{i=1}^{l-1} \text{ev}_i^*[\text{pt}_Y] \quad (3)$$

and local Gopakumar–Vafa invariants [20, 23]

$$\text{GV}_d(E_{Y(D)}) := \sum_{k|d} \frac{\mu(k)}{k^{4-l}} N_d(E_{Y(D)}); \quad (4)$$

- (iii) the quasi-projective Calabi–Yau threefold $\text{Tot}(\mathcal{O}_{Y \setminus \cup_{i < l} D_i}(-D_l))$ equipped with a disjoint union of $l - 1$ Lagrangians L_i fibred over real curves in D_i , $i < l$, as defined in [7, Construction 6.4]:

$$Y^{\text{op}}(D) := (\text{Tot}(\mathcal{O}(-D_l) \rightarrow Y \setminus (D_1 \cup \dots \cup D_{l-1})), L_1 \sqcup \dots \sqcup L_{l-1}),$$

The respective invariants, for a given relative homology degree $d \in H_2(Y^{\text{op}}(D), \mathbb{Z})$, are the open Gromov–Witten counts

$$\begin{aligned} O_{g,d}(Y^{\text{op}}(D)) &:= \int_{[\mathcal{M}_g(Y^{\text{op}}(D),d)]^{\text{vir}}} 1, \\ \mathbb{O}_d(Y^{\text{op}}(D))(\hbar) &:= \sum_{g \geq 0} \hbar^{2g+l-3} O_{g,d}(Y^{\text{op}}(D)) \end{aligned} \quad (5)$$

virtually enumerating genus- g open Riemann surfaces with $l - 1$ connected components of the boundary ending on the Lagrangians L_i , $i = 1, \dots, l-1$. Under relatively lax conditions², $Y^{\text{op}}(D)$ can be deformed to a singular Harvey–Lawson (Aganagic–Vafa) Lagrangian pair with $L_i \simeq \mathbb{R}^2 \times S^1$, for which open GW counts can be defined in the algebraic category [22, 28, 29]. We will also consider the corresponding genus zero/all-genus Labastida–Mariño–Ooguri–Vafa invariants [26, 27, 35]

$$\begin{aligned} \text{LMOV}_{0,d}(Y^{\text{op}}(D)) &= \sum_{k|d} \frac{\mu(k)}{k^{4-l}} O_{0,d/k}(Y^{\text{op}}(D)), \\ \text{LMOV}_d(Y^{\text{op}}(D))(\hbar) &= [1]_q^2 \prod_{i < l} \frac{w_i(d)}{[w_i(d)]_q} \sum_{k|d} \frac{\mu(k)}{k} \mathbb{O}_{d/k}(Y^{\text{op}}(D))(k\hbar), \end{aligned} \quad (6)$$

where $w_i(d)$ denotes the winding number of a relative degree- d open stable map to $Y^{\text{op}}(D)$ around the non-trivial homology circle in L_i , $[n]_q := q^{n/2} - q^{-n/2}$, and $q = e^{\hbar}$;

- (iv) for $l = 2$, a symmetric quiver $Q(Y(D))$ with adjacency matrix determined by $Y(D)$ [7, Thm. 7.3]. For a given charge vector d , the corresponding numbers are the numerical Donaldson–Thomas invariants $\text{DT}_d^{\text{num}}(Q(Y(D)))$, defined as the formal Taylor coefficients of (the plethystic logarithm of) the generating series of Euler characteristics on the stack of representations of $Q(Y(D))$.

The constructions of [7] in particular identify the absolute homology of $Y(D)$, the relative homology of $Y^{\text{op}}(D)$, and the free abelian group over the set of vertices of $Q(Y(D))$,

$$d \in H_2(Y(D), \mathbb{Z}) \simeq H_2(E_{Y(D)}, \mathbb{Z}) \simeq H_2^{\text{rel}}(Y^{\text{op}}(D), \mathbb{Z}) \simeq \mathbb{Z}^{|\text{Q}(Y(D))_0|}.$$

Under these identifications, the authors of [7] propose that the invariants above are essentially the same, as follows.

Conjecture 1.1 (The genus zero log/local/open correspondence). *The genus zero log, local, and open Gromov–Witten invariants associated to a quasi-tame Looijenga pair $Y(D)$ are related as*

$$N_d(E_{Y(D)}) = O_{0,d}(Y^{\text{op}}(D)) = \left(\prod_{j=1}^l \frac{(-1)^{d \cdot D_j - 1}}{d \cdot D_j} \right) N_{0,d}^{\text{log}}(Y(D)), \quad (7)$$

²This was formalised as “Property O” in [7, Definition 6.3]. All quasi-tame pairs with $l = 2$ satisfy Property O, and all non-tame quasi-tame pairs have $l = 2$, so this is safely assumed to hold throughout this paper.

and, for the associated BPS invariants,

$$\mathrm{GV}_d(E_{Y(D)}) = \mathrm{LMOV}_{0,d}(Y^{\mathrm{op}}(D)) \in \mathbb{Z}. \quad (8)$$

Moreover, if $l = 2$,

$$\mathrm{DT}_d^{\mathrm{num}}(\mathrm{Q}(Y(D))) = |\mathrm{GV}_d(E_{Y(D)})|. \quad (9)$$

In [7, Conj. 1.3], the above is further extended to an identity between all-genus Gromov–Witten generating functions.

Conjecture 1.2 (The higher genus log/open correspondence). *The higher genus log and open Gromov–Witten invariants associated to a quasi-tame Looijenga pair $Y(D)$ are related as*

$$\mathbb{O}_d(Y^{\mathrm{op}}(D))(\hbar) = \left(\prod_{j=1}^{l-1} \frac{(-1)^{d \cdot D_j - 1}}{d \cdot D_j} \right) \frac{(-1)^{d \cdot D_l - 1}}{[d \cdot D_l]_q} \mathbb{N}_d^{\mathrm{log}}(Y(D))(\hbar). \quad (10)$$

Moreover,

$$\mathrm{LMOV}_d(Y^{\mathrm{op}}(D))(\hbar) = [1]_q^2 \left(\prod_{i=1}^l \frac{1}{[d \cdot D_i]_q} \right) \sum_{k|d} \frac{(-1)^{d/k \cdot D + l} \mu(k)}{[k]_q^{2-l} k^{2-l}} \mathbb{N}_{d/k}^{\mathrm{log}}(Y(D))(k\hbar) \in \mathbb{Z}[q, q^{-1}]. \quad (11)$$

Conjecture 1.1 was proved in [7, Thm. 1.4–1.6]. Conjecture 1.2 was established in [7, Thm. 1.5 and 1.7] for tame $Y(D)$, and formulated as a conjecture for quasi-tame $Y(D)$ in [7, Conj. 4.8]. Two non-tame, quasi-tame cases of this conjecture were subsequently proved in [25].

In this paper, we establish a stronger statement from which Conjecture 1.2 follows for any quasi-tame Looijenga pair $Y(D)$. We will provide a closed-form calculation and comparison of both sides of (10), returning the two equalities in Conjecture 1.2 as a corollary. Since the tame setting was already dealt with in [7], we may restrict our attention here to non-tame, quasi-tame pairs alone, for which $l = 2$. We give here a slightly discursive version of the main result of this paper (see Propositions 2.2, 2.3 and 2.6 for precise statements).

Theorem 1.3. *Let $Y(D)$ be a non-tame, quasi-tame Looijenga pair $Y(D)$ and $d \in H_2(Y, \mathbb{Z})$. The log and open higher genus generating functions $\mathbb{N}_d^{\mathrm{log}}(Y(D))(\hbar)$ and $\mathbb{O}_d(Y^{\mathrm{op}}(D))(\hbar)$ are explicit rational functions of $q = e^{\hbar}$, with zeroes and poles only at $q = 0, \infty$, or at roots of unity. Furthermore, Conjecture 1.2 holds.*

1.3. Strategy of the proof. Our task is simplified by a number of circumstances, which effectively boil down to considering some explicit calculations of higher genus log Gromov–Witten invariants for *one single example*. As explained in [14] and [7, Prop. 2.2], $Y(D)$ is fully determined by the self-intersections (D_1^2, D_2^2) , and when considering specific examples we will shorten notation by writing $Y(D_1^2, D_2^2)$ for $Y(D)$. Let $\pi : \mathrm{dP}_r \rightarrow \mathbb{P}^2$ be the blow-up of the plane at $r \geq 0$ points $\{P_1, \dots, P_r\}$, and write $H := \pi^* c_1(\mathcal{O}_{\mathbb{P}^2}(1))$, $E_i := [\pi^{-1}(P_i)]$. Up to deformation and permutation of D_1, D_2 and E_1, \dots, E_r , there is a unique non-tame, quasi-tame pair of maximal Picard number given by $Y = \mathrm{dP}_3$ and $D_1 = H - E_1$, $D_2 = 2H - E_2 - E_3$, so that $(D_1^2, D_2^2) = (0, 2)$.

By [7, Prop 2.2], any other non-tame, quasi-tame pair $Y(D)$ is a toric contraction $\pi' : \mathrm{dP}_3(0, 2) \rightarrow Y(D)$. Therefore, as we recall in Proposition 2.2 below, the blow-up formulas for log and open invariants [7, Prop. 4.3 and 6.9] imply that it suffices to check that Conjecture 1.2 holds for the single case $Y(D) = \mathrm{dP}_3(0, 2)$. The l.h.s. of (10) in that case was computed in [7, Sec. 6.3.1] to be

$$\mathbb{O}_d(\mathrm{dP}_3^{\mathrm{op}}(0, 2)) = (-1)^{d_1 + d_2 + d_3} \frac{[d_1]_q}{d_1 [d_0]_q [d_1 + d_2 + d_3 - d_0]_q} \begin{bmatrix} d_3 \\ d_0 - d_1 \end{bmatrix}_q \begin{bmatrix} d_3 \\ d_0 - d_2 \end{bmatrix}_q \begin{bmatrix} d_0 \\ d_3 \end{bmatrix}_q \begin{bmatrix} d_1 + d_2 + d_3 - d_0 \\ d_3 \end{bmatrix}_q, \quad (12)$$

where we decomposed $d = d_0(H - E_1 - E_2 - E_3) + d_1E_1 + d_2E_2 + d_3E_3$ in terms of generators $\{H - E_1 - E_2 - E_3, E_1, E_2, E_3\}$ of $H_2(\mathrm{dP}_3, \mathbb{Z})$, and for non-negative integers n, m we denoted³

$$[n]_q! := \prod_{i=1}^n [i]_q, \quad \begin{bmatrix} n \\ m \end{bmatrix}_q := \begin{cases} \frac{[n]_q!}{[m]_q! [n-m]_q!} & 0 \leq m \leq n, \\ 0 & \text{otherwise.} \end{cases} \quad (13)$$

The first equality (10) in Conjecture 1.2 is then a consequence of the following

Proposition 1.4 (=Propositions 2.3 and 2.6). *With the conventions above, we have*

$$\mathbb{N}_d^{\log}(\mathrm{dP}_3(0, 2))(\hbar) = \frac{[d_1]_q [d_2 + d_3]_q}{[d_0]_q [d_1 + d_2 + d_3 - d_0]_q} \begin{bmatrix} d_3 \\ d_0 - d_1 \end{bmatrix}_q \begin{bmatrix} d_3 \\ d_0 - d_2 \end{bmatrix}_q \begin{bmatrix} d_0 \\ d_3 \end{bmatrix}_q \begin{bmatrix} d_1 + d_2 + d_3 - d_0 \\ d_3 \end{bmatrix}_q. \quad (14)$$

The second statement then also follows from Proposition 1.4 combined with the BPS integrality result of [7, Thm. 8.1] for $l = 2$, whose proof applies identically to this case.

We will show Proposition 1.4 in two main steps. We will first construct a toric model for $\mathrm{dP}_3(0, 2)$ in the sense of [16], and then compute the λ_q -log Gromov–Witten invariants (1) from the corresponding quantum scattering diagram and algebra of quantum broken lines [4–6, 13, 31]. The lack of tameness is epitomised by the existence of infinite scattering when two quantum walls meet, and the resulting sum over quantum broken lines leads to the intricate-looking multi-variate generalised hypergeometric sum in (17). The second step consists of proving that this sum admits a closed-form q -hypergeometric resummation given by (14), which has not to our knowledge previously appeared in the literature, with the exception of the special cases $d_3 = d_2 = d_0$ and $d_3 = d_0 = d_1 + d_2$ considered in [25]. To prove it, we first establish a q -hypergeometric recursion relation satisfied by a 1-parameter deformation of (17), and then show inductively that this recursion has a unique closed-form solution compatible with (17) by repeated use of Jackson’s q -analogue of the Pfaff–Saalschütz summation for the ${}_3\phi_2$ -hypergeometric function.

Acknowledgements. We thank P. Bousseau and M. van Garrel for many enlightening discussions surrounding the topic of this paper. We are also particularly indebted to C. Krattenthaler for an illuminating e-mail correspondence occurred prior to the appearance of [25]. He exposed us to the idea that our sought-for q -hypergeometric identities were too rigid to be tackled directly, whereas suitable parametric refinements might be amenable to an effective recursive strategy. Special thanks are owed to him for this insight, which is key to the arguments of Section 2.2.4.

2. PROOF OF CONJECTURE 1.2

2.1. Log GW invariants from the quantum tropical vertex. We start off by giving a streamlined summary of the combinatorial setup for the calculation of the higher genus log GW invariants (2) from the associated quantum scattering diagram, referring the reader to [7, Sec. 4.2] for a more extensive treatment.

2.1.1. Toric models. Two birational operations on log Calabi–Yau surfaces $Y(D)$ will feature prominently in the foregoing discussion.

- If \tilde{Y} is the blow-up of Y at a node of D and \tilde{D} is the preimage of D in \tilde{Y} we say $\tilde{Y}(\tilde{D})$ is a *corner blow-up* of $Y(D)$.
- In case \tilde{Y} is the blow-up of Y at a smooth point in D and \tilde{D} is the strict transform of D in \tilde{Y} we say $\tilde{Y}(\tilde{D})$ is an *interior blow-up* of $Y(D)$.

³By [7, Prop. 2.5], the effectiveness of d implies that all arguments of the q -binomial expressions in (12) are non-negative integers.

Starting from a Looijenga pair $Y(D)$, we will seek to construct pairs $\widetilde{Y}(\widetilde{D})$ and $\overline{Y}(\overline{D})$ fitting into a diagram

$$\begin{array}{ccc} & \widetilde{Y}(\widetilde{D}) & \\ \varphi \swarrow & & \searrow \pi \\ Y(D) & & \overline{Y}(\overline{D}) \end{array} \quad (15)$$

where φ is a sequence of corner blow-ups and π is a *toric model*, meaning that \overline{Y} is toric, \overline{D} is its toric boundary and π is a sequence of interior blow-ups. By [16, Prop. 1.3] such a diagram always exists. We will write ρ_{D_i} for the generator of the ray in the fan $\overline{\Sigma}$ of \overline{Y} associated to the toric prime divisor which is the push-forward along π of the strict transform of D_i under φ .

Given a toric model $\pi : \widetilde{Y}(\widetilde{D}) \rightarrow \overline{Y}(\overline{D})$ as in (15) we can associate a *consistent quantum scattering diagram* \mathfrak{D} to it, which is what we discuss next.

2.1.2. Quantum scattering and higher genus invariants. The scattering diagram \mathfrak{D} is defined on the lattice $N_{\mathbb{Z}} \cong \mathbb{Z}^2$ of integral points of the fan $\overline{\Sigma}$ of \overline{Y} , as follows: assume that π is a sequence of blow-ups of distinct smooth points P_1, \dots, P_s of \overline{D} and denote $\mathcal{E}_1, \dots, \mathcal{E}_s$ the exceptional divisors in \widetilde{Y} associated to these blow-ups. Further, write ρ_j for the primitive generator of the ray associated to the toric prime divisor which P_j is an element of. For each $j \in \{1, \dots, s\}$ we introduce a wall $\mathfrak{d}_j := \mathbb{R}\rho_j$ decorated with wall-crossing functions $f_{\mathfrak{d}_j} := 1 + t_j z^{-\rho_j} \in \mathbb{C}[[t_1, \dots, t_s]][N_{\mathbb{Z}}]$, where $z^\rho := x^a y^b$ if $\rho = (a, b)$. Then \mathfrak{D} is the unique (up to equivalence) completion of the initial scattering diagram $\mathfrak{D}_{\text{in}} := \{(\mathfrak{d}_j, f_{\mathfrak{d}_j})\}_{j \in \{1, \dots, s\}}$ in the sense of [4, 5, 17, 24]. Such a completion can be found algorithmically by successively adding new rays whenever two walls meet, as we now describe.

First of all, after perturbing the diagram \mathfrak{D}_{in} we may assume that walls intersect in codimension at least one and that no more than two walls meet in a point. Now suppose two walls $\mathfrak{d}^1, \mathfrak{d}^2$ intersect. Denote by $-\rho^i$ a primitive integral direction of \mathfrak{d}^i and assume $f_{\mathfrak{d}^i} = 1 + c_i z^{\rho^i}$. For our purpose the relevant scattering processes are:

- $\det(\rho^1, \rho^2) = \pm 1$ (*simple scattering*): the algorithm tells us to add a ray \mathfrak{d} emanating from the intersection point $\mathfrak{d}^1 \cap \mathfrak{d}^2$ in the direction $-\rho^1 - \rho^2$ decorated with $1 + c_1 c_2 z^{\rho^1 + \rho^2}$.
- $\det(\rho^1, \rho^2) = \pm 2$ (*infinite scattering*): the algorithm creates three types of walls. First, there is a central wall in the direction $-\rho^1 - \rho^2$ decorated with a wall-crossing function whose explicit shape is not of interest in the subsequent analysis and hence omitted. Further – and most relevant for us later – one needs to add walls $\mathfrak{d}_1, \dots, \mathfrak{d}_n, \dots$ with slope $-(n+1)\rho^1 - n\rho^2$ decorated with

$$1 + c_1^{n+1} c_2^n z^{(n+1)\rho^1 + n\rho^2}$$

and last a collection ${}_1\mathfrak{d}, \dots, {}_n\mathfrak{d}, \dots$ of walls respectively having slope $-n\rho^1 - (n+1)\rho^2$ and decorated with

$$1 + c_1^n c_2^{n+1} z^{n\rho^1 + (n+1)\rho^2}.$$

Adding new walls to the scattering diagram possibly creates new intersection points. Perturbing the diagram if necessary and repeating the above described process for each newly created intersection point yields a consistent scattering diagram \mathfrak{D} .

We now introduce the final combinatorial object we require for our computation of log Gromov–Witten invariants. Let $N_{\mathbb{R}} := N_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{R}$. Given $p \in N_{\mathbb{R}}$ and $m \in N_{\mathbb{Z}}$, a *quantum broken line* β with ends p and m consists of

- (i) a continuous piece-wise straight path $\beta : (-\infty, 0] \rightarrow B \setminus \bigcup_{\mathfrak{d} \in \mathfrak{D}} \partial \mathfrak{d} \cup \bigcup_{\mathfrak{d}^1 \neq \mathfrak{d}^2} \mathfrak{d}^1 \cap \mathfrak{d}^2$ intersecting walls transversely;

- (ii) a labelling L_1, \dots, L_n of the successive line segments with L_n ending at p such that each intersection point $L_i \cap L_{i+1}$ lies on a wall;
- (iii) an assignment $a_i z^{m_i}$ to each line segment L_i such that
 - $a_1 z^{m_1} = z^m$,
 - if $L_i \cap L_{i+1} \subset \mathfrak{d}$ with $f_{\mathfrak{d}} = \sum_{r \geq 0} c_r z^{r \rho_{\mathfrak{d}}}$ and $\rho_{\mathfrak{d}}$ chosen primitive then $a_{i+1} z^{m_{i+1}}$ is a monomial occurring in the expansion of

$$a_i z^{m_i} \prod_{\ell = -\frac{1}{2}(|\det(\rho_{\mathfrak{d}}, m_i)| - 1)}^{\frac{1}{2}(|\det(\rho_{\mathfrak{d}}, m_i)| - 1)} \left(\sum_{r \geq 0} c_r q^{r \ell} z^{r \rho_{\mathfrak{d}}} \right), \quad (16)$$

- L_i is directed in direction $-m_i$.

For such a quantum broken line β call $a_{\beta, \text{end}} := a_n$ the end-coefficient and write $m_{\beta, \text{end}} := m_n$. We usually refer to z^m as the asymptotic monomial of β . We remark that for $f_{\mathfrak{d}} = 1 + cz^{\rho_{\mathfrak{d}}}$ the product in (16) takes the form

$$\prod_{\ell = -\frac{1}{2}(|\det(\rho_{\mathfrak{d}}, m_i)| - 1)}^{\frac{1}{2}(|\det(\rho_{\mathfrak{d}}, m_i)| - 1)} (1 + cq^{\ell} z^{\rho_{\mathfrak{d}}}) = \sum_{k=0}^{|\det(\rho_{\mathfrak{d}}, m_i)|} \left[\begin{matrix} |\det(\rho_{\mathfrak{d}}, m_i)| \\ k \end{matrix} \right]_q c^k z^{k \rho_{\mathfrak{d}}}.$$

Now given two lattice vectors $m_1, m_2 \in N_{\mathbb{Z}}$, define

$$C_{m_1, m_2}^{\mathfrak{D}}(q) := \sum_{\substack{\beta_1, \beta_2 \\ \text{Ends}(\beta_i) = (p, m_i) \\ m_{\beta_1, \text{end}} + m_{\beta_2, \text{end}} = 0}} a_{\beta_1, \text{end}} a_{\beta_2, \text{end}}$$

to be the sum of products of all end-coefficients of quantum broken lines β_1 and β_2 with asymptotic monomials z^{m_1} , resp. z^{m_2} , meeting in a common point p and with opposite exponents of their end-monomials. This sum is a polynomial in the variables t_j with coefficients in $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ and is moreover independent of the choice of p .

The higher genus, maximal contact log Gromov–Witten invariants (2) of a Looijenga pair $Y(D)$ can be extracted from the scattering diagram associated to its toric model, as per the following

Proposition 2.1 ([7, 34]). *Let $Y(D = D_1 + D_2)$ be a 2-component Looijenga pair, $\pi : \widetilde{Y}(\widetilde{D}) \rightarrow \overline{Y}(\overline{D})$ a toric model for it as in (15), and \mathfrak{D} the corresponding consistent quantum scattering diagram. For $d \in H_2(Y, \mathbb{Z})$ an effective curve class, set $m_i := (d \cdot D_i) \rho_{D_i}$. Then*

$$\mathbb{N}_d^{\log}(Y(D))(\hbar) = \left[\prod_{j=1}^s t_j^{d \cdot \varphi_* \mathcal{E}_j} \right] C_{m_1, m_2}^{\mathfrak{D}}(q) \Big|_{q=e^{i\hbar}}.$$

2.1.3. Birational invariance. Suppose now that $Y(D)$ is a 2-component quasi-tame Looijenga pair and $\pi : Y'(D') \rightarrow Y(D)$ is a sequence of interior blow-ups such that $Y'(D')$ is also quasi-tame. The following compatibility statement explains how the higher genus log-open correspondence interacts with this type of birational transformations.

Proposition 2.2. [7, Prop. 4.3 and 6.9] *Let $\pi : Y'(D') \rightarrow Y(D)$ be an interior blow-up, with both $Y'(D')$ and $Y(D)$ 2-component quasi-tame Looijenga pairs. Then $\mathbb{N}_d^{\log}(Y(D)) = \mathbb{N}_{\pi^* d}^{\log}(Y'(D'))$, $\mathbb{O}_d(Y^{\text{op}}(D)) = \mathbb{O}_{\pi^* d}(Y'^{\text{op}}(D'))$ for all $d \in H_2(Y, \mathbb{Z})$.*

The comparison statement of Proposition 2.2 for log invariants is easy to visualise in genus 0, where the invariants $\mathbb{N}_{0, d}^{\log}(Y(D))$ are enumerative [34]: since blowing up a point away from the curves does not affect the local geometry, the corresponding counts are invariant, a property also reflected in the broken lines calculations of the scattering diagrams of Section 2.1.2. The corresponding statement for the open

invariants is a non-trivial consequence of the invariance of the topological vertex under flops [21, 30].

By the classification theorem of nef Looijenga pairs in [7, Prop. 2.2], any non-tame, quasi-tame Looijenga pair $Y(D)$ is obtained up to deformation as a sequence of $m \geq 0$ interior blowings-down of $d\mathbb{P}_3(0, 2)$. It follows from Proposition 2.2 that proving Conjecture 1.2 for $Y(D) = d\mathbb{P}_3(0, 2)$ implies that the same statement *a fortiori* holds for any other non-tame, quasi-tame pair (and indeed any 2-component quasi-tame pair with Picard number lower than four).

2.2. The case of maximal Picard number. Let us then specialise to $Y(D) = d\mathbb{P}_3(0, 2)$. Throughout this section we will write $d = d_0(H - E_1 - E_2 - E_3) + \sum_{i=1}^3 d_i E_i$ for a curve class $d \in H_2(d\mathbb{P}_3, \mathbb{Z})$. If $d \cdot D_1$ or $d \cdot D_2$ vanishes, then $\mathbb{N}_d^{\text{log}}(d\mathbb{P}_3(0, 2))(\hbar) = 0$. In case the intersection numbers are strictly positive, we may use the scattering diagram of $d\mathbb{P}_3(0, 2)$ to compute the invariants.

2.2.1. Constructing the toric model. First, we recall from [7, Sec. 4.4] the construction of a toric model of $\mathbb{P}^2(D_1 \cup D_2)$, with D_1 a line and D_2 a smooth conic intersecting D_1 in two distinct points P_1 and P_2 . Let L denote the tangent line through P_1 to D_2 . In the following we will always identify D_1, D_2, L , and some yet to be defined divisors F_1, F_2 with their strict transforms, resp. push-forwards, under blow-ups, resp. blow-downs.

Blow up the point P_1 , and write F_1 for the exceptional divisor, and then blow up the intersection point of F_1 with D_2 and denote the exceptional curve by F_2 . Under these blow-ups, the strict transform of L is a (-1) -curve intersecting F_2 in one point and can therefore be blown down. This results in the log Calabi–Yau surface $(\overline{\mathbb{P}^2(D)}, \overline{D})$ with anti-canonical divisor $\overline{D} = D_1 \cup F_1 \cup F_2 \cup D_2$ where F_1 is a curve with self-intersection -2 , D_2 has self-intersection 2, and D_1, F_2 zero. Therefore, since the irreducible components of \overline{D} form a sequence with the same self-intersections as the toric boundary of \mathbb{F}_2 , we already must have $\overline{\mathbb{P}^2(D)} = \mathbb{F}_2$ with \overline{D} the toric boundary by [14, Lemma 2.10] and hence we have constructed a toric model for $\mathbb{P}^2(D)$. From the discussion of the previous Section, at a tropical level the fact that we blew down L amounts to adding a wall \mathfrak{d}_{F_2} emanating from a *focus-focus singularity* on the ray corresponding to F_2 in the fan of \mathbb{F}_2 .

Now from this we obtain a toric model for $d\mathbb{P}_3(0, 2)$ by blowing up an interior point on D_1 and two points on D_2 and taking the proper transforms, ie. by adding focus-focus singularities on the corresponding rays of the fan of \mathbb{F}_2 . The resulting picture of the fan of $\overline{\mathbb{P}^2(D)} = \mathbb{F}_2(\overline{D})$ decorated with the focus-focus singularities of the toric model of $d\mathbb{P}_3(0, 2)$ is displayed in Figure 1.




FIGURE 1. The toric model of $d\mathbb{P}_3(0, 2)$

2.2.2. *The quantum scattering diagram.* We now follow the construction outlined in Section 2.1.2 to derive the quantum scattering diagram of the toric model of $d\mathbb{P}_3(0, 2)$ (Figure 2). We shoot out walls \mathfrak{d}_{F_2} , \mathfrak{d}_{D_1} , $\mathfrak{d}_{D_{2,1}}$, $\mathfrak{d}_{D_{2,2}}$ emanating from the focus-focus singularities in the direction $-\rho$, where ρ is the respective generator of the ray in the fan. We send the singularities to infinity and perturb the walls as indicated in Figure 2. From these initial walls we now want to construct a consistent scattering diagram. However, since in our subsequent analysis we will only be interested in walls with slope lying in the cone generated by $(0, -1)$ and $(1, 2)$, we will restrict the discussion to such walls only. As $|\rho_{F_2} \wedge \rho_{D_1}| = 2$ there is infinite scattering in the sense of Section 2.1.2 between the walls \mathfrak{d}_{F_2} and \mathfrak{d}_{D_1} . This results in walls $\mathfrak{d}_2, \mathfrak{d}_3, \dots$ with slope $-n\rho_{F_2} - (n-1)\rho_{D_1} = (1, -2(n-1))$ decorated with wallcrossing functions $1 + t^n t_1^{n-1} x^{-1} y^{2(n-1)}$ where $n > 1$. For conformity, let us write $\mathfrak{d}_1 := \mathfrak{d}_{F_2}$. Now for all $n \geq 1$ each wall \mathfrak{d}_n intersects both $\mathfrak{d}_{D_{2,1}}$ and $\mathfrak{d}_{D_{2,2}}$. Luckily, in this case we only have simple scattering resulting in walls with slope $(1, -2n+3)$ and wallcrossing functions $1 + t^n t_1^{n-1} t_i x^{-1} y^{2n-3}$ where $i \in \{2, 3\}$. Lastly, we notice that the wall which is the result of scattering between \mathfrak{d}_n and $\mathfrak{d}_{D_{2,1}}$ intersects $\mathfrak{d}_{D_{2,2}}$ thus producing a wall with slope $(1, -2n+4)$ and wallcrossing function $1 + t^n t_1^{n-1} t_2 t_3 x^{-1} y^{2n+4}$ attached to it. Let us call this wall $\mathfrak{d}_n^{D_2, D_2}$. The whole construction is summarised in Figure 2.





FIGURE 2. The quantum scattering diagram of $d\mathbb{P}_3(0, 2)$.

FIGURE 3. The n th 4-tuple of walls.

We can collect the walls constructed above into repeatedly occurring 4-tuples labelled by an integer $n > 0$ as depicted in Figure 3. The n th tuple consists of the wall \mathfrak{d}_n , the walls which are the result of scattering between \mathfrak{d}_n and $\mathfrak{d}_{D_2,1}$, resp. $\mathfrak{d}_{D_2,2}$, and lastly the wall $\mathfrak{d}_{n+1}^{D_2, D_2}$.

2.2.3. *Higher genus log GW invariants.* We now apply the algorithm of Section 2.1.2 to obtain the following Proposition.

Proposition 2.3. *Let d be an effective curve class with $d \cdot D_1, d \cdot D_2 > 0$. Then*

$$\mathbb{N}_d^{\log}(\mathrm{dP}_3(0, 2))(\hbar) = \sum_{\substack{\forall (i,n) \in \{1,2,3,4\} \times \mathbb{Z}_{>0}: k_{i,n} \geq 0 \\ d_0 = \sum_{n \geq 1} \sum_{i=1}^4 (n + \delta_{i,1}) k_{i,n} \\ d_1 = \sum_{n \geq 1} \sum_{i=1}^4 k_{i,n} \\ d_0 - d_2 = \sum_{n \geq 1} (k_{1,n} + k_{4,n}) \\ d_0 - d_3 = \sum_{n \geq 1} (k_{1,n} + k_{3,n})}} \prod_{n \geq 1} \prod_{i=1}^2 \left[\begin{array}{c} d_2 + d_3 - \sum_{m \geq 1} (2m(k_{1,n+m} + k_{2,n+m}) + (2m-1)(k_{3,n+m} + k_{4,n+m})) \\ k_{i,n} \end{array} \right]_q \times \left[\begin{array}{c} d_2 + d_3 - \sum_{m \geq 0} ((2m+1)(k_{1,n+m} + k_{2,n+m}) + 2m(k_{3,n+m} + k_{4,n+m})) \\ k_{2+i,n} \end{array} \right]_q. \quad (17)$$

Proof. In order to compute $\mathbb{N}_d^{\log}(\mathrm{dP}_3(0, 2))$ we choose a point p sufficiently far into the first quadrant and consider quantum broken lines β_i with ends $(p, z^{(d \cdot D_i) \rho_{D_i}})$, where $i \in \{1, 2\}$, so that the sum of the exponents of their end-monomials at p vanishes. As stated in Proposition 2.1, we then obtain the desired log GW invariant by taking the product of the two end-coefficients, summing this over all such quantum broken lines and extracting the correct monomial in the variables t, t_1, t_2, t_3 .

Let us then analyse a quantum line β_2 coming from the direction D_2 with asymptotic monomial $y^{d \cdot D_2} = y^{-d_2 - d_3}$ ending at p , as shown in Figure 2. We assume that this quantum broken line passes all 4-tuples of walls for $n > N$ without picking up any contribution for some $N \geq 1$. Then after crossing the four walls

belonging to the N th tuple we will find a monomial attached to β_2 of the form

$$\begin{aligned}
y^{-d_2-d_3} &\stackrel{\textcircled{1}}{\mapsto} \begin{bmatrix} d_2 + d_3 \\ k_{1,N} \end{bmatrix}_q t^{(N+1)k_{1,N}} t_1^{Nk_{1,N}} t_2^{k_{1,N}} t_3^{k_{1,N}} x^{-k_{N,1}} y^{-d_2-d_3+2(N-1)k_{N,1}} \\
&\stackrel{\textcircled{2}}{\mapsto} \begin{bmatrix} d_2 + d_3 \\ k_{1,N} \end{bmatrix}_q \begin{bmatrix} d_2 + d_3 \\ k_{2,N} \end{bmatrix}_q t^{(N+1)k_{1,N}+Nk_{2,N}} t_1^{Nk_{1,N}+(N-1)k_{2,N}} t_2^{k_{1,N}} t_3^{k_{1,N}} \\
&\quad \times x^{-(k_{N,1}+k_{N,2})} y^{-d_2-d_3+2(N-1)(k_{N,1}+k_{N,2})} \\
&\stackrel{\textcircled{3}}{\mapsto} \begin{bmatrix} d_2 + d_3 \\ k_{1,N} \end{bmatrix}_q \begin{bmatrix} d_2 + d_3 \\ k_{2,N} \end{bmatrix}_q \begin{bmatrix} d_2 + d_3 - (k_{N,1} + k_{N,2}) \\ k_{3,N} \end{bmatrix}_q \\
&\quad \times t^{k_{1,N}+N \sum_{i=1}^3 k_{i,N}} t_1^{k_{1,N}+(N-1) \sum_{i=1}^3 k_{i,N}} t_2^{k_{1,N}+k_{3,N}} t_3^{k_{1,N}+k_{3,N}} \\
&\quad \times x^{-\sum_{i=1}^3 k_{i,N}} y^{-d_2-d_3+2(N-1)(k_{N,1}+k_{N,2})+(2N-3)k_{3,N}} \\
&\stackrel{\textcircled{4}}{\mapsto} \begin{bmatrix} d_2 + d_3 \\ k_{1,N} \end{bmatrix}_q \begin{bmatrix} d_2 + d_3 \\ k_{2,N} \end{bmatrix}_q \begin{bmatrix} d_2 + d_3 - (k_{N,1} + k_{N,2}) \\ k_{3,N} \end{bmatrix}_q \begin{bmatrix} d_2 + d_3 - (k_{N,1} + k_{N,2}) \\ k_{4,N} \end{bmatrix}_q \\
&\quad \times t^{k_{1,N}+N \sum_{i=1}^4 k_{i,N}} t_1^{k_{1,N}+(N-1) \sum_{i=1}^4 k_{i,N}} t_2^{k_{1,N}+k_{4,N}} t_3^{k_{1,N}+k_{3,N}} \\
&\quad \times x^{-\sum_{i=1}^4 k_{i,N}} y^{-d_2-d_3+2(N-1)(k_{N,1}+k_{N,2})+(2N-3)(k_{3,N}+k_{4,N})}.
\end{aligned}$$

Then, after crossing the $(N-1)$ remaining 4-tuples of walls until reaching the point p , we find that the end-monomial $a_{\beta_2, \text{end}} z^{m_{\beta_2, \text{end}}}$ is of the form

$$\begin{aligned}
y^{-d_2-d_3} &\mapsto \dots \mapsto \prod_{n=1}^N \prod_{i=0}^1 \left[\begin{bmatrix} d_2 + d_3 - \sum_{m=1}^{N-n} (2m(k_{1,n+m} + k_{2,n+m}) + (2m-1)(k_{3,n+m} + k_{4,n+m})) \\ k_{i,n} \end{bmatrix}_q \right. \\
&\quad \times \left. \begin{bmatrix} d_2 + d_3 - \sum_{m=0}^{N-n} ((2m+1)(k_{1,n+m} + k_{2,n+m}) + 2m(k_{3,n+m} + k_{4,n+m})) \\ k_{2+i,n} \end{bmatrix}_q \right] \quad (18) \\
&\quad \times t^{\sum_{n=1}^N (k_{1,n}+n \sum_{i=1}^4 k_{i,n})} t_1^{\sum_{n=1}^N (k_{1,n}+(n-1) \sum_{i=1}^4 k_{i,n})} t_2^{\sum_{n=1}^N (k_{1,n}+k_{4,n})} t_3^{\sum_{n=1}^N (k_{1,n}+k_{3,n})} \\
&\quad \times x^{-\sum_{n=1}^N \sum_{i=1}^4 k_{i,n}} y^{-d_2-d_3+\sum_{n=1}^N (2(n-1)(k_{1,n}+k_{2,n})+(2n-3)(k_{3,n}+k_{4,n}))}.
\end{aligned}$$

Note that we are only interested in such quantum broken lines for which $z^{-m_{\beta_2, \text{end}}} = (xy^2)^{d \cdot D_1} = x^{d_1} y^{2d_1}$ which means that we need to impose

$$d_1 = \sum_{n=1}^N \sum_{i=1}^4 k_{i,n}, \quad (19)$$

$$d_2 + d_3 = \sum_{n=1}^N (2n(k_{1,n} + k_{2,n}) + (2n-1)(k_{3,n} + k_{4,n})). \quad (20)$$

Now [7, Prop. 4.2] tells us that $\mathbb{N}_d^{\log}(\text{dP}_3(0, 2))$ is the coefficient of

$$t^{d \cdot L} t_1^{d \cdot E_1} t_2^{d \cdot E_2} t_3^{d \cdot E_3} = t^{d_0} t_1^{d_0-d_1} t_2^{d_0-d_2} t_3^{d_0-d_3} \quad (21)$$

in the sum over all end-coefficients $a_{\beta_{2,\text{end}}}$ satisfying (19) and (20). In turn, picking the coefficient of (21) amounts to imposing

$$d_0 = \sum_{n=1}^N \left(k_{1,n} + n \sum_{i=1}^4 k_{i,n} \right), \quad (22)$$

$$d_0 - d_1 = \sum_{n=1}^N \left(k_{1,n} + (n-1) \sum_{i=1}^4 k_{i,n} \right), \quad (23)$$

$$d_0 - d_2 = \sum_{n=1}^N (k_{1,n} + k_{4,n}), \quad (24)$$

$$d_0 - d_3 = \sum_{n=1}^N (k_{1,n} + k_{3,n}) \quad (25)$$

in the sum over the end-coefficients. Notice here that (22), (24), and (25) together imply (20), and that (23) is a consequence of (19) and (22). Moreover, we see that (22) forces $k_{i,n} = 0$ for all $n > d_0$ from which we learn that the quantum broken lines we are considering may only pick up a non-trivial contribution for finitely many walls. Note that this also justifies our initial assumption to choose $N > 0$ finite.

Thus, summing over all end-monomials (18) satisfying (19), (20) and extracting the coefficient of (21) we indeed arrive at expression (17) for the log GW invariants. \square

Remark 2.4. It is easy to convince oneself that there are actually only finitely many summands contributing to (17), due to the finite number of choices $(k_{i,n})$ satisfying the summation conditions. Moreover, the product in each of these summands is well-defined since the first sum condition forces $k_{i,n} = 0$ for all $n > d_0$. Thus, only a finite number of binomials can be different from one and therefore the whole expression becomes well-defined.

2.2.4. Recursion relations. To prove that the right-hand side of (17) returns (14), we consider a 1-parameter deformation of (17), and then seek a suitable q -hypergeometric difference equation in the additional parameter that is satisfied by it, broadly following the lead of [25]. For integers a, b, c, d, e , we write

$$G(a, b, c, d, e) := \sum_{\substack{\forall (i,n) \in \{1,2,3,4\} \times \mathbb{Z}_{>0}: k_{i,n} \geq 0 \\ a = \sum_{n \geq 1} \sum_{i=1}^4 (n + \delta_{i,1}) k_{i,n} \\ b = \sum_{n \geq 1} \sum_{i=1}^4 k_{i,n} \\ c = \sum_{n \geq 1} (k_{1,n} + k_{4,n}) \\ d = \sum_{n \geq 1} (k_{1,n} + k_{3,n})}} \prod_{n \geq 1} \prod_{i=1}^2 \left[\frac{e - \sum_{m \geq 1} (2m(k_{1,n+m} + k_{2,n+m}) + (2m-1)(k_{3,n+m} + k_{4,n+m}))}{k_{i,n}} \right]_q \quad (26)$$

$$\times \left[\frac{e - \sum_{m \geq 0} ((2m+1)(k_{1,n+m} + k_{2,n+m}) + 2m(k_{3,n+m} + k_{4,n+m}))}{k_{2+i,n}} \right]_q.$$

Our sought-for log Gromov–Witten generating function is obtained from (26) via the restriction

$$\mathbb{N}_d^{\log}(\text{dP}_3(0, 2)) = G(d_0, d_1, d_0 - d_2, d_0 - d_3, d_2 + d_3). \quad (27)$$

Lemma 2.5. For $a > 0, b > 0, c \geq 0, d \geq 0$, $G(a, b, c, d, e)$ satisfies the q -hypergeometric recursion

$$G(a, b, c, d, e) = \sum_{k_{1,1}, k_{2,1}, k_{3,1}, k_{4,1} \geq 0} \prod_{i=1}^2 \left[\frac{e - 2a + 2b + c + d - k_{3,1} - k_{4,1}}{k_{i,1}} \right]_q \left[\frac{e - 2a + b + c + d}{k_{i+2,1}} \right]_q \quad (28)$$

$$\times G(a - b - k_{1,1}, b - k_{1,1} - k_{2,1} - k_{3,1} - k_{4,1}, c - k_{1,1} - k_{4,1}, d - k_{1,1} - k_{3,1}, e).$$

Proof. Since $a > 0$, $b > 0$, we may rewrite (26) as

$$\begin{aligned}
G(a, b, c, d, e) = & \sum_{k_{1,1}, k_{2,1}, k_{3,1}, k_{4,1} \geq 0} \prod_{i=1}^2 \left[e^{-2a+2b+c+d-k_{3,1}-k_{4,1}} \right]_q \left[e^{-2a+b+c+d} \right]_q \\
& \times \sum_{\substack{\forall (i,n) \in \{1,2,3,4\} \times \mathbb{Z}_{>1}: k_{i,n} \geq 0 \\ a-b-k_{1,1} = \sum_{n \geq 2} \sum_{i=1}^4 (n-1+\delta_{i,1}) k_{i,n} \\ b-k_{1,1}-k_{2,1}-k_{3,1}-k_{4,1} = \sum_{n \geq 2} \sum_{i=1}^4 k_{i,n} \\ c-k_{1,1}-k_{4,1} = \sum_{n \geq 2} (k_{1,n}+k_{4,n}) \\ d-k_{1,1}-k_{3,1} = \sum_{n \geq 2} (k_{1,n}+k_{3,n})}} \prod_{m \geq 2} \prod_{i=1}^2 \left[e^{-\sum_{n \geq 1} (2n(k_{1,m+n} + k_{2,m+n}) + (2n-1)(k_{3,m+n} + k_{4,m+n}))} \right]_q \\
& \times \left[e^{-\sum_{n \geq 0} ((2n+1)(k_{1,m+n} + k_{2,m+n}) + 2n(k_{3,m+n} + k_{4,m+n}))} \right]_q. \tag{29}
\end{aligned}$$

Here we used that the sum conditions imply that

$$e - \sum_{n=1}^{a-2} (2n(k_{1,1+n} + k_{2,1+n}) + (2n-1)(k_{3,1+n} + k_{4,1+n})) = e - 2a + 2b + c + d - k_{3,1} - k_{4,1}$$

and

$$e - \sum_{n=0}^{a-1} ((2n+1)(k_{1,1+n} + k_{2,1+n}) + 2n(k_{3,1+n} + k_{4,1+n})) = e - 2a + b + c + d.$$

Now comparing (29) with (26) we obtain (28). □

Proposition 2.6. *We have that*

$$\begin{aligned}
G(a, b, c, d, e) = & \left[\begin{matrix} b-a+e \\ b-c \end{matrix} \right]_q \left[\begin{matrix} c-a+d+e \\ c \end{matrix} \right]_q \left[\begin{matrix} a-c \\ d \end{matrix} \right]_q \left[\begin{matrix} a-d \\ b-d \end{matrix} \right]_q \\
& - \left[\begin{matrix} b-a+e-1 \\ b-c \end{matrix} \right]_q \left[\begin{matrix} c-a+d+e-1 \\ c \end{matrix} \right]_q \left[\begin{matrix} a-c-1 \\ d \end{matrix} \right]_q \left[\begin{matrix} a-d-1 \\ b-d \end{matrix} \right]_q. \tag{30}
\end{aligned}$$

Proof. When at least one of the integers a, b, c, d is strictly negative, the right-hand sides of both (26) and (30) vanish. Thus, it remains to prove formula (30) for $a, b, c, d \geq 0$. We will do so by recursion in a , using Lemma 2.5.

Let us first establish the base case $a = 0$. The sum conditions in (26) in this case lead to

$$G(0, b, c, d, e) = \delta_{b,0} \delta_{c,0} \delta_{d,0},$$

in agreement with (30), as for $a = 0$ the second line must always vanish while the first one is non-zero only if $b = c = d = 0$.

Let us now move on to the induction step and consider $a > 0$. Suppose first that $b = 0$: then clearly $G(a, 0, c, d, e) = 0$. At the same time we can only expect the right-hand side of (30) to be non-zero for $b = 0$ if $c = d = 0$. However, both contributions from the two summands cancel each other, and thus in every case

$$\begin{aligned}
& \left[\begin{matrix} e-a \\ -c \end{matrix} \right]_q \left[\begin{matrix} c-a+d+e \\ c \end{matrix} \right]_q \left[\begin{matrix} a-c \\ d \end{matrix} \right]_q \left[\begin{matrix} a-d \\ -d \end{matrix} \right]_q \\
& - \left[\begin{matrix} e-a-1 \\ -c \end{matrix} \right]_q \left[\begin{matrix} c-a+d+e-1 \\ c \end{matrix} \right]_q \left[\begin{matrix} a-c-1 \\ d \end{matrix} \right]_q \left[\begin{matrix} a-d-1 \\ -d \end{matrix} \right]_q = \delta_{c,0} \delta_{d,0} - \delta_{c,0} \delta_{d,0} = 0
\end{aligned}$$

as required. We can therefore now assume that $b > 0$.

Define

$$\tilde{G}(a, b, c, d, e) := \begin{bmatrix} b - a + e \\ b - c \end{bmatrix}_q \begin{bmatrix} c - a + d + e \\ c \end{bmatrix}_q \begin{bmatrix} a - c \\ d \end{bmatrix}_q \begin{bmatrix} a - d \\ b - d \end{bmatrix}_q, \quad (31)$$

so that the right-hand side of (30) equates to $\tilde{G}(a, b, c, d, e) - \tilde{G}(a - 1, b, c, d, e - 2)$. Then a sufficient condition for (30) to hold is to show that $\tilde{G}(a, b, c, d, e)$ satisfies the same recursion (28) as $G(a, b, c, d, e)$, i.e. that

$$\begin{aligned} \tilde{G}(a, b, c, d, e) = & \sum_{k_1, k_2, k_3, k_4 \geq 0} \prod_{i=1}^2 \begin{bmatrix} e - 2a + 2b + c + d - k_3 - k_4 \\ k_i \end{bmatrix}_q \begin{bmatrix} e - 2a + b + c + d \\ k_{i+2} \end{bmatrix}_q \\ & \times \tilde{G}(a - b - k_1, b - k_1 - k_2 - k_3 - k_4, c - k_1 - k_4, d - k_1 - k_3, e). \end{aligned} \quad (32)$$

Indeed, since the coefficients of the recursion are invariant under the shift $(a, e) \rightarrow (a - 1, e - 2)$, if $\tilde{G}(a, b, c, d, e)$ satisfies (28) then so will $\tilde{G}(a - 1, b, c, d, e - 2)$. As their difference returns the claimed expression for $G(a, b, c, d, e)$ in (30), and (28) is linear in G , this would conclude the induction step.

In order to prove the stronger equality (32), we shall use the q -analogue of the Pfaff–Saalschütz summation in the form [38]

$$\sum_{k \geq 0} \frac{[A + B + C + D - k]_q!}{[k]_q! [A - k]_q! [B - k]_q! [C - k]_q! [D + k]_q!} = \begin{bmatrix} A + B + D \\ B \end{bmatrix}_q \begin{bmatrix} A + C + D \\ A \end{bmatrix}_q \begin{bmatrix} B + C + D \\ C \end{bmatrix}_q. \quad (33)$$

To use this relation we start from the right-hand side of (32), plug in the definition of \tilde{G} given in (31), expand the binomials, and collect all factorials involving k_1 and k_2 . Doing so one finds for the r.h.s. of (32) that

$$\begin{aligned} & \sum_{k_3, k_4 \geq 0} \begin{bmatrix} b - 2a + c + d + e \\ k_3 \end{bmatrix}_q \begin{bmatrix} b - 2a + c + d + e \\ k_4 \end{bmatrix}_q \\ & \times \frac{[a - b - d + k_3]_q! [2b - 2a + c + d + e - k_3 - k_4]_q!^2 [a - b - c + k_4]_q!}{[b - a + d + e - k_3]_q! [b - a + c + e - k_4]_q!} \\ & \times \sum_{k_1 \geq 0} \frac{[b - a + c + d + e - k_1 - k_3 - k_4]_q!}{[k_1]_q! [c - k_1 - k_4]_q! [d - k_1 - k_3]_q!} \\ & \quad \times \frac{1}{[2b - 2a + c + d + e - k_1 - k_3 - k_4]_q! [a - b - c - d + k_1 + k_3 + k_4]_q!} \\ & \times \sum_{k_2 \geq 0} \frac{[2b - a + e - k_2 - k_3 - k_4]_q!}{[k_2]_q! [b - c - k_2 - k_3]_q! [b - d - k_2 - k_4]_q!} \\ & \quad \times \frac{1}{[2b - 2a + c + d + e - k_2 - k_3 - k_4]_q! [a - 2b + k_2 + k_3 + k_4]_q!}. \end{aligned}$$

We can now use (33) to perform the sum over k_1 and k_2 . Collecting the factorials depending on k_3 and k_4 in the resulting expression, we get

$$\begin{aligned} & \frac{[a-b]_q! [a-c-d]_q! [b-2a+c+d+e]_q!^2}{[b-a+e]_q! [c-a+d+e]_q!} \\ & \times \sum_{k_3 \geq 0} \frac{[b-a+d+e-k_3]_q!}{[k_3]_q! [d-k_3]_q! [b-c-k_3]_q! [b-2a+c+d+e-k_3]_q! [a-b-d+k_3]_q!} \\ & \times \sum_{k_4 \geq 0} \frac{[b-a+c+e-k_4]_q!}{[k_4]_q! [c-k_4]_q! [b-d-k_4]_q! [b-2a+c+d+e-k_4]_q! [a-b-c+k_4]_q!}. \end{aligned}$$

Thus, we can use (33) again to carry out the sums over k_3 and k_4 to deduce that the right-hand side of (32) equals

$$\frac{[a-c]_q! [a-d]_q! [b-a+e]_q! [c-a+d+e]_q!}{[a-b]_q! [b-c]_q! [c]_q! [b-d]_q! [a-c-d]_q! [d]_q! [c-a+e]_q! [d-a+e]_q!}.$$

The above is exactly the expansion of $\tilde{G}(a, b, c, d, e)$ into factorials, proving (32). □

Corollary 2.7. *Conjecture 1.2 holds for any quasi-tame pair $Y(D)$.*

Proof. The tame case having been treated in [7], it suffices to restrict to non-tame, quasi-tame pairs. Taking the specialisation (27) of (30) leads to (14), which together with (12) establishes the first equality of Conjecture 1.2 for $Y(D) = \mathrm{dP}_3(0, 2)$. Since $\mathbb{O}_d(\mathrm{dP}_3^{\mathrm{op}}(0, 2)) = \mathbb{O}_d(\mathrm{dP}_3^{\mathrm{op}}(1, 1))$ [7, Sec. 6.3.1], the BPS integrality statement in the second equality further follows without any modification from the proof of [7, Thm. 8.1] for $l = 2$. Finally, since every non-tame, quasi-tame pair $Y(D)$ is related to $\mathrm{dP}_3(0, 2)$ by a series of $m \geq 0$ iterated interior blow-ups at general points of D [7, Prop. 2.2], Proposition 2.2 further implies that Conjecture 1.2 holds for any such pair. □

REFERENCES

- [1] D. ABRAMOVICH AND Q. CHEN, Stable logarithmic maps to Deligne–Faltings pairs II, *Asian J. Math.*, 18 (2014), 465–488.
- [2] S. BARDWELL-EVANS, M-W. M. CHEUNG, H. HONG, AND Y-S. LIN, Scattering Diagrams from Holomorphic Discs in Log Calabi–Yau Surfaces. Preprint available at 2110.15234.
- [3] L. J. BARROTT, Explicit equations for mirror families to log Calabi–Yau surfaces, *Bull. Korean Math. Soc.*, 57 (2020), 139–165.
- [4] P. BOUSSEAU, Quantum mirrors of log Calabi–Yau surfaces and higher-genus curve counting, *Compos. Math.*, 156 (2020), 360–411.
- [5] ———, The quantum tropical vertex, *Geom. Topol.*, 24 (2020), 1297–1379.
- [6] ———, Strong positivity for the skein algebras of the 4-punctured sphere and of the 1-punctured torus. Preprint available at 2009.02266.
- [7] P. BOUSSEAU, A. BRINI, AND M. VAN GARREL, Stable maps to Looijenga pairs. Preprint available at 2011.08830.
- [8] ———, Stable maps to Looijenga pairs: orbifold examples, *Lett. Math. Phys.*, 111 (2021), 109. Preprint available at 2012.10353.
- [9] ———, On the log-local principle for the toric boundary, *Bull. Lond. Math. Soc.*, in press (2022). Preprint available at 1908.04371.
- [10] Q. CHEN, Stable logarithmic maps to Deligne–Faltings pairs I, *Ann. of Math. (2)*, 180 (2014), 455–521.
- [11] T.M. CHIANG, A. KLEMM, S-T. YAU, AND E. ZASLOW, Local mirror symmetry: Calculations and interpretations, *Adv. Theor. Math. Phys.*, 3 (1999), 495–565. Preprint available at hep-th/9903053.
- [12] T. COATES AND A. GIVENTAL, Quantum Riemann–Roch, Lefschetz and Serre, *Ann. of Math. (2)*, 165 (2007), 15–53.
- [13] B. DAVISON AND T. MANDEL, Strong positivity for quantum theta bases of quantum cluster algebras. Preprint available at 1910.12915.
- [14] R. FRIEDMAN, On the geometry of anticanonical pairs. Preprint available at 1502.02560.
- [15] M. GROSS, P. HACKING, AND S. KEEL, Birational geometry of cluster algebras, *Algebr. Geom.*, 2 (2015), 137–175.
- [16] ———, Mirror symmetry for log Calabi–Yau surfaces I, *Publ. Math. Inst. Hautes Études Sci.*, 122 (2015), 65–168.
- [17] M. GROSS, R. PANDHARIPANDE, AND B. SIEBERT, The tropical vertex, *Duke Math. J.*, 153 (2010), 297–362.
- [18] M. GROSS AND B. SIEBERT, Logarithmic Gromov–Witten invariants, *J. Amer. Math. Soc.*, 26 (2013), 451–510.

- [19] P. HACKING AND A. KEATING, Homological mirror symmetry for log Calabi–Yau surfaces. Preprint available at 2005.05010.
- [20] E. N. IONEL AND T. H. PARKER, The Gopakumar–Vafa formula for symplectic manifolds, *Ann. of Math. (2)*, 187 (2018), 1–64.
- [21] A. IQBAL AND A-K. KASHANI-POOR, The vertex on a strip, *Adv. Theor. Math. Phys.*, 10 (2006), 317–343. Preprint available at hep-th/0410174.
- [22] S. H. KATZ AND C-C. M. LIU, Enumerative geometry of stable maps with Lagrangian boundary conditions and multiple covers of the disc, *Geom. Topol. Monogr.*, 8 (2002), 1–47. Preprint available at math/0103074.
- [23] A. KLEMM AND R. PANDHARIPANDE, Enumerative geometry of Calabi–Yau 4-folds, *Commun. Math. Phys.*, 281 (2008), 621–653. Preprint available at math/0702189.
- [24] M. KONTSEVICH AND Y. SOIBELMAN, Affine structures and non-Archimedean analytic spaces, pp. 321–385, 2006.
- [25] C. KRATTENTHALER, Proof of two multivariate q -binomial sums arising in Gromov–Witten theory. Preprint available at 2102.02360.
- [26] J. M. F. LABASTIDA AND M. MARINO, Polynomial invariants for torus knots and topological strings, *Commun. Math. Phys.*, 217 (2001), 423–449. Preprint available at hep-th/0004196.
- [27] J. M. F. LABASTIDA, M. MARINO, AND C. VAFA, Knots, links and branes at large N , *JHEP*, 11 (2000), 007. Preprint available at hep-th/0010102.
- [28] J. LI, C-C. M. LIU, K. LIU, AND J. ZHOU, A mathematical theory of the topological vertex, *Geom. Topol.*, 13 (2009), 527–621. Preprint available at math/0408426.
- [29] J. LI AND Y. S. SONG, Open string instantons and relative stable morphisms, *Adv.Theor.Math.Phys.*, 5 (2002), 67–91. Preprint available at hep-th/0103100.
- [30] M. MANABE, Topological open string amplitudes on local toric del Pezzo surfaces via remodeling the B-model, *Nucl. Phys.*, B819 (2009), 35–75. Preprint available at 0903.2092.
- [31] T. MANDEL, Scattering diagrams, theta functions, and refined tropical curve counts. Preprint available at 1503.06183.
- [32] ———, Tropical theta functions and log Calabi–Yau surfaces, *Selecta Math. (N.S.)*, 22 (2016), 1289–1335.
- [33] ———, Classification of rank 2 cluster varieties, *SIGMA Symmetry Integrability Geom. Methods Appl.*, 15 (2019), Paper 042, 32.
- [34] ———, Theta bases and log Gromov–Witten invariants of cluster varieties. Preprint available at 1903.03042.
- [35] H. OOGURI AND C. VAFA, Knot invariants and topological strings, *Nucl. Phys.*, B577 (2000), 419–438. Preprint available at hep-th/9912123.
- [36] T. Y. YU, Enumeration of holomorphic cylinders in log Calabi–Yau surfaces. I, *Math. Ann.*, 366 (2016), 1649–1675.
- [37] ———, Enumeration of holomorphic cylinders in log Calabi–Yau surfaces. II. Positivity, integrality and the gluing formula, *Geom. Topol.*, 25 (2021), 1–46. Preprint available at 1608.07651.
- [38] D. ZEILBERGER, A q -Foata proof of the q -Saalschütz identity, *European J. Combin.*, 8 (1987), 461–463.
- [39] Y. ZHOU, Weyl groups and cluster structures of families of log Calabi–Yau surfaces. Preprint available at 1910.05762.