

Knot cobordism and Lee's Perturbation of Khovanov homology

Zipei Zhuang

January 7, 2022

Abstract

For a connected cobordism S between two knots K_1, K_2 in S^3 , we establish an inequality involving the number of local maxima, the genus of S , and the torsion orders of $Kh_t(K_1), Kh_t(K_2)$, where Kh_t denotes Lee's perturbation of Khovanov homology. This shows that the torsion order gives a lower bound for the band-unlinking number.

If K_0, K_1 are knots in S^3 , a cobordism from K_0 to K_1 is a smooth compact oriented surface F embedded in $S^3 \times [0, 1]$ with boundary $K_0 \times \{0\} \cup K_1 \times \{1\}$. If F is an annulus, it is called a concordance. Any cobordism $F \subset S^3 \times [0, 1]$ is called *ribbon* if the projection from $S^3 \times [0, 1]$ to $[0, 1]$ restricts to a Morse function on F with only index 0 and 1 critical points. A knot K_0 is *ribbon concordant* to K_1 if there exists a ribbon concordance from K_0 to K_1 .

In the past few years there has been several works on applications of knot homology theories to knot cobordisms. For Khovanov homology Kh , it is shown [9] that a ribbon concordance between two links (F, K_0, K_1) induces an injective map $\phi_F : Kh(K_0) \rightarrow Kh(K_1)$. Furthermore, Sarkar [11] showed that if F has k saddles, then $(2x)^d Kh_t(K_0) \cong (2x)^d Kh_t(K_1)$, where Kh_t is Lee's perturbation of Khovanov homology. Similar results are obtained in the context of knot Floer homology (see [12] [5]).

The main result of this paper is the following:

Theorem 1. *Suppose that (F, K_0, K_1) is a connected knot cobordism with m births, b saddles and M deaths. Denote by \bar{F} the cobordism from K_1 to K_0 obtained by horizontally mirroring \bar{F} . Then up to a sign*

$$(2x)^M \cdot \phi_{\bar{F}} \circ \phi_F = (2x)^{b-m} \cdot id_{Kh_t(K_0)} \quad (1)$$

This is analogous to the main result of [5], which is for knot Floer homology.

In particular, if F is a ribbon concordance, then $M = b - m = 0$, so $\phi_{\bar{F}} \circ \phi_F = id_{Kh_t(K_0)}$, ϕ_F is injective.

Let $\mathcal{T}(K)$ be the torsion part of $Kh_t(K)$, and $xo(K)$ be the *extortion order* of K , i.e. the minimal n such that $x^n \mathcal{T}(K) = 0$. The same argument as in [5] gives the following corollary:

Corollary 1.

$$xo(K_0) \leq \max\{M, xo(K_1)\} + 2g(F) \quad (2)$$

where $g(F)$ is the genus of F .

The *band-unlinking number* $ul_b(K)$ of a knot K is the minimum number of oriented band moves necessary to reduce K to an unlink. Corollary 1 further gives

Corollary 2.

$$xo(K) \leq ul_b(K) \quad (3)$$

1 Lee's perturbation of Khovanov homology

We will work over a field \mathbb{F} in which 2 is invertible. Let \mathbf{A} be the 2-dimensional Frobenius algebra over $\mathbb{F}[t]$ with basis $\{1, x\}$, with the multiplication map m , comultiplication map Δ , and counit map ϵ defined as follows:

$$\begin{aligned} 1 \otimes 1 &\xrightarrow{m} 1 & 1 \otimes x &\xrightarrow{m} x \\ x \otimes 1 &\xrightarrow{m} x & x \otimes x &\xrightarrow{m} t \\ 1 &\xrightarrow{\Delta} 1 \otimes x + x \otimes 1 & & \\ x &\xrightarrow{\Delta} x \otimes x + t \cdot 1 \otimes 1 & & \\ 1 &\xrightarrow{\epsilon} 0 & x &\xrightarrow{\epsilon} 1 \end{aligned} \quad (4)$$

Applying this Frobenius algebra to the Kauffman cube of resolutions of a knot diagram, we get a chain complex over $\mathbb{F}[t]$, whose homology is called Lee's perturbation of Khovanov homology [3]. In particular, setting $t = 0$, we get Khovanov's original knot homology; taking $t = 1$, it becomes Lee homology [8].




Figure 1: The elementary cobordisms

A knot cobordism (F, K_0, K_1) induces a map (see [6] [4] [3])

$$\phi_F : Kh_t(K_0) \longrightarrow Kh_t(K_1) \quad (5)$$

by decomposing F into a sequence of elementary cobordisms, each represented by a Reidemeister move or a Morse move of the link diagrams. The Reidemeister moves induce isomorphisms on homology. The 0-Morse move (corresponding to a birth), 1-Morse move (saddle), 2-Morse move (death) induce maps on Kh_t according to the unit map, the multiplication/comultiplication map, the counit map, respectively. The map induced by F is the composition of the maps induced by these elementary cobordisms. Furthermore, one can consider dotted cobordisms, where a dot induces a multiplication by x on homology. Such assignment is well-defined and functorial, **up to a sign**.

The cobordism maps satisfy the following local properties(see [11]):

(1) Suppose F has an 1-handle h . Let F' be obtained from F by surgery along h , and let \dot{F}_1 and \dot{F}_2 be obtained by adding a dot to F' at either of the feet of h . Then (see Fig.2) up to a sign,

$$\phi_F = \phi_{\dot{F}_1} + \phi_{\dot{F}_2} \quad (6)$$




Figure 2:

(2) If F is a cobordism which can be decomposed into a sequence F_1, \dots, F_n in which F_i, F_{i+1} are two adjacent saddle moves reverse to each other, let S be the cobordism obtained from F by deleting F_i, F_{i+1} , and let \dot{S}_1, \dot{S}_2 be the cobordism obtained by adding a dot on one of the two sides of S (see Fig.3). Then up to a sign

$$\phi_F = \phi_{\dot{S}_1} + \phi_{\dot{S}_2} \quad (7)$$

2 The cobordism distance

Proof of Theorem 1.

Proof. Let F be a connected oriented knot (dotted) cobordism from K_0 to K_1 in $S^3 \times [0, 1]$ with m births, b saddles and M deaths. We can rearrange the critical points of F so that F can be decomposed into a sequence of elementary cobordisms ordered as below (we omit the cobordisms of Reidemeister moves since they induce isomorphisms on homology):




Figure 3:

- (1) m births, which add m unknots U_1, \dots, U_m ;
- (2) m fusion saddle moves, which merge U_1, \dots, U_m with K_0 ;
- (3) $b - m$ additional saddle moves;
- (4) M deaths, deleting unknots U'_1, \dots, U'_M .

So the cobordism S decomposes as $K_1 \xrightarrow{F_1} K_1 \sqcup U^m \xrightarrow{F_2} K' \xrightarrow{F_3} K_2 \sqcup U^M \xrightarrow{F_4} K_2$, where the piece F_i comes from item (i) above, and U^m denotes the m -component planar unlink. The cobordism \bar{F} decomposes as $K_2 \xrightarrow{\bar{F}_4} K_2 \sqcup U^M \xrightarrow{\bar{F}_3} K' \xrightarrow{\bar{F}_2} K_1 \sqcup U^m \xrightarrow{\bar{F}_1} K_1$. Then $\phi_{\bar{F}} \circ \phi_F$ is the composition of 8 maps. The composition of the fourth and fifth step is to delete the M unknots via M deaths, then add them back with M births. Let S be the cobordism obtained by deleting the fourth and fifth step. Then $\bar{F} \circ F$ is obtained from S by sugerying along the M tubes. By 6, we have

$$\phi_S = (2x)^M \phi_{\bar{F} \circ F} \quad (8)$$

since $\bar{F} \circ F$ is connected.

Let T be the cobordism obtained from S by deleting the third and sixth steps, i.e. T is the composition $K_1 \xrightarrow{F_1} K_1 \sqcup U^m \xrightarrow{F_2} K' \xrightarrow{\bar{F}_2} K_1 \sqcup U^m \xrightarrow{\bar{F}_1} K_1$. By 7, we have

$$\phi_S = (2x)^{b-m} \cdot \phi_T \quad (9)$$

Note that $F_2 \circ F_1$ is a ribbon concordance, by [9] Theorem 1,

$$\phi_T = id_{K_{h_t}(K_0)} \quad (10)$$

Theorem 1 now follows from equation 8 9 10. □

Proof of Corollary 1

Proof. Let F be an oriented knot cobordism from K_0 to K_1 with m births, b saddles, M deaths. For $l \geq 0, \alpha \in Kh_t(K_0)$, by Theorem 1 we have

$$\phi_{\bar{F} \circ F}((2x)^{l+M} \cdot \alpha) = (2x)^M \phi_{\bar{F} \circ F}((2x)^l \cdot \alpha) = (2x)^{l+b-m} \cdot \alpha \quad (11)$$

This shows that if $l \geq \max\{0, xo(K_1) - M\}$, then

$$(2x)^{l+b-m} \cdot \alpha = \phi_{\bar{F}} \circ \phi_F((2x)^{l+M} \cdot \alpha) = 0 \quad (12)$$

Therefore $xo(K_0) \leq b - m + \max\{0, xo(K_1) - M\} = \max\{M, xo(K_1)\} + 2g$ (recall that we work over a base field in which 2 is invertible) since $2g(F) = -\chi(F) = b - m - M$. □

Proof of Corollary 2

Proof. By definition, after attaching $ul_b(K)$ oriented bands to K , we obtain an unlink L . Suppose L has M components. By performing $M - 1$ deaths, we obtain a cobordism S from K to an unknot U with 0 births, $ul_B(K)$ saddles and $M - 1$ deaths. Then by Corollary 1 we have

$$xo(K) \leq \max\{M - 1, 0\} + 2g(S) = M - 1 + ul_b(K) - M + 1 = ul_b(K). \quad (13)$$

□

The following corollary generalizes [11] Theorem 1.1:

Corollary 3. *If F is a knot concordance from K_0 to K_1 with b saddles. Then*

$$(2x)^b Kh_t(K_0) \cong (2x)^b Kh_t(K_1) \quad (14)$$

Proof. Suppose F has m births and M deaths. Since F is an annulus, $\chi(F) = m - b + M = 0, M = b - m$. By Theorem 1,

$$(2x)^M \phi_{\bar{F}} \circ \phi_F = (2x)^{b-m} \cdot Id_{Kh_t(K_0)} = (2x)^M Id_{Kh_t(K_0)} \quad (15)$$

Note that $b \geq M$, for any $\alpha \in Kh_t(K_0)$,

$$\phi_{\bar{F}} \circ \phi_F((2x)^b \cdot \alpha) = (2x)^M \phi_{\bar{F}} \circ \phi_F((2x)^{b-M} \cdot \alpha) = (2x)^b \cdot \alpha \quad (16)$$

Therefore $\phi_{\bar{F}} \circ \phi_F$ is the identity map on $(2x)^b \cdot Kh_t(K_0)$. Reversing the role of F and \bar{F} , we get $\phi_F \circ \phi_{\bar{F}}$ is the identity map on $(2x)^b \cdot Kh_t(K_1)$. So $\phi_F : (2x)^b \cdot Kh_t(K_0) \rightarrow (2x)^b \cdot Kh_t(K_1)$ is an isomorphism. □

References

- [1] Akram Alishahi. Unknotting number and Khovanov homology. *Pacific J. Math.*, 301(1):15–29, 2019.
- [2] Akram Alishahi and Eaman Eftekhary. Knot Floer homology and the unknotting number. *Geom. Topol.*, 24(5):2435–2469, 2020.
- [3] Dror Bar-Natan. Khovanov’s homology for tangles and cobordisms. *Geom. Topol.*, 9:1443–1499, 2005. 1, 1
- [4] Magnus Jacobsson. An invariant of link cobordisms from Khovanov homology. *Algebr. Geom. Topol.*, 4:1211–1251, 2004. 1
- [5] András Juhász, Maggie Miller, and Ian Zemke. Knot cobordisms, bridge index, and torsion in Floer homology. *J. Topol.*, 13(4):1701–1724, 2020. (document)
- [6] Mikhail Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000. 1
- [7] Mikhail Khovanov. An invariant of tangle cobordisms. *Trans. Amer. Math. Soc.*, 358(1):315–327, 2006.
- [8] Eun Soo Lee. An endomorphism of the Khovanov invariant. *Adv. Math.*, 197(2):554–586, 2005. 1
- [9] Adam Simon Levine and Ian Zemke. Khovanov homology and ribbon concordances. *Bull. Lond. Math. Soc.*, 51(6):1099–1103, 2019. (document), 2
- [10] Jacob Rasmussen. Khovanov homology and the slice genus. *Invent. Math.*, 182(2):419–447, 2010.
- [11] Sucharit Sarkar. Ribbon distance and Khovanov homology. *Algebr. Geom. Topol.*, 20(2):1041–1058, 2020. (document), 1, 2
- [12] Ian Zemke. Knot Floer homology obstructs ribbon concordance. *Ann. of Math. (2)*, 190(3):931–947, 2019. (document)