MAXLA-5/21, "GenSol SE" January 10, 2022

General solution of the Schrödinger equation

MIKHAIL N. SERGEENKO

Institute of Radiobiology of the National Academy of Sciences of Belarus BY-246007, Gomel, Belarus msergeen@gmail.com

Abstract

The wave equation in quantum mechanics and its general solution in the phase space are obtained.

 $\mathrm{Belarus}-2021$

The Schrödinger wave (SW) equation in quantum mechanics (QM) is usually solved in terms of special functions or numerically. We use here the abbreviation SW which means "Schrödinger's wave" known as ψ -function. The general approach to solve the SW equation is to reduce its to the equation for hypergeometric function or some special function. To do that one needs to find first a special transformation for the wave function (w.f.) and its arguments to reduce the original equation to the hypergeometric form. However, the solution of the SW equation can be obtained in elementary functions.

The static one-dimensional (1D) SW equation for a free particle of mass m [1–3],

$$\frac{1}{2m} \left(-i\hbar \frac{d}{dx} \right)^2 \psi(x) = E\psi(x), \tag{1}$$

has a general solution in the form of a superposition of two plane waves in configuration space,

$$\psi(x) = C_1 e^{ikx} + C_2 e^{-ikx}.$$
(2)

Here C_1 and C_2 are (in general) complex constants, $k = p/\hbar$ is the wave number defined as $k = 2\pi/\lambda_B$ for the de Broglie wavelength $\lambda_B = h/p$ with the particle constant momentum $p = \sqrt{2mE}$, $h = 2\pi\hbar$ is the Plank's constant.

There is another approach to solving the SW equation. It is known that the SW equation can be derived with the help of the Bohr's correspondence principle [1]. This fundamental principle has been used at the stage of creation of quantum theory. It is used to establish correspondence between classical functions and operators of QM, and to derive the apparent form of the operators. Moreover, the correspondence principle points out the way to a simplest solution of the SW equation.

The correspondence principle states that the laws of quantum physics must be so chosen that in the classical limit, where many quanta are involved, the quantum laws lead to the classical equations as an average. In this way, in [4] this principle has been used to derive the non-relativistic quasi-classical (QC) wave equation appropriate in the QC region and the relativistic QC wave equation [5].

We transform (1), which is a problem on eigenvalues of energy E, to the equivalent form

$$\left(-i\hbar\frac{d}{dx}\right)^2\psi(x) = p_E^2\psi(x),\tag{3}$$

which is a problem on eigenvalues of $p_E^2 = (\hbar k)^2 = 2mE$, and introduce the dimensionless

phase variable $\phi = kx$. This gives the SW equation in the phase space,

$$\psi_{\phi\phi}^{\prime\prime} + \psi = 0, \tag{4}$$

which is the linear homogeneous second-order differential (LHD₂) equation in canonical form. The general solution of this equation is given by a superposition of two plane waves in the phase space $\{\phi\}$,

$$\psi(\phi) = C_1 e^{i\phi} + C_2 e^{-i\phi}.$$
 (5)

The dimensionless phase variable $\phi = W/\hbar$ is written in terms of the reduced *classical* action, W = px, for a free particle. The purpose of this letter is to show that the general solution of the SW equation for the case of an interacting particle has the same form (5) for the corresponding *reduced classical action* W.

In QM, the concept of "action in QM" is used. In the semiclassical approximation, the "action in QM" is used in the form of an expansion in terms of the Plank's constant \hbar : $S = S_0 + \hbar S_1 + \ldots$ [6]. But, in the quantization condition the *classical* action S_0 is used, and the corrections S_1, S_2, \ldots have no physical meaning, but lead to artificial problems such as divergence of the solution at the turning points. In our opinion, the concept of "action in QM" is NOT necessary for the theory. Let us show that the SW equation and its solution can be obtained from the *classical* action.

In the general case of an interacting particle, we consider a conservative system when the Hamiltonian $H(x, p) = p^2/2m + V(x)$ is not an explicit function of time t. The SW equation for the interaction potential V(x),

$$\left[\frac{1}{2m}\left(-i\hbar\frac{d}{dx}\right)^2 + V(x)\right]\psi(x) = E\psi(x),\tag{6}$$

can be written in the Sturm-Liouville canonical form

$$\psi_{xx}'' + [k^2 - U^2(x)]\psi = 0, \tag{7}$$

where $k^2 = 2mE/\hbar^2$, $U^2(x) = 2mV(x)/\hbar^2$, $p^2(x) = k^2 - U^2(x)$.

In this case the Hamilton's principal function (the generating function) is

$$\psi(t, x) = Ce^{i[-Et+W(x)]/\hbar}, \quad W(x) = \int^x p(x)dx.$$
 (8)

The second derivative of this function with respect to the variable x is

$$\psi_{xx}'' = \left[\left(\frac{i}{\hbar} p \right)^2 + \frac{i}{\hbar} \frac{dp}{dx} \right] \psi, \tag{9}$$

where p(x) = (dW/dx). Multiplying (9) by the constant $(\hbar/i)^2$, we get

$$\left(\frac{\hbar}{i}\frac{d}{dx}\right)^2\psi = \left[2m[E-V(x)] + \frac{\hbar}{i}\frac{dp}{dx}\right]\psi.$$
(10)

The equality (10) gives the SW equation (6) if

$$\frac{\hbar}{i}\frac{dp}{dx} = 0 \quad \Rightarrow \quad \bar{p} = \text{const}, \tag{11}$$

What does this mean? The operators in QM are Hermitian, i.e. their eigenvalues are real. The operator \hat{p}^2 in the left hand side of (10) is Hermitian, if the condition (11) is met. The Plank's constant is *constant* value, not zero, hence, the derivative $p'_x = 0$. Here in (11) $\bar{p} = \bar{k}\hbar$ is momentum eigenvalue (constant value) and $\phi = \bar{k}x$ is the phase variable.

The condition (11) is the key one in solving the SW equation and supplies the *Hermiticity* of the squared momentum operator, \hat{p}^2 , in (10). We emphasize that only if the condition (11) is satisfied, it is possible to obtain from (10) the SW equation (6). Functions in mathematics can take both continuous and discrete values for specific values of argument. The requirement (11) is a prerequisite for permitted movements in QM; it defines *allowed motions* or stationary states of a quantum system, i.e., the generalized momentum p(x) can only take some constant discrete values (momentum eigenvalues \bar{p}). Thus, equality (11) leads from (10) to the SW Eq. (6) and its solution of the form (8).

Transform (7) to the dimensionless phase variable $\phi(x) = W(x)/\hbar$. This can be done using the identity

$$\frac{d}{dx}\left(f\frac{dy}{dx}\right) = \left(\sqrt{f}\frac{d^2}{dx^2} - \frac{d^2}{dx^2}\sqrt{f}\right)\left(\sqrt{f}y\right) \tag{12}$$

that gives the equation

$$\Psi_{\phi\phi}'' + [1 - \delta(\phi)]\Psi = 0, \tag{13}$$

where $\Psi = \sqrt{p}\psi$. The function (functional)

$$\delta(\phi) = \frac{1}{\sqrt{p}} \frac{d^2 \sqrt{p}}{d\phi^2} = \frac{1}{2} \frac{d\epsilon}{d\phi} + \frac{1}{4} \epsilon^2, \tag{14}$$

$$\epsilon = \frac{\hbar}{p^2} \frac{dp}{dx} \tag{15}$$

has the meaning of a potential in the phase space with the properties: $1 \delta(\phi) = 0$ for $V(x) = 0, 2 \delta(\phi) = \infty$ at the turning points (TPs) given by the Eq. $p(x) = 0, 3 \delta(\phi)$ is a small quantity of higher order at other points [6]. We can call (13) the SW equation in

the phase space. The fulfillment of the inequality $\epsilon \ll 1$ is a necessary condition for the application of the QC approximation in QM.

The roots of the equation p(x) = 0 separate the classically allowed region where $p(x) \ge 0$ from the classically forbidden region where p(x) < 0. The function $\delta(\phi)$ in (13) has the properties of the δ -function: according to (11) the quantity $\epsilon = k'_x/k^2 = p'_x/p^2 = 0$, therefore, $\delta(\phi) = 0$ excluding the TPs determined by the equality (11); in the TPs $\delta(\phi) = \infty$. The corresponding equations in these regions follow from (13) and are given by the system in the phase space { ϕ [7]

$$\begin{cases} \Psi_{\phi\phi}'' + \Psi = 0, \quad p(x) > 0, \\ \Psi_{\phi\phi}'' - \Psi = 0, \quad p(x) < 0. \end{cases}$$
(16)

The general solutions of these equations are

$$\begin{cases} \Psi(\phi) = Ae^{i\phi} + Be^{-i\phi}, \quad p(x) > 0, \\ \Psi(\phi) = Ce^{\phi} + De^{-\phi}, \quad p(x) < 0. \end{cases}$$
(17)

Solution in QM (w.f.) must be continuous and finite in the entire range $(-\infty, \infty)$. To build the physical solution in the entire range we need to merge the oscillating solution(s) in classically allowed region where $p(x) \ge 0$ with the exponentially decaying solution(s) in classically inaccessible regions where p(x) < 0 [4].

The functions (17) should smoothly merge into each other at the turning points. Matching these functions and their first derivatives at the turning point x_k gives two equalities

$$\begin{cases}
A+B=C+D, \\
iA-iB=-C+D,
\end{cases}$$
(18)

which yields

$$\begin{cases} A = \left(Ce^{i\pi/4} + De^{-i\pi/4} \right) / \sqrt{2}, \\ B = \left(Ce^{-i\pi/4} + De^{i\pi/4} \right) / \sqrt{2}. \end{cases}$$
(19)

The connection formulas (19) supply the continuous transition of the general solutions (17) into each other at the turning point x_k .

The most popular and important in applications are the two-turning point (2TP) problems [4, 7]. For the 2TP problem, the entire interval $(-\infty, \infty)$ is divided by the TPs x_1 and x_2 into three regions. This leads with the help of the connection formulas (19) to the quantization condition [4, 7]

$$\int_{x_1}^{x_2} \sqrt{2m[E - V(x)]} dx = \pi \hbar \left(n + \frac{1}{2} \right).$$
 (20)

The final solution for the 2TP problems (the state function) in the phase space is [7]

$$\Psi_n(\phi) = C_n \begin{cases} e^{\phi - \phi_1}, & x < x_1, \\ \sqrt{2}\cos(\phi - \phi_1 - \frac{\pi}{4}), & x_1 \le x \le x_2, \\ (-1)^n e^{-\phi + \phi_2}, & x > x_2, \end{cases}$$
(21)

where $\phi(n, x) = k_n x$, $\phi_1 = \phi(x_1) = -\pi (n + \frac{1}{2})/2$, $\phi_2 = \phi(x_2) = \pi (n + \frac{1}{2})/2$ [4]. Here we have took into account the fact that, for the stationary states, the phase-space variable $\phi(n, x)$ at the TPs x_1 and x_2 depends on quantum number n and does not depend on the form of the potential. The normalization coefficient,

$$C_n = \sqrt{\frac{k_n}{\pi(n+\frac{1}{2})+1}},$$
(22)

is calculated from the normalization condition $\int_{-\infty}^{\infty} |\psi_n(x)|^2 dx = 1.$

The solution (21) describes *free motion* of a particle-wave in the enclosure (the enclosure being the interaction potential). Therefore, in bound state region, the interaction of the particle-wave with the potential reduces to reflection of the wave by "walls of the potential". The "classical" solution (21) is general for all types of 2TP problems and allows to solve multi-turning point problems which represent a class of the "insoluble" (by standard methods) problems with more than two turning points [5].

The oscillating part of (21),

$$\Psi_n(x) = \sqrt{\frac{2k_n}{\pi(n+\frac{1}{2})+1}} \cos\left(k_n x + \frac{\pi}{2}n\right),$$
(23)

has the form of a standing wave. The form of the phase variable $\phi(n, x) = k_n x + \pi n/2$ guaranties that the state functions $\Psi_n(\phi)$ are necessarily either symmetric (n = 0, 2, 4, ...)or antisymmetric (n = 1, 3, 5, ...). The function (23) corresponds to the principal term of the asymptotic series in theory of the LHD₂ equations, which in QM gives the asymptote of the exact solution of the SW equation. The quantization condition (20) in our solution of the 1D SW equation (6) is not approximate. It is exact, i.e. (20) reproduces the exact energy spectra for all known solvable 2TP problems in QM [4, 7].

The case of three-dimensional (3D) problems was studied in our works [4, 8]. The derivation of the 3D wave equation (30) can be performed siminarly to 1D case given above. Consider the case of a central potential. In spherical coordinates, the variables are separated and the generating function is

$$\psi(t, \mathbf{r}) = C e^{-i[Et - W(\mathbf{r})]/\hbar},\tag{24}$$

where $W(\mathbf{r}) = W(r) + W(\theta) + W(\varphi)$. The first derivative gives

$$\vec{\nabla}\psi(t,\,\mathbf{r}) = \frac{i}{\hbar}(\vec{\nabla}W)\psi(t,\,\mathbf{r}).$$
(25)

The second derivative results in the SW equation,

$$\left[\frac{(-i\hbar\vec{\nabla})^2}{2m} + V(r)\right]\psi(\vec{r}) = E\psi(\vec{r}).$$
(26)

A QC analysis of (26) was performed [8]. The SW equation (26) was reduced to the form of the classical HJ equation. Separation of the resulting equation using the correspondence principle results in the three 1D equations in canonical form (7),

$$\left[\hbar^2 \frac{d^2}{dr^2} + 2m(E - V) - \frac{\vec{M}^2}{r^2}\right] \mathbf{R}(r) = 0,$$
(27)

$$\left[\hbar^2 \frac{d^2}{d\theta^2} + \vec{M}^2 - \frac{M_z^2}{\sin^2 \theta}\right] \Theta(\theta) = 0, \qquad (28)$$

$$\left[\hbar^2 \frac{d^2}{d\varphi^2} + M_z^2\right] \Phi(\varphi) = 0, \qquad (29)$$

where \vec{M}^2 , M_z^2 are the constants of separation and, at the same time, integrals of motion [4]. These equations are equivalent to the 3D equation

$$\left[(-i\hbar)^2 \Delta^c + U(r) \right] \Psi(\vec{r}) = p_E^2 \Psi(\vec{r}), \tag{30}$$

$$\Delta^{c} = \frac{\partial^{2}}{\partial r^{2}} + \frac{1}{r^{2}} \frac{\partial^{2}}{\partial \theta^{2}} + \frac{1}{r^{2} \sin^{2} \theta} \frac{\partial^{2}}{\partial \varphi^{2}}.$$
(31)

Here (30) is the wave equation in canonical form for the eigenvalue of the square of the momentum $p_E^2 = 2mE$; Δ^c is the canonical operator, U(r) = 2mV(r). Note that (30) is not the SW equation.

Solution of (30) reproduces the exact energy spectra for *all* known solvable 2TP problems in QM [4, 7] and "insoluble" potentials with more than two turning points [5]. More examples of application of our solution are in [9].

Summarizing, we have shown that the generating function (8), expressed in terms of the Hamilton's principal function, is the key object in the solution of the SW equation. In our solution, there is no place for the "action in quantum mechanics". This means we can

start consideration with the classical action and continue by its quantization. The general solution of the SW equation is given by the system (17).

In our solution method, we use the same technique for all types of problems. The same simple rules and the general solution (17) formulated for 2TP problems work for multi-TP problems, as well. This approach can be easily generalized for the non-separable problems. In this case the classical action can not be written as sum of actions for each variable (degree of freedom). As a result, we cannot have a unique quantum number for each degree of freedom. We will have one quantum number for the non-separable variables, and unique quantum number for each separable variable. The quantization condition will be a multidimensional integral over the non-separable variables. In this sense, our approach can be considered as a general method for finding the solution of the SW equation.

- [1] D. Bohm, *Quantum Theory* (Dover Publications, 1979).
- [2] L. Schiff, Quantum mechanics, second ed. ed. (McGraw-Hill, New York, 1955).
- [3] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics (Non-Relativistic Theory)*, 3rd ed., Vol. 3 (Pergamon, 1977) iSBN: 978-0-08-020940-1.
- [4] M. N. Sergeenko, Phys. Rev. A53, 3798 (1996), [arXiv: quant-ph/9911075].
- [5] M. N. Sergeenko, Mod. Phys. Lett. A12, 2859 (1997), [arXiv:quant-ph/9911081].
- [6] N. Fröman and P. O. Fröman, JWKB Approximation: Contributions to the Theory (North Holland, Amsterdam, 1965).
- [7] M. N. Sergeenko, Int. J. Mod. Phys. A18, 3041 (2003), [arXiv:quant-ph/0010084v2].
- [8] M. N. Sergeenko, Mod. Phys. Lett. A15, 83 (2000), [arXiv:quant-ph/9912069].
- [9] M. N. Sergeenko, Eur. Phys. J. C72, 2128 (2012), [arXiv:hep-ph/1206.7099].