

Revisiting the gluino mass limits in the pMSSM in the light of the latest LHC data and Dark Matter constraints

Abhi Mukherjee ^{*1}, Saurabh Niyogi ^{†2}, Sujoy Poddar ^{‡3}

¹Department of Physics, University of Kalyani, Kalyani 741235, India

²Gokhale Memorial Girls' College, 1/1 Harish Mukherjee Road, Kolkata 700 020, India

³Department of Physics, Diamond Harbour Women's University, Diamond Harbour Road, Sarisha, South 24 Parganas, West Bengal 743368, India

Abstract

The purpose of this paper is to examine the model dependence of the stringent constraints on the gluino mass obtained from the Large Hadron Collider (LHC) experiments by analyzing the Run II data using specific simplified models based on several ad hoc sparticle spectra which cannot be realized even in the fairly generic pMSSM models. We first revisit the bounds on the gluino mass placed by the ATLAS collaboration using the $1l + jets + \cancel{E}_T$ data. We show that the exclusion region in the $M_{\tilde{g}} - M_{\tilde{\chi}_1^0}$ plane in the pMSSM scenario sensitively depends on the mass hierarchy between the left and right squarks and composition of the lighter electroweakinos and, to a lesser extent, other parameters. Most importantly, for higgsino type lighter electroweakinos (except for the LSP), the bound on the gluino mass from this channel practically disappears. However, if such models are confronted by the ATLAS $jets + \cancel{E}_T$ data, fairly strong limits are regained. Thus in the pMSSM an analysis involving a small number of channels may provide more reliable mass limits. We have also performed detailed analyses on neutralino dark matter (DM) constraints in the models we have studied and have found that for a significant range of

*abhiphys18@klyuniv.ac.in

†saurabhphys@gmail.com

‡sujoy.phy@gmail.com

LSP masses, the relic density constraints from the WMAP/PLANCK data are satisfied and LSP-gluino coannihilation plays an important role in relic density production. We have also checked the simultaneous compatibility of the models studied here with the direct DM detection, the IceCube data, and the LHC constraints.

1 Introduction

Supersymmetry (SUSY) [1–8], which allows inter-conversion of fermions and bosons, is one of the most well motivated and widely studied extension of the Standard Model (SM). It offers solutions to several short-comings of the SM like, the hierarchy problem [9, 10], the unification of gauge couplings [11, 12] at a scale, $M_G \simeq 10^{16}$ GeV and it contains a viable Dark Matter (DM) candidate [13–21]. SUSY can also trigger Electro-Weak Symmetry Breaking (EWSB) [22–24] which is an ad hoc phenomenon within the realm of the SM. SUSY being the most popular and attractive extension of the SM, the ATLAS and the CMS collaborations at the Large Hadron Collider (LHC) experiments have been trying to probe various SUSY scenarios in different channels in order to pin down the physics beyond the SM (BSM). However, so far the LHC has not been able to come up with any hint of SUSY up to center-of-mass energy of 13 TeV yet.

Sparticles like squarks and gluinos belonging to the strong sector of SUSY have been searched with great enthusiasm because of their large production cross sections. Consequently stringent bounds on the masses of these sparticles have been obtained by both ATLAS and CMS collaborations from RUN I [25–31] and RUN II [32–46] data. However, these bounds are obtained on the basis of certain ad hoc assumptions which are often hard to realize even in the generic pMSSM which is a popular and broader framework for studying SUSY. Hence relaxation of the above bounds is possible, for example, in compressed SUSY scenarios [47–53].

Recently the ATLAS collaboration has published their analyses for squark/gluino searches [54] based on integrated luminosity of 139 fb^{-1} at the LHC RUN II which, as expected puts stronger mass bounds on squarks and gluinos. In this paper we begin with the channel $1l + jets + \cancel{E}_T$. The results are interpreted in terms of a simplified model. In the ATLAS analysis the relevant sparticles are assumed to be the gluino (\tilde{g}), the lighter chargino ($\tilde{\chi}_1^\pm$) and the lightest neutralino ($\tilde{\chi}_1^0$), where the masses of \tilde{g} and $\tilde{\chi}_1^0$ are free parameters. Only the first two generations of squarks of L -type (the bosonic counterpart of left handed quarks) are assumed to mediate the gluino decays, while

the R -type squarks are decoupled. In addition $\tilde{\chi}_1^\pm$ mass is assumed to be the arithmetic mean of the two free parameters. This optimizes the phase space of gluino decays. It is further assumed that the lightest supersymmetric particle (LSP) is bino like and the lighter charginos are wino like. All other sparticle masses are set beyond the reach of the LHC. It is also assumed that the gluinos decay into the wino like lighter charginos in association with two light quarks with 100% branching ratio (BR). The lighter charginos subsequently decay into $W^\pm \tilde{\chi}_1^0$ with 100% BR. The final state lepton arises from the decay of the W^\pm according to the SM.

It is obvious that many of the above assumptions are designed to obtain very strong bounds on the gluino mass and even more stronger statements are made on the basis of these bounds. Perhaps the practice of thinking critically on results involving large model dependent uncertainties is no longer in vogue. Even if we leave this issue aside, one can still make several remarks on this approach. First these model dependent limits should be cross checked in a more generic framework like the phenomenological Minimal Supersymmetric Standard Model (pMSSM) [55]. We will show that in this case there are less contrived models where the above limits weaken significantly. In some cases they may even disappear. Secondly, within the narrow framework of a simplified model implications of LHC data for various DM search results can not be reliably studied.

The purpose of this paper is to explore the above issues within the broader framework of the pMSSM. First we examine whether the above limits on the gluino mass can be significantly relaxed in some pMSSM scenarios. To begin with the pMSSM parameters are so chosen that the models resemble some features of the ATLAS simplified model. However, some differences are inevitable. For example, if the lighter chargino is wino like, the $\tilde{\chi}_2^0$ is also wino like and almost degenerate with the former and contributes significantly to gluino decays which affects the signal. The special choice of only relatively light L -type squarks, as in the ATLAS work, allows the gluino to dominantly decay into wino like sparticles. In addition the presence of relatively light R -type squarks enhances the fraction of direct gluino decays into the bino like LSP resulting in a further suppression of the leptonic signal. In summary in the pMSSM gluinos may decay into all lighter electroweakinos none with 100 % BR as assumed in the ATLAS analysis. Phase space suppression due to modified gluino-chargino-LSP mass hierarchies may further reduce the mass limits which we will also explore.

Finally, the impact of changing the compositions of the lighter charginos and neutralinos (to-

gether called lighter electroweakinos) also affects the mass limits. In fact, higgsino like charginos have not received the due attention in sparticle searches at the LHC. Phenomenological studies [56–60] for probing Electro-Weak (EW) sectors through electroweakinos have also been performed in the light of ATLAS/CMS data [61–64].

Since the bounds on the gluino mass obtained by ATLAS is based on wino like lighter charginos, it would be interesting to see the results corresponding to a higgsino like $\tilde{\chi}_1^\pm$.

In this way we identify different classes of models where the gluino mass limits are significantly reduced. In the most dramatic cases we find that in some generic models the bounds from the $1l + jets + \cancel{E}_T$ channels are completely washed out.

However, the class of models with suppressed $1l + jets + \cancel{E}_T$ signal are expected to yield more $jets + \cancel{E}_T$ events. We, therefore, revisit these models with reduced gluino mass limits using the ATLAS $jets + \cancel{E}_T$ data [65]. In many cases stringent exclusion contours in the $M_{\tilde{g}} - M_{\tilde{\chi}_1^0}$ plane are restored. The lesson of this exercise is that even in the popular and more general pMSSM model with 19 parameters more realistic bounds may be obtained if they are derived from data collected from a small number of channels. A similar observation was made using LHC Run I data [56, 57]. One can, therefore, go beyond rather contrived simplified models while analyzing LHC data without making the analysis involving many channels unmanageably complicated.

Various SUSY models with R-parity conservation [7, 8] provide a stable LSP. This sparticle, usually chosen as the lightest neutralino, is a popular DM candidate. This weakly interacting massive particle (WIMP) may explain the observed DM in the universe [13–21, 66–75]. The first stringent constraint on the pMSSM came from the WMAP measurement [76] of the DM relic density. However, we shall use the more recent and slightly improved relic density data obtained by the PLANCK experiment [77].

In ref. [78] it was discussed that the physics of DM relic density depends on a multitudes of scenarios in the pMSSM. Apparently DM relic density production is expected to be almost insensitive to parameters in the strong sector. However, quite often a mass relation among the LSP-lighter chargino and gluino masses (like the one assumed by ATLAS mentioned above) is assumed to simplify the analysis. As a result the exclusion contours in the LSP-gluino mass plane from LHC data get further restricted by the DM data as will be shown in the upcoming sections. More interestingly gluino-LSP coannihilation [79–83] directly plays an important role in DM relic

density production in a wide class of models studied by us. This role of the gluino especially for high LSP mass has not been discussed in the context of recent LHC data.

We will further constrain the allowed parameter space (APS) of the above models using data from direct searches of DM from the LUX [84] and, subsequently, by the XENON1T [85] experiments. It should, however, be borne in mind that the computation of the LSP-nucleon scattering cross section, which is a crucial ingredient in this exercise, involves sizable uncertainties both theoretical and experimental. For example, it is not known whether the flux of DM in the neighbourhood of the Earth is strong enough to lead to an observable signal. On the other hand the theory of LSP-nucleon scattering in a low energy experiment involves several uncertainties. Nevertheless, this terrestrial experiment for DM search remains popular.

Additional constraint on pMSSM models which we have studied, can be obtained from the DM capture in the SUN as measured by the IceCube experiment [86]. This may also constrain the pMSSM models we have studied.

In sec.(2) we shall briefly discuss the LHC and DM constraints and the strategy for simulating the LHC signals. Descriptions of the models under study and the corresponding constraints from various data will be discussed in sec.(3) and (4). Subsequently in secs.(5) and (6) we will discuss the above models in the light of the direct detection of DM and the IceCube data. We discuss the possibility of discriminating among several models, if SUSY signals show up in future LHC runs, in terms of relative signal strengths in sec.(7). Finally, we will make concluding remarks in sec.(8).

2 Methodology

2.1 Constraints

2.1.1 Higgs mass at 125 GeV

In the framework of pMSSM one can easily accommodate the lighter Higgs mass, M_h to be around 125 GeV [87, 88]. Loop corrections for Higgs masses due to top-squarks demand large values of soft trilinear coupling A_t (around 5 TeV) [89] which in turn reduces the average mass scale of SUSY. In our analysis M_h is required to be within the mass window between 122 GeV to 128 GeV. This window is considered to take care of theoretical uncertainty in computing the lighter Higgs mass. This theoretical uncertainty in computing the lighter Higgs mass comes into picture

due to renormalization scheme, scale dependence, radiative corrections to Higgs mass up to three loops [90–95].

2.1.2 Constraints from ATLAS $1l + jets + \cancel{E}_T$ and $jets + \cancel{E}_T$ analyses

In ref. [54], pair productions of gluinos and squarks at 13 TeV LHC with an integrated luminosity of 139 fb^{-1} were considered with subsequent decay of gluinos into $\tilde{\chi}_1^\pm$ and appropriate quarks with 100% BR. The lighter charginos further decay into W^\pm and the LSP with 100% BR leading to the signal comprising of $1l + jets + \cancel{E}_T$. The analysis was based on simplified model with certain assumptions mentioned in ref. [54]. In this analysis ATLAS chose the mass hierarchy among \tilde{g} , $\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^0$ defined by $x = \frac{M_{\tilde{\chi}_1^\pm} - M_{\tilde{\chi}_1^0}}{M_{\tilde{g}} - M_{\tilde{\chi}_1^0}}$, where $M_{\tilde{g}}$, $M_{\tilde{\chi}_1^\pm}$ and $M_{\tilde{\chi}_1^0}$ are the masses of gluino, lighter charginos and LSP respectively. Leptons considered here are only electrons or muons. The sets of cuts implemented were given in Table 2 in [54] with respect to 4 different exclusive signal regions (SR) defined as SR2J, SR4J low-x, SR4J high-x, SR6J. Although SR2J and SR6J are pertinent for our analysis in which x is set to be 1/2, i.e., the mass of $\tilde{\chi}_1^\pm$ is placed midway between $M_{\tilde{g}}$ and $M_{\tilde{\chi}_1^0}$. The upper limit on visible cross sections for different SRs for this study are adopted from Table 11 of ref. [54]. From the exclusion contour in the $M_{\tilde{g}} - M_{\tilde{\chi}_1^0}$ plane shown in the top-left pane of Fig. 8 in ref. [54], it is observed that for negligible mass of the LSP $M_{\tilde{g}} \lesssim 2.2 \text{ TeV}$ is excluded. Moreover, for $m_{\tilde{\chi}_1^0} \gtrsim 1.26 \text{ TeV}$ there is no lower bound on mass of gluino.

Next we briefly discuss the gluino search by the ATLAS collaboration through gluino pair production leading to the final state comprising of $jets + \cancel{E}_T$ at 13 TeV LHC with an integrated luminosity of 139 fb^{-1} . In simplified models only the LSP, the lighter charginos and the gluino are within the kinematic reach of the LHC whereas the other sparticles and heavy Higgses are set beyond the LHC reach. The mass of $\tilde{\chi}_1^\pm$ is considered as the arithmetic mean of the LSP and the gluino masses. In this analysis ATLAS assumed gluinos to be decaying into i) $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ or in ii) $\tilde{g} \rightarrow q\bar{q}'\tilde{\chi}_1^\pm$ both with 100% BR. These cascade decays finally lead to $jets + \cancel{E}_T$ signature. We use the limit which is given for the direct decays of the gluino ($\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$). Our analysis is based on this channel.

ATLAS collaboration consider ten inclusive SRs for studying this simplified model. The analysis cuts are given in Table 8 and Table 9 and the upper bound on effective cross sections of BSM physics

for different SRs are given in Table 12 of ref. [65]. In Fig.14 of ref. [65] it has been observed from the exclusion contour that, for negligible LSP mass $M_{\tilde{g}} \lesssim 2.3$ TeV is excluded. Moreover, for $M_{\tilde{\chi}_1^0} \gtrsim 1.2$ TeV there is practically no bound on $M_{\tilde{g}}$.

2.1.3 Constraints from Dark Matter

Although the PLANCK data [77] for satisfying the observed DM relic density has a tiny observational uncertainty (0.120 ± 0.001), we note that there is about 10% theoretical uncertainty in computing the SUSY DM relic density [96, 97]. This 10% theoretical uncertainty arising from the renormalization scheme and the scale variations due to higher order SUSY-QCD corrections amounts to approximately six times the observational uncertainty as mentioned in refs.[149,150]. Several (see for example, refs. [98,99]) analyses included this theoretical error or even more in their works. This prompts us to consider $\Omega_\chi h^2 = 0.120 \pm 0.006$ that leads to the following bounds

$$0.114 < \Omega_\chi h^2 < 0.126$$

where $h = 0.733 \pm 0.018$ [100] is the expansion rate of the Universe, the Hubble constant, in the units of 100 km/Mpc-s. Apart from the above direct DM constraints we further impose the following indirect searches for the DM candidates.

DM direct detection constraints like bound on spin-independent (SI) LSP-proton scattering cross section $\sigma_{\tilde{\chi}p}^{SI}$ is imposed using the the LUX [84] and the XENON1T [85] data.

This analysis is also subjected to the indirect detection bound on spin-dependent (SD) LSP-proton scattering cross section $\sigma_{\tilde{\chi}p}^{SD}$ obtained from the IceCube experiment [86].

2.2 Simulation using LHC RUN-II data

We generate the sparticle mass spectra and decays in SUSY-HIT [101]. Gluino pairs are then produced using MG5_aMC@NLO [102] at 13 TeV centre of mass energy with the PDF set NNPDF2.3LO [103]. Next the showering and hadronization are processed via Pythia 8.2 [104]. Matched events are generated using the CKKW-L scheme [105]. Delphes 3.4 [106] is employed for the reconstruction of dressed objects like jets, leptons, missing p_T etc. and implementation of analysis cuts. The jets and leptons are pre-selected with certain loose quality requirements and are termed as ‘baseline’ by the ATLAS collaboration. Signal objects, on the other hand, are further selected with tighter

identification criteria applied on the baseline objects. Furthermore, the next to next to leading order (NNLO) cross sections of the gluino pair production are taken from the ATLAS-Twiki [107], for simulation purpose.

In order to validate our simulation we compute the exclusion contour given by the ATLAS collaboration [54, 65] using `Delphes 3.4` [106] after generating the events in `MG5 aMC@NLO` [102]. We have followed the electron/muon-jets isolation criteria for reconstructing and identifying those objects according to the ATLAS prescription. We have implemented the detector simulation using ATLAS card given in the `Delphes 3.4` code. The signal regions segmented by the ATLAS are characterized by a number of kinematical cuts and N_{BSM} (= production cross section \times luminosity \times cut efficiency \times acceptance) which is the number of signal events coming from the BSM scenario only. Any pMSSM parameter space point is excluded in our simulation if the yield exceeds at least one of the values of N_{BSM} . For the purpose of obtaining exclusion contour for our models we have used the N_{BSM} number as the limiting value. We vary $M_{\tilde{\chi}_1^0}$ keeping $M_{\tilde{g}}$ fixed in order to obtain the number of signal events in a particular model very close to N_{BSM} value. The value of $M_{\tilde{\chi}_1^0}$ for which the yield becomes closest to N_{BSM} marks a limiting point for a particular $M_{\tilde{g}}$ in a particular model. Next we vary the gluino mass to construct the entire exclusion contour in the $M_{\tilde{g}} - M_{\tilde{\chi}_1^0}$ plane. The validation results are in agreement with the ATLAS contour within 10% alteration. Throughout this paper we consider the validated exclusion contour as the ATLAS exclusion contour for $1l + jets + \cancel{E}_T$ and $jets + \cancel{E}_T$ channels.

3 Models

This section has two subsections corresponding to different compositions of $\tilde{\chi}_1^\pm/\tilde{\chi}_2^0$. These sparticles are assumed to be dominantly wino (higgsino) in subsec.3.1(3.2). In both the subsections the LSP is assumed to be bino like. The heavier electroweakinos are assumed to be decoupled in both the discussions.

The masses of the LSP and gluino are free parameters and are determined by U(1) and SU(3) gaugino mass parameters M_1 and M_3 , respectively. The SU(2) gaugino mass parameter M_2 is chosen in such a way to produce $M_{\tilde{\chi}_1^\pm}$ exactly halfway between $M_{\tilde{g}}$ and $M_{\tilde{\chi}_1^0}$. In the pMSSM scenario, M_2 sets the masses of both the lighter chargino as well as the second lightest neutralino

which are nearly mass degenerate. All slepton mass parameters are set at 3.0 TeV, and hence they are inaccessible at the LHC RUN II.

Third generation squarks are assumed to be decoupled. All other soft breaking trilinear terms are considered to be zero except the trilinear term for top quark. $A_t = 5$ TeV has been chosen in order to get SM-like Higgs mass at around $M_h \simeq 125$ GeV [90, 91]. Throughout this study $\tan\beta$ is set at 10 and the values of the pseudo scalar Higgs mass M_A is 3 TeV and M_{H^\pm} is obtained around 3.002 TeV for wino model (discussed in subsec.3.1) and for higgsino model (discussed in subsec.3.2).

3.1 The Wino Model

In wino model $\tilde{\chi}_1^\pm/\tilde{\chi}_2^0$ is considered to be wino dominated. In this model we further assume $\tilde{\chi}_1^0$ to be bino dominated in most of the parameter space. This can be achieved by considering a large value of higgsino mass parameter, μ ($\mu \simeq 2M_2$), which satisfies the hierarchy $M_1 < M_2 \ll \mu$, where M_1 and M_2 are the U(1) and SU(2) electroweak gaugino mass parameters respectively.

In this analysis, M_2 is not a free parameter rather it is constrained by the relation $x = \frac{M_{\tilde{\chi}_1^\pm} - M_{\tilde{\chi}_1^0}}{M_{\tilde{g}} - M_{\tilde{\chi}_1^0}}$ (see subsec.(2.1.2)). On top of it, we have a number of variant scenarios assuming different hierarchy of masses of the L -type and R -type squarks, which we describe in the following subsections.

3.1.1 LLRLW: Left Light Right Light Wino model

In the Left Light Right Light Wino (LLRLW) model, it is assumed that the diagonal entries of the first two generations of squark mass matrices take the value around 2.5 TeV, which are heavier than the gluino masses we study. As a result gluino has only three body decay modes and in particular, it has moderately large BR in the decay mode $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$. This is because of the fact that the intermediate off-shell squarks have equally large doublet and singlet contributions, where the latter favorably couples to the bino like LSP. Therefore, the BR ($\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm$) is depleted compared to the simplified version considered by the ATLAS which assumed this BR to be 100%. As a result, the yield of $1l + jets + \cancel{E}_T$ signal events which diminishes appreciably the LSP mass compared to that of the ATLAS simplified scenario corresponding to the same gluino mass.

In Fig.(1) the parameter space excluded by the ATLAS collaboration is displayed by the solid black line and the corresponding degraded exclusion contour in respect of LLRLW model is shown

by the green dashed line. From the ATLAS analysis [54] it has been observed that for negligible LSP mass $M_{\tilde{g}} < 2.2$ TeV is excluded. Furthermore, there is no bound on $M_{\tilde{g}}$ where $M_{\tilde{\chi}_1^0}$ exceeds 1.26 TeV. It is evident from the figure that the limit on gluino mass reduces appreciably by an amount around (200 – 250) GeV with respect to the corresponding ATLAS bound. For negligible LSP mass the excluded region is now relaxed to $M_{\tilde{g}} \approx 2$ TeV, whereas, for $M_{\tilde{\chi}_1^0} \approx 970$ GeV the bound on $M_{\tilde{g}}$ ceases to exist.

Throughout the work, we have considered the compression factor $x = 0.5$ (see subsec.(2.1.2) for details). In this model the limits on the $M_{\tilde{g}}$ comes down from 2 TeV to 1.7 TeV as x increased from 0.5 to 0.7 for negligible $M_{\tilde{\chi}_1^0}$. It may be noted that as we increase x from 0.5 to 0.7, the BR $\tilde{g} \rightarrow \tilde{\chi}_1^0 q \bar{q}$ leaps from 35% to 80%.




Figure 1: Exclusion contour for the Left Light Right Light Wino (LLRLW) model is shown in the $M_{\tilde{g}} - M_{\tilde{\chi}_1^0}$ plane. The solid black line corresponds to the observed limit as given by ATLAS from the $1l + jets + \cancel{E}_T$ data at 13 TeV center-of-mass energy in the simplified model [54]. The green dashed curve is the modified excluded region for the LLRLW model where relatively low L -type and R -type squark masses (but still heavier than the gluino mass) are considered along with wino type lighter charginos and second lightest neutralino. The cyan blue dots signify masses of the LSP which are allowed by the PLANCK relic density data [77]. The region above the red diagonal line is disallowed as gluino becomes the LSP.

The two distinct branches in agreement with the PLANCK data [77] are presented in Fig.(1).

The corresponding allowed regions are shown in cyan blue points. In the lower branch the cyan blue points imply DM producing mechanisms corresponding to LSP-pair annihilation through a s -channel light h -resonance of mass ≈ 125 GeV. Moreover, there are points in that branch which correspond to Z -resonances also. This branch extends up to $M_{\tilde{g}} \lesssim 800$ GeV. For higher gluino masses the tiny higgsino component of the dominantly bino like LSP is too small to continue this mechanism.

The upper branch is approximately parallel to the red line. The lower part of this branch represents the mass region where the LSP pair annihilation is the dominant DM relic density production mechanism. In the upper part of this branch the LSP coannihilation with a nearly mass degenerate sparticle like the \tilde{g} , $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ are the main contribution to the relic density production. For very high gluino masses beyond the LHC reach, LSP coannihilation with gluino plays a very important role for DM relic density production.

3.1.2 LLRHW: Left Light Right Heavy Wino model

This model is closer to the ATLAS simplified model compared to the LLRLW model. In the Left Light Right Heavy Wino (LLRHW) model, the SU(2) doublet L -type squark mass parameters of first two generations continue to be ~ 2.5 TeV as in the previous case. However, the masses of the R -type squarks are considered to be at large value around 6.5 TeV. This means that the R -type squarks play insignificant role in gluino decay processes for this scenario. Since the intermediate off-shell squarks are mostly SU(2) doublet, the BR of the decay mode, $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm$ is higher than that of the LLRLW model. However, in this scenario gluino also decays to $\tilde{\chi}_2^0 q\bar{q}$ with a moderate to large BR. $\tilde{\chi}_2^0$ being wino like, further, predominantly decays to $h\tilde{\chi}_1^0$ along with $Z\tilde{\chi}_1^0$ decay mode with negligible BR. The decay mode $h\tilde{\chi}_1^0$ or $Z\tilde{\chi}_1^0$ does not contribute to the $1l + jets + \cancel{E}_T$ final state. Consequently, the number of $1l + jets + \cancel{E}_T$ signal events decreases appreciably compared to the simplified model considered by the ATLAS collaboration. In this model, the exclusion contour shrinks by a modest amount $\sim (100 - 150)$ GeV in comparison to the exclusion contour of the ATLAS experiment. For negligible LSP mass the exclusion region degrades to $M_{\tilde{g}} \approx 2.05$ TeV, whereas, for $M_{\tilde{\chi}_1^0} \approx 1100$ GeV the bound on $M_{\tilde{g}}$ goes away. As expected the exclusion limit in the LLRHW model gets depleted by a smaller amount compared to the previous LLRLW model.

For the LLRHW model the shrinkage of the exclusion is not so significant due to compression

as compared to the LLRLW model. Here, the limit on the $M_{\tilde{g}}$ relaxes to 1.93 TeV from 2.05 TeV as x changes to 0.7 from 0.5 for the negligible $M_{\tilde{\chi}_1^0}$.

In Fig.(2) we present the PLANCK data [77] satisfying points over the LHC exclusion plot similar to that discussed in subsec.(3.1.1) for Fig.(1).




Figure 2: Exclusion contour (green dashed line) is obtained for the Left Light Right Heavy Wino (LLRHW) model where L -type squark masses are small, and R -type squark masses are quite heavy ($M_{\tilde{g}} < M_{\tilde{q}_L} \ll M_{\tilde{q}_R}$). The lighter charginos and second lightest neutralino are wino type. The ATLAS exclusion limits on gluino masses in their simplified model is displayed by solid black line. The cyan points satisfy relic density data obtained from the PLANCK data [77]. The color conventions are the same as given in Fig.(1).

In another variant of wino model namely, Left Heavy Right Light Wino (LHRLW) model, this $1l + jets + \cancel{E}_T$ signal disappears as the gluino predominantly decays into $jets + \cancel{E}_T$ channel.

3.2 Higgsino model

We now discuss the scenario where the lighter chargino ($\tilde{\chi}_1^\pm$), the second and third lightest neutralinos ($\tilde{\chi}_2^0, \tilde{\chi}_3^0$) are higgsino dominated and $\tilde{\chi}_1^0$ being still bino dominated. This model can be realized by considering large M_2 . The preferred hierarchy of the gaugino and higgsino mass parameters are to be set as $M_1 < \mu \ll M_2$. In the following subsections we will study this model for various scenarios depending on where L -type or R -type squark masses are placed with respect to the gluino

mass.

3.2.1 LLRLH: Left Light Right Light Higgsino model

In the Left Light Right Light Higgsino (LLRLH) model, it is assumed that both the L -type and R -type squark mass parameters of the first two generations are fixed at $\simeq 2.5$ TeV. For this part of the parameter space, the gluino pair production dominantly gives $jets + \cancel{E}_T$ signal. The BR of gluino decaying into $\tilde{\chi}_1^\pm/\tilde{\chi}_2^0$ in association with quarks is almost negligible, resulting in yield of a meager amount of $1l + jets + \cancel{E}_T$ final states. As a result, the bound on $M_{\tilde{g}}$ from $1l + jets + \cancel{E}_T$ [54] search by ATLAS almost vanishes for higgsino model.




Figure 3: In this figure we present a scatter plot of the points satisfying the DM relic density data from the PLANCK [77] experiment in one of the higgsino models (LLRLH model). Since all the higgsino models are allowed by the $1l + jets + \cancel{E}_T$ analysis from the ATLAS collaboration [54], we have shown for reference, the ATLAS exclusion contour for simplified model by the solid black line. Colours and conventions are the same as Fig.(1)

We also present the PLANCK data [77] allowed points for the LLRLH model in Fig.(3). In the figure we observe three separate branches. The lowest one produces the right relic abundance through the LSP-pair annihilation via s -channel h/Z resonances. The upper branches primarily consists of three relic density producing regions namely the LSP-pair annihilation for relatively low $M_{\tilde{g}}$ (as discussed before) and the A and H^\pm resonance regions and a very important coannihilation

region. This region has the same characteristics discussed for LLRLW model. The nearly horizontal branches corresponding to $M_{\tilde{\chi}_1^0} \simeq M_{A/2} \simeq M_{H^\pm/2}$ represents LSP pair annihilation via A and H^\pm resonances.

3.2.2 LLRHH: Left Light Right Heavy Higgsino model

In the Left Light Right Heavy Higgsino (LLRHH) model, the mass of the L -type squarks of first two generations are assumed to have values around 2.5 TeV, whereas, the R -type squarks are set at 6.5 TeV. Therefore, the R -type squarks have less chance to appear in the off-shell propagator in gluino decay. The dominant decay modes of the gluino for this case is $q\bar{q}\tilde{\chi}_1^0$, resulting in very depleted number of $1l + jets + \cancel{E}_T$ [54] final states. For such scenario, therefore, the ATLAS bounds practically the same as in the previous case.

We note in passing that for the case where the R -type squarks are made to be lighter and L -type squarks are heavier, the model is equivalent to the ATLAS simplified model for search in the $jets + \cancel{E}_T$ [65] final state which we will discuss in the next section.

4 Higgsino model in the light of $jets + \cancel{E}_T$ data

As noted in sec.(3), in all the variants of higgsino type models, that practically all the parameter space are allowed by $1l + jets + \cancel{E}_T$ data [54]. This happens because, the gluino dominantly decays into the LSP and jets in such models making the one lepton signal vanishingly small. Therefore, the viability of such models need to be analyzed again by $jets + \cancel{E}_T$ data [65].

For the purpose of illustration we consider the LLRLH model which is unbounded from the ATLAS $1l + jets + \cancel{E}_T$ [54] constraint. In Fig.(4) we present the exclusion contour for ATLAS $jets + \cancel{E}_T$ data in respect of simplified model by black solid line. In the same figure the green dashed line corresponds to exclusion contour for LLRLH model. It is also seen from the figure that the gluino mass bounds do not alter appreciably for that obtained from the $1l + jets + \cancel{E}_T$ analysis for negligible LSP mass. The disallowed region of parameter space (near around $M_{\tilde{g}} = M_{\tilde{\chi}_1^0}$ line) extends a little above the ATLAS $jets + \cancel{E}_T$ exclusion contour thereby making the $M_{\tilde{g}}$ -bound stronger. Hence it is observed that the overall excluded region of parameter space does not change significantly in comparison with the exclusion contour drawn from $jets + \cancel{E}_T$ data [65]. It should




Figure 4: Exclusion contour (green dashed line) is obtained for the Left Light Right Light Higgsino (LLRLH) model where both L -type and R -type squark masses are small but larger than the gluino mass ($M_{\tilde{g}} < M_{\tilde{q}_L} = M_{\tilde{q}_R}$). The lighter charginos and second lightest neutralino are higgsino type. The ATLAS exclusion limit on gluino mass for the search of $jets + \cancel{E}_T$ [65] at 13 TeV for simplified model is displayed in solid black line. The color conventions are the same as given in Fig.(1).

be kept in mind that the whole region of parameter space is allowed by $1l + jets + \cancel{E}_T$ data.

We observe from Fig.(1) (Fig.(2)) that for negligible LSP mass the limit on $M_{\tilde{g}}$ is obtained around 2 TeV for LLRLW and LLRHW models, constrained by $1l + jets + \cancel{E}_T$ data. Compared to these numbers, in LLRLH model the bound on $M_{\tilde{g}}$ becomes 2.3 TeV obtained from $jets + \cancel{E}_T$ data. Thus the bounds on $M_{\tilde{g}}$ in LLRLH model (see Fig.(4)) is quite comparable to that for wino type LLRLW/LLRHW model (see Fig.(1)/(2)).

5 Direct Detection through Spin-independent scattering

In addition to the constraint from the PLANCK results, we compute the spin-independent (SI) LSP-proton scattering cross section ($\sigma_{\chi p}^{SI}$) in relation to the earlier LUX data [84] and the more recent XENON1T data [85]. The cross section is sensitive to the mass of the LSP and the compositions of the LSP [13,108]. We must keep in mind various sizable uncertainties which plague the computation of $\sigma_{\chi p}^{SI}$. For example, they may arise due to low energy hadronic physics [109–112], the local DM density which has not been measured directly [113–115], non-Maxwellian velocity distribution of

the WIMPs [116,117] etc. Combining all these the rate of direct detection may involve an order of magnitude of uncertainty or more.

Within the detector nucleus, SI interaction occurs between the lightest neutralino (LSP) and the quarks inside the nucleon through s -channel squark exchange and t -channel Higgs exchange processes. The dominant contribution to $\sigma_{\chi p}^{SI}$ comes from the t -channel Higgs exchange as squarks are assumed to be considerably heavy throughout this analysis. The effective couplings are dependent on the nature of composition of the LSP. The products of the gaugino and the higgsino components of the neutralino diagonalizing matrix have a major contribution in the $h(H)\tilde{\chi}_1^0\tilde{\chi}_1^0$ coupling where H is the CP even heavy neutral higgs boson. Assuming the LSP to be bino dominated ($M_1 \ll M_2, \mu$), the effective couplings can be approximated as [118],

$$C_{h\tilde{\chi}\tilde{\chi}} \simeq \frac{M_Z s_W t_W}{M_1^2 - \mu^2} [M_1 + \mu \sin 2\beta] \qquad C_{H\tilde{\chi}\tilde{\chi}} \simeq -\frac{M_Z s_W t_W}{M_1^2 - \mu^2} \mu \cos 2\beta$$

where, $s_W = \sin \theta_W$ with θ_W as the Weinberg angle and M_Z is the mass of the Z boson. From the couplings we can easily interpret that a significant amount of bino-higgsino mixing i.e. $M_1 \simeq \mu$ results in a large value of the SI cross section $\sigma_{\chi p}^{SI}$, whereas, a bino dominated LSP has small coupling and hence the XENON1T [85] experiment can probe such models. In the following subsections the computed results are presented for different wino and higgsino models.

5.1 LLRLW and LLRHW

We now consider the points in Figs.(1) and (2) which are consistent with both LHC and PLANCK data and test their viability vis-a-vis the XENON1T [85] data.

In Fig.(5a) we present the plot in the $M_{\tilde{\chi}_1^0} - \sigma_{\chi p}^{SI}$ plane for LLRLW model. The value of $\sigma_{\chi p}^{SI}$ is computed using `micrOMEGAs 5.2` [119]. The exclusion limits for the LUX and the XENON1T results are shown by the brown and orange curves respectively. Dark blue region of parameter space is allowed by the ATLAS data [54] as well as the PLANCK [77] data. The cyan points denote the region of parameter space allowed by the PLANCK data only. Whereas the dark blue region of parameter space consistent with both LHC and PLANCK data satisfies the latest direct detection constraints [84, 85]. The similar observations are found in LLRHW model which is displayed in




Figure 5: Spin-Independent direct detection cross section for the a) Left Light Right Light Wino (LLRLW) model and b) Left Light Right Heavy Wino (LLRHW) model. Upper limits on the spin-independent elastic WIMP-nucleon cross section at 90% C.L provided by the LUX [84] and the XENON1T [85] are shown as brown and orange lines respectively. The cyan points only satisfy the relic density data from the PLANCK [77] experiment. The points which are allowed by both the LHC and PLANCK data are shown by the dark blue points.

Fig.(5b).

We also observe in Fig.(5a) that the points which satisfy the condition $M_{\tilde{\chi}_1^0} \gtrsim 1.1$ TeV are allowed by the ATLAS data [54,65] (see Figs.(1)–(4)), the PLANCK constraint [77] and as well as the DM direct detection results [84,85]. From Fig.(5b) for LLRHW model we further infer that the parameter space points with $M_{\tilde{\chi}_1^0} \gtrsim 1.1$ TeV are also allowed by all the constraints discussed above.

5.2 Higgsino models

For the sake of completeness, we plot the $M_{\tilde{\chi}_1^0} - \sigma_{\chi p}^{SI}$ for the LLRLH model in Fig.(6). It appears that the points consistent with the LHC and PLANCK data are in conflict with XENON1T data. However in view of the theoretical and experimental uncertainties discussed above this conflict is inconclusive.




Figure 6: Spin-Independent direct detection results for the Left Light Right Light Higgsino (LL-RLH) model. Colours and conventions are the same as in Fig.(5)

6 Spin-Dependent cross section from IceCube experiment:

The data obtained from the IceCube experiment [86] is based on the indirect detection technology of the DM particles. Here a pair of DM particles annihilate to produce various SM particles in pairs. The production of the SM particles in the final state depends on the mass of the corresponding DM particles and can lead to interesting signals for indirect search of the DM.

The spin-dependent (SD) $\tilde{\chi}_1^0 - p$ scattering process occurs via s -channel squark mediated and t -channel Z boson mediated feynman diagrams [17, 108, 120]. The larger value of the above scattering cross section, $\sigma_{\chi p}^{SD}$ results from a large $Z\chi\chi$ coupling given by $C_{Z\chi\chi} \propto |N_{13}^2 - N_{14}^2|$ where N_{13} and N_{14} stands for the higgsino components present in the LSP. For the higgsino type models, $\sigma_{\chi p}^{SI}$ is relatively large because of the small difference between M_1 and μ as discussed in sec.(5). Enhancement of both the SI and SD $\tilde{\chi} - p$ cross sections results in loss of energy of the corresponding DM particle so that its velocity falls below the escape velocity. This leads to the gravitational capture of the DM within the Sun (e.g., see [13, 121–124]).

The gravitationally captured LSPs undergo pair annihilation resulting in final states comprising of highly energetic SM particles. Out of these particles W^\pm can decay into neutrinos which can

reach the Earth's atmosphere. The muon flux produced by these neutrinos can be measured by the IceCube experiment [86]. From this data $\sigma_{\chi p}^{SD}$ can be computed [125, 126].

6.1 LLRLW and LLRHW

In Figs.(7a) and (7b) we present $\sigma_{\chi p}^{SD}$ for the LLRLW and LLRHW models against the LSP mass with respect to the IceCube data [86]. The spin-dependent cross section, $\sigma_{\chi p}^{SD}$, is computed by using micrOMEGAs 5.2 [119]. The cyan points in the plot satisfy the relic density observed by the PLANCK experiment [77]. The present upper limit obtained from the IceCube experiment [86] is shown by the orange line. The dark blue points in both the models (see Figs.(7a) and (7b)) are allowed by all data.




Figure 7: (a) Plot of spin-dependent direct detection cross section $\sigma_{\chi p}^{SD}$ vs $M_{\tilde{\chi}_1^0}$ for LLRLW model. (b) Plot of spin-dependent direct detection cross section $\sigma_{\chi p}^{SD}$ vs $M_{\tilde{\chi}_1^0}$ for LLRHW model. The orange line corresponds to the hard annihilation channels ($DM DM \rightarrow W^+W^-$) obtained from the IceCube experiment [86]. The cyan points satisfy the PLANCK data [77] and similarly dark blue points satisfy to both the collider as well as the PLANCK data.

In Fig.(7a), as $M_{\tilde{\chi}_1^0}$ increases $\sigma_{\chi p}^{SD}$ decreases and eventually it dips at around $M_{\tilde{\chi}_1^0} \simeq 1.35$ TeV and thereafter $\sigma_{\chi p}^{SD}$ further starts enhancing as $M_{\tilde{\chi}_1^0}$ increases. This can be explained in the following way. The spin-dependent elastic scattering of $\tilde{\chi}_1^0$ with proton takes place through s -channel squark mediated or t -channel Z boson mediated process. For low LSP mass Z boson mediated process contributes significantly in $\sigma_{\chi p}^{SD}$ but as $M_{\tilde{\chi}_1^0}$ increases the squark mediated process begins

to dominate over the former one. The enhancement of squark mediated process takes place in the region of parameter space where $M_{\tilde{\chi}_1^0}$ is comparable to the mass of the squarks. However, these two processes can interfere destructively resulting in a remarkably small value of $\sigma_{\chi p}^{SD}$ near $M_{\tilde{\chi}_1^0} \simeq 1.35$ TeV.

6.2 Higgsino models




Figure 8: Plot of spin-dependent direct detection cross section $\sigma_{\chi p}^{SD}$ vs $M_{\tilde{\chi}_1^0}$ for LLRLH model. Colours and conventions are the same as described in Fig.(7).

For the purpose of illustration we display the $\sigma_{\chi p}^{SD}$ against $M_{\tilde{\chi}_1^0}$ in Fig.(8) for LLRLH model. All the points in the plot are found to be satisfied by the IceCube data [86]. Here it is noted that the dark blue points satisfy the LHC data [54,65] and PLANCK data. Therefore, these models can be easily probed by the present spin-dependent DM-proton direct detection experiment. In this figure we still see the decrease in $\sigma_{\chi p}^{SD}$ around $M_{\tilde{\chi}_1^0} \simeq 1.5$ TeV due to the reason discussed earlier in subsec.(6.1)

7 Relative signal strengths of various models

From our analyses in two different gluino search channels, we sketch the feasibility of distinguishing different pMSSM scenarios discussed in sec.(3). If gluino signals are observed in at least two

channels, this method might be helpful to identify the underlying model. In order to keep the production cross section fixed, we choose $M_{\tilde{g}} = 1.3$ TeV (allowed by all data), while the other mass parameters are chosen from various models. To carry out our analysis we choose three benchmark points (BPs) from Fig.(1)–(3) (see Table(1)). BP1, BP2 and BP3 are chosen from LLRLW, LLRHW and LLRLH models respectively.

The decay modes and their branching fractions relevant for the gluino signals for $M_{\tilde{g}} = 1.3$ TeV are presented in Table(2). It may be noted that in this table the BRs of the gluino may not add up to 100%. This is due to the fact that in some scenarios the gluino also decays into the other channels with small but non-negligible BRs. However, all modes are taken into account while simulating the gluino signal.

Benchmark point	M_1	M_2	M_3	μ	$M_{\tilde{g}}$	$M_{\tilde{\chi}_1^0}$	$M_{\tilde{\chi}_1^\pm}$	$\Omega_{\tilde{\chi}} h^2$	σ_{SI} (pb)	σ_{SD} (pb)
BP1 (form Fig.1)	1269	1236	982	2472	1300	1238	1269	0.116	3.052×10^{-11}	3.702×10^{-10}
BP2 (from Fig. 2)	1262	1232	946	2464	1300	1231	1265	0.124	2.844×10^{-11}	7.571×10^{-9}
BP3 (form Fig.3)	1310	2492	983	1246	1300	1238	1268	0.122	1.737×10^{-8}	6.022×10^{-6}

Table 1: The sparticle mass spectra corresponding to different benchmark points chosen from Fig.1 to Fig.3. The mass parameters are in GeV.

Decay Modes	BP1	BP2	BP3
$\tilde{g} \rightarrow \tilde{\chi}_1^0 q \bar{q}$	80.86	28.08	64.34
$\rightarrow \tilde{\chi}_2^0 q \bar{q}$	6.16	23.20	–
$\rightarrow \tilde{\chi}_1^\pm q \bar{q}'$	12.68	47.60	–
$\rightarrow \tilde{\chi}_1^0 g$	–	–	29.68
$\rightarrow \tilde{\chi}_2^0 g$	–	–	5.78
$\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 W^{*\pm}$	100	100	100
$\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 q \bar{q}$	82.14	84.13	67.56
$\rightarrow \tilde{\chi}_1^0 l^+ l^-$	3.52	3.67	32.07
$\rightarrow \tilde{\chi}_1^0 \gamma$	12.67	10.49	0.33

Table 2: The BR (%) of the dominant decay modes of the sparticles for the different BPs. Here q and q' denote first two generation of quarks . ‘–’ denotes a negligible branching fraction and $W^{*\pm}$ denotes the off-shell W^\pm .

Points	S_1	S_0	R
BP1	2.03	15.82	0.77
BP2	7.11	14.80	0.35
BP3	0.44	25.11	0.97

Table 3: The table displays S_0 and S_1 and R for different BPs. S_0 and S_1 represent the number of signal events corresponding to $jets + \cancel{E}_T$ data and $1l + jets + \cancel{E}_T$ data respectively for an integrated luminosity 139 fb^{-1} . Here R stands for $\frac{S_0 - S_1}{S_0 + S_1}$.

We now use the ATLAS gluino searches for $\sqrt{s} = 13 \text{ TeV}$ with integrated luminosity $\mathcal{L} = 139 \text{ fb}^{-1}$ in $1l + jets + \cancel{E}_T$ and $jets + \cancel{E}_T$ channels, as described in subsec.(2.1.2) and in sec.(4). We define an observable called relative signal strength, $R = \frac{S_0 - S_1}{S_0 + S_1}$, where S_0 and S_1 denote the number of signal events of $jets + \cancel{E}_T$ and $1l + jets + \cancel{E}_T$ channels respectively. It is worth mentioning that the ratio R is almost free from theoretical uncertainties like the choice of QCD scale and the PDFs

etc.

In Table(3) we display the magnitudes of S_1 , S_0 and R corresponding to different BPs. We find that S_0 is larger than S_1 in all models. However, the ratio R , distinguishes different models. This technique of discriminating among various pMSSM models using multichannel analysis may be a useful tool if significant signals in different channels are found in future LHC experiments. A similar method was used to illustrate this possibility of discrimination using RUN-I data [56, 57, 127].

8 Conclusion

The main purpose of this paper is to reexamine the limits on $M_{\tilde{g}}$ obtained by the ATLAS collaboration using rather contrived simplified models [54, 65]. We analyze the same data in the more generic pMSSM model and study the variation of the limits with parameters which are most relevant for the signal being analyzed.

We first revisit the ATLAS $1l + jets + \cancel{E}_T$ data [54]. Here we assume LSP to be bino like and $\tilde{\chi}_1^\pm$, $\tilde{\chi}_2^0$ to be wino like (see subsec.(3.1)). Other electroweakinos are assumed to be decoupled. In the subsec.(3.1.1) both L -type and R -type squarks are assumed to be light and mass degenerate ($M_{\tilde{q}_L} = M_{\tilde{q}_R} \approx 2.5$ TeV). The resulting exclusion contour is displayed in the $M_{\tilde{g}} - M_{\tilde{\chi}_1^0}$ mass plane in Fig.(1). The modest change in the exclusion contour with respect to the ATLAS contour (due to reasons discussed in the text) can be seen in Fig.(1). However, if the compression factor x (defined in the subsec.(2.1.2)) is increased the same contour shrinks even more. For example if $x = 0.7$, the limit is $M_{\tilde{g}} \gtrsim 1.7$ TeV for negligible LSP mass.

On the other hand if $M_{\tilde{q}_R} \gg M_{\tilde{q}_L} \approx 2.5$ TeV, the pMSSM model is closer to the ATLAS simplified model. This is reflected by the fact that the new exclusion contour shrinks very little compared to the ATLAS result (see Fig.(2)).

The results change dramatically if $\tilde{\chi}_1^\pm$, $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ are higgsino like (see subsec.(3.2)). In this scenario the gluino dominantly decays directly into the LSP and jets. Thus irrespective of the squark mass hierarchy the $1l + jets + \cancel{E}_T$ signature is highly depleted.

We next analyze the higgsino models in the light of ATLAS $jets + \cancel{E}_T$ data. As an illustration we present in Fig.(4) a comparative study of the constraints in the LLRLH model and that in the ATLAS simplified model [65]. There is no significant change observed between the two. From

Figs.(1)–(4) we find that even in the compressed scenario ($M_{\tilde{g}} \simeq M_{\tilde{\chi}_1^0}$), $M_{\tilde{g}} \gtrsim 1.5$ TeV is a fairly conservative limit [128,129]. For smaller LSP mass $M_{\tilde{g}} \gtrsim 2$ TeV is a fairly reliable bound. More importantly it appears that the bounds from a single channel in a simplified model is not reliable. This we have seen in the case of the constraint from the $1l + jets + \cancel{E}_T$ data (see the higgsino models in subsec.(3.2)). However, data analysis in more than one channel may lead to reliable limits even in the more generic pMSSM model with 19 parameters.

As discussed in the introduction one can make a useful analysis of the interplay between the LHC and PLANCK data on dark matter relic density within the framework of the pMSSM. From Figs.(1)–(4) we find that coannihilation of the LSP with other spin half gauginos is an important DM producing mechanism in the parameter space allowed by the LHC data. The coannihilation of the gluinos with the LSP is the dominant among other coannihilation channels. For the higgsino models there is another interesting process for DM relic density production. They are LSP annihilation via H^+ or A resonance (see Fig.(3)).

In Fig.(5a)((5b)) we show the points in the LLRLW (LLRHW) model consistent with both LHC and PLANCK data by the dark blue points. All presently available constraints are satisfied by these points for $M_{\tilde{\chi}_1^0} \gtrsim 1.1$ TeV (see Fig.(4)).

We have displayed the direct detection results in connection with XENON1T data for LLRLH model in Fig.(6). The points satisfying LHC and PLANCK constraints are found to be apparently in conflict with XENON1T data. This disagreement, however, is not conclusive if theoretical and experimental uncertainties in deriving the limits are taken into consideration.

We have also computed the $\sigma_{\chi p}^{SD}$ (see sec.(6)) which is determined by the IceCube experiment [86]. We find that, (see Figs.(7a), (7b) and (8)) in all the models examined by us there are points represented by dark blue dots which are consistent with all data.

Finally, we have carried out a small exercise in sec.(7). This illustrates that if future LHC experiments discover signals in both $jets + \cancel{E}_T$ and $1l + jets + \cancel{E}_T$ channels then the underlying wino and higgsino type models can be distinguished from each other by measuring an observable, R defined in sec.(7).

Acknowledgments

Authors are grateful to Prof. Amitava Datta for his guidance and valuable suggestions on the work and careful reading of the manuscript. Authors acknowledge the help of Dr. Nabanita Ganguly in computation. AM is also thankful to Dr. Jyoti Prasad Saha and Md. Raju for discussions. AM would like to thank DST for providing the INSPIRE Fellowship [IF170693].

References

- [1] H. P. Nilles, “Supersymmetry, Supergravity and Particle Physics”, Phys. Rept. **110** (1984) 1.
- [2] H. E. Haber and G. L. Kane, “The Search for Supersymmetry: Probing Physics Beyond the Standard Model”, Phys. Rept. **117** (1985) 75.
- [3] J. Bagger and J. Wess, *Supersymmetry and Supergravity: Revised Ed*, Princeton University Press Mar 23, 1992.
- [4] J. D. Lykken, “Introduction to supersymmetry”, in *Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 96): Fields, Strings, and Duality 1996*, [arXiv:hep-th/9612114](https://arxiv.org/abs/hep-th/9612114).
- [5] S. P. Martin, “A Supersymmetry primer”, Adv. Ser. Direct. High Energy Phys. **18** (1998) 1, [arXiv:hep-ph/9709356](https://arxiv.org/abs/hep-ph/9709356).
- [6] D. J. H. Chung, L. L. Everett, G. L. Kane, S. F. King, J. D. Lykken and L.-T. Wang, “The Soft supersymmetry breaking Lagrangian: Theory and applications”, Phys. Rept. **407** (2005) 1, [arXiv:hep-ph/0312378](https://arxiv.org/abs/hep-ph/0312378).
- [7] M. Drees, R. Godbole and P. Roy, *Theory And Phenomenology Of Sparticles: An Account Of Four-dim* World Scientific Publishing Company 2005.
- [8] H. Baer and X. Tata, *Weak scale supersymmetry: From superfields to scattering events*, Cambridge University Press 2006.
- [9] E. Gildener and S. Weinberg, “Symmetry Breaking and Scalar Bosons”, Phys. Rev. D **13** (1976) 3333.

- [10] E. Gildener, “Gauge-symmetry hierarchies”, Phys. Rev. D **14** (1976) 1667.
- [11] J. R. Ellis, S. Kelley and D. V. Nanopoulos, “Probing the desert using gauge coupling unification”, Phys. Lett. B **260** (1991) 131.
- [12] U. Amaldi, W. de Boer and H. Furstenau, “Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP”, Phys. Lett. B **260** (1991) 447.
- [13] G. Jungman, M. Kamionkowski and K. Griest, “Supersymmetric dark matter”, Phys. Rept. **267** (1996) 195, [arXiv:hep-ph/9506380](#).
- [14] A. B. Lahanas, N. E. Mavromatos and D. V. Nanopoulos, “WMAPing the universe: Supersymmetry, dark matter, dark energy, proton decay and collider physics”, Int. J. Mod. Phys. D **12** (2003) 1529, [arXiv:hep-ph/0308251](#).
- [15] W. L. Freedman and M. S. Turner, “Measuring and understanding the universe”, Rev. Mod. Phys. **75** (2003) 1433, [arXiv:astro-ph/0308418](#).
- [16] L. Roszkowski, “Particle dark matter: A Theorist’s perspective”, Pramana **62** (2004) 389, [arXiv:hep-ph/0404052](#).
- [17] G. Bertone, D. Hooper and J. Silk, “Particle dark matter: Evidence, candidates and constraints”, Phys. Rept. **405** (2005) 279, [arXiv:hep-ph/0404175](#).
- [18] K. A. Olive, “TASI lectures on astroparticle physics”, in Theoretical Advanced Study Institute in Elementary Particle Physics: Physics in $D \geq 4$ 2005, [arXiv:astro-ph/0503065](#).
- [19] H. Baer and X. Tata, Dark matter and the LHC 2009, [arXiv:0805.1905 \[hep-ph\]](#).
- [20] M. Drees and G. Gerbier, “Mini-Review of Dark Matter: 2012”, [arXiv:1204.2373 \[hep-ph\]](#).
- [21] S. Arrenberg et al., “Working Group Report: Dark Matter Complementarity”, in Community Summer Study 2013: Snowmass on the Mississippi 2013, [arXiv:1310.8621 \[hep-ph\]](#).
- [22] L. E. Ibanez and G. G. Ross, “SU(2)-L x U(1) Symmetry Breaking as a Radiative Effect of Supersymmetry Breaking in Guts”, Phys. Lett. B **110** (1982) 215.

- [23] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, “Renormalization of Supersymmetry Breaking Parameters Revisited”, Prog. Theor. Phys. **71** (1984) 413.
- [24] L. E. Ibanez and C. Lopez, “N=1 Supergravity, the Weak Scale and the Low-Energy Particle Spectrum”, Nucl. Phys. B **233** (1984) 511.
- [25] “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum and 20.3 fb⁻¹ of $\sqrt{s} = 8$ TeV proton-proton collision data”, .
- [26] “Search for strong production of supersymmetric particles in final states with missing transverse momentum and at least three b-jets using 20.1 fb⁻¹ of pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector.”, .
- [27] “Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at $\sqrt{s} = 8$ TeV with the ATLAS detector”, .
- [28] “Search for strongly produced superpartners in final states with two same sign leptons with the ATLAS detector using 21 fb⁻¹ of proton-proton collisions at $\sqrt{s}=8$ TeV.”, .
- [29] G. Aad et al. (ATLAS), “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum at $\sqrt{s}=8$ TeV proton-proton collisions using the ATLAS experiment”, JHEP **10** (2013) 130, [Erratum: JHEP 01, 109 (2014)], [arXiv:1308.1841 \[hep-ex\]](#).
- [30] “Search for Supersymmetry in Events with Large Missing Transverse Momentum, Jets, and at Least One Tau Lepton in 21 fb⁻¹ of $\sqrt{s} = 8$ TeV Proton-Proton Collision Data with the ATLAS Detector”, .
- [31] “Search for supersymmetry using razor variables in events with b-jets in pp collisions at 8 TeV”, .
- [32] G. Aad et al. (ATLAS), “Search for pair production of gluinos decaying via stop and sbottom in events with b-jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **94**[3] (2016) 032003, [arXiv:1605.09318 \[hep-ex\]](#).

- [33] G. Aad et al. (ATLAS), “Search for gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**[10] (2016) 565, [arXiv:1605.04285 \[hep-ex\]](#).
- [34] M. Aaboud et al. (ATLAS), “Search for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. C **76**[7] (2016) 392, [arXiv:1605.03814 \[hep-ex\]](#).
- [35] G. Aad et al. (ATLAS), “Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector”, Eur. Phys. J. C **76**[5] (2016) 259, [arXiv:1602.09058 \[hep-ex\]](#).
- [36] M. Aaboud et al. (ATLAS), “Search for squarks and gluinos in events with an isolated lepton, jets, and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **96**[11] (2017) 112010, [arXiv:1708.08232 \[hep-ex\]](#).
- [37] M. Aaboud et al. (ATLAS), “Search for a scalar partner of the top quark in the jets plus missing transverse momentum final state at $\sqrt{s}=13$ TeV with the ATLAS detector”, JHEP **12** (2017) 085, [arXiv:1709.04183 \[hep-ex\]](#).
- [38] M. Aaboud et al. (ATLAS), “Search for supersymmetry in final states with two same-sign or three leptons and jets using 36 fb^{-1} of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, JHEP **09** (2017) 084, [Erratum: JHEP 08, 121 (2019)], [arXiv:1706.03731 \[hep-ex\]](#).
- [39] M. Aaboud et al. (ATLAS), “Search for squarks and gluinos in final states with jets and missing transverse momentum using 36 fb^{-1} of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, Phys. Rev. D **97**[11] (2018) 112001, [arXiv:1712.02332 \[hep-ex\]](#).
- [40] M. Aaboud et al. (ATLAS), “Search for supersymmetry in final states with missing transverse momentum and multiple b -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **06** (2018) 107, [arXiv:1711.01901 \[hep-ex\]](#).
- [41] M. Aaboud et al. (ATLAS), “Search for new phenomena using the invariant mass distribution of same-flavour opposite-sign dilepton pairs in events with missing transverse momentum in

- $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **78**[8] (2018) 625, [arXiv:1805.11381 \[hep-ex\]](#).
- [42] M. Aaboud et al. (ATLAS), “Search for squarks and gluinos in final states with hadronically decaying τ -leptons, jets, and missing transverse momentum using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. D **99**[1] (2019) 012009, [arXiv:1808.06358 \[hep-ex\]](#).
- [43] G. Aad et al. (ATLAS), “Search for squarks and gluinos in final states with same-sign leptons and jets using 139 fb^{-1} of data collected with the ATLAS detector”, JHEP **06** (2020) 046, [arXiv:1909.08457 \[hep-ex\]](#).
- [44] G. Aad et al. (ATLAS), “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s} = 13$ TeV proton-proton collisions recorded by ATLAS in Run 2 of the LHC”, JHEP **10** (2020) 062, [arXiv:2008.06032 \[hep-ex\]](#).
- [45] G. Aad et al. (ATLAS), “Search for new phenomena in events with an energetic jet and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, [arXiv:2102.10874 \[hep-ex\]](#).
- [46] G. Aad et al. (ATLAS), “Search for new phenomena in final states with b -jets and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, [arXiv:2101.12527 \[hep-ex\]](#).
- [47] T. J. LeCompte and S. P. Martin, “Compressed supersymmetry after 1/fb at the Large Hadron Collider”, Phys. Rev. D **85** (2012) 035023, [arXiv:1111.6897 \[hep-ph\]](#).
- [48] H. K. Dreiner, M. Kramer and J. Tattersall, “How low can SUSY go? Matching, monojets and compressed spectra”, EPL **99**[6] (2012) 61001, [arXiv:1207.1613 \[hep-ph\]](#).
- [49] B. Bhattacharjee and K. Ghosh, “Degenerate SUSY search at the 8 TeV LHC”, [arXiv:1207.6289 \[hep-ph\]](#).
- [50] B. Bhattacharjee, A. Choudhury, K. Ghosh and S. Poddar, “Compressed supersymmetry at 14 TeV LHC”, Phys. Rev. D **89**[3] (2014) 037702, [arXiv:1308.1526 \[hep-ph\]](#).

- [51] T. Cohen, T. Golling, M. Hance, A. Henrichs, K. Howe, J. Loyal, S. Padhi and J. G. Wacker, “SUSY Simplified Models at 14, 33, and 100 TeV Proton Colliders”, JHEP **04** (2014) 117, [arXiv:1311.6480 \[hep-ph\]](#).
- [52] S. Mukhopadhyay, M. M. Nojiri and T. T. Yanagida, “Compressed SUSY search at the 13 TeV LHC using kinematic correlations and structure of ISR jets”, JHEP **10** (2014) 012, [arXiv:1403.6028 \[hep-ph\]](#).
- [53] M. Low and L.-T. Wang, “Neutralino dark matter at 14 TeV and 100 TeV”, JHEP **08** (2014) 161, [arXiv:1404.0682 \[hep-ph\]](#).
- [54] G. Aad et al. (ATLAS), “Search for squarks and gluinos in final states with one isolated lepton, jets, and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, [arXiv:2101.01629 \[hep-ex\]](#).
- [55] A. Djouadi et al. (MSSM Working Group), “The Minimal supersymmetric standard model: Group summary report”, in GDR (Groupement De Recherche) - Supersymetrie 1998, [arXiv:hep-ph/9901246](#).
- [56] M. Chakraborti, U. Chattopadhyay, A. Choudhury, A. Datta and S. Poddar, “The Electroweak Sector of the pMSSM in the Light of LHC - 8 TeV and Other Data”, JHEP **07** (2014) 019, [arXiv:1404.4841 \[hep-ph\]](#).
- [57] M. Chakraborti, U. Chattopadhyay, A. Choudhury, A. Datta and S. Poddar, “Reduced LHC constraints for higgsino-like heavier electroweakinos”, JHEP **11** (2015) 050, [arXiv:1507.01395 \[hep-ph\]](#).
- [58] A. Datta, N. Ganguly and S. Poddar, “New Limits on Heavier Electroweakinos and their LHC Signatures”, Phys. Lett. B **763** (2016) 213, [arXiv:1606.04391 \[hep-ph\]](#).
- [59] M. Chakraborti, A. Datta, N. Ganguly and S. Poddar, “Multilepton signals of heavier electroweakinos at the LHC”, JHEP **11** (2017) 117, [arXiv:1707.04410 \[hep-ph\]](#).
- [60] A. Datta and N. Ganguly, “The past, present and future of the heavier electroweakinos in the light of LHC and other data”, JHEP **01** (2019) 103, [arXiv:1809.05129 \[hep-ph\]](#).

- [61] G. Aad et al. (ATLAS), “Search for direct production of charginos, neutralinos and sleptons in final states with two leptons and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **05** (2014) 071, [arXiv:1403.5294 \[hep-ex\]](#).
- [62] G. Aad et al. (ATLAS), “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, JHEP **04** (2014) 169, [arXiv:1402.7029 \[hep-ex\]](#).
- [63] G. Aad et al. (ATLAS), “Search for the direct production of charginos, neutralinos and staus in final states with at least two hadronically decaying taus and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **10** (2014) 096, [arXiv:1407.0350 \[hep-ex\]](#).
- [64] G. Aad et al. (ATLAS), “Search for direct pair production of a chargino and a neutralino decaying to the 125 GeV Higgs boson in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. C **75**[5] (2015) 208, [arXiv:1501.07110 \[hep-ex\]](#).
- [65] G. Aad et al. (ATLAS), “Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb^{-1} of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, JHEP **02** (2021) 143, [arXiv:2010.14293 \[hep-ex\]](#).
- [66] U. Chattopadhyay, D. Das, P. Konar and D. P. Roy, “Looking for a heavy wino LSP in collider and dark matter experiments”, Phys. Rev. D **75** (2007) 073014, [arXiv:hep-ph/0610077](#).
- [67] U. Chattopadhyay, D. Das, A. Datta and S. Poddar, “Non-zero trilinear parameter in the mSUGRA model: Dark matter and collider signals at Tevatron and LHC”, Phys. Rev. D **76** (2007) 055008, [arXiv:0705.0921 \[hep-ph\]](#).
- [68] R. M. Godbole, M. Guchait and D. P. Roy, “Using Tau Polarization to probe the Stau Co-annihilation Region of mSUGRA Model at LHC”, Phys. Rev. D **79** (2009) 095015, [arXiv:0807.2390 \[hep-ph\]](#).
- [69] U. Chattopadhyay and D. Das, “Higgs funnel region of SUSY dark matter for small $\tan \beta$, RG effects on pseudoscalar Higgs boson with scalar mass non-universality”, Phys. Rev. D **79** (2009) 035007, [arXiv:0809.4065 \[hep-ph\]](#).

- [70] U. Chattopadhyay, D. Das and D. P. Roy, “Mixed Neutralino Dark Matter in Nonuniversal Gaugino Mass Models”, Phys. Rev. D **79** (2009) 095013, [arXiv:0902.4568 \[hep-ph\]](#).
- [71] U. Chattopadhyay, D. Das, D. K. Ghosh and M. Maity, “Probing the light Higgs pole resonance annihilation of dark matter in the light of XENON100 and CDMS-II observations”, Phys. Rev. D **82** (2010) 075013, [arXiv:1006.3045 \[hep-ph\]](#).
- [72] A. Choudhury and A. Datta, “New limits on top squark NLSP from LHC 4.7 fb^{-1} data”, Mod. Phys. Lett. A **27** (2012) 1250188, [arXiv:1207.1846 \[hep-ph\]](#).
- [73] S. Mohanty, S. Rao and D. P. Roy, “Predictions of a Natural SUSY Dark Matter Model for Direct and Indirect Detection Experiments”, JHEP **11** (2012) 175, [arXiv:1208.0894 \[hep-ph\]](#).
- [74] S. Bhattacharya, S. Chakdar, K. Ghosh and S. Nandi, “Non-universal SUGRA at LHC: Prospects and Discovery Potential”, Phys. Rev. D **89**[1] (2014) 015004, [arXiv:1309.0036 \[hep-ph\]](#).
- [75] K. Harigaya, K. Kaneta and S. Matsumoto, “Gaugino coannihilations”, Phys. Rev. D **89**[11] (2014) 115021, [arXiv:1403.0715 \[hep-ph\]](#).
- [76] Y. Huang, G. E. Addison, J. L. Weiland and C. L. Bennett, “Assessing Consistency Between WMAP 9-year and Planck 2015 Temperature Power Spectra”, Astrophys. J. **869**[1] (2018) 38, [arXiv:1804.05428 \[astro-ph.CO\]](#).
- [77] N. Aghanim et al. (Planck), “Planck 2018 results. VI. Cosmological parameters”, Astron. Astrophys. **641** (2020) A6, [arXiv:1807.06209 \[astro-ph.CO\]](#).
- [78] A. Choudhury and A. Datta, “Many faces of low mass neutralino dark matter in the unconstrained MSSM, LHC data and new signals”, JHEP **06** (2012) 006, [arXiv:1203.4106 \[hep-ph\]](#).
- [79] S. Profumo and C. E. Yaguna, “Gluino coannihilations and heavy bino dark matter”, Phys. Rev. D **69** (2004) 115009, [arXiv:hep-ph/0402208](#).
- [80] D. Feldman, Z. Liu and P. Nath, “Gluino NLSP, Dark Matter via Gluino Coannihilation, and LHC Signatures”, Phys. Rev. D **80** (2009) 015007, [arXiv:0905.1148 \[hep-ph\]](#).

- [81] J. Ellis, F. Luo and K. A. Olive, “Gluino Coannihilation Revisited”, JHEP **09** (2015) 127, [arXiv:1503.07142 \[hep-ph\]](#).
- [82] N. Nagata, H. Otono and S. Shirai, “Probing bino–gluino coannihilation at the LHC”, Phys. Lett. B **748** (2015) 24, [arXiv:1504.00504 \[hep-ph\]](#).
- [83] J. Ellis, J. L. Evans, F. Luo and K. A. Olive, “Scenarios for Gluino Coannihilation”, JHEP **02** (2016) 071, [arXiv:1510.03498 \[hep-ph\]](#).
- [84] D. S. Akerib et al. (LUX), “Results from a search for dark matter in the complete LUX exposure”, Phys. Rev. Lett. **118**[2] (2017) 021303, [arXiv:1608.07648 \[astro-ph.CO\]](#).
- [85] E. Aprile et al. (XENON), “Dark Matter Search Results from a One Ton-Year Exposure of XENON1T”, Phys. Rev. Lett. **121**[11] (2018) 111302, [arXiv:1805.12562 \[astro-ph.CO\]](#).
- [86] M. G. Aartsen et al. (IceCube), “Improved limits on dark matter annihilation in the Sun with the 79-string IceCube detector and implications for supersymmetry”, JCAP **04** (2016) 022, [arXiv:1601.00653 \[hep-ph\]](#).
- [87] G. Aad et al. (ATLAS), “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, Phys. Lett. B **716** (2012) 1, [arXiv:1207.7214 \[hep-ex\]](#).
- [88] S. Chatrchyan et al. (CMS), “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC”, Phys. Lett. B **716** (2012) 30, [arXiv:1207.7235 \[hep-ex\]](#).
- [89] A. Djouadi, “The Anatomy of electro-weak symmetry breaking. II. The Higgs bosons in the minimal supersymmetric model”, Phys. Rept. **459** (2008) 1, [arXiv:hep-ph/0503173](#).
- [90] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, “Towards high precision predictions for the MSSM Higgs sector”, Eur. Phys. J. C **28** (2003) 133, [arXiv:hep-ph/0212020](#).
- [91] B. Allanach, A. Djouadi, J. Kneur, W. Porod and P. Slavich, “Precise determination of the neutral Higgs boson masses in the MSSM”, JHEP **09** (2004) 044, [arXiv:hep-ph/0406166](#).

- [92] S. P. Martin, “Three-loop corrections to the lightest Higgs scalar boson mass in supersymmetry”, Phys. Rev. D **75** (2007) 055005, [arXiv:hep-ph/0701051](#).
- [93] R. V. Harlander, P. Kant, L. Mihaila and M. Steinhauser, “Higgs boson mass in supersymmetry to three loops”, Phys. Rev. Lett. **100** (2008) 191602, [arXiv:0803.0672 \[hep-ph\]](#).
- [94] A. Arbey, M. Battaglia, A. Djouadi and F. Mahmoudi, “The Higgs sector of the phenomenological MSSM in the light of the Higgs boson discovery”, JHEP **09** (2012) 107, [arXiv:1207.1348 \[hep-ph\]](#).
- [95] S. Heinemeyer, O. Stal and G. Weiglein, “Interpreting the LHC Higgs Search Results in the MSSM”, Phys. Lett. B **710** (2012) 201, [arXiv:1112.3026 \[hep-ph\]](#).
- [96] J. Harz, B. Herrmann, M. Klasen, K. Kovarik and P. Steppeler, “Theoretical uncertainty of the supersymmetric dark matter relic density from scheme and scale variations”, Phys. Rev. D **93**[11] (2016) 114023, [arXiv:1602.08103 \[hep-ph\]](#).
- [97] M. Klasen, K. Kovarik and P. Steppeler, “SUSY-QCD corrections for direct detection of neutralino dark matter and correlations with relic density”, Phys. Rev. D **94**[9] (2016) 095002, [arXiv:1607.06396 \[hep-ph\]](#).
- [98] G. Bertone, F. Calore, S. Caron, R. Ruiz, J. S. Kim, R. Trotta and C. Weniger, “Global analysis of the pMSSM in light of the Fermi GeV excess: prospects for the LHC Run-II and astroparticle experiments”, JCAP **04** (2016) 037, [arXiv:1507.07008 \[hep-ph\]](#).
- [99] M. Badziak, M. Olechowski and P. Szczerbiak, “Spin-dependent constraints on blind spots for thermal singlino-higgsino dark matter with(out) light singlets”, JHEP **07** (2017) 050, [arXiv:1705.00227 \[hep-ph\]](#).
- [100] K. C. Wong et al., “H0LiCOW – XIII. A 2.4 per cent measurement of H0 from lensed quasars: 5.3 σ tension between early- and late-Universe probes”, Mon. Not. Roy. Astron. Soc. **498**[1] (2020) 1420, [arXiv:1907.04869 \[astro-ph.CO\]](#).
- [101] A. Djouadi, M. Muhlleitner and M. Spira, “Decays of supersymmetric particles: The Program SUSY-HIT (SUSpect-SdecaY-Hdecay-InTerface)”, Acta Phys. Polon. B **38** (2007) 635, [arXiv:hep-ph/0609292](#).

- [102] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, JHEP **07** (2014) 079, [arXiv:1405.0301 \[hep-ph\]](#).
- [103] R. D. Ball et al., “Parton distributions with LHC data”, Nucl. Phys. B **867** (2013) 244, [arXiv:1207.1303 \[hep-ph\]](#).
- [104] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen and P. Z. Skands, “An introduction to PYTHIA 8.2”, Comput. Phys. Commun. **191** (2015) 159, [arXiv:1410.3012 \[hep-ph\]](#).
- [105] L. Lönnblad and S. Prestel, “Merging Multi-leg NLO Matrix Elements with Parton Showers”, JHEP **03** (2013) 166, [arXiv:1211.7278 \[hep-ph\]](#).
- [106] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens and M. Selvaggi (DELPHES 3), “DELPHES 3, A modular framework for fast simulation of a generic collider experiment”, JHEP **02** (2014) 057, [arXiv:1307.6346 \[hep-ex\]](#).
- [107] “See the website <https://twiki.cern.ch/twiki/bin/view/LHCPhysics>”, .
- [108] M. Drees and M. Nojiri, “Neutralino - nucleon scattering revisited”, Phys. Rev. D **48** (1993) 3483, [arXiv:hep-ph/9307208](#).
- [109] J. R. Ellis, K. A. Olive and C. Savage, “Hadronic Uncertainties in the Elastic Scattering of Supersymmetric Dark Matter”, Phys. Rev. D **77** (2008) 065026, [arXiv:0801.3656 \[hep-ph\]](#).
- [110] H. Ohki, H. Fukaya, S. Hashimoto, T. Kaneko, H. Matsufuru, J. Noaki, T. Onogi, E. Shintani and N. Yamada, “Nucleon sigma term and strange quark content from lattice QCD with exact chiral symmetry”, Phys. Rev. D **78** (2008) 054502, [arXiv:0806.4744 \[hep-lat\]](#).
- [111] J. Giedt, A. W. Thomas and R. D. Young, “Dark matter, the CMSSM and lattice QCD”, Phys. Rev. Lett. **103** (2009) 201802, [arXiv:0907.4177 \[hep-ph\]](#).
- [112] M. Perelstein and B. Shakya, “XENON100 implications for naturalness in the MSSM, NMSSM, and λ -supersymmetry model”, Phys. Rev. D **88**[7] (2013) 075003, [arXiv:1208.0833 \[hep-ph\]](#).

- [113] C. Beskidt, W. de Boer, D. I. Kazakov and F. Ratnikov, “Where is SUSY?”, JHEP **05** (2012) 094, [arXiv:1202.3366](#) [hep-ph].
- [114] C. Beskidt, W. de Boer, D. I. Kazakov and F. Ratnikov, “Constraints on Supersymmetry from LHC data on SUSY searches and Higgs bosons combined with cosmology and direct dark matter searches”, Eur. Phys. J. C **72** (2012) 2166, [arXiv:1207.3185](#) [hep-ph].
- [115] J. Bovy and S. Tremaine, “On the local dark matter density”, Astrophys. J. **756** (2012) 89, [arXiv:1205.4033](#) [astro-ph.GA].
- [116] P. Bhattacharjee, S. Chaudhury, S. Kundu and S. Majumdar, “Sizing-up the WIMPs of Milky Way : Deriving the velocity distribution of Galactic Dark Matter particles from the rotation curve data”, Phys. Rev. D **87** (2013) 083525, [arXiv:1210.2328](#) [astro-ph.GA].
- [117] M. Fairbairn, T. Douce and J. Swift, “Quantifying Astrophysical Uncertainties on Dark Matter Direct Detection Results”, Astropart. Phys. **47** (2013) 45, [arXiv:1206.2693](#) [astro-ph.CO].
- [118] J. Hisano, K. Nakayama and M. Yamanaka, “Implications of CDMS II result on Higgs sector in the MSSM”, Phys. Lett. B **684** (2010) 246, [arXiv:0912.4701](#) [hep-ph].
- [119] G. Belanger, A. Mjallal and A. Pukhov, “Recasting direct detection limits within micrOMEGAs and implication for non-standard Dark Matter scenarios”, Eur. Phys. J. C **81**[3] (2021) 239, [arXiv:2003.08621](#) [hep-ph].
- [120] J. R. Ellis, A. Ferstl and K. A. Olive, “Reevaluation of the elastic scattering of supersymmetric dark matter”, Phys. Lett. B **481** (2000) 304, [arXiv:hep-ph/0001005](#).
- [121] W. H. Press and D. N. Spergel, “Capture by the sun of a galactic population of weakly interacting massive particles”, Astrophys. J. **296** (1985) 679.
- [122] J. Silk, K. A. Olive and M. Srednicki, “The Photino, the Sun and High-Energy Neutrinos”, Phys. Rev. Lett. **55** (1985) 257.
- [123] A. Peter, “Dark matter bound to the Solar System: consequences for annihilation searches”, in 44th Rencontres de Moriond on Electroweak Interactions and Unified Theories, p. 359–366 2009, [arXiv:0905.2456](#) [astro-ph.HE].

- [124] A. Ibarra, M. Totzauer and S. Wild, “Higher order dark matter annihilations in the Sun and implications for IceCube”, JCAP **04** (2014) 012, [arXiv:1402.4375 \[hep-ph\]](#).
- [125] A. Ibarra, M. Totzauer and S. Wild, “High-energy neutrino signals from the Sun in dark matter scenarios with internal bremsstrahlung”, JCAP **12** (2013) 043, [arXiv:1311.1418 \[hep-ph\]](#).
- [126] R. Catena, “Dark matter signals at neutrino telescopes in effective theories”, JCAP **04** (2015) 052, [arXiv:1503.04109 \[hep-ph\]](#).
- [127] A. Chatterjee, A. Choudhury, A. Datta and B. Mukhopadhyaya, “Gluino mass limits with sbottom NLSP in coannihilation scenarios”, JHEP **01** (2015) 154, [arXiv:1411.6467 \[hep-ph\]](#).
- [128] J. L. Feng, K. T. Matchev and T. Moroi, “Focus points and naturalness in supersymmetry”, Phys. Rev. D **61** (2000) 075005, [arXiv:hep-ph/9909334](#).
- [129] J. L. Feng, “Naturalness and the Status of Supersymmetry”, Ann. Rev. Nucl. Part. Sci. **63** (2013) 351, [arXiv:1302.6587 \[hep-ph\]](#).