



function  $\tilde{\phi}$  via

$$\phi(\mathbf{p}) = R^{3/2} \tilde{\phi}(R\mathbf{p}), \quad (5)$$

where  $R$  denotes the characteristic size of the wave packet with  $R \rightarrow 0$  corresponding to a sharp localization. The charge density distribution in this state has the form

$$\begin{aligned} \rho_\phi(\mathbf{X} + \mathbf{r}) &\equiv \langle \Phi, \mathbf{X} | \hat{\rho}(\mathbf{r}, 0) | \Phi, \mathbf{X} \rangle \\ &= \int \frac{d^3 p d^3 p'}{(2\pi)^3 \sqrt{4EE'}} (E + E') F(q^2) \phi^*(\mathbf{p}') \phi(\mathbf{p}) \\ &\quad \times e^{i\mathbf{q} \cdot (\mathbf{X} + \mathbf{r})}, \end{aligned} \quad (6)$$

where  $\mathbf{q} = \mathbf{p}' - \mathbf{p}$  and  $q^2 = (E' - E)^2 - \mathbf{q}^2$ . Without loss of generality we choose  $\mathbf{X} = 0$  to place the system at the origin. Finally, introducing the total and relative momentum variables via  $\mathbf{p} = \mathbf{P} - \mathbf{q}/2$  and  $\mathbf{p}' = \mathbf{P} + \mathbf{q}/2$ , the charge density is written as

$$\begin{aligned} \rho_\phi(\mathbf{r}) &= \int \frac{d^3 P d^3 q}{(2\pi)^3 \sqrt{4EE'}} (E + E') F[(E - E')^2 - \mathbf{q}^2] \\ &\quad \times \phi\left(\mathbf{P} - \frac{\mathbf{q}}{2}\right) \phi^*\left(\mathbf{P} + \frac{\mathbf{q}}{2}\right) e^{i\mathbf{q} \cdot \mathbf{r}}, \end{aligned} \quad (7)$$

where the energies are  $E = \sqrt{m^2 + \mathbf{P}^2 - \mathbf{P} \cdot \mathbf{q} + \mathbf{q}^2/4}$  and  $E' = \sqrt{m^2 + \mathbf{P}^2 + \mathbf{P} \cdot \mathbf{q} + \mathbf{q}^2/4}$ .

The traditional (“naive”) interpretation of the charge density in terms of the Fourier transform of the form factor in the Breit frame,  $F(q^2) = F(-\mathbf{q}^2)$ , emerges by first taking the static limit (i.e.,  $m \rightarrow \infty$ ) by substituting  $E = E' = m$  in the integrand in Eq. (7),

$$\begin{aligned} \rho_{\phi, \text{naive}}(\mathbf{r}) &= \int \frac{d^3 P d^3 q}{(2\pi)^3} \phi\left(\mathbf{P} - \frac{\mathbf{q}}{2}\right) \phi^*\left(\mathbf{P} + \frac{\mathbf{q}}{2}\right) \\ &\quad \times F(-\mathbf{q}^2) e^{i\mathbf{q} \cdot \mathbf{r}}, \end{aligned} \quad (8)$$

and subsequently localizing the wave packet by taking the limit  $R \rightarrow 0$  [11, 12]. This can be done without specifying the functions  $F(q^2)$  and  $\phi(\mathbf{p})$  using the method of dimensional counting [14] or, alternatively, the strategy of regions [15]. For  $F(q^2)$  decreasing at large  $q^2$  faster than  $1/q^2$ , the only non-vanishing contribution to  $\rho_{\phi, \text{naive}}(\mathbf{r})$  in the  $R \rightarrow 0$  limit is obtained by substituting  $\mathbf{P} = \tilde{\mathbf{P}}/R$ , expanding the integrand in Eq. (8) in  $R$  around  $R = 0$  and keeping the zeroth order term. The resulting naive charge density has the familiar form

$$\begin{aligned} \rho_{\text{naive}}(r) &= \int \frac{d^3 \tilde{P} d^3 q}{(2\pi)^3} F(-\mathbf{q}^2) |\tilde{\phi}(\tilde{\mathbf{P}})|^2 e^{i\mathbf{q} \cdot \mathbf{r}} \\ &= \int \frac{d^3 q}{(2\pi)^3} F(-\mathbf{q}^2) e^{i\mathbf{q} \cdot \mathbf{r}}, \end{aligned} \quad (9)$$

where in the second equality we made use of Eq. (4). Here and in what follows,  $r \equiv |\mathbf{r}|$ . We have dropped the subscript  $\phi$  to indicate that the above expression is independent of the wave packet shape.

On the other hand, the method of dimensional counting allows one to take the  $R \rightarrow 0$  limit in Eq. (7) without employing the static approximation. Following the same steps as before but for arbitrary  $m$ , we obtain

$$\rho_\phi(\mathbf{r}) = \int \frac{d^3 \tilde{P} d^3 q}{(2\pi)^3} F\left[\frac{(\tilde{\mathbf{P}} \cdot \mathbf{q})^2}{\tilde{\mathbf{P}}^2} - \mathbf{q}^2\right] |\tilde{\phi}(\tilde{\mathbf{P}})|^2 e^{i\mathbf{q} \cdot \mathbf{r}}. \quad (10)$$

The resulting density depends on the shape of the wave packet unless it is spherically symmetric. Since there is no preferred direction in the rest frame of the system, we *define* the charge density distribution in the rest frame by employing spherically symmetric wave packets with  $\tilde{\phi}(\tilde{\mathbf{P}}) = \tilde{\phi}(|\tilde{\mathbf{P}}|)$ . Then, using spherical coordinates to perform the integration over  $\tilde{\mathbf{P}}$  in Eq. (10), we arrive at the final form of the charge density distribution in the rest frame of a particle

$$\rho(r) = \int \frac{d^3 q}{(2\pi)^3} e^{i\mathbf{q} \cdot \mathbf{r}} \int_{-1}^{+1} d\alpha \frac{1}{2} F[(\alpha^2 - 1) \mathbf{q}^2]. \quad (11)$$

While it is argued in Ref. [11] that the traditional result  $\rho_{\text{naive}}(r)$  is valid for the hierarchy of scales  $\Delta \gg 1/m$ , comparing the approximate and exact expressions in Eqs. (9) and (11), respectively, shows that the accuracy of the static approximation leading to  $\rho_{\text{naive}}(r)$  does not depend upon the particle mass  $m$ . It is also clear that the validity of Eq. (11), which provides an unambiguous relationship between the matrix element of the local charge density operator  $\hat{\rho}(0)$  in a quantum system and the experimentally measurable form factor  $F(q^2)$ , does not depend on the relation between the intrinsic size of the system  $\Delta$  and its Compton wavelength  $1/m$  (in contrast to what is claimed in Ref. [11]).

*Discussion:* A striking feature of the obtained result for  $\rho(r)$  is its independence of the particle’s mass. This implies that the traditional expression for the charge density,  $\rho_{\text{naive}}(r)$ , does *not* emerge from  $\rho(r)$  by taking the static limit:  $\rho_{\text{naive}}(r) \neq \lim_{m \rightarrow \infty} \rho(r)$ . At first glance this seems puzzling as one expects the conventional static result to be a better approximation for heavy systems like atoms or atomic nuclei [11, 12]. The reason for this mismatch is the non-commutativity of the  $R \rightarrow 0$  and  $m \rightarrow \infty$  limits of  $\rho_\phi(\mathbf{r})$  in Eq. (7), as implicitly shown in Figs. 1-3 of Ref. [11]. While the static limit and, more generally, the non-relativistic approximation is perfectly valid when calculating the form factor in Eq. (2) provided  $-q^2 \ll m^2$ , *it is violated in certain momentum regions when performing the integration in Eq. (7).*

To have a simple example demonstrating the non-commutativity of the  $m \rightarrow \infty$  and  $R \rightarrow 0$  limits consider the wave packet in one spatial dimension with

$$\phi(p) = \sqrt{\frac{2R}{\pi}} \frac{1}{1 + R^2 p^2}, \quad (12)$$

and the form factor

$$F(q_0^2 - q^2) = \frac{2}{2 - \Delta^2(q_0^2 - q^2)}, \quad (13)$$

so that  $F(0) = 1$  and  $F'(0) = \Delta^2/2$ . We calculate the second order moment of the charge distribution using the version of Eq. (7) in one spatial dimension,

$$\langle x^2 \rangle_\phi = \int_{-\infty}^{+\infty} dx x^2 \int_{-\infty}^{+\infty} \frac{dP dq}{2\pi \sqrt{4EE'}} (E + E') \quad (14)$$

$$\times F[(E - E')^2 - k q^2] \phi\left(P - \frac{q}{2}\right) \phi^*\left(P + \frac{q}{2}\right) e^{iqx}.$$

For demonstration purposes, we have introduced a control parameter  $k$  to be set to  $k = 1$  in the final result. The integral in Eq. (14) can be easily calculated by writing the factors of  $x$  as derivatives acting on the exponential function. The resulting expression has the form

$$\langle x^2 \rangle_\phi = k\Delta^2 - \frac{\Delta^2}{(1 + mR)^2} + \frac{R^2}{2} - \frac{R}{4m(1 + mR)^3}. \quad (15)$$

Taking the limit  $R \rightarrow 0$  in Eq. (15) leads to

$$\langle x^2 \rangle = (k - 1)\Delta^2 = 0, \quad (16)$$

which does not depend on the mass  $m$ . On the other hand, taking first the static limit  $m \rightarrow \infty$  and subsequently the  $R \rightarrow 0$  limit we obtain a different result

$$\langle x^2 \rangle_{\text{naive}} = k\Delta^2 = \Delta^2. \quad (17)$$

The method of dimensional counting reproduces exactly Eq. (16), while Eq. (17) is obtained by first taking the static limit in the integrand of Eq. (14).

We now turn to the interpretation of our result in Eq. (11). Clearly, the dependence of  $\rho(r)$  on the angle-averaged form factor  $\frac{1}{2} \int_{-1}^{+1} d\alpha F[(\alpha^2 - 1)\mathbf{q}^2]$  rather than the Breit frame expression  $F(-\mathbf{q}^2)$  affects the radial profile of the charge density. To quantify the magnitude of this effect, we compare in Fig. 1  $\rho(r)$  and  $\rho_{\text{naive}}(r)$  for a charged and a neutral particle. For illustrative purposes we employ here simple parametrizations of the nucleon form factors, namely the dipole proton form factor  $F_p(q^2) = G_D(q^2) = (1 - q^2/\Lambda^2)^{-2}$  with  $\Lambda^2 = 0.71 \text{ GeV}^2$  and the Galster-type parametrization of the neutron form factor  $F_n(q^2) = A\tau/(1+B\tau) G_D(q^2)$  from Ref. [16], where  $\tau = -q^2/(4m_p^2)$ ,  $A = 1.70$  and  $B = 3.30$ .

To gain further insights into the relationship between the charge density and the form factor it is instructive to rewrite Eq. (11) in coordinate independent form as

$$\rho(r) = \frac{1}{4\pi} \int d^2\hat{n} \rho_{\hat{n}}(\mathbf{r}), \quad (18)$$

where  $\hat{n} \equiv \mathbf{n}/|\mathbf{n}|$  is a unit vector and

$$\rho_{\hat{n}}(\mathbf{r}) = \int \frac{d^3q}{(2\pi)^3} F(-\mathbf{q}_\perp^2) e^{i\mathbf{q}\cdot\mathbf{r}} = \rho_{\hat{n}}(r_\parallel) \rho_{\hat{n}}(r_\perp). \quad (19)$$

Here,  $\mathbf{q}_\perp = \hat{n} \times (\mathbf{q} \times \hat{n})$ ,  $\mathbf{r}_\perp = \hat{n} \times (\mathbf{r} \times \hat{n})$ ,  $r_\parallel = \mathbf{r} \cdot \hat{n}$ ,  $r_\perp \equiv |\mathbf{r}_\perp|$ , and the one- and two-dimensional densities in




FIG. 1: Radial charge density distributions  $4\pi r^2 \rho(r)$  (solid lines) and  $4\pi r^2 \rho_{\text{naive}}(r)$  (dashed lines) for a charged and a neutral particle using the dipole and Galster-type parametrizations of the proton and neutron electric form factors, respectively.

the  $\hat{n}$  and  $\mathbf{r}_\perp$  directions are given by

$$\rho_{\hat{n}}(r_\parallel) = \int \frac{dq_\parallel}{2\pi} e^{iq_\parallel r_\parallel} = \delta(r_\parallel), \quad (20)$$

$$\rho_{\hat{n}}(r_\perp) = \int \frac{d^2q_\perp}{(2\pi)^2} F(-\mathbf{q}_\perp^2) e^{i\mathbf{q}_\perp \cdot \mathbf{r}_\perp},$$

with  $q_\parallel = \mathbf{q} \cdot \hat{n}$ . These expressions make it clear that  $F(q^2)$  effectively measures the deviation from the point-like limit in *two* rather than three spatial dimensions. This in turn explains the squeezing of the radial charge density relative to the naive result as shown in Fig. 1.

To further elaborate on this point we compute radial moments of the charge distributions in  $d \in \mathbb{N}$  spatial dimensions. We start with the inverse Fourier transform of Eq. (9)

$$F(-\mathbf{q}^2) = \int d^d r \rho_{\text{naive}}(r) e^{-i\mathbf{q}\cdot\mathbf{r}}. \quad (21)$$

Taking the  $k^{\text{th}}$  derivative of this expression at  $-\mathbf{q}^2 = 0$ ,  $F^{(k)}(0)$ , we find for  $\langle r^{2k} \rangle_{\text{naive}}^{(d)} \equiv \int d^d r \rho_{\text{naive}}(r) r^{2k}$ :

$$\langle r^{2k} \rangle_{\text{naive}}^{(d)} = \frac{2^{2k} \Gamma(d/2 + k)}{\Gamma(d/2)} F^{(k)}(0). \quad (22)$$

For  $d = 3$ , this reduces to the well-known expression

$$\langle r^{2k} \rangle_{\text{naive}} = \frac{(2k + 1)!}{k!} F^{(k)}(0). \quad (23)$$

On the other hand, using Eqs. (18)-(20) generalized to  $d$  dimensions and noting that  $\langle r^{2k} \rangle_{\hat{n}}^{(d)} \equiv \int d^d r \rho_{\hat{n}}(\mathbf{r}) r^{2k}$  does not depend of  $\hat{n}$ , we obtain for  $d \geq 2$

$$\langle r^{2k} \rangle^{(d)} = \langle r^{2k} \rangle_{\text{naive}}^{(d-1)}, \quad (24)$$

so that in  $d = 3$  spatial dimensions,

$$\langle r^{2k} \rangle = 2^{2k} k! F^{(k)}(0). \quad (25)$$

Notice further that in one spatial dimension,  $\rho(r) = \delta(r)$  independently of the form factor. This explains the vanishing result for the second moment in the considered one-dimensional example, see Eq. (16).

*The charge density in moving frames:* While the static approximation  $\rho_{\text{naive}}(r)$  does not depend on the frame, the expressions for  $\rho(r)$  in Eqs. (11) and (18) are valid in the rest frame of the system. It is straightforward to generalize these results to a boosted frame.

We start with the general expression for  $\rho_\phi(\mathbf{r})$  in Eq. (7) and replace  $\phi(\mathbf{p})$  with  $\phi_{\mathbf{v}}(\mathbf{p})$ , where  $\mathbf{v}$  denotes the boost velocity. Differently to  $\phi(\mathbf{p})$ , we cannot regard the function  $\phi_{\mathbf{v}}(\mathbf{p})$  to be spherically symmetric. Thus, we need to express  $\phi_{\mathbf{v}}(\mathbf{p})$  in terms of the rest frame quantity  $\phi(\mathbf{p})$  in order to obtain a wave packet independent definition for the charge density in the  $R \rightarrow 0$  limit. Using Eq. (3) and the Lorentz transformation properties of the momentum eigenstates  $|p\rangle$ , one finds [17]

$$\phi_{\mathbf{v}}(\mathbf{p}) = \sqrt{\gamma \left(1 - \frac{\mathbf{v} \cdot \mathbf{p}}{E}\right)} \phi[\mathbf{p}_\perp + \gamma(\mathbf{p}_\parallel - \mathbf{v}E)], \quad (26)$$

where  $\gamma = (1 - v^2)^{-1/2}$ ,  $\mathbf{p}_\parallel = (\mathbf{p} \cdot \hat{\mathbf{v}})\hat{\mathbf{v}}$ ,  $\mathbf{p}_\perp = \mathbf{p} - \mathbf{p}_\parallel$  and  $E = \sqrt{m^2 + \mathbf{p}^2}$ . We note in passing that Eq. (26) ensures the invariance of the normalization of the wave packet [17]. Then, following the same steps as in the case of the rest frame and using the method of dimensional counting to evaluate the  $R \rightarrow 0$  limit we arrive at

$$\rho_{\phi, \mathbf{v}}(\mathbf{r}) = \int \frac{d^3 \tilde{P} d^3 q}{(2\pi)^3} \frac{\gamma(\tilde{P} - \mathbf{v} \cdot \tilde{P})}{\tilde{P}} F\left[\frac{(\tilde{P} \cdot \mathbf{q})^2}{\tilde{P}^2} - \mathbf{q}^2\right] \times \left| \tilde{\phi}[\tilde{P}_\perp + \gamma(\tilde{P}_\parallel - \mathbf{v}\tilde{P})] \right|^2 e^{i\mathbf{q} \cdot \mathbf{r}}, \quad (27)$$

where  $\tilde{P} \equiv |\tilde{\mathbf{P}}|$ . We now change the integration variable  $\tilde{\mathbf{P}} \rightarrow \tilde{\mathbf{P}}' = \tilde{\mathbf{P}}_\perp + \gamma(\tilde{\mathbf{P}}_\parallel - \mathbf{v}\tilde{P})$ . Using the relations  $\tilde{\mathbf{P}}_\parallel = \gamma(\tilde{\mathbf{P}}'_\parallel + \mathbf{v}\tilde{P}')$  and  $\tilde{P} = \gamma(\tilde{P}' + v\tilde{P}'_\parallel)$ , it is easy to verify that the Jacobian of the change of variables  $\tilde{\mathbf{P}} \rightarrow \tilde{\mathbf{P}}'$  cancels the first factor in the integrand in Eq. (27), yielding

$$\rho_{\phi, \mathbf{v}}(\mathbf{r}) = \int \frac{d^3 \tilde{P}' d^3 q}{(2\pi)^3} |\tilde{\phi}(\tilde{\mathbf{P}}')|^2 e^{i\mathbf{q} \cdot \mathbf{r}} \times F\left\{ \frac{[\tilde{\mathbf{P}}'_\perp \cdot \mathbf{q}_\perp + \gamma(\tilde{\mathbf{P}}'_\parallel + \mathbf{v}\tilde{P}') \cdot \mathbf{q}_\parallel]^2}{\gamma^2(\tilde{P}' + v\tilde{P}'_\parallel)^2} - \mathbf{q}^2 \right\}. \quad (28)$$

Using Eq. (4) and the spherical symmetry of  $\tilde{\phi}(\tilde{\mathbf{P}}')$ , the integration over  $\tilde{P}'$  becomes trivial. The remaining angular integration over  $\hat{\tilde{\mathbf{P}}}'$  can be done in spherical coordinates. We align the  $z$ - and  $x$ -axes along the  $\mathbf{v}$  and  $\mathbf{q}_\perp$  directions, respectively, and denote  $\eta = \cos\theta$ . Our final result then reads:

$$\rho_{\mathbf{v}}(\mathbf{r}) = \int \frac{d^3 q}{(2\pi)^3} \bar{F}(q_\parallel, q_\perp) e^{i\mathbf{q} \cdot \mathbf{r}}, \quad (29)$$

with  $q_\parallel \equiv \hat{\mathbf{v}} \cdot \mathbf{q}$ ,  $q_\perp \equiv |\mathbf{q}_\perp|$  and

$$\bar{F}(q_\parallel, q_\perp) = \frac{1}{4\pi} \int_{-1}^{+1} d\eta \int_0^{2\pi} d\phi \times F\left\{ \frac{[\sqrt{1 - \eta^2} \cos\phi q_\perp + \gamma(\eta + v)q_\parallel]^2}{\gamma^2(1 + v\eta)^2} - \mathbf{q}^2 \right\}. \quad (30)$$

In the IMF with  $v \rightarrow 1$  and  $\gamma \rightarrow \infty$ , the charge density turns into the usual two-dimensional distribution in the transverse plane,  $\rho_{\text{IMF}}(\mathbf{r}) = \delta(r_\parallel) \rho_{\text{IMF}}(r_\perp)$  with

$$\rho_{\text{IMF}}(r_\perp) = \int \frac{d^2 q_\perp}{(2\pi)^2} F(-\mathbf{q}_\perp^2) e^{i\mathbf{q}_\perp \cdot \mathbf{r}_\perp}. \quad (31)$$

One can also verify that Eq. (29) reduces to the rest frame expression in Eq. (11) in the limit  $v \rightarrow 0$ , albeit this relationship appears somewhat obscured.

Again, it is instructive to rewrite Eqs. (29,30) in a coordinate independent form similar to the rest frame expressions in Eqs. (18)-(20). We introduce a unit vector  $\hat{\mathbf{m}} \equiv \hat{\tilde{\mathbf{P}}}'$  and define a vector valued function

$$\mathbf{n}(\mathbf{v}, \hat{\mathbf{m}}) = \hat{\mathbf{v}} \times (\hat{\mathbf{m}} \times \hat{\mathbf{v}}) + \gamma(\hat{\mathbf{m}} \cdot \hat{\mathbf{v}} + v)\hat{\mathbf{v}}. \quad (32)$$

Then, the charge density  $\rho_{\mathbf{v}}(\mathbf{r})$  can be written as

$$\rho_{\mathbf{v}}(\mathbf{r}) = \frac{1}{4\pi} \int d^2 \hat{m} \rho_{\hat{\mathbf{n}}(\mathbf{v}, \hat{\mathbf{m}})}(\mathbf{r}), \quad (33)$$

where  $\rho_{\hat{\mathbf{n}}(\mathbf{v}, \hat{\mathbf{m}})}(\mathbf{r}) \equiv \rho_{\hat{\mathbf{n}}}(\mathbf{r})$  is defined in Eqs. (19), (20). In this form, both extreme limits for the boosting velocity become particularly transparent by using the relations  $\hat{\mathbf{n}}(\mathbf{v}, \hat{\mathbf{m}}) \xrightarrow{v \rightarrow 0} \hat{\mathbf{m}}$  and  $\hat{\mathbf{n}}(\mathbf{v}, \hat{\mathbf{m}}) \xrightarrow{v \rightarrow 1} \hat{\mathbf{v}}$ , leading evidently to Eqs. (18) and (31), respectively.

Last but not least, we emphasize that radial moments of the charge distribution are, in fact, frame independent, i.e.  $\langle r^{2k} \rangle_{\mathbf{v}} = \langle r^{2k} \rangle$ , in spite of  $\rho_{\mathbf{v}}(\mathbf{r})$  being not spherically symmetric for  $v \neq 0$ . This remarkable feature follows from Eq. (33) by noting that  $\int d^3 r \rho_{\hat{\mathbf{n}}(\mathbf{v}, \hat{\mathbf{m}})}(\mathbf{r}) r^{2k}$  does not depend on  $\mathbf{v}$  and  $\hat{\mathbf{m}}$ . It can also be verified by showing that radial moments of  $\rho_\phi(\mathbf{r})$  in Eq. (10) do not depend on  $\tilde{\phi}(\tilde{\mathbf{P}})$  even if this function is not spherically symmetric.

*Summary and conclusions:* In summary, we introduced an unambiguous definition of a spatial distribution of the expectation values of local operators in spin-0 systems independent of the specific form of the wave packet in which the state was prepared. Our definition also applies to systems whose intrinsic size is comparable or even smaller than the Compton wavelength. We found remarkably simple relationships between the electric form factor and the charge density in the rest and moving frames, thereby reproducing the well-known result in the infinite momentum frame. We have also demonstrated that radial moments of the charge distribution are frame independent.

Our results suggest that form factors effectively measure the deviation from the point-like limit in two rather

than three spatial dimensions. This implies, in particular, that the second moment of the charge distribution, the quantity that should be interpreted as the mean square charge radius of the system, is related to the form factor slope via  $\langle r^2 \rangle = 4F'(0)$  in contrast to the usual relationship  $\langle r^2 \rangle_{\text{naive}} = 6F'(0)$  motivated by the Breit frame distribution  $\rho_{\text{naive}}(r)$ . Thus, the actual size of e.g. the proton measured by the charge distribution is  $\sqrt{\langle r_p^2 \rangle} = 0.6866(3)$  fm rather than  $\sqrt{\langle r_p^2 \rangle_{\text{naive}}} = 0.8409(4)$  fm [18] as commonly accepted. Differently to Refs. [7–13], our results show that the approximation  $\rho_{\text{naive}}(r)$  does not emerge in the static limit of the exact expression for  $\rho(r)$ , and its accuracy is independent of the particle’s mass.

We note that for heavy systems with  $(m\Delta)^{-1} \equiv \epsilon \ll 1$ , one may alternatively attempt to define  $\rho(\mathbf{r})$  using wave packets with  $\epsilon \ll (mR)^{-1} \ll 1$  as suggested in [11], by choosing e.g.  $(mR)^{-1} \sim \mathcal{O}(\epsilon^{1/2})$ . While this leads to an unambiguous definition of the charge density in the static limit  $\epsilon \rightarrow 0$  with  $\rho(\mathbf{r}) \rightarrow \rho_{\text{naive}}(r)$ , corrections beyond this limit are wave packet dependent.

Our analysis can be straightforwardly generalized to systems with non-vanishing spin and to spatial distributions introduced in Refs. [1–6].

*Acknowledgements:* We thank Bob Jaffe for useful comments on the manuscript. This work was supported in part by BMBF (Grant No. 05P18PCFP1), by DFG and NSFC through funds provided to the Sino-German CRC 110 “Symmetries and the Emergence of Structure in QCD” (NSFC Grant No. 11621131001, DFG Project-ID 196253076 - TRR 110), by ERC NuclearTheory (grant No. 885150) and ERC EXOTIC (grant No. 101018170), by CAS through a President’s International

Fellowship Initiative (PIFI) (Grant No. 2018DM0034), by the VolkswagenStiftung (Grant No. 93562), by the EU Horizon 2020 research and innovation programme (STRONG-2020, grant agreement No. 824093), and by the Heisenberg-Landau Program 2021.

---

\* Deceased.

- [1] R. Hofstadter, F. Bumiller, and M. R. Yearian, *Rev. Mod. Phys.* **30**, 482 (1958).
- [2] F. J. Ernst, R. G. Sachs and K. C. Wali, *Phys. Rev.* **119**, 1105-1114 (1960).
- [3] R. G. Sachs, *Phys. Rev.* **126**, 2256-2260 (1962).
- [4] M. V. Polyakov and A. G. Shuvaev, [arXiv:hep-ph/0207153 [hep-ph]].
- [5] M. V. Polyakov, *Phys. Lett. B* **555**, 57 (2003).
- [6] M. V. Polyakov and P. Schweitzer, *Int. J. Mod. Phys. A* **33** (2018) no.26, 1830025.
- [7] M. Burkardt, *Phys. Rev. D* **62** (2000), 071503(R), [erratum: *Phys. Rev. D* **66** (2002), 119903(E)].
- [8] G. A. Miller, *Phys. Rev. Lett.* **99**, 112001 (2007).
- [9] G. A. Miller, *Phys. Rev. C* **79**, 055204 (2009).
- [10] G. A. Miller, *Ann. Rev. Nucl. Part. Sci.* **60** (2010), 1-25.
- [11] R. L. Jaffe, *Phys. Rev. D* **103** (2021) no.1, 016017.
- [12] G. A. Miller, *Phys. Rev. C* **99**, no.3, 035202 (2019).
- [13] A. Freese and G. A. Miller, *Phys. Rev. D* **103**, 094023 (2021).
- [14] J. Gegelia, G. S. Japaridze and K. S. Turashvili, *Theor. Math. Phys.* **101**, 1313-1319 (1994).
- [15] M. Beneke and V. A. Smirnov, *Nucl. Phys. B* **522**, 321-344 (1998).
- [16] J. J. Kelly, *Phys. Rev. C* **70**, 068202 (2004).
- [17] S. E. Hoffmann, [arXiv:1804.00548 [quant-ph]].
- [18] P. A. Zyla *et al.* [Particle Data Group], *PTEP* **2020**, no.8, 083C01 (2020).