

Recycled entanglement detection by arbitrarily many sequential and independent pairs of observers

Mahasweta Pandit,¹ Chirag Srivastava,² and Ujjwal Sen²

¹*Institute of Theoretical Physics and Astrophysics, Faculty of Mathematics,
Physics and Informatics, University of Gdańsk, 80-308 Gdańsk, Poland*

²*Harish-Chandra Research Institute, HBNI, Chhatnag Road, Jhansi, Allahabad 211 019, India*

We investigate the witnessing of two-qubit entangled states by sequential and independent pairs of observers, with both observers of each pair acting independently on their part of the shared state from spatially separated laboratories, and subsequently passing their qubits to the next pair in the sequence. It has previously been conjectured that not more than one pair of observers can detect Clauser-Horne-Shimony-Holt “Bell-nonlocal” correlations in a similar set-up. This is intriguing since it is possible to have an arbitrarily long sequence of Bell-nonlocal correlations when only a single observer is allowed to share a bipartite state with multiple observers at the other end. It is therefore interesting to ask whether such restrictions are also present when entangled correlations are considered in the scenario of multiple pairs of observers. We find that a two-qubit entangled state can be used to witness entanglement arbitrarily many times, by pairs of observers, acting sequentially and independently. We prove the statement to be true when the initial pair of observers in the sequence share any pure entangled state or when they share a state from a class of mixed entangled states. We demonstrate that the phenomenon can also be observed for a certain class of entangled states in which an arbitrarily long sequence of observer pairs witnessing entanglement is reached in the limit of the initial entanglement content tending to a vanishing amount.

I. INTRODUCTION

Quantum entanglement [1, 2] is among the most crucial resources in quantum information processing and communication. It plays a key role in numerous applications, such as quantum cryptography [3, 4], quantum dense coding [5], quantum teleportation [6], entanglement swapping [7, 8] and device-independent tasks like key distribution [9–11], randomness amplification [12], randomness expansion [13–15], etc. It is therefore useful to learn about different techniques of characterizing, quantifying, and utilizing entanglement. A valuable tool to analyze and characterize entanglement exploits a class of functionals called *entanglement witnesses* (EWs) [1, 2, 16]. Given an entangled state, there always exists an EW [2, 17–20], and for example, Bell inequalities [21, 22], such as the Clauser-Horne-Shimony-Holt (CHSH) Bell inequality [23], that detect the so-called “Bell-nonlocal” (or “nonlocal”) states are also EWs.

In [24], the authors introduced a scenario where, given an initial bipartite entangled state, a single observer, Alice, owns one subsystem while the other subsystem is passed among multiple sequential and independent observers, Bobs. The task is to obtain a sequential violation of the CHSH Bell inequality. Initially, an unbounded number of recycled nonlocal correlations was detected only for “biased” measurement strategies [24]. Later the same was attained for a certain family of nonlocal states using a measurement strategy [25] that is unbiased. However, the question whether the same holds for all nonlocal states remains open. Nonlocal states are certainly entangled, and thus this result also answers the question about arbitrary recyclability of entanglement for Bell-nonlocal states. Recently, it was also observed that it is possible to witness and recycle entanglement an arbitrary number of

times using a CHSH Bell-local entangled state [26]. For other theoretical works in the direction of recyclability of entangled and nonlocal correlations, see [27–40]. For experimental works, see [41–45].

In the present work, we consider the scenario in which both the subsystems of a bipartite quantum state are passed on to multiple pairs of independent observers [37–40]. See Fig. 1. It has been conjectured in Ref. [37], with analytical and numerical evidence, that recycling of nonlocal correlations, via violation of the CHSH Bell inequality, is impossible in this scenario. That is, not more than one pair of Alice and Bob can share a CHSH Bell nonlocal state. This is interesting, since a single Alice can share CHSH Bell nonlocal correlations with an arbitrary number of sequential and independent Bobs [25]. But allowing multiple Alices and Bobs seems to restrict such sharability. It is therefore interesting to ask whether entanglement correlations also has such limitations. Contrary to the case of Bell nonlocality, we show that entanglement can be detected and recycled in this scenario, an arbitrary number of times. Therefore, where only one pair could detect CHSH Bell nonlocality in shared two-qubit states, an arbitrary number of pairs can detect entanglement in at least specific instances of the same. We provide a measurement strategy using which, an unbounded number of detections of entanglement can be achieved for any pure entangled two-qubit state and for a class of mixed states. We also show that such a property can be unleashed by a class of states whose entanglement content vanishes in the limit of the number of observing pairs growing unboundedly.

II. ENTANGLEMENT WITNESSES AND UNSHARP MEASUREMENTS

Entanglement witnesses [1, 2, 16] use expectation values of hermitian operators which separate the set of separable states from some of the entangled states. This method of entanglement detection utilizes the Hahn-Banach theorem [46, 47], which says that corresponding to any element falling outside a closed and convex set of a normed linear space, there always exists a functional on that space which “separates” the element with the closed and convex set. An entanglement witness operator is thus an operator W , such that $\langle W \rangle_{\rho_s} \geq 0$, for all separable states, ρ_s , and there exist at least one entangled state, ρ_e , for which $\langle W \rangle_{\rho_e} < 0$, where $\langle W \rangle_{\rho}$ represents the expectation value of the hermitian operator, W , with respect to a state ρ . For example, entanglement of the bipartite state $|\psi^+\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$ can be detected by the hermitian operator $|\phi^-\rangle\langle\phi^-|^T$, where $|\phi^-\rangle = \frac{1}{\sqrt{2}}(|00\rangle - |11\rangle)$, T represents partial transposition of operators, i.e., transposition with respect to any of the local parties, and $|0\rangle$ and $|1\rangle$ represent the eigenstates of the Pauli operator σ_z with eigenvalues 1 and -1, respectively. The expectation values required for witnessing entangled states can also be computed locally, since hermitian operators acting on a joint Hilbert space can be decomposed in hermitian operators acting on the local Hilbert spaces of the joint Hilbert space. For example, an entanglement witness operator, W_{ψ^+} , corresponding to $|\psi^+\rangle$ is decomposed as [20]

$$W_{\psi^+} = \frac{1}{4} [\mathbb{I}_d + \sigma_z \otimes \sigma_z - \sigma_x \otimes \sigma_x - \sigma_y \otimes \sigma_y], \quad (1)$$

where \mathbb{I}_d represents identity operator acting on a d -dimensional Hilbert space and σ_x , σ_y , and σ_z are the Pauli matrices.

While the expectation value of W_{ψ^+} will validate the entanglement in $|\psi^+\rangle$, the projective measurements involved to evaluate the expectation value will destroy the entanglement present in the state. But it is possible to detect entanglement as well as preserve some amount of entanglement at the same time, if one performs an unsharp version of the required projective measurements [24, 25]. Assume an observable, P , acting on the space of qubits and whose expectation value has to be estimated in some qubit ρ . Let $\{P_0, P_1\}$ denotes the projective measurement such that $P_0 + P_1 = \mathbb{I}_2$ and $P = P_0 - P_1$. This measurement is enough to estimate the required expectation value. One can also define an unsharp version of this measurement for sharpness parameter, λ , as

$$\begin{aligned} E_0^\lambda &= \frac{1}{2} (\mathbb{I}_2 + \lambda P), \\ E_1^\lambda &= \frac{1}{2} (\mathbb{I}_2 - \lambda P), \end{aligned} \quad (2)$$

where $0 \leq \lambda \leq 1$. Note that for $\lambda = 1$, the positive operator-valued measure (POVM), $\{E_0^\lambda, E_1^\lambda\}$, reduces to




Figure 1. Sequentially witnessing entanglement by multiple independent pairs of observers. ρ_{AB_1} is the initial state shared between Alice₁ and Bob₁.

the projective measurement, $\{P_0, P_1\}$, whereas for $\lambda = 0$, measurement operators are identity. Therefore, disturbance in the state is the most when $\lambda = 1$ whereas $\lambda = 0$ leads to a trivial measurement, in the sense that the state is totally unaffected. Notice that the unsharp version of the measurement will lead to an expectation value of P multiplied by the sharpness parameter, since $E_0^\lambda - E_1^\lambda = \lambda P$. The post-measurement state is given by the von Neumann-Lüder's rule [48] as

$$\rho \rightarrow \sqrt{E_0^\lambda} \rho \sqrt{E_0^\lambda} + \sqrt{E_1^\lambda} \rho \sqrt{E_1^\lambda}. \quad (3)$$

III. SCENARIO

We consider a generalized version of the sequential scenario presented in [24]. It involves a bipartite entangled state that is shared between a pair of observers, namely the 1st Alice-Bob pair (AB_1). The pair performs their tasks in spatially separated labs. To witness entanglement, the first Alice (A_1) and first Bob (B_1) each performs one of three local measurements at random on their qubit. Both the post-measurement subsystems are then passed to A_2 and B_2 who run the same task on their respective qubits independently and with no prior knowledge of the outcomes attained by their previous observers. This process continues until the state reaches to a pair who fails to detect entanglement. The ultimate aim of this task is to maximize the number of pairs that can witness entanglement independently. We will prove that using the weak measurement strategy presented below, one can achieve an arbitrarily long sequence of Alice-Bob pairs who can successfully detect entanglement for any pure entangled states, a class of mixed entangled states, and weakly entangled states.

Adopted measurement strategy and entanglement witness

Let the k^{th} Alice-Bob (AB_k) pair share a state ρ_{AB_k} and the witness operator used by the pair is given as

follows,

$$W_k = \frac{1}{4} [\mathbb{I}_4 + \sigma_z \otimes \sigma_z - \lambda_k \sigma_x \otimes \lambda_k \sigma_x - \lambda_k \sigma_y \otimes \lambda_k \sigma_y], \quad (4)$$

where λ_k is the sharpness parameter. Therefore, each observer has three measurement settings, i.e., $\{\sigma_x, \sigma_y, \sigma_z\}$. For simplicity, we will denote Pauli matrices, $\{\sigma_x, \sigma_y, \sigma_z\}$ as $\{\sigma_1, \sigma_2, \sigma_3\}$ and sharpness parameter, corresponding to any k^{th} local observer and the setting σ_i as $\lambda_k^{(i)}$, where $i = 1, 2, 3$. One can see from Eq. (4) that

$$\lambda_k^{(3)} = 1 \quad \text{and} \quad \lambda_k^{(1)} = \lambda_k^{(2)} = \lambda_k.$$

$$\begin{aligned} \rho_{AB_k} &= \frac{1}{9} \sum_{i,j=1}^3 \sum_{a,b=0}^1 \sqrt{\mathcal{A}_a^{\lambda_k^{(i)}}} \otimes \sqrt{\mathcal{B}_b^{\lambda_k^{(j)}}} \rho_{AB_{k-1}} \sqrt{\mathcal{A}_a^{\lambda_k^{(i)}}} \otimes \sqrt{\mathcal{B}_b^{\lambda_k^{(j)}}} \\ &= \frac{1}{36} \sum_{i,j=1}^3 \left[(1 + \Lambda_{k-1}^i) (1 + \Lambda_{k-1}^j) \rho_{AB_{k-1}} + (1 + \Lambda_{k-1}^i) (1 - \Lambda_{k-1}^j) \mathbb{I}_2 \otimes \sigma_j \rho_{AB_{k-1}} \mathbb{I}_2 \otimes \sigma_j \right. \\ &\quad \left. + (1 - \Lambda_{k-1}^i) (1 + \Lambda_{k-1}^j) \sigma_i \otimes \mathbb{I}_2 \rho_{AB_{k-1}} \sigma_i \otimes \mathbb{I}_2 + (1 - \Lambda_{k-1}^i) (1 - \Lambda_{k-1}^j) \sigma_i \otimes \sigma_j \rho_{AB_{k-1}} \sigma_i \otimes \sigma_j \right], \end{aligned} \quad (5)$$

where $\Lambda_k^i = \sqrt{1 - \lambda_k^{(i)2}}$, therefore $\Lambda_k^i = \sqrt{1 - \lambda_k^2}$ for $i = 1, 2$ and will be denoted as Λ_k . Now, the expectation values of each term present in the witness operator W_k with respect to ρ_{AB_k} can be expressed in terms of their expectation values with respect to state ρ_{AB_1} , i.e.,

$$\begin{aligned} \text{Tr}[\sigma_z \otimes \sigma_z \rho_{AB_k}] &= \text{Tr}[\sigma_z \otimes \sigma_z \rho_{AB_1}] \Pi_{l=1}^{k-1} \frac{(1 + 2\Lambda_l)^2}{9}, \\ \text{Tr}[\sigma_x \otimes \sigma_x \rho_{AB_k}] &= \text{Tr}[\sigma_x \otimes \sigma_x \rho_{AB_1}] \Pi_{l=1}^{k-1} \frac{(1 + \Lambda_l)^2}{9}. \end{aligned} \quad (6)$$

Note that $\text{Tr}[\sigma_x \otimes \sigma_x \rho_{AB_k}] = \text{Tr}[\sigma_y \otimes \sigma_y \rho_{AB_k}]$ for each $k \in \mathbb{N}$.

IV. SEQUENTIAL ENTANGLEMENT DETECTION ARBITRARILY MANY TIMES

In this section we study the considered scenario with the first pair of observers in the sequence, sharing classes of state.

A. Maximally entangled state

Let the pair AB_1 shares a maximally entangled state, i.e., $\rho_{AB_1} = |\psi^+\rangle\langle\psi^+|$. Therefore, the first pair will observe entanglement if $\text{Tr}[\rho_{AB_1} W_1] < 0$,

$$\implies \lambda_1^2 > 0. \quad (7)$$

Note that $\langle W_k \rangle \geq 0$ for all separable states as $0 \leq \lambda_k^{(i)} \leq 1$ [26]. Let $\{\mathcal{A}_0^{\lambda_k^{(i)}}, \mathcal{A}_1^{\lambda_k^{(i)}}\}$ and $\{\mathcal{B}_0^{\lambda_k^{(i)}}, \mathcal{B}_1^{\lambda_k^{(i)}}\}$ denote the measurements performed by the observers A_k and B_k respectively and their forms are taken as given in Eq. (2). Since it is assumed that all the measurement settings applied by each Alice-Bob pair is equally probable, i.e., unbiased, the state at the hand of the k^{th} pair in sequence is given by

And the pair AB_k will be able to witness the entanglement if $\text{Tr}[\rho_{AB_k} W_k] < 0$,

$$\implies \lambda_k^2 > \frac{1 - \Pi_{l=1}^{k-1} \frac{(1+2\Lambda_l)^2}{9}}{2\Pi_{l=1}^{k-1} \frac{(1+\Lambda_l)^2}{9}}. \quad (8)$$

Therefore, let us define the sequence, λ_k^2 , for $k \in \mathbb{N}$ as follows,

$$\lambda_k^2 = \begin{cases} (1 + \epsilon) \frac{1 - \Pi_{l=1}^{k-1} \frac{(1+2\Lambda_l)^2}{9}}{2\Pi_{l=1}^{k-1} \frac{(1+\Lambda_l)^2}{9}} & \text{if } \lambda_{k-1}^2 \in (0, 1) \\ \infty, & \text{otherwise,} \end{cases} \quad (9)$$

with $0 < \lambda_1^2 < 1$ and where $\epsilon > 0$. Note that $\lambda_k^2 \in (0, 1)$ imply that the pair AB_k will be able to witness the entanglement, whereas $\lambda_k^2 = \infty$ implies that the pair AB_k will not be able to witness entanglement. Now for $\lambda_k^2 \in (0, 1)$,

$$\frac{\lambda_{k+1}^2}{\lambda_k^2} = \frac{9}{(1 + \Lambda_k)^2} \frac{1 - \Pi_{l=1}^k \frac{(1+2\Lambda_l)^2}{9}}{1 - \Pi_{l=1}^{k-1} \frac{(1+2\Lambda_l)^2}{9}} > 1, \quad (10)$$

since $\Lambda_k = \sqrt{1 - \lambda_k^2} \in (0, 1)$. Therefore, the sequence λ_k^2 in Eq. (9) is positive and increasing sequence. The next important observation about this sequence is that as $\lambda_1^2 \rightarrow 0$ implies $\lambda_k^2 \rightarrow 0$ for all $k \in \mathbb{N}$. Thus, it is proved that arbitrary sequence of Alice-Bob pair will be able to witness starting from a maximally entangled state.

Interestingly, one can check that the state received by each pair during the process is a CHSH Bell-nonlocal

state when the initial state is maximally entangled. Therefore, an arbitrary number of pairs can share Bell nonlocality in the process of witnessing entanglement. However, the measurement settings required to witness such Bell nonlocality is different from the measurement settings that are required to witness the entanglement present in the state. Thus, this observation definitely does not overturn the conjecture made in [37].

B. Pure entangled states and a class of mixed entangled states

Any pure entangled state, upto a local unitary, can be written as $|\psi_\alpha\rangle = \sqrt{\alpha}|01\rangle + \sqrt{1-\alpha}|10\rangle$ for $\alpha \in (0, \frac{1}{2}]$, and in Hilbert-Schmidt decomposition can be expressed as

$$\frac{1}{4} \left[\mathbb{I}_4 - \sigma_z \otimes \sigma_z + 2\sqrt{\alpha(1-\alpha)}\sigma_x \otimes \sigma_x + 2\sqrt{\alpha(1-\alpha)}\sigma_y \otimes \sigma_y \right]. \quad (11)$$

Now consider a mixed entangled state shared by AB_1 ,

$$\rho_{AB_1} = p_1|\psi_\alpha\rangle\langle\psi_\alpha| + p_2|01\rangle\langle 01| + p_3|10\rangle\langle 10|, \quad (12)$$

where $p_1 > 0$, $p_2, p_3 \geq 0$ and $p_1 + p_2 + p_3 = 1$. In this case, using the same measurement strategy and witness operator, the sequence of sharpness parameters for the pair of observers will come out to be

$$\lambda_k^2 = \begin{cases} (1+\epsilon) \frac{1 - \prod_{l=1}^{k-1} \frac{(1+2\Lambda_l)^2}{9}}{4p_1 \sqrt{\alpha(1-\alpha)} \prod_{l=1}^{k-1} \frac{(1+\Lambda_l)^2}{9}} & \text{if } \lambda_{k-1}^2 \in (0, 1) \\ \infty, & \text{otherwise,} \end{cases} \quad (13)$$

with $0 < \lambda_1^2 < 1$ and where $\epsilon > 0$. This sequence of sharpness parameters is also positive, increasing and goes to zero as $\lambda_1^2 \rightarrow 0$. Therefore, an unbounded number of entanglement detection is possible by sequential Alice-Bob pairs where the first sequential pair may share all pure entangled states or particular class of mixed entangled states. Note that the unbounded detection of these entangled states can be shown, in similar fashion, even for the scenario where single Alice shares state with multiple Bobs, acting sequentially [24]. But since, it is already known that these states can produce unbounded chain of nonlocal states [25], which in turn imply entanglement and hence a separate analysis is not needed.

V. ARBITRARY SEQUENCE OF OBSERVERS WITH ENTANGLEMENT OF INITIAL STATE TENDING TO ZERO

In the scenario where multiple Bobs could witness and recycle entanglement with a single Alice, an arbitrary number of entanglement detection is possible even with entanglement tending to zero [26]. While the lesser

entanglement imply less amount of resource in various quantum information tasks, nevertheless, it still is a resource and gives quantum advantage in some tasks [49–51]. Therefore it is also important to investigate whether recycling of states is possible only with highly entangled states or even a weakly entangled states can allow such feat. In this section, we investigate the scenario of multiple pair of observers, detecting and recycling entangled states with the initial pair sharing the state,

$$\rho_{AB_1} = \frac{1}{4} [\mathbb{I}_2 \otimes \mathbb{I}_2 - \cos \theta \sigma_z \otimes \sigma_z + \alpha \sin \theta \sigma_x \otimes \sigma_x + \alpha \sin \theta \sigma_y \otimes \sigma_y], \quad (14)$$

where $\theta \in (0, \frac{\pi}{4}]$ and $\frac{1-\cos \theta}{2 \sin \theta} < \alpha \leq 1$. The k^{th} observer pair, AB_k , uses the same witness operator as given in Eq. (4). Therefore, the sequence of the sharpness parameter after applying the condition, $\text{Tr}[\rho_{AB_k} W_k] < 0$, appears as follows:

$$\lambda_k^2 = \begin{cases} (1+\epsilon) \frac{1 - \cos \theta \prod_{l=1}^{k-1} \frac{(1+2\Lambda_l)^2}{9}}{2\alpha \sin \theta \prod_{l=1}^{k-1} \frac{(1+\Lambda_l)^2}{9}}, & \text{if } \lambda_{k-1}^2 \in (0, 1) \\ \infty, & \text{otherwise,} \end{cases} \quad (15)$$

with $\lambda_1^2 > \frac{1-\cos \theta}{2\alpha \sin \theta}$ and where $\epsilon > 0$. We define another sequence for $\theta \in (0, \frac{\pi}{4}]$, which upper bounds the sequence in Eq. (15):

$$\gamma_k^2 = \begin{cases} (1+\epsilon) \frac{1 - \left(1 - \frac{\theta^2}{2}\right) \prod_{l=1}^{k-1} \left(1 - \frac{2\gamma_k^2}{3}\right)^2}{\theta \prod_{l=1}^{k-1} \left(\frac{2-\gamma_k^2}{3}\right)^2}, & \text{if } \gamma_{k-1}^2 \in (0, 1) \\ \infty, & \text{otherwise,} \end{cases} \quad (16)$$

with $\gamma_1^2 = (1+\epsilon) \frac{\theta}{2\alpha} > \lambda_1^2$. It is easy to show that $\gamma_k^2 > \lambda_k^2$ for all k such that λ_k^2 is finite, since $\sqrt{1-x^2} > 1-x^2$ for $0 < x < 1$, $\cos \theta > \left(1 - \frac{\theta^2}{2}\right)$, and $\sin \theta > \frac{\theta}{2}$ for $\theta \in (0, \frac{\pi}{4}]$.

Again, similar to the sequences given in Eqs. (9) and (13), this sequence, γ_k^2 , is also strictly positive and increasing. Also, for $\theta \rightarrow 0$ and for $k \in \mathbb{N}$, $\gamma_k^2 \rightarrow 0$. This implies that $\lambda_k^2 \rightarrow 0$ as $\theta \rightarrow 0$. Therefore, detection and recycling of entanglement is possible when $\theta \rightarrow 0$, an arbitrary number of times. It is important to notice that as $\theta \rightarrow 0$, entanglement of the state ρ_{AB_1} , given in Eq. (14), tends to zero. Surprisingly therefore, an arbitrary number of pairs of observers is possible even in a situation when the initial state has an entanglement that becomes vanishingly small with the number of pairs growing unboundedly.

VI. CONCLUSION

It is found that an arbitrary number of pairs of sequential and independent observers can witness entanglement, acting locally on two-qubit entangled states. This result

is potentially of importance, given that it has been conjectured that not more than a single pair of observers can detect CHSH Bell-nonlocal correlations in the same scenario [37–39]. Assuming that the conjecture in the literature is true, our finding demonstrates a clear distinction between “entanglement nonlocality” and “Bell nonlocality”, with respect to sequential witnessing by pairs of observers. It is to be noted that in the case when a single observer share a bipartite quantum state with multiple sequential observers at the other end, entanglement nonlocality and Bell nonlocality behave similarly. There, it is possible to have an arbitrary long sequence for both entangled and Bell-nonlocal correlations [25, 26].

Here, we identified classes of quantum states which can produce arbitrarily long sequences of entangled correlations. Specifically, all pure entangled states and a certain class of mixed entangled states are helpful in complet-

ing the task. It is also shown that one can succeed in such tasks even with the entanglement of initial shared state tending to zero. But whether other mixed entangled states can also produce such arbitrarily long chains remains an open question.

ACKNOWLEDGMENTS

The research of CS was supported in part by the INFOSYS scholarship. MP acknowledges the NCN (Poland) grant (grant number 2017/26/E/ST2/01008). The authors from Harish-Chandra Research Institute acknowledge partial support from the Department of Science and Technology, Government of India through the QuEST grant (grant number DST/ICPS/QUST/Theme-3/2019/120).

-
- [1] Ryszard Horodecki, Paweł Horodecki, Michał Horodecki, and Karol Horodecki, “Quantum entanglement,” *Rev. Mod. Phys.* **81**, 865–942 (2009).
 - [2] Otfried Gühne and Géza Tóth, “Entanglement detection,” *Physics Reports* **474**, 1–75 (2009).
 - [3] Artur K. Ekert, “Quantum cryptography based on bell’s theorem,” *Phys. Rev. Lett.* **67**, 661–663 (1991).
 - [4] Charles H Bennett and Gilles Brassard, “Proceedings of the ieee international conference on computers, systems and signal processing,” (1984).
 - [5] Charles H. Bennett and Stephen J. Wiesner, “Communication via one- and two-particle operators on einstein-podolsky-rosen states,” *Phys. Rev. Lett.* **69**, 2881–2884 (1992).
 - [6] Charles H. Bennett, Gilles Brassard, Claude Crépeau, Richard Jozsa, Asher Peres, and William K. Wootters, “Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels,” *Phys. Rev. Lett.* **70**, 1895–1899 (1993).
 - [7] M. Żukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, ““event-ready-detectors” bell experiment via entanglement swapping,” *Phys. Rev. Lett.* **71**, 4287–4290 (1993).
 - [8] S. Bose, V. Vedral, and P. L. Knight, “Multiparticle generalization of entanglement swapping,” *Phys. Rev. A* **57**, 822–829 (1998).
 - [9] D. Mayers and A. Yao, “Quantum cryptography with imperfect apparatus,” Proceedings 39th Annual Symposium on Foundations of Computer Science (Cat. No.98CB36280) , 503–509 (1998).
 - [10] Jonathan Barrett, Lucien Hardy, and Adrian Kent, “No signaling and quantum key distribution,” *Phys. Rev. Lett.* **95**, 010503 (2005).
 - [11] Antonio Acín, Nicolas Brunner, Nicolas Gisin, Serge Massar, Stefano Pironio, and Valerio Scarani, “Device-independent security of quantum cryptography against collective attacks,” *Phys. Rev. Lett.* **98**, 230501 (2007).
 - [12] Roger Colbeck and Renato Renner, “Free randomness can be amplified,” *Nature Physics* **8**, 450–453 (2012).
 - [13] Roger Colbeck, “Quantum and relativistic protocols for secure multi-party computation,” (2011), [arXiv:0911.3814 \[quant-ph\]](https://arxiv.org/abs/0911.3814).
 - [14] Roger Colbeck and Adrian Kent, “Private randomness expansion with untrusted devices,” *Journal of Physics A: Mathematical and Theoretical* **44**, 095305 (2011).
 - [15] S. Pironio, A. Acín, S. Massar, A. Boyer de la Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning, and C. Monroe, “Random numbers certified by bell’s theorem,” *Nature* **464**, 1021–1024 (2010).
 - [16] Dariusz Chruściński and Gniewomir Sarbicki, “Entanglement witnesses: construction, analysis and classification,” **47**, 483001 (2014).
 - [17] Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki, “Separability of mixed states: necessary and sufficient conditions,” *Physics Letters A* **223**, 1–8 (1996).
 - [18] Barbara M. Terhal, “Bell inequalities and the separability criterion,” *Physics Letters A* **271**, 319–326 (2000).
 - [19] Dagmar Bruß, J. Ignacio Cirac, Paweł Horodecki, Florian Hulpke, Barbara Kraus, Maciej Lewenstein, and Anna Sanpera, “Reflections upon separability and distillability,” *Journal of Modern Optics* **49**, 1399–1418 (2002), <https://doi.org/10.1080/09500340110105975>.
 - [20] O. Gühne, P. Hyllus, D. Bruss, A. Ekert, M. Lewenstein, C. Macchiavello, and A. Sanpera, “Experimental detection of entanglement via witness operators and local measurements,” *Journal of Modern Optics* **50**, 1079–1102 (2003).
 - [21] J. S. Bell, “On the Einstein Podolsky Rosen paradox,” *Physics Physique Fizika* **1**, 195–200 (1964).
 - [22] Nicolas Brunner, Daniel Cavalcanti, Stefano Pironio, Valerio Scarani, and Stephanie Wehner, “Bell nonlocality,” *Rev. Mod. Phys.* **86**, 419–478 (2014).
 - [23] John F. Clauser, Michael A. Horne, Abner Shimony, and Richard A. Holt, “Proposed experiment to test local hidden-variable theories,” *Phys. Rev. Lett.* **23**, 880–884 (1969).
 - [24] Ralph Silva, Nicolas Gisin, Yelena Guryanova, and Sandu Popescu, “Multiple observers can share the nonlocality of half of an entangled pair by using optimal weak measurements,” *Phys. Rev. Lett.* **114**, 250401 (2015).

- [25] Peter J. Brown and Roger Colbeck, “Arbitrarily many independent observers can share the nonlocality of a single maximally entangled qubit pair,” *Phys. Rev. Lett.* **125**, 090401 (2020).
- [26] Chirag Srivastava, Mahasweta Pandit, and Ujjwal Sen, “Entanglement witnessing by arbitrarily many independent observers recycling a local quantum shared state,” (2021), [arXiv:2109.10310 \[quant-ph\]](#).
- [27] Shiladitya Mal, Archan S. Majumdar, and Dipankar Home, “Sharing of nonlocality of a single member of an entangled pair of qubits is not possible by more than two unbiased observers on the other wing,” *Mathematics* **4** (2016), 10.3390/math4030048.
- [28] Anindita Bera, Shiladitya Mal, Aditi Sen(De), and Ujjwal Sen, “Witnessing bipartite entanglement sequentially by multiple observers,” *Phys. Rev. A* **98**, 062304 (2018).
- [29] Debarshi Das, Arkaprabha Ghosal, Souradeep Sasmal, Shiladitya Mal, and A. S. Majumdar, “Facets of bipartite nonlocality sharing by multiple observers via sequential measurements,” *Phys. Rev. A* **99**, 022305 (2019).
- [30] Sutapa Saha, Debarshi Das, Souradeep Sasmal, Debasis Sarkar, Kaushiki Mukherjee, Arup Roy, and Some Sankar Bhattacharya, “Sharing of tripartite nonlocality by multiple observers measuring sequentially at one side,” *Quantum Information Processing* **18**, 42 (2019).
- [31] Ananda G. Maity, Debarshi Das, Arkaprabha Ghosal, Arup Roy, and A. S. Majumdar, “Detection of genuine tripartite entanglement by multiple sequential observers,” *Phys. Rev. A* **101**, 042340 (2020).
- [32] Chirag Srivastava, Shiladitya Mal, Aditi Sen(De), and Ujjwal Sen, “Sequential measurement-device-independent entanglement detection by multiple observers,” *Phys. Rev. A* **103**, 032408 (2021).
- [33] Saptarshi Roy, Asmita Kumari, Shiladitya Mal, and Aditi Sen De, “Robustness of higher dimensional nonlocality against dual noise and sequential measurements,” (2020), [arXiv:2012.12200 \[quant-ph\]](#).
- [34] Adán Cabello, “Bell nonlocality between sequential pairs of observers,” (2021), [arXiv:2103.11844 \[quant-ph\]](#).
- [35] Changliang Ren, Xiaowei Liu, Wenlin Hou, Tianfeng Feng, and Xiaoqi Zhou, “Non-locality sharing for a three-qubit system via multilateral sequential measurements,” (2021), [arXiv:2105.03709 \[quant-ph\]](#).
- [36] Tinggui Zhang and Shao-Ming Fei, “Sharing quantum nonlocality and genuine nonlocality with independent observables,” *Phys. Rev. A* **103**, 032216 (2021).
- [37] Shuming Cheng, Lijun Liu, Travis J. Baker, and Michael J. W. Hall, “Limitations on sharing bell nonlocality between sequential pairs of observers,” *Phys. Rev. A* **104**, L060201 (2021).
- [38] Shuming Cheng, Lijun Liu, Travis J. Baker, and Michael J. W. Hall, “Recycling qubits for the generation of Bell nonlocality between independent sequential observers,” (2021), [arXiv:2109.03472 \[quant-ph\]](#).
- [39] Jie Zhu, Meng-Jun Hu, Guang-Can Guo, Chuan-Feng Li, and Yong-Sheng Zhang, “Einstein-podolsky-rosen steering in two-sided sequential measurements with one entangled pair,” (2021), [arXiv:2102.02550 \[quant-ph\]](#).
- [40] Arun Kumar Das, Debarshi Das, Shiladitya Mal, Dipankar Home, and A. S. Majumdar, “Resource theoretic efficacy of the single copy of a two-qubit entangled state in a sequential network,” (2021), [arXiv:2109.11433 \[quant-ph\]](#).
- [41] Matteo Schiavon, Luca Calderaro, Mirko Pittaluga, Giuseppe Vallone, and Paolo Villoresi, “Three-observer bell inequality violation on a two-qubit entangled state,” *Quantum Science and Technology* **2**, 015010 (2017).
- [42] Meng-Jun Hu, Zhi-Yuan Zhou, Xiao-Min Hu, Chuan-Feng Li, Guang-Can Guo, and Yong-Sheng Zhang, “Observation of non-locality sharing among three observers with one entangled pair via optimal weak measurement,” *npj Quantum Information* **4**, 63 (2018).
- [43] Giulio Fioletto, Luca Calderaro, Armin Tavakoli, Matteo Schiavon, Francesco Picciariello, Adán Cabello, Paolo Villoresi, and Giuseppe Vallone, “Experimental certification of sustained entanglement and nonlocality after sequential measurements,” *Phys. Rev. Applied* **13**, 044008 (2020).
- [44] Yeon-Ho Choi, Seongjin Hong, Tanumoy Pramanik, Hyang-Tag Lim, Yong-Su Kim, Hojoong Jung, Sang-Wook Han, Sung Moon, and Young-Wook Cho, “Demonstration of simultaneous quantum steering by multiple observers via sequential weak measurements,” *Optica* **7**, 675–679 (2020).
- [45] Tianfeng Feng, Changliang Ren, Yuling Tian, Maolin Luo, Haofei Shi, Jingling Chen, and Xiaoqi Zhou, “Observation of nonlocality sharing via not-so-weak measurements,” *Phys. Rev. A* **102**, 032220 (2020).
- [46] G. F. Simmons, *Introduction to Topology and Modern Analysis* (McGraw-Hill, 1963).
- [47] P. D. Lax, *Functional Analysis* (Wiley-Interscience, 2002).
- [48] Paul Busch, Pekka J. Lahti, and Peter Mittelstaedt, *The Quantum Theory of Measurement*, 2nd ed. (Springer, Berlin, Heidelberg, 1996).
- [49] Charles H. Bennett, Gilles Brassard, Claude Crépeau, Richard Jozsa, Asher Peres, and William K. Wootters, “Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels,” *Phys. Rev. Lett.* **70**, 1895–1899 (1993).
- [50] Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki, “General teleportation channel, singlet fraction, and quasidistillation,” *Phys. Rev. A* **60**, 1888–1898 (1999).
- [51] Michael A Nielsen, “A simple formula for the average gate fidelity of a quantum dynamical operation,” *Physics Letters A* **303**, 249–252 (2002).