

Completeness of derived interleaving distances and sheaf quantization of non-smooth objects*

Tomohiro Asano Yuichi Ike

April 20, 2022

Abstract

We investigate sheaf-theoretic methods to deal with non-smooth objects in symplectic geometry. We show the completeness of a derived category of sheaves with respect to the interleaving distance and construct a sheaf quantization of a Hamiltonian homeomorphism. We also develop Lusternik–Schnirelmann theory in the microlocal theory of sheaves. With these new tools, we prove an Arnold-type theorem for the image of the zero-section under a Hamiltonian homeomorphism by a purely sheaf-theoretic method.

1 Introduction

The microlocal theory of sheaves due to Kashiwara and Schapira [KS90] has been effectively applied to symplectic and contact geometry after the pioneering work by Nadler–Zaslow [NZ09, Nad09] and Tamarkin [Tam18], both of which appeared on arXiv in 2008. In this paper, we apply the theory to symplectic geometry for non-smooth objects, in particular, limits of smooth objects. For this purpose, we prove the completeness of the derived category of sheaves with respect to the interleaving distance, which was introduced by Kashiwara–Schapira [KS18a] and later generalized by Petit–Schapira [PS20].

1.1 Our results

In this work, we use the Tamarkin category, which was introduced by Tamarkin [Tam18] to prove his non-displaceability theorem. Let M be a connected C^∞ -manifold without boundary and let $\pi: T^*M \rightarrow M$ denote its cotangent bundle equipped with the canonical symplectic structure. We also let $(t; \tau)$ denote the homogeneous coordinate system on $T^*\mathbb{R}_t$ and fix a field \mathbf{k} . The Tamarkin category $\mathcal{D}(M)$ is defined to be the left orthogonal of the triangulated subcategory consisting of objects microsupported in $\{\tau \leq 0\} \subset T^*(M \times \mathbb{R}_t)$ in $\mathbf{D}(\mathbf{k}_{M \times \mathbb{R}_t})$, the derived category of sheaves on $M \times \mathbb{R}_t$. To generalize Tamarkin’s non-displaceability theorem to a quantitative version, the authors introduced the interleaving-like pseudo-distance $d_{\mathcal{D}(M)}$ on $\mathcal{D}(M)$ in [AI20], motivated by the convolution distance on $\mathbf{D}(\mathbf{k}_{\mathbb{R}^n})$ due to Kashiwara–Schapira [KS18a]. See Subsection 3.2 for the details of Tamarkin category and the distance $d_{\mathcal{D}(M)}$.

In this paper, we first establish the completeness of the Tamarkin category $\mathcal{D}(M)$ with respect to $d_{\mathcal{D}(M)}$ in Section 4. In fact, we prove the completeness result for a wider

*2020 Mathematics Subject Classification: 37J12, 55N31, 18G80, 35A27

Keywords: interleaving distance, C^0 -symplectic geometry, microlocal theory of sheaves

class of distances associated with thickening kernels, which was introduced by Petit–Schapira [PS20]. For a topological space X , a thickening kernel on X is a monoidal presheaf \mathfrak{K} on $(\mathbb{R}_{\geq 0}, +)$ with values in $\mathbf{D}(\mathbf{k}_{X \times X})$. Given a thickening kernel \mathfrak{K} on X , one can define the interleaving distance $d_{\mathfrak{K}}(F, G)$ between two objects $F, G \in \mathbf{D}(\mathbf{k}_X)$. See Subsection 3.1 for details. The pseudo-distance $d_{\mathcal{D}(M)}$ can be regarded as a special case of this interleaving distance $d_{\mathfrak{K}}$. The first result in this paper is the following.

Theorem 1.1 (see Corollary 4.5). *The derived category of sheaves $\mathbf{D}(\mathbf{k}_X)$ is complete with respect to the pseudo-distance $d_{\mathfrak{K}}$. In particular, the Tamarkin category $\mathcal{D}(X)$ is complete with respect to the pseudo-distance $d_{\mathcal{D}(X)}$.*

Armed with the completeness result, we consider sheaf quantization of Hamiltonian homeomorphisms in Section 5. Let I be an open interval containing the closed interval $[0, 1]$. A homeomorphism $\varphi_{\infty}: T^*M \rightarrow T^*M$ is called a *Hamiltonian homeomorphism* if there exists a sequence of compactly supported Hamiltonian functions $(H_n: T^*M \times I \rightarrow \mathbb{R})_{n \in \mathbb{N}}$ that converges to some continuous function $H: T^*M \times I \rightarrow \mathbb{R}$ in C^0 -sense and the sequence of the time-1 maps of the associated Hamiltonian flows $(\phi_1^{H_n})_{n \in \mathbb{N}} \subset \text{Ham}_c(T^*M)$ converges to φ_{∞} in C^0 -sense (see Definition 5.16 for the precise definition). With the Hamiltonian isotopy $\phi^{H_n}: T^*M \times I \rightarrow T^*M$, one can associate a canonical object $K^{H_n} \in \mathbf{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t \times I})$, called the sheaf quantization or the GKS kernel [GKS12]. The convergence of $(H_n)_{n \in \mathbb{N}}$ and the energy estimate obtained in [AI20] show that the sequence $(K^{H_n}|_{M^2 \times \mathbb{R}_t \times \{1\}})_{n \in \mathbb{N}}$ is a Cauchy sequence with respect to the interleaving distance. Hence, applying Theorem 1.1, we obtain a limit object $K_{\infty} \in \mathbf{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$. We also prove that K_{∞} is uniquely determined by the continuous function H , which gives the notion of the sheaf quantization of Hamiltonian homeomorphism φ_{∞} . One of the advantages of this approach is that we may directly use the limit object without explicitly dealing with a limit of some sequence to study C^0 -symplectic geometry. Remark that we can actually associate a sheaf with an element of the completion of $\text{Ham}_c(T^*M)$ with respect to the Hofer metric (Definition 5.15).

In the course of the construction, we also investigate sheaf quantization of compactly supported Hamiltonian isotopies. For the sheaf quantization K of a compactly supported Hamiltonian isotopy $\phi = (\phi_s)_{s \in I}$, we prove that the restriction $K|_{M^2 \times \mathbb{R}_t \times \{1\}}$ depends only on the time-1 map of ϕ_1 . This allows us to define the sheaf quantization $K^{\varphi} \in \mathbf{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ of a Hamiltonian diffeomorphism φ . We also generalize our previous energy estimate in terms of the distance between K^{φ} and $K^{\varphi'}$ for Hamiltonian diffeomorphisms φ and φ' .

Next, we develop Lusternik–Schnirelmann theory in the microlocal theory of sheaves. More precisely, for an object $F \in \mathcal{D}(M)$, we define the set of spectral invariants $\text{Spec}(F) \subset \mathbb{R}$ and prove the following. Here, $\text{cl}(M)$ denotes the cup-length of M over \mathbf{k} .

Theorem 1.2 (see Theorem 6.4 for a more precise statement). *Let $t: M \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ be the projection. Let $F \in \mathcal{D}(M)$ and assume that*

- (i) *t is proper on $\text{Supp}(F)$,*
- (ii) *$\text{R}\Gamma_{M \times [-c, \infty)}(M \times \mathbb{R}; F) \simeq \text{R}\Gamma(M; \mathbf{k}_M)$ and $F|_{M \times (c, \infty)}$ is locally constant for $c \gg 0$.*

*If $\#\text{Spec}(F) \leq \text{cl}(M)$, then $\pi(\text{SS}(F) \cap \Gamma_{dt})$ is cohomologically non-trivial, where $\Gamma_{dt} \subset T^*M$ is the graph of the 1-form dt . That is, for any open neighborhood U of $\pi(\text{SS}(F) \cap \Gamma_{dt})$, the restriction map $\bigoplus_{n \geq 1} H^n(M; \mathbf{k}) \rightarrow \bigoplus_{n \geq 1} H^n(U; \mathbf{k})$ is non-zero.*

The theorem above was announced to appear in Humilière–Vichery [HV], which motivated our work.

As a byproduct, we give a purely sheaf-theoretic proof of the following Arnold-type theorem for non-smooth objects, which was originally proved by Buhovsky–Humilière–Seyfaddini [BHS19] in a more general setting. Assume that M is compact and let 0_M denote the zero-section of T^*M . Using the sheaf quantization of φ_∞ and Theorem 1.2, we obtain the following.

Theorem 1.3 (see Theorem 7.2). *Let φ_∞ be a Hamiltonian homeomorphism of T^*M . Then, if the number of spectral invariants of $\varphi_\infty(0_M)$ is smaller than $\text{cl}(M) + 1$, then $0_M \cap \varphi_\infty(0_M)$ is cohomologically non-trivial, hence it is infinite.*

Combining the argument above with the C^0 -continuity of the spectral norm, which is obtained in the field of symplectic geometry, we also prove a stronger result than Theorem 1.3, an Arnold principle for a C^0 -limit of Hamiltonian diffeomorphisms (see Proposition 7.3).

We also give a sheaf-theoretic proof of a Legendrian analog of Theorem 1.3. More precisely, we recover an Arnold-type theorem for Hausdorff limits of Legendrian submanifolds in a 1-jet bundle (cf. [BHS19, Thm. 1.5]).

1.2 Related work

The completeness of a persistence category was studied by Cruz [Cru19] and Scoccola [Sco20]. They showed that if the category admits any sequential (co)limit, then there is a limit object for any Cauchy sequence with respect to the interleaving distance. In this paper, we work with derived categories and need a different argument. Our completeness result (Theorem 1.1) also holds in a triangulated category endowed with a persistence structure. See also Biran–Cornea–Zhang [BCZ21] for persistence structures for triangulated categories. During the preparation of this paper, the authors learned from Stéphane Guillermou that he and Claude Viterbo had independently obtained a similar completeness result in a derived category of sheaves [GV22].

The persistence method has been widely applied to symplectic and contact geometry. After the pioneering work by Polterovich–Shelukhin [PS16], persistence modules have been used to study barcodes of Floer cohomology complexes in Usher–Zhang [UZ16] and the study of spectral norms in Kislev–Shelukhin [KS18b], to name a few. In this paper, we also investigate the relation between the interleaving-like distance $d_{\mathcal{D}(M)}$ and the spectral norm (see Proposition 6.8). Similar results are independently discovered by Guillermou and Viterbo [GV22].

Kashiwara–Schapira [KS18a] studied persistence modules from the point of view of the microlocal theory of sheaves. Motivated by the work, Asano–Ike [AI20] introduced the interleaving-like distance $d_{\mathcal{D}(M)}$ and showed that the distance between an object and its Hamiltonian deformation is upper bounded by the Hofer norm. The result was effectively used in Chiu [Chi21] and Li [Li21]. See also Zhang [Zha20] for the interleaving-like distance in the Tamarkin category.

Buhovsky–Humilière–Seyfaddini [BHS18] constructed a counterexample of Arnold’s conjecture for a Hamiltonian homeomorphism of a closed symplectic manifold M with $\dim M \geq 4$. However, one still obtains an Arnold-type theorem for a Hamiltonian homeomorphism if one reformulates the conjecture with the notion of spectral invariants, as in [BHS21, Kaw19, BHS19]. In these studies, the C^0 -continuity of persistence modules associated with Hamiltonian diffeomorphisms was effectively used (see also [RSV18]). Guillermou [Gui13] (see also [Gui19, Part 7]) applied the microlocal theory of sheaves to C^0 -symplectic geometry. He gave a purely sheaf-theoretic proof of the Gromov–Eliashberg theorem, using the involutivity of microsupports.

1.3 Organization

This paper is structured as follows. In Section 2, we recall some basics of the microlocal theory of sheaves. In Section 3, we first recall the interleaving distance for sheaves associated with a thickening kernel. We then briefly review the Tamarkin category $\mathcal{D}(M)$ and sheaf quantization of Hamiltonian isotopies. In Section 4, we prove Theorem 1.1, the completeness of the derived category of sheaves $\mathbf{D}(\mathbf{k}_X)$ with respect to $d_{\mathfrak{R}}$. In Section 5, we first show that the restriction of the sheaf quantization of a Hamiltonian isotopy to $s = 1$ depends only on the time-1 map, which allows us to define the sheaf quantization of a Hamiltonian diffeomorphism. We then construct a sheaf quantization of a Hamiltonian homeomorphism using the completeness result. In Section 6, we develop Lusternik–Schnirelmann theory for the Tamarkin category and prove Theorem 1.2. We prove Theorem 1.3 in Section 7 and its Legendrian analog in Section 8.

Acknowledgments

The authors would like to express their gratitude to Vincent Humilière for fruitful discussion on many parts of this paper. They also thank Stéphane Guillermou and Takahiro Saito for helpful discussion, and Yusuke Kawamoto for drawing their attention to applications of the microlocal theory of sheaves to C^0 -symplectic geometry. They are also grateful to Wenyuan Li, Tatsuki Kuwagaki, and Morimichi Kawasaki for their helpful comments. TA was supported by Innovative Areas Discrete Geometric Analysis for Materials Design (Grant No. 17H06461). YI was supported by JSPS KAKENHI (Grant No. 21K13801) and ACT-X, Japan Science and Technology Agency (Grant No. JPMJAX1903).

2 Microlocal theory of sheaves

We mainly follow the notation of [KS90]. Throughout this paper, let \mathbf{k} be a field. In this section, let X be a C^∞ -manifold without boundary.

2.1 Geometric notions

For a locally closed subset Z of X , we denote by \overline{Z} its closure and by $\text{Int}(Z)$ its interior. We also denote by $\delta_X: X \rightarrow X \times X, x \mapsto (x, x)$ the diagonal map and by $\Delta_X := \delta_X(X)$ the diagonal of $X \times X$. We often write Δ for Δ_X for simplicity. We denote by TX the tangent bundle and by T^*X the cotangent bundle of X . We write $\pi_X: T^*X \rightarrow X$ or simply π for the projection. For a submanifold M of X , we denote by T_M^*X the conormal bundle to M in X . In particular, T_X^*X denotes the zero-section of T^*X , which we often simply write 0_X . We set $\mathring{T}^*X := T^*X \setminus 0_X$.

With a morphism of manifolds $f: X \rightarrow Y$, we associate the following commutative diagram of morphisms of manifolds:

$$\begin{array}{ccccc}
 T^*X & \xleftarrow{f_d} & X \times_Y T^*Y & \xrightarrow{f_\pi} & T^*Y \\
 \pi_X \downarrow & & \downarrow & & \downarrow \pi_Y \\
 X & \xlongequal{\quad} & X & \xrightarrow{f} & Y,
 \end{array} \tag{2.1}$$

where f_π is the projection and f_d is induced by the transpose of the tangent map $f': TX \rightarrow X \times_Y TY$.

We denote by $(x; \xi)$ a local homogeneous coordinate system on T^*X . The cotangent bundle T^*X is an exact symplectic manifold with the Liouville 1-form $\alpha_{T^*X} = \langle \xi, dx \rangle$. Thus the symplectic form on T^*X is defined to be $\omega = d\alpha_{T^*X}$. We denote by $a: T^*X \rightarrow T^*X, (x; \xi) \mapsto (x; -\xi)$ the antipodal map. For a subset A of T^*X , A^a denotes its image under the antipodal map a . A subset A of T^*M is said to be *conic* if it is invariant under the action of $\mathbb{R}_{>0}$ on T^*M , that is, the scaling of the fibers.

2.2 Microsupports of sheaves

We write \mathbf{k}_X for the constant sheaf with stalk \mathbf{k} and $\text{Mod}(\mathbf{k}_X)$ for the abelian category of sheaves of \mathbf{k} -vector spaces on X . Moreover, we denote by $\text{D}(\mathbf{k}_X)$ the unbounded derived category of \mathbf{k} -vector spaces on X . Although all the results are stated for bounded derived categories in [KS90], we can apply most of them for unbounded categories, which we shall state in this subsection. We refer to [Spa88, KS06, RS18]. One can define Grothendieck's six operations $\text{RHom}, \otimes, \text{R}f_*, f^{-1}, \text{R}f_!, f^!$ for a continuous map $f: X \rightarrow Y$ with suitable conditions. For a locally closed subset Z of X , we denote by \mathbf{k}_Z the zero extension of the constant sheaf with stalk \mathbf{k} on Z to X , whose stalk is 0 on $X \setminus Z$. Moreover, for a locally closed subset Z of X and $F \in \text{D}(\mathbf{k}_X)$, we define $F_Z, \text{R}\Gamma_Z(F) \in \text{D}(\mathbf{k}_X)$ by

$$F_Z := F \otimes \mathbf{k}_Z, \quad \text{R}\Gamma_Z(F) := \text{RHom}(\mathbf{k}_Z, F). \quad (2.2)$$

Let us recall the definition of the *microsupport* $\text{SS}(F)$ of an object $F \in \text{D}(\mathbf{k}_X)$.

Definition 2.1 ([KS90, Def. 5.1.2]). Let $F \in \text{D}(\mathbf{k}_X)$ and $p \in T^*X$. One says that $p \notin \text{SS}(F)$ if there is a neighborhood U of p in T^*X such that for any $x_0 \in X$ and any C^∞ -function φ on X (defined on a neighborhood of x_0) satisfying $d\varphi(x_0) \in U$, one has $\text{R}\Gamma_{\{x \in X | \varphi(x) \geq \varphi(x_0)\}}(F)_{x_0} \simeq 0$. One also sets $\text{SS}(F) := \text{SS}(F) \cap T^*X$.

The following is called the microlocal Morse lemma.

Proposition 2.2 ([KS90, Prop. 5.4.17] and [RS18, Thm. 4.1]). *Let $F \in \text{D}(\mathbf{k}_X)$ and $\varphi: X \rightarrow \mathbb{R}$ be a C^∞ -function. Let moreover $a, b \in \mathbb{R}$ with $a < b$. Assume*

- (1) φ is proper on $\text{Supp}(F)$,
- (2) $d\varphi(x) \notin \text{SS}(F)$ for any $x \in \varphi^{-1}([a, b])$.

Then the canonical morphism

$$\text{R}\Gamma(\varphi^{-1}((-\infty, b)); F) \rightarrow \text{R}\Gamma(\varphi^{-1}((-\infty, a)); F) \quad (2.3)$$

is an isomorphism.

We consider the behavior of the microsupports with respect to functorial operations.

Proposition 2.3 ([KS90, Prop. 5.4.4, Prop. 5.4.13, and Prop. 5.4.5]). *Let $f: X \rightarrow Y$ be a morphism of manifolds, $F \in \text{D}(\mathbf{k}_X)$, and $G \in \text{D}(\mathbf{k}_Y)$.*

- (i) *Assume that f is proper on $\text{Supp}(F)$. Then $\text{SS}(\text{R}f_*F) \subset f_\pi f_d^{-1}(\text{SS}(F))$. Moreover, if f is a closed embedding, the inclusion is an equality.*
- (ii) *Assume that f is non-characteristic for $\text{SS}(G)$ (see [KS90, Def. 5.4.12] for the definition). Then $\text{SS}(f^{-1}G) \cup \text{SS}(f^!G) \subset f_d f_\pi^{-1}(\text{SS}(G))$.*

For closed conic subsets A and B of T^*X , let us denote by $A + B$ the fiberwise sum of A and B , that is,

$$A + B := \left\{ (x; a + b) \left| \begin{array}{l} x \in \pi(A) \cap \pi(B), \\ a \in A \cap \pi^{-1}(x), b \in B \cap \pi^{-1}(x) \end{array} \right. \right\} \subset T^*X. \quad (2.4)$$

Proposition 2.4 ([KS90, Prop. 5.4.14]). *Let $F, G \in \mathbf{D}(\mathbf{k}_X)$.*

- (i) *If $\mathrm{SS}(F) \cap \mathrm{SS}(G)^a \subset 0_X$, then $\mathrm{SS}(F \otimes G) \subset \mathrm{SS}(F) + \mathrm{SS}(G)$.*
- (ii) *If $\mathrm{SS}(F) \cap \mathrm{SS}(G) \subset 0_X$, then $\mathrm{SS}(\mathrm{R}\mathcal{H}om(F, G)) \subset \mathrm{SS}(F)^a + \mathrm{SS}(G)$.*

We need an estimate for the microsupport of a kind of limit object in $\mathbf{D}(\mathbf{k}_X)$. For that purpose, we use the following estimates.

Lemma 2.5 (cf. [KS90, Exe. V.7]). *Let I be an index set and $F_i \in \mathbf{D}(\mathbf{k}_X)$ for $i \in I$. Then, one has*

$$\mathrm{SS} \left(\prod_{i \in I} F_i \right), \mathrm{SS} \left(\bigoplus_{i \in I} F_i \right) \subset \overline{\bigcup_{i \in I} \mathrm{SS}(F_i)}. \quad (2.5)$$

2.3 Composition and convolution

We recall the operation called the composition of sheaves.

For $i = 1, 2, 3$, let X_i be a manifold. We write $X_{ij} := X_i \times X_j$ and $X_{123} := X_1 \times X_2 \times X_3$ for short. We denote by q_{ij} the projection $X_{123} \rightarrow X_{ij}$. Similarly, we denote by p_{ij} the projection $T^*X_{123} \rightarrow T^*X_{ij}$. We also denote by p_{12^a} the composite of p_{12} and the antipodal map on T^*X_2 .

Let $A \subset T^*X_{12}$ and $B \subset T^*X_{23}$. We set

$$A \circ B := p_{13}(p_{12^a}^{-1}A \cap p_{23}^{-1}B) \subset T^*X_{13}. \quad (2.6)$$

We define a composition operation of sheaves by

$$\begin{aligned} \circ_{X_2} : \mathbf{D}(\mathbf{k}_{X_{12}}) \times \mathbf{D}(\mathbf{k}_{X_{23}}) &\rightarrow \mathbf{D}(\mathbf{k}_{X_{13}}), \\ (K_{12}, K_{23}) &\mapsto K_{12} \circ_{X_2} K_{23} := \mathrm{R}q_{13!}(q_{12}^{-1}K_{12} \otimes q_{23}^{-1}K_{23}). \end{aligned} \quad (2.7)$$

If there is no risk of confusion, we simply write \circ instead of \circ_{X_2} . By Propositions 2.3 and 2.4, we have the following.

Proposition 2.6. *Let $K_{ij} \in \mathbf{D}(\mathbf{k}_{X_{ij}})$ and set $\Lambda_{ij} := \mathrm{SS}(K_{ij}) \subset T^*X_{ij}$ ($ij = 12, 23$). Assume*

- (1) *q_{13} is proper on $q_{12}^{-1} \mathrm{Supp}(K_{12}) \cap q_{23}^{-1} \mathrm{Supp}(K_{23})$,*
- (2) *$p_{12^a}^{-1} \Lambda_{12} \cap p_{23}^{-1} \Lambda_{23} \cap (T_{X_1}^*X_1 \times T^*X_2 \times T_{X_3}^*X_3) \subset 0_{X_{123}}$.*

Then

$$\mathrm{SS}(K_{12} \circ_{X_2} K_{23}) \subset \Lambda_{12} \circ \Lambda_{23}. \quad (2.8)$$

In this work, we often use sheaves on $X \times \mathbb{R}$. We introduce the operation of convolution for objects of $D(\mathbf{k}_{X \times \mathbb{R}})$. Define the maps

$$\begin{aligned} \tilde{q}_1, \tilde{q}_2, m: X \times \mathbb{R} \times \mathbb{R} &\rightarrow X \times \mathbb{R}, \\ \tilde{q}_1(x, t_1, t_2) &= (x, t_1), \quad \tilde{q}_2(x, t_1, t_2) = (x, t_2), \quad m(x, t_1, t_2) = (x, t_1 + t_2). \end{aligned} \quad (2.9)$$

We define a convolution operation by

$$\begin{aligned} \star: D(\mathbf{k}_{X \times \mathbb{R}}) \times D(\mathbf{k}_{X \times \mathbb{R}}) &\rightarrow D(\mathbf{k}_{X \times \mathbb{R}}), \\ (F, G) &\mapsto F \star G := \mathrm{R}m_!(\tilde{q}_1^{-1}F \otimes \tilde{q}_2^{-1}G). \end{aligned} \quad (2.10)$$

We also introduce a right adjoint to the convolution functor. Set $i: X \times \mathbb{R} \rightarrow X \times \mathbb{R}, (x, t) \mapsto (x, -t)$.

Definition 2.7. For $F, G \in D(\mathbf{k}_{X \times \mathbb{R}})$, one sets

$$\mathrm{Hom}^*(F, G) := \mathrm{R}\tilde{q}_{1*} \mathrm{R}\mathcal{H}om(\tilde{q}_2^{-1}F, m^!G) \quad (2.11)$$

$$\simeq \mathrm{R}m_* \mathrm{R}\mathcal{H}om(\tilde{q}_2^{-1}i^{-1}F, \tilde{q}_1^!G). \quad (2.12)$$

Lemma 2.8. Let $F, G \in D(\mathbf{k}_{X \times \mathbb{R}})$ and assume that there exist two closed cones $A, B \subset \mathbb{R}$ such that $\mathrm{SS}(F) \subset T^*X \times \mathbb{R} \times A$ and $\mathrm{SS}(G) \subset T^*X \times \mathbb{R} \times B$. Then, one has

$$\begin{aligned} \mathrm{SS}(F \star G) &\subset T^*X \times \mathbb{R} \times (A \cap B), \\ \mathrm{SS}(\mathrm{Hom}^*(F, G)) &\subset T^*X \times \mathbb{R} \times (A \cap B). \end{aligned} \quad (2.13)$$

It is useful to define a more general operation, which combines composition and convolution. Set

$$\tilde{q}_{12}: X_{123} \times \mathbb{R}^2 \rightarrow X_{12} \times \mathbb{R}, (x_1, x_2, x_3, t_1, t_2) \mapsto (x_1, x_2, t_1), \quad (2.14)$$

$$\tilde{q}_{23}: X_{123} \times \mathbb{R}^2 \rightarrow X_{23} \times \mathbb{R}, (x_1, x_2, x_3, t_1, t_2) \mapsto (x_2, x_3, t_2), \quad (2.15)$$

$$m_{13}: X_{123} \times \mathbb{R}^2 \rightarrow X_{13} \times \mathbb{R}, (x_1, x_2, x_3, t_1, t_2) \mapsto (x_1, x_3, t_1 + t_2) \quad (2.16)$$

and define

$$\begin{aligned} \bullet_{X_2}: D(\mathbf{k}_{X_{12} \times \mathbb{R}}) \times D(\mathbf{k}_{X_{23} \times \mathbb{R}}) &\rightarrow D(\mathbf{k}_{X_{13} \times \mathbb{R}}), \\ (K_{12}, K_{23}) &\mapsto K_{12} \bullet_{X_2} K_{23} := \mathrm{R}m_{13}!(\tilde{q}_{12}^{-1}K_{12} \otimes \tilde{q}_{23}^{-1}K_{23}). \end{aligned} \quad (2.17)$$

If there is no risk of confusion, we simply write \bullet instead of \bullet_{X_2} .

2.4 Homotopy colimits

Here, we recall the definition of homotopy colimits (cf. [BN93] and [KS06]) and estimate their microsupports.

Let $(F_n)_{n \in \mathbb{N}}$ be a sequence of objects of $D(\mathbf{k}_X)$ together with morphisms $f_n: F_n \rightarrow F_{n+1}$ ($n \in \mathbb{N} = \mathbb{Z}_{\geq 0}$), that is, $(F_n, f_n)_{n \in \mathbb{N}}$ is an inductive system in $D(\mathbf{k}_X)$. Define a morphism $s: \bigoplus_{n \in \mathbb{N}} F_n \rightarrow \bigoplus_{n \in \mathbb{N}} F_n$ as the composite

$$\bigoplus_{n \in \mathbb{N}} F_n \xrightarrow{\bigoplus_n f_n} \bigoplus_{n \in \mathbb{N}} F_{n+1} \simeq \bigoplus_{n \in \mathbb{Z}_{\geq 1}} F_n \rightarrow \bigoplus_{n \in \mathbb{N}} F_n. \quad (2.18)$$

Then one can define the homotopy colimit of the inductive system $(F_n, f_n)_{n \in \mathbb{N}}$ as the cone of the morphism

$$\text{id} - s: \bigoplus_{n \in \mathbb{N}} F_n \rightarrow \bigoplus_{n \in \mathbb{N}} F_n, \quad (2.19)$$

which we write $\text{hocolim}(F_n) \in \mathbf{D}(\mathbf{k}_X)$. We have a canonical morphism $\rho_n: F_n \rightarrow \text{hocolim}(F_n)$. Given a sequence of morphisms $g_n: F_n \rightarrow G$ ($n \in \mathbb{N}$) such that $g_{n+1} \circ f_n = g_n$ for any n , we get a morphism $g: \text{hocolim}(F_n) \rightarrow G$ satisfying $g_n = g \circ \rho_n$.

Lemma 2.9. *Let $(F_n, f_n)_{n \in \mathbb{N}}$ be an inductive system in $\mathbf{D}(\mathbf{k}_X)$. Then*

$$\text{SS}(\text{hocolim}(F_n)) \subset \bigcap_{N \in \mathbb{N}} \overline{\bigcup_{n \geq N} \text{SS}(F_n)}. \quad (2.20)$$

Proof. By Lemma 2.5, $\text{SS}(\bigoplus_{n \in \mathbb{N}} F_n) \subset \overline{\bigcup_n \text{SS}(F_n)}$. Note that we have $\text{hocolim}_n(F_n) \simeq \text{hocolim}_n(G_n)$ with $G_n = F_{n+N}$ for any $N \in \mathbb{N}$. Hence, the result follows from the triangle inequality for microsupports and the definition of homotopy colimits. \square

3 Interleaving distance for sheaves and sheaf quantization

In this section, we review the interleaving distance for sheaves following Petit–Schapira [PS20]. We also briefly review the Tamarkin category [Tam18] and sheaf quantization of Hamiltonian isotopies [GKS12]. See also Guillermou–Schapira [GS14] and Zhang [Zha20] for details of the Tamarkin category.

3.1 Interleaving distance for sheaves

We recall some notions from Petit–Schapira [PS20]. A topological space is said to be *good* if it is Hausdorff, locally compact, countable at infinity, and finite flabby dimension.

Definition 3.1. Let X be a good topological space. A *thickening kernel* on X is a monoidal presheaf \mathfrak{K} on $(\mathbb{R}_{\geq 0}, +)$ with values in the monoidal category $\mathbf{D}(\mathbf{k}_{X \times X})$. In other words, it is a family of kernels $\mathfrak{K}_a \in \mathbf{D}(\mathbf{k}_{X \times X})$ with a morphism $\rho_{b,a}: \mathfrak{K}_b \rightarrow \mathfrak{K}_a$ for $a \leq b$ together with isomorphisms

$$\mathfrak{K}_a \circ \mathfrak{K}_b \simeq \mathfrak{K}_{a+b}, \quad \mathfrak{K}_0 \simeq \mathbf{k}_{\Delta_X} \quad (3.1)$$

satisfying the compatibility conditions (see [PS20, Def. 1.2.2] for details).

For a thickening kernel \mathfrak{K} on X and $F \in \mathbf{D}(\mathbf{k}_X)$, we write $\rho_{b,a}(F)$ for the morphism $\rho_{b,a} \circ \text{id}_F: \mathfrak{K}_b \circ F \rightarrow \mathfrak{K}_a \circ F$ ($a \leq b$).

Definition 3.2. Let \mathfrak{K} be a thickening kernel on X , $F, G \in \mathbf{D}(\mathbf{k}_X)$, and $a, b \in \mathbb{R}_{\geq 0}$.

(i) The pair (F, G) is said to be *weakly (a, b) -isomorphic* if there exist morphisms $\alpha, \delta: \mathfrak{K}_a \circ F \rightarrow G$ and $\beta, \gamma: \mathfrak{K}_b \circ G \rightarrow F$ such that

- (1) the composite $\mathfrak{K}_{a+b} \circ F \xrightarrow{\mathfrak{K}_a \circ \alpha} \mathfrak{K}_b \circ G \xrightarrow{\beta} F$ is equal to $\rho_{a+b,0}(F)$,
- (2) the composite $\mathfrak{K}_{a+b} \circ G \xrightarrow{\mathfrak{K}_b \circ \gamma} \mathfrak{K}_a \circ F \xrightarrow{\delta} G$ is equal to $\rho_{a+b,0}(G)$,
- (3) $\alpha \circ \rho_{2a,a}(F) = \delta \circ \rho_{2a,a}(F)$ and $\beta \circ \rho_{2b,b}(G) = \gamma \circ \rho_{2b,b}(G)$.

(ii) The pair (F, G) is said to be (a, b) -isomorphic if there exist morphisms $\alpha: \mathfrak{K}_a \circ F \rightarrow G$ and $\beta: \mathfrak{K}_a \circ G \rightarrow F$ such that

- (1) the composite $\mathfrak{K}_{a+b} \circ F \xrightarrow{\mathfrak{K}_a \circ \alpha} \mathfrak{K}_b \circ G \xrightarrow{\beta} F$ is equal to $\rho_{a+b,0}(F)$,
- (2) the composite $\mathfrak{K}_{a+b} \circ G \xrightarrow{\mathfrak{K}_b \circ \beta} \mathfrak{K}_a \circ F \xrightarrow{\alpha} G$ is equal to $\rho_{a+b,0}(G)$.

In this case, the pair of morphisms (α, β) is called an (a, b) -isomorphism.

If (F, G) is (a, a) -isomorphic then F and G are called a -isomorphic.

(iii) One sets

$$d_{\mathfrak{K}}(F, G) := \inf \left\{ a + b \in \mathbb{R}_{\geq 0} \mid \begin{array}{l} a, b \in \mathbb{R}_{\geq 0}, \\ (F, G) \text{ is weakly } (a, b)\text{-isomorphic} \end{array} \right\}, \quad (3.2)$$

which defines a pseudo-distance on the category $\mathbf{D}(\mathbf{k}_X)$

(iv) The object F is said to be a -torsion or a -trivial if $\rho_{a,0}(F): \mathfrak{K}_a \circ F \rightarrow F$ is the zero morphism.

Remark 3.3. (i) One can see that

$$(a, b)\text{-isomorphic} \Rightarrow \text{weakly } (a, b)\text{-isomorphic} \quad (3.3)$$

and

$$\begin{aligned} \text{weakly } (a, b)\text{-isomorphic} &\Rightarrow (2a, 2b)\text{-isomorphic} \\ &\Rightarrow 2 \max(a, b)\text{-isomorphic.} \end{aligned} \quad (3.4)$$

(ii) In [PS20], the authors define

$$\text{dist}_{\mathfrak{K}}(F, G) := \inf \{ a \in \mathbb{R}_{\geq 0} \mid F \text{ and } G \text{ are } a\text{-isomorphic} \} \quad (3.5)$$

and call it the *interleaving distance* associated with \mathfrak{K} . By (i), the pseudo-distances $d_{\mathfrak{K}}$ and $\text{dist}_{\mathfrak{K}}$ are equivalent. That is,

$$d_{\mathfrak{K}}(F, G) \leq 2 \text{dist}_{\mathfrak{K}}(F, G) \leq 4d_{\mathfrak{K}}(F, G). \quad (3.6)$$

(iii) The interleaving distance above is a generalization of the convolution distance d_C on $\mathbf{D}(\mathbf{k}_{\mathbb{R}})$ introduced by Kashiwara–Schapira [KS18a] and later investigated by [KS21, BG21, BP21, BGO19] and others. Indeed, when $X = \mathbb{R}$ and $\mathfrak{K}_a = \mathbf{k}_{\Delta_a}$, where $\Delta_a := \{(x, y) \in \mathbb{R}^2 \mid \|x - y\| \leq a\} \subset \mathbb{R}^2$ is the thickened diagonal, we find that $d_C = \text{dist}_{\mathfrak{K}}$.

In what follows, let \mathfrak{K} be a thickening kernel on X . The statement of the following lemma is stronger than [AI20, Lem. 4.14], but the proof itself is almost the same. We need the stronger result in this paper, so we reproduce the proof for the convenience of the reader.

Lemma 3.4 (cf. [AI20, Lem. 4.14]). *Let $F \xrightarrow{u} G \xrightarrow{v} H \xrightarrow{w} F[1]$ be an exact triangle in $\mathbf{D}(\mathbf{k}_X)$ and assume that F is c -torsion. Then (G, H) is weakly $(0, c)$ -isomorphic.*

Proof. By assumption, we have $w \circ \rho_{c,0}(H) = \rho_{c,0}(F[1]) \circ (\mathfrak{K}_c \circ_X w) = 0$. Hence, we get a morphism $\gamma: \mathfrak{K}_c \circ H \rightarrow G$ satisfying $\rho_{c,0}(H) = v \circ \gamma$:

$$\begin{array}{ccccccc}
\mathfrak{K}_c \circ F & \xrightarrow{\mathfrak{K}_c \circ u} & \mathfrak{K}_c \circ G & \xrightarrow{\mathfrak{K}_c \circ v} & \mathfrak{K}_c \circ H & \xrightarrow{\mathfrak{K}_c \circ w} & \mathfrak{K}_c \circ F[1] \\
\downarrow & & \downarrow & \swarrow \gamma & \downarrow & & \downarrow 0 \\
F & \xrightarrow{u} & G & \xrightarrow{v} & H & \xrightarrow{w} & F[1].
\end{array} \tag{3.7}$$

On the other hand, since $\rho_{c,0}(G) \circ (\mathfrak{K}_c \circ_X u) = u \circ \rho_{c,0}(F) = 0$, there exists a morphism $\beta: \mathfrak{K}_c \circ H \rightarrow G$ satisfying $\rho_{c,0}(G) = \beta \circ (\mathfrak{K}_c \circ_X v)$:

$$\begin{array}{ccccccc}
\mathfrak{K}_c \circ F & \xrightarrow{\mathfrak{K}_c \circ u} & \mathfrak{K}_c \circ G & \xrightarrow{\mathfrak{K}_c \circ v} & \mathfrak{K}_c \circ H & \xrightarrow{\mathfrak{K}_c \circ w} & \mathfrak{K}_c \circ F[1] \\
0 \downarrow & & \downarrow & \swarrow \beta & \downarrow & & \downarrow \\
F & \xrightarrow{u} & G & \xrightarrow{v} & H & \xrightarrow{w} & F[1].
\end{array} \tag{3.8}$$

Moreover, we obtain

$$\begin{aligned}
\beta \circ \rho_{2c,c}(H) &= \beta \circ (\mathfrak{K}_c \circ_X \rho_{c,0}(H)) \\
&= \beta \circ (\mathfrak{K}_c \circ_X v) \circ (\mathfrak{K}_c \circ_X \gamma) \\
&= \rho_{c,0}(G) \circ (\mathfrak{K}_c \circ_X \gamma) \\
&= \gamma \circ (\mathfrak{K}_c \circ_X \rho_{c,0}(H)) = \gamma \circ \rho_{2c,c}(H),
\end{aligned} \tag{3.9}$$

which proves the lemma. \square

In particular, if $F \rightarrow G \rightarrow H \xrightarrow{+1}$ is an exact triangle, F is a -torsion, and G is b -torsion, then H is $(a+b)$ -torsion.

In our later applications, we mainly focus on the interleaving distance on the derived category $D(\mathbf{k}_{X \times \mathbb{R}_t})$. For $c \in \mathbb{R}_{\geq 0}$, define

$$\mathfrak{K}_c := \mathbf{k}_{\Delta_X \times \Delta_c} \in D(\mathbf{k}_{(X \times \mathbb{R})^2}), \tag{3.10}$$

where $\Delta_c := \{(t_1, t_2) \mid \|t_1 - t_2\| \leq c\} \subset \mathbb{R}^2$ is the thickened diagonal of $\Delta_{\mathbb{R}} \subset \mathbb{R}^2$. Then the assignment $c \mapsto \mathfrak{K}_c$ defines a thickening kernel on $X \times \mathbb{R}_t$. Hence, we can consider the pseudo-distance $d_{\mathfrak{K}}$ on $D(\mathbf{k}_{X \times \mathbb{R}_t})$ associated with \mathfrak{K} , which we denote by $d_{X \times \mathbb{R}_t}$. This is a slight modification of the relative distance $\text{dist}_{X \times \mathbb{R}_t / X}$ studied in [PS20]. Note also that $\mathfrak{K}_c \circ F \simeq \mathbf{k}_{X \times [-c, c]} \star F$. With the notation in Subsection 2.3, for $F, F' \in D(\mathbf{k}_{X_{12} \times \mathbb{R}_t})$ and $G, G' \in D(\mathbf{k}_{X_{23} \times \mathbb{R}_t})$,

$$d_{X_{13} \times \mathbb{R}_t}(F \bullet G, F' \bullet G') \leq d_{X_{12} \times \mathbb{R}_t}(F, F') + d_{X_{23} \times \mathbb{R}_t}(G, G'). \tag{3.11}$$

3.2 Tamarkin category

Let X be a manifold without boundary. We let $(t; \tau)$ denote the homogeneous coordinate system on $T^*\mathbb{R}_t$. It is prove by Tamarkin [Tam18] that the functor $P_t := \mathbf{k}_{X \times [0, \infty)} \star (*): D(\mathbf{k}_{X \times \mathbb{R}_t}) \rightarrow D(\mathbf{k}_{X \times \mathbb{R}_t})$ defines a projector onto ${}^\perp D_{\{\tau \leq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$, where $\{\tau \leq 0\} = \{(x, t; \xi, \tau) \mid \tau \leq 0\} \subset T^*(M \times \mathbb{R}_t)$ and ${}^\perp(*)$ denotes the left orthogonal.

Definition 3.5. One defines

$$\mathcal{D}(X) := {}^\perp D_{\{\tau \leq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t}), \tag{3.12}$$

and call it the *Tamarkin category* of X .

By the remark above, for an object $F \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$, $F \in \mathcal{D}(M)$ if and only if $P_t(F) \simeq F$. Note also that $\mathcal{D}(X) \subset \mathcal{D}_{\{\tau \geq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$ by Lemma 2.8.

Definition 3.6. One defines $d_{\mathcal{D}}(X)$ as the restriction of the pseudo-distance $d_{X \times \mathbb{R}_t}$ on $\mathcal{D}(\mathbf{k}_{X \times \mathbb{R}_t})$ to $\mathcal{D}(X)$.

We will describe the pseudo-distance using the translation to the \mathbb{R}_t -direction. For $c \in \mathbb{R}$, let $T_c: X \times \mathbb{R}_t \rightarrow X \times \mathbb{R}_t$, $(x, t) \mapsto (x, t + c)$ be the translation map by c to the \mathbb{R} -direction. In what follows, we write T_c instead of T_{c*} for simplicity. Recall that we have set $\mathfrak{K}_c := \mathbf{k}_{\Delta_X \times \Delta_c} \in \mathcal{D}(\mathbf{k}_{(X \times \mathbb{R})^2})$.

Lemma 3.7. *Let $F \in \mathcal{D}_{\{\tau \geq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$. Then $\mathfrak{K}_c \circ F \simeq T_{-c}F$.*

Proof. First we recall that $\mathfrak{K}_c \circ F \simeq \mathbf{k}_{X \times [-c, c]} \star F$. Since there exist an exact triangle $\mathbf{k}_{X \times (-c, c]} \rightarrow \mathbf{k}_{X \times [-c, c]} \rightarrow \mathbf{k}_{X \times \{-c\}} \xrightarrow{+1}$ and an isomorphism $T_{-c}F \simeq \mathbf{k}_{X \times \{-c\}} \star F$, it suffices to show that $\mathbf{k}_{X \times (-c, c]} \star F \simeq 0$. By Lemma 5.2, there exists $H \in \mathcal{D}(\mathbf{k}_X)$ such that $\mathbf{k}_{X \times (-\infty, 0]} \star F \simeq H \boxtimes \mathbf{k}_{\mathbb{R}_t}$. Thus, we obtain

$$\begin{aligned} \mathbf{k}_{X \times (-c, c]} \star F &\simeq \mathbf{k}_{X \times (-c, c]} \star \mathbf{k}_{X \times (-\infty, 0]} \star F \\ &\simeq \mathbf{k}_{X \times (-c, c]} \star (H \boxtimes \mathbf{k}_{\mathbb{R}}) \\ &\simeq H \boxtimes (\mathbf{k}_{(-c, c]} \star \mathbf{k}_{\mathbb{R}}) \simeq 0, \end{aligned} \tag{3.13}$$

which completes the proof. \square

Let $F \in \mathcal{D}_{\{\tau \geq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$. Then, we have an isomorphism

$$\mathrm{R}m_*(\tilde{q}_1^{-1} \mathbf{k}_{X \times [0, \infty)} \otimes \tilde{q}_2^{-1} F) \xrightarrow{\sim} \mathrm{R}m_*(\tilde{q}_1^{-1} \mathbf{k}_{X \times \{0\}} \otimes \tilde{q}_2^{-1} F) \simeq F. \tag{3.14}$$

Hence, for $c, d \in \mathbb{R}_{\geq 0}$ with $c \leq d$, the canonical morphism $\mathbf{k}_{X \times [c, \infty)} \rightarrow \mathbf{k}_{X \times [d, \infty)}$ induces a canonical morphism

$$\begin{aligned} \tau_{c,d}(F): T_c F &\simeq \mathrm{R}m_*(\tilde{q}_1^{-1} \mathbf{k}_{X \times [c, \infty)} \otimes \tilde{q}_2^{-1} F) \\ &\rightarrow \mathrm{R}m_*(\tilde{q}_1^{-1} \mathbf{k}_{X \times [d, \infty)} \otimes \tilde{q}_2^{-1} F) \simeq T_d F. \end{aligned} \tag{3.15}$$

By Lemma 3.7, the morphism is identified with

$$T_{c+d} \rho_{d,c}(F): T_{c+d}(\mathfrak{K}_d \circ F) \rightarrow T_{c+d}(\mathfrak{K}_c \circ F). \tag{3.16}$$

Hence, a pair (F, G) of objects of $\mathcal{D}(X)$ is weakly (a, b) -isomorphic if and only if there exist morphisms $\alpha, \delta: F \rightarrow T_a G$ and $\beta, \gamma: G \rightarrow T_b F$ such that

- (1) $F \xrightarrow{\alpha} T_a G \xrightarrow{T_a \beta} T_{a+b} F$ is equal to $\tau_{0, a+b}(F): F \rightarrow T_{a+b} F$,
- (2) $G \xrightarrow{\gamma} T_b F \xrightarrow{T_b \delta} T_{a+b} G$ is equal to $\tau_{0, a+b}(G): G \rightarrow T_{a+b} G$,
- (3) $\tau_{a, 2a}(G) \circ \alpha = \tau_{a, 2a}(G) \circ \delta$ and $\tau_{b, 2b}(F) \circ \beta = \tau_{b, 2b}(F) \circ \gamma$.

In this form, we can see that $d_{\mathcal{D}(X)}$ is similar to the pseudo-distance introduced in [AI20] (see Remark 3.8 below).

Remark 3.8. The terminology has been changed from that in [AI20]. In that paper, “weakly (a, b) -isomorphic” in this paper was called “ (a, b) -isomorphic”. Moreover, we defined the notion of “ (a, b) -interleaved” as follows: a pair (F, G) of objects of $\mathcal{D}(X)$ is said to be (a, b) -interleaved if there exist morphisms $\alpha, \delta: F \rightarrow T_a G$ and $\beta, \gamma: G \rightarrow T_b F$ satisfying the conditions (1) and (2) above. One can see that

(a, b) -isomorphic \Rightarrow weakly (a, b) -isomorphic \Rightarrow (a, b) -interleaved.

We also remark that the distance $d_{\mathcal{D}(X)}$ in [AI20] is defined by the relation “ (a, b) -interleaved” instead of “weakly (a, b) -isomorphic”, and hence it is different from that in Definition 3.6. However, all the results in [AI20] holds also for the modified $d_{\mathcal{D}(X)}$ thanks to Lemma 3.4. See also Proposition 3.9 below.

The following proposition is stronger than the similar results in the published version of [AI20].

Proposition 3.9 (cf. [AI20, Prop. 4.15]). *Let I be an open interval containing the closed interval $[0, 1]$ and $\mathcal{H} \in \mathcal{D}_{\{\tau \geq 0\}}(X \times \mathbb{R}_t \times I)$. Assume that there exist continuous functions $f, g: I \rightarrow \mathbb{R}_{\geq 0}$ satisfying*

$$\text{SS}(\mathcal{H}) \subset T^*X \times \{(t, s; \tau, \sigma) \mid -f(s) \cdot \tau \leq \sigma \leq g(s) \cdot \tau\}. \quad (3.17)$$

Then $(\mathcal{H}|_{X \times \mathbb{R}_t \times \{0\}}, \mathcal{H}|_{X \times \mathbb{R}_t \times \{1\}})$ is weakly $(\int_0^1 g(s)ds + \varepsilon, \int_0^1 f(s)ds + \varepsilon)$ -isomorphic for any $\varepsilon \in \mathbb{R}_{>0}$. In particular, $d_{X \times \mathbb{R}_t}(\mathcal{H}|_{X \times \mathbb{R}_t \times \{0\}}, \mathcal{H}|_{X \times \mathbb{R}_t \times \{1\}}) \leq \int_0^1 (f + g)(s)ds$.

Proof. The proof is similar to that of [AI20, Prop. 4.15]. We only need to replace [AI20, Lem. 4.14] with Lemma 3.4. \square

3.3 Sheaf quantization of Hamiltonian isotopies

In this subsection, we first recall the existence and uniqueness result of a sheaf quantization of a Hamiltonian isotopy due to Guillermou–Kashiwara–Schapira [GKS12]. Then we give an energy estimate on the sheaf quantization, which is a slight generalization of our previous result in [AI20].

Let M be a connected manifold without boundary and I an open interval of \mathbb{R} containing the closed interval $[0, 1]$. A compactly supported *Hamiltonian isotopy* is a flow of the Hamiltonian vector field of a compactly supported C^∞ -function $H = (H_s)_{s \in I}: T^*M \times I \rightarrow \mathbb{R}$. In this paper, the isotopy associated with H is denoted by $\phi^H = (\phi_s^H)_{s \in I}: T^*M \times I \rightarrow T^*M$. Remark that $(\phi_s^H)^{-1} = \phi_s^{\overline{H}}$ with $\overline{H}_s(p) := -H_s(\phi_s^H(p))$. Moreover, for two compactly supported functions $H, H': T^*M \times I \rightarrow \mathbb{R}$, we have

$$\phi_s^H \circ \phi_s^{H'} = \phi_s^{H \sharp H'}, \quad (3.18)$$

where $(H \sharp H')_s(p) := H_s(p) + H'_s((\phi_s^H)^{-1}(p))$. In particular, for two compactly supported functions $H, H': T^*M \times I \rightarrow \mathbb{R}$,

$$(\phi_s^H)^{-1} \circ \phi_s^{H'} = \phi_s^{\overline{H} \sharp H'}, \quad (3.19)$$

where $(\overline{H} \sharp H')_s(p) = (H' - H)_s(\phi_s^H(p))$.

Definition 3.10. Let $\phi^H = (\phi_s^H)_{s \in I}: T^*M \times I \rightarrow T^*M$ be the compactly supported Hamiltonian isotopy associated with a compactly supported function $H: T^*M \times I \rightarrow \mathbb{R}$.

(i) One defines $\widehat{H}: \mathring{T}^*(M \times \mathbb{R}_t) \times I \rightarrow \mathbb{R}$ by

$$\widehat{H}((x, t; \xi, \tau), s) := \begin{cases} \tau H((x; \xi/\tau), s) & (\tau \neq 0) \\ 0 & (\tau = 0) \end{cases} \quad (3.20)$$

and $\widehat{\phi} = (\widehat{\phi}_s)_{s \in I}: \mathring{T}^*(M \times \mathbb{R}_t) \times I \rightarrow \mathring{T}^*(M \times \mathbb{R}_t)$ to be the homogeneous Hamiltonian flow of \widehat{H} .

(ii) One defines a conic Lagrangian submanifold $\Lambda_{\widehat{\phi}}$ of $\mathring{T}^*(M \times \mathbb{R})^2 \times T^*I$ by

$$\Lambda_{\widehat{\phi}} := \left\{ \left(\widehat{\phi}((x, t; \xi, \tau), s), (x, t; -\xi, -\tau), (s; -\widehat{H}(\widehat{\phi}((x, t; \xi, \tau), s), s)) \right) \mid \right. \\ \left. (x, t; \xi, \tau) \in \mathring{T}^*(M \times \mathbb{R}_t), s \in I \right\}. \quad (3.21)$$

For a Hamiltonian function $H: T^*M \times I \rightarrow \mathbb{R}$, we also define $u = (u_s)_{s \in I}: T^*M \times I \rightarrow \mathbb{R}$ by $u_s(p) = \int_0^s (H_{s'} - \alpha(X_{s'}))(\phi_{s'}^H(p)) ds'$, where $(X_s)_{s \in I}$ is the Hamiltonian vector field for H . Then $\widehat{\phi}$ can be written as

$$\widehat{\phi}_s(x, t; \xi, \tau) = (x', t + u_s(x; \xi/\tau); \xi', \tau), \quad (3.22)$$

where $(x'; \xi'/\tau) = \phi_s(x; \xi/\tau)$ for $\tau \neq 0$, and $\widehat{\phi}_s(x, t; \xi, 0) = (x, t; \xi, 0)$. Hereafter, we use the convention that $\tau H_s(\phi_s(x; \xi/\tau)) = 0$ and $u_s(x; \xi/\tau) = 0$ when $\tau = 0$. Moreover, we write $(x'; \xi'/\tau) = \phi_s(x; \xi/\tau)$ also for $\tau = 0$, in which case it is understood that $(x'; \xi') = (x; \xi)$. We have $du_s = \alpha - (\phi_s)^*\alpha$, which gives the following properties.

Lemma 3.11. (i) *If $\phi_1 = \text{id}_{T^*M}$, then $u_1 \equiv 0$.*

(ii) *If $\phi_1(0_M) = 0_M$, then $u_1|_{0_M} \equiv c$ for some constant $c \in \mathbb{R}$.*

The main theorem of [GKS12] is the following.

Theorem 3.12 ([GKS12, Thm. 4.3]). *Let $\phi: T^*M \times I \rightarrow T^*M$ be a compactly supported Hamiltonian isotopy and $H: T^*M \times I \rightarrow \mathbb{R}$ a C^∞ -function with Hamiltonian flow ϕ . Then there exists a unique simple object $\widetilde{K}^H \in \mathcal{D}(\mathbf{k}_{(M \times \mathbb{R})^2 \times I})$ such that $\mathring{\text{SS}}(\widetilde{K}^H) = \Lambda_{\widehat{\phi}}$ and $\widetilde{K}^H|_{(M \times \mathbb{R})^2 \times \{0\}} \simeq \mathbf{k}_{\Delta_{M \times \mathbb{R}}}$.*

Set $\widetilde{K}_s^H := \widetilde{K}^H|_{(M \times \mathbb{R})^2 \times \{s\}} \in \mathcal{D}(\mathbf{k}_{(M \times \mathbb{R})^2})$. Note that $\mathring{\text{SS}}(\widetilde{K}_s^H) \subset \Lambda_{\widehat{\phi}} \circ T_s^*I$. We also have

$$\widetilde{K}_s^H \circ \widetilde{K}_s^{H'} \simeq \widetilde{K}_s^{H \sharp H'}. \quad (3.23)$$

It is also proved by Guillermou–Schapira [GS14, Prop. 4.29] that the composition with \widetilde{K}_s^H defines a functor

$$\Phi_s^H := \widetilde{K}_s^H \circ (*): \mathcal{D}(M) \longrightarrow \mathcal{D}(M). \quad (3.24)$$

Define $q: (M \times \mathbb{R})^2 \times I \rightarrow M^2 \times \mathbb{R}_t \times I, (x_1, t_1, x_2, t_2, s) \mapsto (x_1, x_2, t_1 - t_2, s)$ and set

$$\Lambda'_H := q_\pi q_d^{-1}(\Lambda_{\widehat{\phi}}) \\ = \left\{ ((x'; \xi'), (x; -\xi), (u_s(x; \xi/\tau); \tau), (s; -\tau H_s(\phi_s(x; \xi/\tau)))) \mid \right. \\ \left. (x; \xi) \in T^*M, s \in I, \tau \in \mathbb{R}, (x'; \xi'/\tau) = \phi_s(x; \xi/\tau) \right\} \\ \subset \mathring{T}^*(M^2 \times \mathbb{R}_t \times I). \quad (3.25)$$

Then the inverse image functor q^{-1} gives an equivalence (see [Gui19, Cor. 2.3.2])

$$\{K \in \mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t \times I}) \mid \mathring{\text{SS}}(K) = \Lambda'_H\} \xrightarrow{\sim} \{\widetilde{K} \in \mathcal{D}(\mathbf{k}_{(M \times \mathbb{R})^2 \times I}) \mid \mathring{\text{SS}}(K) = \Lambda_{\widehat{\phi}}\}. \quad (3.26)$$

Recall that we have a projector $P_l: \mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t}) \rightarrow {}^\perp \mathcal{D}_{\{\tau \leq 0\}}(\mathbf{k}_{M^2 \times \mathbb{R}_t}) = \mathcal{D}(M^2)$ and similarly for $\mathcal{D}(M^2 \times I)$.

Definition 3.13. Let $H: T^*M \times I \rightarrow \mathbb{R}$ be a compactly supported Hamiltonian function.

- (i) One defines $K^H \in \mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t \times I})$ to be the object such that $\mathring{\text{SS}}(K^H) = \Lambda'_H$ and $K^H|_{M^2 \times \mathbb{R}_t \times \{0\}} \simeq \mathbf{k}_{\Delta_M \times \{0\}}$, that is, the object K^H satisfying $q^{-1}K^H \simeq \tilde{K}^H$. One also sets $K_s^H := K^H|_{M^2 \times \mathbb{R}_t \times \{s\}}$ for simplicity.
- (ii) One defines $\mathcal{K}_s^H := P_l(K_s^H) \in \mathcal{D}(M^2)$ and $\mathcal{K}^H := P_l(K^H) \in \mathcal{D}(M^2 \times I)$.

Note that $\tilde{K}_s^H \circ F \simeq K_s^H \bullet F$ for any $F \in \mathcal{D}(M)$ and $s \in I$. By the associativity, for any $F \in \mathcal{D}(M)$ and $s \in I$, we have

$$\begin{aligned} K_s^H \bullet F &\simeq K_s^H \bullet (\mathbf{k}_{M \times [0, \infty)} \star F) \\ &\simeq (K_s^H \bullet_M \mathbf{k}_{M \times [0, \infty)}) \bullet F \\ &\simeq (K_s^H \star \mathbf{k}_{M \times M \times [0, \infty)}) \bullet F \simeq \mathcal{K}_s^H \bullet F. \end{aligned} \quad (3.27)$$

Now we give a generalization of [AI20, Thm. 4.16] in terms of the distance between \mathcal{K}_1^H and \mathcal{K}_1^0 , where 0 is the zero function on $T^*M \times I$. For a compactly supported Hamiltonian function $H: T^*M \times I \rightarrow \mathbb{R}$, we define

$$\begin{aligned} E_+(H) &:= \int_0^1 \max_p H_s(p) ds, & E_-(H) &:= - \int_0^1 \min_p H_s(p) ds, \\ \|H\|_{\text{osc}} &:= E_+(H) + E_-(H) = \int_0^1 \left(\max_p H_s(p) - \min_p H_s(p) \right) ds. \end{aligned} \quad (3.28)$$

Theorem 3.14. *Let $H: T^*M \times I \rightarrow \mathbb{R}$ be a compactly supported Hamiltonian function. Then $(\mathcal{K}_1^0, \mathcal{K}_1^H)$ is weakly $(E_-(H) + \varepsilon, E_+(H) + \varepsilon)$ -isomorphic for any $\varepsilon \in \mathbb{R}_{>0}$, where 0 denotes the zero function on $T^*M \times I$. In particular, $d_{\mathcal{D}(M^2)}(\mathcal{K}_1^0, \mathcal{K}_1^H) \leq \|H\|_{\text{osc}}$.*

Proof. We can apply Proposition 3.9 to \mathcal{K}^H and obtain the result by (3.25). \square

For two compactly supported Hamiltonian functions $H, H': T^*M \times I \rightarrow \mathbb{R}$, by (3.11) and (3.23), we have

$$d_{\mathcal{D}(M^2)}(\mathcal{K}_1^H, \mathcal{K}_1^{H'}) = d_{\mathcal{D}(M^2)}(\mathcal{K}_1^0, \mathcal{K}_1^{\overline{H \# H'}}) \leq \|H - H'\|_{\text{osc}}. \quad (3.29)$$

4 Completeness of derived category of sheaves

In this section, we prove the completeness of the derived category $\mathcal{D}(\mathbf{k}_X)$ with respect to the pseudo-distance $d_{\mathfrak{R}}$ associated with a thickening kernel \mathfrak{R} . If a category with a persistence structure admits any sequential colimit, then the category is complete with respect to the interleaving distance (Cruz [Cru19] and Scozzola [Sco20]). However, the derived category does not admit sequential colimits. Hence, we construct a limit object by using a homotopy colimit instead. Let X be a manifold throughout this section.

In Lemma 3.4, we saw that for an exact triangle $F \rightarrow G \rightarrow H \xrightarrow{+1}$, if H is c -torsion, then (F, G) is weakly $(0, c)$ -isomorphic. Conversely, we obtain Proposition 4.2 below, which is a key to our construction of limit objects.

Lemma 4.1. *Let $F \in \mathcal{D}(\mathbf{k}_X)$ and $a \in \mathbb{R}_{\geq 0}$ and consider the exact triangle*

$$\mathfrak{R}_a \circ F \xrightarrow{\rho_{a,0}(F)} F \rightarrow \text{Cone}(\rho_{a,0}(F)) \xrightarrow{+1}. \quad (4.1)$$

Then $\text{Cone}(\rho_{a,0}(F))$ is $2a$ -torsion.

Proof. Set $C := \text{Cone}(\rho_{a,0})$ and consider the following commutative diagram:

$$\begin{array}{ccccccc}
\mathfrak{K}_{3a} \circ F & \longrightarrow & \mathfrak{K}_{2a} \circ F & \longrightarrow & \mathfrak{K}_{2a} \circ C & \longrightarrow & \mathfrak{K}_{3a} \circ F[1] \longrightarrow \mathfrak{K}_{2a} \circ F[1] \\
& & \downarrow & \nearrow & \downarrow & & \downarrow \\
& & \mathfrak{K}_a \circ F & \xrightarrow{\kappa} & \mathfrak{K}_a \circ C & \longrightarrow & \mathfrak{K}_{2a} \circ F[1] \\
& \nearrow & \downarrow & & \downarrow & & \downarrow \\
\mathfrak{K}_a \circ F & \longrightarrow & F & \longrightarrow & C & \longrightarrow & \mathfrak{K}_a \circ F[1].
\end{array} \tag{4.2}$$

The composite morphism $\mathfrak{K}_{2a} \circ C \rightarrow \mathfrak{K}_{3a} \circ F[1] \rightarrow \mathfrak{K}_{2a} \circ F[1]$ is zero since $\mathfrak{K}_{3a} \circ F \rightarrow \mathfrak{K}_{2a} \circ F \rightarrow \mathfrak{K}_{2a} \circ C \xrightarrow{+1}$ is an exact triangle. By the commutativity, the composite $\mathfrak{K}_{2a} \circ C \rightarrow \mathfrak{K}_a \circ C \rightarrow \mathfrak{K}_{2a} \circ F[1]$ is also zero. Hence, there exists a morphism $\mathfrak{K}_{2a} \circ C \rightarrow \mathfrak{K}_a \circ F$ that makes the lower triangle commutative. Therefore, the morphism $\mathfrak{K}_{2a} \circ C \rightarrow C$ factors the composite $\mathfrak{K}_a \circ F \rightarrow F \rightarrow C$ and hence it is zero. \square

Proposition 4.2. *Let $F, G \in \mathcal{D}(M)$ and assume that the pair (F, G) is (a, b) -isomorphic. Let $(\alpha: \mathfrak{K}_a \circ F \rightarrow G, \beta: \mathfrak{K}_b \circ G \rightarrow F)$ be an (a, b) -isomorphism on (F, G) and consider the exact triangle*

$$\mathfrak{K}_a \circ F \rightarrow G \rightarrow \text{Cone}(\alpha) \xrightarrow{+1} . \tag{4.3}$$

Then $\text{Cone}(\alpha)$ is $3(a+b)$ -torsion.

Proof. Consider the three exact triangles:

$$\begin{array}{ccccccc}
\mathfrak{K}_{a+b} \circ F & \xrightarrow{\mathfrak{K}_b \circ \alpha} & \mathfrak{K}_b \circ G & \longrightarrow & \mathfrak{K}_a \circ \text{Cone}(\alpha) & \xrightarrow{u} & \mathfrak{K}_{a+b} \circ F[1], \\
\mathfrak{K}_{a+b} \circ F & \xrightarrow{\rho_{a+b,0}(F)} & F & \longrightarrow & \text{Cone}(\rho_{a+b,0}(F)) & \longrightarrow & \mathfrak{K}_{a+b} \circ F[1], \\
\mathfrak{K}_b \circ G & \xrightarrow{\beta} & F & \longrightarrow & \text{Cone}(\beta) & \longrightarrow & \mathfrak{K}_b \circ G[1].
\end{array} \tag{4.4}$$

Note that $\beta \circ (\mathfrak{K}_b \circ \alpha) = \rho_{a+b,0}(F)$. By the octahedral axiom, we have the following commutative diagram, where the bottom row is also an exact triangle:

$$\begin{array}{ccccccc}
\mathfrak{K}_{a+b} \circ F & \xrightarrow{\mathfrak{K}_b \circ \alpha} & \mathfrak{K}_b \circ G & \longrightarrow & \mathfrak{K}_b \circ \text{Cone}(\alpha) & \xrightarrow{u} & \mathfrak{K}_{a+b} \circ F[1] \\
\parallel & & \downarrow \beta & & \downarrow v & & \parallel \\
\mathfrak{K}_{a+b} \circ F & \xrightarrow{\rho_{a+b,0}(F)} & F & \longrightarrow & \text{Cone}(\rho_{a+b,0}(F)) & \longrightarrow & \mathfrak{K}_{a+b} \circ F[1] \\
\mathfrak{K}_b \circ \alpha \downarrow & & \parallel & & \downarrow & & \downarrow \\
\mathfrak{K}_b \circ G & \xrightarrow{\beta} & F & \longrightarrow & \text{Cone}(\beta) & \longrightarrow & \mathfrak{K}_b \circ G[1] \\
\downarrow & & \downarrow & & \parallel & & \downarrow \\
\mathfrak{K}_b \circ \text{Cone}(\alpha) & \longrightarrow & \text{Cone}(\rho_{a+b,0}(F)) & \longrightarrow & \text{Cone}(\beta) & \longrightarrow & \mathfrak{K}_b \circ \text{Cone}(\alpha)[1].
\end{array} \tag{4.5}$$

In particular, the morphism $u: \mathfrak{K}_b \circ \text{Cone}(\alpha) \rightarrow \mathfrak{K}_{a+b} \circ F[1]$ factors through $\text{Cone}(\rho_{a+b,0}(F))$. Then the commutative diagram

$$\begin{array}{ccc}
\mathfrak{K}_{2a+3b} \circ \text{Cone}(\alpha) & \xrightarrow{\rho_{2a+3b,b}(\text{Cone}(\alpha))} & \mathfrak{K}_b \circ \text{Cone}(\alpha) \xrightarrow{u} \mathfrak{K}_{a+b} \circ F[1] \\
\mathfrak{K}_{2a+2b} \circ v \downarrow & & \downarrow v \\
\mathfrak{K}_{2a+2b} \circ \text{Cone}(\rho_{a+b,0}(F)) & \xrightarrow{0} & \text{Cone}(\rho_{a+b,0}(F)) \nearrow
\end{array} \tag{4.6}$$

proves that the composite morphism in the first row is zero by Lemma 4.1. Thus, we obtain a morphism $\mathfrak{K}_{3a+3b} \circ \text{Cone}(\alpha) \rightarrow \mathfrak{K}_{a+b} \circ G$ that makes the following diagram commute, where the vertical arrows are the corresponding ρ 's:

$$\begin{array}{ccccc}
& & \mathfrak{K}_{3a+3b} \circ \text{Cone}(\alpha) & & \\
& & \downarrow & & \\
& \mathfrak{K}_{a+b} \circ G & \xrightarrow{\quad} & \mathfrak{K}_{a+b} \circ \text{Cone}(\alpha) & \xrightarrow{\mathfrak{K}_b \circ u} & \mathfrak{K}_{2a+b} \circ F[1] \\
& \swarrow \mathfrak{K}_a \circ \beta & \downarrow & \downarrow & & \\
\mathfrak{K}_a \circ F & \xrightarrow{\alpha} & G & \xrightarrow{\quad} & \text{Cone}(\alpha) & \xrightarrow{\quad} & \mathfrak{K}_a \circ F[1].
\end{array} \tag{4.7}$$

Hence, the morphism $\rho_{3a+3b,0}(\text{Cone}(\alpha))$ factors the composite morphism $\mathfrak{K}_a \circ F \rightarrow G \rightarrow \text{Cone}(\alpha)$ and thus it is zero. \square

Theorem 4.3. *Let $(F_n)_{n \in \mathbb{N}}$ be a sequence of objects in $\mathcal{D}(\mathbf{k}_X)$ and assume that F_n and F_{n+1} are a_n -isomorphic with $\sum_n a_n < \infty$. Set $a_{\geq n} := \sum_{k \geq n} a_k$. Then there exists an object $F_\infty \in \mathcal{D}(\mathbf{k}_X)$ such that (F_n, F_∞) is weakly $(a_{\geq n}, 12a_{\geq n})$ -isomorphic for any $n \in \mathbb{N}$. In particular, $d_{\mathfrak{K}}(F_n, F_\infty) \rightarrow 0$ ($n \rightarrow \infty$).*

Proof. Set $G_n := \mathfrak{K}_{a_{\geq n}} \circ F_n$. Then we have a morphism $\alpha_{n,m}: G_n \rightarrow G_m$ for $n \leq m$ and get an inductive system $(G_n)_{n \in \mathbb{N}}$. Let $F_\infty := \text{hocolim}_n G_n \in \mathcal{D}(\mathbf{k}_X)$.

We fix n and consider the cone $C_{n,m}$ of the morphism $\alpha_{n,m}: G_n \rightarrow G_m$. By composing β_m 's, we obtain $\beta_{m,n}: G_m \rightarrow G_n$. Since $(\alpha_{n,m}, \beta_{m,n})$ gives an $a_{\geq n}$ -isomorphism between G_n and G_m , the cone $C_{n,m}$ is $6a_{\geq n}$ -torsion by Proposition 4.2. Consider the following commutative diagram with solid arrows:

$$\begin{array}{ccccccc}
G_n^{\oplus \mathbb{N}} & \longrightarrow & \bigoplus_{m \geq n} G_m & \longrightarrow & \bigoplus_{m \geq n} C_{n,m} & \longrightarrow & G_n^{\oplus \mathbb{N}}[1] \\
\downarrow & & \downarrow & & \downarrow \text{dotted} & & \downarrow \\
G_n^{\oplus \mathbb{N}} & \longrightarrow & \bigoplus_{m \geq n} G_m & \longrightarrow & \bigoplus_{m \geq n} C_{n,m} & \longrightarrow & G_n^{\oplus \mathbb{N}}[1] \\
\downarrow & & \downarrow & & \downarrow \text{dotted} & & \downarrow \\
\text{hocolim}_m G_n & \cdots \longrightarrow & \text{hocolim}_{m \geq n} G_m & \cdots \longrightarrow & H & \cdots \longrightarrow & \text{hocolim}_m G_n[1] \\
\downarrow & & \downarrow & & \downarrow \text{dotted} & & \downarrow \\
G_n^{\oplus \mathbb{N}}[1] & \longrightarrow & \bigoplus_{m \geq n} G_m[1] & \longrightarrow & \bigoplus_{m \geq n} C_{n,m}[1] & \longrightarrow & G_n^{\oplus \mathbb{N}}[2].
\end{array} \tag{4.8}$$

Then by [KS06, Exercise 10.6], the dotted arrows can be completed so that the right bottom square is anti-commutative, all the other squares are commutative, and all the rows and all the columns are exact triangles. Since $\bigoplus_{m \geq n} C_{n,m}$ is $6a_{\geq n}$ -torsion, H is $12a_{\geq n}$ -torsion. Noticing that $\text{hocolim}_m G_n \simeq G_n$ and $\text{hocolim}_{m \geq n} G_m \simeq F_\infty$, by Lemma 3.4 (G_n, F_∞) is weakly $(0, 12a_{\geq n})$ -isomorphic. Since we set $G_n = \mathfrak{K}_{a_{\geq n}} \circ F_n$, we find that (F_n, F_∞) is weakly $(a_{\geq n}, 12a_{\geq n})$ -isomorphic. \square

Remark 4.4. One can prove a similar completeness result in a more general setting, i.e., for a triangulated category with a persistence structure (cf. persistence triangulated category by Biran–Cornea–Zhang [BCZ21]). Here we do not go into details.

Corollary 4.5. *The derived category of sheaves $\mathcal{D}(\mathbf{k}_X)$ is complete with respect to the pseudo-distance $d_{\mathfrak{K}}$. In particular, the Tamarkin category $\mathcal{D}(X)$ is complete with respect to the pseudo-distance $d_{\mathcal{D}(X)}$.*

Proof. For the latter claim, it suffices to show that for a Cauchy sequence $(F_n)_{n \in \mathbb{N}}$ in the Tamarkin category $\mathcal{D}(X)$, the limit object F_∞ constructed in Theorem 4.3 is also in $\mathcal{D}(X)$. By construction and Lemma 3.7, after taking a subsequence, we have $F_\infty = \text{hocolim}_m T_{-a_{\geq n}} F_n$, where $(a_{\geq n})_{n \in \mathbb{N}}$ is as in Theorem 4.3. Set $G_n := T_{-a_{\geq n}} F_n$. Since the functor $\mathbf{k}_{M \times [0, \infty)} \star (*)$ is defined as the composite of left adjoint functors, it commutes with direct sums. Since each G_n is an object of the left orthogonal ${}^\perp \mathcal{D}_{\{\tau \leq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t}) = \mathcal{D}(X)$, in the following commutative diagram the first and the second vertical arrows are isomorphisms:

$$\begin{array}{ccccc} \mathbf{k}_{M \times [0, \infty)} \star \bigoplus_n G_n & \longrightarrow & \mathbf{k}_{M \times [0, \infty)} \star \bigoplus_n G_n & \longrightarrow & \mathbf{k}_{M \times [0, \infty)} \star F_\infty \xrightarrow{+1} \\ \downarrow \wr & & \downarrow \wr & & \downarrow \\ \bigoplus_n G_n & \longrightarrow & \bigoplus_n G_n & \longrightarrow & F_\infty \xrightarrow{+1} \end{array} \quad (4.9)$$

Thus, we have an isomorphism $\mathbf{k}_{M \times [0, \infty)} \star F_\infty \xrightarrow{\sim} F_\infty$ and find that $F_\infty \in \mathcal{D}(X)$. \square

Remark 4.6. As mentioned in Remark 3.3, $d_{\mathfrak{R}}$ and $\text{dist}_{\mathfrak{R}}$ are equivalent. Hence $\mathcal{D}(\mathbf{k}_X)$ is also complete with respect to $\text{dist}_{\mathfrak{R}}$.

5 Sheaf quantization of Hamiltonian diffeomorphisms and homeomorphisms

In this section, we give an energy estimate on sheaf quantization of Hamiltonian diffeomorphism without projecting it onto the Tamarkin category. We also construct an object in the derived category associated with a limit of Hamiltonian diffeomorphisms with respect to the Hofer metric.

5.1 Sheaf quantization of Hamiltonian diffeomorphisms

In this subsection, we investigate sheaf quantization of Hamiltonian diffeomorphisms. We keep the symbols M for a connected manifold without boundary and I for an open interval containing $[0, 1]$. First, we prove the following, an analog of Theorem 3.14 in the derived category $\mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ not in the Tamarkin category $\mathcal{D}(M^2)$.

Proposition 5.1. *Let $H: T^*M \times I \rightarrow \mathbb{R}$ be a compactly supported Hamiltonian function. Then, one has an inequality*

$$d_{M^2 \times \mathbb{R}_t}(K_1^0, K_1^H) \leq 2 \int_0^1 \|H_s\|_\infty ds \leq 2 \|H\|_{\text{osc}}. \quad (5.1)$$

Similarly to (3.29), for two compactly supported Hamiltonian functions $H, H': T^*M \times I \rightarrow \mathbb{R}$, we have

$$d_{M^2 \times \mathbb{R}_t}(K_1^H, K_1^{H'}) \leq 2 \|H - H'\|_{\text{osc}}. \quad (5.2)$$

To prove the theorem, we prepare some lemmas. Let X be a manifold and let $q: X \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ denote the projection.

Lemma 5.2. *Let $F \in \mathcal{D}_{\{\tau \leq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$ and $G \in \mathcal{D}_{\{\tau \geq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$. Then there exist $H, H' \in \mathcal{D}(\mathbf{k}_X)$ such that $F \star G \simeq H \boxtimes \mathbf{k}_{\mathbb{R}_t}$ and $\mathcal{H}om^*(F, G) \simeq H' \boxtimes \mathbf{k}_{\mathbb{R}_t}$.*

Proof. By Lemma 2.8, we find that $\text{SS}(F \star G) \subset \{\tau = 0\}$ and $\text{SS}(\mathcal{H}om^*(F, G)) \subset \{\tau = 0\}$, which imply the result. \square

For $a, b \in \mathbb{R}_{\geq 0}$, we set

$$D(a, b) := \bigcup_{-a \leq c \leq b} \{(\tau, \sigma) \in \mathbb{R}^2 \mid \sigma = c \cdot \tau\}. \quad (5.3)$$

Note that $D(a, b)$ is a closed cone in \mathbb{R}^2 , which is not necessarily convex. See Figure 5.1. We set $D^+(a, b) := D(a, b) \cap \{(\tau, \sigma) \mid \tau \geq 0\}$ and $D^-(a, b) := D(a, b) \cap \{(\tau, \sigma) \mid \tau \leq 0\}$.




Figure 5.1: $D(a, b)$

Lemma 5.3. Let $F \rightarrow G \rightarrow H \xrightarrow{+1}$ be an exact triangle in $\mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t})$ and $a, b \in \mathbb{R}_{\geq 0}$. Assume

- (1) $F \in \mathbf{D}_{\{\tau \leq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$ and F is a -torsion,
- (2) $G \in \mathbf{D}_{\{\tau \geq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$ and G is b -torsion.

Then H is $\max(a, b)$ -torsion.

Proof. By Lemma 5.2, there exists $H' \in \mathbf{D}(\mathbf{k}_X)$ such that $\mathcal{H}om^*(F, G) \simeq H' \boxtimes \mathbf{k}_{\mathbb{R}}$. Moreover by the isomorphism $T_b \mathcal{H}om^*(F, G) \simeq \mathcal{H}om^*(F, T_b G)$, we find that $\mathcal{H}om^*(F, G)$ is b -torsion. Hence we have $\mathcal{H}om^*(F, G) \simeq 0$ and

$$\begin{aligned} \text{RHom}(F, G) &\simeq \text{RHom}(\mathbf{k}_{X \times \{0\}} \star F, G) \\ &\simeq \text{RHom}(\mathbf{k}_{X \times \{0\}}, \mathcal{H}om^*(F, G)) \simeq 0. \end{aligned} \quad (5.4)$$

Thus $H \simeq G \oplus F[1]$, which proves the result. \square

Next, we give a microlocal cut-off result, which we use to reduce the problem to Proposition 3.9.

Lemma 5.4. Let $\mathcal{H} \in \mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t \times \mathbb{R}})$ and assume that there exist $a, b \in \mathbb{R}_{\geq 0}$ such that

$$\text{SS}(\mathcal{H}) \subset T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times D(a, b). \quad (5.5)$$

Then there exists an exact triangle $\mathcal{H}^- \rightarrow \mathcal{H}^+ \rightarrow \mathcal{H} \xrightarrow{+1}$ in $\mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t \times \mathbb{R}})$ such that $\text{SS}(\mathcal{H}^-) \subset T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times D^-(a, b)$ and $\text{SS}(\mathcal{H}^+) \subset T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times D^+(a, b)$.

Proof. Let $\lambda := \{(\tau, \sigma) \mid \tau \geq 0, \sigma = 0\}$. Then we get an exact triangle

$$\mathbf{k}_{X \times (\lambda \setminus \{0\})} \star \mathcal{H} \longrightarrow \mathbf{k}_{X \times \lambda} \star \mathcal{H} \longrightarrow \mathcal{H} \xrightarrow{+1} \quad (5.6)$$

with $\text{SS}(\mathbf{k}_{X \times \lambda} \star \mathcal{H}) \subset T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times D(a, b) \cap \{(\tau, \sigma) \mid \tau \geq 0\}$. Moreover $\mathbf{k}_{X \times \lambda} \star F \rightarrow F$ is an isomorphism on $T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times \text{Int}(\lambda^\circ)$. Thus, we conclude that $\text{SS}(\mathbf{k}_{\lambda \setminus \{0\}} \star \mathcal{H}) \subset T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times (D(a, b) \setminus \text{Int}(\lambda^\circ)) = T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times D^-(a, b)$. \square

The following is a variant of [AI20, Prop. 4.3].

Proposition 5.5. *Let $\mathcal{H} \in \mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t \times I})$ and $s_1 < s_2 \in I$. Assume that there exist $a, b \in \mathbb{R}_{\geq 0}$ and $r \in \mathbb{R}_{> 0}$ such that*

$$\mathrm{SS}(\mathcal{H}) \cap \pi^{-1}(X \times \mathbb{R}_t \times (s_1 - r, s_2 + r)) \subset T^*X \times (\mathbb{R}_t \times I) \times D(a, b). \quad (5.7)$$

- (i) *Let $q: X \times \mathbb{R}_t \times I \rightarrow X \times \mathbb{R}_t$ be the projection. Then $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}$ and $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times [s_1, s_2)}$ are $(\max(a, b)(s_2 - s_1) + \varepsilon)$ -torsion for any $\varepsilon \in \mathbb{R}_{> 0}$.*
- (ii) *One has $d_{X \times \mathbb{R}_t}(\mathcal{H}|_{X \times \mathbb{R}_t \times \{s_1\}}, \mathcal{H}|_{X \times \mathbb{R}_t \times \{s_2\}}) \leq 2 \max(a, b)(s_2 - s_1)$.*

Proof. (i) Choose a diffeomorphism $\varphi: (s_1 - r, s_2 + r) \xrightarrow{\sim} \mathbb{R}$ satisfying $\varphi|_{[s_1, s_2]} = \mathrm{id}_{[s_1, s_2]}$ and $d\varphi(s) \geq 1$ for any $s \in (s_1 - r, s_2 + r)$. Set $\Phi := \mathrm{id}_{X \times \mathbb{R}_t} \times \varphi: X \times \mathbb{R}_t \times (s_1 - r, s_2 + r) \xrightarrow{\sim} X \times \mathbb{R}_t \times \mathbb{R}$ and $\mathcal{H}' := \Phi_*\mathcal{H} \in \mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t \times \mathbb{R}})$. Then by the assumption on φ , we have

$$\mathrm{SS}(\mathcal{H}') \subset T^*X \times (\mathbb{R}_t \times \mathbb{R}) \times D(a, b) \quad (5.8)$$

and $\mathcal{H}'|_{X \times \mathbb{R}_t \times [s_1, s_2]} \simeq \mathcal{H}|_{X \times \mathbb{R}_t \times [s_1, s_2]}$. Hence, we may assume $I = \mathbb{R}$ from the beginning.

Applying Lemma 5.4, we have an exact triangle $\mathcal{H}^- \rightarrow \mathcal{H}^+ \rightarrow \mathcal{H} \xrightarrow{+1}$ in $\mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t \times \mathbb{R}})$ with $\mathrm{SS}(\mathcal{H}^-) \subset T^*X \times (\mathbb{R} \times \mathbb{R}) \times D^-(a, b)$ and $\mathrm{SS}(\mathcal{H}^+) \subset T^*X \times (\mathbb{R} \times \mathbb{R}) \times D^+(a, b)$. By Proposition 3.9, $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}^+$ is $(b(s_2 - s_1) + \varepsilon)$ -torsion. Similarly we find that $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}^-$ is $(a(s_2 - s_1) + \varepsilon)$ -torsion. Here we have an exact triangle

$$\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}^- \longrightarrow \mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}^+ \longrightarrow \mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]} \xrightarrow{+1} \quad (5.9)$$

with $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}^+ \in \mathbf{D}_{\{\tau \geq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$ and $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}^- \in \mathbf{D}_{\{\tau \leq 0\}}(\mathbf{k}_{X \times \mathbb{R}_t})$. Hence, by Lemma 5.3 we find that $\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]}$ is $(\max(a, b)(s_2 - s_1) + \varepsilon)$ -torsion. The proof for the other case is similar.

(ii) We have the following two exact triangles

$$\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times (s_1, s_2]} \longrightarrow \mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times [s_1, s_2]} \longrightarrow \mathcal{H}|_{X \times \mathbb{R}_t \times \{s_1\}} \xrightarrow{+1}, \quad (5.10)$$

$$\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times [s_1, s_2]} \longrightarrow \mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times [s_1, s_2]} \longrightarrow \mathcal{H}|_{X \times \mathbb{R}_t \times \{s_2\}} \xrightarrow{+1}. \quad (5.11)$$

Hence, by the result of (i) and Lemma 3.4, the two pairs $(\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times [s_1, s_2]}, \mathcal{H}|_{X \times \mathbb{R}_t \times \{s_1\}})$ and $(\mathrm{R}q_*\mathcal{H}_{X \times \mathbb{R}_t \times [s_1, s_2]}, \mathcal{H}|_{X \times \mathbb{R}_t \times \{s_2\}})$ are weakly $(0, (\max(a, b)(s_2 - s_1) + \varepsilon))$ -isomorphic. Hence, the result follows from the triangle inequality. \square

The following theorem is a variant of Proposition 3.9.

Proposition 5.6. *Let I be an open interval containing the closed interval $[0, 1]$ and $\mathcal{H} \in \mathbf{D}(\mathbf{k}_{X \times \mathbb{R}_t \times I})$. Assume that there exist continuous functions $f, g: I \rightarrow \mathbb{R}_{\geq 0}$ satisfying*

$$\mathrm{SS}(\mathcal{H}) \subset T^*X \times \{(t, s; \tau, \sigma) \mid (\tau, \sigma) \in D(f(s), g(s))\}. \quad (5.12)$$

Then $d_{X \times \mathbb{R}_t}(\mathcal{H}|_{X \times \mathbb{R}_t \times \{0\}}, \mathcal{H}|_{X \times \mathbb{R}_t \times \{1\}}) \leq 2 \int_0^1 \max(f, g)(s) ds$.

Proof. We can apply an argument similar to [AI20, Prop. 4.15]. We only need to replace [AI20, Prop. 4.3] with Proposition 5.5. \square

Proof of Proposition 5.1. The result follows from (3.25) and Proposition 5.6. \square

Now we prove that the restriction of the sheaf quantization of a Hamiltonian isotopy to $s = 1$ depends only on the time-1 map.

Proposition 5.7. *Let $H: T^*M \times I \rightarrow \mathbb{R}$ be a compactly supported Hamiltonian function. Then the objects $\tilde{K}_1^H, K_1^H, \mathcal{K}_1^H$ are determined by the time-1 map ϕ_1^H .*

Proof. We shall prove $K_1^H = K_1^{H'}$ assuming that $\phi_1^{H'} = \phi_1^H$. By (3.23), it suffices to show that $K_1^{\overline{H} \sharp H'} \simeq \mathbf{k}_{\Delta_M \times \{0\}}$. Hence, we may assume that $H' \equiv 0$ and $\phi_1^H = \text{id}_{T^*M}$.

Since $u_s \equiv 0$ by Lemma 3.11, we find that $\mathring{\text{SS}}(K_1^H) = \mathring{T}_{\Delta_M \times \{0\}}^*(M^2 \times \mathbb{R}_t)$ and K_1^H is simple along the subset. Moreover, since $K_0^H \simeq \mathbf{k}_{\Delta_M \times \{0\}}$ and u is compactly supported and hence bounded, $K^H|_{M^2 \times \{R\} \times I}$ and $K^H|_{M^2 \times \{-R\} \times I}$ are 0 for sufficiently large R . Hence, there exists a rank one local system L on M such that $K_1^H \simeq \delta_{M*} L \boxtimes \mathbf{k}_{\{0\}}[m]$, where m is some integer. By Proposition 5.1, we have

$$d_{M^2 \times \mathbb{R}_t}(\mathbf{k}_{\Delta_M \times \{0\}}, \delta_{M*} L \boxtimes \mathbf{k}_{\{0\}}[m]) = d_{M^2 \times \mathbb{R}_t}(K_1^0, K_1^H) \leq 2\|H\|_{\text{osc}} < \infty. \quad (5.13)$$

By restricting to $\{(x, x)\} \times \mathbb{R} \subset M^2 \times \mathbb{R}_t$ for some $x \in M$, we obtain $d_{\mathbb{R}_t}(\mathbf{k}_{\{0\}}, \mathbf{k}_{\{0\}}[m]) < \infty$ and find that $m = 0$. Then, we have

$$\text{R}\Gamma(M; L) \simeq \text{R}\Gamma(M^2 \times \mathbb{R}_t; K_1^H) \simeq \text{R}\Gamma(M^2 \times \mathbb{R}_t; K_1^0) \simeq \text{R}\Gamma(M; \mathbf{k}_M). \quad (5.14)$$

In particular, $H^0(M; L) \simeq H^0(M; \mathbf{k}_M)$, which implies that L is trivial. \square

The proposition above shows the well-definedness in the following definition.

Definition 5.8. (i) A diffeomorphism $\varphi: T^*M \rightarrow T^*M$ is said to be a compactly supported *Hamiltonian diffeomorphism* if it is the time-1 map of some compactly supported Hamiltonian isotopy ϕ^H , that is $\varphi = \phi_1^H$. The set of compactly supported Hamiltonian diffeomorphisms is denoted by $\text{Ham}_c(T^*M, \omega)$.

(ii) The *Hofer metric* between Hamiltonian diffeomorphisms is defined by

$$d_H(\varphi, \varphi') := \inf \{ \|H\|_{\text{osc}} \mid \phi_1^H = \varphi^{-1}\varphi' \} \quad (5.15)$$

for $\varphi, \varphi' \in \text{Ham}_c(T^*M, \omega)$.

(iii) For $\varphi \in \text{Ham}_c(T^*M, \omega)$, one sets $\tilde{K}^\varphi := \tilde{K}_1^H, K^\varphi := K_1^H$, and $\mathcal{K}^\varphi := \mathcal{K}_1^H$, where H is any compactly supported function with $\varphi = \phi_1^H$.

By (3.29) and (5.2), we have

$$\begin{aligned} d_{\mathcal{D}(M^2)}(\mathcal{K}^\varphi, \mathcal{K}^{\varphi'}) &\leq d_H(\varphi, \varphi'), \\ d_{M^2 \times \mathbb{R}_t}(K^\varphi, K^{\varphi'}) &\leq 2d_H(\varphi, \varphi'). \end{aligned} \quad (5.16)$$

An argument similar to Proposition 5.7 proves the following lemma.

Lemma 5.9. *Let $\varphi, \varphi' \in \text{Ham}_c(T^*M, \omega)$ be two compactly supported Hamiltonian diffeomorphisms and assume that $\varphi(0_M) = \varphi'(0_M)$. Then there exists some constant $c \in \mathbb{R}$ such that $\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \simeq T_c \mathcal{K}^{\varphi'} \bullet \mathbf{k}_{M \times [0, \infty)}$.*

Proof. It suffices to show that $\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \simeq \mathbf{k}_{M \times [c, \infty)}$ for some $c \in \mathbb{R}$ assuming that $\varphi(0_M) = 0_M$. By Lemma 3.11, we have

$$\mathring{\text{SS}}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}) = 0_M \times \{(c; \tau) \mid \tau > 0\} \quad (5.17)$$

for some constant $c \in \mathbb{R}$. Hence, there exist a rank one local system L on M and some integer $d \in \mathbb{Z}$ such that $\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \simeq L \boxtimes \mathbf{k}_{[c, \infty)}[m]$. By Theorem 3.14, we have

$$\begin{aligned} & d_{\mathcal{D}(M)}(\mathbf{k}_{\Delta_M \times [0, \infty)}, L \boxtimes \mathbf{k}_{[c, \infty)}[m]) \\ &= d_{\mathcal{D}(M)}(\mathcal{K}_1^0 \bullet \mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}) \\ &\leq \|H\|_{\text{osc}} < \infty. \end{aligned} \tag{5.18}$$

Hence, we find that $m = 0$ and L is trivial. \square

5.2 Sheaf quantization of Hamiltonian homeomorphisms

In this subsection, we construct a sheaf quantization of a limit of Hamiltonian diffeomorphisms with respect to the Hofer metric.

Proposition 5.10. *Let $(\varphi_n)_{n \in \mathbb{N}} \subset \text{Ham}_c(T^*M, \omega)$ be a sequence of compactly supported Hamiltonian diffeomorphisms and $K_n := K^{\varphi_n} \in \mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ the sheaf quantization of φ_n . Assume that it is a Cauchy sequence with respect to the Hofer metric d_H . Then there exists an object $K_\infty \in \mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ such that $d_{M^2 \times \mathbb{R}_t}(K_n, K_\infty) \rightarrow 0$ ($n \rightarrow \infty$). Moreover, such an object is unique up to isomorphism, and the endofunctor $K_\infty \bullet (*)$ on $\mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ gives an equivalence of categories.*

Proof. By (5.2), we have

$$d_{M^2 \times \mathbb{R}_t}(K_n, K_m) \leq 2d_H(\varphi_n, \varphi_m), \tag{5.19}$$

which proves that $(K_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $\mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ with respect to $d_{M^2 \times \mathbb{R}_t}$. Hence, Corollary 4.5 shows the existence of a limit object.

Let K'_∞ be another limit object that satisfies $d_{M^2 \times \mathbb{R}_t}(K_\infty, K'_\infty) = 0$. There exists $\overline{K}_n \in \mathcal{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ such that $\overline{K}_n \bullet K_n \simeq K_n \bullet \overline{K}_n \simeq \mathbf{k}_{\Delta_M \times \{0\}}$ for each n . Then we find that the sequence $(\overline{K}_n)_{n \in \mathbb{N}}$ is also Cauchy, and we can take a limit object \overline{K}_∞ . The Cauchy sequence $(\overline{K}_n \bullet K_n)_{n \in \mathbb{N}}$ converges to both $\overline{K}_\infty \bullet K_\infty$ and $\mathbf{k}_{\Delta_M \times \{0\}}$. Hence we have $d_{M^2 \times \mathbb{R}_t}(\overline{K}_\infty \bullet K_\infty, \mathbf{k}_{\Delta_M \times \{0\}}) = 0$. Similarly, we have $d_{M^2 \times \mathbb{R}_t}(K'_\infty \bullet \overline{K}_\infty, \mathbf{k}_{\Delta_M \times \{0\}}) = 0$. By Lemma 5.14 below, we have $\overline{K}_\infty \bullet K_\infty \simeq \mathbf{k}_{\Delta_M \times \{0\}} \simeq K'_\infty \bullet \overline{K}_\infty$. Hence, we conclude that $K'_\infty \simeq K'_\infty \bullet \overline{K}_\infty \bullet K_\infty \simeq K_\infty$ and $\overline{K}_\infty \bullet (*)$ gives the inverse. \square

Note that we have an inequality similar to (3.11) for $d_{X \times \mathbb{R}_t}$.

Lemma 5.11. *Let $F, G \in \mathcal{D}_{\{\tau \geq 0\}}(\mathbf{k}_{\mathbb{R}_t})$. If $d_{\mathbb{R}_t}(F, G) = 0$, then $\mathring{\text{SS}}(F) = \mathring{\text{SS}}(G)$.*

Proof. We prove $\mathring{\text{SS}}(F) \subset \mathring{\text{SS}}(G)$ by contradiction. Choose $(t_0; 1) \in \mathring{\text{SS}}(F) \setminus \mathring{\text{SS}}(G)$. There exists a neighborhood (a, b) of t_0 in \mathbb{R}_t such that $(a, b) \cap \pi(\mathring{\text{SS}}(G)) = \emptyset$. There also exist $t_1, t_2 \in \mathbb{R}$ such that $a < t_1 < t_2 < b$ and the restriction map $\text{R}\Gamma((-\infty, t_2); F) \rightarrow \text{R}\Gamma((-\infty, t_1); F)$ is not an isomorphism. We may choose t_1 and t_2 arbitrarily close to t_0 . On the other hand, $\text{R}\Gamma((-\infty, t''); G) \rightarrow \text{R}\Gamma((-\infty, t'); G)$ is an isomorphism for any $a < t' < t'' < b$ by the microlocal Morse lemma. Thus, an interleaving between $(\text{R}\Gamma((-\infty, t); F))_{t \in \mathbb{R}}$ and $(\text{R}\Gamma((-\infty, t); G))_{t \in \mathbb{R}}$ leads to a contradiction. \square

Lemma 5.12. *Let $F, G \in \mathcal{D}(\mathbf{k}_{\mathbb{R}_t})$ and assume that $d_{\mathbb{R}_t}(F, G) = 0$. Then $\text{SS}(F) = \text{SS}(G)$ and hence $\text{Supp}(F) = \text{Supp}(G)$.*

Proof. Since $d_{\mathbb{R}_t}(F \star \mathbf{k}_{[0,\infty)}, G \star \mathbf{k}_{[0,\infty)}) = 0$, we have

$$\mathrm{SS}(F) \cap \{\tau > 0\} = \mathring{\mathrm{SS}}(F \star \mathbf{k}_{[0,\infty)}) = \mathring{\mathrm{SS}}(G \star \mathbf{k}_{[0,\infty)}) = \mathrm{SS}(G) \cap \{\tau > 0\} \quad (5.20)$$

by Lemma 5.11. Similarly, we obtain

$$\mathrm{SS}(F) \cap \{\tau < 0\} = \mathring{\mathrm{SS}}(F \star \mathbf{k}_{(0,\infty)}) = \mathring{\mathrm{SS}}(G \star \mathbf{k}_{(0,\infty)}) = \mathrm{SS}(G) \cap \{\tau < 0\} \quad (5.21)$$

and hence $\mathring{\mathrm{SS}}(F) = \mathring{\mathrm{SS}}(G)$. Note that $U := \mathbb{R}_t \setminus \pi(\mathring{\mathrm{SS}}(F)) = \mathbb{R}_t \setminus \pi(\mathring{\mathrm{SS}}(G))$ is an open subset of \mathbb{R}_t , and both $F|_U$ and $G|_U$ are locally constant. Thus $d_{X \times \mathbb{R}_t}(F, G) = 0$ implies $F|_U \simeq G|_U$. \square

Lemma 5.13. *For $F, G \in \mathrm{D}(\mathbf{k}_{X \times \mathbb{R}_t})$, $d_{X \times \mathbb{R}_t}(F, G) = 0$ implies $\mathrm{Supp}(F) = \mathrm{Supp}(G)$.*

Proof. Let $(x, t) \in X \times \mathbb{R}_t \setminus \mathrm{Supp}(G)$. Take an open neighborhood $x \in U$ in X and $\varepsilon > 0$ so that $U \times (t - \varepsilon, t + \varepsilon) \subset X \times \mathbb{R}_t \setminus \mathrm{Supp}(G)$. Note that for any $y \in X$, $0 \leq d_{\mathbb{R}_t}(F|_{\{y\} \times \mathbb{R}_t}, G|_{\{y\} \times \mathbb{R}_t}) \leq d_{X \times \mathbb{R}_t}(F, G) = 0$ and $\mathrm{Supp}(G|_{\{y\} \times \mathbb{R}_t}) \subset \mathrm{Supp}(G) \cap (\{y\} \times \mathbb{R}_t)$. By Lemma 5.12, $F|_{\{y\} \times (t - \varepsilon, t + \varepsilon)} \simeq 0$ for any $y \in U$. Hence, we obtain $F|_{U \times (t - \varepsilon, t + \varepsilon)} \simeq 0$ and $\mathrm{Supp}(F) \subset \mathrm{Supp}(G)$. We obtain the converse inclusion $\mathrm{Supp}(F) \supset \mathrm{Supp}(G)$ similarly. \square

Lemma 5.14. *Let $F, G \in \mathrm{D}(\mathbf{k}_{X \times \mathbb{R}_t})$ and assume that $d_{X \times \mathbb{R}_t}(F, G) = 0$ and $\mathrm{Supp}(G) \subset X \times \{0\}$. Then $F \simeq G$.*

Proof. By Lemma 5.13, we have $\mathrm{Supp}(F) = \mathrm{Supp}(G) \subset X \times \{0\}$. We may write $F \simeq F' \boxtimes \mathbf{k}_{\{0\}}$ and $G \simeq G' \boxtimes \mathbf{k}_{\{0\}}$ with some $F', G' \in \mathrm{D}(\mathbf{k}_X)$. Take $\varepsilon > 0$ arbitrarily. Note that $\mathfrak{R}_\varepsilon \circ F \simeq F' \boxtimes \mathbf{k}_{[-\varepsilon, \varepsilon]}$ and $\mathfrak{R}_\varepsilon \circ G \simeq G' \boxtimes \mathbf{k}_{[-\varepsilon, \varepsilon]}$. Hence, there exist morphisms $\alpha: F' \boxtimes \mathbf{k}_{[-\varepsilon, \varepsilon]} \rightarrow G' \boxtimes \mathbf{k}_{\{0\}}$ and $\beta: G' \boxtimes \mathbf{k}_{[-\varepsilon, \varepsilon]} \rightarrow F' \boxtimes \mathbf{k}_{\{0\}}$ such that (α, β) is a ε -isomorphism of (F, G) . Restricting α and β on $X \times \{0\}$, we obtain isomorphisms between F' and G' . \square

Note that Petit–Schapira–Waas [PSW21] proved that $\mathrm{dist}(F, G) = 0$ if and only if $F \simeq G$ when F and G are constructible sheaves up to infinity on a real analytic manifold. This result, as well as its proof, is different from ours and is not related to the Tamarkin category.

Proposition 5.10 shows that we can associate a sheaf with an element of the metric completion of $\mathrm{Ham}_c(T^*M, \omega)$ with respect to the Hofer metric d_H as follows. Let $(\varphi_n)_{n \in \mathbb{N}}$ be a Cauchy sequence of Hamiltonian diffeomorphisms and $K_n := K^{\varphi_n}$ for $n \in \mathbb{N}$. By Proposition 5.10, we obtain a limit object K_∞ of the sequence $(K_n)_{n \in \mathbb{N}}$. The argument in the proof of the proposition also shows that another Cauchy sequence equivalent to $(\varphi_n)_{n \in \mathbb{N}}$ gives the same limit object K_∞ up to isomorphism.

Definition 5.15. Let $[(\varphi_n)_{n \in \mathbb{N}}]$ be an element of the completion of $\mathrm{Ham}_c(T^*M, \omega)$ with respect to d_H . The limit sheaf K_∞ defined as above is denoted by $K^{[(\varphi_n)_n]}$ and called the sheaf quantization of $[(\varphi_n)_{n \in \mathbb{N}}]$.

We use the above construction to obtain a sheaf quantization of a Hamiltonian homeomorphism of T^*M .

Definition 5.16 (Oh–Müller [OM07]). Let $\phi = (\phi_s)_s: T^*M \times I \rightarrow T^*M$ be an isotopy of homeomorphisms of T^*M . The isotopy ϕ is said to be a *continuous Hamiltonian isotopy* if there exists a compact subset $C \subset T^*M$ and a sequence of smooth Hamiltonian functions $H_n: T^*M \times I \rightarrow \mathbb{R}$ supported in C satisfying the following two conditions.

- (1) The sequence of flows $(\phi^{H_n})_{n \in \mathbb{N}}$ C^0 -converges to ϕ , uniformly in $s \in I$.

- (2) The sequence $(H_n)_{n \in \mathbb{N}}$ converges uniformly to a continuous function $H: T^*M \times I \rightarrow \mathbb{R}$. That is, $\|H_n - H\|_\infty \rightarrow 0$.

In this case, H is said to generate ϕ . A homeomorphism of T^*M is called a *Hamiltonian homeomorphism* if it is the time-1 map of a continuous Hamiltonian isotopy.

The following uniqueness theorems hold.

- (i) A continuous Hamiltonian function generates a unique continuous Hamiltonian isotopy (Oh–Müller [OM07]).
- (ii) A continuous Hamiltonian isotopy can be generated by a unique continuous Hamiltonian function up to addition of a function of time (Viterbo [Vit06]).

A continuous Hamiltonian isotopy ϕ defines an element of the metric completion of $\text{Ham}_c(T^*M, \omega)$ with respect to d_H . Indeed, for a sequence $(H_n)_{n \in \mathbb{N}}$ satisfying the condition (2) in Definition 5.16, $(\phi_1^{H_n})_{n \in \mathbb{N}}$ forms a Cauchy sequence with respect to the Hofer metric d_H . Moreover, the element in the metric completion is independent of the choice of a sequence $(H_n)_{n \in \mathbb{N}}$. Thus we obtain a sheaf K_∞ as in Definition 5.15. We give a bound of the microsupport of K_∞ . Set $\varphi_n = \phi_1^{H_n}$ and $K_n := K^{\varphi_n}$. By construction, after taking a subsequence of $(K_n)_{n \in \mathbb{N}}$ if necessary, $K_\infty \simeq \text{hocolim}_n \mathbf{k}_{M \times [-a_{\geq n}, a_{\geq n}]} \star K_n$, where $(a_{\geq n})_{n \in \mathbb{N}}$ is as in Theorem 4.3. Hence, we have

$$\mathring{\text{SS}}(K_\infty) \subset \left\{ ((x'; \xi'), (x; -\xi), (t; \tau)) \left| \begin{array}{l} (x; \xi) \in T^*M, \\ (x'; \xi'/\tau) = \varphi_\infty(x; \xi/\tau) \end{array} \right. \right\} \quad (5.22)$$

by the condition (1) and Lemma 2.9.

Remark 5.17. For the microsupport estimate above, we do not need the full convergence of flows, but only the convergence of the time-1 maps, that is, $\varphi_n \rightarrow \varphi_\infty$ ($n \rightarrow \infty$).

Definition 5.18. Let ϕ be a continuous Hamiltonian isotopy associated with a continuous function $H: T^*M \times I \rightarrow \mathbb{R}$ and $\varphi_\infty := \phi_1$ a Hamiltonian homeomorphism. The limit sheaf K_∞ defined as above is denoted by $K_1^H = K^{\varphi_\infty}$ and called the sheaf quantization of the Hamiltonian homeomorphism φ_∞ .

We shall prove that the object K_1^H depends only on the time-1 map ϕ_1 , which justifies the notation K^{φ_∞} . It suffices to show that if $\phi_1 = \text{id}_{T^*M}$ then $K_1^H \simeq \mathbf{k}_{\Delta_M \times \{0\}}$.

Lemma 5.19. *Let $G \in \text{D}(\mathbf{k}_{\mathbb{R}_t})$ with $d_{\mathbb{R}_t}(G, \mathbf{k}_{\{0\}}) < \infty$. Assume that there exists $\varepsilon > 0$ such that for any $a < b$ with $b - a < \varepsilon$, one has $G \star \mathbf{k}_{[a,b]} \simeq \mathbf{k}_{[a,b]}$ and $G \star \mathbf{k}_{(a,b)} \simeq \mathbf{k}_{(a,b)}$. Then, $G \simeq \mathbf{k}_{\{0\}}$.*

Proof. First, we prove $\mathring{\text{SS}}(G \star \mathbf{k}_{[0,\infty)}) \subset \{(0; \tau) \mid \tau > 0\}$ by contradiction. Note that we already know that $\text{SS}(G \star \mathbf{k}_{[0,\infty)}) \subset \{\tau \geq 0\}$ by Lemma 2.8. Assume that there exists $(t_0; 1) \in \mathring{\text{SS}}(G \star \mathbf{k}_{[0,\infty)})$ with $t_0 \neq 0$. Then, there exist $t_1, t_2 \in \mathbb{R}$ such that $t_1 < t_2$ and the restriction map $\text{R}\Gamma((-\infty, t_2); G \star \mathbf{k}_{[0,\infty)}) \rightarrow \text{R}\Gamma((-\infty, t_1); G \star \mathbf{k}_{[0,\infty)})$ is not an isomorphism. We may choose t_1 and t_2 arbitrary close to t_0 and hence may assume $t_2 - t_1 < \min\{t_1, \varepsilon\}$ if $t_0 > 0$ and $t_2 < 0, t_2 - t_1 < \varepsilon$ if $t_0 < 0$. Applying $G \star (*)$ to the exact triangle $\mathbf{k}_{[0, t_2 - t_1]} \rightarrow \mathbf{k}_{[0, \infty)} \rightarrow \mathbf{k}_{[t_2 - t_1, \infty)} \xrightarrow{+1}$, we have an exact triangle

$$\mathbf{k}_{[0, t_2 - t_1]} \rightarrow G \star \mathbf{k}_{[0, \infty)} \rightarrow T_{t_2 - t_1} G \star \mathbf{k}_{[0, \infty)} \xrightarrow{+1}. \quad (5.23)$$

Since $\mathrm{R}\Gamma((-\infty, t_2); \mathbf{k}_{[0, t_2 - t_1]}) = 0$, we get an isomorphism $\mathrm{R}\Gamma((-\infty, t_2); G \star \mathbf{k}_{[0, \infty)}) \xrightarrow{\sim} \mathrm{R}\Gamma((-\infty, t_1); G \star \mathbf{k}_{[0, \infty)})$. The morphism coincides with the restriction map by the construction, which is a contradiction.

Similarly, we obtain $\mathring{\mathrm{S}}\mathrm{S}(G \star \mathbf{k}_{(0, \infty)}) \subset \{(0; \tau) \mid \tau < 0\}$.

We get $\mathring{\mathrm{S}}\mathrm{S}(G) \subset T_0^* \mathbb{R}_t$ by applying $G \star (*)$ to the exact triangle $\mathbf{k}_{(0, \infty)} \rightarrow \mathbf{k}_{[0, \infty)} \rightarrow \mathbf{k}_{\{0\}} \xrightarrow{+1}$. Hence, we obtain the desired isomorphism from $d_{\mathbb{R}_t}(G, \mathbf{k}_{\{0\}}) < \infty$. \square

Now let ϕ be a continuous Hamiltonian isotopy generated by a continuous function $H: T^*M \times I \rightarrow \mathbb{R}$ and assume that $\phi_1 = \mathrm{id}_{T^*M}$. By the microsupport estimate (5.22), we get $\pi(\mathring{\mathrm{S}}\mathrm{S}(K_1^H)) \subset \Delta_M \times \mathbb{R}$. By construction, $d_{M^2 \times \mathbb{R}_t}(\mathbf{k}_{\Delta_M \times \{0\}}, K_1^H) < \infty$. Restricting to $(M^2 \setminus \Delta_M) \times \mathbb{R}_t$, we find that $d_{(M^2 \setminus \Delta_M) \times \mathbb{R}_t}(0, K_1^H|_{(M^2 \setminus \Delta_M) \times \mathbb{R}_t}) < \infty$, which implies $K_1^H|_{(M^2 \setminus \Delta_M) \times \mathbb{R}_t} \simeq 0$. Hence, we can write $K_1^H \simeq (\delta_M \times \mathrm{id}_{\mathbb{R}_t})_* K'$ with $K' \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$. Again by the estimate (5.22) and Proposition 2.3(i), we find that $\mathring{\mathrm{S}}\mathrm{S}(K') \subset 0_M \times T^* \mathbb{R}_t$.

Let us keep the notation of a compact set $C \subset T^*M$ and $K_n = K^{\varphi_n}$ as above (see also Definition 5.16). For $F \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ such that

$$\mathring{\mathrm{S}}\mathrm{S}(F) \cap \{(x, t; \xi, \tau) \mid \tau \neq 0, (x, ; \xi/\tau) \in C\} = \emptyset, \quad (5.24)$$

we have $K_n \bullet F \simeq F$ for any $n \in \mathbb{N}$, which implies $d_{M \times \mathbb{R}_t}(K_1^H \bullet F, F) = 0$. For each $x_0 \in M$ and $a < b \in \mathbb{R}$ with sufficiently small $b - a$, there exists $F \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ such that (1) its microsupport does not intersect the cone $\{(x, t; \xi, \tau) \mid \tau \neq 0, (x, ; \xi/\tau) \in C\}$ of C and (2) $F|_{\{x_0\} \times \mathbb{R}_t} \simeq \mathbf{k}_{[a, b]}$. For example, such F is obtained as the sheaf quantization of an exact Lagrangian immersion of a sphere. Therefore, $d_{\mathbb{R}_t}(K'|_{\{x_0\} \times \mathbb{R}_t} \star \mathbf{k}_{[a, b]}, \mathbf{k}_{[a, b]}) \leq d_{M \times \mathbb{R}_t}(K_1^H \bullet F, F) = 0$ if $b - a$ is sufficiently small. Applying Lemma 5.12 to $K'|_{\{x_0\} \times \mathbb{R}_t} \star \mathbf{k}_{[a, b]}$ and $\mathbf{k}_{[a, b]}$, we get $K'|_{\{x_0\} \times \mathbb{R}_t} \star \mathbf{k}_{[a, b]} \simeq \mathbf{k}_{[a, b]}$. Similarly, we obtain $K'|_{\{x_0\} \times \mathbb{R}_t} \star \mathbf{k}_{(a, b]} \simeq \mathbf{k}_{(a, b]}$ for any $x_0 \in M$ and sufficiently small $b - a$.

Hence, for any $x_0 \in M$, $K'|_{\{x_0\} \times \mathbb{R}_t} \in \mathcal{D}(\mathbf{k}_{\mathbb{R}_t})$ satisfies the condition of Lemma 5.19. Combining this with the estimate of $\mathring{\mathrm{S}}\mathrm{S}(K')$, we may write $K' = L \boxtimes \mathbf{k}_{\{0\}}$, where L is a rank one local system. Again by the fact that $d_{M^2 \times \mathbb{R}_t}(\mathbf{k}_{\Delta_M \times \{0\}}, K_1^H) < \infty$, we conclude that $L = \mathbf{k}_M$, which proves $K_1^H \simeq \mathbf{k}_{\Delta_M \times \{0\}}$.

6 Spectral invariants in Tamarkin category

In this section, we define spectral invariants for an object of the Tamarkin category and develop Lusternik–Schnirelmann theory. Most of the definitions and the results were announced to appear in Humilière–Vichery [HV].

Definition 6.1. Let $F \in \mathcal{D}(M)$.

(i) For $n \in \mathbb{Z}$, one defines

$$Q_c^n(F) := \mathrm{Hom}(\mathbf{k}_{M \times [0, \infty)}, T_c F[n]) \quad (6.1)$$

$$\simeq H^n \mathrm{RHom}(\mathbf{k}_{M \times [-c, \infty)}, F),$$

$$Q_\infty^n(F) := \varinjlim_{c \rightarrow \infty} Q_c^n(F). \quad (6.2)$$

Set $Q_c^*(F) := \bigoplus_{n \in \mathbb{Z}} Q_c^n(F)$ and similarly for $Q_\infty^*(F)$. Denote the canonical map by $i_c: Q_c^*(F) \rightarrow Q_\infty^*(F)$.

(ii) For $\alpha \in Q_\infty^*(F) \setminus \{0\}$, one defines

$$c(\alpha; F) := \inf\{c \in \mathbb{R} \mid \alpha \in \text{Im } i_c\} \quad (6.3)$$

and calls it the *spectral invariant* of F for α .

(iii) One defines

$$\text{Spec}(F) := \{c(\alpha; F) \in \mathbb{R} \mid \alpha \in Q_\infty^*(F) \setminus \{0\}\}. \quad (6.4)$$

Let $t: M \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ be the projection and let Γ_{dt} denote the graph of the 1-form dt :

$$\Gamma_{dt} := \{(x, t; 0, 1) \mid (x, t) \in M \times \mathbb{R}_t\}. \quad (6.5)$$

Note that

$$\text{Spec}(F) \subset \{-c \in \mathbb{R}_t \mid c \in t\pi(\text{SS}(F) \cap \Gamma_{dt})\}. \quad (6.6)$$

In order to state Lusternik–Schnirelmann theory for sheaves, we recall the algebraic counterpart of cup-length, which is studied in [AI22].

Definition 6.2. Let R be an associative (not necessarily commutative) non-unital ring over \mathbf{k} . For a right R -module A , one defines

$$\text{cl}_R(A) := \inf \left\{ k - 1 \mid \begin{array}{l} k \in \mathbb{N}, a_0 \cdot r_1 \cdots r_k = 0 \\ \text{for any } a_0 \in A \text{ and } (r_1, \dots, r_k) \in R^k \end{array} \right\} \in \mathbb{Z}_{\geq -1} \cup \{\infty\}. \quad (6.7)$$

We note that $\text{cl}_R(A) = -1$ if and only if $A = 0$. If there is no risk of confusion, we simply write $\text{cl}(A)$ for $\text{cl}_R(A)$. By definition, we have the following lemma.

Lemma 6.3. For an exact sequence of right R -modules $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$, one has

$$\text{cl}(B) \leq \text{cl}(A) + \text{cl}(C) + 1. \quad (6.8)$$

Let $F \in \mathcal{D}(M)$. Then, we have a right action of $\text{End}(\mathbf{k}_{M \times [0, \infty)}, \mathbf{k}_{M \times [0, \infty)}) \simeq H^*(M; \mathbf{k})$ on $Q_c^*(F)$, which induces an action on $Q_\infty^*(F)$. Hereafter we set $R := \bigoplus_{n \geq 1} H^n(M; \mathbf{k})$ and consider the cup-length over R .

Theorem 6.4. Assume that t is proper on $\text{Supp}(F)$. Let $F \in \mathcal{D}(M)$ and assume that there exists $c \ll 0$ satisfying $i_c = 0$. If $\#\text{Spec}(F) \leq \text{cl}(Q_\infty^*(F))$, then $\pi(\text{SS}(F) \cap \Gamma_{dt})$ is cohomologically non-trivial. That is, for any open neighborhood U of $\pi(\text{SS}(F) \cap \Gamma_{dt})$, the restriction map $\bigoplus_{n \geq 1} H^n(M; \mathbf{k}) \rightarrow \bigoplus_{n \geq 1} H^n(U; \mathbf{k})$ is non-zero.

For $F \in \mathcal{D}(M)$, if $F|_{M \times (c, \infty)}$ is locally constant for $c \gg 0$, then $i_c = 0$ for $c \ll 0$.

For the proof of the theorem, we prepare some notation. For $d \in \mathbb{R}$, we define

$$Q_{\infty, d}^*(F) := \text{Im}(i_d: Q_d^*(F) \rightarrow Q_\infty^*(F)) \subset Q_\infty^*(F). \quad (6.9)$$

Then we get the following properties:

- (1) If $d < d'$, then $Q_{\infty, d}^*(F) \subset Q_{\infty, d'}^*(F)$.
- (2) If $[d, d'] \cap \text{Spec}(F) = \emptyset$, then $Q_{\infty, d}^*(F) \simeq Q_{\infty, d'}^*(F)$.

(3) For $d < d'$, there exists an exact sequence of right $H^*(M; \mathbf{k})$ -modules

$$0 \rightarrow Q_{\infty, d}^*(F) \rightarrow Q_{\infty, d'}^*(F) \rightarrow Q_{\infty, d'}^*(F)/Q_{\infty, d}^*(F) \rightarrow 0. \quad (6.10)$$

Moreover, we have

$$\mathrm{cl}(H_{M \times [-d', -d]}^*(M \times \mathbb{R}; F)) \geq \mathrm{cl}(Q_{\infty, d'}^*(F)/Q_{\infty, d}^*(F)). \quad (6.11)$$

Proof of Theorem 6.4. If $Q_{\infty}^*(F) \simeq 0$, then $\mathrm{cl}(Q_{\infty}^*(F)) = -1$ and there is nothing to prove.

Assume that $Q_{\infty}^*(F) \neq 0$ and set $\mathrm{Spec}(F) = \{c_1, \dots, c_N\}$ with $c_1 < c_2 < \dots < c_N$. Let $d_0, d_1, \dots, d_N \in \mathbb{R}$ such that $d_0 < c_1 < d_1 < \dots < d_{N-1} < c_N < d_N$. Note that $Q_{\infty, d_0}^*(F) = 0$ by the assumption and $Q_{\infty, d_N}^*(F) = Q_{\infty}^*(F)$. Applying Lemma 6.3 to the exact sequence (6.10) with $d = d_{i-1}, d' = d_i$, by induction we get

$$\mathrm{cl}(Q_{\infty}^*(F)) \leq N - 1 + \sum_{i=1}^N \mathrm{cl}(Q_{\infty, d_i}^*(F)/Q_{\infty, d_{i-1}}^*(F)). \quad (6.12)$$

Hence if $\#\mathrm{Spec}(F) = N \leq \mathrm{cl}(Q_{\infty}^*(F))$, there exists $i \in \{1, \dots, N\}$ such that

$$\mathrm{cl}(Q_{\infty, d_i}^*(F)/Q_{\infty, d_{i-1}}^*(F)) \geq 1. \quad (6.13)$$

For such i above, we claim that $\pi(\mathrm{SS}(F) \cap \Gamma_{dt} \cap \pi^{-1}t^{-1}(-c_i))$ is cohomologically non-trivial. For $c \in \mathbb{R}$ and $I \subset \mathbb{R}$, set

$$K_c := \pi_M(\mathrm{SS}(F) \cap \Gamma_{dt} \cap \pi^{-1}t^{-1}(-c)), \quad K_I := \bigcup_{c \in I} K_c \subset M. \quad (6.14)$$

Let U be any open neighborhood of K_{c_i} in M . We take $K_{c_i} \subset U_0 \Subset U_1 \Subset U$ and a C^∞ -function $\rho: M \rightarrow \mathbb{R}$ such that

- (1) $\rho(x) \in [-1, 1]$ for any $x \in M$,
- (2) $\rho|_{U_0} \equiv 1$,
- (3) $\rho|_{M \setminus U_1} \equiv -1$.

Then there exists $\varepsilon > 0$ such that $K_{[c_i - \varepsilon, c_i + \varepsilon]} \subset U_0$. Taking $0 < \varepsilon' \ll \varepsilon$ and setting $\rho' := \varepsilon' \rho: M \rightarrow \mathbb{R}$, we may assume

$$(x, t; s\rho'(x), 1) \notin \mathrm{SS}(F) \quad (6.15)$$

for $x \in U_1 \setminus U_0, t \in [-c_i - \varepsilon', -c_i + \varepsilon']$, and $s \in [0, 1]$. Hence, we can apply the microlocal Morse lemma to F and $(U_s)_{s \in [0, 1]}$, where

$$U_s := \{(x, t) \in M \times \mathbb{R}_t \mid t < -c_i + (s - 1)\varepsilon' - s\rho'(x)\}, \quad (6.16)$$

and obtain an isomorphism

$$\mathrm{R}\Gamma(M \times (-\infty, -c_i - \varepsilon')) = \mathrm{R}\Gamma(U_0; F) \xrightarrow{\sim} \mathrm{R}\Gamma(U_1; F). \quad (6.17)$$

We set $X = M \times \mathbb{R}$ and

$$Z := M \times (-\infty, -c_i + \varepsilon') \setminus U_1 = \{(x, t) \in M \times \mathbb{R}_t \mid -c_i - \rho'(x) \leq t < -c_i + \varepsilon'\}. \quad (6.18)$$

By the above isomorphism, we get a morphism of exact triangles

$$\begin{array}{ccccccc}
\mathrm{R}\Gamma_{M \times [-c_i + \varepsilon', \infty)}(X; F) & \rightarrow & \mathrm{R}\Gamma_{M \times [-c_i + \varepsilon', \infty) \cup Z}(X; F) & \longrightarrow & \mathrm{R}\Gamma_Z(X; F) & \longrightarrow & \\
\downarrow \wr & & \downarrow \wr & & \downarrow & & \\
\mathrm{R}\Gamma_{M \times [-c_i + \varepsilon', \infty)}(X; F) & \longrightarrow & \mathrm{R}\Gamma_{M \times [-c_i - \varepsilon', \infty)}(X; F) & \longrightarrow & \mathrm{R}\Gamma_{M \times [-c_i - \varepsilon', -c_i + \varepsilon']}(X; F) & \rightarrow & ,
\end{array} \tag{6.19}$$

where the middle vertical morphism is an isomorphism by the above argument. Hence, by the five lemma, we have an isomorphism

$$\mathrm{R}\Gamma_{M \times [-c_i - \varepsilon', -c_i + \varepsilon']}(M \times \mathbb{R}_t; F) \xleftarrow{\sim} \mathrm{R}\Gamma_Z(M \times \mathbb{R}_t; F) \simeq \mathrm{R}\mathrm{Hom}(\mathbf{k}_Z, F). \tag{6.20}$$

Since $\mathrm{Supp}(\mathbf{k}_Z) = \overline{Z} \subset \overline{U}_1 \times [-c_i - \varepsilon', -c_i + \varepsilon']$, the action of $H^*(M; \mathbf{k})$ on $H_Z^*(M \times \mathbb{R}_t; F)$ factors through $H^*(\overline{U}_1)$. Hence, we have

$$\begin{aligned}
\mathrm{cl}(H^*(\overline{U}_1)) &\geq \mathrm{cl}(H_Z^*(M \times \mathbb{R}_t; F)) \\
&= \mathrm{cl}(H_{M \times [-c_i - \varepsilon', -c_i + \varepsilon']}^*(M \times \mathbb{R}_t; F)) \\
&\geq \mathrm{cl}(Q_{\infty, c + \varepsilon'}^*(F)/Q_{\infty, c - \varepsilon'}^*(F)) \geq 1.
\end{aligned} \tag{6.21}$$

Thus, we conclude that $\mathrm{cl}(H^*(U)) \geq 1$, which implies that U is cohomologically non-trivial. \square

We consider the spectral invariants for the sheaf associated with a Hamiltonian diffeomorphism. For a compactly supported Hamiltonian diffeomorphism $\varphi \in \mathrm{Ham}_c(T^*M, \omega)$, let $\mathcal{K}^\varphi \in \mathcal{D}(M^2)$ be the associated sheaf quantization. By Lemma 5.9, if two Hamiltonian diffeomorphisms $\varphi, \varphi' \in \mathrm{Ham}_c(T^*M, \omega)$ satisfy $\varphi(0_M) = \varphi'(0_M)$, then there exists some constant $c \in \mathbb{R}$ such that

$$\mathrm{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}) = \mathrm{Spec}(\mathcal{K}^{\varphi'} \bullet \mathbf{k}_{M \times [0, \infty)}) + c. \tag{6.22}$$

We relate the set of spectral invariants $\mathrm{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$ to the set of spectral invariants $\mathrm{Spec}(\varphi, 0_M)$ of the Lagrangian submanifold $\varphi(0_M)$ for a Hamiltonian diffeomorphism $\varphi \in \mathrm{Ham}_c(T^*M, \omega)$, which is defined in Viterbo [Vit92]. The set of spectral invariants $\mathrm{Spec}(\varphi, 0_M)$ is defined as the set of spectral invariants of a generating function quadratic at infinity $S: E = M \times \mathbb{R}^n \rightarrow \mathbb{R}$ of $\varphi(0_M)$.

Proposition 6.5. *Assume that M is compact. There exists a constant $c \in \mathbb{R}$ such that*

$$\mathrm{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}) = \mathrm{Spec}(\varphi, 0_M) + c. \tag{6.23}$$

Proof. Let $S: E = M \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a generating function quadratic at infinity of $\varphi(0_M)$. It suffices to show that $\mathrm{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$ is equal to the set of spectral invariants associated with S .

Set $\Gamma_S^+ := \{(x, v, t) \in E \times \mathbb{R}_t \mid S(x, v) + t \geq 0\}$ and let $p: E \times \mathbb{R}_t = M \times \mathbb{R}^n \times \mathbb{R}_t \rightarrow M \times \mathbb{R}_t$ be the projection. Then $F_S := \mathrm{Rp}_! \mathbf{k}_{\Gamma_S^+} \in \mathrm{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ satisfies $\mathring{\mathrm{S}}\mathrm{S}(F_S) = \Lambda$, where $\Lambda \subset \mathring{T}^*(M \times \mathbb{R}_t)$ is the conification of $\varphi(0_M)$ with respect to some primitive f :

$$\Lambda := \{(x, t; \xi, \tau) \in T^*(M \times \mathbb{R}_t) \mid \tau > 0, (x; \xi/\tau) \in \varphi(0_M), t = -f(x; \xi/\tau)\}. \tag{6.24}$$

On the other hand, $\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \in \mathcal{D}(M)$ satisfies the same microsupport estimation for some primitive of $\varphi(0_M)$. Hence, by the uniqueness of such object [Gui19, Thm. 12.4.3], there exists $d \in \mathbb{Z}$ and $c' \in \mathbb{R}$ such that

$$\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \simeq T_{c'} F_S[m]. \quad (6.25)$$

Let $j: E \times \mathbb{R}_t = M \times \mathbb{R}^n \times \mathbb{R}_t \hookrightarrow M \times S^n \times \mathbb{R}_t$ be the open embedding that comes from the one-point compactification of \mathbb{R}^n and $\tilde{p}: M \times S^n \times \mathbb{R}_t \rightarrow M \times \mathbb{R}_t$ the projection. Then, $F_S \simeq R\tilde{p}_* j! \mathbf{k}_{\Gamma_S^+}$, $Rp_* \mathbf{k}_{\Gamma_S^+} \simeq R\tilde{p}_* Rj_* \mathbf{k}_{\Gamma_S^+}$, and $R\tilde{p}! \text{Cone}(j! \mathbf{k}_{\Gamma_S^+} \rightarrow Rj_* \mathbf{k}_{\Gamma_S^+})$ is locally constant on $M \times \mathbb{R}_t$. Thus we have

$$\begin{aligned} \text{RHom}(\mathbf{k}_{M \times [-c, \infty)}, F_S) &\simeq \text{RHom}(\mathbf{k}_{M \times [-c, \infty)}, Rp_* \mathbf{k}_{\Gamma_S^+}) \\ &\simeq \text{RHom}(\mathbf{k}_{p^{-1}(M \times [-c, \infty))}, \mathbf{k}_{\Gamma_S^+}) \\ &\simeq \text{R}\Gamma_{\{(x, v) \in E \mid S(x, v) \leq c\}}(E; \mathbf{k}_E). \end{aligned} \quad (6.26)$$

The isomorphism shows that $\text{Spec}(F_S)$ coincides with the set of spectral invariants associated with S , which completes the proof. \square

The spectral norm for Lagrangian submanifolds is defined as follows.

Definition 6.6. Let $\varphi: T^*M \rightarrow T^*M$ be a Hamiltonian diffeomorphism. One defines

$$\gamma(\varphi(0_M)) := \max \text{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}) + \max \text{Spec}(\mathcal{K}^{\varphi^{-1}} \bullet \mathbf{k}_{M \times [0, \infty)}) \quad (6.27)$$

and calls it the *spectral norm* of $\varphi(0_M)$.

Note that the spectral norm $\gamma(\varphi(0_M))$ depends only on the image $\varphi(0_M)$. This can be seen from Lemma 5.9 and Proposition 6.8 below.

We describe this spectral norm in terms of the distance on the Tamarkin category. For that purpose, we introduce an interleaving distance up to shift.

Definition 6.7. For $F, G \in \mathcal{D}(M)$, one defines

$$\overline{d_{\mathcal{D}(M)}}(F, G) := \inf_{c \in \mathbb{R}} d_{\mathcal{D}(M)}(F, T_c G). \quad (6.28)$$

Proposition 6.8. For a Hamiltonian diffeomorphism $\varphi: T^*M \rightarrow T^*M$, one has

$$\gamma(\varphi(0_M)) = \overline{d_{\mathcal{D}(M)}}(\mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}). \quad (6.29)$$

Proof. Let Tor be the full triangulated subcategory of $\mathcal{D}(M)$ consisting of torsion objects $\{F \mid d_{\mathcal{D}(M)}(F, 0) < \infty\}$. Then, the Hom set of the localized category $\mathcal{D}(M)/\text{Tor}$ is computed as

$$\text{Hom}_{\mathcal{D}(M)/\text{Tor}}(F, G) \simeq \varinjlim_{c \rightarrow \infty} \text{Hom}_{\mathcal{D}(M)}(F, T_c G). \quad (6.30)$$

Hence, $Q_\infty^*(F)$ is isomorphic to $\bigoplus_i \text{Hom}_{\mathcal{D}(M)/\text{Tor}}(\mathbf{k}_{M \times [0, \infty)}, F[i])$.

For an object $F \in \mathcal{D}(M)$ with $d(\mathbf{k}_{M \times [0, \infty)}, F) < \infty$, F and $\mathbf{k}_{M \times [0, \infty)}$ are isomorphic in $\mathcal{D}(M)/\text{Tor}$. On the other hand, any isomorphism $\bar{\alpha} \in \text{Hom}_{\mathcal{D}(M)/\text{Tor}}(\mathbf{k}_{M \times [0, \infty)}, F) \simeq Q_\infty^0(F)$ gives an isomorphism $Q_\infty^*(F) \simeq \bigoplus_i \text{Hom}_{\mathcal{D}(M)/\text{Tor}}(\mathbf{k}_{M \times [0, \infty)}, \mathbf{k}_{M \times [0, \infty)}[i]) \simeq H^*(M)$ of right $H^*(M)$ -modules that sends $\bar{\alpha} \in Q_\infty^0(F)$ to $1 \in H^0(M) \simeq \mathbf{k}$. Note that

$$Q_{\infty, c}^0(F) \simeq \begin{cases} \mathbf{k} & (c > c(\bar{\alpha}, F)) \\ 0 & (c < c(\bar{\alpha}, F)) \end{cases} \quad (6.31)$$

by definition. Since $Q_{\infty,c}^*(F) \subset Q_{\infty}^*(F)$ is an $H^*(M)$ -submodule for any c , $Q_{\infty,c}^0(F) \neq 0$ if and only if $Q_{\infty,c}^*(F) = Q_{\infty}^*(F)$. Hence we obtain

$$\max \text{Spec}(F) = c(\bar{\alpha}, F). \quad (6.32)$$

(i) Let $a, b \in \mathbb{R}$ and

$$\alpha: \mathbf{k}_{M \times [0, \infty)} \rightarrow T_a \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}, \quad (6.33)$$

$$\beta: \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \rightarrow T_b \mathbf{k}_{M \times [0, \infty)} \quad (6.34)$$

such that α descends to an isomorphism $\bar{\alpha} \in \text{Hom}_{\mathcal{D}(M)/\text{Tor}}(\mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$ and β descends to its inverse $\bar{\beta}$. Since $\bar{\alpha} \in Q_{\infty,a}^0(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$ is non-zero, $a \geq \max \text{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$. On the other hand, β gives a non-zero element of $Q_{\infty,b}^0(\mathcal{K}^{\varphi^{-1}} \bullet \mathbf{k}_{M \times [0, \infty)})$ since $\mathcal{K}^{\varphi^{-1}} \bullet (*)$ is the inverse functor of $\mathcal{K}^\varphi \bullet (*)$. Hence, we obtain $b \geq \max \text{Spec}(\mathcal{K}^{\varphi^{-1}} \bullet \mathbf{k}_{M \times [0, \infty)})$. This shows $\gamma(\varphi(0_M)) \leq \overline{d_{\mathcal{D}(M)}}(\mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$.

(ii) For any $a > \max \text{Spec}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$, there exists $\alpha: \mathbf{k}_{M \times [0, \infty)} \rightarrow T_a \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}$ that descends to an isomorphism $\bar{\alpha} \in \text{Hom}_{\mathcal{D}(M)/\text{Tor}}(\mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$. For any $b > \max \text{Spec}(\mathcal{K}^{\varphi^{-1}} \bullet \mathbf{k}_{M \times [0, \infty)})$, there also exists $\beta: \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)} \rightarrow T_b \mathbf{k}_{M \times [0, \infty)}$ that descends to an isomorphism $\bar{\beta}$. Since $\text{Hom}_{\mathcal{D}(M)/\text{Tor}}(\mathbf{k}_{M \times [0, \infty)}, \mathbf{k}_{M \times [0, \infty)}) \simeq \mathbf{k}$, we may assume that $\bar{\beta}$ is the inverse of $\bar{\alpha}$ after multiplying a non-zero element of \mathbf{k} to β . The composite $\bar{\beta} \circ \bar{\alpha} = \text{id}_{\mathbf{k}_{M \times [0, \infty)}}$ can be regarded as a non-zero element of $Q_{\infty,a+b}^0(\mathbf{k}_{M \times [0, \infty)})$. Noting that

$$Q_c^0(\mathbf{k}_{M \times [0, \infty)}) \simeq Q_{\infty,c}^0(\mathbf{k}_{M \times [0, \infty)}) \simeq \begin{cases} \mathbf{k} & (c \geq 0) \\ 0 & (c < 0), \end{cases} \quad (6.35)$$

we obtain $a + b \geq 0$ and the preimage $T_a \beta \circ \alpha \in Q_{a+b}^0(\mathbf{k}_{M \times [0, \infty)})$ of $\text{id}_{\mathbf{k}_{M \times [0, \infty)}} = \bar{\beta} \circ \bar{\alpha}$ is $\tau_{a+b}(\mathbf{k}_{M \times [0, \infty)})$. Similarly, we obtain $T_b \alpha \circ \beta = \tau_{a+b}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$ using

$$\text{Hom}_{\mathcal{D}(M)}(\mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}, T_c \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)}) \simeq Q_c^0(\mathbf{k}_{M \times [0, \infty)}) \quad (6.36)$$

and (6.35). This proves $\gamma(\varphi(0_M)) \geq \overline{d_{\mathcal{D}(M)}}(\mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^\varphi \bullet \mathbf{k}_{M \times [0, \infty)})$. \square

Remark 6.9. One can define Hamiltonian spectral invariants in a sheaf-theoretic way as follows. Let $H: T^*M \times I \rightarrow \mathbb{R}$ be a compactly supported Hamiltonian function and $K^H \in \text{D}(\mathbf{k}_{M^2 \times \mathbb{R}_t})$ the associated sheaf quantization. Then, we define

$$\text{Spec}(H) := \text{Spec}(P_l(\mathcal{H}om^*(\mathbf{k}_{\Delta_M \times [0, \infty)}, K_1^H))). \quad (6.37)$$

The Hamiltonian spectral norm is also defined by

$$\gamma(H) := \max \text{Spec}(H) + \max \text{Spec}(\overline{H}) \quad (6.38)$$

and we obtain

$$\gamma(H) = \overline{d_{\mathcal{D}(M^2)}}(\mathbf{k}_{\Delta_M \times [0, \infty)}, \mathcal{K}_1^H). \quad (6.39)$$

We conjecture that $\gamma(H)$ coincides with the Hamiltonian spectral norm of H defined by Frauenfelder–Schlenk [FS07] for a compact manifold M .

7 Arnold-type principle for Hamiltonian homeomorphisms

In this section, we use the previous results to prove an Arnold-type theorem for a Hamiltonian homeomorphism of a cotangent bundle in a purely sheaf-theoretic method. Throughout this section, we assume that M is compact.

Let φ_∞ be a Hamiltonian homeomorphism and $(H_n)_{n \in \mathbb{N}}$ a sequence of Hamiltonian functions that satisfies the condition in Definition 5.16. For any $n \in \mathbb{N}$, we set $\varphi_n := \phi_1^{H_n}$ and $K_n := K^{\varphi_n}$. As discussed just before Definition 5.18, the sequence $(K_n)_{n \in \mathbb{N}}$ forms a Cauchy sequence with respect to $d_{M \times \mathbb{R}_t}$ and gives the sheaf quantization $K_\infty = K^{\varphi_\infty}$ of φ_∞ . Using the object K_n and K_∞ , we define

$$F_n := K_n \bullet \mathbf{k}_{M \times [0, \infty)}, \quad (7.1)$$

$$F_\infty := K_\infty \bullet \mathbf{k}_{M \times [0, \infty)}. \quad (7.2)$$

Then, by (5.22) we have

$$\text{SS}(F_\infty) \cap \{\tau > 0\} \subset \{(x, t; \xi, \tau) \mid \tau > 0, (x, \xi/\tau) \in \varphi_\infty(0_M)\}. \quad (7.3)$$

Moreover, by construction, we get $Q_\infty^*(F_\infty) \simeq H^*(M; \mathbf{k}_M)$.

For a Hamiltonian homeomorphism φ_∞ , the set of spectral invariants $\text{Spec}(\varphi_\infty, 0_M)$ is defined as the limit of those of Hamiltonian diffeomorphisms through the C^0 -continuity. That is, taking a sequence of Hamiltonian function $(H_n)_{n \in \mathbb{N}}$ as above, one defines

$$\text{Spec}(\varphi_\infty, 0_M) = \lim_{n \rightarrow \infty} \text{Spec}(\varphi_n, 0_M) \quad (7.4)$$

with $\varphi_n = \phi_1^{H_n}$. Now we compare $\text{Spec}(F_\infty)$ with $\text{Spec}(\varphi_\infty, 0_M)$.

Lemma 7.1. *There exists a constant $c \in \mathbb{R}$ such that*

$$\text{Spec}(\varphi_\infty, 0_M) = \text{Spec}(F_\infty) + c. \quad (7.5)$$

Proof. Since $d_{\mathcal{D}(M^2)}(\mathcal{K}_n, \mathcal{K}_\infty) \rightarrow 0$ ($n \rightarrow \infty$), we have $d_{\mathcal{D}(M)}(F_n, F_\infty) \rightarrow 0$ ($n \rightarrow \infty$). Thus, we get $\text{Spec}(F_\infty) = \lim_{n \rightarrow \infty} \text{Spec}(F_n)$ and obtain the result by Proposition 6.5. \square

Combining the previous results, we obtain the following result by a purely sheaf-theoretic method, which was originally proved by [BHS19, Thm. 1.1] in a more general setting.

Theorem 7.2. *Let φ_∞ be a Hamiltonian homeomorphism of T^*M . If $\#\text{Spec}(\varphi_\infty, 0_M) \leq \text{cl}(M)$, then $0_M \cap \varphi_\infty(0_M)$ is cohomologically non-trivial, in particular it is infinite.*

Proof. Let $t: M \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ be the projection and Γ_{dt} the graph of the 1-form dt . Then, by (7.3), we have $\pi(\text{SS}(F_\infty) \cap \Gamma_{dt}) \subset 0_M \cap \varphi_\infty(0_M)$. Thus we obtain the result by applying Theorem 6.4 and Lemma 7.1 to F_∞ . \square

Using the spectral norm γ and its C^0 -continuity, we can prove the following stronger result, which is the same as [BHS19, Thm. 1.1]. Remark that the proof is not purely sheaf-theoretic.

Proposition 7.3. *Let $\varphi_\infty: T^*M \rightarrow T^*M$ be a compactly supported homeomorphism. Assume that there exist a compact subset $C \subset T^*M$ and a sequence of Hamiltonian diffeomorphisms $(\varphi_n)_{n \in \mathbb{N}} \subset \text{Ham}_c(T^*M, \omega)$ supported in C that C^0 -converges to φ_∞ for some Riemannian metric.*

(i) There exists an object $F_\infty \in \mathcal{D}(M)$ such that $d_{\mathcal{D}(M)}(\mathbf{k}_{M \times [0, \infty)}, F_\infty) < \infty$ and

$$\text{SS}(F_\infty) \cap \{\tau > 0\} \subset \{(x, t; \xi, t) \mid \tau > 0, (x; \xi/\tau) \in \varphi_\infty(0_M)\}. \quad (7.6)$$

(ii) If $\#\text{Spec}(\varphi_\infty, 0_M) \leq \text{cl}(M)$, then $0_M \cap \varphi_\infty(0_M)$ is cohomologically non-trivial, in particular it is infinite.

Proof. (i) By Proposition 6.5, our $\gamma(\psi(0_M))$ coincides with $\gamma(\psi(0_M))$ in [BHS19] for any compactly supported Hamiltonian diffeomorphism ψ . By [BHS19, Thm. 4,1], for any $\varepsilon > 0$, there exists $\delta > 0$ such that $d_{C^0}(\psi, \text{id}_{T^*M}) < \delta$ implies $\gamma(\psi(0_M)) < \varepsilon$. By the C^0 -convergence of $(\varphi_n)_{n \in \mathbb{N}}$, for any $\delta > 0$, there exists $N \in \mathbb{N}$ such that if $n, m \geq N$, then $d_{C^0}(\varphi_n^{-1}\varphi_m, \text{id}_{T^*M}) < \delta$. Hence, for any $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that if $n, m \geq N$, then $\gamma(\varphi_n^{-1}\varphi_m(0_M)) < \varepsilon$.

We define $F_n := \mathcal{K}^{\varphi_n} \bullet \mathbf{k}_{M \times [0, \infty)} \in \mathcal{D}(M)$. By Proposition 6.8,

$$\gamma(\varphi_n^{-1}\varphi_m(0_M)) = \overline{d'}(\mathbf{k}_{M \times [0, \infty)}, \mathcal{K}^{\varphi_n^{-1}} \bullet \mathcal{K}^{\varphi_m} \bullet \mathbf{k}_{M \times [0, \infty)}) = \overline{d'}(F_n, F_m). \quad (7.7)$$

Hence, the sequence $(F_n)_{n \in \mathbb{N}}$ is a Cauchy sequence with respect to $\overline{d_{\mathcal{D}(M)}}$. This implies that there exists a sequence (c_n) of real numbers such that $(T_{c_n} F_n)_{n \in \mathbb{N}}$ is Cauchy with respect to $d_{\mathcal{D}(M)}$. Thus, there exists a limit object F_∞ of $(T_{c_n} F_n)_{n \in \mathbb{N}}$ by Corollary 4.5. By construction, we obtain $d_{\mathcal{D}(M)}(\mathbf{k}_{M \times [0, \infty)}, F_\infty) < \infty$ and the microsupport estimate.

(ii) Since the set of spectral invariants $\text{Spec}(\varphi_\infty, 0_M)$ is also defined as the limit of $\text{Spec}(\varphi_n, 0_M)$, Lemma 7.1 also holds in this setting. Thus, applying Theorem 6.4 to F_∞ , we obtain the result. \square

8 Arnold-type principle for Hausdorff limits of Legendrians

In this section, we briefly discuss how to prove a Legendrian analog of Theorem 7.2 (cf. [BHS19, Thm. 1.5]) by a sheaf-theoretic method. Again, we assume that M is compact.

Denote by $J^1M = T^*M \times \mathbb{R}$ the 1-jet bundle. For a compact Legendrian submanifold L of J^1M , we define a conic Lagrangian submanifold $c(L)$ of $T^*(M \times \mathbb{R}_t)$ by

$$c(L) := \{(x, t; \xi, \tau) \mid \tau > 0, (x, \xi/\tau, t) \in L\}. \quad (8.1)$$

Let L be a Legendrian submanifold without Reeb chords. Then by the results in [Gui19, Part 12], we can construct $F_L \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ such that $\mathring{\text{SS}}(F_L) = c(L)$.

Consider a sequence $(L_n)_{n \in \mathbb{N}}$ of compact Legendrian submanifolds without Reeb chords of J^1M . Assume that $(L_n)_{n \in \mathbb{N}}$ converges to a compact subset L_∞ of J^1M with respect to the Hausdorff distance. For each L_n , we can construct $F_n := F_{L_n} \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ as above. Following [Gui13] (see also [Gui19, Part 7]), we define $F_\infty \in \mathcal{D}(\mathbf{k}_{M \times \mathbb{R}_t})$ by the exact triangle

$$\bigoplus_{n \in \mathbb{N}} F_n \longrightarrow \prod_{n \in \mathbb{N}} F_n \longrightarrow F_\infty \xrightarrow{+1}. \quad (8.2)$$

Then we get $Q_\infty^*(F_\infty) \simeq C \otimes H^*(M; \mathbf{k}_M)$, where $C := \text{Coker}(\bigoplus_{n \in \mathbb{N}} \mathbf{k} \rightarrow \prod_{n \in \mathbb{N}} \mathbf{k})$. Applying Lemma 2.5 and arguing as in the proof of Lemma 2.9, we have

$$\mathring{\text{SS}}(F_\infty) \subset c(L_\infty) = \lim_{n \rightarrow \infty} c(L_n) \subset T^*(M \times \mathbb{R}_t). \quad (8.3)$$

Let $t: M \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ and $\pi_{\mathbb{R}}: J^1M = T^*M \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ be the projections. Then we find that

$$\text{SS}(F_{\infty}) \cap \Gamma_{dt} \subset \{(x, t; 0, 1) \mid (x, 0, t) \in L\} = (L_{\infty} \cap (0_M \times \mathbb{R}_t)) \times \{1\} \quad (8.4)$$

and hence

$$t\pi(\text{SS}(F_{\infty}) \cap \Gamma_{dt}) \subset \pi_{\mathbb{R}}(L_{\infty} \cap (0_M \times \mathbb{R}_t)). \quad (8.5)$$

We set $\text{Spec}(L_{\infty}) := \text{Spec}(F_{\infty})$.

Proposition 8.1. *In the situation as above, if $\text{Spec}(L_{\infty}) \leq \text{cl}(M)$, then $L_{\infty} \cap (0_M \times \mathbb{R}_t)$ is cohomologically non-trivial, hence it is infinite.*

Proof. By applying Theorem 6.4 to the object $F_{\infty} \in \mathcal{D}(M)$, we obtain the result by (8.3) and $Q_{\infty}^*(F_{\infty}) \simeq C \otimes H^*(M; \mathbf{k}_M)$. \square

References

- [AI20] T. Asano and Y. Ike, Persistence-like distance on Tamarkin’s category and symplectic displacement energy, *J. Symplectic Geom.*, **18** (2020), no. 3, 613–649.
- [AI22] T. Asano and Y. Ike, Sheaf quantization and intersection of rational Lagrangian immersions, to appear in *Annales de l’Institut Fourier*, arXiv:2005.05088 [math.SG], 2022.
- [BCZ21] P. Biran, O. Cornea, and J. Zhang, Triangulation and persistence: Algebra 101, arXiv preprint, arXiv:2104.12258 [math.AT], 2021.
- [BG21] N. Berkouk and G. Ginot, A derived isometry theorem for sheaves, *Advances in Mathematics*, (2021), 108033.
- [BGO19] N. Berkouk, G. Ginot, and S. Oudot, Level-sets persistence and sheaf theory, arXiv preprint, arXiv:1907.09759 [math.AT], 2019.
- [BHS18] L. Buhovsky, V. Humilière, and S. Seyfaddini, A C^0 counterexample to the Arnold conjecture, *Inventiones mathematicae*, **213** (2018), no. 2, 759–809.
- [BHS19] L. Buhovsky, V. Humilière, and S. Seyfaddini, An Arnold-type principle for non-smooth objects, arXiv preprint, arXiv:1909.07081 [math.SG], 2019.
- [BHS21] L. Buhovsky, V. Humilière, and S. Seyfaddini, The action spectrum and C^0 symplectic topology, *Mathematische Annalen*, **380** (2021), no. 1, 293–316.
- [BN93] M. Bökstedt and A. Neeman, Homotopy limits in triangulated categories, *Compositio Mathematica*, **86** (1993), no. 2, 209–234.
- [BP21] N. Berkouk and F. Petit, Ephemeral persistence modules and distance comparison, *Algebraic & Geometric Topology*, **21** (2021), no. 1, 247–277.
- [Chi21] S.-F. Chiu, Quantum speed limit and categorical energy relative to microlocal projector, arXiv preprint, arXiv:2111.05144 [math.SG], 2021.
- [Cru19] J. Cruz, Metric limits in categories with a flow, arXiv preprint, arXiv:1901.04828 [math.CT], 2019.

- [FS07] U. Frauenfelder and F. Schlenk, Hamiltonian dynamics on convex symplectic manifolds, *Israel J. Math.*, **159** (2007), 1–56.
- [GKS12] S. Guillermou, M. Kashiwara, and P. Schapira, Sheaf quantization of Hamiltonian isotopies and applications to nondisplaceability problems, *Duke Math. J.*, **161** (2012), no. 2, 201–245.
- [GS14] S. Guillermou and P. Schapira, Microlocal theory of sheaves and Tamarkin’s non displaceability theorem, In *Homological mirror symmetry and tropical geometry*, Vol. **15** of *Lect. Notes Unione Mat. Ital.* 43–85, Springer, Cham, 2014.
- [Gui13] S. Guillermou, The Gromov-Eliashberg theorem by microlocal sheaf theory, arXiv preprint, [arXiv:1311.0187](https://arxiv.org/abs/1311.0187) [math.SG], 2013.
- [Gui19] S. Guillermou, Sheaves and symplectic geometry of cotangent bundles, arXiv preprint, [arXiv:1905.07341v2](https://arxiv.org/abs/1905.07341v2) [math.SG], 2019.
- [GV22] S. Guillermou and C. Viterbo, The singular support of sheaves is γ -coisotropic, arXiv preprint, [arXiv:2203.12977](https://arxiv.org/abs/2203.12977) [math.SG], 2022.
- [HV] V. Humilière and N. Vichery, Cuplength estimates via microlocal theory of sheaves, in preparation.
- [Kaw19] Y. Kawamoto, On C^0 -continuity of the spectral norm for symplectically non-aspherical manifolds, arXiv preprint, [arXiv:1905.07809](https://arxiv.org/abs/1905.07809) [math.SG], 2019.
- [KS90] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Vol. **292** of *Grundlehren der Mathematischen Wissenschaften*, Springer-Verlag, Berlin, 1990.
- [KS06] M. Kashiwara and P. Schapira, *Categories and sheaves*, Vol. **332** of *Grundlehren der Mathematischen Wissenschaften*, Springer-Verlag, Berlin, 2006.
- [KS18a] M. Kashiwara and P. Schapira, Persistent homology and microlocal sheaf theory, *Journal of Applied and Computational Topology*, **2** (2018), no. 1–2, 83–113.
- [KS18b] A. Kislev and E. Shelukhin, Bounds on spectral norms and barcodes, arXiv preprint, [arXiv:1810.09865](https://arxiv.org/abs/1810.09865) [math.SG], 2018.
- [KS21] M. Kashiwara and P. Schapira, Piecewise linear sheaves, *International Mathematics Research Notices*, **2021** (2021), no. 15, 11565–11584.
- [Li21] W. Li, Estimating Reeb chords using microlocal sheaf theory, arXiv preprint, [arXiv:2106.04079](https://arxiv.org/abs/2106.04079) [math.SG], 2021.
- [Nad09] D. Nadler, Microlocal branes are constructible sheaves, *Selecta Math. (N.S.)*, **15** (2009), no. 4, 563–619.
- [NZ09] D. Nadler and E. Zaslow, Constructible sheaves and the Fukaya category, *J. Amer. Math. Soc.*, **22** (2009), no. 1, 233–286.
- [OM07] Y.-G. Oh and S. Müller, The group of hamiltonian homeomorphisms and C^0 -symplectic topology, *Journal of Symplectic Geometry*, **5** (2007), no. 2, 167–219.
- [PS16] L. Polterovich and E. Shelukhin, Autonomous Hamiltonian flows, Hofer’s geometry and persistence modules, *Selecta Math. (N.S.)*, **22** (2016), no. 1, 227–296.

- [PS20] F. Petit and P. Schapira, Thickening of the diagonal and interleaving distance, arXiv preprint, [arXiv:2006.13150](https://arxiv.org/abs/2006.13150) [math.AT], 2020.
- [PSW21] F. Petit, P. Schapira, and L. Waas, A property of the interleaving distance for sheaves, arXiv preprint, [arXiv:2108.13018](https://arxiv.org/abs/2108.13018) [math.AG], 2021.
- [RS18] M. Robalo and P. Schapira, A lemma for microlocal sheaf theory in the ∞ -categorical setting, *Publications of the Research Institute for Mathematical Sciences*, **54** (2018), no. 2, 379–391.
- [RSV18] F. L. Roux, S. Seyfaddini, and C. Viterbo, Barcodes and area-preserving homeomorphisms, arXiv preprint, [arXiv:1810.03139](https://arxiv.org/abs/1810.03139) [math.SG], 2018.
- [Sco20] L. N. Scoccola, Locally persistent categories and metric properties of interleaving distances, Ph.D thesis, The University of Western Ontario, 2020.
- [Spa88] N. Spaltenstein, Resolutions of unbounded complexes, *Compositio Math.*, **65** (1988), no. 2, 121–154.
- [Tam18] D. Tamarkin. Microlocal condition for non-displaceability. In M. Hitrik, D. Tamarkin, B. Tsygan, and S. Zelditch, editors, *Algebraic and Analytic Microlocal Analysis* 99–223, Cham, 2018. Springer International Publishing.
- [UZ16] M. Usher and J. Zhang, Persistent homology and Floer-Novikov theory, *Geom. Topol.*, **20** (2016), no. 6, 3333–3430.
- [Vit92] C. Viterbo, Symplectic topology as the geometry of generating functions, *Mathematische Annalen*, **292** (1992), no. 1, 685–710.
- [Vit06] C. Viterbo, On the uniqueness of generating hamiltonian for continuous limits of hamiltonians flows, *International Mathematics Research Notices*, **2006** (2006), 34028.
- [Zha20] J. Zhang, *Quantitative Tamarkin Theory*, CRM Short Courses, Springer International Publishing, 2020.

Tomohiro Asano: Faculty of Electrical, Information and Communication Engineering, Institute of Science and Engineering, Kanazawa University, Kakumamachi, Kanazawa, 920-1192, Japan.

E-mail address: tasano@se.kanazawa-u.ac.jp, tomoh.asano@gmail.com

Yuichi Ike: Graduate School of Information Science and Technology, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan.

E-mail address: ike@mist.i.u-tokyo.ac.jp, yuichi.ike.1990@gmail.com