

Dark sectors in neutron-shining-through-a-wall and nuclear absorption signals

Matheus Hostert,^{1,2,3,*} David McKeen,^{4,†} Maxim Pospelov,^{1,2,‡} and Nirmal Raj^{4,§}

¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA*

²*William I. Fine Theoretical Physics Institute, School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA*

³*Perimeter Institute for Theoretical Physics, Waterloo, ON N2J 2W9, Canada*

⁴*TRIUMF, 4004 Wesbrook Mall, Vancouver, BC V6T 2A3, Canada*

(Dated: January 10, 2022)

We propose new searches for n' , a dark baryon that can mix with the Standard Model neutron. We show that IsoDAR, a proposal to place an intense cyclotron near a large-volume neutrino detector deep underground, can look for $n \rightarrow n' \rightarrow n$ transitions with much lower backgrounds than surface experiments. This neutron-shining-through-a-wall search would be possible without any modifications to the experiment and would provide the strongest laboratory constraints on the n - n' mixing for a wide range of mass splittings. We also consider dark neutrons as dark matter and show that their nuclear absorption at deep-underground detectors such as SNO and Borexino places some of the strongest limits in parameter space. Finally, we describe other n' signatures, such as neutrons shining through walls at spallation sources, reactors, and the disappearance of ultracold neutrons.

I. INTRODUCTION

The existence of new baryons in a hidden sector has been a topic of great theoretical and phenomenological interest. One particularly interesting possibility is that of a dark neutron n' , a new fundamental or composite dark particle that mixes with the Standard Model (SM) neutron. In addition to appearing in mirror sectors or brane world theories, n' is interesting on its own due to the potential impact on a number of observables. In cosmology and astrophysics, dark neutrons have been invoked to explain dark matter [1–4], the baryon asymmetry of the Universe [1, 5–8], and help realize asymmetric inflation [9, 10]. They were also shown to modify the physics of cosmic rays, the cosmic microwave background, Big Bang nucleosynthesis, and neutron stars (NS) [11–15]. In the laboratory, dark neutrons can appear in a variety of exotic particle physics processes [11, 16–21]. Among these are hydrogen decays, recently linked to the XENON1T excess [18, 22], and, most notably, n - n' transitions that can explain the long-standing discrepancy between bottle [23–30] and beam [31–33] measurements of the neutron lifetime.

In this paper, we choose to focus on a generic n' neutral state carrying unit baryon number, requiring only that it mixes with the neutron with an arbitrary mixing amplitude $\epsilon_{nn'}$. The low-energy two-state Hamiltonian of the n - n' system is

$$H = \begin{pmatrix} m_n + \Delta E & \epsilon_{nn'} \\ \epsilon_{nn'} & m_n + \delta m \end{pmatrix}, \quad (1)$$

where m_n is the neutron mass and ΔE is the energy contributed to the neutron by some matter-induced poten-

tial. The parameter $\delta m \equiv m_{n'} - m_n$ is the in-vacuum n - n' mass splitting that persists in the limit $\Delta E, \epsilon_{nn'} \rightarrow 0$. Facilities with ultracold neutrons (UCN) [34–38] and cold neutron beams [39] have looked for $n \rightarrow n'$ transitions between n and n' , with rates determined by the parameters in Eq. (1).

We propose exploring a complementary neutron-shining-through-a-wall signature at an accelerator setup located underground. Motivated by recent experimental activity in this area, we consider IsoDAR [40, 41], a proposal for a high-intensity cyclotron to be placed near a large volume detector underground, currently being considered for operation at Yemilab in Korea. The IsoDAR setup would emerge as the most sensitive laboratory probe of dark neutrons to date, partly due to the copious number of neutrons produced by the accelerator, but also due to its underground location, where atmospheric backgrounds are minimal. IsoDAR would provide a marked improvement over the strongest laboratory constraints set by STEREO [42], cold neutron beams [39], and by UCNs at the nEDM experiment [38], all of which constitute surface experiments.

In recasting the UCN nEDM, we discuss how to interpret n - n' oscillation searches in terms of the more general model above, providing complementary coverage in a wider parameter space. We encourage collaborations to go beyond the assumption of a mirror symmetry lifted only by the magnetic or mirror magnetic field, and quote results in terms of the more general model in Eq. (1) (see section II B for details).

For some parameters the dark neutron is stable on cosmological time scales and could be the dark matter of the Universe, an attractive scenario wherein baryon number ensures the stability of both normal and dark matter [1–4, 14]. We place strong limits on this scenario by using the measured absorption rate of single neutrons at the underground large-volume neutrino detectors SNO and Borexino. These limits further complement the probes mentioned above. All in all, our strategies probe new re-

* mhostert@perimeterinstitute.ca

† mckeen@triumf.ca

‡ pospelov@umn.edu

§ nraj@triumf.ca




FIG. 1. The neutron-shining-through-a-wall signature at IsoDAR and the detection processes in the Yemilab liquid-scintillator detector [40, 41]. In the figure, we show the ${}^7\text{Li} - {}^9\text{Be}$ -enriched sleeve enveloping the ${}^9\text{Be}$ target. The sleeve is designed to increase the number of neutrons produced as well as their lifetime inside the sleeve [43], therefore also maximizing the number of $n \rightarrow n'$ conversions. On the right we show two detection possibilities: neutron capture on H or C. Capture on Gd can also be considered, producing ~ 8 MeV photons.

gions of parameter space not previously constrained by UCN searches and astrophysics.

This paper is laid out as follows. In section II we describe our signals, for both neutron-shining-through-a-wall setups and the absorption of dark neutron dark matter, and estimate constraints and future sensitivities. In section III we discuss alternative probes of dark neutrons that may also be of interest, and sketch their sensitivities. We then conclude and provide some discussion on our findings in section IV.

II. SIGNALS AND CONSTRAINTS

In this section we investigate neutron-shining-through-a-wall ($n \rightarrow n' \rightarrow n$) probes and derive limits on n' as dark matter from neutron absorption signals, $n' \rightarrow n$. The resulting limits and sensitivities derived in our study are displayed in fig. 2. We compare them with existing limits from NS overheating (red region) derived in Ref. [44], applicable for δm up to $\mathcal{O}(10)$ MeV as set by the NS nuclear potential. The gray dot-dashed line shows the upper bound on $\epsilon_{nn'}$ derived by Goldman *et al.* [45] from the change in the rotation speed of NSs induced by the production of n' in their cores. Also shown are limits from searches for $n \rightarrow n'$ oscillations at the PSI UCN facility [38] (magenta region) which we describe later.

We now describe the signatures in more detail. We start with the neutron-shining-through-a-wall rate. The net probability of $n \rightarrow n'$ conversion after time t [46, 47] is

$$P_{nn'}(t) = \frac{t}{t_f} \sin^2 2\theta \sin^2 \left[\frac{\sqrt{(\delta m - \Delta E)^2 + 4\epsilon_{nn'}^2 t_f}}{2} \right], \quad (2)$$

where the in-medium mixing angle is given by $\tan 2\theta = 2\epsilon_{nn'}/(\delta m - \Delta E)$ and t_f is the time-of-flight between collisions with the target material. Note that the above expression is valid for t_f much smaller than the neutron lifetime, which in the case of decays in vacuum is 879.4 s, but could be shorter due to absorption in matter. This condition is satisfied in the setups that we examine.

The pre-factor of t/t_f can be intuitively interpreted as the number of collisions N_{coll} , which gets multiplied with the oscillation probability to give the net conversion probability. The number of collisions can be estimated as $N_{\text{coll}} = \Sigma_{\text{el}}/\Sigma_{\text{abs}}$, where $\Sigma_{\text{el}}(\Sigma_{\text{abs}})$ is the column density of the material for elastic scattering (absorption). Therefore, the conversion probability is maximized for materials with low absorption cross section, such as heavy water D_2O .

The effective splitting ΔE could receive contributions from multiple sources:

$$\Delta E = \boldsymbol{\mu}_n \cdot \mathbf{B} + V_F + \dots, \quad (3)$$

where the first term is the Zeeman splitting induced by an external magnetic field, with $|\boldsymbol{\mu}_n| = 1.91e/(2m_p)$, and the second term is the Fermi pseudopotential from neutron forward scattering in matter. The latter is given by

$$V_F = \sum_i \frac{2\pi}{m_n} n_i f_{\text{scat},i}, \quad (4)$$

where n_i is the number density of nuclide i in the scattering target material, and f_{scat} is the neutron scattering length for a given target [48].

We now discuss the calculation of the signal rate for neutrons shining through a wall. At a distance d away from a neutron point source, the flux of dark neutrons can be estimated as

$$\Phi_{n'} = P_{nn'} \frac{R_n}{4\pi d^2}, \quad (5)$$

where R_n is the neutron production rate and the $n \rightarrow n'$ conversion probability is given in Eq. (2). The rate of detection of regenerated neutrons is then given by

$$\Gamma_{n' \rightarrow n}^{\text{det}} = \sum_i N_i \theta_{\text{det}}^2 (\sigma_i^{\text{el}} \epsilon^{\text{el}} + \sigma_i^{\text{abs}} \epsilon^{\text{abs}}) \Phi_{n'} \quad (6)$$

where $i = \{\text{C}, \text{H}\}$ labels the target nuclide, N_i is the number of target nuclides in the detector, and θ_{det} is the mixing angle inside the detector material. The neutron cross section and efficiency of detection are denoted by σ and ϵ , respectively. In brackets we have two terms, the first corresponding to the conversion of $n' \rightarrow n$ in an elastic scattering process followed by the capture of n , and the second corresponding to the conversion of $n' \rightarrow n$ in the capture process. The first term often dominates since $\sigma^{\text{el}} \gg \sigma^{\text{abs}}$, unless the liquid scintillator (LS) is doped with Gd, whose capture cross section can be large enough to compensate for the typical 0.1% concentration.




FIG. 2. Limits on the neutron-dark neutron transition amplitude $\epsilon_{nn'}$ as a function of the absolute mass splitting $|\delta m|$. The solid (dashed) curves correspond to $\delta m > 0$ ($\delta m < 0$), and labels marked with a star correspond to limits derived in this work. **Left.** We show the 10 events/year sensitivity of the near-future IsoDAR experiment at YemiLAB as a black band. The top curve corresponds to the pessimistic scenario (capture on C only) while the bottom curve corresponds to the optimistic one (capture on H or 0.1% Gd). The recent limits posed by the STEREO experiment are shown as cyan regions, which can be improved by IsoDAR by one to two orders of magnitude. These probes are complemented at small mass splittings by constraints from the non-observation of disappearance of ultracold neutrons (UCN) from their traps: the magenta region is our recasting of a search at PSI. Also shown are recently derived limits from overheating of neutron stars [red region] and a bound by Goldman *et al.* using pulsar rotation periods [gray dot-dashed line]. **Right.** Limits for the case of the dark neutron constituting the Galactic dark matter [purple regions]. The upper bound comes from the non-observation of extra neutrons in the measurement of the solar boron-8 neutrino flux at SNO and in a search for exotic decays of carbon at Borexino. The lower bound comes from demanding that the lifetime of the dark neutron exceeds the age of the universe. Also shown are limits from CMB observables on the decay $n' \rightarrow p e \nu$. The grey region in the background is the combination of limits shown on the left panel. See text for further details.

Throughout this work, we use

$$\begin{aligned} \sigma_{\text{H}}^{\text{el}} &= 82 \text{ b}, \quad \sigma_{7\text{Li}}^{\text{el}} = 1.4 \text{ b}, \\ \sigma_{9\text{Be}}^{\text{el}} &= 7.6 \text{ b}, \quad \sigma_{12\text{C}}^{\text{el}} = 5.6 \text{ b}, \end{aligned} \quad (7)$$

for thermal-neutron elastic scattering and

$$\begin{aligned} \sigma_{12\text{C}}^{\text{abs}}(E_{\gamma} = 4.95 \text{ MeV}) &= 3.5 \times 10^{-3} \text{ b}, \\ \sigma_{\text{H}}^{\text{abs}}(E_{\gamma} = 2.22 \text{ MeV}) &= 3.3 \times 10^{-1} \text{ b}, \\ \sigma_{155\text{Gd}}^{\text{abs}}(E_{\gamma} = 8.54 \text{ MeV}) &= 6.1 \times 10^4 \text{ b}, \\ \sigma_{157\text{Gd}}^{\text{abs}}(E_{\gamma} = 7.94 \text{ MeV}) &= 2.5 \times 10^5 \text{ b}, \\ \sigma_{7\text{Li}}^{\text{abs}} &= 4.5 \times 10^{-2} \text{ b}, \quad \sigma_{9\text{Be}}^{\text{abs}} = 7.6 \times 10^{-3} \text{ b}, \end{aligned} \quad (8)$$

for thermal-neutron capture cross section [49]. For simplicity, we assume that the efficiency of detection in elas-

tic and absorption conversions are the same, though the former is expected to be slightly lower due to the possibility of losing the converted neutron. In practice we set $\epsilon^{\text{el}} = \epsilon^{\text{abs}} = 0.3$ [42]. We also assume that $N_{\text{C}} = N_{\text{H}}$, as is the case for benzene and mineral oil.

A. Neutrons-shining-through-a-wall at IsoDAR

IsoDAR is a proposal to place a high-power cyclotron near a large-volume neutrino detector [50]. The proton beam creates neutrons in large numbers, which in turn create ${}^8\text{Li}$, that finally decay at rest to electron antineutrinos. This source can be used for a high-precision and high-intensity program for neutrino physics, including $\bar{\nu}_e$ disappearance searches as a test of light sterile

neutrinos [51, 52] as well as electroweak precision measurements [53]. While the cyclotron concept has been extensively studied [43, 54], the detector choice and siting options are to be determined. Previous proposals considered placing IsoDAR near the KamLAND detector [55, 56] as well as pairing it with water-based liquid scintillator detectors such as THEIA [57]. Recently, however, the possibility of siting IsoDAR at the Yemilab underground laboratories in South Korea has been under active consideration [40, 41]. Motivated by this renewed interest, we consider the sensitivity of IsoDAR@Yemilab to dark neutrons. When appropriate, we also rescale our estimates for IsoDAR@KamLAND, noting that both locations benefit from a low rate of cosmogenic backgrounds.

There are several advantages of considering IsoDAR in searches for new physics coupled to neutrons. It provides a source of neutrons that is as intense as reactors or modern neutron spallation facilities while being underground. This reduces cosmogenic neutrons to negligible levels as compared to the surface. In addition, the proximity to a kt-scale detector maximizes acceptance to new particles, as well as leaking neutrons. The latter, however, can be efficiently shielded with a few meters of material.

We now turn to the experimental details. The IsoDAR cyclotron delivers a 10 mA current of 60 MeV protons on a ^9Be target, corresponding to 6.25×10^{16} protons per second [56]. The neutron production efficiency is approximately 10% yielding an intense source of $R_n \simeq 6.25 \times 10^{15}$ neutrons/second. In the current design [40], the beam target is surrounded by a sleeve with 30% to 70% mass ratio of high-purity ^7Li and ^9Be , respectively, which upon absorbing neutrons, produces ^8Li in large numbers. The ^8Li isotopes subsequently undergo beta-decay at rest to produce electron-antineutrinos $\bar{\nu}_e$ with an endpoint energy of $E_\nu \sim 15$ MeV.

The sleeve is designed to maximize the number of neutrons by reducing absorption on ^6Li impurities and increasing the production of secondary neutrons [43]. Fortunately, this also serves to maximize the rate of n' production; see fig. 1 for a schematic of our setup. In Eq. (2) we replace $t/t_f \rightarrow N_{\text{coll}}$, and approximate the number of collisions as $N_{\text{coll}} = (\Sigma^{\text{el}}/\Sigma^{\text{abs}})_{\text{Li-Be}} \approx 300$. We note, however, that in a full experimental analysis, it would also be important to include the finite size of the sleeve and the collisions of neutrons in the shielding that surrounds it.

For the detector, we assume a 2.5 kt liquid-scintillator (LS) detector with a fiducial mass of 1.16 kt, with its center situated 17 m away from the target [41]. For LS, neutrons can be captured on H, emitting 2.2 MeV gammas, as well as on C, emitting 4.95 MeV gammas. The former is $f_{\text{H}}/f_{\text{C}} \times \sigma_{n\text{H}}/\sigma_{n\text{C}} \simeq 94$ times more common, where f_i is the fraction of H and C in the scintillator, which we assume to be the same. Nevertheless backgrounds for the latter are expected to be much smaller. We also note that a water-based liquid scintillator (WbLS) detector [58, 59] is under consideration at Yemilab [60]. This may present

a more challenging situation for a single neutron measurement due to the larger energy resolution and smaller C concentration.

In all cases, in particular, for a WbLS detector, addition of Gd to the detector volume would be greatly advantageous. Capture on Gd would dominate over that of hydrogen or carbon, and produce 7.9 MeV and 8.54 MeV photons that are much harder to mistake for environmental backgrounds. In fact, for a typical 0.1% Gd concentration, the ratio between capture on Gd and H is approximately

$$\sum_{i=155,157} 0.1\% \times N_{\text{Gd-}i} \sigma_{\text{Gd-}i}^{\text{abs}} \simeq N_{\text{H}} \sigma_{\text{H}}^{\text{abs}}. \quad (9)$$

We exploit this correspondence below.

1. Backgrounds

We now discuss potential backgrounds to a search for reappearing neutrons inside the Yemilab detector. A background study for single-hit $\bar{\nu}_e e^- \rightarrow \bar{\nu}_e e^-$ events at IsoDAR@Yemilab was performed in Ref. [41] under the assumption of a LS detector. In contrast to neutrino-electron scattering, our neutrons-shining-through-a-wall signature consists of a single mono-energetic photon from neutron capture, narrowing the energy region of interest. In addition, if the Yemilab detector technology allows for angular reconstruction (see, for example, progress in angular reconstruction in LS recently reported by Borexino [61, 62], as well as studies for WbLS in Ref. [59]), solar and beam-related neutrino-electron scattering backgrounds could be further reduced since the neutron capture emits gammas isotropically.

a. Neutrino-induced backgrounds The first background to consider comes from mis-identified inverse beta decay (IBD) events. When the positron or the neutron is undetected, IBD can fake a single-hit event. Reference [41] finds that over 5 years about 0.25% of the 1.67×10^6 IBD events at IsoDAR@Yemilab would appear as a single hit. This estimate considers only the detection of a single positron with a missing neutron. In the energy range of interest for capture on H, it is negligible, while for capture on ^{12}C , it constitutes a total of 2.8 events. We note, however, that the alternative case of IBD events with a missing positron and a single neutron could also present a potentially more serious background to our search. If the scattering takes place inside the active or veto volume of the detector, it could be efficiently identified due to the two 511 keV photons from the positron annihilation inside the detector. However, if the positron is produced in a blind region, the neutron can leak inside the fiducial volume and mimic our signal. An evaluation of this background is left to future work in the hope that it be carried out with more sophisticated simulations. In case the rates are large, one should consider smaller fiducial volumes.

In addition to IBD, neutral-current (NC) neutrino-nucleus interactions can also produce a single neutron inside the detector. For ^{12}C , the neutrino energy threshold for such a process is $E_\nu > 18.7$ MeV, which is larger than the energies of both ^8Li and solar ^8B decays. However, $\nu + ^{13}\text{C} \rightarrow \nu + ^{12}\text{C}(\text{g.s.})$ has a lower threshold of $E_\nu > 4.95$ MeV [63–66]. This process is expected to take place in the Yemilab LS, albeit at a small rate due to the natural abundance of ^{13}C of $\sim 1\%$. Using the cross section in Ref. [66], we calculate the ratio between the IBD and $\bar{\nu}_e$ -induced single-neutron-knockout NC interactions to be 6.7×10^{-6} . For the 2.26 kt fiducial volume of the IBD analysis, Ref. [41] quotes 1.67×10^6 IBD events in 5 years of running, which corresponds to 5.7 free neutrons in the 1.16 kt fiducial volume of the single-hit analysis. An additional 2.6 free neutrons are expected from ^8B solar neutrinos. Similar processes also produce the excited state $^{12}\text{C}(2+)$, but at a lower rate and effectively tagged due to the presence of $E_\gamma \sim 4.44$ MeV de-excitation photons. Finally, the LS may also contain some concentration of ^2H , which has a much lower threshold for NC dissociation. Taking, for example, the concentration reported by KamLAND [67], 10^{-4} , and the neutrino-deuteron cross section from Ref. [68], we find a total of 12 and 3.1 events in 5 years from ^8Li antineutrinos and ^8B neutrinos, respectively. Although an irreducible background, the total 23 NC interactions are sub-dominant with respect to environmental backgrounds discussed below and are unlikely to produce signals of neutron capture on C.

While a signal in Ref. [41], $\bar{\nu}_e - e$ interactions represent a background for us. This component has a shape that is well-known and a normalization that would be constrained by *in-situ* measurements. Using a Gaussian energy resolution of $6.4\%/\sqrt{E}$, we find a total of 16 events within two standard deviations of the C capture energy, 4.95 MeV. For capture on H, we find 63 events.

b. Environmental sources Another source of backgrounds comes from radioactivity, spontaneous fission, and spallation in and around the detector. Most backgrounds are smoothly distributed in visible energy and therefore can be constrained outside the peak energy for neutron capture on H, C, or Gd. From Fig. 5 of Ref. [41], we can directly read off the backgrounds to a single-hit n capture on carbon to be 16 events in 5 years of operation. Therefore, provided that IBD events with a missing positron are sufficiently rare, neutron capture on C would constitute a low-background search.

For 2.2 MeV energies no information available. It is fair to assume, however, that this rate is much larger, especially due to low-energy environmental gammas. For instance, the backgrounds at KamLAND increased by two to three orders of magnitude below $E_{\text{vis}} < 3$ MeV, mostly due to radioactivity inside the detector [69, 70]. Therefore, a detailed evaluation of the radiopurity of the detector is necessary to obtain a realistic prediction for backgrounds in the low-energy region. We proceed, however, noting that by exploiting the characteristic wave-

form of the signal, Borexino was able to achieve very low neutron rates at 2.2 MeV, with less than 57 events in 485 days for 533 t of LS [71].

In summary, we find two possibilities for a LS detector: neutron capture on C with $E_\gamma = 4.95$ MeV and an estimated background of 32 events, and neutron capture on H with backgrounds that are expected to be larger, mostly driven by environmental factors. A more detailed evaluation of radiogenic backgrounds at around 2.2 MeV would help confirm if a search for neutron capture on H is at all competitive. We conclude emphasizing that, by virtue of eq. (9), it is possible to interpret our constraints in the H case as an estimate for a Gd-doped detector, where backgrounds would be much smaller.

2. Sensitivity

Having discussed some of the channels of interest, we now discuss the resulting sensitivity of IsoDAR@Yemilab to the n - n' mixing angle. We draw two curves: a pessimistic sensitivity based on neutron capture on C only, and an optimistic sensitivity based on neutron capture on H. In both cases we require $\Gamma_n^{\text{det}} = 10$ events/yr. While this is realistic for the former scenario, it may seem too strong of a requirement for capture on H. This is justified, however, if the detector is doped with Gd 0.1 % concentration. In this case, the capture rate is as large as the one in Hydrogen, and backgrounds can be kept at or under the levels of those of capture on C.

The resulting sensitivities in $\epsilon_{nn'}$ as a function of δm are shown in fig. 2 for both $\delta m > 0$ and $\delta m < 0$. The band corresponds to the difference between optimistic and pessimistic sensitivities. We cut off the $\delta m > 0$ curves at 0.03 eV as that is the kinetic energy down to which neutrons are refrigerated by scatters in the shielding; above this mass splitting, $n \rightarrow n'$ conversions are kinematically forbidden for thermal neutrons. The dips in the curves correspond to resonances at the Fermi pseudopotential, with $V_F = 116$ neV (17 neV) for the Li-Be sleeve (liquid scintillator). Up to background considerations, the corresponding sensitivities for IsoDAR@KamLAND would not be all that different; with a 0.897 kt fiducial mass of liquid-scintillator and a detector center located 16.1 m from the target, the event rates are ~ 1.15 times smaller. As the rate $\propto \epsilon_{nn'}^4$, this results in a reach in $\epsilon_{nn'}$ only 1.03 times weaker than shown for IsoDAR@Yemilab in fig. 2.

Although producing about 10^{16} neutrons per second, comparable in intensity to IsoDAR, STEREO was limited by backgrounds from cosmogenic neutrons. Thanks to IsoDAR's placement deep underground this background can be effectively mitigated and one can improve the sensitivity to $\epsilon_{nn'}$ by 1–2 orders of magnitude, as seen in fig. 2.

B. UCN disappearance

The fact that significant losses of ultracold neutrons from their traps have not been observed can be exploited to place limits on $n \rightarrow n'$ conversions since the feebly interacting n' can escape through the walls. References [34–38] report results on searches for neutron disappearance as a function of a mirror magnetic field B' and assuming $\delta m = 0$. As discussed, we can reinterpret these as upper limits on $\epsilon_{nn'}$ as a function of a non-zero δm , which we display in the magenta region of fig. 2. Here we describe our procedure for reinterpreting these limits.

First, we note that UCN facilities confine neutrons to below a kinetic energy $\text{KE}_{\text{max}} = 300$ neV [72], corresponding to the optical potential of the trap material. This corresponds to a time-of-flight $t_f \lesssim \mathcal{O}(10^{-2})$ s for meter-scale traps [38]. The UCNs receive an additional effective mass of $|\Delta E| \simeq 10^{-12}$ eV $\simeq 1500$ Hz sourced by an externally applied field.

Thus for $|\Delta E - \delta m| \gg t_f^{-1}$ we are in the “fast oscillation” regime, so that in Eq. (2) we can set the second squared sine to $1/2$, giving

$$P_{nn'} \rightarrow \frac{t_s}{t_f} \frac{2\epsilon_{nn'}^2}{4\epsilon_{nn'}^2 + (\delta m - \Delta E)^2}, \quad (10)$$

where t_s is the storage time in UCN traps. Conversely, for $|\Delta E - \delta m| \ll t_f^{-1}$ (in the limit $\epsilon_{nn'} \ll t_f^{-1}$, which will shortly be seen to be true), we are in the “slow oscillation” regime and we get

$$P_{nn'} \rightarrow \epsilon_{nn'}^2 t_s t_f. \quad (11)$$

Thus we get a finite probability on resonance ($\delta m = \Delta E$) that depends on the parametric splitting between the mass eigenstates, $\epsilon_{nn'}$.

Next, we use the data obtained in Ref. [38] for $B' = 0$ to estimate our limits. In particular, the 95% C.L. upper limit on the transition amplitude for two different applied magnetic fields $B = 10$ μT and $B = 20$ μT was found to be

$$\epsilon_{nn'}^{95\%, \delta m=0} < 1.9 \times 10^{-18} \text{ eV}. \quad (12)$$

This is used to infer the corresponding upper limit on the $n \rightarrow n'$ transition probability for ΔE for either value of the magnetic field fixing $\delta m = 0$. We use this value to derive an upper limit on $\epsilon_{nn'}$ using the transition probability with $\delta m \neq 0$, setting t_f to the mean value of 0.065 s in Ref. [38] and properly accounting for the behavior near resonance.

In the left panel of fig. 2 we show the strongest of the two limits obtained for either B ; the two peaks correspond to the resonances at the Zeeman energies. The solid (dashed) curve depicts $\delta m > 0$ ($\delta m < 0$). The $\delta m > 0$ curve is cut off at $\delta m = \text{KE}_{\text{max}}$ as $n \rightarrow n'$ conversions are kinematically forbidden beyond that. No such cutoff exists for the $\delta m < 0$ case, where downscatters are kinematically always allowed.

Our results show that the UCN disappearance search at PSI is generally competitive with and often outperforms the STEREO limit, especially for $\delta m < 0$. We call on UCN collaborations to perform further dedicated analyses involving variations of applied magnetic fields and storage times à la Ref. [38] in order to probe $\epsilon_{nn'}$ vs δm space. By varying the magnitude and direction of external fields, low values of $\epsilon_{nn'}$ may be probed by exploiting resonant dips for both $\delta m > 0$ and $\delta m < 0$. These analyses, along with existing limits by PSI searches, can thus complement IsoDAR for small δm as the $n \rightarrow n'$ disappearance rate goes as only $\epsilon_{nn'}^2$, whereas the $n \rightarrow n' \rightarrow n$ regeneration rate at IsoDAR goes as $\epsilon_{nn'}^4$.

C. Dark neutrons as dark matter: absorption signals

Dark neutrons were likely in chemical equilibrium with SM states in the early universe and can be expected to have primordial abundances comparable to that of standard baryons [15]. If cosmologically long-lived, they could then constitute the dark matter of the universe. We call this scenario dark neutron dark matter (DNDM).

We can probe this scenario by detection of excess neutrons via $n' \rightarrow n$ conversion of the DNDM flux. Reference [4] estimated future sensitivities at neutrino and dark matter experiments using nuclear capture signals. Here we show that extensive limits on DNDM may be already placed by considering past measurements at SNO and Borexino.

Taking the local DM density $\rho_\odot = 0.3$ GeV/cm³ and average speed $v_\chi = 270$ km/s [78], the mean DM flux $\Phi_\chi = (\rho_\odot/m_{n'})v_\chi = 8.6 \times 10^6$ /cm²/s. For keV kinetic energies, the neutron-proton scattering cross section $\sigma_{np} = 4$ b [79], thus the rate-per-proton with which neutrons would be detected from the DNDM flux is

$$R_{n'/p} = \Phi_\chi \times \theta^2 \sigma_{np} = \theta^2 \times 3.5 \times 10^{-17} \text{ s}^{-1}, \quad (13)$$

We compare this rate with the measured rate of single neutrons in underground neutrino detectors.

a. Single neutrons at SNO We first consider the total number of neutrons produced by ⁸B solar neutrinos in $\nu D \rightarrow \nu np$, as measured by SNO [77]. In particular, we use the fact that the ν flux measured by SNO, $\simeq (5.21 \pm 0.67) \times 10^6$ /cm²/s, is in agreement with predictions. Taking the NC cross sections at the relevant energies from Ref. [80], and the ⁸B ν_χ fluxes from Ref. [81], we get the rate per deuteron, (equivalent to the number of neutrons produced per deuteron) as

$$R_{\nu/D} = 2.2 \times 10^{-36} \text{ s}^{-1}. \quad (14)$$

We then demand that not more than 25% (approximately a 95% C.L. constraint) of these SNO neutrons are sourced by our regeneration mechanism, i.e. $R_{n'/p} \leq 0.25 R_{\nu/D}$, using $V_F = 166$ neV for heavy water when computing the in-medium mixing angle.

Type	Experiment	ΔE (eV)	Limit on $\theta_{\text{vac}} \equiv \theta(\Delta E = 0)$
shining-through-a-wall	MARS @ SNS [73]	$\sim 10^{-8}$	6.3×10^{-5}
	MURMUR [74]	$\sim 10^{-8} - 10^{-7}$	1.4×10^{-5}
	STEREO [42]	$\sim 10^{-8} - 10^{-7}$	3.9×10^{-6}
	IsoDAR@YemiLAB [40, 41]	$\sim 10^{-8}$	1.3×10^{-6} (capture on H) 4.0×10^{-7} (capture on C)
ultra cold neutron disappearance	nEDM [38]	$\sim \pm 10^{-12}$	1.8×10^{-6}
resonant neutron regeneration	SNS <i>Broussard et al</i> [39]	–	$[1, 10] \times 10^{-5}$ ^a
	HFIR [75, 76]	–	$\epsilon_{nn'} < 4.4 \times 10^{-17}$ eV ^b
atmospheric neutrons	SNO [77]	10^{-12}	5.9×10^{-4}
	Borexino [71]		6.0×10^{-4}
dark neutron dark matter (nuclear absorption)	SNO [77]	1.7×10^{-7}	2.7×10^{-10}
	Borexino [71]	1.7×10^{-8}	3.3×10^{-10}

^a Valid in the range $\delta m = [50, 400]$ neV.

^b No information available for the $\delta m \gg \Delta E$ regime.

TABLE I. Summary of the exclusion limits imposed by experiments discussed in this work. Bold entries represent future projections. Experimental limits are quoted at the 95% C.L. level, except for Borexino, which are 90% C.L. The IsoDAR projection corresponds to 10 events/year. Here, ΔE is the effective mass contribution to the neutron from some potential sourced by the scattering medium or ambient magnetic fields; see Eq. (1). Limits are quoted on the in-vacuum mixing angle ($\theta_{\text{vac}} = \epsilon_{nn'}/\delta m$ for $\delta m \gg \Delta E$).

b. Single neutrons at Borexino Another experiment we consider is Borexino. In particular, a search for “Pauli Exclusion Principle violation” in $^{12}\text{C} \rightarrow ^{11}\tilde{\text{C}} + n$ transitions was performed by counting the 2.2 MeV photons produced via by neutron capture on H [71]. Using a fiducial mass of 100 tonnes, a 90% C.L. of < 57 signal events was obtained in this channel. This can be compared to Eq. (13) to set limits, where now $V_F = 17$ neV for pseudocumene LS.

Both the SNO and Borexino measurements result in the upper limits on $\epsilon_{nn'}$ versus δm shown in fig. 2 right panel. Also shown in the gray background is the combined region excluded by other probes as displayed on the left panel. The solid (dashed) curve corresponds to $\delta m > 0$ ($\delta m < 0$). The lower limits (i.e., the ceiling of the exclusion region) are set by requiring that dark neutrons do not decay via the beta channel $n' \rightarrow p e \nu$ within the age of the universe:

$$\tau_{n'} \geq \tau_U = 13.7 \text{ Gyr} , \quad (15)$$

with the decay rate¹ given by [15]

$$\tau_{n'}^{-1} = \frac{1}{12 \text{ Gyr}} \left(\frac{\epsilon_{nn'}/\delta m}{5 \times 10^{-8}} \right)^2 , \quad (16)$$

¹ For $\delta m > 0$ the decay channel $n' \rightarrow n \gamma$ is also open. However, this is subdominant to $n' \rightarrow p e \nu$ for $\delta m < 3$ eV.

where we have assumed that the n - n' mass splitting is small compared to the Q value of neutron beta decay of 782 keV.

We have also displayed in fig. 2 right panel the region that would be excluded by cosmic microwave background (CMB) observations of the reionization history, impacted by the injection of electromagnetic energy from the final state electron in n' decay following the recombination epoch. This limit applies for n' lifetimes in the range $10^{12} \text{ s} \lesssim \tau_{n'} \lesssim 10^{26} \text{ s}$ [82, 83]; see Ref. [15] for further details.

We see that in the DNDM scenario we are able to probe parameter space that is complementary to the regions constrained in the previous two sub-sections. In particular, we can limit (a) $\epsilon_{nn'}$ up to three orders of magnitude smaller than STEREO and IsoDAR, due to the detection rate being proportional now only to two powers of the mixing angle as opposed to four, and (b) $\delta m \gg 0.03$ eV as, unlike in the case of searches initiated by $n \rightarrow n'$ conversions, keV dark neutrons are not kinematically limited by in-medium thermalization of neutrons.

We also note that the DM can be comprised of anti- n' particles. In that case, their scattering on nuclei would lead to the appearance of anti-neutrons, which subsequently annihilate with neutrons and deposit ~ 2 GeV energy in detectors. The signature of such DM will be very similar to the di-nucleon decay signal tightly constrained by the Super-Kamiokande collaboration [84]. Ref. [85] determined that $\sim 10^6$ signal events would be registered in the multi-pion channel post kinematic cuts

for a benchmark $\theta = 6 \times 10^{-11}$. About 20 background events are expected in this region, thus this translates to a limit of $\theta \lesssim 4 \times 10^{-12}$ eV, which is more than an order of magnitude stronger than the SNO limit on n' DM (see Table I). This is not surprising given the spectacular signature of dinucleon decay at Super-K resulting in greater energy deposition and fewer backgrounds than nuclear capture at SNO. Moreover, the CMB limits will also be correspondingly stronger, as the energy injection will be $O(10^3)$ times larger compared to the beta decay of n' .

III. OTHER PROBES

In this section, we discuss a few more probes of dark neutrons. As we shall see, these are not as sensitive as the probes described in the previous section, but they still provide a significant amount of neutrons that may be leveraged with future experimental progress. In addition, they may also be relevant for dark baryon models that are more complicated than the one considered here. In Table I we enumerate these strategies alongside those discussed above and, to facilitate comparison, show their reaches in vacuum mixing angle, $\theta_{\text{vac}} = \theta(\Delta E = 0) \simeq \epsilon_{nn'}/\delta m$.

A. Surface neutron sources

1. Reactors

Recently, new limits on n - n' mixing were placed by the STEREO experiment [42] at the Institut Laue-Langevin (ILL) research reactor in France. This neutron-shining-through-a-wall search was originally proposed in Ref. [86] and improved on previous limits by the MURMUR experiment [74, 87], located at BR2 nuclear reactor in Belgium. STEREO searched for neutrons sourced by the ILL reactor converting to dark neutrons via thermalizing scatters in the surrounding heavy water, and subsequently reappearing inside the liquid scintillator detector modules about 10 m away. The search is limited by atmospheric backgrounds, which can be subtracted by a measurement of the neutron rates during the reactor-off periods.

We note that Ref. [42] shows limits for up to $\delta m = 10^4$ eV. However, most neutrons at the STEREO source should thermalize down to room temperatures in the D_2O shielding, analogous to the thermalization of Iso-DAR neutrons in the surrounding sleeve and shielding described above. This would result in an analogous cut-off in δm on the limits; we have drawn a vertical cyan line at $\delta m = 0.03$ eV to roughly indicate what we believe is the boundary of validity of STEREO's limits. The STEREO and MURMUR limits in the vacuum case are shown in table I.

2. ORNL Spallation Neutron Source

The Spallation Neutron Source (SNS) at the Oak Ridge National Laboratory (ORNL) is the world's most intense neutron spallation source. Neutrons are produced by a 1 GeV proton beam impinging on a Hg target. The beam current is $I = 1$ mA with a repetition rate of 60 Hz, corresponding to 1.04×10^{14} protons per pulse. The estimated neutron rate is around 1.5×10^{17} neutrons per second assuming 25 neutrons are produced for every proton. These neutrons can serve as a n' source for the neutrino and neutron detectors used by the COHERENT collaboration [88, 89]. In particular, MARS [73], a neutron detector, was located at a distance of 19.5 m away from the target and was used to understand the flux of fast neutrons at the location of the neutrino alley, where coherent neutrino-nucleus scattering experiments are located.

The neutrino alley is located in a basement with a depth of 8 m.w.e., which provides even less shielding against cosmic ray backgrounds than STEREO's location. This source, however, benefits from the pulsed nature of the beam, allowing for the further reduction of cosmic backgrounds by looking for signals in coincidence with the narrow beam pulses.

The measurement in MARS was performed using a double-hit signature of an elastic neutron scattering on hydrogen followed by a delayed capture on Gd. The flux of prompt neutrons, defined as $t < 2\mu\text{s}$ from the beginning of the beam pulse, was measured for neutrons in the energy range of 3.5 MeV to a few tens of MeV. The final measurement was quoted in terms of a neutron flux, $\Phi_n = 1.20 \pm 0.56$ neutrons/m²/MWh. This quantity can be used to place a limit on neutrons-shining-through-a-wall by demanding that the rate of $n \rightarrow n' \rightarrow n$ conversion do not exceed the observed rate. As an approximation, we neglect multiple scattering at the production and assume all neutrons produced to be within the relevant energy range used in the measurement. The resulting limit on the vacuum mixing angle θ_{vac} is then

$$\theta_{\text{vac}} < 6.3 \times 10^{-5}, \quad (17)$$

at 95% C.L. assuming an integrated power of 7000 GWh per year.

It is also possible to derive limits using delayed thermal neutrons. This would benefit from the increased number of neutron collisions around the target and from the efficacy of the shielding against the thermal visible neutrons. However, the cosmogenic backgrounds would increase, and the rate of observable double-hit events would be more suppressed.

3. J-PARC Spallation Neutron Source

The J-PARC Spallation Neutron Source (JSNS) is the second most intense neutron source in the world. Neutrons are obtained from a 3 GeV proton beam from the

Rapid Cycle Synchrotron (RCS) at J-PARC that strike a Hg target. The beam has 1 MW power and comes in pulses of 8×10^{13} protons/ μs at a 25 Hz frequency. Extrapolating the neutron multiplicity due to the spallation processes up to $E_p = 3$ GeV energies, we find a rate of $\lesssim 50$ neutrons per proton. Overall, we estimate that JSNS produces approximately 1.5×10^{17} neutrons per second.

The J-PARC Sterile Neutrino Search at JSNS (JSNS²) experiment [90] is currently operational and expects to collect several tens of thousands IBD events in the best-fit regions of the sterile-neutrino interpretation of the LSND results in the course of 3 years of operation. The detector [91] is located 24 m away from the Hg target (see also future plans for a second detector [92]), and consists of a cylindrical volume containing a gamma-catcher and a veto volume on the outermost layers with 31 t of LS (no Gd), and a neutrino target with 17 tons of Gd-loaded LS in the innermost volume. The IBD neutrons are captured by Gd after $\sim 30 \mu\text{s}$, yielding an ~ 8 MeV photon, to be compared with the 200 μs -delayed photons with ~ 2.2 MeV energy from capture on H. The former is a preferred signal as the background of beam-related gammas are expected to be too severe up to energies of 2.6 MeV [93].

Single hit events, both prompt and delayed, are copious in the detector and are produced by cosmic rays as well as beam-related photons. The rate for a 25 t detector was found to be $\sim 5.2 \times 10^5$ events/9 μs , quoted for the duration of the “prompt” signal region. This rate assumes a PID rejection larger than 100 to differentiate photons from neutrons, yielding more than 10^7 events in the prompt window. Although there are techniques that can be explored to reduce the number of backgrounds, such as using the pulsed nature of the beam to reject beam-related backgrounds and reducing the detector volume to shield against fast neutrons, cosmic-induced neutrons provide a continuous source of backgrounds that cannot be avoided. A subtraction technique similar to the one employed by STEREO [42] could be explored. We encourage the collaboration to pursue further studies in this direction.

In addition to JSNS and SNS, the European Spallation Source will be well poised to make progress in this field given that its neutron fluence is expected to surpass that of both SNS and JSNS [94].

B. Atmospheric neutrons shining through the ground

Neutrons produced in the collisions of cosmic rays with atmospheric nuclei undergo several nuclear scatters before thermalizing down to non-relativistic speeds. Operating in the interaction basis, these scatters serve as measurements of the interaction eigenstate during which $n \rightarrow n'$ conversions can occur. For small mixing angles, the n' flux produced is unlikely to scatter further,

could go through the Earth’s solid rock, and arrive at an underground detector, where it could regenerate as neutrons that may be detected. This process is similar in some respects to the one described in section II C, with two important differences: the event rate now goes as *four* powers of the mixing versus two, due to two extra powers in the rate of n' production, and the average n' energy is expected to be $\mathcal{O}(\text{eV})$, much lower than the keV energy of DNDM.

We will estimate the sensitivity of this probe at SNO and Borexino as in section II C. To that end, we first take the atmospheric neutron flux from Ref. [95], which we denote by Φ_n^{atm} . Next, in Eq. (2) we set t/t_f (the number of elastic scatters at the top of the atmosphere before the neutron gets absorbed) to 100, roughly the ratio of scattering vs absorption cross sections of the average nuclide in the atmosphere. We then make a simplification by recognizing that the Zeeman splitting due to the Earth’s magnetic field, $\Delta E_{B_\oplus} = 3 \times 10^{-12}$ eV is much greater than the Fermi pseudopotential of air at the top of the atmosphere ($\Delta E_{\text{air}} = 10^{-17}$ eV), where the neutron flux is highest. Thus we set $\Delta E = \Delta E_{B_\oplus}$ in Eq. (2).

The dark neutron flux is now $P_{nn'} \Phi_n^{\text{atm}}$. The neutron appearance rate at SNO and Borexino can now be computed as done in section II C, from which we obtain the limit $\theta_{\text{vac}} \lesssim 6 \times 10^{-4}$ for $\delta m \gg \Delta E$.

C. Resonant cold neutron regeneration

In more controlled environments, like in cold neutron beams, a neutron-shinning-through-a-wall experiment can be carried out with the help of magnetic fields. By adjusting the spatial configuration of the magnetic field, one can arrange for $n \rightarrow n'$ as well as $n' \rightarrow n$ transitions to occur as neutrons and dark neutron beam passes through a series of resonances. Depending on the geometry, magnetic fields, and wavelength of the neutrons, this can drive neutrons to shine through walls. Most recently, this was explored at the SNS with a cold neutron beam passing through a 4.6 T magnetic field [39]. Due to the resonant nature of the signal, the constraints obtained are valid only within a specific range of δm . The range of the cold neutron limits are quoted in table I, applicable only to the δm range of 20 neV to 400 neV.

A similar experiment can also be carried out at ORNL using the High Flux Isotope Reactor (HFIR), which produces even more neutrons than the spallation source. The original proposals in Refs. [75, 76] quote a best sensitivity of $\epsilon_{nn'} > 4.4 \times 10^{-17}$ eV in the degenerate region, $\delta m \ll \Delta E$. We refrain from translating this sensitivity to θ_{vac} since this would depend on the specifics of the experimental apparatus, but note that it should improve upon the SNS limits above due to the larger neutron intensity.

IV. DISCUSSION

We have shown that the near-future IsoDAR@Yemilab setup is highly suitable for a neutron-shining-through-a-wall search for dark neutrons resulting in the sensitivities in fig. 2. These sensitivities complement limits from NS heating and UCN disappearance. While the NS limits are stronger at low δm , it is desirable to probe this parameter space in laboratory conditions as NS temperature measurements come with specific assumptions on nuclear astrophysics and astronomical uncertainties. The NS heating band in fig. 2 is limited from below by the rate at which $n \rightarrow n'$ conversions overheat neutron stars; UCN experiments are uniquely poised to test this region for small δm due to their small ΔE arising from magnetic fields, and disappearance rates that scale only as two powers of $\epsilon_{nn'}$. The NS heating band is also limited from above by the suppression of $n \rightarrow n'$ conversion within the lifetime of the youngest NS observed due to saturation of the n' Fermi sea [44]; IsoDAR is already seen to probe higher $\epsilon_{nn'}$ for $\delta m > 10^{-3}$ eV. IsoDAR is also seen to outperform UCN disappearance searches across a vast range of δm , simply because the population of UCNs in traps cannot compete with the high intensity of neutrons produced in accelerator/cyclotron setups.

One of the main goals of the IsoDAR proposal is the study of sterile neutrinos, which can be viewed as an example of a broader perspective of dark sector physics [96]. Application of the IsoDAR experiment to the search of dark neutrons represents another way this experiment can contribute to the studies of dark sectors. Other applications, that are yet to be studied in the IsoDAR setup, may include the production of exotic unstable particles via nuclear reactions [97] and production of light dark matter with its subsequent interaction in a neutrino detector [98].

Finally, we comment on the implications of our results on ultraviolet physics. A natural UV completion of our setup with sub-eV mass splittings is a mirror sector of the SM giving rise to a dark neutron n' mixing with the

SM neutron via the dimension-9 operator

$$\begin{aligned} \mathcal{L} \supset & \frac{1}{\Lambda^5} u d d u' d' d' + \text{h.c.} \\ & \rightarrow \frac{(4\pi f_\pi^3)^2}{\Lambda^5} n n' + \text{h.c.} \\ & = 10^{-10} \text{ eV} \left(\frac{\text{TeV}}{\Lambda} \right)^5 n n' + \text{h.c.}, \end{aligned} \quad (18)$$

where we have used naive dimensional analysis in moving to the effective Lagrangian at the hadronic level in agreement with lattice computations [99, 100]. This higher dimensional operator arises from renormalizable couplings involving new states—for instance, a color triplet R -parity-violating squark (or anti-diquark) φ and a SM singlet χ with couplings to quarks and mirror quarks of the form $d^{(\prime)} \chi \varphi^*$ and $u^{(\prime)} d^{(\prime)} \varphi$ (see, e.g., [5, 101, 102] for possible UV completions). Necessarily, some of these new states, like φ , carry QCD charge and can be produced at hadron colliders like the LHC. As we can see in Eq. (18), putting these new states at the TeV scale, the rough LHC limit on new QCD-charged particles, can easily lead to a mixing amplitude $\epsilon_{nn'}$ of about 10^{-10} eV. Obtaining larger mixing amplitudes that are above the neutron star cooling limits is possible if the SM-singlet states integrated out to get (18) are pushed down to the GeV scale, i.e. $m_\chi \ll m_\varphi$, increasing $\epsilon_{nn'}$ by $\sim 10^3$ or more. Thus much of the parameter space we propose to probe can arise in plausible UV completions leading to the effective Hamiltonian in (1) that we study.

ACKNOWLEDGMENTS

We gratefully acknowledge fruitful discussion with Adriana Bangau, Janet Conrad, Shmuel Nussinov, Joshua Spitz, and Sandra Zavatarelli. The work of D. M. and N. R. is supported by the Natural Sciences and Engineering Research Council of Canada. TRIUMF receives federal funding via a contribution agreement with the National Research Council Canada. M.P. is supported in part by U.S. Department of Energy Grant No. desc0011842. The research of M.H. was supported in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Research, Innovation and Science.

-
- [1] D. McKeen and A. E. Nelson, *Phys. Rev. D* **94**, 076002 (2016), [arXiv:1512.05359 \[hep-ph\]](#).
 [2] G. K. Karananas and A. Kasseridis, *JCAP* **09**, 036 (2018), [arXiv:1805.03656 \[hep-ph\]](#).

- [3] J. M. Cline and J. M. Cornell, *JHEP* **07**, 081 (2018), [arXiv:1803.04961 \[hep-ph\]](#).
 [4] B. Fornal, B. Grinstein, and Y. Zhao, *Phys. Lett. B* **811**, 135869 (2020), [arXiv:2005.04240 \[hep-ph\]](#).

- [5] K. Aitken, D. McKeen, T. Neder, and A. E. Nelson, *Phys. Rev. D* **96**, 075009 (2017), arXiv:1708.01259 [hep-ph].
- [6] G. Elor, M. Escudero, and A. Nelson, *Phys. Rev. D* **99**, 035031 (2019), arXiv:1810.00880 [hep-ph].
- [7] G. Alonso-Álvarez, G. Elor, A. E. Nelson, and H. Xiao, *JHEP* **03**, 046 (2020), arXiv:1907.10612 [hep-ph].
- [8] A. E. Nelson and H. Xiao, *Phys. Rev. D* **100**, 075002 (2019), arXiv:1901.08141 [hep-ph].
- [9] Z. G. Berezhiani, A. D. Dolgov, and R. N. Mohapatra, *Phys. Lett. B* **375**, 26 (1996), arXiv:hep-ph/9511221.
- [10] K. S. Babu and R. N. Mohapatra, (2021), arXiv:2112.11443 [hep-ph].
- [11] Z. Berezhiani and L. Bento, *Phys. Rev. Lett.* **96**, 081801 (2006), arXiv:hep-ph/0507031.
- [12] Z. Berezhiani and L. Bento, *Phys. Lett. B* **635**, 253 (2006), arXiv:hep-ph/0602227.
- [13] Z. Berezhiani and A. Gazizov, *Eur. Phys. J. C* **72**, 2111 (2012), arXiv:1109.3725 [astro-ph.HE].
- [14] D. McKeen, A. E. Nelson, S. Reddy, and D. Zhou, *Phys. Rev. Lett.* **121**, 061802 (2018), arXiv:1802.08244 [hep-ph].
- [15] D. McKeen, M. Pospelov, and N. Raj, *Phys. Rev. D* **103**, 115002 (2021), arXiv:2012.09865 [hep-ph].
- [16] M. Klopff, E. Jericha, B. Märkisch, H. Saul, T. Soldner, and H. Abele, *Phys. Rev. Lett.* **122**, 222503 (2019), arXiv:1905.01912 [hep-ex].
- [17] D. McKeen and M. Pospelov, (2020), arXiv:2003.02270 [hep-ph].
- [18] D. McKeen, M. Pospelov, and N. Raj, *Phys. Rev. Lett.* **125**, 231803 (2020), arXiv:2006.15140 [hep-ph].
- [19] B. Fornal and B. Grinstein, *Mod. Phys. Lett. A* **35**, 2030019 (2020), arXiv:2007.13931 [hep-ph].
- [20] J. Heck, *Phys. Lett. B* **813**, 136043 (2021), arXiv:2009.01256 [hep-ph].
- [21] A. Strumia, (2021), arXiv:2112.09111 [hep-ph].
- [22] E. Aprile *et al.* (XENON), *Phys. Rev. D* **102**, 072004 (2020), arXiv:2006.09721 [hep-ex].
- [23] A. Serebrov *et al.*, *Phys. Lett. B* **605**, 72 (2005), arXiv:nucl-ex/0408009.
- [24] A. Pichlmaier, V. Varlamov, K. Schreckenbach, and P. Geltenbort, *Phys. Lett. B* **693**, 221 (2010).
- [25] A. Steyerl, J. M. Pendlebury, C. Kaufman, S. S. Malik, and A. M. Desai, *Phys. Rev. C* **85**, 065503 (2012).
- [26] V. F. Ezhov *et al.*, *JETP Lett.* **107**, 671 (2018), arXiv:1412.7434 [nucl-ex].
- [27] S. Arzumanov, L. Bondarenko, S. Chernyavsky, P. Geltenbort, V. Morozov, V. V. Nesvizhevsky, Y. Panin, and A. Strepetov, *Phys. Lett. B* **745**, 79 (2015).
- [28] R. W. Pattie, Jr. *et al.*, *Science* **360**, 627 (2018), arXiv:1707.01817 [nucl-ex].
- [29] A. P. Serebrov *et al.*, *Phys. Rev. C* **97**, 055503 (2018), arXiv:1712.05663 [nucl-ex].
- [30] F. M. Gonzalez *et al.* (UCN τ), *Phys. Rev. Lett.* **127**, 162501 (2021), arXiv:2106.10375 [nucl-ex].
- [31] J. Byrne and P. G. Dawber, *Europhys. Lett.* **33**, 187 (1996).
- [32] J. S. Nico *et al.*, *Phys. Rev. C* **71**, 055502 (2005), arXiv:nucl-ex/0411041.
- [33] A. T. Yue, M. S. Dewey, D. M. Gilliam, G. L. Greene, A. B. Laptev, J. S. Nico, W. M. Snow, and F. E. Wietfeldt, *Phys. Rev. Lett.* **111**, 222501 (2013), arXiv:1309.2623 [nucl-ex].
- [34] G. Ban *et al.*, *Phys. Rev. Lett.* **99**, 161603 (2007), arXiv:0705.2336 [nucl-ex].
- [35] A. P. Serebrov *et al.*, *Nucl. Instrum. Meth. A* **611**, 137 (2009), arXiv:0809.4902 [nucl-ex].
- [36] I. Altarev *et al.*, *Phys. Rev. D* **80**, 032003 (2009), arXiv:0905.4208 [nucl-ex].
- [37] Z. Berezhiani, R. Biondi, P. Geltenbort, I. A. Krasnoshchekova, V. E. Varlamov, A. V. Vassiljev, and O. M. Zhrebtsov, *Eur. Phys. J. C* **78**, 717 (2018), arXiv:1712.05761 [hep-ex].
- [38] C. Abel *et al.* (nEDM), *Phys. Lett. B* **812**, 135993 (2021), arXiv:2009.11046 [hep-ph].
- [39] L. J. Broussard *et al.*, (2021), arXiv:2111.05543 [nucl-ex].
- [40] J. R. Alonso *et al.*, (2021), arXiv:2110.10635 [physics.ins-det].
- [41] J. Alonso, C. A. Argüelles, J. M. Conrad, Y. D. Kim, D. Mishins, S. H. Seo, M. Shaevitz, J. Spitz, and D. Winklehner, (2021), arXiv:2111.09480 [hep-ex].
- [42] H. Almazán *et al.*, (2021), arXiv:2111.01519 [hep-ex].
- [43] A. Bungau, J. Alonso, L. Bartoszek, J. Conrad, M. Shaevitz, and J. Spitz, *JINST* **14**, P03001 (2019), arXiv:1805.00410 [physics.ins-det].
- [44] D. McKeen, M. Pospelov, and N. Raj, *Phys. Rev. Lett.* **127**, 061805 (2021), arXiv:2105.09951 [hep-ph].
- [45] I. Goldman, R. N. Mohapatra, and S. Nussinov, *Phys. Rev. D* **100**, 123021 (2019), arXiv:1901.07077 [hep-ph].
- [46] R. Cowsik and S. Nussinov, *Phys. Lett. B* **101**, 237 (1981).
- [47] B. Kerbikov and O. Lychkovskiy, *Phys. Rev. C* **77**, 065504 (2008), arXiv:0804.0559 [hep-ph].
- [48] V. F. Sears, *Neutron News* **3**, 26 (1992), <https://doi.org/10.1080/10448639208218770>.
- [49] <https://link.springer.com/content/pdf/bbm%3A978-3-642-87614-1%2F1.pdf>.
- [50] A. Bungau *et al.*, *Phys. Rev. Lett.* **109**, 141802 (2012), arXiv:1205.4419 [hep-ex].
- [51] A. Diaz, C. A. Argüelles, G. H. Collin, J. M. Conrad, and M. H. Shaevitz, *Phys. Rept.* **884**, 1 (2020), arXiv:1906.00045 [hep-ex].
- [52] B. Dasgupta and J. Kopp, *Phys. Rept.* **928**, 63 (2021), arXiv:2106.05913 [hep-ph].
- [53] J. M. Conrad, M. H. Shaevitz, I. Shimizu, J. Spitz, M. Troups, and L. Winslow, *Phys. Rev. D* **89**, 072010 (2014), arXiv:1307.5081 [hep-ex].
- [54] J. R. Alonso, R. Barlow, J. M. Conrad, and L. H. Waites, *Nature Reviews Physics* **1**, 533 (2019), arXiv:2003.07931 [physics.acc-ph].
- [55] M. Abs *et al.*, (2015), arXiv:1511.05130 [physics.acc-ph].
- [56] J. R. Alonso and K. Nakamura (IsoDAR), (2017), arXiv:1710.09325 [physics.ins-det].
- [57] G. D. Orebi Gann (THEIA Interest Group), (2015), arXiv:1504.08284 [physics.ins-det].
- [58] J. R. Alonso *et al.*, (2014), arXiv:1409.5864 [physics.ins-det].
- [59] B. J. Land, Z. Bagdasarian, J. Caravaca, M. Smiley, M. Yeh, and G. D. Orebi Gann, *Phys. Rev. D* **103**, 052004 (2021), arXiv:2007.14999 [physics.ins-det].
- [60] S.-H. Seo, (2019), arXiv:1903.05368 [physics.ins-det].
- [61] M. Agostini *et al.* (Borexino), (2021), arXiv:2109.04770 [hep-ex].
- [62] M. Agostini *et al.* (Borexino), (2021), arXiv:2112.11816 [hep-ex].

- [63] J. Arafune, M. Fukugita, Y. Kohyama, and K. Kubodera, *Phys. Lett. B* **217**, 186 (1989).
- [64] M. Fukugita, Y. Kohyama, K. Kubodera, and T. Kuramoto, *Phys. Rev. C* **41**, 1359 (1990).
- [65] T. Suzuki, A. B. Balantekin, and T. Kajino, *Phys. Rev. C* **86**, 015502 (2012), arXiv:1204.4231 [nucl-th].
- [66] T. Suzuki, A. B. Balantekin, T. Kajino, and S. Chiba, *J. Phys. G* **46**, 075103 (2019), arXiv:1904.11291 [nucl-th].
- [67] N. R. Tolich, *Experimental study of terrestrial electron anti-neutrinos with KamLAND*, Ph.D. thesis, Stanford U. (2005).
- [68] S. Nakamura, T. Sato, V. P. Gudkov, and K. Kubodera, *Phys. Rev. C* **63**, 034617 (2001), [Erratum: Phys.Rev.C 73, 049904 (2006)], arXiv:nucl-th/0009012.
- [69] S. Abe *et al.* (KamLAND), *Phys. Rev. C* **81**, 025807 (2010), arXiv:0907.0066 [hep-ex].
- [70] C. Grant, *A Monte Carlo approach to ${}^7\text{Be}$ solar neutrino analysis with KamLAND*, Ph.D. thesis, Alabama U., Alabama U. (2012).
- [71] G. Bellini *et al.* (Borexino), *Phys. Rev. C* **81**, 034317 (2010), arXiv:0911.0548 [hep-ex].
- [72] B. Lauss and B. Blau, *SciPost Phys. Proc.*, 4 (2021).
- [73] D. Akimov *et al.* (COHERENT), (2021), arXiv:2112.02768 [physics.ins-det].
- [74] C. Stasser, G. Terwagne, J. Lamblin, O. Méplan, G. Pignol, B. Coupé, S. Kalcheva, S. Van Dyck, and M. Sarrazin, *Eur. Phys. J. C* **81**, 17 (2021), arXiv:2007.11335 [nucl-ex].
- [75] L. J. Broussard *et al.*, in *Meeting of the APS Division of Particles and Fields* (2017) arXiv:1710.00767 [hep-ex].
- [76] L. J. Broussard *et al.*, *EPJ Web Conf.* **219**, 07002 (2019), arXiv:1912.08264 [physics.ins-det].
- [77] S. N. Ahmed *et al.* (SNO), *Phys. Rev. Lett.* **92**, 181301 (2004), arXiv:nucl-ex/0309004.
- [78] D. Baxter *et al.*, *Eur. Phys. J. C* **81**, 907 (2021), arXiv:2105.00599 [hep-ex].
- [79] <https://nn-online.org/>.
- [80] M. Butler, J.-W. Chen, and X. Kong, *Phys. Rev. C* **63**, 035501 (2001), arXiv:nucl-th/0008032.
- [81] J. Billard, L. Strigari, and E. Figueroa-Feliciano, *Phys. Rev. D* **89**, 023524 (2014), arXiv:1307.5458 [hep-ph].
- [82] T. R. Slatyer, *Physical Review D* **87** (2013), 10.1103/physrevd.87.123513.
- [83] J. M. Cline and P. Scott, *JCAP* **03**, 044 (2013), [Erratum: JCAP 05, E01 (2013)], arXiv:1301.5908 [astro-ph.CO].
- [84] K. Abe *et al.* (Super-Kamiokande), *Phys. Rev. D* **103**, 012008 (2021), arXiv:2012.02607 [hep-ex].
- [85] W.-Y. Keung, D. Marfatia, and P.-Y. Tseng, *JHEP* **09**, 053 (2019), arXiv:1905.03401 [hep-ph].
- [86] M. Sarrazin, G. Pignol, J. Lamblin, F. Petit, G. Terwagne, and V. V. Nesvizhevsky, *Phys. Rev. D* **91**, 075013 (2015), arXiv:1501.06468 [hep-ph].
- [87] M. Sarrazin, G. Pignol, J. Lamblin, J. Pinon, O. Méplan, G. Terwagne, P.-L. Debarsy, F. Petit, and V. V. Nesvizhevsky, *Phys. Lett. B* **758**, 14 (2016), arXiv:1604.07861 [hep-ex].
- [88] D. Akimov *et al.* (COHERENT), *Science* **357**, 1123 (2017), arXiv:1708.01294 [nucl-ex].
- [89] D. Akimov *et al.* (COHERENT), *Phys. Rev. Lett.* **126**, 012002 (2021), arXiv:2003.10630 [nucl-ex].
- [90] S. Ajimura *et al.*, (2017), arXiv:1705.08629 [physics.ins-det].
- [91] S. Ajimura *et al.* (JSNS2), *Nucl. Instrum. Meth. A* **1014**, 165742 (2021), arXiv:2104.13169 [physics.ins-det].
- [92] S. Ajimura *et al.*, (2020), arXiv:2012.10807 [hep-ex].
- [93] M. Harada *et al.*, (2015), arXiv:1502.02255 [physics.ins-det].
- [94] A. Addazi *et al.*, *J. Phys. G* **48**, 070501 (2021), arXiv:2006.04907 [physics.ins-det].
- [95] T. Sato and K. Niita, *Radiation Research* **166**, 544 (2006).
- [96] P. Agrawal *et al.*, *Eur. Phys. J. C* **81**, 1015 (2021), arXiv:2102.12143 [hep-ph].
- [97] E. Izaguirre, G. Krnjaic, and M. Pospelov, *Phys. Lett. B* **740**, 61 (2015), arXiv:1405.4864 [hep-ph].
- [98] E. Izaguirre, G. Krnjaic, and M. Pospelov, *Phys. Rev. D* **92**, 095014 (2015), arXiv:1507.02681 [hep-ph].
- [99] M. I. Buchoff and M. Wagman, *Phys. Rev. D* **93**, 016005 (2016), [Erratum: Phys.Rev.D 98, 079901 (2018)], arXiv:1506.00647 [hep-ph].
- [100] E. Rinaldi, S. Syritsyn, M. L. Wagman, M. I. Buchoff, C. Schroeder, and J. Wasem, *Phys. Rev. D* **99**, 074510 (2019), arXiv:1901.07519 [hep-lat].
- [101] J. M. Arnold, B. Fornal, and M. B. Wise, *Phys. Rev. D* **87**, 075004 (2013), arXiv:1212.4556 [hep-ph].
- [102] Z. Berezhiani, *LHEP* **2**, 118 (2019), arXiv:1812.11089 [hep-ph].