

A NOTE ON LAGRANGIAN SUBMANIFOLDS IN SYMPLECTIC $4m$ -MANIFOLDS

YUGUANG ZHANG

ABSTRACT. This short paper shows a topological obstruction of the existence of certain Lagrangian submanifolds in symplectic $4m$ -manifolds.

1. INTRODUCTION

Let (X, ω) be a symplectic manifold. A submanifold $L \subset X$ is called Lagrangian if $\omega|_L \equiv 0$ and $\dim L = \frac{1}{2} \dim X$. See the textbook and papers [11, 15, 1] for basic properties, general backgrounds, and developments in the study of Lagrangian submanifolds. One question is to understand topological obstructions of embedding manifolds into symplectic manifolds as Lagrangian submanifolds, which has been intensively studied (cf. [1, 3, 4, 6, 12, 13]).

In [16], a phenomenon has been noticed while searching for obstructions of the existence of positive scalar curvature Riemannian metrics on symplectic 4-manifolds. More precisely, Theorem 2.3 in [16] should be interpreted as a simple topological consequence of the presence of certain Lagrangian submanifolds in symplectic 4-manifolds, and this topological constraint is a well-known obstruction of positive scalar curvature metrics by quoting Taubes's theorem on the Seiberg-Witten invariant. The goal of this paper is to reformulate this result in a more suitable context, and to generalise it to the case of $4m$ -dimensional symplectic manifolds, which might have some independent interests.

If X is a compact oriented $4m$ -dimensional manifold, the intersection pairing is the symmetric non-degenerated quadratic form

$$Q : H^{2m}(X, \mathbb{R}) \oplus H^{2m}(X, \mathbb{R}) \rightarrow \mathbb{R}, \quad Q([\alpha], [\beta]) = \int_X \alpha \wedge \beta,$$

which is represented by the diagonal matrix $\text{diag}(1, \dots, 1, -1, \dots, -1)$ under a suitable basis. Denote $b_{2m}^+(X)$ the number of plus ones in the matrix, and $b_{2m}^-(X)$ the number of minus ones. Both of $b_{2m}^\pm(X)$ are topological invariants, and the $2m$ -Betti number $b_{2m}(X) = b_{2m}^+(X) + b_{2m}^-(X)$. If X admits a symplectic form ω , then $[\omega^m] \in H^{2m}(X, \mathbb{R})$, and

$$Q([\omega^m], [\omega^m]) = \int_X \omega^{2m} > 0,$$

which implies

$$b_{2m}^+(X) \geq 1.$$

The main theorem asserts that $b_{2m}^\pm(X)$ provides some constraints of the existence of certain Lagrangian submanifolds.

Theorem 1.1. *Let (X, ω) be a compact symplectic manifold of dimension $4m$, $m \geq 1$, $m \in \mathbb{Z}$, and L be an orientable Lagrangian submanifold in X .*

i) *If the Euler characteristic $\chi(L)$ of L satisfies*

$$(-1)^m \chi(L) > 0, \quad \text{then} \quad b_{2m}^+(X) \geq 2.$$

ii) *If L represents a non-trivial homology class, i.e.*

$$0 \neq [L] \in H_{2m}(X, \mathbb{R}), \quad \text{and} \quad \chi(L) = 0, \\ \text{then} \quad b_{2m}^+(X) \geq 2, \quad \text{and} \quad b_{2m}^-(X) \geq 1.$$

iii) *If*

$$(-1)^m \chi(L) < 0, \quad \text{then} \quad b_{2m}^-(X) \geq 1.$$

iv) *If the $2m$ -Betti number $b_{2m}(X) = 1$, then*

$$\chi(L) = 0, \quad \text{and} \quad 0 = [L] \in H_{2m}(X, \mathbb{R}).$$

When X is 4-dimensional, i) and ii) in the theorem have appeared in the proof of Theorem 2.3 in [16]. There are some immediate applications of Theorem 1.1 that are surely known to experts (cf [4, 3, 6, 7, 12, 13]):

- i) Neither \mathbb{CP}^m nor S^{2m} can be embedded in \mathbb{CP}^{2m} as a Lagrangian submanifold.
- ii) Only possible orientable Lagrangian submanifolds in \mathbb{CP}^2 are 2-tori T^2 . Orientable Lagrangian submanifolds in $\mathbb{CP}^1 \times \mathbb{CP}^1$ are either tori T^2 or spheres S^2 by $b_2^+(\mathbb{CP}^1 \times \mathbb{CP}^1) = 1$.
- iii) There is no Lagrangian sphere S^{4m} in $\mathbb{CP}^{4m-1} \times \Sigma$, where Σ is a compact Riemann surface, since $b_{4m}^+(\mathbb{CP}^{4m-1} \times \Sigma) = 1$.

Theorems in [6, 7, 13] assert that Riemannian manifolds with negative sectional curvature do not admit any Lagrangian embedding into uniruled symplectic manifolds. As a corollary, we show certain constraints of the existence of special metrics on Lagrangian submanifolds.

Corollary 1.2. *Let (X, ω) be a compact symplectic $4m$ -dimensional manifold with*

$$b_{2m}^+(X) = 1.$$

- i) *A compact orientable $2m$ -manifold Y admitting a hyperbolic metric, i.e. a Riemannian metric with sectional curvature -1 , can not be embedded into X as a Lagrangian submanifold.*
- ii) *If $m = 2$, and L is an orientable Lagrangian submanifold in X admitting an Einstein metric g , i.e. the Ricci tensor*

$$\text{Ric}(g) = \lambda g$$

for a constant $\lambda \in \mathbb{R}$, then a finite covering of L is the 4-torus T^4 .

We prove Theorem 1.1 and Corollary 1.2 by using classical differential topology/geometry techniques in the next section.

2. PROOFS

Firstly, we recall basic relevant facts, and provide the detail proofs for the completeness. Let (X, ω) be a compact symplectic $4m$ -dimensional manifold, and L be an orientable Lagrangian submanifold in X . The symplectic form ω induces a natural orientation of X , i.e. given by ω^{2m} . We fix an orientation on L .

Lemma 2.1.

$$[L]^2 = (-1)^m \chi(L),$$

where $[L]^2$ is the self-intersection number of L in X . Consequently, if $\chi(L) \neq 0$, then $[L]^2 \neq 0$, and

$$0 \neq [L] \in H_{2m}(X, \mathbb{R}).$$

Proof. The Weinstein neighbourhood theorem (cf. [15, 11]) asserts that there is a tubular neighbourhood U of L in X diffeomorphic to a neighbourhood of the zero section in the cotangent bundle T^*L , and the restriction of ω on U is the canonical symplectic form on T^*L under the identification. More precisely, if x_1, \dots, x_{2m} are local coordinates on L , then the symplectic form

$$\omega = \sum_{i=1}^{2m} dx_i \wedge dp_i,$$

where p_1, \dots, p_{2m} are coordinates on fibres induced by dx_j , i.e. any 1-form $\sum_{i=1}^{2m} p_i dx_i$ is given by the numbers p_j . We regard L as the zero section of T^*L , and assume that $dx_1 \wedge \dots \wedge dx_{2m}$ gives the orientation of L .

If we denote $\varpi_i = dx_i \wedge dp_i$, $i = 1, \dots, 2m$, then the algebraic relationships are

$$\varpi_i \wedge \varpi_j = \varpi_j \wedge \varpi_i, \quad i \neq j, \quad \text{and} \quad \varpi_i^2 = 0.$$

The orientation inherited from the symplectic manifold is given by

$$\omega^{2m} = (2m)! \varpi_1 \wedge \dots \wedge \varpi_{2m}.$$

The orientation on L induces an orientation on T^*L given by

$$dx_1 \wedge \dots \wedge dx_{2m} \wedge dp_1 \wedge \dots \wedge dp_{2m} = (-1)^m \varpi_1 \wedge \dots \wedge \varpi_{2m}.$$

If L' is a small deformation of L intersecting with L transversally, the intersection number $L \cdot L'$ with respect to the orientation given by ω^{2m} is the self-intersection number of L in X , i.e. $L \cdot L' = [L]^2$. The Euler characteristic of L equals to the intersection number $L \tilde{\cdot} L'$ with respect to the orientation defined by $dx_1 \wedge \dots \wedge dx_{2m} \wedge dp_1 \wedge \dots \wedge dp_{2m}$ (cf. [5]), i.e. $L \tilde{\cdot} L' = \chi(L)$, where we identify the tangent bundle TL with T^*L via a Riemannian metric. Since $L \cdot L' = (-1)^m L \tilde{\cdot} L'$, we obtain $[L]^2 = (-1)^m \chi(L)$. \square

There are analogous formulae in more general contexts, e.g. see [14].

Let g be a Riemannian metric compatible with ω , i.e. $g(\cdot, \cdot) = \omega(\cdot, J\cdot)$ for an almost complex structure J compatible with ω , and let $*$ be the Hodge star operator of g . The volume form dv_g of g is the symplectic volume form, i.e. $dv_g = \frac{1}{(2m)!} \omega^{2m}$. When $*$ acts on $\wedge^{2m} T^*X$, $*^2 = \text{Id}$ holds. If we identify $H^{2m}(X, \mathbb{R})$ with the space of harmonic $2m$ -forms via the Hodge theory (cf. [8]), then $H^{2m}(X, \mathbb{R})$ admits the self-dual/anti-self-dual decomposition

$$H^{2m}(X, \mathbb{R}) = \mathcal{H}_+(X) \oplus \mathcal{H}_-(X), \quad \text{where}$$

$$\mathcal{H}_\pm(X) = \{\alpha \in C^\infty(\wedge^{2m} T^*X) \mid d\alpha = 0, \quad *\alpha = \pm\alpha\}.$$

Note that $b_{2m}^\pm(X) = \dim \mathcal{H}_\pm(X)$.

Lemma 2.2.

$$\omega^m \in \mathcal{H}_+(X).$$

Proof. Since $d\omega^m = 0$, we only need to verify that ω^m is self-dual, i.e. $*\omega^m = \omega^m$, which is a point-wise condition. For any point $x \in X$, we choose coordinates $x_1, \dots, x_{2m}, p_1, \dots, p_{2m}$ on a neighbourhood of x such that on the tangent space $T_x X$,

$$\omega = \sum_{i=1}^{2m} dx_i \wedge dp_i, \quad \text{and} \quad g = \sum_{i=1}^{2m} (dx_i^2 + dp_i^2).$$

The calculation is the same as those in the Kähler case (cf. [8]) because of the locality. If we still write $\varpi_i = dx_i \wedge dp_i$, $i = 1, \dots, 2m$, then the volume form $dv_g = \varpi_1 \wedge \dots \wedge \varpi_{2m}$. Denote the multi-indices sets $I = \{i_1, \dots, i_m\}$, $i_1 < \dots < i_m$, and the complement $I^c = \{1, \dots, 2m\} \setminus I$. Note that

$$\omega^m = (\varpi_1 + \dots + \varpi_{2m})^m = m! \sum_{I \subset \{1, \dots, 2m\}} \varpi_I, \quad \varpi_I = \varpi_{i_1} \wedge \dots \wedge \varpi_{i_m}.$$

By $\varpi_I \wedge * \varpi_I = dv_g = \varpi_I \wedge \varpi_{I^c}$, we obtain $* \varpi_I = \varpi_{I^c}$. Hence $* \omega^m = \omega^m$. \square

Now we are ready to prove the results.

Proof of Theorem 1.1. Note that the assumptions imply $0 \neq [L] \in H_{2m}(X, \mathbb{R})$ in the cases i), ii) and iii). If $PU(L) \in H^{2m}(X, \mathbb{R})$ is the Poincaré dual of $[L]$, then

$$PU(L) = \alpha_+ + \alpha_- \neq 0, \quad \text{and} \quad \alpha_\pm \in \mathcal{H}_\pm(X).$$

Since $\int_X \alpha_+ \wedge \alpha_- = 0$,

$$(-1)^m \chi(L) = [L]^2 = \int_X (\alpha_+ + \alpha_-)^2 = \int_X \alpha_+^2 + \int_X \alpha_-^2.$$

If $(-1)^m \chi(L) \geq 0$, then

$$\int_X \alpha_+^2 \geq - \int_X \alpha_-^2 = \int_X \alpha_- \wedge * \alpha_- = \int_X |\alpha_-|_g^2 dv_g \geq 0.$$

If $\alpha_+ = 0$, then $\alpha_- = 0$ and $PU(L) = 0$, which is a contradiction. Thus $\alpha_+ \neq 0$, and

$$\int_L \alpha_+ = \int_X \alpha_+ \wedge (\alpha_+ + \alpha_-) = \int_X \alpha_+^2 = \int_X |\alpha_+|_g^2 dv_g \neq 0.$$

Since

$$\int_L \omega^m = 0,$$

ω^m and α_+ are linearly independent in $\mathcal{H}_+(X)$. Therefore, we obtain $b_{2m}^+(X) \geq 2$.

If $\chi(L) = 0$, then

$$0 = [L]^2 = \int_X \alpha_+^2 + \int_X \alpha_-^2, \quad \text{and} \quad \int_X |\alpha_-|_g^2 dv_g = \int_X |\alpha_+|_g^2 dv_g.$$

Thus $\alpha_- \neq 0$ and $b_{2m}^-(X) \geq 1$.

If we assume $(-1)^m \chi(L) < 0$, then

$$0 > \int_X \alpha_+^2 + \int_X \alpha_-^2, \quad \text{and} \quad \int_X |\alpha_-|_g^2 dv_g > \int_X |\alpha_+|_g^2 dv_g \geq 0.$$

We obtain the conclusion $b_{2m}^-(X) \geq 1$.

If $b_{2m}(X) = 1$, then $b_{2m}^+(X) = 1$, $b_{2m}^-(X) = 0$, and $H^{2m}(X, \mathbb{R}) = \mathcal{H}_+(X)$. Hence $\chi(L) = 0$ and $[L] = 0$. \square

Proof of Corollary 1.2. Let Y be an orientable compact $2m$ -manifold, and h be a hyperbolic metric on Y . By the Gauss-Bonnet formula of hyperbolic manifolds (cf. [9, 10]),

$$(-1)^m \chi(Y) = \varepsilon_{2m} \text{Vol}_h(Y) > 0,$$

where $\varepsilon_{2m} > 0$ is a constant depending only on m , and $\text{Vol}_h(Y)$ is the volume of the hyperbolic metric. Theorem 1.1 implies i).

Now we assume that $m = 2$. If g is a Riemannian metric on L , then the Gauss-Bonnet-Chern formula for 4-manifolds reads

$$\chi(L) = \frac{1}{8\pi^2} \int_L \left(\frac{R^2}{24} + |W^+|_g^2 + |W^-|_g^2 - \frac{1}{2} |\text{Ric}^o|_g^2 \right) dv_g,$$

where R is the scalar curvature, W^\pm denotes the self-dual/anti-self-dual Weyl curvature of g , and $\text{Ric}^o = \text{Ric} - \frac{R}{4}g$ is the Einstein tensor (cf. [2]). If g is Einstein, then $\text{Ric}^o \equiv 0$ and thus $\chi(L) \geq 0$. By Theorem 1.1, either $\chi(L) < 0$, or $\chi(L) = 0$ and $[L] = 0$ in $H^4(X, \mathbb{R})$ since $b_4^+(X) = 1$. Therefore $\chi(L) = 0$, which implies that

$$R \equiv 0, \quad W^\pm \equiv 0, \quad \text{Ric} \equiv 0.$$

The curvature tensor of g vanishes, i.e. g is a flat metric. Thus a finite covering is the torus T^4 . \square

REFERENCES

- [1] M. Audin, F. Lalonde, L. Polterovich, *Symplectic rigidity: Lagrangian submanifolds*, in *Holomorphic curves in symplectic geometry*, Progress in Mathematics 117, Birkhäuser, (1994), 271–321.
- [2] A.L. Besse, *Einstein manifolds*, Ergebnisse der Math., Springer-Verlag, (1987).
- [3] P. Biran, *Geometry of symplectic intersections*, in *Proceedings of the ICM*, vol. 2 (2002), 241–256.
- [4] P. Biran, K. Cieliebak, *Symplectic topology on subcritical manifolds*, Comment. Math. Helv. 76 (2001), 712–753.
- [5] R. Bott, L. Tu, *Differential forms in algebraic topology*, GTM 82. Springer-Verlag, (1982).
- [6] K. Cieliebak, K. Mohnke, *Punctured holomorphic curves and Lagrangian embeddings*, Invent. Math. vol. 212 (2018), 213–295.
- [7] Y. Eliashberg, A. Givental, H. Hofer, *Introduction to symplectic field theory*, Geom. and Funct. Anal., Special Volume (2000), 560–673.
- [8] P. Griffiths, J. Harris, *Principles of algebraic geometry*, John Wiley and Sons Inc. (1978).
- [9] H. Hopf, *Die curvatura integra Clifford–Kleinscher raumformen*, Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl. (1925), 131–141.
- [10] R. Kellerhals, T. Zehrt, *The Gauss–Bonnet formula for hyperbolic manifolds of finite volume*, Geometriae Dedicata vol. 84, (2001), 49–62.
- [11] D. McDuff, D. Salamon, *Introduction to symplectic topology*, Oxford Mathematical Monographs, Oxford University Press, (1998).
- [12] P. Seidel, *Graded Lagrangian submanifolds*, Bulletin de la S. M. F., tome 128, (2000), 103–149.
- [13] C. Viterbo, *Properties of embedded Lagrange manifolds*, in *First European Congress of Mathematics*, Vol. II (1992), 463–474.
- [14] S. Webster, *Minimal surfaces in a Kähler surface*, J. Diff. Geo. 20, (1984), 463–470.
- [15] A. Weinstein, *Symplectic manifolds and their Lagrangian submanifolds*, Adv. Math. 6, (1971), 329–346.
- [16] Y. Zhang, *Pair-of-pants decompositions of 4-manifolds diffeomorphic to general type hypersurfaces*, arXiv:2102.10037.

INSTITUT FÜR DIFFERENTIALGEOMETRIE, GOTTFRIED WILHELM LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER, DEUTSCHLAND
Email address: yuguangzhang76@yahoo.com