arXiv:2201.02844v2 [hep-th] 17 Jun 2023

# Chiral anomaly in (1+1) dimensions revisited: complementary kinetic perspective and universality

Wei-Han Hsiao and Chiao-Hsuan Wang[1,2,3,4]

[1]*Department of Physics and Center for Theoretical Physics, National Taiwan University, Taipei, 10617, Taiwan*
[2]*Center for Quantum Science and Engineering, National Taiwan University, Taipei, 10617, Taiwan*
[3]*Physics Division, National Center for Theoretical Sciences, Taipei, 10617, Taiwan*
[4]*Pritzker School of Molecular Engineering, University of Chicago, Chicago, IL, 60637, USA*

(Dated: June 2023)

We reinvestigate the classic example of chiral anomaly in (1+1) dimensional spacetime. By reviewing the derivation of charge conservation with the semiclassical Boltzmann equation, we argue that chiral anomalies could emerge in (1+1) dimensions without Berry curvature corrections to the kinetic theory. The pivotal step depends only on the asymptotic behavior of the distribution function of the quasiparticle, and thus its dispersion relation, in the limit of $|\mathbf{p}| \to \pm\infty$ rather than the detailed functional form of the dispersion. We address two subjects motivated by this observation. One concerns reformulating (1+1) dimensional chiral anomaly using kinetic theory with the current algebra approach and the gradient expansion of the Dirac Lagrangian, adding a complementary perspective to the existing approaches. The other demonstrates the universality of chiral anomaly across various quasiparticle dispersions. For two-band models linear in the temporal derivative, with Fujikawa's method we show it is sufficient to have a chirality-odd strictly monotonic dispersion in order to exhibit chiral anomaly.

## I. INTRODUCTION

Anomaly refers to the breakdown of a symmetry of a classical model owing to quantization. One of the most celebrated and vintage examples refers to the chiral anomaly of the (3+1) dimensional massless Dirac fermion in an electromagnetic field [1, 2]. Classically, the two species of Weyl fermions under a massless Dirac fermion decouple. The chiral current $J_5^\mu$, defined as the difference between the currents of the Weyl fermions, is thus conserved $\partial_\mu J_5^\mu = 0$. Nonetheless, quantum mechanically $J_5^\mu$ cannot be conserved if the total U(1) current $J^\mu$, the sum of the two Weyl currents, is computed in a conserved fashion. Historically, this anomaly explains the finite decay rate of pion into photons $\pi^0 \to \gamma\gamma$, and is dubbed triangle anomaly because it appears in a triangle loop diagram when computing this physical process [2]. Ever since its discovery in perturbative calculations, enormous efforts have been dedicated and endowed it with insights from physical and mathematical perspectives. Examples include Fujikawa's path integral measure [3–5] and the index theorem à la Atiyah and Singer [6, 7]. The kinetic equation approach for the triangle anomaly came rather late in the last decade [8–10]. In this formulation, the Berry phase of a Weyl fermion around its Fermi surface induces deformations in phase space volume measure and modifies the equal-time commutation relation between two density operators, leading to the chiral anomaly. This method provides a brief and elegant derivation for the negative magnetoresistance of the Weyl semimetals [11, 12].

This work concentrates on a less popular but technically much simpler model that exhibits chiral anomaly, the (1+1) dimensional Dirac field in the presence of background electromagnetism [1]. It is often considered the warm-up exercise of the triangle anomaly in most educational literature [5, 14, 15]. Nevertheless, there are subtle differences that forbid us from applying every result from the studies on triangle anomaly to this lower dimensional counterpart and vice versa. For instance, the divergence in the calculation of the former occurs in the ultraviolet regime, whereas the ambiguity of the latter turns out to be the proper definition of the charge in the ground state, an infrared quantity. Furthermore, the chiral kinetic theory cannot be defined in parallel because of the nonexistence of Berry curvature owing to the dimensional constraint. The underlying thesis of this work is that a kinetic approach should still exist to bridge the fully quantum electrodynamic treatment and the heuristic argument of spectral flow, where charges are pumped out of or dumped into the Dirac sea depending on their chirality [5, 14–17].

To see how chiral anomaly can arise from an ordinary kinetic theory, let us consider the general Boltzmann equation

$$\frac{\partial n_{\mathbf{p}}}{\partial t} + \mathbf{v} \cdot \frac{\partial n_{\mathbf{p}}}{\partial \mathbf{x}} + \mathbf{f} \cdot \frac{\partial n_{\mathbf{p}}}{\partial \mathbf{p}} = I[\delta n_{\mathbf{p}}], \tag{1}$$

where $n_{\mathbf{p}}(t, \mathbf{x})$, $\mathbf{v}$, $\mathbf{f}$ and $I$ stand for the standard quasiparticle distribution function in the phase space $(\mathbf{x}, \mathbf{p})$, quasiparticle velocity, external force, and collision integral respectively. The conservation laws can be derived by computing the moment averages of $\mathbf{p}$. For instance, the charge conservation $\partial_t \rho + \nabla \cdot \mathbf{j} = 0$ is obtained via integrating the equation with the measure $(d\mathbf{p}) := \int \frac{d^d\mathbf{p}}{(2\pi)^d}$ together with the identifications $\rho = \int (d\mathbf{p}) n_{\mathbf{p}}$ and $\mathbf{j} = \int (d\mathbf{p}) \mathbf{v} n_{\mathbf{p}}$. The external force term would vanishes under the standard integration by part argument [18]. Such consideration is valid for most models and band structures where the regions of large momenta are filled or empty simultaneously and for models whose quasiparticle energy depend only on the magnitude of mementum. It could, however, be violated by an anomalous model such as one chiral sector of the Dirac Lagrangian in (1+1) dimensions. Taking the right-moving fermion, $n_{\mathbf{p}}|_{-\infty}^{\infty} = -1$, and conse-

[1] In this work, we will try to avoid calling it *the Schwinger model* [13] since we do not plan on introducing the dynamical electromagnetism.

quently we have the broken conservation law

$$\partial_t \rho_R + \partial_x j_R = \frac{f}{2\pi}, \tag{2}$$

which becomes the anomalous provided $f = E$, the external electric force. By the same token we can easily see $\partial_t \rho_L + \partial_x j_L = -E/(2\pi)$ and therefore the overall chiral current obeys the anomaly equation

$$\partial_\mu j_5^\mu = \frac{1}{2\pi}\epsilon_{\mu\nu}F^{\mu\nu}. \tag{3}$$

In light of this observation, two topics will be addressed in this work:

1. We first reformulate (1+1) dimensional chiral anomaly in terms of kinetic theory. The first step involves demonstrating the Hamiltonian formulation of the kinetic theory introduced in Ref.[8], together with the asymptotic behavior argument presented in the above, naturally implies the central extension of Kac-Moody algebra for the charge densities of chiral fermions, which in turn implies chiral anomaly. We also consider the conventional gradient expansion of the field theoretical correlation functions, showing each chiral sector of the (1+1) dimensional Dirac fermion satisfies a chirality dependent semiclassical Boltzmann equation. These exercises not only offer an alternative perspective filling the gap between the illustrative spectral-flow picture and other fairly technical approaches such as direct diagrammatics, dispersion relations, or Fujikawa's determinant, but serve as the analogue of the kinetic equation approach of the (3+1) dimensional triangle anomaly.

2. Since this distribution function argument depends only on the asymptotic behavior of $n_{\mathbf{p}}$ as $\mathbf{p} \to \pm\infty$, a strictly monotonic dispersion relation would suffice to yield a finite difference at $T = 0$. To quantify this statement, we will consider a class of (1+1) dimensional Dirac-like models. They are linear in time derivative but allow generic functional forms of the spatial derivative. We show quantum mechanically such models has chiral anomaly so long as its chiral sectors assume the dispersion relations

$$\omega \sim (|\mathbf{q}|)^{2n-1}, n \in \mathbb{N}.$$

The leading contribution is the same value as the one given by the Dirac field, $E/(2\pi)$, regardless of the value of $n$. The argument is then generalized to apply to general strictly monotonic dispersions $\omega \sim \epsilon(-i\partial_x)$.

These themes will be elaborated further in Secs. II and III respectively. A couple of comments will follow to conclude the work.

## II. KINETIC FORMULATION FOR (1+1) DIMENSIONAL CHIRAL ANOMALY

### A. Hamiltonian formulation for anomalous charge commutation relation

The kinetic theory of fermionic quasiparticles is reformulated with the Hamiltonian functional $H[n_{\mathbf{p}}(\mathbf{x})]$ and the commutation relation between the distribution functions $[n_{\mathbf{p}}(\mathbf{x}), n_{\mathbf{q}}(\mathbf{y})]$. To produce Eq (1) via the Louville equation $\partial_t n_{\mathbf{p}}(\mathbf{x}) = i[H, n_{\mathbf{p}}(\mathbf{x})]$ in the $(d+1)$ dimensional spacetime, the commutation relation is postulated to be

$$\begin{aligned} &i[n_{\mathbf{p}}(\mathbf{x}), n_{\mathbf{q}}(\mathbf{y})] \\ =&(2\pi)^d \frac{\partial}{\partial \mathbf{p}}\delta(\mathbf{p}-\mathbf{q}) \cdot \frac{\partial}{\partial \mathbf{x}}\delta(\mathbf{x}-\mathbf{y})[n_{\mathbf{p}}(\mathbf{y}) - n_{\mathbf{q}}(\mathbf{x})] \end{aligned} \tag{4}$$

in Ref.[8]. The bulk of that seminal work is devoted to showing the equal-time commutator between two charge-density operators $n(\mathbf{x}) = \int d\Gamma n_{\mathbf{p}}(\mathbf{x})$ is anomalous when accounting for the effect of Berry phase on the volume measure $d\Gamma$. Consider a model consisting of two species of fermions. In the free-theory limit, they have quasiparticle dispersions $\omega = \sigma|\mathbf{k}|$, where $\sigma = \pm 1$ denotes the *chirality*. As a straightforward evaluation, it can be shown that in $(1+1)$ dimensions, Eq. (4) naturally leads to anomalous density-density commutation relation for a chiral fermion.

$$\begin{aligned} &[n_\sigma(x), n_{\sigma'}(y)] = \int \frac{dp}{(2\pi)}\frac{dq}{(2\pi)}[n_{\sigma p}(x), n_{\sigma' q}(y)] \\ =&\frac{i}{2\pi}\delta_{\sigma\sigma'}\frac{\partial}{\partial x}\delta(x-y)n_p(y)\Big|_{-\infty}^{\infty}. \end{aligned} \tag{5}$$

Accounting for the different asymptotic behaviors of $n_{\sigma p}$ as we did in the introduction, the above reduces to the extended Kac-Moody algebra

$$[n_\sigma(x), n_{\sigma'}(y)] = \frac{\sigma}{2\pi i}\delta_{\sigma\sigma'}\partial_x\delta(x-y). \tag{6}$$

Given this commutation relation, the anomaly can be derived in the same fashion as the hydrodynamic approaches of chiral Luttinger liquids [19]. We consider an inhomogeneous electric potential $\phi(x)$ in the Hamiltonian [2]

$$H = H_0 + \int dx\, n(x)\phi(x). \tag{7}$$

Evaluating the commutator $[H, n(x)]$, $H_0$ part yields the current conservation. The additional contribution from the potential reads

$$\begin{aligned} \partial_t n_\sigma + \partial_x j_\sigma =& i\int dy\, \phi(y)[n_\sigma(y), n_\sigma(x)] \\ =& -\frac{\sigma}{2\pi}\partial_x\phi = \frac{\sigma}{2\pi}E. \end{aligned} \tag{8}$$

[2] $H_0$ is assumed to be the free Hamiltonian. This formulation is seemingly independent of spacetime dimensions, but physically the well-defined quasiparticles are destroyed in (1+1) dimensions as the short-range interaction is turned on.

We then recover the anomalous conservation of charge for chiral fermion. Charge conservation and chiral anomaly can be derived by taking the sum and the difference of the $\sigma = \pm 1$ sectors as we described earlier. Note that since this derivation is independent of monopole or Berry phase and therefore should not be sensitive to adiabatic assumptions, and the current algebra (6) is unsurprisingly identical to ones given by bosonization and the Bjorken-Johnson-Low (BJL) prescription [5, 20].

### B. kinetic equation from microscopic models

We next evaluate the semi-classical expansion of Dirac fermion in (1+1) dimensions.

$$\mathscr{L} = i\bar{\psi}\gamma^\mu D_\mu \psi + \mu\psi^\dagger\psi \tag{9}$$

with $\gamma^\mu = (\sigma^y, i\sigma^x)$ and $D_\mu = \partial_\mu + iA_\mu$. The chiral operator $\gamma_5$ is $\gamma^0\gamma^1 = \sigma^z$. In the absence of the fermion mass, chiral sectors, or the upper and the lower entries of the spinor $\psi$ decouple. As pointed out earlier, a decent amount of effort has been dedicated to the kinetic theory formulation of triangle anomaly in (3+1) dimensions. Nevertheless, not all arguments and technique there can be adapted when tackling its (1+1) dimensional counterpart such as the Berry phase term emerging in the hard-density loop expansions. As such, here we simply perform the most generic semi-classical approximation on each chiral sectors [18, 21, 22]. To proceed, we rephrase the Lagrangian (9) as the sum of two chiral sectors

$$\mathscr{L} = \sum_{\sigma=\pm} \mathscr{L}_\sigma,\ \mathscr{L}_\sigma = i\psi_\sigma^\dagger[D_t + \sigma D_x]\psi_\sigma + \mu\psi_\sigma^\dagger\psi_\sigma. \tag{10}$$

The classical equations of motion is given trivially by varying $\psi^\dagger$ and $\psi$, which are

$$i\partial_t\psi_\sigma = -i\sigma\partial_x\psi + V_\sigma\psi_\sigma, V_\sigma = A_0 + \sigma A_1 - \mu, \tag{11}$$

and its complex conjugate. To derive the kinetic equations, let us consider the Green's function

$$iG_\sigma(x_1, x_2) = \langle T\psi_\sigma(x_1)\psi_\sigma^\dagger(x_2)\rangle. \tag{12}$$

Here the time-ordered propagator at zero temperature is adopted, but the technique is general and should apply to any kind of Green's functions. Differentiating the Green's function with respect to $x$ and $x'$, and substituting the equations of motion for the time derivatives of the field operators, we obtain

$$(\partial_{t_1} + \sigma\partial_{x_1} + iV_\sigma(x_1))iG_\sigma(x_1, x_2) = \delta(x_1 - x_2), \tag{13a}$$

$$(-\partial_{t_2} - \sigma\partial_{x_2} + iV_\sigma(x_2))iG_\sigma(x_1, x_2) = \delta(x_1 - x_2). \tag{13b}$$

The sum and the difference of these two equations yield the on-shell condition and quantum kinetic equation respectively. To proceed, we introduce the center of mass and relative coordinates $(x^\mu, s^\mu)$

$$x = \frac{x_1 + x_2}{2} \tag{14a}$$

$$s = x_1 - x_2. \tag{14b}$$

In terms of these coordinates, the sum of Eqs. (13a) and (13b) reads

$$\left(i\frac{\partial}{\partial s^0} + i\frac{\partial}{\partial s^1}\sigma - V_\sigma(x)\right) iG_\sigma = i\delta(s), \tag{15}$$

where we have expanded $V_\sigma(x \pm s/2)$ to the first order. Fourier transforming the coordinate $s^\mu$ suggests $iG$ has a pole structure $i(\omega - \sigma p^1 - V_\sigma)^{-1}$. Next we consider the difference of Eqs. (13a) and (13b),

$$\left(i\frac{\partial}{\partial x^0} + i\sigma\frac{\partial}{\partial x^1} - s^\mu\partial_\mu V_\sigma\right) iG_\sigma = 0. \tag{16}$$

Schematically, this equation reduces to the kinetic equation after the Wigner transformation. To maintain the gauge covariance, let us consider the covariant Wigner-transformation

$$i\mathscr{G}(x,p) = \int d^2s\, e^{isp} U(x, x_+) iG(x_+, x_-) U(x_-, x), \tag{17}$$

$$U(x,y) = \exp\left(i\int_x^y d\ell^\mu A_\mu\right),\ x_\pm = x \pm \frac{s}{2}.$$

By plugging the inverse-transformation of the above into Eq. (16), we obtain the quantum Boltzmann equation

$$\left(\partial_t + \partial_x\sigma + E\frac{\partial}{\partial p^1} + E\frac{\partial}{\partial\omega}\sigma\right) i\mathscr{G}_\sigma(x,p) = 0 \tag{18}$$

with the electric field $E = \partial_t A_1 - \partial_x A_0$. The third and fourth term have the obvious physical meanings of external force and power $\mathbf{F}\cdot\mathbf{v}$ respectively. The semi-classical Boltzmann equation follows by putting the quantum version on-shell via the on-shell condition $i\mathscr{G}_\sigma(x,p) = 2\pi\delta(\omega - \epsilon_{\sigma p}(x))n_{\sigma p}(x)$ [22]. We thus have the (1+1)-dimensional version of Eq. (1)

$$\partial_t n_{\sigma p} + \partial_x\sigma n_{\sigma p} + E\partial_p n_{\sigma p} = 0. \tag{19}$$

Let us now integrate the last momentum variable with the measure $\int \frac{dp^1}{2\pi}$. The first two terms are simply $\partial_t n_\sigma + \partial_x j_\sigma$. To evaluate the third term, we note that at equilibrium, the sign of the quantity determines $\sigma p^1 + V_\sigma$ if the quasiparticle state is occupied. Away from equilibrium, this should still entail asymptotic occupation at $p^1 \to \pm\infty$, and accordingly

$$\int_{-\infty}^{\infty}\frac{dp^1}{2\pi}\frac{\partial}{\partial p^1}n_{\sigma p} = -\frac{\sigma}{2\pi}. \tag{20}$$

Putting them altogether, we recover Eq.(8)

$$\partial_t n_\sigma + \partial_x j_\sigma = \frac{\sigma}{2\pi}E. \tag{21}$$

Thereby, in terms of anomalous commutation relation of charges and gradient expansion, we establish kinetic approaches for chiral anomaly in (1+1) dimensions as a complementary perspective about this classic topic.

## III. UNIVERSAL CHIRAL ANOMALY FOR ODD DISPERSION

Let us now change the gear, and consider the quantum mechanical implications by the previous semi-classical argument. Although the Boltzmann equation was derived using a linear dispersion relation, the only dependence on the spectrum enters indirectly via the asymptotic behavior of the particle number distribution function $n_{\sigma p}(x)$. By repeating the arguments in the previous two section and keep temporal derivative linear, a chirality-odd strictly monotonic dispersion relation should be sufficient in order for a model to exhibit chiral anomaly regardless of the spacetime symmetry.

To prove this statement, let us first consider a power-law dispersion relation with an odd power.

$$\mathscr{L}_\ell = \psi^\dagger (i\partial_t - \kappa\sigma^z(-i\partial_x)^\ell)\psi, \tag{22}$$

where $\ell$ is an odd integer and $\kappa$ is a dimensionful parameter to restore the unit of the term. Lagrangian (22) is invariant under the global U(1) transformation $\psi \to e^{i\alpha}\psi$ and chiral transformation $\psi \to e^{i\gamma_5\alpha}\psi = e^{i\sigma_z\alpha}\psi$. To evaluate the chiral anomaly, let us deploy Fujikawa's recipe [3–5] by considering the partition functional

$$\mathcal{Z} = \int \mathscr{D}\psi\mathscr{D}\bar{\psi}\, \exp\left(i\int d^2x\, \mathscr{L}_\ell[\psi,\bar{\psi}]\right). \tag{23}$$

Under a local chiral transformation, $\psi \to (1 + i\alpha(x)\gamma_5)\psi$, the chiral current emerges in the Lagrangian $\mathcal{L}_\ell + \alpha(x)\partial_\mu j_5^\mu$, whereas the path integral measure is scaled by the Jacobian

$$\mathscr{D}\psi\mathscr{D}\bar{\psi} \to \mathscr{J}^{-2}\mathscr{D}\psi\mathscr{D}\bar{\psi}, \tag{24}$$

where the logarithm of $\mathscr{J}$ amounts to evaluating the trace of the chiral operator over the Hilbert space. Using the standard Gaussian regularization,

$$\ln \mathscr{J} = i\int d^2x\, \alpha(x) \lim_{M\to\infty} \langle x|\mathrm{Tr}[\gamma_5 e^{-\hat{H}^2/M^2}]|x\rangle \tag{25}$$

with the *energy* operator $\hat{H}^2$ of the form

$$\begin{aligned} -\hat{H}^2 &= [\gamma^0 iD_0 - \kappa\gamma^1(-iD_1)^\ell]^2 \\ &= -D_0^2 - \kappa^2(-iD_1)^{2\ell} - \kappa\gamma_5 iF_{01}\ell(-iD_1)^{\ell-1} \end{aligned} \tag{26}$$

Expanding the Fujikawa Jacobian to the first non-vanishing order and evaluating the trace,

$$\begin{aligned} \mathscr{J} &= -2i\kappa F_{01}\ell\langle x|e^{-[D_0^2-\kappa^2(-iD_1)^{2\ell}]/M^2}\frac{(-iD_1)^{\ell-1}}{M^2}|x\rangle \\ &= 2F_{01}\int \frac{d^2k}{(2\pi)^2} e^{-k_0^2-\kappa^2k_1^{2\ell}}\ell\kappa k_1^{\ell-1} = \frac{1}{2\pi}F_{01}. \end{aligned} \tag{27}$$

Note that a Wick rotation is performed on the temporal momentum so the integral is properly Euclidean. When evaluating the trace it is evident that no anomaly could remain for even $\ell$ since the momentum integral simply vanishes. By a similar token, this derivation generalizes straightforwardly to more generic dispersion relations so long as it is a strictly increasing function in the covariant derivative $\omega \sim \epsilon(-i\partial_x + A_1)$. By inverse function theorem, the integrand and the integral domain could be mapped bijectively to the case of linear dispersion. Thus, we have shown the chiral anomaly in (1+1) dimensions is not sensitive to the exact shape of the dispersion. It is sufficient to have a strictly monotonic energy dispersion that is odd under chiral transformation in the fermion Lagrangian density in order to produce the chiral anomaly (3). The leading contribution has the universal magnitude $E/\pi$ and an extra minus sign could emerge if the dispersion is not strictly increasing in momentum. Note that in the case where $\epsilon$ is strictly decreasing, the notion of the left-moving and right-moving are swapped.

## IV. CONCLUSION

We showed that in (1+1) dimensions a model requires no *anomalous* new term in its kinetic equation in order to have chiral anomaly in contrast to (3+1) dimensions where the Berry phase associated with each Weyl fermion is essential. Instead, the asymptotic behaviors of the dispersion relations of the chiral fermions fully determine the value of the momentum integral. Exploiting this observation, on one hand, we delivered an extremely simple derivation for the vintage exercise of chiral anomaly in (1+1) dimensions, circumventing the subtle infinities of the ground state. On the other hand, we demonstrated that a broader class of models could exhibit the same chiral anomaly. It is physically interesting to investigate the generic principles and forms of anomalies for generalizations of (1+1) dimensional fermions with higher order derivative couplings including those considered in Refs. [23, 24], in which kinetic terms with higher order derivatives are considered with Lorentz symmetry preserved. Mathematically, these models might be related to or motivate other index theorems beyond the validity of the one by Atiyah and Singer [6, 7].

Finally, It might be worth noting that we deliberately kept the electromagnetic field classical and turned off short-range interaction for fermions mainly for delivering the result with minimal amount of components. It could be an intriguing follow-up exercise to understand the robustness of the main results here against interactions. Naively some key arguments fail. For example, when turning on a four-fermion term, the theory becomes a Luttinger liquid, making the step of putting Green's function on-shell questionable [22], and thus there is no semiclassical Boltzmann equation to start with. Nonetheless, borrowing the hindsights from bosonization [16], a contact interaction should merely renormalize the velocity of the dual boson field rather than its mass, which is dual to the long ranged Coulomb interaction and determines the amount of chiral anomaly in that formulation. If chiral anomaly survives and has the same value in this regard, it should seek an alternative route out to manifest itself in the classical limit.

## ACKNOWLEDGMENTS

We thank Umang Mehta for discussions and professor Yeong-Chuan Kao for comments on the primitive ideas underlying this manuscript and bringing the work Ref.[20] to our attention. CHW is supported, in part, by ARO (W911NF-18-1-0020, W911NF-18-1-0212), AFOSR MURI (FA9550-19-1-0399) and AFRL (FA8649-21-P-0781).



[1] S. L. Adler, Axial-vector vertex in spinor electrodynamics, Phys. Rev. **177**, 2426 (1969).

[2] J. S. Bell and R. Jackiw, A pcac puzzle: $\pi^0 \to \gamma\gamma$ in the $\sigma$-model, Il Nuovo Cimento A (1965-1970) **60**, 47 (1969).

[3] K. Fujikawa, Path-integral measure for gauge-invariant fermion theories, Phys. Rev. Lett. **42**, 1195 (1979).

[4] K. Fujikawa, Path integral for gauge theories with fermions, Phys. Rev. D **21**, 2848 (1980).

[5] K. Fujikawa and H. Suzuki, *Path Integrals and Quantum Anomalies* (Oxford University Press, Oxford, 2004).

[6] M. F. Atiyah and I. M. Singer, The index of elliptic operators: I, Annals of Mathematics **87**, 484 (1968).

[7] M. Atiyah, R. Bott, and V. Patodi, On the heat equation and the index theorem., Inventiones mathematicae **19**, 279 (1973).

[8] D. T. Son and N. Yamamoto, Berry curvature, triangle anomalies, and the chiral magnetic effect in fermi liquids, Phys. Rev. Lett. **109**, 181602 (2012).

[9] D. T. Son and N. Yamamoto, Kinetic theory with berry curvature from quantum field theories, Phys. Rev. D **87**, 085016 (2013).

[10] J.-W. Chen, S. Pu, Q. Wang, and X.-N. Wang, Berry curvature and four-dimensional monopoles in the relativistic chiral kinetic equation, Phys. Rev. Lett. **110**, 262301 (2013).

[11] D. T. Son and B. Z. Spivak, Chiral anomaly and classical negative magnetoresistance of weyl metals, Phys. Rev. B **88**, 104412 (2013).

[12] N. P. Armitage, E. J. Mele, and A. Vishwanath, Weyl and dirac semimetals in three-dimensional solids, Rev. Mod. Phys. **90**, 015001 (2018).

[13] J. Schwinger, Gauge invariance and mass. ii, Phys. Rev. **128**, 2425 (1962).

[14] R. A. Bertlmann, *Anomalies in Quantum Field Theory* (Oxford University Press, Oxford, 2000).

[15] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Westview, Boulder, CO, 1995).

[16] N. Manton, The schwinger model and its axial anomaly, Annals of Physics **159**, 220 (1985).

[17] R. Jackiw, Effects of Dirac's Negative Energy Sea on Quantum Numbers, arXiv e-prints , hep-th/9903255 (1999), arXiv:hep-th/9903255 [hep-th].

[18] E. Lifshitz and L. Pitaevskii, *Physical Kinetics: Volume 10*, Course of theoretical physics (Elsevier Science, 1995).

[19] C. L. Kane and M. P. A. Fisher, Quantized thermal transport in the fractional quantum hall effect, Phys. Rev. B **55**, 15832 (1997).

[20] K. Fujikawa, Characteristics of chiral anomaly in view of various applications, Phys. Rev. D **97**, 016018 (2018).

[21] G. D. Mahan, *Many Particle Physics, Third Edition* (Plenum, New York, 2000).

[22] A. Kamenev, *Field Theory of Non-Equilibrium Systems* (Cambridge University Press, 2011).

[23] J. Barcelos-Neto and C. P. Natividade, Schwinger model with higher derivative couplings, Zeitschrift fÃŒr Physik C Particles and Fields **49**, 511 (1991).

[24] R. L. P. G. Amaral, L. V. Belvedere, N. A. Lemos, and C. P. Natividade, Higher-derivative schwinger model, Phys. Rev. D **47**, 3443 (1993).